

Exotic rational elliptic surfaces without 1–handles

KOICHI YASUI

Harer, Kas and Kirby have conjectured that every handle decomposition of the elliptic surface $E(1)_{2,3}$ requires both 1– and 3–handles. In this article, we construct a smooth 4–manifold which has the same Seiberg–Witten invariant as $E(1)_{2,3}$ and admits neither 1– nor 3–handles by using rational blow-downs and Kirby calculus. Our manifold gives the first example of either a counterexample to the Harer–Kas–Kirby conjecture or a homeomorphic but nondiffeomorphic pair of simply connected closed smooth 4–manifolds with the same nonvanishing Seiberg–Witten invariants.

57R55; 57R65, 57R57, 57N13

1 Introduction

It is a basic problem in 4–dimensional topology to classify smooth structures on 4–manifolds. Constructions of exotic smooth structures on 4–manifolds with small Euler characteristics are currently in rapid progress (see, for example, Park [14], Stipsicz–Szabó [17], Fintushel–Stern [5], Park–Stipsicz–Szabó [15] and Akhmedov–Park [1]). However, it is still unknown whether or not \mathbf{S}^4 and \mathbf{CP}^2 admit an exotic smooth structure. If such a structure exists, then each handle decomposition of it has at least either a 1– or 3–handle (see Proposition 6.4). On the contrary, many classical simply connected closed smooth 4–manifolds are known to admit neither 1– nor 3–handles in their handle decompositions (cf Gompf–Stipsicz [7]). Problem 4.18 in Kirby’s problem list [11] is the following: “Does every simply connected, closed 4–manifold have a handlebody decomposition without 1–handles? Without 1– and 3–handles?” The elliptic surfaces $E(n)_{p,q}$ are candidates of counterexamples to Problem 4.18. It is not known whether or not the simply connected closed smooth 4–manifold $E(n)_{p,q}$ (n arbitrary, $p, q \geq 2$, $\gcd(p, q) = 1$) admits a handle decomposition without 1–handles (cf Gompf [6] and Gompf–Stipsicz [7]). In particular, Harer, Kas and Kirby have conjectured in [9] that every handle decomposition of $E(1)_{2,3}$ requires at least a 1–handle. Note that by considering dual handle decompositions, their conjecture is equivalent to the assertion that $E(1)_{2,3}$ requires both 1– and 3–handles.

In this article we construct the following smooth 4–manifolds by using rational blow-downs and Kirby calculus.

Theorem 1.1 (1) For $q = 3, 5$, there exists a smooth 4–manifold E_q with the following properties:

- (a) E_q is homeomorphic to $E(1)_{2,q}$;
- (b) E_q has the same Seiberg–Witten invariant as $E(1)_{2,q}$;
- (c) E_q admits a handle decomposition without 1–handles, namely,

$$E_q = \text{one } 0\text{-handle} \cup \text{twelve } 2\text{-handles} \cup \text{two } 3\text{-handles} \cup \text{one } 4\text{-handle}.$$

(2) There exists a smooth 4–manifold E'_3 with the following properties:

- (a) E'_3 is homeomorphic to $E(1)_{2,3}$;
- (b) E'_3 has the same Seiberg–Witten invariant as $E(1)_{2,3}$;
- (c) E'_3 admits a handle decomposition without 1– and 3–handles, namely,

$$E'_3 = \text{one } 0\text{-handle} \cup \text{ten } 2\text{-handles} \cup \text{one } 4\text{-handle}.$$

As far as the author knows, E_q and E'_3 are the first examples in the following sense: If E_q (resp. E'_3) is diffeomorphic to $E(1)_{2,q}$ (resp. $E(1)_{2,3}$), then the above handle decomposition of $E(1)_{2,q}$ ($= E_q$ (resp. E'_3)) is the first example which has no 1–handles. Otherwise, ie, if E_q (resp. E'_3) is not diffeomorphic to $E(1)_{2,q}$ (resp. $E(1)_{2,3}$), then E_q (resp. E'_3) and $E(1)_{2,q}$ (resp. $E(1)_{2,3}$) are the first homeomorphic but nondiffeomorphic examples which are simply connected closed smooth 4–manifolds with the same nonvanishing Seiberg–Witten invariants.

An affirmative solution to the Harer–Kas–Kirby conjecture implies that both E_3 and E'_3 are not diffeomorphic to $E(1)_{2,3}$, though these three have the same Seiberg–Witten invariants. In this case, the minimal number of 1–handles in handle decompositions does detect the difference of their smooth structures.

Our construction is inspired by rational blow-down constructions of exotic smooth structures on $\mathbf{CP}^2 \# n \overline{\mathbf{CP}}^2$ ($5 \leq n \leq 8$) by Park [14], Stipsicz–Szabó [17], Fintushel–Stern [5] and Park–Stipsicz–Szabó [15]. Our method is different from theirs since, firstly, we use Kirby calculus to perform rational blow-downs, whereas they used elliptic fibrations on $E(1)$ (and knot surgeries), and secondly, they did not examine handle decompositions.

Acknowledgements The author wishes to express his deeply gratitude to his adviser, Professor Hisaaki Endo, for encouragement and many useful suggestions. He would like to thank Professors Selman Akbulut, Kazunori Kikuchi, Ronald J Stern and Yuichi Yamada for helpful comments and discussions. Kikuchi’s theorem [10, Theorem 4] partially gave him the idea of the construction. Yamada gave him interesting questions (cf Remark 6.2). The author is partially supported by JSPS Research Fellowships for Young Scientists.

2 Rational blow-down

In this section we review the rational blow-down introduced by Fintushel–Stern [4]. For details, see also Gompf–Stipsicz [7].

Let C_p and B_p be the smooth 4–manifolds defined by Kirby diagrams in Figure 1, and u_1, \dots, u_{p-1} elements of $H_2(C_p; \mathbf{Z})$ given by corresponding 2–handles in the figure such that $u_i \cdot u_{i+1} = +1$ ($1 \leq i \leq p-2$). The boundary ∂C_p of C_p is diffeomorphic




Figure 1

to the lens space $L(p^2, 1-p)$ and to the boundary ∂B_p of B_p . The following lemma is well known.

- Lemma 2.1** (1) $\pi_1(C_p) = 0$, $\pi_1(B_p) = \mathbf{Z}_p$ and $\pi_1(L(p^2, 1-p)) = \mathbf{Z}_{p^2}$.
 (2) $H_2(C_p; \mathbf{Z}) = \oplus_{p-1} \mathbf{Z}$ and $H_2(B_p; \mathbf{Z}) = H_2(L(p^2, 1-p); \mathbf{Z}) = 0$.

Suppose that C_p embeds in a smooth 4–manifold X . The smooth 4–manifold $X_{(p)} := (X - \text{int } C_p) \cup_{L(p^2, 1-p)} B_p$ is called the rational blow-down of X along C_p . Note that $X_{(p)}$ is uniquely determined up to diffeomorphism by a fixed pair (X, C_p) . This operation preserves b_2^+ , decreases b_2^- , may create torsions in the first homology group, and has the following relation with the logarithmic transformation.

Theorem 2.2 (Fintushel–Stern [4], cf Gompf–Stipsicz [7]) *Suppose that a smooth 4–manifold X contains a cusp neighborhood, that is, a 0–handle with a 2–handle attached along a 0–framed right trefoil knot. Let X_p be the smooth 4–manifold obtained from X by performing a logarithmic transformation of multiplicity p in the cusp neighborhood. Then there exists a copy of C_p in $X \# (p-1)\overline{\mathbf{C}\mathbf{P}}^2$ such that the rational blow-down of $X \# (p-1)\overline{\mathbf{C}\mathbf{P}}^2$ along the copy of C_p is diffeomorphic to X_p .*

Let $E(n)$ be the simply connected elliptic surface with Euler characteristic $12n$ and with no multiple fibers, and $E(n)_{p_1, \dots, p_k}$ the elliptic surface obtained from

$E(n)$ by performing logarithmic transformations of multiplicities p_1, \dots, p_k . We denote h, e_1, e_2, \dots, e_n as a canonical orthogonal basis of $H_2(\mathbf{CP}^2 \# n \overline{\mathbf{CP}}^2; \mathbf{Z}) = H_2(\mathbf{CP}^2; \mathbf{Z}) \oplus_n H_2(\overline{\mathbf{CP}}^2; \mathbf{Z})$ such that $h^2 = 1$ and $e_1^2 = e_2^2 = \dots = e_n^2 = -1$.

Since there is a diffeomorphism $E(1)_p \rightarrow E(1) = \mathbf{CP}^2 \# 9 \overline{\mathbf{CP}}^2$ which maps the class of a regular fiber of $E(1)_p$ to $p(3h - e_1 - e_2 - \dots - e_9) \in H_2(\mathbf{CP}^2 \# 9 \overline{\mathbf{CP}}^2; \mathbf{Z})$ (cf Etgü-Park [2, page 680] and Gompf-Stipsicz [7]), Theorem 2.2 gives us the following corollary.

Corollary 2.3 *For each natural number p and q , the elliptic surface $E(1)_{p,q}$ is obtained from $\mathbf{CP}^2 \# (8 + q) \overline{\mathbf{CP}}^2$ by rationally blowing down along a certain copy pC_q of C_q such that u_1, \dots, u_{q-1} satisfy*

$$\begin{aligned} u_1 &= e_{7+q} - e_{8+q}, \quad u_2 = e_{6+q} - e_{7+q}, \quad \dots, \quad u_{q-2} = e_{10} - e_{11}, \\ u_{q-1} &= p(3h - e_1 - e_2 - \dots - e_9) - 2e_{10} - e_{11} - e_{12} - \dots - e_{8+q} \end{aligned}$$

as elements of $H_2(\mathbf{CP}^2 \# (8 + q) \overline{\mathbf{CP}}^2; \mathbf{Z})$.

Remark 2.4 $E(1)_{p,q}$ is homeomorphic but nondiffeomorphic to $E(1)$, in the case $p, q \geq 2$ and $\gcd(p, q) = 1$ (cf Gompf-Stipsicz [7]).

3 Construction

In this section we construct E_3, E_5 and E'_3 , and prove Theorem 1.1 (1)(a)(c) and (2)(a)(c). In Kirby diagrams, we write the second homology classes given by 2–handles, instead of usual framings. Note that the square of the homology class given by a 2–handle is equal to the usual framing. We do not draw (whole) Kirby diagrams of E_3, E_5, E'_3 and the other manifolds appeared in the following construction. However, one can easily draw whole diagrams.

We begin with a construction of a cusp neighborhood in $\mathbf{CP}^2 \# 9 \overline{\mathbf{CP}}^2$ such that its embedding into $\mathbf{CP}^2 \# 9 \overline{\mathbf{CP}}^2$ has the same homological properties as that of the regular neighborhood of a cusp fiber of $E(1)_2$. We do not know if these embeddings into $\mathbf{CP}^2 \# 9 \overline{\mathbf{CP}}^2$ are the same up to diffeomorphism.

Lemma 3.1 $\mathbf{CP}^2 \# 9 \overline{\mathbf{CP}}^2$ admits the handle decomposition drawn in Figure 2. Here f denotes $6h - 2e_1 - 2e_2 - \dots - 2e_9 \in H_2(\mathbf{CP}^2 \# 9 \overline{\mathbf{CP}}^2; \mathbf{Z})$.

Proof We first create two 2–handles with framings $2h$ and $4h$ in a Kirby diagram of \mathbf{CP}^2 . Figure 8 is a basic Kirby diagram of \mathbf{CP}^2 . Introducing a 2–handle/3–handle




Figure 2: $\mathbf{CP}^2\#9\overline{\mathbf{CP}}^2$

pair gives Figure 9. Handle slides and isotopies yield Figure 12 (Pairs of bold lines in figures denote “bands”):

$$\text{Figure 9} \xrightarrow{0+h} \text{Figure 10} \xrightarrow{h+h} \text{Figure 11} \xrightarrow{\text{isotopy}} \text{Figure 12}.$$

Creating a 2–handle/3–handle pair gives Figure 13. Handle slides produce Figure 17:

$$\text{Figure 13} \xrightarrow{0+h} \text{Figure 14} \xrightarrow{h+h} \text{Figure 15} \xrightarrow{2h+h} \text{Figure 16} \xrightarrow{3h+h} \text{Figure 17}.$$

We blow up \mathbf{CP}^2 nine times:

$$\text{Figure 17} \xrightarrow{\text{three blow-ups}} \text{Figure 18} \xrightarrow{\text{isotopy}} \text{Figure 19} \xrightarrow{\text{six blow-ups}} \text{Figure 20}.$$

We make a handle addition $(4h - 2e_1 - 2e_2 - 2e_3 - e_4 - e_5 - \dots - e_9) + (2h - e_4 - e_5 - \dots - e_9)$. This leads to Figure 21, and an isotopy gives Figure 2. \square

Proposition 3.2 (1) $\mathbf{CP}^2\#11\overline{\mathbf{CP}}^2$ admits the handle decomposition drawn in Figure 3. In particular $\mathbf{CP}^2\#11\overline{\mathbf{CP}}^2$ contains the copy of C_3 drawn in the figure. The elements $u_1, u_2 \in H_2(\mathbf{CP}^2\#11\overline{\mathbf{CP}}^2; \mathbf{Z})$ given by this copy of C_3 are the same as that given by ${}_2C_3$.

(2) $\mathbf{CP}^2\#13\overline{\mathbf{CP}}^2$ admits the handle decomposition drawn in Figure 4. In particular $\mathbf{CP}^2\#13\overline{\mathbf{CP}}^2$ contains the copy of C_5 drawn in the figure. The elements $u_1, \dots, u_4 \in H_2(\mathbf{CP}^2\#13\overline{\mathbf{CP}}^2; \mathbf{Z})$ given by this copy of C_5 are the same as that given by ${}_2C_5$.

Proof (1) Blowing up in Figure 2 yields Figure 22. The handle slide drawn in Figure 23 gives Figure 24. An additional blow-up yields Figure 25, and an isotopy gives Figure 3.




Figure 3: $\mathbb{C}P^2 \# 11\overline{\mathbb{C}P}^2$





Figure 4: $\mathbb{C}P^2 \# 13\overline{\mathbb{C}P}^2$

(2) Handle slides, isotopies and blow-ups in Figure 25 yield Figure 4:



Proposition 3.3 $\mathbb{C}P^2 \# 13\overline{\mathbb{C}P}^2$ admits the handle decomposition drawn in Figure 5. In particular $\mathbb{C}P^2 \# 13\overline{\mathbb{C}P}^2$ contains the copy of C_5 drawn in the figure.

Proof Recall the construction in the proof of Lemma 3.1. In this construction, we created a 2–handle/3–handle pair twice. Instead of introducing a 2–handle/3–handle




Figure 5: $\mathbf{CP}^2 \# 13\overline{\mathbf{CP}}^2$

pair twice, blowing up twice yields Figure 38:

$$\begin{array}{ccccccc}
 \text{Figure 8} & \xrightarrow{\text{blow-up}} & \text{Figure 33} & \xrightarrow{e_1 + h} & \text{Figure 34} & \xrightarrow{(h + e_1) + h} & \text{Figure 35} \\
 & & & \xrightarrow{\text{isotopy}} & \text{Figure 36} & \xrightarrow{\text{blow-up}} & \text{Figure 37} \xrightarrow{e_2 + h} \text{Figure 38.}
 \end{array}$$

Handle slides and blow-ups as in proofs of Lemma 3.1 and Proposition 3.2 gives Figure 39. Repeating handle slides drawn in Figure 23 yields Figure 40. An additional blow-up gives Figure 41, and an isotopy gives Figure 5. \square

Definition 3.4 Let E_q be the smooth 4–manifold obtained from $\mathbf{CP}^2 \# (8 + q)\overline{\mathbf{CP}}^2$ by rationally blowing down along the copy of C_q obtained in Proposition 3.2, for $q = 3, 5$. Let E'_3 be the smooth 4–manifold obtained from $\mathbf{CP}^2 \# 13\overline{\mathbf{CP}}^2$ by rationally blowing down along the copy of C_5 obtained in Proposition 3.3.

Remark 3.5 It is not known whether or not there exists a copy of C_5 in $\mathbf{CP}^2 \# 13\overline{\mathbf{CP}}^2$ such that the rational blow-down is diffeomorphic to $E(1)_{2,3}$.

In [20] we will construct more examples of exotic $\mathbf{CP}^2 \# 9\overline{\mathbf{CP}}^2$ without 1– and 3–handles, by improving the construction of E'_3 . The author does not know if these examples have the same Seiberg–Witten invariants as the elliptic surfaces $E(1)_{p,q}$.

We prepare the following lemma.

Lemma 3.6 (cf Gompf–Stipsicz [7]) *Suppose that a simply connected closed smooth 4–manifold X has the handle decomposition drawn in Figure 6. Here n is an arbitrary integer, h_2 and h_3 are arbitrary natural numbers. Note that we write usual framings instead of homology classes in the figure.*

Let $X_{(p)}$ be the rational blow-down of X along the copy of C_p drawn in Figure 6. Then $X_{(p)}$ admits a handle decomposition

$$X_{(p)} = \text{one } 0\text{-handle} \cup (h_2 + 1) \text{ } 2\text{-handles} \cup h_3 \text{ } 3\text{-handles} \cup \text{one } 4\text{-handle.}$$

In particular $X_{(p)}$ admits a handle decomposition without 1–handles.




Figure 6: Handle decomposition of X

Proof Draw a Kirby diagram of $X_{(p)}$, following the procedure introduced in [7, Section 8.5] (see also [7, page 516, Solution of Exercise 8.5.1.(a)]). Then the n –framed unknot drawn in Figure 6 changes into a meridian of a unique dotted circle which naturally appears in this procedure. Thus we can cancel the 1–handle/2–handle pair. Note that this procedure does not produce new 3–handles. \square

The following proposition gives Theorem 1.1 (1)(a)(c) and (2)(a)(c).

Proposition 3.7 For $q = 3, 5$, the manifold E_q is homeomorphic to $E(1)_{2,q}$ and admits a handle decomposition without 1–handles, namely,

$$E_q = \text{one } 0\text{-handle} \cup \text{twelve } 2\text{-handles} \cup \text{two } 3\text{-handles} \cup \text{one } 4\text{-handle.}$$

E'_3 is homeomorphic to $E(1)_{2,3}$ and admits a handle decomposition without 1– and 3–handles, namely,

$$E'_3 = \text{one } 0\text{-handle} \cup \text{ten } 2\text{-handles} \cup \text{one } 4\text{-handle.}$$

Proof Lemma 3.6 shows the above properties of E_q and E'_3 about handle decompositions. Thus E_q and E'_3 are simply connected. Since E_q is obtained from $\mathbb{C}\mathbb{P}^2 \# (8 + q)\overline{\mathbb{C}\mathbb{P}^2}$ by rationally blowing down along a copy of C_q , we have

$$b_2^+(E_q) = b_2^+(\mathbb{C}\mathbb{P}^2 \# (8 + q)\overline{\mathbb{C}\mathbb{P}^2}) = 1,$$

$$b_2^-(E_q) = b_2^-(\mathbb{C}\mathbb{P}^2 \# (8 + q)\overline{\mathbb{C}\mathbb{P}^2}) - b_2^-(C_q) = (8 + q) - (q - 1) = 9.$$

Similarly we have $b_2^+(E'_3) = 1$ and $b_2^-(E'_3) = 9$. Therefore Freedman’s theorem together with Rochlin’s theorem shows that E_q and E'_3 are homeomorphic to $\mathbb{C}\mathbb{P}^2 \# 9\overline{\mathbb{C}\mathbb{P}^2}$. Thus E_q and E'_3 are homeomorphic to $E(1)_{2,q}$. \square

4 Seiberg–Witten invariants

In this section, we briefly review facts about the Seiberg–Witten invariants with $b_2^+ = 1$. For details and examples of computations, see Fintushel–Stern [3; 4], [5], Stern [16], Park [13; 14], Ozsváth–Szabó [12], Stipsicz–Szabó [17] and Park–Stipsicz–Szabó [15].

Suppose that X is a simply connected closed smooth 4–manifold with $b_2^+(X) = 1$. Let $\mathcal{C}(X)$ be the set of characteristic elements of $H^2(X; \mathbf{Z})$. Fix a homology orientation on X , that is, orient $H_+^2(X; \mathbf{R}) := \{H \in H^2(X; \mathbf{Z}) \mid H^2 > 0\}$. Then the (small-perturbation) Seiberg–Witten invariant $SW_{X,H}(K) \in \mathbf{Z}$ is defined for every positively oriented element $H \in H_+^2(X; \mathbf{R})$ and every element $K \in \mathcal{C}(X)$ such that $K \cdot H \neq 0$. Let $e(X)$ and $\sigma(X)$ be the Euler characteristic and the signature of X , respectively, and $d_X(K)$ the even integer defined by $d_X(K) = \frac{1}{4}(K^2 - 2e(X) - 3\sigma(X))$ for $K \in \mathcal{C}(X)$. It is known that if $SW_{X,H}(K) \neq 0$ for some $H \in H_+^2(X; \mathbf{R})$, then $d_X(K) \geq 0$. The wall-crossing formula tells us the dependence of $SW_{X,H}(K)$ on H : if $H, H' \in H_+^2(X; \mathbf{R})$ and $K \in \mathcal{C}(X)$ satisfy $H \cdot H' > 0$ and $d_X(K) \geq 0$, then

$$SW_{X,H'}(K) = SW_{X,H}(K) + \begin{cases} 0 & \text{if } K \cdot H \text{ and } K \cdot H' \text{ have the same sign,} \\ (-1)^{\frac{1}{2}d_X(K)} & \text{if } K \cdot H > 0 \text{ and } K \cdot H' < 0, \\ (-1)^{1+\frac{1}{2}d_X(K)} & \text{if } K \cdot H < 0 \text{ and } K \cdot H' > 0. \end{cases}$$

Note that these facts imply that $SW_{X,H}(K)$ is independent of H in the case $b_2^-(X) \leq 9$, in other words, the Seiberg–Witten invariant $SW_X: \mathcal{C}(X) \rightarrow \mathbf{Z}$ is well defined.

We recall the change of the Seiberg–Witten invariants by rationally blowing down. Assume that X contains a copy of C_p . Let $X_{(p)}$ be the rational blow-down of X along the copy of C_p . Suppose that $X_{(p)}$ is simply connected. The following theorems are known.

Proposition 4.1 (Fintushel–Stern [4]) *For every element $K \in \mathcal{C}(X_{(p)})$, there exists an element $\tilde{K} \in \mathcal{C}(X)$ such that $K|_{X_{(p)}-\text{int } B_p} = \tilde{K}|_{X-\text{int } C_p}$ and $d_{X_{(p)}}(K) = d_X(\tilde{K})$. We call such an element $\tilde{K} \in \mathcal{C}(X)$ a lift of K .*

Theorem 4.2 (Fintushel–Stern [4]) *If an element $\tilde{K} \in \mathcal{C}(X)$ is a lift of some element $K \in \mathcal{C}(X_{(p)})$, then $SW_{X_{(p)}, H}(K) = SW_{X, H}(\tilde{K})$ for every positively oriented element $H \in H_+^2(X; \mathbf{R})$ which is orthogonal to the subspace $H_2(C_p; \mathbf{R})$ of $H_2(X; \mathbf{R})$. Note that we view H as a positively oriented element of $H_+^2(X_{(p)}; \mathbf{R})$.*

Theorem 4.3 (Fintushel–Stern [4], cf Park [13]) *If an element $\tilde{K} \in \mathcal{C}(X)$ satisfies that $(\tilde{K}|_{C_p})^2 = 1 - p$ and $\tilde{K}|_{\partial C_p} = mp \in \mathbf{Z}_{p^2} \cong H^2(\partial C_p; \mathbf{Z})$ with $m \equiv p - 1 \pmod{2}$, then there exists an element $K \in \mathcal{C}(X_{(p)})$ such that \tilde{K} is a lift of K .*

Corollary 4.4 *If an element $\tilde{K} \in \mathcal{C}(X)$ satisfies $\tilde{K}(u_1) = \cdots = \tilde{K}(u_{p-2}) = 0$ and $\tilde{K}(u_{p-1}) = \pm p$, then \tilde{K} is a lift of some element $K \in \mathcal{C}(X_{(p)})$.*

5 Computations of SW invariants

In this section we complete the proof of Theorem 1.1. We prepare the following lemma here.

Lemma 5.1 *Let X be a simply connected closed smooth 4–manifold which contains a copy of C_p , and ι the inclusion $X - \text{int } C_p \hookrightarrow X$. Let C_p^\perp be the orthogonal complement of the subspace spanned by $u_1, \dots, u_{p-1} \in H_2(X; \mathbf{Z})$, that is,*

$$C_p^\perp := \{v \in H_2(X; \mathbf{Z}) \mid v \cdot u_1 = \cdots = v \cdot u_{p-1} = 0\}.$$

Suppose that there exists an element $\delta \in H_2(X; \mathbf{Z})$ such that $\delta \cdot u_1 = 1$ and $\delta \cdot u_2 = \delta \cdot u_3 = \cdots = \delta \cdot u_{p-1} = 0$. Then

- (1) $\iota_* H_2(X - \text{int } C_p; \mathbf{Z}) = C_p^\perp$;
- (2) $H_1(X - \text{int } C_p; \mathbf{Z}) = 0$.

Proof (1) Since every element of $H_2(X - \text{int } C_p; \mathbf{Z})$ is represented by a surface, it is clear that $\iota_* H_2(X - \text{int } C_p; \mathbf{Z}) \subset C_p^\perp$.

Let ι' be the inclusion $C_p \hookrightarrow X$. Mayer–Vietoris exact sequence of $(X - \text{int } C_p) \cup C_p = X$ is as follows:

$$0 \rightarrow H_2(X - \text{int } C_p; \mathbf{Z}) \oplus H_2(C_p; \mathbf{Z}) \xrightarrow{\iota_* + \iota'_*} H_2(X; \mathbf{Z}) \xrightarrow{\partial} \mathbf{Z}_{p^2}.$$

Since C_p is negative definite and $\text{Im } \iota_* \subset C_p^\perp$, we have $\text{Im } (\iota_* + \iota'_*) = \text{Im } \iota_* \oplus \text{Im } \iota'_*$.

We determine $\partial(\delta)$ here. There clearly exists an element $n \in \mathbf{Z}$ such that $\partial(n\delta) \equiv 0 \pmod{p^2}$. The above exact sequence ensures the existence of elements $u \in \text{Im } \iota'_*$ and

$v \in C_p^\perp$ such that $n\delta = u + v$. The element u satisfies $u \cdot u_1 = n$ ($= n\delta \cdot u_1$) and $u \cdot u_2 = u \cdot u_3 = \cdots = u \cdot u_{p-1} = 0$ ($= n\delta \cdot u_2$). Since u_1, u_2, \dots, u_n is a basis of $\text{Im } \iota'_*$, we can easily see $n \equiv 0 \pmod{p^2}$ by using the intersection form of C_p . Hence $\partial(\delta)$ is a generator of \mathbf{Z}_{p^2} .

Suppose that some element $w \in C_p^\perp$ satisfies $\partial(w) \not\equiv 0 \pmod{p^2}$. Since $\partial(\delta)$ is a generator of \mathbf{Z}_{p^2} , there exists an element $n' \in \mathbf{Z}$ with $n' \not\equiv 0 \pmod{p^2}$ such that $\partial(n'\delta + w) \equiv 0$. Applying the above argument about $n\delta$ to $n'\delta + w$, we get $n' \equiv 0 \pmod{p^2}$. This is a contradiction. Thus we obtain $\partial(C_p^\perp) = 0$. Therefore $C_p^\perp \subset \text{Ker } \partial = \text{Im } \iota_* \oplus \text{Im } \iota'_* \subset C_p^\perp \oplus \text{Im } \iota'_*$. Thus it is easy to see $C_p^\perp \subset \iota_* H_2(X - \text{int } C_p; \mathbf{Z})$.

(2) Since the above ∂ is onto, we can easily show by using Mayer–Vietoris exact sequence. □

Remark 5.2 (1) Since $\iota_*: H_2(X - \text{int } C_p; \mathbf{Z}) \rightarrow H_2(X; \mathbf{Z})$ is injective, the above lemma allows us to identify $H_2(X - \text{int } C_p; \mathbf{Z})$ with C_p^\perp .

(2) Under the same assumption as that in Lemma 5.1, we can also show $H_1(X_{(p)}; \mathbf{Z}) = 0$. Here $X_{(p)}$ denotes the rational blow-down of X along the copy of C_p . It is not known whether or not the fundamental groups of $X - \text{int } C_p$ and $X_{(p)}$ vanish.

The following proposition gives us Theorem 1.1 (1)(b). In the rest of this section, we denote the symbol R_n as $\mathbf{CP}^2 \# n\overline{\mathbf{CP}}^2$.

Proposition 5.3 E_q has the same Seiberg–Witten invariant as $E(1)_{2,q}$, that is, there exists a homeomorphism between E_q and $E(1)_{2,q}$ which preserves the orientations, the homology orientations and the Seiberg–Witten invariants, for $q = 3, 5$.

Proof We first give a proof for $q = 3$. Let $\alpha_1, \alpha_2, \dots, \alpha_9, \beta \in {}_2C_3^\perp$ be the elements defined by

$$\begin{aligned} \alpha_1 &= 4h - e_1 - e_2 - \cdots - e_9 - 2e_{10} - 2e_{11}, \\ \alpha_i &= 5h - 2e_1 - 2e_2 - e_3 - e_4 - \cdots - e_9 - 2e_{10} - 2e_{11} - e_{i+1} \quad (2 \leq i \leq 8), \\ \alpha_9 &= e_1 - e_2, \quad \beta = 30h - 13e_1 - 10e_2 - 7e_3 - 7e_4 - \cdots - 7e_9 - 12e_{10} - 12e_{11}. \end{aligned}$$

We can view $\alpha_1, \alpha_2, \dots, \alpha_9, \beta$ as elements of $H_2(E(1)_{2,3}; \mathbf{Z})$ by Lemma 5.1.(1), Corollary 2.3 and the following natural identification:

$$H_2(E(1)_{2,3} - \text{int } B_3; \mathbf{Z}) (= H_2(R_{11} - \text{int } C_3; \mathbf{Z})) \subset H_2(E(1)_{2,3}; \mathbf{Z}).$$

This identification preserves cup products. Therefore the elements $\alpha_1, \alpha_2, \dots, \alpha_9, \beta$ of $H_2(E(1)_{2,3}; \mathbf{Z})$ satisfy

$$\begin{aligned}\alpha_1^2 = \alpha_2^2 = \dots = \alpha_8^2 &= -1, \alpha_9^2 = -2, \alpha_i \cdot \alpha_j = 0 \quad (1 \leq i < j \leq 9), \\ \beta^2 = 0, \beta \cdot \alpha_1 = \beta \cdot \alpha_2 = \dots = \beta \cdot \alpha_8 &= 0, \beta \cdot \alpha_9 = 3.\end{aligned}$$

Recall that the intersection form of $E(1)_{2,3}$ is $\langle 1 \rangle \oplus 9\langle -1 \rangle$ (This notation of the intersection form is the same as that in Gompf–Stipsicz [7, Section 1.2]). This implies that either the matrix $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ or $\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ represents the symmetric bilinear form on $\langle \alpha_1, \alpha_2, \dots, \alpha_8 \rangle^\perp$. We here denote the symbol $\langle \alpha_1, \alpha_2, \dots, \alpha_8 \rangle^\perp$ as the orthogonal complement of the subspace spanned by $\alpha_1, \alpha_2, \dots, \alpha_8 \in H_2(E(1)_{2,3}; \mathbf{Z})$. Since α_9 and β are elements of $\langle \alpha_1, \alpha_2, \dots, \alpha_8 \rangle^\perp$, it is easy to check that the matrix $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ represents the symmetric bilinear form on $\langle \alpha_1, \alpha_2, \dots, \alpha_8 \rangle^\perp$. We can easily see that there exists an element $\alpha_{10} \in H_2(E(1)_{2,3}; \mathbf{Z})$ such that $3\alpha_{10} = \beta$, by using a basis of $\langle \alpha_1, \alpha_2, \dots, \alpha_8 \rangle^\perp$. Note that $\alpha_1, \alpha_2, \dots, \alpha_{10}$ is a basis of $H_2(E(1)_{2,3}; \mathbf{Z})$.

Proposition 3.2.(1) allows us to apply the above argument to E_3 . Thus we get a basis $\alpha'_1, \dots, \alpha'_{10}$ of $H_2(E_3; \mathbf{Z})$ which is corresponding to the basis $\alpha_1, \dots, \alpha_{10}$ of $H_2(E(1)_{2,3}; \mathbf{Z})$. Let $\varphi: H^2(E(1)_{2,3}; \mathbf{Z}) \rightarrow H^2(E_3; \mathbf{Z})$ be the isomorphism defined by $PD(\alpha_i) \mapsto PD(\alpha'_i)$ ($1 \leq i \leq 10$). Here PD denotes Poincaré dual. The isomorphism φ preserves the intersection forms and the homology orientations of $E(1)_{2,3}$ and E_3 .

Proposition 4.1 gives us a lift $\tilde{K} \in \mathcal{C}(R_{11})$ of K for every $K \in \mathcal{C}(E(1)_{2,3})$. Lemma 5.1.(2) together with the universal coefficient theorem implies that $\tilde{K}|_{R_{11} - \text{int } C_3}$ and $\varphi(K)|_{E_3 - \text{int } B_3}$ are uniquely determined by their values on $H_2(R_{11} - \text{int } C_3; \mathbf{Z}) = H_2(E_3 - \text{int } B_3; \mathbf{Z})$. Since $\alpha'_i = \alpha_i$ ($1 \leq i \leq 9$) and $3\alpha'_{10} = 3\alpha_{10}$ as elements of ${}_2C_3^\perp$, it is easy to check that \tilde{K} is also a lift of the element $\varphi(K) \in \mathcal{C}(E_3)$. Thus Theorem 4.2 shows $SW_{E_3}(\varphi(K)) = SW_{E(1)_{2,3}}(K)$. Hence the isomorphism φ preserves the Seiberg–Witten invariants of $E(1)_{2,3}$ and E_3 . Freedman's theorem gives us a required homeomorphism $\Phi: E_3 \rightarrow E(1)_{2,3}$ which preserves the orientations and satisfies $\Phi^* = \varphi$.

We briefly give a proof for $q = 5$. Let $\alpha_{5,1}, \alpha_{5,2}, \dots, \alpha_{5,9}, \beta_5 \in {}_2C_5^\perp$ be the elements defined by

$$\begin{aligned}\alpha_{5,i} &= 17h - 3e_1 - 4e_2 - \dots - 4e_9 - 6e_{10} - \dots - 6e_{13} - e_{i+1} \quad (1 \leq i \leq 8), \\ \alpha_{5,9} &= 96h - 19e_1 - 23e_2 - \dots - 23e_9 - 34e_{10} - \dots - 34e_{13}, \\ \beta_5 &= 537h - 104e_1 - 129e_2 - \dots - 129e_9 - 190e_{10} - \dots - 190e_{13}.\end{aligned}$$

Applying the above argument to E_5 , we obtain a proof. \square

To prove Theorem 1.1 (2)(b), we compute the Seiberg–Witten invariant of $E(1)_{2,3}$.

Lemma 5.4 *Let $K_3 \in \mathcal{C}(E(1)_{2,3})$ be the element defined by $K_3 = PD(\alpha_1 + \dots + \alpha_8 - 2\alpha_9 - 4\alpha_{10})$. Here $\alpha_1, \alpha_2, \dots, \alpha_{10}$ denote the elements of $H_2(E(1)_{2,3}; \mathbf{Z})$ defined in the proof of Proposition 5.3. Then K_3 satisfies $SW_{E(1)_{2,3}}(\pm K_3) = \pm 1$ and is the unique element of $\mathcal{C}(E(1)_{2,3})$ up to sign for which $SW_{E(1)_{2,3}}$ is nonzero.*

Proof Let $\tilde{K}_3 \in \mathcal{C}(R_{11})$ and $H \in H_+^2(R_{11}; \mathbf{R})$ be the elements defined by $\tilde{K}_3 = PD(3h - e_1 - e_2 - \dots - e_{11})$ and $H = PD(7h - 2e_1 - 2e_2 - \dots - 2e_{11})$. Note that H is orthogonal to the subspace $H_2(C_3; \mathbf{R})$ of $H_2(R_{11}; \mathbf{R})$. It is well known that $SW_{R_n, PD(h)}(\tilde{K}) = 0$ for every $\tilde{K} \in \mathcal{C}(R_n)$ and every $n \geq 0$. Applying the wall-crossing formula to $\pm \tilde{K}_3$, H and $PD(h)$, we have $SW_{R_{11}, H}(\pm \tilde{K}_3) = \pm 1$. Corollary 4.4 shows that \tilde{K}_3 is a lift of some element $K_3 \in \mathcal{C}(E(1)_{2,3})$. Thus Theorem 4.2 gives $SW_{E(1)_{2,3}}(\pm K_3) = \pm 1$.

Since \tilde{K}_3 is a lift of K_3 , the element K_3 satisfies $K_3(\alpha_i) = \tilde{K}_3(\alpha_i)$ ($1 \leq i \leq 9$) and $K_3(3\alpha_{10}) = \tilde{K}_3(3\alpha_{10})$. Hence the values of K_3 are as follows: $K_3(\alpha_1) = K_3(\alpha_2) = \dots = K_3(\alpha_8) = -1$, $K_3(\alpha_9) = 0$ and $K_3(\alpha_{10}) = -2$. Therefore we get $K_3 = PD(\alpha_1 + \dots + \alpha_8 - 2\alpha_9 - 4\alpha_{10})$.

Suppose that an element $L \in \mathcal{C}(E(1)_{2,3})$ satisfies $SW_{E(1)_{2,3}}(L) \neq 0$. Proposition 4.1 ensures the existence of a lift $\tilde{L} \in \mathcal{C}(R_{11})$ of L such that $SW_{R_{11}, H}(\tilde{L}) \neq 0$. We put $a := \tilde{L}(h)$. Since L is characteristic and $d_{R_{11}}(\tilde{L}) \geq 0$, the integer a is odd and $|a| \geq 3$. In the case $a \geq 3$, Cauchy–Schwartz inequality $((x_1 y_1 + \dots + x_n y_n)^2 \leq (x_1^2 + \dots + x_n^2)(y_1^2 + \dots + y_n^2))$ for $x_1, \dots, x_n, y_1, \dots, y_n \in \mathbf{R}$ and $d_{R_{11}}(\tilde{L}) = \frac{1}{4}(a^2 - ((\tilde{L}(e_1))^2 + (\tilde{L}(e_2))^2 + \dots + (\tilde{L}(e_{11}))^2) + 2) \geq 0$ show

$$\begin{aligned} \tilde{L} \cdot H &= 7a - 2\tilde{L}(e_1) - 2\tilde{L}(e_2) - \dots - 2\tilde{L}(e_{11}) \\ &\geq 7a - \sqrt{2^2 + 2^2 + \dots + 2^2} \sqrt{(\tilde{L}(e_1))^2 + (\tilde{L}(e_2))^2 + \dots + (\tilde{L}(e_{11}))^2} \\ &\geq 7a - 2\sqrt{11} \sqrt{a^2 + 2}. \end{aligned}$$

Since $SW_{E(1)_{2,3}}(L) \neq 0$ and $a \geq 3$, the wall-crossing formula shows $\tilde{L} \cdot H < 0$. Therefore we get $a = 3$. This together with $\tilde{L} \cdot H < 0$ shows $\tilde{L}(e_i) = 1$ ($1 \leq i \leq 11$). We thus have $\tilde{L} = \tilde{K}_3$. Similarly we have $\tilde{L} = -\tilde{K}_3$ in the case $a \leq -3$. Hence $L = \pm K_3$. □

The following proposition completes the proof of Theorem 1.1.

Proposition 5.5 *E'_3 has the same Seiberg–Witten invariant as $E(1)_{2,3}$, that is, there exists a homeomorphism between E'_3 and $E(1)_{2,3}$ which preserves the orientations, the homology orientations and the Seiberg–Witten invariants.*

Proof Let $\tilde{K}'_3 \in \mathcal{C}(R_{13})$ and $H' \in H^2_+(R_{13}; \mathbf{R})$ be the elements defined by $\tilde{K}'_3 = PD(3h + e_1 + e_2 - e_3 - \cdots - e_{13})$ and $H' = PD(23h + 6e_1 + 6e_2 - 6e_3 - \cdots - 6e_{13})$. Note that H' is orthogonal to the subspace $H_2(C_5; \mathbf{R})$ of $H_2(R_{13}; \mathbf{R})$. Applying the wall-crossing formula to $\pm \tilde{K}'_3$, H' and $PD(h)$, we get $SW_{R_{13}, H'}(\pm \tilde{K}'_3) = \pm 1$. Corollary 4.4 shows that \tilde{K}'_3 is a lift of some element $K'_3 \in \mathcal{C}(E'_3)$. Thus Theorem 4.2 gives $SW_{E'_3}(\pm K'_3) = \pm 1$. The same argument as that in the proof of Lemma 5.4 shows that K'_3 is the unique element up to sign for which $SW_{E'_3}$ is nonzero.

Let $\alpha' \in H_2(R_{13}; \mathbf{Z})$ be the element defined by $\alpha' = 3h + e_1 - e_3 - e_4 - \cdots - e_7 - e_{10} - e_{11} - e_{12} - e_{13}$. Lemma 5.1.(1) allows us to view α' as an element of $H_2(E'_3; \mathbf{Z})$. We set $L'_3 \in H^2(E'_3; \mathbf{Z})$ by $L'_3 = K'_3 - PD(\alpha')$. The element L'_3 is a characteristic element of $\langle PD(\alpha') \rangle^\perp$ and satisfies $L'^2_3 = 1$ and $K'_3 = L'_3 + PD(\alpha')$, because $K'^2_3 = 0$, $K_3 \cdot PD(\alpha') = -1$ and $(PD(\alpha'))^2 = -1$. We here denote the symbol $\langle PD(\alpha') \rangle^\perp$ as the orthogonal complement of the subspace spanned by $PD(\alpha') \in H^2(E'_3; \mathbf{Z})$. Since the symmetric bilinear form on $\langle PD(\alpha') \rangle^\perp$ is $\langle 1 \rangle \oplus 8\langle -1 \rangle$, the following lemma together with the above property of L'_3 gives us an orthogonal basis v_1, \dots, v_{10} of $H^2(E'_3; \mathbf{Z})$ such that $v^2_1 = 1$, $v^2_2 = \cdots = v^2_{10} = -1$ and $K'_3 = 3v_1 - v_2 - \cdots - v_{10}$.

Lemma 5.6 (Stipsicz–Szabó [17, The proof of Proposition 4.3], cf Wall [18, The proof of 1.6]) *Let M be a free \mathbf{Z} -module equipped with a symmetric bilinear form $\langle 1 \rangle \oplus 8\langle -1 \rangle$. If a characteristic element K of M satisfies $K^2 = 1$, then there exists an automorphism of M which preserves the symmetric bilinear form on M and maps K to $3v_1 - v_2 - \cdots - v_9$. Here v_1, \dots, v_9 denotes an arbitrary orthogonal basis of M such that $v^2_1 = 1$ and $v^2_2 = \cdots = v^2_9 = -1$.*

Similarly the above lemma together with Lemma 5.4 gives us an orthogonal basis w_1, \dots, w_{10} of $H^2(E(1)_{2,3}; \mathbf{Z})$ such that $w^2_1 = 1$, $w^2_2 = \cdots = w^2_{10} = -1$ and $K_3 = 3w_1 - w_2 - \cdots - w_{10}$. Let $\varphi' : H^2(E'_3; \mathbf{Z}) \rightarrow H^2(E(1)_{2,3}; \mathbf{Z})$ be the isomorphism defined by $v_i \mapsto w_i$ ($1 \leq i \leq 10$). The isomorphism φ' preserves the intersection forms and the Seiberg–Witten invariants.

Let $H \in H^2_+(R_{11}; \mathbf{R})$ be the element defined in the proof of Lemma 5.4. Recall that we can view H and H' as positively oriented elements of $H^2_+(E(1)_{2,3}; \mathbf{R})$ and $H^2_+(E'_3; \mathbf{R})$, respectively. Note that $(-K_3) \cdot H = 1$ and

$$(-K_3) \cdot \varphi'(H') = \varphi'(-K'_3) \cdot \varphi'(H') = (-K'_3) \cdot H' = 9.$$

We thus have $(-K_3) \cdot H > 0$ and $(-K_3) \cdot \varphi'(H') > 0$. These two inequalities together with the lemma below show $H \cdot \varphi'(H') > 0$. Hence φ' preserves the homology orientations. Freedman’s theorem gives us a required homeomorphism $\Phi' : E(1)_{2,3} \rightarrow E'_3$ which preserves the orientations and satisfies $\Phi'^* = \varphi'$.

Lemma 5.7 *Let V be a vector space of rank n over \mathbf{R} equipped with a symmetric bilinear form such that $b_2^+(V) = 1$ and $b_2^-(V) = n - 1$. Here b_2^+ and b_2^- are the same notation as that in Gompf–Stipsicz [7].*

If elements $u, v, w \in V$ satisfy $u^2 > 0$, $v^2 > 0$, $w^2 \geq 0$, $u \cdot w > 0$ and $v \cdot w > 0$, then $u \cdot v > 0$.

Proof Let $\langle u \rangle$ be the subspace spanned by u , and $\langle u \rangle^\perp$ the orthogonal complement of $\langle u \rangle$. The subspace $\langle u \rangle^\perp$ is negative definite, because $b_2^+(V) = 1$ and $u^2 > 0$. Since $V = \langle u \rangle \oplus \langle u \rangle^\perp$, there exist elements $a, a' \in \mathbf{Z}$, $x, x' \in \langle u \rangle^\perp$ such that $v = au + x$ and $w = a'u + x'$. Cauchy–Schwartz inequality implies $\sqrt{x^2(x')^2} \geq x \cdot x'$. Inequalities $v^2 > 0$ and $w^2 \geq 0$ give us $a^2u^2 > -x^2$ and $(a')^2u^2 \geq -(x')^2$. These three inequalities together with $v \cdot w > 0$ show $aa'u^2 + |aa'|u^2 > 0$. This inequality and $u \cdot w > 0$ give us $a > 0$. Hence $u \cdot v > 0$. \square

This completes the proof of Proposition 5.5. \square

6 Further remarks


We conclude this article by making some remarks.

Remark 6.1 In Figure 15–Figure 17, we used the peculiar bands, that is, bands not in local positions to prove Lemma 3.1. Note that standard bands, that is, bands in local positions are also enough to prove Lemma 3.1. However, the peculiar bands are the key of our construction of exotic $\mathbf{CP}^2 \# n \overline{\mathbf{CP}}^2$ ($5 \leq n \leq 9$) (see our forthcoming paper [20]). In the proof of Lemma 3.1, we used two 2–handles with framings $(2h, 4h)$. Instead of these two 2–handles, we can use two 2–handles with framings both $(h, 5h)$ and $(3h, 3h)$ to prove Lemma 3.1. We can also use a 2–handle with a framing $6h$ to construct Figure 7. In this construction, we can decrease the number of 3–handles of E_3 . Precisely E_3 admits a handle decomposition

$$E_3 = \text{one } 0\text{-handle} \cup \text{eleven } 2\text{-handles} \cup \text{one } 3\text{-handle} \cup \text{one } 4\text{-handle}.$$

We do not know if choices of the above bands and the above 2–handles affect diffeomorphism types of E_3 and E_5 .

Remark 6.2 Yamada asked the author if a topologically trivial but smoothly nontrivial h -cobordism between E_q and $E(1)_{2,q}$ exists. Following the argument in Gompf–Stipsicz [7, Example 9.2.15], we can prove that such an h -cobordism exists. Note that the same argument also shows that a topologically trivial but smoothly nontrivial h -cobordism between $E(1)_{2,q}$ and itself exists.

Figure 7: $\mathbf{CP}^2 \# 9\overline{\mathbf{CP}^2}$

Remark 6.3 Let X be a simply connected closed smooth 4–manifold which contains a copy of C_p , and $X_{(p)}$ the rational blow-down of X along the copy of C_p . Suppose that $X_{(p)}$ is simply connected. Do the following two conditions, X and the homomorphism $H_2(C_p; \mathbf{Z}) \rightarrow H_2(X; \mathbf{Z})$ induced by the copy of C_p , suffice to determine the (small perturbation) Seiberg–Witten invariant of $X_{(p)}$?

The proofs of Proposition 5.3 and Proposition 5.5 give an affirmative answer to this question in some cases. In a forthcoming paper, we will give a more general result for this question.

We here give a proof of the following proposition referred in the introduction of this article.

Proposition 6.4 *If a smooth 4–manifold is homeomorphic to \mathbf{S}^4 (resp. \mathbf{CP}^2) and admits neither 1– nor 3–handles in a handle decomposition, then the 4–manifold is diffeomorphic to \mathbf{S}^4 (resp. \mathbf{CP}^2).*

Proof Note that if a simply connected closed smooth 4–manifold has neither 1– nor 3–handles in a handle decomposition, then the number of 2–handles appeared in the handle decomposition is equal to the rank of the second homology group of the 4–manifold.

Suppose that a smooth 4–manifold is homeomorphic to \mathbf{S}^4 and has neither 1– nor 3–handles in a handle decomposition. Then this handle body consists of a 0–handle and a 4–handle. Since attaching a 4–handle is unique (see Gompf–Stipsicz [7]), the 4–manifold is diffeomorphic to \mathbf{S}^4 .

Suppose that a smooth 4–manifold is homeomorphic to \mathbf{CP}^2 and has neither 1– nor 3–handles in a handle decomposition. Then this handle body consists of a 0–handle, a

2–handle and a 4–handle. Thus the attaching circle of the 2–handle produces S^3 by a Dehn surgery with coefficient $+1$. Since such a knot is unknot (see Gordon–Luecke [8]), the 4–manifold is diffeomorphic to CP^2 . \square

Contrary to the above proposition, many simply connected closed topological 4–manifolds are known to admit at least two different smooth structures without 1– and 3–handles (cf Gompf–Stipsicz [7]). As far as the author knows, S^4 and CP^2 are the only known exceptions. Thus the following problem is natural.

Problem 6.5 Which simply connected closed topological 4–manifold has a unique smooth structure without 1– and 3–handles?

Finally we refer to further constructions.

Remark 6.6 This article is based on the author’s announcement [21]. In [19], we will give the rest of examples announced in [21]. In addition to these examples, we will construct a smooth 4–manifold which has the same Seiberg–Witten invariant as $E(1)_{2,3}$ and admits no 1–handles as follows: We “naturally” construct Figure 7 and perform a logarithmic transformation of multiplicity 3 in the cusp neighborhood.

In [20], we will construct examples of exotic $CP^2 \# n \overline{CP}^2$ ($5 \leq n \leq 9$) by using rational blow-downs and Kirby calculus. We also prove that our examples admit a handle decomposition without 1– and 3–handles in the case $7 \leq n \leq 9$.




Figure 8: CP^2




Figure 9: CP^2




Figure 10: CP^2




Figure 11: CP^2




Figure 12: $\mathbb{C}P^2$




Figure 13: $\mathbb{C}P^2$




Figure 14: $\mathbb{C}P^2$




Figure 15: $\mathbb{C}P^2$




Figure 16: $\mathbb{C}P^2$




Figure 17: $\mathbb{C}P^2$




Figure 18: $\mathbb{C}P^2 \# 3\overline{\mathbb{C}P}^2$




Figure 19: $\mathbb{C}P^2 \# 3\overline{\mathbb{C}P}^2$




Figure 20: $\mathbb{C}P^2 \# 9\overline{\mathbb{C}P}^2$




Figure 21: $\mathbb{C}P^2 \# 9\overline{\mathbb{C}P}^2$




Figure 22: $\mathbb{C}P^2 \# 10\overline{\mathbb{C}P}^2$




Figure 23: Handle slide




Figure 24: $\mathbb{C}P^2 \# 10\overline{\mathbb{C}P}^2$




Figure 25: $\mathbb{C}P^2 \# 11\overline{\mathbb{C}P}^2$




Figure 26: $\mathbb{C}P^2 \# 11\overline{\mathbb{C}P}^2$




Figure 27: $\mathbb{C}P^2 \# 11\overline{\mathbb{C}P^2}$




Figure 28: $\mathbb{C}P^2 \# 11\overline{\mathbb{C}P^2}$




Figure 29: $\mathbb{C}P^2 \# 12\overline{\mathbb{C}P^2}$




Figure 30: $\mathbb{C}P^2 \# 12\overline{\mathbb{C}P^2}$




Figure 31: $\mathbb{C}P^2 \# 12\overline{\mathbb{C}P^2}$




Figure 32: $\mathbb{C}P^2 \# 13\overline{\mathbb{C}P^2}$




Figure 33: $\mathbb{C}P^2 \# \overline{\mathbb{C}P^2}$




Figure 34: $\mathbb{C}P^2 \# \overline{\mathbb{C}P^2}$




Figure 35: $\mathbb{C}P^2 \# \overline{\mathbb{C}P^2}$




Figure 36: $\mathbb{C}P^2 \# \overline{\mathbb{C}P^2}$




Figure 37: $\mathbf{CP}^2 \# 2\overline{\mathbf{CP}}^2$




Figure 38: $\mathbf{CP}^2 \# 2\overline{\mathbf{CP}}^2$




Figure 39: $\mathbf{CP}^2 \# 12\overline{\mathbf{CP}}^2$




Figure 40: $\mathbb{C}P^2 \# 12\overline{\mathbb{C}P}^2$




Figure 41: $\mathbb{C}P^2 \# 13\overline{\mathbb{C}P}^2$

References

- [1] **A Akhmedov, B D Park**, *Exotic smooth structures on small 4–manifolds* arXiv: math/0701664
- [2] **T Etgü, B D Park**, *Symplectic tori in rational elliptic surfaces*, Math. Ann. 334 (2006) 679–691 MR2207879
- [3] **R Fintushel, R J Stern**, *Six lectures on four 4–manifolds* arXiv:math/0610700
- [4] **R Fintushel, R J Stern**, *Rational blowdowns of smooth 4–manifolds*, J. Differential Geom. 46 (1997) 181–235 MR1484044
- [5] **R Fintushel, R J Stern**, *Double node neighborhoods and families of simply connected 4–manifolds with $b^+ = 1$* , J. Amer. Math. Soc. 19 (2006) 171–180 MR2169045
- [6] **R E Gompf**, *Nuclei of elliptic surfaces*, Topology 30 (1991) 479–511 MR11113691
- [7] **R E Gompf, A I Stipsicz**, *4–manifolds and Kirby calculus*, Grad. Studies in Math. 20, Amer. Math. Soc. (1999) MR1707327

- [8] **C M Gordon, J Luecke**, *Knots are determined by their complements*, J. Amer. Math. Soc. 2 (1989) 371–415 MR965210
- [9] **J Harer, A Kas, R Kirby**, *Handlebody decompositions of complex surfaces*, Mem. Amer. Math. Soc. 62 (1986) iv+102 MR849942
- [10] **K Kikuchi**, *Positive 2–spheres in 4–manifolds of signature $(1, n)$* , Pacific J. Math. 160 (1993) 245–258 MR1233354
- [11] **R Kirby, editor**, *Problems in low-dimensional topology*, from: “Geometric topology (Athens, GA, 1993)”, AMS/IP Stud. Adv. Math. 2, Amer. Math. Soc. (1997) 35–473 MR1470751
- [12] **P Ozsváth, Z Szabó**, *On Park’s exotic smooth four-manifolds*, from: “Geometry and topology of manifolds”, Fields Inst. Commun. 47, Amer. Math. Soc. (2005) 253–260 MR2189937
- [13] **J Park**, *Seiberg-Witten invariants of generalised rational blow-downs*, Bull. Austral. Math. Soc. 56 (1997) 363–384 MR1490654
- [14] **J Park**, *Simply connected symplectic 4-manifolds with $b_2^+ = 1$ and $c_1^2 = 2$* , Invent. Math. 159 (2005) 657–667 MR2125736
- [15] **J Park, A I Stipsicz, Z Szabó**, *Exotic smooth structures on $\mathbb{C}P^2\#5\overline{\mathbb{C}P^2}$* , Math. Res. Lett. 12 (2005) 701–712 MR2189231
- [16] **R J Stern**, *Will we ever classify simply-connected smooth 4–manifolds?*, from: “Floer homology, gauge theory, and low-dimensional topology”, Clay Math. Proc. 5, Amer. Math. Soc. (2006) 225–239 MR2249255
- [17] **A I Stipsicz, Z Szabó**, *An exotic smooth structure on $\mathbb{C}P^2\#6\overline{\mathbb{C}P^2}$* , Geom. Topol. 9 (2005) 813–832 MR2140993
- [18] **C T C Wall**, *On the orthogonal groups of unimodular quadratic forms. II*, J. Reine Angew. Math. 213 (1963/1964) 122–136 MR0155798
- [19] **K Yasui**, *Exotic rational elliptic surfaces without 1–handles II*, in preparation
- [20] **K Yasui**, *Small exotic rational surfaces without 1– and 3–handles*, in preparation
- [21] **K Yasui**, *An exotic rational surface without 1– or 3–handles*, from: “Intelligence of low dimensional topology 2006”, Ser. Knots Everything 40, World Sci. Publ., Hackensack, NJ (2007) 375–382 MR2371749

Department of Mathematics, Graduate School of Science, Osaka University
Toyonaka, Osaka 560-0043, Japan

kyasui@cr.math.sci.osaka-u.ac.jp

Received: 20 September 2007