

All 2–dimensional links in 4–space live inside a universal 3–dimensional polyhedron

CHERRY KEARTON
VITALIY KURLIN

The hexabasic book is the cone of the 1–dimensional skeleton of the union of two tetrahedra glued along a common face. The universal 3–dimensional polyhedron UP is the product of a segment and the hexabasic book. We show that any closed 2–dimensional surface in 4–space is isotopic to a surface in UP. The proof is based on a representation of surfaces in 4–space by marked graphs, links with double intersections in 3–space. We construct a finitely presented semigroup whose central elements uniquely encode all isotopy classes of 2–dimensional surfaces.

[57Q45](#), [57Q35](#), [57Q37](#)

1 Introduction

1.1 Brief summary

This is research on the interface between geometric topology, singularity theory and semigroups. A 2–link is a closed 2–dimensional surface in 4–dimensional space \mathbb{R}^4 . We study 2–links up to isotopy, that is, a smooth deformation of the ambient 4–dimensional space. We prove that any 2–link is isotopic to a surface embedded into the universal 3–dimensional polyhedron UP. We also reduce the isotopy classification of 2–links in 4–space to a word problem in a finitely presented semigroup.

1.2 The universal polyhedron containing 2–dimensional links

First we define the universal 3–dimensional polyhedron UP.

Definition 1.1 The *theta* graph TG consists of 3 edges connecting 2 vertices. The *circled theta* graph CT is $TG \cup S^1$, where the circle S^1 meets each edge of TG in one point; see [Figure 1](#). Then CT is the 1–dimensional skeleton of two tetrahedra glued along a common face. The *hexabasic book* HB is the cone of CT. Being embedded in 3–space, the book HB divides a neighbourhood of the central vertex into 6 parts. The *universal 3–dimensional polyhedron* is $UP = HB \times [-1, 1]$.




Figure 1: The theta graph TG, circled theta graph CT, book HB

We will work in the smooth category, ie all diffeomorphisms are C^∞ -smooth. We will make necessary comments on similar constructions in the PL case.

Definition 1.2 An *embedding* is a diffeomorphism onto its image. A *2-link* is a closed (possibly disconnected or nonorientable) smooth surface S embedded into \mathbb{R}^4 . An *isotopy* between 2-links S and S' is a smooth family of diffeomorphisms $F^u: \mathbb{R}^4 \rightarrow \mathbb{R}^4$, $u \in [0, 1]$, such that $F^0 = \text{id}_{\mathbb{R}^4}$, $F^1(S) = S'$.

Fix the 4-th coordinate t in 4-space $\mathbb{R}^3 \times \mathbb{R}$. Then a 2-link in $\mathbb{R}^3 \times \mathbb{R}$ can be studied in terms of its *cross-sections* $S_t = S \cap (\mathbb{R}^3 \times \{t\})$; see Fox and Milnor [6]. Any 2-link can be isotopically deformed to a surface $S \subset \mathbb{R}^3 \times [-1, 1]$ such that the projection $\text{pr}: S \rightarrow [-1, 1]$ has distinct nondegenerate critical values. A general cross-section S_t is a classical link in $\mathbb{R}^3 \times \{t\}$, while a cross-section containing a saddle is a link with a double point. When t passes through a saddle, the cross-section $S_t = S \cap (\mathbb{R}^3 \times \{t\})$ changes by the Morse modification in the left picture of Figure 2.




Figure 2: Resolving a singular point and a band in \mathbb{R}^3

A PL analogue of the smooth approach is to decompose a 2-link $S \subset \mathbb{R}^3 \times [-1, 1]$ into handles located in different sections $\mathbb{R}^3 \times \{t_j\}$. The 1-handles of S will be represented by bands that have a distinguished core and are attached to a classical link in 3-space. Any attached band can be retracted to a singular point marked by a bridge encoding the core of the band. The cross-sections of S below and above every 1-handle locally look like the right picture of Figure 2.

1.3 Main results

The hexabasic book HB is closely related to the 3-page book TB, the cone of the *theta* graph TG consisting of 3 edges connecting 2 vertices; see Figure 7. The binding segment of TB is the cone of the 2 vertices of TG. From another point of view, the 3-page book TB can be considered as $\mathbb{R} \times T$, where T is the *triod* consisting of 3 edges connecting the central vertex O to other 3 vertices; here the binding axis α is $\mathbb{R} \times O$. The hexabasic book HB is obtained from TB by adding 3 half-disks whose 6 boundary radii are attached to the 3 edges of $\{0\} \times T$; see Figure 1.

Theorem 1.3 Any 2-dimensional link $S \subset \mathbb{R}^4$ is isotopic to a surface embedded into the universal 3-dimensional polyhedron $UP = HB \times [-1, 1]$.

The key idea of Theorem 1.3 is to put a given surface S in general position and consider its cross-sections S_t through saddles of $pr: S \rightarrow [-1, 1]$; see Claim 2.3. Such a cross-section S_t is a link with exactly one singular point, so S_t can be embedded into the 3-page book TB using the technique of 3-page embeddings developed by Kurlin and Vershinin [9; 11]; see Proposition 3.2. Both resolutions of the singular point of S_t can be realised in TB, ie the embedding extends to a regular neighbourhood of S_t in S . It remains to embed the complement of the regular neighbourhoods of all saddles into $HB \times [-1, 1]$ realising any isotopy of classical links in HB; see Lemma 3.4.

We will develop a 1-dimensional calculus for 2-links as follows. Any 2-link S in general position in $\mathbb{R}^3 \times [-1, 1]$ can be represented by a banded link BL whose bands are associated to the saddles of $pr: S \rightarrow [-1, 1]$; see Proposition 2.6(i). Retracting each band to a point, we get a marked graph whose singular points are marked by bridges encoding the cores of bands. There is a complete set of moves on marked graphs generating any isotopy of 2-links in 4-space; see Proposition 4.2. Any marked graph can be embedded into the 3-page book TB and can be encoded by a word in the alphabet of 15 letters. The moves on marked graphs are translated into relations on words, which leads to the universal semigroup SL of 2-links in 4-space.

Consider the universal semigroup SL generated by the letters a_i, b_i, c_i, d_i, x_i subject to relations (1-1)–(1-8), where $i \in \mathbb{Z}_3 = \{0, 1, 2\}$, eg $0 - 1 = 2 \pmod{3}$.

$$(1-1) \quad d_0 d_1 d_2 = 1, \quad b_i d_i = 1 = d_i b_i$$

$$(1-2) \quad a_i = a_{i+1} d_{i-1}, \quad b_i = a_{i-1} c_{i+1}, \quad c_i = b_{i-1} c_{i+1}, \quad d_i = a_{i+1} c_{i-1}$$

$$(1-3) \quad uv = vu, \quad \begin{aligned} u &\in \{a_i b_i, d_i c_i, b_{i-1} d_i d_{i-1} b_i, d_i x_i b_i\}, \\ v &\in \{a_{i+1}, b_{i+1}, c_{i+1}, b_i d_{i+1} d_i, x_{i+1}\} \end{aligned}$$

$$(1-4) \quad x_{i-1} = b_{i+1} x_i d_{i+1}, \quad b_i x_i b_i = a_i (b_i x_i b_i) c_i, \quad d_i x_i d_i = a_i (d_i x_i d_i) c_i$$

$$(1-5) \quad (d_i x_i b_i) d_i^2 d_{i+1}^2 d_{i-1}^2 = d_i^2 d_{i+1}^2 d_{i-1}^2 (d_i x_i b_i)$$

$$(1-6) \quad a_i x_i = a_i, \quad a_i b_i x_i d_i c_i = 1$$

$$(1-7) \quad d_i x_i b_i c_i x_i = b_i x_i d_i c_i x_i$$

$$(1-8) \quad \begin{aligned} w_i d_{i+1} d_i^2 d_{i-1} a_{i+1} b_{i+1} x_i b_i d_{i+1} b_i^2 b_{i+1} d_i^2 \\ = w_i b_{i-1} b_i a_i b_{i+1} a_{i+1} d_i^2 c_{i-1} b_i x_i b_i, \end{aligned} \quad \text{where } w_i = a_i b_i x_i b_i c_i$$

One of the 6 relations $b_i d_i = 1 = d_i b_i$ is superfluous and can be deduced from the remaining relations in (1-1). Moreover, the commutativity of $d_i c_i$ with a_{i+1}, b_{i+1} follows from the other relations in (1-3); see more details in Kurlin [9]. So the semigroup SL is generated by not more than 15 letters and 96 relations.

Theorem 1.4 Any 2-link $S \subset \mathbb{R}^4$ is encoded by an element $w_S \in \text{SL}$ in such a way that 2-links S and S' are isotopic if and only if their encoding elements w_S and $w_{S'}$ are equal in SL. An element $w \in \text{SL}$ encodes a 2-link if and only if w is central in SL.

Outline In Section 2, we represent 2-links in 4-space by banded links and marked graphs in 3-space. Theorem 1.3 and Theorem 1.4 are proved in Section 3 and Section 4, respectively. Banded links are more convenient for deriving a complete set of moves generating any isotopy of 2-links. Marked graphs will be used to prove our main results on embedding and encoding 2-links up to isotopy.

Acknowledgements The authors thank S Carter, F Tari and the anonymous referee for useful suggestions.

2 Representing 2-links by banded links and marked graphs

2.1 Critical level embeddings of 2-links in 4-space

Here we describe the PL approach where a 2-link is isotopically deformed to a nice embedding with handles at different levels. The smooth version of crucial Claim 2.3(ii) is a standard statement on general position proved in Section 5.

Definition 2.1 A handle of dimension n and index k is $D^k \times D^{n-k}$. A handle decomposition of a manifold M^n is a sequence of submanifolds $M_0 \subset M_1 \subset \dots \subset M_l = M$, where M_0 is a disjoint union of n -dimensional disks, each M_{i+1} is obtained from M_i by adding a handle of some index k_i . One can write $M_{i+1} = M_i \cup_{\varphi_i} (D^{k_i} \times D^{n-k_i})$, where $\varphi_i: \partial D^{k_i} \times D^{n-k_i} \rightarrow \partial M_i$ is an embedding. If before and after each handle addition one inserts a collar, the product of the attaching area and a segment, then one gets a collared handle decomposition; see Kearton and Lickorish [8, p 416].




Figure 3: A critical level embedded torus and its banded link $BL \subset \mathbb{R}^3$

A 2-link with a collared handle decomposition can be nicely embedded in \mathbb{R}^4 . The left picture of Figure 3 shows a similar embedding, where the standard 2-torus in \mathbb{R}^3 has the collared handle decomposition consisting of 4 handles and 3 collars:

- (1) the lowest handle is a 0-handle (a disk) at the level $t = -1$;
- (2) the 2 intermediate handles are 1-handles (bands) at the levels $t = \pm 1/2$;
- (3) the highest handle is a 2-handle (a disk) at the level $t = +1$.

Definition 2.2 A *critical level PL embedding* is a PL embedding of a 2-link $S \subset \mathbb{R}^3 \times [-1, 1]$ with a collared handle decomposition satisfying (i), (ii); see Kearton and Lickorish [8, p 417]:

- (i) the handles are in different sections $\mathbb{R}^3 \times \{t_j\}$, where $-1 < t_1 < \dots < t_n < 1$;
- (ii) each collar between adjacent handles of S is embedded as the direct product $A \times [t_j, t_{j+1}] \subset \mathbb{R}^3 \times [t_j, t_{j+1}]$, where $A \subset \mathbb{R}^3$ is the attaching area of the handles.

A smooth embedding $S \subset \mathbb{R}^3 \times [-1, 1]$ is called a smooth *critical level embedding* if the projection $\text{pr}: S \rightarrow [-1, 1]$ has all its critical points in different sections $\mathbb{R}^3 \times \{t_j\}$. This is a general position assumption.

Claim 2.3 (i) (Kearton and Lickorish [8, Theorem 1, p 420]) Any 2-dimensional PL link in 4-space is isotopic to the image of a critical level PL embedding $S \subset \mathbb{R}^3 \times [-1, 1]$.

(ii) Any smooth 2-link is smoothly isotopic to a surface $S \subset \mathbb{R}^3 \times [-1, 1]$ such that all critical points of $\text{pr}: S \rightarrow [-1, 1]$ are nondegenerate and have distinct values.

We will use the smooth version of [Claim 2.3\(ii\)](#), which will be deduced from the transversality theorem of Thom in [Section 5](#). [Claim 2.3\(i\)](#) is worth keeping in mind when one associates a banded link to a 2–link in [Proposition 2.6\(i\)](#).

2.2 Representing 2–links in 4–space by banded links in 3–space

We define banded links, links with bands, which will represent 2–links in 4–space.

Definition 2.4 A *banded link* is a collection of circles and bands in \mathbb{R}^3 such that

- (i) the circles and bands are nonoriented and non-self-intersecting;
- (ii) the circles and bands are disjoint except for each band having a pair of opposite sides *attached* to disjoint arcs in the circles, the other sides are called *free*.

In every band we mark its *core*, an arc connecting its attached opposite sides; see [Figure 3](#). Banded links are considered up to isotopy of \mathbb{R}^3 . The bands of a banded link will represent 1–handles of a 2–link. In every band B of a banded link BL consider the opposite free sides not connected by the core of B . Replace B by its free sides, the resulting usual nonoriented link in \mathbb{R}^3 is called the *positive resolution* BL_+ of the banded link BL ; see the right picture of [Figure 3](#). Similarly define the *negative resolution* BL_- replacing every band B by the opposite attached sides connected by the core of B . A banded link BL is *admissible*, if both resolutions BL_{\pm} are trivial links.

If a PL 2–link $S \subset \mathbb{R}^3 \times [-1, 1]$ has all its 1–handles in the zero section $\mathbb{R}^3 \times \{t = 0\}$, then the cross-section $S_0 = S \cap (\mathbb{R}^3 \times \{t = 0\})$ is a banded link. We will use much weaker assumptions and construct a banded link for any critical level embedding. [Proposition 2.6](#) leads to a calculus for 2–links in [Proposition 4.2](#) and provides a function from the set of 2–links to the set of admissible banded links.

Definition 2.5 Given a 2–dimensional surface S , consider the space of all smooth functions $f: S \rightarrow \mathbb{R}^4$ with the Whitney topology; see [Definition 5.2](#). The space CS of all 2–links $S \subset \mathbb{R}^4$ has the induced topology. Points in CS will be classified using the projection $\text{pr}: S \rightarrow \mathbb{R}$ to the 4–th coordinate t . A 2–link $S \in \text{CS}$ is called

- *generic* if all critical points of pr are nondegenerate and have distinct values;
- an $A_1^+ A_1^+$ –*singularity* if S fails to be generic because of 2 nondegenerate extrema of $\text{pr}: S \rightarrow \mathbb{R}$ that have the same value;
- an $A_1^+ A_1^-$ –*singularity* if S fails to be generic because of a nondegenerate saddle and extremum of $\text{pr}: S \rightarrow \mathbb{R}$ that have the same value;

- an $A_1^- A_1^-$ -singularity if S fails to be generic because of 2 nondegenerate saddles of $\text{pr}: S \rightarrow \mathbb{R}$ that have the same value;
- an A_2 -singularity if S fails to be generic because of a singularity of $\text{pr}: S \rightarrow \mathbb{R}$ having the form $\text{pr}(x, y) = x^2 - y^3$ in local coordinates x, y .

The sign in the notation above is the sign of the determinant $\text{pr}_{xx}\text{pr}_{yy} - \text{pr}_{xy}^2$ of the Jacobi matrix of 2nd order derivatives at a critical point. Denote by $\Sigma_{++}, \Sigma_{+-}, \Sigma_{--}$ and Σ_2 the *subspaces* of the corresponding singularities in the space CS. Introduce the *singular* subspace $\Sigma = \Sigma_{++} \cup \Sigma_{+-} \cup \Sigma_{--} \cup \Sigma_2$. An isotopy of 2-links can be considered as a path in CS. In Proposition 2.6 we consider paths nicely meeting the singular subspace Σ .

Proposition 2.6 (i) *To any a critical level embedding $S \subset \mathbb{R}^3 \times [-1, 1]$ we associate a banded link BL well-defined up to the slide/swim moves in Figure 4.*




Figure 4: Cup/cap moves and slide/swim moves of banded links

(ii) *If 2-links S, S' are isotopic through generic 2-links, then the associated banded links BL, BL' are related by the slide/swim moves in Figure 4.*

(iii) If 2-links S, S' are isotopic through generic 2-links and one of $A_1^+ A_1^{+-}, A_1^+ A_1^{-}, A_1^- A_1^{-}$ -singularities, then BL, BL' are related by the slide/swim moves in [Figure 4](#).

(iv) If 2-links S, S' are isotopic through generic 2-links and exactly one A_2 -singularity, then BL, BL' are related by the cap/cup and slide/swim moves in [Figure 4](#).

Proof (i) The lowest critical point of a generic 2-link S with respect to $\text{pr}: S \rightarrow [-1, 1]$ at $t = t_1$ is a minimum, so the cross-section $S_{t_1+\varepsilon}$ is a trivial knot for some $\varepsilon > 0$. The section $S_{t_1+\varepsilon}$ is a prototype of a future banded link BL , which will be located in a fixed copy of \mathbb{R}^3 . The key idea in constructing BL is to watch the current cross-section $S_t = S \cap (\mathbb{R}^3 \times \{t\})$ simultaneously adding bands and trivial knots corresponding to new saddles and minima, respectively. The left column of [Figure 5](#) contains cross-sections S_t for different values of t . The right column shows successive stages of constructing BL whose final form is the top right.

While t is increasing, we isotopically deform the current banded link $BL \subset \mathbb{R}^3$ following $S_t = S \cap (\mathbb{R}^3 \times \{t\})$; see [Figure 5](#). The existing bands of BL can be deformed to avoid intersections with the rest of BL . For each new minimum of S in $\mathbb{R} \times \{t_j\}$, add a trivial knot from $S_{t_j+\varepsilon}$ to the current banded link $BL \subset \mathbb{R}^3$.

For each new saddle of S , attach a small band B to BL . The band B has 2 opposite sides attached to branches of the previous link BL . While t passes the critical value, the attached sides of B are retracted to a point and are replaced by the free sides of B . The band B can not meet the attached sides of other bands of BL since these sides are not included into the current cross-section of S . So there are only 2 cases when the new link with bands does not satisfy [Definition 2.4](#).

(a) One (or two) of the attached sides of B may meet a free side of another band B' of BL ; see the upper picture of [Figure 6](#). Then slide B along the free side of B' in any of the two directions so that in the end the attached side of B does not meet B' .

(b) The band B intersects the interior of another band B' of BL , see the lower picture of [Figure 6](#). Then B swims through any of the attached sides of B' , so B, B' fall apart. The band B can not swim through the free sides of B' as they belong to the current cross-section of S . For each new 2-handle (a maximum), we keep the corresponding trivial knot of BL , although it disappears from $S_t = S \cap (\mathbb{R}^3 \times \{t\})$.

After we have passed all critical values of $\text{pr}: S \rightarrow \mathbb{R}$, the associated banded link $BL \subset \mathbb{R}^3$ has been constructed.

(ii) The construction above is not affected by an isotopy of S keeping the order of critical points of $\text{pr}: S \rightarrow [-1, 1]$. Indeed all cross-sections S_t are replaced by isotopic




Figure 5: Cross-sections and a banded link of the spun 2-knot of the trefoil




Figure 6: A band slides or swims to remove an intersection

links, so the resulting banded link is isotopic to the original one provided that we remove intersections of bands in [Figure 6](#) in the same way.

(iii) The given isotopy of S is a smooth path passing through one of $A_1^+ A_1^{+-}$, $A_1^+ A_1^-$, $A_1^- A_1^-$ -singularities in the space CS of 2-links. For $A_1^+ A_1^+$ or $A_1^+ A_1^-$, an extremum and another singularity swap their heights, so we add a new trivial knot (passing a minimum) or keep an existing trivial knot (passing a maximum) that does not affect the other singularity. For an $A_1^- A_1^-$ -singularity, two saddles of S swap their heights, so we add 2 bands to BL in the reverse order. Consider the critical moment when both saddles are in the same section $\mathbb{R}^3 \times \{t_j\}$. If the associated bands do not intersect each other, then the new banded link is isotopic to the original one. In (i) we listed the only cases (a), (b) when one band may intersect another, which led to the moves in [Figure 6](#) so the banded links are equivalent through the slide/swim moves.

(iv) If an isotopy of S passes through an A_2 -singularity, then around this moment a nondegenerate saddle and extremum appear in a 2-link, see [Claim 4.3\(iv\)](#). In the case of a minimum, one adds a trivial knot to the current banded link BL and a band attached to the trivial knot and to an existing branch of BL as shown in the cup move of [Figure 4](#). In the case of a maximum, one adds a band attached by both sides to a branch of the current banded link BL as shown in the cap move of [Figure 4](#). Recall that we keep the trivial knot when t passes a maximum. The leftmost and rightmost columns of [Figure 4](#) show projections of 2-links to \mathbb{R}^3 around singular moments. The 4-th axis of $\mathbb{R}^3 \times \mathbb{R}$ projects to the vertical axis of \mathbb{R}^3 . \square

Conversely, any admissible banded link will give rise to a 2-link in 4-space; see [Lemma 2.8](#). One can describe all moves of banded links generating any isotopy of 2-links in 4-space. Banded links were called *knots with bands* in Swenton [10].

2.3 Representing 2-links in 4-space by marked graphs in 3-space

[Theorem 1.4](#) is easier to prove representing 2-links by marked graphs, which are singular links with bridges at singular points.

Definition 2.7 After deformation retracting each band of a banded link BL to a point, we get a *singular link* (see Kurlin and Vershinin [11]), a collection of closed curves with finitely many double transversal intersections; see [Figure 2](#) and [Figure 5](#). The core of each retracted band defines a bridge at the singular point, a straight arc in a small plane neighbourhood of each singular point. We consider the resulting *marked graph* MG up to isotopy in \mathbb{R}^3 keeping a neighbourhood of each singular point in a (moving) plane.

In the smooth approach, the zero section $S \cap (\mathbb{R}^3 \times \{0\})$ containing all saddles of $p: S \rightarrow [-1, 1]$ is a marked graph whose bridges show how to resolve the singular points for $t > 0$ (along bridges) and $t < 0$ (across bridges); see [Figure 2](#) and [Figure 3](#). An abstract marked graph MG , ie a singular link with bridges, can be converted into a banded link BL replacing each bridge by a small rectangle whose core coincides with the bridge. So there is a 1-1 correspondence between banded links and marked graphs. [Lemma 2.8](#) provides a unique function from the set of admissible banded links to the set of 2-links, which is the inverse of the function from [Proposition 2.6](#).

Lemma 2.8 Any admissible banded link $BL \subset \mathbb{R}^3$ gives rise to a 2-link $S \subset \mathbb{R}^4$ that can be represented by BL as in [Proposition 2.6\(i\)](#).

Proof Take the marked graph $MG \subset \mathbb{R}^3$ associated to the given banded link. Isotopically deform MG in such a way that neighbourhoods of all singular points of MG are contained in a single hyperplane of $\mathbb{R}^3 \times \{0\}$.

Resolving the singular points along the bridges for $t > 0$ and across the bridges for $t < 0$, extend the embedding $MG \subset \mathbb{R}^3 \times \{0\}$ to a surface $S' \subset \mathbb{R}^3 \times [-\varepsilon, \varepsilon]$ for some $\varepsilon > 0$, such that the boundary $\partial S'$ consists of trivial links in $\mathbb{R}^3 \times \{\pm\varepsilon\}$.

Since both sections $S'_{\pm\varepsilon} = S' \cap (\mathbb{R}^3 \times \{t = \pm\varepsilon\})$ are unlinks, one can find isotopies $\varphi_t^\pm: \mathbb{R}^3 \rightarrow \mathbb{R}^3$, $t \in [\varepsilon, 1-\varepsilon]$, such that each $\varphi_{1-\varepsilon}^\pm(S'_{\pm\varepsilon})$ is a collection of small disjoint circles in a plane. The isotopies φ_t^\pm define the embedding of a 2-link S without small

disks into $\mathbb{R}^3 \times [\varepsilon - 1, 1 - \varepsilon]$, one disk for each component of ∂S . Attaching a disc to each boundary circle gives a closed surface $S \subset \mathbb{R}^3 \times [-1, 1]$.

The zero section $S \cap (\mathbb{R}^3 \times \{0\})$ is the original marked graph MG . A small isotopy deformation makes S generic. The construction of Proposition 2.6(i) gives a banded link equivalent to MG as all bands may be chosen small and nonintersecting. \square

3 Three-page embeddings of marked graphs

3.1 Any marked graph can be embedded into the 3–page book

Recall that the 3–page book is $TB = \mathbb{R} \times T$, where T is the triod consisting of 3 edges E_0, E_1, E_2 joining the vertex O to the other 3 vertices. The line $\alpha = \mathbb{R} \times O$ is said to be the *binding axis*, $P_i = \mathbb{R} \times E_i$ are called the *pages*, $i = 0, 1, 2$.




Figure 7: The encoding letters for 3–page embeddings of marked graphs

Definition 3.1 An embedding of a marked graph G into the 3–page book TB is called a 3–page embedding, if conditions (i)–(v) hold:

- (i) the intersection $G \cap \alpha$ of G and the binding axis α is a finite set of points;
- (ii) the arcs at every point of $G \cap \alpha$ lie in 2 pages P_i, P_j , $i \neq j$ (see Figure 7);
- (iii) all singular points of G lie in α , a neighbourhood of each singular point lies in a broken plane consisting of two pages and looks locally like a cross \times ;
- (iv) the bridge at each singular point lies in the binding axis α ;
- (v) every connected component of $G \cap P_i$ is projected monotonically to α .

The arcs in the page P_2 are dashed in Figure 7 and Figure 8. All classical and singular links can be embedded into TB in the sense of Definition 3.1; see Figure 8.

The pictures in each vertical column of Figure 7 are obtained from each other by rotation around α . The rotation corresponds to the shift $i \mapsto i + 1$ of indices, $i \in \mathbb{Z}_3 = \{0, 1, 2\}$. A 3-page embedding can be encoded by a word in the alphabet of 15 letters describing the local behaviour of G near the intersection points $G \cap \alpha$; see Figure 7. The 3-page embedding in Figure 8 is encoded by $w_G = a_0 a_1 (b_2 b_0 b_1)^2 d_0 a_1 (x_1 b_1)^2 c_1 d_1 b_0 (d_1 d_0 d_2)^2 c_1 c_0$. So a 3-page embedding of the marked graph G_S of a 2-link S is a 1-dimensional representation of $S \subset \mathbb{R}^4$.




Figure 8: A 3-page embedding of the marked graph from Figure 9

We give a proof of the embedding result from Kurlin and Vershinin [11], because this construction plays an important role in further considerations.

Proposition 3.2 (Kurlin and Vershinin [11]) *Any marked graph $G \subset \mathbb{R}^3$ is isotopic to a 3-page embedding $G \subset TB$ in the sense of Definition 3.1.*

Proof Consider a plane diagram D of $G \subset \mathbb{R}^3$ in general position with finitely many double crossings. At each crossing in the diagram D mark a small overcrossing arc. Recall that, at each singular point of G , there is a marked bridge transversally intersecting both branches of G passing through the singular point.

In the plane containing the diagram D , draw a continuous path α such that

- (1) the path α passes through each marked arc and bridge exactly once;
- (2) α transversally intersects the rest of D , the endpoints of α are away from D .




Figure 9: How to construct a 3–page embedding of a marked graph

Isotopically deform the plane containing D in such a way that α becomes a straight line containing all marked arcs and bridges of D . Denote the upper half-plane and lower half-plane of $\mathbb{R}^2 - \alpha$ by P_0 and P_2 , respectively. Notice that a neighbourhood of each singular point looks like a cross \times with a centre in the axis α ; see Figure 9.

Attach the third half-plane P_1 to α and push all marked arcs into P_1 ; see Figure 8. If both (say) upper arcs at some singular point $v \in G$ go to points on one side of the point $v \in \alpha$, then make an additional couple of crossings in the intersection $\alpha \cap D$ like in Reidemeister move II; see Figure 13. For instance, in the embedding $a_2b_2x_2$ both upper arcs go to the right, see the lower right picture of Figure 15, more details are in Kurlin and Vershinin [11]. Then the intersection $G \cap P_i$ is a finite collection of disjoint arcs, which can be made monotonic with respect to the projection $TB \rightarrow \alpha$, $i = 0, 1, 2$. \square

3.2 Any isotopy of links can be realised in the hexabasic book

The following lemma is a keystone of the 3–page approach to knot theory and was originally proved by I Dynnikov [2; 3].

Lemma 3.3 (Kurlin [9]) *Any isotopy of 3–page embeddings of classical links is decomposed into finitely moves in Figure 10 and their images under $i \mapsto i + 1$, $i \in \mathbb{Z}_3$.*

The algebraic form of the moves in Figure 10 is below, $i \in \mathbb{Z}_3 = \{0, 1, 2\}$; see Kurlin [9]:

$$(3-1) \quad d_0d_1d_2 = 1, \quad b_id_i = 1 = d_ib_i$$

$$(3-2) \quad a_i = a_{i+1}d_{i-1}, \quad b_i = a_{i-1}c_{i+1}, \quad c_i = b_{i-1}c_{i+1}, \quad d_i = a_{i+1}c_{i-1}$$

$$(3-3) \quad uv = vu, \quad \text{where} \quad \begin{aligned} u &\in \{a_ib_i, d_ic_i, b_{i-1}d_id_{i-1}b_i\}, \\ v &\in \{a_{i+1}, b_{i+1}, c_{i+1}, b_id_{i+1}d_i\} \end{aligned}$$

Lemma 3.4 is the crucial step in Theorem 1.3.




Figure 10: Finitely many moves generating any isotopy of classical links

Lemma 3.4 *The moves in Figure 10 are realised in the hexabasic book HB.*

Proof All the moves in Figure 10, apart from the commutativity of $a_i, b_i, c_i, b_{i-1} d_i d_{i-1}$ with $b_{i+1} d_{i-1} d_{i+1} b_{i-1}$, can be realised in the 3-page book TB. For instance, the relation $b_2 d_2 = 1$ is realised by compressing the slice between the 2 intersection points and removing the resulting point from α . The other relations are realised in HB; see a geometric realisation of $(b_1 d_2 d_1 b_2) a_0 = a_0 (b_1 d_2 d_1 b_2)$ in Figure 11. \square

3.3 Any 2-link is isotopic to a surface in the universal polyhedron

Here we finish the proof of Theorem 1.3.




Figure 11: Realising a commutative relation in the hexabasic book HB

Proof of Theorem 1.3 By Claim 2.3 any 2–link in 4–space is isotopic to a surface $S \subset \mathbb{R}^3 \times [-1, 1]$ having all maxima, minima and saddles in different sections $\mathbb{R}^3 \times \{t_j\}$ for some $-1 < t_1 < \dots < t_n < 1$. In Step 1 we embed each cross-section S_{t_j} into the 3–page book. In Step 2 we extend this embedding to a regular neighbourhood of S_{t_j} . In Step 3 we embed the complement of the neighbourhoods into $\text{HB} \times [-1, 1]$.

Step 1 Choose $\varepsilon > 0$ such that the closed ε –neighbourhoods $N_\varepsilon(S_{t_j})$ of S_{t_j} in S are disjoint and each of them contains exactly one critical point of $\text{pr}: S \rightarrow [-1, 1]$, $j = 1, \dots, n$. Then the boundaries $\partial N_\varepsilon(S_{t_j})$ are classical links. By Proposition 3.2 there is an isotopy $f_j^u: \mathbb{R}^3 \times \{t_j\} \rightarrow \mathbb{R}^3 \times \{t_j\}$, $u \in [0, 1]$, moving S_{t_j} into TB, ie $f_j^0 = \text{id}_{\mathbb{R}^3}$, $f_j^1(S_{t_j}) \subset \text{TB} \times \{t_j\}$ is a 3–page embedding. Take smooth functions $g_j: [t_j - \varepsilon, t_j + \varepsilon] \rightarrow [0, 1]$ such that $g_j(t_j) = 1$ and $g_j(t_j \pm \varepsilon) = 0$. Extend f_j^u to

$$F_j^u: \mathbb{R}^3 \times [t_j - \varepsilon, t_j + \varepsilon] \rightarrow \mathbb{R}^3 \times [t_j - \varepsilon, t_j + \varepsilon], \quad u \in [0, 1],$$

$$F_j^u(x, t) = (f_j^{ug_j(t)}(x), t), \quad \text{where } x \in \mathbb{R}^3, t \in [t_j - \varepsilon, t_j + \varepsilon].$$

Then $F_j^u = f_j^u$ for $t = t_j$ and $F_j^u = \text{id}$ for $t = t_j \pm \varepsilon$. Hence $\partial N_\varepsilon(S_{t_j})$ are pointwise fixed and we may combine F_j^u together to form a smooth isotopy $F^u: \mathbb{R}^3 \times [-1, 1] \rightarrow \mathbb{R}^3 \times [-1, 1]$ moving each S_{t_j} into $\text{TB} \times \{t_j\}$. Denote the resulting surface by S' .

Step 2 If a singular cross-section S'_{t_j} has a double intersection, then both positive and negative resolutions of S'_{t_j} can be embedded into TB. Indeed the positive and negative resolutions of the singular point x_i are encoded by 1 and $c_i a_i$, respectively; see Figure 12. Given an encoding word w_j of $S'_{t_j} \subset \text{TB}$, the positive resolution of S'_{t_j} is encoded by w_j after removing the letter x_i representing the double point of S'_{t_j} .

The argument below with the sign \pm covers 2 cases when either $+$ or $-$ is taken in all formulae. If S'_{t_j} contains a maximum or minimum, $S'_{t_j \pm \varepsilon/2}$ can be embedded into TB. So there are isotopies $h_{\pm j}^u: \mathbb{R}^3 \times \{t_j \pm \varepsilon/2\} \rightarrow \mathbb{R}^3 \times \{t_j \pm \varepsilon/2\}$, $u \in [0, 1]$, moving each $S'_{t_j \pm \varepsilon/2}$ into $\text{TB} \times \{t_j \pm \varepsilon/2\}$. Take smooth functions $\tilde{g}_j: [t_j - \varepsilon, t_j + \varepsilon] \rightarrow [0, 1]$ such that $\tilde{g}_j(t_j) = 0 = \tilde{g}_j(t_j \pm \varepsilon)$ and $\tilde{g}_j(t_j \pm \varepsilon/2) = 1$. Extend $h_{\pm j}^u$ to

$$H_j^u: \mathbb{R}^3 \times [t_j - \varepsilon, t_j + \varepsilon] \rightarrow \mathbb{R}^3 \times [t_j - \varepsilon, t_j + \varepsilon], \quad u \in [0, 1],$$




Figure 12: Resolving a singular point in the 3-page book TB

$$H_j^u(x, t) = (h_{\pm j}^{u\tilde{g}_j(t)}(x), t) \text{ for } x \in \mathbb{R}^3, t \text{ between } t_j \text{ and } t_j \pm \varepsilon.$$

Then $H_j^u = h_{\pm j}^u$ for $t = t_j \pm \varepsilon/2$ and $H_j^u = \text{id}$ for $t = t_j, t = t_j \pm \varepsilon$. Hence S'_{t_j} and $\partial N_\varepsilon(S'_{t_j})$ are pointwise fixed and we may combine H_j^u together to form a smooth isotopy $H^u: \mathbb{R}^3 \times [-1, 1] \rightarrow \mathbb{R}^3 \times [-1, 1]$ moving each $N_{\varepsilon/2}(S'_{t_j})$ into $\text{TB} \times [t_j - \varepsilon/2, t_j + \varepsilon/2]$. Denote the resulting surface by S'' .

Step 3 The cross-sections $S''_{t_j + \varepsilon/2}$ and $S''_{t_{j+1} - \varepsilon/2}$ are isotopic classical links, $j = 1, \dots, n - 1$. By Lemma 3.3 and Lemma 3.4 any isotopy of classical links can be realised in HB. Then the layers $S'' \cap (\mathbb{R}^3 \times [t_j + \varepsilon/2, t_{j+1} - \varepsilon/2])$ can be replaced by an isotopy of links in $\text{HB} \times [t_j + \varepsilon/2, t_{j+1} - \varepsilon/2]$. It remains to extend the embedding to the neighbourhoods of the lowest minimum and highest maximum of S'' shrinking their boundaries in HB. So the final surface is embedded into $\text{HB} \times [-1, 1]$. \square

4 The universal semigroup of 2-dimensional links

4.1 Local moves of marked graphs generate any isotopy of 2-links

Here we derive a complete set of moves of banded links and marked graphs, that generate any isotopy of 2-links in 4-space. Marked graphs can be represented by plane diagrams with small straight arcs denoting bridges over singular points; see Figure 2 and Figure 9. In particular, the cyclic order of edges at each singular point is invariant.

Lemma 4.1 [7] *Marked graphs are isotopic in \mathbb{R}^3 if and only if their plane diagrams can be obtained from each other by finitely many Reidemeister moves in Figure 13, where all symmetric images of the moves should be considered.*

The moves in Figure 13 are *local* in the sense, that a small disk in the left part of each move is replaced by another small disk in the right part of the move, while the rest of




Figure 13: Reidemeister moves for rigid isotopy of marked graphs

the diagram remains unchanged. The singular points in moves IV and V of Figure 13 can be equipped with arbitrary corresponding bridges. The proof is a direct application of the transversality theorem of Thom similarly to a proof of the Reidemeister theorem for plane diagrams of classical links; see Fiedler and Kurlin [5, Section 2; 4, Section 2].

Proposition 4.2 *Marked graphs represent isotopic 2–links in 4–space if and only if they can be obtained from each other by finitely many moves in Figure 14.*




Figure 14: Moves of marked graphs generating isotopy of 2–links

Symmetric images of the moves in Figure 14 are skipped as they can be reduced to the standard moves using an isotopy in \mathbb{R}^3 . Proposition 4.2 was conjectured by K Yoshikawa in [12]. F Swenton [10] claimed a proof of Proposition 4.2 using banded links and the equivalent moves in Figure 4. M Saito wrote in his review for the MathSciNet: “It is claimed that this set of moves is equivalent to Yoshikawa’s moves. It might be beneficial of some more detailed accounts, for example, those for the above claim, are discussed further and presented elsewhere in the literature.” The authors were asked by S Carter to fill in these details, so we give a more detailed proof of Proposition 4.2 for banded links. Recall that the singular subspace Σ of the space CS of 2–links was introduced in Definition 2.5. The following result will be formally deduced in Section 5 using the transversality theorem of Thom.

- Claim 4.3** (i) *The closure of the subspace Σ has codimension 1 in the space CS.*
- (ii) *The complement of the closure $\overline{\Sigma}$ in CS consists of generic 2-links.*
- (iii) *Any isotopy of 2-links can be deformed in such a way that all intermediate 2-links are generic apart from finitely many singularities of [Definition 2.5](#).*
- (iv) *If an isotopy passes through an A_2 -singularity, then a nondegenerate saddle and extremum collide and disappear as shown in the top picture of [Figure 18](#).*

[Claim 4.3](#)(i)–(ii) say that any point of CS can be removed from Σ by a small perturbation, ie a 2-link can be made generic, which implies [Claim 2.3](#)(ii). [Claim 4.3](#)(iii) says that the singularities of [Definition 2.5](#) are the only singularities occurring in any isotopy of 2-links in general position.

Proof of [Proposition 4.2](#) By [Claim 4.3](#)(iii) any isotopy of 2-links can be deformed into a smooth path transversal to the subspace $\Sigma \subset \text{CS}$. When the path passes through one of the singularities, the associated banded link changes according to [Proposition 2.6](#)(iii),(iv), which led to the moves in [Figure 4](#) as required. \square

4.2 A 1-dimensional encoding 2-links up to isotopy in 4-space

Here we reduce the isotopy classification of 2-links in 4-space to a word problem in the finitely presented semigroup SL, the universal semigroup of 2-links. Recall that moves (1-1)–(1-8) on 3-page embeddings were defined in [Section 1.3](#). [Theorem 1.4](#) follows from the following generalisation of [Lemma 3.3](#) to singular links.

Proposition 4.4 (Kurlin and Vershinin [[11](#)]) *Consider the semigroup SK generated by $a_i, b_i, c_i, d_i, x_i, i \in \mathbb{Z}_3$, subject to relations (1-1)–(1-5) from [Section 1.3](#). Then any singular link $G \subset \mathbb{R}^3$ is encoded by an element $w_G \in \text{SK}$ in such a way that singular links G, G' are isotopic if and only if the elements w_G and $w_{G'}$ are equal in SK. An element $w \in \text{SK}$ encodes a singular link if and only if w is central in SK.*

Proof of [Theorem 1.4](#) Any 2-link can be represented by its marked graph G whose 3-page embedding is encoded by a word in the letters $a_i, b_i, c_i, d_i, x_i, i \in \mathbb{Z}_3$, as described before [Proposition 3.2](#). All encoding elements form the centre of SL as the same result holds for the universal semigroup SK of singular links, ie relations (1-1)–(1-5) imply that any encoding element commutes with the generators.

The remaining part of [Theorem 1.4](#) states that two 3-page embeddings of marked graphs represent isotopic 2-links in 4-space if and only if they can be related by algebraic

moves (1–1)–(1–8) in Section 1.3. By Lemma 3.3, Lemma 3.4 and Proposition 4.2 it suffices to realise moves VI, VII, VIII in Figure 14 by 3–page embeddings.

In moves VI, VII, VIII a small disk in the left part is replaced by another small disk in the right part. Similarly to the construction of a 3–page embedding, choose a path α passing through overcrossing arcs and bridges at singular points; see Figure 15, Figure 16 and Figure 17. Deform the diagrams in such a way that α becomes a straight line and push all overcrossing arcs into the half-plane P_1 , all bridges remain in α .




Figure 15: Realising moves VI of Figure 14 in terms of 3–page embeddings

In Figure 15 moves VI are encoded by $a_1x_1 = a_1$ and $a_1b_1x_1d_1c_1 = 1$ equivalent to (1–6) for $i = 1$. We made additional intersections of α with the diagram to decompose the resulting embedding into local 3–page embeddings from Figure 7.




Figure 16: Realising move VII of Figure 14 in terms of 3–page embeddings

In Figure 16 move VII is encoded by $d_1x_1b_1c_1x_1 = b_1x_1d_1c_1x_1$, which is (1-7) for $i = 1$. Numbers 1, 2, 3, 4, 5, 6 denote arcs going out of the small disk replaced by move VII, eg the path α starts between arcs 1, 4 and ends between arcs 3, 6.




Figure 17: Realising move VIII of Figure 14 in terms of 3-page embeddings

In Figure 17 move VIII is encoded by

$$(a_1b_1x_1b_1c_1)d_2d_1(b_2d_2)d_1d_0a_2b_2x_1b_1d_2b_1(b_2d_2)b_1b_2d_1^2 = (a_1b_1x_1b_1c_1)b_0b_1(b_2d_2)a_1b_2a_2d_1(b_2d_2)d_1c_0b_1x_1b_1,$$

which is equivalent to (1-8) for $i = 1$ after removing $b_2d_2 = 1$ by relation (1-1). The relations for other $i \in \mathbb{Z}_3$ were added to make the presentation symmetric. \square

5 Appendix: the multi-jet transversality theorem of Thom

Here we follow Arnold, Varchenko and Gusein-Zade [1, Sections I.2, I.8].

Let $\xi, \eta: M \rightarrow N$ be smooth maps between finite dimensional manifolds with Riemannian metrics ρ_M, ρ_N , respectively.

Definition 5.1 The maps ξ and η have the tangency of order k at a point $z \in M$ if k is the maximal integer such that $\rho_N(\xi(w), \eta(w))/(\rho_M(z, w))^k \rightarrow 0$ as $w \in M$ tends to z , eg the curve $\xi(w) = w^{k+1}$ has the tangency of order k with $\eta(w) = 0$.

The l -tuple k -jet of the map ξ at $(z_1, \dots, z_l) \in M^l$ is the equivalence class of smooth maps $\eta: M \rightarrow N$ up to tangency of order k at the points $z_1, \dots, z_l \in M$, eg the 1-tuple 1-jet $j_{[1]}^1 \xi(z)$ of a map $\xi: \mathbb{R} \rightarrow \mathbb{R}$ is determined by $z, \xi(z), \dot{\xi}(z)$.

Denote by $J_{[l]}^k(M, N)$ the space of all l -tuple k -jets of smooth maps $\xi: M \rightarrow N$ for all $(z_1, \dots, z_l) \in M^l$. Let (x_1, \dots, x_m) and (y_1, \dots, y_n) be local coordinates in M and N , respectively. If ξ is defined locally by $y_j = \xi_j(x_1, \dots, x_m)$, $j = 1, \dots, n$, then the l -tuple k -jet of ξ at (z_1, \dots, z_l) is determined by l arrays of the data below:

$$\{x_1, \dots, x_m\}; \quad \{y_1, \dots, y_n\}; \quad \left\{ \frac{\partial \xi_j}{\partial x_i} \right\}; \quad \dots \quad \left\{ \frac{\partial^k \xi_j}{\partial x_{i_1} \dots \partial x_{i_s}} \right\}, \quad i_1 + \dots + i_s = k.$$

The quantities above define local coordinates in $J_{[l]}^k(M, N)$. The l -tuple k -jet $j_{[l]}^k \xi$ of a smooth map $\xi: M \rightarrow N$ can be considered as the map $j_{[l]}^k \xi: M^l \rightarrow J_{[l]}^k(M, N)$, namely (z_1, \dots, z_l) goes to the l -tuple k -jet of ξ at (z_1, \dots, z_l) .

The manifold $J_{[l]}^k(M, N)$ is finite dimensional, eg $J_{[l]}^0(M, N) = (M \times N)^l$,

$$\dim J_{[l]}^1(M, N) = (m + n + mn)l, \quad \dim J_{[l]}^2(M, N) = (m + n + mn + \frac{m(m+1)}{2}n)l.$$

Definition 5.2 Take an open set $W \subset J_{[l]}^k(M, N)$. The set of smooth maps $f: M \rightarrow N$ with l -tuple k -jets from W is *open*. These sets for all open $W \subset J_{[l]}^k(M, N)$ over all k, l form a basis of the *Whitney topology* in $C^\infty(M, N)$. The space CS of all 2-links $S \subset \mathbb{R}^4$ inherits the *Whitney topology* from $C^\infty(S, \mathbb{R}^4)$.

So maps are close in the Whitney topology if they are close with all derivatives.

Definition 5.3 Let M be a finite dimensional smooth manifold. A subspace $\Lambda \subset M$ is called a *stratified space* if Λ is the union of disjoint smooth submanifolds Λ^i (*strata*) such that the boundary of each stratum is a finite union of strata of less dimensions.

Let N be a finite dimensional manifold. A smooth map $\xi: M \rightarrow N$ is *transversal* to a smooth submanifold $U \subset N$ if the spaces $\xi_*(T_z M)$ and $T_{\xi(z)}U$ generate $T_{\xi(z)}N$ for each $z \in M$. A smooth map is $\eta: M \rightarrow V$ *transversal* to a stratified space $\Lambda \subset V$ if the the map η is transversal to each stratum of Λ .

Briefly [Theorem 5.4](#) says that any map can be approximated by “a nice map”.

Theorem 5.4 (Multi-jet *transversality* theorem of Thom, see Arnold, Varchenko and Gusein-Zade [[1](#), Section I.2]) *Let M, N be compact smooth manifolds, $\Lambda \subset J_{[l]}^k(M, N)$ be a stratified space. Given a smooth map $\xi: M \rightarrow N$, there is a smooth map $\eta: M \rightarrow N$ such that*


- *the map η is arbitrarily close to ξ with respect to the Whitney topology;*
- *the l -tuple k -jet $j_{[l]}^k \eta: M^l \rightarrow J_{[l]}^k(M, N)$ is transversal to $\Lambda \subset J_{[l]}^k(M, N)$.*

Proof of Claim 4.3 (i) For any critical point of $\text{pr}: S \rightarrow \mathbb{R}$, fix local coordinates $(x, y) \in S$ such that the derivatives $\text{pr}_x = \text{pr}_y = 0$. The closures of the subspaces $\overline{\Sigma_{++} \cup \Sigma_{+-} \cup \Sigma_{--}}$ and $\overline{\Sigma}_2$ from [Definition 2.5](#) can be mapped onto the subspaces of the finite-dimensional spaces $J_{[2]}^1(S, \mathbb{R})$ and $J_{[1]}^2(S, \mathbb{R})$ given by the equations $\text{pr}(x_1, y_1) = \text{pr}(x_2, y_2)$ and $\text{pr}_{xx}\text{pr}_{yy} - \text{pr}_{xy}^2 = 0$, respectively. The resulting subspaces of jets have codimension 1 as preimages of 0 under smooth functions, eg the image of $\overline{\Sigma}_2$ in $J_{[1]}^2(S, \mathbb{R})$ is $(\text{pr}_{xx}\text{pr}_{yy} - \text{pr}_{xy}^2)^{-1}(0)$. Hence the closures $\overline{\Sigma_{++} \cup \Sigma_{+-} \cup \Sigma_{--}}$ and $\overline{\Sigma}_2$ have codimension 1 in the space CS of 2-links.

(ii) If a 2-link is not generic, then either some critical points of the projection $\text{pr}: S \rightarrow \mathbb{R}$ are degenerate or have the same value. The singularities of [Definition 2.5](#) are all multi local codimension 1 singularities of smooth functions $\mathbb{R}^2 \rightarrow \mathbb{R}$; see Arnold, Varchenko and Gusein-Zade [[1](#)].

(iii) By [Theorem 5.4](#) any smooth isotopy of 2-links is a path in CS and can be made transversal to the singular subspace $\overline{\Sigma}$, which has codimension 1 by (i), hence the new path will contain only finitely many isolated singularities of [Definition 2.5](#).

(iv) The normal form of an A_2 -singularity of a function $\mathbb{R}^2 \rightarrow \mathbb{R}$ is $\text{pr}(x, y) = x^2 - y^3$, ie the projection $\text{pr}: S \rightarrow \mathbb{R}$ has the form above in suitable local coordinates $(x, y) \in S$. A 2-link S , its cross-sections around the singularity and the graph of y^3 look like the middle pictures of [Figure 18](#). The versal deformation of an A_2 -singularity is $\text{pr}(x, y; \varepsilon) = x^2 - y^3 + \varepsilon y$ (see Arnold, Varchenko and Gusein-Zade [[1](#)]) ie any smooth deformation of $\text{pr}(x, y)$ can be expressed as $f_1(x, y; \varepsilon) \cdot \text{pr}(f_2(x, y; \varepsilon), f_3(x, y; \varepsilon); f_4(\varepsilon))$, where f_1, f_2, f_3, f_4 are smooth, $f_1(0, 0; 0) \neq 0$, $f_2(x, y; 0) \equiv x$, $f_3(x, y; 0) \equiv y$ and $f_4(0) = 0$.

Figure 18: Transformation of a 2-link near an A_2 -singularity

For $\varepsilon < 0$, a 2-link S , its cross-sections around the singularity and the graph of $y^3 - \varepsilon y$ look like the left pictures of Figure 18. For $\varepsilon > 0$, a 2-link S , its cross-sections around the singularity and the graph of $y^3 - \varepsilon y$ look like the right pictures of Figure 18. For instance, 2-links for $\varepsilon > 0$ have a nondegenerate saddle at $x = 0$, $y = \sqrt{\varepsilon/3}$ and a local extremum at $x = 0$, $y = -\sqrt{\varepsilon/3}$. \square

References

- [1] **VI Arnol'd, AN Varchenko, SM Gusein-Zade**, *Osobennosti differentsiruemykh otobrazhenii [Singularities of differentiable mappings]*, Nauka, Moscow (1982) [MR685918](#)
- [2] **IA Dynnikov**, *Three-page approach to knot theory. Coding and local motions*, Funktsional. Anal. i Prilozhen. 33 (1999) 25–37, 96 [MR1746427](#)
- [3] **IA Dynnikov**, *A three-page approach to knot theory. The universal semigroup*, Funktsional. Anal. i Prilozhen. 34 (2000) 29–40, 96 [MR1756732](#)
- [4] **T Fiedler, V Kurlin**, *Fiber quadrisecants in knot isotopies*, to appear in J. Knot Theory Ramifications [arXiv:math.GT/0701878](#)
- [5] **T Fiedler, V Kurlin**, *A one-parameter approach to links in solid torus* [arXiv:math.GT/0606381](#)
- [6] **RH Fox, JW Milnor**, *Singularities of 2-spheres in 4-space and cobordism of knots*, Osaka J. Math. 3 (1966) 257–267 [MR0211392](#)

- [7] **L H Kauffman**, *Invariants of graphs in three-space*, Trans. Amer. Math. Soc. 311 (1989) 697–710 [MR946218](#)
- [8] **C Kearton, W B R Lickorish**, *Piecewise linear critical levels and collapsing*, Trans. Amer. Math. Soc. 170 (1972) 415–424 [MR0310899](#)
- [9] **V Kurlin**, *Three-page encoding and complexity theory for spatial graphs*, J. Knot Theory Ramifications 16 (2007) 59–102 [MR2300429](#)
- [10] **F J Swenton**, *On a calculus for 2-knots and surfaces in 4-space*, J. Knot Theory Ramifications 10 (2001) 1133–1141 [MR1871221](#)
- [11] **V V Vershinin, V A Kurlin**, *Three-page embeddings of singular knots*, Funktsional. Anal. i Prilozhen. 38 (2004) 16–33, 95 [MR2061788](#)
- [12] **K Yoshikawa**, *An enumeration of surfaces in four-space*, Osaka J. Math. 31 (1994) 497–522 [MR1309400](#)

Department of Mathematical Sciences, Durham University
Durham DH1 3LE, United Kingdom

cherry.kearton@durham.ac.uk, vitaliy.kurlin@durham.ac.uk

<http://www.maths.dur.ac.uk/~dma0ck/>, <http://maths.dur.ac.uk/~dma0vk/>

Received: 7 April 2008 Revised: 7 June 2008