

The topological cyclic Deligne conjecture

PAOLO SALVATORE

Let O be a cyclic topological operad with multiplication. In the framework of the cosimplicial machinery by McClure and Smith, we prove that the totalization of the cosimplicial space associated to O has an action of an operad equivalent to the framed little 2–discs operad.

18D50; 55P48

1 Introduction

The Deligne conjecture, proved by several authors, states that the little 2–discs operad \mathcal{D}_2 acts on the chain level on the Hochschild complex of any associative algebra. The question was motivated by a well known action on the homological level (Gerstenhaber [5]). The natural generalization of the conjecture is about the Hochschild complex of an operad with multiplication, that is an operad with a map from the associative operad. The original statement is recovered taking the endomorphism operad of an associative algebra. McClure and Smith proved a topological version of the Deligne conjecture [17; 18; 19], where the Hochschild complex is replaced by the totalization of a cosimplicial space associated to a topological operad with multiplication. A major example, constructed by Sinha [24], occurs when the operad with multiplication is a version of the little n –discs operad, the Kontsevich operad \mathcal{K}_n . The (homotopy) totalization of \mathcal{K}_n has as factor the space Emb_n of long knots in \mathbb{R}^n for $n > 3$. We proved in [22] that Emb_n itself has an action of the little 2–discs operad. The *cyclic* Deligne conjecture states that the operad of framed little 2–discs $f\mathcal{D}_2$ acts on the chain level on the Hochschild complex of a Frobenius algebra. The conjecture was proved in this form by Kaufmann [11], and Kontsevich–Soibelman [14]. Since the endomorphism operad of a Frobenius algebra is cyclic, the natural generalization of the conjecture claims an action of the chains of $f\mathcal{D}_2$ on the Hochschild complex of a cyclic operad with multiplication. This statement was proved on the homological level by Menichi [20]. The topological counterpart of the statement says that the framed little 2–discs operad $f\mathcal{D}_2$ acts on the totalization of a topological cyclic operad with multiplication. This is the main result of our paper (Theorem 7.7). An equivalent

statement was conjectured by McClure and Smith [19]. Hu [8] stated a related result and applied it to the Hochschild complex of commutative Frobenius algebras.

We will revisit the proof by McClure and Smith of the non-cyclic statement showing that the model of \mathcal{D}_2 they used, denoted here MS , is strongly related to Kaufmann's spineless cacti operad. In fact the elements of MS can be thought of as spineless cacti together with a monotone parametrization. As far as we know this has not been observed before. The parametrizations are necessary if we want to include the constant base point at the 0 level of the operad.

We will then construct an operad fMS , a model of $f\mathcal{D}_2$, whose elements correspond to Voronov's cacti with a parametrization, and prove the main theorem. The proof is conceptually similar to that of the non-cyclic statement, except that the triple defining cyclic operads rather than non-cyclic operads comes into play.

The main motivation of this paper is a joint project of Ryan Budney and the author to endow the space $f\text{Emb}_n$ of framed long knots in \mathbb{R}^n with a meaningful action of the framed little 2-discs operad. The action is expected because $f\text{Emb}_n$ is the (homotopy) totalization of an operad with multiplication equivalent to the framed little n -discs operad $f\mathcal{D}_n$ [22], and $f\mathcal{D}_n$ is equivalent to a cyclic operad [2].

The paper is relatively self-contained. In Section 2 and Section 3 we recall the definitions and give several examples of cosimplicial spaces and (non-symmetric) operads. In Section 4 we define geometrically the operad MS and its framed version fMS . Then we reprove in Section 5 the theorem of McClure and Smith (Theorem 5.4) from this geometric point of view. In Section 6 we recall the definition of the cyclic category and of a cocyclic space. In Section 7 we define cyclic operads, and we prove the main Theorem 7.7. For example the theorem gives an action of fMS on based loop groups. Another important example we recover is the string topology action of fMS on a free loop space, that makes sense according to Voronov [25] in the category of correspondences. This paper grew out of my lecture notes at the Lille Summer School on Algebraic Topology in June 2007. I am grateful to the Laboratoire Painlevé and in particular to Sadok Kallel for support and hospitality.

2 Cosimplicial spaces

We recall the definition of a cosimplicial space. Let Δ be the category with the standard ordered sets $[k] = \{0 < \dots < k\}$ as objects ($k \in \mathbb{N}$) and monotone maps as morphisms. For a fixed $n \in \mathbb{N}$ and $0 \leq i \leq n + 1$ let $\delta_i: [n] \rightarrow [n + 1]$ be the morphism such that $\delta_i(j) = j$ if $j < i$ and $\delta_i(j) = j + 1$ if $j \geq i$.

For $0 \leq i \leq n$ let $\sigma_i: [n+1] \rightarrow [n]$ be the morphism such that $\sigma_i(j) = j$ if $j \leq i$ and $\sigma_i(j) = j-1$ if $j > i$. It is well known that Δ is the category generated by these morphisms under the relations

$$\begin{aligned} \delta_j \delta_i &= \delta_i \delta_{j-1} & i < j \\ \sigma_j \sigma_i &= \sigma_i \sigma_{j+1} & i \leq j \\ \sigma_j \delta_i &= \delta_i \sigma_{j-1} & i < j \\ \sigma_j \delta_i &= \text{id} & j \leq i \leq j+1 \\ \sigma_j \delta_i &= \delta_{i-1} \sigma_j & i > j+1. \end{aligned}$$

Definition 2.1 A *cosimplicial space* is a covariant functor from the category Δ to the category of topological spaces. The image under such functor of a morphism δ_i is a *coface* denoted d^i and the image of a morphism σ_j is a *codegeneracy* denoted s^j .

The definition is dual to that of a *simplicial space*, that is a *contravariant* functor from Δ to the category of topological spaces.

Example 2.2 Let us consider the k -simplex

$$\Delta^k = \{(x_1, \dots, x_k) \mid 0 \leq x_1 \leq \dots \leq x_k \leq 1\}.$$

The collection Δ^* forms a cosimplicial space with the cofaces

$$\begin{aligned} d^i(x_1, \dots, x_k) &= (x_1, \dots, x_i, x_i, \dots, x_k) & 1 \leq i \leq k \\ d^0(x_1, \dots, x_k) &= (0, x_1, \dots, x_k) \\ d^{k+1}(x_1, \dots, x_k) &= (x_1, \dots, x_k, 1) \end{aligned}$$

and with the codegeneracies

$$s^i(x_1, \dots, x_k) = (x_1, \dots, \widehat{x_{i+1}}, \dots, x_k) \quad 0 \leq i \leq k-1.$$

Definition 2.3 The totalization $\text{Tot}(S^*)$ of a cosimplicial space S^* is the space of natural transformations of functors $\Delta^* \rightarrow S^*$. This means that

$$\text{Tot}(S^*) \subset \prod_{k \in \mathbb{N}} \text{Map}(\Delta^k, S^k)$$

is the space of sequences of maps $f^k: \Delta^k \rightarrow S^k$ commuting with the codegeneracies and the cofaces.

The definition is dual to that of the *realization* of a simplicial space.

We recall two standard examples of cosimplicial spaces.

Example 2.4 (The cosimplicial model for the based loop space ΩX) For a given topological space X with base point $*$ $\in X$ let us consider the cosimplicial space ωX such that $(\omega X)^k = X^k$, with

$$\begin{aligned} d^0(x_1, \dots, x_k) &= (*, x_1, \dots, x_k) \\ d^i(x_1, \dots, x_k) &= (x_1, \dots, x_i, x_i, \dots, x_k) \quad 1 \leq i \leq k \\ d^{k+1}(x_1, \dots, x_k) &= (x_1, \dots, x_k, *) \\ s^i(x_1, \dots, x_{k+1}) &= (x_1, \dots, \widehat{x_{i+1}}, \dots, x_{k+1}) \quad 0 \leq i \leq k. \end{aligned}$$

It is well known that the evaluation maps $\Omega X \times \Delta^k \rightarrow X^k$

$$(\gamma, (x_1, \dots, x_k)) \mapsto (\gamma(x_1), \dots, \gamma(x_k))$$

induce by adjointness a homeomorphism $\Omega X \cong \text{Tot}(\omega X)$.

Example 2.5 (The cosimplicial model for the free loop space LX) For a topological space X , let us consider the cosimplicial space lX such that

$$\begin{aligned} (lX)^k &= X^{k+1} = \{(x_0, \dots, x_k) \mid x_i \in X\} \\ d^i(x_0, \dots, x_k) &= (x_0, \dots, x_i, x_i, \dots, x_k) \quad 0 \leq i \leq k \\ d^{k+1}(x_0, \dots, x_k) &= (x_0, \dots, x_k, x_0) \\ s^i(x_0, \dots, x_{k+1}) &= (x_0, \dots, \widehat{x_{i+1}}, \dots, x_{k+1}) \quad 0 \leq i \leq k. \end{aligned}$$

The evaluation map $LX \times \Delta^k \rightarrow X^{k+1}$ sending

$$(\gamma, (x_1, \dots, x_k)) \mapsto (\gamma(0), \gamma(x_1), \dots, \gamma(x_k))$$

induces by adjointness a homeomorphism $LX \cong \text{Tot}(lX)$. Actually, modulo this homeomorphism, the cosimplicial inclusion $i: \omega X \rightarrow lX$ defined by $i^k(x_1, \dots, x_k) = (*, x_1, \dots, x_k)$ corresponds to the inclusion $\Omega X \rightarrow LX$.

3 Operads

We recall some material on operads. For more details we refer to Markl–Shnider–Stasheff [15]. Roughly speaking an element of an operad is an operation with many inputs and one output.

Definition 3.1 A non-symmetric topological operad O is a collection of spaces $O(k)$, $k \in \mathbb{N}$, together with composition maps

$$\circ_t: O(k) \times O(l) \rightarrow O(k+l-1) \quad 1 \leq t \leq k$$

such that for $a \in O(n)$, $b \in O(p)$, $c \in O(q)$

$$\begin{aligned} (a \circ_i b) \circ_{j+p-1} c &= (a \circ_j c) \circ_i b & 1 \leq i < j \leq n \\ a \circ_i (b \circ_j c) &= (a \circ_i b) \circ_{i+j-1} c & 1 \leq i \leq n, 1 \leq j \leq p. \end{aligned}$$

In addition there is a point $\iota \in O(1)$, the *unit*, that is a bi-sided unit for the \circ_t -operations.

The operad is *symmetric* if for each natural number k the symmetric group Σ_k acts on $O(k)$ on the right, compatibly with the composition maps [15], in the sense that for $x \in O(m)$, $y \in O(n)$, $\sigma \in \Sigma_m$ and $\tau \in \Sigma_n$

$$(x\sigma) \circ_i (y\tau) = (x \circ_i y)(\sigma \circ_i \tau)$$

where $\sigma \circ_i \tau \in \Sigma_{m+n-1}$ is the permutation exchanging by σ m consecutive blocks having all one element except the i th having n elements, and acting by τ on the i th block. Here are some examples of topological operads.

Example 3.2 For a topological space X , the *endomorphism* operad $\text{End}(X)$ is the symmetric operad defined by $\text{End}(X)(n) = \text{Map}(X^n, X)$, such that the composition \circ_i is insertion in the i th variable.

For $f \in \text{End}(X)(k)$ and $g \in \text{End}(X)(l)$

$$f \circ_i g \in \text{End}(X)(k+l-1) = \text{Map}(X^{k+l-1}, X)$$

is

$$(f \circ_i g)(x_1, \dots, x_{k+l-1}) = f(x_1, \dots, x_{i-1}, g(x_i, \dots, x_{i+l-1}), x_{i+l}, \dots, x_{k+l-1}).$$

The identity of X is the unit of $\text{End}(X)$.

We say that a space A is *acted on* by a (symmetric) operad O , or that it is an O -*algebra*, if we are given a morphism of (symmetric) operads $O \rightarrow \text{End}(A)$.

The concepts of (symmetric) operads and their algebras can be defined likewise in any (symmetric) monoidal category.

Example 3.3 Dually to Example 3.2, for a topological space X the *coendomorphism* operad $\text{Coend}(X)$ is the symmetric operad defined by

$$\text{Coend}(X)(n) = \text{Map}(X, X^n) = \text{Map}(X, X)^n.$$

Given maps $f = (f_1, \dots, f_k): X \rightarrow X^k$ and $g = (g_1, \dots, g_l): X \rightarrow X^l$ the composition is defined by

$$f \circ_i g = (f_1, \dots, f_{i-1}, g_1 \circ f_i, \dots, g_l \circ f_i, f_{i+1}, \dots, f_k): X \rightarrow X^{k+l-1}.$$


Example 3.4 Let M be a topological monoid. Then there is an operad \underline{M} with $\underline{M}(n) = M^n$. The composition is

$$(x_1, \dots, x_m) \circ_i (y_1, \dots, y_n) = (x_1, \dots, x_{i-1}, x_i y_1, \dots, x_i y_n, x_{i+1}, \dots, x_m).$$

Actually the coendomorphism operad of Example 3.3 is the special case with $M = \text{Map}(X, X)$ endowed with the multiplication $f \cdot g = g \circ f$.

Example 3.5 The associative operad Ass is the operad such that $\text{Ass}(n)$ is a point for each n . This is a non-symmetric operad and is the final object in the category of non-symmetric topological operads. An algebra over Ass is exactly a topological monoid.

Example 3.6 The *little n -discs* operad \mathcal{D}_n by Boardman and Vogt is the main example that motivated historically the introduction of operads [1] (May [16]). The space $\mathcal{D}_n(k)$ is the space of k -tuples of self-maps of the unit n -disc $f_i: D_n \rightarrow D_n$, $i = 1, \dots, k$ obtained by composing translations and dilatations, ie $f_i(z) = \lambda_i z + z_i$, $\lambda_i \in \mathbb{R}_+$, $z_i \in \mathbb{R}^n$, and such that $f_i(D_n) \cap f_j(D_n) = \emptyset$ for $i \neq j$. This is a symmetric suboperad of the coendomorphism operad of the disc D_n . We can associate a picture to an element of \mathcal{D}_n drawing the images of the maps.



The n -fold loop space $\Omega^n(X) \cong \text{Map}(D_n, S^{n-1}; X, x_0)$ on a based space (X, x_0) is an algebra over the symmetric operad \mathcal{D}_n . The structure map is adjoint to the map

$$\theta_n: \mathcal{D}_n(k) \times \Omega^n(X)^k \rightarrow \Omega^n(X)$$

sending $(f_1, \dots, f_k; \omega_1, \dots, \omega_k)$, for $f_i: D_n \rightarrow D_n$ and $\omega_j: D_n \rightarrow X$, to the n -fold loop

$$\theta_n(f_1, \dots, f_k; \omega_1, \dots, \omega_k): D_n \rightarrow X$$

coinciding with $\omega_i \circ (f_i)^{-1}$ on the image $f_i(D_n)$ for $i = 1, \dots, k$ and sending any point that is not in the image of any f_i to the base point $x_0 \in X$.

A partial converse holds, the *recognition principle*.

Proposition 3.7 (Boardman–Vogt [1] May [16]) *Let A be an algebra over \mathcal{D}_n , with $n \geq 1$. If A is connected then A is weakly homotopy equivalent to a n -fold loop space.*

More generally this is true if A is *group-like*. This means that the set of components $\pi_0(A)$, that is always a monoid by the action of \mathcal{D}_n , is also a group.

Example 3.8 The *framed little n -discs operad* $f\mathcal{D}_n$ introduced by Getzler in [6] contains \mathcal{D}_n , and is defined similarly as \mathcal{D}_n except that the self-maps in the definition can be composed by oriented rotations.


Precisely $f\mathcal{D}_n(k)$ is the space of k -tuples of self-maps $f_i: D_n \rightarrow D_n$, $i = 1, \dots, k$ such that $f_i(z) = \lambda_i A_i z + z_i$ with $\lambda_i \in \mathbb{R}_+$, $A_i \in SO(n)$, $z_i \in \mathbb{R}^n$, and such that $f_i(D_n) \cap f_j(D_n) = \emptyset$ for $i \neq j$. Also $f\mathcal{D}_n$ is a suboperad of the coendomorphism operad of the disc D_n . There is an obvious inclusion of operads $\mathcal{D}_n \subset f\mathcal{D}_n$.

The operad $f\mathcal{D}_n$ acts on the n -fold loop space $\Omega^n(X)$ of any space X with based $SO(n)$ -action.

The recognition principle for algebras over $f\mathcal{D}_n$ is the following.

Proposition 3.9 (Salvatore–Wahl [23]) *Let A be an algebra over $f\mathcal{D}_n$. If A is group-like then it is weakly equivalent to $\Omega^n(X)$, where X has a based $SO(n)$ -action, via a zig-zag of based $SO(n)$ -equivariant maps.*

Definition 3.10 (The language of trees) Here we follow the terminology of the book by Markl, Shnider and Stasheff [15], with the important difference that we work with non-symmetric rather than symmetric operads, and with planar rather than non-planar trees. An important point about non-symmetric operads is that it makes sense to compose elements of the operad along a *rooted planar tree*. This is a planar graph with no cycles, where we allow some open half-edges. The *root* is one of these half-edges, and the remaining ones are the *leaves*. All other edges are called *internal* edges. There is a unique structure of directed graph on a rooted planar tree such that there is a directed path from each point to the root. Given a rooted planar tree T , the arity $|v|$ of a vertex v is the number of incoming edges. The counterclockwise ordering induces an ordering on the incoming edges of a vertex. It is customary to draw the tree so that the root is on the bottom and the edges are directed downward. Here is an example of a tree with 4 vertices of arity $|x| = 0$, $|y| = 1$, $|z| = 2$ and $|t| = 3$.



Given a topological operad O , let us consider the product $O(T) = \prod_v O(|v|)$ indexed over all vertices v of T . This product can be thought of as the space of trees of shape T with vertices labeled by the operad O . If a tree T' is obtained from T by contracting an internal edge $e: v \rightarrow w$, and e is the i th incoming edge of w , then the composition map

$$\circ_i: O(|w|) \times O(|v|) \rightarrow O(|w| + |v| - 1)$$

induces a map $O(T) \rightarrow O(T')$. The axioms of Definition 3.1 guarantee that for iterated contractions the result does not depend on the order. In the end there is a well defined composition map $O(T) \rightarrow O(n)$, where n is the number of leaves of the tree T . This can be expressed by the notion of a *triple*. We recall that a triple (F, μ, u) in a category \mathcal{C} is a monoid in the monoidal category of endofunctors of \mathcal{C} , ie a functor $F: \mathcal{C} \rightarrow \mathcal{C}$ together with natural transformations $\mu: F \circ F \rightarrow F$ and $u: \text{id}_{\mathcal{C}} \rightarrow F$ satisfying associative and unital properties. An algebra over the triple is an object A of \mathcal{C} with a morphism $\rho: F(A) \rightarrow A$ compatible in an appropriate sense with μ and u . In our case \mathcal{C} is the category of sequences of topological spaces $X = (X(n))$ ($n \in \mathbb{N}$) with a distinguished base point $\iota \in X(1)$. The functor F sends a sequence X to a sequence $F(X)$, where $F(X)(n)$ is the space of rooted planar trees on n leaves of all possible shapes, labelled by elements of X , modulo the relation deleting arity 1 vertices labeled by the base point $\iota \in X(1)$. The natural transformation $\mu: F \circ F \rightarrow F$ sends a tree T with labels that are themselves labeled trees to the labeled tree obtained cutting a neighbourhood of each vertex of T and gluing in its label. The unit transformation $u: \text{id} \rightarrow F$ identifies a point to a corolla labeled by the same point, where a *corolla* is a tree with a single vertex. An algebra over this triple is exactly a non-symmetric topological operad. In other words $F(X)$ is the sequence underlying the free operad generated by X .

4 Cacti operads

We claim that the operads used by various authors dealing with the Deligne conjecture are very similar.


We recall that two topological operads A, B are said to be *equivalent* if there is a zig-zag of operad morphisms that are levelwise homotopy equivalences connecting A and B . An E_n operad is an operad equivalent to the little n -discs \mathcal{D}_n . Voronov introduced in [25] the *cacti operad* Cacti that is equivalent to the framed little 2-discs operad $f\mathcal{D}_2$ if we forget constants, i.e. the arity 0 part. A variation of the cacti operad that is instead equivalent to the little 2-discs \mathcal{D}_2 , if we forget constants, is the *spineless cacti operad*, introduced by Kaufmann [10].

We will describe in this section an E_2 operad constructed by McClure and Smith in [18], that we call MS . This operad is closely related to the spineless cacti operad.

The space $MS(n)$ splits as a product of a finite CW-complex $\mathcal{F}(n)$ and a contractible factor. We will give a geometric description of $\mathcal{F}(n)$.

Definition 4.1 An element $x \in \mathcal{F}(n)$ is a partition of the circle S^1 into n closed 1-manifolds $I_j(x)$ ($j = 1, \dots, n$) with equal length (=measure), and pairwise disjoint interiors, such that there is no cyclically ordered 4-tuple (z_1, z_2, z_3, z_4) in S^1 with $z_1, z_3 \in \overset{\circ}{I}_j(x)$, $z_2, z_4 \in \overset{\circ}{I}_k(x)$ and $j \neq k$.

The figure shows an element of $\mathcal{F}(4)$.



We can define on $\mathcal{F}(n)$ a topology induced by a metric such that two partitions $x, y \in \mathcal{F}(n)$ are close if the manifolds $I_j(x)$ and $I_j(y)$ are close for each j . For example the metric $d(x, y) = \sum_{j=1}^n l(I_j(x) \setminus \overset{\circ}{I}_j(y))$ does the job, where l denotes the measure. The symmetric group Σ_n acts on $\mathcal{F}(n)$ by reindexing the labels.

Clearly $\mathcal{F}(1)$ is a point. Let $\pi: I \rightarrow I/\partial I$ be the projection. The space $\mathcal{F}(2)$ is Σ_2 -equivariantly homeomorphic to a circle with antipodal Σ_2 -action. The homeomorphism $I/\partial I \cong \mathcal{F}(2)$ sends $\pi(x)$ into the pair of intervals $\{\pi[x, x + \frac{1}{2}], \pi[x + \frac{1}{2}, x + 1]\}$.

The cell structure The pullback of a partition $x \in \mathcal{F}(n)$ to I via $\pi: I \rightarrow I/\partial I \cong S^1$ determines a partition of I into say k intervals labelled by natural numbers between 1 and n . Any component of $\pi^{-1}(I_j(x))$ is labelled by j . This gives an ordered sequence of boundary points $0 = x_0 < x_1 < \dots < x_{k-1} < x_k = 1$, together with the sequence of labels $X = (X_1, \dots, X_k)$, such that X_i is a label of the interval $[x_{i-1}, x_i]$.

It follows that $\mathcal{F}(n)$ is a regular CW-complex with one cell for each fixed sequence of labels X . The closed cell indexed by a sequence X of length k corresponds to the subspace of the $(k-1)$ -simplex containing the elements

$$0 = x_0 \leq x_1 \leq \dots \leq x_{k-1} \leq x_k = 1$$

such that

$$\sum_{i|X_i=j} (x_i - x_{i-1}) = 1/n$$

for each $j = 1, \dots, n$. This subspace is homeomorphic to the product of simplices $\prod_{j=1}^n \Delta^{d(j)}$, where $d(j)$ is one less than the number of components (intervals) of $\pi^{-1}(I_j(x))$.

The upshot is that cells of $\mathcal{F}(n)$ correspond to those finite sequences $X = (X_i)$ with values in $\{1, \dots, n\}$ such that:

- (1) all values between 1 and n appear in X ,
- (2) two adjacent values in X are distinct,
- (3) the sequence X does not have a subsequence of the form $\{i, j, i, j\}$ with $i \neq j$.


Given a partition $x \in \mathcal{F}(n)$ and an index j , let us consider the quotient space of S^1 under the relation that identifies points in the same component (closed interval) of $S^1 \setminus I_j(x)$. The result is a circle with a base point, the image of the base point of S^1 . If we rescale this circle by the factor n and identify it to S^1 , then we see the quotient map as a based self-map $\pi_j: S^1 \rightarrow S^1$.

We say that $c(x) := (\pi_1, \dots, \pi_n): S^1 \rightarrow (S^1)^n$ is the *cactus map* and its image is the *cactus* C_x associated to x .


The map $c: \mathcal{F}(n) \rightarrow \text{Map}(S^1, (S^1)^n)$ is an embedding, but the image does not form a suboperad of the coendomorphism operad $\text{Coend}(S^1)$ of Example 3.3.

The cactus C_x is the union of n circles, the *lobes* $S_j(x)$ ($j = 1, \dots, n$), that are the images under $c(x)$ of the submanifolds $I_j(x)$. The base point $P \in C_x$ is the image of the base point $1 \in S^1$ under the cactus map $c(x): S^1 \rightarrow C_x$.

Here is an example of a cactus with the indexes of the lobes.



Definition 4.2 (The associated tree) One can associate to each partition $x \in \mathcal{F}(n)$ a rooted planar tree (constant on open cells) constructed as follows. The tree is the graph that has a vertex for each lobe (ie for each number between 1 and n), a vertex for each intersection point of distinct lobes, and an edge between the vertex of an intersection point and the vertex of each lobe it belongs to. Let $P \in C_x$ be the base point. If P is an intersection point then the *root vertex* is its vertex. Otherwise P belongs to a unique lobe and the root vertex is the vertex of that lobe. The *root* is an additional half-open edge incident to the root vertex. The tree is a ribbon graph with a cyclic ordering of the edges incident to a given vertex. The ordering is induced by the standard orientation of the circle via the map $c(x): S^1 \rightarrow C_x$. The tree admits a planar embedding such that the cyclic ordering is compatible with the standard orientation of the plane (Voronov [25]). It is possible to embed the cactus C_x into the plane, in a way that this is compatible with the embedding of the tree on the intersection points, as shown in the next figure. The tree is the union of corollas inscribed in the lobes.



More operads McClure and Smith constructed in [17] first an operad \mathcal{C}' that is equivalent to little 2-discs \mathcal{D}_2 away from the arity 0 part, because $\mathcal{C}'(0)$ is empty but $\mathcal{D}_2(0)$ is a point. As a Σ_n -space $\mathcal{C}'(n) = \mathcal{F}(n) \times \Delta_{n-1}$.

The space $\mathcal{C}'(n)$ is naturally homeomorphic to a space of partitions defined similarly as in Definition 4.1, except that one drops the condition that all manifold must have the same length. In this setting there is also a cactus map $\mathcal{C}' \rightarrow \text{Coend}(S^1)$ that is an embedding onto a symmetric suboperad of $\text{Coend}(S^1)$ and defines an operad structure on \mathcal{C}' .

The spineless cacti operad Cact by Kaufmann [10] is actually a pseudo-operad, in the sense that is associative, non-unital, and is isomorphic to $\mathcal{C}' \times R$, where R is the constant pseudo-operad $R(n) = \mathbb{R}_+$ with composition $\mathbb{R}_+ \times \mathbb{R}_+ \rightarrow \mathbb{R}_+$ equal to the projection onto the first factor. The extra parameter corresponds to the length of the parametrizing circle in [10].

McClure and Smith extended in [17] the operad \mathcal{C}' , that has $\mathcal{C}'(0)$ empty, to an operad \mathcal{C} equivalent to the little 2-discs operad such that $\mathcal{C}(0)$ is a base point. The operad \mathcal{C} is the smallest suboperad of $\text{Coend}(S^1)$ containing \mathcal{C}' and the base point in arity 0.

Later on McClure and Smith defined a larger operad that is denoted \mathcal{D}_2 in [18]. We present a geometric construction of such operad.

Definition 4.3 Let $\text{Mon}(I, \partial I)$ be the space of self-maps of the circle $S^1 \cong I/\partial I$ induced by weakly monotone self-maps of the interval I restricting to the identity on the boundary ∂I . We define $MS(n) = \mathcal{F}(n) \times \text{Mon}(I, \partial I)$ for $n > 0$. We also define $MS(0)$ to be a point.

Remark 4.4 The space $\text{Mon}(I, \partial I)$ is homeomorphic to the contractible space $\text{Tot}(\Delta^*)$, ie the monoid of cosimplicial endomorphisms of the cosimplicial space Δ^* . The homeomorphism $T: \text{Mon}(I, \partial I) \rightarrow \text{Tot}(\Delta^*)$ sends $f: I \rightarrow I$ to the sequence of maps $T(f)^k: \Delta^k \rightarrow \Delta^k$ with

$$T(f)^k: (0 \leq x_1 \leq \dots \leq x_k \leq 1) \mapsto (0 \leq f(x_1) \leq \dots \leq f(x_k) \leq 1).$$

There is an embedding $\zeta_n: MS(n) \rightarrow \text{Coend}(S^1)(n)$ sending, for $n > 0$, $x \in \mathcal{F}(n)$ and $f \in \text{Mon}(I, \partial I)$ to the composition $c(x) \circ f: S^1 \rightarrow (S^1)^n$. The following characterization is immediate.

Proposition 4.5 *The image $\zeta_n(MS(n))$ is the space of based maps $g: S^1 \rightarrow (S^1)^n$ such that*

- (1) $\pi_i \circ g$ is a weakly monotone degree 1 map for each $i = 1, \dots, n$,
- (2) there is a partition of S^1 into closed intervals intersecting at the boundary such that on each interval $\pi_i \circ g$ is constant for all indexes i except one special index,

- (3) the counterclockwise sequence of special indices does not contain a subsequence of the form $\{i, j, i, j\}$ with $i \neq j$.

It follows that the image of the embedding ζ is a symmetric suboperad of $\text{Coend}(S^1)$. This gives an operad structure on MS .

Proposition 4.6 *The operad MS is isomorphic to the E_2 operad denoted \mathcal{D}_2 in [18].*

A proof will be presented in the appendix.

We build next an operad fMS that is equivalent to the framed little 2–discs. It is related to MS in the same way as Voronov’s cacti operad is related to Kaufmann’s spineless cacti operad.

Definition 4.7 Let us consider the Σ_n –space $fMS(n) = MS(n) \times (S^1)^n$ for any natural number n . There is an embedding $c': fMS(n) \rightarrow \text{Coend}(S^1)(n)$ sending

$$(x, f, (z_1, \dots, z_n)) \mapsto (L_{z_1} \times \dots \times L_{z_n}) \circ c(x) \circ f$$

where $L_z: S^1 \rightarrow S^1$ is multiplication by $z \in S^1$. A map $g: S^1 \rightarrow (S^1)^n$ belongs to the image $c'(fMS(n))$ if and only if it is an *unbased* map satisfying the conditions 1,2,3 of Proposition 4.5. It follows that the image of c' is a symmetric suboperad of $\text{Coend}(S^1)$, and this defines an operad structure on fMS .

In order to prove that fMS is equivalent to $f\mathcal{D}_2$ we need a lemma. The lemma is exposed as [3, Theorem 2.2.2] and follows from [23, Theorem 7.3], that is a generalization of a similar result by Fiedorowicz for the little 2–discs operad.

Lemma 4.8 *Let O be a symmetric operad together with a Σ_k –equivariant homotopy equivalence $O(k) \simeq f\mathcal{D}_2(k)$ for each k , and together with an operad map $E \rightarrow O(k)$ from an E_1 –operad E . Then O is equivalent to $f\mathcal{D}_2$.*

Proposition 4.9 *The operad fMS is equivalent to the framed little 2–discs operad $f\mathcal{D}_2$.*

Proof We know from Proposition 4.6 that MS is an E_2 –operad, and this gives for each k a Σ_k –equivariant homotopy equivalence

$$fMS(k) = MS(k) \times (S^1)^k \simeq \mathcal{D}_2(k) \times (S^1)^k = f\mathcal{D}_2(k).$$

In addition $MS \subset fMS$ contains an E_1 –operad MS_1 defined in the appendix. By Lemma 4.8 this concludes the proof. \square

For $y = (x, f, (z_1, \dots, z_n)) \in fMS(n)$ we denote by C_y and call *cactus* the image of $(L_{z_1} \times \dots \times L_{z_n}) \circ c(x): S^1 \rightarrow (S^1)^n$. We also call *lobe* and denote $S_j(y)$ the image of $I_j(x)$ via the same map. The lobe has a standard identification to S^1 via the projection $(S^1)^n \rightarrow S^1$ onto the j th coordinate. The base point of the cactus is the image of the base point of S^1 . In general the base points of the lobes are not intersection points or the base point of the cactus as in the spineless case.

The connection to Voronov's cacti operad \mathbf{Cacti} , that actually is a pseudo-operad, is as follows. The operad fMS contains an operad $f\mathcal{C}'$ with $f\mathcal{C}'(n) = \mathcal{C}'(n) \times (S^1)^n$. In particular $f\mathcal{C}'(0)$ is empty. The inclusion $f\mathcal{C}'(n) \rightarrow fMS(n)$ is a Σ_n -equivariant homotopy equivalence for all $n > 0$, and there is an isomorphism of pseudo-operads $\mathbf{Cacti} \cong f\mathcal{C}' \times R$.

5 The non-cyclic Deligne conjecture

We shall reprove in this section the result by McClure and Smith. We need the following definition.

Definition 5.1 A topological operad with multiplication O is a non-symmetric operad O together with a morphism $\text{Ass} \rightarrow O$ from the associative operad.

Equivalently O is a non-symmetric operad in the monoidal category of based spaces with the cartesian product as tensor product. Also equivalently O is a topological operad with points $m \in O(2)$ (the multiplication) and $u \in O(0)$ (the unit of the multiplication) such that

$$\begin{aligned} m \circ_1 m &= m \circ_2 m \in O(3) \\ m \circ_1 u &= m \circ_2 u = \iota \in O(1). \end{aligned}$$

Remark 5.2 There is a triple F_* in the category of sequences of based spaces such that an operad with multiplication is exactly an algebra over F_* . The triple F_* is closely related to the triple F of Definition 3.10. Namely, for a sequence X , $F_*(X)$ is the quotient of $F(X)$ under the identification relation induced by collapsing internal edges with both vertices labeled by base points.

We can relate now cosimplicial spaces and operads.

Definition 5.3 An operad O with multiplication defines a cosimplicial space O^* sending $[k]$ to $O(k)$. The coface operator $d^i: O(k) \rightarrow O(k+1)$ is defined by

$$\begin{aligned} d^i(x) &= x \circ_i m \text{ for } 1 \leq i \leq k \\ d^0(x) &= m \circ_2 x \\ d^{k+1}(x) &= m \circ_1 x. \end{aligned}$$

The codegeneracies $s^i: O(k) \rightarrow O(k-1)$ are defined by

$$s^i(x) = x \circ_{i+1} u.$$

Theorem 5.4 (McClure–Smith [18]) *Let O be an operad with multiplication. Then the totalization $\text{Tot}(O^*)$ admits the action of the E_2 operad MS .*

Proof

We explain geometrically the action of MS on $\text{Tot}(O^*)$. We need to describe the action of MS on $\text{Tot}(O^*)$ via maps

$$\theta_m: MS(m) \times \text{Tot}(O^*)^m \rightarrow \text{Tot}(O^*).$$

Given $(x, f) \in MS(m) = \mathcal{F}(m) \times \text{Tot}(\Delta^*)$ and m cosimplicial maps

$$a_1^*, \dots, a_m^* \in \text{Tot}(O^*)$$

we want to define a cosimplicial map

$$a^* = \theta_m(x, f; a_1^*, \dots, a_m^*) \in \text{Tot}(O^*),$$

so we need compatible maps $a^k: \Delta^k \rightarrow O(k)$ for every non-negative integer k . A point of the simplex Δ^k is a sequence of points of the unit interval

$$0 = x_0 \leq x_1 \leq \dots \leq x_k \leq 1.$$

The cactus map induces a curve

$$\gamma = c(x) \circ f \circ \pi: I \rightarrow C_x \subset (S^1)^m.$$


We construct a rooted planar tree T associated to these data. We start with the tree associated to $x \in \mathcal{F}(m)$ in Definition 4.2 with its root removed. This tree has

- a vertex for each lobe,
- a vertex for each intersection point,
- edges between vertices of intersection points and vertices of their lobes.

We call the points of the form $\gamma(x_j)$ ($0 \leq j \leq k$) *special points*. For each special point that is not an intersection point we add a new vertex, and an edge connecting it to the vertex of the lobe containing $\gamma(x_j)$.

Finally for each point of the form x_j ($0 \leq j \leq k$) we add a leaf (half-open edge) incident to the vertex of $\gamma(x_j)$. The edge incident to the vertex of the base point $\gamma(x_0)$ is the root. The difference with the tree in Definition 4.2 is that we subdivide the former root inserting a vertex.

The tree T is a planar rooted tree: the ordering of the edges incident to a vertex is induced by the standard orientation of the interval via γ and by the ordering of the indices. Both the tree T and the cactus C_x can be embedded together consistently into the plane, so that the embeddings are compatible on points of the form $\gamma(x_j)$ and on intersection points. An example with $k = 6$ and $m = 4$ is given in the next figure.



Next we label each vertex v of T by an element of the operad of the same arity.

If v corresponds to an intersection point of lobes, or to a point of the form $\gamma(x_j)$, then we label it by the base point of $O(|v|)$. Let $\text{Int} \subset C_x$ be the set of all intersection points.

If v corresponds to a lobe S_j , then let us take the set

$$H_j = (\text{Int} \cup \gamma(\{x_1, \dots, x_k\})) \cap (S_j - *)$$

of special points and intersection points in the lobe that are not its base point.


Suppose that H_j has l_j points. The arity of v is exactly $|v| = l_j$. There is a standard identification of the lobe S_j with S^1 via inclusion $S_j(x) \subset C_x \subset (S^1)^m$ followed by projection onto the j th factor. Pulling back H_j via $I \rightarrow I/\partial I \cong S^1 \cong S_j$ gives l_j points in the unit interval I , ie a point y_j of the simplex Δ^{l_j} . Actually y_i is

in the interior of the simplex. The cosimplicial map $a_j \in \text{Tot}(O^*)$ in degree l_j $a_j^{l_j}: \Delta^{l_j} \rightarrow O(l_j)$ evaluated on y_j gives an element of the operad $a_j^{l_j}(y_j) \in O(l_j)$ that we assign as label of v . In the terminology of Definition 3.10 our labelled tree is some $\Gamma \in O(T)$. The generalised composition $O(T) \rightarrow O(k)$ applied to the labelled tree Γ defines the desired element $a^k(x_1, \dots, x_k) \in O(k)$.

The resulting map $a^*: \Delta^* \rightarrow O^*$ is a cosimplicial map essentially because a_1^*, \dots, a_m^* are all cosimplicial maps. The maps θ_m define a MS -algebra structure on $\text{Tot}(O^*)$. To see this we need to look at the labelled trees whose composition defines the structure maps, and use the fact that the ordering of composition by contracting edges is not relevant. More precisely we are using the fact that O is an algebra over the triple F_* of Remark 5.2. □

Example 5.5 Suppose that $x \in MS(2)$, so that C_x has two lobes, that the base point is in S_1 , and that $k = 1$. We wish to compute the value of the map $a^1: \Delta^1 \rightarrow O(1)$ on $x_1 \in I$. We suppose that the intersection point $Q = S_1 \cap S_2$ coincides with the special point $Q = \gamma(x_1)$. We also assume that the curve γ at x_1 goes from the lobe S_1 into S_2 , ie $\gamma([0, x_1]) \subset S_1$. Then $l_1 = 1, l_2 = 0$ and y_1 is the point mapping to Q via $I \rightarrow S^1 \cong S_1$. It follows that

$$a^1(x_1) = (a_1^1(y_1) \circ_1 m_2) \circ_2 a_2^0(*).$$



Theorem 5.4 and the recognition principle (Proposition 3.7) immediately imply the following corollary.

Corollary 5.6 *If O is an operad with multiplication and $\text{Tot}(O^*)$ is group-like then $\text{Tot}(O^*)$ has the weak homotopy type of a double loop space.*

Example 5.7 Let M be a topological monoid. The operad \underline{M} of Example 3.4 has a multiplication by choosing $m_2 = (1, 1) \in M^2$, and $m_0 = * \in M^0$. Theorem 5.4 tells us that the operad MS acts on $\text{Tot}(\omega M) \cong \Omega M \simeq \Omega^2(BM)$. This is consistent with Corollary 5.6.

Example 5.8 (Knots from operads) We recall that a *long knot* is an embedding $\mathbb{R} \rightarrow \mathbb{R}^n$ that is fixed near infinity, and a *framed long knot* is a long knot with a trivialization of its tubular neighbourhood that is fixed near infinity. The Kontsevich operad \mathcal{K}_n is an operad with multiplication that is equivalent to the little discs \mathcal{D}_n , which instead does not admit a multiplication. Sinha proved in [24], using Goodwillie calculus, that the *homotopy totalization* $\widetilde{\text{Tot}}(\mathcal{K}_n^*)$ is weakly equivalent to $\text{Emb}_n \times \Omega^2 S^{n-1}$ for $n > 3$, where Emb_n is the space of long knots in \mathbb{R}^n . The homotopy totalization of a cosimplicial space is weakly equivalent to the homotopy limit of the cosimplicial space as a diagram. One cannot apply directly Theorem 5.4 to the homotopy totalization, but there is a derived version of Theorem 5.4 by McClure and Smith ([19, Theorem 15.3]) constructing an E_2 operad that acts on the homotopy totalization of any operad with multiplication. By Corollary 5.6 the space $\text{Emb}_n \times \Omega^2 S^{n-1}$, which is connected by Whitney's theorem, is a double loop space for $n > 3$. We proved in [22] that Emb_n itself is a double loop space for $n > 3$. We also proved that the space of framed long knots $f\text{Emb}_n$ is a double loop space, because it is weakly equivalent to $\widetilde{\text{Tot}}(f\mathcal{K}_n^*)$, where $f\mathcal{K}_n$ is the *framed* Kontsevich operad, that is equivalent to the framed little n -discs $f\mathcal{D}_n$.

6 The cyclic category

The cyclic category Λ was introduced by Connes, and is an extension of the category Δ by adding for each n an automorphism $\tau_n: [n] \rightarrow [n]$ of order $n + 1$, such that

$$\begin{aligned}\tau_n \delta_i &= \delta_{i-1} \tau_{n-1} & 1 \leq i \leq n \\ \tau_n \sigma_i &= \sigma_{i-1} \tau_{n+1} & 1 \leq i \leq n.\end{aligned}$$

As a consequence of the axioms $\tau_n \delta_0 = \delta_n$ and $\tau_n \sigma_0 = \sigma_n (\tau_{n+1})^2$. We refer to Drinfeld [4] for a discussion of the cyclic category.

A *cocyclic space* is a covariant functor from Λ to the category of topological spaces. In particular by restriction to $\Delta \subset \Lambda$ a cocyclic space has an underlying cosimplicial space, that we denote with the same name. We denote by t_n the image of τ_n under the functor defining a cocyclic space.

Proposition 6.1 (Jones [9]) *Let C be a cocyclic space. Then the space $\text{Tot}(C)$ has a natural circle action.*

Example 6.2 ([9]) The cosimplicial model of the free loop space lX of Example 2.5 has a compatible cocyclic space structure defined by maps $t_n: X^{n+1} \rightarrow X^{n+1}$ with

$$t_n(x_0, \dots, x_n) = (x_1, \dots, x_n, x_0).$$

The resulting circle action on $\text{Tot}(lX)$, modulo the identification $\text{Tot}(lX) \cong LX$, coincides with the reparameterization action on the free loop space $S^1 \times LX \rightarrow LX$ defined by $(z, \gamma) \mapsto \gamma \circ L_z$, where $L_z: S^1 \rightarrow S^1$ is multiplication by z .

Example 6.3 Let G be a topological group. The cosimplicial space ωG of Example 2.4 has a compatible cocyclic space structure defined by maps $t_n: G^n \rightarrow G^n$ with

$$t_n(g_1, \dots, g_n) = (g_1^{-1} g_2, \dots, g_1^{-1} g_n, g_1^{-1}).$$

The homeomorphism $\Omega G \cong \text{Tot}(\omega G)$ is S^1 -equivariant, where the circle action $\theta: S^1 \times \Omega G \rightarrow \Omega G$ on $\Omega G = \text{Map}_*(S^1, G)$

$$\theta(z, \gamma)(w) = \gamma(z)^{-1} \gamma(zw)$$

is given by normalized reparameterization.

Example 6.4 The cocyclic space Λ^* is defined so that $\Lambda^n \subset (S^1)^n$ is the subspace of those $(n + 1)$ -tuples (x_0, \dots, x_n) that are cyclically ordered with respect to the counterclockwise cyclic ordering.

Similarly as for Δ^* (Example 2.2) the i th coface doubles the coordinate x_i , the i th codegeneracy removes the coordinate x_{i+1} and the automorphism t_n acts cyclically on the coordinates by

$$t_n(x_0, \dots, x_n) = (x_1, \dots, x_n, x_0).$$

There is a homeomorphism $\Delta^n \times S^1 \cong \Lambda^n$ sending

$$((x_1, \dots, x_n), z) \mapsto (z, z + x_1, \dots, z + x_n)$$

for $z \in \mathbb{R}/\mathbb{Z} \cong [0, 1]/\{0, 1\} \cong S^1$. The inclusion $\Delta^n \times \{1\} \rightarrow \Lambda^n$ makes Δ^* into a cosimplicial subspace of Λ^* .

The totalization $\text{Tot}(X^*)$ of a cocyclic space X^* is naturally isomorphic to the space $\text{Mor}(\Lambda^*, X^*)$ of natural transformations $\Lambda^* \rightarrow X^*$ of cocyclic spaces. In particular for any space Y there is a homeomorphism $LY \cong \text{Mor}(\Lambda^* \rightarrow lY)$ sending a loop γ to the sequence of maps $(x_0, \dots, x_k) \mapsto (\gamma(x_0), \dots, \gamma(x_k))$. Jones' circle action on $\text{Tot}(X^*)$ of Proposition 6.1 is induced by the circle action on $\Lambda^* \cong S^1 \times \Delta^*$.

7 Cyclic operads and the Deligne conjecture

Roughly speaking a cyclic operad exchanges the roles of inputs and outputs. We give the definition by default in the non-symmetric sense.

Definition 7.1 A cyclic operad O is an operad such that $O(n)$ has an action of the cyclic group \mathbb{Z}_{n+1} for each n , with the generator acting by $t_n: O(n) \rightarrow O(n)$, such that for $f \in O(m)$, $g \in O(n)$

$$\begin{aligned} t_{m+n-1}(f \circ_1 g) &= t_n(g) \circ_n t_m(f) \\ t_{m+n-1}(f \circ_i g) &= t_m(f) \circ_{i-1} g \quad 1 < i \leq m \\ t_1(\iota) &= \iota. \end{aligned}$$

A *symmetric* cyclic operad is a cyclic operad such that $O(n)$ admits the action of the symmetric group Σ_{n+1} , extending the action of \mathbb{Z}_{n+1} and of Σ_n .

The simplest example of a cyclic operad is the associative operad Ass with trivial action of the cyclic groups.

Example 7.2 A non trivial example of non-symmetric cyclic operad is the A_∞ operad by Stasheff [15]. In particular \mathbb{Z}_5 acts on the pentagon $A_\infty(4)$ by rotations.

Example 7.3 An example of symmetric cyclic operad is the operad \mathcal{M} of moduli space of Riemann surfaces with boundary. Here $\mathcal{M}(n)$ is the moduli space of the surfaces with $n + 1$ boundary components. The operadic composition goes by gluing surfaces at the boundary components, and Σ_{n+1} acts on the labels of the components. The genus zero part \mathcal{M}_0 is a cyclic operad equivalent to the framed little discs operad $f\mathcal{D}_2$.

Example 7.4 Budney defines in [2] the operad of *conformal balls* \mathcal{CB}_n that is symmetric cyclic, and equivalent to the framed little n -discs $f\mathcal{D}_n$.

Definition 7.5 (The triple for cyclic operads) Getzler and Kapranov described in [7] the triple for symmetric cyclic operads. We describe the analog triple in the non-symmetric case. A *cyclic sequence* is a sequence of spaces $X(n)$, $n \in \mathbb{N}$ together with an action of \mathbb{Z}_{n+1} on $X(n)$ for each n . We also assume that $X(1)$ has a base point in $X(1)$ fixed by the \mathbb{Z}_2 -action. There is a triple F^+ on the category of cyclic sequences. Given a cyclic sequence X , if we forget the actions of the cyclic groups then $F^+(X)$ agrees with $F(X)$. By definition $F(X)(n)$ is the space of rooted labelled trees with n leaves, or equivalently with $n + 1$ half edges. The group action of \mathbb{Z}_{n+1} on $F(X)(n)$ switches the roots. The action of $[m] \in \mathbb{Z}_{n+1}$ on some $T \in F(X)(n)$ sets the new root as the m th half-edge in the counterclockwise cyclic ordering after the old root. The new root defines a new structure of directed graph on the labelled tree T . For each vertex v of arity p , or equivalently of valence $p + 1$, there is a unique outgoing edge v_o (respectively v_n) in the old (respectively new) directed graph structure. Suppose

that v_n is the q th edge after v_o in the counterclockwise ordering of the edges incident to v . The new label of v is the result of the action of $[q] \in \mathbb{Z}_{p+1}$ on the old label in $X(p)$. Given a cyclic operad O' , by virtue of Definition 7.1, the action of the triple $F(O') \rightarrow O'$ respects the action of the cyclic groups. This shows that O' is an algebra over the triple F^+ . Vice versa an algebra O over the triple F^+ is a cyclic operad. The Definition 7.1 is satisfied by restricting the action $F(O) \rightarrow O$ to labelled trees with one internal edge.

Definition 7.6 A cyclic operad with multiplication is an operad O together with a morphism of cyclic operads $\text{Ass} \rightarrow O$.

Equivalently O is an operad with multiplication $m_2 \in O(2)$, and a cyclic operad, such that $t_2(m_2) = m_2$.

There is a triple F_*^+ describing cyclic operads with multiplication. The triple F_*^+ is defined on the category of cyclic sequences of based spaces, with the base points fixed by the cyclic group action. If we ignore the cyclic group action, then F_*^+ agrees with the triple F_* of Remark 5.2. The action of the cyclic groups is defined similarly as for F^+ in Definition 7.5.

We are now ready to state the main theorem.

Theorem 7.7 Let O be a cyclic operad with multiplication. Then the operad fMS acts on $\text{Tot}(O^*)$.

Proof We need to describe the action of fMS on $\text{Tot}(O^*)$ via maps

$$\theta_m: fMS(m) \times \text{Tot}(O^*)^m \rightarrow \text{Tot}(O^*).$$

Given $x \in fMS(m)$, and cocyclic maps

$$a_1^*, \dots, a_m^* \in \text{Tot}(O^*) = \text{Mor}(\Lambda^*, O^*),$$

we want to define a cocyclic map


$$a^* := \theta_m(x; a_1^*, \dots, a_m^*) \in \text{Tot}(O^*) = \text{Mor}(\Lambda^*, O^*)$$

so we need compatible maps $a^k: \Lambda^k \rightarrow O(k)$ for $k \in \mathbb{N}$.

The construction is similar to the one for non-cyclic operads in the proof of Theorem 5.4.

Let $\gamma = c'(x): S^1 \rightarrow (S^1)^m$ be the parametrization of the cactus C_x associated to x in Definition 4.7. Given a system of cyclically ordered points $(x_0, \dots, x_k) \in \Lambda^k$ on the

circle S^1 we obtain via $\gamma: S^1 \rightarrow C_x$ a system of special points $(\gamma(x_0), \dots, \gamma(x_k))$ on the cactus C_x . We build a rooted labelled tree Γ almost exactly as in the proof of Theorem 5.4. There is only a slight difference in the definition of the label of a vertex associated to a lobe. Let S_j be a lobe of C_x and let $H_j \subset S_j$ be the finite subset of all intersection points of S_j with other lobes and of all special points of the form $\gamma(x_i)$ in the lobe. We have a preferred based identification of S_j with S^1 , but in general the base point is not an intersection point or $\gamma(x_0)$ as in the non-cyclic case. In the next figure we give an example for $m = 3$ and $k = 3$. The base point of the cactus is P and the base points of the lobes are P_1, P_2, P_3 . Only the vertices of the lobes are indicated.



Under the identification $S^1 \cong S_j$, the subset $H_j \subset S_j$ corresponds to a cyclically ordered sequence of points on S^1 , that is an element $y_j \in \Lambda^{l_j} = \Delta^{l_j} \times S^1$ (actually y_j is in the interior), where l_j is one less than the cardinality of H_j . The map $a_j^{l_j}: \Lambda^{l_j} \rightarrow O(l_j)$ defines the label $a_j^{l_j}(y_j) \in O(l_j)$ of the vertex of S_j . We define $a^k(x_0, \dots, x_k) \in O(k)$ as the composition of all labels of the rooted labelled tree Γ . The crucial point is that the map $a^k: \Lambda^k \rightarrow O(k)$ is equivariant with respect to the action of \mathbb{Z}_{k+1} , because O is a cyclic operad. Moreover the map $a^*: \Lambda^* \rightarrow O^*$ respects cofaces and codegeneracies because each a_i^* does so. Finally the structure maps θ_* make $\text{Tot}(O^*)$ into a fMS -algebra, since O is an algebra over the triple F_*^+ . \square

Example 7.8 For a group G the operad \underline{G} is a cyclic operad with multiplication. The operad with multiplication is in Example 3.4 with $M = G$, and the cyclic group action in Example 6.2. These are compatible because

$$t_2(m_2) = t_2(1, 1) = (1, 1) = m_2.$$

By Theorem 7.7

$$\text{Tot}(\underline{G}^*) \cong \Omega G \simeq \Omega^2 BG$$

has the action of an operad equivalent to the little framed 2-discs. By Proposition 3.9 BG is equivalent to a based space with a based S^1 -action. We expect the action to be trivial, up to a zig-zag of based equivariant maps that are non-equivariant weak equivalences.

Actually there is a specific based circle action on BG , that we also expect to be equivalent. A *cyclic space* is a *contravariant* functor from Λ to the category of topological spaces, and in particular it has the underlying structure of a simplicial space. The realization of a cyclic space is equipped with a circle action [9]. We recall from [9] that the space LBG is equivalent (by a S^1 -equivariant map that is a non-equivariant equivalence) to the realization of a cyclic space lbG with $(lbG)_n = G^{n+1}$, and

$$\begin{aligned} d_i(x_0, \dots, x_n) &= (x_0, \dots, x_{i-1}, x_i x_{i+1}, x_{i+2}, \dots, x_n) \\ s_i(x_0, \dots, x_n) &= (x_0, \dots, x_{i-1}, e, x_{i+1}, x_{i+2}, \dots, x_n) \\ t_n(x_0, \dots, x_n) &= (x_n, x_0, x_1, \dots, x_{n-1}). \end{aligned}$$

Here e is the unit of G . The realization $|lbG|$ of lbG can be thought of as the space of finite configurations of points on the circle with summable labels in G [21]. There is a cyclic subspace $bG \subset lbG$ that at level n is the space $(bG)_n$ of n -tuples (g_0, \dots, g_n) such that the ordered product satisfies $g_0 \cdots g_n = 1$. The homeomorphism $(bG)_n \cong G^n$ induced forgetting g_0 gives an isomorphism between bG and the classical simplicial construction B_*G with realization $|B_*G| = BG$. The category Λ is well known to be non-canonically isomorphic to its opposite Λ^{op} . The choice of an isomorphism gives a bijective correspondence between cyclic spaces and cocyclic spaces. In particular under the isomorphism described in [9, Section 1] the cyclic space bG corresponds to the cocyclic space ωG of Example 2.4.

Example 7.9 (String topology in the category of correspondences) The cocyclic space lX does not come from a cyclic operad because of course the totalization LX is not a double loop space in general. However if we work in the category Cor of correspondences this works. Objects in Cor are topological spaces and a morphisms $X \rightarrow Y$ is a pair of maps $X \xleftarrow{f} Z \xrightarrow{g} Y$. The composition is defined via pullbacks. When the backward map f is an inclusion we can view a morphism as a map defined only partially. If f is a homeomorphism then we identify the morphism to the honest map $g \circ f^{-1}: X \rightarrow Y$. The cartesian product induces a monoidal structure on Cor and it makes sense to consider operads in Cor .

It turns out that the collection lX forms a cyclic operad with multiplication in the category of correspondences.

The composition map

$$\circ_i: (lX)_m \times (lX)_n \rightarrow lX_{m+n-1}$$

is defined on each pair $((x_0, \dots, x_m), (y_0, \dots, y_n))$ such that $x_i = y_0$, sending it to

$$(x_0, \dots, x_{i-1}, y_1, \dots, y_n, x_{i+1}, \dots, x_m).$$

The operad map from the associative operad $\text{Ass} \rightarrow lX$ is defined at level n by the correspondence $* \leftarrow X \xrightarrow{\Delta} X^{n+1}$, where Δ is the inclusion of the thin diagonal. The associated cosimplicial object is the honest cosimplicial space lX . A small variant of Theorem 7.7 shows that the totalization $LX = \text{Tot}(lX)$ admits the action of the operad fMS in the category of correspondences. This is exactly the action constructed by Sasha Voronov [25], at least on the suboperad $\text{Cacti} \subset fMS$, and that motivated him to work in the category of correspondences.

Further directions Kaufmann [12; 13] and Kontsevich–Soibelman [14] proved that the Hochschild complex of a Frobenius algebra has an action on the chain level of the operad \mathcal{M} of moduli spaces of Riemann surfaces with boundary. The restriction of this action to the operad of genus 0 surfaces is the cyclic Deligne conjecture. The generalization works because the endomorphism operad of a Frobenius algebra is a *modular operad* [15] with multiplication. Let O be a modular operad in based spaces. It is natural to ask whether the totalization $\text{Tot}(O^*)$ admits an action of the operad \mathcal{M} .

8 Appendix: comparison with the McClure–Smith construction

We give the proof of Proposition 4.6. More generally we produce for any m a geometric construction of an operad MS_m that is isomorphic to the E_m operad called \mathcal{D}_m by McClure–Smith in [18]. In this section \mathcal{D}_m will denote such operad and not the little m -discs operad.

Definition 8.1 Let $MS_m(n) = \mathcal{F}_m(n) \times \text{Mon}(I, \partial I)$, where $\mathcal{F}_m(n)$ is defined similarly as $\mathcal{F}(n)$ in Definition 4.1 except that the 4-tuple (z_1, z_2, z_3, z_4) is replaced by a $(m+2)$ -tuple (z_1, \dots, z_{m+2}) with $z_{2i-1} \in I_j(x)$ and $z_{2i} \in I_k(x)$ for all i . The space $\mathcal{F}_m(n)$ is a regular CW-complex with a cell for each sequence assuming all values between 1 and n , with no equal adjacent values, and with no subsequence of

length $m + 1$ of the form $abab \dots$, for $a \neq b$. The cell is identified to the product $\prod_{j=1}^n \Delta^{d(j)}$, where $d(j)$ is one less than the number of occurrences of j in the sequence. Clearly $\mathcal{F}(n) = \mathcal{F}_2(n)$ and $MS = MS_2$.

The direct limits are denoted $MS_\infty = \lim_m MS_m$ and $\mathcal{F}_\infty(n) = \lim_m \mathcal{F}_m(n)$. There is a symmetric operad structure on MS_m that is defined as for MS in Section 4 by an embedding onto a suboperad of $\text{Coend}(S^1)$.

Proposition 8.2 *There is an isomorphism between the operad MS_∞ and the E_∞ operad \mathcal{D} in [18]. This restricts for each m to an isomorphism between MS_m and the E_m operad denoted \mathcal{D}_m in [18].*

Proof By definition $\mathcal{D}(k) = \text{Tot}(Y_k^*)$, where $Y_k^* = \Xi_k(\Delta^*, \dots, \Delta^*)$ is the cosimplicial space described in [18, Section 6]. Given cosimplicial spaces X_1^*, \dots, X_k^* , the degree s space $\Xi_k(X_1^*, \dots, X_k^*)^s$ is the colimit of a functor $G_1: A_1 \rightarrow \text{Top}$ to the category of topological spaces. Here A_1 is the category whose objects are pairs (f, h) consisting of a monotone map $f: [t] \rightarrow [s]$ and a map $h: [t] \rightarrow \{1, \dots, k\}$, for some t . A morphism from (f, h) to (f', h') is an ordered map $g: [t] \rightarrow [t']$ between the respective domains such that $f' \circ g = f$ and $h' \circ g = h$. The functor G_1 sends a pair (f, h) to $\prod_{1 \leq i \leq k} X^{|f^{-1}(i)|-1}$.

In the special case $X_1^* = \dots = X_k^* = \Delta^*$ [18, Corollary 12.3] states that Y_k^s is a regular CW complex with a cell for each pair (f, h) as above such that f is surjective and for $0 \leq j < t$ either $f(j) \neq f(j + 1)$ or $h(j) \neq h(j + 1)$. Such cell has dimension $t + 1 - k$ and is $G_1(f, h) = \prod_{1 \leq i \leq k} \Delta^{|f^{-1}(i)|-1}$. Thus there is a natural identification between the cell of Y_k^0 indexed by $(f, *)$ and the cell of $\mathcal{F}_\infty(k)$ indexed by the sequence $f(0), \dots, f(t)$. Here $*: [t] \rightarrow [0]$ is the unique map. These identifications are compatible with the inclusion of boundary cells and give a cellular Σ_k -equivariant homeomorphism $\alpha: \mathcal{F}_\infty(k) \cong Y_k^0$. In [18, Proposition 12.7] McClure–Smith construct an isomorphism of cosimplicial spaces $\omega: Y_k^* \cong \Delta^* \times Y_k^0$. We describe geometrically the homeomorphism

$$\Phi = \omega^{-1} \circ (\Delta^s \times \alpha): \Delta^s \times \mathcal{F}_\infty(k) \cong Y_k^s.$$

A partition $x \in \mathcal{F}_\infty(k)$ is a subdivision of S^1 into k 1-manifolds. An element of the simplex $z \in \Delta^s$ determines s points on the circle and generically s intervals that we call *coarse* intervals, inducing a subdivision of the 1-manifolds into intervals that we call *fine* intervals. We number by $0, \dots, t$ in the counterclockwise ordering the fine intervals. Let $f: [t] \rightarrow \{1, \dots, k\}$ be the map sending the index of a fine interval to the index of the 1-manifold containing it. Similarly let $h: [t] \rightarrow [s]$ be the map sending the index of a fine interval to the index of the coarse interval containing it. The cell of $\Phi(z, x) \in Y_k^s$

is indexed by the pair (f, h) . The point $\Phi(z, x)$ of the cell $\prod_{1 \leq i \leq k} \Delta^{|f^{-1}(i)|-1}$ is determined by the lengths of the fine intervals: the i th component corresponds to the lengths of the fine intervals contained in the i th 1-manifold, that sum up to 1 and thus determine an element of $\Delta^{|f^{-1}(i)|-1}$. The homeomorphisms Φ induce a Σ_k -equivariant homeomorphism

$$\Psi : \mathcal{D}(k) = \text{Tot}(Y_k^*) \cong \text{Tot}(\Delta^*) \times \mathcal{F}_\infty(k) = MS_\infty(k).$$

Given $f \in \mathcal{D}(a)$ and $g \in \mathcal{D}(b)$, ie cosimplicial maps $f^* : \Delta^* \rightarrow \Xi_a(\Delta^*, \dots, \Delta^*) = Y_a^*$ and $g^* : \Delta^* \rightarrow \Xi_b(\Delta^*, \dots, \Delta^*) = Y_b^*$, we recall that the operadic composition $f \circ_i g \in \mathcal{D}(a+b-1)$ is the cosimplicial map

$$\begin{array}{ccc} \Delta^* & \xrightarrow{f^*} & Y_a^* \xrightarrow{\Xi_a(\text{id}, \dots, g^*, \dots, \text{id})} \Xi_a(\Delta^*, \dots, \Xi_b(\Delta^*, \dots, \Delta^*), \dots, \Delta^*) \\ & & \cong \Xi_{a+b-1}(\Delta^*, \dots, \Delta^*) \\ & & \uparrow \Gamma \\ & & \Xi_{a+b-1}(\Delta^*, \dots, \Delta^*) \end{array}$$

where Γ is the cosimplicial isomorphism described in [18, Section 6]. Via the geometric correspondence Φ it is straightforward to check that for any $x \in \Delta^s$

$$\Psi(f \circ_i g)(x) = [\Psi(f) \circ_i \Psi(g)](x) \in \Delta^s \times \mathcal{F}_\infty(a+b-1),$$

and so $\Psi : \mathcal{D} \cong MS_\infty$ is an operad isomorphism.

For a fixed m let $Z_*^k \subset Y_*^k$ be the cosimplicial subspace in section 13 of [18]. The Σ_k -equivariant homeomorphism $\alpha : \mathcal{F}_\infty(k) \cong Y_0^k$ restricts to a Σ_k -equivariant homeomorphism $\mathcal{F}_m(k) \cong Z_0^k$, because the cells of $\mathcal{F}_m(k)$ correspond to the cells of Y_0^k whose indexing sequence has *complexity* not exceeding m , ie the cells of Z_0^k . The splitting $\omega : Y_*^k \cong \Delta^* \times Y_k^0$ restricts to a splitting $Z_*^k \cong \Delta^* \times Z_k^0$. The E_m -operad \mathcal{D}_m in [18] has as k th space $\mathcal{D}_m(k) = \text{Tot}(Z_*^k)$. Thus the operad isomorphism $\Psi : \mathcal{D} \cong MS_\infty$ restricts to an isomorphism between the suboperad $\mathcal{D}_m \subset \mathcal{D}$ in [18] and the suboperad $MS_m \subset MS$. \square

Proposition 4.6 follows as a special case for $m = 2$.

References

- [1] **JM Boardman, RM Vogt**, *Homotopy invariant algebraic structures on topological spaces*, Lecture Notes in Mathematics, Vol. 347, Springer, Berlin (1973) MR0420609
- [2] **R Budney**, *The operad of framed discs is cyclic*, J. Pure Appl. Algebra 212 (2008) 193–196 MR2355044

- [3] **R L Cohen, A A Voronov**, *Notes on string topology*, from: “String topology and cyclic homology”, Adv. Courses Math. CRM Barcelona, Birkhäuser, Basel (2006) 1–95 MR2240287
- [4] **V Drinfeld**, *On the notion of geometric realization*, Mosc. Math. J. 4 (2004) 619–626, 782 MR2119142
- [5] **M Gerstenhaber**, *The cohomology structure of an associative ring*, Ann. of Math. (2) 78 (1963) 267–288 MR0161898
- [6] **E Getzler**, *Batalin–Vilkovisky algebras and two-dimensional topological field theories*, Comm. Math. Phys. 159 (1994) 265–285 MR1256989
- [7] **E Getzler, M M Kapranov**, *Cyclic operads and cyclic homology*, from: “Geometry, topology, & physics”, Conf. Proc. Lecture Notes Geom. Topology, IV, Int. Press, Cambridge, MA (1995) 167–201 MR1358617
- [8] **P Hu**, *The Hochschild cohomology of a Poincaré algebra* arXiv:0707.4118
- [9] **J D S Jones**, *Cyclic homology and equivariant homology*, Invent. Math. 87 (1987) 403–423 MR870737
- [10] **R Kaufmann**, *On several varieties of cacti and their relations*, Algebr. Geom. Topol. 5 (2005) 237–300 MR2135554
- [11] **R Kaufmann**, *A proof of a cyclic version of Deligne’s conjecture via cacti* arXiv: math.QA/0403340
- [12] **R Kaufmann**, *Moduli space actions on the Hochschild co-chains of a Frobenius algebra I* arXiv:math.AT/0606064
- [13] **R Kaufmann**, *Moduli space actions on the Hochschild co-chains of a Frobenius algebra II* arXiv:math.AT/0606065
- [14] **M Kontsevich, Y Soibelman**, *Notes on A -infinity algebras, A -infinity categories and non-commutative geometry. I* arXiv:math.RA/0606241
- [15] **M Markl, S Shnider, J Stasheff**, *Operads in algebra, topology and physics*, Mathematical Surveys and Monographs 96, American Mathematical Society, Providence, RI (2002) MR1898414
- [16] **P May**, *The geometry of iterated loop spaces*, Lectures Notes in Mathematics 271, Springer (1972)
- [17] **J E McClure, J H Smith**, *A solution of Deligne’s Hochschild cohomology conjecture*, from: “Recent progress in homotopy theory (Baltimore, MD, 2000)”, Contemp. Math. 293, Amer. Math. Soc., Providence, RI (2002) 153–193 MR1890736
- [18] **J E McClure, J H Smith**, *Cosimplicial objects and little n -cubes. I*, Amer. J. Math. 126 (2004) 1109–1153 MR2089084

- [19] **J E McClure, J H Smith**, *Operads and cosimplicial objects: an introduction*, from: “Axiomatic, enriched and motivic homotopy theory”, NATO Sci. Ser. II Math. Phys. Chem. 131, Kluwer Acad. Publ., Dordrecht (2004) 133–171 MR2061854
- [20] **L Menichi**, *Batalin–Vilkovisky algebras and cyclic cohomology of Hopf algebras*, *K–Theory* 32 (2004) 231–251 MR2114167
- [21] **P Salvatore**, *Configuration spaces with summable labels*, from: “Cohomological methods in homotopy theory (Bellaterra, 1998)”, *Progr. Math.* 196, Birkhäuser, Basel (2001) 375–395 MR1851264
- [22] **P Salvatore**, *Knots, operads, and double loop spaces*, *Int. Math. Res. Not.* (2006) Art. ID 13628, 22 MR2276349
- [23] **P Salvatore, N Wahl**, *Framed discs operads and Batalin–Vilkovisky algebras*, *Q. J. Math.* 54 (2003) 213–231 MR1989873
- [24] **D P Sinha**, *Operads and knot spaces*, *J. Amer. Math. Soc.* 19 (2006) 461–486 MR2188133
- [25] **A A Voronov**, *Notes on universal algebra*, from: “Graphs and patterns in mathematics and theoretical physics”, *Proc. Sympos. Pure Math.* 73, Amer. Math. Soc. (2005) 81–103 MR2131012

Università di Roma Tor Vergata, Dipartimento di Matematica, Via della Ricerca Scientifica, Roma 00133, Italy,

salvator@mat.uniroma2.it

<http://www.mat.uniroma2.it/~salvator/>

Received: 25 June 2008 Revised: 5 November 2008