

Stein fillable Seifert fibered 3–manifolds

ANA G LECUONA

PAOLO LISCA

We characterize the closed, oriented, Seifert fibered 3–manifolds which are oriented boundaries of Stein manifolds. We also show that for this class of 3–manifolds the existence of Stein fillings is equivalent to the existence of symplectic fillings.

57R17; 53D10

1 Introduction and statement of results

The most important dichotomy in 3–dimensional contact topology is the one introduced by Eliashberg between tight and overtwisted contact structures (see eg Etnyre [5] and Geiges [6]). Nowadays there are several different ways to prove that a contact structure is tight, but for a long time the only systematic way to construct tight contact structures on a closed 3–manifold Y was to show Y to be orientation preserving diffeomorphic to the oriented boundary of a Stein manifold and then appeal to a theorem of Eliashberg [3] and Gromov [10]. This naturally led to the question of which 3–manifolds carry tight contact structures, as well as to the related question of which 3–manifolds admit *Stein fillings*, ie are orientation preserving diffeomorphic to the boundary of a Stein manifold. The first example of an oriented 3–manifold admitting no Stein fillings was provided by the second author in [12], and infinitely many examples were found by the second author and Stipsicz in [16, Theorem 4.2] and [19, Proposition 4.1]. While the second author and Stipsicz [20] recently achieved the classification of the closed, Seifert fibered 3–manifolds carrying tight contact structures, the classification of the Stein fillable ones was still missing. The purpose of the present paper is to fill this gap. Our main result, Theorem 1.5 below, identifies explicitly the family of closed, oriented, Seifert fibered 3–manifolds which are orientation preserving diffeomorphic to the boundary of a Stein manifold.

We need some preliminaries in order to state our results. Eliashberg [4] proved that smooth, even-dimensional manifolds carrying Stein structures can be characterized as having suitable handle decompositions. Gompf [9, Theorem 5.4] elaborated on Eliashberg’s result to show that a closed, oriented, Seifert fibered 3–manifold Y

admits a Stein filling unless Y is orientation preserving diffeomorphic to the oriented 3–manifold $Y(e_0; r_1, \dots, r_k)$ given by the surgery description of Figure 1, where

$$e_0 = -1, \quad k \geq 3 \quad \text{and} \quad 1 > r_1 \geq \dots \geq r_k > 0.$$




Figure 1: The Seifert 3–manifold $Y(e_0; r_1, \dots, r_k)$

Gompf also discovered a sufficient condition for the existence of Stein fillings of a 3–manifold of the form $Y(-1; r_1, \dots, r_k)$. We describe and use this condition in Section 2.

Definition 1.1 A k –tuple $(r_1, \dots, r_k) \in (\mathbb{Q} \cap (0, 1))^k$ with $k \geq 3$ and $r_1 \geq r_2 \geq \dots \geq r_k$ is *realizable* if there exist coprime integers $n > h > 0$ such that

$$\frac{h}{n} > r_1, \quad \frac{n-h}{n} > r_2 \quad \text{and} \quad \frac{1}{n} > r_3, \dots, r_k.$$

Definition 1.2 A closed, oriented, Seifert fibered 3–manifold is *of special type* if it is orientation preserving diffeomorphic to $Y(-1; r_1, \dots, r_k)$, where $k \geq 3$, $1 > r_1 \geq r_2 \geq \dots \geq r_k > 0$ and the following conditions both hold:

- (1) (r_1, \dots, r_k) is not realizable.
- (2) $r_1 + r_2 < 1$.

In Section 2 we use Gompf’s sufficient condition for the existence of Stein fillings of $Y(-1; r_1, \dots, r_k)$ to establish the following:

Theorem 1.3 *Let Y be a closed, oriented, Seifert fibered 3–manifold which is not of special type. Then, Y is orientation preserving diffeomorphic to the boundary of a Stein surface.*

Recall (see eg Geiges [6]) that a *symplectic filling* of a contact 3–manifold (Y, ξ) is a pair (X, ω) , where X is a smooth 4–manifold with boundary oriented by a symplectic form $\omega \in \Omega^2(X)$, and such that there is an orientation preserving diffeomorphism $\varphi: Y \rightarrow \partial X$ with $\omega|_{d\varphi(\xi)} \neq 0$ at each point of ∂X . A Stein filling is a symplectic filling

but the converse is not true, since there are examples of symplectically fillable contact 3-manifolds which are not Stein fillable; see Ghiggini [7]. Similarly, there exist several examples of tight, contact Seifert fibered 3-manifolds which are not symplectically fillable; see Ghiggini, Lisca and Stipsicz [8] and Lisca and Stipsicz [15; 16; 17]. In Section 3 we apply Donaldson's "Theorem A" on the intersection forms of definite 4-manifolds to prove the following:

Theorem 1.4 *A closed, oriented, Seifert fibered 3-manifold of special type admits no symplectic fillings.*

Combining Theorems 1.3 and 1.4 immediately gives our main result:

Theorem 1.5 *Let Y be a closed, oriented, Seifert fibered 3-manifold. Then, the following conditions are equivalent:*

- (1) Y admits Stein fillings.
- (2) Y admits symplectic fillings.
- (3) Y is not of special type.

Proof A Stein filling is a symplectic filling, therefore (1) implies (2). By Theorem 1.4, (2) implies (3), and by Theorem 1.3, (3) implies (1). \square

Theorem 1.5 implies Corollary 1.6 below, which shows that the condition that a 3-manifold is Seifert fibered of special type can be reformulated in terms of open book decompositions. Recall that an open book decomposition of a closed 3-manifold is called *positive* if its monodromy can be written as a product of right-handed Dehn twists. Loi and Piergallini [21, Theorem 4] proved that a smooth, closed, oriented 3-manifold Y is the boundary of a Stein surface if and only if Y admits a positive open book decomposition. Combining this result with Theorem 1.5 immediately yields the following:

Corollary 1.6 *A closed, oriented, Seifert fibered 3-manifold Y admits a positive open book decomposition if and only if Y is not of special type.* \square

We should perhaps say something about the history of the realizability condition of Definition 1.1. Such condition originated in the literature on smooth foliations from the early 1980's. Indeed, the papers of Eisenbud, Hirsch and Neumann [2], Jankins and Neumann [11] and Naimi [22] make use of the realizability condition to give a classification of the closed, oriented, Seifert fibered 3-manifolds carrying smooth foliations transverse to the Seifert fibration (see Lisca and Matić [14, Theorem 1.2])

for a statement summarizing the results). Such classification was used in [14] to characterize the closed, oriented, Seifert fibered 3-manifolds which carry positive contact structures transverse to the fibration. The realizability condition also appears in the characterizations of the Seifert fibered Heegaard Floer L -spaces in terms of transverse contact structures, transverse foliations and taut foliations given by the second author and Stipsicz in [18, Theorem 1.1].

We would like to end this introduction with a few remarks about Seifert 3-manifolds of special type. It should be clear from the definition that there exist infinitely many oriented Seifert 3-manifolds of special type. In fact, infinitely many examples of closed, oriented Seifert fibered 3-manifolds without symplectic fillings are known [16, Theorem 4.2; 19, Proposition 4.1]. According to Theorem 1.5 such examples must be all of special type, and indeed this fact can be verified directly. The infinitely many oriented Seifert 3-manifolds which do not carry tight contact structures [19; 20] are also of special type. Moreover, if $Y = Y(-1; r_1, \dots, r_k)$ is of special type then $-Y$ is not, because it is of the form $Y(-1 + k; 1 - r_k, \dots, 1 - r_1)$. This is consistent with the general fact that each oriented Seifert fibered 3-manifold admits a Stein filling after possibly reversing its orientation [9, Corollary 5.5(a)]. Also, Condition (1) from Definition 1.2 together with the results of [18] imply that an oriented Seifert fibered 3-manifold of special type is an L -space in the sense of Ozsváth and Szabó [25, Definition 1.1]. In fact, it follows from [18] that if Y is an oriented Seifert fibered 3-manifold which is an L -space, then after possibly reversing orientation Y is of the form $Y(e_0; r_1, \dots, r_k)$, with $k \geq 3$, $1 > r_1 \geq \dots \geq r_k > 0$ and either (i) $e_0 \geq 0$ or (ii) $e_0 = -1$ and (r_1, \dots, r_k) is not realizable. Therefore, the oriented Seifert fibered 3-manifolds of special type are precisely the oriented, Seifert fibered L -spaces of the form $Y(-1; r_1, \dots, r_k)$ with $k \geq 3$, $1 > r_1 \geq \dots \geq r_k > 0$ and (r_1, \dots, r_k) satisfying Condition (2) from Definition 1.2.

The organization of the paper is straightforward: in Section 2 we prove Theorem 1.3 and in Section 3 we prove Theorem 1.4.

2 Existence of Stein fillings

The purpose of this section is to prove Theorem 1.3. We start with recalling Gompf's sufficient condition from [9] for the existence of a Stein filling of $Y(-1; r_1, \dots, r_k)$.

Given a rational number $r \in \mathbb{Q}$ we can define an integer $\llbracket r \rrbracket \in \mathbb{Z}$ by setting $r = \llbracket r \rrbracket + \text{frac}(r)$, where $\text{frac}(r) \in [0, 1)$. Let

$$r'_i := -\frac{1}{r_i}, \quad i = 1, \dots, k,$$

and $s \in (-\infty, -1)$ be such that $1/s := -1 - 1/r'_1$. If $s \neq r'_2$ then it is easy to check that there is a map

$$A: \mathbb{Q} \cup \{\infty\} \rightarrow \mathbb{Q} \cup \{\infty\}$$

of the form $A(r) = (c + dr)/(a + br)$ with $a, b, c, d \in \mathbb{Z}$, such that

$$(2-1) \quad ad - bc = \pm 1,$$

$$(2-2) \quad A(s) \in (-1, 0],$$

$$(2-3) \quad A(r'_2) \in [-\infty, -1).$$

Let

$$t := \begin{cases} 0 & \text{if } A(0) \in [0, +\infty], \\ 1/A(s) & \text{if } A(0) \in [-1, 0), \\ A(r'_2) & \text{if } A(0) \in (-\infty, -1). \end{cases}$$

Set $M := \max(|a|, |c|)$, $m := \min(|a|, |c|)$ and

$$n_A(r'_1, r'_2) := -m(\lceil t \rceil + 1) - M.$$

Finally, let

$$n(r'_1, r'_2) := \begin{cases} 0 & \text{if } s = r'_2, \\ \sup_A n_A(r'_1, r'_2) & \text{if } s \neq r'_2, \end{cases}$$

where the supremum is taken over the set of matrices A as above, satisfying Conditions (2-1), (2-2) and (2-3). Gompf [9] shows that $Y(-1; r_1, \dots, r_k)$ is the boundary of a Stein surface if

$$(2-4) \quad n(r'_1, r'_2) > r'_3, \dots, r'_k.$$

Observe that when $s = r'_2$ Condition (2-4) is automatically satisfied.

In order to prove Theorem 1.3 we need two results. The first result is Theorem 2.1, which establishes the existence of a Stein filling for $Y(-1; r_1, \dots, r_k)$ under the assumption that the k -tuple (r_1, \dots, r_k) is *realizable*, that is to say that there exist coprime integers $n > h > 0$ such that $r_1 < h/n$, $r_2 < (n - h)/n$ and $r_3, \dots, r_k < 1/n$.

Theorem 2.1 *Suppose that $k \geq 3$, $1 > r_1 \geq r_2 \geq \dots \geq r_k > 0$ and (r_1, \dots, r_k) is realizable. Then, $Y(-1; r_1, \dots, r_k)$ is orientation preserving diffeomorphic to the boundary of a Stein surface.*

Proof Recall that we defined $r'_i := -1/r_i$, $i = 1, \dots, k$. We will prove that there is a map $A: \mathbb{Q} \cup \{\infty\} \rightarrow \mathbb{Q} \cup \{\infty\}$ satisfying Properties (2-1), (2-2), (2-3) above, and such that

$$n_A(r'_1, r'_2) > r'_3, \dots, r'_k.$$

In view of Gompf's condition (2-4), this clearly suffices to prove the statement.

By the realizability assumption, there is a positive integer n_0 such that, for some integer h_0 coprime with n_0 and satisfying $n_0 > h_0 > 0$ we have

$$r'_2 < -\frac{n_0}{h_0} < s \quad \text{and} \quad -n_0 > r'_3, \dots, r'_k.$$

Denote by n the smallest positive integer such that, for some integer h coprime with n with $n > h > 0$, we have

$$n \leq n_0 \quad \text{and} \quad r'_2 < -\frac{n}{h} < s.$$

Notice that, since $n \leq n_0$, $-n \geq -n_0 > r'_3, \dots, r'_k$. Moreover, n and h being coprime, there exist $a, b \in \mathbb{Z}$ such that

$$(2-5) \quad ah - bn = 1.$$

If the pair (a, b) solves Equation (2-5), so does the pair $(a + zn, b + zh)$ for each $z \in \mathbb{Z}$. Therefore, we can choose a solution (a, b) such that $0 \leq a < n$. Indeed, since $a = 0$ would imply $n = 1$, which is not the case because $n > h > 0$, we can assume $0 < a < n$. From Equation (2-5) we get

$$b = \frac{ah}{n} - \frac{1}{n},$$

hence $-1/n < b < h - 1/n$, which is equivalent to

$$0 \leq b < h.$$

Soon it will be convenient to have $b > 0$, therefore we deal now with the special case $b = 0$. By Equation (2-5), $b = 0$ implies $a = h = 1$, therefore $r'_2 < -n < s$. Moreover, by the minimality of n we must have $-(n-1) \geq s$. Define the map A by

$$A(r) := r + n - 1.$$

This map is of the form $(c + dr)/(a + br)$ with $a = 1$, $b = 0$, $c = n - 1$ and $d = 1$, therefore $ad - bc = 1$. Clearly A is monotone increasing, $A(-n) = -1$ and $A(-n + 1) = 0$. Therefore $A(r'_2) \in (-\infty, -1)$ and $A(s) \in (-1, 0]$, thus A satisfies the required Properties (2-1), (2-2) and (2-3). Since $A(0) = n - 1 \in [0, +\infty]$ we have $t = 0$, $m = 1$ and $M = n - 1$, therefore

$$n_A(r'_1, r'_2) = -m - M = -1 - n + 1 = -n > r'_3, \dots, r'_k.$$

From now on we assume $0 < b < h$. Observe that Equation (2-5) is equivalent to

$$\frac{a}{b} = \frac{n}{h} + \frac{1}{hb},$$

which implies

$$-\frac{a}{b} < -\frac{n}{h}.$$

In fact, by our choice of n we must have

$$-\frac{a}{b} \leq r'_2.$$

Now we define A by

$$A(r) := \frac{(n-a) + (h-b)r}{a+br} = -1 + \frac{n+hr}{a+br}.$$

For this map we have $c = n - a$ and $d = h - b$, therefore

$$ad - bc = a(h - b) - b(n - a) = ah - bn = 1.$$

The map A is monotone increasing for every $r \neq -a/b$, because

$$\frac{dA}{dr}(r) = \frac{1}{(a+br)^2}.$$

Equation (2-5) implies $-(n-a)/(h-b) > -n/h$, thus by the choice of n we have $s \leq -(n-a)/(h-b)$. We conclude

$$A\left(-\frac{a}{b}\right) = -\infty \leq A(r'_2) < A\left(-\frac{n}{h}\right) = -1 < A(s) \leq A\left(-\frac{n-a}{h-b}\right) = 0.$$

Therefore A satisfies Properties (2-1), (2-2) and (2-3). Since

$$A(0) = -1 + \frac{n}{a} = \frac{n-a}{a} \in (0, +\infty)$$

we have $t = 0$, thus

$$n_A(r'_1, r'_2) = -m - M = -|a| - |c| = -a - (n - a) = -n > r'_3, \dots, r'_k. \quad \square$$

We can now move on to the second result needed for the proof of Theorem 1.3, that is Theorem 2.5 below. This result will establish the existence of a Stein filling for $Y(-1; r_1, \dots, r_k)$ under the assumption $r_1 + r_2 \geq 1$. The proof of Theorem 1.3 will then follow combining Theorems 2.1 and 2.5.

Consider the standard Farey tessellation of the hyperbolic plane. Figure 2 illustrates some of the arcs of the tessellation with both endpoints in the interval $[-\infty, -1]$. We shall refer to any such arc with endpoints $\alpha < \beta$ as to the *Farey arc* $\widehat{\alpha\beta}$.

Observe that, given a Farey arc $\widehat{\alpha\gamma}$, there is a unique point β such that $\alpha < \beta < \gamma$ and there exist Farey arcs $\widehat{\alpha\beta}$ and $\widehat{\beta\gamma}$. In what follows, we shall refer to the unique point β as to the *middle point* of $\widehat{\alpha\gamma}$, and denote it by $m(\alpha, \gamma)$.




Figure 2: Arcs of the Farey tessellation

Lemma 2.2 Let $s, r'_2 \in (-\infty, -1)$, with $s < r'_2$. Then, there exists a Farey arc $\widehat{\alpha\gamma}$ with middle point $\beta = m(\alpha, \gamma)$ satisfying

$$-\infty \leq \alpha < s \leq \beta \leq r'_2 < \gamma \leq -1.$$

In other words, there is a configuration of Farey arcs as in Figures 3(A), 3(B) or 3(C).

Proof We provisionally define $\alpha = -\infty$ and $\gamma = -1$. If $s \leq \beta = m(\alpha, \gamma) \leq r'_2$ then α, β and γ already satisfy the statement and the lemma is proved. Otherwise we have either $s > \beta$ or $r'_2 < \beta$ (but not both, because $s < r'_2$). If $s > \beta$ we redefine $\alpha = \beta$, while if $r'_2 < \beta$ we redefine $\gamma = \beta$, and in both cases we set β equal to the new middle point $m(\alpha, \gamma)$. As before, if $s \leq \beta = m(\alpha, \gamma) \leq r'_2$ we are done, otherwise either $s > \beta$ or $r'_2 < \beta$ (but not both). Continuing in this fashion, after a finite number of steps we necessarily arrive at a configuration satisfying the statement of the lemma. \square




Figure 3: The configurations of Farey arcs of Lemma 2.2

Lemma 2.3 Let $\widehat{\alpha\beta}$ be a Farey arc with $\alpha, \beta \in [-\infty, -1]$, and let $s \in (\alpha, \beta)$. Then, there exist Farey arcs $\widehat{\alpha\beta'}$ and $\widehat{\alpha'\beta}$ such that

$$m(\alpha, \beta') \leq s < \beta' \leq \beta \quad \text{and} \quad \alpha \leq \alpha' < s \leq m(\alpha', \beta).$$

In other words, there are configurations of Farey arcs as in Figures 4(A) and 4(B).




Figure 4: The configurations of Farey arcs of Lemma 2.3

Proof We only prove the existence of the arc $\widehat{\alpha\beta'}$ – the existence of the arc $\widehat{\alpha'\beta}$ can be established in the same way. If $m(\alpha, \beta) \leq s$ we define $\beta' = \beta$, and the statement is proved. If $s < m(\alpha, \beta)$, we define (temporarily) $\beta' = m(\alpha, \beta)$. If $m(\alpha, \beta') \leq s$ the arc $\widehat{\alpha\beta'}$ satisfies the statement. Otherwise $s < m(\alpha, \beta')$, we redefine $\beta' = m(\alpha, \beta')$ and we keep going in the same way. In a finite number of steps we are bound to find the arc $\widehat{\alpha\beta'}$ with the stated property. \square

Lemma 2.4 Let $s, r'_2 \in (-\infty, -1)$, with $s < r'_2$. Then, there exist Farey arcs $\widehat{\alpha\beta}, \widehat{\beta\gamma}, \widehat{\gamma\delta}, \widehat{x\gamma}$, with $\widehat{x\gamma} \in \{\widehat{\alpha\gamma}, \widehat{\beta\delta}\}$, satisfying

$$-\infty \leq \alpha < s \leq \beta < \gamma \leq r'_2 < \delta \leq -1.$$

In other words, there is a configuration of Farey arcs as in Figures 5(A) or 5(B).




Figure 5: The configurations of Farey arcs of Lemma 2.4

Proof By Lemma 2.2 there is a configuration of Farey arcs as in Figures 3(A), 3(B) or 3(C). If Figure 3(A) holds we can apply Lemma 2.3 to the point s and the arc $\widehat{\alpha\beta}$ of Figure 3(A) to find a configuration as in Figure 4(B). If we set $m = m(\alpha', \beta)$, the Farey arcs $\widehat{\alpha'm}, \widehat{m\beta}, \widehat{\alpha'\beta}$ and $\widehat{\beta\gamma}$ provide a configuration as in Figure 5(B). If Figure 3(B) holds we can apply Lemma 2.3 to the point r'_2 and the arc $\widehat{\beta\gamma}$ of Figure 3(B) to find a configuration as in Figure 4(A). In other words, there exists a Farey arc $\widehat{\beta\beta'}$ such that $m(\beta, \beta') \leq r'_2 < \beta'$. Setting $m = m(\beta, \beta')$, the Farey arcs $\widehat{\alpha\beta}, \widehat{\beta m}, \widehat{m\beta'}$ and $\widehat{\beta\beta'}$ provide a configuration as in Figure 5(A). Finally, if Figure 3(C) holds we can apply Lemma 2.3 to the point s and the arc $\widehat{\alpha\beta}$ of Figure 3(C) to find a configuration as in Figure 4(B). If we set $m = m(\alpha', \beta)$, the Farey arcs $\widehat{\alpha'm}, \widehat{m\beta}, \widehat{\alpha'\beta}$ and $\widehat{\beta\gamma}$ provide a configuration as in Figure 5(B). \square

Theorem 2.5 *Suppose that $k \geq 3$, $1 > r_1 \geq r_2 \geq \dots \geq r_k > 0$ and $r_1 + r_2 \geq 1$. Then, $Y(-1; r_1, \dots, r_k)$ is orientation preserving diffeomorphic to the boundary of a Stein surface.*

Proof The inequality $r_1 + r_2 \geq 1$ is equivalent to $s \leq r'_2$. If $s = r'_2$ Condition (2-4) is automatically satisfied, therefore we may assume $s < r'_2$. By Lemma 2.4 there is a configuration of Farey arcs as in Figures 5(A) or 5(B). Let us suppose that the first case occurs. In view of the action of $SL(2, \mathbb{Z})$ on the Farey tessellation, there is a unique map $A: \mathbb{Q} \cup \{\infty\} \rightarrow \mathbb{Q} \cup \{\infty\}$ of the form $A(r) = (c + dr)/(a + br)$ satisfying Condition (2-1) and such that $A(\beta) = 0$, $A(\gamma) = \infty$ and $A(\delta) = -1$. By construction, A satisfies Conditions (2-2) and (2-3) as well, $c/a = A(0) \in (-1, 0)$ and $A(s) \in (-1, 0)$. According to Gompf's condition (2-4), in order to prove that $Y(-1; r_1, \dots, r_k)$ carries Stein fillable contact structures it suffices to show that

$$n_A(r'_1, r'_2) = -m(\llbracket t \rrbracket + 1) - M \geq -1,$$

where $t = 1/A(s) \in (-\infty, -1)$, $M = \max(|a|, |c|)$ and $m = \min(|a|, |c|)$. Observe that, since $A(0) = c/a \in (-1, 0)$, $M = |a|$ and $m = |c|$. Condition (2-1) implies that if $m = 0$ then $M = 1$, therefore we may assume without loss of generality that $m > 0$. An easy calculation shows that the condition $-m(\llbracket t \rrbracket + 1) - M \geq -1$ is equivalent to

$$\llbracket t \rrbracket + 1 - \frac{1}{m} \leq -\frac{M}{m} = \frac{1}{A(0)}.$$

In order to prove this inequality it suffices to show that there is an integer N strictly greater than $1/A(s)$ and less than or equal to $1/A(0)$, ie such that

$$\llbracket t \rrbracket + 1 \leq N \leq \frac{1}{A(0)}.$$

This condition is satisfied if and only if there exists a Farey arc $\widehat{\infty x}$ with $1/A(s) < x \leq 1/A(0)$ or, equivalently, if and only if there exists a Farey arc $\widehat{y 0}$ with $A(0) \leq y < A(s)$. Setting $y := A(\alpha)$, such an arc is provided by the image under A of the Farey arc $\widehat{\alpha \beta}$ from Lemma 2.4, because by construction $A(\beta) = 0$ and $0 < \infty = -\infty \leq \alpha < s$. This concludes the proof under the assumption that when at the beginning of the argument we apply Lemma 2.4 we end up with a configuration of Farey arcs as in Figure 5(A). In case the configuration is the one given by Figure 5(B) we can argue in a similar way, so we just describe the steps where there is a difference. We choose the unique map A so that $A(\alpha) = -1$, $A(\beta) = 0$ and $A(\gamma) = +\infty = -\infty$. Then, $c/a = A(0) \in (-\infty, -1)$, $t = A(r'_2) \in (-\infty, -1)$, $M = |c|$, $m = |a|$ and as before we may assume without loss that $m > 0$. The same calculation as in the previous case shows that Gompf's condition

is equivalent to

$$\llbracket A(r'_2) \rrbracket + 1 - \frac{1}{m} \leq -\frac{M}{m} = A(0).$$

As before, this condition is satisfied if there exists a Farey arc $-\infty z$ with $A(r'_2) < z \leq A(0)$. Setting $z := A(\delta)$, such an arc is provided by the image under A of the Farey arc $\widehat{\gamma\delta}$ from Lemma 2.4, because by construction $A(\gamma) = -\infty$ and $r'_2 < \delta \leq -1 < 0$. \square

We are now ready to prove Theorem 1.3. We restate the result for the reader's convenience:

Theorem 2.6 *Let Y be a closed, oriented, Seifert fibered 3-manifold which is not of special type. Then, Y is orientation preserving diffeomorphic to the boundary of a Stein surface.*

Proof Gompf showed [9, Theorem 5.4] that a closed, oriented, Seifert fibered 3-manifold Y admits a Stein filling unless it is of the form $Y(-1; r_1, \dots, r_k)$ with $k \geq 3$ and $1 > r_1 \geq r_2 \geq \dots \geq r_k > 0$. Thus, if Y is not of special type then it is diffeomorphic to the boundary of a Stein surface unless Y is of the form $Y(-1; r_1, \dots, r_k)$ with $k \geq 3$, $1 > r_1 \geq r_2 \geq \dots \geq r_k > 0$ and either (i) (r_1, \dots, r_k) is realizable or (ii) $r_1 + r_2 \geq 1$. But Conditions (i) and (ii) are the assumptions of, respectively, Theorem 2.1 and Theorem 2.5, therefore if Y is not of special type then Y is necessarily the boundary of a Stein surface. \square

3 Nonexistence of symplectic fillings

The purpose of this section is to establish Theorem 1.4. We start with some preliminaries, then we prove three auxiliary lemmas. After that we prove Theorem 1.4.

Let $Y = Y(e_0; r_1, \dots, r_k)$ denote the oriented, Seifert fibered 3-manifold given by the surgery description of Figure 1, where $e_0 \in \mathbb{Z}$, $r_i \in (0, 1) \cap \mathbb{Q}$ and $r_1 \geq r_2 \geq \dots \geq r_k$. The oriented 3-manifold Y is the oriented boundary of the 4-dimensional plumbing P_Γ of D^2 -bundles over 2-spheres described by the star-shaped weighted graph Γ with k legs illustrated in Figure 6.

The weights of the vertices in the leg L_i , which form the string $(a^i_1, \dots, a^i_{h_i})$, are given by the unique continued fraction expansion such that $a^i_j \leq -2$ for every $j \in \{1, \dots, h_i\}$:

$$-\frac{1}{r_i} = [a^i_1, \dots, a^i_{h_i}] := a^i_1 - \frac{1}{a^i_2 - \frac{1}{\dots - \frac{1}{a^i_{h_i-1} - \frac{1}{a^i_{h_i}}}}}, \quad i = 1, \dots, k.$$




Figure 6: The weighted star-shaped graph Γ

We can associate to Γ the intersection lattice $(\mathbb{Z}^{|\Gamma|}, Q_\Gamma)$ of the plumbing P_Γ . In the proof of Theorem 1.4 we will show that if Y admits a symplectic filling then the intersection lattice of the plumbing associated to $-Y$ admits an isometric embedding into a standard diagonal lattice. Our present aim will be to prepare the ground for Lemma 3.3, which shows that under a certain assumption such an embedding does not exist.

We shall need Riemenschneider’s point rule [26], which we now recall. Let $p > q > 0$ be coprime integers, and suppose

$$-\frac{p}{q} = [a_1, \dots, a_l], \quad a_i \leq -2, \quad -\frac{p}{p-q} = [b_1, \dots, b_m], \quad b_j \leq -2.$$

Then, the coefficients a_1, \dots, a_l and b_1, \dots, b_m are related by a “point diagram” as in Figure 7, where the i -th row contains $|a_i| - 1$ “points” for $i = 1, \dots, l$, and the first point of each row is vertically aligned with the last point of the previous row. The




Figure 7: The point diagram underlying Riemenschneider’s point rule

point rule says that there are m columns, and the j -th column contains $|b_j| - 1$ points for every $j = 1, \dots, m$. For example if $-7/5 = [-2, -2, -3]$ and $-7/2 = [-4, -2]$

the corresponding diagram is given by



Let Γ be either a star-shaped or a linear weighted graph. If $(\mathbb{Z}^{|\Gamma|}, Q_\Gamma)$ admits an embedding into a standard diagonal lattice $(\mathbb{Z}^k, -\text{Id})$ with basis e_1, \dots, e_k , then we will write, for every subset S of the set of vertices of Γ ,

$$U_S := \{e_i \mid e_i \cdot v \neq 0 \text{ for some } v \in S\}.$$

We can now start to work towards Lemma 3.3 and Theorem 1.4.

Lemma 3.1 *Suppose $-1/r = [a_1, \dots, a_n]$ and $-1/s = [b_1, \dots, b_m]$ where $r + s = 1$. Consider a weighted linear graph Ψ having two connected components, Ψ_1 and Ψ_2 , where Ψ_1 consists of n vertices v_1, \dots, v_n with weights a_1, \dots, a_n and Ψ_2 of m vertices w_1, \dots, w_m with weights b_1, \dots, b_m . Moreover, suppose that there is an embedding of the lattice $(\mathbb{Z}^{n+m}, Q_\Psi)$ into $(\mathbb{Z}^k, -\text{Id})$, with basis e_1, \dots, e_k , such that $e_1 \in U_{v_1} \cap U_{w_1}$ and $U_\Psi = \{e_1, \dots, e_k\}$. Then,*

- (1) $U_{\Psi_1} = U_{\Psi_2}$ and
- (2) $k = n + m$.

Proof We start showing that (1) implies (2). In fact, since $r + s = 1$, by [13, Lemma 2.6], we have

$$\sum_{i=1}^n (-a_i - 3) + \sum_{j=1}^m (-b_j - 3) = -2$$

and therefore

$$(3-1) \quad |\text{tr}(Q_\Psi)| = \sum |a_i| + \sum |b_j| = 3(n + m) - 2.$$

The matrix Q_Ψ is not singular [13, Remark 2.1] so we necessarily have $k \geq n + m$. Let us write $k = n + m + x$ for some $x \geq 0$. Since we are assuming (1), each vector of the basis e_1, \dots, e_k satisfies $e_i \in U_{\Psi_1} \cap U_{\Psi_2}$ and therefore $|\text{tr}(Q_\Psi)| \geq 2k$. Moreover, since the graph Ψ has $n + m - 2$ edges, it follows that $|\text{tr}(Q_\Psi)| \geq 2k + n + m - 2 = 3(n + m) - 2 + 2x$. Hence, by (3-1), $x = 0$ and (2) holds.

We now assume that (1) does not hold and show that this assumption leads to a contradiction. Start by defining the sets $E_1 := U_{\Psi_1} \setminus U_{\Psi_2}$, $E_2 := U_{\Psi_2} \setminus U_{\Psi_1}$ and $E_{12} := U_{\Psi_1} \cap U_{\Psi_2}$. Since we are assuming that (1) does not hold, we have $E_1 \neq \emptyset$

or $E_2 \neq \emptyset$. By symmetry, we may assume that $E_1 \neq \emptyset$. It follows that there exists a smallest index $n_0 \in \{1, \dots, n\}$ such that $U_{v_{n_0}} \cap E_1 \neq \emptyset$. This condition allows us to construct a new connected linear graph $\bar{\Psi}_1$ with n_0 vertices and associated string of weights $(a_1, \dots, a_{n_0-1}, \bar{a}_{n_0})$, where

$$\bar{a}_{n_0} := a_{n_0} + \sum_{e_\ell \in E_1} (v_{n_0} \cdot e_\ell)^2.$$

Notice that since the intersection lattice associated with Ψ admits an embedding into a diagonal lattice, there is a naturally induced analogous embedding of the intersection lattice associated with $\bar{\Psi}_1 \cup \Psi_2$. We claim that $\bar{a}_{n_0} \leq -2$. In fact, if $n_0 = 1$ the assumption $e_1 \in U_{v_1} \cap U_{w_1}$ and the equality $v_1 \cdot w_1 = 0$ imply $|E_{12} \cap U_{v_1}| \geq 2$ and therefore in this case $\bar{a}_1 \leq -2$. On the other hand, if $n_0 > 1$ then, by definition of n_0 , it holds that $U_{v_{n_0-1}} \subseteq E_{12}$. The equalities $v_{n_0-1} \cdot v_{n_0} = 1$ and $v_{n_0} \cdot w_\ell = 0$ for every $\ell \in \{1, \dots, m\}$ force $|E_{12} \cap U_{v_{n_0}}| \geq 2$ and therefore $\bar{a}_{n_0} \leq -2$ as claimed.

Now, since $-2 \geq \bar{a}_{n_0} > a_{n_0}$ we have, by standard facts on continued fractions, that $-1/\bar{r} := [a_1, \dots, a_{n_0-1}, \bar{a}_{n_0}]$ satisfies $\bar{r} + s > 1$ (since $r + s = 1$). Let \bar{r}' be such that $\bar{r} + \bar{r}' = 1$. Using Riemenschneider's point diagram it is not difficult to check that $r + s = 1$ and $-2 \geq \bar{a}_{n_0} > a_{n_0}$ imply that there is some $t < m$ such that $-1/\bar{r}' = [b_1, \dots, b_t]$. Let us call $\bar{\Psi}'_1 \subseteq \Psi_2$ the linear connected subgraph with associated string of weights (b_1, \dots, b_t) . There are two possibilities, either $U_{\bar{\Psi}_1} = U_{\bar{\Psi}'_1}$ or $U_{\bar{\Psi}_1} \neq U_{\bar{\Psi}'_1}$.

Notice that by construction $|\bar{\Psi}_1| \leq |\Psi_1|$ and $|\bar{\Psi}'_1| < |\Psi_2|$. Moreover, if $|\bar{\Psi}_1| = 1$ [resp. $|\bar{\Psi}'_1| = 1$] then $\bar{\Psi}'_1$ [resp. $\bar{\Psi}_1$] is a (-2) -chain and it is immediate to check that in this case, since $e_1 \in U_{v_1} \cap U_{w_1}$, it holds $U_{\bar{\Psi}_1} = U_{\bar{\Psi}'_1}$. Since $\bar{r} + \bar{r}' = 1$, if $U_{\bar{\Psi}_1} \neq U_{\bar{\Psi}'_1}$ we can repeat the above construction with $\bar{\Psi}_1$ and $\bar{\Psi}'_1$ playing the role of Ψ_1 and Ψ_2 . It follows that, after a finite number of steps, we necessarily obtain from Ψ_1 and Ψ_2 two linear weighted graphs, which we still call $\bar{\Psi}_1$ and $\bar{\Psi}'_1$, such that $U_{\bar{\Psi}_1} = U_{\bar{\Psi}'_1}$ and either $\bar{\Psi}_1 \subseteq \Psi_1$ or $\bar{\Psi}'_1 \subseteq \Psi_2$. By symmetry we may assume that $\bar{\Psi}'_1 \subseteq \Psi_2$.

Since the strings of weights associated to $\bar{\Psi}_1$ and $\bar{\Psi}'_1$ are related to one another by Riemenschneider's point rule, we know, by the first part of this proof and using the same notation, that $|U_{\bar{\Psi}_1} \cup U_{\bar{\Psi}'_1}| = n_0 + t$. Consider the vector w_{t+1} . Since $U_{\bar{\Psi}_1} = U_{\bar{\Psi}'_1}$, $w_t \cdot w_{t+1} = 1$ and $w_{t+1} \cdot v_\ell = 0$ for every $\ell \in \{1, \dots, n_0\}$, the vector

$$\bar{w}_{t+1} := w_{t+1} + \sum_{e_i \notin U_{\bar{\Psi}_1}} (e_i \cdot w_{t+1}) e_i$$

satisfies $\bar{w}_{t+1} \cdot \bar{w}_{t+1} \leq -2$. It follows that the disconnected linear graph $\bar{\Psi}_1 \cup \bar{\Psi}'_1 \cup \{w_{t+1}\}$, which has $n_0 + t + 1$ vertices admits an embedding into a diagonal lattice of rank $|U_{\bar{\Psi}_1}| = n_0 + t$ which contradicts [13, Remark 2.1]. □

Lemma 3.2 Let $-1/r = [a_1, \dots, a_n]$ and $-1/s = [b_1, \dots, b_m]$ be such that $r + s > 1$. Then there exists $n_0 \leq n$ and $m_0 \leq m$ such that $-1/r_0 = [a_1, \dots, a_{n_0}]$ and $-1/s_0 = [b_1, \dots, b_{m_0}]$ satisfy $r_0 + s_0 = 1$.

Proof Let r' be such that $r + r' = 1$ and suppose $-1/r' = [a'_1, \dots, a'_{n'}]$. Since $s > r'$, by standard facts on continued fractions there are two possibilities: either $b_i = a'_i$ for all $i \in \{1, \dots, n'\}$ and $m > n'$ or there is a smallest index k such that $b_k > a'_k$. In the first case we set $n_0 = n$ and $m_0 = n'$. In the second case let us consider the first k columns of dots in the Riemenschneider's point diagram obtained from (a_1, \dots, a_n) . Then, n_0 equals the number of rows in this diagram minus $b_k - a'_k$ and $m_0 = k$. Note that in this way $[a_1, \dots, a_{n_0}]$ and $[b_1, \dots, b_{m_0}]$ are related to one another by Riemenschneider's point rule and therefore $r_0 + s_0 = 1$. \square

Lemma 3.3 Suppose $k \geq 3$ and $1 > r_1 \geq \dots \geq r_k > 0$ and $r_{k-1} + r_k > 1$. Then, the intersection lattice of the plumbing associated to $Y := Y(-k + 1; r_1, \dots, r_k)$ cannot be embedded into a negative diagonal standard lattice.

Proof Let Γ be the plumbing graph of Figure 6 associated to Y , and suppose by contradiction that there exists an embedding of $(\mathbb{Z}^{|\Gamma|}, Q_\Gamma)$ into $(\mathbb{Z}^d, -\text{Id})$ with basis e_1, \dots, e_d for some $d \geq |\Gamma|$. We will use the following notation for the vertices of Γ : v_0 for the central vertex and v_j^i for the vertices in the legs, where i indicates the leg to which v_j^i belongs and j the position in the leg, with $j = 1$ being the index of the vertex connected to the central vertex.

Since Γ has k legs connected to the central vertex which has weight $-k + 1$, there must exist some basis vector, say e_1 , and two legs, say L_i and L_j , such that the products $v_0 \cdot e_1, v_1^i \cdot e_1$ and $v_1^j \cdot e_1$ are not 0.

Let $-1/r_i = [a_1^i, \dots, a_{p_i}^i]$ and $-1/r_j = [a_1^j, \dots, a_{q_j}^j]$. Since $r_i + r_j > 1$, by Lemma 3.2 there exist $p_0 \leq p$ and $q_0 \leq q$ such that the strings $(a_1^i, \dots, a_{p_0}^i)$ and $(a_1^j, \dots, a_{q_0}^j)$ are related to one another by Riemenschneider's point rule. Moreover, since $e_1 \in U_{v_1^i} \cap U_{v_1^j}$ Lemma 3.1 applies and therefore the disconnected subgraph $\Psi \subseteq \Gamma$ consisting of the vertices $v_1^i, \dots, v_{p_0}^i, v_1^j, \dots, v_{q_0}^j$ satisfies $|U_\Psi| = p_0 + q_0$. Furthermore, writing $\Psi = \Psi_1 \cup \Psi_2$ where Ψ_1 [resp. Ψ_2] consists of the vertices $v_1^i, \dots, v_{p_0}^i$ [resp. $v_1^j, \dots, v_{q_0}^j$] we have, by Lemma 3.1(1), $U_{\Psi_1} = U_{\Psi_2}$.

Now, since $r_i + r_j > 1$ we have $(p_0, q_0) \neq (p, q)$ so we can assume without loss of generality $p > p_0$. The vector $v_{p_0+1}^i$ satisfies $v_{p_0+1}^i \cdot v_{p_0}^i = 1$ and the equality $U_{\Psi_1} = U_{\Psi_2}$ implies that $|U_\Psi \cap U_{v_{p_0+1}^i}| \geq 2$. Consider the vector

$$\bar{v}_{p_0+1} := v_{p_0+1}^i + \sum_{e_i \notin U_\Psi} (e_i \cdot v_{p_0+1}^i) e_i,$$

which by construction satisfies $U_{\bar{v}_{p_0+1}} \subseteq U_\Psi$ and $\bar{v}_{p_0+1} \cdot \bar{v}_{p_0+1} \leq -2$. It follows that the linear graph $\Psi \cup \{v_{p_0+1}\}$, which has $p_0 + q_0 + 1$ vertices admits an embedding into a diagonal lattice of rank $|U_\Psi| = p_0 + q_0$ which contradicts [13, Remark 2.1]. \square

We are now ready to prove Theorem 1.4. We restate the result for the reader's convenience:

Theorem 3.4 *A closed, oriented, Seifert fibered 3–manifold of special type admits no symplectic fillings.*

Proof Suppose that the oriented 3–manifold Y is orientation preserving diffeomorphic to $Y(-1; r_1, \dots, r_k)$, where $k \geq 3$, $1 > r_1 \geq r_2 \geq \dots \geq r_k > 0$ and (r_1, \dots, r_k) satisfies Conditions (1) and (2) of Definition 1.2. The fact that (r_1, \dots, r_k) is not realizable implies that Y is an L –space [18]. Therefore, if Y admits a symplectic filling W then $b_2^+(W) = 0$ [24, Theorem 1.4]. Moreover, since by [2; 22] (see [22, pages 157–158]), if $r_1 + \dots + r_k \leq 1$ then (r_1, \dots, r_k) is realizable, we must have $r_1 + \dots + r_k > 1$. Consider the space $-Y = Y(1-k; \bar{r}_1, \dots, \bar{r}_k)$ where $\bar{r}_i := 1 - r_{k-i+1}$ for $i \in \{1, \dots, k\}$. Since

$$e(-Y) := 1 - k + \sum_i \bar{r}_i = 1 - \sum_i r_i < 0,$$

by [23, Theorem 5.2] there is a negative definite plumbing graph Γ such that $-Y = \partial P_\Gamma$. Consider the 4–manifold X obtained gluing together P_Γ and W along their common boundary. By construction X is smooth, closed and negative, therefore by Donaldson's celebrated “Theorem A” [1] its associated intersection form must be diagonalizable. It follows that if Y admitted a symplectic filling then the intersection lattice of P_Γ would admit an embedding into a diagonal, negative standard lattice. The assumption $1 > r_1 + r_2$ for Y reads $\bar{r}_{k-1} + \bar{r}_k > 1$ for $-Y$, which by Lemma 3.3 implies that the intersection lattice of P_Γ does not admit an embedding into a diagonal lattice. Therefore we conclude that Y admits no symplectic fillings. \square

References

- [1] **S K Donaldson**, *The orientation of Yang–Mills moduli spaces and 4–manifold topology*, J. Differential Geom. 26 (1987) 397–428 MR910015
- [2] **D Eisenbud, U Hirsch, W Neumann**, *Transverse foliations of Seifert bundles and self-homeomorphism of the circle*, Comment. Math. Helv. 56 (1981) 638–660 MR656217
- [3] **Y Eliashberg**, *Filling by holomorphic discs and its applications*, from: “Geometry of low-dimensional manifolds, 2 (Durham, 1989)”, (S K Donaldson, C B Thomas, editors), London Math. Soc. Lecture Note Ser. 151, Cambridge Univ. Press (1990) 45–67 MR1171908

- [4] **Y Eliashberg**, *Topological characterization of Stein manifolds of dimension > 2* , Internat. J. Math. 1 (1990) 29–46 MR1044658
- [5] **JB Etnyre**, *Introductory lectures on contact geometry*, from: “Topology and geometry of manifolds (Athens, GA, 2001)”, (G Matic, C McCrory, editors), Proc. Sympos. Pure Math. 71, Amer. Math. Soc. (2003) 81–107 MR2024631
- [6] **H Geiges**, *An introduction to contact topology*, Cambridge Studies in Advanced Math. 109, Cambridge Univ. Press (2008) MR2397738
- [7] **P Ghiggini**, *Strongly fillable contact 3-manifolds without Stein fillings*, Geom. Topol. 9 (2005) 1677–1687 MR2175155
- [8] **P Ghiggini, P Lisca, A I Stipsicz**, *Tight contact structures on some small Seifert fibered 3-manifolds*, Amer. J. Math. 129 (2007) 1403–1447 MR2354324
- [9] **RE Gompf**, *Handlebody construction of Stein surfaces*, Ann. of Math. (2) 148 (1998) 619–693 MR1668563
- [10] **M Gromov**, *Pseudoholomorphic curves in symplectic manifolds*, Invent. Math. 82 (1985) 307–347 MR809718
- [11] **M Jankins, WD Neumann**, *Rotation numbers of products of circle homeomorphisms*, Math. Ann. 271 (1985) 381–400 MR787188
- [12] **P Lisca**, *Symplectic fillings and positive scalar curvature*, Geom. Topol. 2 (1998) 103–116 MR1633282
- [13] **P Lisca**, *Lens spaces, rational balls and the ribbon conjecture*, Geom. Topol. 11 (2007) 429–472 MR2302495
- [14] **P Lisca, G Matic**, *Transverse contact structures on Seifert 3-manifolds*, Algebr. Geom. Topol. 4 (2004) 1125–1144 MR2113899
- [15] **P Lisca, A I Stipsicz**, *An infinite family of tight, not semi-fillable contact three-manifolds*, Geom. Topol. 7 (2003) 1055–1073 MR2026538
- [16] **P Lisca, A I Stipsicz**, *Ozsváth–Szabó invariants and tight contact three-manifolds. I*, Geom. Topol. 8 (2004) 925–945 MR2087073
- [17] **P Lisca, A I Stipsicz**, *Tight, not semi-fillable contact circle bundles*, Math. Ann. 328 (2004) 285–298 MR2030378
- [18] **P Lisca, A I Stipsicz**, *Ozsváth–Szabó invariants and tight contact 3-manifolds. III*, J. Symplectic Geom. 5 (2007) 357–384 MR2413308
- [19] **P Lisca, A I Stipsicz**, *Ozsváth–Szabó invariants and tight contact three-manifolds. II*, J. Differential Geom. 75 (2007) 109–141 MR2282726
- [20] **P Lisca, A I Stipsicz**, *On the existence of tight contact structures on Seifert fibered 3-manifolds*, Duke Math. J. 148 (2009) 175–209 MR2524494
- [21] **A Loi, R Piergallini**, *Compact Stein surfaces with boundary as branched covers of B^4* , Invent. Math. 143 (2001) 325–348 MR1835390

- [22] **R Naimi**, *Foliations transverse to fibers of Seifert manifolds*, Comment. Math. Helv. 69 (1994) 155–162 MR1259611
- [23] **W D Neumann, F Raymond**, *Seifert manifolds, plumbing, μ -invariant and orientation reversing maps*, from: “Algebraic and geometric topology (Proc. Sympos., Univ. California, Santa Barbara, Calif., 1977)”, (K C Millett, editor), Lecture Notes in Math. 664, Springer, Berlin (1978) 163–196 MR518415
- [24] **P Ozsváth, Z Szabó**, *Holomorphic disks and genus bounds*, Geom. Topol. 8 (2004) 311–334 MR2023281
- [25] **P Ozsváth, Z Szabó**, *On knot Floer homology and lens space surgeries*, Topology 44 (2005) 1281–1300 MR2168576
- [26] **O Riemenschneider**, *Deformationen von Quotientensingularitäten (nach zyklischen Gruppen)*, Math. Ann. 209 (1974) 211–248 MR0367276

*Dipartimento di Matematica “L Tonelli”, Università di Pisa
Largo Bruno Pontecorvo 5, I-56127 Pisa, Italy*

lecuona@mail.dm.unipi.it, lisca@dm.unipi.it

Received: 23 September 2010