

Toda brackets and congruences of modular forms

GERD LAURES

This paper investigates the relation between Toda brackets and congruences of modular forms. It determines the f -invariant of Toda brackets and thereby generalizes the formulas of JF Adams for the classical e -invariant to the chromatic second filtration.

11F33, 55N34, 55T15; 55Q45

1 Introduction

In this work congruences between modular forms are used to compute Toda brackets in the second Adams–Novikov filtration. The translation from stable homotopy classes between spheres and modular forms is based on the tertiary invariant

$$f: \pi_{2n-2}^{\text{st}} \longrightarrow (D/M_0 + M_n)_{\mathbb{Q}/\mathbb{Z}}$$

defined by the author in [19] with values in Katz's ring of divided congruences. The f -invariant generalizes the classical e invariant of Frank Adams. The purpose of this work is an extension of Adams' formulas [1, Section 11] for the e -invariant of Toda brackets to the second filtration:

$$\begin{aligned} e \langle \alpha, p\iota, \beta \rangle &= -pe(\alpha)e(\beta) \\ e \langle p\iota, \alpha, \beta \rangle &= -p\delta e(\alpha)e(\beta) \\ e \langle \alpha, \beta, p\iota \rangle &= -p\delta e(\alpha)e(\beta) \end{aligned}$$

modulo the indeterminacy. Here, δ is a certain number which only depends on the dimensions.

The e -invariant is not strong enough to detect elements in the second filtration. The reason is the following: the e -invariant is based on K -theory and in the K -based Adams–Novikov spectral sequence there aren't any classes in the second line other than classes in the image of J . In [19] K -theory was replaced by elliptic cohomology theory of level N (see Brylinski [11] and Franke [14]). It was shown that the corresponding tertiary invariant f is injective on the 2-line. So, it is natural to ask for a precise relationship between the f -invariant and Toda brackets.

The 2–line of the classical Adams–Novikov spectral sequence was computed for odd primes by Miller, Ravenel and Wilson in [24] with the help of the chromatic spectral sequence and by Shimomura for $p = 2$ in [27]. In contrast to the 1–line it is not cyclic. The number of generators is unbounded when considered as a function of the dimension. Its complicated structure makes it hard to determine classes such as Massey products. The f –invariant simplifies the 2–line in two steps: first the complex orientation gives an injective map into the elliptic based E_2 term and in the second step the extension group is mapped to Katz’s ring of divided congruences. In this paper it is reinterpreted as a coboundary of a resolution which can be viewed as an integral version of a chromatic resolution.

The paper starts with a recollection on higher homotopy invariants in general. Here, the edge homomorphisms are used to construct invariants for stable homotopy classes with the help of any theory E . Under mild conditions the first few invariants take values in the E –based Adams–Novikov E_2 –term. Then the relation between Toda brackets and Massey products is explained. Adams considered Massey products for extensions of exact sequences. However, it turns out that the cobar complex is better suited to keep the homological algebra small for higher filtrations. This holds especially when a \mathbb{Q}/\mathbb{Z} –reduction is used to come down by one filtration. It follows a computational section where the commutation between the coboundary operator and Toda brackets is investigated. Proposition 5.2 gives a precise translation of the elliptic Adams–Novikov E_2 term by primitives in the ring of divided congruences. It comes from a resolution which is closely related to the chromatic resolution. Finally, the desired formulas for the f –invariants of Toda brackets are given. In the last section some examples are calculated including the Kervaire class in dimension 30 as 4–fold Toda bracket.

2 Higher Homotopy invariants

Suppose E is a flat ring spectrum. Let \bar{E} be the fibre of the unit map $S^0 \rightarrow E$. Then there is spectral sequence (E_r, d_r) with the associated filtration of π_*^S given for $s \geq 0$ by

$$F^s = \text{im}(\pi_* \bar{E}^s \rightarrow \pi_* S^0)$$

The differential d_r raises the filtration by r . Hence there are well defined ‘edge homomorphisms’

$$e_s: F^s \rightarrow F^s / F^{s+1} \twoheadrightarrow E_\infty^s \twoheadrightarrow E_{s+1}^s.$$

For a stable class α the value $e_s(\alpha)$ is defined if and only if $e_r(\alpha)$ vanishes for all $r < s$. It associates to α its representative in the E_{s+1}^s –term. The invariants are natural with respect to maps between spectral sequences.

In case E is complex bordism MU this is the classical Adams–Novikov spectral sequence. Its quotient group F^0/F^1 is concentrated in dimension 0 and the invariant

$$e_0: \pi_0^S = F^0 \longrightarrow \pi_0(MU) = \mathbb{Z}$$

takes values in the integers. It associates to a stable self map of S^0 its degree d . In fact we may replace E by any other spectrum with the property that the Hurewicz homomorphism is injective in dimension 0. If E has torsion coefficients the map e_0 can be non trivial in positive dimensions. For instance this happens for real K -theory in dimensions $8k + 1$ and $8k + 2$.

The invariant e_1 has been studied by Adams for real and complex K -theory in [1]. It can be described in terms of exact sequences as follows: Suppose the e_0 invariant of $\alpha \in \pi_n^S$ vanishes and let C_α be the cofibre of α . Then the sequence

$$E_*S^0 \twoheadrightarrow E_*C_\alpha \twoheadrightarrow E_*S^{n+1}$$

of E_*E -comodules is short exact and hence determines an element in

$$E_2^{1,n+1} \cong \text{Ext}_{E_*E}^{1,n+1}(E_*, E_*).$$

Adams determines the extension group for K -theory with the help of a monomorphism

$$\iota: E_2^{1,n+1} \twoheadrightarrow \mathbb{Q}/\mathbb{Z}.$$

In particular, the extension term is cyclic. For K -theory its order in dimension $n = 4t - 1$ is the denominator of the divided Bernoulli number $B_t/2t$. The composite ιe_1 is called the classical e -invariant.

For variations of the topological modular forms spectrum TMF the invariant e_2 has been studied (in chronological order) by the author [19; 20], by Hornborstel–Naumann [15], by Behrens–Laures in [7] and by von Bodecker [9; 8]. For its relation to characteristic classes of manifolds with corners see Laures [20] and for an interpretation as a spectral invariant see von Bodecker [10] and Bunke–Naumann [12].

Before describing the invariant we first take a look at the case of complex bordism. Here, for positive even n the e_1 -invariant vanishes. Moreover, there are no d_2 -differentials hitting the 2-line. Thus the E_3^2 -term injects into the E_2^2 term. This yields a well defined homomorphism

$$e_2: \pi_n^S \longrightarrow E_2^{2,n+2} = \text{Ext}_{E_*E}^{2,n+2}(E_*, E_*).$$

Let us describe this invariant in terms of exact sequences in more detail: observe that any lift $\bar{\alpha}$ of a stable class α is unique since E_* and E_*E are evenly graded. Moreover, the sequence

$$E_*\Sigma\bar{E} \twoheadrightarrow E_*\Sigma C_{\bar{\alpha}} \twoheadrightarrow E_*S^{n+2}$$

is short exact and can be spliced together with

$$E_*S^0 \twoheadrightarrow E_*E \twoheadrightarrow E_*\Sigma\bar{E}$$

to get an extension of degree 2.

Next suppose E is complex oriented with (p, v_1) regular and v_2 invertible mod (p, v_1) . Then locally at p the 2–line of E coincides with the 2–line of $v_2^{-1}MU_{(p)}$ (see Laures [20, 4.3.3]) and there is a well understood injection (see Ravenel [25])

$$E_2^2(MU_{(p)}) \twoheadrightarrow E_2^2(v_2^{-1}MU_{(p)}) \cong E_2^2(E).$$

In case of $\Gamma_1(N)$ topological modular forms $E = TMF_1(N)$ there is an injection

$$E_2^{2,2t} \twoheadrightarrow (D/M_0 + M_t)_{\mathbb{Q}/\mathbb{Z}}$$

where D is the ring of divided congruences and M the ring of modular forms. This map will be reviewed in Section 5. Furthermore, Proposition 5.2 describes its precise image.

The e_3 –invariant is defined for odd dimensional homotopy classes in the cokernel of the J –homomorphism. It takes values in the E_2 –term since all elements in the 1–line are permanent cycles. Hence the invariant

$$e_3: \text{coker } J \longrightarrow E_2^3 = \text{Ext}_{E_*E}^3(E_*, E_*)$$

is well defined. It will not be further investigated here.

3 Toda brackets and Massey products

In the sequel we assume that E is a complex oriented flat ring spectrum. The E_1 –term of the E –based Adams–Novikov spectral sequence is the cobar complex Ω_* which is a differential graded algebra. We use the notation Γ for the comodule E_*E over $A = E_*$ and $\bar{\Gamma}$ for the augmentation ideal $E_*\Sigma\bar{E}$. Then the differential

$$d_1: \bar{\Gamma}^{\otimes s} \longrightarrow \bar{\Gamma}^{\otimes(s+1)}$$

sends $[\alpha_1 | \cdots | \alpha_s]$ to $[1|\alpha_1 | \cdots | \alpha_s] - [\alpha'_1 | \alpha''_1 | \cdots | \alpha_s] + \cdots + (-1)^{s+1}[\alpha_1 | \cdots | \alpha_s | 1]$ and the product is given by concatenation. In fact, each (E_r, d_r) is a differential graded algebra in which Massey products can be defined (see Ravenel [25, Appendix]). Massey products in the cobar complex are related to Toda brackets in stable homotopy. This relation has been studied by Adams, Moss, Lawrence, May, Kochman and others and is well known.

Recall the following definition from Kochman [18].

Definition 3.1 Suppose $u \in E_r^{s,t+s}$ and $w \in E_{r'}^{s',t+s'}$ with $s' < s$. Then $d_{r'}(w)$ is a crossing differential of $d_r(u)$ if $s' + r' > s + r$.

The following result reformulates the result [18, 5.7.5] for the homotopy invariants of the last section.

Proposition 3.2 Suppose that the Toda bracket $\langle \alpha_1, \dots, \alpha_n \rangle$ is strictly defined. Let a_i be a representative of $e_{s(i)}(\alpha_i)$ in the MU -based $E_{r+1}^{s(i)}$ for some r . Suppose further that the Massey product $\langle a_1, \dots, a_n \rangle$ is defined in E_{r+1} and that there are no crossing differentials of $d_r a_{ij}$ for all defining systems (a_{ij}) .

Then for $s = \sum_i s(i)$ there is a class $\alpha \in \langle \alpha_1, \dots, \alpha_n \rangle$ of filtration $s - n + 2$ with

$$e_{s-n+2}(\alpha) \in \langle a_1, \dots, a_n \rangle.$$

Proof Since a_i converges to α_i we can apply of [18, 5.7.5]. Hence $\langle a_1, \dots, a_n \rangle$ consists of infinite cycles which converge to elements of $\langle \alpha_1, \dots, \alpha_n \rangle$. Since all elements have filtration $s - n + 2$ the claim follows. \square

Corollary 3.3 Under the assumptions of Proposition 3.2 we have

$$e_{s-n+2}(\alpha) \in \langle a_1, \dots, a_n \rangle$$

for all spectral sequences based on a complex oriented theory E .

Proof This follows from the naturality of all constructions. \square

The proposition suggests that for the e_k -invariant of n -fold Toda brackets one should consider only sums of filtrations $s = k + n - 2$. The cases for $n = 3$ and $k = 0, 1$ were considered by Adams in [1, Theorem 5.3].

The case $k = 2$ is the object of the rest of the paper. In the next section triple Massey products are considered. Note that in this case there aren't any crossing differentials in the defining system as long as there aren't any r -boundaries in the 3-line for $r \geq 2$. Moreover, if the Toda bracket $\langle \alpha_1, \alpha_2, \alpha_3 \rangle$ is defined then so is the Massey product $\langle a_1, a_2, a_3 \rangle$.

Corollary 3.4 (i) Suppose that the Toda bracket $\langle \alpha_1, \alpha_2, \alpha_3 \rangle$ is defined. Then

$$e_2 \langle \alpha_1, \alpha_2, \alpha_3 \rangle = \langle e_{s_1} \alpha_1, e_{s_2} \alpha_2, e_{s_3} \alpha_3 \rangle$$

with indeterminacy

$$e_{s_1} \alpha_1 H^{s_2+s_3-1, |\alpha_2|+s_2+|\alpha_3|+s_3}(\Omega_*) + H^{s_1+s_1-1, |\alpha_1|+s_1+|\alpha_2|+s_2}(\Omega_*) e_{s_3} \alpha_3.$$

- (ii) Suppose that the Toda bracket $\langle \alpha_1, \alpha_2, \alpha_3, \alpha_4 \rangle$ is strictly defined and that there are no crossing differentials. Then

$$e_2 \langle \alpha_1, \alpha_2, \alpha_3, \alpha_4 \rangle = \langle e_{s_1} \alpha_1, e_{s_2} \alpha_2, e_{s_3} \alpha_3, e_{s_4} \alpha_4 \rangle$$

with indeterminacy

$$\sum_{a,b,c} \langle a, e_{s_3} \alpha_3, e_{s_4} \alpha_4 \rangle + \langle e_{s_1} \alpha_1, b, e_{s_4} \alpha_4 \rangle + \langle e_{s_1} \alpha_1, e_{s_2} \alpha_2, c \rangle.$$

Proof Since the invariant e_2 is multiplicative it maps the indeterminacy of the Toda bracket

$$\alpha_1 \pi_{|\alpha_2|+|\alpha_3|+1}^S + \pi_{|\alpha_1|+|\alpha_2|+1}^S \alpha_3$$

to the indeterminacy of the Massey product stated above. The same holds for the 4-fold products for which the indeterminacy can be found for the Toda bracket in Kochman [17, Theorem 2.3.1] and for the Massey product in May [23, Proposition 2.4]. □

4 The \mathbb{Q}/\mathbb{Z} -reduction

Let (A, Γ) be a Hopf algebroid. In this section we assume that (A, Γ) has the following properties:

- (i) Γ is flat as an A -module.
- (ii) A and Γ are torsion free.
- (iii) The map $\phi: A_{\mathbb{Q}} \otimes A_{\mathbb{Q}} \rightarrow \Gamma_{\mathbb{Q}}$ which sends $a \otimes b$ to $a\eta_R(b)$ is an isomorphism.

Define the cosimplicial abelian group $\Omega_{\mathbb{Q}}^n = A_{\mathbb{Q}}^{n+1}$ with cofaces

$$\partial^i(a_0 \otimes \cdots \otimes a_n) = a_0 \otimes \cdots \otimes a_{i-1} \otimes 1 \otimes a_{i+1} \otimes \cdots \otimes a_n.$$

The notation is justified to the fact that the map $\phi^{\otimes n}$ provides an isomorphism between the complex $\Omega_{\mathbb{Q}}$ and the rationalized cobar complex. Its cohomology is concentrated in dimension 0

$$H^*(\Omega_{\mathbb{Q}}^*) = H^0(\Omega_{\mathbb{Q}}^*) = \mathbb{Q}.$$

Its algebra structure is given by

$$(a_0 \otimes \cdots \otimes a_n) \otimes (b_0 \otimes \cdots \otimes b_m) \mapsto a_0 \otimes \cdots \otimes a_{n-1} \otimes a_n b_0 \otimes b_1 \otimes \cdots \otimes b_m.$$

The short exact sequence

$$\Omega^* \twoheadrightarrow \Omega_{\mathbb{Q}}^* \twoheadrightarrow \Omega_{\mathbb{Q}/\mathbb{Z}}^*$$

yields a connecting isomorphism $\delta: H^{n-1}(\Omega_{\mathbb{Q}/\mathbb{Z}}^*) \rightarrow H^n(\Omega^*)$ in positive dimensions. Choose an augmentation $\tau: \Omega_{\mathbb{Q}} \rightarrow \mathbb{Q}$. Then the following observation is easily verified.

Lemma 4.1 *Let $a \in \Omega^n$ be a cycle and suppose $\sum a_0 \otimes \cdots \otimes a_n \in \Omega_{\mathbb{Q}}^n$ is its rationalization. Then we have*

$$\delta^{-1}[a] = \left[\sum \tau(a_0)a_1 \otimes a_2 \otimes \cdots \otimes a_n \right].$$

Moreover, when writing $b' = \sum \tau(b_0)b_1 \otimes \cdots \otimes b_n$ for $b \in \Omega_{\mathbb{Q}}^n$ we have

$$(db)' = b - db'.$$

We are going to study the image of triple Massey products $\langle x, y, z \rangle$ under δ^{-1} . For $n = 2$ there are three cases of interest:

$$(|y| = 0, |x| + |z| = 3), \quad (|y| = 1, |x| + |z| = 2) \quad \text{and} \quad (|y| = 2, |x| + |z| = 1).$$

We start with the first case.

Definition 4.2 Suppose $\delta[a] \in H^2(\Omega^*)$ is represented by some $a \in A_{\mathbb{Q}} \otimes A_{\mathbb{Q}}$. We say that a is q -adapted for some $q \in \mathbb{Z}$ if $qa \in \Gamma$.

Lemma 4.3 *q -torsion classes admit q -adapted representatives.*

Proof Choose a representative a which is not yet q -adapted. Since $\delta[a]$ is q -torsion the element qa represents a boundary in $\Omega_{\mathbb{Q}/\mathbb{Z}}^1 = \Gamma_{\mathbb{Q}/\mathbb{Z}}$. Hence there is $r \in A_{\mathbb{Q}}$ with

$$\tilde{a} = qa + dr \in \Gamma.$$

and \tilde{a}/q is a q -adapted representative. □

Proposition 4.4 *Suppose that for some $q \in \mathbb{Z}$ the Massey product $\langle \delta[a], q, \delta[c] \rangle$ is defined.*

(i) *For $||a|| = 1, ||c|| = 0$ and a q -adapted we have*

$$\delta[qac] \in \langle \delta[a], q, \delta[c] \rangle.$$

(ii) *For $||a|| = 0, ||c|| = 1$ and c q -adapted we have*

$$\delta[-qac] \in \langle \delta[a], q, \delta[c] \rangle.$$

Proof For (i) observe that $\delta[a]$ is represented by $da \in \Gamma^{\otimes 2}$. Hence $q da$ is the boundary of $qa \in \Gamma$. Similarly, let $dc \in \Gamma$ then $q dc$ is the boundary of $qc \in A$. Thus we obtain the representative of the Massey product

$$[da qc - qa dc] \in \langle \delta[a], q, \delta[c] \rangle.$$

The lemma tells us that the desired class is obtained by the applying τ to the first factor in its tensor product expression. Alternatively, one writes the representative as $d(qac)$ and obtains the result. The second formula follows from a similar calculation. \square

We turn to the other cases.

Lemma 4.5 For $s, t \geq 1$ the product of classes $\delta[a] \in H^s \Omega^*$, $\delta[b] \in H^t \Omega^*$ vanishes if and only if there is an $r \in A_{\mathbb{Q}}^{\otimes s+t-1}$ with

$$(a db + dr) \in \Gamma^{s+t-1}.$$

We write $r = r(a, b)$ for any such element.

Proof If $\delta[a]\delta[b] = 0$ there is $x \in \Gamma^{s+t-1}$ with

$$dx = da db.$$

Applying $\tau 1 \otimes \cdots \otimes 1$ gives

$$x - dx' = (a - da')db$$

and we set $r = x' - a' db$. Conversely, we have

$$d(a db + dr) = da db. \quad \square$$

Proposition 4.6 Suppose that the Massey product $\langle \delta[a], \delta[b], \delta[c] \rangle$ is defined. Then the following holds:

(i) For $|a| = |b| = |c| = 0$,

$$\delta[a(b dc + dr(b, c)) + r(a, b)dc] \in \langle \delta[a], \delta[b], \delta[c] \rangle.$$

(ii) For $|a| = 1$, $|b| = 0$ and $c = q \in \mathbb{Z}$,

$$\delta[q(ab + r(a, b))] \in \langle \delta[a], \delta[b], q \rangle.$$

(iii) For $a = q \in \mathbb{Z}$, $|b| = 0$ and $|c| = 1$,

$$\delta[-qr(b, c)] \in \langle q, \delta[b], \delta[c] \rangle.$$

(iv) If $\langle q, \delta[b], \delta[c] \rangle$ with $|b| = 1, |c| = 0$ is defined and b is q -adapted then

$$\delta[qr(b, c)] \in \langle q, \delta[b], \delta[c] \rangle$$

(v) If $\langle \delta[a], \delta[b], q \rangle$ with $|b| = 1, |a| = 0$ is defined and b is q -adapted then

$$\delta[-q(ab + r(a, b))] \in \langle \delta[a], \delta[b], q \rangle.$$

Proof By the lemma a representative of the first Massey product is

$$da(bdc + dr(b, c)) + (adb + dr(a, b))dc.$$

Applying $\tau 1 \otimes 1$ gives

$$a(bdc + dr(b, c)) + r(a, b)dc$$

as claimed. For (ii) a representative is

$$daqb + (adb + dr(a, b))q.$$

In the last case it is

$$-q(bdc + dr(b, c)) + qbdc.$$

Again applying $\tau 1 \otimes 1$ gives the results up to boundary terms. The other Massey products are obtained with the help of the lemma as before. In the case (iv) one gets

$$q(bdc + dr) + (-qb)dc = qdr$$

and in the last case

$$-daqb + (-adb - dr)q = -q(d(ab) + dr). \quad \square$$

We close this section with an example of a 4-fold Massey product. Note that in principle other types and even higher Massey products can be calculated with the same method.

Proposition 4.7 *Suppose a and b have degree 1 and are q -adapted. Then the following conditions are equivalent:*

- (i) the Massey product $\langle q, \delta[a], q, \delta[b] \rangle$ is defined
- (ii) $0 \in \langle \delta[a], q, \delta[b] \rangle$
- (iii) there are $r \in A_{\mathbb{Q}} \otimes A_{\mathbb{Q}}$ and $dv, dw \in \Gamma$ with $qab + adw + dvb + dr \in \Gamma^{\otimes 2}$.

In this case the 4-fold product contains $\delta(-qr + vdw)$. Moreover, the same assertions hold for $\langle \delta[a], q, \delta[b], q \rangle$.

Proof A defining system can look like

$$\begin{array}{cccc}
 q & & da & & q & & db \\
 & & -qa-dv & & -qa-dv & & -qb-dw \\
 & & 0 & & x & &
 \end{array}$$

In particular a defining system exists if there is a $x \in \Gamma^{\otimes 2}$ with

$$dx = qd(ab) + dv db + da dw.$$

Applying $\tau 1 \otimes 1 \otimes 1$ gives $x = qab + dv b + a dw + dr$ for some r . This shows the equivalences. Finally, if x is as above the missing corner is $-qr + v dw$ up to a boundary. □

5 Invariants in modular forms

For a $\mathbb{Z}[1/N]$ -algebra R let $M_k(\Gamma_1(N))_R$ be the ring of $\Gamma_1(N)$ modular forms of weight k over $R[\zeta_N]$ which are meromorphic at the cusps. Let $TMF_1(N)$ denote the corresponding spectrum of topological modular forms. Its coefficients $\pi_* TMF_1(N)$ are concentrated in even degrees, and we have

$$\pi_{2k} TMF_1(N) \cong M_k(\Gamma_1(N))_{\mathbb{Z}[1/N]}.$$

The spectrum $TMF_1(N)$ is complex orientable with formal group isomorphic to the formal completion of the universal elliptic curve over the ring of $\Gamma_1(N)$ modular forms. In fact, it is Landweber exact (see the author’s paper [19, Theorem 1.4, page 390f]). Here and in some other places of the older literature as Brylinsky [11], Franke [14] and Baker [2] it carried the names $Ell^{\Gamma_1(N)}$ or $E^{\Gamma_1(N)}$. Nowadays one uses the notation $TMF_1(N)$ since it is a complex oriented relative of the spectrum TMF of topological modular forms. The theory TMF can be obtained as global sections of a sheaf of E_∞ spectra over some moduli space of elliptic curves. This interpretation will not be used for the theory $TMF_1(N)$ in the article at hand. However, we mention the articles by Behrens [6] and Lawson–Naumann [22] where this line of thought has been pursued. Since the congruence subgroup $\Gamma_1(N)$ will be fixed once and for all we will remove it from the notation and write TMF instead of $TMF_1(N)$ and M_k for $M_k(\Gamma_1(N))$.

When localized at a prime the theory TMF has the same Adams Novikov E_2 term as the the Johnson–Wilson theories $E(2)$ by Laures [20, Lemma 4.3.3]. Moreover, it has the advantage that it furnishes an interpretation of elements in the Adams–Novikov

E_2 -term in terms of congruences between modular forms. These have been extensively studied in the literature.

Let $D = D(\Gamma_1(N))$ be the ring of divided congruences (see Katz [16]). An element in D is a sum $\sum f_k$ of modular forms $f_k \in (M_k)_{\mathbb{Q}}$ with an integral q -expansion, that is, an expansion with coefficients in $\mathbb{Z}[1/N, \zeta_N]$. The following key result is well known (see Baker [3] for $p \neq 2, 3$ or Laures [20; 21] for the general case.)

Theorem 5.1 *The K -homology of TMF vanishes in odd degrees and there is an isomorphism*

$$\pi_0 K \wedge TMF \cong D.$$

Set $E = TMF$. Consider the resolution of E_* as E_*E comodule

$$\begin{array}{ccccc}
 & E_* \otimes \mathbb{Q} & \dashrightarrow & E_* \Sigma K \otimes \mathbb{Q}/\mathbb{Z} & \dashrightarrow & E_* \Sigma \bar{K} \otimes \mathbb{Q}/\mathbb{Z} \\
 & \nearrow & & \nearrow & & \nearrow \\
 E_* & & & E_* \otimes \mathbb{Q}/\mathbb{Z} & & E_* \Sigma \bar{K} \otimes \mathbb{Q}/\mathbb{Z} \\
 & \searrow & & \searrow & & \searrow \\
 & & & & & E_* \Sigma \bar{K} \otimes \mathbb{Q}/\mathbb{Z}
 \end{array}$$

=

The middle sequence is exact since $E_* = M_*$ is a pure subgroup of $E_*K = D$. The short exact sequences provide connecting homomorphisms δ for the extension groups.

Proposition 5.2 *Let $P(M) = \text{Hom}_{E_*E}(E_*, M)$ denote the primitives of a comodule M . Then the sequence*

$$\pi_* \Sigma \bar{K} \otimes \mathbb{Q}/\mathbb{Z} \longrightarrow P(E_* \Sigma \bar{K} \otimes \mathbb{Q}/\mathbb{Z}) \xrightarrow{\delta^2} \text{Ext}_{E_*E}^2(E_*, E_*) \longrightarrow 0$$

is exact. Moreover the f -invariant is the composite of e_2 with the inclusion

$$\text{Ext}_{E_*E}^{2, 2n}(E_*, E_*) \cong P(E_{2n-1} \bar{K})/\pi_{2n-1} \bar{K} \otimes \mathbb{Q}/\mathbb{Z} \hookrightarrow (D/M_0 + M_n)_{\mathbb{Q}/\mathbb{Z}}$$

Proof It is not hard to see that the cohomology of $E_* \otimes \mathbb{Q}$ is concentrated in degree and dimension 0. In fact, the map τ gives a contracting homotopy. Hence, for positive dimension we have the isomorphism

$$\delta: \text{Ext}^1(E_*, E_* \otimes \mathbb{Q}/\mathbb{Z}) \longrightarrow \text{Ext}^2(E_*, E_*).$$

Furthermore, the middle short exact sequence of the resolution gives the exact sequence

$$P(E_* K \otimes \mathbb{Q}/\mathbb{Z}) \longrightarrow P(E_* \Sigma \bar{K} \otimes \mathbb{Q}/\mathbb{Z}) \xrightarrow{\delta} \text{Ext}^1(E_*, E_* \otimes \mathbb{Q}/\mathbb{Z}).$$

We claim that δ is surjective in positive dimensions. For that it suffices to show that the map into $\text{Ext}^1(E_*, E_*K \otimes \mathbb{Q}/\mathbb{Z})$ vanishes. This map admits a factorization

$$\text{Ext}^1(E_*, E_* \otimes \mathbb{Q}/\mathbb{Z}) \longrightarrow \text{Ext}^1(E_*, E_*E \otimes \mathbb{Q}/\mathbb{Z}) \longrightarrow \text{Ext}^1(E_*, E_*K \otimes \mathbb{Q}/\mathbb{Z})$$

induced by the ring map $\chi_0: E \rightarrow K$ in E -homology. Since the middle term vanishes we have shown that δ is surjective.

It remains to identify the first map. We claim that the map

$$\pi_*K \otimes \mathbb{Q}/\mathbb{Z} \longrightarrow P(E_*K \otimes \mathbb{Q}/\mathbb{Z})$$

is an isomorphism. Let K_T be the elliptic theory associated to the Tate curve. Its coefficients are integral Laurent series in a variable q in all even degrees and they vanish in odd degrees. There is the Miller character, that is, a ring map $\chi: E \rightarrow K_T$ which is the q -expansion map on coefficients. Consider the injective map

$$(\chi \wedge 1)_*: E_*K \otimes \mathbb{Q}/\mathbb{Z} \longrightarrow K_{T*}K \otimes \mathbb{Q}/\mathbb{Z}$$

which takes q -expansions. A primitive in its source gives a primitive in the target if the target is viewed as a comodule over $K_{T*}K_T$. The primitives in the target all lie in the primitive of the extended comodule $K_{T*}K_T \otimes \mathbb{Q}/\mathbb{Z}$ and hence in $\pi_*K_T \otimes \mathbb{Q}/\mathbb{Z}$. Since they also lie in $K_{T*}K \otimes \mathbb{Q}/\mathbb{Z}$ they must lie in $(\pi_*K) \otimes \mathbb{Q}/\mathbb{Z}$. This group coincides with $\pi_*(\Sigma \bar{K}) \otimes \mathbb{Q}/\mathbb{Z}$ in positive dimension and hence the claim follows.

The f -invariant is the composite of e_2 with

$$H^2\Omega_* \twoheadrightarrow \Omega_2/d\Omega_1 \xrightarrow{\tau_1 \otimes 1} \Gamma/d(M_n) \otimes \mathbb{Q}/\mathbb{Z} \xrightarrow{1 \wedge \chi_0} D/(M_0 + M_n)_{\mathbb{Q}/\mathbb{Z}}.$$

As we have seen before the second map gives an inverse of the connecting homomorphism. Hence the claim follows from the commutative diagram

$$\begin{array}{ccc} H^1(\Omega_{\mathbb{Q}/\mathbb{Z}}) & \xleftarrow{\cong \cong} P(E_*\Sigma \bar{K} \otimes \mathbb{Q}/\mathbb{Z})/\sim & \longrightarrow D/(M_0 + M_n)_{\mathbb{Q}/\mathbb{Z}} \\ \uparrow = & \uparrow 1 \wedge \chi_0 & \uparrow 1 \wedge \chi_0 \\ H^1(\Omega_{\mathbb{Q}/\mathbb{Z}}) & \xleftarrow{\cong} P(\bar{\Gamma} \otimes \mathbb{Q}/\mathbb{Z})/\sim & \longrightarrow \Gamma/d(M_n)_{\mathbb{Q}/\mathbb{Z}} \end{array}$$

in which a 1-cycle in the cobar complex of the left lower corner is sent to itself under the composite of the bottom row. □

Remark 5.3 We will not study the primitives of

$$E_*\Sigma \bar{K} = (D/M_n)_{\mathbb{Q}/\mathbb{Z}}$$

here since it is not needed for Toda brackets and the rest of this work. However, we mention that the primitives are eigenforms under the action of the Hecke operations and they are fixed under the action of the Adams operations (see Baker [4, 1.1]). This suggests that the elements of the 2–line in the cokernel of J are related to newforms. A precise relationship locally at primes $p \geq 5$ between the 2–line and certain p –adic modular forms is described in the paper by Behrens [6].

The following result is due to Bunke and Naumann. It is very useful for explicit calculations.

Lemma 5.4 (i) *There exists a \mathbb{Z} –basis $f_0 \dots f_{n_k}$ of M_k such that $q^i(f_j) = \delta_{ij}$ for all i, j .*

(ii) *For a basis as above, the map*

$$(D/M_k)_{\mathbb{Q}/\mathbb{Z}} \longrightarrow \prod_{i \geq n_k} \mathbb{Q}/\mathbb{Z}, \quad f \mapsto \left(q^v \left(f - \sum_{i=0}^{n_k-1} q^i(f) f_i \right) \right)_{v \geq n_k}$$

is injective.

Proof The first part is Bunke–Naumann [13, Lemma 9.2] and the second part is stated there in terms of q –expansions. □

Clearly, it suffices to check finitely many Fourier coefficients for a modular form in the source. There are upper bounds for this number but we do not work them out here.

6 The f –invariant of Toda brackets

In this section we apply our formulas to the f –invariant of Toda brackets. We start with the simplest case.

Theorem 6.1 *Suppose that the Toda bracket $\langle \alpha, p\iota, \beta \rangle$ is defined for some α, β in positive dimensions m, n and some $p \in \mathbb{Z}$.*

(i) *Let the f –invariant of α be defined. Choose a representative of $f(\alpha)$ whose q –expansion is annihilated by p . Then we have*

$$f \langle \alpha, p\iota, \beta \rangle = pe(\beta) f(\alpha)$$

with indeterminacy $e(\beta)H^{0,m+2}(\Omega_{\mathbb{Q}/\mathbb{Z}}^)$.*

- (ii) Let the f -invariant of β be defined. Choose a representative of $f(\alpha)$ whose q -expansion is annihilated by p . Then we have

$$f \langle \alpha, p\iota, \beta \rangle = -pe(\alpha)f(\beta)$$

with indeterminacy $e(\alpha)H^{0,n+2}(\Omega_{\mathbb{Q}/\mathbb{Z}}^*)$.

Proof First note in view of Corollary 3.4 that the indeterminacy in the theorem coincides with the image under the f -invariant of the Massey product indeterminacy. For (i) we can choose a p -adapted representative a with $\delta[a] = e_2(\alpha)$. Its image under χ_*^0 coincides with the normalized representative of $f(\alpha)$ up to a constant and a modular form g of weight $m/2 + 1$ for which pg is a \mathbb{Q}/\mathbb{Z} -cycle. Hence it follows from Proposition 4.4 (i) with $\delta[c] = e_1\beta$

$$f \langle \alpha, p\iota, \beta \rangle \ni \chi_*^0[pac] = pf(\alpha)e(\beta) + pge(\beta)$$

which is the first claim. This shows (i) and (ii) is analogues. □

Other Toda brackets are more complicated. We first need the

Definition 6.2 A divided congruence f in k variables has *virtual weight* n if there is a modular form g of weight n with the same \mathbb{Q}/\mathbb{Z} Fourier coefficients a_{i_1, i_2, \dots, i_k} for all $i_1, i_2, \dots, i_k > 0$. We write $[f]_n$ for such a \mathbb{Q}/\mathbb{Z} -modular form g .

Lemma 6.3 Suppose f has virtual weight n .

- (i) For $k = 1$ any two modular forms in the bracket $[f]_n$ only differ by cycles in $\Omega_{\mathbb{Q}/\mathbb{Z}}^{0,2n}$.
- (ii) For $k = 2$ any two modular forms in the bracket only differ by cycles in $\Omega_{\mathbb{Q}/\mathbb{Z}}^1$ and modular forms of the form $g \otimes 1$ with g of weight n .

Proof For (i) two elements in $[f]$ differ by modular forms h of weight n with integral expansion except the 0-coefficient. Any such h is a cycle in $\Omega_{\mathbb{Q}/\mathbb{Z}}^{0,2n}$. We remark for n even and level 1 that the highest denominator in the constant coefficient appearing among all such h is the divided Bernoulli number which happens if h is the divided Eisenstein series \bar{E}_n (see Serre [26]). For the case of two variables one observes that for any difference modular form $h = \sum h_1 \otimes h_2$ it holds

$$\sum h_1 \otimes h_2 - h_1 h_2 \otimes 1 + h_2 \otimes h_1 - 1 \otimes h_1 h_2 = 0$$

(see Laures [19, Equation 3.2]). Hence, $z = h - \sum h_1 h_2 \otimes 1$ is antisymmetric with vanishing $q_L^i q_R^j$ -coefficients for $i, j > 0$. This implies that z is a cycle as one easily verifies. □

In the following we write $e_M(\alpha)$ for any representative of $\delta^{-1}e_1(\alpha)$ in $\Omega_{\mathbb{Q}/\mathbb{Z}}^0$ and

$$e(\alpha) = q^0(e_M(\alpha)).$$

Note that for $|\alpha| = 4n - 1$ and level 1 modular forms with 6 inverted we have

$$e_M(\alpha) = e(\alpha)E_n$$

and e is the classical e -invariant.

Theorem 6.4 *Suppose that the Toda bracket $\langle \alpha, \beta, \gamma \rangle$ is defined.*

- (i) *Let $|\alpha| = 2k - 1$, $|\beta| = 2l - 1$ and $|\gamma| = 2m - 1$. Then the modular forms $f(\alpha, \beta) = [e_M(\alpha)e(\beta)]_{k+l}$ and $f(\beta, \gamma) = [e_M(\beta)e(\gamma)]_{l+m}$ exist and we have*

$$f \langle \alpha, \beta, \gamma \rangle = e_M(\alpha)(q^0(f(\beta, \gamma)) - e(\beta)e(\gamma)) + e(\gamma)f(\alpha, \beta)$$

with indeterminacy $e_M(\alpha)q^0H^{0,2(l+m)}(\Omega_{\mathbb{Q}/\mathbb{Z}}^) + e(\gamma)H^{0,2(k+l)}(\Omega_{\mathbb{Q}/\mathbb{Z}}^*)$.*

- (ii) *Let $|\alpha| = 2k - 2$, $|\beta| = 2l - 1$. Then $[f(\alpha) \otimes e_M(\beta)]_{k+l}$ exists and we have*

$$f \langle \alpha, \beta, p\iota \rangle = pf(\alpha)e(\beta) + p\chi_*^0[f(\alpha) \otimes e_M(\beta)]_{k+l}$$

with indeterminacy $pP((D/M_{k+l})_{\mathbb{Q}/\mathbb{Z}})$.

- (iii) *Let $|\beta| = 2l - 1$, $|\gamma| = 2m - 2$. Then $[e_M(\beta) \otimes f(\gamma)]_{l+m}$ exists and we have*

$$f \langle p\iota, \beta, \gamma \rangle = -p\chi_*^0[e_M(\beta) \otimes f(\gamma)]_{l+m}$$

with indeterminacy $pP((D/M_{l+m})_{\mathbb{Q}/\mathbb{Z}})$.

- (iv) *Let $|\alpha| = 2k - 1$ and $|\beta| = 2l - 2$. Choose a representative of $f(\beta)$ whose q -expansion is annihilated by p . Then the modular form $[e_M(\alpha) \otimes f(\beta)]_{k+l}$ exists and we have*

$$f \langle \alpha, \beta, p\iota \rangle = -pe_M(\alpha)f(\beta) - p\chi_*^0[e_M(\alpha) \otimes f(\beta)]_{k+l}$$

with indeterminacy $pP((D/M_{k+l})_{\mathbb{Q}/\mathbb{Z}})$.

- (v) *Let $|\beta| = 2l - 2$ and $|\gamma| = 2m - 1$. Choose a representative of $f(\beta)$ whose q -expansion is annihilated by p . Then the modular form $[f(\beta) \otimes e_M(\gamma)]_{l+m}$ exists and we have*

$$f \langle p\iota, \beta, \gamma \rangle = p\chi_*^0[f(\beta) \otimes e_M(\gamma)]_{l+m}$$

with indeterminacy $pP((D/M_{l+m})_{\mathbb{Q}/\mathbb{Z}})$.

Proof For (i) α, β, γ is represented by $a = e_M(\alpha), b = e_M(\beta), c = e_M(\gamma)$ in the \mathbb{Q}/\mathbb{Z} -cobar complex. Since the product $da db$ vanishes there is a modular form r of degree $k + l$ with $a db + dr \in \Gamma = E_*E$. Applying χ_*^0 gives

$$aq^0(b) - ab + q^0(r) - r \in D.$$

Thus $aq^0(b)$ is congruent to the modular form $ab + r$ of degree $k + l$ up to a constant and

$$f(\alpha, \beta) = [e_M(\alpha)e(\beta)]_{k+l} = ab + r$$

modulo cycles of degree $k + l$. The analogues statement for b, c shows that the brackets exist. The f -invariant of the Toda bracket is hence obtained from [Proposition 4.6\(i\)](#) and the computation

$$\begin{aligned} f \langle \alpha, \beta, \gamma \rangle &= \chi_*^0 \delta^{-1} \langle \delta[a], \delta[b], \delta[c] \rangle \\ &= \chi_*^0 [a(bc + dr(b, c)) + r(a, b)dc] \\ &= ab(q^0(c) - c) + a(q^0(r(b, c)) - r(b, c)) + r(a, b)(q^0(c) - c) \\ &= abq^0(c) + aq^0(r(b, c)) + r(a, b)q^0(c) \\ &= aq^0(f(\beta, \gamma) - bc) + f(\alpha, \beta)q^0(c) \\ &= e_M(\alpha)(q^0(f(\beta, \gamma)) - e(\beta)e(\gamma)) + e(\gamma)f(\alpha, \beta) \end{aligned}$$

This is the result.

For (ii) we find with [Lemma 4.5](#) a modular form r of weight $k + l$ with $a db + dr \in \Gamma^2$. This is an expression with Fourier coefficients in variables q_L, q_M and q_R . For $i, k > 0$ we have

$$\begin{aligned} q_L^i q_M^0 q_R^k (a db + dr) &= \sum q^i(a_1)q^0(a_2)q^k(b) - q^i(r_1)q^k(r_2) \\ &= q_L^i q_R^k \left(\sum a_1 q^0(a_2) \otimes b - r_1 \otimes r_2 \right) \end{aligned}$$

which hence is integral. This shows

$$r = [f(\alpha) \otimes e_M(\beta)]_{k+l}.$$

We compute with [Proposition 4.6\(ii\)](#)

$$\begin{aligned} f \langle \alpha, \beta, p\iota \rangle &= \chi_*^0 \delta^{-1} \langle \delta[a], \delta[b], p\iota \rangle \\ &= \chi_*^0 (p(ab + r)) \\ &= pf(\alpha)e(\beta) + p\chi_*^0 [f(\alpha) \otimes e_M(\beta)]_{k+l} \end{aligned}$$

Similarly, for (iii) we have

$$\begin{aligned} f \langle p\iota, \beta, \gamma \rangle &= \chi_*^0 \delta^{-1} \langle p\iota, \delta[b], \delta\gamma \rangle \\ &= \chi_*^0(-pr) \\ &= -p\chi_*^0[e_M(\beta) \otimes f(\gamma)]_{l+m} \end{aligned}$$

and for (iv)

$$\begin{aligned} f \langle \alpha, \beta, p\iota \rangle &= \chi_*^0 \delta^{-1} \langle \delta[a], \delta[b], p\iota \rangle \\ &= -\chi_*^0(p(ab + r)) \\ &= -pe_M(\alpha) f(\beta) - p\chi_*^0[e_M(\alpha) \otimes f(\beta)]_{k+l} \end{aligned}$$

Finally,

$$\begin{aligned} f \langle p\iota, \beta, \gamma \rangle &= \chi_*^0 \delta^{-1} \langle p\iota, \delta[b], \delta\gamma \rangle \\ &= \chi_*^0(pr) \\ &= p\chi_*^0[f(\beta) \otimes e_M(\gamma)]_{l+m} \end{aligned}$$

The indeterminacy is readily verified with [Corollary 3.4](#). □

Theorem 6.5 *Suppose that the Toda bracket $\langle p\iota, \alpha, p\iota, \beta \rangle$ is strictly defined and let $|\alpha| = 2k - 2, |\beta| = 2l - 2$. Then there are representatives of $f(\alpha)$ and $f(\beta)$ which are annihilated by p and for which the bracket $[pf(\alpha) \otimes f(\beta)]_{k+l}$ exists. For any such modular form we have*

$$f \langle p\iota, \alpha, p\iota, \beta \rangle = p\chi_*^0[pf(\alpha) \otimes f(\beta)]_{k+l}$$

with indeterminacy

$$\chi_*^0([H^{0,2k}(\Omega_{\mathbb{Q}/\mathbb{Z}}^*) \otimes f(\beta)]_{k+l} + [f(\alpha) \otimes H^{0,2l}(\Omega_{\mathbb{Q}/\mathbb{Z}}^*)]_{k+l}) + pq^0(H^{0,2k}(\Omega_{\mathbb{Q}/\mathbb{Z}}^*)) f(\beta)$$

and the indeterminacies coming from the 3-fold brackets

$$p(P(D/M_{k+l})_{\mathbb{Q}/\mathbb{Z}}) + H^{0,2k}(\Omega_{\mathbb{Q}/\mathbb{Z}}^*)q^0(H^{0,2l}(\Omega_{\mathbb{Q}/\mathbb{Z}}^*)).$$

The same holds for the bracket $\langle \alpha, p\iota, \beta, p\iota \rangle$.

Proof With [Lemma 4.3](#) we find p -adapted a, b with $e_2(\alpha) = \delta[a]$ and $e_2(\beta) = \delta[b]$. Without loss of generality we can assume $pab + dr \in \Gamma^{\otimes 2}$ (else replace a by $a + dv/p$ and b by $b + dw/p$). In particular for $i, j > 0$ the numbers

$$\begin{aligned} q_L^i q_M^0 q_R^j \left(p \sum a_1 \otimes a_2 b_1 \otimes b_2 + dr \right) &= p \sum q^i(a_1) q^0(a_2 b_1) q^j(b_2) - q^i(r_1) q^j(r_2) \\ &= q_L^i q_R^j \left(p \sum a_1 q^0(a_2) \otimes q^0(b_1) b_2 - r \right) \end{aligned}$$

are integral. Since $f(\alpha) = q^0(a_2)a_1$ and $f(\beta) = -q^0(b_1)b_2$ we conclude that the bracket $[pf(\alpha) \otimes f(\beta)]_{k+l}$ exists. Its indeterminacy is as in Lemma 6.3(ii). Hence we have with Proposition 4.7

$$\begin{aligned} f\langle p\nu, \alpha, p\nu, \beta \rangle &= \chi_*^0 \delta^{-1} \langle p\nu, \delta[a], p\nu, \delta[b] \rangle \\ &= -\chi_*^0 p r \\ &= p \chi_*^0 [pf(\alpha) \otimes f(\beta)]_{k+l} \end{aligned}$$

The indeterminacy makes use of Corollary 3.4 and the calculations of the 3-fold brackets above. □

7 Examples

In dimension 8 there is the Toda bracket $\langle \nu^2, 2, \eta \rangle$ where ν is the Hopf map of dimension 3. We use the formula Theorem 6.1 for level 3 *TMF* to show that this class coincides with β_2 : The f -invariant of the product ν^2 can be computed from the formula (see von Bodecker [9])

$$f(\nu^2) = e(\nu)e_M(\nu) = \frac{E_1^2}{12^2}.$$

which can be normalized to $-1/2((E_1^2 - 1)/12)^2$. Hence we have

$$f\langle \nu^2, 2, \eta \rangle = -\frac{1}{2} \left(\frac{E_1^2 - 1}{12} \right)^2$$

which coincides with the f -invariant of β_2 (see von Bodecker [8]).

Similarly, we have for the dimension 7 Hopf map σ instead of ν

$$f(\sigma^2) = e(\sigma)e_M(\sigma) = \frac{E_4}{240^2} = -\frac{1}{2} \left(\frac{E_4 - 1}{240} \right)^2$$

and hence

$$f\langle \sigma^2, 2, \eta \rangle = -\frac{1}{2} \left(\frac{E_4 - 1}{240} \right)^2.$$

The computation in [9, page 7] shows that this expression is congruent to

$$f(\beta_{4/3}) = \frac{1}{2} \left(\frac{E_1^2 - 1}{4} \right)^4 + \frac{1}{2} \left(\frac{E_1^2 - 1}{4} \right)^3.$$

There is another way to compute this class using the Toda relation

$$\langle \sigma^2, 2, \eta \rangle = \langle \sigma, 2\sigma, \eta \rangle$$

and formula [Theorem 6.4\(i\)](#): First we have

$$f(\sigma, 2\sigma) = \left[2 \frac{E_4}{240^2} \right]_8 = \frac{E_4^2}{240^2} \quad \text{and} \quad f(2\sigma, \eta) = \left[\frac{E_4}{240} \right]_5 = 0.$$

This gives

$$f\langle \sigma, 2\sigma, \eta \rangle = -\frac{E_4}{240^2} + \frac{1}{2} \frac{E_4^2}{240^2} = \frac{1}{2} \left(\frac{E_4 - 1}{240} \right)^2 = f(\beta_{4/3})$$

In dimension 18 there is the class $\langle \sigma, 2\sigma, \nu \rangle$ for which the formula reads

$$f\langle \sigma, 2\sigma, \nu \rangle = \frac{E_4}{240} \left(q^0 \left[\frac{E_4}{1440} \right]_6 - \frac{1}{1440} \right) + \frac{1}{12} \frac{E_4^2}{240^2}$$

To evaluate the bracket observe that

$$d^3 = d^5 \pmod 8$$

for all integers d and hence

$$\frac{E_4 - 1}{240} = \sum_{n \geq 1} \sum_{d|n} d^3 q^n = \sum_{n \geq 1} \sum_{d|n} d^5 q^n = \frac{1 - E_6}{504} \pmod 8.$$

This gives

$$\begin{aligned} f\langle \sigma, 2\sigma, \nu \rangle &= \frac{E_4}{240} \left(-\frac{1}{3024} - \frac{1}{1440} \right) + \frac{1}{12} \frac{E_4^2}{240^2} \\ &= \frac{1}{6} \frac{1}{240^2} \left(-\frac{31}{21} E_4 + \frac{1}{2} E_4^2 \right) \end{aligned}$$

which has order 4 modulo indeterminacy. In fact with [Lemma 5.4](#) one can show that it coincides with $f(-\beta_{4/2,2})$.

The Toda bracket $\langle \sigma^2, 2, \sigma^2, 2 \rangle$ in dimension 30 exists (see Barratt, Jones and Mahowald [\[5, 1.2 and 1.3\]](#)) Compute modulo indeterminacy

$$\begin{aligned} &\left[\frac{1}{2} \left(\frac{E_4 - 1}{240} \right)^2 \otimes \left(\frac{E_4 - 1}{240} \right)^2 \right]_{16} \\ &= \left[\frac{1}{240^4} \left(\frac{E_4^2 \otimes E_4^2}{2} - E_4^2 \otimes E_4 - E_4 \otimes E_4^2 + 2E_4 \otimes E_4 \right) \right]_{16} \\ &= \frac{1}{240^4} \left(\frac{E_4^2 \otimes E_4^2}{2} - \frac{E_4^3 \otimes E_4}{3} - \frac{E_4 \otimes E_4^3}{3} \right) \\ &= \frac{1}{12} \left(\frac{E_4 \otimes 1 - 1 \otimes E_4}{240} \right)^4 \end{aligned}$$

Here we used in the second step that

$$\frac{1}{3} \left(\frac{E_4 - 1}{240} \right)^3 \otimes \frac{E_4 - 1}{240}$$

is integral. This gives with [Theorem 6.5](#)

$$\begin{aligned} f(\sigma^2, 2, \sigma^2, 2) &= 2\chi_*^0 \left[\frac{1}{2} \left(\frac{E_4 - 1}{240} \right)^2 \otimes \left(\frac{E_4 - 1}{240} \right)^2 \right]_{16} \\ &= 2\chi_*^0 \left(\frac{1}{12} \left(\frac{E_4 \otimes 1 - 1 \otimes E_4}{240} \right)^4 \right) \\ &= \frac{1}{2} \left(\frac{E_4 - 1}{240} \right)^4 \end{aligned}$$

which has order 2 and coincides with the f invariant of the Kervaire class $f(\beta_{8/8})$ (compare von Bodecker [\[8\]](#)).

References

- [1] **JF Adams**, *On the groups $J(X)$ IV*, Topology 5 (1966) 21–71 [MR0198470](#)
- [2] **A Baker**, *Elliptic genera of level N and elliptic cohomology*, J. London Math. Soc. (2) 49 (1994) 581–593 [MR1271552](#)
- [3] **A Baker**, *Operations and cooperations in elliptic cohomology I: Generalized modular forms and the cooperation algebra*, New York J. Math. 1 (1994/95) 39–74, electronic [MR1307488](#)
- [4] **A Baker**, *Hecke operations and the Adams E_2 -term based on elliptic cohomology*, Canad. Math. Bull. 42 (1999) 129–138 [MR1692001](#)
- [5] **MG Barratt, JDS Jones, ME Mahowald**, *Relations amongst Toda brackets and the Kervaire invariant in dimension 62*, J. London Math. Soc. (2) 30 (1984) 533–550 [MR810962](#)
- [6] **M Behrens**, *Congruences between modular forms given by the divided β family in homotopy theory*, Geom. Topol. 13 (2009) 319–357 [MR2469520](#)
- [7] **M Behrens, G Laures**, *β -family congruences and the f -invariant*, from: “New topological contexts for Galois theory and algebraic geometry (BIRS 2008)”, Geom. Topol. Monogr. 16, Geom. Topol. Publ., Coventry (2009) 9–29 [MR2544384](#)
- [8] **H von Bodecker**, *The beta family at the prime two and modular forms of level three* [arXiv:0912.3082](#)
- [9] **H von Bodecker**, *On the f -invariant of products* [arXiv:0808.0428](#)
- [10] **H von Bodecker**, *On the geometry of the f invariant* [arXiv:0909.3968](#)

- [11] **J-L Brylinski**, *Representations of loop groups, Dirac operators on loop space, and modular forms*, Topology 29 (1990) 461–480 [MR1071369](#)
- [12] **U Bunke, N Naumann**, *The f -invariant and index theory*, Manuscripta Math. 132 (2010) 365–397 [MR2652438](#)
- [13] **U Bunke, N Naumann**, *The f -invariant and index theory*, Manuscripta Math. 132 (2010) 365–397 [MR2652438](#)
- [14] **J Franke**, *On the construction of elliptic cohomology*, Math. Nachr. 158 (1992) 43–65 [MR1235295](#)
- [15] **J Hornbostel, N Naumann**, *Beta-elements and divided congruences*, Amer. J. Math. 129 (2007) 1377–1402 [MR2354323](#)
- [16] **N M Katz**, *Higher congruences between modular forms*, Ann. of Math. (2) 101 (1975) 332–367 [MR0417059](#)
- [17] **S O Kochman**, *Stable homotopy groups of spheres*, Lecture Notes in Mathematics 1423, Springer, Berlin (1990) [MR1052407](#) A computer-assisted approach
- [18] **S O Kochman**, *Bordism, stable homotopy and Adams spectral sequences*, Fields Institute Monographs 7, American Mathematical Society, Providence, RI (1996) [MR1407034](#)
- [19] **G Laures**, *The topological q -expansion principle*, Topology 38 (1999) 387–425 [MR1660325](#)
- [20] **G Laures**, *On cobordism of manifolds with corners*, Trans. Amer. Math. Soc. 352 (2000) 5667–5688 [MR1781277](#)
- [21] **G Laures**, *$K(1)$ -local topological modular forms*, Invent. Math. 157 (2004) 371–403 [MR2076927](#)
- [22] **T Lawson, N Naumann**, *Topological modular forms of level 3 and E_∞ -structures on truncated Brown–Peterson spectra* [arXiv:1101.3897](#)
- [23] **J P May**, *Matric Massey products*, J. Algebra 12 (1969) 533–568 [MR0238929](#)
- [24] **H R Miller, D C Ravenel, W S Wilson**, *Periodic phenomena in the Adams–Novikov spectral sequence*, Ann. Math. (2) 106 (1977) 469–516 [MR0458423](#)
- [25] **D C Ravenel**, *Complex cobordism and stable homotopy groups of spheres*, Pure and Applied Mathematics 121, Academic Press, Orlando, FL (1986) [MR860042](#)
- [26] **J-P Serre**, *Formes modulaires et fonctions zêta p -adiques*, from: “Modular functions of one variable, III (Proc Internat. Summer School, Univ. Antwerp, 1972)”, Springer, Berlin (1973) 191–268. Lecture Notes in Math., Vol. 350 [MR0404145](#)
- [27] **K Shimomura**, *Novikov’s Ext^2 at the prime 2*, Hiroshima Math. J. 11 (1981) 499–513 [MR635034](#)

*Fakultät für Mathematik, Ruhr-Universität Bochum
NA1/66, D-44780 Bochum, Germany*

gerd@laures.de

<http://www.gerd.laures.de/>

Received: 30 June 2010 Revised: 6 April 2011