

## On sections of hyperelliptic Lefschetz fibrations

SHUNSUKE TANAKA

We construct a relation among right-handed Dehn twists in the mapping class group of a compact oriented surface of genus  $g$  with  $4g + 4$  boundary components. This relation gives an explicit topological description of  $4g + 4$  disjoint  $(-1)$ -sections of a hyperelliptic Lefschetz fibration of genus  $g$  on the manifold  $\mathbb{C}\mathbb{P}^2 \# (4g + 5)\overline{\mathbb{C}\mathbb{P}^2}$ .

57N13; 20F34

### 1 Introduction

Lefschetz fibrations relate the topology of symplectic 4-manifolds to the combinatorics on relations in Dehn twist generators of mapping class groups of surfaces. It is well-known that a Lefschetz fibration of genus 1 on the manifold  $E(1) = \mathbb{C}\mathbb{P}^2 \# 9\overline{\mathbb{C}\mathbb{P}^2}$  constructed by blowing up nine intersections of two generic cubics in  $\mathbb{C}\mathbb{P}^2$  has twelve singular fibers and nine disjoint  $(-1)$ -sections. Korkmaz and Ozbagci [7] constructed a relation among right-handed Dehn twists in the mapping class group of a torus with nine boundary components to locate a set of nine disjoint  $(-1)$ -sections in a Kirby diagram of  $E(1)$ . It is also known to algebraic geometers that a hyperelliptic Lefschetz fibration of genus  $g$  on the manifold  $X_g = \mathbb{C}\mathbb{P}^2 \# (4g + 5)\overline{\mathbb{C}\mathbb{P}^2}$  has  $8g + 4$  singular fibers and  $4g + 4$  disjoint  $(-1)$ -sections for  $g \geq 2$  (see Saitō and Sakakibara [10, Section 3] and Kitagawa and Konno [6, Remark 1.1]).

In this paper we construct a relation among right-handed Dehn twists in the mapping class group of a compact oriented surface of genus  $g$  with  $4g + 4$  boundary components to locate a set of  $4g + 4$  disjoint  $(-1)$ -sections in a Kirby diagram of  $X_g$ . In the case  $g = 2$ , our relation can be considered as an improvement of Onaran's relations [9] in mapping class groups of surfaces of genus two with at most eight boundary components.

In Section 2 we review basic relations in mapping class groups and produce two relations on a torus with eight boundary components. Combining these relations, we construct a new relation on a surface of genus  $g$  with  $4g + 4$  boundary components in Section 3. In Section 4 we apply the relation to visualize  $4g + 4$  disjoint  $(-1)$ -sections in a Kirby diagram of a hyperelliptic Lefschetz fibration of genus  $g$ .

## 2 Building blocks

In this section we review basic relations in mapping class groups and produce two relations on a torus with boundary both used in the next section.

### 2.1 Basic relations in mapping class groups

Let  $\Sigma_{g,r}$  be a compact oriented surface of genus  $g$  with  $r$  boundary components and  $\text{Diff}_+\Sigma_{g,r}$  the group of orientation-preserving diffeomorphisms of  $\Sigma_{g,r}$  fixing the boundary  $\partial\Sigma_{g,r}$  pointwise equipped with the  $C^\infty$ -topology. The group  $\pi_0(\text{Diff}_+\Sigma_{g,r})$  of path-components of  $\text{Diff}_+\Sigma_{g,r}$  is called the *mapping class group* of  $\Sigma_{g,r}$  and we denote it by  $\mathcal{M}_{g,r}$ . We denote by  $\mathcal{F}_{g,r}$  the free group generated by all isotopy classes  $\mathcal{S}_{g,r}$  of simple closed curves in the interior of  $\Sigma_{g,r}$ . There is a natural epimorphism  $\varpi : \mathcal{F}_{g,r} \rightarrow \mathcal{M}_{g,r}$  which sends (the isotopy class of) a simple closed curve  $a$  in the interior of  $\Sigma_{g,r}$  to the right-handed Dehn twist  $t_a$  along  $a$ . We set  $\mathcal{R}_{g,r} := \text{Ker } \varpi$ .

A word in the generators  $\mathcal{S}_{g,r}$  is called *positive* if it includes no negative exponents. We put  $W(c) := t_a^{\varepsilon_r} \cdots t_{a_1}^{\varepsilon_1}(c) \in \mathcal{S}_{g,r}$  for  $c \in \mathcal{S}_{g,r}$  and  $W = a_r^{\varepsilon_r} \cdots a_1^{\varepsilon_1} \in \mathcal{F}_{g,r}$  ( $a_1, \dots, a_r$  in  $\mathcal{S}_{g,r}$ ,  $\varepsilon_1, \dots, \varepsilon_r$  in  $\{\pm 1\}$ ). We often denote  $a^{-1}$  by  $\bar{a}$  for an element  $a$  of  $\mathcal{S}_{g,r}$ . For two words  $W_1, W_2 \in \mathcal{F}_{g,r}$ , we denote  $W_1 \equiv W_2$  if  $\varpi(W_1) = \varpi(W_2)$ . If the relation  $W_1 \equiv W_2$  holds for  $W_1, W_2 \in \mathcal{F}_{g,r}$ , we obtain another relation  $VW_1V^{-1} \equiv VW_2V^{-1}$ , which is called a *conjugate* of  $W_1 \equiv W_2$ , for every  $V \in \mathcal{F}_{g,r}$ .

We recall definitions of basic relations in mapping class groups.

**Definition 2.1** [3] (1) For disjoint simple closed curves  $a, b$  in the interior of  $\Sigma_{g,r}$ , we have a relation  $ab \equiv ba$  in  $\mathcal{F}_{g,r}$  called a *commutativity relation*. A regular neighborhood of  $a \cup b$  is the disjoint union of two annuli.

(2) For simple closed curves  $a, b$  in the interior of  $\Sigma_{g,r}$  which intersect transversely at one point, we have a relation  $aba \equiv bab$  in  $\mathcal{F}_{g,r}$  called a *braid relation*. A regular neighborhood of  $a \cup b$  is a torus with one boundary component.

(3) For simple closed curves  $\alpha, \sigma, \gamma, \delta_1, \delta_2, \delta_3, \delta_4$  in the interior of  $\Sigma_{g,r}$  shown in Figure 1, we have a relation  $\delta_1\delta_2\delta_3\delta_4 \equiv \gamma\sigma\alpha$  in  $\mathcal{F}_{g,r}$  called a *lantern relation*. The union of  $\delta_1, \delta_2, \delta_3, \delta_4$  bounds a sphere with four boundary components in  $\Sigma_{g,r}$ .

(4) An ordered  $n$ -tuple  $(c_1, \dots, c_n)$  of simple closed curves in the interior of  $\Sigma_{g,r}$  is called a *chain* of length  $n$  if  $c_i$  and  $c_{i+1}$  intersect transversely at one point ( $i = 1, \dots, n-1$ ) and other  $c_i$  and  $c_j$  never intersect. For a chain  $(c_1, \dots, c_{2g+1})$  of length  $2g+1$  on  $\Sigma_{g,0}$ , we have a relation  $(c_1 \cdots c_{2g+1} c_{2g+1} \cdots c_1)^2 \equiv 1$  in  $\mathcal{F}_{g,0}$  called a *hyperelliptic relation* (see Figure 2).




Figure 1: Lantern relation




Figure 2: Hyperelliptic relation

**Remark 2.2** Let  $a$  and  $b$  be simple closed curves in the interior of  $\Sigma_{g,r}$  and  $c$  the simple closed curve  $t_b(a) = b(a)$ . Then we have the relation  $c \equiv bab$  in  $\mathcal{F}_{g,r}$ . If  $a$  and  $b$  intersect transversely at one point, we have another relation  $b \equiv ac\bar{a}$ . This relation together with the relation  $c \equiv bab$  yields a braid relation  $aba \equiv bab$ .

## 2.2 Two relations on a torus with boundary

In this subsection we construct two relations on a torus with eight boundary components. The first relation is the following.

**Proposition 2.3** Relation (A) For simple closed curves in the interior of  $\Sigma_{1,8}$  shown in Figure 3, we have the relation

$$a_1 a_2 \delta_1 \delta_2 \delta_3 \delta_4 \delta_5 \delta_6 \equiv a_5 a_4 b_2 \bar{a}_4 \sigma_1 \sigma_4 a_{10} a_3 b_2 \bar{a}_3 \sigma_2 a_5 a_3 a_8 b_2 \bar{a}_8 \bar{a}_3 \sigma_3 \sigma_5 a_{11}.$$

We make use of the five-holed torus relation found by Korkmaz and Ozbagci [7] in order to prove Proposition 2.3.

**Lemma 2.4** (Korkmaz–Ozbagci [7]) For simple closed curves in the interior of  $\Sigma_{1,5}$  shown on the right in Figure 4, we have the relation

$$\delta_2 \delta_1 a_2 \gamma \delta_3 \equiv a_5 b_2 a_3 a_4 a_5 b_2 \sigma_1 a_6 a_3 b_2 \sigma_2 a_8.$$




Figure 3: Relation (A)

**Remark 2.5** The relation in Lemma 2.4 is deduced from the original five-holed torus relation

$$\delta_2 \delta_1 a_2 \gamma \delta_3 \equiv a_5 a_3 a_4 b_2 \sigma_1 a_6 a_3 b_2 \sigma_2 a_8 a_5 b_2$$

(see [7, Section 3.5]) by using commutativity relations and conjugations.




Figure 4: Five-holed torus relation

**Proof of Proposition 2.3** Applying commutativity relations and conjugations to the five-holed torus relation in Lemma 2.4, we obtain

$$\begin{aligned} a_2\delta_1\delta_2\delta_3\gamma &\equiv a_5b_2a_3a_4a_5b_2\sigma_1a_6a_3b_2\sigma_2a_8 \equiv a_8b_2\bar{a}_8a_8a_3a_4a_5b_2\sigma_1a_6a_3b_2\sigma_2a_5 \\ &\equiv a_8a_3a_4a_5b_2\sigma_1a_6a_3b_2\sigma_2a_5a_8b_2\bar{a}_8. \end{aligned}$$

Multiplying both sides of this relation by  $\bar{\gamma}$ , we have

$$a_2\delta_1\delta_2\delta_3 \equiv a_8a_3a_4a_5b_2\sigma_1a_6a_3b_2\sigma_2a_5a_8b_2\bar{a}_8\bar{\gamma}.$$

We embed  $\Sigma_{1,5}$  into  $\Sigma_{1,6}$  and take simple closed curves  $a_1, a_9, \delta_4, \sigma_3$  in the interior of  $\Sigma_{1,6}$  shown in Figure 4. Then we have a lantern relation

$$\delta_4a_1a_3a_8 \equiv \gamma\sigma_3a_9.$$

Combining these relations and applying commutativity relations, we obtain

$$\begin{aligned} a_8a_3a_1a_2\delta_1\delta_2\delta_3\delta_4 &\equiv a_8a_3a_4a_5b_2\sigma_1a_6a_3b_2\sigma_2a_5a_8b_2\bar{a}_8\bar{\gamma}\gamma\sigma_3a_9 \\ &\equiv a_8a_3a_4a_5b_2\sigma_1a_6a_3b_2\sigma_2a_5a_8b_2\bar{a}_8\sigma_3a_9. \end{aligned}$$

Multiplying both sides of this relation by  $\bar{a}_3\bar{a}_8$ , we have a relation

$$(A1) \quad a_1a_2\delta_1\delta_2\delta_3\delta_4 \equiv a_4a_5b_2\sigma_1a_6a_3b_2\sigma_2a_5a_8b_2\bar{a}_8\sigma_3a_9$$

on  $\Sigma_{1,6}$ .

We change the name  $\delta_2$  of a curve in relation (A1) into  $\gamma$  (shown on the right in Figure 5) and apply commutativity relations and conjugations to it to obtain

$$\begin{aligned} a_1a_2\delta_1\delta_3\delta_4\gamma &\equiv a_4a_5b_2\sigma_1a_6a_3b_2\sigma_2a_5a_8b_2\bar{a}_8\sigma_3a_9 \\ &\equiv a_5a_4b_2\bar{a}_4a_4\sigma_1a_6a_3b_2\sigma_2a_5a_8b_2\bar{a}_8\sigma_3a_9 \\ &\equiv a_4a_6a_3b_2\sigma_2a_5a_8b_2\bar{a}_8\sigma_3a_9a_5a_4b_2\bar{a}_4\sigma_1. \end{aligned}$$

Multiplying both sides of this relation by  $\bar{\gamma}$ , we have

$$a_1a_2\delta_1\delta_3\delta_4 \equiv a_4a_6a_3b_2\sigma_2a_5a_8b_2\bar{a}_8\sigma_3a_9a_5a_4b_2\bar{a}_4\sigma_1\bar{\gamma}.$$

We embed  $\Sigma_{1,6}$  into  $\Sigma_{1,7}$  and take simple closed curves  $a_{10}, \delta_2, \delta_5, \sigma_4$  in the interior of  $\Sigma_{1,7}$  shown in Figure 5. Then we have a lantern relation

$$\delta_2\delta_5a_4a_6 \equiv \gamma\sigma_4a_{10}.$$

Combining these relations and applying commutativity relations, we obtain

$$\begin{aligned} a_4a_6a_1a_2\delta_1\delta_2\delta_3\delta_4\delta_5 &\equiv a_4a_6a_3b_2\sigma_2a_5a_8b_2\bar{a}_8\sigma_3a_9a_5a_4b_2\bar{a}_4\sigma_1\bar{\gamma}\gamma\sigma_4a_{10} \\ &\equiv a_4a_6a_3b_2\sigma_2a_5a_8b_2\bar{a}_8\sigma_3a_9a_5a_4b_2\bar{a}_4\sigma_1\sigma_4a_{10}. \end{aligned}$$




Figure 5: Embedding of  $\Sigma_{1,6}$  into  $\Sigma_{1,7}$  (I)

Multiplying both sides of this relation by  $\bar{a}_6\bar{a}_4$ , we have a relation

$$(A2) \quad a_1 a_2 \delta_1 \delta_2 \delta_3 \delta_4 \delta_5 \equiv a_3 b_2 \sigma_2 a_5 a_8 b_2 \bar{a}_8 \sigma_3 a_9 a_5 a_4 b_2 \bar{a}_4 \sigma_1 \sigma_4 a_{10}$$

on  $\Sigma_{1,7}$ .

We change the name  $a_1$  of a curve in relation (A2) into  $\gamma$  (shown on the right in Figure 6) and apply commutativity relations and conjugations to it to obtain

$$\begin{aligned} a_2 \delta_1 \delta_2 \delta_3 \delta_4 \delta_5 \gamma &\equiv a_3 b_2 \sigma_2 a_5 a_8 b_2 \bar{a}_8 \sigma_3 a_9 a_5 a_4 b_2 \bar{a}_4 \sigma_1 \sigma_4 a_{10} \\ &\equiv a_3 b_2 \bar{a}_3 \sigma_2 a_5 a_3 a_8 b_2 \bar{a}_8 \sigma_3 a_9 a_5 a_4 b_2 \bar{a}_4 \sigma_1 \sigma_4 a_{10} \\ &\equiv a_3 b_2 \bar{a}_3 \sigma_2 a_5 a_3 a_8 b_2 \bar{a}_8 \bar{a}_3 \sigma_3 a_3 a_9 a_5 a_4 b_2 \bar{a}_4 \sigma_1 \sigma_4 a_{10} \\ &\equiv a_3 a_9 a_5 a_4 b_2 \bar{a}_4 \sigma_1 \sigma_4 a_{10} a_3 b_2 \bar{a}_3 \sigma_2 a_5 a_3 a_8 b_2 \bar{a}_8 \bar{a}_3 \sigma_3. \end{aligned}$$

Multiplying both sides of this relation by  $\bar{\gamma}$ , we have

$$a_2 \delta_1 \delta_2 \delta_3 \delta_4 \delta_5 \equiv a_3 a_9 a_5 a_4 b_2 \bar{a}_4 \sigma_1 \sigma_4 a_{10} a_3 b_2 \bar{a}_3 \sigma_2 a_5 a_3 a_8 b_2 \bar{a}_8 \bar{a}_3 \sigma_3 \bar{\gamma}.$$

We embed  $\Sigma_{1,7}$  into  $\Sigma_{1,8}$  and take simple closed curves  $a_1, a_{11}, \delta_6, \sigma_5$  in the interior of  $\Sigma_{1,8}$  shown in Figure 6. Then we have a lantern relation

$$\delta_6 a_1 a_3 a_9 \equiv \gamma \sigma_5 a_{11}.$$

Combining these relations and applying commutativity relations, we obtain

$$\begin{aligned} a_3 a_9 a_1 a_2 \delta_1 \delta_2 \delta_3 \delta_4 \delta_5 \delta_6 &\equiv a_3 a_9 a_5 a_4 b_2 \bar{a}_4 \sigma_1 \sigma_4 a_{10} a_3 b_2 \bar{a}_3 \sigma_2 a_5 a_3 a_8 b_2 \bar{a}_8 \bar{a}_3 \sigma_3 \bar{\gamma} \gamma \sigma_5 a_{11} \\ &\equiv a_3 a_9 a_5 a_4 b_2 \bar{a}_4 \sigma_1 \sigma_4 a_{10} a_3 b_2 \bar{a}_3 \sigma_2 a_5 a_3 a_8 b_2 \bar{a}_8 \bar{a}_3 \sigma_3 \sigma_5 a_{11}. \end{aligned}$$




Figure 6: Embedding of  $\Sigma_{1,7}$  into  $\Sigma_{1,8}$  (I)

Multiplying both sides of this relation by  $\bar{a}_9\bar{a}_3$ , we finally obtain relation (A). This completes the proof of Proposition 2.3. □

The second relation constructed in this subsection is the following.

**Proposition 2.6** Relation (B) For simple closed curves in the interior of  $\Sigma_{1,8}$  shown in Figure 7, we have the relation

$$a_1a_2a_7a_8\delta_1\delta_2\delta_3\delta_4 \equiv a_4a_5''\bar{a}_6b_2a_6a_3b_2\bar{a}_3\tau'\tau'''a_5a_4''\bar{a}_3b_2a_3a_6b_2\bar{a}_6\tau\tau''.$$

We make use of the four-holed torus relation found by Korkmaz and Ozbagci [7] in order to prove Proposition 2.6.

**Lemma 2.7** (Korkmaz–Ozbagci [7]) For simple closed curves in the interior of  $\Sigma_{1,4}$  shown on the left in Figure 8, we have the relation

$$a_2a_1a_7\gamma \equiv (a_3a_6b_2a_4a_5b_2)^2.$$

**Remark 2.8** The relation in Lemma 2.7 is not the exact four-holed torus relation but the relation written in a more symmetric form (see [7, Section 3.4, Remark]).

**Proof of Proposition 2.6** We consider the four-holed torus relation reviewed in Lemma 2.7. We then embed  $\Sigma_{1,4}$  into  $\Sigma_{1,5}$  and take simple closed curves  $a'_5, a_8, \delta_1, \tau$  in the interior of  $\Sigma_{1,5}$  shown in Figure 8. Then we have a lantern relation

$$\delta_1a_8a_6a_5 \equiv \gamma\tau a'_5.$$




Figure 7: Relation (B)

Combining this relations with the four-holed torus relation, and applying commutativity relations and conjugations, we obtain a relation

$$\begin{aligned}
 \text{(B1)} \quad a_1 a_2 a_7 a_8 \delta_1 &\equiv \bar{a}_5 a_4 a_5 b_2 a_3 a_6 b_2 a_4 a_5 b_2 a_3 a_6 b_2 \bar{\gamma} \cdot \bar{a}_6 \gamma \tau a'_5 \\
 &\equiv a_4 b_2 a_3 a_6 b_2 a_4 a_5 b_2 a_6 a_3 b_2 \bar{a}_6 \tau a'_5 \\
 &\equiv a_4 a_5 b_2 a_6 a_3 b_2 \bar{a}_6 \tau a'_5 a_4 b_2 a_3 a_6 b_2
 \end{aligned}$$




Figure 8: Four-holed torus relation



on  $\Sigma_{1,5}$ .




Figure 9: Embedding of  $\Sigma_{1,5}$  into  $\Sigma_{1,6}$

We change the name  $a_1$  of a curve in relation (B1) into  $\gamma$  (shown on the right in Figure 9) to obtain

$$\gamma a_2 a_7 a_8 \delta_1 \equiv a_4 a_5 b_2 a_6 a_3 b_2 \bar{a}_6 \tau a'_5 a_4 b_2 a_3 a_6 b_2.$$

We embed  $\Sigma_{1,5}$  into  $\Sigma_{1,6}$  and take simple closed curves  $a_1, a'_4, \delta_2, \tau'$  in the interior of  $\Sigma_{1,6}$  shown in Figure 9. Then we have a lantern relation

$$a_4 a_3 a_1 \delta_2 \equiv \gamma \tau' a'_4.$$

Combining these relations and applying commutativity relations and conjugations, we obtain a relation

$$\begin{aligned} \text{(B2)} \quad a_1 a_2 a_7 a_8 \delta_1 \delta_2 &\equiv \bar{a}_4 a_4 a_5 b_2 a_6 a_3 b_2 \bar{a}_6 \tau a'_5 a_4 b_2 a_3 a_6 b_2 \bar{\gamma} \cdot \bar{a}_3 \gamma \tau' a'_4 \\ &\equiv a_5 b_2 a_6 a_3 b_2 \bar{a}_6 \tau a'_5 a_4 b_2 a_6 a_3 b_2 \bar{a}_3 \tau' a'_4 \\ &\equiv b_2 a_6 a_3 b_2 \bar{a}_3 \tau' a'_4 a_5 b_2 a_6 a_3 b_2 \bar{a}_6 \tau a'_5 a_4. \end{aligned}$$

on  $\Sigma_{1,6}$ .

We change the name  $a_8$  of a curve in relation (B2) into  $\gamma$  (shown on the left in Figure 10) to obtain

$$a_1 a_2 a_7 \gamma \delta_1 \delta_2 \equiv b_2 a_6 a_3 b_2 \bar{a}_3 \tau' a'_4 a_5 b_2 a_6 a_3 b_2 \bar{a}_6 \tau a'_5 a_4.$$




Figure 10: Embedding of  $\Sigma_{1,6}$  into  $\Sigma_{1,7}$  (II)

We embed  $\Sigma_{1,6}$  into  $\Sigma_{1,7}$  and take simple closed curves  $a''_5, a_8, \delta_3, \tau''$  in the interior of  $\Sigma_{1,7}$  shown in Figure 10. Then we have a lantern relation

$$\delta_3 a_8 a_6 a'_5 \equiv \gamma \tau'' a''_5.$$

Combining these relations and applying commutativity relations and conjugations, we obtain a relation

$$\begin{aligned} \text{(B3)} \quad a_1 a_2 a_7 a_8 \delta_1 \delta_2 \delta_3 &\equiv \bar{a}_6 b_2 a_6 a_3 b_2 \bar{a}_3 \tau' a'_4 a_5 b_2 a_6 a_3 b_2 \bar{a}_6 \tau a'_5 a_4 \bar{\gamma} \cdot \bar{a}'_5 \gamma \tau'' a''_5 \\ &\equiv \bar{a}_6 b_2 a_6 a_3 b_2 \bar{a}_3 \tau' a'_4 a_5 b_2 a_6 a_3 b_2 \bar{a}_6 \tau \tau'' a_4 a''_5 \\ &\equiv b_2 a_3 a_6 b_2 \bar{a}_6 \tau \tau'' a_4 a''_5 \bar{a}_6 b_2 a_6 a_3 b_2 \bar{a}_3 \tau' a'_4 a_5. \end{aligned}$$

We change the name  $a_1$  of a curve in relation (B3) into  $\gamma$  (shown on the right in Figure 11) to obtain

$$\gamma a_2 a_7 a_8 \delta_1 \delta_2 \delta_3 \equiv b_2 a_3 a_6 b_2 \bar{a}_6 \tau \tau'' a_4 a''_5 \bar{a}_6 b_2 a_6 a_3 b_2 \bar{a}_3 \tau' a'_4 a_5.$$

We embed  $\Sigma_{1,7}$  into  $\Sigma_{1,8}$  and take simple closed curves  $a_1, a''_4, \delta_4, \tau'''$  in the interior of  $\Sigma_{1,8}$  shown in Figure 11. Then we have a lantern relation

$$\delta_4 a_1 a_3 a'_4 \equiv \gamma \tau''' a''_4.$$




Figure 11: Embedding of  $\Sigma_{1,7}$  into  $\Sigma_{1,8}$  (II)

Combining these relations and applying commutativity relations and conjugations, we finally obtain relation (B):

$$\begin{aligned}
 a_1 a_2 a_7 a_8 \delta_1 \delta_2 \delta_3 \delta_4 &\equiv \bar{a}_3 b_2 a_3 a_6 b_2 \bar{a}_6 \tau \tau'' a_4 a_5'' \bar{a}_6 b_2 a_6 a_3 b_2 \bar{a}_3 \tau' a_4' a_5 \bar{\gamma} \cdot \bar{a}_4' \gamma \tau''' a_4'' \\
 &\equiv \bar{a}_3 b_2 a_3 a_6 b_2 \bar{a}_6 \tau \tau'' a_4 a_5'' \bar{a}_6 b_2 a_6 a_3 b_2 \bar{a}_3 \tau' \tau''' a_5 a_4'' \\
 &\equiv a_4 a_5'' \bar{a}_6 b_2 a_6 a_3 b_2 \bar{a}_3 \tau' \tau''' a_5 a_4'' \bar{a}_3 b_2 a_3 a_6 b_2 \bar{a}_6 \tau \tau'' .
 \end{aligned}$$

This completes the proof of Proposition 2.6. □

**Remark 2.9** Both of relations (A) and (B) are different from the eight-holed torus relation of Korkmaz and Ozbagci [7] though the constructions are similar.

### 3 Constructions

In this section we construct a new relation on a compact oriented surface of genus  $g$  with  $4g + 4$  boundary components by combining copies of relations (A) and (B) obtained in the previous section.

#### 3.1 Higher genus

We assume  $g \geq 3$ . For integers  $m, n$  ( $0 < m \leq n$ ) and words  $W_m, W_{m+1}, \dots, W_n$  in  $\mathcal{F}_{g,r}$ , we denote the product  $W_m W_{m+1} \cdots W_n$  (respectively  $W_n \cdots W_{m+1} W_m$ ) by  $\prod_{i=m}^n W_i$  (respectively  $\prod_{i=n}^m W_i$ ).

**Theorem 3.1** Relation  $(H_g)$  For simple closed curves in the interior of  $\Sigma_{g,4g+4}$  shown in Figure 12, we have the relation

$$\delta_1 \delta_2 \cdots \delta_{4g+3} \delta_{4g+4} \equiv \prod_{i=g-1}^2 \beta_i''' \beta_i \tau'_{i-1} \tau'''_{i-1} \cdot \beta_1 \sigma'_1 \sigma'_4 a_{3g+3} \beta'_1 \sigma'_2 a_1 \beta''_1 \sigma'_3 \sigma'_5 \\ \times \prod_{i=2}^{g-1} \beta_i'' \beta'_i \tau_{i-1} \tau''_{i-1} \cdot \beta_g \sigma_1 \sigma_4 a_{3g} \beta'_g \sigma_2 a_{3g-1} \beta''_g \sigma_3 \sigma_5$$

in  $\mathcal{M}_{g,4g+4}$ , where

$$\beta_1 := a_{3g+4}(b_1), \quad \beta'_1 := a_3(b_1), \quad \beta''_1 := a_{3g+5} a_3(b_1), \\ \beta_g := a_{3g+1}(b_g), \quad \beta'_g := a_{3g-3}(b_g), \quad \beta''_g := a_{3g-3} a_{3g+2}(b_g), \\ \beta_i := a_{3i-3}(b_i), \quad \beta'_i := a_{3i}(b_i), \quad \beta''_i := \bar{a}_{3i-3}(b_i), \quad \beta_i''' := \bar{a}_{3i}(b_i),$$

and  $i = 2, \dots, g - 1$ .

**Proof** We combine two copies of relation (A) and  $g - 2$  copies of relation (B) to obtain the desired relation. We first consider two relations for simple closed curves shown in Figure 13. One is a copy of relation (A):

$$a_5 a'_4 \delta_1 \delta_4 \delta_6 \delta_2 \delta_3 \delta_5 \\ \equiv a_1 a_{3g+4} b_1 \bar{a}_{3g+4} \sigma'_1 \sigma'_4 a_{3g+3} a_3 b_1 \bar{a}_3 \sigma'_2 a_1 a_3 a_{3g+5} b_1 \bar{a}_{3g+5} \bar{a}_3 \sigma'_3 \sigma'_5 a_2$$

Applying commutativity relations and conjugations, we obtain a relation

$$a'_4 a_5 \delta_1 \delta_2 \delta_3 \delta_4 \delta_5 \delta_6 \\ \equiv a_1 a_2 a_{3g+4} b_1 \bar{a}_{3g+4} \sigma'_1 \sigma'_4 a_{3g+3} a_3 b_1 \bar{a}_3 \sigma'_2 a_1 a_3 a_{3g+5} b_1 \bar{a}_{3g+5} \bar{a}_3 \sigma'_3 \sigma'_5 \\ \equiv a_1 a_2 \beta_1 \sigma'_1 \sigma'_4 a_{3g+3} \beta'_1 \sigma'_2 a_1 \beta''_1 \sigma'_3 \sigma'_5.$$

Note that  $\beta_1 \equiv a_{3g+4} b_1 \bar{a}_{3g+4}$ ,  $\beta'_1 \equiv a_3 b_1 \bar{a}_3$  and  $\beta''_1 \equiv a_{3g+5} a_3 b_1 \bar{a}_3 \bar{a}_{3g+5}$  by Remark 2.2. The other is a copy of relation (B):

$$a_1 a_2 a'_7 a_8 \delta_7 \delta_8 \delta_9 \delta_{10} \equiv a_4 a'_5 \bar{a}_6 b_2 a_6 a_3 b_2 \bar{a}_3 \tau'_1 \tau'''_1 a_5 a'_4 \bar{a}_3 b_2 a_3 a_6 b_2 \bar{a}_6 \tau_1 \tau'''_1$$

Applying commutativity relations and conjugations, we obtain a relation

$$a_1 a_2 a'_7 a_8 \delta_7 \delta_8 \delta_9 \delta_{10} \equiv a_5 a'_4 \bar{a}_3 b_2 a_3 a_6 b_2 \bar{a}_6 \tau_1 \tau'''_1 a_4 a'_5 \bar{a}_6 b_2 a_6 a_3 b_2 \bar{a}_3 \tau'_1 \tau'''_1.$$

We embed two copies of  $\Sigma_{1,8}$  in Figure 13 into  $\Sigma_{2,12}$  as shown in Figure 14.






Figure 13: Relations (A) and (B)




Figure 14: Embeddings of two copies of  $\Sigma_{1,8}$  into  $\Sigma_{2,12}$  (I)

We next consider relation (C2) and another copy of relation (B) for simple closed curves shown in Figure 15:

$$a_4 a'_5 a'_{10} a_{11} \delta_{11} \delta_{12} \delta_{13} \delta_{14} \equiv a_8 a'_7 \bar{a}_6 b_3 a_6 a_9 b_3 \bar{a}_9 \tau_2 \tau_2'' a_7 a'_8 \bar{a}_9 b_3 a_9 a_6 b_3 \bar{a}_6 \tau_2' \tau_2'''$$

We embed  $\Sigma_{2,12}$  in Figure 14 and  $\Sigma_{1,8}$  in Figure 15 into  $\Sigma_{3,16}$  as shown in Figure 16.




Figure 15: Another relation (B)




Figure 16: Embeddings of  $\Sigma_{2,12}$  and  $\Sigma_{1,8}$  into  $\Sigma_{3,16}$

Combining these relations and applying commutativity relations and conjugations, we obtain a relation

$$\begin{aligned}
 (C3) \quad & a'_{10} a_{11} \delta_1 \delta_2 \delta_3 \delta_4 \delta_5 \delta_6 \delta_7 \delta_8 \delta_9 \delta_{10} \delta_{11} \delta_{12} \delta_{13} \delta_{14} \\
 & \equiv \bar{a}'_7 \bar{a}_8 a_8 a'_7 \bar{a}_6 b_3 a_6 a_9 b_3 \bar{a}_9 \tau_2 \tau_2'' a_7 a'_8 \bar{a}_9 b_3 a_9 a_6 b_3 \bar{a}_6 \tau_2' \tau_2'' \bar{a}_4 \bar{a}'_5 \\
 & \quad \cdot a_4 a'_5 \bar{a}_6 b_2 a_6 a_3 b_2 \bar{a}_3 \tau_1' \tau_1'' \cdot \beta_1 \sigma_1' \sigma_4' a_{3g+3} \beta_1' \sigma_2' a_1 \beta_1'' \sigma_3' \sigma_5' \\
 & \quad \cdot \bar{a}_3 b_2 a_3 a_6 b_2 \bar{a}_6 \tau_1 \tau_1''
 \end{aligned}$$

$$\begin{aligned} &\equiv \bar{a}_6 b_3 a_6 a_9 b_3 \bar{a}_9 \tau_2 \tau_2'' a_7 a_8' \bar{a}_9 b_3 a_9 a_6 b_3 \bar{a}_6 \tau_2' \tau_2''' \cdot \bar{a}_6 b_2 a_6 a_3 b_2 \bar{a}_3 \tau_1' \tau_1''' \\ &\quad \cdot \beta_1 \sigma_1' \sigma_4' a_{3g+3} \beta_1' \sigma_2' a_1 \beta_1'' \sigma_3' \sigma_5' \cdot \bar{a}_3 b_2 a_3 a_6 b_2 \bar{a}_6 \tau_1 \tau_1'' \\ &\equiv a_7 a_8' \bar{a}_9 b_3 a_9 a_6 b_3 \bar{a}_6 \tau_2' \tau_2''' \cdot \bar{a}_6 b_2 a_6 a_3 b_2 \bar{a}_3 \tau_1' \tau_1''' \\ &\quad \cdot \beta_1 \sigma_1' \sigma_4' a_{3g+3} \beta_1' \sigma_2' a_1 \beta_1'' \sigma_3' \sigma_5' \cdot \bar{a}_3 b_2 a_3 a_6 b_2 \bar{a}_6 \tau_1 \tau_1'' \\ &\quad \cdot \bar{a}_6 b_3 a_6 a_9 b_3 \bar{a}_9 \tau_2 \tau_2'' . \end{aligned}$$

We repeat similar procedures by making use of  $g - 4$  copies of relation (B):

$$\begin{aligned} &a_{3i-5} a'_{3i-4} a_{3i+1} a_{3i+2} \delta_{4i-1} \delta_{4i} \delta_{4i+1} \delta_{4i+2} \\ &\quad \equiv a_{3i-1} a'_{3i-2} \bar{a}_{3i-3} b_i a_{3i-3} a_3 b_i \bar{a}_{3i} \tau_{i-1} \tau_{i-1}'' \\ &\quad \quad \cdot a_{3i-2} a'_{3i-1} \bar{a}_{3i} b_i a_{3i} a_{3i-3} b_i \bar{a}_{3i-3} \tau_{i-1}' \tau_{i-1}''' \end{aligned}$$

for  $i = 4, \dots, g - 1$  to obtain relations (C4), (C5), ... and

$$\begin{aligned} (C(g-1)) \quad &a_{3g-2} a_{3g-1} \delta_1 \delta_2 \cdots \delta_{4g-3} \delta_{4g-2} \\ &\equiv a_{3g-5} a'_{3g-4} \prod_{i=g-1}^2 \bar{a}_{3i} b_i a_{3i} a_{3i-3} b_i \bar{a}_{3i-3} \tau_{i-1}' \tau_{i-1}''' \\ &\quad \cdot \beta_1 \sigma_1' \sigma_4' a_{3g+3} \beta_1' \sigma_2' a_1 \beta_1'' \sigma_3' \sigma_5' \prod_{i=2}^{g-1} \bar{a}_{3i-3} b_i a_{3i-3} a_3 b_i \bar{a}_{3i} \tau_{i-1} \tau_{i-1}'' \end{aligned}$$

for simple closed curves on  $\Sigma_{g-1,4g}$  shown in Figure 17.

We finally consider the other copy of relation (A) for simple closed curves shown in Figure 18:

$$\begin{aligned} &a_{3g-5} a'_{3g-4} \delta_{4g+1} \delta_{4g+2} \delta_{4g+4} \delta_{4g} \delta_{4g+1} \delta_{4g+3} \\ &\quad \equiv a_{3g-1} a_{3g+1} b_g \bar{a}_{3g+1} \sigma_1 \sigma_4 a_{3g} a_{3g-3} b_g \bar{a}_{3g-3} \sigma_2 \\ &\quad \quad \cdot a_{3g-1} a_{3g-3} a_{3g+2} b_g \bar{a}_{3g+2} \bar{a}_{3g-3} \sigma_3 \sigma_5 a_{3g-2} \end{aligned}$$

Applying commutativity relations and conjugations, we obtain a relation

$$\begin{aligned} &a_{3g-5} a'_{3g-4} \delta_{4g-1} \delta_{4g} \delta_{4g+1} \delta_{4g+2} \delta_{4g+3} \delta_{4g+4} \\ &\quad \equiv a_{3g-2} a_{3g-1} a_{3g+1} b_g \bar{a}_{3g+1} \sigma_1 \sigma_4 a_{3g} a_{3g-3} b_g \\ &\quad \quad \cdot \bar{a}_{3g-3} \sigma_2 a_{3g-1} a_{3g-3} a_{3g+2} b_g \bar{a}_{3g+2} \bar{a}_{3g-3} \sigma_3 \sigma_5 \\ &\quad \equiv a_{3g-2} a_{3g-1} \beta_g \sigma_1 \sigma_4 a_{3g} \beta_g' \sigma_2 a_{3g-1} \beta_g'' \sigma_3 \sigma_5 . \end{aligned}$$

Note that

$$\beta_g \equiv a_{3g+1} b_g \bar{a}_{3g+1}, \quad \beta_g' \equiv a_{3g-3} b_g \bar{a}_{3g-3}, \quad \beta_g'' \equiv a_{3g-3} a_{3g+2} b_g \bar{a}_{3g+2} \bar{a}_{3g-3}$$






Figure 17: Relation  $(C(g - 1))$

by Remark 2.2. We embed  $\Sigma_{g-1,4g}$  in Figure 17 and  $\Sigma_{1,8}$  in Figure 18 into  $\Sigma_{g,4g+4}$  as shown in Figure 12.

Combining these relations and applying commutativity relations and conjugations, we obtain relation  $(H_g)$ . Note that  $\beta_i \equiv a_{3i-3}b_i\bar{a}_{3i-3}$ ,  $\beta'_i \equiv a_{3i}b_i\bar{a}_{3i}$ ,  $\beta''_i \equiv \bar{a}_{3i-3}b_i a_{3i-3}$  and  $\beta'''_i \equiv \bar{a}_{3i}b_i a_{3i}$  ( $i = 2, \dots, g - 1$ ) by Remark 2.2. Thus we complete the proof of Theorem 3.1. □

### 3.2 Genus two

In this subsection we construct a relation on  $\Sigma_{2,12}$  similar to relations constructed in the previous subsection.




Figure 18: The other relation (A)

**Theorem 3.2** Relation (H<sub>2</sub>) For simple closed curves in the interior of Σ<sub>2,12</sub> shown in Figure 19, we have the relation

$\delta_1 \delta_2 \delta_3 \delta_4 \delta_5 \delta_6 \delta_7 \delta_8 \delta_9 \delta_{10} \delta_{11} \delta_{12} \equiv \beta_1 \sigma'_1 \sigma'_4 a_{13} \beta'_1 \sigma'_2 a_1 \beta''_1 \sigma'_3 \sigma'_5 \beta_2 \sigma_1 \sigma_4 a_{12} \beta'_2 \sigma_2 a_5 \beta''_2 \sigma_3 \sigma_5$   
in  $\mathcal{M}_{2,12}$ , where

$$\begin{aligned} \beta_1 &:= a_2(b_1), & \beta'_1 &:= a_3(b_1), & \beta''_1 &:= a_3 a_9(b_1), \\ \beta_2 &:= a_4(b_2), & \beta'_2 &:= a_3(b_2), & \beta''_2 &:= a_3 a_8(b_2). \end{aligned}$$

**Proof** We first consider two copies of relation (A) for simple closed curves shown in Figure 20:

$$\begin{aligned} a_5 a_{14} \delta_7 \delta_8 \delta_9 \delta_{10} \delta_{11} \delta_{12} &\equiv a_1 a_2 b_1 \bar{a}_2 \sigma'_1 \sigma'_4 a_{13} a_3 b_1 \bar{a}_3 \sigma'_2 a_1 a_3 a_9 b_1 \bar{a}_9 \bar{a}_3 \sigma'_3 \sigma'_5 a_{15} \\ a_1 a_{15} \delta_1 \delta_2 \delta_3 \delta_4 \delta_5 \delta_6 &\equiv a_5 a_4 b_2 \bar{a}_4 \sigma_1 \sigma_4 a_{12} a_3 b_2 \bar{a}_3 \sigma_2 a_5 a_3 a_8 b_2 \bar{a}_8 \bar{a}_3 \sigma_3 \sigma_5 a_{14}. \end{aligned}$$

Applying commutativity relations and conjugations, we obtain relations

$$\begin{aligned} a_5 a_{14} \delta_7 \delta_8 \delta_9 \delta_{10} \delta_{11} \delta_{12} &\equiv a_{15} a_1 \beta_1 \sigma'_1 \sigma'_4 a_{13} \beta'_1 \sigma'_2 a_1 \beta''_1 \sigma'_3 \sigma'_5 \\ a_1 a_{15} \delta_1 \delta_2 \delta_3 \delta_4 \delta_5 \delta_6 &\equiv a_{14} a_5 \beta_2 \sigma_1 \sigma_4 a_{12} \beta'_2 \sigma_2 a_5 \beta''_2 \sigma_3 \sigma_5. \end{aligned}$$

Note that  $\beta_1 \equiv a_2 b_1 \bar{a}_2$ ,  $\beta'_1 \equiv a_3 b_1 \bar{a}_3$ ,  $\beta''_1 \equiv a_3 a_9 b_1 \bar{a}_9 \bar{a}_3$ ,  $\beta_2 \equiv a_4 b_2 \bar{a}_4$ ,  $\beta'_2 \equiv a_3 b_2 \bar{a}_3$  and  $\beta''_2 \equiv a_3 a_8 b_2 \bar{a}_8 \bar{a}_3$  by Remark 2.2.

Combining these relations and applying commutativity relations and conjugations, we obtain relation (H<sub>2</sub>). Thus we complete the proof of Theorem 3.2. □




Figure 19: Embeddings of two copies of  $\Sigma_{1,8}$  into  $\Sigma_{2,12}$  (II)




Figure 20: Two copies of relation (A)

## 4 Sections of Lefschetz fibrations

In this section we show that the relation constructed in the previous section gives an explicit topological description of  $4g + 4$  disjoint  $(-1)$ -sections of a hyperelliptic Lefschetz fibration of genus  $g$  on the manifold  $\mathbb{C}\mathbb{P}^2 \# (4g + 5)\overline{\mathbb{C}\mathbb{P}^2}$ .

We begin with a definition of Lefschetz fibrations (see [4; 8]).

**Definition 4.1** Let  $M$  be a closed oriented smooth 4–manifold. A smooth map  $f : M \rightarrow S^2$  is called a *Lefschetz fibration* of genus  $g$  if it satisfies the following conditions:

- (i)  $f$  has finitely many critical values  $b_1, \dots, b_n \in S^2$  and  $f$  is a smooth fiber bundle over  $S^2 - \{b_1, \dots, b_n\}$  with fiber  $\Sigma_{g,0}$
- (ii) for each  $i$  ( $i = 1, \dots, n$ ), there exists a unique critical point  $p_i$  in the *singular fiber*  $f^{-1}(b_i)$  such that  $f$  is locally written as  $f(z_1, z_2) = z_1^2 + z_2^2$  with respect to some local complex coordinates around  $p_i$  and  $b_i$  which are compatible with orientations of  $M$  and  $S^2$
- (iii) no fiber contains a  $(-1)$ –sphere.

**Remark 4.2** We always assume that a Lefschetz fibration is relatively minimal, it has at most one critical point on each fiber, and the genus of the base is equal to zero. A more general definition can be found in [4, Chapter 8].

Suppose that  $g \geq 2$ . According to theorems of Kas and Matsumoto, there exists a one-to-one correspondence between the isomorphism classes of Lefschetz fibrations and the equivalence classes of positive relators modulo simultaneous conjugations

$$c_1 \cdots c_n \sim W(c_1) \cdots W(c_n),$$

and elementary transformations

$$c_1 \cdots c_i \cdot c_{i+1} \cdots c_n \sim c_1 \cdots c_{i+1} \cdot c_{i+1}^{-1}(c_i) \cdots c_n,$$

$$c_1 \cdots c_i \cdot c_{i+1} \cdots c_n \sim c_1 \cdots c_i(c_{i+1}) \cdot c_i \cdots c_n,$$

where  $c_1 \cdots c_n \in \mathcal{R}_{g,0}$  is a *positive relator* in the generators  $\mathcal{S}_{g,0}$  and  $W \in \mathcal{F}_{g,0}$ . This correspondence is described by using the holonomy (or monodromy) homomorphism induced by the classifying map of  $f$  restricted on  $S^2 - \{b_1, \dots, b_n\}$  (see [4; 8]).

**Definition 4.3** Let  $f : M \rightarrow S^2$  be a Lefschetz fibration of genus  $g$ . A smooth map  $s : S^2 \rightarrow M$  is called a *section* of  $f$  if it satisfies  $f \circ s = \text{id}_{S^2}$ . A section  $s$  of  $f$  is an embedding of  $S^2$  into  $M$ . The self-intersection number of the homology class  $s_*([S^2]) \in H_2(M; \mathbb{Z})$  is called the *self-intersection number* of  $s$ . A section of  $f$  with self-intersection number  $k$  is often called a  $k$ –*section*.

For a positive integer  $r$ , we attach  $r$  disks to the boundary components of  $\Sigma_{g,r}$  to obtain a closed surface  $\Sigma_{g,0}$  and an embedding  $\Sigma_{g,r} \hookrightarrow \Sigma_{g,0}$ . This embedding induces a natural commutative diagram

$$\begin{array}{ccccccc}
 1 & \longrightarrow & \mathcal{R}_{g,r} & \longrightarrow & \mathcal{F}_{g,r} & \xrightarrow{\varpi} & \mathcal{M}_{g,r} & \longrightarrow & 1 \\
 & & \lambda \downarrow & & \lambda \downarrow & & \downarrow & & \\
 1 & \longrightarrow & \mathcal{R}_{g,0} & \longrightarrow & \mathcal{F}_{g,0} & \xrightarrow{\varpi} & \mathcal{M}_{g,0} & \longrightarrow & 1,
 \end{array}$$

where the two horizontal sequences are exact. If two words  $W_1$  and  $W_2$  in  $\mathcal{F}_{g,r}$  satisfy  $W_1 \equiv W_2$ , then we have  $\lambda(W_1) \equiv \lambda(W_2)$  in  $\mathcal{F}_{g,0}$ . In this case we call the relation  $W_1 \equiv W_2$  a *lift* of the relation  $\lambda(W_1) \equiv \lambda(W_2)$ .

**Lemma 4.4** [1; 2; 11] *Let  $f : M \rightarrow S^2$  be a Lefschetz fibration of genus  $g$  and  $c_1 \cdots c_n \in \mathcal{R}_{g,0}$  a positive relator corresponding to  $f$ . Suppose that there exists a relation  $a_1 \cdots a_n \equiv \delta_1^{k_1} \cdots \delta_r^{k_r}$  ( $a_1, \dots, a_n \in \mathcal{S}_{g,r}$ ,  $k_1, \dots, k_r > 0$ ) in  $\mathcal{F}_{g,r}$  which is a lift of the relation  $c_1 \cdots c_n \equiv 1$  in  $\mathcal{F}_{g,0}$ , where  $\delta_1, \dots, \delta_r$  are simple closed curves parallel to the boundary components of  $\Sigma_{g,r}$ . Then  $f$  admits disjoint  $r$  sections  $s_1, \dots, s_r : S^2 \rightarrow M$  with self-intersection number  $-k_1, \dots, -k_r$ , respectively.*

For a chain  $(c_1, \dots, c_{2g+1})$  of length  $2g + 1$  on  $\Sigma_{g,0}$ , we obtain a Lefschetz fibration  $X_g \rightarrow S^2$  of genus  $g$  associated to the hyperelliptic relation

$$(c_1 \cdots c_{2g+1} c_{2g+1} \cdots c_1)^2 \equiv 1 \quad \text{in } \mathcal{F}_{g,0}.$$

The total space  $X_g$  of this fibration is known to be diffeomorphic to  $\mathbb{C}\mathbb{P}^2 \# (4g+5)\overline{\mathbb{C}\mathbb{P}^2}$  (see [4; 5]).

We denote the positive word on the right-hand side of relation  $(H_g)$  by  $U_g$  for  $g \geq 2$ . We consider the above embedding  $\Sigma_{g,r} \hookrightarrow \Sigma_{g,0}$  and the commutative diagram for  $r = 4g + 4$ . By Theorems 3.1 and 3.2, relation  $(H_g)$ :  $U_g \equiv \delta_1 \delta_2 \cdots \delta_{4g+3} \delta_{4g+4}$  in  $\mathcal{F}_{g,4g+4}$  is a lift of the relation  $\lambda(U_g) \equiv 1$  in  $\mathcal{F}_{g,0}$ . This implies that the Lefschetz fibration  $Y_g \rightarrow S^2$  of genus  $g$  associated to the relation  $\lambda(U_g) \equiv 1$  admits disjoint  $4g + 4$  sections with self-intersection number  $-1$  by virtue of Lemma 4.4.

**Theorem 4.5** *The two Lefschetz fibrations  $X_g$  and  $Y_g$  are isomorphic to each other.*

**Proof** Suppose that  $g \geq 3$ . We set

$$\begin{aligned}
 c_1 &:= \lambda(a_1), & c_{2i} &:= \lambda(b_i) & (i = 1, \dots, g), \\
 c_{2g+1} &:= \lambda(a_{3g-1}), & c_{2i+1} &:= \lambda(a_{3i}) & (i = 1, \dots, g-1).
 \end{aligned}$$

Since  $(a_1, b_1, a_3, b_2, \dots, a_{3g-3}, b_g, a_{3g-1})$  is a chain of length  $2g + 1$  on  $\Sigma_{g,4g+4}$ ,  $(c_1, c_2, c_3, c_4, \dots, c_{2g-1}, c_{2g}, c_{2g+1})$  is a chain of length  $2g + 1$  on  $\Sigma_{g,0}$ . It is easily seen from Figure 12 that

$$\begin{aligned} \lambda(a_{3g+3}) &= \lambda(a_{3g+4}) = \lambda(a_{3g+5}) = \lambda(\sigma'_1) = \lambda(\sigma'_4) = c_1, \\ \lambda(a_{3g}) &= \lambda(a_{3g+1}) = \lambda(a_{3g+2}) = \lambda(\sigma_1) = \lambda(\sigma_4) = c_{2g+1}, \\ \lambda(\sigma'_2) &= \lambda(\sigma'_3) = \lambda(\sigma'_5) = c_3, \\ \lambda(\sigma_2) &= \lambda(\sigma_3) = \lambda(\sigma_5) = c_{2g-1}, \\ \lambda(\tau'_{i-1}) &= \lambda(\tau'''_{i-1}) = c_{2i-1}, \\ \lambda(\tau_{i-1}) &= \lambda(\tau''_{i-1}) = c_{2i+1}, \end{aligned}$$

for  $i = 2, \dots, g - 1$ . Hence we obtain

$$\begin{aligned} \lambda(U_g) &= \prod_{i=g-1}^2 (\bar{c}_{2i+1}(c_{2i})c_{2i-1}(c_{2i}) \cdot c_{2i-1}^2) \cdot c_1(c_2) \cdot c_1^3 \cdot c_3(c_2) \cdot c_3c_1 \cdot c_1c_3(c_2) \cdot c_3^2 \\ &\quad \cdot \prod_{i=2}^{g-1} (\bar{c}_{2i-1}(c_{2i})c_{2i+1}(c_{2i}) \cdot c_{2i+1}^2) \cdot c_{2g+1}(c_{2g}) \cdot c_{2g+1}^3 \cdot c_{2g-1}(c_{2g}) \\ &\quad \cdot c_{2g-1}c_{2g+1} \cdot c_{2g-1}c_{2g+1}(c_{2g}) \cdot c_{2g-1}^2. \end{aligned}$$

We now prove that  $\lambda(U_g) \sim (c_1 \cdots c_{2g+1}c_{2g+1} \cdots c_1)^2$  for  $g \geq 3$ . Applying elementary transformations (including cyclic permutations), we obtain the following sequence of equivalences:

$$\begin{aligned} \lambda(U_g) &\sim c_{2g-1} \cdot \prod_{i=g-1}^2 (\bar{c}_{2i+1}(c_{2i}) \cdot c_{2i-1}c_{2i}c_{2i-1}) \cdot c_1c_2c_1^2c_3c_2c_1c_3 \cdot c_1(c_2) \cdot c_3 \\ &\quad \cdot \prod_{i=2}^{g-1} (\bar{c}_{2i-1}(c_{2i}) \cdot c_{2i+1}c_{2i}c_{2i+1}) \\ &\quad \cdot c_{2g+1}c_{2g}c_{2g+1}^2c_{2g-1}c_{2g}c_{2g+1}c_{2g-1} \cdot c_{2g+1}(c_{2g}) \\ &\sim \prod_{i=g-1}^2 (c_{2i+1} \cdot \bar{c}_{2i+1}(c_{2i}) \cdot c_{2i-1}c_{2i}) \cdot c_3c_1c_2c_1^2c_3c_2c_1c_3 \cdot c_1(c_2) \\ &\quad \cdot \prod_{i=2}^{g-1} (c_{2i-1} \cdot \bar{c}_{2i-1}(c_{2i}) \cdot c_{2i+1}c_{2i}) \\ &\quad \cdot c_{2g-1}c_{2g+1}c_{2g}c_{2g+1}^2c_{2g-1}c_{2g}c_{2g+1}c_{2g-1} \cdot c_{2g+1}(c_{2g}) \\ &\sim c_1c_{2g+1} \cdot \prod_{i=g-1}^2 c_{2i}c_{2i+1}c_{2i-1}c_{2i} \cdot c_3c_2c_1^2c_3c_2c_1c_3 \cdot c_1(c_2) \end{aligned}$$

$$\begin{aligned}
 & \cdot \prod_{i=2}^{g-1} c_{2i} c_{2i-1} c_{2i+1} c_{2i} \cdot c_{2g-1} c_{2g} c_{2g+1}^2 c_{2g-1} c_{2g} c_{2g+1} c_{2g-1} \cdot c_{2g+1} (c_{2g}) \\
 \sim & \prod_{i=2g-2}^3 c_i c_{i+1} \cdot c_3 c_2 c_1^2 c_3 c_2 c_1 c_3 \cdot c_1 (c_2) \cdot c_1 \\
 & \cdot \prod_{i=3}^{2g-2} c_{i+1} c_i \cdot c_{2g-1} c_{2g} c_{2g+1}^2 c_{2g-1} c_{2g} c_{2g+1} c_{2g-1} \cdot c_{2g+1} (c_{2g}) \cdot c_{2g+1} \\
 \sim & \prod_{i=2g-2}^3 c_i c_{i+1} \cdot c_3 c_2 c_3 c_1^2 c_2 c_1^2 c_3 c_2 \\
 & \cdot \prod_{i=3}^{2g-2} c_{i+1} c_i \cdot c_{2g-1} c_{2g} c_{2g-1} c_{2g+1}^2 c_{2g} c_{2g+1} c_{2g-1} c_{2g+1} c_{2g} \\
 \sim & \prod_{i=2g-2}^3 c_i c_{i+1} \cdot c_2 c_3 c_2 c_1 c_2 c_1 c_2 c_1 c_3 c_2 \\
 & \cdot \prod_{i=3}^{2g-2} c_{i+1} c_i \cdot c_{2g} c_{2g-1} c_{2g} c_{2g+1} c_{2g} c_{2g+1} c_{2g} c_{2g+1} c_{2g-1} c_{2g} \\
 \sim & \prod_{i=2g-2}^2 c_i c_{i+1} \cdot c_1 c_2 c_1 c_1 c_2 c_1 \\
 & \cdot \prod_{i=2}^{2g-1} c_{i+1} c_i \cdot c_{2g+1} c_{2g} c_{2g+1} c_{2g+1} c_{2g} c_{2g+1} c_{2g-1} c_{2g} \\
 \sim & c_{2g} c_{2g+1} c_{2g-1} c_{2g} \cdot \prod_{i=2g-2}^1 c_i c_{i+1} \cdot c_1 c_1 \cdot \prod_{i=1}^{2g} c_{i+1} c_i \cdot c_{2g+1} c_{2g+1} \\
 \sim & \prod_{i=2g}^1 c_i c_{i+1} \cdot c_1 c_1 \cdot \prod_{i=1}^{2g} c_{i+1} c_i \cdot c_{2g+1} c_{2g+1} \\
 \sim & \prod_{i=2g}^1 c_i \cdot \prod_{i=2g+1}^2 c_i \cdot c_1 c_1 \cdot \prod_{i=2}^{2g+1} c_i \cdot \prod_{i=1}^{2g} c_i \cdot c_{2g+1} c_{2g+1} \\
 = & \prod_{i=2g}^1 c_i \cdot \prod_{i=2g+1}^1 c_i \cdot \prod_{i=1}^{2g+1} c_i \cdot \prod_{i=1}^{2g+1} c_i \cdot c_{2g+1} \\
 \sim & \prod_{i=2g+1}^1 c_i \cdot \prod_{i=2g+1}^1 c_i \cdot \prod_{i=1}^{2g+1} c_i \cdot \prod_{i=1}^{2g+1} c_i \\
 \sim & \prod_{i=1}^{2g+1} c_i \cdot \prod_{i=2g+1}^1 c_i \cdot \prod_{i=1}^{2g+1} c_i \cdot \prod_{i=2g+1}^1 c_i = (c_1 \cdots c_{2g+1} c_{2g+1} \cdots c_1)^2.
 \end{aligned}$$

Suppose that  $g = 2$ . We set

$$c_1 := \lambda(a_1), \quad c_2 := \lambda(b_1), \quad c_3 := \lambda(a_3), \quad c_4 := \lambda(b_2), \quad c_5 := \lambda(a_5).$$

Since  $(a_1, b_1, a_3, b_2, a_5)$  is a chain of length 5 on  $\Sigma_{2,12}$ ,  $(c_1, c_2, c_3, c_4, c_5)$  is a chain of length 5 on  $\Sigma_{2,0}$ . It is easily seen from Figure 19 that

$$\begin{aligned} \lambda(a_2) &= \lambda(a_9) = \lambda(a_{13}) = \lambda(\sigma'_1) = \lambda(\sigma'_4) = c_1, \\ \lambda(a_4) &= \lambda(a_8) = \lambda(a_{12}) = \lambda(\sigma_1) = \lambda(\sigma_4) = c_5, \\ \lambda(\sigma'_2) &= \lambda(\sigma'_3) = \lambda(\sigma'_5) = \lambda(\sigma_2) = \lambda(\sigma_3) = \lambda(\sigma_5) = c_3. \end{aligned}$$

Hence we obtain

$$\lambda(U_2) = c_1(c_2) \cdot c_1^3 \cdot c_3(c_2) \cdot c_3c_1 \cdot c_3c_1(c_2) \cdot c_3^2 \cdot c_5(c_4) \cdot c_5^3 \cdot c_3(c_4) \cdot c_3c_5 \cdot c_3c_5(c_4) \cdot c_3^2.$$

We now prove that  $\lambda(U_2) \sim (c_1c_2c_3c_4c_5c_4c_3c_2c_1)^2$ . Applying elementary transformations (including cyclic permutations), we obtain the following sequence of equivalences:

$$\begin{aligned} \lambda(U_2) &\sim c_1c_2c_1^2c_3c_2c_1c_3 \cdot c_1(c_2) \cdot c_3 \cdot c_5c_4c_5^2c_3c_4c_5c_3 \cdot c_5(c_4) \cdot c_3 \\ &\sim c_5c_2c_1^2c_3c_2c_1c_3 \cdot c_1(c_2) \cdot c_3 \cdot c_4c_5^2c_3c_4c_5c_3 \cdot c_5(c_4) \cdot c_3c_1 \\ &\sim c_2c_1^2c_3c_2c_1c_3 \cdot c_1(c_2) \cdot c_1c_3 \cdot c_4c_5^2c_3c_4c_5c_3 \cdot c_5(c_4) \cdot c_3c_5 \\ &\sim c_2c_1^2c_3c_2c_1c_3c_1c_2c_3 \cdot c_4c_5^2c_3c_4c_5c_3 \cdot c_5(c_4) \cdot c_5c_3 \\ &\sim c_2c_1^2c_3c_2c_1c_3c_1c_2c_3c_4c_5^2c_3c_4c_5c_3c_5c_4c_3 \\ &\sim c_1c_3c_2c_3c_1c_1c_2c_3c_4c_3c_5^2c_4c_5c_3c_5c_4c_3c_2c_1 \\ &\sim c_1c_2c_3c_2c_1c_1c_2c_4c_3c_4c_5^2c_4c_5c_3c_5c_4c_3c_2c_1 \\ &\sim c_1c_2c_3c_4c_2c_1c_1c_2c_3c_4c_5c_4c_5c_4c_3c_5c_4c_3c_2c_1 \\ &\sim c_1c_2c_3c_4c_2c_1c_1c_2c_3c_5c_4c_5c_5c_4c_3c_5c_4c_3c_2c_1 \\ &\sim c_1c_2c_3c_4c_5c_2c_1c_1c_2c_3c_4c_5c_5c_4c_3c_5c_4c_3c_2c_1 \\ &\sim c_1c_2c_3c_4c_5c_5c_4c_3c_5c_4c_3c_2c_1c_1c_2c_3c_4c_5c_2c_1 \\ &\sim c_1c_2c_3c_4c_5c_5c_4c_3c_2c_1c_1c_2c_3c_4c_5c_5c_4c_3c_2c_1 \\ &= (c_1c_2c_3c_4c_5c_5c_4c_3c_2c_1)^2. \end{aligned}$$

This completes the proof of Theorem 4.5. □

The next corollary immediately follows from the theorem.

**Corollary 4.6** *The Lefschetz fibration  $X_g \rightarrow S^2$  of genus  $g$  associated to the hyper-elliptic relation admits disjoint  $4g + 4$  sections with self-intersection number  $-1$ .*



By virtue of Theorem 3.2, we can even depict disjoint twelve sections of the Lefschetz fibration  $Y_2 \rightarrow S^2$  in a Kirby diagram of  $Y_2 - \nu F$ , where  $\nu F$  is an open fibered neighborhood of a regular fiber of  $Y_2$  (see [7, Section 4]). We first construct a handle decomposition of  $\Sigma_{2,0} \times D^2$  with one 0–handle, four 1–handles, and one 2–handle with framing 0 from a fixed handle decomposition of  $\Sigma_{2,0}$ . We then attach twenty 2–handles to  $\Sigma_{2,0} \times D^2$  along the simple closed curves  $\beta_1, \sigma'_1, \sigma'_4, a_{13}, \beta'_1, \sigma'_2, a_1, \beta''_1, \sigma'_3, \sigma'_5, \beta_2, \sigma_1, \sigma_4, a_{12}, \beta'_2, \sigma_2, a_5, \beta''_2, \sigma_3, \sigma_5$  (see Figure 19) on different fibers of  $\Sigma_{2,0} \times S^1 \rightarrow S^1$  with framing one less than the product framing of  $\Sigma_{2,0} \times S^1$  to obtain a handle decomposition of  $Y_2 - \nu F$ . Thus we have a Kirby diagram of  $Y_2 - \nu F$  shown in Figure 21. The framing coefficient of every component of the link but one with framing 0 is equal to  $-1$ . Twelve disjoint sections coming from the simple closed curves  $\delta_1, \dots, \delta_{12}$  are represented by twelve unknots transverse to each fiber of the fibration  $\Sigma_{2,0} \times S^1 \rightarrow S^1$  and meeting a fiber at twelve points indicated by encircled numbers  $1, \dots, 12$  in Figure 21. Attaching a 2–handle with framing  $-1$  along any one of the twelve unknots together with four 3–handles and a 4–handle to  $Y_2 - \nu F$ , we have a handle decomposition of the closed manifold  $Y_2$ .

By virtue of Theorem 3.1, we can also depict disjoint  $4g + 4$  sections of the Lefschetz fibration  $Y_g \rightarrow S^2$  in a Kirby diagram of  $Y_g - \nu F$  for  $g \geq 3$  in a similar way.

The following proposition implies that the largest possible number of disjoint  $(-1)$ –sections of  $X_g \rightarrow S^2$  is equal to  $4g + 4$  for most  $g$ .

**Proposition 4.7** *If  $g$  is not equal to  $k^2 + k - 1$  for any positive integer  $k$ , then the Lefschetz fibration  $X_g \rightarrow S^2$  cannot admit disjoint  $4g + 5$  sections with self-intersection number  $-1$ .*

**Proof** Suppose that the Lefschetz fibration  $X_g \rightarrow S^2$  admits disjoint  $4g + 5$  sections  $s_1, \dots, s_{4g+5}$  with self-intersection number  $-1$ . The orientation of  $S^2$  induces that of  $S_i := s_i(S^2)$  for  $i = 1, \dots, 4g + 5$ . We orient a regular fiber  $F$  of  $X_g$  so that it satisfies  $[F] \cdot [S_i] = +1$  for  $i = 1, \dots, 4g + 5$ . Blowing down the  $(-1)$ –spheres  $S_1, \dots, S_{4g+5}$  in  $X_g$ , we obtain a 4–manifold  $X'$  and the image  $F'$  of  $F$  under the projection  $X_g \rightarrow X'$ . Since

$$[F] = [F'] - [S_1] - \dots - [S_{4g+5}] \quad \text{in } H_2(X_g; \mathbb{Z}) \cong H_2(X'; \mathbb{Z}) \oplus (4g + 5)H_2(\overline{\mathbb{C}\mathbb{P}^2}; \mathbb{Z})$$

and  $[F]^2 = 0$ , we have  $[F']^2 = 4g + 5$ . On the other hand,  $[F']^2$  must be the square of an integer because  $[F']$  is a multiple of a generator of  $H_2(X'; \mathbb{Z}) \cong \mathbb{Z}$ . It is easy to see that  $4g + 5$  is the square of an integer if and only if  $g$  is equal to  $k^2 + k - 1$  for some positive integer  $k$ . □




Figure 21: A Kirby diagram of  $Y_2 - \nu F$

**Remark 4.8** Two generic degree  $d$  curves in  $\mathbb{C}\mathbb{P}^2$  induce a Lefschetz pencil of genus  $(d-1)(d-2)/2$ . Blowing up the base locus, we obtain a Lefschetz fibration  $M_d \rightarrow S^2$  of the same genus. This fibration has  $d^2$  sections with self-intersection number  $-1$  and the total space  $M_d$  is diffeomorphic to  $\mathbb{C}\mathbb{P}^2 \# d^2 \overline{\mathbb{C}\mathbb{P}^2}$ . It is well-known that the fibration  $M_3 \rightarrow S^2$  is isomorphic to  $X_1 \rightarrow S^2$ , whereas the fibration  $M_d \rightarrow S^2$  for  $d \geq 4$  cannot be isomorphic to  $X_g \rightarrow S^2$  for any  $g$ .

**Acknowledgement** The author is grateful to Professor Kazuhiro Konno for helpful comments on sections of fibrations on algebraic surfaces, and to Professor Yoshihisa Sato for helpful comments (Remark 4.8) on Lefschetz fibrations. And I wish to express my deep gratitude to my mentor, Hisaaki Endo.

## References

- [1] **J Amorós, F Bogomolov, L Katzarkov, T Pantev**, *Symplectic Lefschetz fibrations with arbitrary fundamental groups*, J. Differential Geom. 54 (2000) 489–545 MR1823313
- [2] **H Endo, M Korkmaz, D Kotschick, B Ozbagci, A Stipsicz**, *Commutators, Lefschetz fibrations and the signatures of surface bundles*, Topology 41 (2002) 961–977 MR1923994
- [3] **B Farb, D Margalit**, *A primer on mapping class groups*, Princeton Math. Series 49, Princeton Univ. Press (2012) MR2850125
- [4] **RE Gompf, AI Stipsicz**, *4-manifolds and Kirby calculus*, Graduate Studies in Mathematics 20, American Mathematical Society (1999) MR1707327
- [5] **T Ito**, *Splitting of singular fibers in certain holomorphic fibrations*, J. Math. Sci. Univ. Tokyo 9 (2002) 425–480 MR1930415
- [6] **S Kitagawa, K Konno**, *Fibred rational surfaces with extremal Mordell–Weil lattices*, Math. Z. 251 (2005) 179–204 MR2176471
- [7] **M Korkmaz, B Ozbagci**, *On sections of elliptic fibrations*, Michigan Math. J. 56 (2008) 77–87 MR2433657
- [8] **Y Matsumoto**, *Lefschetz fibrations of genus two—a topological approach*, from: “Topology and Teichmüller spaces”, (S Kojima, Y Matsumoto, K Saito, M Seppälä, editors), World Sci. Publ., River Edge, NJ (1996) 123–148 MR1659687
- [9] **SÇ Onaran**, *On sections of genus two Lefschetz fibrations*, Pacific J. Math. 248 (2010) 203–216 MR2734172
- [10] **M-H Saitō, K-I Sakakibara**, *On Mordell–Weil lattices of higher genus fibrations on rational surfaces*, J. Math. Kyoto Univ. 34 (1994) 859–871 MR1311624

- [11] **I Smith**, *Geometric monodromy and the hyperbolic disc*, *Q. J. Math.* 52 (2001) 217–228  
MR1838364

*Department of Mathematics, Graduate School of Science  
Osaka University, Osaka 560-0043, Japan  
qqx33r5d@almond.ocn.ne.jp*

Received: 6 March 2012      Revised: 10 August 2012