

Satellites of Legendrian knots and representations of the Chekanov–Eliashberg algebra

LENHARD NG

DANIEL RUTHERFORD

We develop a close relation between satellites of Legendrian knots in \mathbb{R}^3 and the Chekanov–Eliashberg differential graded algebra of the knot. In particular, we generalize the well-known correspondence between rulings of a Legendrian knot in \mathbb{R}^3 and augmentations of its DGA by showing that the DGA has finite-dimensional representations if and only if there exist certain rulings of satellites of the knot. We derive several consequences of this result, notably that the question of existence of ungraded finite-dimensional representations for the DGA of a Legendrian knot depends only on the topological type and Thurston–Bennequin number of the knot.

57R17; 53D42, 57M25

1 Introduction

The satellite construction in knot theory produces new knot types from a given knot K by considering the image of a solid torus knot (or link) $L \subset S^1 \times D^2$ inside a tubular neighborhood of K . This construction provides a simple template for producing whole classes of knot invariants of K since we can simply apply existing invariants to the various satellites of K . As a well-known example, when this scheme is applied to the Jones polynomial, the associated class of satellite invariants are typically organized into a sequence of so-called colored Jones polynomials. This wider collection of invariants generalizes the Jones polynomial to a family of quantum \mathfrak{sl}_2 invariants obtained by labeling K by an arbitrary irreducible representation.

An analog of the satellite construction exists for Legendrian knots in \mathbb{R}^3 with its standard contact structure. In this setting, a Legendrian satellite $S(K, L) \subset \mathbb{R}^3$ arises from a Legendrian pattern, $L \subset J^1(S^1)$, and a Legendrian companion, $K \subset \mathbb{R}^3$, where $J^1(S^1) = T^*S^1 \times \mathbb{R}$ denotes the 1-jet space of S^1 . In this article, we show that using this Legendrian satellite operation, one obtains a correspondence between two well-studied invariants of Legendrian knots: the Chekanov–Eliashberg differential graded algebra (DGA) of the companion and the ruling polynomials of the satellite. (All of the relevant definitions are recalled in Section 2.)

In Legendrian knot theory, there is a well-studied correspondence between augmentations of the Chekanov–Eliashberg DGA of a Legendrian knot K and normal rulings of K ; see eg Fuchs [4], Fuchs and Ishkhanov [5] and Sabloff [17] and Theorem 2.25 below. Here we generalize this to (roughly speaking) a correspondence between finite-dimensional representations of the Chekanov–Eliashberg DGA of K and normal rulings of Legendrian satellites of K ; an augmentation is simply a 1–dimensional representation. In addition, by work of the second author [15] (see Theorem 2.15 below), the existence of either an augmentation of the DGA or a normal ruling depends only on the Thurston–Bennequin number and topological knot type of K . We generalize this result as well to finite-dimensional representations of the DGA.

To give a more precise overview of our main results, we introduce some notation. Denote the Chekanov–Eliashberg differential graded algebra of K over $\mathbb{Z}/2$ by $(\mathcal{A}(K, *), \partial)$. In our notation, $*$ refers to a chosen base point on K that corresponds to a distinguished algebra generator t measuring homology classes in $H_1(K)$. Given a divisor d of twice the rotation number of K , a d –graded representation of $(\mathcal{A}(K, *), \partial)$ consists of a \mathbb{Z}/d –graded vector space V together with a homomorphism of differential graded algebras $f: (\mathcal{A}(K, *), \partial) \rightarrow (\text{End}(V), 0)$. The requirements here are that $f \circ \partial = 0$ and f preserves grading mod d . See Definition 2.27 below.

In Theorem 4.8 we provide necessary and sufficient conditions for the existence of d –graded representations of $(\mathcal{A}(K, *), \partial)$ of any fixed dimension n in terms of so-called normal rulings (see Section 2.5 for the definition of normal ruling) of satellites of K that wrap around K n times. We can give a particularly simple statement in the special case of 1–graded (ie ungraded) representations:

Theorem 1.1 (cf Theorem 4.8) *Let $K \subset \mathbb{R}^3$ be a Legendrian knot. Then the DGA $\mathcal{A}(K, *)$ admits a 1–graded representation of dimension n if and only if the satellite of K with an n –stranded Legendrian full twist, tw_n , has a normal ruling.*

See Figure 1 for an illustration.

Topologically, the satellite in Theorem 1.1 is the n –component link given by n parallel copies of K with respect to framing coefficient $tb(K) + 1$. We also generalize Theorem 4.8 to the case of a general pattern L without cusps, where there is an equivalence between the existence of reduced rulings of $S(K, L)$ and finite-dimensional representations of $\mathcal{A}(K, *)$ in which the distinguished generator t is sent to a matrix related to the path matrix introduced by Kálmán [7]. See Theorem 4.7.

In order to prove Theorems 1.1 and 4.7, we need to study normal rulings of general Legendrian satellites $S(K, L)$ in terms of the pattern $L \subset J^1(S^1)$ and companion




Figure 1: The DGA for a Legendrian knot K (left) admits a representation of dimension $n = 2$ if and only if its satellite with an n -strand full twist (right) has a normal ruling. For the pictured case, both are true: see [19] for a 2-dimensional representation of K , and compare Figure 15 for a ruling of the satellite.

$K \subset \mathbb{R}^3$, and this is an interesting subject in its own right. Many normal rulings of a satellite $S(K, L)$ do not reflect any significant aspect of K . Indeed, an explicit construction (Theorem 3.6 below) shows that any normal ruling of the pattern L may be extended to a normal ruling of $S(K, L)$, so the satellite always has at least as many normal rulings as L . (See Corollary 3.7.) If $S(K, L)$ happens to have more normal rulings than L we say that K is L -compatible.

The question of existence of normal rulings of K then naturally generalizes to the question of L -compatibility. This turns out to be related to the existence of representations of $\mathcal{A}(K, *)$: $\mathcal{A}(K, *)$ has a finite-dimensional representation if and only if we can find a pattern $L \subset J^1(S^1)$ such that K is L -compatible, and for particular L we can classify which sorts of representations of $\mathcal{A}(K, *)$ are equivalent to L -compatibility. (See Section 5 for a detailed and concrete discussion in the case where L has two strands, which corresponds to the case of 2-dimensional representations.) The approach is to reduce from the case of an arbitrary pattern L to the case when L is a product (disjoint union) of particularly simple patterns A_k called *basic fronts*. See Theorems 4.7 and 4.11.

We also extend other results regarding augmentations and rulings to the setting of finite-dimensional representations. For instance, we have the following generalization of the result that the existence of an augmentation of the DGA implies maximal Thurston–Bennequin number and depends only on topological knot type and tb :

Theorem 1.2 (cf Theorem 4.9) *If the DGA $(\mathcal{A}(K, *), \partial)$ has an (ungraded) finite-dimensional representation, then K maximizes tb within its topological knot type.*

Furthermore, the existence of such a representation depends only on $tb(K)$ and the topological type of K .

This result again makes use of the connection with the Kauffman and HOMFLY-PT polynomials as in the work of the second author [15]. There is also a relation to a conjecture by the first author [10] about the topological invariance of the so-called abelianized characteristic algebra; see Section 4.2.

We note that there are examples of Legendrian torus knots with maximal tb that admit 2-dimensional representations but not 1-dimensional representations. This was observed by Sivek in [19]; the question of existence of finite-dimensional representations was raised in the same article, and is discussed in this context in Section 5. At this time, it is an open question whether there are knots that admit 3-dimensional representations but have no representations of dimension 1 or 2, although presumably they exist.

Our work indicates that in a certain concrete way, information about the DGA of various satellites of K is already encoded in the Chekanov–Eliashberg algebra of K itself, though in an algebraically complicated manner. (By comparison, note eg that the colored Jones polynomials for a smooth knot are not determined by the Jones polynomial.) For example, Theorem 1.1 shows that augmentations of the satellite of K with a full twist correspond to n -dimensional representations of $\mathcal{A}(K, *)$. It is interesting to ask how much this situation persists to the plethora of other invariants derived from $\mathcal{A}(K, *)$. For instance, can the collection of linearized homology groups of satellites of K be recovered from $\mathcal{A}(K, *)$?

We conclude this section by outlining the rest of the paper. In Section 2, we recall necessary background about the satellite construction, normal rulings, and the Chekanov–Eliashberg algebra. Section 3 focuses on normal rulings of satellite links. A restricted class of *reduced normal rulings* is introduced as they have a particularly close connection with representations of $\mathcal{A}(K, *)$. Section 4 contains theorems connecting finite-dimensional representations and normal rulings of satellites, including most of the results discussed in this introduction. Finally, Section 5 provides a detailed treatment of a special case, the question of the existence of 2-dimensional representations (with particular restrictions on the image of t). A sufficient condition for A_2 -compatibility is given in Theorem 5.4, and the case of knot types with 10 or fewer crossings is addressed completely.

Acknowledgments

We thank Brad Henry, Steven Sivek and Michael Sullivan for interesting conversations related to this work. The first author was partially supported by NSF CAREER

grant number DMS-0846346. The second author thanks the Max-Planck-Institut für Mathematik in Bonn for providing an excellent research atmosphere during a portion of this work. Finally, we thank the referee for many helpful suggestions.

2 Background

In this section, we give background on Legendrian links in \mathbb{R}^3 and $J^1(S^1)$, the Legendrian satellite construction, normal rulings, the Chekanov–Eliashberg differential graded algebra, and assorted other constructions that will be necessary for the remainder of the paper.

2.1 Legendrian links

We consider Legendrian links in \mathbb{R}^3 and in an open solid torus $S^1 \times \mathbb{R}^2$ with the contact structure provided in either case by the kernel of $dz - y dx$. From the point of view of contact geometry, these spaces are perhaps more naturally viewed as the 1–jet spaces of the line and circle respectively. Correspondingly, we will usually use $J^1(S^1)$ to denote $S^1 \times \mathbb{R}^2$ with this contact structure.

A Legendrian link L in a 1–jet space $J^1(M)$ can be recovered from its image in $M \times \mathbb{R}$ under the projection $T^*M \times \mathbb{R} \rightarrow M \times \mathbb{R}$; this image is referred to as the *front projection* or *front diagram* of L . We will use the same notation for a link and its front diagram, but will point out the distinction when necessary. In the case when $L \subset J^1(S^1)$, the x –coordinate is circle-valued, so the front projection is a subset of an annulus. We write this annulus $S^1 \times \mathbb{R}$ as $[0, 1] \times \mathbb{R}$ with the lines $\{0\} \times \mathbb{R}$ and $\{1\} \times \mathbb{R}$ identified, and we will often view the front projection of a Legendrian link in $J^1(S^1)$ as a subset of $[0, 1] \times \mathbb{R}$.

Generically, front projections are unions of closed curves in the xz –plane or annulus which are immersed away from semi-cubical cusp singularities and one-to-one except for transverse double points. In addition, vertical tangencies cannot occur. Conversely, any collection of curves of this type lifts to a Legendrian link. Two Legendrian links, L_0 and L_1 , are *Legendrian isotopic* if there is a smooth isotopy, L_t , connecting them with L_t Legendrian for all $t \in [0, 1]$. The Legendrian isotopy L_t may always be chosen so that the front projections of the L_t are generic except for a finite number of *Legendrian Reidemeister moves*; see Figure 2. A Legendrian isotopy of this type will be referred to as a *generic isotopy*.

A Legendrian link L has a framing arising from the contact structure. There is a Legendrian isotopy invariant $tb(L) \in \mathbb{Z}$, which equals the corresponding framing




Figure 2: The Legendrian Reidemeister moves, from left to right: I, II, III

number in the case that L is null-homologous. In general, we define $tb(L)$ via a generic front projection for L by $tb(L) = w(L) - \frac{1}{2}c(L)$, where $w(L)$ is the writhe of the projection and $c(L)$ is the number of cusps.

For an oriented (connected) Legendrian knot K , a second integer-valued invariant, the rotation number $r(K)$, is provided by the winding number of the tangent to K around 0 in the contact planes. This is computed from a front diagram as $\frac{1}{2}(d(K) - u(K))$ where $d(K)$ (resp. $u(K)$) denotes the number of downward (resp. upward) oriented cusps. For a multi-component link L , we will adopt the convention of taking $r(L)$ to be the greatest common divisor of the rotation numbers of the components of L .

2.2 Maslov potentials

The following additional structure on a front diagram may be viewed as a generalization of an orientation.

Definition 2.1 Let d be a divisor of $2r(K)$. A d -graded Maslov potential, μ , for L is a function from the front diagram of L to \mathbb{Z}/d that is constant except at cusp points where it increases by 1 when moving from the lower branch of the cusp to the upper branch. See Figure 3. If d is even we assume in addition that μ is even along strands where the orientation of L is in the positive x -direction.




Figure 3: Requirements on a Maslov potential near cusps

A d -graded Legendrian link is a pair (L, μ) consisting of a Legendrian link L with chosen d -graded Maslov potential μ . Maslov potentials may be continued in a canonical way during any of the Legendrian Reidemeister moves. Two d -graded Legendrian links (L_i, μ_i) , $i = 0, 1$, are Legendrian isotopic if there is a generic isotopy from L_0 to L_1 that takes μ_0 to μ_1 .

We emphasize that in Definition 2.1 we allow for the cases $d = 1$ and $d = 2$; the notions of d -graded Legendrian isotopy in these cases correspond to the usual notion of isotopy through unoriented and oriented Legendrian links, respectively. Also, note that if d is a divisor of $2r(K)$ then a d -graded Maslov potential for K always exists. For a single-component Legendrian knot K , Maslov potentials are unique up to the addition of an overall constant. In fact, in \mathbb{R}^3 more is true: if μ and μ' are d -graded Maslov potentials on K , then (K, μ) and (K, μ') are Legendrian isotopic, assuming if d is even that μ and μ' determine the same orientation on K . See Theorem 2.4.

2.3 Legendrian satellites

The following construction first appears in the literature in Ng and Traynor [13], to which we refer the reader for additional details. Let $(K, *) \subset \mathbb{R}^3$ be an oriented (connected) Legendrian knot with chosen base point, $*$, and $L \subset J^1(S^1)$ a link. Using this information, we form a new link $S(K, L) \subset \mathbb{R}^3$ whose Legendrian isotopy type depends only on the Legendrian isotopy types of K and L . The knot K is referred to as the *companion*, L is the *pattern* and $S(K, L)$ is the resulting *satellite*.

Say that $(K, *)$ is in *general position* if its front has generic singularities and $*$ lies away from these, and say that $L \subset J^1(S^1)$ is in *general position* if its front, viewed as a subset of $[0, 1] \times \mathbb{R}$ with ends identified, has generic singularities, all away from $x = 0$. Then we can define $S(K, L)$ diagrammatically.

Let n denote the number of intersection points of the front diagram of L with the vertical line $x = 0$. We begin by forming a link whose front projection is obtained by taking n copies of K and shifting the z -coordinate of each successive copy downward by a small amount. This link is referred to as the n -copy of K . Next, we insert the front projection of L into the n -copy of K at the location of the base point. To do this we view the front projection of L as a subset of $[0, 1] \times \mathbb{R}$ and scale appropriately so that the n intersection points of L with $x = 0$ and $x = 1$ line up with the n parallel strands in the n -copy of K . Furthermore, the scaling should be carried out so that L does not intersect other parts of the n -copy of K . If K is oriented to the right at $*$ then we insert L directly into the n -copy; if K is oriented to the left at $*$ we instead insert the reflection of L across a vertical axis (see [13] for further discussion). See Figure 4. Note that in the case $L = \emptyset$, the satellite $S(K, L)$ is empty as well.

Remark 2.2 It will often be convenient to have an enumeration of the n parallel copies of K that make up the satellite. In this article, we adopt the convention of labeling strands of $S(K, L)$ corresponding to a single strand of K from 1 to n from top to bottom, with the index label increasing as the z -coordinate decreases.




Figure 4: The Legendrian satellite construction

If d is a common divisor of $2r(K)$ and $2r(L)$, then the choice of d -graded Maslov potentials, μ and η , for K and L gives rise to a d -graded Maslov potential, $\tilde{\mu}$, for $S(K, L)$ as follows. At the location of the base point $*$ where L is inserted, $\tilde{\mu}$ is the sum of $\mu(*)$ and η . Since K is connected, this uniquely characterizes $\tilde{\mu}$. Indeed, let η_i denote the value of η on the i^{th} strand of L at $x = 0$. Then at the i^{th} strand of $S(K, L)$ corresponding to the point $k \in K$, $\tilde{\mu} = \mu(k) + \eta_i$.

In [13], generic Legendrian isotopies are given to show that the Legendrian isotopy type of $S(K, L)$ depends only on the Legendrian isotopy types of K and L and, in particular, is independent of the choice of base point $*$. Paying attention to Maslov potentials in the proof shows that the d -graded Legendrian isotopy type of $(S(K, L), \tilde{\mu})$ depends only on that of (K, μ) and (L, η) .

Remark 2.3 The analogous construction for smooth knots requires a choice of framing for K in order to produce a satellite $S(K, L)$ whose isotopy type is well defined. If K

and L are Legendrian, such a framing is given by the contact framing for K , which has framing coefficient $tb(K)$ relative to the Seifert framing; thus in this case the smooth knot type of $S(K, L)$ depends only on $tb(K)$ along with the underlying smooth knot types of K and L .

We conclude this subsection by noting that one can use the satellite construction to give an easy proof of the following result, previously alluded to at the end of Section 2.2.

Theorem 2.4 *Let K be a connected Legendrian knot in \mathbb{R}^3 with two d -graded Maslov potentials μ and μ' , where $\mu' - \mu = k$ for some constant k , with the additional stipulation that if d is even, then k is also even. Then (K, μ) and (K, μ') are Legendrian isotopic.*





Figure 5: Changing Maslov potential for the unknot via Legendrian isotopy

Proof It suffices to prove the theorem when $k = 2$, since a general k is a multiple of 2 (note that when d is odd, every integer is even mod d). If in addition K is the standard unknot U , then the theorem holds since we can apply two Legendrian Reidemeister I moves, then undo these moves, as shown in Figure 5.

For a general knot K , note that K can be expressed as a satellite of the unknot, $K = S(U, L)$ for some $L \subset J^1(S^1)$: simply perform a Reidemeister I move somewhere along the knot, and then the new loop is the unknot U . We can choose Maslov potentials on U and L so that the Maslov potential μ on K is the sum of these two, in the sense discussed in this subsection. By the theorem for the unknot, there is a Legendrian isotopy on U that changes the Maslov potential on U by 2; then the induced Legendrian isotopy on K changes μ by 2 as well. \square

2.4 Basic fronts

For $k \geq 1$, let $A_k \subset J^1(S^1)$ denote the Legendrian knot whose front diagram is given by identifying the ends of the m -stranded braid $\sigma_1\sigma_2 \cdots \sigma_{k-1}$, where strands are labeled from top to bottom, composition of braids is from left to right, and $\sigma_1, \dots, \sigma_{k-1}$ represent the standard generators of the braid group B_k . That is, A_k is the closure of the k -strand front



which winds k times around the S^1 factor and has $k - 1$ crossings. We will refer to the A_k as *basic fronts*. Moreover, for $m \in \mathbb{Z}/d$ we write A_k^m for the basic front A_k with d -graded Maslov potential identically equal to m .

Given front diagrams $L_1, L_2 \subset J^1(S^1)$, we define their product $L_1 \cdot L_2$ by stacking L_1 above L_2 . We introduce some notation for d -graded products of basic fronts (see Figure 6 for an illustration). Let $\lambda = (\lambda_1, \dots, \lambda_\ell)$ be a (finite) sequence of positive integers and $\mathbf{m} = (m_1, \dots, m_\ell)$ a sequence of elements of \mathbb{Z}/d . Writing $\Lambda = (\lambda, \mathbf{m})$ for a pair of such sequences, we define $A_\Lambda \subset J^1(S^1)$ as the d -graded Legendrian link $A_\Lambda = A_{\lambda_1}^{m_1} \cdots A_{\lambda_\ell}^{m_\ell}$. Given such a pair Λ , we introduce a function

$$n_\Lambda: \mathbb{Z}/d \rightarrow \mathbb{Z}_{\geq 0}, \quad \text{where } n_\Lambda(k) = \sum_{i \text{ with } m_i=k} \lambda_i.$$

That is, $n_\Lambda(k)$ denotes the total number of strands of a fixed x -coordinate with Maslov potential equal to k .




Figure 6: The product of basic fronts A_Λ , where $\Lambda = (\lambda, \mathbf{m})$ with $\lambda = (3, 2, 2)$ and $\mathbf{m} = (1, 2, 1)$. The function n_Λ satisfies $n_\Lambda(1) = 5$, $n_\Lambda(2) = 2$, and $n_\Lambda(k) = 0$ for $k \neq 1, 2$. Orientations are determined by the reduction of $\mathbf{m} \pmod 2$. (See Definition 2.1.)

Later, we are able to reduce some questions about satellites with arbitrary pattern $L \subset J^1(S^1)$ to the case where L is a product of basic fronts.

2.5 Normal rulings

The notion of normal ruling was developed independently in the works Pushkar' and Chekanov [14] and Fuchs [4]. Let L be a Legendrian knot in \mathbb{R}^3 or $J^1(S^1)$ whose front projection is generic in the sense described earlier in this section. In addition, we now assume that all crossings and cusps have distinct x -coordinates. This can be achieved after a small Legendrian isotopy.

In this section, we unify our notation by viewing $\mathbb{R}^3 \cong J^1(\mathbb{R})$. For $M = \mathbb{R}$ or S^1 , we let $\pi: M \times \mathbb{R} \rightarrow M$ denote the projection $\pi(x, z) = x$, where the domain is viewed as the front projection of $J^1(M)$. Let $\Sigma \subset M$ denote the projection of the set of cusp and crossing points of the front projection of L . Furthermore, for any $x \in M$ let $L_x = \pi^{-1}(x)$.

Definition 2.5 A continuous function f from a subset $N \subset M$ to the front projection of $L \subset M \times \mathbb{R}$ is called a *section* if $\pi \circ f = id_N$.

Definition 2.6 A *normal ruling* of the front projection of a link L in \mathbb{R}^3 or $J^1(S^1)$ is a continuous involution, $\rho: L \setminus \pi^{-1}(\Sigma) \rightarrow L \setminus \pi^{-1}(\Sigma)$, $\rho^2 = id_{L \setminus \pi^{-1}(\Sigma)}$, satisfying the following:

- (i) The involution ρ is fixed point free.
- (ii) The involution satisfies $\pi \circ \rho = \pi|_{L \setminus \pi^{-1}(\Sigma)}$ and therefore restricts to involutions $\rho_x: L_x \rightarrow L_x$ for each $x \in M \setminus \Sigma$.
For a component $N \subset M \setminus \Sigma$, the preimage of N in L is a union of “strands”, $\pi^{-1}(N) = \bigsqcup S_i$, where each S_i is mapped homeomorphically onto N by π . Due to the continuity condition, ρ induces an involution of the collection of strands, and we say ρ pairs S_i with S_j if $\rho(S_i) = S_j$.
- (iii) In a neighborhood of a cusp point, the involution ρ interchanges the upper and lower branch of the cusp. On the remaining strands, the involution induced by ρ_x should be the same on either side of the cusp.
- (iv) Strands meeting at a crossing should not be paired by ρ .
- (v) The involution ρ extends continuously near crossings in the following sense. If $N \subset M$ is such that $\pi^{-1}(N)$ contains a single crossing at x -coordinate x_0 , then we can find sections $f_1, f_2, \dots, f_N: N \rightarrow L \subset M \times \mathbb{R}$ such that every point of $L \cap \pi^{-1}(N)$ is in the image of exactly one of the f_i except for the

crossing point, which is in the image of exactly two of the f_i . Moreover, these sections are preserved by the involution. That is, for any $i = 1, \dots, N$, there exists j such that $\rho \circ f_i = f_j$ on $N \setminus \{x_0\}$. It is clear that, except for their enumeration, the sections are uniquely determined by the involution ρ .

At the crossing point, there are two possibilities. Either the two sections that meet at the crossing follow the diagram and cross in a transverse manner, or they each turn a corner at the crossing. In the latter case, we refer to the crossing as a *switch* of ρ . At a switch, one section covers the upper half of the crossing, and another covers the lower half. Due to requirements (i) and (iv), each of these sections is paired by ρ with a *companion strand* away from the crossing.

- (vi) (*Normality condition*) Near a switch we can produce intervals on the vertical axis by connecting each switching strand with its companion strand. These two intervals should either be disjoint or one should be contained in the other. See Figure 7.




Figure 7: The normality condition

Definition 2.7 Suppose now that (L, μ) is a d -graded Legendrian link. We say that a ruling ρ of L is d -graded with respect to μ if for $(x, z), (x, z') \in L \setminus \pi_x^{-1}(\Sigma)$ with $z < z'$ and $\rho(x, z) = (x, z')$, we have $\mu(x, z') = \mu(x, z) + 1 \pmod{d}$.

Note that every normal ruling is 1-graded. In the literature 1-graded rulings are often called “ungraded” rulings.

Remark 2.8 Alternatively, a normal ruling may be viewed as a global decomposition of the front diagram into pairs of sections, and we will make use of this perspective in our figures and proofs. In the case of a link in $J^1(S^1)$, it is important here that we view the front diagram as a subset of $[0, 1] \times \mathbb{R}$. Then starting at $x = 0$ or at the first left cusp of L and working to the right, we can piece together the sections f_i from condition (v). This allows us to cover the front diagram of L with a collection of sections with maximal domains of definition. The involution then divides the sections into pairs (P_i, Q_i) that begin and end at common cusps or possibly at common components of the boundary of $[0, 1] \times \mathbb{R}$ in the case $L \subset J^1(S^1)$.

Note that in the case $L \subset J^1(S^1)$, a section that begins at $x = 0$ does not necessarily have to end up at the same z -coordinate if it makes it all the way to $x = 1$ without terminating at a cusp. However, the involution ρ is defined on the front diagram L viewed as a subset of $S^1 \times \mathbb{R}$, so the overall division of points of L into pairs at $x = 0$ and $x = 1$ should be the same. See Figure 8 for an example.




Figure 8: A normal ruling of a 2-component link in $J^1(S^1)$

Example 2.9 A normal ruling ρ for a 2-component link $L \subset J^1(S^1)$ is pictured in Figure 8. With respect to the indicated Maslov potential, ρ is 0-graded. However, if additional switches were added at the 2nd and 4th crossings (numbered from left to right), then the resulting normal ruling would only be 1-graded.

For Legendrian links in $J^1(S^1)$, it is appropriate for some purposes (see Lavrov and Rutherford [8]) to relax the fixed point free condition of Definition 2.6(i).

Definition 2.10 Let $L \subset J^1(S^1)$. A *generalized normal ruling* of L is an involution ρ satisfying the requirements of Definition 2.6 except for the following modifications.

- (i) The involution may have fixed points. Near crossings, the locally defined sections f_i are no longer uniquely determined by ρ in the case where both of the crossing strands are fixed by ρ . However, if at least one of the crossing strands is not fixed by ρ , then uniqueness still holds. In particular, it is possible to have a switch where one of the switching strands has a companion strand and the other is a fixed point strand. In this case, the normality condition is extended.
- (ii) (*Generalized normality condition*) Near switches where one of the strands is fixed by ρ , the vertical interval connecting the non-fixed point strand to its companion strand should not intersect the other strand of the switch. See Figure 9.

Definition 2.7 carries over without change to provide a notion of d -graded generalized normal ruling.



Figure 9: The generalized normality condition

Example 2.11 For $k \geq 1$, the basic front A_k does not admit a normal ruling. However, each A_k does admit a generalized normal ruling where the involution ρ is simply the identity.

Remark 2.12 A generalized normal ruling produces a decomposition of the front diagram of L into pairs of sections (P_i, Q_i) and a fixed point subset F that does not contain cusps. (Compare Remark 2.8.) We make use of this perspective in our figures. See Figure 11 below for an example of a generalized ruling.

Given a d -graded Legendrian link (L, μ) in $J^1(\mathbb{R})$ or $L \subset J^1(S^1)$, let $\mathcal{R}^d(L, \mu)$ (resp. $\mathcal{GR}^d(L, \mu)$) denote the set of all normal rulings (resp. generalized normal rulings) of L that are d -graded with respect to μ . Finally, we define the d -graded ruling polynomial, $R^d_{(L, \mu)}$, by

$$R^d_{(L, \mu)}(z) = \sum_{\rho \in \mathcal{R}^d(L, \mu)} z^{j(\rho)}, \quad \text{where } j(\rho) = \#\text{switches} - \#\text{right cusps}.$$

Chekanov and Pushkar' [14] prove the following invariance result.

Theorem 2.13 [14] *If (L_1, μ_1) and (L_2, μ_2) are Legendrian isotopic as d -graded links, then $R^d_{(L_1, \mu_1)}(z) = R^d_{(L_2, \mu_2)}(z)$.*

Remark 2.14 Note that ruling polynomials are unchanged by the addition of an overall constant to the Maslov potential. In particular, if K is a (connected) knot, then the d -graded ruling polynomials are independent of the choice of μ . Moreover, if $K \subset \mathbb{R}^3$ is an oriented knot, then for any d -graded $(L, \eta) \subset J^1(S^1)$ with $d \mid 2r(K)$, the polynomial $R^d_{S(K, L)}(z)$ is a Legendrian isotopy invariant of K .

When $d = 1$ or 2 the ruling polynomials depend only on the underlying framed knot type of L . This follows from:

Theorem 2.15 [15] *For any Legendrian link $L \subset \mathbb{R}^3$, let $F_L, P_L \in \mathbb{Z}[a^{\pm 1}, z^{\pm 1}]$ denote the Kauffman and HOMFLY-PT link polynomials.¹ Then the 1-graded (resp.*

¹We follow here the conventions of [15] for the HOMFLY-PT and Kauffman polynomials. However, our conventions for the power $j(\rho)$ appearing in the ruling polynomial differ by 1 from [15].

2-graded) ruling polynomial $R_L^1(z)$ (resp. $R_L^2(z)$) is equal to z^{-1} times the coefficient of $a^{-tb(L)-1}$ in F_L (resp. P_L).

Remark 2.16 An analogous but more complicated result holds for links in $J^1(S^1)$. See Rutherford [16], and Lavrov and Rutherford [8].

2.6 Chekanov–Eliashberg differential graded algebra

In this subsection we recall the definition of the Chekanov–Eliashberg DGA associated to a Legendrian knot in \mathbb{R}^3 , with some adjustments (related to base points and commutativity) to adapt the standard treatment to the needs of this paper.

For the purposes of defining the DGA, it is more convenient to work in the Lagrangian (xy) projection than in the front (xz) projection used in the first part of this section. One can use an elementary construction called resolution [10] to produce a Lagrangian projection from a front projection: diagrammatically, smooth out all left cusps, and replace right cusps by a loop with a negative crossing.

Let K be an oriented Legendrian knot with a base point $*$, generic in the sense that the Lagrangian projection $\pi_{xy}(K)$ is immersed with only transverse double points as singularities, and $\pi_{xy}(*)$ lies away from the double points. Contact homology associates to $(K, *)$ a differential graded algebra $(\mathcal{A}(K, *), \partial)$, the *Chekanov–Eliashberg algebra*, as we now briefly recall; see eg Chekanov [1] and Etnyre, Ng and Sabloff [3] for more details.

Definition 2.17 Label the crossings (double points) of $\pi_{xy}(K)$ by a_1, \dots, a_n . The algebra $\mathcal{A}(K, *)$ is the associative, noncommutative unital algebra over $\mathbb{Z}/2$ generated by

$$a_1, \dots, a_n, t, t^{-1}$$

with no relations besides $t \cdot t^{-1} = t^{-1} \cdot t = 1$.

The algebra $\mathcal{A}(K, *)$ is generated as a $(\mathbb{Z}/2)$ -vector space by words of the form

$$t^{\alpha_0} a_{i_0} t^{\alpha_1} a_{i_1} \dots a_{i_k} t^{\alpha_k}$$

(including the empty word, which serves as the identity element 1), with multiplication given by concatenation. We note that this definition of $\mathcal{A}(K, *)$ is slightly different from the corresponding definition in [3], even accounting for the fact that we work over $\mathbb{Z}/2$ and not \mathbb{Z} : the algebra considered in [3] is the quotient of ours by allowing powers of t to commute with the a_i . This construction follows Ng [11]; see Ekholm, Etnyre, Ng and Sullivan [2, Section 2.3.2] for further discussion.

We next give $\mathcal{A}(K, *)$ a $\mathbb{Z}/(2r(K))$ -grading (and thus a (\mathbb{Z}/d) -grading for any $d \mid 2r(K)$). If a_i is a crossing of $\pi_{xy}(K)$, let $\gamma_i \subset \mathbb{R}^2$ be a path along $\pi_{xy}(K)$ beginning at the overcrossing and ending at the undercrossing of a_i . Let $r(\gamma_i)$ be the (non-integral) number of counterclockwise revolutions made by the unit tangent vector to the path γ_i from beginning to end, and define $|a_i| = \lfloor 2r(\gamma_i) \rfloor$. Note that the grading is well-defined, independent of the choice of γ_i , modulo $2r(K)$. Extend the grading to all of $\mathcal{A}(K, *)$ by setting $|t| = |t^{-1}| = 0$ and extending in the usual way (the degree of a product is the sum of the degrees).

If $\pi_{xy}(K)$ is a resolution of a front diagram endowed with a Maslov potential, then the degree of crossing a_i is the difference $k - l$ where k (resp. l) is the value of the Maslov potential on the overstrand (resp. understrand) passing through a_i . We can also use this to define a grading on the DGA in the more general case where K is a multi-component link.

We now come to the differential on \mathcal{A} . Attach signs to each corner at every crossing in $\pi_{xy}(K)$ as depicted:



Let D^2 denote the closed disk. For $\ell \geq 0$, write $D_\ell^2 = D^2 \setminus \{r, s_1, \dots, s_\ell\}$, where r, s_1, \dots, s_ℓ are points on ∂D^2 appearing in order as we traverse the boundary counterclockwise.

Definition 2.18 Let a, b_1, \dots, b_ℓ be crossings in the Lagrangian projection $\pi_{xy}(K)$ of a Legendrian knot $K \subset \mathbb{R}^3$. Define $\Delta(a; b_1, \dots, b_\ell)$ to be the space of all orientation-preserving immersions $f: (D_\ell^2, \partial D_\ell^2) \rightarrow (\mathbb{R}^2, \pi_{xy}(K))$, up to reparametrization, such that:

- f sends the boundary punctures of D_ℓ^2 to the crossings of $\pi_{xy}(K)$,
- f sends a neighborhood of r to a corner at a labeled by a $+$,
- f sends a neighborhood of each s_i to a corner at b_i labeled by a $-$.

For $f \in \Delta(a; b_1, \dots, b_\ell)$, the image of ∂D_ℓ^2 maps to a union of $\ell + 1$ paths $\gamma_0, \dots, \gamma_\ell \subset \pi_{xy}(K)$, where each path begins and ends at a crossing, and γ_0 goes from a to b_1 , γ_i from b_i to b_{i+1} for $1 \leq i \leq \ell - 1$, and γ_ℓ from b_ℓ to a . For each of these paths γ_i , we can associate a monomial $w(\gamma_i) \in \mathbb{Z}/2[t^{\pm 1}]$ by $w(\gamma_i) = t^{\alpha_i}$, where α_i is the number of times γ_i passes through the base point $*$, counted with sign according to the orientation of K . Finally, we can associate a monomial $w(f) \in \mathcal{A}(K, *)$ to f , as follows:

$$w(f) = w(\gamma_0)b_1w(\gamma_1)b_2 \cdots b_\ell w(\gamma_\ell).$$

Definition 2.19 Let a be a crossing of K . The differential $\partial(a)$ is defined by:

$$\partial(a) = \sum_{f \in \Delta(a; b_1, \dots, b_\ell)} w(f),$$

where the sum is over all $\ell \geq 0$ and all choices of crossings b_1, \dots, b_ℓ such that $\Delta(a; b_1, \dots, b_\ell)$ is nonempty.

We can extend the map ∂ to all of $\mathcal{A}(K, *)$ by setting $\partial(t) = \partial(t^{-1}) = 0$ and imposing the Leibniz rule.

Theorem 2.20 [1; 3] *The map $\partial: \mathcal{A}(K, *) \rightarrow \mathcal{A}(K, *)$ lowers degree by 1 and is a differential: $\partial^2 = 0$. Up to stable tame isomorphism, the differential graded algebra $(\mathcal{A}(K, *), \partial)$ is an invariant of K under Legendrian isotopy (and choice of base point).*

Here “stable tame isomorphism” is an equivalence relation of differential graded algebras that in particular fixes t and preserves isomorphism type of the homology $H_*(\mathcal{A}(K*), \partial)$; see [1] or [2] for a definition in our setting. For the purposes of this paper, this relation may be treated as a black box.

In Section 4, we will need a slight generalization of the above notion of the Chekanov–Eliashberg DGA, to the setting where we have multiple base points $*_1, \dots, *_k$ on K . As before, we assume that in the xy projection, no base point coincides with a crossing; we also assume that the base points are cyclically ordered along K , ie $*_1, \dots, *_k$ are encountered in that order as we traverse the knot in the direction of its orientation.

Given this data, we define the algebra $\mathcal{A}(K, *_1, \dots, *_k)$ to be the noncommutative unital algebra over $\mathbb{Z}/2$ generated by crossings a_1, \dots, a_n , along with $2k$ additional generators $t_1^{\pm 1}, \dots, t_k^{\pm 1}$, with no relations besides $t_i \cdot t_i^{-1} = t_i^{-1} \cdot t_i = 1$ for all i . (Note in particular that the t_i do not commute with the a ’s, or indeed with each other.) We give $\mathcal{A}(K, *_1, \dots, *_k)$ a $\mathbb{Z}/(2r(K))$ –grading as before, with $|t_i| = |t_i^{-1}| = 0$ for all i .

We can define a differential ∂ on $\mathcal{A}(K, *_1, \dots, *_k)$ analogously to Definition 2.19. Note that in the presence of multiple base points, the monomial $w(\gamma)$ associated to a path γ in $\pi_{xy}(K)$ can involve any or all of $t_1^{\pm 1}, \dots, t_k^{\pm 1}$: it is the product $t_{i_1}^{\pm 1} \dots t_{i_l}^{\pm 1}$, where γ passes through $*_{i_1}, \dots, *_{i_l}$ in succession, and the signs depend on whether the orientation of γ agrees or disagrees with the orientation of K as γ passes through the base point.

The DGA $(\mathcal{A}(K, *_1, \dots, *_k), \partial)$ depends minimally on the choice of base points, and indeed contains no more information than the single-base-pointed DGA $(\mathcal{A}(K, *), \partial)$. More precisely, we have the following results.

Theorem 2.21 Let $*_1, \dots, *_k$ and $*'_1, \dots, *'_k$ denote two collections of base points on K , each of which is cyclically ordered along K . Let $(\mathcal{A}(K, *_1, \dots, *_k), \partial)$ and $(\mathcal{A}(K, *'_1, \dots, *'_k), \partial')$ denote the corresponding multi-pointed DGAs. Then there is a DGA isomorphism $\phi: (\mathcal{A}(K, *_1, \dots, *_k), \partial) \rightarrow (\mathcal{A}(K, *'_1, \dots, *'_k), \partial')$ such that $\phi(t_i) = t_i$ for all i .

Proof It suffices to establish the result when $(*_1, \dots, *_k)$ and $(*'_1, \dots, *'_k)$ are identical except that for some i , $*'_i$ is the result of sliding $*_i$ across a crossing of $\pi_{xy}(K)$. Suppose then that $*_i$ and $*'_i$ lie on opposite sides of a crossing a_l , with the orientation of K pointing from $*_i$ to $*'_i$. We first consider the case where the strand containing $*_i$ and $*'_i$ is the overstrand at a_l . In this case, if f is a disk with a positive corner at a_l and $w(f)$ and $w'(f)$ are the words associated to f in $\mathcal{A}(K, *_1, \dots, *_j, \dots, *_k)$ and $\mathcal{A}(K, *_1, \dots, *'_j, \dots, *_k)$, respectively, then $w'(f) = t_i w(f)$. Furthermore, if f is a disk with a negative corner at a_l , then $w'(f)$ is the result of replacing a_l by $t_i^{-1} a_l$ in $w(f)$. It follows that the map ϕ defined by $\phi(a_l) = t_i^{-1} a_l$, $\phi(a_j) = a_j$ for $j \neq l$, and $\phi(t_j) = t_j$ for all j satisfies $\phi \circ \partial = \partial' \circ \phi$.

If the strand containing $*_i$ and $*'_i$ is the understrand at a_l , a similar argument shows that the map ϕ defined by $\phi(a_l) = a_l t_i$, $\phi(a_j) = a_j$ for $j \neq l$, and $\phi(t_j) = t_j$ for all j satisfies $\phi \circ \partial = \partial' \circ \phi$. □

Theorem 2.22 Let $*_1, \dots, *_k$ be a cyclically ordered collection of base points along K , and let $*$ be a single base point on K . Then there is a DGA homomorphism $\phi: (\mathcal{A}(K, *), \partial) \rightarrow (\mathcal{A}(K, *_1, \dots, *_k), \partial)$ such that $\phi \circ \partial = \partial \circ \phi$ and $\phi(t) = t_1 \cdots t_k$.

Proof By Theorem 2.21, we may assume that $*_1, \dots, *_k$ all lie in a small neighborhood of $*$. In this case, the map ϕ defined by $\phi(a_i) = a_i$ for all crossings a_i and $\phi(t) = t_1 \cdots t_k$ is the desired homomorphism. □

2.7 Augmentations and representations of the DGA

Here we discuss representations of the DGA introduced in the previous subsection. We begin with augmentations, which can be viewed as 1–dimensional representations.

Definition 2.23 Let $d \mid 2r(K)$. A d –graded augmentation of $(\mathcal{A}(K, *), \partial)$ is an algebra map $\epsilon: \mathcal{A}(K, *) \rightarrow \mathbb{Z}/2$ such that:

- $\epsilon(1) = \epsilon(t) = \epsilon(t^{-1}) = 1$
- $\epsilon \circ \partial = 0$
- $\epsilon(a) = 0$ if $a \in \mathcal{A}$ with $|a| \not\equiv 0 \pmod{d}$

Stable tame isomorphism (discussed briefly in the previous subsection) preserves the existence and nonexistence of d –graded augmentations. Theorem 2.20 then immediately implies the following.

Theorem 2.24 *If K and K' are Legendrian isotopic knots with base points $*$ and $*'$, then for any $d \mid 2r(K)$, the Chekanov–Eliashberg DGA $(\mathcal{A}(K, *), \partial)$ has a d –graded augmentation if and only if $(\mathcal{A}(K', *'), \partial)$ does.*

There is a well-known correspondence between augmentations and rulings:

Theorem 2.25 [4; 5; 17] *Let K be a Legendrian knot in \mathbb{R}^3 , and let $d \mid 2r(K)$. Then the front projection of K has a d –graded ruling if and only if the DGA $(\mathcal{A}(K, *), \partial)$ has a d –graded augmentation.*

We need the following more precise statement. Recall that a Legendrian link in \mathbb{R}^3 is said to be in *plat position* if all right cusps have the same x –coordinate as do all left cusps.

Theorem 2.26 [17; 6] *Let K be a d –graded Legendrian link in \mathbb{R}^3 with front diagram in plat position, and denote by (\mathcal{A}, ∂) the Chekanov–Eliashberg DGA associated with the resolution of K .*

- (i) *Given any d –graded augmentation of (\mathcal{A}, ∂) , there exists a d –graded normal ruling ρ of K so that the first switch of ρ occurs at or to the right of the first augmented crossing.*
- (ii) *For any d –graded normal ruling ρ of K , there is a d –graded augmentation of (\mathcal{A}, ∂) so that the first augmented crossing agrees with the first switch of ρ .*

In the statement of Theorem 2.26, we order crossings from left to right. Given an augmentation, ϵ , we say that a crossing, a_i , is augmented if $\epsilon(a_i) \neq 0$.

Proof Statement (i) follows from an algorithm in [17, Section 3.3] that assigns a normal ruling to an augmentation of a plat position front diagram. This algorithm is also presented in Ng and Sabloff [12], and it is easy to see that the first switch of the normal ruling must be to the right of the first augmented crossing.

For (ii), we cite work of Henry [6]. The main objects of study in [6] are “Morse complex sequences,” which consist of sequences of chain complexes assigned to a front diagram of a Legendrian link in \mathbb{R}^3 . In [6, Section 6.5], two different standard forms for a Morse complex sequence (MCS) are introduced, the $S\bar{R}$ –form and the A –form. These standard forms are related to normal rulings and augmentations, respectively.

Given a normal ruling ρ as in (ii), we consider the $S\bar{R}$ -form MCS associated with ρ where none of the returns have handleslides. [6, Theorem 6.20] shows that this MCS may be transformed into an A -form MCS by an algorithm that sweeps handleslide marks from left to right. In particular, the leftmost handleslide in this A -form MCS will be located directly to the left of the first switch of ρ . According to [6, Corollary 6.29], this A -form MCS corresponds to an augmentation of the Chekanov–Eliashberg algebra of the resolution of K where a crossing is augmented if and only if there is a handleslide immediately to the left of the crossing. This augmentation has the desired form. \square

We now generalize our discussion from augmentations to representations of the DGA. Suppose V is a finite-dimensional vector space over $\mathbb{Z}/2$ with a \mathbb{Z}/d -grading,

$$V = \bigoplus_{k \in \mathbb{Z}/d} V_k.$$

Then

$$\text{End}(V) = \bigoplus_{i,j} \text{Hom}_{\mathbb{Z}/2}(V_i, V_j)$$

is a \mathbb{Z}/d -graded algebra where we take each $\text{Hom}_{\mathbb{Z}/2}(V_i, V_j)$ to be homogeneous of degree² $i - j \in \mathbb{Z}/d$.

Definition 2.27 A d -graded representation of (\mathcal{A}, ∂) is a d -graded vector space V over $\mathbb{Z}/2$ along with a DGA map from (\mathcal{A}, ∂) to $(\text{End}(V), 0)$, ie, a grading-preserving algebra map $f: \mathcal{A} \rightarrow \text{End}(V)$ satisfying $f(1) = id_V$ and $f \circ \partial = 0$. The *graded dimension* of the representation, $\dim(V)$, is the function $\mathbf{n}: \mathbb{Z}/d \rightarrow \mathbb{Z}_{\geq 0}$ defined by $\mathbf{n}(k) = \dim V_k$.

Note that a representation of $(\mathcal{A}(K, *), \partial)$ does not need to send t to id_V , but merely to an invertible map on V .

3 Normal rulings of Legendrian satellites

We begin this section by establishing some basic properties of normal rulings of Legendrian satellites. The definition of normal ruling requires working with a front projection of $S(K, L)$ with the property that crossings have distinct x -coordinates. To achieve this, we assume that $(K, *)$ and L are in general position, and then apply a planar isotopy to perturb the x -coordinates of the crossings of the front diagram for

² Note that with this convention, our definition of degree is the *negative* of the standard grading for graded linear maps.

$S(K, L)$ described in Section 2.3. The precise order that the crossings end up in will not be relevant for our arguments. In this section, we continue to use the convention of labeling the parallel translates of a strand of K appearing in $S(K, L)$ from 1 to n with decreasing z -coordinate.

We introduce some terminology associated with a normal ruling ρ of $S(K, L)$. In the following definition we use the term *strand* in a local sense so that it refers to an arc on the front projection that projects homeomorphically to an arc on the x -axis.

Definition 3.1 Outside of neighborhoods of cusps, crossings and the base point, we can assign an involution, ρ_T , of $\{1, \dots, n\}$ to each point $k_0 \in K$ according to:

- (i) Let $\rho_T(i) = j$ if, of the n parallel strands in $S(K, L)$ corresponding to a small neighborhood of k_0 in K , the ruling ρ pairs the i^{th} strand with the j^{th} strand, and
- (ii) let $\rho_T(i) = i$ if ρ pairs the i^{th} strand corresponding to k_0 in $S(K, L)$ with a strand of $S(K, L)$ corresponding to a point other than k_0 in K .

We will refer to ρ_T as the *thin part* of ρ at k_0 . In addition, we refer to strands of $S(K, L)$ that correspond to the same strand of K and are paired by ρ as a pair of *thin strands* of ρ .

Example 3.2 In Figure 10, a portion of a normal ruling ρ of a satellite $S(K, L)$ is pictured. The part of $S(K, L)$ appearing there corresponds to a single left cusp of K . The thin part of ρ satisfies $\rho_T(1) = 4$, $\rho_T(2) = 2$ and $\rho_T(3) = 5$ both before and after the cusp.

In combination, the following two lemmas show that the thin part of ρ is independent of $k_0 \in K$.

Lemma 3.3 (Crossing Lemma) *A pair of thin strands cannot be involved in any switches at crossings of $S(K, L)$ that correspond to a crossing of K .*

Proof Let Y denote the subset of the front diagram $S(K, L)$ corresponding to a neighborhood of a crossing q of K . We prove the more general statement:

Let P and Q be a pair of companion paths of the ruling. If both P and Q pass through the region Y , then neither of them can switch in Y .

The proof is by induction on M , the number of strands lying in the vertical interval between P and Q at such a switch. The base cases of $M = 0$ or 1 are prohibited by (iv) of Definition 2.6 and the normality condition, respectively. We achieve the inductive step by contradiction. By symmetry we may assume that P lies above Q and such a switch occurs along P .

Case 1 The corner along P at the switch points toward Q . Then by the normality condition, the other path at the switch must have its companion path between P and Q . The inductive hypothesis then provides a contradiction.

Case 2 The corner along P at the switch points away from Q . Then heading in the appropriate direction, (left or right, depending on the slope of Q at the switch), P and Q will be on course to intersect within Y . This is prohibited, so there must be another switch along either P or Q . This next switch will be of the type covered by Case 1, and, prior to this switch, the number of strands between P and Q can only decrease since they are angled toward one another in this direction. Thus, the inductive hypothesis may be applied. \square

Lemma 3.4 (Cusp Lemma) *The thin part of ρ does not change when passing a cusp. Furthermore, at those crossings of $S(K, L)$ corresponding to a particular cusp of K the crossing between the i^{th} strand and the j^{th} strand is a switch if and only if $\rho_T(i) = j$ before and after the cusp.*

Proof By symmetry, we may consider the case of a left cusp of K . Let C_1, \dots, C_n denote the corresponding left cusps of $S(K, L)$ numbered from top to bottom. The ruling ρ provides a pair of paths P_i and Q_i emanating from each of the C_i where we assume P_i to have larger z -coordinate than Q_i . Although it need not literally be the case, we will refer to those strands of $S(K, L)$ that correspond to the upper (resp. lower) branches of cusps as positive (resp. negative) sloped. (Compare Figure 10.)

Claim *Among those crossings near the cusp of K , none of the P_i (resp. Q_i) can have a switch that causes the slope of P_i (resp. Q_i) to increase (resp. decrease).*

This is proved by induction on M , the number of strands lying between P_i and Q_i at the offending switch S . As in the proof of the Crossing Lemma, the base cases $M = 0$ or 1 would violate the definition of a normal ruling.

We prove the inductive step by contradiction. By symmetry we may assume S is a switch along P_i . The normality condition forces that there is some $j \neq i$ such that (i) the lower half of S is an arc along the path P_j , and (ii) Q_j lies above Q_i at $x(S)$.

Case 1 Q_j is positively sloped at $x(S)$. Then immediately to the right of S , P_j is negatively sloped and Q_j is positively sloped. To the right of S , one of these paths must switch or they will intersect before leaving the collection of crossings near the cusps C_1, \dots, C_n . However, as we move to the right of S the number of strands between P_j and Q_j can only decrease, therefore this contradicts the inductive hypothesis.

Case 2 Q_j is negatively sloped at $x(S)$. This time follow P_j and Q_j to the left of S . Since they lie between P_i and Q_i at $x(S)$, and because of the way they are sloped, one of P_j and Q_j must switch in order for them to meet at the cusp C_j . However, the number of strands between P_j and Q_j decreases when moving to the left, and thus the inductive hypothesis applies to provide a contradiction at the first such switch.

Finally, we deduce the result from the claim. Near the cusp, each pair of paths P_i and Q_i can have at most one switch combined. Indeed, the claim forbids both P_i and Q_i from individually having more than one switch, and if both P_i and Q_i have a single switch then they will intersect before leaving the collection of crossings near the cusp. Now, in the case that there is no switch along P_i or Q_i , the i^{th} strand will be a fixed point of ρ_T along either branch of the cusp. Any switches that do occur must have upper strand Q_i and lower strand P_j with $i < j$, and such a switch results in $\rho_T(i) = j$ before and after the crossing. Conversely, if $\rho_T(i) = j$ before or after the crossing, then a switch of this type is necessary in order for the corresponding thin strands to meet at a common cusp. \square

3.1 Reduced normal rulings

Definition 3.5 For a non-empty pattern $L \subset J^1(S^1)$, we say that a normal ruling $\rho \in \mathcal{R}^d(S(K, L))$ is *reduced* if

- (i) the front projection of L intersects $x = 0$ and
- (ii) there are no switches at the crossings of $S(K, L)$ that arise from left cusps of K .

Denote by $\widetilde{\mathcal{R}}^d(K, L)$ the set of reduced d -graded normal rulings of $S(K, L)$ and by $\widetilde{R}_{S(K,L)}^d(z)$ the corresponding *reduced ruling polynomial*. When $L = \emptyset$ we make the convention that the empty ruling is reduced so that $\widetilde{\mathcal{R}}^d(K, \emptyset)$ contains a single element.

Any normal ruling of $S(K, L)$ corresponds to a generalized normal ruling τ of L along with a reduced normal ruling of a certain satellite associated to τ . More precisely, we have the following.

Theorem 3.6 Assume that $(K, *) \subset \mathbb{R}^3$ and $L \subset J^1(S^1)$ have front diagrams in general position. There is a bijection

$$\Phi: T \xrightarrow{\cong} \mathcal{R}^1(S(K, L)),$$

where T is the set of ordered pairs (τ, σ) satisfying $\tau \in \mathcal{GR}^1(L)$ and $\sigma \in \widetilde{\mathcal{R}}^1(K, L^\tau)$. Here, $L^\tau \subset J^1(S^1)$ denotes the link whose front diagram corresponds to the portion of L that is fixed by τ . Furthermore, $j(\Phi(\tau, \sigma)) = j(\tau) + j(\sigma)$, and $\Phi(\tau, \sigma)$ is d -graded if and only if τ and σ are.

Proof Given a pair $(\tau, \sigma) \in T$, $\Phi(\tau, \sigma)$ is constructed as follows.

Step 1 Extend τ to a partial ruling $\tilde{\tau}$ of $S(K, L)$ with some fixed point strands: This is done by letting $\tilde{\tau} = \tau$ along the subset $L \subset S(K, L)$ and then extending so that away from cusps the thin part of $\tilde{\tau}$ agrees with the involution τ at the boundary of the front projection of L in $[0, 1] \times \mathbb{R}$. That is, except near cusps, the i^{th} and j^{th} parallel copies of a strand of K in $S(K, L)$ are paired by $\tilde{\tau}$ if and only if on the vertical line $x = 0$, $\tau(i) = j$. All other strands of $S(K, L)$ are fixed point strands for $\tilde{\tau}$. Finally, to piece $\tilde{\tau}$ together, for each such i and j we add one switch at every cusp. Assuming $i < j$, these switches occur where the i^{th} strand corresponding to the lower branch of the cusp passes over the j^{th} strand. See Figure 10.




Figure 10: The thin part of a ruling of $S(K, L)$ near cusps

Step 2 Extend $\tilde{\tau}$ to $\Phi(\tau, \sigma)$: The fixed point strands of $\tilde{\tau}$ (after smoothing near any relevant switches within L) form a front diagram that is combinatorially the same as $S(K, L^\tau)$. We extend by requiring that the restriction to $S(K, L^\tau)$ is the normal ruling σ . See Figure 11.

The switches of $\Phi(\tau, \sigma)$ can be divided into three disjoint types: (A) Switches of τ , (B) switches of σ and (C) switches near cusps added in Step 1.

The normality condition is easily verified for switches of type (B) and (C). For switches of type (A) we need to consider two subcases. If neither of the involved strands are fixed point strands of τ , then the normality condition follows since it holds in τ . If one of the switching strands, say P_0 , is a fixed point strand of τ , then the normality condition for the ruling $\Phi(\tau, \sigma)$ follows from the generalized normality condition for τ provided that we know the companion strand of P_0 in $\Phi(\tau, \sigma)$ lies outside of $L \subset S(K, L)$. This is true for the following reason. The fixed point subset, L^τ , cannot contain cusps, so if there is a pair of thin strands of σ within the subset $L^\tau \subset S(K, L^\tau)$, then we will continue to have a pair of thin strands immediately to the left of L . Following this strand of K to the left, the Crossing Lemma implies that we continue to have a pair of




Figure 11: The bijection from Theorem 3.6. In the pictured example, K is a left-handed trefoil (topologically the mirror of the trefoil pictured in Figure 4). The fixed point subset L^τ is the basic front A_2 .

thin strands until we reach a left cusp of K . Finally, the Cusp Lemma then contradicts the assumption that σ is a *reduced* ruling of $S(K, L^\tau)$.

Next, we verify that $j(\Phi(\tau, \sigma)) = j(\tau) + j(\sigma)$. The cusps of $S(K, L)$ not belonging to the subsets L and $S(K, L^\tau)$ are in two-to-one correspondence with switches of type (C). Recall that the negative term of j counts $\frac{1}{2}$ the total number of cusps. Thus, in the computation of $j(\Phi(\tau, \sigma))$, switches of type (C) precisely cancel these unclaimed cusps and we are left with $j(\tau) + j(\sigma)$. (Here, we used that L^τ does not contain cusps, so that none of the cusps of L are double counted in $S(K, L^\tau)$.)

Now we check that $\Phi(\tau, \sigma)$ is d -graded if τ and σ are. In the construction the only paired paths of $\Phi(\tau, \sigma)$ that do not belong to either L or $S(K, L^\tau)$ are the thin pairs arising from Step 1. Clearly, immediately next to L the requirement of Definition 2.7 continues to hold. Following along K , it is enough to verify that the condition continues to hold for thin pairs after passing a cusp. This is immediate. Near cusps, thin pairs belonging to the upper and lower branch of the cusp are in correspondence, and the Maslov potentials differ by ± 1 .

Finally, we show that Φ is onto. Let $\rho \in \mathcal{R}^1(S(K, L))$ be arbitrary. The restriction of ρ to $L \subset S(K, L)$ produces a generalized normal ruling by making the convention that strands of L mapped outside of L by ρ are fixed point strands of τ . The Cusp and Crossing Lemmas imply that the thin part of ρ is constant and that, precisely as in the construction from Step 1, the only switches outside of L involving thin strands are near cusps of K . Moreover, the complement of the thin strands of $S(K, L)$ will be precisely $S(K, L^\tau)$ and restricting ρ to this subset produces σ such that $\Phi(\tau, \sigma) = \rho$. \square

Corollary 3.7 *For any Legendrians $K \subset \mathbb{R}^3$ and $L \subset J^1(S^1)$ with \mathbb{Z}/d -valued Maslov potential, we have*

$$R_{S(K,L)}^d(z) \geq R_L^d(z),$$

where \geq refers to inequality between all coefficients of corresponding powers of z .

Proof This follows from Theorem 3.6 and the injection $\mathcal{R}^d(L) \hookrightarrow T$ sending $\tau \mapsto (\tau, \sigma_0)$, where σ_0 is the unique element of $\widetilde{\mathcal{R}}^d(K, \emptyset)$. \square

3.2 Normal rulings of satellites and the Thurston–Bennequin number

Corollary 3.7 suggests the following definition.

Definition 3.8 Let $K \subset \mathbb{R}^3$ be a Legendrian knot and $L \subset J^1(S^1)$ a Legendrian link, each equipped with a \mathbb{Z}/d -valued Maslov potential. Then K is L -compatible if $R_{S(K,L)}^d(z) \neq R_L^d(z)$.

In this subsection, we discuss a correlation between L -compatibility and maximal Thurston–Bennequin number; later we show that L -compatibility is related to the existence of representations of the Chekanov–Eliashberg DGA.

Recall that a Legendrian knot $K \subset \mathbb{R}^3$ has positive and negative stabilizations $S_+(K)$ and $S_-(K)$ obtained by inserting a pair of consecutive cusps (a zigzag) into a strand of the front projection of K . The stabilization is positive (resp. negative) if the new cusps have the orientation of K running downward (resp. upward) along the cusp. The

Legendrian isotopy type of $S_{\pm}(K)$ depends only on the Legendrian isotopy type of K , and any Legendrian $K' \subset \mathbb{R}^3$ that is isotopic to $S_{\pm}(K)$ for some $K \subset \mathbb{R}^3$ is said to be *stabilized*.

Remark 3.9 (i) Stabilization may be viewed as a special case of the Legendrian satellite construction since $S_{\pm}(K) = S(K, L_{\pm})$ where $L_{\pm} \subset J^1(S^1)$ are pictured in Figure 12.

(ii) Note that $tb(S_{\pm}(K)) = tb(K) - 1$.




Figure 12: Front projections of L_+ and L_-

Theorem 3.10 If K is stabilized, then for any nonempty $L \subset J^1(S^1)$ with any choice of Maslov potential, $R_{S(K,L)}^d(z) = R_L^d(z)$, ie, K is not L -compatible.

Proof We may assume that $(K, *)$ and L are in general position, that the front diagram of K contains a zigzag, and that the base point $*$ does not lie on the strand, S , that connects the two cusps of the zigzag. On the front diagram of the satellite $S(K, L)$, let C_1, \dots, C_n and D_1, \dots, D_n denote the left and right cusps corresponding to the zigzag on K .

Let ρ be a normal ruling of $S(K, L)$. We will show that along S the thin part of ρ , ρ_T , does not have fixed points. Consider the ruling paths P_i and Q_i originating at a cusp C_i . If P_i switches before leaving the crossing near the C_i , then it follows from the Cusp Lemma that $\rho_T(i) \neq i$. If not, then P_i must end at one of the D_j . Then Q_i would need to end at D_j as well, but this ensures that Q_i must switch at one of the crossings near the C_i . Again, the Cusp Lemma shows that $\rho_T(i) \neq i$.

Now, from the Cusp and Crossing Lemmas we see that the thin part of ρ must be fixed point free everywhere. In particular, ρ restricts to a normal ruling τ (without fixed points) on $L \subset J^1(S^1)$. It follows that $\rho = \Phi(\tau, \sigma_0)$. Thus, the injection of $\mathcal{R}^d(L) \hookrightarrow T$ from Corollary 3.7 composed with Φ is onto $\mathcal{R}^d(K, L)$, and the result follows. \square

Corollary 3.11 If there exists a pattern $L \subset J^1(S^1)$ such that K is L -compatible, then $tb(K)$ is maximal within the smooth knot type of K .

Proof If $tb(K)$ is non-maximal, then there exists a stabilized knot K' with the same smooth type as K and $tb(K) = tb(K')$. Then

$$R_{S(K,L)}^1(z) = R_{S(K',L)}^1(z) = R_L^1(z),$$

where the first equality is a combination of Remark 2.3 with Theorem 2.15 and the second is Theorem 3.10.

If K is L -compatible where L has a d -graded Maslov potential, then it remains L -compatible in the ungraded setting. This contradicts $R_{S(K,L)}^1 = R_L^1$. \square

3.3 Reduced ruling polynomials

Theorem 3.6 shows that

$$(1) \quad R_{S(K,L)}^d(z) = \sum_{\tau \in \mathcal{GR}^d(L)} z^{j(\tau)} \tilde{R}_{S(K,L^\tau)}^d(z).$$

We will use this relation to deduce properties of reduced ruling polynomials from corresponding results about standard ruling polynomials.

Theorem 3.12 *Let $K \subset \mathbb{R}^3$ be a Legendrian knot with $d \mid 2r(K)$. For any fixed d -graded $L \subset J^1(S^1)$ with generic front projection, $\tilde{R}_{S(K,L)}^d$ is a Legendrian isotopy invariant of K .*

Proof First we establish the result for those front projections L that do not have cusps. In this case, the summation on the right side of (1) contains $\tilde{R}_{S(K,L)}^d$ (when τ is the generalized ruling that fixes every strand of A_Λ). The remaining terms on the right hand side are a $\mathbb{Z}[z^{\pm 1}]$ -linear combination of reduced ruling polynomials of the form $\tilde{R}_{S(K,L')}^d$, where L' is again a front projection without cusps and fewer strands than L . The front projections L' that appear in the summation as well as the coefficients depend only on L . Since the left hand side of (1) is a Legendrian isotopy invariant of K (Theorem 2.13), the result follows from inducting on the number of strands of L .

The general case when L has cusps follows from this special case. The reduced ruling polynomial of L arises from restricting the sum on the right hand side of (1) to those generalized rulings τ that are fixed point free on the line $x = 0$. (This is due to the construction of the bijection in Theorem 3.6.) The remaining terms form a linear combination of reduced ruling polynomials of satellites of K where the patterns do not have cusps. (For any generalized ruling, the fixed point set L^τ cannot have cusps.) Moreover, the particular patterns and coefficients appearing in this linear combination only depend on L . These reduced ruling polynomials are all known to be Legendrian isotopy invariants of K , as is the left hand side of (1), so the result follows. \square

Remark 3.13 The reduced ruling polynomials $\tilde{R}_{S(K,L)}^d$ are not invariant under Legendrian isotopy of L . For instance, an isotopy of L that pushes a left cusp from the right side of $x = 0$ to the left side of $x = 1$ causes the reduced ruling polynomial to vanish. Currently, we do not know if reduced ruling polynomials may be reformulated to obtain this property, and we leave the matter as an open question.

Theorem 3.14 For $d = 1$ or 2 , $\tilde{R}_{S(K,L)}^d$ depends only on L , $tb(K)$ and the underlying smooth knot type of K .

Proof The proof follows a similar scheme to that used for Theorem 3.12. In this case, Theorem 2.15 together with Remark 2.3 are used to establish the inductive step. \square

Example 3.15 As an example, we consider the case of 1–graded reduced rulings when the pattern is a product of basic fronts, A_Λ , where $\Lambda = (\lambda, \mathbf{m})$. Notation is as in Section 2.3. In the case of 1–graded rulings, \mathbf{m} is uninteresting, and it is not too hard to use (1) to give a rather explicit relation between the ruling polynomials of satellites $S(K, A_\lambda)$ and their reduced analogs.

Theorem 3.16 Given a partition $\lambda = (\lambda_1, \dots, \lambda_\ell)$, let M_λ^{sym} denote the set of all symmetric $\ell \times \ell$ matrices with nonnegative integer entries with row sums and column sums equal to λ . Then

$$R_{S(K,A_\lambda)}^1(z) = z^{\ell(\ell-1)} \sum_{(b_{ij}) \in M_\lambda^{\text{sym}}} \left(\prod_i z^{-\delta_i} \right) \left(\prod_{i < j} \langle b_{ij} \rangle \right) \tilde{R}_{S(K,A_{(b_{11}, \dots, b_{\ell\ell})})}^1(z),$$

where δ_i is the Kronecker delta $\delta_{b_{ii},0}$, and $\langle m \rangle$ denotes the ruling polynomial $R_{A_m A_m}^1(z)$ if $m \neq 0$ and z^{-2} if $m = 0$.

The proof is similar to that of [8, Theorem 3.4] and is omitted here. (Actually, we will not need such a precise formula.) Inductively, Theorem 3.16 can be used to give a formula for $\tilde{R}_{S(K,A_\lambda)}^1(z)$ in terms of ordinary ruling polynomials.

The following result allows us to reduce questions of L –compatibility for general L to the special case of the products A_Λ .

Theorem 3.17 Let K be a Legendrian knot in \mathbb{R}^3 and d a divisor of $2r(K)$. Then the following are equivalent:

- (i) There exists a nonempty d –graded product of basic fronts, A_Λ , with $\Lambda = (\lambda, \mathbf{m})$, such that $\tilde{R}_{S(K,A_\Lambda)}^d \neq 0$.

- (ii) There exists a nonempty d -graded product of basic fronts, A_Λ , with $\Lambda = (\lambda, \mathbf{m})$, such that K is A_Λ -compatible.
- (iii) There exists $L \subset J^1(S^1)$ with a d -graded Maslov potential, such that K is L -compatible.

Furthermore, any of these conditions imply that K maximizes tb .

Proof The remark about K maximizing tb is Corollary 3.11.

The equivalence of (i) and (ii) follows from Equation (1) with $L = A_\Lambda$. Indeed, all of the summands on the right hand side have nonnegative coefficients, and the terms with $L^\tau = \emptyset$ produce exactly the polynomial $R_{A_\Lambda}^d(z)$. Since L^τ also has the form $A_{\Lambda'}$ for some Λ' , we see that these are the only non-zero terms if $\tilde{R}_{S(K, A_\Lambda)}^d = 0$ for all $\Lambda \neq \emptyset$. Hence, (ii) implies (i). On the other hand, $\tilde{R}_{S(K, A_\Lambda)}^d$ also appears in the sum when τ is the generalized ruling where every strand is a fixed point. It follows that if $\tilde{R}_{S(K, A_\Lambda)}^d \neq 0$, then $R_{S(K, A_\Lambda)}^d(z) > R_{A_\Lambda}^d(z)$. Thus, (i) implies (ii).

That (ii) implies (iii) is immediate. For the converse, we need to recall some results about ruling polynomials.

The ruling polynomial $R_L^d(z)$ satisfies skein relations as in [16, Lemma 6.8] with the following modification from the 2-graded case: The coefficient δ_1 (resp. δ_2) is 1 if the strands that cross in the first (resp. second) term have equal Maslov potential mod d and is 0 otherwise. (The proof is virtually identical.) Moreover, the proof of Lemma 6.10 in the same article provides an algorithm for evaluating the ruling polynomial of an arbitrary Legendrian $L \subset J^1(S^1)$ as a linear combination of ruling polynomials of basic fronts using these skein relations. The algorithm, addressed to the 2-graded case in [16], applies equally well in the d -graded case. In addition, the ruling polynomial of a satellite, $R_{S(K, L)}^d(z)$, satisfies the same skein relations in the factor L . It follows that we can find coefficients $c_\Lambda(z) \in \mathbb{Z}[z^{\pm 1}]$ such that

$$(2) \quad R_L^d(z) = \sum c_\Lambda(z) R_{A_\Lambda}^d(z) \quad \text{and} \quad R_{S(K; L)}^d(z) = \sum c_\Lambda(z) R_{S(K; A_\Lambda)}^d(z).$$

We now prove (the contrapositive of) (iii) implies (ii). Assume that for all Λ ,

$$R_{S(K; A_\Lambda)}^d(z) = R_{A_\Lambda}^d(z).$$

Combining this with (2) shows that $R_L^d(z) = R_{S(K; L)}^d(z)$. □

4 Finite-dimensional representations of $\mathcal{A}(K, *)$

In this section we give necessary and sufficient conditions for the existence of finite-dimensional representations of $\mathcal{A}(K, *)$ in terms of normal rulings of Legendrian satellites of K .

4.1 The path matrix

We begin with some linear algebra. Let $L \subset J^1(S^1)$ be a d -graded Legendrian link without cusps that intersects the line $x = 0$ in n points. In this case, the DGA of the satellite $S(K, L)$ can be described in terms of the DGA of $(K, *)$ and a matrix P_L , known as the path matrix of L , which was introduced in [7]. The definition and properties of P_L discussed here are all contained in [7].

Label the crossings of L from left to right as p_1, p_2, \dots, p_r . As usual each crossing is assigned a degree in \mathbb{Z}/d as the difference of the value of the Maslov potential on the over- and understrands of the crossing, and the p_i are viewed as non-commuting variables.

Definition 4.1 View the front projection of L as a subset of $[0, 1] \times \mathbb{R}$ without identifying the boundary lines, and consider paths within the front projection of L that begin on the i^{th} strand (counting from top to bottom) at $x = 0$ and end on the j^{th} strand (from top to bottom) at $x = 1$. At crossings we allow paths to either go straight through the crossing or turn a corner around the upper quadrant of the crossing. To each such path we assign a word in the p_i that is the product of crossings corresponding to corners of the path ordered from left to right (if there are no corners, the word is 1). The *path matrix* P_L is the matrix whose ij -entry is the sum of words associated with all such paths.

The path matrix is invertible as a matrix with entries in the non-commutative $\mathbb{Z}/2$ -algebra generated by the p_i . To see this note that

$$P_L = C_1 C_2 \cdots C_r,$$

where C_i , $1 \leq i \leq r$, is the invertible matrix equal to the identity matrix except with the 2×2 block

$$\begin{bmatrix} p_i & 1 \\ 1 & 0 \end{bmatrix}$$

placed on the diagonal in rows k and $k + 1$, where $k, k + 1$ are the labels of the strands involved in crossing p_i . Using this perspective, it is also not hard to see that the entries of $(P_L)^{-1}$ correspond to paths from right to left in L that are allowed to turn corners around the lower quadrant of a crossing.

Example 4.2 For the basic front A_n , label the crossings in A_n from left to right as p_1, \dots, p_{n-1} ; then the path matrix P_{A_n} of A_n satisfies:

$$P_{A_n} = \begin{bmatrix} p_1 & p_2 & \cdots & p_{n-1} & 1 \\ 1 & 0 & \cdots & 0 & 0 \\ 0 & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & 1 & 0 \end{bmatrix} \quad \text{and} \quad P_{A_n}^{-1} = \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \\ 1 & p_1 & p_2 & \cdots & p_{n-1} \end{bmatrix}$$

Note that $(P_{A_n}^{-1})^T$ is a matrix in rational canonical form, the so-called companion matrix to the polynomial $\lambda^n + p_{n-1}\lambda^{n-1} + \dots + p_1\lambda + 1 \in (\mathbb{Z}/2)[\lambda]$.

More generally, if $\Lambda = (\lambda, \mathbf{m})$ and $\lambda = (\lambda_1, \dots, \lambda_\ell)$, then $(P_{A_\Lambda}^{-1})^T$ is also a matrix in rational canonical form: it is in block-diagonal form with blocks $(P_{A_{\lambda_1}}^{-1})^T, \dots, (P_{A_{\lambda_\ell}}^{-1})^T$. For future use, we record the following:

Lemma 4.3 Any invertible matrix $M \in GL_n(\mathbb{Z}/2)$ is conjugate to a matrix of the form P_{A_Λ} for some Λ .

Proof Any matrix at all, in particular $(M^{-1})^T$, is conjugate to a block diagonal matrix with blocks of the form $(P_{A_n}^{-1})^T$ except that in some blocks the 1 in the upper right corner may be replaced by 0. (This is the standard rational canonical form found in most introductory algebra texts.) Since $(M^{-1})^T$ is invertible all of these entries must equal 1, so M has the desired form. \square

Example 4.4 Let Δ_n denote a positive half twist of n strands. The path matrix of Δ_n is skew-upper triangular (ie, all entries below the northeast-southwest ‘‘antidiagonal’’ are 0), with 1’s on the antidiagonal and variables s_{ij} in the entries with $i + j \leq n$. See Figure 13.




Figure 13: A positive half twist with $n = 5$ and its path matrix

Given $\mathbf{n}: \mathbb{Z}/d \rightarrow \mathbb{Z}_{\geq 0}$ we let $tw_{\mathbf{n}}$ denote a positive full twist of $n = \sum_{k \in \mathbb{Z}/d} \mathbf{n}(k)$ strands with d -graded Maslov potential μ as follows: The first $\mathbf{n}(0)$ strands have

$\mu = 0$, the next $\mathbf{n}(1)$ strands have $\mu = 1$, and continue in this manner until the last $\mathbf{n}(d - 1)$ strands have $\mu = d - 1$. The full twist is the concatenation of two half twists, $tw_{\mathbf{n}} = \Delta_{\mathbf{n}} * \Delta_{\mathbf{n}}$, so it follows that the path matrix of $tw_{\mathbf{n}}$ is the product of two skew-upper triangular matrices, $S_1 S_2$. Note that in S_1 (resp. S_2) the crossings with degree $0 \pmod d$ are all on the blocks of sizes $\mathbf{n}(0) \times \mathbf{n}(0), \dots, \mathbf{n}(d - 1) \times \mathbf{n}(d - 1)$ running along the antidiagonal and ordered from upper right to lower left (resp. lower left to upper right).

We conclude this discussion by proving two lemmas. The first provides a standard form result related to the path matrices $P_{tw_{\mathbf{n}}}$, and the second involves normal rulings of satellites $S(K, tw_{\mathbf{n}})$.

Lemma 4.5 Any matrix $M \in GL_n(\mathbb{Z}/2)$ is conjugate to a matrix of the form

$$S_1 S_2 U,$$

where U is upper triangular and S_1, S_2 are skew-upper triangular.

Proof We first prove the lemma when M has the form $P_{A_n}^{-1}$ in the notation of Example 4.2. If we define S_1, S_2, U by

$$(S_1)_{ij} = \begin{cases} 1 & i + j = n \text{ or } n + 1, \\ 0 & \text{otherwise,} \end{cases} \quad (S_2)_{ij} = \begin{cases} 1 & i + j \leq n + 1, \\ 0 & \text{otherwise,} \end{cases}$$

$$U_{ij} = \begin{cases} 1 & i = j, \\ 1 + p_{j-1} & i = 1 \text{ and } j > 1, \\ 0 & \text{otherwise,} \end{cases}$$

then it is easy to check that $S_1 S_2 U = P_{A_n}^{-1}$.

For general M , up to conjugation, we may assume that M^T is in rational canonical form, so that M is conjugate to the block-diagonal form with blocks

$$P_{A_{\lambda_1}}^{-1}, \dots, P_{A_{\lambda_\ell}}^{-1}.$$

Apply the lemma to each of these blocks to obtain the desired S_1, S_2, U . (Note that S_1, S_2 are also block-diagonal, with blocks corresponding to blocks of M running along the antidiagonal from top right to bottom left in S_1 and from bottom left to top right in S_2 .) □

Lemma 4.6 Every normal ruling of $S(K, \Delta_n)$ or $S(K, tw_{\mathbf{n}})$ is reduced.

Proof We treat the case of a half twist; the proof applies equally well to a full twist.

Following the labeling scheme for crossings indicated in Figure 13, we modify the front projection of Δ_n by a planar isotopy so that crossings appear from left to right in the order $s_{n-1,1}, s_{n-2,1}, s_{n-2,2}, \dots, s_{1,1}, \dots, s_{1,n-1}$. That is, working from left to right the $(n-1)$ -strand crosses over the n -strand, then the $(n-2)$ -strand crosses over the two strands below it. This continues, so that each strand takes its turn crossing over all of the strands below.

Suppose that we are given a non-reduced normal ruling of $S(K, \Delta_n)$. From the Cusp and Crossing Lemmas, there will be at least one pair of thin strands entering the left side of $\Delta_n \subset S(K, \Delta_n)$. To avoid intersecting each other, this pair of strands must be involved in at least one switch within Δ_n since every pair of strands crosses in Δ_n . Thus we can find a switch, s , within Δ_n with the property that at least one of the companion strands also lies within Δ_n , and we can assume that the number of strands lying between the switching strand and its companion strand near the x -coordinate of the switch is minimized.

Case 1 This switching strand lies above its companion at s . Then the strand must constitute the lower half of the switch. (Suppose not; then, contrary to the minimality assumption, the normality condition would show that the companion strand to the lower half of the switch lies between the upper switching strand and its companion.) According to the way we have arranged Δ_n , heading to the right the switching strand will cross over its companion strand unless one of the two strands switches. Again, the normality condition would force such a switch to contradict the minimality hypothesis.

Case 2 This switching strand lies below its companion at s . The argument is symmetric. This switching strand must be the upper half of the switch. Heading left, it will intersect its companion strand unless a switch occurs. Such a switch would contradict the minimality assumption. \square

4.2 Necessary and sufficient conditions for the existence of representations

We are now in a position to state precisely our main results relating finite-dimensional representations and satellite rulings.

Given a d -graded Legendrian link $L \subset J^1(S^1)$ we define a (\mathbb{Z}/d) -graded $(\mathbb{Z}/2)$ -vector space V_L . Suppose the front projection of L intersects the vertical line $x = 0$ at points with Maslov potential $\eta_1, \dots, \eta_n \in \mathbb{Z}/d$ from top to bottom. Let V_L be the vector space with basis e_1, \dots, e_n with basis vectors assigned the grading $|e_i| = \eta_i$

for $i = 1, \dots, n$. The chosen basis e_1, \dots, e_n provides an isomorphism $\text{End}(V_L) \cong \text{Mat}_{n \times n}(\mathbb{Z}/2)$, and in the following we make use of this identification to view elements of $\text{End}(V_L)$ as matrices. Note that V_L has graded dimension $\mathbf{n}: \mathbb{Z}/d \rightarrow \mathbb{Z}_{\geq 0}$ where $\mathbf{n}(k)$ is the number of strands of L with Maslov potential equal to k at $x = 0$.

Our main technical statement relates reduced normal rulings of $S(K, L)$ to representations of $(\mathcal{A}(K, *), \partial)$ with underlying vector space V_L .

Theorem 4.7 *Let $K \subset \mathbb{R}^3$ be a d -graded Legendrian knot and $L \subset J^1(S^1)$ a d -graded Legendrian link without cusps. Then $S(K, L)$ has a d -graded reduced normal ruling if and only if there exists f such that:*

- f is a d -graded representation from $(\mathcal{A}(K, *), \partial)$ to $(\text{End}(V_L), 0)$, ie f is a $(\mathbb{Z}/2)$ -algebra map $\mathcal{A}(K, *) \rightarrow \text{End}(V_L)$ such that f preserves the \mathbb{Z}/d -grading and $f \circ \partial = 0$,
- $f(t)$ is a matrix of the form $M_L U$, where U is an upper triangular $n \times n$ matrix, and M_L is the image of the path matrix P_L under an algebra map that sends each p_i to some element of $\mathbb{Z}/2$, with p_i sent to 0 unless $|p_i| \equiv 0 \pmod{d}$.

The proof of Theorem 4.7 is given in Section 4.4. First we derive some consequences that are some of the central results of this paper.

Theorem 4.8 *Let K be a Legendrian knot in \mathbb{R}^3 and d a divisor of $2r(K)$. Then for fixed, non-zero $\mathbf{n}: \mathbb{Z}/d \rightarrow \mathbb{Z}_{\geq 0}$, the following are equivalent:*

- (i) *The Chekanov–Eliashberg algebra $\mathcal{A}(K, *)$ has a d -graded representation of graded dimension \mathbf{n} .*
- (ii) *There exists $\Lambda = (\lambda, \mathbf{m})$ with $\mathbf{n}_\Lambda = \mathbf{n}$ such that $S(K, A_\Lambda)$ has a d -graded reduced normal ruling.*
- (iii) *The satellite of K with a full positive twist, $S(K, tw_{\mathbf{n}})$, has a d -graded normal ruling. (Note that this link is topologically the $(tb(K) + 1)$ -twisted n -copy of K .)*

Note that when \mathbf{n} is the map $\mathbf{n}(0) = 1, \mathbf{n}(k) = 0$ for $k \neq 0$, Theorem 4.8 reduces to Theorem 2.25, the correspondence between existence of d -graded augmentations and d -graded rulings.

Proof of Theorem 4.8 The forward direction of Theorem 4.7 in conjunction with Lemma 4.6 shows that either of (ii) or (iii) implies (i).

We now assume that $f: \mathcal{A}(K, *) \rightarrow \text{End}(V)$ is a d -graded representation with $\dim(V) = \mathbf{n}$, and prove that (ii) and (iii) hold. From the definitions, $V = \bigoplus_{k \in \mathbb{Z}/d} V_k$ with $\dim_{\mathbb{Z}/2} V_k = \mathbf{n}(k)$, and since $|t| = 0$, $f(t)$ has the form $\sum_{k \in \mathbb{Z}/d} f(t)_k$ with $f(t)_k \in \text{GL}(V_k)$.

For (ii), we choose bases for each V_k , so that the matrix of $f(t)_k$ has the form described in Lemma 4.3, and then concatenate these bases to produce a basis for V . Now, let $\lambda_1, \dots, \lambda_\ell$ denote the block sizes and $m_1, \dots, m_\ell \in \mathbb{Z}/d$ denote the grading of the corresponding components. Using notation as in the statement of Theorem 4.7, the choice of basis provides a grading preserving isomorphism $V \cong V_{A_\Lambda}$. With this identification, we obtain a representation f

$$f: \mathcal{A}(K, *) \rightarrow \text{End}(V_{A_\Lambda}),$$

and as discussed in Example 4.2 $f(t)$ has the desired form M_{A_Λ} so that we can apply Theorem 4.7 to produce a reduced ruling of $S(K, A_\Lambda)$.

Now we establish (iii). Applying Lemma 4.5, we can choose a basis for each V_k , $k = 0, \dots, d - 1$, such that the matrix of $f(t)_k$ has the form $S_1^k S_2^k U^k$ with S_i^k skew-upper triangular and U^k upper triangular. Concatenating these bases provides a grading preserving isomorphism $V \cong V_{\text{tw}_n}$. With respect to the distinguished basis, $f(t)$ has the form $S_1 S_2 U$ where S_1 (resp. S_2) is obtained from placing the blocks S_1^k (resp. S_2^k) along the antidiagonal from upper right to lower left (resp. lower left to upper right). Consulting Example 4.4 we see that this matrix is of the form M_{tw_n} so that an application of Theorem 4.7 completes the proof. \square

In conjunction with Theorem 3.14 and Corollary 3.11, Theorem 4.8 implies the following:

Theorem 4.9 *The existence of a 1 or 2-graded representation of $\mathcal{A}(K, *)$ of any given dimension depends only on the smooth knot type of K and $tb(K)$. If $\mathcal{A}(K, *)$ admits finite-dimensional representations, then K must have maximal Thurston–Bennequin number.*

Remark 4.10 Shonkwiler and Vela-Vick [18] gave examples of Legendrian knots in the topological knot types $m(10_{145})$ and $m(10_{161})$ with non-maximal Thurston–Bennequin and non-trivial Chekanov–Eliashberg algebras. Sivek [19] observed by a direct argument that these algebras do not admit finite-dimensional representations. This may alternatively be deduced from Theorem 4.9.

The following corollary of Theorem 3.17 and Theorem 4.8 addresses the problem of finding an arbitrary pattern L that is compatible with K .

Theorem 4.11 *The DGA $\mathcal{A}(K, *)$ has a d –graded finite-dimensional representation if and only if there exists a d –graded pattern $L \subset J^1(S^1)$ so that K is L –compatible.*

We conclude this subsection by discussing the relation between Theorem 4.9 and a conjecture from [10] about topological invariance of the abelianized characteristic algebra.

Definition 4.12 [10] Let K be a Legendrian knot with DGA (\mathcal{A}, ∂) . Define $I \subset \mathcal{A}$ to be the two-sided ideal generated by the collection $\{\partial(a_i)\}$ of differentials of generators of \mathcal{A} . The *characteristic algebra* of K is the algebra \mathcal{A}/I ; the *abelianized characteristic algebra* of K is the abelianization of \mathcal{A}/I .

It was observed in [10] that the abelianized characteristic algebra, viewed without grading, often seems to depend only on the smooth knot type of K and $tb(K)$, and it was conjectured that this is always the case (up to a natural equivalence relation; see [10]). We do not resolve this conjecture here, but address a related construction.

Definition 4.13 Let K be a Legendrian knot, and (\mathcal{A}, ∂) , I as above. Define $I' \subset \mathcal{A}$ to be the smallest two-sided ideal containing I such that whenever $x, y \in \mathcal{A}$ satisfy $1 - xy \in I'$, then $1 - yx \in I'$ as well. The *partially abelianized characteristic algebra* of K is defined to be \mathcal{A}/I' .

Note that we have a sequence of successive quotients: characteristic algebra, partially abelianized characteristic algebra, abelianized characteristic algebra. From computations, it appears that the (ungraded) partially abelianized characteristic algebra, like the abelianized version, depends only on smooth type and tb , and perhaps this is a more natural conjecture than the conjecture from [10].

Conjecture 4.14 *Up to equivalence, the ungraded partially abelianized characteristic algebra of a Legendrian knot K depends only on the smooth type and tb of K .*

Theorem 4.9 gives some corroborating evidence:

Corollary 4.15 *Let K_1, K_2 be Legendrian knots with the same smooth type and tb . Then for any n , the ungraded partially abelianized characteristic algebras of K_1 and K_2 either both have, or both do not have, an n –dimensional representation over $\mathbb{Z}/2$.*

It is conceivable that this corollary could be strengthened to give some sort of correspondence between n –dimensional representations of K_1 and K_2 , and that the collection of finite-dimensional representations of a $\mathbb{Z}/2$ –algebra of the type considered here is

enough to determine the $\mathbb{Z}/2$ -algebra, in which case Conjecture 4.14 would follow. However, we do not pursue this direction further in this paper.

The rest of Section 4 is devoted to the proof of Theorem 4.7. We first address some preliminary issues before presenting the proof in Section 4.4.

4.3 The DGA of the satellite $S(K, L)$

For the rest of this section, we let $K \subset \mathbb{R}^3$ and $L \subset J^1(S^1)$ be d -graded Legendrian links such that L has no cusps. We fix a base point $*$ on K that is assumed to be located on a strand of K that is oriented to the right. For the proof of Theorem 4.7, there is no loss of generality from this assumption in view of Theorems 3.12 and 2.22. Furthermore, we may assume that the front diagram of K is in plat position. This can always be achieved by a Legendrian isotopy that will not affect the existence of either reduced rulings (Theorem 3.12) or the type of representation in question. (After an isotopy, $A(K, *)$ changes only by a stable tame isomorphism that maps t to t .)

When K is in plat position it follows that $S(K, L)$ will be plat as well. We now describe the Chekanov–Eliashberg algebra associated with the resolution of this front diagram. We maintain here our convention of, away from L , labeling the n strands running parallel to K from 1 to n as one moves from top to bottom.




Figure 14: Generators b_{ij}^m , q_{ij}^m or c_{ij}^m of $\mathcal{A}(S(K, L))$ corresponding to a left cusp, crossing, or right cusp m of K ; the m superscripts are suppressed

4.3.1 Generators of $\mathcal{A}(S(K, L))$ We first label the crossings and right cusps of $S(K, L)$; see Figure 14 for an illustration. Enumerate the left cusps of K as b_1, \dots, b_{M_1} . Each of these produces a strictly upper triangular matrix worth of generators b_{ij}^m , $1 \leq i < j \leq n$, $m = 1, \dots, M_1$ where b_{ij}^m denotes the crossing of the i^{th} strand over the j^{th} strand at the location of the cusp b_m . Enumerate the

crossings of K as q_1, \dots, q_{M_2} . For $1 \leq m \leq M_2$, there are corresponding generators q_{ij}^m with $1 \leq i, j \leq n$ where the i^{th} strand crosses over the j^{th} strand at q_m . Finally, at each of the right cusps c_1, \dots, c_{M_3} there is a lower triangular matrix worth of generators (note the absence of the term *strictly*) c_{ij}^m with $1 \leq j \leq i \leq n$ where the i^{th} strand crosses over the j^{th} strand. Generators of the form c_{ii}^m correspond to right cusps of $S(K, L)$.

Finally, there are generators corresponding to the crossings of L , which, following our convention for the path matrix, we enumerate as p_1, p_2, \dots, p_r from left to right.

4.3.2 Grading For $j = 1, \dots, n$, let $\eta_j \in \mathbb{Z}/d$ denote the value of the Maslov potential for L on the j^{th} strand at $x = 0$. The generators of $\mathcal{A}(S(K; A_\Lambda))$ have degrees related to the degrees of generators of $\mathcal{A}(K, *)$ as follows:

$$|b_{ij}^m| = \eta_i - \eta_j - 1, \quad |c_{ij}^m| = \eta_i - \eta_j + |c_m|, \quad |q_{ij}^m| = \eta_i - \eta_j + |q_m|.$$

The crossings p_i arising from L have their degrees determined by the Maslov potential of L as discussed above Definition 4.1.

4.3.3 Differential After a short preparation we will describe the differential for $\mathcal{A}(S(K, L))$. We collect generators corresponding to the left cusps, crossings and right cusps of K into matrices B_m, Q_m and C_m , respectively. The matrices C_m (resp. B_m) are lower triangular (resp. strictly upper triangular). Let

$$\Phi_L: \mathcal{A}(K, *) \rightarrow \text{Mat}_{n \times n}(\mathcal{A}(S(K, L)))$$

denote the algebra homomorphism that takes generators q_m and c_m to the corresponding matrices Q_m and C_m and takes t to the path matrix P_L .

In our notation we will use ∂ to denote the differential of $\mathcal{A}(K, *)$, and D for the differential in $\mathcal{A}(S(K, L))$. Moreover, let

$$\bar{D}: \text{Mat}_{n \times n}(\mathcal{A}(S(K, L))) \rightarrow \text{Mat}_{n \times n}(\mathcal{A}(S(K, L)))$$

denote the map resulting from applying D entry by entry.

Theorem 4.16 *The differential $D: \mathcal{A}(S(K, L)) \rightarrow \mathcal{A}(S(K, L))$ satisfies the following matrix formulas:*

- (3)
$$\bar{D}B_m = (B_m)^2,$$
- (4)
$$\bar{D}Q_m = \Phi_L(\partial q_m) + O(B), \text{ and}$$
- (5)
$$\bar{D}C_m = \pi_{\text{low}} \circ \Phi_L(\partial c_m) + O(B).$$

In formulas (4) and (5), $O(B)$ denotes a matrix whose entries belong to the two-sided ideal generated by the b_{ij}^m . In (5), π_{low} denotes the projection which replaces all of the entries above the main diagonal by 0.

In addition, for any crossing p_i arising from L , Dp_i belongs to the two-sided ideal generated by the b_{ij}^m .

Proof The formula (3) is easily seen explicitly.

In establishing (4) and (5) we follow Mishachev [9] and divide the disks involved in computing the differential for $\mathcal{A}(S(K, L))$ into two disjoint sets: *thin disks* and *thick disks*. By definition, thin disks are entirely contained in the neighborhood of the front diagram of K where the satellite construction is carried out, and all remaining disks are considered to be thick. (At each right cusp c_{ii}^m there is also a disk not visible on the front projection that contributes 1 to dc_{ii}^m . These disks arise from the twists added near right cusps when converting a front projection to a Lagrangian projection via the resolution procedure. We will consider such disks to be thick and refer to them later in the proof as *invisible disks*.)

As in [9], we consider a “stick together map” $s: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ that retracts the neighborhood of the front projection of K containing $S(K, L)$ onto the front projection of K itself. The image of a thick disk under s coincides with a disk $f \in \Delta(a; b_1, \dots, b_\ell)$ involved in the computation of the differential in $\mathcal{A}(K, *)$. Moreover, we have the following:

Claim If $f \in \Delta(a; b_1, \dots, b_\ell)$ has $w(f) = t^{\alpha_0} b_1 t^{\alpha_1} \dots b_\ell t^{\alpha_\ell} \in \mathcal{A}(K, *)$, then the collection of thick disks corresponding to f provides precisely the term

$$\Phi_L(t^{\alpha_0} b_1 t^{\alpha_1} \dots b_\ell t^{\alpha_\ell}) \quad (\text{resp. } \pi_{\text{low}} \circ \Phi_L(t^{\alpha_0} b_1 t^{\alpha_1} \dots b_\ell t^{\alpha_\ell}))$$

in $\bar{D}\Phi_L(a)$ when a is a crossing (resp. right cusp) of K .

From this claim it follows that restricting the sum defining D to thick disks produces precisely the first terms on the right hand side of (4) and (5).

To verify the claim, we need to consider “lifts” of

$$f: (D_\ell^2, \partial D_\ell^2) \rightarrow (\mathbb{R}^2, \pi_{xy}(K)) \quad \text{to} \quad \tilde{f}: (D_{\ell'}^2, \partial D_{\ell'}^2) \rightarrow (\mathbb{R}^2, \pi_{xy}(S(K, L)))$$

such that $s \circ \tilde{f} = f|_{D_\ell^2}$. Here, we use the notation from Section 2.6: r, s_1, \dots, s_ℓ are marked points along ∂D^2 , and $D_\ell^2 = D^2 \setminus \{r, s_1, \dots, s_\ell\}$. We need to allow the possibility that $\ell' \geq \ell$ since \tilde{f} may have more negative corners than f because of the additional possibility of negative corners at crossings of L . For concreteness, let us assume that $a = q_m$ and $b_1 = q_{m_1}, \dots, b_\ell = q_{m_\ell}$ are all crossings. Moreover, initially we treat the case in which $f(\partial D_\ell)$ does not contain the base point $*$ of K .

Such a lift \tilde{f} arises from an appropriate choice of corners in $S(K, L)$. If $\tilde{f}(r) = q_{ij}^m$, then since a neighborhood of r must map to a $+$ corner at q_m we see that the arc of ∂D_ℓ^2 extending counter-clockwise from r must initially map to the i^{th} copy of K in $S(K, L)$. (Recall that our subscripts i, j always indicate the i^{th} copy of K crossing over the j^{th} copy.) When we arrive at the crossings of $S(K, L)$ corresponding to q_{m_1} , the boundary of \tilde{f} remains on the i^{th} copy of K and must turn around a $-$ corner of a crossing of the form $q_{i,k_1}^{m_1}$. This puts the next arc of ∂D_ℓ^2 on the k_1 -copy of K . Similarly, the next corner of \tilde{f} must be a $-$ corner at a crossing of the form $q_{k_1,k_2}^{m_2}$. In total, \tilde{f} has negative corners at $q_{i,k_1}^{m_1}, q_{k_1,k_2}^{m_2}, \dots, q_{k_{\ell-1},j}^{m_\ell}$ where the choices of $1 \leq k_1, \dots, k_{\ell-1} \leq n$ can be arbitrary.

We see that lifts of f with initial positive corner at $q_{i,j}^m$ contribute the sum

$$\sum_{k_1, \dots, k_{\ell-1}} q_{i,k_1}^{m_1} q_{k_1,k_2}^{m_2} \cdots q_{k_{\ell-1},j}^{m_\ell}$$

to Dq_{ij}^m . This is precisely the ij entry of the product $Q_{m_1} \cdots Q_{m_\ell} = \Phi_L(q_{m_1} \cdots q_{m_\ell})$, so in this case the claim follows.

Now we consider how the collection of lifts \tilde{f} changes when $f(\partial D_\ell^2)$ is allowed to intersect the base point. Recall the notation $\gamma_0, \gamma_1, \dots, \gamma_\ell$ for the images under f of the circular arcs from r to s_1 , s_1 to s_2 , \dots , and s_ℓ to r , respectively. Suppose γ_i intersects $*$ positively. Since we have assumed $*$ is located on a strand of K that is oriented to the right, it follows that γ_i crosses $*$ from left to right. Moreover, since \tilde{f} is orientation preserving, a neighborhood of $\gamma_i^{-1}(*)$ in D_ℓ^2 maps to the region of \mathbb{R}^2 above $*$. The corresponding portion of \tilde{f} , $\tilde{\gamma}_i$, will travel from left to right through the subset $L \subset S(K, L)$, with possibly some convex negative corners at crossings p_i in L at which the image of \tilde{f} necessarily covers the upper quadrant of p_i . The ij -entry of the path matrix P_L records precisely the products of negative corners that can result if $\tilde{\gamma}_i$ enters the subset L along the i^{th} strand of K and departs along the j^{th} strand of K . This shows that, as desired, such an occurrence of t in $w(f)$ requires placing $P_L = \Phi_L(t)$ between $Q_{m_{i-1}}$ and Q_{m_i} when computing the terms in $\bar{D}Q_m$ that correspond to lifts of f .

A similar argument shows that appearances of t^{-1} in $w(f)$ translate to $\Phi_L(t^{-1})$ in the computation of $\bar{D}Q_m$. Indeed, if γ_i intersects $*$ negatively, then $\tilde{\gamma}_i$ travels from right to left along L with convex corners corresponding to the bottom quadrants of crossings of L . As discussed after Definition 4.1 the matrix whose ij -entry corresponds to such paths is precisely $P_L^{-1} = \Phi_L(t^{-1})$.

The above analysis of thick disks applies equally well to establish the term $\pi_{\text{low}} \circ \Phi_L(t^{\alpha_0} b_1 t^{\alpha_1} \cdots b_\ell t^{\alpha_\ell})$ in $\bar{D}C_m$. The projection π_{low} appears simply because there

are no crossings c_{ij}^m with $i < j$. Note also that the invisible disks at right cusps c_{ii}^m mentioned earlier in the proof contribute the identity matrix I to $\bar{D}C_m$ and this corresponds to $\pi_{\text{low}} \circ \Phi_L$ applied to the 1 in ∂c_m arising from the invisible disk in K at c_m .

The only claim made in the theorem about Dp_i and the remaining terms of $\bar{D}Q_m$ and $\bar{D}C_m$ is that they belong to the two-sided ideal generated by the b_{ij}^m . All of these items correspond to thin disks. Therefore, the proof is completed by showing that any thin disk must have a negative corner at some b_{ij}^m .

In general, any disk other than an invisible disk at a right cusp will attain its minimum x -coordinate at a left cusp. (As a consequence of the resolution construction negative corners can never provide a minimum x -coordinate and only the invisible disks have positive corners at the right quadrant of a crossing.) With our assumption that L has no cusps, the only left cusps of $S(K, L)$ are those corresponding to a left cusp b_m of K . They are accompanied on their right by a collection of crossings of the form b_{ij}^m .

Now, in order to reach its left cusp, the boundary of a thin disk will enter the collection of crossings b_{ij}^m along two parallel strands of $S(K, L)$ that correspond to the same strand of K . For these strands to meet up at a common left cusp of $S(K, L)$, one of them must have a negative corner at one of the b_{ij}^m . □

Remark 4.17 When L consists of n horizontal lines so that $S(K, L)$ is the n -copy, we can be more explicit about the terms $O(B)$. In this case, we have

$$\begin{aligned} \bar{D}Q_m &= B_{m'}Q_m + Q_m B_{m''} + \Phi_L(\partial q_m), \\ \bar{D}C_m &= B_{m'}C_m + C_m B_{m''} + \pi_{\text{low}} \circ \Phi_L(\partial c_m). \end{aligned}$$

Here, m' (resp. m'') are such that, beginning at the upper strand (resp. lower strand) of q_m or c_m and heading left, the first left cusp reached will be $b_{m'}$ (resp. $b_{m''}$).

4.4 Proof of Theorem 4.7

Throughout this subsection we fix notation as in the statement of Theorem 4.7: the d -graded vector space V_L has basis vectors e_1, \dots, e_n where the degree of e_i is given by the Maslov potential of L at $x = 0$ on the i^{th} strand. Also, M_L denotes a matrix obtained from the path matrix P_L by assigning values in $\mathbb{Z}/2$ to the variables p_i with the restriction that $p_i = 0$ unless $|p_i| = 0 \pmod d$.

Lemma 4.18 *The satellite $S(K, L)$ has a d -graded reduced ruling if and only if there exists a d -graded augmentation of $\mathcal{A}(S(K, L))$ such that $\varepsilon(B_m) = 0$ for all m .*

Proof This is an application of Theorem 2.26. □

Consider the Lagrangian projection of K where in addition to our original base point, $* = *_0$, we add base points $*_1, \dots, *_{M_3}$ (cyclically ordered) at all right cusps. We use $\widehat{\partial}$ to denote the differential in the corresponding multi-pointed DGA for K . (See Section 2.6.) The relation of $\widehat{\partial}$ with the differential ∂ of $\mathcal{A}(K, *)$ is simply

$$\widehat{\partial}q_m = \partial q_m \quad \text{and} \quad \widehat{\partial}c_m = \partial c_m + 1 + t_m,$$

provided that we identify t_0 and t to view $\mathcal{A}(K, *)$ as the subalgebra $\mathcal{A}(K, *_0) \subset \mathcal{A}(K, *_0, \dots, *_{M_3})$.

Lemma 4.19 *There exists a d -graded augmentation of $\mathcal{A}(S(K, L))$ such that $\varepsilon(B_m) = 0$ for all m if and only if there exists a d -graded representation,*

$$\widehat{f}: (\mathcal{A}(K, *_0, \dots, *_{M_3}), \widehat{\partial}) \rightarrow (\text{End}(V_L), 0),$$

such that $t_0 \mapsto M_L$ and $t_i \mapsto U_i$ with U_i upper triangular for all $i \geq 1$.

Proof (\Rightarrow) Assume $\varepsilon: \mathcal{A}(S(K, L)) \rightarrow \mathbb{Z}/2$ is such an augmentation. We let

$$\bar{\varepsilon}: \text{Mat}_{n \times n}(\mathcal{A}(S(K, L))) \rightarrow \text{Mat}_{n \times n}(\mathbb{Z}/2) \cong \text{End}(V_L)$$

denote the algebra homomorphism resulting from applying ε entry by entry.

We define the desired representation $\widehat{f}: \mathcal{A}(K, *_0, \dots, *_{M_3}) \rightarrow \text{End}(V_L)$ in two steps. First, we require that on the subalgebra $\mathcal{A}(K, *_0) \subset \mathcal{A}(K, *_0, \dots, *_{M_3})$ we have

$$(6) \quad \widehat{f}|_{\mathcal{A}(K, *_0)} = \bar{\varepsilon} \circ \Phi_L.$$

Explicitly, $\widehat{f}(q_m) = \bar{\varepsilon}(Q_m)$, $\widehat{f}(c_m) = \bar{\varepsilon}(C_m)$ and $\widehat{f}(t_0) = \bar{\varepsilon}(P_L)$. Note that since ε is d -graded $\widehat{f}(t_0)$ has the required form M_L . On the remaining generators t_m with $m = 1, \dots, M_3$ we define

$$\widehat{f}(t_m) = I + \bar{\varepsilon} \circ \Phi_L(\partial c_m).$$

The computation

$$0 = \bar{\varepsilon} \circ \bar{D}(C_m) = \bar{\varepsilon} \circ \pi_{\text{low}} \circ \Phi_L(\partial c_m) = \pi_{\text{low}}(\bar{\varepsilon} \circ \Phi_L(\partial c_m))$$

shows that $\widehat{f}(t_m)$ is non-singular and upper triangular as required.

To check that \widehat{f} is d -graded it suffices to show that $|\widehat{f}(s)| = |s|$ on any generator s . (The grading on $\text{End}(V_L)$ is defined as in Section 2.7.) First we treat the case when s has the form q_m . Note that the same argument applies when $s = c_m$.

On a basis vector e_j we have

$$\widehat{f}(q_m)e_j = \sum_i \varepsilon(q_{i,j}^m)e_i.$$

Since ε is d -graded, if $\varepsilon(q_{i,j}^m) \neq 0$ we have

$$0 = |q_{i,j}^m| = \eta_i - \eta_j + |q_m|,$$

which shows that

$$|e_j| - |e_i| = |q_m|.$$

It follows from the definition that $\widehat{f}(q_m)$ has degree $|q_m|$ in $\text{End}(V)$.

Next, we verify that $|\widehat{f}(t_0)| = 0$. Notice that if the i, j -entry of $\bar{\varepsilon}(P_L)$ is non-zero, then there is a path from $x = 0$ to $x = 1$ that starts on the i^{th} strand, ends on the j^{th} strand, and only turns along crossings which have equal Maslov potential. Clearly, the Maslov potential is constant along such a path, so $\eta_i = \eta_j$, which implies that $\widehat{f}(t_0)$ preserves the grading on V_L .

At this point we have shown that \widehat{f} is degree preserving on the sub-algebra $\mathcal{A}(K, *_0)$. Since ∂ lowers degree by 1 on $\mathcal{A}(K, *_0)$, it follows now that both the terms defining $\widehat{f}(t_m)$ have degree 0.

It remains to show that $\widehat{f} \circ \widehat{\partial} = 0$, and it suffices to verify this equality on generators with the case of t_i being immediate for $i = 0, \dots, M_3$. For a crossing q_m of K , $\widehat{\partial}(q_m) = \partial(q_m) \in \mathcal{A}(K, *_0)$, so using (6), (4) and the hypothesis that $\bar{\varepsilon}(B_m) = 0$, we can compute

$$\widehat{f} \circ \widehat{\partial}(q_m) = \bar{\varepsilon} \circ \Phi_L \circ \partial(q_m) = \bar{\varepsilon} \circ \bar{D}(Q_m) = 0.$$

For a cusp c_m of K , we have $\widehat{\partial}(c_m) = \partial(c_m) + 1 + t_m$, so we compute

$$\widehat{f} \circ \widehat{\partial}(c_m) = \bar{\varepsilon} \circ \Phi_L \circ \partial(c_m) + I + \widehat{f}(t_m) = 0.$$

(The final equality is just the definition of $\widehat{f}(t_m)$.)

(\Leftarrow) Suppose now that the representation \widehat{f} is given. We define $\varepsilon: \mathcal{A}(S(K, L)) \rightarrow \mathbb{Z}/2$ by requiring that the corresponding homomorphism of matrix algebras

$$\bar{\varepsilon}: \text{Mat}_{n \times n}(\mathcal{A}(S(K, L))) \rightarrow \text{Mat}_{n \times n}(\mathbb{Z}/2)$$

satisfies

$$(7) \quad \bar{\varepsilon}(B_m) = 0, \quad \bar{\varepsilon}(Q_m) = \widehat{f}(q_m), \quad \bar{\varepsilon}(C_m) = \pi_{\text{low}}(\widehat{f}(c_m)) \quad \text{and} \quad \bar{\varepsilon}(P_L) = M_L.$$

These formulas uniquely specify ε except possibly on the generators p_i , where the hypothesis on the matrix M_L allows us to fix values for $\varepsilon(p_i)$ that satisfy the matrix

equation and have $\varepsilon(p_i) = 0$ unless $|p_i| = 0$. That ε is d -graded on generators of the form q_{ij}^m and c_{ij}^m is verified in a similar manner to the corresponding portion of the proof of the forward implication.

To complete the proof, we show that $\varepsilon \circ D = 0$. This is immediate on generators of the form b_{ij}^m or p_i since in either case D applied to such a generator belongs to the two-sided ideal generated by the b_{ij}^m . For the remaining generators it suffices to verify that $\bar{\varepsilon} \circ \bar{D}(Q_m) = \bar{\varepsilon} \circ \bar{D}(C_m) = 0$.

Let \mathcal{A}' denote the subalgebra of $\mathcal{A}(K, *_0, \dots, *_{M_3})$ generated by crossings q_m and the original base point $*_0$. Note that since K is in plat position, we have $\partial q_m, \partial c_m \in \mathcal{A}'$ for any m . Furthermore, Equation (7) shows that $\hat{f}|_{\mathcal{A}'} = \bar{\varepsilon} \circ \Phi_L|_{\mathcal{A}'}$. Now, using these observations and (4) we compute

$$\bar{\varepsilon} \circ \bar{D}(Q_m) = \bar{\varepsilon} \circ \Phi_L(\partial q_m) = \hat{f}(\partial q_m) = \hat{f} \circ \hat{\partial} q_m = 0.$$

For a right cusp c_m , since $\partial c_m = \hat{\partial} c_m + 1 + t_m$ we have:

$$\begin{aligned} \bar{\varepsilon} \circ \bar{D}(C_m) &= \bar{\varepsilon} \circ \pi_{\text{low}} \circ \Phi_L(\partial c_m) = \pi_{\text{low}} \circ \hat{f}(\partial c_m) = \pi_{\text{low}}(\hat{f} \circ \hat{\partial} c_m + \hat{f}(1 + t_m)) \\ &= \pi_{\text{low}}(I + U_m) = 0. \quad \square \end{aligned}$$

Lemma 4.20 *The algebra $\mathcal{A}(K, *_0, \dots, *_{M_3})$ has a d -graded representation*

$$\hat{f}: (\mathcal{A}(K, *_0, \dots, *_{M_3}), \partial) \rightarrow (\text{End}(V_L), 0)$$

such that $\hat{f}(t_0) = M_L$ and $\hat{f}(t_i) = U_i$ with U_i upper triangular for all $i \geq 1$, if and only if there exists a d -graded representation $f: \mathcal{A}(K, *) \rightarrow (\text{End}(V_L), 0)$ such that $f(t) = M_L U$ with U upper triangular.

Proof Given such a representation \hat{f} of $\mathcal{A}(K, *_0, \dots, *_{M_3})$ we obtain the required representation of $\mathcal{A}(K, *)$ as the composition $f = \hat{f} \circ \phi$ where

$$\phi: \mathcal{A}(K, *) \rightarrow \mathcal{A}(K, *_0, \dots, *_{M_3})$$

is the homomorphism guaranteed by Theorem 2.22.

For the converse, assume that $f: \mathcal{A}(K, *) \rightarrow \text{End}(V_L)$ is a representation with $f(t) = M_L U$. Place base points a_0, \dots, a_{M_3} in a small neighborhood of the original base point $*$. Then there is a DGA homomorphism $g: \mathcal{A}(K, *) \rightarrow \mathcal{A}(K, a_0, \dots, a_{M_3})$ defined by fixing all generators other than t and setting $g(t) = t_0 \cdots t_{M_3}$. We define an algebra homomorphism $\hat{f}: \mathcal{A}(K, a_0, \dots, a_{M_3}) \rightarrow \text{End}(V_L)$ by

$$\hat{f}(t_0) = M_L, \quad \hat{f}(t_1) = U, \quad \hat{f}(t_i) = I \quad \text{for } i > 1,$$

and $\widehat{f}(s) = f(s)$ on the remaining generators. Note that $\widehat{f} \circ g = f$. We verify that $\widehat{f} \circ \widehat{d} = 0$ by checking the equality on generators with the case of the t_i being immediate since $\widehat{d}(t_i) = 0$. For any other generator, s , we can compute:

$$\widehat{f} \circ \widehat{\partial}(s) = \widehat{f} \circ \widehat{\partial} \circ g(s) = \widehat{f} \circ g \circ \partial(s) = f \circ \partial(s) = 0.$$

Finally, we obtain a representation of $\mathcal{A}(K, *_0, \dots, *_M)$ by composing \widehat{f} with the isomorphism guaranteed by Theorem 2.21. □

5 Ungraded 2–dimensional representations of $\mathcal{A}(K, *)$

As mentioned earlier, Theorem 4.8 generalizes the known result that a Legendrian knot $K \subset \mathbb{R}^3$ has a d –graded ruling if and only if its DGA (\mathcal{A}, ∂) has a d –graded augmentation. In this section, we analyze the next simplest case of Theorem 4.8, which provides a correspondence between rulings of doubles of K and 2–dimensional representations of $(\mathcal{A}(K, *), \partial)$. For the sake of simplicity, we will specialize to the case $d = 1$, in which all representations and all rulings are ungraded, and we will suppress any occurrences of d or of the grading in our notation.

In this case, Theorem 4.8 states that $(\mathcal{A}(K, *), \partial)$ has an ungraded 2–dimensional representation if and only if $\widetilde{R}_{S(K, A_\Lambda)}(z) \neq 0$ for at least one of the two partitions of 2, $\Lambda = (2)$ and $\Lambda = (1, 1)$. For $\Lambda = (1, 1)$, the only generalized rulings of $A_{(1,1)}$ are the trivial ruling with no fixed points (where the two strands of $A_{(1,1)}$ are paired) and the trivial ruling with all fixed points. It follows from Theorem 3.6 or (1) that we have

$$R_{S(K, A_{(1,1)})}(z) = 1 + \widetilde{R}_{S(K, A_{(1,1)})}(z) = R_{A_{(1,1)}}(z) + \widetilde{R}_{S(K, A_{(1,1)})}(z).$$

Thus K is $A_{(1,1)}$ –compatible if and only if $S(K, A_{(1,1)})$ has a reduced ruling.

For $\Lambda = (2)$, the only generalized ruling of $A_{(2)}$ is the trivial ruling where all points are fixed by the involution. From (1), we have

$$R_{S(K, A_{(2)})}(z) = \widetilde{R}_{S(K, A_{(2)})}(z) = R_{A_{(2)}}(z) + \widetilde{R}_{S(K, A_{(2)})}(z).$$

In this case as well, K is $A_{(2)}$ –compatible if and only if $S(K, A_{(2)})$ has a reduced ruling.

Theorem 4.7 then yields the following statement for 2–dimensional representations.

Theorem 5.1 *Let K be a Legendrian knot in \mathbb{R}^3 with DGA $(\mathcal{A}(K, *), \partial)$.*

- (1) *K is $A_{(1,1)}$ –compatible if and only if $(\mathcal{A}(K, *), \partial)$ has a 2–dimensional (ungraded) representation sending t to an upper triangular 2×2 matrix $\begin{pmatrix} 1 & * \\ 0 & 1 \end{pmatrix} \in \text{GL}_2(\mathbb{Z}/2)$.*

- (2) K is $A_{(2)}$ –compatible if and only if $(\mathcal{A}(K, *), \partial)$ has a 2–dimensional representation sending t to a matrix $M_{A_2}U$, where U is upper triangular and M_{A_2} is of the form $\begin{pmatrix} * & 1 \\ 1 & 0 \end{pmatrix}$.

Given a representation of a DGA $(\mathcal{A}(K, *), \partial)$ sending t to a matrix M , we can clearly construct a representation sending t to any matrix conjugate to M . In the group $\text{GL}_2(\mathbb{Z}/2)$, there are three conjugacy classes, represented by

$$I = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad A = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad B = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}.$$

Easy linear algebra yields the following corollary to Theorem 5.1.

Corollary 5.2 *Let K be a Legendrian knot in \mathbb{R}^3 with DGA $(\mathcal{A}(K, *), \partial)$.*

- (1) K is $A_{(1,1)}$ –compatible if and only if $(\mathcal{A}(K, *), \partial)$ has a 2–dimensional representation sending t to either I or A .
- (2) K is $A_{(2)}$ –compatible if and only if $(\mathcal{A}(K, *), \partial)$ has a 2–dimensional representation sending t to either A or B (equivalently, sending t to any invertible matrix besides I).

Furthermore, both conditions depend only on the smooth type and Thurston–Bennequin number of K , and either of these conditions ensures that K maximizes tb .

Proof For the two numbered statements, enumerate the conjugacy classes in $\text{GL}_2(\mathbb{Z}/2)$ represented by matrices of the form U or $M_{A_2}U$, with notation as in Theorem 5.1. For the final statement, refer to Theorem 3.14 and Theorem 3.17. \square

We now discuss $A_{(1,1)}$ –compatibility and $A_{(2)}$ –compatibility separately. Note that any K that has an ungraded ruling is automatically $A_{(1,1)}$ –compatible (just double the ruling in $S(K, A_{(1,1)})$); it also has an ungraded augmentation and thus a (reducible) 2–dimensional representation sending t to I .

Sivek, in his paper [19] along with some other examples posted online, has constructed examples of Legendrian knots that have no rulings but whose DGAs do have a (necessarily irreducible) 2–dimensional representation sending t to I . These include Legendrian versions of the torus knots $T(p, -q)$ with $q > p \geq 3$ as well as $m(9_{42})$, $m(10_{128})$ and $m(10_{136})$. By Corollary 5.2, each of these knots K is $A_{(1,1)}$ –compatible and thus $S(K, A_{(1,1)})$ has a reduced ruling. This last fact can be seen explicitly for the torus knots, independent of Sivek’s work; see Figure 15, which gives a reduced ruling when K is a $(3, -4)$ torus knot, and which can be readily generalized to any maximal– tb Legendrian negative torus knot.




Figure 15: A ruling for the double of a negative torus knot. Dots represent switches. Shown here: $S(K, A_{(1,1)})$, where K is a standard Legendrian $(3, -4)$ torus knot.

Maximal- tb Legendrian representatives of most knots with 10 or fewer crossings satisfy the Kauffman polynomial bound on tb and thus have ungraded rulings and augmentations. The exceptions are the knots $T(3, -4) = m(8_{19})$, $m(9_{42})$, $m(10_{124}) = T(3, -5)$, $m(10_{128})$, $m(10_{132})$ and $m(10_{136})$. Sivek's calculations, along with Corollary 5.2, give the following.

Theorem 5.3 *Let K be a Legendrian representative of a knot with crossing number at most 10. Then K is $A_{(1,1)}$ -compatible if and only if both of the following hold: K maximizes tb , and K is not of topological type $m(10_{132})$.*

Note that the “only if” statement follows from Sivek's calculation in [19] that the DGA for a particular maximal- tb representative of $m(10_{132})$ is trivial.

We remark that from our results, the existence of a ruling for $S(K, A_{(1,1)})$ implies the existence of a 2-dimensional representation of $(\mathcal{A}(K, *), \partial)$, but not necessarily one that sends t to I as in Sivek's examples; the representation might send t to A . However, we do not know of a knot where $S(K, A_{(1,1)})$ has a ruling but $(\mathcal{A}(K, *), \partial)$ has no 2-dimensional representation sending t to I .

Finally, we turn to $A_{(2)}$ –compatibility. Unlike for $A_{(1,1)}$, the existence of an ungraded ruling of K does *not* necessarily imply that K is $A_{(2)}$ –compatible. Indeed, the standard Legendrian unknot with $tb = -1$ is not $A_{(2)}$ –compatible. However, a slightly stronger condition on rulings of K does imply $A_{(2)}$ –compatibility.

Theorem 5.4 *If K is a Legendrian knot in \mathbb{R}^3 with $\deg R_K(z) \geq 0$ (ie, K has a ruling where the number of switches is at least the number of right cusps), then K is $A_{(2)}$ –compatible.*

Proof Consider a ruling of K where the number of switches s is at least the number of right cusps c ; this decomposes the front of K into c unknots. Construct a (planar) graph with c vertices and s edges, where the vertices correspond to the unknots and edges correspond to switches. Since $s \geq c$, this graph has a nonempty closed loop. Thus we may choose some nonempty subset of “distinguished” switches such that every unknot contains an even number of distinguished switches.




Figure 16: Switches at doubled crossings in $S(K, A_{(2)})$: at the distinguished switch D (left), at every other distinguished switch (middle) and at every non-distinguished switch (right)

We use this information to construct a ruling of $S(K, A_{(2)})$. Choose one distinguished switch D of the ruling of K , which we also view as a crossing in the front of K . We construct the front for $S(K, A_{(2)})$ as follows: start with the double of the front for K , so that every crossing in the front for K produces four crossings in the double, then place the extra crossing for $A_{(2)}$ in the middle of the four crossings corresponding to D , as shown in the leftmost diagram of Figure 16.

Now place switches at crossings of $S(K, A_{(2)})$ as follows: do not switch at crossings corresponding to cusps of K , and at crossings corresponding to crossings of K , switch according to Figure 16. We leave it as an exercise for the reader to check that this choice of switches on $S(K, A_{(2)})$ determines a ruling. \square

We can use Theorem 5.4 to address the question of which small Legendrian knots are $A_{(2)}$ –compatible. Recall from [15] that the ungraded ruling polynomial $R_K(z)$ depends only on $tb(K)$ and the Kauffman polynomial of the smooth knot underlying K . An inspection of the Kauffman polynomial for smooth knots with up to 10

crossings shows that if K is a maximal- tb representative of a smooth knot with at most 10 crossings, then $\deg R_K(z) \geq 0$ unless K is of one of the following types: 0_1 , $m(8_{19})$, $m(9_{42})$, $m(9_{46})$, $m(10_{124})$, $m(10_{128})$, $m(10_{132})$, $m(10_{136})$ and $m(10_{140})$. Of these exceptions, a direct computation using *Mathematica* shows that maximal- tb representatives of $m(8_{19})$, $m(9_{42})$, $m(9_{46})$, $m(10_{124})$, $m(10_{128})$ and $m(10_{136})$ are $A_{(2)}$ -compatible: their satellite with $A_{(2)}$ has an ungraded ruling. On the other hand, maximal- tb representatives of 0_1 , $m(10_{132})$ and $m(10_{140})$ are not $A_{(2)}$ -compatible. We summarize these findings in the following result.

Theorem 5.5 *Let K be a Legendrian representative of a knot with crossing number at most 10. Then K is $A_{(2)}$ -compatible if and only if both of the following hold: K has maximal tb , and K is not of topological type 0_1 , $m(10_{132})$ or $m(10_{140})$.*

References

- [1] **Y V Chekanov**, *Differential algebra of Legendrian links*, Invent. Math. 150 (2002) 441–483 MR1946550
- [2] **T Ekholm, J B Etnyre, L Ng, M G Sullivan**, *Knot contact homology*, Geom. Topol. 17 (2013) 975–1112 MR3070519
- [3] **J B Etnyre, L Ng, J M Sabloff**, *Invariants of Legendrian knots and coherent orientations*, J. Symplectic Geom. 1 (2002) 321–367 MR1959585
- [4] **D Fuchs**, *Chekanov–Eliashberg invariant of Legendrian knots: existence of augmentations*, J. Geom. Phys. 47 (2003) 43–65 MR1985483
- [5] **D Fuchs, T Ishkhanov**, *Invariants of Legendrian knots and decompositions of front diagrams*, Mosc. Math. J. 4 (2004) 707–717, 783 MR2119145
- [6] **MB Henry**, *Connections between Floer-type invariants and Morse-type invariants of Legendrian knots*, Pacific J. Math. 249 (2011) 77–133 MR2764942
- [7] **T Kálmán**, *Braid-positive Legendrian links*, Int. Math. Res. Not. 2006 (2006) MR2272097
- [8] **M Lavrov, D Rutherford**, *Generalized normal rulings and invariants of Legendrian solid torus links*, Pacific J. Math. 258 (2012) 393–420 MR2981960
- [9] **K Mishachev**, *The N -copy of a topologically trivial Legendrian knot*, J. Symplectic Geom. 1 (2003) 659–682 MR2039159
- [10] **L Ng**, *Computable Legendrian invariants*, Topology 42 (2003) 55–82 MR1928645
- [11] **L Ng**, *Rational symplectic field theory for Legendrian knots*, Invent. Math. 182 (2010) 451–512 MR2737704
- [12] **L Ng, J M Sabloff**, *The correspondence between augmentations and rulings for Legendrian knots*, Pacific J. Math. 224 (2006) 141–150 MR2231655

- [13] **L Ng, L Traynor**, *Legendrian solid-torus links*, *J. Symplectic Geom.* 2 (2004) 411–443 MR2131643
- [14] **P E Pushkar’, Y V Chekanov**, *Combinatorics of fronts of Legendrian links, and Arnol’d’s 4–conjectures*, *Uspekhi Mat. Nauk* 60 (2005) 99–154 MR2145660 In Russian; translated in *Russian Math. Surveys* 60 (2005) 95–149
- [15] **D Rutherford**, *Thurston–Bennequin number, Kauffman polynomial, and ruling invariants of a Legendrian link: the Fuchs conjecture and beyond*, *Int. Math. Res. Not.* 2006 (2006) MR2219227
- [16] **D Rutherford**, *HOMFLY-PT polynomial and normal rulings of Legendrian solid torus links*, *Quantum Topol.* 2 (2011) 183–215 MR2783131
- [17] **J M Sabloff**, *Augmentations and rulings of Legendrian knots*, *Int. Math. Res. Not.* 2005 (2005) 1157–1180 MR2147057
- [18] **C Shonkwiler, D S Vela-Vick**, *Legendrian contact homology and nondestabilizability*, *J. Symplectic Geom.* 9 (2011) 33–44 MR2787360
- [19] **S Sivek**, *The contact homology of Legendrian knots with maximal Thurston–Bennequin invariant*, *J. Symplectic Geom.* 11 (2013) 167–178

*Department of Mathematics, Duke University
Box 90320, Durham, NC 27708-0320, USA*

*Department of Mathematics, University of Arkansas
301 SCEN, 1 University of Arkansas, Fayetteville, AR 72701, USA*

ng@math.duke.edu, drruther@uark.edu

<http://www.math.duke.edu/~ng/>

Received: 15 June 2012 Revised: 8 April 2013

