

Koszul duality theory for operads over Hopf algebras

OLIVIA BELLIER

The transfer of the generating operations of an algebra to a homotopy equivalent chain complex produces higher operations. The first goal of this paper is to describe precisely the higher structure obtained when the unary operations commute with the contracting homotopy. To solve this problem, we develop the Koszul duality theory of operads in the category of modules over a cocommutative Hopf algebra. This gives rise to a simpler category of homotopy algebras and infinity morphisms, which allows us to get a new description of the homotopy category of algebras over such operads. The main example of this theory is given by Batalin–Vilkovisky algebras.

18D50, 18G55; 16W30, 55P48

1 Introduction

Homotopy transfer In homotopical algebra, one problem is to know how algebraic structures behave under homotopy equivalences. For instance, the product of a differential graded associative algebra induces, on a homotopy equivalent chain complex, a product which is not associative in general. However, Kadeishvili [16] proved that, in this case, the homotopy equivalent chain complex carries higher operations, in addition to the transferred product. These operations endow it with a homotopy associative algebra structure, also known as A_∞ -algebra, defined by Stasheff in [25]. This is one of the first examples of a *homotopy transfer theorem* (HTT). An HTT was also proved for Lie algebras, for commutative algebras, and more generally, for other types of algebras using the theory of operads; see Loday and Vallette [21, Section 10.3].

Batalin–Vilkovisky algebras The theory we develop in this paper is motivated by the example of Batalin–Vilkovisky algebras. These algebras play an important role in geometry, topology and mathematical physics; see for instance Batalin and Vilkovisky [2], Barannikov and Kontsevich [1], Manin [22], Chas and Sullivan [6], Getzler [11] and Lian and Zuckerman [19]. In brief, a BV-algebra structure is given by a commutative product, a Lie bracket and a unary square-zero BV-operator, satisfying some compatibility relations. An HTT for BV-algebras was given by Galvez-Carillo, Tonks and Vallette [10] in terms of a new notion of homotopy BV-algebra. This structure is very rich, and at the same time very intricate.

However, when the BV-operator commutes with the underlying contracting homotopy, it induces a unary operator which still squares to zero. In this case, we have that there are no higher operations arising from the BV-operator when we apply the HTT of [10]. It is natural to ask what is the precise structure to which it reduces. In this paper, we solve this question as follows: we insert the square-zero operator in the underlying category of chain complexes and, then, we work with operads in this category. This implies that we have to work with operads in the category of modules over a *cocommutative Hopf algebra*. For instance, to encode the category of BV-algebras, we use the operad of Gerstenhaber algebras, which is enriched with an action of the dual numbers Hopf algebra.

Relative Koszul duality Salvatore and Wahl [24] began the study of operads with an action of a Hopf algebra. Here, we go even further and we extend the classical Koszul duality theory to this framework in Section 3.3. At each step of the operadic theory, we show that all the objects can be enriched with a compatible action of a Hopf algebra and we prove that the results still hold in this context. In particular, we extend the bar–cobar constructions in Section 3.2. We define a new notion of homotopy algebras and their infinity morphisms in Section 4.1, in order to prove a new homotopy transfer theorem in Section 4.2. We also extend the results of the homotopy theory of algebras over an operad to the context of algebras over an H -operad in Section 4.3.

Theorem A *Let H be a cocommutative Hopf algebra and let \mathcal{P} be a Koszul operad in the category of H -modules. We consider two homotopy equivalent chain complexes in the category of H -modules, such that one of them is endowed with a compatible \mathcal{P} -algebra structure. Then the second chain complex is endowed with a \mathcal{P}_∞ -algebra structure compatible with the H -action.*

The main example of two homotopy equivalent chain complexes is given by a differential graded algebra and its homology groups. In this case, being homotopy equivalent in the category of H -modules means just that the contracting homotopy and the H -action have to commute.

The homotopy category is obtained by localizing the initial category with respect to the quasi-isomorphisms. The ∞ -morphisms provide a different description of the homotopy category of algebras over an operad. It is easier to deal with this description since the ∞ -morphisms admit a homotopy inverse; see Vallette [27]. Encoding a category of algebras with an operad in H -modules, we get a new description of the associated homotopy category, using simpler ∞ -morphisms, as follows.

Theorem B *Let H be a cocommutative Hopf algebra and let \mathcal{P} be a Koszul operad in the category of H -modules. The homotopy category of dg \mathcal{P} -algebras in H -modules is equivalent to the homotopy category of \mathcal{P} -algebras in H -modules, together with their ∞ - H -morphisms.*

In the case of BV-algebras, we define a simpler category of homotopy BV-algebras: a *strict homotopy BV-algebra* is a homotopy Gerstenhaber algebra together with a compatible square-zero unary operator action. [Theorem A](#) implies that, when the BV-operator and the contracting homotopy commute, the transferred homotopy BV-algebra structure reduces exactly to a strict homotopy BV-algebra. [Theorem B](#) gives a new description of the homotopy category of BV-algebras. We also get a new way to prove the existence of a zigzag of quasi-isomorphisms of BV-algebras, which could help to study the mirror symmetry conjecture by Kontsevich [17]; see Cao and Zhou [4; 5].

Related literature The idea to put unary operations in the underlying category is already used by Ginzburg and Kapranov [14], but not exactly in the same way as we do. We both put the algebra of *all* the arity one operations in the underlying category, while removing them from the operad. In [14], they keep track of their action using the tensor product over this algebra. Instead, we also remove from the operad all the operations made up of the unary ones, and we keep track of the action of these operations via the module structure over their algebra. Thus, we can still work with the tensor product over the ground field. So, homological results hold in our context, while, in [14], one needs the algebra of unary operations to be semisimple. Nevertheless, we need this algebra to form a Hopf algebra. Moreover, since Ginzburg and Kapranov remove *only* the unary operations from the operad, their homotopy algebra structure is much bigger than ours.

There are so far two ways of proving an HTT. On the one hand, one can prove it with model category arguments by Berger and Moerdijk [3]. This relies on a compatibility between the monoidal and the model structures of the underlying category. But, to the best of our knowledge, there is not yet a monoidal model category structure on the category of modules over a cocommutative Hopf algebra together with the tensor product over the ground field, such that the weak equivalences are the quasi-isomorphisms. On the other hand, one can prove an HTT with explicit formulae using the Koszul duality theory for operads, which enables one to prove formality results at the level of algebras for instance. So, we chose this latter method to prove an HTT for operads over Hopf algebras.

Layout In the second section, we describe the notion of operads in the category of modules over a Hopf algebra and the associated objects. We give some examples of algebras over such operads. In [Section 3](#), we extend the classical Koszul duality to this

framework. In [Section 4](#), we prove an HTT for algebras over Koszul operads in this context, and we apply it to the example of Batalin–Vilkovisky algebras. Furthermore, we study the homotopy theory of such algebras.

Setting Throughout this paper, we work over a field \mathbb{K} of characteristic zero and unadorned tensor products are over \mathbb{K} . All the \mathbb{S} –modules we consider are reduced, that is the arity zero space is reduced to zero.

2 Operads over Hopf algebras

2.1 The monoidal category of symmetric modules over a cocommutative Hopf algebra

We construct the monoidal category of (differential graded) \mathbb{S} –modules over a cocommutative Hopf algebra, following the example of \mathbb{S} –modules given by Loday and Vallette [[21](#), Section 5.1]. We refer to the book of Sweedler [[26](#)] for more details on Hopf algebras.

2.1.1 Recollection on the symmetric monoidal closed category of modules over a Hopf algebra Let $(H, \mu, \Delta, u, \varepsilon)$ be a cocommutative bialgebra, where (μ, u) and (Δ, ε) are respectively the unital associative algebra structure and the counital cocommutative coalgebra structure. We will denote $1_H := u(1_{\mathbb{K}})$. Then, we have $\varepsilon(1_H) = 1_{\mathbb{K}}$ and $H \cong \text{Ker}(\varepsilon: H \rightarrow \mathbb{K}) \oplus \mathbb{K}1_H$.

We define the iterated comultiplication $\Delta^n: H \rightarrow H^{\otimes n+1}$ by $\Delta^0 = \text{Id}$, $\Delta^1 = \Delta$ and

$$\Delta^n := (\Delta \otimes \text{Id} \otimes \cdots \otimes \text{Id}) \circ \Delta^{n-1}.$$

The category $(H\text{-Mod}, \otimes, \mathbb{K})$ of left– H –modules is a symmetric monoidal category where the structure of the H –module is given as follows:

- On \mathbb{K} , it is defined by the map $H \otimes \mathbb{K} \xrightarrow{\varepsilon \otimes \mathbb{K}} \mathbb{K} \otimes \mathbb{K} \longrightarrow \mathbb{K}$ where the last map is the scalar multiplication.
- On the tensor product of two H –modules M and N , it is defined by

$$h.(m \otimes n) = h_{(0)}.m \otimes h_{(1)}.n, \quad \forall h \in H, \forall m \in M, \forall n \in N,$$

where $\Delta(h) = h_{(0)} \otimes h_{(1)}$, under the sumless Sweedler notation. The Hopf compatibility relation ensures that it is a left H –action.

We assume moreover that H is a Hopf algebra, that is H is equipped with a linear map $S: H \rightarrow H$, called the *antipode*, which is an inverse of the identity for the convolution product \star :

$$S \star \text{Id} = u\varepsilon = \text{Id} \star S.$$

Then, for any H -modules M and N , the vector space $\text{Hom}_{\mathbb{K}}(M, N)$ carries an H -module structure, given by

$$\begin{aligned} H \otimes \text{Hom}_{\mathbb{K}}(M, N) &\rightarrow \text{Hom}_{\mathbb{K}}(M, N), \\ h \otimes f &\mapsto h.f: m \mapsto h_{(0)}.f(S(h_{(1)}).m). \end{aligned}$$

Lemma 2.1 *The category of H -modules is closed.*

Proof For any triple (A, B, C) of H -modules, the map

$$\begin{aligned} \text{Hom}(A \otimes B, C) &\rightarrow \text{Hom}(A, \text{Hom}(B, C)), \\ f &\mapsto (a \mapsto f(a \otimes -)), \end{aligned}$$

is well defined. It is easy to check that it is a natural isomorphism of H -modules. \square

The previous results extend to the category of differential graded modules.

Proposition 2.2 *We have that the category $(\text{dg } H\text{-Mod}, \otimes, \mathbb{K})$ of differential graded left- H -modules is a symmetric monoidal closed category, where \mathbb{K} is concentrated in degree 0.*

Example Consider the graded algebra $D := \mathbb{K}[\delta]/(\delta^2)$ of dual numbers, where δ is of degree $+1$. It is a cocommutative Hopf algebra where the comultiplication $\Delta: D \rightarrow D \otimes D$ and the antipode $S: D \rightarrow D$ are respectively given by

$$\begin{aligned} \Delta(\delta) &:= 1 \otimes \delta + \delta \otimes 1, \\ S(\delta) &:= -\delta. \end{aligned}$$

A D -module is simply a graded vector space endowed with a map of degree $+1$ that squares to zero, *ie* a cochain complex.

2.1.2 The monoidal category of \mathbb{S} - H -modules The notion of \mathbb{S} -modules can be defined in any symmetric monoidal category. We make explicit these objects when the underlying category is equal to the category of H -modules.

Definition 2.3 An \mathbb{S} - H -module M is an \mathbb{S} -module $M = \{M(n)\}_{n \in \mathbb{N}}$, such that each $M(n)$ has a left- H -module structure which commutes with the \mathbb{S}_n -module structure. For $\mu \in M(n)$, the integer n is called the *arity* of μ and μ is called an n -*ary operation*.

A *morphism of \mathbb{S} - H -modules* $f: M \rightarrow N$ is a morphism of \mathbb{S} -modules such that each $f_n: M(n) \rightarrow N(n)$ is H -equivariant.

We denote the associated category by \mathbb{S} - H -Mod.

In particular, each $M(n)$ is a (H, \mathbb{S}_n) -bimodule and each $f_n: M(n) \rightarrow N(n)$ is a morphism of (H, \mathbb{S}_n) -bimodules.

Definition 2.4 For any \mathbb{S} - H -modules M and N , their *tensor product* is the \mathbb{S} - H -module $M \otimes N$ defined by

$$(M \otimes N)(n) := \bigoplus_{i+j=n} \text{Ind}_{\mathbb{S}_i \times \mathbb{S}_j}^{\mathbb{S}_n} M(i) \otimes N(j),$$

where the action of H is induced by the H -module structure on the tensor product of two H -modules. Since H is cocommutative, the tensor product of H -modules is symmetric.

Proposition 2.5 *The tensor product of \mathbb{S} - H -modules is associative with unit the \mathbb{S} - H -module $(\mathbb{K}, 0, 0, \dots)$.*

Proof By [21, Proposition 5.1.5], we have an isomorphism of associativity of the underlying \mathbb{S} -modules. It is a morphism of H -modules by coassociativity of the comultiplication. \square

Definition 2.6 For any \mathbb{S} - H -modules M and N , their *composite product* is the \mathbb{S} - H -module $M \circ N$ defined by

$$M \circ N := \bigoplus_{k \geq 0} M(k) \otimes_{\mathbb{S}_k} N^{\otimes k}.$$

Here $N^{\otimes k}$ stands for the tensor product of k copies of the \mathbb{S} - H -module N .

For any pair of morphisms of \mathbb{S} - H -modules $f: M \rightarrow N$ and $g: M' \rightarrow N'$, their *composite product* is the morphism of \mathbb{S} - H -modules $f \circ g: M \circ N \rightarrow M' \circ N'$ given explicitly by the formula

$$f \circ g(\mu; v_1, \dots, v_k) := (f(\mu); g(v_1), \dots, g(v_k)).$$

In arity n , we have

$$(M \circ N)(n) := \bigoplus_{k \geq 0} M(k) \otimes_{\mathbb{S}_k} \left(\bigoplus_{i_1 + \dots + i_k = n} \text{Ind}_{\mathbb{S}_{i_1} \times \dots \times \mathbb{S}_{i_k}}^{\mathbb{S}_n} N(i_1) \otimes \dots \otimes N(i_k) \right),$$

where the action of H is induced by the H -module structure on tensor products of H -modules.

Proposition 2.7 *The category of \mathbb{S} - H -modules $(\mathbb{S}\text{-}H\text{-Mod}, \circ, I)$ is a monoidal category, where the \mathbb{S} - H -module $I = (0, \mathbb{K}, 0, \dots)$ concentrated in arity 1 is called identity \mathbb{S} - H -module.*

Proof By [21, Proposition 5.1.14], the category of \mathbb{S} -modules $(\mathbb{S}\text{-Mod}, \circ, I)$ is monoidal. Since the composite product of \mathbb{S} - H -modules is defined as the composite product of the underlying \mathbb{S} -modules endowed with the H -action induced by the comultiplication, the coassociativity and the counitariness of the comultiplication of H imply that the isomorphisms of associativity and unit of the underlying \mathbb{S} -modules are compatible with the H -module structure. \square

As for \mathbb{S} -modules, the composite product of two \mathbb{S} - H -modules is not linear on the right-hand side. However, the infinitesimal composite product, defined in [21, Section 6.1.1], extends to the category $\mathbb{S}\text{-}H\text{-Mod}$ to produce a product $\circ_{(1)}$ which is linear on the right-hand side. Recall that the infinitesimal product is defined by

$$M \circ_{(1)} N := M \circ (I; N),$$

where, for any \mathbb{S} -modules P, P_1 and P_2 , $P \circ (P_1; P_2)$, is the \mathbb{S} -module

$$(P \circ (P_1; P_2))(n) := \bigoplus_{k=1}^n P(k) \otimes_{\mathbb{S}_k} \left(\bigoplus_{i_1 + \dots + i_k = n} \bigoplus_{j=1}^k \text{Ind}_{\mathbb{S}_{i_1} \times \dots \times \mathbb{S}_{i_k}}^{\mathbb{S}_n} P_1(i_1) \otimes \dots \otimes \underbrace{P_2(i_j)}_{j^{\text{th}} \text{ position}} \otimes \dots \otimes P_1(i_k) \right).$$

The infinitesimal objects $f \circ_{(1)} g, f \circ' g, \gamma_{(1)}$ and $\Delta_{(1)}$, defined in [21, Section 6.1], also extend to the category $\mathbb{S}\text{-}H\text{-Mod}$.

2.1.3 Differential graded framework

Definition 2.8 *A differential graded \mathbb{S} - H -module, or dg \mathbb{S} - H -module for short, is a dg \mathbb{S} -module (M, d) such that each $M(n)$ is an H - \mathbb{S}_n -bimodule and the differential d is compatible with the H -action.*

A morphism of differential graded \mathbb{S} - H -modules $f: (M, d_M) \rightarrow (N, d_N)$ is a morphism of the underlying dg \mathbb{S} -modules which is H -equivariant. We denote by $\text{dg } \mathbb{S}\text{-}H\text{-Mod}$ the category of dg \mathbb{S} - H -modules with their morphisms.

The objects described in the previous section extend to the differential graded framework. However, they now involve signs in their definition. For more details, we refer the reader to [21, Section 6.2].

We define the *suspension* and the *desuspension* of a graded \mathbb{S} - H -module as the suspension and the desuspension of its underlying graded \mathbb{S} -module respectively, where the action of H is given by the counit of H .

Proposition 2.9 *The category of $(\text{dg } \mathbb{S}\text{-}H\text{-Mod}, \circ, I)$ is a monoidal category.*

Proof This follows by combining Proposition 2.7 and [21, Proposition 6.2.4]. \square

2.2 Operad over a cocommutative Hopf algebra

In [21, Section 5.2], the authors give a monoidal definition of an algebraic operad, that is an operad in the category of vector spaces. More generally, one can define the notion of operad in any monoidal category. Here, we point out the extra structure one gets when one considers operads in the category of \mathbb{S} - H -modules instead of just \mathbb{S} -modules.

2.2.1 Monoidal definition

Definition 2.10 An H -operad is a monoid $(\mathcal{P}, \gamma, \eta)$ in the monoidal category of \mathbb{S} - H -modules. A morphism $\alpha: \mathcal{P} \rightarrow \mathcal{Q}$ of H -operads is a morphism of monoids in the category of \mathbb{S} - H -modules, that is a morphism of \mathbb{S} - H -modules which is compatible with the monoidal structures. We denote the category of H -operads by Op_H .

Remark An H -operad is just an operad $(\mathcal{P}, \gamma, \eta)$, as defined in [21, Section 5.2.1], such that each $\mathcal{P}(n)$ has an H -action which makes \mathcal{P} into an \mathbb{S} - H -module and such that the maps γ and η commute with the H -action. Furthermore, a morphism of H -operads is a morphism of the underlying operads which commutes with the action of H .

When A is an H -module, the \mathbb{S} -module

$$\text{End}_A(n) := \text{Hom}_{\mathbb{K}}(A^{\otimes n}, A)$$

is endowed with an H -module structure, according to the previous section, which makes it into an \mathbb{S} - H -module. This structure is explicitly given by

$$H \otimes \text{End}_A(n) \rightarrow \text{End}_A(n),$$

$$h \otimes f \mapsto h.f := a_1 \otimes \cdots \otimes a_n \mapsto h_{(0)}.f(S(h_{(1)}).a_1, \dots, S(h_{(n)}).a_n).$$

Lemma 2.11 *The triple $(\text{End}_A, \gamma_{\text{End}_A}, \eta_{\text{End}_A}: 1_{\mathbb{K}} \mapsto \text{Id}_A)$, where γ_{End_A} is the usual composition map given by $\gamma_{\text{End}_A}(f; g_1, \dots, g_k) := f \circ (g_1 \otimes \cdots \otimes g_k)$, is an H -operad.*

Proof It is immediate to check End_A is an operad. So, we have to prove γ_{End_A} and η_{End_A} are H -equivariant maps. For $(f; g_1, \dots, g_k) \in \text{End}_A(n)$, $a_1 \otimes \cdots \otimes a_n \in A^{\otimes n}$ and $h \in H$, we have

$$\begin{aligned} & \gamma_{\text{End}_A}(h.(f; g_1, \dots, g_k))(a_1 \otimes \cdots \otimes a_n) \\ &= h_{(0)}.f \left(\bigotimes_{j=1}^k [S(h_{(n+j)(0)})h_{(n+j)(1)}].g_j \left(\bigotimes_{l=1}^{i_j} S(h_{(i_1+\dots+i_{j-1}+l)})a_{i_1+\dots+i_{j-1}+l} \right) \right) \\ &= h_{(0)}.f \left(\bigotimes_{j=1}^k u\varepsilon(h_{(n+j)}).g_j \left(\bigotimes_{l=1}^{i_j} S(h_{(i_1+\dots+i_{j-1}+l)})a_{i_1+\dots+i_{j-1}+l} \right) \right) \\ &= h_{(0)}.(f \circ (g_1 \otimes \cdots \otimes g_k))(S(h_{(1)}).a_1 \otimes \cdots \otimes S(h_{(n)}).a_n) \\ &= (h.\gamma_{\text{End}_A}(f; g_1, \dots, g_k))(a_1 \otimes \cdots \otimes a_n). \end{aligned}$$

The H -equivariance of η_{End_A} follows from the H -module structure of \mathbb{K} and from the fact that the antipode is the inverse of the identity for the convolution product. \square

Recall that the *Hadamard product* $M \otimes_{\mathcal{H}} N$ of two \mathbb{S} -modules M and N is defined to be the following \mathbb{S} -module

$$M \otimes_{\mathcal{H}} N(n) := M(n) \otimes N(n).$$

Proposition 2.12 *The Hadamard product of the underlying \mathbb{S} -modules of two H -operads, endowed with the H -module structure on the tensor product of two \mathbb{S} -modules, is an H -operad.*

Proof Let \mathcal{P} and \mathcal{Q} be two H -operads. By [21, Section 5.3.3], we have to prove that the composition map given there is H -equivariant. This is the case since it is the composite of H -equivariant maps. \square

Definition 2.13 An ideal \mathcal{I} of an H -operad \mathcal{P} is a sub- \mathbb{S} - H -module of \mathcal{P} such that the composition $\gamma_{\mathcal{P}}(\mu; v_1, \dots, v_k)$ is in \mathcal{I} as soon as one of the μ, v_1, \dots, v_k is in \mathcal{I} . The quotient of an H -operad \mathcal{P} by the ideal \mathcal{I} is the H -operad \mathcal{P}/\mathcal{I} given by

$$(\mathcal{P}/\mathcal{I})(n) := \mathcal{P}(n)/\mathcal{I}(n), \quad \forall n \in \mathbb{N},$$

where the composition map $\gamma_{\mathcal{P}/\mathcal{I}}$ is induced by $\gamma_{\mathcal{P}}$.

2.2.2 The free H -operad

Definition 2.14 A free H -operad over an \mathbb{S} - H -module M is an H -operad $\mathcal{F}M$ equipped with a morphism $\eta_M: M \rightarrow \mathcal{F}M$ of \mathbb{S} - H -modules, which satisfies the following universal property.

Any morphism $f: M \rightarrow \mathcal{P}$ of \mathbb{S} - H -modules, where \mathcal{P} is an H -operad, extends uniquely into a morphism $\tilde{f}: \mathcal{F}M \rightarrow \mathcal{P}$ of H -operads:

$$\begin{array}{ccc} M & \xrightarrow{\eta_M} & \mathcal{F}M \\ & \searrow f & \downarrow \tilde{f} \\ & & \mathcal{P} \end{array}$$

In other words, the functor $\mathcal{F}: \mathbb{S}\text{-}H\text{-Mod} \rightarrow \text{Op}_H$ is left adjoint to the forgetful functor \sqcup from Op_H to $\mathbb{S}\text{-}H\text{-Mod}$.

Recall that the free operad $\mathcal{T}M$ over an \mathbb{S} -module M admits the realization

$$(\mathcal{T}M)(n) \cong \bigoplus_{\tau \in T'(n)} \tau(M),$$

where $T'(n)$ denotes a set of representatives of isomorphisms classes of n -trees. For any tree τ , the treewise tensor module $\tau(M)$, defined by Hoffbeck [15, Section 2.6], is given by

$$\tau(M) \cong \bigotimes_{v \in V(\tau)} M(|I_v|),$$

where $V(\tau)$ denotes the set of vertices of τ and where I_v denotes the set of entries of the vertex v . Then, if M is an \mathbb{S} - H -module, we have an H -module structure on the treewise tensor module $\tau(M)$, for any tree τ in $T'(n)$. This action is given by the map

$$\begin{aligned} H \otimes \tau(M) &\rightarrow \tau(M), \\ h \otimes m_{i_1} \cdots m_{i_{|V(\tau)|}} &\mapsto h_{(0)} \cdot m_{i_1} \cdots h_{(|V(\tau)|-1)} \cdot m_{i_{|V(\tau)|}}, \end{aligned}$$

where $\Delta^{|\mathcal{V}(\tau)|-1}(h) := h_{(0)} \otimes \cdots \otimes h_{(|\mathcal{V}(\tau)|-1)}$. It amounts to act on the labeling of each vertex of τ using the comultiplication of H .

Proposition 2.15 *Let M be an \mathbb{S} - H -module and \underline{M} be its underlying \mathbb{S} -module. The free operad $\mathcal{T}\underline{M}$ endowed with the H -action described above is the free H -operad over M .*

Proof The composition map $\gamma_{\mathcal{T}\underline{M}}$, which corresponds to the grafting of trees, is an H -equivariant map. Moreover, if a morphism $f: M \rightarrow \mathcal{P}$ of \mathbb{S} -modules is H -equivariant then so is $\tilde{f}: \mathcal{T}\underline{M} \rightarrow \mathcal{P}$. □

As a consequence, the free H -operad inherits the weight of its underlying free operad. The *weight* of an operation μ in the free operad $\mathcal{T}M$ is defined as follows:

$$\begin{aligned} w(\text{id}) &= 0, \\ w(\mu) &= 1, \quad \forall \mu \in M, \\ w(v; v_1, \dots, v_k) &= w(v) + w(v_1) + \cdots + w(v_k). \end{aligned}$$

We denote by $\mathcal{T}M^{(r)}$ the \mathbb{S} - H -module of operations in $\mathcal{T}(M)$ of weight r .

Proposition 2.16 *Let M be an \mathbb{S} -module and R be a sub- \mathbb{S} -module of $\mathcal{T}M$. Then, the operad $\mathcal{T}M/(R)$, where (R) is the ideal of $\mathcal{T}M$ generated by R , is an H -operad if and only if*

- M is an \mathbb{S} - H -module,
- R is a sub- \mathbb{S} - H -module of $\mathcal{T}M$.

In this case, the H -module structure is induced by the action on each vertex using the comultiplication of H .

Proof The proof by Salvatore and Wahl [24, Proposition 4.2] extends to our linear context. □

2.2.3 Algebra over an H -operad

Definition 2.17 Let \mathcal{P} be an H -operad. An *algebra over \mathcal{P}* , or for short an H - \mathcal{P} -algebra, is an H -module A equipped with an H -equivariant map $\gamma_A: \mathcal{P}(A) \rightarrow A$, which is compatible with the monoidal structure of \mathcal{P} . A *morphism $f: A \rightarrow A'$ of H - \mathcal{P} -algebras* is a morphism of H -modules which is compatible with γ_A and $\gamma_{A'}$. We denote by H - \mathcal{P} -Alg the category of H - \mathcal{P} -algebras.

If A is an H - \mathcal{P} -algebra, then the map γ_A is

$$\gamma_A: \mathcal{P}(A) = \bigoplus_{n \geq 0} \mathcal{P}(n) \otimes_{\mathbb{S}_n} A^{\otimes n} \rightarrow A.$$

For $\mu \in \mathcal{P}(n)$ and $a_1 \otimes \cdots \otimes a_n \in A^{\otimes n}$, we denote $\gamma_A(\mu; a_1, \dots, a_n)$ simply by $\mu(a_1, \dots, a_n)$.

Proposition 2.18 *Let \mathcal{P} be an H -operad. An H - \mathcal{P} -algebra structure on an H -module A is equivalent to a morphism of H -operads $\mathcal{P} \rightarrow \text{End}_A$.*

Proof By [21, Proposition 5.2.13], it remains to prove that $\gamma_A: \mathcal{P}(n) \otimes A^{\otimes n} \rightarrow A$ is H -equivariant if and only if $\alpha: \mathcal{P}(n) \rightarrow \text{End}_A(n)$ is H -equivariant, since the symmetric and the H actions are compatible. This follows from the H -action on the space of morphisms of H -modules. \square

Proposition 2.19 [24] *Let \mathcal{P} be a graded H -operad. There exists a graded operad, the semidirect product $\mathcal{P} \rtimes H$, such that the category of H - \mathcal{P} -algebras, that is H -modules with an action of the H -operad \mathcal{P} , is isomorphic to the category of $\mathcal{P} \rtimes H$ -algebras, that is modules with an action of the operad $\mathcal{P} \rtimes H$.*

The operad $\mathcal{P} \rtimes H$ is given by the \mathbb{S} -module defined by

$$\mathcal{P} \rtimes H(n) := \mathcal{P}(n) \otimes H^{\otimes n},$$

where \mathbb{S}_n acts diagonally acting on $\mathcal{P}(n)$ and permuting the elements of $H^{\otimes n}$, together with the composition map defined by

$$\begin{aligned} \gamma_{\mathcal{P} \rtimes H}((\mu \otimes h); (v_1 \otimes g_1), \dots, (v_k \otimes g_k)) \\ := \gamma_{\mathcal{P}}(\mu; h_{(0)}^1 \cdot v_1, \dots, h_{(0)}^k \cdot v_k) \otimes h_{(1)}^1 \cdot g_1 \otimes \cdots \otimes h_{(1)}^k \cdot g_k, \end{aligned}$$

where $h = h^1 \otimes \cdots \otimes h^k$, $g_i = g_i^1 \otimes \cdots \otimes g_i^{n_i}$ and where h^i acts on g_i via the comultiplication of H . The unit of $\mathcal{P} \rtimes H$ is given by $\text{id} \otimes 1_H$.

2.2.4 Cooperads over a cocommutative Hopf algebra

Definition 2.20 An H -cooperad is a comonoid $(\mathcal{C}, \Delta, \eta)$ in the monoidal category of \mathbb{S} - H -modules. A morphism of H -cooperads $f: \mathcal{C} \rightarrow \mathcal{D}$ is a morphism of the underlying \mathbb{S} - H -modules compatible with the comonoidal structure of \mathcal{C} and \mathcal{D} . We denote by Coop_H the category of H -cooperads. There is an element $\text{id} \in \mathcal{C}(1)$ such that $\eta(\text{id}) = 1_{\mathbb{K}}$ and which is called the *identity cooperation*.

An H -cooperad \mathcal{C} is said to be *coaugmented* if there is a morphism $\nu: I \rightarrow \mathcal{C}$ of H -cooperads such that $\eta\nu = \text{Id}_I$.

The notion of H -cooperad is dual to that of H -operad. From the explicit description of the composite product of two \mathbb{S} - H -modules and the isomorphism between invariants and coinvariants, it follows that Δ is made up of H - \mathbb{S}_n -bimodule morphisms:

$$\Delta(n): \mathcal{C}(n) \rightarrow (\mathcal{C} \circ \mathcal{C})(n) = \bigoplus_{k=1}^n \mathcal{C}(k) \otimes_{\mathbb{S}_k} \left(\bigoplus_{i_1+\dots+i_k=n} \text{Ind}_{\mathbb{S}_{i_1} \times \dots \times \mathbb{S}_{i_k}}^{\mathbb{S}_n} \mathcal{C}(i_1) \otimes \dots \otimes \mathcal{C}(i_k) \right)$$

Proposition 2.21 *The Hadamard product of two H -cooperads carries an H -cooperad structure.*

Proof Let \mathcal{C} and \mathcal{D} be two H -cooperads. By [21, Section 8.3.3], it remains to prove that the decomposition map given there is H -equivariant. This is the case since this decomposition map is the composite of $\Delta_{\mathcal{C}} \otimes \Delta_{\mathcal{D}}$ with maps of the type $\text{Id} \otimes \dots \otimes \tau \otimes \dots \otimes \text{Id}$, which are both H -equivariant. \square

Definition 2.22 Let M be an \mathbb{S} - H -module such that $M(0) = 0$. The *cofree H -cooperad on M* is the H -cooperad, denoted by $\mathcal{T}^c(M)$, which is cofree in the category of conilpotent H -cooperads. It means that $\mathcal{T}^c(M)$ satisfies the following universal property.

For any morphism of \mathbb{S} - H -modules $\Phi: \mathcal{C} \rightarrow M$, from a conilpotent H -cooperad \mathcal{C} , such that $\Phi(\text{id}) = 0$, there exists a unique morphism $\tilde{\Phi}: \mathcal{C} \rightarrow \mathcal{T}^c(M)$ of H -cooperads whose corestriction to M is equal to Φ :

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{\Phi} & M \\ & \searrow \tilde{\Phi} & \uparrow \\ & & \mathcal{T}^c(M) \end{array}$$

2.2.5 From operads to cooperads and vice versa In the classical case, the aritywise linear dual \mathbb{S} -module $\mathcal{C}^* = \{\text{Hom}_{\mathbb{K}}(\mathcal{C}(n), \mathbb{K})\}_{n \in \mathbb{N}}$ associated to a cooperad \mathcal{C} carries an operad structure. The unit is obtained by dualization of the counit $\eta_{\mathcal{C}}$ and the composition map is obtained by dualization of $\Delta_{\mathcal{C}}$ composed with the natural map from invariants to coinvariants given in [21, Section 5.1.21]. Since H is a Hopf algebra, if \mathcal{C} is moreover an H -cooperad then its aritywise dual has an H -operad structure. Equally, if \mathcal{P} is an H -operad, such that each $\mathcal{P}(n)$ is finite dimensional, then its linear dual has an H -cooperad structure. We denote this cooperad by \mathcal{P}^c .

2.2.6 Differential graded framework As for operads, the notion of H -operad extends to the differential graded framework. In particular, a *differential graded H -operad* is a monoid $(\mathcal{P}, d_{\mathcal{P}}, \gamma, \eta)$ in the monoidal category $(\text{dg } \mathbb{S}\text{-}H\text{-Mod}, \circ, I)$, that is $(\mathcal{P}, d_{\mathcal{P}})$ is a dg \mathbb{S} - H -module and $(\mathcal{P}, \gamma, \eta)$ is an H -operad structure on \mathcal{P} , such that γ and η are morphisms of dg \mathbb{S} - H -modules. Moreover, the results on dg operads of [21, Section 6.3] can be extended to dg H -operads.

We denote by \mathcal{S} (resp. \mathcal{S}^{-1}) the endomorphism H -operad associated to the graded \mathbb{S} - H -module \mathbb{K}_s (resp. $\mathbb{K}_{s^{-1}}$), which is nothing but the H -module \mathbb{K} concentrated in degree $+1$ (resp. in degree -1). Note that we will denote by $\mathcal{S}M$ (resp. $\mathcal{S}^{-1}M$) the Hadamard product of \mathcal{S} (resp. \mathcal{S}^{-1}) with any \mathbb{S} - H -module M .

2.3 Examples

2.3.1 Mixed chain complexes A *mixed chain complex* is a graded vector space $V = V_{\bullet}$ endowed with two maps: a degree -1 map d and a degree $+1$ map δ , which both square to zero and anti-commute. One can see any mixed chain complex either as a dg- D -module, or as a dg D - I -algebra, where I is the identity D -operad.

2.3.2 Batalin–Vilkovisky algebras

Definition A *Gerstenhaber algebra* is a differential graded vector space (A, d_A) endowed with

- ◇ a symmetric binary product \bullet of degree 0 ,
- ◇ a symmetric bracket $\langle - ; - \rangle$ of degree $+1$,

such that d_A is a derivation with respect to each of them and such that:

- ▷ The product \bullet is associative:

$$((- \bullet -) \bullet -) = (- \bullet (- \bullet -)).$$

- ▷ The bracket $\langle - ; - \rangle$ satisfies the Jacobi identity

$$\langle \langle - ; - \rangle ; - \rangle + \langle \langle - ; - \rangle ; - \rangle \cdot (123) + \langle \langle - ; - \rangle ; - \rangle \cdot (321) = 0.$$

- ▷ The product \bullet and the bracket $\langle - ; - \rangle$ satisfy the Leibniz relation

$$\langle - ; - \bullet - \rangle = (\langle - ; - \rangle \bullet -) + (- \bullet \langle - ; - \rangle) \cdot (12).$$

A Batalin–Vilkovisky algebra, or *BV-algebra* for short, is a Gerstenhaber algebra A endowed, in addition, with

- ◊ a unary operator Δ of degree $+1$,

such that d_A anticommutes with Δ and such that:

- ▶ The operator satisfies $\Delta^2 = 0$.
- ▶ The bracket is the obstruction to Δ being a derivation with respect to the product \bullet :

$$\langle -; - \rangle = \Delta \circ (- \bullet -) - (\Delta(-) \bullet -) - (- \bullet \Delta(-)).$$

- ▶ The operator Δ is a graded derivation with respect to the bracket $\langle -; - \rangle$:

$$\Delta \circ (\langle -; - \rangle) + \langle \Delta(-); - \rangle + \langle -; \Delta(-) \rangle = 0.$$

Let \mathcal{G} be the operad encoding Gerstenhaber algebras. It is defined by generators and relations as

$$\mathcal{G} := \mathcal{T}(E)/(R),$$

where $E = E(2) := \bullet \mathbb{K}_2 \oplus \langle -; - \rangle \mathbb{K}_2$, with \mathbb{K}_2 the trivial representation of the symmetric group \mathbb{S}_2 . The space of relations R is the sub- \mathbb{S} -module of $\mathcal{T}(E)$ generated by the relations \triangleright . The operad \mathcal{BV} , encoding BV-algebras, is then given by

$$\mathcal{T}(E')/(R'),$$

where $E' := E \oplus \Delta \mathbb{K}$ and where R' is the sub- \mathbb{S} -module of $\mathcal{T}(E')$ generated by the relations \triangleright and \blacktriangleright .

Lemma 2.23 *The action of D on the generators \bullet and $\langle -; - \rangle$ of \mathcal{G} is given by*

$$\begin{aligned} \delta: \bullet &\mapsto \langle -; - \rangle, \\ \langle -; - \rangle &\mapsto 0, \end{aligned}$$

and induces a D -operad structure on \mathcal{G} .

Proof It is clear that E is an \mathbb{S} - D -module. Under the action of δ , the associativity relation is sent to a linear combination of Leibniz relations, the Jacobi relation is sent to 0 and the Leibniz relation is sent to the Jacobi relation. Thus, R is a sub- \mathbb{S} - D -module of $\mathcal{T}(E)$. We conclude with [Proposition 2.16](#). \square

Remark The δ -action on the generators of \mathcal{G} corresponds to the effect in homology of the action of the circle group S^1 on the little discs operad which is given by Getzler in [12, Section 4]. This structure induced in homology is made explicit by Salvatore and Wahl [24, Section 5].

Let Com denote the operad encoding commutative algebras and $\mathcal{L}ie_1$ denote the operad encoding Lie algebras with a bracket of degree $+1$. As a consequence of a result by Markl [23, Theorem 2.7, Example 4.4], we have $\mathcal{G} \cong Com \circ \mathcal{L}ie_1$ as operads, under the distributive law

$$\Lambda: \langle -; - \bullet - \rangle \mapsto (\langle -; - \rangle \bullet -) + (- \bullet \langle -; - \rangle). \quad (12).$$

This isomorphism gives us a way to make explicit the D -operad structure on \mathcal{G} . As \mathbb{S} -modules, we have

$$Com \circ \mathcal{L}ie_1 = \bigoplus_{k=1}^n Com(t) \otimes_{\mathbb{S}_t} \left(\bigoplus_{i_1+\dots+i_k=n} \text{Ind}_{\mathbb{S}_{i_1} \times \dots \times \mathbb{S}_{i_k}}^{\mathbb{S}_n} \mathcal{L}ie_1(i_1) \otimes \dots \otimes \mathcal{L}ie_1(i_t) \right).$$

The action of δ is compatible with the symmetric action, thus it is sufficient to give the action on the elements of $Com(t) \otimes_{\mathbb{S}_t} \mathcal{L}ie_1(i_1) \otimes \dots \otimes \mathcal{L}ie_1(i_t)$. Since $Com(t)$ is given by the trivial representation on \mathbb{K} , an element in $Com(t) \otimes_{\mathbb{S}_t} \mathcal{L}ie_1(i_1) \otimes \dots \otimes \mathcal{L}ie_1(i_t)$ is of the form $L_1 \odot \dots \odot L_t$, with $L_i \in \mathcal{L}ie_1$, for $i = 1, \dots, t$; and where \odot denotes the commutative tensor product, that is the quotient of the tensor product under the permutation of terms.

Proposition 2.24 *Under the isomorphism of operads $\mathcal{G} \cong Com \circ \mathcal{L}ie_1$, the structure of D -operad on the right-hand side is given by*

$$\delta \cdot (L_1 \odot \dots \odot L_t) = \sum_{1 \leq i < j \leq n} (-1)^{\varepsilon_{i,j}} \langle L_i; L_j \rangle \odot L_1 \odot \dots \odot \widehat{L}_i \odot \dots \odot \widehat{L}_j \odot \dots \odot L_t$$

where $\langle L_i; L_j \rangle := \gamma_{\mathcal{L}ie_1}(\langle -; - \rangle; L_i, L_j)$ and where the sign $\varepsilon_{i,j}$, arising from the Koszul sign rule, is given by

$$\varepsilon_{i,j} = (|L_i| + |L_j|)(|L_1| + \dots + |L_{i-1}|) + |L_j|(|L_{i+1}| + \dots + |L_{j-1}|).$$

Proof For $t = 2$, a representative of $L_1 \odot L_2$ is given by $\bullet \otimes L_1 \otimes L_2$, with $L_1, L_2 \in \mathcal{L}ie_1$. Since the bracket is sent to 0 under the action of δ , so is any element of $\mathcal{L}ie_1$. Thus, we get

$$\delta \cdot (\bullet \otimes L_1 \otimes L_2) = \langle -; - \rangle \otimes L_1 \otimes L_2 = \langle L_1; L_2 \rangle.$$

We conclude the proof by induction on $t \geq 2$. □

Remark The action of δ on an element $L_1 \odot \cdots \odot L_t$ in $Com \circ Lie_1$ is exactly the image of $\delta^* \otimes L_1 \odot \cdots \odot L_t$ under the derivation ${}^t d_\phi$, which is given by Galvez-Carillo, Tonks and Vallette [10, Proof of Lemma 5]. In particular, δ acts as the Chevalley–Eilenberg boundary maps defining the homology of Lie algebras.

Proposition 2.25 *The category of D – \mathcal{G} –algebras is isomorphic to the category of \mathcal{BV} –algebras.*

Proof It is a straightforward consequence of Proposition 2.19 conjugated with the following isomorphism of operads proved by Salvatore and Wahl in [24]:

$$\mathcal{BV} \cong \mathcal{G} \rtimes D. \quad \square$$

Remark The algebraic structure of a Batalin–Vilkovisky algebra can be encoded in two different ways, using the operad \mathcal{BV} or the D –operad \mathcal{G} . When using the graded D –operad \mathcal{G} , the unary operator Δ is provided by the underlying category of mixed chain complexes and its relations with the product and the Lie bracket are encoded in the action of δ on those generating operations.

More generally, we can consider algebras over the homology of the framed little n –discs operad $f\mathcal{D}_n$. Batalin–Vilkovisky algebras correspond to the case $n = 2$, by Getzler [11]. In this case, Poisson n –algebras, that is algebras over the homology of the little n –discs operad \mathcal{D}_n , play the role of Gerstenhaber algebras. In the same way, it is proved in [24] that

$$f\mathcal{D}_n \cong \mathcal{D}_n \rtimes \mathrm{SO}(n),$$

as topological operads. Hence, taking the homology, we obtain that

$$H(f\mathcal{D}_n) \cong H(\mathcal{D}_n) \rtimes H(\mathrm{SO}(n)).$$

In other words, a structure of algebra over $H(f\mathcal{D}_n)$ is equivalent to a Poisson n –algebra structure in the category of $H(\mathrm{SO}(n))$ –modules.

However, when $n > 3$, the algebra $H(\mathrm{SO}(n))$ has more than one generator. So, we can choose to keep some in the operad and to put only a part of these generators in the underlying category. We still have a Hopf algebra, which acts on the operad thus obtained.

2.3.3 Differential graded algebras over an operad Let \mathcal{P} be an operad. A differential graded \mathcal{P} –algebra is a chain complex (A, d_A) endowed with a \mathcal{P} –algebra structure such that d_A is a derivation for all the operations of \mathcal{P} . We can encode this structure in the D –operad \mathcal{P} , with the trivial D –action. In particular, for $\mathcal{P} = \mathcal{A}ss$, the semidirect product $\mathcal{A}ss \rtimes D$ is isomorphic to the operad encoding differential graded algebras,

considered by Livernet, Roitzheim and Whitehouse [20] when the bicomplexes are concentrated in horizontal degree 0.

3 Relative Koszul duality theory

In this section, we develop the Koszul duality theory for operads over a cocommutative Hopf algebra H , following the method of Loday and Vallette [21, Chapter 7]. At each step, we prove that the objects can be enriched with an action of H , and we show that the results still hold in this context.

Let $(\mathcal{P}, \gamma, \eta, d_{\mathcal{P}})$ be a dg H -operad and $(\mathcal{C}, \Delta, \eta, d_{\mathcal{C}})$ be a dg H -cooperad.

3.1 Twisting morphisms

We consider the following dg \mathbb{S} - H -module

$$(\text{Hom}(\mathcal{C}, \mathcal{P}) := \{\text{Hom}_{\mathbb{K}}(\mathcal{C}(n), \mathcal{P}(n))\}_{n \geq 0}, \partial),$$

where \mathbb{S}_n acts by conjugation and where $\partial(f) = [d_{\mathcal{P}}, f] := d_{\mathcal{P}} \circ f - (-1)^{|f|} f \circ d_{\mathcal{C}}$.

Proposition 3.1 *The dg \mathbb{S} - H -module $(\text{Hom}(\mathcal{C}, \mathcal{P}), \partial)$ is a dg H -operad, called the convolution H -operad.*

Proof By Berger and Moerdijk [3, Section 1], the \mathbb{S} -module $(\text{Hom}(\mathcal{C}, \mathcal{P}), \partial)$ is an operad with respect to the composition map

$$\gamma_{\text{Hom}(\mathcal{C}, \mathcal{P})}(f; g_1, \dots, g_k) := \gamma_{\mathcal{P}} \circ (f \otimes g_1 \otimes \dots \otimes g_k) \circ \Delta_{\mathcal{C}}.$$

Moreover, in [21, Section 6.4.1], it is proved that $(\text{Hom}(\mathcal{C}, \mathcal{P}), \gamma_{\text{Hom}(\mathcal{C}, \mathcal{P})}, \partial)$ is a dg operad. The map $\gamma_{\text{Hom}(\mathcal{C}, \mathcal{P})}$ is H -equivariant as a composite of H -equivariant maps. And, since $d_{\mathcal{P}}$ and $d_{\mathcal{C}}$ are H -equivariant, then so is the differential ∂ . \square

Recall that the following composite defines a *pre-Lie product* on $\prod_{n \geq 0} \text{Hom}(\mathcal{C}, \mathcal{P})(n)$:

$$f \star g := \mathcal{C} \xrightarrow{\Delta_{(1)}} \mathcal{C} \circ_{(1)} \mathcal{C} \xrightarrow{f \circ_{(1)} g} \mathcal{P} \circ_{(1)} \mathcal{P} \xrightarrow{\gamma_{(1)}} \mathcal{P}.$$

We denote by $\text{Hom}_{H\text{-}\mathbb{S}_n}(\mathcal{C}(n), \mathcal{P}(n))$ the space of H - \mathbb{S}_n -bimodule morphisms from $\mathcal{C}(n)$ to $\mathcal{P}(n)$ and we denote the associated product of \mathbb{S} - H -equivariant maps by

$$\text{Hom}_{\mathbb{S}\text{-}H}(\mathcal{C}, \mathcal{P}) := \prod_{n \geq 0} \text{Hom}_{H\text{-}\mathbb{S}_n}(\mathcal{C}(n), \mathcal{P}(n)).$$

Proposition 3.2 *The space $(\text{Hom}_{\mathbb{S}\text{-}H}(\mathcal{C}, \mathcal{P}), \star, \partial)$ is a dg pre-Lie algebra. The associated Lie bracket induces a dg Lie algebra structure $(\text{Hom}_{\mathbb{S}\text{-}H}(\mathcal{C}, \mathcal{P}), [-, -], \partial)$.*

Proof By [21, Proposition 6.4.5, Lemma 6.4.6], it only remains to prove that if f and g are in $\text{Hom}_{\mathbb{S}\text{-}H}(\mathcal{C}, \mathcal{P})$ then so is $f \star g$. This is the case since the pre-Lie product \star is defined to be the composite of H -equivariant maps. \square

Consider the Maurer–Cartan equation in the dg-pre-Lie algebra $(\text{Hom}_{\mathbb{S}\text{-}H}(\mathcal{C}, \mathcal{P}), \star, \partial)$

$$\partial(\alpha) + \alpha \star \alpha = 0.$$

Definition 3.3 *A solution $\alpha: \mathcal{C} \rightarrow \mathcal{P}$ of degree -1 to the Maurer–Cartan equation is called a twisting H -morphism. We denote by $\text{Tw}_H(\mathcal{C}, \mathcal{P})$ the space of twisting H -morphisms from \mathcal{C} to \mathcal{P} .*

When \mathcal{C} is a coaugmented dg cooperad and \mathcal{P} is an augmented dg operad, we require that the composition of a twisting morphism with respectively the coaugmentation map or the augmentation map vanishes.

3.2 Bar and cobar constructions

Are the two functors $\text{Tw}_H(\mathcal{C}, -)$ and $\text{Tw}_H(-, \mathcal{P})$ representable?

Recall from [21, Section 6.5], that the *bar construction* of an augmented dg operad \mathcal{P} is the dg conilpotent cooperad

$$\text{B}\mathcal{P} := (\mathcal{T}^c(s\bar{\mathcal{P}}), d_{\text{B}\mathcal{P}}),$$

and that the *cobar construction* of a coaugmented dg cooperad \mathcal{C} is the dg augmented operad

$$\Omega\mathcal{C} := (\mathcal{T}(s^{-1}\bar{\mathcal{C}}), d_{\Omega\mathcal{C}}).$$

Proposition 3.4 *Let \mathcal{P} be an H -operad and \mathcal{C} be an H -cooperad. Then, the H -module structure of the free operad over an \mathbb{S} - H -module makes $\text{B}\mathcal{P}$ and $\Omega\mathcal{C}$ into a dg H -cooperad and a dg H -operad respectively.*

Proof By Proposition 2.15, the free operad $\mathcal{T}(s^{-1}\bar{\mathcal{C}})$ is an H -operad. The differential $d_{\Omega\mathcal{C}}$ is equal to the sum $d_1 + d_2$, where d_1 is the unique derivation which extends

$$\varphi: s^{-1}\bar{\mathcal{C}} \xrightarrow{\text{id} \otimes d_{\mathcal{C}}} \mathcal{T}(s^{-1}\bar{\mathcal{C}}),$$

and where d_2 is the unique derivation which extends

$$\begin{aligned} \psi: (\mathbb{K}_s^{-1} \otimes \bar{\mathcal{C}}) &\xrightarrow{\Delta_s \otimes \Delta_{(1)}} (\mathbb{K}_s^{-1} \otimes \mathbb{K}_s^{-1}) \otimes (\bar{\mathcal{C}} \circ_{(1)} \bar{\mathcal{C}}) \\ &\xrightarrow{\text{Id} \otimes \tau \otimes \text{Id}} (\mathbb{K}_s^{-1} \otimes \bar{\mathcal{C}}) \circ_{(1)} (\mathbb{K}_s^{-1} \otimes \bar{\mathcal{C}}) \cong \mathcal{T}(s^{-1}\bar{\mathcal{C}})^{(2)} \twoheadrightarrow \mathcal{T}(s^{-1}\bar{\mathcal{C}}). \end{aligned}$$

The result [21, Proposition 6.3.15] extends to graded \mathbb{S} - H -modules. Indeed, if E is an \mathbb{S} - H -module and $\alpha: E \rightarrow \mathcal{T}(E)$ is a morphism of \mathbb{S} - H -modules, the unique derivation which extends α is a composite of H -equivariant maps. Thus, since φ and ψ are morphisms of \mathbb{S} - H -modules, the derivations d_1 and d_2 , and hence $d_{\Omega\mathcal{C}}$, are H -equivariant maps.

In the same way, we prove that $B\mathcal{P}$ is a dg H -cooperad. □

Proposition 3.5 *The bar and cobar constructions form a pair of adjoint functors*

$$\Omega: \{\text{conil. dg } H\text{-cooperads}\} \rightleftarrows \{\text{aug. dg } H\text{-operads}\} : B,$$


such that the adjunction is given by the set of twisting H -morphisms. That is for every augmented dg H -operad \mathcal{P} and every conilpotent dg H -cooperad \mathcal{C} , there exist natural bijections

$$\text{Hom}_{\text{dgOp}_H}(\Omega\mathcal{C}, \mathcal{P}) \cong \text{Tw}_H(\mathcal{C}, \mathcal{P}) \cong \text{Hom}_{\text{dgCoop}_H}(\mathcal{C}, B\mathcal{P}).$$

Proof It is a consequence of [21, Theorem 6.5.10] and of the definitions of the free H -operad and the cofree H -cooperad over an \mathbb{S} - H -module. □

We denote by $\eta_{\mathcal{C}}: \mathcal{C} \rightarrow \Omega\mathcal{C}$ and $\pi_{\mathcal{P}}: B\mathcal{P} \rightarrow \mathcal{P}$ the universal twisting H -morphisms corresponding respectively to the unit $v: \mathcal{C} \rightarrow B\Omega\mathcal{C}$ and to the counit $\epsilon: \Omega B\mathcal{P} \rightarrow \mathcal{P}$ of the adjunction.

Proposition 3.6 *Any twisting H -morphism $\alpha: \mathcal{C} \rightarrow \mathcal{P}$ factors uniquely through the universal twisting morphisms $\pi_{\mathcal{P}}$ and $\eta_{\mathcal{C}}$ as follows:*



where g_{α} is a morphism of dg H -operads and f_{α} is a morphism of dg H -cooperads.

Proof It is a consequence of the adjunction given in Proposition 3.5. □

When dealing with operads in the monoidal category of dg \mathbb{S} -modules, a twisting morphism α in $\text{Tw}(\mathcal{C}, \mathcal{P})$ is called *Koszul* when a certain chain complex, called the *Koszul complex*, is acyclic; see [21, Section 6.6.1].

Definition 3.7 A twisting H -morphism $\alpha \in \text{Tw}_H(\mathcal{C}, \mathcal{P})$ is said to be *Koszul* when, seen as a morphism of dg \mathbb{S} -modules, it is a Koszul morphism.

So, the property for a twisting morphism to be Koszul is a homological property and only depends on the differential structures of \mathcal{C} and \mathcal{P} . By definition, to be a quasi-isomorphism is also a homological property, so [21, Theorem 6.6.2] extends to H -operads to give the following result.

Theorem 3.8 Let \mathcal{P} be a connected weight graded dg H -operad and let \mathcal{C} be a connected weight graded dg H -cooperad. Let $\alpha: \mathcal{C} \rightarrow \mathcal{P}$ be a twisting H -morphism. The following assertions are equivalent:

- (1) The twisting morphism α is Koszul.
- (2) The morphism of dg H -cooperads $f_\alpha: \mathcal{C} \rightarrow \text{B}\mathcal{P}$ is a quasi-isomorphism.
- (3) The morphism of dg H -operads $g_\alpha: \Omega\mathcal{C} \rightarrow \mathcal{P}$ is a quasi-isomorphism.

Theorem 3.9 The counit $v: \Omega\text{B}\mathcal{P} \rightarrow \mathcal{P}$ and the unit $\epsilon: \mathcal{C} \rightarrow \text{B}\Omega\mathcal{C}$ of the adjunction are H -quasi-isomorphisms of dg H -operads and dg H -cooperads respectively.

The resolution $\Omega\text{B}\mathcal{P}$ is called the *bar-cobar resolution* of \mathcal{P} .

3.3 Koszul duality of H -operads

Definition 3.10 A *quadratic H -datum* consists of a pair (E, R) made up of a graded \mathbb{S} - H -module E and a graded sub- \mathbb{S} - H -module R of $\mathcal{T}(E)$ ⁽²⁾ called respectively the *generating operations* and the *relations*.

The *quadratic H -operad* $\mathcal{P}(E, R)$ associated to a quadratic H -datum (E, R) is the quotient H -operad of $\mathcal{T}(E)$, which satisfies the following universal property: for any H -operad \mathcal{P} of $\mathcal{T}(E)$ such that the composite of morphisms of \mathbb{S} - H -modules

$$R \twoheadrightarrow \mathcal{T}(E) \twoheadrightarrow \mathcal{P}$$

is trivial, there exists a unique morphism of H -operads which makes the following diagram commutative:

$$\begin{array}{ccccc}
 R & \longrightarrow & \mathcal{T}(E) & \longrightarrow & \mathcal{P} \\
 & & & \searrow & \uparrow \\
 & & & & \mathcal{P}(E, R)
 \end{array}$$

Dually, we define the *quadratic H -cooperad* $\mathcal{C}(E, R)$ associated to a quadratic H -datum (E, R) .

Proposition 3.11 *Let (E, R) be a quadratic H -datum. The quadratic H -operad (resp. H -cooperad) associated to (E, R) is given by the quadratic operad (resp. cooperad) associated to (E, R) in the category of \mathbb{S} -modules, endowed with the H -module structure induced by the one on the \mathbb{S} - H -module $\mathcal{T}(E)$.*

Proof We only prove this result for the quadratic H -operad. In the category of operads, $\mathcal{P}(E, R)$ is given by the quotient $\mathcal{T}(E)/(R)$ of the free operad on E by the ideal generated by R . By Proposition 2.16, $\mathcal{T}(E)/(R)$ is an H -operad and, by assumptions on R and on \mathcal{P} , the morphism of operads $\mathcal{P}(E, R) \rightarrow \mathcal{P}$ turns out to be an H -equivariant map. □

We can extend the definition of the Koszul dual (co)operad, given in [21, Section 7.2], to the category of \mathbb{S} - H -modules. In the classical case, recall that the *Koszul dual cooperad* of quadratic operad \mathcal{P} is defined by the cooperad

$$\mathcal{P}^i := \mathcal{C}(sE, s^2R),$$

and the *Koszul dual operad* of \mathcal{P} by the operad

$$\mathcal{P}^! := (\mathcal{S}^c \otimes_{\mathcal{H}} \mathcal{P}^i)^*,$$

where $*$ denotes the aritywise linear dual; see Section 2.2.5.

Proposition 3.12 *If \mathcal{P} is a quadratic H -operad, then the Koszul dual cooperad (resp. operad) of \mathcal{P} is an H -cooperad (resp. H -operad).*

Proof By definition, \mathcal{P}^i is an H -cooperad. Since $\Delta_{\text{End}_{\mathbb{K},s}}$ is H -equivariant, the H -operad structure on $\mathcal{P}^!$ follows from Proposition 2.21 and Section 2.2.5. □

Proposition 3.13 *Let (E, R) be a quadratic H -datum such that E is a finite dimensional \mathbb{S} - H -module. The Koszul dual operad $\mathcal{P}^!$ of the quadratic operad $\mathcal{P} = \mathcal{P}(E, R)$ admits the quadratic presentation*

$$\mathcal{P}^! \cong \mathcal{P}(s^{-1} \mathcal{S}^{-1} \otimes_{\mathcal{H}} E^*, R^\perp),$$

where R^\perp denotes the sub- \mathbb{S} - H -module obtained by proper suspension of the operations indexing the vertices of the trees of the orthogonal module $(s^2 R)^\perp \subset \mathcal{T}(s^{-1} E^*)^{(2)}$, which is the image of $(s^2 R)^*$ under the isomorphism $(\mathcal{T}(E)^{(2)})^* \cong \mathcal{T}(s^{-1} E^*)^{(2)}$.

Proof By [21, Proposition 7.2.4], this result is true in the category of operads. It is left to prove that the isomorphism is an H -equivariant map: the restriction of this isomorphism to the space of generators $s^{-1} \mathcal{S}^{-1} \otimes_{\mathcal{H}} E^*$ is H -equivariant, thus so is the isomorphism. □

Proposition 3.14 *Let (E, R) be a quadratic H -datum and $\mathcal{P} = \mathcal{P}(E, R)$ be its associated quadratic H -operad. The natural H -cooperad inclusion $i: \mathcal{P}^i \hookrightarrow B\mathcal{P}$ induces an isomorphism of graded H -operads*

$$i: \mathcal{P}^i \xrightarrow{\cong} H^0(B^*\mathcal{P}),$$

where $H^\bullet(B^*\mathcal{P})$ denotes the cohomology groups of the syzygy degree cochain complex associated to $B\mathcal{P}$; see [21, Section 7.3.1].

Proof By [21, Proposition 7.3.2], we have an isomorphism of operads and the inclusion $\mathcal{P}^i \hookrightarrow B^0\mathcal{P}$ is exactly the kernel of the differential defining $H^\bullet(B^*\mathcal{P})$, that is $H^0(B^*\mathcal{P})$. Since this differential is H -equivariant, this isomorphism is compatible with the H -module structure. □

Lemma 3.15 *Let (E, R) be a quadratic H -datum. Then, the composite*

$$\kappa: \mathcal{C}(sE, s^2 R) \twoheadrightarrow sE \xrightarrow{s^{-1}} E \hookrightarrow \mathcal{P}(E, R)$$

associated to (E, R) , is a twisting H -morphism.

Proof It is straightforward from [21, Lemma 7.4.2], since κ is a composite of H -equivariant maps. □

Theorem 3.16 (Koszul criterion) *Let (E, R) be a quadratic H -datum and $\mathcal{P} = \mathcal{P}(E, R)$ be its associated quadratic H -operad. The following statements are equivalent:*

- (1) *The inclusion $i: \mathcal{P}^i \hookrightarrow \text{B}\mathcal{P}$ is a quasi-isomorphism of dg H -cooperads.*
- (2) *The projection $p: \Omega\mathcal{P}^i \twoheadrightarrow \mathcal{P}$ is a quasi-isomorphism of dg H -operads.*

When these propositions hold, we say that \mathcal{P} is a Koszul operad.

Proof This follows from [Theorem 3.8](#). □

3.4 Example

Let us now focus on the example of Batalin–Vilkovisky algebras.

Getzler and Jones [13] proved that the operad \mathcal{G} is Koszul. Then, by the present definition, so is the D -operad \mathcal{G} .

Lemma 3.17 *The action of D on \mathcal{G}^i is induced by*

$$\begin{aligned} \delta \cdot \mathcal{G}^i &\twoheadrightarrow \mathbb{K}s \bullet \oplus \mathbb{K}s \langle -; - \rangle, \\ s \bullet &\mapsto -s \langle -; - \rangle, \\ s \langle -; - \rangle &\mapsto 0. \end{aligned}$$

Proof By definition, the action of D is characterized by its value on the cogenerators of \mathcal{G}^i , which is deduced from the one on \mathcal{G} . □

Proposition 3.18 *Under the isomorphism of cooperads $\mathcal{G}^i \cong S^{-1c} \text{Com}_1^c \circ S^{-1c} \mathcal{L}ie^c$, the structure of D -cooperad on the right-hand side is given by*

$$\delta \cdot (L_1 \odot \cdots \odot L_t) = \sum_{k=1}^n (-1)^{\varepsilon_k} L_1 \odot \cdots \odot L'_k \odot L''_k \odot \cdots \odot L_t,$$

where $L'_k \odot L''_k$ is the sumless Sweedler’s notation for the image of L_k under the binary part of $\Delta_{S^{-1c} \mathcal{L}ie^c}$, that is $\Delta_{S^{-1c} \mathcal{L}ie^c}: S^{-1c} \mathcal{L}ie^c \rightarrow S^{-1c} \mathcal{L}ie^c(2) \otimes_{\mathbb{S}_2} (S^{-1c} \mathcal{L}ie^c \otimes S^{-1c} \mathcal{L}ie^c)$ and

$$\varepsilon_k = |L_k| + \cdots + |L_t| + 1.$$

Proof Recall that, for any S – H –module M , $S^{-1^c}M$ (resp. SM) denotes the Hadamard product of S^{-1^c} (resp. S) with M . To make explicit the action of δ , we use the D –operad structure on $(\mathcal{G}^i)^* \cong \mathcal{S}Com_{-1} \circ \mathcal{S}Lie$, which is given by the isomorphism of operads $(\mathcal{G}^i)^* \cong \mathcal{S}^2\mathcal{G} \cong \mathcal{S}^2(Com \circ Lie_1)$. We have

$$\delta \cdot (L_1 \odot \cdots \odot L_t) = \delta \cdot (L_1^* \odot \cdots \odot L_t^*)^*,$$

with $L_1^* \odot \cdots \odot L_t^* \in \mathcal{S}Com_{-1} \circ \mathcal{S}Lie$. The only elements in $\mathcal{S}Com_{-1} \circ \mathcal{S}Lie$ whose images under $\delta \cdot (L_1^* \odot \cdots \odot L_t^*)^*$ are nonzero are of the form

$$L_1^* \odot \cdots \odot (L'_k)^* \odot (L''_k)^* \odot \cdots \odot L_t^*, \quad k \in \{1, \dots, t\},$$

where $(L'_k)^*, (L''_k)^*$ are elements of $\mathcal{S}Lie$ such that $L_k^* = \gamma_{\mathcal{S}Lie}(s^{-1}[-; -]; (L'_k)^*, (L''_k)^*)$. This image is equal to $(-1)^{\varepsilon_k + 1}$, where $\varepsilon_k = |L_k^*| + \cdots + |L_t^*|$. \square

Remark Dually to the D –operad structure on \mathcal{G} , the D –cooperad structure on \mathcal{G}^i is equal, up to sign, to the image of an element in $\mathbb{K}[\delta]_1 \circ S^{-1^c}Com_1^c \circ S^{-1^c}Lie^c$ under the coderivation d_φ , given by Galvez-Carillo, Tonks and Vallette [10, Lemma 5].

4 Homotopy algebras and transfer theorem

In this section, we extend some definitions and results of [21, Section 10] to the category of H –modules, requiring in addition the compatibility with the H –module structure.

4.1 The category of homotopy algebras

Let \mathcal{P} be a Koszul H –operad. By definition, a *homotopy H – \mathcal{P} –algebra*, or an *H – \mathcal{P}_∞ –algebra*, is an algebra over the Koszul resolution $\mathcal{P}_\infty := \Omega\mathcal{P}^i$ of \mathcal{P} in the category of H –operads. So, a structure of algebra over \mathcal{P}_∞ on an H –module A is equivalent to a morphism of H –operads $\mathcal{P}_\infty := \Omega\mathcal{P}^i \rightarrow \text{End}_A$. Notice that a H – \mathcal{P}_∞ –algebra is a \mathcal{P}_∞ –algebra endowed with a compatible H –module structure.

Example Any H – \mathcal{P} –algebra structure on an H –module is a particular case of a H – \mathcal{P}_∞ –algebra structure.

Proposition 4.1 *The set of H – \mathcal{P}_∞ –algebra structures is equivalently given by*

$$\begin{aligned} \text{Hom}_{\text{dg}H\text{-Op}}(\Omega\mathcal{P}^i, \text{End}_A) &\cong \text{Tw}_H(\mathcal{P}^i, \text{End}_A) \\ &\cong \text{Hom}_{\text{dg}H\text{-coOp}}(\mathcal{P}^i, \mathbf{B}\text{End}_A) \cong \text{Codiff}_H(\mathcal{P}^i(A)), \end{aligned}$$

where $\text{Codiff}_H(\mathcal{P}^i(A))$ denotes the set of codifferentials on $\mathcal{P}^i(A)$, that are H –equivariant coderivations on $\mathcal{P}^i(A)$ squaring to zero.

Proof The first two bijections are given by the adjunction of Proposition 3.5. The proof of Proposition 2.18 implies that

$$\text{Hom}_{\mathbb{S}\text{-}H\text{-Mod}}(\mathcal{P}^i, \text{End}_A) \cong \text{Hom}_H(\mathcal{P}^i(A), A).$$

Moreover, [21, Proposition 6.3.17] extends to H -modules. So, if \mathcal{C} is a dg H -cooperad, V is an H -module and $\alpha: \mathcal{C}(V) \rightarrow V$ is H -equivariant, then the unique coderivation of the cofree \mathcal{C} -coalgebra $\mathcal{C}(V)$ which extends α is given by a sum of composites of H -equivariant maps. Thus, we get the isomorphism

$$\text{Hom}_H(\mathcal{P}^i(A), A) \cong \text{Coder}_H(\mathcal{P}^i(A)).$$

We conclude with [21, Proposition 10.1.19], that an element in $\text{Hom}_{\mathbb{S}\text{-}H\text{-Mod}}(\mathcal{P}^i, \text{End}_V)$ is a solution to the Maurer–Cartan equation if and only if the associated coderivation on $\mathcal{P}^i(V)$ squares to zero, which does not depend on the H -module structure. \square

This result provides us with four equivalent definitions of $H\text{-}\mathcal{P}_\infty$ -algebra structures. Thus, when dealing with this algebraic structure, we can make an ad hoc choice of one of those definitions.

By extension of the classical definition of [21, Section 10.2.2], an ∞ - H -morphism of $H\text{-}\mathcal{P}_\infty$ -algebras is a morphism $A \rightsquigarrow B$ of dg $H\text{-}\mathcal{P}^i$ -coalgebras

$$F: (\mathcal{P}^i(A), d_\varphi) \rightarrow (\mathcal{P}^i(B), d_\psi),$$

where φ and ψ are the twisting H -morphisms defining the structure of $H\text{-}\mathcal{P}_\infty$ -algebra on A and B respectively. We denote by $\infty\text{-}\mathcal{P}_\infty\text{-}H\text{-alg}$ the category of $H\text{-}\mathcal{P}_\infty$ -algebras and their ∞ - H -morphisms, where the composite of two ∞ - H -morphisms is defined as the composite of the associated morphisms of dg $H\text{-}\mathcal{P}^i$ -coalgebras.

Equivalently, an ∞ - H -morphism of $H\text{-}\mathcal{P}_\infty$ -algebras is an H -equivariant ∞ -morphism of the underlying \mathcal{P}_∞ -algebras. A ∞ - H -morphism of $H\text{-}\mathcal{P}_\infty$ -algebras is called an ∞ - H -isomorphism (resp. ∞ - H -quasi-isomorphism) if its first component $A \rightarrow B$ is an H -isomorphism (resp. quasi-isomorphism).

4.2 Homotopy transfer theorem

Let (B, d_B) be a homotopy retract of (A, d_A) in the category of H -modules, that is a homotopy retract of chain complexes

$$h \begin{array}{c} \hookrightarrow \\ \circlearrowleft \\ \hookrightarrow \end{array} (A, d_A) \begin{array}{c} \xrightarrow{p} \\ \xleftarrow{i} \end{array} (B, d_B),$$

$$\text{Id}_A - ip = d_A h + h d_A,$$

such that the maps i , p , h are H -equivariant and i is a quasi-isomorphism.

Theorem 4.2 *Let \mathcal{P} be a Koszul H -operad. Let (B, d_B) be a homotopy retract of (A, d_A) in the category of dg - H -modules.*

Any H - \mathcal{P}_∞ -algebra structure on A , defined by generating operations $\{m_\mu: A^{\otimes n} \rightarrow A, \mu \in \mathcal{P}_\infty\}$, can be transferred into a H - \mathcal{P}_∞ -algebra structure on B , which extends the transferred operations $pm_\mu i^{\otimes n}: B^{\otimes n} \rightarrow B$, and such that i extends to an ∞ - H -quasi-isomorphism.

Proof We check each step of the proof of [21, Theorem 10.3.2] extends to H -modules. Since the maps h , i and p are H -equivariant, then the map $\Psi: B \text{ End}_A \rightarrow B \text{ End}_B$, defined in [21, Section 10.3.3], is a morphism of H -cooperads. To prove that Ψ is compatible with the differential structure, we do not care about the H -module structure so it follows from Van der Laan [18, Theorem 5.2]. Then, as a consequence of the bar-cobar adjunction, the composite $\mathcal{P}^i \rightarrow B \text{ End}_A \rightarrow B \text{ End}_B$ defines an H - \mathcal{P}_∞ -algebra structure on B . Similarly, to be an ∞ - H -quasi-isomorphism does not depend on the H -module structure therefore, by [21, Theorem 10.3.11], we just have to prove that the map i_∞ , defined there, commutes with the H -module structure. This is the case since it is a composite of such maps. \square

In particular, the transferred structure and the ∞ - H -quasi-isomorphism are both given by the explicit tree-wise formulae of [21, Section 10.3.10, Theorem 10.3.6].

Theorem 4.3 *Let \mathcal{P} be a Koszul H -operad. Let (A, d_A) be a $\mathcal{P} \rtimes H$ -algebra and (B, d_B) be a homotopy retract of (A, d_A) in the category of dg - H -modules. Then, B inherits a structure of H - \mathcal{P}_∞ -algebra, which extends the transferred operations, and such that i extends to an ∞ -quasi-isomorphism of H - \mathcal{P}_∞ -algebras.*

Proof By Proposition 2.19, the $\mathcal{P} \rtimes H$ -algebra structure on A is equivalent to an H - \mathcal{P} -algebra structure. Then, we apply Theorem 4.2. \square

Remark When the operad $\mathcal{P} \rtimes H$ is Koszul, one can apply the HTT for operads, but the $(\mathcal{P} \rtimes H)_\infty$ -algebra structure can be very complex. If we can do the transfer in the category of H -modules, then the transferred structure reduces to a much simpler structure. On the other hand, when the operad $\mathcal{P} \rtimes H$ is not Koszul, the classical method to transfer an algebraic structure must be improved. While, if the H -operad \mathcal{P} is itself Koszul, then we can apply our version of the HTT, assuming that the homotopy retract is compatible with the H -action.

4.3 Homotopy theory for algebras over an H -operad

The results of [21, Chapter 11] extend *mutatis mutandis* to the framework of H -operads. The objects and the maps defined there are compatible with the H -module structure. For the homological considerations, the results still hold because they do not depend on the H -action. In particular, we obtain the following results.

Theorem 4.4 (Rectification) *Let \mathcal{P} be a Koszul H -operad. Any H - \mathcal{P}_∞ -algebra is naturally ∞ - H -quasi-isomorphic to a dg H - \mathcal{P} -algebra.*

Theorem 4.5 (Equivalence between homotopy categories) *Let \mathcal{P} be a Koszul H -operad. The homotopy category of dg H - \mathcal{P} -algebras is equivalent to the homotopy category of H - \mathcal{P} -algebras with their ∞ - H -morphisms.*

Remark In particular, we obtain a new description for the homotopy category of $\mathcal{P} \rtimes H$ -algebras, using a simpler class of ∞ -morphisms than [21, Theorem 11.4.12].

An advantage of this result lies in the “invertibility” of the ∞ - H -quasi-isomorphisms. Indeed, any ∞ - H -quasi-isomorphism admits an ∞ - H -quasi-isomorphism in the opposite direction, as in [21, Theorem 10.4.7], while it is not the case for H -quasi-isomorphisms. This enables us to prove the following result.

Theorem 4.6 (H -quasi-isomorphism vs ∞ - H -quasi-isomorphism) *Let \mathcal{P} be a Koszul H -operad and A, B be two dg H - \mathcal{P} -algebras. The following assertions are equivalent:*

- (1) *There exists a zigzag of H -quasi-isomorphisms of dg H - \mathcal{P} -algebras:*

$$A \xleftarrow{\sim} \bullet \xrightarrow{\sim} \bullet \xleftarrow{\sim} \bullet \cdots \bullet \xrightarrow{\sim} B$$

- (2) *There exists an ∞ - H -quasi-isomorphism of dg H - \mathcal{P} -algebras $A \xrightarrow{\sim} B$.*

Remark Since an H - \mathcal{P} -algebra structure is equivalent to a $\mathcal{P} \rtimes H$ -algebra one, the theorem gives a new and simpler way to prove formality results for $\mathcal{P} \rtimes H$ -algebras.

4.4 Example

When we encode BV-algebra structures with the operad \mathcal{BV} , a notion of homotopy BV-algebra structure is given by Galvez-Carillo, Tonks and Vallette [10]. We describe the homotopy structure obtained when we use the D -operad \mathcal{G} instead, and compare these two.

Proposition 4.7 A homotopy D - \mathcal{G} -algebra is a mixed chain complex (A, d_A, Δ_A) endowed with a homotopy Gerstenhaber algebra structure such that

$$\Delta_A m_{p_1, \dots, p_t} - (-1)^{t-2} \sum_{i=1}^n m_{p_1, \dots, p_t} \circ_i \Delta_A = \sum_{k=1}^t \sum_{p'_k + p''_k = p_k} (-1)^{\varepsilon_k + 1} m_{p_1, \dots, p'_k, p''_k, \dots, p_t},$$

where $n = p_1 + \dots + p_t$ and where $\Delta_A: A \rightarrow A$ is the unary operator provided by the D -module structure on A .

Proof By Proposition 4.1, a structure of D - \mathcal{G}_∞ -algebra on a mixed chain complex A is given by a twisting D -morphism $\mathcal{G}^i \rightarrow \text{End}_A$. It is a twisting morphism $\mathcal{G}^i \rightarrow \text{End}_A$ between the underlying \mathbb{S} -modules, that is a homotopy Gerstenhaber structure as defined in [10, Proposition 16], which commutes with the action of D . This condition is then a consequence of Proposition 3.18, according to the study of homotopy Gerstenhaber algebras done in [10, Section 2.1]. \square

As conjectured at the end of [10, Section 2.4], we have the following relation with the homotopy BV-algebras.

Proposition 4.8 A homotopy BV-algebra structure (as defined in [10, Theorem 20]), such that all the operations are trivial except the maps m_{p_1, \dots, p_t}^0 and the map m_1^1 , is equivalent to a D - \mathcal{G}_∞ -algebra.

Proof Looking at the explicit description of the homotopy BV-algebra structure given in [10, Theorem 20], all the relations become trivial except the relations R_{p_1, \dots, p_t}^0 , defining the homotopy Gerstenhaber structure, the relations R_{p_1, \dots, p_t}^1 , corresponding to those given in Proposition 4.7, and the relation R_1^2 , which means that m_1^1 squares to zero. \square

From now on, we call a D - \mathcal{G}_∞ -algebra a *strict homotopy BV-algebra*, since a D - \mathcal{G}_∞ -algebra is a homotopy BV-algebra such that the operator Δ strictly satisfies the relations of a BV operator.

Remark A strict homotopy BV-algebra is a homotopy BV-algebra such that any elements in \mathcal{BV}^i (as defined in [10, Section 1]) containing a vertex decorated by Δ , except Δ itself, acts as zero. If we ask, moreover, that Δ acts as zero, we get the notion of *strongly trivialized homotopy BV-algebras*, defined by Drummond-Cole [8]. In that article, the author proved that any strongly trivialized homotopy BV structure on a chain complex induces a structure of homotopy hypercommutative algebra; see Drummond-Cole and Vallette [9], on that complex. Thus, we get another description of the homotopy hypercommutative structure.

Theorem 4.9 *Let (B, d_B, Δ_B) be a homotopy retract of (A, d_A, Δ_A) in the category of mixed chain complexes. If A is endowed with a BV-algebra structure (or with a strict homotopy BV-algebra structure), then B inherits a structure of strict homotopy BV-algebra which extends the naive transferred operations, and such that i extends to an ∞ - D -quasi-isomorphism of strict homotopy BV-algebras.*

Proof By Proposition 2.25, A is a D - \mathcal{G} -algebra. Then we apply Theorem 4.2. \square

When the homotopy retract lives in the category of mixed chain complexes, this theorem shows that the transferred homotopy BV-algebra structure reduces to a strict homotopy BV-algebra, ie a homotopy BV-algebra structure without higher homotopies arising from Δ and its relations with the product and with the bracket.

Proposition 4.10 *Let (B, d_B) be a homotopy retract of (A, d_A) , in the category of chain complexes, which satisfies the following side conditions:*

$$hi = ph = 0.$$

If $(A, d_A, \bullet, \langle -; - \rangle, \Delta)$ is a BV-algebra structure on A such that

$$\Delta h + h\Delta = 0,$$

then B inherits a strict homotopy BV-algebra structure, induced by the BV-algebra structure on A , and such that i extends to an ∞ - D -quasi-isomorphism of strict homotopy BV-algebras.

Proof We endow the chain complex (B, d_B) with the unary operator $\tilde{\Delta}$ induced by Δ , and defined by $\tilde{\Delta} := p\Delta i$. The condition $\Delta h + h\Delta = 0$ implies that $\tilde{\Delta}^2 = 0$ so $(B, d_B, \tilde{\Delta})$ is a mixed chain complex. Moreover, the side conditions imply that the maps p and i are compatible with $\tilde{\Delta}$. Thus, we have a homotopy retract between (A, d_A, Δ) and $(B, d_B, \tilde{\Delta})$ in the category of mixed chain complexes. We conclude by applying Theorem 4.9. \square

This proposition is the result which answers the main question raised at the beginning of this paper.

When B consists of the homology groups $H^\bullet(A, d)$ of the complex (A, d) , it is always possible to build a homotopy retract which satisfies the side conditions: it is called the *Hodge decomposition*; see Loday and Vallette [21, Lemma 9.4.7]. More precisely, the chain complex (A, d) splits into a direct sum of graded spaces as

$$A = H \oplus B \oplus C,$$

where $B = \text{Ker}(d) \cap \text{Im}(d)$, $H \oplus B = \text{Ker}(d)$ and $H \cong H^\bullet(A, d)$. Then, the homotopy retract is defined as follows:


- i is the inclusion of H in A .
- p is the projection of A onto H .
- h is equal to 0 on $H \oplus C$ and is equal to d^{-1} on B .

In this case, the homotopy retract satisfies the aforementioned side conditions.

Example We consider the following nonunital dg commutative algebra A generated by

$$x_3, y_3, t_3, \xi_4, \omega_5, z_7, u_7 \text{ and } v_8,$$


where the subscript gives the homological degree, such that the product by u , by v , by ξ and by ω is equal to zero. The algebra A is finite dimensional and spanned by the elements $x, y, t, \xi, \omega, xy, xt, yt, u, z, v, xz, yz, tz, xyz, xyt$ and $xyzt$. We summarize in the following picture the definition of the differential and of the BV operator on the generators:



where an element is sent to 0 if nothing else is specified.

Proposition 4.11 *The aforementioned algebra (A, d, Δ) is a dg BV-algebra. Moreover, its homology $H^\bullet(A, d)$ is endowed with a strict homotopy BV-algebra structure, such that the inclusion $H^\bullet(A, d) \hookrightarrow A$ extends to an ∞ -quasi-isomorphism of strict homotopy BV-algebras.*

Proof It is straightforward to prove that A together with d and Δ forms a dg BV-algebra. The Hodge decomposition of A is given by



where an element is sent to 0 if nothing else is specified. The contracting homotopy and Δ commute. Then, we apply [Proposition 4.10](#). \square

Note that the product induced in homology is associative, since the differential on $H^\bullet(A, d)$ is 0, but the first homotopy for the associativity relation, given by

$$m_3^0 := p((h(i(-)) \bullet i(-)) \bullet i(-)) - p(i(-) \bullet (h(i(-)) \bullet i(-)))$$

is not equal to 0. Moreover, the BV operator does not vanish in homology.

Remark Introduced by Deligne, Griffiths, Morgan and Sullivan [\[7\]](#) to study the differential forms of compact Kähler manifolds and used by Barranikov and Kontsevich [\[1\]](#) for the Dolbeault complex of Calabi–Yau manifolds, the dd^c -lemma implies the condition $[\Delta, h] = 0$, but is a strong condition in our setting. Indeed, if the dd^c -lemma is satisfied, then the operator Δ vanishes on the homology groups.

Encoding the BV-algebra structure in the D -operad \mathcal{G} allows us to deal with a notion of homotopy BV-algebra which is simpler than the one given by Galvez-Carillo, Tonks and Vallette [\[10\]](#). The associated notion of ∞ -morphism is also simplified.

Proposition 4.12 *An ∞ - D -morphism of (strict homotopy) BV-algebras is an ∞ -morphism of the underlying (homotopy) Gerstenhaber algebras which commutes with the extra action of the unary operator, provided by the (strict homotopy) BV-algebra structures.*

Proof It is a consequence of the definition of an ∞ -morphism in the category of D - \mathcal{G}_∞ -algebras. \square

Corollary 4.13 *The homotopy category of BV-algebras is equivalent to the homotopy category of strict homotopy BV-algebras with their ∞ -morphisms. Moreover, for any BV-algebras A and B , the following assertions are equivalent:*

- (1) *There exists a zigzag of quasi-isomorphisms of BV-algebras:*

$$A \xleftarrow{\sim} \bullet \xrightarrow{\sim} \bullet \xleftarrow{\sim} \bullet \cdots \bullet \xrightarrow{\sim} B$$

- (2) *There exists an ∞ -quasi-isomorphism of Gerstenhaber algebras $A \xrightarrow{\sim} B$, commuting with the unary operator provided by the BV-algebra structures.*

Proof Since the operad \mathcal{G} is Koszul, we can apply [Theorem 4.5](#) and [Theorem 4.6](#), combined with [Proposition 4.12](#). \square

This theorem provides a new way to prove the existence of a chain of quasi-isomorphisms between BV-algebras. In particular, it could help to prove the formality of a BV-algebra. Moreover, we could hope to use it to study the interpretation of the mirror symmetry conjecture of Kontsevich [17] in terms of BV-algebras.

The mirror symmetry conjecture says that a certain A-type theory of a Calabi–Yau manifold can be identified with a B-type theory on a mirror manifold. Cao and Zhou [4] interpret this conjecture as the existence of a quasi-isomorphism of dg BV-algebras from the De Rham complex of a Calabi–Yau manifold \mathcal{M} to the Dolbeault complex of a dual Calabi–Yau manifold $\widetilde{\mathcal{M}}$:

$$(\Omega^{n-\bullet}(\mathcal{M}), d_{DR}, \wedge, \Delta, \langle -; - \rangle) \xrightarrow{\sim} (\Gamma(\widetilde{\mathcal{M}}, \wedge^{\bullet} \bar{T}_{\widetilde{\mathcal{M}}}^* \otimes \wedge^{\bullet} T_{\widetilde{\mathcal{M}}}), \bar{\partial}, \wedge, \text{div}, \langle -; - \rangle_S).$$

Using Corollary 4.13, it is equivalent to the existence of an ∞ -quasi-isomorphism of the underlying Gerstenhaber algebras, which commutes strictly with the action of the unary operators Δ and div .

Acknowledgments This paper constitutes a part of the author’s PhD thesis. I would like to thank my advisor, Bruno Vallette, for being so receptive and involved. I am grateful for our instructive discussions and for his fruitful ideas. I also would like to thank Joan Millès for his reactivity to my many questions and Jim Stasheff for his comments. The author is supported by the ANR grant HOGT, referenced as ANR-11-BS01-0002.

References

- [1] **S Barannikov, M Kontsevich**, *Frobenius manifolds and formality of Lie algebras of polyvector fields*, Internat. Math. Res. Notices (1998) 201–215 [MR1609624](#)
- [2] **IA Batalin, G A Vilkovisky**, *Gauge algebra and quantization*, from: “Quantum gravity”, (M A Markov, P C West, editors), Plenum, New York (1984) 463–480 [MR746994](#)
- [3] **C Berger, I Moerdijk**, *Axiomatic homotopy theory for operads*, Comment. Math. Helv. 78 (2003) 805–831 [MR2016697](#)
- [4] **H-D Cao, J Zhou**, *DGBV algebras and mirror symmetry*, from: “First International Congress of Chinese Mathematicians”, (L Yang, S-T Yau, editors), AMS/IP Stud. Adv. Math. 20, Amer. Math. Soc. (2001) 279–289 [MR1830184](#)
- [5] **H-D Cao, J Zhou**, *On quasi-isomorphic DGBV algebras*, Math. Ann. 326 (2003) 459–478 [MR1992272](#)
- [6] **M Chas, D Sullivan**, *String topology* [arXiv:9911159](#)
- [7] **P Deligne, P Griffiths, J Morgan, D Sullivan**, *Real homotopy theory of Kähler manifolds*, Invent. Math. 29 (1975) 245–274 [MR0382702](#)

- [8] **G C Drummond-Cole**, *Formal formality of the hypercommutative algebras of low dimensional Calabi–Yau varieties* [arXiv:1201.6111](#)
- [9] **G C Drummond-Cole**, **B Vallette**, *The minimal model for the Batalin–Vilkovisky operad*, *Selecta Math.* 19 (2013) 1–47 [MR3029946](#)
- [10] **I Gálvez-Carrillo**, **A Tonks**, **B Vallette**, *Homotopy Batalin–Vilkovisky algebras*, *J. Noncommut. Geom.* 6 (2012) 539–602 [MR2956319](#)
- [11] **E Getzler**, *Batalin–Vilkovisky algebras and two-dimensional topological field theories*, *Comm. Math. Phys.* 159 (1994) 265–285 [MR1256989](#)
- [12] **E Getzler**, *Two-dimensional topological gravity and equivariant cohomology*, *Comm. Math. Phys.* 163 (1994) 473–489 [MR1284793](#)
- [13] **E Getzler**, **J D S Jones**, *Operads, homotopy algebra and iterated integrals for double loop spaces* [arXiv:9403055](#)
- [14] **V Ginzburg**, **M Kapranov**, *Koszul duality for operads*, *Duke Math. J.* 76 (1994) 203–272 [MR1301191](#)
- [15] **E Hoffbeck**, *A Poincaré–Birkhoff–Witt criterion for Koszul operads*, *Manuscripta Math.* 131 (2010) 87–110 [MR2574993](#)
- [16] **T V Kadeishvili**, *The algebraic structure in the homology of an $A(\infty)$ -algebra*, *Soobshch. Akad. Nauk Gruzin. SSR* 108 (1982) 249–252 [MR720689](#)
- [17] **M Kontsevich**, *Homological algebra of mirror symmetry*, from: “Proceedings of the International Congress of Mathematicians, Vol. 1, 2”, (S D Chatterji, editor), Birkhäuser, Basel (1995) 120–139 [MR1403918](#)
- [18] **P Van der Laan**, *Coloured Koszul duality and strongly homotopy operads* [arXiv:0312147](#)
- [19] **B H Lian**, **G J Zuckerman**, *New perspectives on the BRST-algebraic structure of string theory*, *Comm. Math. Phys.* 154 (1993) 613–646 [MR1224094](#)
- [20] **M Livernet**, **C Roitzheim**, **S Whitehouse**, *Derived A_∞ -algebras in an operadic context*, *Algebr. Geom. Topol.* 13 (2013) 409–440 [MR3031646](#)
- [21] **J-L Loday**, **B Vallette**, *Algebraic operads*, *Grundlehr. Math. Wissen.* 346, Springer, Heidelberg (2012) [MR2954392](#)
- [22] **Y I Manin**, *Frobenius manifolds, quantum cohomology, and moduli spaces*, *AMS Coll. Publ.* 47, Amer. Math. Soc. (1999) [MR1702284](#)
- [23] **M Markl**, *Distributive laws and Koszulness*, *Ann. Inst. Fourier (Grenoble)* 46 (1996) 307–323 [MR1393517](#)
- [24] **P Salvatore**, **N Wahl**, *Framed discs operads and Batalin–Vilkovisky algebras*, *Q. J. Math.* 54 (2003) 213–231 [MR1989873](#)
- [25] **J D Stasheff**, *Homotopy associativity of H -spaces: I, II*, *Trans. Amer. Math. Soc.* 108, 275–292; *ibid.* 108 (1963) 293–312 [MR0158400](#)

- [26] **ME Sweedler**, *Hopf algebras*, Mathematics Lecture Note Series, W. A. Benjamin, New York (1969) [MR0252485](#)
- [27] **B Vallette**, *Homotopy theory of homotopy algebras* (2013) Available at <http://math.unice.fr/~brunov/HomotopyTheory.pdf>

Institut de Mathématiques de Toulouse, Université Paul Sabatier
118 route de Narbonne, 31062 Toulouse Cedex 9, France

olivia.bellier@math.univ-toulouse.fr

<http://www.math.univ-toulouse.fr/~obellier>

Received: 25 February 2013 Revised: 6 June 2013