

Classifying spaces of algebras over a prop

SINAN YALIN

We prove that a weak equivalence between cofibrant props induces a weak equivalence between the associated classifying spaces of algebras. This statement generalizes to the prop setting a homotopy invariance result which is well known in the case of algebras over operads. The absence of model category structure on algebras over a prop creates difficulties and we introduce new methods to overcome them. We also explain how our result can be extended to algebras over colored props in any symmetric monoidal model category tensored over chain complexes.

18G55; 18D10, 18D50

Introduction

The notion of a prop was introduced to algebra by MacLane [11]. The name prop is actually an acronym for “product and permutation.” Briefly, a prop P is a double sequence of objects $P(m, n)$ whose elements represent operations with m inputs and n outputs.

Certain categories of algebras, like associative, Poisson or Lie algebras, have a structure which is fully determined by operations with a single output. These categories are associated to props P of a certain form, where operations in components $P(m, 1)$ generate the prop. Boardman and Vogt coined the name “categories of operators of standard form” to refer to props of this particular form [1]. Peter May introduced the axioms of operads to deal with the components $P(m, 1)$ which define the core of such prop structures [14]. The work of these authors was initially motivated by the theory of iterated loop spaces in topology (see Boardman and Vogt [2] and [14]). Operads have now proved to be a powerful device to handle a variety of algebraic structures occurring in many branches of mathematics.

However, if one wants to deal with bialgebras it becomes necessary to use general props instead of operads. Important examples appear in particular in mathematical physics and string topology: the Frobenius bialgebras (whose category is equivalent to the category of two-dimensional topological quantum field theories), the topological conformal field theories (which are algebras over the chain Segal prop), or the Lie

bialgebras introduced by Drinfeld in quantization theory are categories of bialgebras associated to props.

The purpose of this article is to set up a theory for the homotopical study of algebras over a (possibly colored) prop. In a seminal series of papers at the beginning of the 1980s, Dwyer and Kan investigated the simplicial localization of categories. They proved that the simplicial localization gives a good device to capture secondary homology structures usually defined in the framework of Quillen's model categories (see Dwyer and Kan [3]). An important homotopy invariant of a model category is its classifying space, defined as the nerve of its subcategory of weak equivalences. The interest of such a classifying space has been shown in the work of Dwyer and Kan [3], who proved that this classifying space encodes information about the homotopy types of the objects and their internal symmetries, ie their homotopy automorphisms. They also proved that such a classifying space is homotopy invariant under Quillen equivalences of model categories.

The algebras over an operad in a model category themselves form, under suitable assumptions, a model category. A consequence of usual results about model categories is that the classifying space of such a category is homotopy invariant up to the weak homotopy type of the underlying operad. Unfortunately, there is no model category structure on the algebras over a prop in general. We cannot handle our motivating examples of bialgebras occurring in mathematical physics and string topology by using this approach, and we aim to overcome this difficulty.

The basic problem is to compare categories of algebras over a prop. In order to bypass difficulties due to the absence of model structure on these algebras, our overall strategy is to stay at the prop level as far as possible, and to use factorization and lifting properties in the model category of props. The structure of an algebra over a prop P can be encoded by a prop morphism $P \rightarrow \text{End}_A$, where End_A is the endomorphism prop associated to A . One can construct a version of endomorphism props modeling P -algebra structures on diagrams. We can in particular use these diagrams endomorphisms props to define path objects in the category of P -algebras. But we need an analogue of this device for a variable P -algebra A , not a fixed object. The idea is to perform such a construction on the abstract prop P itself before moving to endomorphism props. Combining this "prop of P -diagrams" construction with lifting and factorization techniques, we endow the category of P -algebras with functorial path objects.

Consequently, the first main outcome of our study is the following homotopy invariance theorem. Let $\text{Ch}_{\mathbb{K}}$ be the category of \mathbb{Z} -graded chain complexes over a field \mathbb{K} of characteristic zero. Let $(\text{Ch}_{\mathbb{K}})^P$ be the category of algebras associated to a prop P in this category and $w(\text{Ch}_{\mathbb{K}})^P$ its subcategory obtained by restriction to morphisms which are weak equivalences in $\text{Ch}_{\mathbb{K}}$. Our result reads:

Theorem 0.1 *Let $\varphi: P \xrightarrow{\sim} Q$ be a weak equivalence between two cofibrant props. The map φ gives rise to a functor $\varphi^*: w(\text{Ch}_{\mathbb{K}})^Q \rightarrow w(\text{Ch}_{\mathbb{K}})^P$ which induces a weak equivalence of simplicial sets $\mathcal{N}\varphi^*: \mathcal{N}w(\text{Ch}_{\mathbb{K}})^Q \xrightarrow{\sim} \mathcal{N}w(\text{Ch}_{\mathbb{K}})^P$.*

We can remove the hypothesis about the characteristic of \mathbb{K} if we suppose that P is a prop with nonempty inputs or outputs (see Definition 1.9 and Theorem 1.10). We explain in Section 2.7 how to extend Theorem 0.1 to the case of a category tensored over $\text{Ch}_{\mathbb{K}}$. In Section 3, we also briefly show that the proof of Theorem 0.1 extends readily to the colored props context if we suppose that \mathbb{K} has characteristic zero (this hypothesis is needed to put a model category structure on colored props in $\text{Ch}_{\mathbb{K}}$, see the work of Johnson and Yau[10]). Recall that examples include cofibrant resolutions of the props encoding associative bialgebras, Lie bialgebras, Frobenius bialgebras or Poisson bialgebras for instance. Algebras over a cofibrant resolution of a given prop P are called homotopy P -algebras. Theorem 0.1 implies that the classifying space does not depend on the choice of the cofibrant resolution and thus provides a well-defined homotopy invariant.

Remark 0.2 We do not address the case of simplicial sets. However, the work in [10, Theorem 1.4] endows the algebras over a colored prop in simplicial sets with a model category structure. Moreover, the free algebra functor exists in this case. Therefore one can transpose the methods used in the operadic setting to obtain a simplicial version of Theorem 0.1. We also conjecture that our results have a version in simplicial modules which follows from arguments similar to ours.

Organization The overall setting is reviewed in Section 1. We recall some definitions about symmetric monoidal categories over a base category and axioms of monoidal model categories. Then we introduce the precise definition of props and algebras over a prop. We conclude these preliminaries with a fundamental result, the existence of a model structure on the category of props.

The heart of this paper consists of Section 2, devoted to the proof of Theorem 0.1. The proof of Theorem 0.1 is quite long and has been consequently divided in several steps. Section 2.1 gives a sketch of our main arguments. In Sections 2.3, 2.4 and 2.5, we define particular props called props of P -diagrams, which allow us to build a functorial path object in P -algebras. In Section 2.6, we give a proof of Theorem 0.1. At the end of Section 2 we generalize Theorem 0.1 to categories tensored over $\text{Ch}_{\mathbb{K}}$. Finally, we quickly present in Section 3 the extension of our arguments to colored props.

1 Recollections and general results

1.1 Symmetric monoidal categories over a base category

Definition 1.1 Let \mathcal{C} be a symmetric monoidal category. A *symmetric monoidal category over \mathcal{C}* is a symmetric monoidal category $(\mathcal{E}, \otimes_{\mathcal{E}}, 1_{\mathcal{E}})$ endowed with a symmetric monoidal functor $\eta: \mathcal{C} \rightarrow \mathcal{E}$, that is, an object under \mathcal{C} in the 2–category of symmetric monoidal categories.

This defines on \mathcal{E} an external tensor product $\otimes: \mathcal{C} \times \mathcal{E} \rightarrow \mathcal{E}$ by $C \otimes X = \eta(C) \otimes_{\mathcal{E}} X$ for every $C \in \mathcal{C}$ and $X \in \mathcal{E}$. This external tensor product is equipped with the following natural unit, associativity and symmetry isomorphisms:

- (1) For all $X \in \mathcal{E}$, $1_{\mathcal{C}} \otimes X \cong X$.
- (2) For all $X \in \mathcal{E}$, for all $C, D \in \mathcal{C}$, $(C \otimes D) \otimes X \cong C \otimes (D \otimes X)$.
- (3) For all $C \in \mathcal{C}$, for all $X, Y \in \mathcal{E}$, $C \otimes (X \otimes Y) \cong (C \otimes X) \otimes Y \cong X \otimes (C \otimes Y)$.

The coherence constraints of these natural isomorphisms (associativity pentagons, symmetry hexagons and unit triangles which mix both internal and external tensor products) come from the symmetric monoidal structure of the functor η .

We will implicitly assume throughout the paper that all small limits and small colimits exist in \mathcal{C} and \mathcal{E} , and that each of these categories admit an internal hom bifunctor. We suppose moreover the existence of an external hom bifunctor $\text{Hom}_{\mathcal{E}}(-, -): \mathcal{E}^{\text{op}} \times \mathcal{E} \rightarrow \mathcal{C}$ satisfying an adjunction relation

$$\text{for all } C \in \mathcal{C}, \text{ for all } X, Y \in \mathcal{E}, \quad \text{Mor}_{\mathcal{E}}(C \otimes X, Y) \cong \text{Mor}_{\mathcal{C}}(C, \text{Hom}_{\mathcal{E}}(X, Y))$$

(so \mathcal{E} is naturally an enriched category over \mathcal{C}).

Examples (1) The differential graded \mathbb{K} –modules (where \mathbb{K} is a commutative ring) form a symmetric monoidal category over the \mathbb{K} –modules. This is the main category used in this paper.

(2) Any symmetric monoidal category \mathcal{C} forms a symmetric monoidal category over Set (the category of sets) with an external tensor product defined by

$$\begin{aligned} \otimes: \text{Set} \times \mathcal{C} &\rightarrow \mathcal{C}, \\ (S, C) &\mapsto \bigoplus_{s \in S} C. \end{aligned}$$

(3) Let I be a small category; the I –diagrams in a symmetric monoidal category \mathcal{C} form a symmetric monoidal category over \mathcal{C} . The internal tensor product

is defined pointwise, and the external tensor product is defined by the functor η which associates to $X \in \mathcal{C}$ the constant I -diagram C_X on X . The external hom $\text{Hom}_{\mathcal{C}^I}(-, -): \mathcal{C}^I \times \mathcal{C}^I \rightarrow \mathcal{C}$ is given by

$$\text{Hom}_{\mathcal{C}^I}(X, Y) = \int_{i \in I} \text{Hom}_{\mathcal{C}}(X(i), Y(i)).$$

Proposition 1.2 *Let $F: \mathcal{D} \rightleftarrows \mathcal{E}: G$ be a symmetric monoidal adjunction between two symmetric monoidal categories over \mathcal{C} . If F preserves the external tensor product then F and G satisfy an enriched adjunction relation*

$$\text{Hom}_{\mathcal{E}}(F(X), Y) \cong \text{Hom}_{\mathcal{D}}(X, G(Y))$$


at the level of the external hom bifunctors (see Fresse [6, Proposition 1.1.16]).

We now deal with symmetric monoidal categories equipped with a model structure. We assume that the reader is familiar with the basics of model categories. We refer to the paper of Dwyer and Spalinski [4] for a complete and accessible introduction and to Hirschhorn [8] and Hovey [9] for a comprehensive treatment. We just recall the axioms of symmetric monoidal model categories formalizing the interplay between the tensor and the model structures.

Definition 1.3 Let \mathcal{C} be a category with small colimits and $F: \mathcal{A} \times \mathcal{B} \rightarrow \mathcal{C}$ a bifunctor. The *pushout-product* of two morphisms $f: A \rightarrow B \in \mathcal{A}$ and $g: C \rightarrow D \in \mathcal{B}$ is the morphism

$$(f_*, g_*): F(A, D) \oplus_{F(A, C)} F(B, C) \rightarrow F(B, D)$$

given by the commutative diagram:



Definition 1.4 (1) A *symmetric monoidal model category* is a symmetric monoidal category \mathcal{C} equipped with a model category structure such that the following axiom holds:

MM1 The pushout-product $(i_*, j_*): A \otimes D \oplus_{A \otimes C} B \otimes C \rightarrow B \otimes D$ of cofibrations $i: A \rightarrow B$ and $j: C \rightarrow D$ is a cofibration which is also acyclic as soon as i or j is so.

(2) Suppose that \mathcal{C} is a symmetric monoidal model category. A symmetric monoidal category \mathcal{E} over \mathcal{C} is a symmetric monoidal model category over \mathcal{C} if the axiom MM1 holds for both the internal and external tensor products of \mathcal{E} .

Example The category $\text{Ch}_{\mathbb{K}}$ of chain complexes over a field \mathbb{K} is our main working example of symmetric monoidal model category. The weak equivalences of $\text{Ch}_{\mathbb{K}}$ are the quasi-isomorphisms, that is, the morphisms of chain complexes inducing isomorphisms of graded vector spaces at the homology level. The fibrations are the degreewise surjections and the cofibrations the degreewise injections.

Lemma 1.5 *In a symmetric monoidal model category \mathcal{E} over \mathcal{C} the axiom MM1 for the external tensor product is equivalent to the following one:*

MM1' *The morphism*

$$(i^*, p_*) : \text{Hom}_{\mathcal{E}}(B, X) \rightarrow \text{Hom}_{\mathcal{E}}(A, X) \times_{\text{Hom}_{\mathcal{E}}(A, Y)} \text{Hom}_{\mathcal{E}}(B, Y)$$

induced by a cofibration $i: A \twoheadrightarrow B$ and a fibration $p: X \twoheadrightarrow Y$ is a fibration in \mathcal{C} which is also acyclic as soon as i or p is so (cf [9, Lemma 4.2.2]).

One can use the internal hom bifunctor to see that the axiom MM1 for the internal tensor product is in the same way equivalent to a “dual” axiom MM1'.

1.2 On Σ -bimodules, props and algebras over a prop

Let \mathcal{C} be a symmetric monoidal category admitting all small limits and small colimits, whose tensor product preserves colimits, and which is endowed with an internal hom bifunctor. Let \mathbb{B} be the category having the pairs $(m, n) \in \mathbb{N}^2$ as objects together with morphism sets such that

$$\text{Mor}_{\mathbb{B}}((m, n), (p, q)) = \begin{cases} \Sigma_m^{\text{op}} \times \Sigma_n & \text{if } (p, q) = (m, n), \\ \emptyset & \text{otherwise.} \end{cases}$$

The Σ -bibijects in \mathcal{C} are the \mathbb{B} -diagrams in \mathcal{C} . So a Σ -biobject is a double sequence $\{M(m, n) \in \mathcal{C}\}_{(m,n) \in \mathbb{N}^2}$ where each $M(m, n)$ is equipped with a right action of Σ_m and a left action of Σ_n commuting with each other. Let \mathbb{A} be the discrete category of pairs $(m, n) \in \mathbb{N}^2$. We have an obvious forgetful functor $\phi^*: \mathcal{C}^{\mathbb{B}} \rightarrow \mathcal{C}^{\mathbb{A}}$. This functor has a left adjoint $\phi_! : \mathcal{C}^{\mathbb{A}} \rightarrow \mathcal{C}^{\mathbb{B}}$ defined on objects by

$$\phi_! M(m, n) = 1_{\mathcal{C}}[\Sigma_n \times \Sigma_m^{\text{op}}] \otimes M(m, n) \cong \bigoplus_{\Sigma_n \times \Sigma_m^{\text{op}}} M(m, n)$$

for all $M \in \mathcal{C}^{\mathbb{A}}$ and for all $(m, n) \in \mathbb{N}^2$.

Definition 1.6 Let \mathcal{C} be a symmetric monoidal category. A *prop* in \mathcal{C} is a symmetric monoidal category P , enriched over \mathcal{C} , with \mathbb{N} as object set and the tensor product given by $m \otimes n = m + n$ on objects. Let us unwrap this definition. Firstly we see that a prop is a Σ -biobject. Indeed, the group Σ_m acts on $m = 1 + \dots + 1 = 1^{\otimes m}$ and the group Σ_n^{op} acts on $n = 1 + \dots + 1 = 1^{\otimes n}$ by permuting the variables at the morphisms level. A prop is endowed with horizontal products

$$\circ_h: P(m_1, n_1) \otimes P(m_2, n_2) \rightarrow P(m_1 + m_2, n_1 + n_2)$$

which are defined by the tensor product of homomorphisms, since

$$P(m_1 \otimes m_2, n_1 \otimes n_2) = P(m_1 + m_2, n_1 + n_2)$$

by definition of the tensor product on objects. It also admits vertical composition products

$$\circ_v: P(k, n) \otimes P(m, k) \rightarrow P(m, n)$$

corresponding to the composition of homomorphisms, and units $1 \rightarrow P(n, n)$ corresponding to identity morphisms of the objects $n \in \mathbb{N}$ in P . These operations satisfy relations coming from the axioms of symmetric monoidal categories. We refer the reader to Enriquez and Etingof [5] for an explicit description of props in the context of modules over a ring. We denote by \mathcal{P} the category of props.

Another construction of props is given in [10]: props are defined there as \boxtimes_h -monoids in the \boxtimes_v -monoids of colored Σ -biobjects, where \boxtimes_h and \boxtimes_v denote respectively a horizontal composition product and a vertical composition product.

Fresse [7, Appendix A] provides a construction of the free prop on a Σ -biobject. The free prop functor is left adjoint to the forgetful functor

$$F: \mathcal{C}^{\mathbb{B}} \rightleftarrows \mathcal{P}: U.$$

Definition 1.7 (1) To any object X of \mathcal{C} we can associate an *endomorphism prop* End_X defined by

$$\text{End}_X(m, n) = \text{Hom}_{\mathcal{C}}(X^{\otimes m}, X^{\otimes n}).$$

The actions of the symmetric groups are the permutations of the input variables and of the output variables, the horizontal product is the tensor product of homomorphisms and the vertical composition product is the composition of homomorphisms. The units $1_{\mathcal{C}} \rightarrow \text{Hom}_{\mathcal{C}}(X^{\otimes n}, X^{\otimes n})$ represent $\text{id}_{X^{\otimes n}}$.

(2) An *algebra over a prop* P , or P -algebra, is an object $X \in \mathcal{C}$ equipped with a prop morphism $P \rightarrow \text{End}_X$.

We can also define a P -algebra in a symmetric monoidal category over \mathcal{C} :

Definition 1.8 Let \mathcal{E} be a symmetric monoidal category over \mathcal{C} .

- (1) The endomorphism prop of $X \in \mathcal{E}$ is given by $\text{End}_X(m, n) = \text{Hom}_{\mathcal{E}}(X^{\otimes m}, X^{\otimes n})$, where $\text{Hom}_{\mathcal{E}}(-, -)$ is the external hom bifunctor of \mathcal{E} .
- (2) Let P be a prop in \mathcal{C} . A P -algebra in \mathcal{E} is an object $X \in \mathcal{E}$ equipped with a prop morphism $P \rightarrow \text{End}_X$.

Example We recall from [7] an explicit definition in the case of a diagram category over \mathcal{E} : Let $\{X_i\}_{i \in I}$ be a I -diagram in \mathcal{E} . Then we have

$$\text{End}_{\{X_i\}_{i \in I}} = \int_{i \in I} \text{Hom}_{\mathcal{E}}(X_i^{\otimes m}, X_i^{\otimes n}).$$

This end can equivalently be defined as a coreflexive equalizer

$$\begin{array}{ccc} \text{End}_{\{X_i\}}(m, n) & & \\ \downarrow & & \\ \prod_{i \in I} \text{Hom}_{\mathcal{E}}(X_i^{\otimes m}, X_i^{\otimes n}) & \xrightleftharpoons[d_1]{d_0} & \prod_{i \rightarrow j \in \text{mor}(I)} \text{Hom}_{\mathcal{E}}(X_i^{\otimes m}, X_j^{\otimes n}), \\ & \xleftarrow{s_0} & \end{array}$$

where d_0 is the product of the maps

$$u_*: \text{Hom}_{\mathcal{E}}(X_i^{\otimes m}, X_i^{\otimes n}) \rightarrow \text{Hom}_{\mathcal{E}}(X_i^{\otimes m}, X_j^{\otimes n})$$

induced by the morphisms $u: i \rightarrow j$ of I and d_1 is the product of the maps

$$u^*: \text{Hom}_{\mathcal{E}}(X_j^{\otimes m}, X_j^{\otimes n}) \rightarrow \text{Hom}_{\mathcal{E}}(X_i^{\otimes m}, X_j^{\otimes n}).$$

The section s_0 is the projection on the factors associated to the identities $\text{id}: i \rightarrow i$.

Further, this construction is functorial in I : given a J -diagram $\{X_j\}_{j \in J}$, every functor $\alpha: I \rightarrow J$ gives rise to a prop morphism $\alpha^*: \text{End}_{\{X_j\}_{j \in J}} \rightarrow \text{End}_{\{X_{\alpha(i)}\}_{i \in I}}$.

1.3 The semimodel category of props

Suppose that \mathcal{C} is a cofibrantly generated symmetric monoidal model category. The category of Σ -biobjects $\mathcal{C}^{\mathbb{B}}$ is a diagram category over \mathcal{C} , so it inherits a cofibrantly generated model category structure. The weak equivalences and fibrations are defined componentwise. The generating (acyclic) cofibrations are given by $i \otimes \phi!G(m, n)$,

where $(m, n) \in \mathbb{N}^2$ and i ranges over the generating (acyclic) cofibrations of \mathcal{C} . Here \otimes is the external tensor product of $\mathcal{C}^{\mathbb{B}}$ and $G_{(m,n)}$ is the double sequence defined by

$$G_{(m,n)}(p, q) = \begin{cases} 1_{\mathcal{C}} & \text{if } (p, q) = (m, n), \\ 0 & \text{otherwise.} \end{cases}$$

We can also see this result as a transfer of cofibrantly generated model category structure via the adjunction $\phi_! : \mathcal{C}^{\mathbb{A}} \rightleftarrows \mathcal{C}^{\mathbb{B}} : \phi^*$ (via exactly the same proof as in the case of Σ -objects; see for instance [6, Proposition 11.4.A]). The question is to know whether the adjunction $F : \mathcal{C}^{\mathbb{B}} \rightleftarrows \mathcal{P} : U$ transfer this model category structure to the props. In the general case it works only with the subcategory \mathcal{P}_0 of props with nonempty inputs or outputs and does not give a full model category structure. We give the precise statement in Theorem 1.10.

Definition 1.9 A Σ -biobject M has *nonempty inputs* if it satisfies

$$M(0, n) = \begin{cases} 1_{\mathcal{C}} & \text{if } n = 0, \\ 0 & \text{otherwise.} \end{cases}$$

We define in a symmetric way a Σ -biobject with *nonempty outputs*. The category of Σ -biobjects with nonempty inputs is noted $\mathcal{C}_0^{\mathbb{B}}$.

The composite adjunction

$$\mathcal{C}^{\mathbb{A}} \rightleftarrows \mathcal{C}^{\mathbb{B}} \rightleftarrows \mathcal{P}$$

restricts to an adjunction

$$\mathcal{C}_0^{\mathbb{A}} \rightleftarrows \mathcal{C}_0^{\mathbb{B}} \rightleftarrows \mathcal{P}_0.$$

We define the weak equivalences (respectively fibrations) in \mathcal{P}_0 componentwise, ie their images by the forgetful functor $U : \mathcal{P}_0 \rightarrow \mathcal{C}_0^{\mathbb{A}}$ are weak equivalences (respectively fibrations) in $\mathcal{C}_0^{\mathbb{A}}$. We define the generating (acyclic) cofibrations as the images under the free prop functor of the generating (acyclic) cofibrations of $\mathcal{C}_0^{\mathbb{B}}$. We have the following result:

Theorem 1.10 (cf [7, Theorem 4.9]) *Let \mathcal{C} be a cofibrantly generated symmetric monoidal model category. Suppose moreover that the unit of \mathcal{C} is cofibrant. Then the category \mathcal{P}_0 of props with nonempty inputs (or outputs) equipped with the classes of weak equivalences, fibrations and cofibrations defined above forms a semimodel category. Moreover the forgetful functor $U : \mathcal{P}_0 \rightarrow \mathcal{C}_0^{\mathbb{A}}$ preserves cofibrations with cofibrant domain.*

A semimodel category structure is a slightly weakened version of model category structure: the lifting axioms hold only for cofibrations with cofibrant domain, and the factorization axioms hold only on maps with cofibrant domain (see the relevant section of [7]). The notion of a semimodel category is sufficient to do homotopy theory. In certain categories we recover a full model structure on the whole category of props:

Theorem 1.11 (cf [7, Theorem 5.5]) *If the base category \mathcal{C} is the category of dg-modules over a ring \mathbb{K} such that $\mathbb{Q} \subset \mathbb{K}$, simplicial modules over a ring, simplicial sets or topological spaces, then there is a transfer of model category structure on the whole category of props via the adjunction $\mathcal{C}^{\mathbb{A}} \rightleftarrows \mathcal{P}$.*

2 Homotopy invariance of the classifying space

The purpose of this section is to establish Theorem 0.1. We give the details of our arguments in the case $\mathcal{E} = \mathcal{C} = \text{Ch}_{\mathbb{K}}$ (the \mathbb{Z} -graded chain complexes over a field \mathbb{K} of characteristic zero). Afterwards, we briefly explain the generalization of these arguments when \mathcal{E} is a cofibrantly generated symmetric monoidal model category over $\text{Ch}_{\mathbb{K}}$.

2.1 Statement of the result and outline of the proof

In the work of Dwyer–Kan [3], the classifying space of a category \mathcal{M} equipped with a subcategory of weak equivalences $w\mathcal{M}$ is the simplicial set $\mathcal{N}(w\mathcal{M})$, where \mathcal{N} is the simplicial nerve functor. This simplicial set satisfies the following crucial property:

Theorem 2.1 (Dwyer–Kan) *Let M be a category, W a class of morphisms of M and wM the subcategory of M defined by $\text{ob}(wM) = \text{ob}(M)$ and $\text{mor}(wM) = W$. Then one has a homotopy equivalence*

$$\mathcal{N}wM \sim \coprod_{[X]} \overline{W}LwM(X, X),$$

where \mathcal{N} is the simplicial nerve functor, $[X]$ ranges over the weak equivalence classes of the objects of M , \overline{W} is the simplicial classifying space (see May [13]) and $L(-)$ is the simplicial localization functor. When M is a model category, one has moreover

$$\mathcal{N}wM \sim \coprod_{[X]} \overline{W}\text{haut}(X),$$

where $\text{haut}(X)$ is the simplicial monoid of self weak equivalences on a fibrant–cofibrant resolution of X .

In the case of \mathcal{E}^P (the P -algebras in \mathcal{E} for a prop P defined in \mathcal{C}) we use the name “classifying space” to refer to the simplicial set $\mathcal{N}w(\mathcal{E}^{cf})^P$, where $w(\mathcal{E}^{cf})^P$ is the subcategory of P -algebra morphisms whose underlying morphisms in \mathcal{E} are weak equivalences between fibrant-cofibrant objects.

2.1.1 The operadic case In the operadic context, algebras over operads satisfy the following fundamental property: a weak equivalence between two cofibrant operads induces a weak equivalence between their associated classifying spaces of algebras. The proof of this result is done in three steps. Firstly, one shows the existence of an adjunction between the two categories of algebras: if $\phi: P \rightarrow Q$ is a morphism of operads, it induces an adjunction

$$\phi_! : \mathcal{E}^P \rightleftarrows \mathcal{E}^Q : \phi^*,$$

where ϕ^* is given on each Q -algebra $Q \rightarrow \text{End}_X$ by the precomposition

$$P \xrightarrow{\phi} Q \rightarrow \text{End}_X$$

and $\phi_!$ is obtained via a certain coequalizer for which we refer the reader to [6]. Secondly one proves that if ϕ is a weak equivalence and P and Q are “well-behaved” operads, then this adjunction actually forms a Quillen equivalence, presented in full generality in [6]. This Quillen equivalence is precisely obtained between semimodel categories of algebras over weakly equivalent Σ -cofibrant operads, in a cofibrantly generated symmetric monoidal model category \mathcal{E} over a base category \mathcal{C} (see [6, Theorem 12.5.A proved in Chapter 16]). Finally, according to the results of Dwyer–Kan, a Quillen equivalence induces a weak homotopy equivalence of the classifying spaces (actually it induces much more, that is, a Dwyer–Kan equivalence of the simplicial localizations).


2.1.2 The key statement Such a method fails in the prop setting: one does not know how to construct a left adjoint of the functor ϕ^* . And even if such an adjoint exists, there is no free algebra functor and a model structure does not exist on the category of algebras over a prop except in some particular cases such as simplicial sets (see [10]). So the difficult part is to deal with this absence of model structure to get a similar result for algebras over props. Therefore, our method is entirely different from this one. The crux of our proof is given by the following statement:

Theorem 2.2 *Let P be a cofibrant prop. The mappings*


$$\mathcal{N}\phi^*, \mathcal{N}\psi^* : \mathcal{N}w(\mathcal{E}^{cf})^P \rightrightarrows \mathcal{N}w(\mathcal{E}^{cf})^P$$

associated to homotopic prop morphisms $\phi, \psi : P \rightrightarrows P$ are homotopic in sSet .

Let us outline the main steps of the proof of Theorem 2.2 in the case $\mathcal{E} = \mathcal{C} = \text{Ch}_{\mathbb{K}}$. The idea is to construct a zigzag of natural transformations $\varphi^* \xleftarrow{\sim} Z \xrightarrow{\sim} \psi^*$, where Z is a functorial path object in $\text{Ch}_{\mathbb{K}}^P$. For this, we need in particular to obtain such a functorial path object. The existence of a path object for algebras over props is proved in [7, Section 8], but this path object is not functorial. The main point of our proof is to solve this functoriality problem by “correcting” in a certain sense the P -algebra structure on the path object, and then following arguments similar to those of [7] but with functorial P -actions. We proceed as follows. We use functional notation $\mathcal{Y}(X)$, $\mathcal{Z}(X)$ and $\mathcal{V}(X)$ to refer to diagrams functorially associated to an object X which, in our constructions, ranges within (some subcategory of) $\text{Ch}_{\mathbb{K}}$. We first consider the functorial path object diagram associated to any X in $\text{Ch}_{\mathbb{K}}$,



and its subdiagram $\mathcal{Z}(X) = \{X_0 \xleftarrow{\sim} Z(X) \xrightarrow{\sim} X_1\}$. We prove that the natural P -action on the diagram



extends to a natural P -action on $\mathcal{Y}(X)$. For this, we consider “props of P -diagrams”, which are built by replacing all the operations $X^{\otimes m} \rightarrow X^{\otimes n}$ in the endomorphism prop of a given diagram by operations of $P(m, n)$. We use notation $\text{End}_{\mathcal{Y}(P)}$, $\text{End}_{\mathcal{Z}(P)}$ and $\text{End}_{\mathcal{V}(P)}$ to refer to these props of P -diagrams. We verify that these constructions give rise to props acting naturally on the endomorphism prop of the associated diagram. We use these props of P -diagrams to give a P -action on the zigzag of endofunctors $\text{id} \xleftarrow{\sim} Z \xrightarrow{\sim} \text{id}$. We check that we retrieve the action given by φ and ψ on the extremity of this zigzag. We thus have a zigzag connecting φ^* and ψ^* and yielding the desired homotopy between $\mathcal{N}\varphi^*$ and $\mathcal{N}\psi^*$.

Our argument line is divided into two steps.

Step 1 For every $X \in \text{Ch}_{\mathbb{K}}^P$, we have $\text{End}_{\mathcal{V}(X)} \cong \text{End}_X$ so the morphism $P \rightarrow \text{End}_X$ trivially induces a functorial P -action $P \rightarrow \text{End}_{\mathcal{V}(X)}$. In our first step we build a diagram

$$\begin{array}{ccc}
 & \text{End}_{\mathcal{V}(P)} & \xrightarrow{\text{ev}_X} & \text{End}_{\mathcal{V}(X)} \\
 & \nearrow & \downarrow \pi \sim & \downarrow \\
 P & \xrightarrow{=} & P & \xrightarrow{\text{ev}_X} & \text{End}_{\mathcal{V}(X)}.
 \end{array}$$

In $\text{Ch}_{\mathbb{K}}$, the endomorphism prop $\text{End}_{\mathcal{V}(X)}$ is built via the two pullbacks

$$\begin{array}{ccc}
 \text{End}_{\mathcal{V}(X)} & \longrightarrow & \text{End}_{\mathcal{Z}(X)} \\
 \downarrow & & \downarrow s^* \circ \text{pr} \\
 \text{End}_X & \xrightarrow{s_*} & \text{Hom}_{X, \mathcal{Z}(X)} \\
 \\
 \text{End}_{\mathcal{Z}(X)} & \longrightarrow & \text{End}_{X_0} \times \text{End}_{X_1} \\
 \downarrow & & \downarrow d_0^* \times d_1^* \\
 \text{End}_{\mathcal{Z}(X)} & \xrightarrow{(d_0, d_1)_*} & \text{Hom}_{\mathcal{Z}(X), X_0} \times \text{Hom}_{\mathcal{Z}(X), X_1}
 \end{array}$$

where s_* and $(d_0, d_1)_*$ are maps induced by the composition by s and (d_0, d_1) , and s^* , d_0^* and d_1^* are maps induced by the precomposition by s , d_0 and d_1 . The projection $\text{pr}: \text{End}_{\mathcal{Z}(X)} \rightarrow \text{End}_{\mathcal{Z}(X)}$ is induced by the inclusion of diagrams $\{Z(X)\} \hookrightarrow \{X_0 \xleftarrow{\sim} Z(X) \xrightarrow{\sim} X_1\}$ (see [7, Section 8]). The idea is to define a prop of P -diagrams $\text{End}_{\mathcal{V}(P)}$ with a form similar to that of $\text{End}_{\mathcal{V}(X)}$, in order to get the prop morphism $\text{ev}_X: \text{End}_{\mathcal{V}(P)} \rightarrow \text{End}_{\mathcal{V}(X)}$ induced by the morphism $P \rightarrow \text{End}_X$ for each $X \in \text{Ch}_{\mathbb{K}}^P$. For this aim we use two pullbacks similar to those above with props of P -diagrams and Σ -biodjects replacing the usual ones.

Step 2 In our second step, we show that π is an acyclic fibration in \mathcal{P} in order to obtain the desired lifting $P \rightarrow \text{End}_{\mathcal{V}(P)}$ inducing natural P -actions

$$P \rightarrow \text{End}_{\mathcal{V}(P)} \rightarrow \text{End}_{\mathcal{V}(X)}$$

for every $X \in \text{Ch}_{\mathbb{K}}^P$, which respect the P -algebra structures on the diagrams $\mathcal{V}(X)$. It endows the category of P -algebras with a functorial path object. Finally, we prove Theorem 2.2 in Section 2.6, by using lifting properties in the category of props and providing the desired zigzag of natural transformations $\varphi^* \xleftarrow{\sim} Z \xrightarrow{\sim} \psi^*$. Then we show how to deduce Theorem 0.1.

Remark 2.3 We can also wonder about the homotopy invariance of the classifying space up to Quillen equivalences. Let P be a prop in \mathcal{E}_1 . Let $F: \mathcal{E}_1 \rightleftarrows \mathcal{E}_2 : G$ be a symmetric monoidal adjunction. The prop $F(P)$ is defined by applying the functor F entry-wise to P : the fact that F is symmetric monoidal ensures the preservation of the composition products of P , giving to $F(P)$ a prop structure. Further, [10, Lemma 7.1] says that the adjoint pair (F, G) induces an adjunction $\bar{F}: \mathcal{E}_1^P \rightleftarrows \mathcal{E}_2^{F(P)} : \bar{G}$. Now suppose that (F, G) forms a Quillen adjunction. By Brown’s lemma, the functor F preserves weak equivalences between cofibrant objects and the functor G preserves weak equivalences between fibrant objects. If all the objects of \mathcal{E}_1 and \mathcal{E}_2 are fibrant and cofibrant, then the adjoint pair (\bar{F}, \bar{G}) restricts to an adjunction $\bar{F}: w(\mathcal{E}_1)^P \rightleftarrows w(\mathcal{E}_2)^{F(P)} : \bar{G}$ and thus gives rise to a homotopy equivalence $\mathcal{N}w(\mathcal{E}_1)^P \sim \mathcal{N}w(\mathcal{E}_2)^{F(P)}$.

2.2 The path object $Z(X) = Z \otimes X$

Recall that in the model category structure of $\text{Ch}_{\mathbb{K}}$, the fibrations are the degreewise surjections, the cofibrations are the degreewise injections, and the weak equivalences are the morphisms inducing isomorphisms in homology. The category $\text{Ch}_{\mathbb{K}}$ has moreover the simplifying feature that finite products and coproducts coincide. Let Z be the chain complex defined by

$$Z = \mathbb{K}\rho_0 \oplus \mathbb{K}\rho_1 \oplus \mathbb{K}\sigma_0 \oplus \mathbb{K}\sigma_1 \oplus \mathbb{K}\tau.$$

The elements τ, ρ_0 and ρ_1 are three generators of degree 0 and σ_0, σ_1 two generators of degree -1 . The differential d_Z is defined by $d_Z(\sigma_0) = d_Z(\sigma_1) = 0, d_Z(\tau) = 0, d_Z(\rho_0) = \sigma_0$ and $d_Z(\rho_1) = \sigma_1$.

Lemma 2.4 *The chain complex $Z \otimes X$ defines a path object on X in $\text{Ch}_{\mathbb{K}}$, fitting in a factorization $X \xrightarrow{\tilde{s}} Z \otimes X \xrightarrow{(d_0, d_1)} X \oplus X$ of the diagonal*

$$\Delta = (\text{id}_X, \text{id}_X): X \rightarrow X \oplus X$$

such that s is an acyclic cofibration and (d_0, d_1) a fibration.

Proof Let $s: X \rightarrow Z \otimes X$ be the map defined by $s(x) = \tau \otimes x$. Given the differential of Z , the map s is clearly an injective morphism of $\text{Ch}_{\mathbb{K}}$, ie a cofibration. We can also write $Z \otimes X \cong (\tilde{Z} \otimes X) \oplus X$, where

$$\tilde{Z} = \mathbb{K}\rho_0 \oplus \mathbb{K}\rho_1 \oplus \mathbb{K}\sigma_0 \oplus \mathbb{K}\sigma_1$$

is an acyclic complex. The acyclicity of \tilde{Z} implies that s is an acyclic cofibration. We now define a map $(d_0, d_1): Z \otimes X \rightarrow X \oplus X$ such that $(d_0, d_1) \circ s = (\text{id}_X, \text{id}_X)$ and (d_0, d_1) is a fibration. The map d_0 is determined for every $x \in X$ by $d_0(\tau \otimes x) = x$

and $d_0(\sigma_0 \otimes x) = d_0(\sigma_1 \otimes x) = d_0(\rho_0 \otimes x) = d_0(\rho_1 \otimes x) = 0$. The map d_1 is determined for every $x \in X$ by $d_1(\rho_0 \otimes x) = x$, $d_1(\tau \otimes x) = x$, $d_1(\sigma_0 \otimes x) = d_1(\sigma_1 \otimes x) = d_1(\rho_1 \otimes x) = 0$. The map (d_0, d_1) is clearly a surjective chain complexes morphism, ie a fibration and satisfies the equality $(d_0, d_1) \circ s = (\text{id}_X, \text{id}_X)$. \square

The two advantages of this path object on X are its writing in the form of a tensor product with X and its decomposition into a direct sum of X with an acyclic complex.

2.3 The prop $\text{End}_{Z(P)}$

Consider the endomorphism prop of $Z(X)$:

$$\begin{aligned} \text{End}_{Z(X)}(m, n) &= \text{Hom}_{\text{Ch}_{\mathbb{K}}}(Z(X)^{\otimes m}, Z(X)^{\otimes n}) \\ &\cong \text{Hom}_{\text{Ch}_{\mathbb{K}}}(Z^{\otimes m} \otimes X^{\otimes m}, Z^{\otimes n} \otimes X^{\otimes n}) \\ &\cong (Z^{\otimes m})^* \otimes Z^{\otimes n} \otimes \text{End}_X(m, n). \end{aligned}$$

We define a prop of P -diagrams such that

$$\begin{aligned} \text{End}_{Z(P)}(m, n) &= (Z^{\otimes m})^* \otimes Z^{\otimes n} \otimes P(m, n) \\ &= \bigoplus t_1^* \otimes \cdots \otimes t_m^* \otimes t_1 \otimes \cdots \otimes t_n \otimes P(m, n), \end{aligned}$$

where $t_i \in \{\rho_0, \rho_1, \sigma_0, \sigma_1, \tau\}$, together with the following structure maps:

Vertical composition product Let

$$\begin{aligned} \alpha &\in t_1^* \otimes \cdots \otimes t_k^* \otimes t_1 \otimes \cdots \otimes t_n \otimes P(k, n), \\ \beta &\in u_1^* \otimes \cdots \otimes u_m^* \otimes u_1 \otimes \cdots \otimes u_k \otimes P(m, k). \end{aligned}$$

We set

$$\alpha \circ_v \beta = \begin{cases} \alpha \circ_v^P \beta & \text{if } (u_1, \dots, u_k) = (t_1, \dots, t_k), \\ 0 & \text{otherwise,} \end{cases}$$

where \circ_v^P is the vertical composition product of P .

Horizontal product Let

$$\begin{aligned} \alpha &\in t_1^* \otimes \cdots \otimes t_{m_1}^* \otimes t_1 \otimes \cdots \otimes t_{n_1} \otimes P(m_1, n_1), \\ \beta &\in u_1^* \otimes \cdots \otimes u_{m_2}^* \otimes u_1 \otimes \cdots \otimes u_{n_2} \otimes P(m_2, n_2). \end{aligned}$$

We set

$$\begin{aligned} \alpha \circ_h \beta &= t_1^* \otimes \cdots \otimes t_{m_1}^* \otimes u_1^* \otimes \cdots \otimes u_{m_2}^* \\ &\quad \otimes t_1 \otimes \cdots \otimes t_{n_1} \otimes u_1 \otimes \cdots \otimes u_{n_2} \otimes (\alpha |_{P(m_1, n_1)} \circ_h^P \beta |_{P(m_2, n_2)}) \\ &\in t_1^* \otimes \cdots \otimes t_{m_1}^* \otimes u_1^* \otimes \cdots \otimes u_{m_2}^* \\ &\quad \otimes t_1 \otimes \cdots \otimes t_{n_1} \otimes u_1 \otimes \cdots \otimes u_{n_2} \otimes P(m_1 + n_1, m_2 + n_2), \end{aligned}$$

where \circ_h^P is the horizontal product of P .

Actions of the symmetric groups Let $\alpha = t_1^* \otimes \cdots \otimes t_m^* \otimes t_1 \otimes \cdots \otimes t_n \otimes \alpha_P \in \text{End}_{Z(P)}(m, n)$ with $\alpha_P \in P(m, n)$. The action of a permutation $\sigma \in \Sigma_m$ on the right of this prop element is given by $\alpha \cdot \sigma = t_{\sigma(1)}^* \otimes \cdots \otimes t_{\sigma(m)}^* \otimes t_1 \otimes \cdots \otimes t_n \otimes \alpha_P \cdot \sigma$. The action of a permutation $\tau \in \Sigma_n$ on the left of this prop element is given by $\tau \cdot \alpha = t_1^* \otimes \cdots \otimes t_m^* \otimes t_{\tau^{-1}(1)} \otimes \cdots \otimes t_{\tau^{-1}(n)} \otimes \tau \cdot \alpha_P$.

Let $X \in \text{Ch}_{\mathbb{K}}^P$ be a P -algebra. From the definition of $\text{End}_{Z(P)}(m, n)$, we easily see that the prop morphism $P \rightarrow \text{End}_X$ induces a prop morphism

$$\text{ev}_X: \text{End}_{Z(P)} \rightarrow \text{End}_{Z(X)}$$

satisfying the appropriate functoriality diagrams (see Section 2.1.3).

2.4 The prop $\text{End}_{Z(P)}$

2.4.1 The pullback defining $\text{End}_{Z(X)}$ and its explicit maps For every $(m, n) \in \mathbb{N}^2$, we have a pullback:

$$\begin{array}{ccc} \text{End}_{Z(X)}(m, n) & \longrightarrow & \text{End}_{X_0}(m, n) \oplus \text{End}_{X_1}(m, n) \\ \downarrow & & \downarrow (d_0^{\otimes m})^* \oplus (d_1^{\otimes m})^* \\ \text{End}_{Z(X)}(m, n) & \searrow (d_0^{\otimes n}, d_1^{\otimes n})^* & \text{Hom}_{Z(X), X_0}(m, n) \oplus \text{Hom}_{Z(X), X_1}(m, n) \end{array}$$

For every $X \in \text{Ch}_{\mathbb{K}}^P$ and $(m, n) \in \mathbb{N}^2$ we have the isomorphisms

$$\begin{aligned} \text{Hom}_{X, Z(X)}(m, n) &= \text{Hom}_{\text{Ch}_{\mathbb{K}}}(X^{\otimes m}, Z(X)^{\otimes n}) \\ &\cong \text{Hom}_{\text{Ch}_{\mathbb{K}}}(X^{\otimes m}, Z^{\otimes n} \otimes X^{\otimes n}) \cong Z^{\otimes n} \otimes \text{End}_X(m, n) \end{aligned}$$

and

$$\begin{aligned} \text{Hom}_{Z(X), X_i}(m, n) &= \text{Hom}_{\text{Ch}_{\mathbb{K}}}(Z(X)^{\otimes m}, X^{\otimes n}) \\ &\cong \text{Hom}_{\text{Ch}_{\mathbb{K}}}(Z^{\otimes m} \otimes X^{\otimes m}, X^{\otimes n}) \cong (Z^{\otimes m})^* \otimes \text{End}_{X_i}(m, n). \end{aligned}$$

Applying these isomorphisms, we get a pullback:

$$\begin{array}{ccc} \text{End}_{Z(X)}(m, n) & \longrightarrow & \text{End}_{X_0}(m, n) \oplus \text{End}_{X_1}(m, n) \\ \downarrow & & \downarrow \overline{(d_0^{\otimes m})^* \oplus (d_1^{\otimes m})^*} \\ (Z^{\otimes m})^* \otimes Z^{\otimes n} \otimes \text{End}_X(m, n) & \searrow \overline{(d_0^{\otimes n}, d_1^{\otimes n})^*} & (Z^{\otimes m})^* \otimes \text{End}_{X_0}(m, n) \\ & & \oplus (Z^{\otimes m})^* \otimes \text{End}_{X_1}(m, n) \end{array}$$

We have to make explicit the maps $\overline{(d_0^{\otimes n}, d_1^{\otimes n})^*}$ and $\overline{(d_0^{\otimes m})^* \oplus (d_1^{\otimes m})^*}$ and replace $\text{End}_{X_0}(m, n)$, $\text{End}_{X_1}(m, n)$ and $\text{End}_X(m, n)$ by $P_0(m, n)$, $P_1(m, n)$ and $P(m, n)$ to obtain a prop of P -diagrams $\{\text{End}_{Z(P)}(m, n)\}_{(m,n) \in \mathbb{N}^2}$ acting naturally on the chain complexes $\text{End}_{Z(X)}(m, n)$, $X \in \text{Ch}_{\mathbb{K}}^P$. Then we apply the same method to build a prop of P -diagrams $\text{End}_{Y(P)}$ acting naturally on $\text{End}_{Y(X)}$, $X \in \text{Ch}_{\mathbb{K}}^P$.

Lemma 2.5 *Let $\{\underline{z}_i\}_{i \in I}$ be a basis of $Z^{\otimes m}$. The map*

$$\overline{(d_1^{\otimes m})^*}: \text{End}_X(m, n) \rightarrow (Z^{\otimes m})^* \otimes \text{End}_X(m, n)$$

is defined by the formula

$$\overline{(d_1^{\otimes m})^*}(\xi) = \sum_{j \in J} (\underline{z}_j^* \otimes \xi) = \left(\sum_{j \in J} \underline{z}_j^* \right) \otimes \xi,$$

where J is the subset of I such that $d_1^{\otimes m}(\underline{z}_j \otimes \underline{x}) = \underline{x}$ for $\underline{x} \in X^{\otimes m}$ and $j \in J$.

Proof First we give an explicit inverse to the well-known isomorphism

$$\begin{aligned} \lambda: U^* \otimes \text{Hom}_{\text{Ch}_{\mathbb{K}}}(V, V') &\xrightarrow{\cong} \text{Hom}_{\text{Ch}_{\mathbb{K}}}(U \otimes V, V'), \\ \varphi \otimes f &\mapsto [u \otimes v \mapsto \varphi(u).f(v)], \end{aligned}$$

where U is supposed to be of finite dimension. Let $\{u_i\}_{i \in I}$ be a basis of U . We have $\lambda = \sum_{i \in I} \lambda_i$, where

$$\begin{aligned} \lambda_i: \mathbb{K}u_i^* \otimes \text{Hom}_{\text{Ch}\mathbb{K}}(V, V') &\rightarrow \text{Hom}_{\text{Ch}\mathbb{K}}(\mathbb{K}u_i \otimes V, V'), \\ u_i^* \otimes f &\mapsto u_i^*.f: u_i \otimes v \mapsto u_i^*(u_i).f(v) = f(v), \end{aligned}$$

so

$$\begin{aligned} \lambda^{-1}: \text{Hom}_{\text{Ch}\mathbb{K}}(U \otimes V, V') &\rightarrow U^* \otimes \text{Hom}_{\text{Ch}\mathbb{K}}(V, V'), \\ f &\mapsto \sum_{i \in I} (u_i^* \otimes f|_{\mathbb{K}u_i \otimes V}). \end{aligned}$$

Let $\sigma: Z^{\otimes m} \otimes X^{\otimes m} \rightarrow (Z \otimes X)^{\otimes m}$ be the map permuting the variables. Recall that the map d_1 is determined for every $x \in X$ by $d_1(\rho_0 \otimes x) = x$, $d_1(\tau \otimes x) = x$, $d_1(\sigma_0 \otimes x) = d_1(\sigma_1 \otimes x) = d_1(\rho_1 \otimes x) = 0$. The map

$$\begin{aligned} \overline{(d_1^{\otimes m})^*}: \text{Hom}_{\text{Ch}\mathbb{K}}(X^{\otimes m}, X^{\otimes n}) &\rightarrow \text{Hom}_{\text{Ch}\mathbb{K}}(Z^{\otimes m} \otimes X^{\otimes m}, X^{\otimes n}) \\ &\xrightarrow{\cong} (Z^{\otimes m})^* \otimes \text{Hom}_{\text{Ch}\mathbb{K}}(X^{\otimes m}, X^{\otimes n}) \end{aligned}$$

is defined by

$$\xi \mapsto \xi \circ d_1^{\otimes m} \circ \sigma \mapsto \sum_{i \in I} (\underline{z}_i^* \otimes (\xi \circ d_1^{\otimes m} \circ \sigma)|_{\mathbb{K}\underline{z}_i \otimes V})$$

We obtain finally

$$\begin{aligned} \overline{(d_1^{\otimes m})^*}: \text{End}_X(m, n) &\rightarrow (Z^{\otimes m})^* \otimes \text{End}_X(m, n), \\ \xi &\mapsto \sum_{j \in J} (\underline{z}_j^* \otimes \xi) = \left(\sum_{j \in J} \underline{z}_j^* \right) \otimes \xi, \end{aligned}$$

where J is the subset of I such that $d_1^{\otimes m}(\underline{z}_j \otimes \underline{x}) = \underline{x}$ for $\underline{x} \in X^{\otimes m}$ and $j \in J$. If $j \notin J$ then $d_1^{\otimes m}|_{\mathbb{K}\underline{z}_j \otimes X^{\otimes m}} = 0$. □

Recall that the map $d_0: Z \otimes X \rightarrow X$ is defined for every $x \in X$ by

$$d_0(\tau \otimes x) = x \quad \text{and} \quad d_0(\sigma_0 \otimes x) = d_0(\sigma_1 \otimes x) = d_0(\rho_0 \otimes x) = d_0(\rho_1 \otimes x) = 0.$$

As before, the map $\overline{(d_0^{\otimes m})^*}$ has a form similar to that of the map $\overline{(d_1^{\otimes m})^*}$, and we have determined

$$\overline{(d_0^{\otimes m})^*} \oplus \overline{(d_1^{\otimes m})^*}.$$

Lemma 2.6 *The map $\overline{(d_0^{\otimes n}, d_1^{\otimes n})_*}$ is determined by*

$$\overline{(d_0^{\otimes n}, d_1^{\otimes n})_*}: \underline{z}_j^* \otimes \underline{z}'_i \otimes \xi \mapsto \sum_{k \in I} (\underline{z}_k^* \otimes ((d_0^{\otimes n}, d_1^{\otimes n}) \circ \underline{z}_j^*(-) \cdot \underline{z}'_i \otimes \xi) |_{\mathbb{K}_{\underline{z}_k} \otimes X^{\otimes m}}).$$

Proof Let $\{\underline{z}'_i\}_{i \in I'}$ be the basis of $Z^{\otimes n}$. We have the isomorphism

$$(Z^{\otimes m})^* \otimes Z^{\otimes n} \otimes \text{Hom}_{\text{Ch}\mathbb{K}}(X^{\otimes m}, X^{\otimes n}) \rightarrow \text{Hom}_{\text{Ch}\mathbb{K}}(Z^{\otimes m} \otimes X^{\otimes m}, Z^{\otimes n} \otimes X^{\otimes n}),$$

$$\underline{z}_j^* \otimes \underline{z}'_i \otimes \xi \mapsto \underline{z}_j^*(-) \cdot \underline{z}'_i \otimes \xi,$$

that we compose with

$$(d_0^{\otimes n}, d_1^{\otimes n}): Z^{\otimes n} \otimes X^{\otimes n} \rightarrow X_0^{\otimes n} \oplus X_1^{\otimes n},$$

$$\underline{z}_j \otimes x \mapsto \begin{cases} x \oplus x & \text{if } j \in J', \\ x \oplus 0 \text{ or } 0 \oplus x & \text{otherwise,} \end{cases}$$

where J' is the subset of I such that $d_0 |_{\mathbb{K}_{\underline{z}_j} \otimes X^{\otimes n}} \neq 0$ and $d_1 |_{\mathbb{K}_{\underline{z}_j} \otimes X^{\otimes n}} \neq 0$ for $j \in J'$. Finally we compose with the isomorphism

$$\text{Hom}_{\text{Ch}\mathbb{K}}(Z^{\otimes m} \otimes X^{\otimes m}, X_0^{\otimes n} \oplus X_1^{\otimes n}) \xrightarrow{\cong} (Z^{\otimes m})^* \otimes \text{Hom}_{\text{Ch}\mathbb{K}}(X^{\otimes m}, X_0^{\otimes n} \oplus X_1^{\otimes n}),$$

$$f \mapsto \sum_{i \in I} (\underline{z}_i^* \otimes f |_{\mathbb{K}_{\underline{z}_i} \otimes X^{\otimes m}}),$$

and get the map

$$\overline{(d_0^{\otimes n}, d_1^{\otimes n})_*}: \underline{z}_j^* \otimes \underline{z}'_i \otimes \xi \mapsto \sum_{k \in I} (\underline{z}_k^* \otimes ((d_0^{\otimes n}, d_1^{\otimes n}) \circ \underline{z}_j^*(-) \cdot \underline{z}'_i \otimes \xi) |_{\mathbb{K}_{\underline{z}_k} \otimes X^{\otimes m}}).$$

This completes the proof. □

2.4.2 The associated prop of P -diagrams The key observation is that the two maps

$$\overline{(d_0^{\otimes m})^*} \oplus \overline{(d_1^{\otimes m})^*} \quad \text{and} \quad \overline{(d_0^{\otimes n}, d_1^{\otimes n})_*}$$

that fix the prop structure on $\text{End}_{\mathcal{Z}(X)}(m, n)$ in term of those of $(Z^{\otimes m})^* \otimes Z^{\otimes n} \otimes \text{End}_X(m, n)$ and $\text{End}_{X_0}(m, n) \oplus \text{End}_{X_1}(m, n)$ do not modify the operations ξ in $\text{End}_X(m, n)$ themselves. Therefore, we can replace $\text{End}_{X_0}(m, n)$, $\text{End}_{X_1}(m, n)$ and

$\text{End}_X(m, n)$ by $P_0(m, n)$, $P_1(m, n)$ and $P(m, n)$ to get the new pullback:

$$\begin{array}{ccc}
 \text{End}_{\mathcal{Z}(P)}(m, n) & \longrightarrow & P_0(m, n) \oplus P_1(m, n) \\
 \downarrow & & \downarrow \\
 (Z^{\otimes m})^* \otimes Z^{\otimes n} \otimes P(m, n) & & \overline{(d_0^{\otimes m})^* \oplus (d_1^{\otimes m})^*} \\
 & \searrow & \downarrow \\
 & & \overline{(d_0^{\otimes n}, d_1^{\otimes n})_*} \\
 & & (Z^{\otimes m})^* \otimes P_0(m, n) \oplus (Z^{\otimes m})^* \otimes P_1(m, n)
 \end{array}$$

The explicit formulae of the applications defining this pullback, given by Lemmas 2.5 and 2.6, show that these replacements do not break the prop structure transfer. Thus we get the desired prop of P -diagrams $\text{End}_{\mathcal{Z}(P)}$ having the same shape as that of $\text{End}_{\mathcal{Z}(X)}$ and thus acting naturally on the associated diagram of P -algebras via the evaluation morphism

$$\text{ev}_X: \text{End}_{\mathcal{Z}(P)} \rightarrow \text{End}_{\mathcal{Z}(X)}.$$

2.5 The prop $\text{End}_{\mathcal{Y}(P)}$ and the functorial path object in P -algebras

Now let us define $\text{End}_{\mathcal{Y}(P)}$. For every $(m, n) \in \mathbb{N}^2$, the pullback

$$\begin{array}{ccc}
 \text{End}_{\mathcal{Y}(X)}(m, n) & \longrightarrow & \text{End}_{\mathcal{Z}(X)}(m, n) \\
 \downarrow & & \downarrow (s^{\otimes m})^* \text{opr} \\
 \text{End}_X(m, n) & \xrightarrow{(s^{\otimes n})_*} & \text{Hom}_{X, \mathcal{Z}(X)}(m, n)
 \end{array}$$

induces via the isomorphisms explained at the beginning of Section 2.3 and Section 2.4.1 a pullback:

$$\begin{array}{ccc}
 \text{End}_{\mathcal{Y}(X)}(m, n) & \longrightarrow & \text{End}_{\mathcal{Z}(X)}(m, n) \\
 \downarrow & & \downarrow \overline{(s^{\otimes m})^* \text{opr}} \\
 \text{End}_X(m, n) & \xrightarrow{(s^{\otimes n})_*} & Z^{\otimes n} \otimes \text{End}_X(m, n)
 \end{array}$$

In the same manner as before, given that $s: X \rightarrow Z \otimes X$ sends every $x \in X$ to $\tau \otimes x$, the map $\overline{(s^{\otimes m})^*}$ is of the form

$$\begin{aligned}
 (Z^{\otimes m})^* \otimes Z^{\otimes n} \otimes \text{End}_X(m, n) &\rightarrow Z^{\otimes n} \otimes \text{End}_X(m, n), \\
 \underline{z}_j^* \otimes \underline{z}'_i \otimes \xi &\mapsto \begin{cases} \underline{z}'_i \otimes \xi & \text{if } j \in K, \\ 0 & \text{otherwise,} \end{cases}
 \end{aligned}$$

where K is a certain subset of I and $\overline{(s^{\otimes n})_*}$ is of the form

$$\begin{aligned} \text{End}_X(m, n) &\rightarrow Z^{\otimes n} \otimes \text{End}_X(m, n), \\ \xi &\mapsto \sum_{i \in K'} z'_i \otimes \xi, \end{aligned}$$

where K' is a certain subset of I' . These two maps $\overline{(s^{\otimes m})^* \circ \text{pr}}$ and $\overline{(s^{\otimes n})_*}$, fixing the prop structure on $\text{End}_{\mathcal{Y}(X)}(m, n)$ in function of those of $\text{End}_{\mathcal{Z}(X)}(m, n)$ and $\text{End}_X(m, n)$, do not modify the operations $\xi \in \text{End}_X(m, n)$ themselves. Therefore, we replace $\text{End}_X(m, n)$ by $P(m, n)$ and $\text{End}_{\mathcal{Z}(X)}(m, n)$ by $\text{End}_{\mathcal{Z}(P)}(m, n)$ to get this new pullback:

$$\begin{array}{ccc} \text{End}_{\mathcal{Y}(P)}(m, n) & \longrightarrow & \text{End}_{\mathcal{Z}(P)}(m, n) \\ \downarrow & & \downarrow \overline{(s^{\otimes m})^* \circ \text{pr}} \\ P(m, n) & \xrightarrow{\overline{(s^{\otimes n})_*}} & Z^{\otimes n} \otimes P(m, n) \end{array}$$

The explicit formulae of the applications defining this pullback show that these replacements do not break the prop structure transfer. Thus we get the desired prop of P -diagrams $\text{End}_{\mathcal{Y}(P)}$ having the same shape as that of $\text{End}_{\mathcal{Y}(X)}$ and thus acting naturally on the associated diagram of P -algebras via the evaluation morphism

$$\text{ev}_X: \text{End}_{\mathcal{Y}(P)} \rightarrow \text{End}_{\mathcal{Y}(X)}.$$

We finally obtain the following lemma:

Lemma 2.7 *There is a commutative diagram of props:*

$$\begin{array}{ccc} \text{End}_{\mathcal{Y}(P)} & \xrightarrow{\text{ev}_X} & \text{End}_{\mathcal{Y}(X)} \\ \pi \downarrow & & \downarrow \\ P \xrightarrow{=} P & \longrightarrow & \text{End}_{\mathcal{Y}(X)} \end{array}$$

Now we want to prove that the morphism $P \rightarrow \text{End}_{\mathcal{Y}(X)}$ lifts to a morphism

$$P \rightarrow \text{End}_{\mathcal{Y}(P)} \xrightarrow{\text{ev}_X} \text{End}_{\mathcal{Y}(X)}.$$

Lemma 2.8 *The map π is an acyclic fibration in the category of props.*

Proof According to the model category structure on \mathcal{P} , it is sufficient to prove that for every $(m, n) \in \mathbb{N}^2$, $\pi(m, n)$ is an acyclic fibration of chain complexes. The map


$\pi(m, n)$ is given by the base extension

$$\pi(m, n) = P(m, n) \times_{\text{Hom}_{P, Z(P)}(m, n)} \phi(m, n) \times_{\text{Hom}_{Z(P), P_0}(m, n) \oplus \text{Hom}_{Z(P), P_1}(m, n)} (P_0(m, n) \oplus P_1(m, n))$$


where

$$\phi(m, n): \text{End}_{Z(P)}(m, n) \rightarrow \text{Hom}_{P, Z(P)}(m, n) \times_{P_0(m, n) \oplus P_1(m, n)} (\text{Hom}_{Z(P), P_0}(m, n) \oplus \text{Hom}_{Z(P), P_1}(m, n))$$

comes from the diagram:



That is:



We have an isomorphism

$$P_0(m, n) \oplus P_1(m, n) \xrightarrow{\cong} (\mathbb{K} p_0 \oplus \mathbb{K} p_1) \otimes P(m, n),$$

$$p \oplus p' \mapsto p_0 \otimes p + p_1 \otimes p,$$

where p_0 and p_1 are two generators of degree 0. The previous computations give

$$\overline{(d_0^{\otimes n}, d_1^{\otimes n})_*}: Z^{\otimes n} \otimes P(m, n) \rightarrow (\mathbb{K}p_0 \oplus \mathbb{K}p_1) \otimes P(m, n),$$

$$\underline{z}'_i \otimes p \mapsto \begin{cases} (p_0 \oplus p_1) \otimes p & \text{if } i \in J', \\ p_0 \otimes p \text{ or } p_1 \otimes p & \text{otherwise,} \end{cases}$$

and the map

$$\overline{(s^{\otimes m})^*} \oplus \overline{(s^{\otimes m})_*}: (Z^{\otimes m})^* \otimes (\mathbb{K}p_0 \oplus \mathbb{K}p_1) \otimes P(m, n) \rightarrow (\mathbb{K}p_0 \oplus \mathbb{K}p_1) \otimes P(m, n)$$

is defined by

$$\underline{z}^*_j \otimes (\lambda p_0 \oplus \mu p_1) \otimes p \mapsto \begin{cases} (\lambda p_0 \oplus \mu p_1) \otimes p \text{ or } \lambda p_0 \otimes p \text{ or } \mu p_1 \otimes p & \text{if } j \in K, \\ 0 = 0 \otimes p & \text{otherwise.} \end{cases}$$

We have similar results for the two maps starting from $(Z^{\otimes m})^* \otimes Z^{\otimes n} \otimes P(m, n)$. We deduce that the previous diagram is the image under the functor $- \otimes P(m, n)$ of the dual pushout-product

$$\begin{array}{ccc} \text{Hom}_{\text{Ch}_{\mathbb{K}}}(Z^{\otimes m}, Z^{\otimes n}) & \xrightarrow{(g_{d_0, d_1})_*} & \text{Hom}_{\text{Ch}_{\mathbb{K}}}(Z^{\otimes m}, \mathbb{K}p_0 \oplus \mathbb{K}p_1) \\ \downarrow (f_s^*, (g_{d_0, d_1})_*) & \searrow \text{pullback} & \downarrow f_s^* \\ \text{Hom}_{\text{Ch}_{\mathbb{K}}}(\mathbb{K}, Z^{\otimes n}) & \xrightarrow{(g_{d_0, d_1})_*} & \text{Hom}_{\text{Ch}_{\mathbb{K}}}(\mathbb{K}, \mathbb{K}p_0 \oplus \mathbb{K}p_1) \\ \downarrow f_s^* & & \downarrow f_s^* \end{array}$$

modulo the isomorphisms

$$\begin{aligned} Z^{\otimes n} &\cong \text{Hom}_{\text{Ch}_{\mathbb{K}}}(\mathbb{K}, Z^{\otimes n}), \\ (Z^{\otimes m})^* \otimes Z^{\otimes n} &\cong \text{Hom}_{\text{Ch}_{\mathbb{K}}}(Z^{\otimes m}, Z^{\otimes n}), \\ (Z^{\otimes m})^* \otimes (\mathbb{K}p_0 \oplus \mathbb{K}p_1) &\cong \text{Hom}_{\text{Ch}_{\mathbb{K}}}(Z^{\otimes m}, \mathbb{K}p_0 \oplus \mathbb{K}p_1), \\ \mathbb{K}p_0 \oplus \mathbb{K}p_1 &\cong \text{Hom}_{\text{Ch}_{\mathbb{K}}}(\mathbb{K}, \mathbb{K}p_0 \oplus \mathbb{K}p_1). \end{aligned}$$

The map $g_{d_0, d_1}: Z^{\otimes n} \rightarrow \mathbb{K}p_0 \oplus \mathbb{K}p_1$ is surjective so it is a fibration of chain complexes. Recall that we have a decomposition of Z into $Z = \tilde{Z} \oplus \mathbb{K}\tau$, where \tilde{Z} is acyclic, which implies a decomposition of $Z^{\otimes m}$ of the form $Z^{\otimes m} \cong S_m \oplus \mathbb{K}(\tau^{\otimes n})$, where S_m is acyclic because it is a sum of tensor products containing \tilde{Z} . The map f_s is an injection sending \mathbb{K} on $\mathbb{K}(\tau^{\otimes n})$ so it is a cofibration, and S_m is acyclic so f_s is an acyclic cofibration. Applying the axiom MM1' in $\text{Ch}_{\mathbb{K}}$ we conclude that $(f_s^*, (g_{d_0, d_1})_*)$ is an acyclic fibration. Therefore $\phi(m, n) = (f_s^*, (g_{d_0, d_1})_*) \otimes \text{id}_{P(m, n)}$ is an acyclic

fibration, and so is $\pi(m, n)$, given that the class of acyclic fibrations is stable by base extension. □

We have proved the following result:

Proposition 2.9 *There is a functorial P -action $P \rightarrow \text{End}_{\mathcal{V}(X)}$, and consequently a functorial path object $Z: (\text{Ch}_{\mathbb{K}})^P \rightarrow (\text{Ch}_{\mathbb{K}})^P$ in the category of cofibrant–fibrant P -algebras $(\text{Ch}_{\mathbb{K}})^P$.*

2.6 Proof of the final result

Consider now the square of inclusions of diagrams

$$\begin{array}{ccc} \mathcal{T}(X) & \xhookrightarrow{t} & \mathcal{V}(X) \\ u \downarrow & & \downarrow v \\ \mathcal{Z}(X) & \xhookrightarrow{w} & \mathcal{Y}(X), \end{array}$$

where $\mathcal{V}(X)$, $\mathcal{Z}(X)$ and $\mathcal{Y}(X)$ are the diagrams defined previously and $\mathcal{T}(X)$ is the diagram $\{X_0, X_1\}$ consisting of two copies of X and no arrows between them. This square of inclusions induces the commutative square of endomorphism props

$$\begin{array}{ccc} \text{End}_{\mathcal{V}(X)} & \xrightarrow{w_*} & \text{End}_{\mathcal{Z}(X)} \\ v_* \downarrow & & \downarrow u^* \\ \text{End}_{\mathcal{Y}(X)} & \xrightarrow{t^*} & \text{End}_{\mathcal{T}(X)}, \end{array}$$

where u^* , v^* , t^* and w^* are the maps induced by the inclusions of the associated diagrams of P -algebras. We have a commutative diagram of props of P -diagrams reflecting this square:

$$\begin{array}{ccc} \text{End}_{\mathcal{Y}(P)} & \xrightarrow{w_*} & \text{End}_{\mathcal{Z}(P)} \\ v_* \downarrow & & \downarrow u^* \\ \text{End}_{\mathcal{V}(P)} = P & \xrightarrow{t^*} & \text{End}_{\mathcal{T}(P)} = P_0 \times P_1, \end{array}$$

where v^* is the acyclic fibration π of Lemma 2.8 and u^* is a fibration because it is clearly surjective in each biarity (recall that the surjective morphisms are the fibrations of $\text{Ch}_{\mathbb{K}}$ and that the fibrations of \mathcal{P} are determined componentwise). Now we can use this commutative square to prove the final result:

Theorem 2.10 *Let P be a cofibrant prop and $\varphi, \psi: P \rightarrow P$ two homotopic prop morphisms. Then there exists a diagram of functors*

$$\varphi^* \xleftarrow{\sim} Z \xrightarrow{\sim} \psi^*,$$

where Z is the path object functor defined in Proposition 2.9 and the natural transformations are pointwise acyclic fibrations.

Proof This proof follows the arguments of the proof of [7, Theorem 8.4]. We consider a cylinder object of P fitting in a diagram of the form

$$P \vee P \xrightarrow{(d_0, d_1)} \tilde{P} \xrightarrow{s_0} P.$$

The components d_0 and d_1 of the morphism (d_0, d_1) are acyclic cofibrations because P is cofibrant by assumption (see [4, Lemma 4.4]) and s_0 an acyclic fibration. The fact that φ and ψ are homotopic implies the existence of a lifting in

$$\begin{array}{ccc} P \vee P & \xrightarrow{(\varphi, \psi)} & P \\ \downarrow (d_0, d_1) & \nearrow h & \downarrow \\ \tilde{P} & \longrightarrow & 0. \end{array}$$

We produce the lifting

$$\begin{array}{ccc} I & \longrightarrow & \text{End}_{\mathcal{Y}(P)} \\ \downarrow & \nearrow k & \downarrow \sim v^* \\ P & \xrightarrow{\varphi} & P \end{array}$$

(by the axiom MC4 of model categories; see [4]) and form $(\varphi \circ s_0, h): \tilde{P} \rightarrow P_0 \times P_1$ to get the following commutative diagram:

$$\begin{array}{ccccc} P & \xrightarrow{k} & \text{End}_{\mathcal{Y}(P)} & \xrightarrow{w^*} & \text{End}_{\mathcal{Z}(P)} \\ \downarrow d_0 \sim & & \nearrow l & & \downarrow u^* \\ \tilde{P} & \xrightarrow{(\varphi \circ s_0, h)} & P_0 \times P_1 & & \end{array}$$

We have $(\varphi \circ s_0, h) \circ d_0 = (\varphi \circ s_0 \circ d_0, h \circ d_0) = (\varphi, \varphi)$ and $u^* \circ w^* \circ k = t^* \circ v^* \circ k = t^* \circ \varphi = (\varphi, \varphi)$ so this diagram is indeed commutative and there exists a lifting (axiom MC4) $l: \tilde{P} \rightarrow \text{End}_{\mathcal{Z}(P)}$. Then we form $l \circ d_1: P \rightarrow \text{End}_{\mathcal{Z}(P)}$ and observe that

$u^* \circ l \circ d_1 = (\varphi \circ s_0, h) \circ d_1 = (\varphi \circ s_0 \circ d_1, h \circ d_1) = (\varphi, \psi)$, ie we obtain the diagram

$$\begin{array}{ccc}
 & \text{End}_{\mathcal{Z}}(P) & \xrightarrow{\text{ev}_X} & \text{End}_{\mathcal{Z}}(X) \\
 & \nearrow l \circ d_1 & & \downarrow \\
 P & \xrightarrow{(\varphi, \psi)} & P_0 \times P_1 & \xrightarrow{\text{ev}_X} & \text{End}_{\mathcal{T}}(X) \\
 & & \downarrow u^* & & \downarrow
 \end{array}$$

and consequently a diagram of functors $\varphi^* \xleftarrow{\sim} Z \xrightarrow{\sim} \psi^*$. The functorial path object Z on $\text{Ch}_{\mathbb{K}}$ preserves weak equivalences and restrict to an endofunctor of $w\text{Ch}_{\mathbb{K}}$, so the associated functorial path object Z on $\text{Ch}_{\mathbb{K}}^P$ do the same. Moreover, the natural transformations are weak equivalences in each component, so this diagram restricts to the desired diagram of endofunctors of $w\text{Ch}_{\mathbb{K}}^P$. \square

Now we can conclude the proof of Theorem 0.1 in the case $\mathcal{E} = \text{Ch}_{\mathbb{K}}$:

Theorem 2.11 *Let $\text{Ch}_{\mathbb{K}}$ be the category of \mathbb{Z} -graded chain complexes over a field \mathbb{K} of characteristic zero. Let $\varphi: P \xrightarrow{\sim} Q$ be a weak equivalence between two cofibrant props. The map φ gives rise to a functor $\varphi^*: w(\text{Ch}_{\mathbb{K}})^{\mathcal{Q}} \rightarrow w(\text{Ch}_{\mathbb{K}})^P$ which induces a weak equivalence of simplicial sets $\mathcal{N}\varphi^*: \mathcal{N}w(\text{Ch}_{\mathbb{K}})^{\mathcal{Q}} \xrightarrow{\sim} \mathcal{N}w(\text{Ch}_{\mathbb{K}})^P$.*

Proof Recall that \mathcal{P} is the category of props in $\text{Ch}_{\mathbb{K}}$. Let us suppose first that $\varphi: P \xrightarrow{\sim} Q$ is an acyclic cofibration between two cofibrant props of \mathcal{P} . All objects in $\text{Ch}_{\mathbb{K}}$ are fibrant, so by definition of the model category structure on \mathcal{P} the prop P is fibrant and thus we have the lifting

$$\begin{array}{ccc}
 P & \xrightarrow{=} & P \\
 \downarrow \varphi & \nearrow s & \downarrow \\
 Q & \longrightarrow & \text{pt},
 \end{array}$$

where $s: Q \xrightarrow{\sim} P$ satisfies

$$\begin{cases} s \circ \varphi = \text{id}_P, \\ \varphi \circ s \sim \text{id}_Q; \end{cases}$$

the relation \sim is the homotopy relation for the model category structure of \mathcal{P} . These maps induce functors $\varphi^*: (w\mathcal{E}^{cf})^{\mathcal{Q}} \rightarrow (w\mathcal{E}^{cf})^P$ and $s^*: (w\mathcal{E}^{cf})^P \rightarrow (w\mathcal{E}^{cf})^{\mathcal{Q}}$. Applying the simplicial nerve functor, we obtain

$$\begin{cases} \mathcal{N}(s \circ \varphi)^* = \mathcal{N}\varphi^* \circ \mathcal{N}s^* = \text{id}_{(w\mathcal{E}^{cf})^P}, \\ \mathcal{N}(\varphi \circ s)^* = \mathcal{N}s^* \circ \mathcal{N}\varphi^* \sim \text{id}_{(w\mathcal{E}^{cf})^{\mathcal{Q}}}, \end{cases}$$

so $\mathcal{N}\varphi^*$ is a homotopy equivalence in \mathbf{sSet} , which implies that it is a weak equivalence of simplicial sets. The functor

$$\begin{aligned} \mathcal{P} &\rightarrow \mathbf{sSet}, \\ P &\mapsto \mathcal{N}w(\mathcal{E}^{cf})^P, \end{aligned}$$


is defined between two model categories and maps the acyclic cofibrations between cofibrant objects to weak equivalences, so it preserves weak equivalences between cofibrant objects according to Brown’s lemma. \square

2.7 The general case of a category \mathcal{E} tensored over $\mathbf{Ch}_{\mathbb{K}}$


To complete our results we explain how the proof of Theorem 2.10 extends to a category \mathcal{E} tensored over $\mathbf{Ch}_{\mathbb{K}}$.

Theorem 2.12 *Let \mathcal{E} be a cofibrantly generated symmetric monoidal model category over $\mathbf{Ch}_{\mathbb{K}}$. Let $\varphi: P \xrightarrow{\sim} Q$ be a weak equivalence between two cofibrant props defined in $\mathbf{Ch}_{\mathbb{K}}$. This morphism φ gives rise to a functor $\varphi^*: w(\mathcal{E}^c)^Q \rightarrow w(\mathcal{E}^c)^P$ which induces a weak equivalence of simplicial sets $\mathcal{N}\varphi^*: \mathcal{N}w(\mathcal{E}^c)^Q \rightarrow \mathcal{N}w(\mathcal{E}^c)^P$.*

Proof The chain complex Z defined previously is itself the path object on C^0 , so we have the following commutative diagram:



Given that C^0 is the unit of $\mathbf{Ch}_{\mathbb{K}}$, for any $X \in \mathcal{E}$ we have $C^0 \otimes X \cong X$, thus by applying the functor $- \otimes X$ we get the following commutative diagram:



The axiom MM1 for the external tensor product \otimes implies that if X is cofibrant, then the functor $- \otimes X$ preserves acyclic cofibrations of $\text{Ch}_{\mathbb{K}}$ (all the objects of $\text{Ch}_{\mathbb{K}}$ are cofibrant) and thus, by Brown’s lemma, it preserves the weak equivalences. Therefore $s \otimes \text{id}_X$ is still an acyclic cofibration and $d_0 \otimes \text{id}_X, d_1 \otimes \text{id}_X$ are weak equivalences. Moreover, given the properties of \otimes and the fact that endomorphism props in $\text{Ch}_{\mathbb{K}}$ for objects of \mathcal{E} are defined with the external hom bifunctor $\text{Hom}_{\mathcal{E}}(-, -)$ of \mathcal{E} , we have the following isomorphisms:

$$\begin{aligned} \text{End}_{Z \otimes X}(m, n) &= \text{Hom}_{\mathcal{E}}((Z \otimes X)^{\otimes m}, (Z \otimes X)^{\otimes n}) \\ &\cong \text{Hom}_{\mathcal{E}}(Z^{\otimes m} \otimes X^{\otimes m}, Z^{\otimes n} \otimes X^{\otimes n}) \\ &\cong (Z^{\otimes m})^* \otimes Z^{\otimes n} \otimes \text{End}_X(m, n), \\ \text{Hom}_{X, Z \otimes X}(m, n) &= \text{Hom}_{\mathcal{E}}(X^{\otimes m}, (Z \otimes X)^{\otimes n}) \\ &\cong \text{Hom}_{\mathcal{E}}(X^{\otimes m}, Z^{\otimes n} \otimes X^{\otimes n}) \\ &\cong Z^{\otimes n} \otimes \text{End}_X(m, n), \\ \text{Hom}_{Z \otimes X, X_i}(m, n) &= \text{Hom}_{\mathcal{E}}((Z \otimes X)^{\otimes m}, X^{\otimes n}) \\ &\cong \text{Hom}_{\mathcal{E}}(Z^{\otimes m} \otimes X^{\otimes m}, X^{\otimes n}) \\ &\cong (Z^{\otimes m})^* \otimes \text{End}_{X_i}(m, n). \end{aligned}$$

The proofs in Sections 2.3, 2.4 and 2.5 extend without changes to the case of a category \mathcal{E} tensored over $\text{Ch}_{\mathbb{K}}$: we still work in $\text{Ch}_{\mathbb{K}}$, and as before the operations associated to $s \otimes \text{id}_X, d_0 \otimes \text{id}_X$ and $d_1 \otimes \text{id}_X$ in the pullbacks do not transform the elements of $\text{End}_X(m, n)$ themselves, so that the replacement of $\text{End}_X(m, n)$ by $P(m, n)$ does not break the transfer of prop structure in these pullbacks. We obtain a diagram of endofunctors $\varphi^* \leftarrow Z \xrightarrow{\sim} \psi^*$ of $(\mathcal{E}^c)^P$ where the natural transformations are weak equivalences in each component, so this diagram restricts to the desired diagram of endofunctors of $w(\mathcal{E}^c)^P$. Theorem 0.1 is proved in the general case. \square

3 Extension of the results in the colored prop setting

Definition 3.1 Let C be a nonempty set, called the *set of colors*, and \mathcal{C} be a symmetric monoidal category.

(1) A C -colored Σ -biproject M is a double sequence of objects $\{M(m, n) \in \mathcal{E}\}_{(m, n) \in \mathbb{N}^2}$ where each $M(m, n)$ admits commuting left Σ_m -action and right Σ_n -action as well as a decomposition

$$M(m, n) = \text{colim}_{c_i, d_i \in C} M(c_1, \dots, c_m; d_1, \dots, d_n)$$

compatible with these actions. The objects $M(c_1, \dots, c_m; d_1, \dots, d_n)$ should be thought as spaces of operations with colors c_1, \dots, c_m indexing the m inputs and colors d_1, \dots, d_n indexing the n outputs.

(2) A C -colored prop P is a C -colored Σ -biobject endowed with a horizontal composition

$$\begin{aligned} \circ_h: P(c_{11}, \dots, c_{1m_1}; d_{11}, \dots, d_{1n_1}) \otimes \cdots \otimes P(c_{k1}, \dots, c_{km_k}; d_{k1}, \dots, d_{kn_k}) \\ \rightarrow P(c_{11}, \dots, c_{km_k}; d_{k1}, \dots, d_{kn_k}) \subseteq P(m_1 + \cdots + m_k, n_1 + \cdots + n_k) \end{aligned}$$

and a vertical composition

$$\begin{aligned} \circ_v: P(c_1, \dots, c_k; d_1, \dots, d_n) \otimes P(a_1, \dots, a_m; b_1, \dots, b_k) \\ \rightarrow P(a_1, \dots, a_m; d_1, \dots, d_n) \subseteq P(m, n) \end{aligned}$$

which is equal to zero unless $b_i = c_i$ for $1 \leq i \leq k$. These two compositions satisfy associativity axioms (we refer the reader to [10] for details).

Definition 3.2 (1) Let $\{X_c\}_C$ be a collection of objects of \mathcal{E} . The C -colored endomorphism prop $\text{End}_{\{X_c\}_C}$ is defined by

$$\text{End}_{\{X_c\}_C}(c_1, \dots, c_m; d_1, \dots, d_n) = \text{Hom}_{\mathcal{E}}(X_{c_1} \otimes \cdots \otimes X_{c_m}, X_{d_1} \otimes \cdots \otimes X_{d_n})$$

with horizontal composition given by the tensor product of homomorphisms and vertical composition given by the composition of homomorphisms with matching colors.

(2) Let P be a C -colored prop. A P -algebra is the data of a collection of objects $\{X_c\}_C$ and a C -colored prop morphism $P \rightarrow \text{End}_{\{X_c\}_C}$.

Example 3.3 Let I be a small category, P a prop in \mathcal{C} . We can build an $\text{ob}(I)$ -colored prop P_I such that the P_I -algebras are the I -diagrams of P -algebras in \mathcal{E} in the same way as that of Markl [12].

To endow the category of colored props with a model category structure, the cofibrantly generated symmetric monoidal model structure on \mathcal{C} is not sufficient. We have to suppose moreover that the domains of the generating cofibrations and acyclic generating cofibrations are small (cf [8, 10.4.1]), that is to say, the model structure is strongly cofibrantly generated:

Theorem 3.4 (cf [10, Theorem 1.1]) *Let C be a nonempty set. Let \mathcal{C} be a strongly cofibrantly generated symmetric monoidal model category with a symmetric monoidal fibrant replacement functor, and either*

- (1) *a cofibrant unit and a cocommutative interval, or*
- (2) *functorial path data.*

Then the category \mathcal{P}_C of C -colored props in \mathcal{C} forms a strongly cofibrantly generated model category with fibrations and weak equivalences defined componentwise in \mathcal{C} .

This theorem works in particular with the categories of simplicial sets, simplicial modules over a commutative ring and chain complexes over a ring of characteristic 0 (our main category in this paper).

This model structure is similar to that of 1-colored props, and we can define C -colored endomorphism props of morphisms (see [10, Section 4]) and more generally of any kind of diagram, so the lifting properties used in the previous section work in the C -colored case. Moreover, in the proof of Theorem 0.1, the replacement of the operations $X^{\otimes m} \rightarrow X^{\otimes n}$ by $P(m, n)$ can be done using a C -colored prop P instead of a 1-colored one without changing anything to the proof, therefore we finally get the C -colored version of Theorem 0.1. We do not have to change anything to Theorem 0.1, given that $\text{Ch}_{\mathbb{K}}$ satisfies the hypotheses of Theorem 3.4.

4 Afterword

Let \mathcal{E} be a symmetric monoidal model category over $\text{Ch}_{\mathbb{K}}$. Let P be a cofibrant prop defined in $\text{Ch}_{\mathbb{K}}$ and X an object of \mathcal{E} . One can consider the moduli space $P\{X\}$ of P -algebra structures on X , which is a simplicial set whose 0-simplices are prop morphisms $P \rightarrow \text{End}_X$ representing all the P -algebra structures on X . More precisely, the moduli space of P -algebra structures on X is the simplicial set such that

$$P\{X\} = \text{Mor}_{\mathcal{P}_0}(P \otimes \Delta[-], \text{End}_X),$$

where $(-) \otimes \Delta[-]$ is a cosimplicial resolution of P . This space is a Kan complex which is homotopy invariant under weak equivalences of cofibrant props at the source (it follows from general arguments on simplicial mapping spaces in model categories; see [8, Chapter 16]). Moreover, its connected components are exactly the homotopy classes of P -algebra structures on X .

As a consequence of Theorem 0.1, one can follow arguments similar to those of Rezk in [15] to characterize such a moduli space as a homotopy fiber of a map between classifying spaces. To be explicit, we have a homotopy pullback of simplicial sets

$$\begin{array}{ccc} P\{X\} & \longrightarrow & \mathcal{N}(fw(\mathcal{E}^c)^P) \\ \downarrow & & \downarrow \\ \{X\} & \longrightarrow & \mathcal{N}(w\mathcal{E}^c), \end{array}$$

where $fw(\mathcal{E}^c)^P$ is the subcategory of morphisms of P -algebras being acyclic fibrations in \mathcal{E} .

We will prove in a followup paper that the classifying space of acyclic fibrations $\mathcal{N}(fw(\mathcal{E}^c)^P)$ is actually weakly equivalent to the whole classifying space $\mathcal{N}(w(\mathcal{E}^c)^P)$. We accordingly have a homotopy pullback which extends to the setting of algebras over dg (colored) props the result obtained by Rezk [15, Theorem 1.1.5] in the operadic case:

$$\begin{array}{ccc} P\{X\} & \longrightarrow & \mathcal{N}(w(\mathcal{E}^c)^P) \\ \downarrow & & \downarrow \\ \{X\} & \longrightarrow & \mathcal{N}(w\mathcal{E}^c) \end{array}$$

This result implies in particular that the moduli space admits a decomposition in classifying spaces of homotopy automorphisms

$$P\{X\} \sim \coprod_{[X]} \overline{W}Lw(\mathcal{E}^c)^P(X, X),$$

where $[X]$ ranges over the weak equivalence classes of P -algebras having X as underlying object. An interesting set-theoretic consequence is that the homotopy automorphisms of P -algebras $Lw(\mathcal{E}^c)^P(X, X)$ are homotopically small in the sense of Dwyer–Kan [3].

Acknowledgements

I would like to thank my advisor Benoit Fresse for his useful and constant help. I am grateful to Jim Stasheff and Eric Hoffbeck for their relevant remarks on the first version of this paper. I also wish to thank David Chataur for stimulating discussions about these moduli spaces. I thank the referee for his thorough report and detailed suggestions.

References

- [1] **J M Boardman, R M Vogt**, *Homotopy-everything H-spaces*, Bull. Amer. Math. Soc. 74 (1968) 1117–1122 MR0236922
- [2] **J M Boardman, R M Vogt**, *Homotopy invariant algebraic structures on topological spaces*, Lecture Notes in Mathematics 347, Springer (1973) MR0420609
- [3] **W G Dwyer, D M Kan**, *Function complexes in homotopical algebra*, Topology 19 (1980) 427–440 MR584566

- [4] **W G Dwyer, J Spaliński**, *Homotopy theories and model categories*, from: “Handbook of algebraic topology”, North-Holland (1995) 73–126 MR1361887
- [5] **B Enriquez, P Etingof**, *On the invertibility of quantization functors*, J. Algebra 289 (2005) 321–345 MR2142375
- [6] **B Fresse**, *Modules over operads and functors*, Lecture Notes in Mathematics 1967, Springer (2009) MR2494775
- [7] **B Fresse**, *Props in model categories and homotopy invariance of structures*, Georgian Math. J. 17 (2010) 79–160 MR2640648
- [8] **P S Hirschhorn**, *Model categories and their localizations*, Mathematical Surveys and Monographs 99, Amer. Math. Soc. (2003) MR1944041
- [9] **M Hovey**, *Model categories*, Mathematical Surveys and Monographs 63, Amer. Math. Soc. (1999) MR1650134
- [10] **M W Johnson, D Yau**, *On homotopy invariance for algebras over colored PROPs*, J. Homotopy Relat. Struct. 4 (2009) 275–315 MR2559644
- [11] **S Mac Lane**, *Categorical algebra*, Bull. Amer. Math. Soc. 71 (1965) 40–106 MR0171826
- [12] **M Markl**, *Homotopy algebras are homotopy algebras*, Forum Math. 16 (2004) 129–160 MR2034546
- [13] **J P May**, *Simplicial objects in algebraic topology*, Van Nostrand Mathematical Studies 11, D Van Nostrand (1967) MR0222892
- [14] **J P May**, *The geometry of iterated loop spaces*, Lectures Notes in Mathematics 271, Springer (1972) MR0420610
- [15] **C W Rezk**, *Spaces of algebra structures and cohomology of operads*, PhD thesis, Massachusetts Institute of Technology (1996) MR2716655 Available at <http://search.proquest.com/docview/304311860>

Laboratoire Paul Painlevé, Université de Lille 1
Cité Scientifique, 59655 Villeneuve d’Ascq Cedex, France
sinan.yalin@math.univ-lille1.fr

Received: 9 October 2012 Revised: 19 December 2013

