

Surgery along star-shaped plumbings and exotic smooth structures on 4–manifolds

ÇAĞRI KARAKURT
LAURA STARKSTON

We define a new 4–dimensional symplectic cut and paste operation which is analogous to Fintushel and Stern’s rational blow-down. We use this operation to produce multiple constructions of symplectic smoothly exotic complex projective spaces blown up eight, seven, and six times. We also show how this operation can be used in conjunction with knot surgery to construct an infinite family of minimal exotic smooth structures on the complex projective space blown-up seven times.

53Dxx; 57R57

1 Introduction

Fintushel and Stern [14] introduced a cut and paste operation for 4–manifolds called rational blow-down. They used it to compute the Donaldson polynomial of the logarithmic transforms of the elliptic surfaces. Since then, the rational blow-down operation has proven to be very useful in 4–dimensional topology. It was a useful constructional tool in the exotic copies of blown-up complex projective plane; see Park [36], Park, Stipsicz and Szabó [37], Stipsicz and Szabó [42], Fintushel and Stern [16], and Michalogiorgaki [30]. It can be used to construct symplectic manifolds, as shown by Symington [44; 45] and Gay and Stipsicz [22], and in the presence of a certain Lefschetz fibration structure one can reinterpret it as a monodromy substitution; see Endo and Gurtas [10] and Endo, Mark and Van Horn-Morris [11].

The purpose of the present paper is to define a new cut and paste operation, called *star surgery*, which is a strong generalization of Fintushel and Stern’s rational blow-down. Just like rational blow-down, our operation reduces b_2^- of the manifold to which it is applied. Moreover it can be performed symplectically and can be seen as a monodromy substitution.

The original rational blow-down operation, and its generalizations — Park [35], Stipsicz, Szabó and Wahl [43], Bhupal and Stipsicz [7] — amount to removing the neighborhood of a union of spheres which intersect according to a particular plumbing tree and

regluing a rational ball which has the same boundary as the neighborhood. Our star surgery operation is similarly defined. First identify (symplectic) spheres which intersect according to a star-shaped graph with a negativity condition on the central vertex. The star surgery operation cuts out a neighborhood of these spheres and replaces it by an alternate symplectic filling of the induced contact boundary. It is shown by the second author [40] that these alternate fillings always have smaller Euler characteristic than the neighborhood of spheres, and are negative-definite. Unlike the rational blow-down we do not require the alternate filling to be a rational homology ball. This greatly generalizes the set of configurations of spheres which we can consider for these operations.

By reinterpreting this operation as a monodromy substitution, one can show that some star surgeries are obtained by a sequence of rational blow-downs. The spheres to rationally blow down after the first step in the sequence are not all visible in the original configuration, and would be difficult to find. The star surgery bypasses the need to find these spheres by performing a single symplectic cut and paste operation that performs the entire sequence of rational blow-downs simultaneously.

However, there are other star surgeries which are inequivalent to any sequence of symplectic rational blow-downs as proven by a family of examples in [40]. It is expected that many of these star surgery operations cannot be obtained from sequences of rational blow-downs. This contrasts with the operations one would obtain by replacing linear plumbings of spheres by alternate fillings, which were shown by Bhupal and Özbağcı [6] to all be equivalent to sequences of rational blow-downs.

Using star surgery, we construct many examples of exotic 4-manifolds. These constructions involve two steps. First we must find a configuration of symplectic spheres inside a well understood 4-manifold. In our examples we do this by looking at blow-ups of elliptic fibrations $E(1)$ using varying types of singular fibers to find symplectic spheres with the required intersection data. We explicitly construct many elliptic fibrations by blowing up various Lefschetz pencils on $\mathbb{C}P^2$. Then we apply the star surgery operation which replaces this neighborhood of spheres with the smallest symplectic filling of the induced contact boundary Seifert fibered space. By keeping track of the homology classes of all of the spheres in the elliptic fibration, we are able to compute the small perturbation Seiberg–Witten invariants of the resulting manifold.

Using this technique, we construct a minimal symplectic 4-manifold which is an exotic copy of $\mathbb{C}P^2 \# 8 \mathbb{C}P^2$.

Theorem 1.1 *There is a minimal symplectic 4-manifold X which is homeomorphic but not diffeomorphic to $\mathbb{C}P^2 \# 8 \mathbb{C}P^2$ and which is obtained by a star surgery. The symplectic Kodaira dimension of X is 2.*

We push our techniques further with different examples of star surgery operations. These examples yield exotic (potentially nonminimal) symplectic copies of $\mathbb{C}P^2 \# k \overline{\mathbb{C}P^2}$, for $k = 6, 7$.

Theorem 1.2 *There are constructions of symplectic exotic copies of $\mathbb{C}P^2 \# 7 \overline{\mathbb{C}P^2}$ and $\mathbb{C}P^2 \# 6 \overline{\mathbb{C}P^2}$ obtained by performing star surgery operations on blow-ups of $E(1)$.*

Other star surgery operations, including the star surgery which is known to be inequivalent to any sequence of rational blow-downs, can be used for similar constructions. In particular, we show that this star surgery can be used to construct an exotic $\mathbb{C}P^2 \# 8 \overline{\mathbb{C}P^2}$, and related star surgeries can be used to improve these constructions to manifolds with $b_2^- = 6, 7$.

While the star surgery operations are inspired by symplectic topology, they can also be used smoothly in the absence of a symplectic structure. By first performing Fintushel and Stern's knot surgery in a double node neighborhood [16] (which destroys the symplectic structure), and then performing the star surgery as in Theorem 1.1, we prove the following result.

Theorem 1.3 *For every $n \geq 2$ there exist smooth minimal mutually nondiffeomorphic 4-manifolds Y_n which are all homeomorphic to $\mathbb{C}P^2 \# 7 \overline{\mathbb{C}P^2}$. These manifolds are obtained by a star surgery.*

The same technique of first performing knot surgery in a double node neighborhood can be used with the star surgery constructions in the proof of Theorem 1.2 to produce infinitely many smooth but not symplectic nondiffeomorphic manifolds in the homeomorphism classes of $\mathbb{C}P^2 \# 6 \overline{\mathbb{C}P^2}$ and $\mathbb{C}P^2 \# 5 \overline{\mathbb{C}P^2}$.

Examples of (minimal) exotic copies of $\mathbb{C}P^2 \# k \overline{\mathbb{C}P^2}$, for $k = 6, 7, 8$ have previously been constructed using the rational blow-down technique. The exotic structure was detected by calculating the effect of the rational blow-down on the Seiberg–Witten invariants; see Park [36], Stipsicz and Szabó [42], Fintushel and Stern [16], and Michalogiorgaki [30]. The effect of star surgery on Seiberg–Witten invariants is similar to the effect of rational blow-down [30]. The main reason is that the boundary of the star shaped configuration is an L -space. In other words the monopole Floer homology of the boundary of the neighborhood of these configurations of spheres is the simplest group it could be.

Finding exotic copies of $\mathbb{C}P^2 \# k \overline{\mathbb{C}P^2}$ for small k is a problem which has been studied for many years. In the 1980s, gauge theoretic techniques were used to distinguish Dolgachev surfaces from $\mathbb{C}P^2 \# 9 \overline{\mathbb{C}P^2}$, as in Donaldson [9] and Friedman and Morgan [20], and the Barlow surface from $\mathbb{C}P^2 \# 8 \overline{\mathbb{C}P^2}$; see Kotschick [25]. Significant

progress was made using the rational blow-down to construct an exotic $\mathbb{C}P^2 \#_k \overline{\mathbb{C}P^2}$ for $k \geq 5$; see Park [36], Stipsicz and Szabó [42], Park, Stipsicz and Szabó [37], Fintushel and Stern [16], and Michalogiorgaki [30]. Later this was improved to $k \geq 2$ using different techniques; see Akhmedov and Park [4; 3], Akhmedov [1], Akhmedov, Baykur and Park [2], Baldridge and Kirk [5], Fintushel and Stern [18] and Fintushel, Park and Stern [13]. Because these star surgery operations greatly increase the possible configurations of surfaces which can be cut out and replaced, we hope that more star surgery constructions will be found and can be used to improve this bound or exhibit other new phenomena in smooth 4–manifold topology.

The organization is as follows: in Section 2 we define our star surgery operation, and describe the explicit examples which we will use in constructions of exotic 4–manifolds. In Section 3, we determine properties of these star surgery operations by computing algebraic topological invariants of the fillings. In Section 4, we construct three explicit elliptic fibrations which we will use to embed configurations of symplectic spheres on which to perform star surgery. Theorems 1.1 and 1.2 are proven in Section 5, where we use star surgery to construct manifolds and compute their homeomorphism invariants, Kodaira dimension, and Seiberg–Witten invariants. Finally, Theorem 1.3 is proven in Section 6, by using knot surgery and star surgery together.

Acknowledgments We would like to thank Kouichi Yasui and Tian-Jun Li for helpful correspondences. In the course of this work, the first author was supported by the National Science Foundation FRG Grant DMS-1065178 and a TUBITAK grant BIDEB 2232. The second author was supported by a National Science Foundation Graduate Research Fellowship under Grant No. DGE-1110007.

2 Star surgery

2A Description

Rational blow-downs of plumblings of spheres were shown to be symplectic operations by Symington [44; 45] proving that both the plumbing of spheres and the rational homology ball support symplectic structures with convex boundary inducing the same contact structures. One may ask more generally what can replace a neighborhood of spheres in this symplectic cut and paste manner. This question is reduced to understanding symplectic fillings of certain contact structures by the following result of Gay and Stipsicz.

Theorem 2.1 [23, Theorem 1.2] *If $C = C_1 \cup \dots \cup C_n \subset (X, \omega)$ is a collection of symplectic surfaces in a symplectic 4-manifold (X, ω) intersecting each other ω -orthogonally according to the negative-definite plumbing graph Γ and $\nu C \subset X$ is an open set containing C , then C admits an ω -convex neighborhood $U_C \subset \nu C \subset (X, \omega)$.*

Note that the ω -orthogonal condition can be achieved by any configuration of spheres which intersect positively and transversely by an isotopy through symplectic spheres.

The contact structures induced on the boundaries can be understood through an open book decomposition by results of Gay and Mark (under the additional assumption that the plumbing graph contains no bad vertices, ie $w_j + e_j < 0$, where w_j is the weight of a vertex v_j , and e_j is the number of edges emanating from v_j). Let C be the union of symplectic surfaces intersecting ω -orthogonally according to such a graph. Form a surface Σ from the plumbing graph as follows. Start with the surfaces corresponding to each vertex v_j and take the connected sum with $|w_j + e_j|$ disks. Then take the connected sum of the resulting surfaces according to the edges of the graph. Take one simple closed curve around each connected sum neck, and denote these curves by c_1, \dots, c_k .

Theorem 2.2 [21, Theorem 1.1] *Any neighborhood of C contains a neighborhood of C with strongly convex boundary that admits a symplectic Lefschetz fibration having regular fibers Σ and exactly one singular fiber. The vanishing cycles are c_1, \dots, c_k and C is the union of the closed components of the singular fiber. The induced contact structure on the boundary is supported by the induced open book (Σ, τ) , where τ is a composition of positive Dehn twists around the curves c_1, \dots, c_k .*

Note that since the curves c_1, \dots, c_k are disjoint from each other, the order of the Dehn twists does not matter in defining τ .

Remark 2.3 It was shown by Park and Stipsicz [34] that this contact structure is in fact the canonical contact structure on the boundary Seifert fibered space (given by the complex tangencies on the link of the corresponding normal surface singularity). Their result holds more generally for the boundary contact structure of any convex negative-definite plumbing of surfaces.

In the case that the symplectic spheres intersect according to a star-shaped graph, additional tools are available to search for alternate convex fillings of the same convex boundary. Classifications of such fillings were studied in [41]. While most plumbings do not share the same convex boundary with a rational homology ball, many share convex boundary with a symplectic filling of significantly smaller Euler characteristic.

We will call *star surgery* the operation of cutting out the neighborhood of spheres which intersect according to a star-shaped graph, and replacing it with an alternative convex symplectic filling of strictly smaller Euler characteristic.

It is not clear that any contactomorphism of the boundary extends over the alternate convex filling, so the star surgery will depend on an identification of the convex contact boundary of the filling with the concave contact boundary of the complement of the star-shaped plumbing. In our cases, this identification will be made using the open book decomposition defined by [Theorem 2.2](#) and an equivalent open book decomposition on the boundary of the alternate convex filling.

In each case, the neighborhood of spheres will be replaced by an alternate symplectic filling supported by a Lefschetz fibration. The fibers of this Lefschetz fibration will agree with the fibers of the Lefschetz fibration constructed by Gay and Mark on the plumbing neighborhood of spheres. However, the vanishing cycles will differ. We will show that the induced contact structures on the boundary agree by showing that the open book decompositions have equal monodromy. In order to do this, we will require knowledge of relations in the mapping class group of planar surfaces.

2B Conventions on mapping class elements and handle diagrams for Lefschetz fibrations

A Lefschetz fibration naturally induces an open book decomposition on the boundary where the fibers of the open book are the same as the fibers of the Lefschetz fibration, and the monodromy is given by a product of positive (right-handed) Dehn twists about the vanishing cycles. Since mapping class groups of surfaces are nonabelian, the order of the vanishing cycles generally matters. Conventions in the literature vary, but we will use a fixed set of conventions throughout this paper which are consistent with each other, which we describe here.

Suppose c_1, \dots, c_n are simple closed curves on the fiber. Denote by D_{c_i} a positive Dehn twist around c_i . The product $D_{c_1} D_{c_2} \cdots D_{c_n}$ means first Dehn twist along c_1 , then c_2 , and so on until finally along c_n , meaning we are using group notation as opposed to functional notation. When the fiber is a disk with holes, we can place the holes along a circle concentric with the boundary of the disk. Labeling the holes $\{1, \dots, m\}$ counterclockwise, we use the notation D_{i_1, \dots, i_k} for $i_1, \dots, i_k \in \{1, \dots, m\}$ to indicate a positive Dehn twist about a curve which convexly contains the holes i_1, \dots, i_k .

Any factorization of the monodromy of an open book decomposition into a product of positive Dehn twists corresponds to a Lefschetz fibration. When the fibers are




Figure 1: The Lefschetz fibration corresponding to the monodromy factorization $D_{1,3}D_{2,5}D_{4,7} = D_{c_1}D_{c_2}D_{c_3}$. The tops and bottoms of the dotted segments are identified to form a closed trivial braid.

disks with holes, we have the natural handlebody decomposition for this Lefschetz fibration where the holes are represented by dotted circles forming a trivial braid corresponding to 1–handles and the vanishing cycles correspond to 2–handles. We view the holed disk fibers as orthogonal to the dotted circles, oriented so that the outward normal points *downward* (ie turn the holed disk upside-down). Then the monodromy factorization $D_{c_1} \cdots D_{c_n}$ corresponds to the Lefschetz fibration where the vanishing cycles appear as curves, each lying in a disk transverse to the trivial braid of dotted circles such that c_1 is at the top of the diagram and c_n at the bottom.

To draw the handlebody, we will isotope the holes on the disk so that they all lie on the bottom half of the disk along a circle concentric to the boundary. Then using the upside-down disk convention, the holes, ordered counterclockwise on the downward pointing disk, correspond to dotted trivial braid components labeled *left to right*. A curve which convexly encloses holes i_1, \dots, i_k will appear in this diagram as a circle, half of which passes in front of all of the dotted circles, and the other half passes behind the dotted circles corresponding to i_1, \dots, i_k , but in front of all the other circles. An example, using the top to bottom convention where the outward normal to the disk points downward, is in [Figure 1](#).

The mapping class group on a disk with holes is generated by Dehn twists. Dehn twists about disjoint curves commute. If we place the holes on a circle concentric with the boundary, we can order them counter-clockwise. Suppose A , B , and C are collections of holes such that all the holes of A precede all the holes of B which precede all the holes of C going around the circle counterclockwise. Then the *lantern relation* states

$$(2-1) \quad D_{AUBUC}D_AD_BD_C = D_{AUB}D_{AUC}D_{BUC}.$$

The Dehn twists on the right-hand side can be cyclically permuted.

By combining a sequence of lantern relations one obtains the *daisy relation*, which is given as follows. Suppose B_0, B_1, \dots, B_m are disjoint subsets of the k holes on the disk labeled counter-clockwise ($m \geq 2$). Then

$$(2-2) \quad D_{B_0 \cup B_1 \cup \dots \cup B_m} D_{B_0}^{m-1} D_{B_1} \cdots D_{B_m} = D_{B_0 \cup B_1} D_{B_0 \cup B_2} \cdots D_{B_0 \cup B_m} D_{B_1 \cup \dots \cup B_m}.$$

The daisy relations were shown to correspond to Fintushel and Stern’s rational blow-down operations in [11].

We will use one more combination of lantern moves, corresponding to one of Park’s generalized rational blow-downs which starts with a linear plumbing with weights $(-2, -5, -3)$. It was first worked out in [11] that the relation is given by

$$(2-3) \quad D_{AUBUCUDUE} D_{AUB} D_A D_B D_C^2 D_D D_E \\ = D_{AUC} D_{BUC} D_{AUBUD} D_{AUBUE} D_{CUDUE}$$

for a disk whose holes are grouped into sets A, B, C, D, E , labeled counter-clockwise. This equality can be shown by performing one daisy relation (2-2) where $B_0 = A \cup B$ introducing some negative Dehn twists, followed by a lantern relation (2-1).

2C The family of star surgeries $(\mathcal{S}_i, \mathcal{T}_i)$

A particularly nice family of star-shaped surgeries is given by symplectically replacing a neighborhood of a configurations of spheres, \mathcal{S}_i by its smallest filling \mathcal{T}_i (a specific filling of minimal Euler characteristic). The configurations \mathcal{S}_i are made up of symplectic spheres which intersect according to star-shaped graphs with $i + 2$ arms. Each arm contains $i - 1$ spheres of square -2 , and the central vertex is a sphere of square $-i - 3$; see Figure 2(a). Note that \mathcal{S}_1 is just a -4 -sphere, and the replacement \mathcal{T}_1 is the rational blow-down of this -4 -sphere. However for $i > 1$, the graphs are star-shaped but not linear, and the replacement fillings \mathcal{T}_i are not obtained by a rational blow-down of a subgraph of the spheres shown in the original configuration. Handlebody diagrams for the fillings \mathcal{T}_i for $i = 1, 2, 3$ are shown in Figure 3. In general, a handlebody diagram for \mathcal{T}_i (see Figure 9(a)) has $i + 2$ 1-handles represented by dotted circles, and $\frac{1}{2}(i + 1)(i + 2)$ 2-handles, one passing through each distinct pair of 1-handles. The corresponding monodromy factorization is

$$(D_{1,2} D_{1,3} D_{1,4} \cdots D_{1,i+2})(D_{2,3} D_{2,4} \cdots D_{2,i+2}) \cdots (D_{i,i+1} D_{i,i+2})(D_{i+1,i+2}).$$

We can replace \mathcal{S}_i by \mathcal{T}_i symplectically due to the following proposition.




Figure 2: (a) Plumbing graph for S_i and (b) fibers and vanishing cycles for S_i .

Proposition 2.4 *The contact structure induced on the convex boundary of the plumbing of spheres S_i is the same as the contact structure induced on the convex boundary of \mathcal{T}_i (whose symplectic structure is determined by the supporting Lefschetz fibration described above).*

Proof For the plumbing of spheres S_i , Gay and Mark’s construction implies that the fibers of the supporting Lefschetz fibration are $(i+2)$ -holed disks, and the vanishing cycles consist of a single curve parallel to the outer boundary component, and i boundary parallel curves around each of the holes; see Figure 2(b).

In order to show that the contact structures induced on the boundaries of S_i and \mathcal{T}_i agree, we will show that the open book decompositions induced on the boundary of the corresponding Lefschetz fibrations are the same. Since the pages are the same, it suffices to show the monodromies are equal, which amounts to the following relation:

$$\begin{aligned}
 (2-4) \quad D_{1,2,\dots,i+1,i+2} D_1^i \cdots D_{i+2}^i &= (D_{1,2} D_{1,3} D_{1,4} \cdots D_{1,i+2}) (D_{2,3} D_{2,4} \cdots D_{2,i+2}) \cdots \\
 &\quad \cdots (D_{i,i+1} D_{i,i+2}) (D_{i+1,i+2}).
 \end{aligned}$$

This relation is sometimes referred to in the literature as the *generalized lantern relation*.

To see these are equal in the mapping class group, proceed by induction on i . When $i = 1$ this is the standard lantern relation. By a relabeled version of the $i - 1$ case, we can use the inductive hypothesis to say that the right-hand side is equal to

$$(D_{1,2} D_{1,3} D_{1,4} \cdots D_{1,i+2}) D_{2,3,\dots,i+1,i+2} D_2^{i-1} D_3^{i-1} \cdots D_{i+1}^{i-1} D_{i+2}^{i-1}.$$

Applying a daisy relation to this then gives the left-hand side. □




Figure 3: The first three configurations of spheres and their replacement symplectic fillings in the family $(\mathcal{S}_i, \mathcal{T}_i)$. Top left: $\mathcal{S}_1, \mathcal{T}_1$. Top right: $\mathcal{S}_2, \mathcal{T}_2$. Bottom: $\mathcal{S}_3, \mathcal{T}_3$.

Remark 2.5 For the negative-definite star plumblings we consider, the induced contact structure is supported by a planar open book. By work of Wendl [47], any other convex filling is supported by a planar Lefschetz fibration inducing the same open book decomposition on the boundary. Therefore the two Lefschetz fibrations correspond to positive factorizations of the same monodromy. Equivalent elements in a planar mapping class group are always related by some sequence of lantern relations and commutation, but this sequence may pass through factorizations involving negative Dehn twists. If one can obtain one positive factorization from another through a sequence of relations so that at each stage we remain in a positive factorization then the overall symplectic operation is broken down into a sequence of other symplectic operations.




Figure 4: A star-shaped plumbing graph for \mathcal{Q} , and an alternate symplectic filling \mathcal{R} .

For example, this proof of the generalized lantern relation shows that it can be obtained by performing a sequence of daisy and lantern relations, so that after each relation, we still have a positive factorization. Endo, Mark, and Van Horn-Morris [11] showed that daisy relations correspond to Fintushel–Stern rational blow-downs. Therefore these particular star surgeries are equivalent to sequences of rational blow-downs. However, it is not easy to see the existence of all the configurations which are rationally blown-down at each stage, so in applications it would be difficult to find all of these rational blow-downs to perform. Instead, we can just perform the sequence all at once with a single star surgery.

A question one can ask is whether all such star surgery operations arise as sequences of rational blow-downs. Surprisingly, this was shown in the linear case in [6], but it was suspected that star surgery was more general. After the appearance of the first draft of this paper, the second author proved that certain examples of a star surgery cannot be realized as any sequence of symplectic rational blow-downs [40, Theorem 1.2].

2D The star surgery $(\mathcal{Q}, \mathcal{R})$

The following star surgery is related to the $\mathcal{S}_2, \mathcal{T}_2$ star surgery but improves it in the sense that it reduces the Euler characteristic by a larger amount. Let \mathcal{Q} denote the configuration of spheres indicated on the left-hand side of Figure 4. Let ξ_{can} be the canonical contact structure $\partial\mathcal{Q}$.

Proposition 2.6 *There exists a symplectic manifold \mathcal{R} of Euler characteristic 3 with convex boundary, such that the induced contact manifold on the boundary is contactomorphic to $(\partial\mathcal{Q}, \xi_{\text{can}})$.*

Proof Let \mathcal{R} be the 4–manifold given by the handlebody diagram on the right of Figure 4. This particular diagram makes apparent a Lefschetz fibration structure on \mathcal{R} . The fibers of the Lefschetz fibration are 6–holed disks, and the base is a disk. The fibers near the boundary are disks perpendicular to the dotted circles which give the holes. The vanishing cycles are given by the -1 –framed 2–handles. We can verify that the induced open book decomposition on the boundary agrees with the one that is induced on the boundary of the symplectic plumbing given by [21].

We choose a standard 6–holed disk, such that the holes are centered at the vertices of a regular hexagon on the disk. Label the holes with numbers $1, \dots, 6$ going around counter-clockwise. The construction of Gay and Mark [21] indicates that the open book induced on the boundary of the plumbing \mathcal{Q} has pages which are 6–holed disks, with monodromy given by positive Dehn twists about disjoint curves enclosing holes as follows:

$$D_{123456} D_{12} D_1 D_2 D_3^2 D_{45}^2 D_4 D_5 D_6^3.$$

The monodromy induced by the Lefschetz fibration on \mathcal{R} (reading the vanishing cycles from top to bottom) is

$$D_{46} D_{56} D_{145} D_{245} D_{345} D_{123} D_{126} D_{36}.$$

Commuting when needed and then performing a Park relation (2-3) on the plumbing monodromy, where $A = \{4\}$, $B = \{5\}$, $C = \{6\}$, $D = \{1, 2\}$ and $E = \{3\}$, we get an intermediate factorization:

$$D_{46} D_{56} D_{1245} D_{345} D_{1236} D_{45} D_6 D_1 D_2 D_3.$$

Note that this corresponds to a symplectic filling obtained from the original plumbing by rationally blowing down the configuration that comes from $u_{4,1}$, u_0 and the symplectic resolution of the union of $u_{3,1}$ with $u_{3,2}$. Continuing, by commuting terms and performing a lantern relation (2-1), where $A = \{45\}$, $B = \{1\}$ and $C = \{2\}$, we obtain the factorization

$$D_{46} D_{56} D_{145} D_{245} D_{12} D_{345} D_{1236} D_6 D_3.$$

Note this corresponds to rationally blowing down a -4 –sphere which was not visible until after the first rational blow-down. We commute terms and perform one more lantern relation (2-1) (corresponding to blowing down another -4 –sphere), where




Figure 5: A star-shaped plumbing graph for \mathcal{U} , and an alternate symplectic filling \mathcal{V} .

$A = \{12\}$, $B = \{3\}$ and $C = \{6\}$, to obtain the factorization corresponding to the Lefschetz fibration on T :

$$(2-5) \quad D_{46} D_{56} D_{145} D_{245} D_{345} D_{123} D_{126} D_{36}. \quad \square$$

Remark 2.7 The proof makes it clear that \mathcal{R} is obtained from \mathcal{Q} through a sequence of rational blow-downs: one Park rational blow-down of a $(-2, -5, -3)$ configuration, followed by two consecutive rational blow-downs of -4 -spheres.

2E The star surgery $(\mathcal{U}, \mathcal{V})$

Let \mathcal{U} denote the plumbing according to the graph in Figure 5. Let ξ denote the (canonical) contact structure induced on the convex boundary of the plumbing.

Proposition 2.8 *There exists a symplectic manifold \mathcal{V} of Euler characteristic 3 with convex boundary, such that the induced contact manifold on the boundary is contactomorphic to $(\partial\mathcal{U}, \xi_{\text{can}})$.*

Proof Gay and Mark’s construction gives an open book on the boundary of the Lefschetz fibration for the plumbing \mathcal{U} whose pages are 8-holed disks and whose monodromy is given as follows:

$$D_{12345678} D_{12}^3 D_1 D_2 D_{34}^2 D_3 D_4 D_{56}^2 D_5 D_6 D_{78} D_7 D_8.$$




Figure 6: The \mathcal{K} plumbing and alternate filling \mathcal{L} , providing a star surgery operation which is inequivalent to a sequence of rational blow-downs [40].

Using the monodromy equivalence corresponding to the Park $(-2, -5, -3)$ rational blow-down, where $A = \{1\}$, $B = \{2\}$, $C = \{3, 4\}$, $D = \{5, 6\}$, $E = \{7, 8\}$, we get the following monodromy:

$$D_{134}D_{234}D_{1256}D_{1278}D_{345678}D_{12}^2D_3D_4D_{56}D_5D_6D_7D_8.$$

After commuting Dehn twists about disjoint curves, we can perform two lantern relations, one where $A = \{1, 2\}$, $B = \{5\}$, and $C = \{6\}$ and the other where $A = \{1, 2\}$, $B = \{7\}$, and $C = \{8\}$, which results in the following factorization:

$$D_{134}D_{234}D_{125}D_{126}D_{56}D_{127}D_{128}D_{78}D_{345678}D_3D_4D_{56}.$$

After commuting D_{56} and D_{78} towards the end, we can use a daisy relation with $B_0 = \{5, 6\}$, $B_1 = \{7, 8\}$, $B_2 = \{3\}$, and $B_3 = \{4\}$. The resulting monodromy corresponds to that of the Lefschetz fibration for \mathcal{V} :

(2-6) $D_{134}D_{234}D_{125}D_{126}D_{127}D_{128}D_{5678}D_{356}D_{456}D_{3478}.$ □

Remark 2.9 Note that this proof shows that this operation is also obtained as a sequence of rational blow-downs.

2F The star surgeries $(\mathcal{K}, \mathcal{L})$, $(\mathcal{M}, \mathcal{N})$ and $(\mathcal{O}, \mathcal{P})$

In [40], it was shown that a configuration of symplectic spheres intersecting according to a graph as in Figure 6 can be replaced by a symplectic filling of Euler characteristic two, whose Lefschetz fibration handlebody is shown in Figure 6.

The corresponding monodromy substitution for this \mathcal{K}, \mathcal{L} star surgery is

(2-7) $D_1^2D_2^2D_3D_4^2D_5^2D_{12345} = D_{123}D_{14}D_{15}D_{24}D_{25}D_{345}.$




Figure 7: The \mathcal{M}, \mathcal{N} star surgery. The dotted circles are labeled left to right as $1, 2_a, 2_b, 3, 4_a, 4_b, 5$.

The equivalence of these elements was shown directly in [40] and it was also shown that this substitution is not equivalent to any sequence of symplectic rational blow-downs/ups. From this star surgery, we can generate two other useful star surgeries, such that the alternate fillings also have Euler characteristic two. The plumbing and filling diagrams are shown in Figures 7 and 8.

These monodromy substitutions are obtained from the \mathcal{K}, \mathcal{L} substitution (2-7) together with an additional lantern relation applied according to [11, Lemma 2.1], where some of the original holes split into two holes (eg hole 2 splits into holes 2_a and 2_b). When applying the \mathcal{K}, \mathcal{L} monodromy substitution, treat the split holes as a single joined hole, but when applying the lantern relations, treat them as separate holes. Performing two splittings and applying [11, Lemma 2.1] each time, we get a monodromy substitution

$$D_1^2 D_{2_a 2_b}^2 D_{2_a} D_{2_b} D_3 D_{4_a 4_b} D_{4_a} D_{4_b} D_5^3 D_{12_a 2_b 3 4_a 4_b 5} \\ = D_{12_a 2_b 3} D_{14_a 4_b} D_{15} D_{2_a 2_b 4_a} D_{2_a 2_b 4_b} D_{2_a 5} D_{2_b 5} D_{3 4_a 4_b 5}$$

corresponding to the \mathcal{M}, \mathcal{N} star surgery of Figure 7. Here, the holes are labeled $1, 2_a, 2_b, 3, 4_a, 4_b, 5$ counter-clockwise around the disk. Performing two more splits, we get the monodromy substitution

$$D_{1_a 1_b}^2 D_{1_a} D_{1_b} D_{2_a 2_b}^2 D_{2_a} D_{2_b} D_3 D_{4_a 4_b}^2 D_{4_a} D_{4_b} D_{5_a 5_b}^2 D_{5_a} D_{5_b} D_{1_a 1_b 2_a 2_b 3 4_a 4_b 5_a 5_b} \\ = D_{1_a 1_b 2_a 2_b 3} D_{1_a 4_a 4_b} D_{1_b 4_a 4_b} D_{1_a 1_b 5_a} D_{1_a 1_b 5_b} D_{2_a 2_b 4_a} D_{2_a 2_b 4_b} \\ D_{2_a 5_a 5_b} D_{2_b 5_a 5_b} D_{3 4_a 4_b 5_a 5_b}$$




Figure 8: The \mathcal{O}, \mathcal{P} star surgery. The dotted circles are labeled left to right as $1_a, 1_b, 2_a, 2_b, 3, 4_a, 4_b, 5_a, 5_b$.

corresponding to the \mathcal{O}, \mathcal{P} star surgery, as shown in Figure 8 on the disk with holes labeled $1_a, 1_b, 2_a, 2_b, 3, 4_a, 4_b, 5_a, 5_b$ counter-clockwise. Note that the proof that these factorizations are equivalent involves applying the substitution from Equation (2-7) and then lantern relations, but the first step will introduce negative Dehn twists into the factorization, so there is not a simple way to understand these operations as a sequence of known operations (though it would be more difficult to prove that they are not equivalent to sequences of known operations; see [41; 40] for an idea of how this might be proven).

3 Algebraic topology of the star surgery fillings

In this section we will compute the fundamental group of the fillings $\mathcal{T}_i, \mathcal{R}, \mathcal{V}, \mathcal{L}$, and various other algebraic topology invariants which will be needed to understand the homeomorphism type, Kodaira dimension, and Seiberg–Witten invariants of the manifolds constructed by star surgeries using these fillings. The computations are reasonably straightforward given the handlebody descriptions of these manifolds and the Lefschetz fibration structure. The most thorough computations will be given for \mathcal{T}_2 , as this will be our model example used to show how star surgery can be applied to create exotic manifolds whose Kodaira dimension and Seiberg–Witten invariants can be fully computed.

3A Properties of the fillings \mathcal{T}_i

We will compute basic algebraic topological properties of \mathcal{T}_i , and specifically \mathcal{T}_2 since we will use the $\mathcal{S}_2, \mathcal{T}_2$ star surgery to construct an exotic copy of $\mathbb{C}P^2 \# 8 \overline{\mathbb{C}P^2}$.

Proposition 3.1 *The fillings \mathcal{T}_i satisfy $\pi_1(\mathcal{T}_i) = \mathbb{Z}/2\mathbb{Z}$. The generator can be represented by the meridian of any of the -2 -framed surgery curves on the ends of the arms in $\partial\mathcal{S}_i$.*

Proof A handlebody diagram for \mathcal{T}_i is given as in Figure 9(a) by $(i+2)$ 1-handles corresponding to generators $\{y_1, \dots, y_{i+2}\}$ of $\pi_1(\mathcal{T}_i)$, and $\frac{1}{2}(i+1)(i+2)$ 2-handles corresponding to the relations $\{y_j y_k = 1\}_{j \neq k \in \{1, \dots, i+2\}}$. We can easily compute the fundamental group:

$$\pi_1(\mathcal{T}_i) = \langle y_1, \dots, y_{i+2} : y_j = y_k^{-1}, j \neq k \rangle = \langle y_1 : y_1^2 = 1 \rangle.$$

Now we will track a curve representing the generator y_1 of $\pi_1(\mathcal{T}_i)$ through a sequence of equivalent surgery diagrams, to show that it restricts to a nontrivial element in $\partial\mathcal{T}_i = \partial\mathcal{S}_i$ represented by the meridian of a -2 -framed surgery curve on the end of the first arm in the standard handlebody diagram for \mathcal{S}_i . Note that π_1 is equivalently generated by any of the y_j with the relation $y_j^2 = 1$, and that y_j restricts in the same way to the meridian of the last curve in the j^{th} arm.

Start with the handlebody diagram in Figure 9(a). The generator of the fundamental group, y_1 , is represented by the dashed red curve. By rotating the plane of projection about a vertical axis, observe this handlebody diagram is isotopic to that given by Figure 9(b) (the direction into the page in Figure 9(a) corresponds to the right side of the page in Figure 9(b)). Now, exchange the dotted circles for 0-framed circles and treat the handlebody diagram as a surgery diagram for its boundary 3-manifold. After blowing down all the -1 -framed 2-handles, we obtain a surgery diagram as in Figure 9(c) containing $i+2$ unknotted $(i+1)$ -framed circles, twisted together with a full positive twist. Blowing up negatively once at a common intersection point of their Seifert surfaces, and then i times along each individual curve, the curves become untwisted and 0-framed again, so we switch the 0-framings to dotted circles as in Figure 9(d). Perform handle slides by first sliding the dashed red curve over the top -1 -framed 2-handle on that dotted circle, then sliding that 2-handle over the one below it, and so on until there is a chain of -2 curves linked to a single -1 curve. Do this for each arm (without the red reference curve on the other arms), and then finally slide the -1 -framed handle that links all of the dotted circles over each of the remaining -1 -framed 2-handles. Then we see that after canceling 1, 2-handle pairs, the dashed red curve appears as the meridian of the last -2 -framed curve as in Figure 9(e). \square




Figure 9: Relating diagrams for $\partial\mathcal{T}_i$ and $\partial\mathcal{S}_i$. We display (a) \mathcal{T}_i , (b) an isotopic diagram in a different projection, (c) the surgery diagram for the common boundary, (d) \mathcal{S}_i Lefschetz fibration, and (e) an \mathcal{S}_i equivalent diagram showing the plumbing.

Proposition 3.2 *The second homology of the fillings is given by $H_2(\mathcal{T}_2; \mathbb{Z}) = \mathbb{Z} \oplus \mathbb{Z}$. The generators can be represented by tori. Moreover, $\chi(\mathcal{T}_2) = 3$ and $\sigma(\mathcal{T}_2) = -2$.*

Proof Using the handlebody diagram as in Figure 9(a) but ignoring the reference curve, we can compute the CW chain complex. Let y_j denote the j^{th} 1-handle and let x_{jk} denote the 2-handle whose attaching circle passes through the j^{th} and k^{th} 1-handles. Then the relevant chain groups are $C_2(\mathcal{T}_i) = \langle x_{jk} : j \neq k \in \{1, \dots, i+2\} \rangle$, $C_1(\mathcal{T}_i) = \langle y_j : j \in \{1, \dots, i+2\} \rangle$. The boundary map is determined by $\partial x_{jk} = y_j + y_k$. In particular, the Euler characteristic is

$$\chi(\mathcal{T}_i) = \frac{1}{2}(i+1)(i+2) - (i+2) + 1.$$

When $i = 2$, $\chi(\mathcal{T}_2) = 6 - 4 + 1 = 3$ and the 2-cycles are generated freely by $x_{12} + x_{34} - x_{13} - x_{24}$ and $x_{14} + x_{23} - x_{12} - x_{34}$. Note that each x_{jk} has square -1 so the intersection form with respect to the above basis is given by the matrix

$$\begin{bmatrix} -4 & 2 \\ 2 & -4 \end{bmatrix} \sim_{\mathbb{R}} \begin{bmatrix} -4 & 0 \\ 0 & -3 \end{bmatrix}.$$

Therefore \mathcal{T}_2 is negative-definite.

We can see that the homology class $x_{12} + x_{34} - x_{13} - x_{24}$ can be represented by tori by examining the handlebody diagram. Take the cores of the 2-handles x_{12} and x_{34} , and the cores with opposite orientation for x_{13} and x_{14} . We connect these up to a closed torus by adding in the twice punctured disks whose outer boundary coincides with the attaching circle for x_{ij} which does not intersect any of the dotted circles, and tubing together the holes with a tube encircling the dotted circle so the orientations match up as in Figure 10. One can see this surface is indeed a torus directly or check by calculating its Euler characteristic: $4\chi(D^2) + 4\chi(2\text{-holed } D^2) + 4\chi(\text{tube}) = 4 - 4 + 0 = 0$. A similar surface represents the other generator $x_{14} + x_{23} - x_{12} - x_{34}$. \square

Proposition 3.3 *The canonical class of the filling vanishes: $K|_{\mathcal{T}_2} = 0$.*

Proof We will use the obstruction theoretic interpretation of the first Chern class. See [24, Proposition 2.3] and [12, Section 3.1] for related discussions. Figure 9(a) suggests that each \mathcal{T}_i admits a positive allowable Lefschetz fibration whose fibers are disks with $i+2$ punctures, which are indicated by the circles with dots, and $\frac{1}{2}(i+2)(i+1)$ vanishing cycles which are indicated by -1 -framed 2-handles. This Lefschetz fibration defines an almost complex structure which is compatible with the symplectic structure of \mathcal{T}_i . We will show that $c_1(T\mathcal{T}_i, J) = 0$ for $i = 2$.

Let x_{jk} be a 2-cell which is a generator of the chain complex $C_2(\mathcal{T}_i)$ as indicated in the proof of Proposition 3.2. The attaching curve \tilde{x}_{jk} of each 2-cell x_{jk} can be put on a fiber F_{jk} of the Lefschetz fibration. Drawing the regular fibers on planes




Figure 10: A torus representing a generator of the homology of \mathcal{T}_2 .

induces trivializations of their tangent bundles which in turn induce a trivialization of $(T\mathcal{T}_i, J)$ over 1–skeleton. Now $c_1(T\mathcal{T}_i, J)(x_{jk})$ is the obstruction to extending this trivialization over the 2–cell x_{jk} . This obstruction is precisely the winding number of \tilde{x}_{jk} measured with respect to the trivialization of F_{jk} . Each \tilde{x}_{jk} is an embedded planar curve, so its winding number is one. For $i = 2$ the proof of Proposition 3.2 tells us that the generators of $H_2(\mathcal{T}_2, \mathbb{Z})$ are $x_{12} + x_{34} - x_{13} - x_{24}$ and $x_{14} + x_{23} - x_{12} - x_{34}$. Hence $c_1(T\mathcal{T}_i, J)$ evaluates as zero on both of these generators. \square

We will need to understand $H^2(\mathcal{T}_2)$ and $H^2(\partial\mathcal{T}_2)$, as well as the restriction map between them. By Poincaré duality, $H^2(\partial\mathcal{T}_2) \cong H_1(\partial\mathcal{T}_2)$. We can compute the first homology of the boundary of the filling explicitly from the surgery diagram obtained from the handlebody diagram of \mathcal{T}_2 by switching the dotted circles to 0–framed circles as in Figure 11.

Proposition 3.4 *Let $\lambda_1, \lambda_2, \lambda_3, \lambda_4$ denote the meridians of the four 0–framed curves (from left to right) in the surgery diagram in Figure 11. Then*

$$\begin{aligned}
 H_1(\partial\mathcal{T}_2; \mathbb{Z}) &= \langle \lambda_1, \lambda_2 - \lambda_1, \lambda_3 - \lambda_1 \mid 2(\lambda_2 - \lambda_1) = 2(\lambda_3 - \lambda_1) = 12\lambda_1 = 0 \rangle \\
 &\cong \mathbb{Z}/12 \oplus \mathbb{Z}/2 \oplus \mathbb{Z}/2.
 \end{aligned}$$

Proof Let μ_{ij} denote the meridian of the -1 –framed curve which links the i^{th} and j^{th} 0–framed curves ($i \neq j \in \{1, 2, 3, 4\}$).




Figure 11: A surgery diagram for $\partial\mathcal{T}_2$.

A presentation for $H_1(\partial\mathcal{T}_2)$ has generators $(\lambda_1, \lambda_2, \lambda_3, \lambda_4, \mu_{12}, \mu_{13}, \mu_{14}, \mu_{23}, \mu_{24}, \mu_{34})$ with relations given by the linking matrix:

$$\begin{aligned} -\mu_{ij} + \lambda_i + \lambda_j &= 0, \\ \mu_{12} + \mu_{13} + \mu_{14} &= 0, \\ \mu_{12} + \mu_{23} + \mu_{24} &= 0, \\ \mu_{13} + \mu_{23} + \mu_{34} &= 0, \\ \mu_{14} + \mu_{24} + \mu_{34} &= 0. \end{aligned}$$

Eliminate the μ_{ij} using the first relations and then solve for λ_4 . The resulting presentation for $H_1(\partial\mathcal{T}_2; \mathbb{Z})$ is

$$\langle \lambda_1, \lambda_2, \lambda_3 \mid 2(\lambda_2 - \lambda_1) = 2(\lambda_3 - \lambda_1) = -8\lambda_1 - 2\lambda_2 - 2\lambda_3 = 0 \rangle.$$

Equivalently $\langle \lambda_1, \lambda_2 - \lambda_1, \lambda_3 - \lambda_1 \mid 2(\lambda_3 - \lambda_1) = 2(\lambda_2 - \lambda_1) = 12\lambda_1 = 0 \rangle$. □

Proposition 3.5 $H^2(\mathcal{T}_2) \cong \mathbb{Z} \oplus \mathbb{Z} \oplus \mathbb{Z}/2$

Proof The d -cells of \mathcal{T}^2 are generators for the d^{th} CW homology chain complex. Their duals freely generate $C^d(\mathcal{T}_2)$. Let y_1, y_2, y_3, y_4 denote the 1-handles in the diagram for \mathcal{T}_2 (from left to right), and let x_{ij} denote the 2-handle whose attaching circle passes over y_i and y_j . Let ϕ^i denote the dual of y_i , and ψ^{ij} denote the dual of x_{ij} so $C^1(\mathcal{T}_2) = \langle \phi^1, \phi^2, \phi^3, \phi^4 \rangle$ and $C^2(\mathcal{T}_2) = \langle \psi^{12}, \psi^{13}, \psi^{14}, \psi^{23}, \psi^{24}, \psi^{34} \rangle$ and $C^k(\mathcal{T}^2) = 0$ for $k > 2$. Then

$$(\delta\phi^i)(x_{jk}) = \phi^i(\partial x_{jk}) = \phi^i(y_j + y_k) = \delta_j^i + \delta_k^i = \left(\sum_{\ell \neq i} \psi^{i\ell} \right)(x_{jk}).$$

Therefore $\delta\phi^i = \sum_{\ell \neq i} \psi^{i\ell}$. (Note we identify $\psi^{ij} = \psi^{ji}$.)

We conclude

$$H^2(\mathcal{T}_2) = \langle \psi^{12}, \psi^{13}, \psi^{14}, \psi^{23}, \psi^{24}, \psi^{34} \mid \psi^{12} + \psi^{13} + \psi^{14} = \psi^{12} + \psi^{23} + \psi^{24} \\ = \psi^{13} + \psi^{23} + \psi^{34} = \psi^{14} + \psi^{24} + \psi^{34} = 0 \rangle.$$

By eliminating variables we obtain

$$H^2(\mathcal{T}_2) = \langle \psi^{12}, \psi^{13}, \psi^{12} + \psi^{13} + \psi^{23} \mid 2(\psi^{12} + \psi^{13} + \psi^{23}) = 0 \rangle \cong \mathbb{Z} \oplus \mathbb{Z} \oplus \mathbb{Z}/2. \quad \square$$

Proposition 3.6 *The image of the restriction map $i: H^2(\mathcal{T}_2) \rightarrow H^2(\partial\mathcal{T}_2)$ has index 2 (therefore order 24).*

Proof The restriction map $i: H^2(\mathcal{T}_2) \rightarrow H^2(\partial\mathcal{T}_2)$ composed with Poincaré duality yields a map $\rho = \text{PD} \circ i: H^2(\mathcal{T}_2) \rightarrow H_1(\partial\mathcal{T}_2)$. On generators ψ^{ij} we have $\rho(\psi^{ij}) = \mu_{ij}$, where μ_{ij} is the meridian of the surgery curve corresponding to the attaching circle for x_{ij} . In $H_1(\partial\mathcal{T}_2)$ we had the relation $\mu_{ij} = \lambda_i + \lambda_j$ so

$$\rho(\psi^{12}) = (\lambda_2 - \lambda_1) + 2\lambda_1, \quad \rho(\psi^{13}) = (\lambda_3 - \lambda_1) + 2\lambda_1, \\ \rho(\psi^{12} + \psi^{13} + \psi^{23}) = 2(\lambda_2 - \lambda_1) + 2(\lambda_3 - \lambda_1) + 6\lambda_1 = 6\lambda_1 \in H_1(\partial\mathcal{T}_2).$$

Therefore the image of ρ (which equals the image of i) is generated by $(\lambda_2 - \lambda_1, \lambda_3 - \lambda_1, 2\lambda_1)$, which has index 2 in $H_1(\partial\mathcal{T}_2)$. □

3B Properties of \mathcal{R}

Lemma 3.7 *The filling \mathcal{R} is simply connected.*

Proof Using the handlebody decomposition, a presentation for $\pi_1(\mathcal{R})$ is

$$\langle y_1, \dots, y_6 \mid y_3y_6 = y_4y_6 = y_5y_6 = y_1y_4y_5 = y_2y_4y_5 \\ = y_3y_4y_5 = y_1y_2y_3 = y_1y_2y_6 = 1 \rangle.$$

We can eliminate y_1, y_2, y_3, y_4, y_5 with the first five relations, resulting in the presentation

$$\langle y_6 \mid y_6^{-3} = y_6^3 = y_6^5 = 1 \rangle.$$

Since 3 and 5 are relatively prime, this is a presentation for the trivial group. □

Lemma 3.8 *The intersection form on $H_2(\mathcal{R}; \mathbb{Z})$ is negative-definite. In fact, there exist homology classes α and β which freely generate $H_2(\mathcal{R}; \mathbb{Z}) \cong \mathbb{Z}^2$, such that the intersection form with respect to the basis $\langle \alpha, \beta \rangle$ for $H_2(\mathcal{R}; \mathbb{Z})$ is given by*

$$\begin{bmatrix} -10 & -23 \\ -23 & -79 \end{bmatrix}.$$

Proof Labeling the 2–handles from top to bottom as x_1, \dots, x_8 , the cycles generating homology from the CW complex given by the handlebody decomposition are $\alpha = x_1 + x_2 - x_3 - x_4 + x_5 + x_6 - 2x_8$ and $\beta = x_1 + x_2 - 3x_3 - 3x_4 + 5x_5 + 3x_7 - 5x_8$. Since $x_i^2 = -1$ and the x_i are orthogonal with respect to the intersection pairing, the intersection form can be computed directly. \square

3C Properties of \mathcal{V}

Lemma 3.9 *The filling \mathcal{V} is simply connected.*

Proof Using the handlebody decomposition, a presentation for $\pi_1(\mathcal{V})$ is

$$\langle y_1, \dots, y_8 \mid y_1 y_3 y_4 = y_2 y_3 y_4 = y_1 y_2 y_5 = y_1 y_2 y_6 = y_1 y_2 y_7 = y_1 y_2 y_8 \\ = y_5 y_6 y_7 y_8 = y_3 y_5 y_6 = y_4 y_5 y_6 = y_3 y_4 y_7 y_8 = 1 \rangle.$$

Using the first six relations to eliminate $y_1, y_2, y_5, y_6, y_7, y_8$, this simplifies to

$$\langle y_3, y_4 \mid (y_3 y_4)^8 = y_3 (y_3 y_4)^4 = y_4 (y_3 y_4)^4 = (y_3 y_4)^5 = 1 \rangle.$$

Since 5 and 8 are relatively prime, $y_3 y_4 = 1$. Thus $y_3 = y_4 = 1$ so the group is trivial. \square

Lemma 3.10 *The intersection form on $H_2(\mathcal{V}; \mathbb{Z})$ is negative-definite. Moreover, $H_2(\mathcal{V}; \mathbb{Z}) \cong \mathbb{Z}^2$ and the intersection form with respect to a basis is*

$$\begin{bmatrix} -30 & 5 \\ 5 & -49 \end{bmatrix}.$$

Proof Labeling the 2–handles from top to bottom as x_1, \dots, x_{10} , two cycles generating homology from the CW complex given by the handlebody decomposition are $-2x_1 - 2x_2 + x_3 + x_4 - 3x_7 + x_8 + x_9 + 3x_{10}$ and $-3x_3 - 3x_4 + 3x_5 + 3x_6 - x_7 + 2x_8 + 2x_9 - 2x_{10}$. Since $x_i^2 = -1$ and the x_i are orthogonal with respect to the intersection pairing, the intersection form can be computed directly. \square

3D Properties of \mathcal{L}

We compute the fundamental group and second homology of the filling \mathcal{L} of Figure 6.

Proposition 3.11 *$\pi_1(\mathcal{L}) \cong \mathbb{Z}/4$, and the generator restricts to the boundary Seifert fibered space as a meridian of any of the -2 –surgery curves in the plumbing diagram.*

Proof Using the handlebody decomposition of Figure 6, we obtain a presentation for $\pi_1(\mathcal{L})$ generated by the five 1–handles with relations given by the 2–handles as follows:

$$\langle y_1, y_2, y_3, y_4, y_5 \mid y_1 y_2 y_3 = y_1 y_4 = y_1 y_5 = y_2 y_4 = y_2 y_5 = y_3 y_4 y_5 = 1 \rangle,$$

which simplifies to

$$\langle y_1, y_2, y_3 \mid y_1 y_2 y_3 = y_2 y_1^{-1} = y_3 y_2^{-2} = 1 \rangle,$$

which again simplifies to

$$\langle y_1 \mid y_1^4 = 1 \rangle.$$

Note that the relations set y_1, y_2, y_4^{-1} and y_5^{-1} all equal. Each of these curves can be isotoped into the boundary of \mathcal{L} . By performing blow-downs, handle-slides, and handle cancellations, in a similar manner to Proposition 3.1, we see that these curves in $\partial\mathcal{L}$ are meridians of the -2 –surgery curves in the diagram for $\partial\mathcal{K}$. \square

Proposition 3.12 $H_2(\mathcal{L}) \cong \mathbb{Z}$. It is generated by an element $x_{14} - x_{15} - x_{24} + x_{25}$ of square -4 , represented by a torus obtained by gluing tubes to the cores of the specified 2–handles x_{ij} which link the i^{th} and j^{th} dotted circles in Figure 6.

The proof is a direct computation from the handlebody decomposition of Figure 6.

Proposition 3.13 Using the symplectic structure induced by the Lefschetz fibration on \mathcal{L} , we have $c_1(\mathcal{L}) = 0$, therefore the canonical class is trivial.

Proof This follows from the winding number interpretation of $c_1(\mathcal{L})$ for Lefschetz fibrations, and the fact that the generator of $H_2(\mathcal{L})$ passes over two Lefschetz 2–handles with $+1$ multiplicity and two 2–handles with -1 multiplicity. \square

4 Elliptic fibrations

In order to find symplectic embeddings of the plumbings into well-understood symplectic 4–manifolds, we will use many different elliptic fibrations exhibiting various types of singular fibers. These fibrations were classified by Persson in [38], providing a full list of possible configurations of singular fibers. However, in order to keep track of homology classes of symplectic spheres, we need to explicitly construct these elliptic fibrations by blowing up a special Lefschetz pencil in $\mathbb{C}P^2$.

Lemma 4.1 *There is an elliptic fibration on $E(1) = \mathbb{C}P^2 \# 9 \overline{\mathbb{C}P^2}$ with one I_3 fiber, one I_0^* fiber, three fishtail fibers, and a section. Labeling the components of the I_0^* fiber as S_1, \dots, S_5 where S_5 intersects the E_9 section, and S_4 intersects S_1, S_2, S_3 and S_5 , and labeling the components of the I_3 fiber as V_1, V_2, V_3 where V_1 intersects the E_9 section, the homology classes are as follows:*

$$\begin{aligned} [S_1] &= h - e_1 - e_2 - e_3, & [V_1] &= h - e_1 - e_8 - e_9, \\ [S_2] &= h - e_1 - e_4 - e_5, & [V_2] &= h - e_2 - e_4 - e_6, \\ [S_3] &= h - e_1 - e_6 - e_7, & [V_3] &= h - e_3 - e_5 - e_7, \\ [S_4] &= e_1 - e_8, \\ [S_5] &= e_8 - e_9. \end{aligned}$$

Proof Let C_1 be a reducible degree-three algebraic curve in $\mathbb{C}P^2$ defined by a homogeneous polynomial p_1 and made up of three complex projective lines L_1, L_2 and L_3 , which share a single common intersection point $p \in \mathbb{C}P^2$. Let C_2 be a reducible degree-three algebraic curve defined by a homogeneous polynomial p_2 and made up of three lines L_4, L_5 and L_6 , such that L_4 passes through p , and L_5 and L_6 each intersect all other L_i generically in double points; see Figure 12(a). Note that the homology class of L_i for $i = 1, \dots, 6$ is the generator of $H_2(\mathbb{C}P^2; \mathbb{Z})$, h . Define a Lefschetz pencil on $\mathbb{C}P^2$ by setting $C_{[t_1:t_2]} = \{t_1 p_1 + t_2 p_2 = 0\}$. Note the base locus is the set of points where C_1 intersects C_2 .

Blow up at p , and let the exceptional sphere represent the homology class e_1 . Then the proper transforms of L_1, L_2, L_3 and L_4 represent $h - e_1$ in homology, and L_5 and L_6 are unchanged; see Figure 12(b). Therefore the proper transform \tilde{C}_1 represents $3h - 3e_1$ in homology and the proper transform \tilde{C}_2 represents $3h - e_1$. We redefine C_1 to be the curve given by \tilde{C}_1 together with the exceptional class E_1 with multiplicity two, and let $C_2 = \tilde{C}_2$; see Figure 12(c). Then the curves C_1 and C_2 represent the same class in homology so they define a new Lefschetz pencil as before.

Now blow up at the six intersection points of L_1, L_2, L_3 with L_5, L_6 , so that the exceptional classes are labeled e_2, \dots, e_7 ; see Figure 12(d). Redefine C_1 and C_2 as the proper transforms, and note that these curves both represent the homology class $3h - e_1 - e_2 - \dots - e_7$ and thus define a Lefschetz pencil.

There is still a nonempty base locus since the exceptional sphere E_1 is now part of C_1 which intersects the proper transform of L_4 . We blow up at this point to obtain a new exceptional sphere E_8 ; see Figure 12(e). The proper transform of E_1 represents $e_1 - e_8$, so the proper transform of C_1 represents $3h - e_1 - e_2 - \dots - e_7 - 2e_8$ whereas the proper transform of C_2 represents $3h - e_1 - e_2 - \dots - e_8$. Redefining C_1 as its




Figure 12: Blowing up the Lefschetz pencil to a fibration. The orange curves represent C_1 and the purple curves represent C_2 .

proper transform together with the exceptional sphere E_8 with multiplicity one, and C_2 as its proper transform, the two curves again define a Lefschetz pencil, but still intersect where E_8 meets L_4 ; see Figure 12(f). We blow up one more time at this point, and the resulting proper transforms of C_1 and C_2 are homologous and do not intersect: Figure 12(g).

The resulting elliptic fibration has singular fibers C_1 and C_2 which are of type I_0^* and I_3 respectively, as well as other singular fibers, which we can perturb to generic fishtail fibers. An Euler characteristic computation implies that there are three fishtail fibers. A schematic for the singular fibers in this elliptic fibration together with the section E_9 is given by Figure 12(h). □

Next, we will produce an elliptic fibration on $E(1)$ with two I_2 fibers and two I_4 fibers, and specify the homology classes of the spheres in the singular fibers.

Lemma 4.2 *There exists an elliptic fibration on $E(1) \cong \mathbb{C}P^2 \# 9 \overline{\mathbb{C}P^2}$, with two I_2 fibers and two I_4 fibers such that the spheres in these singular fibers represent homology classes as follows:*

- The first I_2 fiber is made up of two -2 -spheres \tilde{C}_1 and \tilde{L}_1 such that

$$[\tilde{C}_1] = 2h - e_1 - e_2 - e_3 - e_4 - e_7 - e_8, \quad [\tilde{L}_1] = h - e_5 - e_6 - e_9.$$

- The second I_2 fiber is made up of two -2 -spheres \tilde{C}_2 and \tilde{L}_2 such that

$$[\tilde{C}_2] = 2h - e_1 - e_2 - e_3 - e_4 - e_5 - e_6, \quad [\tilde{L}_2] = h - e_7 - e_8 - e_9.$$

- The first I_4 fiber is made up of four -2 -spheres, $\tilde{X}_3, \tilde{Y}_3, \tilde{Z}_3, \tilde{E}_3$ with homology classes

$$[\tilde{X}_3] = h - e_3 - e_6 - e_8, \quad [\tilde{Y}_3] = h - e_3 - e_5 - e_7, \quad [\tilde{Z}_3] = h - e_1 - e_2 - e_9, \quad [\tilde{E}_3] = e_3 - e_4.$$

- The second I_4 fiber is made up of four -2 -spheres, $\tilde{X}_4, \tilde{Y}_4, \tilde{Z}_4, \tilde{E}_4$ with homology classes

$$[\tilde{X}_4] = h - e_1 - e_6 - e_7, \quad [\tilde{Y}_4] = h - e_1 - e_5 - e_8, \quad [\tilde{Z}_4] = h - e_3 - e_4 - e_9, \quad [\tilde{E}_4] = e_1 - e_2.$$

Moreover, this fibration admits a section whose homology class is e_2 . See Figure 13 for an illustration.

Proof Consider the degree-three homogeneous polynomials in three complex variables $p_1(x, y, z) = (yz - x^2)(y - z)$ and $p_2(x, y, z) = (yz + x^2)(y + z)$. Now define $p_{(t_1, t_2)}(x, y, z) = t_1 p_1(x, y, z) + t_2 p_2(x, y, z)$. Then the degree-three curves $T_{[t_1: t_2]} := \{p_{(t_1, t_2)} = 0\} \subset \mathbb{C}P^2$ are the fibers of a Lefschetz pencil on $\mathbb{C}P^2$. Observe that $T_{[1:0]} = \{p_1 = 0\}$ and $T_{[0:1]} = \{p_2 = 0\}$ are reducible curves each made up of a linear part and an irreducible quadratic part. Let $C_1 = \{yz - x^2 = 0\}$ denote the quadratic part of $T_{[1:0]}$ and $L_1 = \{y - z = 0\}$ denote the linear part. Similarly, let $C_2 = \{yz + x^2 = 0\}$ and $L_2 = \{y + z = 0\}$ denote the quadratic and linear parts of $T_{[0:1]}$ respectively.

We will keep track of two more fibers in this Lefschetz fibration:

$$T_{[-1:1]} = \{(x + iz)(x - iz)y = 0\}, \quad T_{[1:1]} = \{(y + ix)(y - ix)z = 0\}.$$

These are reducible curves, each made up of three linear parts which we will label as $X_3 = \{x + iz = 0\}$, $Y_3 = \{x - iz = 0\}$, $Z_3 = \{y = 0\}$, $X_4 = \{x + iy = 0\}$, $Y_4 = \{x - iy = 0\}$, and $Z_4 = \{z = 0\}$.




Figure 13: An elliptic fibration with two I_2 fibers and two I_4 fibers.

	C_2	L_2	X_3	Y_3	Z_3	X_4	Y_4	Z_4
C_1	$[0:0:1]_2$ $[0:1:0]_2$	$[i:-1:1]$ $[-i:-1:1]$	$[0:1:0]$ $[-i:-1:1]$	$[0:1:0]$ $[i:-1:1]$	$[0:0:1]_2$	$[0:0:1]$ $[i:-1:1]$	$[0:0:1]$ $[-i:-1:1]$	$[0:1:0]_2$
L_1	$[i:1:1]$ $[-i:1:1]$	$[1:0:0]$	$[-i:1:1]$	$[i:1:1]$	$[1:0:0]$	$[-i:1:1]$	$[i:1:1]$	$[1:0:0]$
C_2			$[0:1:0]$ $[-i:1:1]$	$[0:1:0]$ $[i:1:1]$	$[0:0:1]_2$	$[0:0:1]$ $[-i:1:1]$	$[0:0:1]$ $[i:1:1]$	$[0:1:0]_2$
L_2			$[-i:-1:1]$	$[i:-1:1]$	$[1:0:0]$	$[i:-1:1]$	$[-i:-1:1]$	$[1:0:0]$
X_3				$[0:1:0]$	$[-i:0:1]$	$[-i:1:1]$	$[-i:-1:1]$	$[0:1:0]$
Y_3					$[i:0:1]$	$[i:-1:1]$	$[i:1:1]$	$[0:1:0]$
Z_3						$[0:0:1]$	$[0:0:1]$	$[1:0:0]$
X_4							$[0:0:1]$	$[-i:1:0]$
Y_4								$[i:1:0]$

Table 1: The intersection data of the curves we track through the pencil. Multiplicities greater than one are indicated by subscripts.

Now we will blow up at the intersection points of $T_{[1:0]}$ and $T_{[0:1]}$, and we would like to know how all of these curves intersect at those points and with what multiplicities, in order to determine their homology classes after blowing up the pencil. We summarize the relevant intersection data in Table 1.

To obtain an elliptic fibration, we blow up at the intersection points of $C_1 \cup L_1$ with $C_2 \cup L_2$. By Table 1, the relevant points are

$$\begin{aligned}
 [0:0:1] &= C_1 \cap C_2 \cap Z_3 \cap X_4 \cap Y_4, & [0:1:0] &= C_1 \cap C_2 \cap X_3 \cap Y_3 \cap Z_4, \\
 [i:1:1] &= L_1 \cap C_2 \cap Y_3 \cap Y_4, & [-i:1:1] &= L_1 \cap C_2 \cap X_3 \cap X_4, \\
 [i:-1:1] &= C_1 \cap L_2 \cap Y_3 \cap X_4, & [-i:-1:1] &= C_1 \cap L_2 \cap X_3 \cap Y_4 \\
 [1:0:0] &= L_1 \cap L_2 \cap Z_3 \cap Z_4.
 \end{aligned}$$

Observe that $[0:0:1]$ and $[0:1:0]$ appear as intersection points of multiplicity two in C_1, C_2, Z_3 and C_1, C_2, Z_4 respectively, but all other intersections are transverse. We will need to blow up twice at the multiplicity two points, and once at each other point, to eliminate the base locus of the pencil.

We will denote the generator of $H_2(\mathbb{C}P^2; \mathbb{Z})$ by h . Note that the homology class represented by one of the listed curves is h if the curve is linear, and $2h$ if the curve is quadratic.

Blow up, introducing the exceptional sphere E_1 (with homology class e_1), at $[0 : 0 : 1]$. Then the proper transforms \tilde{X}_4 and \tilde{Y}_4 intersect E_1 at distinct points, and \tilde{C}_1 , \tilde{C}_2 , and \tilde{Z}_3 all intersect at a common third point on E_1 . Note that while $X_4 \cup Y_4 \cup Z_4$ was a fiber of the Lefschetz pencil, the homology class e_1 appears with multiplicity two in its proper transform, while it appears with multiplicity one in the proper transforms of the other fibers. Therefore the Lefschetz pencil on the blown-up manifold now has a fiber $\tilde{X}_4 \cup \tilde{Y}_4 \cup \tilde{Z}_4 \cup E_1$ which includes the exceptional sphere with multiplicity one so that the fibers all represent the same homology class.

Next, we blow up at $\tilde{C}_1 \cap \tilde{C}_2 \cap \tilde{Z}_3 \cap E_1$, introducing a new exceptional sphere E_2 which intersects the proper transforms \tilde{C}_1 , \tilde{C}_2 , \tilde{Z}_3 , and \tilde{E}_1 . At this point the homology classes of all proper transforms are as follows:

$$\begin{aligned} [\tilde{C}_1] &= 2h - e_1 - e_2, & [\tilde{L}_1] &= h, & [\tilde{E}_1] &= e_1 - e_2, \\ [\tilde{C}_2] &= 2h - e_1 - e_2, & [\tilde{L}_2] &= h, & & \\ [\tilde{X}_3] &= h, & [\tilde{Y}_3] &= h, & [\tilde{Z}_3] &= h - e_1 - e_2, \\ [\tilde{X}_4] &= h - e_1, & [\tilde{Y}_4] &= h - e_1, & [\tilde{Z}_4] &= h. \end{aligned}$$

Note that \tilde{E}_1 is included with multiplicity one in a fiber with $\tilde{X}_4 \cup \tilde{Y}_4 \cup \tilde{Z}_4$.

A similar situation occurs at $[0 : 1 : 0]$. We blow up two new exceptional spheres represented homology classes e_3 and e_4 at this point. This time we must include \tilde{E}_3 with multiplicity one in a fiber with $\tilde{X}_3 \cup \tilde{Y}_3 \cup \tilde{Z}_3$.

Finally we blow up once at the points $[i : 1 : 1]$, $[-i : 1 : 1]$, $[i : -1 : 1]$, $[-i : -1 : 1]$, and $[1 : 0 : 0]$, introducing exceptional homology classes e_5, e_6, e_7, e_8 , and e_9 respectively. The homology classes of the proper transforms of the relevant curves in the four singular fibers are given as in the statement of the proposition. \square

Finally, we construct an elliptic fibration with two I_5 fibers and two fishtails.

Lemma 4.3 *There is an elliptic fibration on $E(1) = \mathbb{C}P^2 \# 9 \overline{\mathbb{C}P^2}$ with two I_5 fibers two fishtails and a section. Both I_5 fibers are made up of five -2 -spheres C_1, \dots, C_5 , D_1, \dots, D_5 with homology classes given below:*

$$\begin{aligned} [C_1] &= e_1 - e_6, & [D_1] &= e_4 - e_9, \\ [C_2] &= h - e_1 - e_4 - e_9, & [D_2] &= h - e_3 - e_4 - e_5, \\ [C_3] &= h - e_2 - e_5 - e_8, & [D_3] &= e_5 - e_8, \\ [C_4] &= e_2 - e_7, & [D_4] &= h - e_1 - e_5 - e_6, \\ [C_5] &= h - e_1 - e_2 - e_3, & [D_5] &= h - e_2 - e_4 - e_7. \end{aligned}$$




Figure 14: Initial configuration of lines.




Figure 15: The pencil after first five blow-ups.




Figure 16: Elliptic fibration.

Proof Start with a configuration of six complex projective lines, intersecting as shown in Figure 14. We can view these lines as two degenerate cubic curves indicated by distinct colors in Figure 14, which generate a Lefschetz pencil on $\mathbb{C}P^2$.

Blowing up this pencil along the five intersection points between the orange and purple curves yields the configuration in Figure 15. In order to keep the homology classes of the two curves the same so they continue to define a pencil, two of the exceptional spheres must be included with multiplicity one in the orange curve, and two must be included with multiplicity one in the purple curve. The resulting curves defining a pencil on $\mathbb{C}P^2 \# 5 \overline{\mathbb{C}P^2}$ intersect in four distinct points, and blowing up at each of these points yields an elliptic fibration shown in Figure 16.

The proper transforms of the original curves defining the pencil become the I_5 fibers representing the specified homology classes. Any other singular fibers can be perturbed to be generic nodal (fishtail) singular fibers, and an Euler characteristic computation indicates there are two of these. \square

5 Constructions of symplectic exotic 4-manifolds

In this section we put together the information from the previous sections, to construct small exotic symplectic manifolds using star surgery. In the first construction, we will perform a full analysis of the smooth invariants of the resulting manifold. In the subsequent examples, we provide abridged computations that suffice to prove that the examples are exotic copies of $\mathbb{C}P^2 \# k \overline{\mathbb{C}P^2}$.

5A A symplectic exotic $\mathbb{C}P^2 \# 8 \overline{\mathbb{C}P^2}$

5A1 The construction Our first construction of an exotic $\mathbb{C}P^2 \# 8 \overline{\mathbb{C}P^2}$ uses the elliptic fibration given in [Lemma 4.1](#), which has an I_3 fiber and I_0^* fiber, three fishtail fibers, and a section, with homology classes specified in the proof of the lemma.

Lemma 5.1 *The configuration S_2 symplectically embeds into $\mathbb{C}P^2 \# 11 \overline{\mathbb{C}P^2}$ such that its vertices represent the following homology classes, (where $f = 3h - (e_1 + \dots + e_9)$):*

$$[u_0] = 2f + e_1 - 2e_{10} - 2e_{11},$$

$$[u_1] = h - e_1 - e_2 - e_3,$$

$$[u_2] = h - e_1 - e_4 - e_5,$$

$$[u_3] = h - e_1 - e_6 - e_7,$$

$$[u_4] = h - e_1 - e_8 - e_9.$$

Proof Consider the elliptic fibration constructed in [Lemma 4.1](#). In this fibration we see the following symplectic spheres:

- The components S_1, \dots, S_5 of the I_0^* fiber, with homology classes $[S_1] = h - e_1 - e_2 - e_3$, $[S_2] = h - e_1 - e_4 - e_5$, $[S_3] = h - e_1 - e_6 - e_7$, $[S_4] = e_1 - e_8$ and $[S_5] = e_8 - e_9$.
- The component V_1 of the I_3 fiber with homology class $[V_1] = h - e_1 - e_8 - e_9$.
- The exceptional sphere E_9 which is a section.

Take two fishtail fibers and blow up their double points. The proper transforms F_1 and F_2 are now symplectic spheres in $\mathbb{C}P^2 \# 11 \overline{\mathbb{C}P^2}$. Their cohomology classes are $[F_1] = f - 2e_{10}$ and $[F_2] = f - 2e_{11}$.

Take the union of spheres $u_0 = F_1 \cup F_2 \cup E_9 \cup S_4 \cup S_5$. After symplectically smoothing its double points, u_0 gives a symplectic sphere whose homology class is $[u_0] = 2f + e_1 - 2e_{10} - 2e_{11}$. Let $u_1 = S_1$, $u_2 = S_2$, $u_3 = S_3$, $u_4 = V_1$. Then the union $\bigcup_{i=0}^4 u_i$ gives the required embedding of S_2 . \square




Figure 17: A blow-up of the elliptic fibration of Lemma 4.1 shown in Figure 12. The symplectic resolution of the union of the green curves is the sphere u_0 , and the four red curves are the spheres u_1, \dots, u_4 .

Using this embedding of S_2 , perform star surgery on $\mathbb{C}P^2 \# 11 \overline{\mathbb{C}P^2}$ by cutting out this embedded S_2 and replacing it with \mathcal{T}_2 resulting in a manifold

$$X = ((\mathbb{C}P^2 \# 11 \overline{\mathbb{C}P^2}) \setminus S_2) \cup_{\partial} \mathcal{T}_2.$$

Lemma 5.2 *The manifold X is homeomorphic to $\mathbb{C}P^2 \# 8 \overline{\mathbb{C}P^2}$.*

Proof First we show the manifold X is simply connected. By Proposition 3.1, it suffices to show that the dual circle of one of the -2 -spheres on ∂S_2 bounds a disk in $\mathbb{C}P^2 \# 11 \overline{\mathbb{C}P^2} \setminus S_2$. By our construction the sphere u_4 is a part of the I_3 fiber. Hence the meridian of u_4 bounds a disk D in the complement of S_2 which is contained in a sphere component of the I_3 fiber transversely intersecting u_4 .

Next we show that the intersection form of X is isomorphic to the intersection form of $\mathbb{C}P^2 \# 8 \overline{\mathbb{C}P^2}$. We calculate the Euler characteristic and signature to get

$$\begin{aligned} \chi(X) &= \chi(\mathbb{C}P^2 \# 11 \overline{\mathbb{C}P^2}) - \chi(S_2) + \chi(\mathcal{T}_2) = 11, \\ \sigma(X) &= \sigma(\mathbb{C}P^2 \# 11 \overline{\mathbb{C}P^2}) - \sigma(S_2) + \sigma(\mathcal{T}_2) = -7. \end{aligned}$$

These imply that $b_2^+(X) = 1$ and $b_2(X) = 9$. The intersection form cannot be even, otherwise it would be written as a direct sum of hyperbolic pieces and E_8 s, but the values of $b_2^+(X)$ and $b_2(X)$ say that such a decomposition is impossible. Hence X is homeomorphic to $\mathbb{C}P^2 \# 8 \overline{\mathbb{C}P^2}$ by Freedman’s theorem [19]. \square

5A2 Kodaira dimension In this section we compute the symplectic Kodaira dimension [27] of X . Along the way, we distinguish X from $\mathbb{C}P^2 \# 8 \overline{\mathbb{C}P^2}$. Our argument is similar to Park’s [36].

Definition 5.3 [27] For a minimal symplectic 4-manifold (M, ω) with symplectic canonical class K_ω , the Kodaira dimension of (M, ω) is defined in the following way:

$$\kappa(M, \omega) = \begin{cases} -\infty & \text{if } K_\omega \cdot [\omega] < 0 \text{ or } K_\omega \cdot K_\omega < 0, \\ 0 & \text{if } K_\omega \cdot [\omega] = 0 \text{ and } K_\omega \cdot K_\omega = 0, \\ 1 & \text{if } K_\omega \cdot [\omega] > 0 \text{ and } K_\omega \cdot K_\omega = 0, \\ 2 & \text{if } K_\omega \cdot [\omega] > 0 \text{ and } K_\omega \cdot K_\omega > 0. \end{cases}$$

The Kodaira dimension of a nonminimal manifold is defined to be that of any of its symplectic minimal models.

It is known that the Kodaira dimension is well-defined for every closed symplectic 4-manifold and depends only on the diffeomorphism type of the manifold.

Let K and ω denote the canonical class and the symplectic class of $\mathbb{C}P^2 \# 11 \overline{\mathbb{C}P^2}$ respectively. Let $X_0 = \mathbb{C}P^2 \# 11 \overline{\mathbb{C}P^2} \setminus S_2$. Then

$$\begin{aligned} K &= K|_{X_0} + K|_{S_2}, \\ \omega &= \omega|_{X_0} + \omega|_{S_2}. \end{aligned}$$

Hence

$$K \cdot \omega = K|_{X_0} \cdot \omega|_{X_0} + K|_{S_2} \cdot \omega|_{S_2}.$$

Let K_X , and ω_X denote the canonical class and the symplectic class of X respectively. Then

$$\begin{aligned} K_X &= K|_{X_0} + K|_{\mathcal{T}_2}, \\ \omega_X &= \omega|_{X_0} + \omega|_{\mathcal{T}_2}. \end{aligned}$$

Since $K|_{\mathcal{T}_2} = 0$ by Proposition 3.3, we have

$$K_X \cdot \omega_X = K|_{X_0} \cdot \omega|_{X_0} = K \cdot \omega - K|_{S_2} \cdot \omega|_{S_2}.$$

Lemma 5.4 For every $k > 0$, the manifold $\mathbb{C}P^2 \# k \overline{\mathbb{C}P^2}$ admits a symplectic structure whose cohomology class is given by $\omega = ah - b_1 e_1 - \dots - b_k e_k$ for some rational numbers a, b_1, \dots, b_k with $a > b_1 > \dots > b_k$ and $a > b_1 + \dots + b_k$.

Proof Note that a is the symplectic area of $\mathbb{C}P^1 \subset \mathbb{C}P^2$ and b_j is the symplectic area of the exceptional sphere e_j for all $j = 1, \dots, k$. By [29, Chapter 7], we have $b_j = \lambda^2$ where λ is the weight of the corresponding blow-up. In other words λ is the radius of the Darboux ball which will be removed during the blow-up process. Since the weights of the blow-ups can be chosen to be arbitrarily small, the result follows. \square

Lemma 5.5 $K_X \cdot \omega_X > 0$

Proof Write

$$K = -3h + e_1 + \cdots + e_{11},$$

$$\omega = ah - b_1e_1 - \cdots - b_{11}e_{11},$$

where a, b_1, \dots, b_{11} are rational numbers such that $a > b_1 > \cdots > b_{11}$ and

$$(5-1) \quad a > b_1 + \cdots + b_{11}.$$

Then we have

$$K \cdot \omega = -3a + b_1 + \cdots + b_{11}.$$

Let $\gamma_0, \dots, \gamma_4$ be the basis of $H^2(S_2, \mathbb{Q})$ which is dual to u_0, \dots, u_4 . Then the adjunction formula implies

$$K|_{S_2} = (K \cdot u_0)\gamma_0 + \cdots + (K \cdot u_4)\gamma_4 = 3\gamma_0.$$

We calculate the restriction of the symplectic class on S_2 using Lemma 5.1:

$$\begin{aligned} \omega|_{S_2} &= (\omega \cdot u_0)\gamma_0 + \cdots + (\omega \cdot u_4)\gamma_4 \\ &= (6a - b_1 - 2b_2 - \cdots - 2b_{11})\gamma_0 + (a - b_1 - b_2 - b_3)\gamma_1 \\ &\quad + (a - b_1 - b_4 - b_5)\gamma_2 + (a - b_1 - b_6 - b_7)\gamma_3 + (a - b_1 - b_8 - b_9)\gamma_4. \end{aligned}$$

Let P denote the intersection matrix for S_2 . Then

$$P^{-1} = -\frac{1}{12} \begin{pmatrix} 4 & 2 & 2 & 2 & 2 \\ 2 & 7 & 1 & 1 & 1 \\ 2 & 1 & 7 & 1 & 1 \\ 2 & 1 & 1 & 7 & 1 \\ 2 & 1 & 1 & 1 & 7 \end{pmatrix}.$$

Hence

$$\begin{aligned} K|_{S_2} \cdot \omega|_{S_2} &= -\frac{1}{4}[4(6a - b_1 - 2b_2 - \cdots - 2b_{11}) + 2(a - b_1 - b_2 - b_3) \\ &\quad + 2(a - b_1 - b_4 - b_5) + 2(a - b_1 - b_6 - b_7) + 2(a - b_1 - b_8 - b_9)] \\ &= -\frac{1}{4}[32a - 12b_1 - 10b_2 - 10b_3 - \cdots - 10b_9 - 8b_{10} - 8b_{11}]. \end{aligned}$$

Therefore

$$\begin{aligned} K_X \cdot \omega_X &= K|_{X_0} \cdot \omega|_{X_0} = K \cdot \omega - K|_{S_2} \cdot \omega|_{S_2} \\ &= -3a + b_1 + \cdots + b_{11} + [8a - 3b_1 - \frac{5}{2}b_2 - \cdots - \frac{5}{2}b_9 - 2b_{10} - 2b_{11}] \\ &= 5a - 2b_1 - \frac{3}{2}b_2 - \cdots - \frac{3}{2}b_9 - b_{10} - b_{11} \\ &> 0 \quad (\text{by Equation (5-1)}). \end{aligned}$$

□

Remark 5.6 The important input which ensures there is a symplectic form on X with $K_X \cdot \omega_X > 0$ is the high multiplicity of the homology class h in the classes $[u_i]$ of the spheres in the embedding of S_2 . Note that if the embedding of S_2 were completely disjoint from the $\mathbb{C}P^1$ representing h , then by McDuff's theorem, the resulting star surgered manifold would be a standard blow-up of $\mathbb{C}P^2$. This computation provides a quantitative way of ensuring that when the plumbing spheres intersect $\mathbb{C}P^1$ enough for the star surgery to produce an exotic smooth structure.

Proposition 5.7 *The manifold X is not diffeomorphic to $\mathbb{C}P^2 \#_8 \overline{\mathbb{C}P^2}$.*

Proof The standard symplectic form on $\mathbb{C}P^2 \#_k \overline{\mathbb{C}P^2}$ satisfies $K \cdot \omega < 0$. According to [28, Theorem D], there is a unique symplectic structure on $\mathbb{C}P^2 \#_k \overline{\mathbb{C}P^2}$ for $2 \leq k \leq 9$ up to diffeomorphism and deformation. Hence $\mathbb{C}P^2 \#_8 \overline{\mathbb{C}P^2}$ does not admit a symplectic structure with $K \cdot \omega > 0$. We have just seen that $K_X \cdot \omega_X > 0$. Hence X is not diffeomorphic to $\mathbb{C}P^2 \#_8 \overline{\mathbb{C}P^2}$. \square

Proposition 5.8 *The manifold X is minimal.*

The proof is postponed until the next section. With this in hand, we are ready to determine the Kodaira dimension of X .

Proposition 5.9 *The manifold X has symplectic Kodaira dimension 2.*

Proof Recall that the symplectic Kodaira dimension is defined on the minimal model. Since X is minimal, $K_X \cdot \omega_X > 0$, and $K_X^2 = 3\sigma(X) + 2\chi(X) = 1 > 0$, its symplectic Kodaira dimension is 2. \square

This proposition shows that X can also be distinguished from $\mathbb{C}P^2 \#_8 \overline{\mathbb{C}P^2}$ using the symplectic Kodaira dimension, since $\kappa(\mathbb{C}P^2 \#_8 \overline{\mathbb{C}P^2}) = \kappa(\mathbb{C}P^2) = -\infty$.

5A3 Minimality Throughout this section, we assume that the reader is familiar with the Seiberg–Witten invariants of manifolds with $b_2^+ = 1$. See [31; 17; 42] for excellent expositions.

In order to prove the minimality of X one needs to know the effect of star surgery on Seiberg–Witten invariants. It suffices to know all the Seiberg–Witten basic classes of X . The effect of such cut and paste operations on Seiberg–Witten invariants was studied by Michalोगιorgaki in [30] in a more general framework; see also [39] for an analogous result from the perspective of Heegaard Floer theory.

Theorem 5.10 [30, Theorem 1] *Suppose Y is a rational homology sphere which is a monopole L -space. Let P and B be negative-definite 4-manifolds with $b_1(P) = b_1(B) = 0$ and $\partial P = \partial B = Y$. Let $X = Z \cup_Y P$ and $X' = Z \cup_Y B$, for some 4-manifold Z . If $\mathfrak{s} \in \text{Spin}^c(X)$, $\mathfrak{s}' \in \text{Spin}^c(X')$, $d_X(\mathfrak{s}), d_{X'}(\mathfrak{s}') \geq 0$ and $\mathfrak{s}|_Z = \mathfrak{s}'|_Z$, then $\text{SW}_X(\mathfrak{s}) = \text{SW}_{X'}(\mathfrak{s}')$.*

In the case that $b_2^+(X) = 1$, $\text{SW}_{X,a_1}(\mathfrak{s}) = \text{SW}_{X',a_2}(\mathfrak{s}')$, where $a_1 \in H_2(X, \mathbb{Z})$, $a_2 \in H_2(X', \mathbb{Z})$ specify chambers such that $a_1|_P = a_2|_B = 0$ and $a_1|_Z = a_2|_Z$.

This result and the wall-crossing formula reduces the problem of determining basic classes of X to a cohomology computation. In this section, we incorporate a search method invented by Ozsváth and Szabó to find all the basic classes of X . First we find a homology class that determines a common chamber for the manifolds before and after star surgery.

Lemma 5.11 *The element $V \in H_2(\mathbb{C}P^2 \# 11 \overline{\mathbb{C}P^2})$ defined by*

$$V = 86h - 36e_1 - 25e_2 - 25e_3 - 25e_4 - 25e_5 - 25e_6 - 25e_7 - 19e_8 - 31e_9 - 20e_{10} - 20e_{11}$$

satisfies the following conditions:

- (1) $V \cdot [u_i] = 0$ for $i = 0, \dots, 4$ (ie V is orthogonal to each embedded sphere of S_2),
- (2) $V \cdot V > 0$,
- (3) $V \cdot h > 0$,
- (4) $V \cdot K > 0$, where $K = -3h + e_1 + \dots + e_{11}$ is the canonical class of $\mathbb{C}P^2 \# 11 \overline{\mathbb{C}P^2}$.

Proof The proof is a direct computation. □

Lemma 5.12 *The small perturbation Seiberg–Witten invariant of X at the canonical class $\tilde{K} \in H^2(X, \mathbb{Z})$ is nonzero.*

Proof We first compute the Seiberg–Witten invariant of $\mathbb{C}P^2 \# 11 \overline{\mathbb{C}P^2}$ in the chamber determined by the element V in Lemma 5.11. Orient $H_2^+(\mathbb{C}P^2 \# 11 \overline{\mathbb{C}P^2})$ with h . Note that the homology class h gives the chamber of the positive scalar curvature metric hence all Seiberg–Witten invariants of $\mathbb{C}P^2 \# 11 \overline{\mathbb{C}P^2}$ are zero in this chamber. In particular $\text{SW}_{\mathbb{C}P^2 \# 11 \overline{\mathbb{C}P^2}, h}(K) = 0$ for the canonical class K . By part (2) of Lemma 5.11, $V \in H_2^+(\mathbb{C}P^2 \# 11 \overline{\mathbb{C}P^2})$, and by part (3) V has the correct orientation. Hence V determines a chamber. Since $K \cdot h < 0$, part (4) says that there is a wall between the chambers of h and V with respect to the canonical class. By the wall-crossing formula [28], $\text{SW}_{\mathbb{C}P^2 \# 11 \overline{\mathbb{C}P^2}, V}(K) = \pm 1$.

Next we relate the Seiberg–Witten invariants of $\mathbb{C}P^2 \# 11 \overline{\mathbb{C}P^2}$ and X . The canonical class \tilde{K} of X satisfies $\tilde{K}_{X \setminus \mathcal{T}_2} = K_{\mathbb{C}P^2 \# 11 \overline{\mathbb{C}P^2} \setminus S_2}$. Part (1) says V determines a cham-

ber in X . If we can show that $\partial\mathcal{T}_2$ is a monopole L -space then by [Theorem 5.10](#), noting that $d_X(\tilde{K}) = d_{\mathbb{C}P^2 \#_{11} \overline{\mathbb{C}P^2}}(K) = 0$, we have $\text{SW}_{\mathbb{C}P^2 \#_{11} \overline{\mathbb{C}P^2}, \nu}(K) = \text{SW}_{X, \nu}(\tilde{K})$. Since there is a unique small perturbation chamber for manifolds with $b_2^- \leq 9$, the latter is equal to $\text{SW}_X(\tilde{K})$.

It remains to see that $\partial\mathcal{T}_2 = \partial\mathcal{S}_2$ is a monopole L -space. In fact this holds for $\partial\mathcal{S}_i$ for all i . The plumbing graph of \mathcal{S}_i satisfies $|m(v)| > d(v)$ at each vertex v , where $m(v)$ is the weight of v and $d(v)$ is the number of edges connected to v . In [\[33, Theorem 7.1\]](#), Ozsváth and Szabó show that the boundary of such a plumbing is a Heegaard Floer L -space. Their proof uses only the surgery exact triangle and the formal properties of Heegaard Floer homology. Hence it can be repeated to show that boundaries of such plumbings are also monopole L -spaces. Alternatively, one can refer to the recently established equivalence of the Heegaard Floer homology and the monopole Floer homology to see that every Heegaard Floer L -space is also a monopole L -space [\[26; 8\]](#). \square

Remark 5.13 [Lemma 5.12](#) gives an alternative proof of [Proposition 5.7](#). Indeed, the small perturbation Seiberg–Witten invariant is well defined for manifolds with $b_2^- \leq 9$ since there is a unique chamber for such manifolds. For $\mathbb{C}P^2 \#_k \overline{\mathbb{C}P^2}$ with $k \leq 9$ the unique chamber is the one given by the positive scalar curvature metric. Hence $\text{SW}_{\mathbb{C}P^2 \#_k \overline{\mathbb{C}P^2}}(L) = 0$ for all characteristic cohomology class L , for all $k \leq 9$.

Proof of Proposition 5.8 By the blow-up formula for the Seiberg–Witten invariant, it suffices to show that the only basic classes of X are $\pm\tilde{K}$. Therefore, we want to check which integral characteristic cohomology classes (representing Spin^c structures) on X are Seiberg–Witten basic classes. While we will show how to use [Theorem 5.10](#) and the wall-crossing formula to compute the Seiberg–Witten invariant on a given cohomology class, there are infinitely many classes to check. The strategy of Ozsváth and Szabó (which they used to prove minimality of a rationally blown-down manifold) is to check only the finitely many *adjunctive classes* and then use the information about the adjunctive basic classes to rule out the possibility of nonadjunctive basic classes.

The fact that the homology of \mathcal{T}_2 has nontrivial rank makes the search somewhat more complicated than the case of rational blow-down. Even so, the computations can be handled by a simple computer program. First we find a basis for the subspace $H_2(\mathbb{C}P^2 \#_{11} \overline{\mathbb{C}P^2})$ which is orthogonal to the homology of the configuration \mathcal{S}_2 . The following elements form such a basis:

$$\begin{aligned} A_1 &= h - e_3 - e_5 - e_7 - e_9 + e_{11}, & A_2 &:= -3h + 2e_1 + e_3 + e_5 + e_7 + e_9 + 2e_{10} + 2e_{11}, \\ A_3 &= e_2 - e_3, & A_4 &= e_4 - e_5, & A_5 &= e_6 - e_7, & A_6 &= e_{10} - e_{11}, & A_7 &= e_8 - e_9. \end{aligned}$$

Note that all of these homology classes can be represented by embedded spheres, excepting A_2 which can be represented by an embedded torus. Moreover, all of these surfaces can be chosen in the complement of S_2 in $\mathbb{C}P^2 \# 11 \overline{\mathbb{C}P^2}$. Additionally we let A_8 and A_9 be the homology generators of \mathcal{T}_2 , which can be represented by embedded tori in \mathcal{T}_2 , as indicated in Proposition 3.2. Then A_1, \dots, A_9 represent a basis for $H_2(X; \mathbb{Z})$ so we can represent cohomology classes in $H^2(X; \mathbb{Z})$ by a tuple of integers representing $[\langle L, A_1 \rangle, \dots, \langle L, A_9 \rangle]$.

First we will determine which of these tuples represent *integral*, *characteristic*, and *adjunctive* cohomology classes. Because the inverse of the intersection matrix has rational coefficients, some tuples of integers could represent rational, but not integral, homology classes. We check a mod 2 equivalence to see if the cohomology class is characteristic, and an adjunctive inequality for each A_i . Computationally, we find that there are exactly 243000 cohomology classes $L \in H^2(X, \mathbb{Q})$ which satisfy these conditions for all $i = 1, \dots, 9$:

$$\langle L, A_i \rangle \in \mathbb{Z}, \quad \langle L, A_i \rangle \equiv A_i^2 \pmod{2}, \quad |\langle L, A_i \rangle| \leq -A_i^2.$$

In order to test whether a characteristic class is basic for X , we first check whether the expected dimension is nonnegative and even (necessary conditions for the Seiberg–Witten invariant to be nonzero). Let $d_X(L) := (L^2 - 3\sigma(X) - 2\chi(X))/4 = (L^2 - 1)/4$. In the second round of our search, we check how many of these cohomology classes satisfy $d_X(L) \in \mathbb{Z}$, $d_X(L) \geq 0$ and $d_X(L) \equiv 0 \pmod{2}$. It turns out that there are 25040 such classes.

We will use Theorem 5.10 and the wall-crossing formula to calculate the Seiberg–Witten invariant evaluated on each of these 25040 classes. To do this, we must relate each class on X to a class on $\mathbb{C}P^2 \# 11 \overline{\mathbb{C}P^2}$ so that the two classes have equal restrictions to $X \setminus \mathcal{T}_2 = (\mathbb{C}P^2 \# 11 \overline{\mathbb{C}P^2}) \setminus S_2$. The difficulty is, once we restrict the chosen class to $X \setminus \mathcal{T}_2$, we must find a class on S_2 which extends it over $\mathbb{C}P^2 \# 11 \overline{\mathbb{C}P^2}$. For this, we compare the classes on ∂S_2 which occur as the restriction of a class on S_2 to the classes on $\partial \mathcal{T}_2 = \partial S_2$ that occur as the restriction of a class on \mathcal{T}_2 . We showed that the image of the map $H^2(\mathcal{T}_2, \mathbb{Z}) \rightarrow H^2(\partial \mathcal{T}_2, \mathbb{Z})$ is an index-2 subgroup, hence it has order 24. Our aim is to find a numerical criterion for a characteristic cohomology class $L \in H^2(S_2, \mathbb{Z})$ to extend to \mathcal{T}_2 when restricted to $\partial S_2 = \partial \mathcal{T}_2$. First observe that the set of Spin^c structures on ∂S_2 as a $H^2(\partial S_2, \mathbb{Z})$ torsor is isomorphic to $2H^2(S_2, \partial S_2, \mathbb{Z})$ –orbits of characteristic elements in $H^2(S_2, \mathbb{Z})$. Hence two characteristic classes $L_1, L_2 \in H^2(S_2, \mathbb{Z})$ restrict to the same Spin^c structure on the boundary if and only if $L_1 = L_2 + 2 \text{PD}(Z)$ for some $Z \in H_2(S_2, \mathbb{Z})$. We will use d –invariants to determine which orbits in $H^2(S_2, \mathbb{Z})$ restrict to a Spin^c structure on $\partial S_2 = \partial \mathcal{T}_2$ which appears in the boundary

of a Spin^c structure on \mathcal{T}_2 and to choose distinguished representatives of these orbits which maximize the expected dimension of the Seiberg–Witten moduli space. Define $d_{S_2}(L) = (L^2 - 3\sigma(S_2) - 2\chi(S_2))/4 = (L^2 + 3)/4$. Observe that the $d_{S_2}(L) \pmod 2$ is constant in the orbit $L + 2H^2(S_2, \partial S_2, \mathbb{Z})$. Each orbit has a representative L' satisfying $[u_i]^2 + 2 \leq \langle L', [u_i] \rangle \leq -[u_i]^2$ for all $i = 0, \dots, 4$, where each u_i is a sphere appearing as a vertex in the star-shaped plumbing graph of S_2 . Computing d_{S_2} for all of these representatives, we see that the possible mod 2 reductions of d_{S_2} for characteristic cohomology classes on S_2 are $\{\frac{-2}{3}, \frac{-1}{3}, \frac{-1}{4}, \frac{-1}{12}, 0, \frac{1}{4}, \frac{2}{3}, \frac{-11}{12}, 1\}$.

In the proof of Proposition 3.2, we computed the intersection form of \mathcal{T}_2 . From this, we see that the possible mod 2 reductions of $d_{\mathcal{T}_2}$ for characteristic cohomology classes on \mathcal{T}_2 are $\{\frac{-1}{3}, 0, \frac{2}{3}, 1\}$. We observe that exactly 24 of the $2H^2(S_2, \partial S_2, \mathbb{Z})$ orbits have the mod 2 reductions of their d_{S_2} in this set. Hence we conclude that a characteristic cohomology class $L \in H^2(S_2, \mathbb{Z})$ extends to \mathcal{T}_2 if and only if $d_{S_2}(L) \in \{\frac{-1}{3}, 0, \frac{2}{3}, 1\} \pmod 2$. We can explicitly write distinguished representatives of these 24 orbits. They are the elements of the following set:

$$\begin{aligned} \Phi = \{ & [1, 0, 0, 0, 0], [-3, 2, 2, 2, 2], [-1, 2, 0, 0, 2], [-3, 0, 2, 2, 0], \\ & [1, 2, 0, 2, 0], [3, 0, 0, 0, 0], [-1, 2, 2, 0, 0], [-3, 2, 0, 2, 0], [-1, 0, 2, 0, 2], \\ (5-2) \quad & [-3, 0, 0, 2, 2], [1, 0, 2, 2, 0], [1, 0, 0, 2, 2], [-1, 0, 2, 2, 0], [-3, 0, 2, 0, 2], \\ & [-3, 0, 0, 0, 0], [1, 2, 0, 0, 2], [5, 0, 0, 0, 0], [-3, 2, 0, 0, 2], [-1, 0, 0, 2, 2], \\ & [1, 0, 2, 0, 2], [-1, 2, 0, 2, 0], [-3, 2, 2, 0, 0], [1, 2, 2, 0, 0], [-1, 0, 0, 0, 0] \}. \end{aligned}$$

Here each cohomology class $L \in H^2(S_2, \mathbb{Z})$ is represented by the tuple $[\langle L, [u_0] \rangle, \dots, \langle L, [u_4] \rangle]$. Elements of Φ maximize d_{S_2} in their respective $2H^2(S_2, \partial S_2, \mathbb{Z})$ orbits.

We continue our search for basic classes of X . Recall that in round two, we got 25,040 potential adjunctive basic classes. We restrict each one to $X \setminus \mathcal{T}_2$ (ie we forget the intersections with A_8 and A_9). Each of these characteristic cohomology classes glues to exactly one element of the set Φ to define a characteristic cohomology class on $\mathbb{C}P^2 \# 11 \overline{\mathbb{C}P^2}$. There are 600,960 triples (A, B, C) where A is an adjunctive class on \mathcal{T}_2 , B is an adjunctive class on $X \setminus \mathcal{T}_2 = \mathbb{C}P^2 \# 11 \overline{\mathbb{C}P^2} \setminus S_2$, C is one of the 24 distinguished cohomology classes on S_2 , and the pair (B, A) is one of the 25,040 potential adjunctive basic classes on X . We now restrict our attention to all triples where the combination (B, C) represents a cohomology class on $\mathbb{C}P^2 \# 11 \overline{\mathbb{C}P^2}$ with $d \in \mathbb{Z}$, $d \geq 0$ and $d \equiv 2 \pmod 2$. This leaves us with 219,064 possible triples. In only 122,212 of these triples does (B, C) represent an integral and characteristic cohomology class on $\mathbb{C}P^2 \# 11 \overline{\mathbb{C}P^2}$. Finally, we check whether the chamber determined by V in $\mathbb{C}P^2 \# 11 \overline{\mathbb{C}P^2}$ is the same as the chamber of the positive scalar curvature metric with respect to each of the characteristic cohomology classes (B, C) in the remaining

triples by comparing the signs of $(B, C) \cdot V$ with $(B, C) \cdot H$. When the chambers agree, B will not descend to a basic class on X , but when they disagree the wall-crossing formula and [Theorem 5.10](#) ensure that (A, B) is a basic class on X . It turns out that only two of the remaining triples (A, B, C) have the property that V is in a different chamber than H with respect to (B, C) . These are necessarily the canonical class and its negative. Hence \tilde{K} and $-\tilde{K}$ are the only adjunctive basic classes of X .

Lastly we argue that there can be no nonadjunctive basic class. If there was such an L , then the adjunction relations [\[32\]](#) imply that adding or subtracting twice the Poincaré dual of any surface with negative self-intersection where the adjunction inequality fails, we would obtain another basic class L' with $d_X(L') > d_X(L)$.

Since X has only finitely many basic classes this process eventually stops at an adjunctive basic class, \bar{L} with $d_X(\bar{L}) > d_X(L) \geq 0$, but this is a contradiction because \tilde{K} and $-\tilde{K}$ are the only adjunctive basic classes and $d_X(\tilde{K}) = d_X(-\tilde{K}) = 0$. \square

Proof of [Theorem 1.1](#) Our theorem follows from [Lemma 5.2](#), [Proposition 5.7](#), [Proposition 5.8](#), and [Proposition 5.9](#). \square

Remark 5.14 The Seiberg–Witten invariants of X agree up to isomorphism with all known examples of minimal exotic $\mathbb{C}P^2 \# 8 \overline{\mathbb{C}P^2}$ [\[36; 42; 16; 30\]](#). Hence we cannot distinguish X from these manifolds using Seiberg–Witten invariants.

5B An exotic $\mathbb{C}P^2 \# 7 \overline{\mathbb{C}P^2}$

Lemma 5.15 *There is an embedding of the star-shaped plumbing \mathcal{Q} into $\mathbb{C}P^2 \# 12 \overline{\mathbb{C}P^2}$ such that the spheres represent the following homology classes:*

$$\begin{aligned} [u_0] &= 6h - 2e_1 - e_2 - 2e_3 - \cdots - 2e_9 - 2e_{11} - 2e_{12}, \\ [u_{1,1}] &= 2h - e_1 - e_2 - e_3 - e_4 - e_5 - e_6, \\ [u_{2,1}] &= h - e_1 - e_2 - e_9, \\ [u_{2,2}] &= h - e_3 - e_5 - e_7, \\ [u_{3,1}] &= h - e_1 - e_6 - e_7, \\ [u_{3,2}] &= h - e_3 - e_4 - e_9 - e_{10}, \\ [u_{4,1}] &= h - e_1 - e_5 - e_8 - e_{10}. \end{aligned}$$

Proof In [Lemma 4.2](#), we showed that there is an elliptic fibration with two I_2 fibers, two I_4 fibers, and a section, and we specified the homology classes of the spheres making up the singular fibers and the section. According to that notation, the spheres

in the two I_2 fibers were \tilde{C}_1, \tilde{L}_1 and \tilde{C}_2, \tilde{L}_2 , respectively. The spheres in the I_4 fibers were denoted $\tilde{X}_3, \tilde{Y}_3, \tilde{Z}_3, \tilde{E}_3$, and $\tilde{X}_4, \tilde{Y}_4, \tilde{Z}_4, \tilde{E}_4$, respectively. $\tilde{C}_1, \tilde{C}_2, \tilde{Z}_3$ and \tilde{E}_4 intersected the section E_2 .

To construct the embedding of \mathcal{Q} , first blow up at the intersection of \tilde{Y}_4 and \tilde{Z}_4 to produce two adjacent -3 -spheres. Then we can perturb the fibration near one of the I_2 fibers to split it into two fishtail fibers. Blow up each the self-intersection points of each of these fishtail fibers to create two -4 -spheres. To produce the central -5 -sphere, take the symplectic resolution of the section E_2 , the two blown-up fishtail fibers, and E_4 (the sphere in the blown-up I_4 fiber which intersects the section). The remaining spheres can be taken to be $u_{1,1} = \tilde{C}_2, u_{2,1} = \tilde{Z}_3, u_{2,2} = \tilde{Y}_3, u_{3,1} = \tilde{X}_4, u_{3,2} = \tilde{Z}_4, u_{4,1} = \tilde{Y}_4$, where \bar{C} denotes the proper transform under blow-up. Using the homology computation indicated in Lemma 4.2, the homology classes for the $u_{i,j}$ follows. □

Lemma 5.16 *The element $R \in H_2(\mathbb{C}P^2 \#_{12} \overline{\mathbb{C}P^2})$ defined by*

$$R = 533\tilde{h} - 188e_1 - 186e_2 - 192e_3 - 126e_4 - 185e_5 - 189e_6 - 156e_7 - 104e_8 - 159e_9 - 56e_{10} - 0e_{11} - 151e_{12}$$

satisfies the following conditions:

- (1) $R \cdot [u_{i,j}] = 0$ for all i and j ,
- (2) $R \cdot R > 0$,
- (3) $R \cdot h > 0$,
- (4) $R \cdot K > 0$, where $K = -3h + e_1 + \dots + e_{12}$ is the canonical class of $\mathbb{C}P^2 \#_{12} \overline{\mathbb{C}P^2}$.

Proof All of the above claims can be verified by a direct computation. □

Proposition 5.17 *The result of star surgery on this embedding of \mathcal{Q} into $\mathbb{C}P^2 \#_{12} \overline{\mathbb{C}P^2}$ is an exotic copy of $\mathbb{C}P^2 \#_7 \overline{\mathbb{C}P^2}$ which supports a symplectic structure.*

Proof Let $X' = ((\mathbb{C}P^2 \#_{12} \overline{\mathbb{C}P^2}) \setminus \mathcal{Q}) \cup \mathcal{R}$ be the result of star surgery. Then $\chi(X') = \chi(\mathbb{C}P^2 \#_{12} \overline{\mathbb{C}P^2}) - \chi(\mathcal{Q}) + \chi(\mathcal{R}) = 15 - 8 + 3 = 10$. Since \mathcal{Q} and \mathcal{R} are both negative-definite, $\sigma(X') = -6$. Since \mathcal{R} is simply connected, X' is simply connected. Therefore X' is homeomorphic to $\mathbb{C}P^2 \#_7 \overline{\mathbb{C}P^2}$ by Freedman’s theorem [19]. Recall that for 4-manifolds with $b_2^+ = 1$, the Seiberg–Witten invariants depend upon a choice of a chamber. Let h denote the homology class of $\mathbb{C}P^1 \subset \mathbb{C}P^2 \#_{12} \overline{\mathbb{C}P^2}$. This class gives the chamber of positive scalar curvature metric, so $\text{SW}_{\mathbb{C}P^2 \#_{12} \overline{\mathbb{C}P^2}, h}(L) = 0$ for every characteristic class L . In particular for the canonical class K , we have




Figure 18: Elliptic fibration on $E(1)$ with section.

$SW_{\mathbb{C}P^2 \#_{12} \overline{\mathbb{C}P^2}, h}(K) = 0$. By Lemma 5.16, the class R determines a chamber and there is a wall between this chamber and the chamber of h with respect to K . Hence by the wall-crossing formula [28], we have $SW_{\mathbb{C}P^2 \#_{12} \overline{\mathbb{C}P^2}, R}(K) = \pm 1$. We will show that X' also has a nonzero Seiberg–Witten invariant. Indeed the 3–manifold ∂S is a monopole L –space. Let $K_{X'}$ denote the canonical class of X' . Clearly we have

$$K_{X'}|_{X' \setminus \mathcal{R}} = K|_{\mathbb{C}P^2 \#_{12} \overline{\mathbb{C}P^2} \setminus \mathcal{Q}} \quad \text{and} \quad d_{X'}(K_{X'}) = d_{\mathbb{C}P^2 \#_{12} \overline{\mathbb{C}P^2}}(K) = 0.$$

Hence by [30], we have

$$SW_{X', R}(K_{X'}) = SW_{\mathbb{C}P^2 \#_{12} \overline{\mathbb{C}P^2}, R}(K) = \pm 1.$$

For manifolds with $b_2^- \leq 9$, the choice of a chamber is unique. Therefore X' is not diffeomorphic to $\mathbb{C}P^2 \#_7 \overline{\mathbb{C}P^2}$ whose Seiberg–Witten invariants are all zero in the unique chamber of positive scalar curvature. \square

5C An exotic $\mathbb{C}P^2 \#_6 \overline{\mathbb{C}P^2}$

For this construction we use the elliptic fibration with two I_5 fibers, two fishtails, and a section, with homology classes specified in Lemma 4.3

Lemma 5.18 *The configuration of spheres \mathcal{U} embeds into $\mathbb{C}P^2 \#_{13} \overline{\mathbb{C}P^2}$.*

Proof By blowing up at one singular point in each I_5 fiber, we obtain four -3 –spheres which will make up the ends of each of the arms in the configuration \mathcal{U} . The central sphere of square -5 is obtained by taking the symplectic resolution of a section with two blown-up fishtail fibers as well as the -2 –spheres in each I_5 fiber which intersect this section.




Figure 19: Embedding of \mathcal{U} into $\mathbb{C}P^2 \# 13 \overline{\mathbb{C}P^2}$.

The configuration is shown in Figure 19, and the homology classes are given below:

$$\begin{aligned}
 [u_0] &= 2[F] - 2e_{12} - 2e_{13} + [C_2] + [D_1] + e_9, \\
 [u_{1,1}] &= [D_2], & [u_{2,1}] &= [C_3], & [u_{3,1}] &= [C_1], & [u_{4,1}] &= [D_5] - e_{11}. \\
 [u_{1,2}] &= [D_3], & [u_{2,2}] &= [C_4] - e_{10}, & [u_{3,2}] &= [C_5] - e_{10}, \\
 [u_{1,3}] &= [D_4] - e_{11}, & & & & & & \square
 \end{aligned}$$

Lemma 5.19 *The element $R \in H_2(\mathbb{C}P^2 \# 13 \overline{\mathbb{C}P^2})$ defined by*

$$\begin{aligned}
 R &= 5656h - 1728e_1 - 1846e_2 - 1836e_3 - 1915e_4 - 1905e_5 - 1728e_6 - 1600e_7 \\
 &\quad - 1905e_8 - 1890e_9 - 246e_{10} - 295e_{11} - 393e_{12} - 1241e_{13}
 \end{aligned}$$

satisfies the following conditions:

- (1) $R \cdot [u_{i,j}] = 0$ for all i and j (ie R is orthogonal to each embedded sphere of \mathcal{U}),
- (2) $R \cdot R > 0$,
- (3) $R \cdot h > 0$,
- (4) $R \cdot K > 0$, where $K = -3h + e_1 + \dots + e_{13}$ is the canonical class of $\mathbb{C}P^2 \# 13 \overline{\mathbb{C}P^2}$.

Proof The proof is a direct check. □

Proposition 5.20 *The result of the $(\mathcal{U}, \mathcal{V})$ star surgery on this embedding of \mathcal{U} into $\mathbb{C}P^2 \# 12 \overline{\mathbb{C}P^2}$ is an exotic copy of $\mathbb{C}P^2 \# 6 \overline{\mathbb{C}P^2}$ which supports a symplectic structure.*

Proof The proof is similar to the proof of Proposition 5.17. □

5D Another symplectic exotic $\mathbb{C}P^2 \#8 \overline{\mathbb{C}P^2}$

Proposition 5.21 *It is possible to symplectically embedded the configuration \mathcal{K} into $\mathbb{C}P^2 \#12 \overline{\mathbb{C}P^2}$ so that the homology classes of the central sphere C_0 and the four spheres in the arms C_1, C_2, C_3, C_4 are given as follows:*

$$\begin{aligned} [C_0] &= 2f + e_1 - 2e_{10} - 2e_{11} - e_{12}, \\ [C_1] &= h - e_1 - e_2 - e_3, \\ [C_2] &= h - e_1 - e_4 - e_5, \\ [C_3] &= h - e_1 - e_6 - e_7, \\ [C_4] &= h - e_1 - e_8 - e_9, \end{aligned}$$

where $f = 3h - (e_1 + \dots + e_9)$.

Proof This proposition follows immediately from blowing up the embedding of S_2 into $\mathbb{C}P^2 \#11 \overline{\mathbb{C}P^2}$ given in Lemma 5.1 along a single point on the section. \square

Note the other elliptic fibrations constructed in this paper can be used to find other embeddings of S_2 and \mathcal{K} into blow-ups of $E(1)$. It is an interesting question to ask whether the same star surgery operation performed on different embeddings of the plumbing into the same manifold can result in nondiffeomorphic manifolds.

Theorem 5.22 *The manifold resulting from star surgery on this embedding of \mathcal{K} into $\mathbb{C}P^2 \#12 \overline{\mathbb{C}P^2}$ is an exotic copy of $\mathbb{C}P^2 \#8 \overline{\mathbb{C}P^2}$.*

Proof The manifold is homeomorphic to $\mathbb{C}P^2 \#8 \overline{\mathbb{C}P^2}$ by Freedman’s theorem once we show that it is simply connected and has Euler characteristic 11 and signature -7 . The Euler characteristic and signature computations follow from the fact that $\chi(\mathcal{L}) = 2$, $\sigma(\mathcal{L}) = -1$, $\chi(\mathcal{K}) = 6$, and $\sigma(\mathcal{K}) = -5$. The manifold is simply connected as in the proof of Lemma 5.2 because the generator of $\pi_1(\mathcal{L})$ is isotopic to the meridian of C_4 in the embedding which is homotopically trivial because C_4 is one sphere in an I_3 -fiber where the other transversally intersecting spheres are not cut out in the star surgery.

The diffeomorphism type can be distinguished from $\mathbb{C}P^2 \#8 \overline{\mathbb{C}P^2}$ either by showing that the resulting symplectic manifold has Kodaira dimension two as in Lemma 5.5, or by showing that the canonical class and its negation are basic classes in the small perturbation chamber of the star surgered manifold as in Lemma 5.12. In fact the value of $K_X \cdot \omega_X$ to compute the Kodaira dimension comes out to be exactly the same value as for the star surgery using S_2 and \mathcal{T}_2 , and the element V defining the chamber on $\mathbb{C}P^2 \#12 \overline{\mathbb{C}P^2}$ which descends to the small perturbation chamber in the star surgered manifold can be defined identically as in Lemma 5.11. \square

Similar constructions could yield applications of the $(\mathcal{M}, \mathcal{N})$ and $(\mathcal{O}, \mathcal{P})$ star surgeries, using the embeddings yielding the exotic copies of $\mathbb{C}P^2 \# 7 \overline{\mathbb{C}P^2}$ and $\mathbb{C}P^2 \# 5 \overline{\mathbb{C}P^2}$ of sections 5B and 5C.

6 Infinitely many exotic smooth structures on $\mathbb{C}P^2 \# 7 \overline{\mathbb{C}P^2}$

Using a different embedding of S_2 , we can produce other examples of exotic 4-manifolds using star surgery. Here we find an embedding which produces an exotic $\mathbb{C}P^2 \# 7 \overline{\mathbb{C}P^2}$ by combining our star surgery operation with a knot surgery in the double node neighborhood which was introduced by Fintushel and Stern [16]. Thanks to the knot surgery we will have a better control over the Seiberg–Witten invariants of our manifold. The price we pay is that our manifold is no longer symplectic.

Proof of Theorem 1.3 We will start with the elliptic fibration with two I_2 fibers, and two I_4 fibers whose homology classes are specified in Lemma 4.2. The homology classes of these spheres in $\mathbb{C}P^2 \# k \overline{\mathbb{C}P^2}$ will be crucial to our computation of the Seiberg–Witten invariant of the manifold resulting from star surgery along this embedding.

We follow the same steps as in [16]. Let K_n denote the n -twist knot. Recall that K_n admits a Seifert surface of genus one and its symmetrized Alexander polynomial is given by

$$\Delta_{K_n}(t) = nt - (2n + 1) + nt^{-1}.$$

Consider the elliptic fibration described in Lemma 4.2. We pick one of the I_2 fibers, say the one consisting of \tilde{C}_1 and \tilde{L}_1 , and perturb the fibration locally so the I_2 fiber turns into a double node neighborhood. Then we pick a regular fiber in the double node neighborhood and do knot surgery on it. The knot surgery operation does not preserve the fibration structure inside the double node neighborhood. In particular the section represented by the exceptional sphere E_2 does not survive after the knot surgery. On the other hand if we choose the gluing map in the knot surgery carefully, then E_2 turns into a *pseudosection*, an immersed sphere with one transverse self-intersection which is a section outside of the double node neighborhood.

Before describing our construction further, we would like to make a couple of observations about the manifold $E(1)_{K_n}$ which is the result of the knot surgery described above. First note that $E(1)_{K_n}$ is simply connected: the fundamental group of the complement of a regular fiber is generated by a normal circle which bounds a disk in E_2 . The second observation is that the only Seiberg–Witten basic classes of $E(1)_{K_n}$ are $\pm \text{PD}([F])$, where $[F]$ is the fiber class, and the small perturbation Seiberg–Witten




Figure 20: $E(1)_{K_n}$.




Figure 21: Configuration S_2 inside $E(1)_{K_n} \# \overline{\mathbb{C}P^2}$.

invariants at these classes are n [46; 15]. Hence $E(1)_{K_n}$ is homeomorphic but not diffeomorphic to $E(1) = \mathbb{C}P^2 \# 9 \overline{\mathbb{C}P^2}$.

We continue with the construction. Blow up $E(1)_{K_n}$ at the double point of the pseudo-section E_2 . The proper transform of the pseudo-section is an embedded sphere S whose homology class is represented by $e_2 - 2e_{10}$. Now we see the configuration S_2 embedded in $E(1)_{K_n} \# \overline{\mathbb{C}P^2}$ using Lemma 4.2:

$$\begin{aligned}
 [u_0] &= [S] + [\tilde{E}_4] = e_1 - 2e_{10}, \\
 [u_1] &= [\tilde{C}_2] = 2h - e_1 - e_2 - e_3 - e_4 - e_5 - e_6, \\
 [u_2] &= [\tilde{Z}_3] = h - e_1 - e_2 - e_9, \\
 [u_3] &= [\tilde{X}_4] = h - e_1 - e_6 - e_7, \\
 [u_4] &= [\tilde{Y}_4] = h - e_1 - e_5 - e_8.
 \end{aligned}$$

Let Y_n denote the result of star surgery of $E(1)_{K_n} \# \overline{\mathbb{C}P^2}$ along the configuration S_2 described above. We claim that Y_n is homeomorphic to $\mathbb{C}P^2 \# 7 \overline{\mathbb{C}P^2}$. First we must see that Y_n is simply connected. By Proposition 3.1, the generator of $\pi_1(\mathcal{T}_2)$ is represented by the curve in $\partial\mathcal{T}_2 = \partial S_2$ given by the boundary of the normal disk to any of the -2 -spheres in S_2 . We will take the representative curve which bounds the normal disk to u_2 . Since u_2 is one -2 -sphere in the I_4 fiber, and there are other -2 -spheres in that fiber which intersect u_2 transversally but are otherwise disjoint from the embedding of S_2 (\tilde{X}_3 or \tilde{E}_3), this curve bounds a disk in $E(1)_{K_n} \# \overline{\mathbb{C}P^2} \setminus S_2 = Y_n \setminus \mathcal{T}_2$. Therefore Y_n

is simply connected. Next we check that Y_n has the same Euler characteristic, signature, and parity of $\mathbb{C}P^2 \# 7\overline{\mathbb{C}P^2}$. A simple computation shows

$$\chi(Y_n) = \chi(E(1)_{K_n} \# \overline{\mathbb{C}P^2}) - \chi(\mathcal{S}_2) + \chi(\mathcal{T}_2) = 10,$$

and

$$\sigma(Y_n) = \sigma(E(1)_{K_n} \# \overline{\mathbb{C}P^2}) - \sigma(\mathcal{S}_2) + \sigma(\mathcal{T}_2) = -6.$$

Since $b_2(Y_n) = 8$ and $b_2^+(Y_n) = 1$, the intersection form cannot be a direct sum of hyperbolic pieces and E_8 's, so the parity of Y_n is odd. Therefore Y_n is homeomorphic to $\mathbb{C}P^2 \# 7\overline{\mathbb{C}P^2}$ by Freedman's theorem.

Finally we compute the Seiberg–Witten invariants of Y_n . By the blow-up formula $E(1)_{K_n} \# \overline{\mathbb{C}P^2}$ has exactly four Seiberg–Witten basic classes $\pm \text{PD}([F]) \pm e_{10}$. The small perturbation Seiberg–Witten invariant in the chamber of h evaluates as $\pm n$. We need to translate this information to a chamber whose representative homology class is orthogonal \mathcal{S}_2 . Consider the following homology class:

$$H := 50h - 32e_1 - 14e_2 - 12e_3 - 21e_4 - 5e_5 - 15e_6 - 3e_7 - 12e_8 - 4e_9 - 16e_{10}.$$

It can be checked that $H \cdot H > 0$, $H \cdot h > 0$, and $H \cdot [u_i] = 0$ for all $i = 0, \dots, 4$. Let $K = -\text{PD}([F]) - e_{10} = -3h + e_1 + \dots + e_9 - e_{10}$. We have $K \cdot H < 0$ and $K \cdot h < 0$. Hence there is no wall between the chambers determined by H and h with respect to K . Note that $K|_{\mathcal{S}_2}$ is the canonical class of \mathcal{S}_2 , so K descends to Y_n as a characteristic class \tilde{K} with $\tilde{K}|_{\mathcal{T}_2} = 0$.

Let $X_n = E(1)_{K_n} \# \overline{\mathbb{C}P^2}$, we will check if $d_{X_n}(K) \geq 0$ and $d_{Y_n}(\tilde{K}) \geq 0$. Clearly

$$d_{X_n}(K) = \frac{K^2 - 3\sigma(X_n) - 2\chi(X_n)}{4} = \frac{-1 - 3(-9) - 2(13)}{4} = 0.$$

On the other hand

$$\begin{aligned} d_{Y_n}(\tilde{K}) &= \frac{\tilde{K}^2 - 3\sigma(Y_n) - 2\chi(Y_n)}{4} \\ &= \frac{(K)^2 - (K|_{\mathcal{S}_2})^2 + (\tilde{K}|_{\mathcal{T}_2})^2 - 3\sigma(Y_n) - 2\chi(Y_n)}{4} \\ &= \frac{(-1) - (-3) + (0) - 3(-6) - 2(10)}{4} \\ &= 0. \end{aligned}$$

Hence by [Theorem 5.10](#), we have

$$|\text{SW}_{Y_n, H}(\pm \tilde{K})| = |\text{SW}_{E(1)_{K_n} \# \overline{\mathbb{C}P^2}, H}(\pm K)| = n.$$

Since the small perturbation Seiberg–Witten invariant is well-defined for those manifolds with $b_2^- \leq 9$, we conclude that Y_n has at least two basic classes. In particular Y_n is not diffeomorphic to $\mathbb{C}P^2 \# 7 \overline{\mathbb{C}P^2}$, which does not have any basic classes.

It remains to prove the minimality of Y_n for $n \geq 2$. By the blow-up formula, it suffices to show that there are exactly two basic classes whose Seiberg–Witten invariants are $\pm n$. We will show that $\pm \tilde{K}$ are the only Seiberg–Witten basic classes of Y_n satisfying $|\text{SW}_{Y_n}(\pm \tilde{K})| = n$. In other words, we will prove that the cohomology class $P := -\text{PD}([F]) + e_{10}$ (the only other basic class up to sign of X_n), does not descend to a basic class of Y_n . Suppose, to the contrary, that there is a basic class \tilde{P} of Y_n such that $\tilde{P}|_{Y_n - \mathcal{T}_2} = P|_{X_n \setminus \mathcal{S}_2}$. Then

$$\begin{aligned} d_{Y_n}(\tilde{P}) &= \frac{\tilde{P}^2 - 3\sigma(Y_n) - 2\chi(Y_n)}{4} \\ &= \frac{(P)^2 - (P|_{\mathcal{S}_2})^2 + (\tilde{P}|_{\mathcal{T}_2})^2 - 3\sigma(Y_n) - 2\chi(Y_n)}{4} \\ &= \frac{(-1) - (-1/3) + (\tilde{P}|_{\mathcal{T}_2})^2 - 3(-6) - 2(10)}{4} \\ &= -\frac{13}{6} + \frac{1}{4}(\tilde{P}|_{\mathcal{T}_2})^2 < 0. \end{aligned}$$

The last inequality follows from the fact that the intersection form of \mathcal{T}_2 is negative-definite. This contradicts the assumption that \tilde{P} is a basic class. \square

References

- [1] **A Akhmedov**, *Small exotic 4–manifolds*, *Algebr. Geom. Topol.* 8 (2008) 1781–1794 [MR2448872](#)
- [2] **A Akhmedov, R İ Baykur, B D Park**, *Constructing infinitely many smooth structures on small 4–manifolds*, *J. Topol.* 1 (2008) 409–428 [MR2399137](#)
- [3] **A Akhmedov, B D Park**, *Exotic smooth structures on small 4–manifolds*, *Invent. Math.* 173 (2008) 209–223 [MR2403396](#)
- [4] **A Akhmedov, B D Park**, *Exotic smooth structures on small 4–manifolds with odd signatures*, *Invent. Math.* 181 (2010) 577–603 [MR2660453](#)
- [5] **S Baldridge, P Kirk**, *A symplectic manifold homeomorphic but not diffeomorphic to $\mathbb{C}P^2 \# 3 \overline{\mathbb{C}P^2}$* , *Geom. Topol.* 12 (2008) 919–940 [MR2403801](#)
- [6] **M Bhupal, B Özbağcı**, *Symplectic fillings of lens spaces as Lefschetz fibrations* (2013) [arXiv:1307.6935](#)
- [7] **M Bhupal, A I Stipsicz**, *Weighted homogeneous singularities and rational homology disk smoothings*, *Amer. J. Math.* 133 (2011) 1259–1297 [MR2843099](#)

- [8] **V Colin, P Ghiggini, K Honda**, *Equivalence of Heegaard Floer homology and embedded contact homology via open book decompositions*, Proc. Natl. Acad. Sci. USA 108 (2011) 8100–8105 [MR2806645](#)
- [9] **S K Donaldson**, *Irrationality and the h -cobordism conjecture*, J. Differential Geom. 26 (1987) 141–168 [MR892034](#)
- [10] **H Endo, Y Z Gurtas**, *Lantern relations and rational blowdowns*, Proc. Amer. Math. Soc. 138 (2010) 1131–1142 [MR2566578](#)
- [11] **H Endo, T E Mark, J Van Horn-Morris**, *Monodromy substitutions and rational blowdowns*, J. Topol. 4 (2011) 227–253 [MR2783383](#)
- [12] **J B Etnyre, B Özbağcı**, *Invariants of contact structures from open books*, Trans. Amer. Math. Soc. 360 (2008) 3133–3151 [MR2379791](#)
- [13] **R Fintushel, B D Park, R J Stern**, *Reverse engineering small 4-manifolds*, Algebr. Geom. Topol. 7 (2007) 2103–2116 [MR2366188](#)
- [14] **R Fintushel, R J Stern**, *Rational blowdowns of smooth 4-manifolds*, J. Differential Geom. 46 (1997) 181–235 [MR1484044](#)
- [15] **R Fintushel, R J Stern**, *Knots, links, and 4-manifolds*, Invent. Math. 134 (1998) 363–400 [MR1650308](#)
- [16] **R Fintushel, R J Stern**, *Double node neighborhoods and families of simply connected 4-manifolds with $b^+ = 1$* , J. Amer. Math. Soc. 19 (2006) 171–180 [MR2169045](#)
- [17] **R Fintushel, R J Stern**, *Six lectures on four 4-manifolds*, from: “Low dimensional topology”, (T S Mrowka, P S Ozsváth, editors), IAS/Park City Math. Ser. 15, Amer. Math. Soc. (2009) 265–315 [MR2503498](#)
- [18] **R Fintushel, R J Stern**, *Pinwheels and nullhomologous surgery on 4-manifolds with $b^+ = 1$* , Algebr. Geom. Topol. 11 (2011) 1649–1699 [MR2821436](#)
- [19] **M H Freedman**, *The topology of four-dimensional manifolds*, J. Differential Geom. 17 (1982) 357–453 [MR679066](#)
- [20] **R Friedman, J W Morgan**, *On the diffeomorphism types of certain algebraic surfaces. I*, J. Differential Geom. 27 (1988) 297–369 [MR925124](#)
- [21] **D Gay, T E Mark**, *Convex plumbings and Lefschetz fibrations*, J. Symplectic Geom. 11 (2013) 363–375 [MR3100798](#)
- [22] **D T Gay, A I Stipsicz**, *Symplectic rational blow-down along Seifert fibered 3-manifolds*, Int. Math. Res. Not. 2007 (2007) Art. ID rnm084, 20 [MR2376205](#)
- [23] **D T Gay, A I Stipsicz**, *Symplectic surgeries and normal surface singularities*, Algebr. Geom. Topol. 9 (2009) 2203–2223 [MR2558309](#)
- [24] **R E Gompf**, *Handlebody construction of Stein surfaces*, Ann. of Math. 148 (1998) 619–693 [MR1668563](#)

- [25] **D Kotschick**, *On manifolds homeomorphic to $\mathbb{C}P^2\#8\overline{\mathbb{C}P^2}$* , Invent. Math. 95 (1989) 591–600 [MR979367](#)
- [26] **Ç Kutluhan, Y-J Lee, C H Taubes**, *$HF = HM I$: Heegaard Floer homology and Seiberg–Witten Floer homology* (2010) [arXiv:1007.1979](#)
- [27] **T-J Li**, *The Kodaira dimension of symplectic 4–manifolds*, from: “Floer homology, gauge theory, and low-dimensional topology”, (D A Ellwood, P S Ozsváth, A I Stipsicz, Z Szabó, editors), Clay Math. Proc. 5, Amer. Math. Soc. (2006) 249–261 [MR2249257](#)
- [28] **T J Li, A Liu**, *Symplectic structure on ruled surfaces and a generalized adjunction formula*, Math. Res. Lett. 2 (1995) 453–471 [MR1355707](#)
- [29] **D McDuff, D Salamon**, *Introduction to symplectic topology*, Oxford Mathematical Monographs, The Clarendon Press Oxford University Press, New York (1995) [MR1373431](#)
- [30] **M Michalogiorgaki**, *Rational blow-down along Wahl type plumbing trees of spheres*, Algebr. Geom. Topol. 7 (2007) 1327–1343 [MR2350284](#)
- [31] **J W Morgan**, *The Seiberg–Witten equations and applications to the topology of smooth four-manifolds*, Mathematical Notes 44, Princeton Univ. Press (1996) [MR1367507](#)
- [32] **P Ozsváth, Z Szabó**, *The symplectic Thom conjecture*, Ann. of Math. 151 (2000) 93–124 [MR1745017](#)
- [33] **P Ozsváth, Z Szabó**, *Holomorphic triangle invariants and the topology of symplectic four-manifolds*, Duke Math. J. 121 (2004) 1–34 [MR2031164](#)
- [34] **H Park, A I Stipsicz**, *Smoothings of singularities and symplectic surgery* (2012) [arXiv:1211.6830v1](#)
- [35] **J Park**, *Seiberg–Witten invariants of generalised rational blow-downs*, Bull. Austral. Math. Soc. 56 (1997) 363–384 [MR1490654](#)
- [36] **J Park**, *Simply connected symplectic 4–manifolds with $b_2^+ = 1$ and $c_1^2 = 2$* , Invent. Math. 159 (2005) 657–667 [MR2125736](#)
- [37] **J Park, A I Stipsicz, Z Szabó**, *Exotic smooth structures on $\mathbb{C}P^2\#5\overline{\mathbb{C}P^2}$* , Math. Res. Lett. 12 (2005) 701–712 [MR2189231](#)
- [38] **U Persson**, *Configurations of Kodaira fibers on rational elliptic surfaces*, Math. Z. 205 (1990) 1–47 [MR1069483](#)
- [39] **L P Roberts**, *Rational blow-downs in Heegaard–Floer homology*, Commun. Contemp. Math. 10 (2008) 491–522 [MR2444846](#)
- [40] **L Starkston**, *Comparing star surgery to rational blow-down* (2014) [arXiv:1407.3293](#)
- [41] **L Starkston**, *Symplectic fillings of Seifert fibered spaces*, Trans. Amer. Math. Soc. 367 (2015) 5971–6016 [MR3347194](#)

- [42] **A I Stipsicz, Z Szabó**, *An exotic smooth structure on $\mathbb{C}P^2\#6\overline{\mathbb{C}P^2}$* , *Geom. Topol.* 9 (2005) 813–832 [MR2140993](#)
- [43] **A I Stipsicz, Z Szabó, J Wahl**, *Rational blowdowns and smoothings of surface singularities*, *J. Topol.* 1 (2008) 477–517 [MR2399141](#)
- [44] **M Symington**, *Symplectic rational blowdowns*, *J. Differential Geom.* 50 (1998) 505–518 [MR1690738](#)
- [45] **M Symington**, *Generalized symplectic rational blowdowns*, *Algebr. Geom. Topol.* 1 (2001) 503–518 [MR1852770](#)
- [46] **Z Szabó**, *Exotic 4-manifolds with $b_2^+ = 1$* , *Math. Res. Lett.* 3 (1996) 731–741 [MR1426531](#)
- [47] **C Wendl**, *Strongly fillable contact manifolds and J -holomorphic foliations*, *Duke Math. J.* 151 (2010) 337–384 [MR2605865](#)

Department of Mathematics, Bogazici University
34342 Istanbul, Turkey

Mathematics Department, Stanford University
450 Serra Mall, Building 380, Stanford, CA 78712-1202, United States

cagri.karakurt@boun.edu.tr, lstarkst@stanford.edu

Received: 8 October 2014 Revised: 15 September 2015

