

Center of the Goldman Lie algebra

ARPAN KABIRAJ

We show that the center of the Goldman Lie algebra associated to a closed orientable surface is generated by the class of the trivial loop. For an orientable nonclosed surface of finite type, the center is generated by closed curves which are either homotopically trivial or homotopic to boundary components or punctures.

57M50; 57M07, 57M05

1 Introduction

Let F be an oriented surface. Given two free homotopy classes of oriented closed curves α and β , consider two oriented closed curves x and y representing α and β , respectively. Performing a small homotopy if necessary, we can assume that x and y intersect transversally in double points. Goldman [10] defined the bracket of α and β as the sum,

$$[\alpha, \beta] = \sum_{p \in x \cap y} \iota(p) \langle x *_p y \rangle,$$

where $x \cap y$ denotes the set of all intersection points between x and y , $\iota(p)$ denotes the sign of the intersection between x and y at p , $(x *_p y)$ denotes the loop product of x and y at p , and $\langle z \rangle$ denotes the free homotopy class of a curve z .

Let \mathcal{C} be the set of all free homotopy classes of oriented closed curves in F . This bracket is extended by linearity to $\mathbb{Z}(\mathcal{C})$, the free module generated by \mathcal{C} . Goldman [10] showed that this bracket is well defined, skew-symmetric and satisfies the Jacobi identity. Therefore, this is a Lie bracket, and it gives a Lie algebra structure on $\mathbb{Z}(\mathcal{C})$, which we denote by $\mathcal{L}(F)$. Recall that the *center of a Lie algebra* \mathcal{L} is the set of all elements x in \mathcal{L} such that $[x, y] = 0$ for all y in \mathcal{L} . The main object of this paper is to study the center of $\mathcal{L}(F)$.

The structure of the Goldman Lie algebra for surfaces of nonnegative Euler characteristic is either trivial or well understood; see Chas [3, Lemma 7.6] for the torus case.

Chas and Sullivan conjectured that, for a closed surface F , the center of the Goldman algebra is generated by the trivial loop. It is natural to conjecture (see Chas [4,

Open Problem 1], [3, Problem 13.1] and Kawazumi and Kuno [12, Section 8.3]) that, for a surface F with nonempty boundary, the center of $\mathcal{L}(F)$ is generated by the free homotopy classes of oriented closed curves which are either homotopic to a point, homotopic to a boundary, or homotopic to a puncture. In this paper, we prove these conjectures.

Main Theorem *The center of the Goldman Lie algebra of any closed orientable surface F is one-dimensional, and is generated by the class of the trivial loop. If F is an orientable surface of finite type with boundary, then the center of $\mathcal{L}(F)$ is generated by the set of all free homotopy classes of oriented closed curves which are homotopic to either a point, a boundary component, or a puncture.*

Remark 1 The closed case was done by Etingof in [8] using representation theory, but that proof did not address the case of surfaces with boundary. Our proof of both cases uses different ideas from hyperbolic geometry.

Goldman discovered this bracket while studying the Weil–Petersson symplectic form on Teichmüller spaces. Using Wolpert’s [15] result on length and twist flow, he showed that if the Goldman bracket between two closed curves is zero and one of them has a simple representative, then their geometric intersection number is zero. The combinatorial structure of $\mathcal{L}(F)$ has also been studied. Using combinatorial topology and group theory, Chas [3] proved a stronger version of Goldman’s result, namely if one of the curves has a simple representative, then the number of terms in the Goldman bracket is the same as their geometric intersection number. Chas and Krongold [6] proved that, for a compact surface with nonempty boundary, $[x, x^3]$ determines the self-intersection number of x . Using hyperbolic geometry, Chas and Gadgil [5] proved that there exists a positive integer m_0 such that, for all $m \geq m_0$, the geometric intersection number between x and y is the number of terms in $[x^m, y]$ divided by m . There is a Lie cobracket defined by Turaev [14] on $\mathbb{Z}(\mathcal{C})$ which is the dual object of the Goldman bracket. This structure has been studied by Chas and Krongold [7; 2].

Idea of the proof Our proof is based on hyperbolic geometry. Given an oriented surface of negative Euler characteristic, we fix a hyperbolic metric on it with geodesic boundary. There are two key ideas behind our proof.

The first idea is from [5]. Given two closed oriented curves x and y intersecting transversally, we construct lifts of $(x *_p y)$ in the hyperbolic plane \mathbb{H} for each intersection point p . By [5, Lemma 7.1], the lifts are quasigeodesics. Hence they are homotopic to unique geodesics. Therefore, if two terms $(x *_p y)$ and $(x *_q y)$ cancel each other, then the corresponding geodesics will be the same. By [5, Main Theorem],

there exists m_0 such that if we take a power $m \geq m_0$ of x , then we can ensure that if the geodesics are the same, then the quasigeodesics are also the same, and hence the terms have the same sign. Therefore, there is no cancellation between the terms of $[x^m, y]$.

The second key idea is that all lifts of a simple closed geodesic are disjoint. Now, if an element $y = \sum_{i=1}^k y_i$ of $\mathcal{L}(F)$ belongs to the center, then we consider a simple closed curve x which intersects at least one of the curves y_i nontrivially. Taking sufficiently large powers of x we can ensure that the same terms of $[x^m, y_i]$ have the same sign. Then, using that the lifts of x are disjoint, we show that if one term of $[x^m, y_i]$ and another term of $[x^m, y_j]$ are the same, then y_i and y_j are conjugate.

Therefore, if $[x, y]$ is zero for all closed curves x , then each y_i is disjoint from every simple closed curve, and hence each y_i is either homotopic to a point or to a boundary component or to a puncture.

Organization of the paper Throughout the paper we follow the notation and definitions from [5].

In Section 2, we recall some basic facts about hyperbolic surfaces and closed curves on hyperbolic surfaces. We also mention a well-known result about hyperbolic elements of the fundamental group of a hyperbolic surface.

In Section 3, we recall from [5] the algebraic definition of the Goldman bracket between conjugacy classes of elements. Throughout the paper, we use this as the definition of Goldman bracket.

In Section 4, we recall from [5] the description of the lifts of the terms of Goldman bracket. We also state the lemma that these lifts are quasigeodesic and, therefore, in a neighborhood of a geodesic, following [5].

In Section 5, we show that if we take a sufficiently high power of a simple closed curve, then there is no cancellation between the terms of the Goldman bracket with any other closed curve.

In Section 6, we mention a classical theorem and prove the main theorems.

Acknowledgements The author would like to thank Siddhartha Gadgil for his encouragement and enlightening conversations. The author would like to thank Bidyut Sanki and Divakaran D. for all the discussions which helped him to understand the problem. The author would also like to thank Moira Chas for her valuable suggestions and comments. The author is supported by NET-CSIR (India) Senior Research Fellowship.

2 Closed curves on a hyperbolic surface

In this section, we recall some basic facts about hyperbolic surfaces and closed curves on hyperbolic surfaces. We use the same notation as [5]. References for the results mentioned in this section are [1; 5; 11; 13].

Let F be an orientable surface of finite type with negative Euler characteristic; ie F is a surface of genus g with b boundary components and n punctures such that $2-2g-b-n < 0$. By [9, Theorem 1.2], we can endow F with a hyperbolic metric. By a hyperbolic surface we mean an orientable surface with negative Euler characteristic and with a given hyperbolic metric. Given a hyperbolic surface F , we identify the fundamental group $\pi_1(F)$ of F with a discrete subgroup of $\mathrm{PSL}_2(\mathbb{R})$, the group of orientation preserving isometries of the upper half plane \mathbb{H} . The action of $\pi_1(F)$ on \mathbb{H} is properly discontinuous without fixed points, and the quotient space is isometric to F . Henceforth, by an isometry of \mathbb{H} we mean an orientation preserving isometry, and by a closed curve we mean an oriented closed curve.

A homotopically nontrivial closed curve in F is called *essential* if it is not homotopic to a puncture. A closed curve is called *peripheral* if it is homotopic to a power of a simple closed curve bounding a once-punctured disc. By a *lift* of a closed curve γ to \mathbb{H} , we mean the image of a lift $\mathbb{R} \rightarrow \mathbb{H}$ of the map $\gamma \circ \pi$, where $\pi: \mathbb{R} \rightarrow S^1$ is the usual covering map.

There is a bijective correspondence between the set of all free homotopy classes of oriented closed curves in F and the set of all conjugacy classes in $\pi_1(F)$ [5, Theorem 2.3]. Given an oriented closed curve γ in F , we denote both its free homotopy class and the corresponding conjugacy class in $\pi_1(F)$ by $\langle \gamma \rangle$. Abusing notation, we sometimes denote the conjugacy class of γ by γ itself. Given $a, g \in \pi_1(F)$, we denote gag^{-1} by a^g and the translation length of a by τ_a . If a is hyperbolic, then a^g is also hyperbolic with $\tau_{a^g} = \tau_a$ and $A_{a^g} = gA_a$ for all $g \in \pi_1(F)$, where A_a denotes the axis of a .

The *geometric intersection number* between two free homotopy classes of closed curves x and y , denoted by $i(x, y)$, is defined to be the minimal number of intersection points between a representative curve in the class $\langle x \rangle$ and a representative curve in the class $\langle y \rangle$ which intersect transversally in double points.

Every free homotopy class of an essential closed curve contains a unique closed geodesic whose length is the same as the translation length of any element of the corresponding conjugacy class. By a slight abuse of notation, we denote the free homotopy classes of essential closed curves by their geodesic representatives.

The following lemma is a well-known result. See [5, Corollary 6.3] for a proof.

Lemma 2.1 *Let G be a discrete subgroup of $\mathrm{PSL}_2(\mathbb{R})$. Given two nonzero positive numbers L and C , there exists a constant $M > 0$ such that for every pair of hyperbolic elements x and y in G with $\tau_x \leq L$ and $\tau_y \leq L$, the set*

$$\{x \in A_x : d(x, A_y) < C\}$$

is either empty or a geodesic segment of length at most M .

3 Goldman bracket

In this section, we recall from [5] the algebraic definition of the Goldman bracket between two curves intersecting transversally (not necessarily in double points). For the equivalence of this definition with the previous one, see [5, Section 3].

Given two hyperbolic transformations x and y whose axes A_x and A_y , respectively, intersect in a point P , let $I(x, y)$ denote the segment of A_x of length τ_x starting from P in the positive direction of A_x , containing P but not containing xP .

Definition 3.1 Let $\langle x \rangle$ and $\langle y \rangle$ be two nontrivial conjugacy classes in $\pi_1(F)$. Define

$$[\langle x \rangle, \langle y \rangle] = \begin{cases} \sum_{gY \in J(x,y)} \iota(x, y^g) \langle xy^g \rangle & \text{if both } x \text{ and } y \text{ are hyperbolic,} \\ 0 & \text{if either } x \text{ or } y \text{ is parabolic,} \end{cases}$$

where Y is the cyclic subgroup generated by y ,

$$J(x, y) = \{gY \in \pi_1(F)/Y : I(x, y) \cap gA_y \neq \emptyset\},$$

and $\iota(x, y)$ denotes the sign of intersection between the axes of x and y if they intersect and 0 otherwise.

Remark 2 This definition is independent of the type of the intersection points of the representative curves. Therefore, we can use the geodesic representatives of the corresponding conjugacy classes (which intersect transversally but not necessarily in double points). Henceforth, we use this as the definition of the Goldman bracket.

4 Terms of the Goldman bracket

In this section we recall the description of the lifts of the terms of the Goldman bracket from [5, Section 7].

Let x and y be two hyperbolic elements in $\pi_1(F)$ whose axes intersect at the point P . Denote the projections of A_x and A_y in F by x_1 and y_1 respectively. Let p be the




Figure 1: Lift of a term in the Goldman bracket

projection of P in F . By [5, Remark 3.2], there exists a unique $g \in \pi_1(F)$ such that $gY \in J(x, y)$ and gY corresponds to p . A lift of $(x_1 *_p y_1)$ is a bi-infinite piecewise geodesic passing through P , which we denote by $\gamma(x, y^g)$; see Figure 1.

Let $h = xy^g$. If we denote the geodesic arc from P to $x(P)$ by I_x^0 and the geodesic segment from $x(P)$ to $h(P)$ by $I_{y^g}^0$, then $\gamma(x, y^g)$ consists of geodesic segments of the form $h^k(I_x^0)$ and $h^k(I_{y^g}^0)$ occurring alternately.

Remark 3 Denote $h^k(I_x^0)$ by I_x^k and $h^k(I_{y^g}^0)$ by $I_{y^g}^k$. From the definition, the length of I_x^k is τ_x and the length of $I_{y^g}^k$ is τ_y for all $k \in \mathbb{Z}$. Hence, by the description of the axis of the product of two isometries given in [5, Remark 3.4], A_h intersects I_x^k and $I_{y^g}^k$ in their midpoint for all $k \in \mathbb{Z}$.

For the definition of quasigeodesic and the proof of the following lemma, see [5, Section 7, Lemmas 7.1 and 7.2].

Lemma 4.1 Given $L > 0$, there exist $K \geq 1$ and $C > 0$, depending on $\pi_1(F)$, such that if x and y are two hyperbolic elements in $\pi_1(F)$ whose axes are distinct with $\tau_x \leq L$ and $\tau_y \leq L$, then for any $g \in \pi_1(F)$ and $m \in \mathbb{N}$, if A_x and A_{y^g} intersect, then:

- (1) $\gamma(x^m, y^g)$ is a K -quasigeodesic, and $\gamma(x^m, y^g)$ is homotopic to A_h , where $h = x^m y^g$;
- (2) $\gamma(x^m, y^g) \subset N_{C/2}(A_h)$ and $A_h \subset N_{C/2}(\gamma(x^m, y^g))$, where $N_C(A_h)$ denotes the C neighborhood of A_h .

5 Noncancellation lemma

Denote the length of a curve x by $l(x)$. For the proof of the following lemma, see [5, Lemma 7.3].

Lemma 5.1 *Let L, K and C be as in Lemma 4.1. For hyperbolic elements $x, y \in \pi_1(F)$ with $\tau_x \leq L$ and $\tau_y \leq L$, let m be a positive integer such that $m\tau_x > 6KC$.*

Let S and R be the points in $I_{x^m}^0$ at distance $3KC$ from P and $x^m P$ (see Figure 2). Let s (respectively r) be the open half-plane bounded by the line perpendicular to A_x through S (respectively R), containing the point $x^m P$ (respectively P).

Set $U = s \cap r \cap N_C(I_{x^m}^0)$. Then U contains an open segment J of $I_{x^m}^0$ such that $N_C(I_{x^m}^0) \setminus U$ is disconnected, $l(J) \geq m\tau_x - 6KC$, $U \subset N_C(I_{x^m}^0)$, $\bar{U} \cap N_C(I_{x^m}^k) = \emptyset$ for all $k \neq 0$ and $\bar{U} \cap N_C(I_y^k) = \emptyset$ for all $k \in \mathbb{Z}$.

The following lemma is the main lemma of this paper. The proof is based on the proof of [5, Lemma 7.4, Claims 1 and 2] and the idea that lifts of simple closed geodesics are disjoint.

Lemma 5.2 *Let x be a hyperbolic element in $\pi_1(F)$ such that the geodesic representative in the free homotopy class of x is simple. Let $x_1 = x^h$ for some $h \in \pi_1(F)$. Suppose y and y_1 are two distinct hyperbolic elements in $\pi_1(F)$ whose axes are distinct and intersect the axes of x and x_1 , respectively. Let L be a positive number such that the translation lengths of x, y and y_1 are bounded above by L . Then there exists m_0 such that for any $m > m_0$, we have $\gamma(x^m, y) = \gamma(x_1^m, y_1)$ whenever $x^m y = x_1^m y_1$. Moreover, there exists $u \in \pi_1(F)$ such that $x_1 = x^u$ and $y_1 = y^u$.*




Figure 2: The open segment J as in Lemma 5.2

Proof Let τ_0 be the systole, ie the length of a shortest length closed geodesic of F .

Let C, K and M be the constants defined in Lemmas 4.1 and 2.1 and let $m > m_0$. Define $m_0 = K(3M + 10C)/\tau_0$.

Since $x^m y = x_1^m y_1$, we have $A_{x^m y} = A_{x_1^m y_1}$. Let $g = x^m y = x_1^m y_1$. By Lemma 5.1,

$$(1) \quad \gamma(x_1^m, y_1) \subset N_{C/2}(A_g) \subset N_C(\gamma(x^m, y)).$$

Let J and U be as in Lemma 5.1 corresponding to $\gamma(x^m, y)$. Therefore, $J \subset U$, $J \subset I_{x^m}^0 \subset \gamma(x^m, y)$, and $\text{length}(J) \geq m\tau_x - 6KC$.

Claim 1 *The curve $\gamma(x_1^m, y_1)$ intersects U and does not intersect the part of the boundary of U contained in the boundary of $N_C(\gamma(x^m, y))$.*

Proof of claim If $\gamma(x_1^m, y_1)$ does not intersect U , by (1), $\gamma(x_1^m, y_1)$ is contained in $N_C(\gamma(x^m, y)) \setminus U = N_C(\gamma(x^m, y) \setminus J)$, which is disconnected. Hence by (1), $\gamma(x_1^m, y_1)$ should intersect both components, which contradicts that $\gamma(x_1^m, y_1)$ is connected. By Lemma 5.1, $\bar{U} \cap N_C(I_{x^m}^k) = \emptyset$ for all $k \neq 0$ and $\bar{U} \cap N_C(I_y^k) = \emptyset$ for all $k \in \mathbb{Z}$. Therefore, $\gamma(x_1^m, y_1)$ does not intersect the part of the boundary of U contained in the boundary of $N_C(\gamma(x^m, y))$. □

Therefore, any component of $U \cap \gamma(x_1^m, y_1)$ consists of piecewise geodesic arcs starting and ending at the sides of U of length $2C$.

Claim 2 *Let β be a component of $\gamma(x_1^m, y_1) \cap U$. Then β contains a geodesic segment l of length greater than M .*

Proof of claim Case 1 Suppose β contains more than three vertices. Then β contains $I_{x_1^m}^k$ for some $k \in \mathbb{Z}$ and $\text{length}(I_{x_1^m}^k) = m\tau_{x_1} > m_0\tau_0 = K(3M + 10C) > M$.

Case 2 Suppose β contains at most three vertices. Then β consists of at most three segments. Let v be the longest segment of β and let $r = l(v)$. By hypothesis, $l(J) \geq m\tau_x - 6KC > 3KM + 4KC$. Using the triangle inequality and the properties of m and K , we have

$$(3M + 4C) \leq K(3M + 4C) < m\tau_x - 6KC \leq l(J) \leq 2C + 3r + 2C = 3r + 4C.$$

Hence $r > M$ which proves Claim 2. □

The geodesic segment v is contained in $\gamma(x_1^m, y_1)$ and $v \subset N_C(I_{x^m}^0)$. Therefore, by Lemma 2.1, v intersects $I_{x^m}^0$ in a geodesic segment. Hence $I_{x^m}^0$ and $\gamma(x_1^m, y_1)$ intersect in a geodesic segment.

Claim 3 If $\gamma(x^m, y)$ and $\gamma(x_1^m, y_1)$ intersect in a geodesic segment contained in $I_{x^m}^0$, then they are equal, and there exists $u \in \pi_1(F)$ such that $x_1 = x^u$ and $y_1 = y^u$.

Proof of claim As $I_{x^m}^0$ intersects $\gamma(x_1^m, y_1)$ in a geodesic segment, $I_{x^m}^0$ intersects either $I_{x_1^m}^k$ or $I_{y_1}^k$ in a geodesic segment.

Now $l(I_{x_1^m}^k) = m\tau_{x_1} = m\tau_x = l(I_{x^m}^0)$ for all $k \in \mathbb{Z}$, and A_g intersects $I_{x^m}^0$ and $I_{x_1^m}^k$ in their midpoints. So if $I_{x^m}^0$ intersects $I_{x_1^m}^k$ in a geodesic segment, then they are equal.

If $I_{x^m}^0$ intersects $I_{y_1}^k$ in a geodesic segment for some $k \in \mathbb{Z}$, then by the construction of $\gamma(x^m, y)$, we see that A_x intersects $I_{x_1^m}^{k+1}$, which lies in a translate of the geodesic A_{x_1} and hence in a translate of A_x (as x and x_1 are conjugates). As the geodesic representative in the free homotopy class of x is simple, all translates of A_x are disjoint. Hence $I_{x^m}^0$ cannot intersect $I_{y_1}^k$ for any $k \in \mathbb{Z}$.

Since $I_{x^m}^0$ can not intersect $I_{y_1}^k$ in a geodesic segment, $I_{x^m}^0$ intersects $I_{x_1^m}^k$ for some $k \in \mathbb{Z}$. Thus $I_{x^m}^0 = I_{x_1^m}^k$. Since I_y^0 and $I_{y_1}^k$ are the unique geodesic segments joining the end point of $I_{x^m}^0 = I_{x_1^m}^k$ with the image of the starting point of $I_{x^m}^0 = I_{x_1^m}^k$ under g , we see that $I_y^0 = I_{y_1}^k$. By the periodic property of the definition of $\gamma(x^m, y)$ and $\gamma(x_1^m, y_1)$, they are equal. Since $g^n I_{x^m}^0 = I_{x_1^m}^0$ and $g^n I_y^0 = I_{y_1}^0$ for some n , taking $u = g^n$, we have $x_1 = x^u$ and $y_1 = y^u$. This proves the claim and thus the lemma. \square

6 Center of the Goldman Lie algebra

Lemma 6.1 Let F be a hyperbolic surface. Suppose x is an essential simple closed curve and y is an essential closed curve. If $i(x, y) \neq 0$, then there exists m_0 such that $[x^m, y] \neq 0$ for all $m > m_0$.

Proof Let $L = \max\{\tau_x, \tau_y\}$ and m_0 be as in Lemma 5.2. If $m > m_0$, then

$$[x^m, y] = m \left(\sum_{kB \in J(x^m, y)} \iota(x^m, y^k) \langle x^m y^k \rangle \right).$$

Suppose $\langle x^m y^k \rangle = \langle x^m y^{k_1} \rangle$. Then for some $g \in \pi_1(F)$,

$$x^m y^k = (x^m y^{k_1})g = (x^m)g(y)^{k_1}g.$$

By Lemma 5.2, there exists $u \in \pi_1(F)$ such that x is conjugate to x^g and y^k is conjugate to $y^{k_1}g$ by the element u . Therefore,

$$\iota(x^m, y^k) = \iota((x^m)^u, (y^k)^u) = \iota(x^{mg}, y^{k_1}g) = \iota(x^m, y^{k_1}).$$

Hence $[x^m, y] \neq 0$. \square

The following lemma is a classical result.

Lemma 6.2 *Let F be a hyperbolic surface of finite type with geodesic boundary. Let γ be a closed curve whose geometric intersection number with any other nontrivial simple closed geodesic is zero. Then γ is either homotopically trivial or homotopic to a boundary curve or peripheral.*

Theorem 6.3 *Let F be a hyperbolic surface of finite type with geodesic boundary. Let $y = \sum_{i=1}^n c_i y_i \in \mathcal{L}(F)$, where each y_i is a geodesic and $y_i \neq y_j$ for $i \neq j$. If y belongs to the center of $\mathcal{L}(F)$, then $i(x, y_i) = \emptyset$ for every simple closed geodesic x and for all $i \in \{1, 2, \dots, n\}$.*

Proof We show that, given any simple closed geodesic x , if $i(x, y_i) \neq \emptyset$ for some $i \in \{1, 2, \dots, n\}$, then there exists $m \in \mathbb{N}$ such that $[x^m, y] \neq 0$.

Let x be a simple closed geodesic which intersects at least one y_i . If some y_k is disjoint from x , then the Goldman bracket between x and y_k is zero; therefore, without loss of generality, assume that x intersects y_j for all $j \in \{1, 2, \dots, n\}$. Let $L = \max\{\tau_x, \tau_{y_1}, \tau_{y_2}, \dots, \tau_{y_n}\}$. Hence by Lemma 5.2, there exists m_i for every $i \in \{1, 2, \dots, n\}$ such that, if $m > \max\{m_i\}$ with $x_1^m = (x^m)^h$ for some $h \in \pi_1(F)$, then $\gamma(x^m, y_i) = \gamma(x_1^m, y_j)$ whenever $x^m y_i = x_1^m y_j$. Also there exists $g \in \pi_1(F)$ such that $x_1 = x^g$ and $y_j = y_i^g$. Since $[x^m, \sum_{i=1}^n c_i y_i] = \sum_{i=1}^n c_i [x^m, y_i]$, by Lemma 6.1, it is enough to show that the terms of $[x^m, y_i]$ are distinct from the terms of $[x^m, y_j]$ for $i \neq j$.

Suppose $\langle x^m y_i^{k_i} \rangle = \langle x^m y_j^{k_j} \rangle$. Hence there exists $h \in \pi_1(F)$ such that

$$x^m y_i^{k_i} = (x^m y_j^{k_j})^h = (x^m)^h (y_j)^{k_j h}.$$

By Lemma 5.2, $y_i^{k_i}$ and $y_j^{k_j}$ are conjugates of each other in $\pi_1(F)$. Therefore, y_i and y_j are freely homotopic to each other. Hence the geodesic representative corresponding to y_i and y_j are the same, which contradicts the assumption. \square

The Main Theorem follows at once from Lemma 6.2 and Theorem 6.3.

References

- [1] **A F Beardon**, *The geometry of discrete groups*, Graduate Texts in Mathematics 91, Springer, New York (1983) MR
- [2] **M Chas**, *Combinatorial Lie bialgebras of curves on surfaces*, Topology 43 (2004) 543–568 MR

- [3] **M Chas**, *Minimal intersection of curves on surfaces*, *Geom. Dedicata* 144 (2010) 25–60 MR
- [4] **M Chas**, *The Goldman bracket and the intersection of curves on surfaces*, from “Geometry, groups and dynamics” (C S Aravinda, W M Goldman, K Gongopadhyay, A Lubotzky, M Mj, A Weaver, editors), *Contemp. Math.* 639, Amer. Math. Soc., Providence, RI (2015) 73–83 MR
- [5] **M Chas, S Gadgil**, *The Goldman bracket determines intersection numbers for surfaces and orbifolds*, preprint (2012) arXiv
- [6] **M Chas, F Krongold**, *An algebraic characterization of simple closed curves on surfaces with boundary*, *J. Topol. Anal.* 2 (2010) 395–417 MR
- [7] **M Chas, F Krongold**, *Algebraic characterization of simple closed curves via Turaev’s cobracket*, *J. Topol.* 9 (2016) 91–104 MR
- [8] **P Etingof**, *Casimirs of the Goldman Lie algebra of a closed surface*, *Int. Math. Res. Not.* 2006 (2006) Art. ID 24894 MR
- [9] **B Farb, D Margalit**, *A primer on mapping class groups*, Princeton Mathematical Series 49, Princeton Univ. Press (2012) MR
- [10] **W M Goldman**, *Invariant functions on Lie groups and Hamiltonian flows of surface group representations*, *Invent. Math.* 85 (1986) 263–302 MR
- [11] **S Katok**, *Fuchsian groups*, University of Chicago Press (1992) MR
- [12] **N Kawazumi, Y Kuno**, *The Goldman–Turaev Lie bialgebra and the Johnson homomorphisms*, preprint (2013) arXiv
- [13] **J G Ratcliffe**, *Foundations of hyperbolic manifolds*, 2nd edition, Graduate Texts in Mathematics 149, Springer, New York (2006) MR
- [14] **V G Turaev**, *Skein quantization of Poisson algebras of loops on surfaces*, *Ann. Sci. École Norm. Sup.* 24 (1991) 635–704 MR
- [15] **S Wolpert**, *On the symplectic geometry of deformations of a hyperbolic surface*, *Ann. of Math.* 117 (1983) 207–234 MR

Department of Mathematics, Indian Institute of Science
Bangalore 560012, India

arpan.into@math.iisc.ernet.in

Received: 19 February 2015 Revised: 13 December 2015

