

Spectral sequences in smooth generalized cohomology

DANIEL GRADY
HISHAM SATI

We consider spectral sequences in smooth generalized cohomology theories, including differential generalized cohomology theories. The main differential spectral sequences will be of the Atiyah–Hirzebruch (AHSS) type, where we provide a filtration by the Čech resolution of smooth manifolds. This allows for systematic study of torsion in differential cohomology. We apply this in detail to smooth Deligne cohomology, differential topological complex K-theory and to a smooth extension of integral Morava K-theory that we introduce. In each case, we explicitly identify the differentials in the corresponding spectral sequences, which exhibit an interesting and systematic interplay between (refinements of) classical cohomology operations, operations involving differential forms and operations on cohomology with $U(1)$ coefficients.

55N15, 55T10, 55T25; 53C05, 55S05, 55S35

1 Introduction

Spectral sequences are very useful algebraic tools that often allow for efficient computations that would otherwise require brute force; see McCleary [54] for a broad survey. The Atiyah–Hirzebruch spectral sequence (henceforth AHSS) for K-theory and any generalized cohomology theory, in the topological sense, was introduced by Atiyah and Hirzebruch in [3]. An excellent introduction to the generalized cohomology AHSS can also be found in Hilton [38] and Adams [1, Section III.7]. Other useful references on the subject include Switzer [67] (Section 15, from a homology point of view, including the Gysin sequence from AHSS), and interesting remarks in relation to spectra are given in Rudyak [59]: Theorem 3.45 (homology), Remark 4.24 (sheaves and Čech), Remark 4.34 (Postnikov) and Corollary 7.12. A description with an eye for applications is given in Husemöller, Joachim, Jurčo and Schottenloher [42, Chapter 21].

The goal of this paper is to systematically study the spectral sequence in the context of smooth or differential cohomology; see Cheeger and Simons [20], Freed [27], Hopkins and Singer [41], Simons and Sullivan [66], Bunke [13], Bunke and Schick [17] and Schreiber [63]. Existence and interesting aspects of the AHSS in twisted forms of

such differential cohomology theories have been considered briefly by Bunke and Nikolaus [15], where the main interest was the effect of the geometric part of the twist on the spectral sequence. In this paper, we take a step back and consider untwisted differential generalized cohomology to systematically study the corresponding AHSS in generality and determine the differentials explicitly as cohomology operations. From the geometric point of view, one might expect on general grounds that the geometric information carried by the differential cohomology theory should somehow manifest itself within the spectral sequence. On the other hand, from an algebraic point of view, one might a priori expect not much of that information to be retained, or even expect it to be totally stripped out while running through the homological algebra machine. We will show that the answer lies somewhat in between, and both intuitions are to some extent correct: the differentials in the spectral sequence will be essentially refinements of classical ones, but with additional operations on differential forms. We recently characterized such operations in [33], and so this paper is a natural continuation of that work.

Just as generalized cohomology theories are represented by spectra, differential cohomology theories are represented by certain sheaves of smooth spectra called *differential function spectra*. The original definition of differential function spectra was due to Hopkins and Singer [41], generalized by Bunke, Nikolaus and Völkl [16], and reformulated in terms of cohesion by Schreiber [63]. The terms *smooth cohomology* and *differential cohomology* seem to be used interchangeably in some of the literature; see eg Bunke and Schick [18]. However, we will find it useful for us to provide a specific and precise usage, where the first is viewed as being more general than the second. We also present most of our ∞ -categories as combinatorial, simplicial model categories, rather than quasicategories. We believe that this way, nice objects are more easily and explicitly identifiable, which is desirable when dealing with differential cohomology. Indeed, our discussion will be very explicit, and the results will be readily utilizable.

Ordinary cohomology has smooth extension with various different realizations, including those of Cheeger and Simons [20], Gajer [30], Brylinski [11], Dupont and Ljungmann [23], Hopkins and Singer [41] and Bunke, Kreck and Schick [14]. All these realizations are in fact isomorphic [66; 18]. A description of K-theory with coefficients that combines vector bundles, connections and differential forms into a topological context was initiated by Karoubi [45]. Using Karoubi's description, Lott introduced \mathbb{R}/\mathbb{Z} -valued K-theory [49] as well as differential flat K-theory [50]. Currently, there are various geometric models of differential K-theory; see Lott [49], Bunke and Schick [17], Simons and Sullivan [66], Freed and Lott [28], and Tradler, Wilson and Zeinalian [69; 70]. As in the case of ordinary differential cohomology, these models should be equivalent. Indeed, explicit isomorphisms between various models have been demonstrated: for instance, between the differential K-theory group

of Hopkins and Singer [41] and that of Freed and Lott [28] in Klonoff [46], between Lott's \mathbb{R}/\mathbb{Z} K-theory and Lott–Freed differential K-theory in [28], between Bunke–Schick differential K-theory and Lott(–Freed) differential K-theory in Ho [40], and between Simons–Sullivan [66] and Freed–Lott [28] in Ho [39].

The group structure of differential K-theory splits into odd and even-degree parts; thus the refinement preserves the grading. However, the odd part turns out to be more delicate than the even part. In particular, while any two differential extensions of even K-theory are isomorphic by the uniqueness results in [18], odd K-theory requires extra data in order to obtain uniqueness. There are various concrete models in the odd case: using smooth maps to the unitary group [69], via loop bundles (see Hekmati, Murray, Schlegel and Vozzo [37]) and via Hilbert bundles (see Gorokhovsky and Lott [31]). Our results in both even and odd K-theory will, of course, not depend on the particular model chosen.

Suppose \mathcal{E} is a spectrum and X is a space of the homotopy type of a CW-complex. Then there is a half-plane spectral sequence (AHSS)

$$E_2^{p,q} \cong H^p(X; \mathcal{E}^q(*)),$$

converging conditionally to $\mathcal{E}^*(X)$. An immediate matter that we encounter in setting up the spectral sequence which calculates the generalized differential cohomology of a smooth manifold X is how to deal with filtrations. Classically, Maunder [52] gave two approaches to any generalized cohomology theory. The first is by filtering over the q -skeletons X^q of the topological space X , and the second by filtering over the Postnikov systems of spaces Y_q , which are the layers of an Ω -spectrum associated to the cohomology theory. Maunder also gives an isomorphism between the two approaches. While we expect this to be the case in the differential setting, the proof might require considerable work. Hence we leave this as an open problem. Maunder sets up his construction in the simplicial complex setting, which is equivalent to doing so in the CW-complex setting, as the geometric realization of a simplicial set is a CW-complex. Simplicial and Čech spectral sequences are discussed by May and Sigurdsson [53, Chapter 22].

We will prefer the filtration of the spaces/manifolds rather than of the corresponding spectra, as this will naturally bring out the geometry desired in the smooth setting. We first would like to replace a topological space with skeletal filtration by a smooth manifold and then view this manifold as a stack. Hence, in doing this, we need an analogue of a skeleton in stacks. This will be done via Čech resolution of smooth spaces, and the replacement of skeletons of a space X will be the various intersections of open sets covering the smooth manifold X .

We will use $\text{diff}(\Sigma^n \mathcal{E}, \text{ch})$ to denote the differential refinement in degree n of a cohomology theory \mathcal{E} . This was the notation used in [41] and carries more data than other notation, such as $\mathcal{E}(n)$. It also avoids possible confusion with other notations, eg when dealing with Morava K-theory $K(n)$ at chromatic level n . The axiomatic approach is very useful for characterizing a smooth cohomology theory, but one still needs the model of [41] for actually constructing examples of such smooth spectra. We will be using features of two main approaches at once, namely from [41] with $I: \text{diff}(\Sigma^n \mathcal{E}, \text{ch}) \rightarrow \underline{\mathcal{E}}$ and from [16; 63] with $I: \mathcal{E} \rightarrow \Pi \mathcal{E}$. Note that $\underline{\mathcal{E}}$ is not discrete while $\Pi \mathcal{E}$ is, but both are equivalent as smooth spectra: $\underline{\mathcal{E}} \simeq \Pi \mathcal{E}$. This essentially boils down to the fact that since $\Pi \mathcal{E}$ is locally constant, the underlying theory satisfies $\Pi \mathcal{E}^*(U) = \Pi \mathcal{E}^*(*)$ on contractible open sets. On the other hand, the homotopy invariance of the theory $\underline{\mathcal{E}}$ implies the same thing: namely, $\underline{\mathcal{E}}(U) \simeq \underline{\mathcal{E}}(*)$ for a contractible U . These relationships are discussed in further detail in [16].

We will be interested in how the differentials look in our spectral sequences. One might a priori suspect that the differentials in the refined theories should at least loosely be connected to the differentials of the underlying topological theory. We will make this precise below, and so it seems appropriate to understand the form and structure of the differentials in the topological case. To illustrate the point, we will focus on what might perhaps be the most prominent example, namely the first differential $d_3: H^*(X, K^0(*)) \rightarrow H^*(X, K^0(*))$ in complex topological K-theory $K(X)$ of a topological space X . This is given by $\text{Sq}_{\mathbb{Z}}^3$; see Atiyah and Hirzebruch [3; 4]. There are exactly two stable cohomology operations $H^*(X; \mathbb{Z}) \rightarrow H^{*+3}(X; \mathbb{Z})$, since $H^{n+3}(K(\mathbb{Z}, n)) = \mathbb{Z}/2$ for n sufficiently large. One of these is zero and the other is $\beta \circ \text{Sq}^2 \circ \rho_2$, where β is the Bockstein associated to the sequence $\mathbb{Z} \xrightarrow{\times 2} \mathbb{Z} \xrightarrow{\rho_2} \mathbb{Z}_2$ with ρ_2 denoting both the mod 2 reduction and its effect on cohomology with these as coefficients, ie $\rho_2: H^i(X; \mathbb{Z}) \rightarrow H^i(X; \mathbb{Z}/2)$.

The above class, which is a priori in mod 2 cohomology, turned out to be a class in integral cohomology. One could work at any prime [4] by noting the following; see eg Fomenko, Fuchs and Gutenmacher [26] or Hatcher [36]. For any class $x \in H^n(X; \mathbb{Z}/p)$, and with β_p the Bockstein associated with the sequence $\mathbb{Z}_p \xrightarrow{\times p} \mathbb{Z}_{p^2} \xrightarrow{\rho_p} \mathbb{Z}_p$, the element $\beta_p(x)$ is an integral class in $H^{n+1}(X; \mathbb{Z}/p)$; ie it belongs to the image of the reduction homomorphism $\rho_p: H^{n+1}(X; \mathbb{Z}) \rightarrow H^{n+1}(X; \mathbb{Z}/p)$. This can be used to prove the integrality of the class $d \in H^3(K(\mathbb{Z}/p, 2); \mathbb{Z}/p)$ as follows; see [26]. The cohomology Serre spectral sequence for the path-loop fibration $\Omega K(\mathbb{Z}, 2) \rightarrow PK(\mathbb{Z}, 3) \rightarrow K(\mathbb{Z}, 3)$ gives that $H^*(K(\mathbb{Z}, 3); \mathbb{Z}/p)$ has a single additive generator \bar{d} in dimension $\leq 2p$. Now we have a map $\beta: K(\mathbb{Z}/p, 2) \rightarrow K(\mathbb{Z}, 3)$ such that $\beta^*(\bar{d}) = d \in H^3(K(\mathbb{Z}/p, 2); \mathbb{Z}/p)$, constructed via the Serre spectral sequence of the path-loop fibration $K(\mathbb{Z}/p, 1) \rightarrow PK(\mathbb{Z}/p, 2) \rightarrow K(\mathbb{Z}/p, 2)$. The map β induces

a map of loop spaces which are also Serre fibrations:

$$\begin{array}{ccccc}
 K(\mathbb{Z}/p, 1) & \longrightarrow & PK(\mathbb{Z}/p, 2) & & PK(\mathbb{Z}, 3) \longleftarrow K(\mathbb{Z}/p, 2) \\
 & & \downarrow & & \downarrow \\
 & & K(\mathbb{Z}/p, 2) & \longrightarrow & K(\mathbb{Z}, 3)
 \end{array}$$

The induced homomorphism on the special sequences sends \bar{d} to d by the construction of β . Now we have $H^3(K(\mathbb{Z}/p, 2); \mathbb{Z}/p) = \mathbb{Z}/p$; hence d is contained in the image of the homomorphism $\rho_p: H^3(K(\mathbb{Z}/p, 2); \mathbb{Z}) \rightarrow H^3(K(\mathbb{Z}/p, 2); \mathbb{Z}/p)$. Therefore, d is an integral class. This is attractive as it makes it readily amenable to differential refinement.

Such statements, and generalizations to other primes and to other generalized cohomology theories, can be made at the level of spectra; see eg Schwede [64]. The first nontrivial k -invariant of connective complex K-theory spectrum ku is a morphism $k_2(ku) \in H^2(H\mathbb{Z}, \mathbb{Z})$, which is equal to $\beta \circ \text{Sq}^2$, where $\beta: H\mathbb{Z}/2 \rightarrow \Sigma(H\mathbb{Z})$ is the Bockstein operator associated to the extension $\mathbb{Z} \xrightarrow{\times 2} \mathbb{Z} \rightarrow \mathbb{Z}/2$, and $\text{Sq}^2_{\mathbb{Z}}$ is the pullback of the Steenrod operation $\text{Sq}^2 \in H^2(H\mathbb{Z}/2, \mathbb{Z}/2)$ along the projection morphism $\rho_2: H\mathbb{Z} \rightarrow H\mathbb{Z}/2$ given by mod 2 reduction. Since ku is a symmetric ring spectrum, then by [64, Proposition 8.8], the k -invariants are derivations. The only derivations (up to units) in the mod p Steenrod algebra \mathcal{A}_p are the Milnor primitives $Q_n \in H^{2p^n-1}(H\mathbb{Z}/p, \mathbb{Z}/p)$. At the lowest level, we have $Q_0 = \beta_p$, the mod p Bockstein, and the others are realized as k -invariants of symmetric spectra, the connective Morava K-theory spectra $k(n)$. That is, we have $Q_n = k_{2p^n-2}(k(n))$. We will consider refinements of integral lifts of these.

The classical AHSS collapses already at the first page if the generalized cohomology theory is rational. In fact, it can be shown that for any reasonably behaved spectrum like all the ones we consider, all the differentials in the AHSS are torsion, ie zero when rationalized; see [59, Corollary 7.12]. The differentials in the AHSS in the topological case are analyzed by systematically by Arlettaz [2]. Using the structure of the integral homology of the Eilenberg–Mac Lane spectra, it is proved there that for any connected space X , there are integers R_r such that $R_r d_r^{s,t} = 0$ for all $r \geq 2$ and for all s, t . Some aspects of this general feature will continue to hold in the differential setting. From a homotopy point of view, there is not much difference between the localizations at \mathbb{R} and at \mathbb{Q} . However, from a geometric point of view there is a considerable difference. Nevertheless, we will still use the term “rationalize” when we discuss localization at \mathbb{R} , as is customary in the homotopy theory literature. We stress that the distinction is needed in certain geometric settings (see Griffiths and Morgan [35]), but it will not be an issue for us in this paper.

Note that although the differential cohomology diamond, ie the diagram that characterizes such theories (see Remark 12), certainly detects torsion classes in the flat part of the theory, it does not distinguish between torsion at various primes. As a by-product, our analysis can be seen as a systematic method for addressing p -primary torsion in differential theories. In [33], we found that the Deligne–Beilinson squaring operation admits lower-degree operations refining the Steenrod squares. We have the familiar pattern

$$DD, \widehat{Sq}^1, \widehat{Sq}^2, \widehat{Sq}^3, \dots, DD^2, \dots,$$

where DD is the Dixmier–Douady class: a nontorsion differential cohomology operation. The refined squares \widehat{Sq}^{2k+1} , as the classical squares Sq^{2k} , are operations that are 2-torsion. In this paper, we get \widehat{Sq}^{2k+1} as we expect, but also differentials d_{2m} at lowest degree for every m :

$$(1-1) \quad d_{2m}: \prod_k \Omega^{2k}(M) \rightarrow H^{2m}(M; U(1)).$$

We consider this as a cohomology operation, which can be viewed as first projecting on to the homogeneous component ch_{2m} of the Chern character. A $U(1)$ -valued Čech cocycle is obtained by restricting to $2m$ -fold intersections of an open cover, pairing with an appropriate simplex of degree $2m$ and exponentiating; this will be spelled out in detail in Section 4. If indeed the form ch_{2m} arises as the curvature of a bundle, it must represent a closed form with *integral* periods. The differential d_{2m} can therefore be understood as the obstruction to this condition. Similar results hold for the odd part, ie for differentially refined K^1 -theory, where the refined Steenrod square takes the same form as in differential K^0 -theory, while the differentials arising from forms — the analogues of those in (1-1) — are now of odd degrees.

The paper is organized as follows. In Section 2, we start by carefully setting up the background in smooth and differential cohomology, preparing the scene for our constructions. In particular, in Section 2.1, we adapt abstract general results on stacks (or simplicial sheaves) to our context and spell out specific definitions and constructions that will be useful for us in later sections; more general and comprehensive accounts can be found in Jardine [43], Lurie [51] and Schreiber [63]. Then in Section 2.2, we take the approach to differential cohomology that allows for a direct generalization. Our main constructions will be in Section 3. In particular, in Section 3.1, we provide the filtration via Čech resolutions; then we construct the AHSS for smooth spectra in Section 3.2 and compare to the AHSS of the underlying topological theory. This refinement will depend on whether the degree is positive, negative or zero. Then we explore the compatibility of the differentials with the product structure in Section 3.3.

Having given the main construction, our main applications of the general spectral sequence to various differential cohomology theories will be presented in Section 4.

The construction is general enough to apply to any structured cohomology theory whose coefficients are known. We will explicitly emphasize three main examples: ordinary differential cohomology, differential K-theory and a differential version of integral Morava K-theory that we introduce. As a test of our method, in Section 4.1, we recover the usual hypercohomology spectral sequence for the Deligne complex (see [11] and Esnault and Viehweg [24, Appendix]), and we do so for manifolds, then products of these, and then more generally for smooth fiber bundles. Then the AHSS for K-theory is generalized in Section 4.2 to differential K-theory, where the differentials involve refinements of Steenrod squares, in the sense of [33], as well as operations on forms, as indicated above around expression (1-1). We also show that the odd case, ie smooth extension of K^1 , leads to a similar construction, but with the differentials now involving odd forms. Then in Section 4.3, we first introduce a refinement of the integral form of Morava K-theory, discussed in Kriz and Sati [47], Sati [60] and Sati and Westerland [62], and then characterize the corresponding differentials, which turn out to have a similar pattern as in K-theory, where the operation that gets refined is the Milnor primitive Q_n encountered above. We end with an application to an example from M-theory and string theory.

Notation We have the following morphism that we will use repeatedly throughout. Denote by $\rho_p: \mathbb{Z} \rightarrow \mathbb{Z}/p$ the mod p reduction on coefficients with corresponding morphism using the same notation on the cohomology groups with these as coefficients. We will denote by β , β_p and $\tilde{\beta}$ the Bockstein homomorphisms associated with the coefficient sequences

$$\begin{aligned} 0 \rightarrow \mathbb{Z} \rightarrow \mathbb{R} \xrightarrow{\exp} U(1) \rightarrow 0, \\ 0 \rightarrow \mathbb{Z}/p \xrightarrow{\times p} \mathbb{Z}/p^2 \xrightarrow{\rho_p} \mathbb{Z}/p \rightarrow 0, \\ 0 \rightarrow \mathbb{Z} \xrightarrow{\times p} \mathbb{Z} \xrightarrow{\rho_p} \mathbb{Z}/p \rightarrow 0, \end{aligned}$$

respectively. We will let $\Gamma_2: \mathbb{Z}/2 \hookrightarrow U(1)$ denote the representation as the square roots of unity, and also the induced map $\Gamma_2: H^n(-; \mathbb{Z}/2) \rightarrow H^n(-; U(1))$ on cohomology. We will also use more refined Bockstein homomorphisms associated with spectra, and these will be defined as we need them.

2 Smooth cohomology

2.1 Smooth cohomology and the stable category of smooth stacks

In this section, we adapt abstract general results on stacks (or simplicial sheaves) to our context and spell out specific definitions and constructions that will be useful for

us in later sections. The interested reader can find more general and comprehensive accounts in [43; 51; 63]. For the reader who is more interested in the applications to differential cohomology theories, this section can be skipped. However, we would like to emphasize that although the language used in this section is rather abstract, the generality gained from this formalism is far reaching and allows this machinery to be used for a wide variety theories beyond just differential cohomology theories.

Essentially, the axioms characterizing a smooth cohomology theory are not much different from the axioms characterizing usual cohomology theories. The big difference is where the theory takes place. More precisely, we want to consider homotopical functors on the category of pointed *smooth stacks* $\text{Sh}_\infty(\text{CartSp})_+$ with CartSp the category of Cartesian spaces, rather than the category of pointed topological spaces Top_+ . Let $\mathcal{A}b_{\text{gr}}$ be the category of graded abelian groups.

Definition 1 (smooth cohomology) Let $\mathcal{E}^*: \text{Sh}_\infty(\text{CartSp})_+^{\text{op}} \rightarrow \mathcal{A}b_{\text{gr}}$ be a functor satisfying the following axioms:

- (1) **Invariance** \mathcal{E}^* sends equivalences to isomorphisms.
- (2) **Additivity** For small coproducts (ie ones forming sets) of pointed stacks, $\bigvee_\alpha X_\alpha$, we have

$$\mathcal{E}^*(\bigvee_\alpha X_\alpha) = \prod_\alpha \mathcal{E}^*(X_\alpha).$$

- (3) **Mayer–Vietoris** For any homotopy pushout of pointed stacks

$$\begin{array}{ccc} Z & \longrightarrow & Y \\ \downarrow & & \downarrow \\ X & \longrightarrow & X \cup_Z Y \end{array}$$

the induced sequence

$$\mathcal{E}^*(X \cup_Z Y) \rightarrow \mathcal{E}^*(X) \oplus \mathcal{E}^*(Y) \rightarrow \mathcal{E}^*(Z)$$

is exact.

- (4) **Suspension** For any stack X , there is an isomorphism $\mathcal{E}^{n+1}(\Sigma X) \simeq \mathcal{E}^n(X)$.

Then we call \mathcal{E}^* a *smooth cohomology theory*.

Remark 2 Note that the Mayer–Vietoris axiom implies the usual Mayer–Vietoris sequence. Indeed, let M be a manifold and let V be a local chart of M . Let U be an

open set such that $\{U, V\}$ is a cover of M . Then the strict pushout

$$\begin{array}{ccc} U \cap V & \longrightarrow & V \\ \downarrow & & \downarrow \\ U & \longrightarrow & U \cup V \end{array}$$

is actually a homotopy pushout. We can equivalently write this diagram as a homotopy coequalizer

$$U \cap V \rightrightarrows U \sqcup V \rightarrow U \cup V$$

in which the homotopy cofiber of the second map can be identified with $\Sigma U \cap V$. By iterating this argument and applying \mathcal{E}^* to the resulting diagram, one obtains the long exact sequence

$$\dots \rightarrow \mathcal{E}^*(U \cap V) \rightarrow \mathcal{E}^*(M) \rightarrow \mathcal{E}^*(U) \oplus \mathcal{E}^*(V) \rightarrow \mathcal{E}^{*+1}(U \cap V) \rightarrow \dots,$$

which is the familiar Mayer–Vietoris sequence.

The above axioms can be taken as a generalization of the Eilenberg–Steenrod axioms (see [1; 38]), where the Mayer–Vietoris axiom subsumes both the excision axiom and the long exact sequence axiom. It is interesting to note that the axioms *do not* require homotopy invariance. Namely, if two manifolds M and N are *homotopic*, they may fail to be equivalent as stacks. In fact, an equivalence of stacks requires, in particular, that for every sheaf F (embedded as a stack), we have an isomorphism

$$F(N) \simeq \pi_0 \text{Map}(N, F) \simeq \pi_0 \text{Map}(M, F) \simeq F(M).$$

In particular, we can take the sheaf of smooth \mathbb{R} -valued functions on a manifold. Then if every homotopy equivalence $f: M \rightarrow N$ induced an equivalence of stacks, we would have an induced isomorphism

$$f^*: C^\infty(N; \mathbb{R}) \rightarrow C^\infty(M; \mathbb{R}).$$

Taking $N = *$ and $M = \mathbb{R}^n$ immediately gives a contradiction. On the other hand, every equivalence of stacks does produce a weak homotopy equivalence of geometric realizations. To see this, simply note that the geometric realization functor

$$\Pi: \text{Sh}_\infty(\text{CartSp}) \rightarrow s\text{Set},$$

being a Quillen functor, has a derived functor by Ken Brown’s lemma [10]. It therefore preserves weak equivalences between fibrant objects. But these objects are exactly those that satisfy descent, namely stacks (eg manifolds) [63; 22].

Remark 3 Given a smooth cohomology theory \mathcal{E}^* , we always get a presheaf of graded abelian groups on the site $\mathcal{C}art\mathcal{S}p$ by precomposing with the Yoneda embedding:

$$\mathcal{E}^*: \mathcal{C}art\mathcal{S}p \xrightarrow{Y} \mathit{Sh}(\mathcal{C}art\mathcal{S}p) \xrightarrow{sk_0} \mathit{Sh}_\infty(\mathcal{C}art\mathcal{S}p) \xrightarrow{+} \mathit{Sh}_\infty(\mathcal{C}art\mathcal{S}p)_+ \xrightarrow{\mathcal{E}^*} \mathcal{A}b_{gr},$$

where sk_0 embeds a sheaf as a discrete simplicial sheaf. We will use this fact later in the construction of the spectral sequence in Theorem 25.

Just as all cohomology theories are representable by Ω -spectra, via Brown representability, all smooth cohomology theories are representable by smooth spectra. This follows from the version of Brown representability formulated by Jardine in [43] applied to the stable homotopy category of smooth stacks. We will quickly review the basic properties of this category (see [51; 44]) to establish where our objects of interest live.

We first recall some operations on stacks that are counterparts to standard operations on topological spaces. Let X and Y be two pointed stacks.

- (i) The wedge product $X \vee Y$ is defined via the pushout diagram:

$$\begin{array}{ccc} Y & \longrightarrow & Y \vee X \\ \uparrow & & \uparrow \\ * & \longrightarrow & X \end{array}$$

- (ii) The smash product $X \wedge Y$ is defined as the quotient $X \wedge Y := X \times Y / X \vee Y$ of the Cartesian product by the wedge product.

- (iii) The suspension ΣX is defined via the homotopy pushout diagram:

$$\begin{array}{ccc} X & \longrightarrow & * \\ \downarrow & & \downarrow \\ * & \longrightarrow & \Sigma X \end{array}$$

- (iv) The looping, ie loop space, ΩX is defined via the homotopy pullback:

$$\begin{array}{ccc} \Omega X & \longrightarrow & * \\ \downarrow & & \downarrow \\ * & \longrightarrow & X \end{array}$$

Definition 4 We define the *stabilization* $\mathit{Stab}(\mathit{Sh}_\infty(\mathcal{C}art\mathcal{S}p)_+)$ of smooth pointed stacks to be the following category:

- The objects of $\text{Stab}(\text{Sh}_\infty(\text{CartSp})_+)$ are sequences of pointed stacks

$$\{\mathcal{E}_n\} \subset \text{Sh}_\infty(\text{CartSp})_+, \quad n \in \mathbb{Z},$$

equipped with maps $\sigma_n: \Sigma \mathcal{E}_n \rightarrow \mathcal{E}_{n+1}$.

- The morphisms between \mathcal{E} and \mathcal{F} are defined to be the levelwise morphisms $\mathcal{E}_n \rightarrow \mathcal{F}_n$ commuting with the σ_n .

This category carries a stable model structure given by first taking the projective model structure on sequences of stacks and then performing Bousfield localization with respect to stable weak equivalences in the usual way. This process is described in detail in [43; 51; 44], and we summarize the relevant results found there. The category $\text{Stab}(\text{Sh}_\infty(\text{CartSp})_+)$ admits a stable, closed, simplicial model structure with the following properties:

- The weak equivalences are *stable* weak equivalences. That is, a morphism of smooth spectra $f: \mathcal{E}_\bullet \rightarrow \mathcal{F}_\bullet$ is a weak equivalence if and only if it induces a weak equivalence

$$Q(f): \lim_{i \rightarrow \infty} \Omega^i \mathcal{E}_{n+i} \rightarrow \lim_{j \rightarrow \infty} \Omega^j \mathcal{F}_{n+j}.$$

- The fibrant objects are precisely the smooth Ω -Spectra, that is, the sequence of stacks X_\bullet whose structure maps

$$\sigma_n: \Sigma \mathcal{E}_n \rightarrow \mathcal{E}_{n+1}$$

induce equivalences $\mathcal{E}_n \xrightarrow{\sim} \Omega \mathcal{E}_{n+1}$.

Remark 5 We will refer to the stable model category $\text{Stab}(\text{Sh}_\infty(\text{CartSp})_+)$ as the category of *smooth spectra* and denote it by

$$\text{Sh}_\infty(\text{CartSp}; \text{Sp}) := \text{Stab}(\text{Sh}_\infty(\text{CartSp})_+).$$

Example 6 Let $M \in \text{Sh}_\infty(\text{CartSp})_+$ be a manifold, viewed a stack and equipped with a basepoint. We can define the smooth spectrum $\Sigma^\infty M$ in the usual way, as the sequence of suspensions of the manifold M . Given a smooth Ω -spectrum \mathcal{E} , we can define a smooth cohomology theory \mathcal{E}^* , by setting

$$\mathcal{E}^q(M) \simeq \pi_0 \text{Map}(\Sigma^{-q} \Sigma^\infty M, \mathcal{E}).$$

Differential cohomology theories are examples of the theories introduced above, although it may not be immediately apparent where the differential cohomology “diamond” diagram [66] fits into this context. In fact, it was observed by Bunke, Nikolaus and Völkl in [16] that the diamond provides a further characterization of *all* smooth cohomology theories in terms of refinement of topological theories. This characterization happens

in addition to the Brown representability described above, and it happens only when the category of stacks exhibits so-called *cohesion*. We now review the properties of the cohesive structure on smooth stacks [63] that we need, along with the characterization of smooth cohomology theories described in [16]. It is shown in [63] that the category $\text{Sh}_\infty(\text{CartSp})$ admits a quadruple ∞ -categorical adjunction $(\Pi \dashv \text{disc} \dashv \Gamma \dashv \text{codisc})$

$$(2-1) \quad \text{Sh}_\infty(\text{CartSp}) \begin{array}{c} \xrightarrow{\Pi} \\ \xleftarrow{\text{disc}} \\ \xleftarrow{\Gamma} \\ \xrightarrow{\text{codisc}} \end{array} s\text{Set},$$

where Π preserves finite ∞ -limits, and the functors disc and codisc are fully faithful.

One implication of this is that $s\text{Set}$ embeds into $\text{Sh}_\infty(\text{CartSp})$ as an ∞ -subcategory in two different ways: one reflective, the other reflective and coreflective. From the reflectors, one can produce two monads and one comonad defined as follows:

$$\Pi := \Pi \circ \text{disc}, \quad \flat := \text{disc} \circ \Gamma, \quad \sharp := \text{codisc} \circ \Gamma.$$

These monads fit into a triple ∞ -adjunction $(\Pi \dashv \flat \dashv \sharp)$ which is called a *cohesive adjunction*.

Remark 7 Each monad in the cohesive adjunction picks out a different part of the nature of a smooth stack. This nature is perhaps best exemplified by how the adjoints behave on smooth manifolds (viewed as stacks). More precisely, if M is a smooth manifold, then for instance:

- (i) The comonad \flat takes the underlying set of points of the manifold and then embeds this set back into stacks as a discrete object. This functor therefore misses the smooth structure of the manifold and treats it instead as a discrete object.
- (ii) The monad Π essentially takes the singular nerve of the manifold using *smooth* paths and higher smooth simplices on the manifold. It therefore retains the geometry of the manifold and “knows” that the points of the manifold ought to be connected together in a smooth way.

The following observation on lifting from simplicial sets to spectra is known [63, Proposition 4.1.9], but we supply a proof for completeness.

Proposition 8 *The ∞ -adjunction (2-1) lifts to an ∞ -adjunction*

$$\text{Sh}_\infty(\text{CartSp}; \mathcal{S}\text{p}) \begin{array}{c} \xrightarrow{\Pi^s} \\ \xleftarrow{\text{disc}^s} \\ \xleftarrow{\Gamma^s} \\ \xrightarrow{\text{codisc}^s} \end{array} \mathcal{S}\text{p}$$

on the stable ∞ -category of smooth spectra. Moreover, the adjoints satisfy the same condition as the ∞ -adjunction (2-1) does.

Proof The category of smooth stacks is presented by the combinatorial simplicial model category

$$\text{Sh}_\infty(\text{CartSp}) = [\text{CartSp}, s\text{Set}]_{\text{loc,proj}},$$

where *loc* denotes the Bousfield localized model structure at the maps out of Čech nerves. The quadruple adjunction is presented by Quillen adjoints $(\Pi \dashv \text{disc} \dashv \Gamma \dashv \text{codisc})$ [63]. We need to show that this adjunction holds on the stable model category of smooth spectra. The adjunction immediately gives an underlying categorical adjunction by simply applying the functors degreewise. In the projective model structure, the right adjoints are Quillen by definition, and the closed model axioms imply that the left adjoints are also Quillen.

Now the functors (in the global model structure on $\mathcal{S}p$) *disc* and *codisc* both preserve homotopy limits. Hence for a local weak equivalence $f: \mathcal{E} \rightarrow \mathcal{F}$ of spectra, we have

$$\begin{aligned} \lim_{i \rightarrow \infty} \Omega^i \text{disc}(\mathcal{E})_{n+i} &\simeq \text{disc}\left(\lim_{i \rightarrow \infty} \Omega^i \mathcal{F}_{n+i}\right) \\ &\simeq \text{disc}\left(\lim_{j \rightarrow \infty} \Omega^j \mathcal{F}_{n+j}\right) \\ &\simeq \lim_{j \rightarrow \infty} \Omega^j \text{disc}(\mathcal{F})_{n+j}, \end{aligned}$$

and *disc*(*f*) induces a weak equivalence $Q(\text{disc}(f))$. Hence *disc*(*f*) is a weak equivalence. In the same way, *codisc* preserves local weak equivalences. It follows by the basic properties of Bousfield localization that *disc* and *codisc* are right Quillen adjoints. Again, by the axioms of a closed model category, it follows that the entire adjunction holds as a Quillen adjunction of stable model categories. □

Remark 9 The proof of the previous proposition implies that both *disc* and *codisc* preserve Ω -spectra. However, Π and Γ need not take Ω -spectra to Ω -spectra. This problem can be remedied by taking Π^s (or Γ^s) to be the composition $R \circ \Pi$ (or $R \circ \Gamma$), where *R* is the fibrant replacement in spectra. Since *R* defines a left ∞ -adjoint to the inclusion of fibrant objects (and preserves finite ∞ -limits), we will still have an adjunction at the level of ∞ -categories (although this is not presented by a Quillen adjunction).

As in the case of smooth stacks, the quadruple adjunction in Proposition 8 produces adjoint monads $(\Pi^s \dashv \flat^s \dashv \sharp^s)$ exhibiting *stable* cohesion. The main observation in [16], recast in the cohesive setting in [63], is the following. Let $j: \flat^s \rightarrow \text{id}$ be the

count of the comonad \flat^s , and let $I: \text{id} \rightarrow \Pi^s$ be the unit of the monad Π^s . Let $\mathcal{E} \in \text{Sh}_\infty(\text{CartSp}; \mathcal{S}p)$ be a smooth spectrum. Then \mathcal{E} sits inside a hexagon diagram

$$(2-2) \quad \begin{array}{ccccc} & & \text{fib}(\eta)(\mathcal{E}) & \longrightarrow & \text{cofib}(\epsilon)(\mathcal{E}) \\ & \nearrow & & \searrow & \\ & \Sigma^{-1} \Pi^s \text{cofib}(\epsilon)(\mathcal{E}) & & & \\ & \searrow & & \nearrow & \\ & & \mathcal{E} & & \Pi^s \text{cofib}(\epsilon)(\mathcal{E}) \\ & & \nearrow \scriptstyle j & \searrow \scriptstyle I & \\ & & \flat^s \mathcal{E} & \longrightarrow & \Pi^s \mathcal{E} \end{array}$$

where the diagonals are fiber sequences (by definition), the top and bottom sequences are fiber sequences, and the two squares in the hexagon are homotopy Cartesian; ie both are homotopy pullback squares and hence homotopy pushouts (via the equivalence of the two in the stable setting). The latter property is key because it is a homotopy Cartesian square, as on the right of the hexagon, which Hopkins and Singer [41] took as the definition of differential cohomology (for a specific choice of the object of differential forms). Bunke, Nikolaus and Völkl [16] observed that by the hexagon, every smooth spectrum satisfies this kind of Hopkins–Singer definition, if one just allows more general objects of differential forms, which is the object $\text{cofib}(\epsilon)(\mathcal{E})$ in our notation above.

It often happens in practice that the smooth spectra $\text{fib}(\eta)(\mathcal{E})$ and $\text{cofib}(\epsilon)(\mathcal{E})$ contain no information away from degree 0. In particular, it often happens that for $n > 0$,

$$(2-3) \quad \pi_n \text{Map}(M, \text{cofib}(\epsilon)(\mathcal{E})) \simeq 0,$$

$$(2-4) \quad \pi_{-n} \text{Map}(M, \text{fib}(\eta)(\mathcal{E})) \simeq 0.$$

In this case, the \mathcal{E} –cohomology of a manifold can be calculated as either the flat cohomology or the underlying topological cohomology in all degrees but 0. This is summarized as the following result.

Proposition 10 *Let \mathcal{E} be a smooth spectrum such that (2-3) and (2-4) are satisfied. Then the \mathcal{E} –theory of a manifold M is given by*

$$\mathcal{E}^n(M) := \begin{cases} (\Pi^s \mathcal{E})^n(M), & n > 0, \\ (\flat^s \mathcal{E})^n(M), & n < 0, \end{cases}$$

where $\mathcal{E}(M)$ is already characterized in degree 0 by the diamond (2-2).

Proof Since the diagonals of the diamond are fiber sequences, they induce long exact sequences in cohomology. Let n be a positive integer. The sequence

$$\flat^s \mathcal{E} \rightarrow \mathcal{E} \rightarrow \text{cofib}(\epsilon)(\mathcal{E})$$

gives the section of the long sequence

$$\pi_{n+1} \text{Map}(M, \text{cofib}(\epsilon)(\mathcal{E})) \rightarrow b^s \mathcal{E}^{-n}(M) \rightarrow \mathcal{E}^{-n}(M) \rightarrow \pi_n \text{Map}(M, \text{cofib}(\epsilon)(\mathcal{E})).$$

By assumption, the leftmost and rightmost groups are 0. Thus we have an isomorphism

$$(b^s \mathcal{E})^{-n}(M) \simeq \mathcal{E}^{-n}(M).$$

Similarly, the sequence

$$\text{fib}(\eta)(\mathcal{E}) \rightarrow \mathcal{E} \rightarrow \Pi^s \mathcal{E}$$

gives the long sequence

$$\pi_{-n} \text{Map}(M, \text{fib}(\eta)(\mathcal{E})) \rightarrow \mathcal{E}^n(M) \rightarrow (\Pi^s \mathcal{E})^n(M) \rightarrow \pi_{-n-1} \text{Map}(M, \text{fib}(\eta)(\mathcal{E})),$$

and again we get the desired isomorphism. □

2.2 Differential cohomology and differential function spectra

The main applications we have in mind, as we indicated in the introduction, concern *differential cohomology theories*. In this section, we review some of the concepts established in [13; 16; 63] (which generalize [66]), adapted to our context.

Definition 11 Let \mathcal{E}^* be a cohomology theory. A *differential refinement* $\widehat{\mathcal{E}}^*$ of \mathcal{E}^* consists of the following data:

- (1) a functor $\widehat{\mathcal{E}}^*: \text{Sh}_\infty(\text{CartSp}_+)^{\text{op}} \rightarrow \text{Ab}_{\text{gr}}$;
- (2) three natural transformations:
 - (a) **Integration** $I: \widehat{\mathcal{E}}^* \rightarrow \mathcal{E}^*$;
 - (b) **Curvature** $R: \widehat{\mathcal{E}}^* \rightarrow Z_*(\Omega^* \otimes \mathcal{E}^*(*))$;
 - (c) **Secondary Chern character** $a: \Omega^* \otimes \mathcal{E}^*(*)[1] / \text{im}(d) \rightarrow \widehat{\mathcal{E}}^*$;

such that the following axioms hold:

- o **Chern–Weil** We have a commutative diagram

$$\begin{array}{ccc} \widehat{\mathcal{E}}^* & \xrightarrow{R} & Z_*(\Omega^* \otimes \mathcal{E}^*(*)) \\ \downarrow I & & \downarrow q \\ \mathcal{E}^* & \xrightarrow{\text{ch}} & H_*(\Omega^* \otimes \mathcal{E}^*(*)) \end{array}$$

where *ch* is the Chern character map.

- **Secondary Chern–Weil** We have a commutative diagram

$$\begin{array}{ccc}
 \Omega^* \otimes \mathcal{E}^*(*)[1]/\text{im}(d) & \xrightarrow{d} & Z_*(\Omega^* \otimes \mathcal{E}^*(*)) \\
 & \searrow a & \nearrow R \\
 & & \widehat{\mathcal{E}}^*
 \end{array}$$

and an exact sequence

$$\dots \rightarrow \mathcal{E}^*[1] \rightarrow \Omega^* \otimes \mathcal{E}^*(*)[1]/\text{im}(d) \rightarrow \widehat{\mathcal{E}}^* \rightarrow \mathcal{E}^* \rightarrow \dots$$

Note that in the Chern–Weil axiom above, $H_*(\Omega^* \otimes \mathcal{E}^*(*))$ appears as the codomain of the Chern character. As explained in [16], this becomes a locally constant stack equivalent to just the locally constant stack on the rationalization of \mathcal{E}^* ; ie ch is equivalent to $\text{ch}: \mathcal{E}^* \rightarrow \mathcal{E}^* \wedge H\mathbb{R}$ (or $M\mathbb{R}$).

Remark 12 The above characterization can ultimately be summarized by saying that differential cohomology fits into an exact diamond

$$\begin{array}{ccccc}
 & & \Omega^* \otimes \mathcal{E}^*(*)[1]/\text{im}(d) & \xrightarrow{d} & Z_*(\Omega^* \otimes \mathcal{E}^*(*)) \\
 & \nearrow & & \searrow a & \nearrow R \\
 \mathcal{E}^{*-1} \otimes \mathbb{R} & & & & \\
 & \searrow & & & \searrow \\
 & & \mathcal{E}_{\mathbb{R}/\mathbb{Z}}^{*-1} & \xrightarrow{\beta_{\mathcal{E}}} & \mathcal{E}^* \\
 & \nearrow & & \searrow I & \nearrow \text{ch} \\
 & & \widehat{\mathcal{E}}^* & & \mathcal{E}^* \otimes \mathbb{R}
 \end{array}$$

where the diagonal, top and bottom sequences are all part of long exact sequences. The bottom sequence is obtained by observing that the cofiber of the rationalization map is an $MU(1)$ (Eilenberg–Moore spectrum), where we identify \mathbb{R}/\mathbb{Z} with $U(1)$ throughout. That is, we have a cofiber sequence involving the unit map from the sphere spectrum $\mathbb{S} = M\mathbb{Z}$:

$$\mathbb{S} \rightarrow M\mathbb{R} \rightarrow MU(1).$$

Smashing on the left with the theory \mathcal{E} , we obtain a “Bockstein sequence”

$$\mathcal{E} \rightarrow \mathcal{E} \wedge M\mathbb{R} \rightarrow \mathcal{E} \wedge MU(1) \xrightarrow{\beta_{\mathcal{E}}} \Sigma\mathcal{E}.$$

We define the flat theory as

$$\mathcal{E}_{U(1)} := \mathcal{E} \wedge MU(1)$$

and the rational theory as

$$\mathcal{E}_{\mathbb{R}} := \mathcal{E} \wedge M\mathbb{R}.$$

Remark 13 Differential cohomology theories are special cases of smooth cohomology theories, while differential function spectra are special cases of smooth spectra. Thus, this section can be viewed as describing a special case of the previous section.

Since differential cohomology theories will arise as certain homotopy pullbacks (in Definition 17 below), we will first need to establish the components of the pullback. We begin with the following lemma that can be found in [13, Lemma 6.10], which explains how we can transition from a topological cohomology theory to a smooth one, in a process whose direction is opposite to that of the map I .

Lemma 14 *Let \mathcal{E} be a spectrum and define the smooth presheaf of spectra $\underline{\mathcal{E}}$ via the assignments*

$$\text{objects: } U \mapsto \text{Map}(\Sigma^\infty U, \mathcal{E}),$$

$$\text{morphisms: } (f: U \rightarrow V) \mapsto (f^*: \text{Map}(\Sigma^\infty V, \mathcal{E}) \rightarrow \text{Map}(\Sigma^\infty U, \mathcal{E})).$$

Then $\underline{\mathcal{E}}$ satisfies descent.

Proof Let $C^\bullet(\{U_\alpha\})$ denote the Čech nerve of a good open cover $\{U_\alpha\}$ of some manifold M . The Yoneda lemma and basic properties of the mapping space functor imply that we have the sequence of equivalences

$$\begin{aligned} \underline{\mathcal{E}}(M) &:= \text{Map}(\Sigma^\infty M, \mathcal{E}) \\ &\simeq \text{Map}(\Sigma^\infty \text{hocolim}_{\Delta^{\text{op}}} C^\bullet(\{U_\alpha\}), \mathcal{E}) \\ &\simeq \text{Map}(\text{hocolim}_{\Delta^{\text{op}}} \Sigma^\infty C^\bullet(\{U_\alpha\}), \mathcal{E}) \\ &\simeq \text{holim}_{\Delta^{\text{op}}} \text{Map}(\Sigma^\infty C^\bullet(\{U_\alpha\}), \mathcal{E}) \\ &\simeq \text{holim} \left\{ \cdots \begin{array}{c} \xleftarrow{\quad} \\ \xleftarrow{\quad} \\ \xleftarrow{\quad} \end{array} \prod_{\alpha\beta\gamma} \text{Map}(\Sigma^\infty U_{\alpha\beta\gamma}, \mathcal{E}) \begin{array}{c} \xleftarrow{\quad} \\ \xleftarrow{\quad} \\ \xleftarrow{\quad} \end{array} \prod_{\alpha\beta} \text{Map}(\Sigma^\infty U_{\alpha\beta}, \mathcal{E}) \\ \xleftarrow{\quad} \prod_{\alpha} \text{Map}(\Sigma^\infty U_{\alpha}, \mathcal{E}) \right\} \\ &\simeq \text{holim} \left\{ \cdots \begin{array}{c} \xleftarrow{\quad} \\ \xleftarrow{\quad} \\ \xleftarrow{\quad} \end{array} \prod_{\alpha\beta\gamma} \underline{\mathcal{E}}(U_{\alpha\beta\gamma}) \begin{array}{c} \xleftarrow{\quad} \\ \xleftarrow{\quad} \\ \xleftarrow{\quad} \end{array} \prod_{\alpha\beta} \underline{\mathcal{E}}(U_{\alpha\beta}) \xleftarrow{\quad} \prod_{\alpha} \underline{\mathcal{E}}(U_{\alpha}) \right\}, \end{aligned}$$

and so $\underline{\mathcal{E}}$ satisfies descent. □

The other components of the pullback we want to establish are presented by sheaves of chain complexes. There is a general functorial construction by which one can turn an unbounded chain complex into a spectrum, which we now describe; see [65] for details. This functor is called the *Eilenberg–Mac Lane* functor

$$(2-5) \quad H: \text{Ch} \rightarrow \text{Sp},$$

and acts on objects as follows. Let C_\bullet be an unbounded chain complex, and let Z_n denote the subgroup of cycles in degree n . The functor H takes C_\bullet and forms the sequence $C_\bullet(\bullet)$ of truncated bounded chain complexes:

$$\begin{aligned} C_\bullet(0) &= (\cdots \rightarrow C_n \rightarrow C_{n-1} \rightarrow \cdots \rightarrow C_1 \rightarrow Z_0), \\ C_\bullet(1) &= (\cdots \rightarrow C_n \rightarrow C_{n-1} \rightarrow \cdots \rightarrow C_0 \rightarrow Z_{-1}), \\ C_\bullet(2) &= (\cdots \rightarrow C_n \rightarrow C_{n-1} \rightarrow \cdots \rightarrow C_{-1} \rightarrow Z_{-2}), \\ &\vdots \\ C_\bullet(k) &= (\cdots \rightarrow C_n \rightarrow C_{n-1} \rightarrow \cdots \rightarrow C_{-k+1} \rightarrow Z_{-k}), \\ &\vdots \end{aligned}$$

The reason for the group of cycles appearing in degree 0 comes from using the *right* adjoint to the inclusion $i: \mathcal{C}h^+ \rightarrow \mathcal{C}h$ (as opposed to the left). The left adjoint simply truncates the complex in degree 0, while the right adjoint truncates and then takes only the cycles in degree 0.

Continuing with our discussion, at each level in the sequence, H applies the Dold–Kan functor $DK: \mathcal{C}h^+ \rightarrow s\mathcal{S}et$ to the bounded chain complex in that degree. This gives a sequence $DK(C_\bullet(\bullet))$ of spaces

$$DK(C_\bullet(0)), DK(C_\bullet(1)), DK(C_\bullet(2)), \dots, DK(C_\bullet(k)), \dots$$

Since DK preserves looping (being a right Quillen adjoint) and equivalences (being a Quillen equivalence of model categories), we get induced equivalences

$$\sigma_k: DK(C_\bullet(k)) \rightarrow \Omega DK(C_\bullet(k - 1)),$$

which turns $DK(C_\bullet(\bullet))$ into a spectrum.

Example 15 Consider the unbounded chain complex $\mathbb{Z}[0]$, with \mathbb{Z} concentrated in degree 0. Then

$$H(\mathbb{Z}[0]) \simeq H\mathbb{Z},$$

where the right-hand side denotes the Eilenberg–Mac Lane spectrum.

Example 16 Fix a manifold M and consider the de Rham complex

$$\Omega^* := (\cdots \rightarrow 0 \rightarrow \Omega^0(M) \rightarrow \Omega^1(M) \rightarrow \cdots \rightarrow \Omega^k(M) \rightarrow \cdots),$$

where the nonzero terms are concentrated in negative degrees. Then H takes Ω^* to the spectrum:

$$H(\Omega^*(M)) = \begin{cases} \text{DK}(\cdots \rightarrow 0 \rightarrow \Omega_{\text{cl}}^0(M)), \\ \text{DK}(\cdots \rightarrow 0 \rightarrow \Omega^0(M) \rightarrow \Omega_{\text{cl}}^1(M) \rightarrow \cdots), \\ \text{DK}(\cdots \rightarrow 0 \rightarrow \Omega^0(M) \rightarrow \Omega^1(M) \rightarrow \Omega_{\text{cl}}^2(M) \rightarrow \cdots), \\ \vdots \\ \text{DK}(\cdots \rightarrow 0 \rightarrow \Omega^0(M) \rightarrow \Omega^1(M) \rightarrow \cdots \rightarrow \Omega_{\text{cl}}^k(M) \rightarrow \cdots), \\ \vdots \end{cases}$$

By the basic properties of the Dold–Kan functor, the stable homotopy groups of this spectrum are computed as

$$\begin{aligned} \pi_n^s H(\Omega^*(M)) &\simeq \lim_{k \rightarrow \infty} \pi_{k+n} \text{DK}(\cdots \rightarrow 0 \rightarrow \Omega^0(M) \rightarrow \Omega^1(M) \rightarrow \cdots \rightarrow \Omega_{\text{cl}}^k(M)) \\ &\simeq \lim_{k \rightarrow \infty} H_{k+n}(\cdots \rightarrow 0 \rightarrow \Omega^0(M) \rightarrow \Omega^1(M) \rightarrow \cdots \rightarrow \Omega_{\text{cl}}^k(M)). \end{aligned}$$

For $n > 0$, these groups are 0. For $n \leq 0$, they are the n^{th} de Rham groups $H_{\text{dR}}^n(M)$.

Now the functor H in (2-5) prolongs to a functor on prestacks

$$H: [\text{CartSp}, \text{Ch}] \rightarrow [\text{CartSp}, \text{Sp}].$$

In fact, using the properties of the Dold–Kan correspondence, it is fairly straightforward to show that this functor preserves local weak equivalences [10]. We therefore get a functor of smooth stacks

$$(2-6) \quad H: \text{Sh}_\infty(\text{CartSp}; \text{Ch}) \rightarrow \text{Sh}_\infty(\text{CartSp}; \text{Sp}).$$

Recall that for an Ω –spectrum \mathcal{E} , we always have a rational equivalence

$$r: \mathcal{E} \wedge M\mathbb{R} \rightarrow H(\pi_*(\mathcal{E}) \otimes \mathbb{R}),$$

where $M\mathbb{R}$ denotes an Eilenberg–Moore spectrum. Now, since we are working over the site of Cartesian spaces, the Poincaré lemma implies that the inclusion $j: \mathbb{R}[0] \rightarrow \Omega^*$ induces an equivalence

$$\text{id} \otimes j: \pi_*(\mathcal{E}) \otimes \mathbb{R}[0] \rightarrow \pi_*(\mathcal{E}) \otimes \Omega^*,$$

where $\pi_*(\mathcal{E}) = \mathcal{E}(\ast)$ (which follows from suspension).

Definition 17 Let \mathcal{E} be a spectrum. For an unbounded chain complex C_\bullet , let $\tau_{\leq 0}C_\bullet$ denote the truncated complex

$$\tau_{\leq 0}C_\bullet = (\cdots \rightarrow 0 \rightarrow C_0 \rightarrow C_{-1} \rightarrow \cdots \rightarrow C_{-n} \rightarrow \cdots).$$

A differential function spectrum $\text{diff}(\mathcal{E}, \text{ch})$ is a homotopy pullback

$$\begin{array}{ccc} \text{diff}(\mathcal{E}, \text{ch}) & \longrightarrow & H(\tau_{\leq 0}\Omega^* \otimes \pi_*(\mathcal{E})) \\ \downarrow & & \downarrow \\ \mathcal{E} & \xrightarrow{\text{ch}} & H(\Omega^* \otimes \pi_*(\mathcal{E})) \end{array}$$

where $\text{ch} = j \circ r$, and j induces an equivalence $j: \pi_*(\mathcal{E}) \otimes \mathbb{R}[0] \xrightarrow{\simeq} \pi_*(\mathcal{E}) \otimes \Omega^*$.

Remark 18 In our definition, we have chosen the complex $\Omega^* \otimes \pi_*(\mathcal{E})$ as the de Rham complex modeling our rational theory. In general, the differential function spectrum depends on this choice and on the equivalence j [13]. For the purposes of clarity and utility, we will always choose this model, although other models can be treated analogously. We do, however, keep the dependence on the map ch explicit to emphasize this fact.

Example 19 (Deligne cohomology) Let $\mathcal{E} = H(\mathbb{Z}[n]) \simeq \Sigma^n H\mathbb{Z}$ be the n -fold suspension of the Eilenberg–Mac Lane spectrum. In unbounded chain complexes, we have a natural isomorphism

$$\mathbb{Z}[n] \otimes \Omega^* \simeq \Omega^*[n],$$

where $\mathbb{Z}[n]$ is the sheaf of locally constant integer-valued functions in degree n , and the complex on the right-hand side has been shifted up n units. That is, Ω^n is in degree 0, while Ω^0 is in degree n . Since $\Sigma^n H\mathbb{Z}$ is in the image of the Eilenberg–Mac Lane functor H , and H preserves homotopy pullbacks, the homotopy pullback

$$\begin{array}{ccc} \text{diff}(\Sigma^n H\mathbb{Z}, \text{ch}) & \longrightarrow & H(\tau_{\leq 0}\Omega^*[n]) \\ \downarrow & & \downarrow \\ \Sigma^n H\mathbb{Z} & \xrightarrow{\text{ch}} & H(\Omega^*[n]) \end{array}$$

is presented by the homotopy pullback of unbounded chain complexes:

$$\begin{array}{ccc} \mathbb{Z}[n] \times_{\Omega^*[n]}^h \tau_{\geq 0}\Omega^*[n] & \longrightarrow & \tau_{\leq 0}\Omega^*[n] \\ \downarrow & & \downarrow \\ \mathbb{Z}[n] & \longrightarrow & \Omega^*[n] \end{array}$$

By stability, we can identify the homotopy pullback with the shifted mapping cone:

$$\mathbb{Z}[n] \times_{\Omega^*[n]}^h \tau_{\leq 0}\Omega^*[n] \simeq \text{cone}(\mathbb{Z}[n] \oplus \tau_{\leq 0}\Omega^* \rightarrow \Omega^*[n])[-1].$$

The right-hand side is precisely the Deligne complex $\mathbb{Z}_{\mathcal{D}}^{\infty}(n + 1)$. We therefore have an equivalence

$$H(\mathbb{Z}_{\mathcal{D}}^{\infty}(n + 1)) \simeq \text{diff}(\Sigma^n H\mathbb{Z}, \text{ch}).$$

The underlying theory this spectrum represents is precisely Deligne cohomology. In fact, by the Dold–Kan correspondence, we have an isomorphism of graded abelian groups

$$\pi_0 \text{hom}_{\text{ch}}(N(C(\{U_i\}), \mathbb{Z}_{\mathcal{D}}^{\infty}(n + 1))) \simeq \pi_0 \text{Map}(\Sigma^{\infty} M, \text{diff}(\Sigma^n H\mathbb{Z}, \text{ch})).$$

Here N denotes the normalized Moore complex (adjoint to the Dold–Kan functor DK) and $C(\{U_i\})$ denotes the Čech nerve of some good open cover of X . The right-hand side is simply the definition of $\text{diff}(\Sigma^n H\mathbb{Z}, \text{ch})^0(M)$, while the left-hand side is the shifted total complex of the Čech–Deligne double complex. It therefore computes the degree- n Deligne cohomology $H^n(M; \mathbb{Z}_{\mathcal{D}}^{\infty}(n + 1))$.

The above example illustrates what exactly differential function spectra have to do with differential cohomology theories. The following definition can be found in [16].

Definition 20 Let \mathcal{E} be a spectrum, and let

$$\text{ch}: \mathcal{E} \rightarrow H(\tau_{\leq 0}\Omega^* \otimes \pi_*(\mathcal{E}))$$

be the Chern character map as in Definition 17. The *differential \mathcal{E} -cohomology of a manifold* is the smooth cohomology theory with degree- n component

$$\widehat{\mathcal{E}}^n(M) \simeq \text{diff}(\Sigma^n \mathcal{E}, \text{ch})^0(M).$$

Since $\text{diff}(\Sigma^n \mathcal{E}, \text{ch})$ is a smooth spectrum for each n , it fits into a diamond diagram of the form (2-2), as established in [16; 64]. In [16], it was shown that the form that this diamond takes is precisely the differential cohomology diamond in Remark 12. In particular, Proposition 10 allows us to calculate the $\text{diff}(\Sigma^n \mathcal{E}, \text{ch})$ cohomology in degrees away from 0 as

$$\text{diff}(\Sigma^n \mathcal{E}, \text{ch})^q(M) = \begin{cases} \mathcal{E}^{n+q}(M), & q > 0, \\ \mathcal{E}_{U(1)}^{n-1+q}(M), & q < 0. \end{cases}$$

3 The smooth Atiyah–Hirzebruch spectral sequence (AHSS)

In this section, we describe general machinery to construct an Atiyah–Hirzebruch spectral sequence (AHSS) from a smooth spectrum \mathcal{E} . We also describe how to compare this spectral sequence to the classical AHSS spectral sequence for the underlying theory $\Pi\mathcal{E}$, in nice cases.

3.1 Construction of the spectral sequence via Čech resolutions

The trick to describing the spectral sequence is to choose the right filtration on a fixed manifold. In the local (projective) model structure on smooth stacks, a natural choice arises: namely, the Čech-type filtration on good open covers. This is indeed the most natural choice since the maps which are weakly inverted in the local model structure are precisely those arising from taking the Čech nerve of a good open cover of a manifold. That is, we have a weak equivalence

$$w: \operatorname{hocolim} \left\{ \cdots \begin{array}{c} \rightrightarrows \\ \rightrightarrows \\ \rightrightarrows \\ \rightrightarrows \\ \rightrightarrows \end{array} \coprod_{\alpha\beta\gamma} U_{\alpha\beta\gamma} \begin{array}{c} \rightrightarrows \\ \rightrightarrows \\ \rightrightarrows \\ \rightrightarrows \\ \rightrightarrows \end{array} \coprod_{\alpha\beta} U_{\alpha\beta} \begin{array}{c} \rightrightarrows \\ \rightrightarrows \\ \rightrightarrows \\ \rightrightarrows \\ \rightrightarrows \end{array} \coprod_{\alpha} U_{\alpha} \right\} \rightarrow X.$$

We now explicitly describe a filtration on $C(\{U_i\})$. Recall that any simplicial diagram $J: \Delta^{\text{op}} \rightarrow \operatorname{Sh}_{\infty}(\operatorname{CartSp})$ can be filtrated by skeleta. More precisely, let $i: \Delta_{\leq k} \hookrightarrow \Delta$ denote the embedding of the full subcategory of linearly ordered sets $[r]$ such that $r \leq k$. Then i induces a restriction between functor categories (the k^{th} truncation)

$$\tau_{\leq k}: [\Delta^{\text{op}}, \operatorname{Sh}_{\infty}(\operatorname{CartSp})] \rightarrow [\Delta_{\leq k}^{\text{op}}, \operatorname{Sh}_{\infty}(\operatorname{CartSp})].$$

By general abstract nonsense (the existence of left and right Kan extensions), there are left and right adjoints ($\operatorname{sk}_k \dashv \tau_{\leq k} \dashv \operatorname{cosk}_k$)

$$[\Delta^{\text{op}}, \operatorname{Sh}_{\infty}(\operatorname{CartSp})] \begin{array}{c} \xleftarrow{\operatorname{sk}_k} \\ \xrightarrow{\tau_{\leq k}} \\ \xleftarrow{\operatorname{cosk}_k} \end{array} [\Delta_{\leq k}^{\text{op}}, \operatorname{Sh}_{\infty}(\operatorname{CartSp})].$$

Furthermore, by composing adjoints, we have an adjunction ($\operatorname{sk}_k \dashv \operatorname{cosk}_k$)

$$[\Delta^{\text{op}}, \operatorname{Sh}_{\infty}(\operatorname{CartSp})] \begin{array}{c} \xrightarrow{\operatorname{sk}_k} \\ \xleftarrow{\operatorname{cosk}_k} \end{array} [\Delta^{\text{op}}, \operatorname{Sh}_{\infty}(\operatorname{CartSp})].$$

The functor sk_k freely fills in degenerate simplices above level k , while cosk_k probes a simplicial object with simplices only up to level k (the singular k -skeleton).

Proposition 21 *Let Y_{\bullet} be a simplicial object in $\operatorname{Sh}_{\infty}(\operatorname{CartSp})$. Then we can filter Y_{\bullet} by skeleta*

$$\operatorname{sk}_0 Y_{\bullet} \rightarrow \operatorname{sk}_1 Y_{\bullet} \rightarrow \cdots \rightarrow \operatorname{sk}_k Y_{\bullet} \rightarrow \cdots \rightarrow Y_{\bullet}.$$

The homotopy colimit over Y_{\bullet} is presented by the ordinary colimit

$$\operatorname{hocolim}_{\Delta^{\text{op}}} (Y_{\bullet}) \simeq \operatorname{colim}_{k \rightarrow \infty} \mathbb{L}\operatorname{colim}_{\Delta^{\text{op}}} (\operatorname{sk}_k Y_{\bullet}),$$

where $\mathbb{L}\operatorname{colim}$ is the left derived functor of the colimit, hence computable upon suitable cofibrant replacement of the diagram.¹

¹We take this particular model of the homotopy colimit in order to ensure that taking the colimit of the resulting diagram makes sense. The claim will also hold for other presentations of the homotopy colimit.

Proof Since $\text{Sh}_\infty(\text{CartSp})$ is presented by a combinatorial simplicial model category, the homotopy colimit over a filtered diagram is presented by the ordinary colimit, and the canonical map

$$\mathbb{L}\text{colim}_{k \rightarrow \infty} \mathbb{L}\text{colim}_{\Delta^{\text{op}}}(\text{sk}_k Y_\bullet) \rightarrow \text{colim}_{k \rightarrow \infty} \mathbb{L}\text{colim}_{\Delta^{\text{op}}}(\text{sk}_k Y_\bullet)$$

is an equivalence. Since homotopy colimits commute with homotopy colimits, we also have an equivalence

$$\mathbb{L}\text{colim}_{k \rightarrow \infty} \mathbb{L}\text{colim}_{\Delta^{\text{op}}}(\text{sk}_k Y_\bullet) \simeq \mathbb{L}\text{colim}_{\Delta^{\text{op}}} \mathbb{L}\text{colim}_{k \rightarrow \infty}(\text{sk}_k Y_\bullet).$$

Again, using the fact that the ordinary colimit over a filtered diagram presents the homotopy colimit, we have an equivalence

$$\mathbb{L}\text{colim}_{\Delta^{\text{op}}} \mathbb{L}\text{colim}_{k \rightarrow \infty}(\text{sk}_k Y_\bullet) \rightarrow \mathbb{L}\text{colim}_{\Delta^{\text{op}}} \text{colim}_{k \rightarrow \infty}(\text{sk}_k Y_\bullet) \simeq \mathbb{L}\text{colim}_{\Delta^{\text{op}}}(Y_\bullet). \quad \square$$

Remark 22 The above proposition says that the homotopy colimit over the simplicial object is filtered by homotopy colimits of its skeleta. In particular, if M is a paracompact manifold, we can fix a good open cover on M and form the simplicial object given by its Čech nerve

$$C(\{U_i\}) := \cdots \begin{array}{c} \rightrightarrows \\ \rightrightarrows \\ \rightrightarrows \\ \rightrightarrows \end{array} \coprod_{\alpha\beta\gamma} U_{\alpha\beta\gamma} \begin{array}{c} \rightrightarrows \\ \rightrightarrows \\ \rightrightarrows \\ \rightrightarrows \end{array} \coprod_{\alpha\beta} U_{\alpha\beta} \begin{array}{c} \rightrightarrows \\ \rightrightarrows \\ \rightrightarrows \\ \rightrightarrows \end{array} \coprod_{\alpha} U_{\alpha}.$$

The homotopy colimit over this object is then filtered by its skeleta.

Let us see exactly what the skeleta look like in this case. To this end, we recall that in $\text{Sh}_\infty(\text{CartSp})$, the full homotopy colimit is presented by the local homotopy formula

$$\text{hocolim}_{\Delta^{\text{op}}} C(\{U_i\}) = \int^{n \in \Delta} \coprod_{\alpha_0 \cdots \alpha_n} U_{\alpha_0 \cdots \alpha_n} \odot \Delta[n].$$

The filtration on this object is given by first truncating the Čech nerve and then freely filling in degenerate simplices. As a consequence, in degree k , we can forget about the simplices of dimension higher than k . The homotopy colimit over this skeleton is then given by a strict colimit over the diagram

$$(3-1) \quad \coprod_{\alpha_0 \cdots \alpha_k} U_{\alpha_0 \cdots \alpha_k} \odot \Delta[k] \cdots \begin{array}{c} \rightrightarrows \\ \rightrightarrows \\ \rightrightarrows \\ \rightrightarrows \end{array} \coprod_{\alpha\beta\gamma} U_{\alpha\beta\gamma} \odot \Delta[2] \\ \begin{array}{c} \rightrightarrows \\ \rightrightarrows \\ \rightrightarrows \\ \rightrightarrows \end{array} \coprod_{\alpha\beta} U_{\alpha\beta} \odot \Delta[1] \begin{array}{c} \rightrightarrows \\ \rightrightarrows \\ \rightrightarrows \\ \rightrightarrows \end{array} \coprod_{\alpha} U_{\alpha} \odot \Delta[0],$$

where the face and degeneracy maps are induced by the face and degeneracy maps of $\Delta[k]$. Taking $k \rightarrow \infty$, we do indeed reproduce the coend representing the full homotopy colimit $C(\{U_i\})$.

We would like to eventually use this filtration to define a Mayer–Vietoris like spectral sequence for general cohomology theory \mathcal{E} . To get to this step, however, we will need to identify the successive quotients of the filtration. To simplify notation in what follows, we will fix a manifold M with Čech nerve $C(\{U_i\})$, and we set

$$X_k := \operatorname{hocolim}_{\Delta^{\text{op}}}(\operatorname{sk}_k C(\{U_i\})).$$

Then the quotient X_k/X_{k-1} can be identified from the previous discussion by quotienting out the face maps at level k described in diagram (3-1). Since the tensor of a simplicial set and a stack is given by the product of the stack with the discrete inclusion of the simplicial set, we can identify the quotient from the pushout of coends

$$\begin{array}{ccc} \int^{n < k} \coprod_{\alpha_0 \cdots \alpha_n} U_{\alpha_0 \cdots \alpha_n} \times \operatorname{disc}(\Delta[n]) & \longrightarrow & * \\ & \downarrow \partial & \\ \int^{m \leq k} \coprod_{\alpha_0 \cdots \alpha_m} U_{\alpha_0 \cdots \alpha_m} \times \operatorname{disc}(\Delta[m]) & & \end{array}$$

where ∂ denotes the boundary inclusion. At the level of points (or elements), a simplex in $\int^{n < k} \coprod_{\alpha_0 \cdots \alpha_n} U_{\alpha_0 \cdots \alpha_n} \times \operatorname{disc}(\Delta[n])$ is given by a pair

$$(\rho, \sigma) \in \coprod_{\alpha_0 \cdots \alpha_{k-1}} U_{\alpha_0 \cdots \alpha_{k-1}} \times \operatorname{disc}(\Delta[k-1]),$$

which is glued to lower simplices via the face and degeneracy relations.

Let us identify where the boundary inclusion takes a generic simplex. Then the quotient X_k/X_{k-1} will be obtained by gluing these simplices together to a single point. Note that the face and degeneracy relations imply that simplices of the form $(\rho, s_{j+1}\sigma)$ are sent by d_j to $(d_j\rho, \sigma)$. Since simplices in the image of the face maps are precisely those which are collapsed to a point, we see that

$$(d_j\rho, \sigma) \sim * \quad \text{for every } \sigma.$$

We therefore see that each term of the coproduct $\coprod_{\alpha_0 \cdots \alpha_k} U_{\alpha_0 \cdots \alpha_k}$ is joined to another by the inclusion into a lower intersection. These lower intersections are then collapsed to a point yielding the wedge product

$$\bigvee_{\alpha_0 \cdots \alpha_k} U_{\alpha_0 \cdots \alpha_k} \subset X_k/X_{k-1}.$$

Similarly, the simplex $(s_{j+1}\rho, \sigma)$ is sent to $(\rho, d_j\sigma)$ under d_j . We therefore identify the discrete simplicial sphere in the quotient

$$\operatorname{disc}(\Delta[k]/\partial\Delta[k]) \subset X_k/X_{k-1}.$$

Finally, the relations imposed by the coend imply that a simplex of the form $(s_j \rho, \sigma)$ is glued to $(\rho, d_j \sigma)$. The former are precisely those simplices in the simplicial sphere, while the latter are glued to the point. Similarly, $(\rho, s_j \sigma)$ is glued to the point. Thus we have the following.

Lemma 23 *We can identify the quotient with the smash product:*

$$X_k / X_{k-1} \simeq \text{disc}(\Delta[k] / \partial \Delta[k]) \wedge \bigvee_{\alpha_0 \cdots \alpha_k} U_{\alpha_0 \cdots \alpha_k} \simeq \Sigma^k (\bigvee_{\alpha_0 \cdots \alpha_k} U_{\alpha_0 \cdots \alpha_k}).$$

Remark 24 (the filtration as a natural choice) Another way to think of our filtration above is the following. Let us form a Čech nerve of a manifold, then contract all the patches and intersections in that Čech nerve as points, so we obtain a simplicial set. Then *Borsuk's nerve theorem* (see [6] for a survey, [36, Corollary 4G.3] or [57, Theorem 3.21]) says that this simplicial set is equivalent — weak homotopy equivalent — to the singular simplicial complex of the manifold, hence to its homotopy type. Moreover, that singular simplicial complex (or rather its geometric realization) in turn gives a CW-complex realization of the original manifold. So with this in mind, one may view our filtration above as the natural smooth refinement of the filtration by CW-stages of the manifold. That is, in taking the Čech nerve *without* contracting all its patches to points, we retain exactly the smooth information that, via Borsuk's theorem, corresponds to each cell in the canonical CW-complex incarnation of the manifold. So in this sense, our refinement can be viewed as the canonical smooth refinement of the traditional filtering by CW-stages.

We are now ready to describe the spectral sequence.

Theorem 25 (AHSS for general smooth spectra) *Let M be a compact smooth manifold, and let \mathcal{E} be a smooth spectrum. There is a spectral sequence with*

$$E_2^{p,q} = H^p(M, \mathcal{E}^q) \implies \mathcal{E}^{p+q}(M).$$

Here H^p denotes the p^{th} Čech cohomology with coefficients in the presheaf \mathcal{E}^q . Moreover, the differential on the E_1 -page is given by the differential in Čech cohomology.

Proof The proof is almost immediate from the definitions. Recall that we have identified the quotients in Lemma 23. By the axioms for a smooth cohomology theory, we have that the \mathcal{E} -cohomology of the quotient is given by

$$\begin{aligned} \mathcal{E}^*(X_k / X_{k-1}) &\simeq \mathcal{E}^*(\Sigma^k (\bigvee_{\alpha_0 \cdots \alpha_k} U_{\alpha_0 \cdots \alpha_k})) \\ &\simeq \mathcal{E}^{*-k} (\bigvee_{\alpha_0 \cdots \alpha_k} U_{\alpha_0 \cdots \alpha_k}) \\ &\simeq \bigoplus_{\alpha_0 \cdots \alpha_k} \mathcal{E}^{*-k} (U_{\alpha_0 \cdots \alpha_k}). \end{aligned}$$

Applying \mathcal{E}^{p+q} to the cofiber sequence $X_p \hookrightarrow X_{p+1} \twoheadrightarrow X_{p+1}/X_p$ gives the long exact sequence in \mathcal{E} -cohomology

$$(3-2) \quad \dots \rightarrow \mathcal{E}^{p+q}(X_{p+1}/X_p) \rightarrow \mathcal{E}^{p+q}(X_{p+1}) \rightarrow \mathcal{E}^{p+q}(X_p) \rightarrow \mathcal{E}^{p+q+1}(X_{p+1}/X_p) \rightarrow \dots$$

Forming the corresponding exact triangle, we get a spectral sequence with $E_1^{p,q}$ -term

$$E_1^{p,q} = \bigoplus_{\alpha_0, \dots, \alpha_p} \mathcal{E}^q(U_{\alpha_0 \dots \alpha_p}).$$

Now we want to show that the differential on this page is given by the Čech differential

$$\delta: E_1^{p,q} = \bigoplus_{\alpha_0 \dots \alpha_p} \mathcal{E}^q(U_{\alpha_0 \dots \alpha_p}) \rightarrow \bigoplus_{\alpha_0 \dots \alpha_{p+1}} \mathcal{E}^q(U_{\alpha_0 \dots \alpha_{p+1}}) = E_1^{p+1,q}.$$

To this end, note that differential on the E_1 -page, by definition, comes from the exact sequence

$$\dots \rightarrow \mathcal{E}^{p+q}(X_{p+1}/X_p) \xrightarrow{j} \mathcal{E}^{p+q}(X_{p+1}) \xrightarrow{i} \mathcal{E}^{p+q}(X_p) \xrightarrow{\partial} \mathcal{E}^{p+q+1}(X_{p+1}/X_p) \rightarrow \dots$$

We need to show that $\partial j = d_1 = \delta$ is the Čech differential. By naturality of the connecting homomorphism ∂ , we have a commutative diagram

$$\begin{array}{ccc} \check{C}^{p-1}(M; \mathcal{E}^q) & \xrightarrow{d_1} & \check{C}^p(M; \mathcal{E}^q) \\ \cong \downarrow & & \downarrow \cong \\ \bigoplus_{\alpha_0 \dots \alpha_{p-1}} \mathcal{E}^q(U_{\alpha_0 \dots \alpha_{p-1}}) & \xrightarrow{\quad} & \bigoplus_{\alpha_0 \dots \alpha_p} \mathcal{E}^q(U_{\alpha_0 \dots \alpha_p}) \\ \cong \downarrow & & \downarrow \cong \\ \mathcal{E}^{p+q-1}(X_{p-1}/X_{p-2}) & \xrightarrow{j} \mathcal{E}^{p+q-1}(X_{p-1}) \xrightarrow{\partial} & \mathcal{E}^{p+q}(X_p/X_{p-1}) \\ \downarrow & \downarrow & \downarrow \\ \mathcal{E}^{p+q-1}(\partial\Delta[p] \times U_{\alpha_0 \dots \alpha_{p-1}}) & & \mathcal{E}^{p+q}(\Delta[p]/\partial\Delta[p] \wedge U_{\alpha_0 \dots \alpha_p}) \\ & \searrow \text{id} \quad \downarrow \quad \nearrow \partial & \\ & \mathcal{E}^{p+q-1}(\partial\Delta[p] \times U_{\alpha_0 \dots \alpha_{p-1}}) & \end{array}$$

where the vertical bottom maps are induced from the inclusion of a factor

$$(3-3) \quad \begin{array}{ccc} \Delta[p] \times U_{\alpha_0 \dots \alpha_p} & \hookrightarrow & X_p \\ \uparrow & & \uparrow \\ \partial\Delta[p] \times U_{\alpha_0 \dots \alpha_{p-1}} & \hookrightarrow & X_{p-1} \\ \uparrow & & \uparrow \\ \emptyset & \longrightarrow & X_{p-2} \end{array}$$

into the p -level of the filtration. Comparing the top and bottom composite morphisms in the big diagram, we see that on $(p-1)$ -fold intersections $U_{\alpha_0 \dots \alpha_{p-1}}$, the map d_1 is forced to map a section to the alternating sum of restrictions, as this is precisely the map induced by the boundary inclusion in (3-3).

All that remains is the convergence. To establish that, we simply note that compactness implies that, for large values of p , we have an equivalence $X_p \simeq X$. Moreover, there are only finitely many diagonal entries at each page of the sequence. With this assumption, the convergence to the corresponding graded complex

$$E_{\infty}^{p,q} = \frac{\ker(\mathcal{E}^{p+q}(X) \rightarrow \mathcal{E}^{p+q}(X_p))}{\ker(\mathcal{E}^{p+q}(X) \rightarrow \mathcal{E}^{p+q}(X_{p+1}))} = \frac{F_p \mathcal{E}^{p+q}(X)}{F_{p+1} \mathcal{E}^{p+q}(X)}$$

follows exactly as in the classical case in [3]. □

Fiber bundles We can also construct a spectral sequence for a fiber bundle

$$F \rightarrow N \xrightarrow{p} M,$$

where each map is a smooth map of manifolds and M is compact. To that end, we note that for a fixed good open cover $\{U_i\}$ of M , the pullbacks $\{p^{-1}(U_i)\}$ define a good open cover of N . By local triviality, we have that each $p^{-1}(U_i) \simeq F \times U_i$. Then, using the filtration

$$X_k = \text{hocolim}_{\Delta^{\text{op}}}(\text{sk}_k C(\{p^{-1}(U_i)\}))$$

on the total space N , we identify the successive quotients

$$X_k / X_{k-1} \simeq \Sigma^k \bigvee_{\alpha_0 \dots \alpha_k} U_{\alpha_0 \dots \alpha_k} \wedge F.$$

A similar argument as in the proof of Theorem 25 gives:

Theorem 26 (smooth AHSS for fiber bundles) *Let M, N and F be manifolds with M compact. Let $F \rightarrow N \xrightarrow{p} M$ be a fiber bundle. Let \mathcal{E} be a sheaf of spectra. Then there is a spectral sequence*

$$E_2^{p,q} = H^p(M, \mathcal{E}^q(- \wedge F)) \implies \mathcal{E}^{p+q}(N).$$

Here H^p denotes the p^{th} Čech cohomology with coefficients in the presheaf $\mathcal{E}^{-q}(- \wedge F)$.

Remark 27 (unreduced theories) Note that the smooth spectral sequence works for reduced theories. One can treat unreduced theories similarly by setting

$$\mathcal{E}^q(M, *) := \tilde{\mathcal{E}}^q(M_+),$$

where the tilde denotes the reduced theory and M_+ is the pointed stack with basepoint $*$. In this case, we have the slight modification on the second spectral sequence, which takes the form

$$E_2^{p,q} = H^p(M, \mathcal{E}^q(- \times F)) \implies \mathcal{E}^{p+q}(N).$$

3.2 Morphisms of smooth spectral sequences and refinement of the AHSS

Our next task will be to show that these spectral sequences do indeed refine the classical Atiyah–Hirzebruch spectral sequence (AHSS) [3]. Since any smooth theory \mathcal{E} comes as a refinement of the underlying topological theory $\Pi\mathcal{E}$, we will immediately get a morphism of spectral sequences induced by the morphism of spectra

$$I: \mathcal{E} \rightarrow \Pi\mathcal{E}.$$

Unfortunately, this morphism does not allow us to compare the differentials of the spectral sequences in the way that we would ideally hope for. However, as we will progressively see, the situation can be remedied by constructing a slightly different morphism of spectral sequences. This morphism is related to the *boundary map* of spectral sequences which occurs when a morphism of spectra induces the 0 map on corresponding spectral sequences; see [55] for a discussion in the case of the Adams spectral sequence. We first discuss the morphism induced by I , then construct this “boundary-type” map, and prove that it indeed defines a morphism of spectral sequences.

Definition 28 Let $E_n^{p,q}$ and $F_n^{p,q}$ be spectral sequences, that is, a sequence of bigraded complexes $E_n^{p,q}$ and $F_n^{p,q}$, $n \in \mathbb{N}$. A *morphism of spectral sequences* is a morphism of bigraded complexes

$$f_n: E_n^{p,q} \rightarrow F_n^{p,q},$$

defined for all $n > N$, where N is some fixed positive integer. Furthermore, we require the map f_{n+1} to be the map on homology induced by f_n . We call the smallest integer N such that f_n are defined for $n > N$ the *rank* of the morphism.

We now apply this to the smooth AHSS. The next result should follow from general principles, but we emphasize it explicitly for clarity and for subsequent use.

Proposition 29 Let \mathcal{E} and \mathcal{F} be smooth spectra. Then a map $f: \mathcal{E} \rightarrow \mathcal{F}$ induces a morphism of corresponding smooth AHSSs

$$E_n^{p,q} \rightarrow F_n^{p,q}.$$

Proof Fix a manifold X and a good open cover $\{U_i\}$. Let X_p denote the p^{th} filtration of the Čech nerve as before. It is clear by naturality that a map of spectra $f: \mathcal{E} \rightarrow \mathcal{F}$ induces a morphism of long exact sequences (see (3-2))

$$\begin{array}{cccccccc} \dots & \mathcal{E}^{p+q}(X_{p+1}/X_p) & \longrightarrow & \mathcal{E}^{p+q}(X_{p+1}) & \longrightarrow & \mathcal{E}^{p+q}(X_p) & \longrightarrow & \mathcal{E}^{p+q+1}(X_{p+1}/X_p) \cdots \\ & \downarrow & & \downarrow & & \downarrow & & \downarrow \\ \dots & \mathcal{F}^{p+q}(X_{p+1}/X_p) & \longrightarrow & \mathcal{F}^{p+q}(X_{p+1}) & \longrightarrow & \mathcal{F}^{p+q}(X_p) & \longrightarrow & \mathcal{F}^{p+q+1}(X_{p+1}/X_p) \cdots \end{array}$$

It follows immediately from the construction of the corresponding exact triangles that this morphism commutes with the differentials. □

This now allows us to compare the topological and the smooth theories.

Corollary 30 *Let \mathcal{E} be a smooth spectrum and $\Pi\mathcal{E}$ the underlying topological theory. Let E_n and F_n denote the spectral sequences corresponding to \mathcal{E} and $\Pi\mathcal{E}$, respectively. The natural map $I: \mathcal{E} \rightarrow \Pi\mathcal{E}$ induces a morphism of classical AHSSs²*

$$I: E_n^{p,q} \rightarrow F_n^{p,q}.$$

Remark 31 It is interesting that the smooth spectrum $\Pi\mathcal{E}$ is, by definition, locally constant. From the discussion around (2-1), this means that we have an isomorphism

$$\Pi\mathcal{E}^q(U) \simeq \pi_{-q} \text{Map}(U, \Pi\mathcal{E}) \simeq \pi_{-q} \text{Map}(*, \Pi\mathcal{E}) \simeq \pi_{-q} \Pi\mathcal{E} \simeq \Pi\mathcal{E}^q(*)$$

for every element of a good open cover (or higher intersection) U . This connects, via Borsuk’s theorem mentioned in Remark 24 above, the “smooth AHSS for locally constant coefficients” with the classical AHSS: the locally constant coefficients see each (contractible) patch as a point, and hence by Borsuk’s theorem, they see our “Čech filtration” to be the classical CW-cell filtration.

From the construction of our smooth AHSS, it directly follows that the spectral sequence associated to the smooth spectrum is a refinement of the classical topological AHSS.

Corollary 32 *The spectral sequence $F_n^{p,q}$ is precisely the AHSS for the cohomology theory $\Pi\mathcal{E}$.*

²Here we have an unfortunate conflict of notation. We are using the same symbols for the pages in the spectral sequences for both the classical and the refined theories. We will aim to make the context explicit whenever a possible ambiguity arises.

We now would like to apply the above machinery to differential cohomology theories. In particular, we note that for a differential function spectrum $\text{diff}(\mathcal{E}, \text{ch})$, the natural map

$$I: \text{diff}(\mathcal{E}, \text{ch}) \rightarrow \underline{\mathcal{E}},$$

which strips the differential theory of the differential data and maps to the bare underlying theory, is precisely the map induced by the unit $I: \text{id} \rightarrow \Pi$. In the above discussion, we observed that this map always induces a morphism of spectral sequences. Moreover, the target spectral sequence is exactly the AHSS for the underlying topological theory. One might hope to be able to use this map to compare the differentials in the refined theory with those differentials in the classical AHSS.

Unfortunately, this does not work in practice, as we will see when we discuss applications in Section 4. The core issue is that the spectral sequence for the refined theory usually ends up shifted with respect to the classical AHSS. As a consequence, the nonzero terms in each sequence are interlaced with respect to one another, and the map I ends up killing all the nonzero terms. This, in turn, stems from the appearance of the Bockstein map (which raises degree by 1) in the differential cohomology diagram.

However, there is often a different map between the *lower quadrants* of the two spectral sequences corresponding to $\text{diff}(\mathcal{E}, \text{ch})$ and \mathcal{E} , which lowers the degree as to match the corresponding nonzero entries. This map is related to the so-called *boundary map* between spectral sequences studied in [55]. The next proposition concerns this map and will be essential for comparing the differentials in the refined theory to those of the classical theory.

Proposition 33 (i) *Let \mathcal{E} be a spectrum such that $\pi_*(\mathcal{E})$ is concentrated in degrees which are a multiple of some integer $n \geq 2$ (eg K-theory, Morava K-theory). Suppose, moreover, that $\pi_*(\mathcal{E})$ is projective in those degrees. Then the sequence of spectra*

$$\mathcal{E} \rightarrow \mathcal{E} \wedge M\mathbb{R} \rightarrow \mathcal{E} \wedge MU(1) \xrightarrow{\beta_{\mathcal{E}}} \Sigma\mathcal{E}$$

induces a short exact sequence on coefficients

$$(3-4) \quad 0 \rightarrow \pi_*(\mathcal{E}) \rightarrow \pi_*(\mathcal{E}) \otimes \mathbb{R} \rightarrow \pi_*(\mathcal{E}) \otimes U(1) \rightarrow 0.$$

(ii) *Let β denote the connecting homomorphism (ie the Bockstein) for the coefficient sequence (3-4). Let $E_n^{p,q}$ denote the spectral sequence corresponding to $\Sigma^{-1}\mathcal{E} \wedge MU(1)$ and let $F_n^{p,q}$ denote the spectral sequence corresponding to \mathcal{E} . Then*

$$\beta: E_n^{p,q} \rightarrow F_n^{p,q}$$

induces a morphism of spectral sequences of rank 2.

Proof Consider the long Bockstein sequence

$$\dots \rightarrow \mathcal{E} \xrightarrow{r} \mathcal{E} \wedge M\mathbb{R} \xrightarrow{e} \mathcal{E} \wedge MU(1) \xrightarrow{\beta_{\mathcal{E}}} \Sigma\mathcal{E} \rightarrow \dots$$

induced by the cofiber sequence

$$\mathbb{S} \rightarrow M\mathbb{R} \rightarrow MU(1).$$

Fix a manifold M and let X_p denote the p -level of the Čech filtration. Now each spectrum in the above sequence has a long exact sequence induced by the cofiber sequences

$$X_{p-1} \rightarrow X_p \rightarrow X_p/X_{p-1},$$

from which one builds the exact couple for the corresponding spectral sequence. Using the properties of $\pi_*(\mathcal{E})$ along with this sequence, we can fit the long exact sequences into a diagram

$$\begin{array}{ccccccc} \check{C}^p(X; \pi_{-q-1}(\mathcal{E})) & \xrightarrow{q^*} & \mathcal{E}^{p+q-1}(X_p) & \xrightarrow{i^*} & \mathcal{E}^{p+q-1}(X_{p-1}) & \xrightarrow{\partial} & 0 \\ \downarrow r & & \downarrow r & & \downarrow r & & \downarrow \\ \check{C}^p(X; \pi_{-q-1}(\mathcal{E}_{\mathbb{R}})) & \xrightarrow{q^*} & \mathcal{E}_{\mathbb{R}}^{p+q-1}(X_p) & \xrightarrow{i^*} & \mathcal{E}_{\mathbb{R}}^{p+q-1}(X_{p-1}) & \xrightarrow{\partial} & 0 \\ \downarrow e & & \downarrow e & & \downarrow e & & \downarrow \\ \check{C}^p(X; \pi_{-q-1}(\mathcal{E}_{U(1)})) & \xrightarrow{q^*} & \mathcal{E}_{U(1)}^{p+q}(X_p) & \xrightarrow{i^*} & \mathcal{E}_{U(1)}^{p+q}(X_{p-1}) & \xrightarrow{\partial} & 0 \\ \downarrow \beta_{\mathcal{E}} & & \downarrow \beta_{\mathcal{E}} & & \downarrow \beta_{\mathcal{E}} & & \downarrow \\ 0 & \longrightarrow & \mathcal{E}^{p+q}(X_{p+1}) & \longrightarrow & \mathcal{E}^{p+q}(X_p) & \longrightarrow & \check{C}^p(X; \pi_{-q+1}(\mathcal{E})) \end{array}$$

where both the rows and columns are part of exact sequences, and $\check{C}^p(X; A)$ denotes the group of Čech p -cochains with coefficients in A . Since everything commutes, this induces a corresponding short exact sequence of E_1 -pages. At each (p, q) -entry, this sequence is given by

$$0 \rightarrow C^p(X; \pi_{-q}(\mathcal{E})) \rightarrow C^p(X; \pi_{-q}(\mathcal{E}) \otimes \mathbb{R}) \rightarrow C^p(X; \pi_{-q}(\mathcal{E}) \otimes U(1)) \rightarrow 0.$$

Since the differentials on the E_1 -page are precisely the Čech differentials, the construction of the Bockstein map in Čech cohomology will produce a map of E_2 -pages

$$\beta: H^p(X; \pi_{-q}(\mathcal{E}) \otimes U(1)) \rightarrow H^{p+1}(X; \pi_{-q}(\mathcal{E})).$$

We need to show that this map commutes with the differential. Choose a representative x of a class in $H^p(X; \pi_{-q}(\mathcal{E}) \otimes U(1))$. By definition, $y = \beta(x)$ is a class such that $r(y) = \delta(\bar{x})$, where \bar{x} is such that $e(\bar{x}) = x$. Then

$$r(d_2y) = d_2r(y) = d_2\delta(\bar{x}).$$

We want to show that there is a lift z of d_2x such that $\delta(z) = d_2\delta(\bar{x})$. Indeed, if this is the case, then d_2y represents $\beta(d_2x)$ and we are done.

To construct z , recall that d_2x is defined by first pulling back by the quotient q , which lies in the image of the map induced by the inclusion $i: X_p \hookrightarrow X_{p+1}$, and then applying the boundary to an element of the preimage. Let w be such that

$$i^*(w) = q^*(x).$$

Chasing the diagram

$$\begin{array}{ccccc} \check{C}^p(X; \pi_{-q-1}(\mathcal{E}_{\mathbb{R}})) & \xrightarrow{q^*} & \mathcal{E}_{\mathbb{R}}^{p+q-1}(X_p) & \xrightarrow{e} & \mathcal{E}_{\mathbb{R}}^{p+q-1}(X_{p-1}) \\ \downarrow e & & \downarrow e & & \downarrow e \\ \check{C}^p(X; \pi_{-q-1}(\mathcal{E}_{U(1)})) & \xrightarrow{q^*} & \mathcal{E}_{U(1)}^{p+q}(X_p) & \xrightarrow{i^*} & \mathcal{E}_{U(1)}^{p+q}(X_{p-1}) \\ \downarrow \beta_{\mathcal{E}} & & \downarrow \beta_{\mathcal{E}} & & \downarrow \beta_{\mathcal{E}} \\ 0 & \longrightarrow & \mathcal{E}^{p+q}(X_p) & \xrightarrow{i^*} & \mathcal{E}^{p+q}(X_{p-1}) \end{array}$$

we see that $0 = \beta_{\mathcal{E}}q^*(x) = \beta_{\mathcal{E}}i^*(w) = i^*(\beta_{\mathcal{E}}w)$. By exactness of the rows, this implies that $\beta_{\mathcal{E}}w = 0$. Therefore, there is a class $\bar{w} \in \mathcal{E}_{\mathbb{R}}^{p+q+1}(X_{p+1})$ such that $e(\bar{w}) = w$.

Now, by definition of the differential, we have

$$e(\partial\bar{w}) = \partial(e(\bar{w})) = \partial w = d_2x,$$

and $z := \partial\bar{w}$ is a lift of d_2x . Using the fact that $\delta = d_1 = \partial q^*$, we have

$$\delta(z) = \delta(\partial\bar{w}) = \partial(q^*\partial\bar{w}).$$

By exactness, we have

$$i^*(q^*\partial\bar{w}) = 0 = q^*\partial q^*(\bar{x}) = q^*(\delta(\bar{x})),$$

and it follows from the definition that $\delta(z) = d_2(\delta(\bar{x}))$.

To show that $H^*(\beta)$ commutes with the higher differentials, we proceed by induction. The above discussion proves the base case. Suppose β induces a map $H_n(\beta)$ on E_n which commutes with d_n . Then $H^n(\beta)$ induces a well-defined map $H^{n+1}(\beta)$ on the E_{n+1} -page. Let $x \in \bigcap_{i=1}^n \ker(d_{n+1})$ be a representative of a class on the E_n -page. Then by definition, $H^{n+1}(\beta)(x) = \beta(x)$, and the exact same argument as before (replacing d_2 with d_{n+1}) gives the result. \square

Having done the heavy lifting in the above proposition, we will now apply this to straightforwardly relate the differentials of the refined theory to those of the underlying topological theory. This will use an explicit alternative to the map I , along the lines of the discussion preceding Proposition 33.

Theorem 34 (refinement of differentials) *Let \mathcal{E} be a spectrum satisfying the properties of Proposition 33, and let $\text{diff}(\mathcal{E}, \text{ch})$ be a differential function spectrum refining \mathcal{E} . Let E_n and F_n denote the smooth AHSSs corresponding to $\text{diff}(\mathcal{E}, \text{ch})$ and \mathcal{E} , respectively. Then the Bockstein β defines a rank-2 morphism of fourth quadrant spectral sequences*

$$\beta: E_n^{p,q} \rightarrow F_n^{p,q}, \quad q < 0.$$

Proof Recall that for $q < 0$, Proposition 10 implies that $\text{diff}(\mathcal{E}, \text{ch})^q(M) \simeq \mathcal{E}_{U(1)}^{q-1}(M)$. The claim then follows from the previous proposition. \square

3.3 Product structure and the differentials

Let \mathcal{E} be an E_∞ ring spectrum. Then the associative graded-commutative product on \mathcal{E}^* induces a product (associative and graded-commutative) on the refinement $\text{diff}(\Sigma^n \mathcal{E}, \text{ch})^*$, that is, a map

$$(3-5) \quad \cup: \text{diff}(\Sigma^n \mathcal{E}, \text{ch})^k \otimes \text{diff}(\Sigma^m \mathcal{E}, \text{ch})^j \rightarrow \text{diff}(\Sigma^{n+m} \mathcal{E}, \text{ch})^{k+j}$$

(see [13; 71]). The goal of this section will be to establish the following very useful property, in analogy with the classical case.

Proposition 35 (compatibility with products) *The product*

$$\cup: \text{diff}(\Sigma^n \mathcal{E}, \text{ch})^k \otimes \text{diff}(\Sigma^m \mathcal{E}, \text{ch})^j \rightarrow \text{diff}(\Sigma^{n+m} \mathcal{E}, \text{ch})^{k+j}$$

induces a morphism of spectral sequences

$$\cup: E_*(n) \times E_*(m) \rightarrow E_*(n+m).$$

Moreover, the differentials satisfy the Leibniz rule

$$d(xy) = d(x)y + (-1)^{p+q} x d(y).$$

Let us first work out what the cup product pairing is on the E_1 -page. Recall from the construction of the spectral sequence that $E_1^{p,q}$ is given by

$$E_1^{p,q} = \bigoplus_{\alpha_0 \cdots \alpha_p} \text{diff}(\Sigma^n \mathcal{E}, \text{ch})^q(U_{\alpha_0 \cdots \alpha_p}) \simeq \check{C}^p(M; \text{diff}(\Sigma^n \mathcal{E}, \text{ch})^q).$$

Using the product (3-5), we get a cross product map

$$(3-6) \quad \times: \bigoplus_{\alpha_0 \cdots \alpha_p} \text{diff}(\Sigma^n \mathcal{E}, \text{ch})^q(U_{\alpha_0 \cdots \alpha_p}) \times \bigoplus_{\alpha_0 \cdots \alpha_r} \text{diff}(\Sigma^m \mathcal{E}, \text{ch})^t(U_{\alpha_0 \cdots \alpha_r}) \\ \rightarrow \bigoplus_{\alpha_0 \cdots \alpha_p} \bigoplus_{\alpha_0 \cdots \alpha_r} \text{diff}(\Sigma^{n+m} \mathcal{E}, \text{ch})^{q+t}(U_{\alpha_0 \cdots \alpha_p} \times U_{\alpha_0 \cdots \alpha_r}).$$

We also have an isomorphism

$$\begin{aligned} \bigoplus_{\alpha_0 \cdots \alpha_s} \text{diff}(\Sigma^{n+m} \mathcal{E}, \text{ch})^{q+t} ((U \times U)_{\alpha_0 \cdots \alpha_s}) & \\ \simeq \text{diff}(\Sigma^{n+m} \mathcal{E}, \text{ch})^{q+t} (\bigvee_{\alpha_0 \cdots \alpha_s} (U \times U)_{\alpha_0 \cdots \alpha_s}) & \\ \simeq \text{diff}(\Sigma^{n+m} \mathcal{E}, \text{ch})^{q+t} (\bigvee_{\alpha_0 \cdots \alpha_p} \bigvee_{\alpha_0 \cdots \alpha_r} \bigvee_{p+r=s} (U_{\alpha_0 \cdots \alpha_p} \times U_{\alpha_0 \cdots \alpha_r})) & \\ \simeq \bigoplus_{\alpha_0 \cdots \alpha_p} \bigoplus_{\alpha_0 \cdots \alpha_r} \bigoplus_{p+r=s} \text{diff}(\Sigma^{n+m} \mathcal{E}, \text{ch})^{q+t} (U_{\alpha_0 \cdots \alpha_p} \times U_{\alpha_0 \cdots \alpha_r}) & \end{aligned}$$

given by decomposing the product of the cover $\{U_\alpha\}$ with itself. Finally, we can pullback by the diagonal map

$$\begin{aligned} \Delta^*: \bigoplus_{\alpha_0 \cdots \alpha_s} \text{diff}(\Sigma^{n+m} \mathcal{E}, \text{ch})^{q+t} ((U \times U)_{\alpha_0 \cdots \alpha_s}) & \\ \rightarrow \bigoplus_{\alpha_0 \cdots \alpha_s} \text{diff}(\Sigma^{n+m} \mathcal{E}, \text{ch})^{q+t} (U_{\alpha_0 \cdots \alpha_s}) \simeq \check{C}^{p+r}(M; \text{diff}(\Sigma^{n+m} \mathcal{E}, \text{ch})^{q+t}). & \end{aligned}$$

The cup product on the E_1 -page is defined by the composite map $\Delta^* \times$.

Lemma 36 *The differential d_1 on the E_1 -page satisfies the Leibniz rule.*

Proof The construction of the cup product on the E_1 -page is precisely the cup product structure for Čech cohomology. The Čech differential satisfies the Leibniz rule, and this is precisely d_1 by construction. □

We are now ready to prove Proposition 35.

Proof The proof follows by induction on the pages of the spectral sequence. The base case is satisfied by Lemma 36. Now suppose we have a cup product map

$$\cup: E(n)_k \times E(n)_k \rightarrow E(n+m)_k$$

such that d_k satisfies Leibniz. By definition, we have

$$E(n)_{k+1}^{p,q} = \frac{\ker(d_k: E(n)_k^{p,q} \rightarrow E(n)_k^{p+k,q+k-1})}{\text{im}(d_k: E(n)_{k-1}^{p-k,q-k+1} \rightarrow E(n)_k^{p,q})},$$

and we define the cup product

$$\cup: E(n)_{k+1}^{p,q} \times E(m)_{k+1}^{r,s} \rightarrow E(n+m)_{k+1}^{p+r,q+s}$$

by restricting to elements in the kernel of d_k . The product is well defined since d_k satisfies the Leibniz rule. At this stage, the problem looks formally like the classical problem. Hence, analogously to the classical discussion in [36], it is tedious but straightforward to show that d_{k+1} also satisfies the Leibniz rule. □

4 Applications to differential cohomology theories

In this section, we would like to apply the spectral sequence constructed in the previous section to various differential cohomology theories. The construction is general enough to apply to any structured cohomology theory whose coefficients are known. We will explicitly emphasize three main examples. The first two are to known theories, namely ordinary differential cohomology and differential K-theory. We take this opportunity to explicitly develop the third theory, which is differential Morava K-theory, and then apply our smooth AHSS construction to it.

4.1 Ordinary differential cohomology theory

We begin by recovering the usual hypercohomology spectral sequence for the Deligne complex (see [11; 24, Appendix]) using our methods. We will first look at manifolds, then products of these, and then more generally to smooth fiber bundles.

Let us consider the smooth spectrum $\text{diff}(\Sigma^n H\mathbb{Z}, \text{ch})$ representing differential cohomology in degree n . We would like to see what our smooth AHSS gives in this case. We recall that $\text{diff}(\Sigma^n H\mathbb{Z}, \text{ch})$ is represented by Deligne cohomology of the sheaf of chain complexes $\mathbb{Z}_{\mathcal{D}}^{\infty}(n)$ via the Eilenberg–Mac Lane functor $H: \text{Sh}_{\infty}(\text{CartSp}; \text{Ch}) \rightarrow \text{Sh}_{\infty}(\text{CartSp}; \text{Sp})$ (2-6). It follows from the general properties of this functor that the homotopy groups are given by

$$\pi_k \text{diff}(\Sigma^n H\mathbb{Z}, \text{ch}) \simeq H_k \mathbb{Z}_{\mathcal{D}}^{\infty}(n).$$

In this case, we have the immediate corollary to Theorem 25.

Corollary 37 *The spectral sequence for Deligne cohomology takes the form*

$$E_2^{p,q} = H^p(X; H_{-q} \mathbb{Z}_{\mathcal{D}}^{\infty}(n)) \implies H^{p+q}(X; \mathbb{Z}_{\mathcal{D}}^{\infty}(n)),$$

which is essentially the hypercohomology spectral sequence for the Deligne complex, but shifted as a fourth quadrant spectral sequence.

For the sake of completeness, we work out this spectral sequence and recover the differential cohomology diamond (2-2) from the sequence. This will help to illustrate how the general spectral sequence behaves and how it can be used to calculate general differential cohomology groups.

Now over the site of Cartesian spaces, the Poincaré lemma implies that we have an isomorphism of presheaves $d: \Omega^{n-1} / \text{im}(d) \xrightarrow{\cong} \Omega_{\text{cl}}^n$. Since Ω_{cl}^n is a sheaf over the site of smooth manifolds, the gluing condition allows us to calculate the relevant terms on the E_2 -page of the spectral sequence in Figure 1.




Figure 1: $E_2^{p,q}$ from Corollary 37

The term $H^{n-1}(M; U(1))$ will survive to the E_∞ -page, and we have an isomorphism

$$H^{n-1}(M; U(1)) \simeq F_{n-1} \hat{H}^n(M; \mathbb{Z}) / F_n \hat{H}^n(M; \mathbb{Z}).$$

In fact, it is not hard to see that the definition of the filtration gives $F_n \hat{H}^n(M; \mathbb{Z}) \simeq 0$, and we have an injection

$$H^{n-1}(M; U(1)) \simeq F_{n-1} \hat{H}^n(M; \mathbb{Z}) \hookrightarrow \hat{H}^n(M; \mathbb{Z}).$$

On the E_n -page, we get one possibly nonzero differential

$$d_n: \Omega^n(M)_{cl} \rightarrow H^n(M; U(1)).$$

Proposition 38 *The differential d_n for the AHSS for Deligne cohomology can be identified with the composition*

$$\Omega_{cl}^n(M) \rightarrow H_{dR}^n(M) \xrightarrow{J_{\Delta^n}} H^n(M; \mathbb{R}) \xrightarrow{\exp} H^n(M; U(1)),$$

and the kernel is precisely those forms which have integral periods.

Proof We will unpack the definition of the differential in the AHSS in detail. This in turn will require unpacking the connecting homomorphism in the Deligne model of ordinary differential cohomology; see [11]. Denote by X_p the Čech filtration, and let

$$\partial: \text{diff}(\Sigma^n H\mathbb{Z}, \text{ch})^q(X_p) \rightarrow \text{diff}(\Sigma^n H\mathbb{Z}, \text{ch})^{q+1}(X_{p+1}/X_p)$$

denote the connecting homomorphism in the long exact sequence associated to the cofiber sequence $X_p \hookrightarrow X_{p+1} \twoheadrightarrow X_{p+1}/X_p$ in the usual way. In what follows, we will denote Čech–Deligne cochains on the p -level of the filtration X_p as a p -tuple

$$(z_0, z_1, \dots, z_p) \in \hat{C}^q(X_p),$$

where z_i is a $(q-i)$ -form defined on i -fold intersections.

Now, by definition, $d_n: E_n^{0,0} \rightarrow E_n^{n,0}$ is given by $d_n = \partial(j^*)^{-1}$, where $(j^*)^{-1}$ denotes a choice of element in the preimage of the restriction j^* induced by $j: X_0 \hookrightarrow X_{n-1}$.³ Since we have $d_k = 0$ for $k < n$, the differential d_n is defined on all elements $z \in \Omega_{cl}^n(M)$. Let g_0 be a locally defined $(n-1)$ -form trivializing z . Then we can choose $(j^*)^{-1}z$ to be the Čech–Deligne cocycle

$$(4-1) \quad (j^*)^{-1}z = \underbrace{(g_0, g_1, g_2, \dots, g_{n-2})}_{n-1} \in \widehat{C}^0(X_{n-1}),$$

where each g_k is a $(n-k-1)$ -form that satisfies the cocycle condition $\delta(g_k) = (-1)^k dg_{k+1}$. To see where the boundary map takes this element, let y be a Čech–Deligne cochain given by

$$y = \underbrace{(g_0, g_1, g_2, \dots, g_{n-2}, \exp(2\pi i g_{n-1}))}_n \in \widehat{C}^0(X_n),$$

where g_{n-1} is any smooth \mathbb{R} -valued function satisfying $d(g_{n-1}) = (-1)^{n-1}\delta(g_{n-2})$.⁴ Now y is not Čech–Deligne closed in general since

$$Dy = (d + (-1)^{n-1}\delta)y = (0, 0, \dots, \exp((-1)^{n-1}2\pi i \cdot \delta(g_{n-1}))),$$

and g_{n-1} may not satisfy the cocycle condition $\delta(g^{n-1}) = 0$. However, by the Čech–de Rham isomorphism (see for example [7]), this element in the Čech–de Rham double complex is isomorphic to an \mathbb{R} -valued Čech cocycle on n -fold intersections. Explicitly, there is a constant \mathbb{R} -valued cocycle r_n such that $\delta(g^{n-1}) = r_n$. It follows from the isomorphisms between the Čech, de Rham, and singular cohomologies that the class of r_n can be represented by the singular cocycle given by the pairing $\int_\sigma z$ for any cycle σ in M . Since the class $\int_\sigma z$ was just an unraveling of the boundary $\partial((j^*)^{-1}z)$, we have proved the claim. \square

In the next section, we will need to make use of a differential refinement of the Chern character. To this end, we briefly discuss differential cohomology with rational coefficients $\widehat{H}^n(-; \mathbb{Q})$. These groups are obtained via the differential function spectra $\text{diff}(\Sigma^n H\mathbb{Q}, \text{ch})$ which fit into the homotopy cartesian square:

$$\begin{array}{ccc} \text{diff}(H\mathbb{Q}, \text{ch}) & \longrightarrow & H(\tau_{\leq 0}\Omega^*[n]) \\ \downarrow & & \downarrow \\ \Sigma^n H\mathbb{Q} & \longrightarrow & H(\Omega^*[n]) \end{array}$$

³Note that the differential only takes this form at the $(0, 0)$ -entry. In general, the differential formed from the n^{th} derived couple will be more complicated.

⁴Note that this cocycle condition is necessary for y to be an lift of $(j^*)^{-1}z$ to the n -level of the filtration.

As a consequence of Proposition 10, the cohomology groups with values in this spectrum are calculated as

$$\text{diff}(\Sigma^n H\mathbb{Q}, \text{ch})^q(M) = \begin{cases} H^{n+q}(M), & q > 0, \\ \hat{H}^n(M; \mathbb{Q}), & q = 0, \\ H^{n-1+q}(M; \mathbb{R}/\mathbb{Q}), & q < 0. \end{cases}$$

The explicit calculation of the differential in Proposition 38 can be easily modified to get the following.⁵

Proposition 39 *The differential d_n on the E_n -page for the AHSS spectral sequence for $\text{diff}(\Sigma^n H\mathbb{Q}, \text{ch})$ is given by*

$$\Omega_{\text{cl}}^n(M) \rightarrow H_{\text{dR}}^n(M) \xrightarrow{\int_{\Delta^n}} H^n(M; \mathbb{R}) \rightarrow H^n(M; \mathbb{Q}/\mathbb{Z}),$$

and the kernel is precisely those forms which have rational periods.

We will make use of this result when we discuss the differentials in smooth K-theory in the next section. For now, from Proposition 38, we immediately get the following characterization of closed forms with integral periods and forms with rational periods using our smooth AHSS.

Corollary 40 (i) *The group of closed forms with integral periods on a manifold M is given by*

$$\Omega_{\text{cl}, \mathbb{Z}}^n(M) \simeq \hat{H}^n(M; \mathbb{Z}) / F_1 \hat{H}^n(M; \mathbb{Z}).$$

(ii) *The group of closed forms with rational periods on a manifold M is given by*

$$\Omega_{\text{cl}, \mathbb{Q}}^n(M) \simeq \hat{H}^n(M; \mathbb{Q}) / F_1 \hat{H}^n(M; \mathbb{Q}).$$

4.2 Differential K-theory

In this section, we examine the smooth AHSS for the differential function spectrum $\text{diff}(K, \text{ch})$, corresponding to complex K-theory. Proposition 10 allows us to calculate the cohomology groups on a paracompact manifold M as (see [49; 17; 66; 28])

$$(4-2) \quad \text{diff}(K, \text{ch})^q(M) = \begin{cases} K^q(M), & q > 0, \\ \hat{K}^0(M), & q = 0, \\ K_{U(1)}^q(M), & q < 0. \end{cases}$$

Both groups K and $K_{U(1)}$ are periodic. Indeed, $K_{U(1)}(M)$ fits into an exact sequence

$$\dots \rightarrow K^{-1}(M) \otimes \mathbb{R} \rightarrow K_{U(1)}^{-1}(M) \rightarrow K(M) \rightarrow K(M) \otimes \mathbb{R} \rightarrow \dots .$$

⁵The exact argument in the proof of Proposition 38 applies, with \mathbb{R}/\mathbb{Q} in place of $\mathbb{R}/\mathbb{Z} \simeq U(1)$.

Consequently, the periodicity of both integral and rational K-theory, along with an application of the five lemma, imply that $K_{U(1)}$ is 2–periodic. In particular, we have

$$K_{U(1)}^{2q}(\ast) \simeq U(1) \quad \text{and} \quad K_{U(1)}^{2q+1}(\ast) \simeq 0, \quad q \in \mathbb{Z}.$$

Given (4-2), we see that for a contractible open set U , we have an isomorphism

$$\text{diff}(K, \text{ch})^{2q+1}(U) \simeq K_{U(1)}^{2q}(\ast) \simeq U(1)$$

for $q < 0$. For degree 0, the differential cohomology diamond in this case takes the form:

$$\begin{array}{ccccc}
 & \prod_{2k-1} \Omega^{2k-1} / \text{im}(d) & \xrightarrow{d} & \prod_{2k} \Omega_{\text{cl}}^{2k} & \\
 & \searrow a & & \nearrow R & \\
 K_{\mathbb{R}}^{-1} & & \hat{K}^0 & & K_{\mathbb{R}}^0 \\
 & \nearrow & \searrow I & & \nearrow \text{ch} \\
 & K_{U(1)}^{-1} & \xrightarrow{\beta_K} & K^0 & \\
 & & & &
 \end{array}$$

This implies that for a contractible open set U , differential K-theory $\hat{K}^0(U)$ fits into the short exact sequence

$$0 \rightarrow \prod_{2k-1} \Omega^{2k-1} / \text{im}(d)(U) \rightarrow \hat{K}^0(U) \rightarrow \mathbb{Z} \rightarrow 0.$$

Hence, over the site of Cartesian spaces, we have a naturally split short exact sequence of presheaves

$$0 \rightarrow \prod_{2k-1} \Omega^{2k-1} / \text{im}(d) \rightarrow \hat{K}^0 \rightarrow \underline{\mathbb{Z}} \rightarrow 0.$$

Over that site, the presheaf on the left-hand side is actually a sheaf and is naturally isomorphic (by the Poincaré lemma) to the sheaf $\prod_{2k} \Omega_{\text{cl}}^{2k}$. We therefore make the identification

$$(4-3) \quad \hat{K}^0 \simeq \prod_{2k} \Omega_{\text{cl}}^{2k} \oplus \underline{\mathbb{Z}}.$$

Remark 41 It is important to note that the identification (4-3) is only true on the site of *Cartesian spaces*, which is to say that it holds only locally. On the site of smooth manifolds, this is of course not the case.

Next, since both Ω_{cl}^{2k} and $\underline{\mathbb{Z}}$ are sheaves on the site of smooth manifolds, we can identify the degree-0 Čech cohomology with these coefficients with the value of this




Figure 2: $E_2^{p,q}$ for even differential K-theory

sheaf on M . Isolating the terms on the E_2 -page which converge to $\widehat{K}^0(M)$, we get Figure 2.

We see that all the differentials are zero except for the map labeled d_2 above. On the E_3 -page, we get Figure 3.

The higher pages will fall into cases depending on the parity. We observe that for each even page E_{2m} , there is one nonzero differential given by d_{2m} . For the odd pages the differentials are given by an odd-degree $U(1)$ -cohomology operation.

Note that, in the diagrams, we are interested in the case $p + q = 0$, corresponding to diagonal entries. Now $p \geq 0$, as the Čech filtrations are of nonnegative degrees, which implies that $q \leq 0$. Hence the entries go down the diagonal. Our first goal will be to identify the even differentials d_{2m} . In order to do this, let us recall that there is a differential Chern character map (see [13; 63]) which is stably given by a morphism of smooth spectra

$$\widehat{\text{ch}}: \text{diff}(K, \text{ch}) \rightarrow \prod_{2k} \text{diff}(\Sigma^{2k} H\mathbb{Q}, \text{ch}).$$

Postcomposing this map with the projection pr_{2m} onto the $2m$ -component gives a map of smooth spectra

$$\text{pr}_{2m} \widehat{\text{ch}}: \text{diff}(K, \text{ch}) \rightarrow \text{diff}(\Sigma^{2m} H\mathbb{Q}, \text{ch}).$$

Using this map, we can prove the following analogue of Proposition 39.

Proposition 42 *The group of permanent cycles in bidegree $(0, 0)$ in the AHSS for $\text{diff}(K, \text{ch})$ is a subgroup of even-degree closed forms with rational periods. That is, we have*

$$E_\infty^{0,0} \subset \prod_k \Omega_{\text{cl},\mathbb{Q}}^{2k}(M) \oplus \mathbb{Z}.$$




Figure 3: $E_3^{p,q}$ for even differential K-theory

Proof We prove by induction on the even pages⁶ of the spectral sequence that, for all n , $E_{2n}^{0,0}$ must be a subgroup of

$$\prod_{2k \leq 2n} \Omega_{cl, \mathbb{Q}}^{2k}(M) \oplus \prod_{2k > 2n} \Omega_{cl}^{2k}(M) \oplus \mathbb{Z}.$$

For the base case, observe that the map $\text{pr}_2 \widehat{\text{ch}}$ induces a rank-1 morphism of AHSSs and therefore commutes with d_2 . It is straightforward to check, using the definitions, that this leads to the following commutative diagram:

$$\begin{array}{ccc} \prod_{2k} \Omega_{cl}^{2k}(M) \oplus \mathbb{Z} & \xrightarrow{\text{pr}_2} & \Omega_{cl}^2(M) \\ \downarrow d_2 & & \downarrow d'_2 \\ H^2(M; \mathbb{R}/\mathbb{Z}) & \xrightarrow{q} & H^2(M; \mathbb{R}/\mathbb{Q}) \end{array}$$

We see that the kernel of d_2 must be a subgroup of $\Omega_{cl, \mathbb{Q}}^2(M) \oplus \prod_{2k > 2} \Omega^{2k}(M) \oplus \mathbb{Z}$ by Proposition 38.

Now suppose the claim is true for d_{2n} . Again, we have that $\text{pr}_{2n+2} \widehat{\text{ch}}$ commutes with d_{2n+2} , and we have the following commutative diagram:

$$\begin{array}{ccc} \ker(d_{2n}) & \xrightarrow{\text{pr}_{2n+2}} & \Omega_{cl}^{2n+2}(M) \\ \downarrow d_{2n+2} & & \downarrow d'_{2n+2} \\ H^{2n+2}(M; \mathbb{R}/\mathbb{Z}) & \xrightarrow{q} & H^{2n+2}(M; \mathbb{R}/\mathbb{Q}) \end{array}$$

⁶The differential is 0 for the odd pages, and so no generality is lost by restricting to the even pages.

By the induction hypothesis,

$$\ker(d_{2n}) \subset \prod_{2k \leq 2n} \Omega_{\text{cl}, \mathbb{Q}}^{2k}(M) \oplus \prod_{2k > n} \Omega_{\text{cl}}^{2k}(M) \oplus \mathbb{Z},$$

and the kernel of d_{2n+2} is as claimed. □

We now turn to the first odd differential d_3 . Recall that β and $\tilde{\beta}$ denote the Bockstein homomorphisms corresponding to the sequences $0 \rightarrow \mathbb{Z} \rightarrow \mathbb{R} \xrightarrow{\text{exp}} U(1) \rightarrow 0$ and $0 \rightarrow \mathbb{Z} \rightarrow \mathbb{Z} \rightarrow \mathbb{Z}/2 \rightarrow 0$, respectively. We still also denote by $\Gamma_2: H^n(-; \mathbb{Z}/2) \rightarrow H^n(-, U(1))$ the map induced by the representation of $\mathbb{Z}/2$ as the square roots of unity and $\rho_2: \mathbb{Z} \rightarrow \mathbb{Z}/2$ as the mod 2 reduction.

Proposition 43 (degree-3 differential) *The first odd-degree differential in the AHSS for differential K-theory is given by*

$$d_3 = \begin{cases} \widehat{\text{Sq}}^3 := \Gamma_2 \text{Sq}^2 \rho_2 \beta, & q < 0, \\ \text{Sq}_{\mathbb{Z}}^3 := \tilde{\beta} \text{Sq}^2 \rho_2, & q > 0, \\ 0, & q = 0. \end{cases}$$

Proof The case for $q = 0$ is obvious. For $q > 0$, this follows from the fact that the integration map defines an isomorphism $I: \text{diff}(K, \text{ch})^q(M) \xrightarrow{\cong} K^q(M)$ for $q > 0$. Since the differential d_3 for the classical AHSS is given by $\text{Sq}_{\mathbb{Z}}^3$, and the integration map defines an isomorphism of corresponding first quadrant spectral sequences, the case $q > 0$ is settled.

For $q < 0$, Corollary 30 implies that the Bockstein β commutes with the differentials on the E_3 -page. We therefore have

$$(4-4) \quad \beta d_3 = \text{Sq}_{\mathbb{Z}}^3 \beta = \tilde{\beta} \text{Sq}^3 \rho_2 \beta.$$

Rephrasing, we have the commuting diagram:

$$\begin{array}{ccc} H^{n-1}(M; U(1)) & \xrightarrow{d_3} & H^{n+3-1}(M; U(1)) \\ \beta \downarrow & & \downarrow \beta \\ H^n(M; \mathbb{Z}) & \xrightarrow{\text{Sq}_{\mathbb{Z}}^3} & H^{n+3}(M; \mathbb{Z}) \end{array}$$

We now claim that $\tilde{\beta} = \beta \circ \Gamma_2$. Indeed, we have a morphism of short exact sequences:

$$\begin{array}{ccccc} \mathbb{Z} & \xrightarrow{\times 2} & \mathbb{Z} & \xrightarrow{\rho_2} & \mathbb{Z}/2 \\ \downarrow \text{id} & & \downarrow \times \pi i & & \downarrow \Gamma_2 \\ \mathbb{Z} & \xrightarrow{\times 2\pi i} & \mathbb{R} & \xrightarrow{\text{exp}} & U(1) \end{array}$$

This morphism induces a morphism on the associated long exact sequences on cohomology. After delooping once to extend to the left, the homotopy commutativity of the resulting diagram

$$\begin{array}{ccc} \mathbb{Z}/2 & \xrightarrow{\tilde{\beta}} & B\mathbb{Z} \\ \Gamma_2 \downarrow & & \parallel \\ U(1) & \xrightarrow{\beta} & B\mathbb{Z} \end{array}$$

immediately establishes the claim.

Now it follows from (4-4) that $d_3 - \Gamma_2 \text{Sq}^3 \rho_2 \beta$ is in the kernel of β . By exactness of the Bockstein, this implies that it must be in the image of the exponential map $\text{exp}: H^*(-; \mathbb{R}) \rightarrow H^*(-; U(1))$. Hence there is an operation $\psi: H^*(-; U(1)) \rightarrow H^*(-; \mathbb{R})$ such that

$$\phi := \text{exp} \circ \psi = \text{exp}(\psi) = d_3 - \Gamma_2 \text{Sq}^3 \rho_2 \beta.$$

Equivalently, we have a factorization:

$$\begin{array}{ccc} H^*(-; U(1)) & \xrightarrow{\phi} & H^*(-; U(1)) \\ & \searrow \psi & \nearrow \text{exp} \\ & H^*(-; \mathbb{R}) & \end{array}$$

We expect to have $\text{hom}(H^*(-; U(1)), H^*(-; \mathbb{R})) = 0$ since $U(1)$ is almost completely torsion (and since the second argument is an \mathbb{R} -vector space). However, we need to be slightly careful here, since not all elements $a \in H^*(M; U(1))$ represent torsion classes. In fact, identifying $U(1) \simeq \mathbb{R}/\mathbb{Z}$, such an element will be torsion if and only if it represents an element in $H^*(M; \mathbb{Q}/\mathbb{Z}) \hookrightarrow H^*(M; \mathbb{R}/\mathbb{Z})$. To fix this issue, we observe that for any abelian group A , we have an isomorphism

$$r: \text{hom}(H^*(M; \mathbb{R}), A) \rightarrow \text{hom}(H^*(M; \mathbb{Q}), A)$$

given by restricting a map to the rationals $\mathbb{Q} \subset \mathbb{R}$. The inverse is given by restricting a map to the generators and extending with real coefficients. This implies, in turn, that we have an isomorphism

$$r: \text{hom}(H^*(M; \mathbb{R}/\mathbb{Z}), A) \rightarrow \text{hom}(H^*(M; \mathbb{Q}/\mathbb{Z}), A);$$

ie \mathbb{R}/\mathbb{Z} and \mathbb{Q}/\mathbb{Z} behave equivalently when taken as coefficients of cohomology inside the hom. Finally, since $\text{hom}(H^*(M; \mathbb{Q}/\mathbb{Z}), H^*(M; \mathbb{R})) = 0$, we must have $\text{hom}(H^*(M; \mathbb{R}/\mathbb{Z}), H^*(M; \mathbb{R})) = 0$, which forces $\psi = 0$. Consequently, $\text{exp} \circ \psi = 0$, so that $\phi = 0$. Therefore, indeed we have

$$d_3 = \Gamma_2 \text{Sq}^3 \rho_2 \beta. \quad \square$$

Remark 44 The above proposition suggests that these operations are related to some sort of *differential Steenrod squares*. Indeed, this is the case, which has been investigated by the authors in [33], with \widehat{Sq}^3 being one such operation.

Now that we have established the algebraic construction, we turn to investigating the convergence of the spectral sequence from a geometric point of view. In particular, we immediately observe that the only terms in the spectral sequence which contain information about differential forms are at $q = 0$. These terms converge to elements in the filtered graded complex (since $q = 0$)

$$\widehat{K}(M)/F_1 \widehat{K}(M).$$

Since the filtration is given by the Čech-type filtration on M , we see that this quotient contains elements which have nontrivial data on all open sets, intersections and higher intersections. For the degrees $q < 0$, the filtration quotients

$$F_p \widehat{K}(M)/F_{p+1} \widehat{K}(M)$$

have trivial data below p -intersections.

In fact, it is not too surprising that this occurs. There is a geometric model for reduced \widehat{K}^0 which is given by the moduli stack $\mathbf{BU}_{\text{conn}}$ of unitary vector bundles, equipped with connection. Let \mathbf{Vect}_{∇} be the moduli stack of vector bundles with connections. It was shown in [16] that there is a cycle map

$$\text{cycl: } \pi_0 \text{Map}(M, \mathbf{Vect}_{\nabla}) \rightarrow \widehat{K}^0(M),$$

which induces an isomorphism upon group completion. In our construction, this is equivalent to

$$\text{cycl: } \pi_0 \text{Map}(M, \mathbf{BU}_{\text{conn}}) \rightarrow \widehat{K}^0(M).$$

Now the stack $\mathbf{BU}_{\text{conn}}$ can be identified with the moduli stack obtained by taking the nerve of the action groupoid $C^\infty(-, U)/\Omega^1(-; \mathfrak{u})$ with the action given by gauge transformations, where \mathfrak{u} is the Lie algebra of the unitary group. Let $\{U_\alpha\}$ be a good open cover of M . Then a map $M \rightarrow \mathbf{BU}_{\text{conn}}$ is given by the following data:

- a choice of smooth $U(n)$ -valued function $g_{\alpha\beta}$ on intersections $U_\alpha \cap U_\beta$,
- a choice of local connection 1-form $\mathcal{A}_{\alpha\beta}$ on open sets U_α .

This is precisely the data needed to define a unitary vector bundle on M .

Remark 45 More relevant to our needs, though, is the fact that the effects of the filtration become transparent when taking $\mathbf{BU}_{\text{conn}}$ as a model for \widehat{K}^0 . We now see that the $q = 0$ terms converge to terms which involve the data of the connection, while the $q < 0$ terms contain data about bundles with trivializable connections (in particular, flat connections).

Differential K^1 -theory We now consider odd differential K-theory, K^1 . In this case, the representing spectrum is the unitary group U itself. Viewing this as a classifying space, we can write $U = B\Omega U$. Of course we are interested in the corresponding stacks. Unfortunately, we do not have the analogue of the above group-loop group relation in stacks; ie $U_{\text{conn}} \not\cong \mathbf{B}\Omega U_{\text{conn}}$. Nevertheless, the machinery that we set up will work equally well for differential K^1 -theory, as far as the third differential goes; ie we still have $d_3 = \widehat{S}q^3$. However, the even differentials are now transgressed in degree by one, so that they are also of odd degree. This is expected as the Chern character in this case is a map to cohomology of odd degree.

The story for \widehat{K}^1 can be worked out similarly as we indicated above. Let us expand on this in more detail. In the odd case, the differential cohomology diamond takes the form

$$\begin{array}{ccccc}
 & \prod_{2k} \Omega^{2k} / \text{im}(d) & \xrightarrow{d} & \prod_{2k+1} \Omega_{\text{cl}}^{2k+1} & \\
 & \nearrow a & & \nearrow R & \\
 K_{\mathbb{R}}^0 & & \widehat{K}^1 & & K_{\mathbb{R}}^1 \\
 & \searrow & \nearrow I & \searrow \text{ch} & \\
 & K_{U(1)}^0 & \xrightarrow{\beta_K} & K^1 &
 \end{array}$$

and we get a short exact sequence of presheaves (on the site of Cartesian spaces)

$$0 \rightarrow \mathbb{Z} \rightarrow \prod_{2k} \Omega^{2k} / \text{im}(d) \rightarrow \widehat{K}^1 \rightarrow 0.$$

It is straightforward to show that the map $\mathbb{Z} \rightarrow \prod_{2k} \Omega^{2k} / \text{im}(d)$ is zero. Consequently, we have the isomorphism

$$\widehat{K}^1 \simeq \prod_{2k} \Omega^{2k} / \text{im}(d) \simeq \prod_{2k+1} \Omega_{\text{cl}}^{2k+1}.$$

Using the same type of argument as in the even K-theory K^0 , we likewise get a refinement of the differential of the underlying topological theory. More precisely, we see that the first nonzero differentials appear on the E_3 -page as in Figure 4.

Proposition 46 *Proposition 43 holds for differential K^1 -theory. That is, the degree-3 differential in \widehat{K}^1 is given by the refinement of the Steenrod square of dimension three.*

Also, using the same argument as in the proof of Proposition 42, we see that the permanent cycles in bidegree $(0, 0)$ are a subgroup of odd-degree forms with rational periods.




Figure 4: $E_2^{p,q}$ for odd differential K-theory

Proposition 47 *The group of permanent cycles in bidegree $(0, 0)$ in the AHSS for $\text{diff}(\Sigma K, \text{ch})$ is a subgroup of odd degree closed forms with rational periods. That is, we have*

$$E_\infty^{0,0} \subset \prod_k \Omega_{cl, \mathbb{Q}}^{2k-1}(M) \oplus \mathbb{Z}.$$

Example 48 (fields in string theory and M-theory) In the string theory and M-theory literature, one encounters settings where cohomology classes are compared to K-theory elements, in the sense of asking when a cohomology class arises from or “lifts to” a K-theory class. This involves, in a sense, a physical modeling of the process of building the AHSS. One such obstruction is Sq^3 , viewed as the first nontrivial differential d_3 in K-theory, so that the condition $Sq^3 x = 0$ on a cohomology class x amounts to saying that the class lifts to K-theory. This is desirable in the study of the partition function of the fields in type-IIA string theory; see [21; 47]. On the other hand, it is desirable to have differential refinements for physical purposes. Therefore, now that we have the differential AHSS at our disposal, it is natural to consider expressions such as $d_3(\widehat{x}) := \widehat{Sq}^3 \widehat{x} = 0$ on the differential cohomology class \widehat{x} that refines the topological class x . This can be viewed as a condition on cohomology with $U(1)$ -coefficients (or flat n -bundles), in order that they lift to flat elements in \widehat{K} .⁷ If the degree of the class x is even, then we are in type-IIA string theory, and we lift to differential K^0 -theory. On the other hand, being in type-IIB string theory means the degree of x is odd, and we are lifting to differential K^1 -theory. The new differentials d_{2m} and d_{2m+1} arising from differential forms will correspond to even- and odd-degree closed differential forms as the particular forms representing the physical fields F_{2m} and F_{2m+1} via the Chern character.

⁷This could end up being stronger in the sense that it is a condition for lifting differential cohomology classes to differential K-theory, but we will leave that for future investigations.

Example 49 (D-brane charges) The charges of D-branes can, a priori, be taken to be given as a class in cohomology $\mathcal{Q}_H \in H^*(X; \mathbb{Q})$. Quantum effects requires some of these charges to be (up to shifts) to be in integral cohomology. However, in order to not discuss isomorphism classes of such physical objects but pinning down a particular physical object, one considers the charges to take values in differential cohomology, with Deligne cohomology being one such presentation: $\mathcal{Q}_{\widehat{H}} \in \widehat{H}^*(X; \mathbb{Z})$; see [19]. On the other hand, careful analysis reveals that the charges take values in K-theory rather than in cohomology $\mathcal{Q}_K \in K^i(X)$ for $i = 0, 1$ for type IIB/IIA; see [56; 29; 9]. Such a class exists if the cohomology charge satisfies $\text{Sq}^3 \mathcal{Q}_H = 0$. Again, at this stage, adding in the geometry requires the charges to take values in differential K-theory $\mathcal{Q}_{\widehat{K}} \in \widehat{K}^i(X)$. Our construction now allows for a characterization of when charges in Deligne cohomology lift to charges in differential K-theory, namely when they are annihilated by the third differential in the smooth AHSS, ie when $\widehat{\text{Sq}}^3 \mathcal{Q}_{\widehat{H}} = 0$.

4.3 Differential Morava K-theory

There are various interesting generalized cohomology theories that descend from complex cobordism, among which are Morava K-theory and Morava E-theory. Such theories can be defined using their coefficient rings, which in general are polynomials over finite or p -adic fields on generators whose dimension depends on the chromatic level and the prime p . As such, these kind of theories do not lend themselves directly to immediate geometric interpretation in contrast to the case of K-theory, which can be formulated via stable isomorphism classes of vector bundles.

However, recent work in [48] (generalizing some aspects of [5]) seems to give hope in that direction. Nevertheless, just because an entity is defined over a finite field does not automatically make it ineligible for differential refinement. In fact, we have recently demonstrated this [33] for the case of Steenrod cohomology operations, which are, a priori, \mathbb{Z}/p -valued operations. The main point there was that as long as these admit integral lifts, they do have a chance at a differential refinement. What we will seek here is something analogous: integral refinements of such generalized cohomology theories.

We will consider the integral Morava K-theory $\widetilde{K}(n)$ highlighted in [47; 60; 62]. Morava K-theory $K(n)$ is the mod p reduction of an integral (or p -adic) lift $\widetilde{K}(n)$ with coefficient ring $\widetilde{K}(n)_* = \mathbb{Z}_p[v_n, v_n^{-1}]$. This theory more closely resembles complex K-theory than is the case for the mod p versions (for $n = 1$, it is the p -completion of K-theory). The integral theory is much more suited to applications in physics [47; 60; 12; 62].

The Atiyah–Hirzebruch spectral sequence for Morava K-theory has been studied by Yagita in [72]; see also [47]. There is a spectral sequence converging to $K(n)^*(X)$ with E_2 -term $E_2^{p,q} = H^p(X, K(n)^q)$. While this can be done for any prime, we will

focus on the prime 2. In this case, the first possibly nontrivial differential is $d_{2^{n+1}-1}$; this is given by [72] as

$$d_{2^{n+1}-1}(xv_n^k) = Q_n(x)v_n^{k-1}.$$

Here Q_n is the n^{th} Milnor primitive at the prime 2, which we define inductively as $Q_0 = \text{Sq}^1$, the Bockstein operation, and $Q_{j+1} = \text{Sq}^{2^j} Q_j - Q_j \text{Sq}^{2^j}$, where $\text{Sq}^j: H^n(X; \mathbb{Z}_2) \rightarrow H^{n+j}(X; \mathbb{Z}_2)$ is the j^{th} Steenrod square. These operations are derivations

$$Q_j(xy) = Q_j(x)y + (-1)^{|x|} x Q_j(y).$$

The signs are of course irrelevant at $p = 2$, but will become important in the integral version. Extensive discussion of the mod p Steenrod algebra in terms of these operations is given in [68].

The integral theory is also computable via an AHSS, which can be deduced from [47; 62]. There is an AHSS converging to $\tilde{K}(n)^*(X)$ with $E_2^{p,q} = H^p(X, \tilde{K}(n)^q)$. The first possibly nontrivial differential is $d_{2^{n+1}-1}$; this is given by

$$d_{2^{n+1}-1}(xv_n^k) = \tilde{Q}_n(x)v_n^{k-1}.$$

Here $\tilde{Q}_k: H^*(X; \mathbb{Z}) \rightarrow H^{*+2^{k+1}-1}(X; \mathbb{Z})$ is an integral cohomology operation lifting the Milnor primitive Q_k .

In order to consider differential refinement of Morava K-theory, we need geometric information encoded in differential forms, hence rational information. The rationalization of Morava K-theory $\tilde{K}(n)$, like any reasonable spectrum, exists and can be thought of as localization at $\tilde{K}(0) = H\mathbb{Q}$; see [8; 58]. We can, in the same way, localize at \mathbb{R} . More precisely, the localized theory is given by

$$\tilde{K}_{\mathbb{R}}(n) = \tilde{K}(n) \wedge M\mathbb{R},$$

where $M\mathbb{R}$ is an Eilenberg–Moore spectrum. We have an equivalence

$$\tilde{K}_{\mathbb{R}}(n) \simeq H(\mathbb{Z}[v_n, v_n^{-1}] \otimes \mathbb{R})$$

and a Chern character map

$$\text{ch}: \tilde{K}(n) \rightarrow H(\mathbb{Z}[v_n, v_n^{-1}] \otimes \Omega^*).$$

Thus we can form the differential function spectrum $\text{diff}(\tilde{K}(n), \text{ch})$, and we can form the associated AHSS. To see what form the spectral sequence takes, we need to discuss the flat Morava K-theory $\tilde{K}_{U(1)}(n)$, defined by the fiber sequence

$$\tilde{K}(n) \rightarrow \tilde{K}(n) \wedge M\mathbb{R} \rightarrow \tilde{K}_{U(1)}(n) := \tilde{K}(n) \wedge MU(1).$$




Figure 5: $E_{2(2^n-1)}^{p,q}$ for Morava K-theory

This theory is periodic with period $2(2^n - 1)$. Indeed, both $\tilde{K}(n)$ and its rationalization are periodic, and we have a long exact sequence

$$\dots \rightarrow \tilde{K}(n)^m(M) \rightarrow (\tilde{K}(n) \wedge M\mathbb{R})^m(M) \rightarrow \tilde{K}_{U(1)}^m(n)(M) \rightarrow \tilde{K}(n)^{m+1}(M) \rightarrow \dots$$

relating the flat theory to both the rational and integral theories. This, in particular, gives the following identification.

Lemma 50 *The coefficients of flat Morava K-theory are given by*

$$\tilde{K}_{U(1)}(n)^m(*) \simeq \begin{cases} U(1), & m = 2(2^n - 1), \\ 0, & \text{otherwise.} \end{cases}$$

Knowing the coefficients of the flat theory, we can write down the relevant nonzero terms on the $E_{2(2^n-1)}$ -page of the corresponding spectral sequence in Figure 5, and the only nonzero differential is given by

$$d_{2(2^n-1)}: \prod_k \Omega_{cl}^{k2(2^n-1)}(M) \oplus \mathbb{Z} \rightarrow H^{2(2^n-1)}(M; \mathbb{R}/\mathbb{Z}).$$

Just as in the case for differential K-theory (see Propositions 42 and 47), we have:

Proposition 51 *The group of permanent cycles in bidegree $(0, 0)$ in the AHSS for $\text{diff}(\tilde{K}(n), \text{ch})$ is a subgroup of certain closed forms with rational periods. More precisely, we have*

$$E_{\infty}^{0,0} \subset \prod_k \Omega_{cl, \mathbb{Q}}^{2k(2^n-1)}(M) \oplus \mathbb{Z}.$$

To identify the Čech cohomology groups with coefficients in $\widehat{K}(n)^0$, we make the identification (as we did for differential K-theory)

$$\widehat{K}(n)^0 \simeq \prod_k \Omega_{\text{cl}}^{2k(2^n-1)} \oplus \mathbb{Z}$$

on the site of Cartesian spaces. Again, using the sheaf condition over smooth manifolds, we have

$$H^P(M; \widehat{K}(n)^0) \simeq \prod_k \Omega_{\text{cl}}^{2k(2^n-1)}(M) \oplus \mathbb{Z}.$$

We now consider the differential refinement of the (integrally lifted) Milnor primitive. As before, let $\Gamma_2: H^n(-; \mathbb{Z}/2) \rightarrow H^n(-; U(1))$ denote the map induced by the representation of $\mathbb{Z}/2$ as the square roots of unity, and let $\rho_2: \mathbb{Z} \rightarrow \mathbb{Z}/2$ denote the mod 2 reduction.

Lemma 52 *The integral Milnor primitive \widetilde{Q}_n factors through the representation $\Gamma_2: \mathbb{Z}/2 \hookrightarrow U(1)$. That is, there exists an operation \widehat{Q}_n such that*

$$Q_n \rho_2 = \rho_2 \widetilde{Q}_n = \rho_2 \beta \Gamma_2 \widehat{Q}_n,$$

where β is the Bockstein for the exponential sequence.

Proof Recall first that $\rho_2 \beta \Gamma_2 = \rho_2 \widetilde{\beta} = \text{Sq}^1$, where $\widetilde{\beta}$ is the Bockstein for the mod 2 reduction sequence. We can therefore rewrite the above equation as

$$Q_n \rho_2 = \rho_2 \widetilde{Q}_n = \rho_2 \beta \Gamma_2 \widehat{Q}_n = \text{Sq}^1 \widehat{Q}_n,$$

and the existence of the class \widehat{Q}_n holds if and only if $\text{Sq}^1 Q_n \rho_2 = 0$. On the other hand, the existence of the integral lift \widetilde{Q}_n immediately implies this condition. \square

Again, let β and $\widetilde{\beta}$ denote the Bockstein homomorphism corresponding to the sequences $0 \rightarrow \mathbb{Z} \rightarrow \mathbb{R} \rightarrow \mathbb{R}/\mathbb{Z} \rightarrow 0$ and $0 \rightarrow \mathbb{Z} \rightarrow \mathbb{Z} \rightarrow \mathbb{Z}/2 \rightarrow 0$, respectively. Then the following can be proved in a similar way as we did for Proposition 43 in the case of differential K-theory.

Proposition 53 (odd differentials for Morava AHSS) *The $(2^{n+1}-1)$ -differential in the AHSS for differential Morava K-theory is given by*

$$d_{2^{n+1}-1} = \begin{cases} \Gamma_2 \widehat{Q}_n \rho_2 \beta, & q < 0, \\ \widetilde{Q}_n, & q > 0, \\ 0, & q = 0. \end{cases}$$

Remark 54 (odd primes) The above discussion has been for the prime 2; that is, we are considering integral Morava K-theory as arising from lifting of the $p = 2$ Morava

K-theory. We can do the same for odd primes, leading to integral Morava K-theory lifted from an odd prime p . A similar discussion follows and we have an integral lift of the Milnor primitive at odd primes, as in Lemma 52. The differentials will be again given by these refinements of the Milnor primitive; ie Proposition 53 holds except that the primitives are defined using the Steenrod reduced power operations P^j . Precisely, Q_0 is the Bockstein homomorphism associated to reduction mod p sequence, and inductively $Q_{i+1} = P^{p^i} Q_i - Q_i P^{p^i}$. The operations P^j have been differentially refined in [33]. Hence the refinement of the Milnor primitives at odd primes will also follow. Then the $(p^{n+1}-1)$ -differential in the AHSS for differential Morava K-theory is given by

$$d_{p^{n+1}-1} = \begin{cases} \Gamma_p \widehat{Q}_n \rho_p \beta, & q < 0, \\ \widetilde{Q}_n, & q > 0, \\ 0, & q = 0. \end{cases}$$

Example 55 (lifting fields to differential Morava K-theory) We build on Example 48 and aim to lift the cohomology classes beyond K-theory. In particular, for $x = \lambda = \frac{1}{2} p_1$ the first Spin characteristic class, we have $\widehat{x} = \widehat{\lambda}$ the differential refinements of λ [61; 25] (which can be viewed as a lifted Wu class [41]), and we would have $\widehat{Sq}^3 \widehat{\lambda} = 0$. This condition in differential cohomology can be viewed as a refinement of the condition $W_7 = Sq^3 \lambda = 0$ leading to orientation with respect to integral Morava $K(2)$ -theory (lifted from the prime $p = 2$) as shown in [47] and elaborated further in [12]. From the structure of the smooth AHSS in relation to the classical AHSS, one can extend various results to the differential case. For instance, one can generalize the statement in [47] on orientation to state that: *an oriented smooth 10-dimensional manifold is oriented with respect to differential (integrally lifted from $p = 2$) Morava $K(2)$ -theory $\widehat{K}(2)$ if the class $\widehat{W}_7 := \widehat{Sq}^3 \widehat{\lambda}$ is equal to 0.* The development of this, as well as the relation to refinements of characteristic classes, deserves a separate treatment and will be addressed elsewhere.

Remark 56 (i) Note that our construction allows for an AHSS for other spectra beyond the particular ones we discussed above. This holds for any spectrum which admits a rationalization, whose coefficients are known, and which can be lifted integrally in the sense that we discussed at the beginning of this section.

(ii) All the cohomology theories that we used in this paper can be twisted. Indeed, the construction in this paper can be generalized to construct an AHSS for twisted differential spectra [32], in the sense of [15], and using [34].

Acknowledgements The authors would like to thank Ulrich Bunke, Thomas Nikolaus and Craig Westerland for interesting discussions at the early stages of this project

and Urs Schreiber for very useful comments on the first draft. We are grateful to the anonymous referee for a careful reading and for useful suggestions.

References

- [1] **J F Adams**, *Stable homotopy and generalised homology*, Univ. of Chicago Press (1974) MR
- [2] **D Arlettaz**, *The order of the differentials in the Atiyah–Hirzebruch spectral sequence*, *K-Theory* 6 (1992) 347–361 MR
- [3] **M F Atiyah, F Hirzebruch**, *Vector bundles and homogeneous spaces*, from “Differential Geometry” (C B Allendoerfer, editor), *Proc. Sympos. Pure Math.* 3, Amer. Math. Soc., Providence, RI (1961) 7–38 MR
- [4] **M F Atiyah, F Hirzebruch**, *Analytic cycles on complex manifolds*, *Topology* 1 (1962) 25–45 MR
- [5] **N A Baas, B I Dundas, J Rognes**, *Two-vector bundles and forms of elliptic cohomology*, from “Topology, geometry and quantum field theory” (U Tillmann, editor), *London Math. Soc. Lecture Note Ser.* 308, Cambridge Univ. Press (2004) 18–45 MR
- [6] **A Björner**, *Topological methods*, from “Handbook of combinatorics, II” (R L Graham, M Grötschel, L Lovász, editors), *Elsevier Sci. B. V.*, Amsterdam (1995) 1819–1872 MR
- [7] **R Bott, L W Tu**, *Differential forms in algebraic topology*, *Graduate Texts in Mathematics* 82, Springer (1982) MR
- [8] **A K Bousfield**, *The localization of spectra with respect to homology*, *Topology* 18 (1979) 257–281 MR
- [9] **J Brodzki, V Mathai, J Rosenberg, R J Szabo**, *D-branes, RR-fields and duality on noncommutative manifolds*, *Comm. Math. Phys.* 277 (2008) 643–706 MR
- [10] **K S Brown**, *Abstract homotopy theory and generalized sheaf cohomology*, *Trans. Amer. Math. Soc.* 186 (1973) 419–458 MR
- [11] **J-L Brylinski**, *Loop spaces, characteristic classes and geometric quantization*, *Progress in Mathematics* 107, Birkhäuser, Boston (1993) MR
- [12] **L Buhné**, *Properties of integral Morava K-theory and the asserted application to the Diaconescu–Moore–Witten anomaly*, PhD thesis, Hamburg Univ. (2011)
- [13] **U Bunke**, *Differential cohomology*, preprint (2012) arXiv
- [14] **U Bunke, M Kreck, T Schick**, *A geometric description of differential cohomology*, *Ann. Math. Blaise Pascal* 17 (2010) 1–16 MR
- [15] **U Bunke, T Nikolaus**, *Twisted differential cohomology*, preprint (2014) arXiv
- [16] **U Bunke, T Nikolaus, M Völkl**, *Differential cohomology theories as sheaves of spectra*, *J. Homotopy Relat. Struct.* 11 (2016) 1–66 MR

- [17] **U Bunke, T Schick**, *Smooth K-theory*, Astérisque 328, Soc. Math. France, Paris (2009) 45–135 MR
- [18] **U Bunke, T Schick**, *Uniqueness of smooth extensions of generalized cohomology theories*, *J. Topol.* 3 (2010) 110–156 MR
- [19] **A L Carey, S Johnson, M K Murray**, *Holonomy on D-branes*, *J. Geom. Phys.* 52 (2004) 186–216 MR
- [20] **J Cheeger, J Simons**, *Differential characters and geometric invariants*, from “Geometry and topology” (J Alexander, J Harer, editors), *Lecture Notes in Math.* 1167, Springer (1985) 50–80 MR
- [21] **D-E Diaconescu, G Moore, E Witten**, *E_8 gauge theory, and a derivation of K-theory from M-theory*, *Adv. Theor. Math. Phys.* 6 (2002) 1031–1134 MR
- [22] **D Dugger**, *Combinatorial model categories have presentations*, *Adv. Math.* 164 (2001) 177–201 MR
- [23] **J L Dupont, R Ljungmann**, *Integration of simplicial forms and Deligne cohomology*, *Math. Scand.* 97 (2005) 11–39 MR
- [24] **H Esnault, E Viehweg**, *Lectures on vanishing theorems*, DMV Seminar 20, Birkhäuser, Basel (1992) MR
- [25] **D Fiorenza, U Schreiber, J Stasheff**, *Čech cocycles for differential characteristic classes: an ∞ -Lie theoretic construction*, *Adv. Theor. Math. Phys.* 16 (2012) 149–250 MR
- [26] **A T Fomenko, D B Fuchs, V L Gutenmacher**, *Homotopic topology*, Akadémiai Kiadó, Budapest (1986) MR
- [27] **D S Freed**, *Dirac charge quantization and generalized differential cohomology*, from “Surveys in differential geometry” (S-T Yau, editor), *Surv. Differ. Geom.* 7, International Press, Somerville, MA (2000) 129–194 MR
- [28] **D S Freed, J Lott**, *An index theorem in differential K-theory*, *Geom. Topol.* 14 (2010) 903–966 MR
- [29] **D S Freed, E Witten**, *Anomalies in string theory with D-branes*, *Asian J. Math.* 3 (1999) 819–851 MR
- [30] **P Gajer**, *Geometry of Deligne cohomology*, *Invent. Math.* 127 (1997) 155–207 MR
- [31] **A Gorokhovskiy, J Lott**, *A Hilbert bundle description of differential K-theory*, preprint (2015) arXiv
- [32] **D Grady, H Sati**, *AHSS for twisted differential spectra*, in preparation
- [33] **D Grady, H Sati**, *Primary operations in differential cohomology*, preprint (2016) arXiv
- [34] **D Grady, H Sati**, *Massey products in differential cohomology via stacks*, *J. Homotopy Relat. Struct.* (online publication May 2017)
- [35] **P A Griffiths, J W Morgan**, *Rational homotopy theory and differential forms*, *Progress in Mathematics* 16, Birkhäuser, Boston (1981) MR

- [36] **A Hatcher**, *Algebraic topology*, Cambridge Univ. Press (2002) MR
- [37] **P Hekmati, M K Murray, V S Schlegel, R F Vozzo**, *A geometric model for odd differential K-theory*, *Differential Geom. Appl.* 40 (2015) 123–158 MR
- [38] **P Hilton**, *General cohomology theory and K-theory*, London Math. Soc. Lecture Note Ser. 1, Cambridge Univ. Press (1971) MR
- [39] **M-H Ho**, *The differential analytic index in Simons–Sullivan differential K-theory*, *Ann. Global Anal. Geom.* 42 (2012) 523–535 MR
- [40] **M-H Ho**, *Remarks on flat and differential K-theory*, *Ann. Math. Blaise Pascal* 21 (2014) 91–101 MR
- [41] **M J Hopkins, I M Singer**, *Quadratic functions in geometry, topology, and M-theory*, *J. Differential Geom.* 70 (2005) 329–452 MR
- [42] **D Husemöller, M Joachim, B Jurčo, M Schottenloher**, *Basic bundle theory and K-cohomology invariants*, *Lecture Notes in Physics* 726, Springer (2008) MR
- [43] **J F Jardine**, *Stable homotopy theory of simplicial presheaves*, *Canad. J. Math.* 39 (1987) 733–747 MR
- [44] **J F Jardine**, *Local homotopy theory*, Springer (2015) MR
- [45] **M Karoubi**, *Homologie cyclique et K-théorie*, *Astérisque* 149, Soc. Math. France, Paris (1987) MR
- [46] **K R Klonoff**, *An index theorem in differential K-theory*, PhD thesis, The University of Texas at Austin (2008) MR Available at <http://search.proquest.com/docview/230710076>
- [47] **I Kriz, H Sati**, *M-theory, type IIA superstrings, and elliptic cohomology*, *Adv. Theor. Math. Phys.* 8 (2004) 345–394 MR
- [48] **J A Lind, H Sati, C Westerland**, *A higher categorical analogue of topological T-duality for sphere bundles*, preprint (2016) arXiv
- [49] **J Lott**, *R/Z index theory*, *Comm. Anal. Geom.* 2 (1994) 279–311 MR
- [50] **J Lott**, *Secondary analytic indices*, from “Regulators in analysis, geometry and number theory” (A Reznikov, N Schappacher, editors), *Progr. Math.* 171, Birkhäuser, Boston (2000) 231–293 MR
- [51] **J Lurie**, *Higher algebra*, prepublication book draft (2011) Available at <http://www.math.harvard.edu/~lurie/papers/HA.pdf>
- [52] **C R F Maunder**, *The spectral sequence of an extraordinary cohomology theory*, *Proc. Cambridge Philos. Soc.* 59 (1963) 567–574 MR
- [53] **J P May, J Sigurdsson**, *Parametrized homotopy theory*, *Mathematical Surveys and Monographs* 132, Amer. Math. Soc., Providence, RI (2006) MR
- [54] **J McCleary**, *A user’s guide to spectral sequences*, 2nd edition, *Cambridge Studies in Advanced Mathematics* 58, Cambridge Univ. Press (2001) MR

- [55] **H R Miller**, *On relations between Adams spectral sequences, with an application to the stable homotopy of a Moore space*, J. Pure Appl. Algebra 20 (1981) 287–312 MR
- [56] **R Minasian, G Moore**, *K-theory and Ramond–Ramond charge*, J. High Energy Phys. (1997) art. id. 002, 7 pages MR
- [57] **V V Prasolov**, *Elements of combinatorial and differential topology*, Graduate Studies in Mathematics 74, Amer. Math. Soc., Providence, RI (2006) MR
- [58] **D C Ravenel**, *Localization with respect to certain periodic homology theories*, Amer. J. Math. 106 (1984) 351–414 MR
- [59] **Y B Rudyak**, *On Thom spectra, orientability, and cobordism*, Springer (1998) MR
- [60] **H Sati**, *Geometric and topological structures related to M-branes*, from “Superstrings, geometry, topology, and C^* -algebras” (R S Doran, G Friedman, J Rosenberg, editors), Proc. Sympos. Pure Math. 81, Amer. Math. Soc., Providence, RI (2010) 181–236 MR
- [61] **H Sati, U Schreiber, J Stasheff**, *Twisted differential string and fivebrane structures*, Comm. Math. Phys. 315 (2012) 169–213 MR
- [62] **H Sati, C Westerland**, *Twisted Morava K-theory and E-theory*, J. Topol. 8 (2015) 887–916 MR
- [63] **U Schreiber**, *Differential cohomology in a cohesive infinity-topos*, preprint (2013) arXiv
- [64] **S Schwede**, *Symmetric spectra*, unpublished manuscript (2012) Available at <http://www.math.uni-bonn.de/people/schwede/SymSpec-v3.pdf>
- [65] **B Shipley**, *$H\mathbb{Z}$ -algebra spectra are differential graded algebras*, Amer. J. Math. 129 (2007) 351–379 MR
- [66] **J Simons, D Sullivan**, *Axiomatic characterization of ordinary differential cohomology*, J. Topol. 1 (2008) 45–56 MR
- [67] **R M Switzer**, *Algebraic topology — homotopy and homology*, Grundle. Math. Wissen. 212, Springer (1975) MR
- [68] **H Tamanoi**, *\mathcal{Q} -subalgebras, Milnor basis, and cohomology of Eilenberg–Mac Lane spaces*, J. Pure Appl. Algebra 137 (1999) 153–198 MR
- [69] **T Tradler, S O Wilson, M Zeinalian**, *An elementary differential extension of odd K-theory*, J. K-Theory 12 (2013) 331–361 MR
- [70] **T Tradler, S O Wilson, M Zeinalian**, *Differential K-theory as equivalence classes of maps to Grassmannians and unitary groups*, New York J. Math. 22 (2016) 527–581 MR
- [71] **M Upmeyer**, *Algebraic structure and integration maps in cocycle models for differential cohomology*, Algebr. Geom. Topol. 15 (2015) 65–83 MR
- [72] **N Yagita**, *On the Steenrod algebra of Morava K-theory*, J. London Math. Soc. 22 (1980) 423–438 MR

*Department of Mathematics, New York University, Abu Dhabi
Abu Dhabi, United Arab Emirates*

*Department of Mathematics, University of Pittsburgh
Pittsburgh, PA, United States*

djg9@nyu.edu, hsati@pitt.edu

Received: 3 June 2016 Revised: 11 October 2016