

Second mod 2 homology of Artin groups

TOSHIYUKI AKITA
YE LIU

In this paper, we compute the second mod 2 homology of an arbitrary Artin group, without assuming the $K(\pi, 1)$ conjecture. The key ingredients are (A) Hopf’s formula for the second integral homology of a group and (B) Howlett’s result on the second integral homology of Coxeter groups.

20F36, 20J06; 20F55

1 Introduction

An Artin group (or an Artin–Tits group) is a finitely presented group with at most one simple relation between a pair of generators. Examples includes finitely generated free abelian groups, free groups of finite rank, Artin’s braid groups with finitely many strands, right-angled Artin groups, etc. Artin groups appear in diverse branches of mathematics such as singularity theory, low-dimensional topology, geometric group theory, the theory of hyperplane arrangements, etc.

Artin groups are closely related to Coxeter groups. For a Coxeter graph Γ and the corresponding Coxeter system $(W(\Gamma), S)$, we associate an Artin group $A(\Gamma)$ obtained by, informally speaking, dropping the relations that each generator has order 2 from the standard presentation of $W(\Gamma)$. The symmetric group \mathfrak{S}_n is the Coxeter group associated to the Coxeter graph of type A_{n-1} , and the braid group $\text{Br}(n)$ is the corresponding Artin group. The Coxeter group $W(\Gamma)$ can be realized as a reflection group acting on a convex cone U (called Tits cone) in \mathbb{R}^n with $n = \#S$ the rank of W . Let \mathcal{A} be the collection of reflection hyperplanes. The complement

$$M(\Gamma) = (\text{int}(U) + \sqrt{-1}\mathbb{R}) \setminus \bigcup_{H \in \mathcal{A}} H \otimes \mathbb{C}$$

admits the free $W(\Gamma)$ -action, and the resulting orbit space $N(\Gamma) = M(\Gamma)/W(\Gamma)$ has the fundamental group isomorphic to $A(\Gamma)$; see van der Lek [20]. The celebrated $K(\pi, 1)$ conjecture states that $N(\Gamma)$ is a $K(A(\Gamma), 1)$ space. See Section 2.3 for a list of Γ for which the $K(\pi, 1)$ conjecture is proved.

Existing results about (co)homology of Artin groups all focus on particular types Artin groups for which the $K(\pi, 1)$ conjecture has been proved. There are very few properties that can be said for (co)homology of *all* Artin groups (except for their first integral homology, which is simply the abelianization). In this paper, we compute the second mod 2 homology of *all* Artin groups without assuming an affirmative solution of the $K(\pi, 1)$ conjecture. Our main tools are Hopf's formula on the second homology (or the Schur multiplier) of groups, together with Howlett's theorem (Theorem 3.2) on the second integral homology of Coxeter groups. We are inspired by Korkmaz and Stipsicz [19], who computed the second integral homology of the mapping class groups of oriented surfaces using Hopf's formula.

Our main result is the following.

Theorem 1.1 *Let $A(\Gamma)$ be the Artin group associated to a Coxeter graph Γ . Then*

$$H_2(A(\Gamma); \mathbb{Z}_2) \cong \mathbb{Z}_2^{p(\Gamma)+q(\Gamma)},$$

where $p(\Gamma)$ and $q(\Gamma)$ are nonnegative integers associated to Γ (see Theorem 2.6 for definitions).

As a corollary, we obtain a sufficient condition that the classifying map $c: N(\Gamma) \rightarrow K(A(\Gamma), 1)$ induces an isomorphism

$$c_*: H_2(N(\Gamma); \mathbb{Z}) \rightarrow H_2(A(\Gamma); \mathbb{Z}).$$

Furthermore, we conclude that the induced homomorphism

$$c_* \otimes \text{id}_{\mathbb{Z}_2}: H_2(N(\Gamma); \mathbb{Z}) \otimes \mathbb{Z}_2 \rightarrow H_2(A(\Gamma); \mathbb{Z}) \otimes \mathbb{Z}_2$$

is always an isomorphism. This gives affirmative evidence for the $K(\pi, 1)$ conjecture.

A part of this paper is based on the second author's PhD thesis.

2 Preliminaries

We collect relevant definitions and properties of Coxeter groups and Artin groups. We refer to [2; 17] for Coxeter groups and [23; 24; 25] for Artin groups.

2.1 Coxeter groups

Let S be a finite set. A *Coxeter matrix* over S is a symmetric matrix $M = (m(s, t))_{s, t \in S}$ such that $m(s, s) = 1$ for all $s \in S$, and $m(s, t) = m(t, s) \in \{2, 3, \dots\} \cup \{\infty\}$ for distinct $s, t \in S$. It is convenient to represent M by a labeled graph Γ , called the *Coxeter graph* of M , defined as follows:

- the vertex set is $V(\Gamma) = S$;
- the edge set is $E(\Gamma) = \{\{s, t\} \subset S \mid m(s, t) \geq 3\}$;
- the edge $\{s, t\}$ is labeled by $m(s, t)$ if $m(s, t) \geq 4$.

Let Γ_{odd} be the subgraph of Γ with $V(\Gamma_{\text{odd}}) = V(\Gamma)$ and $E(\Gamma_{\text{odd}}) = \{\{s, t\} \in E(\Gamma) \mid m(s, t) \text{ odd}\}$ inheriting labels from Γ . By abuse of notation, we frequently regard Γ (hence also Γ_{odd}) as its underlying 1–dimensional CW-complex.

Definition 2.1 Let Γ be a Coxeter graph and S its vertex set. The *Coxeter system* associated to Γ is the pair $(W(\Gamma), S)$, where the *Coxeter group* $W(\Gamma)$ is defined by the standard presentation

$$W(\Gamma) = \langle S \mid (st)^{m(s,t)} = 1 \text{ for all } s, t \in S \text{ with } m(s, t) \neq \infty \rangle.$$

Each generator $s \in S$ of W has order 2. For distinct $s, t \in S$, the order of st is precisely $m(s, t)$ if $m(s, t) \neq \infty$. In the case where $m(s, t) = \infty$, the element st has infinite order.

However, in this paper, we adopt an equivalent definition. For two letters s, t and an integer $m \geq 2$, we shall use the following notation of the word of length m consisting of s and t in an alternating order:

$$(st)_m := \overbrace{stst \cdots}^m.$$

For example, $(st)_2 = st$, $(st)_3 = sts$ and $(st)_4 = stst$.

Definition 2.2 Let Γ be a Coxeter graph and S its vertex set. The *Coxeter group* associated to Γ is the group defined by the presentation

$$W(\Gamma) = \langle S \mid \overline{R_W} \cup Q_W \rangle.$$

The sets of relations are $\overline{R_W} = \{R(s, t) \mid m(s, t) < \infty\}$ and $Q_W = \{Q(s) \mid s \in S\}$, where $R(s, t) := (st)_{m(s,t)}(ts)_{m(s,t)}^{-1}$ and $Q(s) := s^2$.

Note that since $R(s, t) = R(t, s)^{-1}$, we may reduce the relation set $\overline{R_W}$ by introducing a total order on S and put $R_W := \{R(s, t) \mid m(s, t) < \infty, s < t\}$. We have the following presentation with fewer relations:

$$W(\Gamma) = \langle S \mid R_W \cup Q_W \rangle.$$

We shall omit the reference to Γ if there is no ambiguity. The rank of W is defined to be $\#S$.

Let (W, S) be a Coxeter system. For a subset $T \subset S$, let W_T denote the subgroup of W generated by T , called a *parabolic subgroup* of W . In particular, $W_S = W$ and $W_\emptyset = \{1\}$. It is known that (W_T, T) is the Coxeter system associated to the Coxeter graph Γ_T (the full subgraph of Γ spanned by T inheriting labels); see Théorème 2 in Chapter IV of [2].

2.2 Artin groups

The Artin group $A(\Gamma)$ associated to a Coxeter graph Γ is obtained from the presentation of $W(\Gamma)$ by dropping the relation set Q_W .

Definition 2.3 Given a Coxeter graph Γ (hence a Coxeter system (W, S)), we introduce a set $\Sigma = \{a_s \mid s \in S\}$ in one-to-one correspondence with S . Then the Artin system associated to Γ is the pair $(A(\Gamma), \Sigma)$, where $A(\Gamma)$ is the *Artin group* of type Γ defined by the presentation

$$A(\Gamma) = \langle \Sigma \mid \overline{R_A} \rangle,$$

where $\overline{R_A} = \{R(a_s, a_t) \mid m(s, t) < \infty\}$ and $R(a_s, a_t) = (a_s a_t)_{m(s,t)} (a_t a_s)_{m(s,t)}^{-1}$.

As in the Coxeter group case, we introduce a total order on S and put $R_A := \{R(a_s, a_t) \mid m(s, t) < \infty, s < t\}$. We have the following presentation with fewer relations:

$$A(\Gamma) = \langle \Sigma \mid R_A \rangle.$$

There is a canonical projection $p: A(\Gamma) \rightarrow W(\Gamma)$, $a_s \mapsto s$ ($s \in S$), whose kernel is called the *pure Artin group* of type Γ .

We say that an Artin group $A(\Gamma)$ is of finite type (or spherical type) if the associated Coxeter group $W(\Gamma)$ is finite; otherwise, $A(\Gamma)$ is of infinite type (or nonspherical type).

2.3 $K(\pi, 1)$ conjecture

Consider a Coxeter graph Γ and the associated Coxeter system (W, S) with rank $\#S = n$. Recall that W can be realized as a reflection group acting on a Tits cone $U \subset \mathbb{R}^n$; see [24]. Let \mathcal{A} be the collection of the reflection hyperplanes. Put

$$M(\Gamma) := (\text{int}(U) + \sqrt{-1}\mathbb{R}^n) \setminus \bigcup_{H \in \mathcal{A}} H \otimes \mathbb{C}.$$

Then W acts on $M(\Gamma)$ freely and properly discontinuously. Denote the orbit space by

$$(2-1) \quad N(\Gamma) := M(\Gamma)/W.$$

It is known that:

Theorem 2.4 [20] *The fundamental group of $N(\Gamma)$ is isomorphic to the Artin group $A(\Gamma)$.*

In general, $N(\Gamma)$ is only conjectured to be a classifying space of $A(\Gamma)$.

Conjecture 2.5 *Let Γ be an arbitrary Coxeter graph. Then the orbit space $N(\Gamma)$ is a $K(\pi, 1)$ space, hence a classifying space of the Artin group $A(\Gamma)$.*

This conjecture is proved to hold for a few classes of Artin groups. Here is a list of such classes known so far:

- Artin groups of finite type [12].
- Artin groups of large type [14].
- 2-dimensional Artin groups [7].
- Artin groups of FC type [7].
- Artin groups of affine types \tilde{A}_n, \tilde{C}_n [21].
- Artin groups of affine type \tilde{B}_n [5].
- Artin groups $A(\Gamma)$ such that the $K(\pi, 1)$ conjecture holds for all $A(\Gamma_T)$ where $T \subset S$ and Γ_T does not contain ∞ -labeled edges [13].
- Artin groups whose corresponding Coxeter system (W, S) has the following property: every finite irreducible parabolic subgroup W_T is either $\mathfrak{S}_4, \mathbb{Z}_2$ or a dihedral group [6].

2.4 First and second homology of $N(\Gamma)$

Clancy and Ellis [8] computed the second integral homology of $N(\Gamma)$ using the Salvetti complex for an Artin group. We recall their result and follow their notation.

Let us first fix some notation. Let Γ be a Coxeter graph with vertex set S . Define $Q(\Gamma) = \{\{s, t\} \subset S \mid m(s, t) \text{ is even}\}$ and $P(\Gamma) = \{\{s, t\} \subset S \mid m(s, t) = 2\}$. Write $\{s, t\} \equiv \{s', t'\}$ if two such pairs in $P(\Gamma)$ satisfy $s = s'$ and $m(t, t')$ is odd. This generates an equivalence relation on $P(\Gamma)$, denoted by \sim . Let $P(\Gamma)/\sim$ be the set of equivalence classes. An equivalence class is called a *torsion* class if it is represented by a pair $\{s, t\} \in P(\Gamma)$ such that there exists a vertex $v \in S$ with $m(s, v) = m(t, v) = 3$. In the above situation, Clancy and Ellis proved the following theorem.

Theorem 2.6 [8] *Let Γ be a Coxeter graph and $N(\Gamma)$ as in (2-1). Then*

$$H_2(N(\Gamma); \mathbb{Z}) \cong \mathbb{Z}_2^{p(\Gamma)} \oplus \mathbb{Z}^{q(\Gamma)},$$

where

$$p(\Gamma) := \text{number of torsion classes in } P(\Gamma)/\sim,$$

$$q_1(\Gamma) := \text{number of nontorsion classes in } P(\Gamma)/\sim,$$

$$q_2(\Gamma) := \#(Q(\Gamma) - P(\Gamma)) = \#\{\{s, t\} \subset S \mid m(s, t) \geq 4, m(s, t) \text{ is even}\},$$

$$q_3(\Gamma) := \text{rank } H_1(\Gamma_{\text{odd}}; \mathbb{Z}),$$

$$q(\Gamma) := q_1(\Gamma) + q_2(\Gamma) + q_3(\Gamma).$$

Remark Note that $H_1(N(\Gamma); \mathbb{Z}) \cong H_1(A(\Gamma); \mathbb{Z})$ is isomorphic to the abelianization of $A(\Gamma)$, which is a free abelian group with rank equal to $\text{rank } H_0(\Gamma_{\text{odd}}; \mathbb{Z})$, the number of connected components of Γ_{odd} .

3 Second mod 2 homology of Artin groups

The (co)homology of the orbit space $N(\Gamma)$ coincides with that of the Artin group $A(\Gamma)$, provided the $K(\pi, 1)$ conjecture for $A(\Gamma)$ holds. There are many results about (co)homology of $N(\Gamma)$ in the literature, for example, [10; 9; 4; 5]. The $K(\pi, 1)$ conjecture is known to hold in these cases.

In this section, nevertheless, we shall work on the second homology of *arbitrary* Artin groups without assuming that the $K(\pi, 1)$ conjecture holds. Our main result is the following theorem.

Theorem 3.1 *Let Γ be an arbitrary Coxeter graph and $A(\Gamma)$ the associated Artin group. Then the second mod 2 homology of $A(\Gamma)$ is*

$$H_2(A(\Gamma); \mathbb{Z}_2) \cong \mathbb{Z}_2^{p(\Gamma)+q(\Gamma)},$$

where $p(\Gamma)$ and $q(\Gamma)$ are as in Theorem 2.6.

The outline of our proof is as follows. In Section 3.1, we state Howlett's theorem on the second integral homology group $H_2(W(\Gamma); \mathbb{Z})$ of the Coxeter group $W(\Gamma)$. Next, in Section 3.2, we recall Hopf's formula of the second homology of a group. The key of the proof is that, by virtue of Hopf's formula, we are able to find explicitly a set $\Omega(W)$ of generators of $H_2(W(\Gamma); \mathbb{Z})$ (Section 3.3), as well as a set $\Omega(A)$

of generators of $H_2(A(\Gamma); \mathbb{Z})$ (Section 3.4). On the other hand, Howlett’s theorem implies that $\Omega(W)$ forms a basis of $H_2(W(\Gamma); \mathbb{Z})$, which is an elementary abelian 2–group of rank $p(\Gamma) + q(\Gamma)$. Furthermore, we will show that the homomorphism $p_*: H_2(A(\Gamma); \mathbb{Z}) \rightarrow H_2(W(\Gamma); \mathbb{Z})$ induced by the projection $p: A(\Gamma) \rightarrow W(\Gamma)$ maps $\Omega(A)$ onto $\Omega(W)$. Hence p_* is actually an epimorphism and becomes an isomorphism when tensored with \mathbb{Z}_2 .

3.1 Howlett’s theorem

As previously mentioned, we shall study the homomorphism $p_*: H_2(A(\Gamma); \mathbb{Z}) \rightarrow H_2(W(\Gamma); \mathbb{Z})$ induced by the projection $p: A(\Gamma) \rightarrow W(\Gamma)$. A reason for doing so is that we have the following theorem of Howlett:

Theorem 3.2 [16] *The second integral homology of the Coxeter group $W(\Gamma)$ associated to a Coxeter graph Γ is*

$$H_2(W(\Gamma); \mathbb{Z}) \cong \mathbb{Z}_2^{p(\Gamma)+q(\Gamma)},$$

where $p(\Gamma)$ and $q(\Gamma)$ are as in Theorem 2.6.

Remark The original statement in [16] was

$$H_2(W(\Gamma); \mathbb{Z}) \cong \mathbb{Z}_2^{-n_1(\Gamma)+n_2(\Gamma)+n_3(\Gamma)+n_4(\Gamma)},$$

where

$$\begin{aligned} n_1(\Gamma) &:= \#S, \\ n_2(\Gamma) &:= \#\{\{s, t\} \in E(\Gamma) \mid m(s, t) < \infty\}, \\ n_3(\Gamma) &:= \#P(\Gamma)/\sim, \\ n_4(\Gamma) &:= \text{rank } H_0(\Gamma_{\text{odd}}; \mathbb{Z}). \end{aligned}$$

For a Coxeter graph Γ , the above numbers are related to those used by Clancy–Ellis as follows:

$$-n_1(\Gamma) + n_2(\Gamma) + n_3(\Gamma) + n_4(\Gamma) = p(\Gamma) + q(\Gamma).$$

In fact, $n_1(\Gamma) = \#V(\Gamma_{\text{odd}})$, $n_2(\Gamma) = q_2(\Gamma) + \#E(\Gamma_{\text{odd}})$ and $n_3(\Gamma) = p(\Gamma) + q_1(\Gamma)$. The above equation follows from the Euler–Poincaré theorem applied to Γ_{odd} :

$$\#V(\Gamma_{\text{odd}}) - \#E(\Gamma_{\text{odd}}) = \text{rank } H_0(\Gamma_{\text{odd}}; \mathbb{Z}) - \text{rank } H_1(\Gamma_{\text{odd}}; \mathbb{Z}).$$

Example 3.3 We shall make use of the following example later.

Let $\Gamma = I_2(m)$. Thus $W(\Gamma) = D_{2m}$ is the dihedral group of order $2m$. Theorem 3.2 shows that

$$H_2(W(\Gamma); \mathbb{Z}) \cong \begin{cases} \mathbb{Z}_2 & \text{if } m \text{ is even,} \\ 0 & \text{if } m \text{ is odd.} \end{cases}$$

See Corollary 10.1.27 of [18] for a complete list of integral homology of dihedral groups.

3.2 Hopf’s formula

Hopf’s formula gives a description of the second integral homology of a group. We first recall some notation. For a group G , the commutator of $x, y \in G$ is the element $[x, y] = xyx^{-1}y^{-1}$. The commutator subgroup $[G, G]$ of G is the subgroup of G generated by all commutators. In general, for any subgroups H and K of G , we define $[H, K]$ as the subgroup of G generated by all $[h, k]$ for $h \in H, k \in K$.

Theorem 3.4 (Hopf’s formula) *If a group G has a presentation $\langle S \mid R \rangle$, then*

$$H_2(G; \mathbb{Z}) \cong \frac{N \cap [F, F]}{[F, N]},$$

where $F = F(S)$ is the free group generated by S and $N = N(R)$ is the normal closure of R (the subgroup of F normally generated by the relation set R).

See Section II.5 of [3] for a topological proof. Moreover, Hopf’s formula admits the following naturality; see Section II.6, Exercise 3(b) of [3].

Proposition 3.5 *Let $G = F/N = \langle S \mid R \rangle$ and $G' = F'/N' = \langle S' \mid R' \rangle$ as in Theorem 3.4. Suppose a homomorphism $\alpha: G \rightarrow G'$ lifts to $\tilde{\alpha}: F \rightarrow F'$. Then the following diagram commutes:*

$$\begin{array}{ccc} H_2(G; \mathbb{Z}) & \xrightarrow{\cong} & N \cap [F, F]/[F, N] \\ H_2(\alpha) \downarrow & & \downarrow \alpha_* \\ H_2(G'; \mathbb{Z}) & \xrightarrow{\cong} & N' \cap [F', F']/[F', N'] \end{array}$$

where α_* is induced by $\tilde{\alpha}$.

For simplicity, we denote by $\langle x \rangle_G = x[F, N] \in F/[F, N]$ the coset of $[F, N]$ represented by $x \in F$ and $\langle x, y \rangle_G = [x, y][F, N] \in [F, F]/[N, F]$ for $x, y \in F$. Thanks to Hopf's formula, second homology classes of G can be regarded as $\langle x \rangle_G$ for $x \in N \cap [F, F]$.

To see what the representatives look like, we make the following simple observations, which we learned from [19].

Lemma 3.6 *The group $N/[F, N]$ is abelian.*

Proof Note that $N/[F, N]$ is a quotient group of $N/[N, N]$ and the latter is the abelianization of N . □

Thus we write the group $N/[F, N]$ additively. It is clear $\langle n \rangle_G = -\langle n^{-1} \rangle_G$ for $n \in N$.

Lemma 3.7 *In the abelian group $N/[F, N]$, for $n \in N$ and $f \in F$, we have*

$$\langle n \rangle_G = \langle fnf^{-1} \rangle_G.$$

Proof Since $[f, n] \in [F, N]$, we have

$$\langle f, n \rangle_G = \langle fnf^{-1}n^{-1} \rangle_G = \langle fnf^{-1} \rangle_G - \langle n \rangle_G = 0. \quad \square$$

Therefore, a coset in $N/[F, N]$ is represented by an element of the form $\prod_{r \in R} r^{n(r)}$ for $n(r) \in \mathbb{Z}$. Hopf's formula implies that a second homology class of G can be represented by an element $\prod_{r \in R} r^{n(r)} \in [F, F]$.

Lemma 3.8 *Let $G = F/N$ as in Theorem 3.4. If $x, y, z \in F$ such that $[x, y], [x, z] \in N \cap [F, F]$, then*

$$\langle x, yz \rangle_G = \langle x, y \rangle_G + \langle x, z \rangle_G, \quad \langle x, y^{-1} \rangle_G = -\langle x, y \rangle_G.$$

Proof Note that $[x, yz] = [x, y]y[x, z]y^{-1}$. Then in the abelian group $N/[F, N]$,

$$\langle x, yz \rangle_G = \langle x, y \rangle_G + \langle y[x, z]y^{-1} \rangle_G.$$

For the last term, we have $\langle y[x, z]y^{-1} \rangle_G = \langle x, z \rangle_G$ since

$$[x, z]^{-1}y[x, z]y^{-1} = [[x, z]^{-1}, y] \in [N, F].$$

Hence the first equality holds. The second follows immediately from the first. □

3.3 Hopf’s formula applied to Coxeter groups

The aim of this subsection is to construct an explicit set $\Omega(W)$ of generators of $H_2(W(\Gamma); \mathbb{Z})$. Combined with Howlett’s theorem (Theorem 3.2), we show that $\Omega(W)$ is a basis of $H_2(W(\Gamma); \mathbb{Z})$.

Let us describe the construction of $\Omega(W)$. Let Γ be a Coxeter graph and (W, S) the associated Coxeter system with S totally ordered. Then $W = \langle S \mid R_W \cup Q_W \rangle$ is as in Definition 2.2. Let $F_W = F(S)$ be the free group on S and $N_W = N(R_W \cup Q_W)$ the normal closure of $R_W \cup Q_W$. Therefore, $W = F_W/N_W$. Using Hopf’s formula, we identify $H_2(W; \mathbb{Z}) \cong (N_W \cap [F_W, F_W])/[F_W, N_W]$. We shall construct three sets $\Omega_i(W) \subset (N_W \cap [F_W, F_W])/[F_W, N_W]$ ($i = 1, 2, 3$). In view of Lemma 3.6 and Lemma 3.7, a second homology class of W is of the form $\langle x \rangle_W$ with x expressed by a word $\prod_{R(s,t) \in R_W} R(s,t)^{n(s,t)} \prod_{Q(s) \in Q_W} Q(s)^{n(s)} \in [F_W, F_W]$. We decompose $\langle x \rangle_W = \langle x_1 \rangle_W + \langle x_2 \rangle_W + \langle x_3 \rangle_W$, as in the proof of Theorem 3.15, such that $\langle x_i \rangle_W$ is generated by $\Omega_i(W)$. Then $\Omega(W) = \Omega_1(W) \cup \Omega_2(W) \cup \Omega_3(W)$ generates $H_2(W; \mathbb{Z})$. Now we exhibit respectively the constructions of $\Omega_i(W)$ ($i = 1, 2, 3$).

3.3.1 Construction of $\Omega_1(W)$ Let

$$\Omega_1(W) = \{ \langle s, t \rangle_W \mid s, t \in S, s < t, m(s, t) = 2 \}.$$

Recall that $\langle s, t \rangle_W = [s, t][F_W, N_W] \in (N_W \cap [F_W, F_W])/[F_W, N_W]$ and $R(s, t) = [s, t]$ when $m(s, t) = 2$. Note that the above expression may have repetitions. In fact, we have the following.

Proposition 3.9 $\# \Omega_1(W) \leq p(\Gamma) + q_1(\Gamma)$.

Proof We shall show that $\langle s, t \rangle_W = \langle s, t' \rangle_W$ in $\Omega_1(W)$ if $\{s, t\} \equiv \{s, t'\}$ in $P(\Gamma)$. Suppose $s < t$ and $s < t'$ with $\{s, t\} \equiv \{s, t'\}$ in $P(\Gamma)$; that is, $m(s, t) = m(s, t') = 2$ and $m(t, t')$ is odd. Then in $N_W/[F_W, N_W]$,

$$\langle s, R(t, t') \rangle_W = \langle s, (tt' \cdots t)(t't \cdots t')^{-1} \rangle_W = \langle s, t \rangle_W - \langle s, t' \rangle_W,$$

where the latter equality follows from Lemma 3.8. On the other hand,

$$\begin{aligned} \langle s, R(t, t') \rangle_W &= \langle sR(t, t')s^{-1}R(t, t')^{-1} \rangle_W \\ &= \langle sR(t, t')s^{-1} \rangle_W + \langle R(t, t')^{-1} \rangle_W \\ &= \langle R(t, t') \rangle_W - \langle R(t, t') \rangle_W = 0, \end{aligned}$$

where the second equality follows from Lemma 3.8 and the third from Lemma 3.7. Thus we conclude $\langle s, t \rangle_W = \langle s, t' \rangle_W$. Similarly, $\langle s, t \rangle_W = \langle s', t \rangle_W$ in $\Omega_1(W)$ if $\{s, t\} \equiv \{s', t\}$ in $P(\Gamma)$. Hence $\#\Omega_1(W) \leq \#(P(\Gamma)/\sim) = p(\Gamma) + q_1(\Gamma)$. \square

3.3.2 Construction of $\Omega_2(W)$ Let

$$\Omega_2(W) = \{ \langle R(s, t) \rangle_W \mid s, t \in S, s < t, m(s, t) \geq 4, m(s, t) \text{ is even} \}.$$

Recall that $R(s, t) = (st)_{m(s,t)}(ts)_{m(s,t)}^{-1}$. Note that when $m(s, t)$ is even, $R(s, t)$ is in the kernel of the abelianization map $\text{Ab}: F_W \rightarrow F_W/[F_W, F_W]$, and hence $R(s, t) \in [F_W, F_W]$. The following result is obvious.

Proposition 3.10 $\#\Omega_2(W) \leq q_2(\Gamma)$.

3.3.3 Construction of $\Omega_3(W)$ The construction of $\Omega_3(W)$ requires more preparation. Recall that Γ_{odd} is the subgraph of Γ considered as a 1–dimensional CW-complex with 0–cells S and 1–cells $\{ \langle s, t \rangle \mid s, t \in S, s < t, m(s, t) \text{ odd} \}$ oriented by $\partial \langle s, t \rangle = t - s$. We define a group

$$\mathcal{C}_W = \{ (\alpha, \beta) \in C_1(\Gamma_{\text{odd}}) \oplus 2C_0(\Gamma_{\text{odd}}) \mid \partial \alpha = \beta \},$$

where $2C_0(\Gamma_{\text{odd}}) = \{ 2\gamma \mid \gamma \in C_0(\Gamma_{\text{odd}}) \}$ is the group of 0–chains with all coefficients even, and \mathcal{D}_W is the subgroup of \mathcal{C}_W generated by $(2 \langle s, t \rangle, -2s + 2t)$ for all 1–cells $\langle s, t \rangle$.

Consider the homomorphism

$$\Phi_W: \mathcal{C}_W \rightarrow \frac{N_W \cap [F_W, F_W]}{[F_W, N_W]}$$

defined by

$$\Phi_W \left(\sum_{\substack{s < t \\ m(s,t) \text{ odd}}} n(s, t) \langle s, t \rangle, \sum_{s \in S} 2n(s) s \right) = \left\langle \prod_{\substack{s < t \\ m(s,t) \text{ odd}}} R(s, t)^{n(s,t)} \prod_{s \in S} Q(s)^{n(s)} \right\rangle_W.$$

The definition is indeed valid by the following easy lemma.

Lemma 3.11 *The following are equivalent:*

- (A) $\left(\sum_{\substack{s < t \\ m(s,t) \text{ odd}}} n(s, t) \langle s, t \rangle, \sum_{s \in S} 2n(s) s \right) \in \mathcal{C}_W,$
- (B) $\prod_{\substack{s < t \\ m(s,t) \text{ odd}}} R(s, t)^{n(s,t)} \prod_{s \in S} Q(s)^{n(s)} \in [F_W, F_W].$

Proof We suppress the ranges since they should be clear:

$$\begin{aligned} \text{(A)} &\iff \partial\left(\sum n(s, t)\langle s, t \rangle\right) = \sum 2n(s)s \\ &\iff \sum 2n(s)s + \sum n(s, t)(s - t) = 0 \\ &\iff \text{Ab}\left(\prod R(s, t)^{n(s, t)} \prod Q(s)^{n(s)}\right) = 0 \iff \text{(B)}, \end{aligned}$$

where Ab is the abelianization map as above, and we write $F_W/[F_W, F_W]$ additively. Note that $\text{Ab}(R(s, t)) = s - t$ if $m(s, t)$ is odd. □

The following is a consequence of Example 3.3.

Proposition 3.12 \mathcal{D}_W lies in the kernel of Φ_W .

Proof It suffices to show that any generator $(2\langle s, t \rangle, -2s + 2t)$ of \mathcal{D}_W is mapped to the identity by Φ_W , or equivalently, the word $((st)_m(ts)_m^{-1})^2(s^2)^{-1}t^2$ lies in $[F_W, N_W]$ when m is odd. Let $s, t \in S$ with $m := m(s, t)$ odd; consider the parabolic subgroup $W' := W_{\{s, t\}}$ of W , which is isomorphic to the dihedral group D_{2m} of order $2m$. From Example 3.3, we know $H_2(W'; \mathbb{Z}) = 0$. On the other hand, Hopf’s formula applied to W' shows that $H_2(W'; \mathbb{Z}) \cong (N_{W'} \cap [F_{W'}, F_{W'}])/[F_{W'}, N_{W'}]$. Therefore, the word $((st)_m(ts)_m^{-1})^2(s^2)^{-1}t^2 \in N_{W'} \cap [F_{W'}, F_{W'}]$ represents the trivial homology class. That is to say,

$$((st)_m(ts)_m^{-1})^2(s^2)^{-1}t^2 \in [F_{W'}, N_{W'}] \subset [F_W, N_W]. \quad \square$$

As a consequence, the homomorphism Φ_W factors through

$$\mathcal{C}_W \twoheadrightarrow \mathcal{C}_W/\mathcal{D}_W \rightarrow (N_W \cap [F_W, F_W])/[F_W, N_W].$$

Let $Z_1(\Gamma_{\text{odd}}; \mathbb{Z})$ denote the group of 1–cycles of Γ_{odd} with integral coefficients and $Z_1(\Gamma_{\text{odd}}; \mathbb{Z}_2)$ the group of 1–cycles of Γ_{odd} with coefficients in \mathbb{Z}_2 . Define a homomorphism $\Xi_W: \mathcal{C}_W \rightarrow Z_1(\Gamma_{\text{odd}}; \mathbb{Z}_2)$ by $(\alpha, \partial\alpha) \mapsto \bar{\alpha}$, where $\alpha \in C_1(\Gamma_{\text{odd}})$ such that $\partial\alpha \in 2C_0(\Gamma_{\text{odd}})$ and $\bar{\alpha} \in C_1(\Gamma_{\text{odd}}; \mathbb{Z}_2)$ is the mod 2 reduction of α . The condition $\partial\alpha \in 2C_0(\Gamma_{\text{odd}})$ asserts that $\bar{\alpha}$ is indeed a 1–cycle of Γ_{odd} with coefficients in \mathbb{Z}_2 .

Proposition 3.13 The homomorphism $\Xi_W: \mathcal{C}_W \rightarrow Z_1(\Gamma_{\text{odd}}; \mathbb{Z}_2)$ factors through an isomorphism:

$$\begin{array}{ccc} \mathcal{C}_W & & \\ \downarrow & \searrow \Xi_W & \\ \mathcal{C}_W/\mathcal{D}_W & \xrightarrow{\cong} & Z_1(\Gamma_{\text{odd}}; \mathbb{Z}_2) \end{array}$$

Proof The homomorphism Ξ_W is obviously an epimorphism, and

$$\begin{aligned} (\alpha, \partial\alpha) \in \text{Ker } \Xi_W &\iff \bar{\alpha} = 0 \in Z_1(\Gamma_{\text{odd}}; \mathbb{Z}_2) \\ &\iff \alpha \in 2C_1(\Gamma_{\text{odd}}) \\ &\iff (\alpha, \partial\alpha) \in \mathcal{D}_W. \end{aligned}$$

Hence $\text{Ker } \Xi_W = \mathcal{D}_W$. □

Via the isomorphism in Proposition 3.13, we obtain a homomorphism

$$\Psi_W: Z_1(\Gamma_{\text{odd}}; \mathbb{Z}_2) \rightarrow \frac{N_W \cap [F_W, F_W]}{[F_W, N_W]},$$

which fits into the following commutative diagram:

$$(3-1) \quad \begin{array}{ccc} C_W & \xrightarrow{\Phi_W} & N_W \cap [F_W, F_W] / [F_W, N_W] \\ \downarrow & \nearrow & \uparrow \Psi_W \\ C_W / \mathcal{D}_W & \xrightarrow{\cong} & Z_1(\Gamma_{\text{odd}}; \mathbb{Z}_2) \end{array}$$

We fix a basis $\Omega(\Gamma_{\text{odd}}; \mathbb{Z})$ of $Z_1(\Gamma_{\text{odd}}; \mathbb{Z}) \cong \mathbb{Z}^{q_3(\Gamma)}$ once and for all and denote by $\Omega(\Gamma_{\text{odd}}; \mathbb{Z}_2)$ the basis of $Z_1(\Gamma_{\text{odd}}; \mathbb{Z}_2) \cong \mathbb{Z}_2^{q_3(\Gamma)}$ obtained from $\Omega(\Gamma_{\text{odd}}; \mathbb{Z})$ by mod 2 reduction $Z_1(\Gamma_{\text{odd}}; \mathbb{Z}) \twoheadrightarrow Z_1(\Gamma_{\text{odd}}; \mathbb{Z}_2)$.

Define $\Omega_3(W)$ to be the image of $\Omega(\Gamma_{\text{odd}}; \mathbb{Z}_2)$ under Ψ_W :

$$\Omega_3(W) = \Psi_W(\Omega(\Gamma_{\text{odd}}; \mathbb{Z}_2)).$$

To be precise,

$$\Omega_3(W) = \left\{ \left\langle \prod_{\substack{s < t \\ m(s,t) \text{ odd}}} R(s,t)^{n(s,t)} \right\rangle_W \mid \sum_{\substack{s < t \\ m(s,t) \text{ odd}}} n(s,t) \langle s,t \rangle \in \Omega(\Gamma_{\text{odd}}; \mathbb{Z}_2) \right\}.$$

Proposition 3.14 $\#\Omega_3(W) \leq q_3(\Gamma)$.

Let $\Omega(W) = \Omega_1(W) \cup \Omega_2(W) \cup \Omega_3(W)$. We conclude that:

Theorem 3.15 $\Omega(W)$ is a basis of $H_2(W; \mathbb{Z})$.

Proof Since $\#\Omega(W) \leq p(\Gamma) + q(\Gamma)$ and $H_2(W; \mathbb{Z}) \cong \mathbb{Z}_2^{p(\Gamma)+q(\Gamma)}$ (Theorem 3.2), it suffices to show that $\Omega(W)$ generates $H_2(W; \mathbb{Z})$. An arbitrary homology class

in $H_2(W; \mathbb{Z})$ is represented by the coset $\langle x_1 x_2 x_3 \rangle_W$ with

$$\begin{aligned} x_1 &= \prod_{s < t, m(s,t)=2} [s, t]^{n(s,t)}, \\ x_2 &= \prod_{\substack{s < t, m(s,t) \geq 4 \\ m(s,t) \text{ even}}} R(s, t)^{n(s,t)}, \\ x_3 &= \prod_{s < t, m(s,t) \text{ odd}} R(s, t)^{n(s,t)} \prod_{Q(s) \in Q_W} Q(s)^{n(s)}, \end{aligned}$$

with $x_1, x_2, x_3 \in [F_W, F_W]$. Thus $\langle x_1 x_2 x_3 \rangle_W = \langle x_1 \rangle_W + \langle x_2 \rangle_W + \langle x_3 \rangle_W$. We claim that $\langle x_i \rangle_W$ is generated by $\Omega_i(W)$. In fact, the claim for $i = 1, 2$ is straightforward. For $i = 3$, let

$$\alpha = \sum_{s < t, m(s,t) \text{ odd}} n(s, t) \langle s, t \rangle, \quad \beta = \sum_{s \in S} 2n(s)s.$$

Thus $(\alpha, \beta) \in \mathcal{C}_W$ by Lemma 3.11 with $\Phi_W(\alpha, \beta) = \langle x_3 \rangle_W$. By the commutative diagram (3-1), the mod 2 reduction $\bar{\alpha} \in Z_1(\Gamma_{\text{odd}}; \mathbb{Z}_2)$ of α is mapped to $\langle x_3 \rangle_W$ by Ψ_W . This proves the claim. \square

Remark It is worth noting that in the previous proof, we have managed to get rid of the relations $Q(s)$ without altering the homology class $\langle x_3 \rangle_W$. This will be crucial in the proof of Theorem 3.20.

3.4 Hopf’s formula applied to Artin groups

Now we turn to the Artin group case. The arguments here are parallel to those in the Coxeter group case.

Let Γ be a Coxeter graph with the vertex set S totally ordered and $A = A(\Gamma)$ the Artin group of type Γ with the presentation $A = \langle \Sigma \mid R_A \rangle$ given in Definition 2.3. Let $F_A = F(\Sigma)$ be the free group on Σ and N_A the normal closure of R_A . Hopf’s formula yields $H_2(A; \mathbb{Z}) \cong (N_A \cap [F_A, F_A]) / [F_A, N_A]$. For the same reason as before, a second homology class of A is represented by a coset $\langle x \rangle_A$ with x of the form $\prod_{R(a_s, a_t) \in R_A} R(a_s, a_t)^{n(s,t)} \in [F_A, F_A]$.

We construct a set $\Omega(A)$ of generators of $H_2(A; \mathbb{Z})$ using the same method as in the previous subsection.

3.4.1 Constructions of $\Omega_1(A)$ and $\Omega_2(A)$ The constructions of $\Omega_1(A)$ and $\Omega_2(A)$ are exactly parallel to those in the Coxeter case. Let

$$\begin{aligned} \Omega_1(A) &= \{ \langle a_s, a_t \rangle_A \mid s, t \in S, s < t, m(s, t) = 2 \}, \\ \Omega_2(A) &= \{ \langle R(a_s, a_t) \rangle_A \mid s, t \in S, s < t, m(s, t) \geq 4, m(s, t) \text{ is even} \}. \end{aligned}$$

The same reasoning shows that:

Proposition 3.16 $\#\Omega_1(A) \leq p(\Gamma) + q_1(\Gamma), \#\Omega_2(A) \leq q_2(\Gamma).$

3.4.2 Construction of $\Omega_3(A)$ Consider the homomorphism

$$\Psi_A: Z_1(\Gamma_{\text{odd}}; \mathbb{Z}) \rightarrow \frac{N_A \cap [F_A, F_A]}{[F_A, N_A]}$$

defined by

$$\Psi_A \left(\sum_{\substack{s < t \\ m(s,t) \text{ odd}}} n(s, t) \langle s, t \rangle \right) = \left\langle \prod_{\substack{s < t \\ m(s,t) \text{ odd}}} R(a_s, a_t)^{n(s,t)} \right\rangle_A.$$

The definition is valid by the following lemma.

Lemma 3.17 *The following are equivalent:*

- (A) $\sum_{\substack{s < t \\ m(s,t) \text{ odd}}} n(s, t) \langle s, t \rangle \in Z_1(\Gamma_{\text{odd}}; \mathbb{Z}),$
- (B) $\prod_{\substack{s < t \\ m(s,t) \text{ odd}}} R(a_s, a_t)^{n(s,t)} \in [F_A, F_A].$

Proof We suppress again the ranges:

$$\begin{aligned} \text{(A)} &\iff \partial \left(\sum n(s, t) \langle s, t \rangle \right) = 0 \\ &\iff \sum n(s, t) (t - s) = 0 \\ &\iff \text{Ab} \left(\prod R(a_s, a_t)^{n(s,t)} \right) = 0 \iff \text{(B)}, \end{aligned}$$

where $\text{Ab}: F_A \rightarrow F_A/[F_A, F_A]$ is the abelianization map. Note that $\text{Ab}(R(a_s, a_t)) = a_s - a_t$ if $m(s, t)$ is odd. □

Recall that we have chosen a basis $\Omega(\Gamma_{\text{odd}}; \mathbb{Z})$ for $Z_1(\Gamma_{\text{odd}}; \mathbb{Z})$. Let $\Omega_3(A)$ be the image of $\Omega(\Gamma_{\text{odd}}; \mathbb{Z})$ under Ψ_A :

$$\Omega_3(A) = \Psi_A(\Omega(\Gamma_{\text{odd}}; \mathbb{Z})).$$

To be precise,

$$\Omega_3(A) = \left\{ \left\langle \prod_{\substack{s < t \\ m(s,t) \text{ odd}}} R(a_s, a_t)^{n(s,t)} \right\rangle_A \mid \sum_{\substack{s < t \\ m(s,t) \text{ odd}}} n(s,t) \langle s, t \rangle \in \Omega(\Gamma_{\text{odd}}; \mathbb{Z}) \right\}.$$

Proposition 3.18 $\# \Omega_3(A) \leq q_3(\Gamma).$

Let $\Omega(A) = \Omega_1(A) \cup \Omega_2(A) \cup \Omega_3(A)$; hence $\# \Omega(A) \leq p(\Gamma) + q(\Gamma).$

Theorem 3.19 $\Omega(A)$ is a set of generators of $H_2(A; \mathbb{Z}).$

Proof The proof is similar to that of Theorem 3.15, so we omit it. □

3.5 Proof of main results

Theorem 3.1 will follow from the next more precise theorem.

Theorem 3.20 The projection $p: A(\Gamma) \rightarrow W(\Gamma)$ induces an epimorphism between the second integral homologies:

$$p_*: H_2(A(\Gamma); \mathbb{Z}) \twoheadrightarrow H_2(W(\Gamma); \mathbb{Z}).$$

Proof The epimorphism $p: A \rightarrow W$ defined by $p(a_s) = s$ lifts to $\tilde{p}: F_A \rightarrow F_W$. Then by Proposition 3.5, we obtain the explicit formulation

$$p_*: H_2(A; \mathbb{Z}) \cong \frac{N_A \cap [F_A, F_A]}{[F_A, N_A]} \rightarrow \frac{N_W \cap [F_W, F_W]}{[F_W, N_W]} \cong H_2(W; \mathbb{Z}),$$

$$\left\langle \prod_{R(a_s, a_t) \in R_A} R(a_s, a_t)^{n(s,t)} \right\rangle_A \mapsto \left\langle \prod_{R(s,t) \in R_W} R(s,t)^{n(s,t)} \right\rangle_W.$$

We claim that p_* maps $\Omega_i(A)$ onto $\Omega_i(W)$. The claim is obvious for $i = 1, 2$. As for the case $i = 3$, consider the following diagram:

$$\begin{array}{ccc} \Omega(\Gamma_{\text{odd}}; \mathbb{Z}) & \xrightarrow{\Psi_A} & \Omega_3(A) \\ \downarrow \text{mod } 2 & & \downarrow p_* \\ \Omega(\Gamma_{\text{odd}}; \mathbb{Z}_2) & \xrightarrow{\Psi_W} & \Omega_3(W) \end{array}$$

Take $\alpha = \sum_{s < t, m(s,t) \text{ odd}} n(s, t) \langle s, t \rangle \in \Omega(\Gamma_{\text{odd}}; \mathbb{Z})$; then

$$p_* \circ \Psi_A(\alpha) = \left\langle \prod_{\substack{s < t \\ m(s,t) \text{ odd}}} R(s, t)^{n(s,t)} \right\rangle_W \in \frac{N_W \cap [F_W, F_W]}{[F_W, N_W]}.$$

Recall the construction of Φ_W ; we have $\Phi_W(\alpha, \partial\alpha) = p_* \circ \Psi_A(\alpha)$. Thus we obtain

$$\Psi_W(\bar{\alpha}) = p_* \circ \Psi_A(\alpha)$$

by the commutative diagram (3-1). This proves that p_* maps $\Omega_3(A)$ into $\Omega_3(W)$, and the above diagram commutes. Since the mod 2 reduction restricts to a bijection $\Omega(\Gamma_{\text{odd}}; \mathbb{Z}) \rightarrow \Omega(\Gamma_{\text{odd}}; \mathbb{Z}_2)$, and since the horizontal maps are onto by definition, $p_*: \Omega_3(A) \rightarrow \Omega_3(W)$ is onto. The proof is complete. \square

Proof of Theorem 3.1 Consider the composition of epimorphisms

$$(3-2) \quad \mathbb{Z}^{\Omega(A)} \xrightarrow{\phi} H_2(A; \mathbb{Z}) \xrightarrow{p_*} H_2(W; \mathbb{Z}),$$

where $\mathbb{Z}^{\Omega(A)}$ is the free abelian group generated by $\Omega(A)$ and $\phi(\omega) = \omega$ for $\omega \in \Omega(A)$. Taking the tensor product with \mathbb{Z}_2 for terms in the above sequence (3-2), we have

$$\mathbb{Z}_2^{\Omega(A)} \xrightarrow{\phi \otimes \text{id}_{\mathbb{Z}_2}} H_2(A; \mathbb{Z}) \otimes \mathbb{Z}_2 \xrightarrow{p_* \otimes \text{id}_{\mathbb{Z}_2}} H_2(W; \mathbb{Z}) \otimes \mathbb{Z}_2,$$

where $\mathbb{Z}_2^{\Omega(A)}$ is the elementary abelian 2–group generated by $\Omega(A)$ with rank $\#\Omega(A) \leq p(\Gamma) + q(\Gamma)$ and $H_2(W; \mathbb{Z}) \otimes \mathbb{Z}_2 \cong \mathbb{Z}_2^{p(\Gamma)+q(\Gamma)}$ (Theorem 3.2). Since tensoring with \mathbb{Z}_2 preserves surjectivity, this forces $\#\Omega(A) = p(\Gamma) + q(\Gamma)$, and both maps are in fact isomorphisms. Thus $H_2(A(\Gamma); \mathbb{Z}) \otimes \mathbb{Z}_2 \cong \mathbb{Z}_2^{p(\Gamma)+q(\Gamma)}$. On the other hand, we have the following exact sequence by the universal coefficient theorem:

$$0 \rightarrow H_2(A; \mathbb{Z}) \otimes \mathbb{Z}_2 \rightarrow H_2(A; \mathbb{Z}_2) \rightarrow \text{Tor}(H_1(A; \mathbb{Z}), \mathbb{Z}_2) \rightarrow 0,$$

where $\text{Tor}(H_1(A(\Gamma); \mathbb{Z}), \mathbb{Z}_2) = 0$ since $H_1(A(\Gamma); \mathbb{Z})$ is torsion free (Theorem 2.6 and the Remark following). Now we conclude $H_2(A(\Gamma); \mathbb{Z}_2) \cong \mathbb{Z}_2^{p(\Gamma)+q(\Gamma)}$ and finish the proof of Theorem 3.1. \square

As a byproduct of the proof, we have the following corollaries. Recall that $M(\Gamma)$ is the complement of the complexified arrangement of reflection hyperplanes associated to the Coxeter group $W(\Gamma)$. The orbit space $N(\Gamma) = M(\Gamma)/W(\Gamma)$ has fundamental

group $\pi_1(N(\Gamma)) \cong A(\Gamma)$. Let $c: N(\Gamma) \rightarrow K(A(\Gamma), 1)$ be the classifying map. Then c always induces an isomorphism $c_*: H_1(N(\Gamma); \mathbb{Z}) \rightarrow H_1(A(\Gamma); \mathbb{Z})$ and an epimorphism $c_*: H_2(N(\Gamma); \mathbb{Z}) \rightarrow H_2(A(\Gamma); \mathbb{Z})$. We give a sufficient condition on Γ such that c induces an isomorphism $c_*: H_2(N(\Gamma); \mathbb{Z}) \rightarrow H_2(A(\Gamma); \mathbb{Z})$.

Corollary 3.21 *If Γ is such that*

- $P(\Gamma)/\sim$ consists of torsion classes,
- $\Gamma = \Gamma_{\text{odd}}$,
- Γ is a tree,

then

$$H_2(A(\Gamma); \mathbb{Z}) \cong \mathbb{Z}_2^{p(\Gamma)}.$$

Hence c induces an isomorphism $c_*: H_2(N(\Gamma); \mathbb{Z}) \rightarrow H_2(A(\Gamma); \mathbb{Z})$.

Proof Since $N(\Gamma)$ is path-connected and has fundamental group $\pi_1(N(\Gamma)) \cong A(\Gamma)$, there is an exact sequence (see for example, Section II.5, Theorem 5.2 of [3])

$$(3-3) \quad \pi_2(N(\Gamma)) \xrightarrow{h_2} H_2(N(\Gamma); \mathbb{Z}) \xrightarrow{c_*} H_2(A(\Gamma); \mathbb{Z}) \rightarrow 0,$$

where h_2 is the Hurewicz homomorphism. Suppose that Γ satisfies the three conditions; then $q_1(\Gamma) = q_2(\Gamma) = q_3(\Gamma) = 0$. Theorem 2.6 implies that $H_2(N(\Gamma); \mathbb{Z}) \cong \mathbb{Z}_2^{p(\Gamma)}$. Then by Theorem 3.20, $H_2(A(\Gamma); \mathbb{Z})$ sits in the sequence

$$\mathbb{Z}_2^{p(\Gamma)} \twoheadrightarrow H_2(A(\Gamma); \mathbb{Z}) \twoheadrightarrow \mathbb{Z}_2^{p(\Gamma)}.$$

The composition must be an isomorphism; hence $H_2(A(\Gamma); \mathbb{Z}) \cong \mathbb{Z}_2^{p(\Gamma)}$. As a result, c_* must be an isomorphism. □

Corollary 3.22 *If the three conditions in Corollary 3.21 are satisfied, then $p: A \rightarrow W$ induces an isomorphism*

$$p_*: H_2(A; \mathbb{Z}) \rightarrow H_2(W; \mathbb{Z}).$$

Proof This follows from Howlett’s Theorem 3.2, Theorem 3.20 and Corollary 3.21. □

Corollary 3.23 *For any Coxeter graph Γ , the induced map $c_*: H_2(N(\Gamma); \mathbb{Z}) \rightarrow H_2(A(\Gamma); \mathbb{Z})$ becomes an isomorphism after tensoring with \mathbb{Z}_2 .*

Proof By right-exactness of the tensor functor, taking the tensor product with \mathbb{Z}_2 preserves the exactness of (3-3):

$$\pi_2(N(\Gamma)) \otimes \mathbb{Z}_2 \xrightarrow{h_2 \otimes \text{id}_{\mathbb{Z}_2}} H_2(N(\Gamma); \mathbb{Z}) \otimes \mathbb{Z}_2 \xrightarrow{c_* \otimes \text{id}_{\mathbb{Z}_2}} H_2(A(\Gamma); \mathbb{Z}) \otimes \mathbb{Z}_2 \rightarrow 0.$$

Note that $c_* \otimes \text{id}_{\mathbb{Z}_2}$ is an isomorphism as a consequence of Theorem 3.1 and Clancy–Ellis’ Theorem 2.6. □

Example 3.24 The Coxeter graphs of affine type \tilde{D}_n ($n \geq 4$) and \tilde{E}_i ($i = 6, 7, 8$) all satisfy the conditions in Corollary 3.21. Therefore, we compute the second integral homology of the associated Artin groups as follows:

$$H_2(A(\tilde{D}_n); \mathbb{Z}) \cong \begin{cases} \mathbb{Z}_2^6 & \text{for } n = 4, \\ \mathbb{Z}_2^3 & \text{for } n \geq 5, \end{cases} \quad H_2(A(\tilde{E}_i); \mathbb{Z}) \cong \mathbb{Z}_2 \quad \text{for } i = 6, 7, 8.$$

Besides the above cases, the Coxeter graphs of certain hyperbolic Coxeter groups also provide plenty of examples satisfying the conditions in Corollary 3.21. We point out that to the best of the authors’ knowledge, the $K(\pi, 1)$ conjecture has not been proved in the above-mentioned cases.

3.6 Homological stability

We mention a corollary concerning homological stability in the end of this paper. Consider a family of Coxeter graphs $\{\Gamma_i\}_{i \geq 1}$ starting from Γ_1 with a base vertex s_1 , and each Γ_i ($i \geq 2$) is obtained by adding a vertex s_i connected to s_{i-1} by an unlabeled edge. The embedding $\Gamma_i \hookrightarrow \Gamma_{i+1}$ of Coxeter graphs induces an inclusion of Coxeter groups $W(\Gamma_i) \hookrightarrow W(\Gamma_{i+1})$, as well as an inclusion of Artin groups $A(\Gamma_i) \hookrightarrow A(\Gamma_{i+1})$; see [20; 22]. It is known that the families of Artin groups $\{A(A_n)\}$, $\{A(B_n)\}$ and $\{A(D_n)\}$ possess integral cohomological stability [1; 11]. Hepworth proved a more general result for Coxeter groups.

Theorem 3.25 [15] *The map $H_k(W(\Gamma_{n-1})) \rightarrow H_k(W(\Gamma_n))$ is an isomorphism for $2k \leq n$ with arbitrary constant coefficient.*

As for the sequence of Artin groups $\{A(\Gamma_i)\}$, it is not difficult to see that the first integral homology admits stability. We prove a stability result for the second mod 2 homology of the sequence $\{A(\Gamma_i)\}$.

Theorem 3.26 *The map $H_2(A(\Gamma_{n-1}); \mathbb{Z}_2) \rightarrow H_2(A(\Gamma_n); \mathbb{Z}_2)$ is an isomorphism for $n \geq 4$.*

Proof Consider the commutative diagram

$$\begin{array}{ccc}
 H_2(A(\Gamma_{n-1}); \mathbb{Z}_2) & \longrightarrow & H_2(A(\Gamma_n); \mathbb{Z}_2) \\
 \uparrow & & \uparrow \\
 H_2(A(\Gamma_{n-1}); \mathbb{Z}) \otimes \mathbb{Z}_2 & \longrightarrow & H_2(A(\Gamma_n); \mathbb{Z}) \otimes \mathbb{Z}_2 \\
 \downarrow p_* \otimes \text{id}_{\mathbb{Z}_2} & & \downarrow p_* \otimes \text{id}_{\mathbb{Z}_2} \\
 H_2(W(\Gamma_{n-1}); \mathbb{Z}) \otimes \mathbb{Z}_2 & \longrightarrow & H_2(W(\Gamma_n); \mathbb{Z}) \otimes \mathbb{Z}_2
 \end{array}
 \tag{3-4}$$

where the commutativity of the upper square follows from the naturality of the universal coefficient theorem and that of the lower from tensoring the following commutative diagram with \mathbb{Z}_2 :

$$\begin{array}{ccc}
 H_2(A(\Gamma_{n-1}); \mathbb{Z}) & \longrightarrow & H_2(A(\Gamma_n); \mathbb{Z}) \\
 \downarrow p_* & & \downarrow p_* \\
 H_2(W(\Gamma_{n-1}); \mathbb{Z}) & \longrightarrow & H_2(W(\Gamma_n); \mathbb{Z})
 \end{array}
 \tag{3-5}$$

Since all vertical maps in (3-4) are isomorphisms and the bottom horizontal map is an isomorphism when $n \geq 4$ (Theorem 3.25), the top horizontal map is an isomorphism when $n \geq 4$. □

Corollary 3.27 *If Γ_n satisfies the three conditions in Corollary 3.21, then the map $H_2(A(\Gamma_{n-1}); \mathbb{Z}) \rightarrow H_2(A(\Gamma_n); \mathbb{Z})$ is an isomorphism for $n \geq 4$.*

Proof This follows from Corollary 3.22, Theorem 3.25 and the diagram (3-5). □

Acknowledgement Akita was partially supported by JSPS KAKENHI Grant Number 26400077. Liu was supported by JSPS KAKENHI Grant Number 16J00125. They also thank the editor and the referee for valuable suggestions.

References

- [1] **VI Arnol'd**, *The cohomology ring of the colored braid group*, Mat. Zametki 5 (1969) 227–231 MR In Russian; translated in Mathematical Notes 5 (1969), 138–140
- [2] **N Bourbaki**, *Éléments de mathématique, XXXIV: Groupes et algèbres de Lie, Chapitres 4–6*, Actualités Scientifiques et Industrielles 1337, Hermann, Paris (1968) MR

- [3] **K S Brown**, *Cohomology of groups*, Graduate Texts in Mathematics 87, Springer (1982) MR
- [4] **F Callegaro, D Moroni, M Salvetti**, *Cohomology of affine Artin groups and applications*, Trans. Amer. Math. Soc. 360 (2008) 4169–4188 MR
- [5] **F Callegaro, D Moroni, M Salvetti**, *The $K(\pi, 1)$ problem for the affine Artin group of type \tilde{B}_n and its cohomology*, J. Eur. Math. Soc. 12 (2010) 1–22 MR
- [6] **R Charney**, *The Deligne complex for the four-strand braid group*, Trans. Amer. Math. Soc. 356 (2004) 3881–3897 MR
- [7] **R Charney, M W Davis**, *The $K(\pi, 1)$ -problem for hyperplane complements associated to infinite reflection groups*, J. Amer. Math. Soc. 8 (1995) 597–627 MR
- [8] **M Clancy, G Ellis**, *Homology of some Artin and twisted Artin groups*, J. K-Theory 6 (2010) 171–196 MR
- [9] **C De Concini, C Procesi, M Salvetti**, *Arithmetic properties of the cohomology of braid groups*, Topology 40 (2001) 739–751 MR
- [10] **C De Concini, C Procesi, M Salvetti, F Stumbo**, *Arithmetic properties of the cohomology of Artin groups*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. 28 (1999) 695–717 MR
- [11] **C De Concini, M Salvetti**, *Stability for the cohomology of Artin groups*, Adv. Math. 145 (1999) 291–305 MR
- [12] **P Deligne**, *Les immeubles des groupes de tresses généralisés*, Invent. Math. 17 (1972) 273–302 MR
- [13] **G Ellis, E Sköldbberg**, *The $K(\pi, 1)$ conjecture for a class of Artin groups*, Comment. Math. Helv. 85 (2010) 409–415 MR
- [14] **H Hendriks**, *Hyperplane complements of large type*, Invent. Math. 79 (1985) 375–381 MR
- [15] **R Hepworth**, *Homological stability for families of Coxeter groups*, Algebr. Geom. Topol. 16 (2016) 2779–2811 MR
- [16] **R B Howlett**, *On the Schur multipliers of Coxeter groups*, J. London Math. Soc. 38 (1988) 263–276 MR
- [17] **J E Humphreys**, *Reflection groups and Coxeter groups*, Cambridge Studies in Advanced Mathematics 29, Cambridge Univ. Press (1990) MR
- [18] **G Karpilovsky**, *Group representations, Vol 2*, North-Holland Mathematics Studies 177, North-Holland, Amsterdam (1993) MR
- [19] **M Korkmaz, A I Stipsicz**, *The second homology groups of mapping class groups of oriented surfaces*, Math. Proc. Cambridge Philos. Soc. 134 (2003) 479–489 MR
- [20] **H van der Lek**, *The homotopy type of complex hyperplane complements*, PhD thesis, University of Nijmegen (1983)

- [21] **C Okonek**, *Das $K(\pi, 1)$ -Problem für die affinen Wurzelsysteme vom Typ A_n, C_n* , Math. Z. 168 (1979) 143–148 MR
- [22] **L Paris**, *Parabolic subgroups of Artin groups*, J. Algebra 196 (1997) 369–399 MR
- [23] **L Paris**, *Braid groups and Artin groups*, from “Handbook of Teichmüller theory, Vol II” (A Papadopoulos, editor), IRMA Lect. Math. Theor. Phys. 13, Eur. Math. Soc., Zürich (2009) 389–451 MR
- [24] **L Paris**, *$K(\pi, 1)$ conjecture for Artin groups*, Ann. Fac. Sci. Toulouse Math. 23 (2014) 361–415 MR
- [25] **L Paris**, *Lectures on Artin groups and the $K(\pi, 1)$ conjecture*, from “Groups of exceptional type, Coxeter groups and related geometries” (N S N Sastry, editor), Springer Proc. Math. Stat. 82, Springer (2014) 239–257 MR

*Department of Mathematics, Hokkaido University
Sapporo, 060-0810, Japan*

akita@math.sci.hokudai.ac.jp, liu@math.sci.hokudai.ac.jp

Received: 19 February 2017 Revised: 28 June 2017