

# Noncharacterizing slopes for hyperbolic knots

KENNETH L BAKER  
KIMIHIKO MOTEGI

A nontrivial slope  $r$  on a knot  $K$  in  $S^3$  is called a characterizing slope if whenever the result of  $r$ -surgery on a knot  $K'$  is orientation-preservingly homeomorphic to the result of  $r$ -surgery on  $K$ , then  $K'$  is isotopic to  $K$ . Ni and Zhang ask: for any hyperbolic knot  $K$ , is a slope  $r = p/q$  with  $|p| + |q|$  sufficiently large a characterizing slope? In this article, we prove that if we can take an unknot  $c$  so that  $(0, 0)$ -surgery on  $K \cup c$  results in  $S^3$  and  $c$  is not a meridian of  $K$ , then  $K$  has infinitely many noncharacterizing slopes. As the simplest known example, the hyperbolic, two-bridge knot  $8_6$  has no integral characterizing slopes. This answers the above question in the negative. We also prove that any L-space knot never admits such an unknot  $c$ .

57M25

## 1 Introduction

Let  $K$  be a knot in the oriented 3-sphere  $S^3$ . Denote by  $K(p/q)$  the 3-manifold obtained by  $p/q$ -Dehn surgery on  $K$  which has the orientation induced from  $S^3$ . We call  $p/q \in \mathbb{Q}$  a *characterizing slope* for  $K$  if whenever  $K'(p/q)$  is orientation-preservingly homeomorphic to  $K(p/q)$ , then  $K'$  is isotopic to  $K$ . For the trivial knot, Gordon [10] conjectured that every nontrivial slope  $p/q \in \mathbb{Q}$  is a characterizing slope. Kronheimer, Mrowka, Ozsváth and Szabó [19] proved this conjecture in the positive using Seiberg–Witten monopoles. See Ozsváth and Szabó [28; 31] for alternative proofs using Heegaard Floer homology. Furthermore, Ozsváth and Szabó [30] showed that for the trefoil knot and the figure-eight knot, every nontrivial slope is a characterizing slope.

On the other hand, it is known that many knots have noncharacterizing slopes. The first such example was given by Lickorish [21]. Some torus knots have noncharacterizing slopes. For instance, 21-surgeries on  $T_{5,4}$  and  $T_{11,2}$  produce the same oriented 3-manifold, and hence 21 is a noncharacterizing slope for both  $T_{5,4}$  and  $T_{11,2}$  [26]. However, Ni and Zhang [26] prove that for a torus knot  $T_{r,s}$  with  $r > s > 1$ , a slope  $p/q$  is a characterizing slope if  $p/q > 30(r^2 - 1)(s^2 - 1)/67$ . Later, McCoy [23]

lowers the bound to  $43(rs - r - s)/4$ . See also McCoy [24]. This suggests that for a given knot  $K$ , sufficiently large slopes should be characterizing ones. For hyperbolic knots, Ni and Zhang ask the following:

**Question 1.1** [26] *Let  $K$  be a hyperbolic knot. Is a slope  $r = p/q$  with  $|p| + |q|$  sufficiently large a characterizing slope of  $K$ ?*

**Remark 1.2** For any given hyperbolic knot  $K$ , there is a number  $N_K > 0$  such that a slope  $p/q$  with  $|p| + |q| > N_K$  has a special geometric meaning due to Thurston's hyperbolic Dehn surgery theorem; see Benedetti and Petronio [4], Boileau and Porti [5], Petronio and Porti [32] and Thurston [34; 35]. For such a slope  $p/q$ , the 3-manifold  $K(p/q)$  is hyperbolic, and the surgery dual to  $K$  is the unique shortest closed geodesic in  $K(p/q)$ . Hence for any finite family of hyperbolic knots  $\mathcal{K}$ , there is a number  $N_{\mathcal{K}} > 0$  such that any slope  $p/q$  with  $|p| + |q| > N_{\mathcal{K}}$  is a characterizing slope for every knot  $K \in \mathcal{K}$ .

The purpose in this article is to answer [Question 1.1](#) in the negative. To this end, we need to construct a hyperbolic knot with infinitely many noncharacterizing slopes. The theorem below gives a sufficient condition for a knot  $K$  to have infinitely many noncharacterizing slopes.

**Theorem 1.3** *Let  $K$  be a knot in  $S^3$ . Suppose that we can take an unknot  $c$  disjoint from  $K$  so that  $(0, 0)$ -surgery on  $K \cup c$  results in  $S^3$  and  $c$  is not a meridian of  $K$ . Then  $K$  has infinitely many noncharacterizing slopes.*

Note that the condition that  $(0, 0)$ -surgery on  $K \cup c$  results in  $S^3$  implies that  $|\text{lk}(K, c)| = 1$  for homological reasons, where  $\text{lk}(K, c)$  denotes the linking number between  $K$  and  $c$ . Of course, if  $c$  is a meridian of  $K$ , then the result of  $(0, 0)$ -surgery on  $K \cup c$  is always  $S^3$ ; see [Lemma 2.4](#).

As shown by [Theorem 2.5](#), if we find a link  $K \cup c$  which satisfies the condition in [Theorem 1.3](#), then we have infinitely many distinct knots, each of which has infinitely many noncharacterizing slopes. We apply this to present explicit examples.

First, for comparison, recall that every nontrivial slope is a characterizing slope for a trefoil knot and the figure-eight knot [30], which are genus-one fibered knots. If we drop one of these conditions, we have:

- Example 1.4** (1) Let  $K$  be the hyperbolic, fibered knot  $9_{42}$  in Rolfsen's table, which has genus two. Then every integer except possibly 2 is not a characterizing slope for  $K$ .
- (2) Let  $K$  be the hyperbolic, genus-one pretzel knot  $P(-3, 3, 5)$ , which is not fibered. Then every integer except possibly 0 is not a characterizing slope for  $K$ .

A modification of the above examples leads us to demonstrate:

**Theorem 1.5** *There exists a hyperbolic knot for which every integral slope is a non-characterizing slope. In particular, every integral slope is not a characterizing slope for the hyperbolic, two-bridge knot  $8_6$  in Rolfsen's table.*

Such a phenomenon can occur for prime satellite knots and composite knots as well. More precisely, we are able to prove:

- Theorem 1.6** (1) *Given a nontrivial knot  $k$ , there exists a prime satellite knot with companion knot  $k$  for which every integral slope is a noncharacterizing slope.*
- (2) *Given a nontrivial knot  $k$ , there exists a composite knot with  $k$  a connected summand for which every integral slope is a noncharacterizing slope.*

Among known examples, the knot  $8_6$  is the simplest knot (with respect to crossing numbers) which has infinitely many noncharacterizing slopes. So we would like to ask:

**Question 1.7** *Are there any knots of crossing number less than 8 that have infinitely many noncharacterizing slopes?*

It is natural to ask which knots  $K$  admit an unknot  $c$  that satisfies the condition in [Theorem 1.3](#).

**Theorem 1.8** *Let  $K \cup c$  be a two-component link in  $S^3$  with unknotted component  $c$  which is not a meridian of  $K$ . Suppose that  $(0, 0)$ -surgery on  $K \cup c$  results in  $S^3$ . Then  $K$  is not an  $L$ -space knot.*

In the last section, we will give further questions concerning characterizing and non-characterizing slopes for knots.

Throughout the paper we will use  $N(*)$  to denote a tubular neighborhood of  $*$  and  $\mathcal{N}(*)$  to denote the interior of  $N(*)$  for notational simplicity.

**Acknowledgements** We would like to thank the referee for a careful reading and useful comments.

Baker was partially supported by a grant from the Simons Foundation (#209184 to Kenneth L Baker). Motegi was partially supported by JSPS KAKENHI Grant Number JP26400099 and Joint Research Grant of Institute of Natural Sciences at Nihon University for 2016.

## 2 Noncharacterizing slopes and twist families of surgeries

In this section, we establish the general principle [Theorem 2.1](#) and its extension [Theorem 2.5](#), from which [Theorem 1.3](#) follows. Throughout this article, for two oriented 3-manifolds  $M$  and  $N$ , by  $M \cong N$  we mean that  $M$  is orientation-preservingly homeomorphic to  $N$ .

**Theorem 2.1** *Let  $k \cup c$  be a two-component link in  $S^3$  such that  $c$  is unknotted. Suppose that  $(0, 0)$ -surgery on  $k \cup c$  results in  $S^3$ . Let  $K$  be the knot in  $S^3$  which is surgery dual to  $c$ , the image of  $c$ , in the surgered  $S^3$ , and let  $k_n$  be the knot obtained from  $k$  by twisting  $n$  times along  $c$ . Then  $K(n) \cong k_n(n)$  for all integers  $n$ .*

*Moreover, if  $c$  is not a meridian of  $k$ , then  $K \not\cong k_n$  for all but finitely many integers  $n$ .*

**Proof** Since  $(0, 0)$ -surgery on  $k \cup c$  is  $S^3$ , a homology calculation shows that  $|\text{lk}(k, c)| = 1$ . Performing  $(-1/n)$ -surgery along  $c$  takes the knot  $k$  with the surgery slope 0 to a knot  $k_n$  with a surgery slope  $n = 0 + n(\text{lk}(k, c))^2$ ; ie an  $n$ -twist along  $c$  converts a knot-slope pair  $(k, 0)$  into another knot-slope pair  $(k_n, n)$ . Thus we obtain a twist family of knot-slope pairs  $\{(k_n, n)\}$ . Let  $V$  be the solid torus  $S^3 - \mathcal{N}(c)$  which contains  $k$  in its interior. Observe that  $V(k; 0) \cong V(k_n; n)$  for all  $n$ .

Let  $(\mu_c, \lambda_c)$  be a preferred meridian-longitude pair of  $c \subset S^3$ , oriented with the right-handed orientation (so that if  $c$  is oriented in the same direction as  $\lambda_c$  in  $\mathcal{N}(c)$ , then  $\text{lk}(\mu_c, c) = 1$ ). Note that  $\lambda_c$  represents the 0-slope on  $\mathcal{N}(c)$ , and  $\lambda_c$  bounds a meridian disk of the solid torus  $V$ . Let  $c_n$  be the surgery dual to the  $(-1/n)$ -surgery on  $c$  (ie a core of the filled solid torus) with meridian  $\mu_n$ , the  $(-1/n)$ -surgery slope of  $c$  in  $\partial V$ . These curves  $\mu_n$  are each longitudes of  $V$  and satisfy  $[\mu_n] = -[\mu_c] + n[\lambda_c] \in H_1(\partial V)$ ; hence  $[\mu_0] = -[\mu_c]$ .

Since  $k$  wraps algebraically once in  $V$ , a preferred longitude of  $k \subset V \subset S^3$  is homologous to  $\mu_c$  in  $V - \mathcal{N}(k)$ . Hence  $\mu_c$  is null-homologous in  $V(k; 0)$ .

Let  $K$  be the surgery dual to  $c$  with respect to  $\lambda_c$ -surgery. (Adapting the above notation,  $K$  may be regarded as  $c_\infty$ .) Since  $(0, 0)$ -surgery on  $k \cup c$  results in  $S^3$ , we have that  $K$  is a knot in this surgered  $S^3$  with exterior  $S^3 - \mathcal{N}(K) = V(k; 0)$  and meridian  $\lambda_c$ . Because  $\mu_c$  is null-homologous in  $V(k; 0)$ , it is the boundary of a Seifert surface for  $K$ .

With right-handed orientation, a preferred meridian-longitude pair for  $K$  in  $S^3$  is given by  $(\lambda_c, -\mu_c)$ . Thus  $[\mu_n] = -[\mu_c] + n[\lambda_c] = n[\lambda_c] + (-[\mu_c])$  corresponds to a slope  $n$  with respect to the preferred meridian-longitude pair  $(\lambda_c, -\mu_c)$ . Therefore,  $k_n(n) = K(n)$  for all integers  $n$ .

If  $c$  is not a meridian of  $k$ , then since  $\text{lk}(k, c) \neq 0$ , any disk bounded by  $c$  intersects  $k$  more than once. Then it follows from [18] that there are only finitely many  $n$  such that  $k_n$  is isotopic to  $K$ . □

**Remark 2.2** Gompf and Miyazaki [9] had previously utilized the mirror of the knot  $K$  associated to  $k$  as described in Theorem 2.1 for a satellite construction of ribbon knots that generalizes the connected sum of a knot and its mirror.

Let  $k \cup c$  be a link as in Theorem 2.1; ie  $c$  is unknotted and the result of  $(0, 0)$ -surgery on  $k \cup c$  is  $S^3$ . In Theorem 2.1,  $K$  denotes the surgery dual to  $c$ . Similarly we denote by  $C$  the surgery dual to  $k$ . Thus we have the surgery dual link  $C \cup K$  to  $k \cup c$  in the surgered  $S^3$ .

**Lemma 2.3** *Let  $k \cup c$  be a link as in Theorem 2.1 with surgery dual link  $C \cup K$ . Then  $C$  is unknotted in  $S^3$ .*

**Proof** After 0-surgery on  $c$ , we have that  $k$  becomes some knot in  $c(0) = S^1 \times S^2$ . Since a nontrivial surgery (corresponding to the 0-surgery) on  $k \subset S^1 \times S^2$  yields  $S^3$ , due to Gabai [8, Corollary 8.3], it turns out that  $k$  (as a knot in  $S^1 \times S^2$ ) is an  $S^1$ -fiber in some product structure  $\mathcal{P}$  of  $S^1 \times S^2$ , and intersects an  $S^2$ -fiber in  $\mathcal{P}$  exactly once. As usual, we may isotope an  $S^2$ -fiber in  $\mathcal{P}$  to  $S = S^2 \times \{0\}$  in the original product structure; the knot  $k$  is simultaneously isotoped to a knot intersecting  $S$  in a single point. Then a further ambient isotopy, possibly with “light bulb” moves which are accomplished by an ambient isotopy of the type illustrated in Figure 1 (see [33, page 257]), enables us to deform  $k$  to an  $S^1$ -fiber in the original product structure of  $c(0) = S^1 \times S^2$ . Thus the surgery dual  $C$  to  $k$  in  $(k \cup c)(0, 0) = S^3$  is an unknot while the surgery dual  $K$  to  $c$  is not necessarily unknotted in this  $S^3$ . Figure 2 illustrates such a situation. □




Figure 1: The “light bulb” move in  $S^1 \times S^2$

In the special case where  $c$  is a meridian of  $k$ , we have:

**Lemma 2.4** *Let  $k \cup c$  be a two-component link in  $S^3$  such that  $c$  is a meridian of  $k$ . Then  $(0, 0)$ –surgery on  $k \cup c$  results in  $S^3$  with its surgery dual link  $C \cup K$ , for which  $C$  is a meridian of  $K$ , and  $K$  is isotopic to  $k$  in  $S^3$ .*

**Proof** This is essentially shown in [9, page 119] without a proof. So for completeness, we give a proof. Since  $c$  is a meridian of  $k$ , we may straighten  $k$  in  $c(0) = S^1 \times S^2$  using light bulb moves and isotopies; the framing 0 of  $k$  is changed into some even integer, and the image  $K$  of  $c$  in  $c(0) = S^1 \times S^2$  intersects  $\{x\} \times S^2$  once for some  $x \in S^1$ . Then we see that  $(0, 0)$ –surgery on  $k \cup c$  results in  $S^3$  with its surgery dual  $C \cup K$  in which the dual  $C$  to  $k$  is a meridian of  $K$  in  $S^3$ . Let us see that  $K$  is isotopic to  $k$ . Since  $c$  is a meridian of  $k$ , the exterior  $S^3 - \mathcal{N}(k \cup c)$  is the union of the 2–fold composing space  $X$  (ie  $[\text{disk with 2–holes}] \times S^1$ ) and a knot space  $E$  which is homeomorphic to  $S^3 - \mathcal{N}(k)$ . Note that a regular fiber  $t$  of  $X$  which lies in  $\partial N(c)$  intersects a meridian  $\mu_c$  exactly once, and a regular fiber  $t$  of  $X$  which lies in  $\partial N(k)$  coincides with a meridian  $\mu_k$ . The former condition implies  $X \cup N(c) \cong S^1 \times S^1 \times [0, 1]$ . After  $(0, 0)$ –surgery on  $k \cup c$ , we obtain the dual link  $C \cup K$  in this surgered  $S^3$ . Observe that the regular fiber  $t$  which lies in  $\partial N(K)$  coincides with a meridian  $\mu_K$ , and the regular fiber  $t$  of  $X$  which lies in  $\partial N(C)$  intersects a meridian  $\mu_C$  exactly once. The latter condition implies that  $X \cup N(C) \cong S^1 \times S^1 \times [0, 1]$ . Hence

$$S^3 - \mathcal{N}(K) = (k \cup c)(0, 0) - \mathcal{N}(K) = E \cup (X \cup N(C)) \cong E \cong S^3 - \mathcal{N}(k).$$

Thus Gordon and Luecke [11, Theorem 1] show  $K$  is isotopic to  $k$ . □

In the proof of [Theorem 2.1](#), we observe that  $(k \cup c)(0, -\frac{1}{n}) \cong (C \cup K)(\frac{1}{0}, n)$ ,  $(k \cup c)(0, -\frac{1}{n}) \cong k_n(n)$  and  $(C \cup K)(\frac{1}{0}, n) \cong K(n)$ . Starting with  $m$ –surgery instead of 0–surgery on  $k$ , the argument in the proof of [Theorem 2.1](#) leads us to the following extension. Recall that the surgery dual  $C$  to  $k$  is unknotted in  $S^3$  by [Lemma 2.3](#). In

what follows,  $K_m$  denotes the knot obtained from  $K$  by twisting  $m$  times along the unknot  $C$ .

**Theorem 2.5** *Let  $k \cup c$  be a link as in Theorem 2.1 with surgery dual link  $C \cup K$ , where  $K$  is dual to  $c$  and  $C$  is dual to  $k$ . Then for any integers  $m$  and  $n$ ,*

$$K_m(n + m) \cong k_n(m + n).$$

*Moreover, if  $c$  is not a meridian of  $k$ , then each family  $\{K_m\}$  and  $\{k_n\}$  contains infinitely many distinct knots, each of which has only finitely many integral characterizing slopes.*

**Proof** Observe that  $S^3 - \mathcal{N}(k \cup c) = S^3 - \mathcal{N}(C \cup K)$  and the meridian-longitude pairs  $(\mu_k, \lambda_k)$  for  $k$  and  $(\mu_c, \lambda_c)$  for  $c$  become meridian-longitude pairs  $(\lambda_k, -\mu_k)$  for  $C$  and  $(\lambda_c, -\mu_c)$  for  $K$ . The latter correspondence was shown in the proof of Theorem 2.1. For the former correspondence, by definition,  $\lambda_k$  becomes a meridian of  $C$ , the surgery dual to  $k$ . Observe also that  $\mu_k$  is homologous to  $\lambda_c$  (because  $|\text{lk}(k, c)| = 1$ ), which bounds a disk of the filled solid torus after 0–surgery on  $c$ . Thus  $\mu_k$  is a preferred longitude of  $C$ . Now the orientation convention gives the desired result. We note here that the above observation shows that  $(0, 0)$ –surgery on  $C \cup K$  yields  $S^3$  with surgery dual  $k \cup c$ . In particular,  $|\text{lk}(K, C)| = 1$ .

Then we have the surgery relation

$$K_m(n + m) \cong (C \cup K)\left(-\frac{1}{m}, n\right) \cong (k \cup c)\left(m, -\frac{1}{n}\right) \cong k_n(m + n)$$

as claimed.

Following Lemma 2.4, if  $C$  is a meridian of  $K$ , then  $c$  is a meridian of  $k$ . Thus if  $c$  is not a meridian of  $k$ , then  $C$  is not a meridian of  $K$  either. Since  $|\text{lk}(k, c)| = 1$  and  $|\text{lk}(K, C)| = 1$ , the wrapping numbers of  $k$  about  $c$  and  $K$  about  $C$  are at least 2. Then [18, Theorem 3.2] implies that each twist family of knots  $\{k_n\}$  and  $\{K_m\}$  partitions into infinitely many distinct knot types containing finitely many members. Therefore, since  $K_m(n + m) \cong k_n(m + n)$ , each knot in these two families has only finitely many characterizing slopes.  $\square$

**Proof of Theorem 1.3** By the assumption, the link  $K \cup c$  satisfies the condition in Theorem 2.5, where  $K$  should be read as  $k$ ; ie notationally  $K$  and  $k$  are exchanged. Then we have  $k_m(m) \cong K(m)$  by putting  $n = 0$ . Since  $\{k_m\}$  contains infinitely many distinct knots,  $K$  has infinitely many noncharacterizing slopes.  $\square$

### 3 Alexander polynomials of knots in twist families

We take  $\Delta_{A \cup B}(x, y)$  to be the *symmetrized* multivariable Alexander polynomial of the *oriented* two-component link  $A \cup B$ , where  $x$  corresponds to the oriented meridian  $\mu_A$  of  $A$  and  $y$  corresponds to the oriented meridian  $\mu_B$  of  $B$ . Due to the symmetrization,

$$\Delta_{A \cup B}(x, y) = \Delta_{A \cup B}(x^{-1}, y^{-1}) = \Delta_{-A \cup -B}(x, y).$$

However, in general,  $\Delta_{A \cup B}(x, y) \neq \Delta_{A \cup -B}(x, y)$ .

Recall that if  $k \cup c$  is a link in  $S^3$  such that  $c$  is unknotted and  $(0, 0)$ -surgery on  $k \cup c$  yields  $S^3$  with surgery dual link  $C \cup K$ , then  $C$  is also unknotted and  $|\text{lk}(K, C)| = 1$ .

**Proposition 3.1** *Assume  $k \cup c$  is an oriented two-component link with  $\text{lk}(k, c) = 1$  such that  $c$  is an unknot. Further assume  $(0, 0)$ -surgery on  $k \cup c$  results in  $S^3$  with surgery dual  $C \cup K$ , where  $K$  is dual to  $c$  and  $C$  is dual to  $k$ , oriented so that  $\text{lk}(K, C) = 1$ . Then  $\Delta_{K \cup C}(x, y) = \Delta_{k \cup c}(x, y^{-1})$ , or equivalently,  $\Delta_{k \cup c}(x, y) = \Delta_{K \cup C}(x, y^{-1})$ .*

**Proof** Let us write  $\mu_J$  and  $\lambda_J$  for the meridian and preferred longitude of an oriented knot  $J$  in  $S^3$  which we view as oriented curves in  $\partial\mathcal{N}(J)$  such that  $\text{lk}(J, \mu_J) = 1$  and  $\lambda_J$  is homologous to  $J$ . Let  $X = S^3 - \mathcal{N}(k \cup c)$  be the exterior of the link  $k \cup c$ . Since the linking number of  $k \cup c$  is 1, in  $H_1(X; \mathbb{Z})$ , we have that  $[\mu_k] = [\lambda_c]$  and  $[\mu_c] = [\lambda_k]$ . Furthermore, these homologies are realized by oriented Seifert surfaces  $\Sigma_c$  and  $\Sigma_k$  that are each punctured once by  $k$  and  $c$ , respectively. In particular, restricting to  $X$ , we have that  $\partial\Sigma_c = \lambda_c - \mu_k$  and  $\partial\Sigma_k = \lambda_k - \mu_c$ .

Since  $K$  is the surgery dual to  $c$  with respect to 0-surgery on  $c$ , and  $C$  is the surgery dual to  $k$  with respect to 0-surgery on  $k$ , we have  $X = S^3 - \mathcal{N}(K \cup C)$ . Upon surgery, the punctured Seifert surfaces  $\Sigma_k$  and  $\Sigma_c$  respectively cap off to oriented Seifert surfaces  $\Sigma_K$  and  $\Sigma_C$  for  $K$  and  $C$ . Using these surfaces to orient  $K$  and  $C$  and thus their meridians and longitudes, we obtain that  $(\mu_K, \lambda_K) = (\lambda_c, -\mu_c)$  and  $(\mu_C, \lambda_C) = (\lambda_k, -\mu_k)$ . Therefore,  $[\mu_K] = [\mu_k]$  and  $[\mu_C] = [\mu_c]$  in  $H_1(X; \mathbb{Z})$ . However, since  $[\lambda_C] = -[\mu_k] = -[\mu_K]$ , we find that  $\text{lk}(K, C) = -1$ . To orient  $K$  and  $C$  so that  $\text{lk}(K, C) = 1$ , we must flip the orientation on  $C$ , say. Then for this correctly oriented  $C$ , we have  $[\mu_C] = -[\mu_c]$ . Hence  $\Delta_{K \cup C}(x, y) = \Delta_{k \cup c}(x, y^{-1})$ .  $\square$

We recall also the following twisting formula for Alexander polynomials.

**Proposition 3.2** [1, Theorem 2.1] *Let  $k \cup c$  be an oriented two-component link such that  $c$  is an unknot and  $\omega = \text{lk}(k, c) > 0$ . Denote by  $k_n$  a knot obtained from  $k$  by an  $n$ -twist along  $c$ . Then  $\Delta_{k_n}(t) = \Delta_{k \cup c}(t, t^{n\omega})$ .*

Propositions 3.1 and 3.2 lead us some symmetry among Alexander polynomials of  $k_n$  and  $K_n$ .

**Corollary 3.3** *Let  $k \cup c$  be a link as in Theorem 2.1 with surgery dual link  $C \cup K$ , where  $K$  is dual to  $c$  and  $C$  is dual to  $k$ . Then for the twist families of knots  $\{k_n\}$  and  $\{K_n\}$ , we have  $\Delta_{k_n}(t) = \Delta_{K_{-n}}(t)$ . In particular,  $\Delta_k(t) = \Delta_K(t)$ .*

**Proof** We may orient  $k$  and  $c$  so that  $\text{lk}(k, c) = 1$ . Then Propositions 3.1 and 3.2 show that  $\Delta_{k_n}(t) = \Delta_{k \cup c}(t, t^n) = \Delta_{K \cup C}(t, t^{-n}) = \Delta_{K_{-n}}(t)$ . In particular, putting  $n = 0$ , we have  $\Delta_k(t) = \Delta_K(t)$ . □

## 4 Examples

In this section, we will provide examples which satisfy the condition in Theorem 2.1, and hence Theorem 2.5. Example 1.4 follows from Examples 4.1 and 4.3. A slight modification gives a nonhyperbolic example, Example 4.5, that demonstrates Theorem 1.6. We will make a further modification of the first example to present Example 4.6 which implies Theorem 1.5.

Let us take a two-component link  $k \cup c$  in  $S^3$  with  $|\text{lk}(k, c)| = 1$  as in Figure 2. To perform 0-surgery on the unknot  $c$ , we first remove  $N(c)$  and glue it back to  $V = S^3 - N(c)$  so that a meridian of  $N(c)$  is identified with a meridian of  $V$ , a preferred longitude of  $c$ . Then the union of meridian disks of  $N(c)$  and  $V$  forms a nonseparating 2-sphere  $S$  in  $c(0) = S^1 \times S^2$ . The second picture from the left of Figure 2 describes  $c(0) = S^1 \times S^2$  in which the bottom 2-sphere and the top 2-sphere are identified (without twisting) to result in the nonseparating 2-sphere  $S$ . From the second to the seventh picture, since the total space is  $S^1 \times S^2$  rather than  $S^3$ , we do not put extra labels to corresponding components. We apply light bulb moves from the third to the fourth and from the fifth to the sixth picture of Figure 2. In the second picture from the right of Figure 2, the straight knot is the image of  $k$  in  $c(0) = S^1 \times S^2$ , and 0-surgery on this knot gives  $S^3$ . This  $S^3$  resulting from  $(0, 0)$ -surgery on  $k \cup c$  is shown with the surgery dual link  $C \cup K \subset S^3$  in the rightmost picture of Figure 2. Thus  $k \cup c$  satisfies the condition in Theorem 2.1, and  $K(n) \cong k_n(n)$  does hold for all integers  $n$ .



Hence by Proposition 3.2, we have

$$(\star) \quad \Delta_{k_n}(t) = \Delta_{k \cup c}(t, t^n) = -(t^{-1} - 2 + t)t^{-n} + 1 - (t^{-1} - 2 + t)t^n.$$

In particular, since the Alexander polynomial of  $k_n$  varies depending on  $n$ , we have that  $c$  is not a meridian of  $k$ .

Let us generalize this following Theorem 2.5. Let  $K_m$  be a knot obtained from  $K$  by an  $m$ -twist along  $C$ . Then Theorem 2.5 asserts that  $K_m(n + m) \cong k_n(m + n)$  for any integers  $m$  and  $n$ . Figure 3 demonstrates this fact pictorially.

Let us choose an integer  $m$  arbitrarily. Observe that in this example, we have  $K_m = k_m$ ; see Figure 3. Hence if  $k_n = K_m$  for some integer  $n$ , then  $k_n = k_m$ . Thus  $\Delta(k_n) \doteq \Delta(k_m)$ , and  $(\star)$  implies that  $n = \pm m$ . Thus at most  $k_m$  and  $k_{-m}$  can be isotopic to  $K_m$ . Since  $K_m(n + m) \cong k_n(m + n)$  for all integers  $m$  and  $n$ , we have the following:

- For a given integer  $m$ , every integral slope except possibly 0 and  $2m$  fails to be a characterizing slope for  $K_m$ .
- If furthermore  $K_{-m} \neq K_m$ , then 0 will fail to be a characterizing slope as well.

**Example 4.1** (genus-one, nonfibered knot) Let us choose  $m = 0$  in the above. Then  $K_0(n) = k_n(n)$  for all integers  $n$ , and as mentioned above, every nonzero integral slope fails to be a characterizing slope for  $K_0$ . In Figure 4, we identify  $K_0 = k_0$  as the pretzel knot  $P(-5, 3, -3)$ , which is known to be hyperbolic by [27]. The pretzel knot  $P(-5, 3, -3)$  is a genus-one knot, but it is not fibered. Seifert’s algorithm easily produces a genus-one Seifert surface of  $P(-5, 3, -3)$ .

**Remark 4.2** Notably, the (mirror of the) knot  $P(-5, 3, -3)$  was the basic example of the first two families of nonstrongly invertible knots with a small Seifert fibered space surgery [22]. Indeed,  $(-1)$ -surgery on  $P(-5, 3, -3)$  is the Seifert fibered space  $S^2(-\frac{2}{5}, \frac{3}{4}, -\frac{1}{3})$ .

Since  $P(-5, 3, -3)$  is the knot  $K_0$ , and  $K_0(n) = k_n(n)$  for all integers  $n$ , we have  $K_0(-1) = k_{-1}(-1) = K_{-1}(-1)$ . Thus  $(-1)$ -surgery on  $K_{-1}$  is the same Seifert fibered space. SnapPy recognizes the complement of  $K_{-1}$  as the mirror of the census manifold  $o9_{34801}$ . Furthermore, SnapPy reports this manifold as asymmetric, implying that  $K_{-1}$  is neither strongly invertible nor cyclically periodic, and hence cannot be embedded in a genus-2 Heegaard surface; see [7, Lemma 7.4].




Figure 5: The knot  $K_1$  in Figure 3 is isotoped into a presentation as the 9 crossing Montesinos knot  $M(\frac{1}{3}, -\frac{1}{2}, \frac{2}{5})$  which may be recognized as the knot  $9_{42}$  in Rolfsen’s table [33].

**Example 4.3** (fibered, genus-two knot) By choosing  $m = 1$  instead of 0, we obtain a knot  $K_1$  for which we have  $K_1(n + 1) = k_n(1 + n)$  for all integers  $n$ . As we mentioned, every integral slope other than 0 or 2 are noncharacterizing slope for  $K_1$ . In Figure 5, we recognize the knot  $K_1$  as the 9–crossing Montesinos knot  $M(\frac{1}{3}, -\frac{1}{2}, \frac{2}{5})$  which is the knot  $9_{42}$  in Rolfsen’s table [33]. Following [27],  $K_1$  is a hyperbolic knot. The knot  $9_{42}$  is a fibered knot, but it has genus two [13, Theorem 3.2].

Now let us show that 0–slope is also a noncharacterizing slope for  $K_1$ . Since  $K_1(0) \cong k_{-1}(0)$ , it is sufficient to see that  $K_1 \neq k_{-1}$ . Recall that  $K_m = k_m$  for any  $m$ . Alexander polynomials distinguish  $k_1$  from  $k_n$  for all  $n \neq \pm 1$ ; see (★). The Jones polynomial<sup>2</sup> will however distinguish  $k_1 = K_1$  and  $k_{-1}$ :

$$V_{k_1}(q) = q^{-3} - q^{-2} + q^{-1} - 1 + q - q^2 + q^3,$$

while

$$V_{k_{-1}}(q) = q^{-1} + q^{-3} - q^{-6} - q^{-8} + q^{-9} - q^{-10} + q^{-11}.$$

(As noted in Remark 4.2, SnapPy also identifies the complement of  $K_{-1} = k_{-1}$  as distinct from the complement of  $K_1 = 9_{42}$ , thereby distinguishing these knots.) Hence all integers except possibly 2 are noncharacterizing slopes for the hyperbolic knot  $K_1 = 9_{42}$ .

**Question 4.4** *Is 0 a characterizing slope for  $P(-3, 3, 5)$ ? Is 2 a characterizing slope for  $9_{42}$ ?*

Next we provide examples of nonhyperbolic knots such that all integral slopes are noncharacterizing slopes, from which Theorem 1.6 follows.

<sup>2</sup>Kodama’s software KNOT [17] was used confirm the Jones polynomials of knots.




Figure 6: The sum of 1–string tangles  $\tau'$  and  $\tau''$  is the connected sum  $k = k' \# k''$ .

**Example 4.5** (nonhyperbolic example) Given any nontrivial knot  $k''$ , let us take a two-component link  $k \cup c$  as in Figure 6, where  $k$  is a connected sum of a knot  $k'$  (which is  $k$  in Figure 2, the closure of the 1–string tangle  $\tau'$ ) and the nontrivial knot  $k''$  (the closure of the 1–string tangle  $\tau''$ ).

Then as in Figure 2, we see that  $(0, 0)$ –surgery on  $k \cup c$  gives  $S^3$  with the surgery dual  $C \cup K$ . Actually, we follow the isotopy and light bulb moves as indicated in Figure 2 to obtain the sixth figure, in which  $k$  is almost an  $S^1$  fiber, but it has the connected summand  $k''$  (ie the knotted arc  $\tau''$ ). Then we apply further light bulb moves to  $k$  so that it becomes an  $S^1$  fiber;  $K$  becomes a satellite knot with  $k''$  as a companion knot; see Lemma 2.4. Then by Theorem 2.5,  $K_m(n + m) \cong k_n(m + n)$  for all integers  $m$  and  $n$ .

It is easy to observe that  $k_n$  is a connected sum  $k'_n \# k''$ , where  $k'_n$  is a knot obtained from  $k'$  by an  $n$ –twist along  $c$ . For instance,  $k_0 = P(-5, 3, -3) \# k''$  and  $k_1 = 9_{42} \# k''$ . Since  $k'_n$  is nontrivial for all integers  $n$  by  $(\star)$ ,  $k_n$  is not prime for all integers  $n$ .

On the other hand, we show that  $K_m$  is prime for all integers  $m$ . (We note that, by construction,  $K_m$  has  $k''$  as a companion knot for every integer  $m$ .) In the following, we fix an integer  $m$  arbitrarily. First we observe that  $k_n(m + n)$  is obtained by gluing  $E(k'_n)$  and  $E(k'')$  along their boundary tori. Recall that the exterior  $E(k_n)$  may be expressed as the union of the 2–fold composing space  $X$  (ie  $[\text{disk with 2–holes}] \times S^1$ ) and two knot spaces  $E(k'_n)$  and  $E(k'')$ . We note that  $\partial X$  consists of  $\partial E(k_n)$ ,  $\partial E(k'_n)$  and  $\partial E(k'')$ , and a regular fiber in  $\partial X \cap \partial E(k_n)$  is a meridian of  $k_n$ . Since the surgery slope  $m + n$  is integral, the corresponding Dehn filling of  $X$  results in  $S^1 \times S^1 \times [0, 1]$ , and  $k_n(m + n)$  can be viewed as the union of  $E(k'_n)$  and  $E(k'')$ .

Hence  $K_m(n+m) \cong k_n(m+n) = E(k'_n) \cup E(k'')$  for all integers  $n$ . It should be noted here that  $E(k'')$  is independent of  $n$ , but the topological type of  $E(k'_n)$  depends on  $n$ .

Now assume for a contradiction that  $K_m$  is not prime and express  $K_m = t_1 \# \dots \# t_p$ , where  $t_i$  is a prime knot for  $1 \leq i \leq p$ . Then  $E(K_m)$  is the union of the  $p$ -fold composing space  $Y = [\text{disk with } p\text{-holes}] \times S^1$  and  $p$  knot spaces  $E(t_1), \dots, E(t_p)$ , where a regular fiber in  $\partial Y \cap \partial E(K_m)$  is a meridian of  $K_m$ . Since the surgery slope  $n+m$  is integral, the corresponding Dehn filling of  $Y$  results in  $(p-1)$ -fold composing space  $Y' = [\text{disk with } (p-1)\text{-holes}] \times S^1$ . Hence  $K_m(n+m)$  is expressed as the union  $Y' \cup E(t_1) \cup \dots \cup E(t_p)$ . If necessary, decomposing each  $E(t_i)$  further by essential tori, we obtain a torus decomposition of  $K_m(n+m)$  in the sense of Jaco, Shalen and Johannson [14; 15]. Note that identifications of  $Y'$  and  $E(t_i)$  ( $1 \leq i \leq p$ ) depends on  $n$ , but the topological type of  $E(t_i)$  ( $1 \leq i \leq p$ ) does not depend on  $n$ . To be precise, let us focus on the case of  $n = 0, 1$ . Then  $K_m(n+m) \cong k_n(m+n) = E(k'_n) \cup E(k'')$ , and  $E(k'_n)$  admits a hyperbolic structure in its interior:  $E(k'_0)$  is the exterior of the hyperbolic knot  $P(-5, 3, -3)$  and  $E(k'_1)$  is the exterior of the hyperbolic knot  $9_{42}$ . If  $E(k'')$  is neither hyperbolic nor Seifert fibered, we decompose  $E(k'')$  by essential tori to obtain a torus decomposition of  $K_m(n+m) \cong k_n(m+n)$  in the sense of Jaco, Shalen and Johannson. Since  $E(k'_0) \not\cong E(k'_1)$ , uniqueness of the torus decomposition of  $K_m(n+m)$  shows that some  $E(t_i)$  changes according as  $n = 0, 1$ . This is a contradiction. It follows that  $K_m$  is a prime knot.

Since the knot  $K_m$  is prime, while  $k_n$  is not prime for all integers  $m$  and  $n$ , we have  $\{K_m\} \cap \{k_n\} = \emptyset$ . Thus every integral slope fails to be a characterizing slope for a prime satellite knot  $K_m$  (with a given knot  $k''$  a companion knot) for any integer  $m$ , establishing Theorem 1.6(1). Similarly, every integral slope fails to be a characterizing slope for a composite knot  $k_n$  (with a given knot  $k''$  a connected summand) for any integer  $n$ . This establishes Theorem 1.6(2).

**Example 4.6** (Proof of Theorem 1.5) Figure 7 shows a sequence of transformations relating  $(m+n, \infty)$ -surgery on a link  $k_n \cup c$  to  $(\infty, n+m)$ -surgery on a link  $C \cup K_m$ . In particular, it gives two twist families of knots  $\{k_n\}$  and  $\{K_m\}$  such that  $k_n(m+n) = K_m(n+m)$ .

Replace  $(m+n, \infty)$ -surgery on  $k_n \cup c$  by  $(0, 0)$ -surgery on  $k_0 \cup c$ , and follow isotopies and light bulb moves as indicated in Figure 7 to see that  $(0, 0)$ -surgery on  $k_0 \cup c$  yields  $S^3$  with surgery dual  $C \cup K_0$ , where  $K_0$  is dual to  $c$  and  $C$  is dual to  $k_0$ .




Figure 7: Two families of knots  $\{k_n\}$  and  $\{K_m\}$  such that  $k_n(m+n) = K_m(n+m)$




Figure 8: The knot  $k_1$  of the family depicted in Figure 7 is isotoped into the 8-crossing Montesinos knot  $M(3, \frac{1}{3}, \frac{1}{2})$  which is the two-bridge knot  $\frac{23}{10}$  and also the knot  $8_6$  in Rolfsen's table.

As Figure 8 demonstrates, the knot  $k_1$  is the hyperbolic 8-crossing Montesinos knot  $M(3, \frac{1}{3}, \frac{1}{2})$ . It is the knot  $8_6$  in Rolfsen’s table, the two-bridge knot  $\frac{23}{10}$ . Following [27], (compare [25; 12]) it is a hyperbolic knot.

Using  $n = 0$ , we may calculate that

$$(\star\star) \Delta_{k_0 \cup c}(x, y) = (x^{-1} - 2 + x)y^{-1} + (x^{-2} - 4x^{-1} + 5 - 4x + x^2) + (x^{-1} - 2 + x)y,$$

which is equal to  $\Delta_{K_0 \cup c}(x, y^{-1})$  by Proposition 3.1. Note also that  $\Delta_{k_0 \cup c}(x, y) = \Delta_{k_0 \cup c}(x, y^{-1})$ ; see  $(\star\star)$ . Hence  $\Delta_{k_0 \cup c}(t, t^n) = \Delta_{k_0 \cup c}(t, t^{-n}) = \Delta_{K_0 \cup c}(t, t^n)$ , and it follows from Proposition 3.2 that

$$\Delta_{k_n}(t) = \Delta_{K_n}(t) = (t^{-1} - 2 + t)t^{-n} + (t^{-2} - 4t^{-1} + 5 - 4t + t^2) + (t^{-1} - 2 + t)t^n.$$

Thus Alexander polynomials distinguish  $k_1$  from  $K_m$  for all integers  $m \neq \pm 1$ .

We further calculate the Jones polynomials of  $k_1, K_1$ , and  $K_{-1}$  to be

$$\begin{aligned} V_{k_1}(q) &= \frac{1}{q^7} - \frac{2}{q^6} + \frac{3}{q^5} - \frac{4}{q^4} + \frac{4}{q^3} - \frac{4}{q^2} + \frac{3}{q} - 1 + q, \\ V_{K_1}(q) &= -\frac{1}{q^{15}} + \frac{1}{q^{14}} + \frac{1}{q^{11}} - \frac{1}{q^8} + \frac{1}{q^7} - \frac{3}{q^6} + \frac{3}{q^5} - \frac{4}{q^4} + \frac{5}{q^3} - \frac{4}{q^2} + \frac{3}{q}, \\ V_{K_{-1}}(q) &= -\frac{1}{q^{21}} + \frac{1}{q^{20}} + \frac{1}{q^{17}} - \frac{1}{q^{14}} + \frac{1}{q^{13}} - \frac{2}{q^{12}} + \frac{1}{q^{11}} - \frac{1}{q^{10}} \\ &\quad + \frac{1}{q^9} - \frac{1}{q^8} + \frac{1}{q^7} - \frac{1}{q^6} + \frac{2}{q^5} - \frac{3}{q^4} + \frac{4}{q^3} - \frac{3}{q^2} + \frac{2}{q}, \end{aligned}$$

to conclude that  $k_1 \neq K_{\pm 1}$ . Thus  $k_1$  is an 8-crossing hyperbolic knot for which every integral slope is not a characterizing slope.

### 5 Further discussions

Let  $K \cup c$  be a two-component link such that  $c$  is unknotted and not a meridian of  $K$ . If  $(0, 0)$ -surgery on  $K \cup c$  yields  $S^3$ , then Theorem 1.3 assures that  $K$  has infinitely many noncharacterizing slopes. Moreover, by Theorem 2.5, each knot  $K_n$  (obtained from  $K$  by an  $n$ -twist along  $c$ ) has also infinitely many noncharacterizing slopes.

**Proposition 5.1** *The  $(0, 0)$ -surgery on  $K_n \cup c$  results in  $S^3$ .*

**Proof** Note that  $(K_n \cup c)(0, 0) \cong (K \cup c)(-n, 0)$ . Since  $(K \cup c)(0, 0) \cong S^3$ , viewing  $K \subset c(0) = S^1 \times S^2$ , it is isotopic to an  $S^1$ -fiber in  $c(0)$ ; see the proof of Lemma 2.3 and Figure 2 for an illustration. Hence, we see that  $(K \cup c)(-n, 0)$  is also  $S^3$ . This then implies that  $(0, 0)$ -surgery on  $K_n \cup c$  results in  $S^3$  as well. □

Toward characterization of knots with infinitely many noncharacterizing slopes, we would like to ask:

**Question 5.2** Assume that  $K$  is a knot with infinitely many noncharacterizing slopes. Then can we take an unknot  $c$  so that  $c$  is not a meridian of  $K$  and  $(0, 0)$ -surgery on  $K \cup c$  yields  $S^3$ ?

**Question 5.3** For which knot  $K$  can we take an unknot  $c$  so that the link  $K \cup c$  enjoys the following properties:

- $c$  is not a meridian of  $K$ , and
- the result of the  $(0, 0)$ -surgery on  $K \cup c$  is  $S^3$ ?

Following Lemma 2.4, if  $c$  is a meridian of  $K$ , then  $(0, 0)$ -surgery on  $K \cup c$  always results in  $S^3$  independent of the knot  $K$ . However, if  $c$  is not a meridian of  $K$ , the second condition in Question 5.3 imposes a strong restriction on  $K$ . It follows from Ni and Zhang [26] and McCoy [23] that if  $K$  is a torus knot  $T_{r,s}$  with  $r > s > 1$  (resp.  $-r > s > 1$ ), then sufficiently positive (resp. negative) slopes are characterizing slopes. Hence any nontrivial torus knot does not admit an unknot  $c$  described in Question 5.3.

A knot  $K$  is an  $L$ -space knot if for some nonzero slope  $p/q \in \mathbb{Q}$  the manifold  $K(p/q)$  is an  $L$ -space, a rational homology 3-sphere for which  $\text{rk } \widehat{\text{HF}}(K(p/q)) = |H_1(K(p/q))|$ ; see [29]. If  $p/q > 0$ , then  $K$  is called a *positive*  $L$ -space knot, and if  $p/q < 0$ , then it is called a *negative*  $L$ -space knot. Motivated by the fact that torus knots are fundamental examples of  $L$ -space knots, we can prove:

**Theorem 1.8** Let  $K \cup c$  be a link with  $c$  a trivial knot. If  $c$  is not a meridian of  $K$  and the result of the  $(0, 0)$ -surgery on  $K \cup c$  is  $S^3$ , then  $K$  is not an  $L$ -space knot.

**Proof** Since  $(0, 0)$ -surgery on  $K \cup c$  results in  $S^3$ , the linking number between  $K$  and  $c$  must be  $\pm 1$ . Now suppose for a contradiction that  $K = K_0$  is an  $L$ -space knot. Then  $K_0(m)$  is an  $L$ -space for infinitely many integers  $m$  [29, Proposition 2.1]; more precisely, if  $K_0$  is a positive (resp. negative)  $L$ -space knot, then  $K_0(m)$  is an  $L$ -space for  $m \geq 2g(K_0) - 1$  (resp.  $m \leq -2g(K_0) + 1$ ); see [31]. By Theorem 2.5,  $K_0(m) = k_m(m)$  for all integers  $m$ ; hence the twist family  $\{(k_m, m)\}$  contains infinitely many  $L$ -space surgeries. Recall that the linking number between  $k$  and  $C$  is also  $\pm 1$ ; see the proof of Theorem 2.5. Furthermore, it follows from [1, Proposition 1.10] that

the  $k_m$  have the same Alexander polynomial for all  $m \in \mathbb{Z}$ , and  $g(k_m)$  is constant for infinitely many integers  $m$ . On the other hand, since  $|\text{lk}(K, C)| = 1$ , a recent work of Baker and Taylor [2] shows that  $g(k_m) \rightarrow \infty$  when  $|m| \rightarrow \infty$ , a contradiction.  $\square$

So we may expect a positive answer to the following:

**Question 5.4** *Does an L-space knot have only finitely many noncharacterizing slopes?*

Although the construction given by Theorem 2.1 and Theorem 2.5 provides infinitely many knots with infinitely many noncharacterizing slopes, we still expect that these knots have characterizing slopes as well.

**Question 5.5** *Does every knot  $K$  have a characterizing slope? More strongly, does every knot have infinitely many characterizing slopes?*

Our technique does not work directly for nonintegral slopes. So we would like to propose a modified version of Ni and Zhang's question:

**Question 5.6** *For a hyperbolic knot  $K$ , is a **nonintegral** slope  $p/q$  with  $|p| + |q|$  sufficiently large a characterizing slope?*

**Remark 5.7** Lackenby [20] shows that for each atoroidal, homotopically trivial knot  $K$  in a 3-manifold  $Y$  with  $H_1(Y; \mathbb{Q}) \neq \{0\}$ , there exists a number  $C(Y, K)$  such that  $p/q$  is a characterizing slope for  $K$  if  $|q| > C(Y, K)$ .

Ni and Zhang ask if every rational number is a noncharacterizing slope for some knot [26, Question 1.5]. We ask the opposite:

**Question 5.8** *Is there a rational number  $r$  which is a characterizing slope for all knots?*

More strongly and specifically, we would like to ask:

**Question 5.9** *Let  $r$  be a rational number which cannot be written in the form  $m + \frac{1}{n}$  for any integers  $m$  and  $n$ . Then is  $r$  a characterizing slope for all knots?*

It should be noted here that Kawachi [16] demonstrates that if  $r$  is written as  $m + \frac{1}{n}$  for some nonzero integers  $m$  and  $n$ , then it is not a characterizing slope for some hyperbolic knot. More precisely, he demonstrates that for any integer  $N > 1$  and any such an  $r$ , there are hyperbolic knots  $K_1, \dots, K_N$  whose  $r$ -surgery result in the same oriented 3-manifold.

## References

- [1] **KL Baker, K Motegi**, *Twist families of L-space knots, their genera, and Seifert surgeries*, preprint (2015) [arXiv](#) To appear in Comm. Anal. Geom.
- [2] **KL Baker, SA Taylor**, *Dehn filling and the Thurston norm*, preprint (2016) [arXiv](#) To appear in J. Differential Geom.
- [3] **D Bar-Natan**, *KnotTheory* Mathematica package <http://tinyurl.com/MKnTh>
- [4] **R Benedetti, C Petronio**, *Lectures on hyperbolic geometry*, Springer (1992) [MR](#)
- [5] **M Boileau, J Porti**, *Geometrization of 3-orbifolds of cyclic type*, Astérisque 272, Soc. Math. France, Paris (2001) [MR](#)
- [6] **M Culler, NM Dunfield, M Goerner, JR Weeks**, *SnapPy, a computer program for studying the geometry and topology of 3-manifolds* <http://snappy.computop.org>
- [7] **A Deruelle, K Miyazaki, K Motegi**, *Networking Seifert surgeries on knots*, Mem. Amer. Math. Soc. 1021, Amer. Math. Soc., Providence, RI (2012) [MR](#)
- [8] **D Gabai**, *Foliations and the topology of 3-manifolds, III*, J. Differential Geom. 26 (1987) 479–536 [MR](#)
- [9] **RE Gompf, K Miyazaki**, *Some well-disguised ribbon knots*, Topology Appl. 64 (1995) 117–131 [MR](#)
- [10] **CM Gordon**, *Some aspects of classical knot theory*, from “Knot theory” (JC Hausmann, editor), Lecture Notes in Math. 685, Springer (1978) 1–60 [MR](#)
- [11] **CM Gordon, J Luecke**, *Knots are determined by their complements*, J. Amer. Math. Soc. 2 (1989) 371–415 [MR](#)
- [12] **A Hatcher, W Thurston**, *Incompressible surfaces in 2-bridge knot complements*, Invent. Math. 79 (1985) 225–246 [MR](#)
- [13] **M Hirasawa, K Murasugi**, *Genera and fibredness of Montesinos knots*, Pacific J. Math. 225 (2006) 53–83 [MR](#)
- [14] **WH Jaco, PB Shalen**, *Seifert fibered spaces in 3-manifolds*, Mem. Amer. Math. Soc. 220, Amer. Math. Soc., Providence, RI (1979) [MR](#)
- [15] **K Johannson**, *Homotopy equivalences of 3-manifolds with boundaries*, Lecture Notes in Mathematics 761, Springer (1979) [MR](#)
- [16] **A Kawachi**, *Mutative hyperbolic homology 3-spheres with the same Floer homology*, Geom. Dedicata 61 (1996) 205–217 [MR](#)
- [17] **K Kodama**, *KNOT* Sather program <http://www.artsci.kyushu-u.ac.jp/~sumi/C/knot/>
- [18] **M Kouno, K Motegi, T Shibuya**, *Twisting and knot types*, J. Math. Soc. Japan 44 (1992) 199–216 [MR](#)
- [19] **P Kronheimer, T Mrowka, P Ozsváth, Z Szabó**, *Monopoles and lens space surgeries*, Ann. of Math. 165 (2007) 457–546 [MR](#)

- [20] **M Lackenby**, *Dehn surgery on knots in 3-manifolds*, J. Amer. Math. Soc. 10 (1997) 835–864 [MR](#)
- [21] **W B R Lickorish**, *Surgery on knots*, Proc. Amer. Math. Soc. 60 (1976) 296–298 [MR](#)
- [22] **T Mattman, K Miyazaki, K Motegi**, *Seifert-fibered surgeries which do not arise from primitive/Seifert-fibered constructions*, Trans. Amer. Math. Soc. 358 (2006) 4045–4055 [MR](#)
- [23] **D McCoy**, *Surgeries, sharp 4-manifolds and the Alexander polynomial*, preprint (2014) [arXiv](#)
- [24] **D McCoy**, *Non-integer characterizing slopes for torus knots*, preprint (2016) [arXiv](#)
- [25] **W Menasco**, *Closed incompressible surfaces in alternating knot and link complements*, Topology 23 (1984) 37–44 [MR](#)
- [26] **Y Ni, X Zhang**, *Characterizing slopes for torus knots*, Algebr. Geom. Topol. 14 (2014) 1249–1274 [MR](#)
- [27] **U Oertel**, *Closed incompressible surfaces in complements of star links*, Pacific J. Math. 111 (1984) 209–230 [MR](#)
- [28] **P Ozsváth, Z Szabó**, *Holomorphic disks and genus bounds*, Geom. Topol. 8 (2004) 311–334 [MR](#)
- [29] **P Ozsváth, Z Szabó**, *On knot Floer homology and lens space surgeries*, Topology 44 (2005) 1281–1300 [MR](#)
- [30] **P Ozsváth, Z Szabó**, *The Dehn surgery characterization of the trefoil and the figure eight knot*, preprint (2006) [arXiv](#)
- [31] **PS Ozsváth, Z Szabó**, *Knot Floer homology and rational surgeries*, Algebr. Geom. Topol. 11 (2011) 1–68 [MR](#)
- [32] **C Petronio, J Porti**, *Negatively oriented ideal triangulations and a proof of Thurston’s hyperbolic Dehn filling theorem*, Expo. Math. 18 (2000) 1–35 [MR](#)
- [33] **D Rolfsen**, *Knots and links*, Mathematics Lecture Series 7, Publish or Perish, Houston (1990) [MR](#)
- [34] **WP Thurston**, *The geometry and topology of three-manifolds*, lecture notes, Princeton University (1979) <http://msri.org/publications/books/gt3m>
- [35] **WP Thurston**, *Three-dimensional manifolds, Kleinian groups and hyperbolic geometry*, Bull. Amer. Math. Soc. 6 (1982) 357–381 [MR](#)

Department of Mathematics, University of Miami  
Coral Gables, FL, United States

Department of Mathematics, Nihon University  
Tokyo, Japan

[k.baker@math.miami.edu](mailto:k.baker@math.miami.edu), [motegi@math.chs.nihon-u.ac.jp](mailto:motegi@math.chs.nihon-u.ac.jp)

Received: 21 November 2016      Revised: 15 May 2017