

# Algebraic and topological properties of big mapping class groups

PRIYAM PATEL  
NICHOLAS G VLAMIS

Let  $S$  be an orientable, connected topological surface of infinite type (that is, with infinitely generated fundamental group). The main theorem states that if the genus of  $S$  is finite and at least 4, then the isomorphism type of the pure mapping class group associated to  $S$ , denoted by  $\text{PMap}(S)$ , detects the homeomorphism type of  $S$ . As a corollary, every automorphism of  $\text{PMap}(S)$  is induced by a homeomorphism, which extends a theorem of Ivanov from the finite-type setting. In the process of proving these results, we show that  $\text{PMap}(S)$  is residually finite if and only if  $S$  has finite genus, demonstrating that the algebraic structure of  $\text{PMap}(S)$  can distinguish finite- and infinite-genus surfaces. As an independent result, we also show that  $\text{Map}(S)$  fails to be residually finite for any infinite-type surface  $S$ . In addition, we give a topological generating set for  $\text{PMap}(S)$  equipped with the compact-open topology. In particular, if  $S$  has at most one end accumulated by genus, then  $\text{PMap}(S)$  is topologically generated by Dehn twists, otherwise it is topologically generated by Dehn twists along with handle shifts.

20E26, 37E30, 57M07, 57S05

## 1 Introduction

A surface is of finite type if its fundamental group is finitely generated; otherwise, it is of infinite type. Throughout, all surfaces are assumed to be connected, orientable, and to have compact (possibly empty) boundary.

The *mapping class group*, denoted by  $\text{Map}(S)$ , of a surface  $S$  is the group of orientation-preserving homeomorphisms of  $S$  up to isotopy, where we require all homeomorphisms as well as every stage of an isotopy to fix  $\partial S$  pointwise. The algebraic and geometric structure of mapping class groups of finite-type surfaces is generally well understood. In contrast, very little is known about *big* mapping class groups, ie mapping class groups of infinite-type surfaces.

Big mapping class groups arise naturally from the study of group actions on surfaces (eg Calegari [12]), taut foliations of 3-manifolds (eg Cantwell and Conlon [14]), and the Artinization of groups (eg Funar and Kapoudjian [18; 19]). (See Calegari [13] for a detailed discussion of these connections.) There has been a recent trend aimed at understanding infinite-type surfaces and their mapping class groups. For instance, recent work has investigated big mapping class group actions on hyperbolic graphs (eg Aramayona, Fossas and Parlier [2], Bavard [6] and Durham, Fanoni and Vlamis [16]).

This article focuses on the algebraic structure of big mapping class groups. Our methods use the language of topological groups and initiates the study of big mapping class groups in this category. The motivation for our work is a type of algebraic rigidity question for mapping class groups:

**Question 1.1** Does the isomorphism type of  $\text{Map}(S)$  determine the topology of  $S$ ? In particular, if  $\text{Map}(S)$  is isomorphic to  $\text{Map}(S')$ , then are  $S$  and  $S'$  homeomorphic?

**Remark** Since the original submission of this article, Question 1.1 has been answered in the affirmative by Bavard, Dowdall and Rafi [7]. Their techniques are different than those used in this article in the proof of Theorem 1. In particular, we use a topological-group-theoretic characterization of Dehn twists to prove Theorem 1, and Bavard, Dowdall and Rafi use an algebraic characterization of Dehn twists to obtain their result.

With the exception of finitely many sporadic cases in low complexity, we can see that Question 1.1 has a positive answer in the finite-type setting. To do so, observe that the rank of the center of  $\text{Map}(S)$ , the virtual cohomological dimension of  $\text{Map}(S)$  (see Harer [21]), and the rank of a maximal abelian subgroup in  $\text{Map}(S)$  (see Birman, Lubotzky and McCarthy [11]) together determine the genus, number of punctures, and the number of boundary components of  $S$  and thus its topological type.

Observe that  $\text{Map}(S)$  is countable if and only if  $S$  is of finite type. Therefore, the isomorphism type of  $\text{Map}(S)$  can distinguish between finite- and infinite-type surfaces. This reduces the question to considering specifically infinite-type surfaces, where little is known. The invariants mentioned for the finite-type setting, with the exception of the rank of the center, are all infinite for big mapping class groups.

In progress towards this question, we change our focus to a natural subgroup of  $\text{Map}(S)$ . In particular, we will work with the *pure mapping class group*, denoted by  $\text{PMap}(S)$ ,

consisting of mapping classes acting trivially on the topological ends of  $S$  (see Section 2.2 for the definition of ends). In this setting we address the same question and provide a partial answer:

**Theorem 1** *Let  $S$  be a surface whose genus is finite and at least 4. If  $S'$  is a surface, then any isomorphism between  $\text{PMap}(S)$  and  $\text{PMap}(S')$  is induced by a homeomorphism.*

If  $S$  has empty boundary, then the *extended* mapping class group, denoted by  $\text{Map}^\pm(S)$ , is the degree-2 extension of  $\text{Map}(S)$  that includes orientation-reversing mapping classes; otherwise, we set  $\text{Map}^\pm(S) = \text{Map}(S)$ . Setting  $S' = S$  in Theorem 1 yields a generalization of a celebrated result of Ivanov [25] for finite-type mapping class groups:

**Corollary 2** *If  $S$  is a borderless surface whose genus is finite and at least 4, then the natural monomorphism from  $\text{Map}^\pm(S)$  to  $\text{Aut}(\text{PMap}(S))$  is an isomorphism.*

An essential aspect of the proof of Theorem 1, just as in Ivanov's work, is to give a characterization of Dehn twists (Proposition 3.7), which allows us to conclude that an isomorphism of pure mapping class groups must send Dehn twists to Dehn twists. To obtain this characterization, we introduce a nonstandard topology  $\tau_w$  on  $\text{PMap}(S)$ , described in detail in Section 3. In Corollary 5.4, we discuss the structure of  $\text{Out}(\text{PMap}(S))$ ; in particular, it is isomorphic to the automorphism group of a countable boolean algebra.

Because of Theorem 1, it is natural for one to expect that  $\text{PMap}(S)$  is a characteristic subgroup of  $\text{Map}(S)$ . If this is the case, then Corollary 2 can be extended to  $\text{Map}(S)$  to again match the finite-type setting. We state these as conjectures with the same assumptions as in Theorem 1; however, we expect these conjectures and Corollary 2 to hold in general.

**Conjecture 1.2** *Let  $S$  be a surface whose genus is finite and at least 4.*

- (1)  *$\text{PMap}(S)$  is a characteristic subgroup of  $\text{Map}(S)$  and  $\text{Map}^\pm(S)$ .*
- (2) *If  $S$  is borderless, then  $\text{Aut}(\text{Map}^\pm(S)) = \text{Aut}(\text{Map}(S)) = \text{Map}^\pm(S)$ .*

(This conjecture has been subsequently proven — in greater generality — by Bavard, Dowdall and Rafi [7].)

The first step in proving Theorem 1 is to distinguish finite- and infinite-genus surfaces via the algebraic structure of their associated pure mapping class groups. In order to do this, we investigate whether big mapping class groups are residually finite. As these results are of independent interest, we include Theorem 3 as a summary of the results in Section 4. Recall that a group is *residually finite* if the intersection of its proper finite-index normal subgroups is trivial.

**Theorem 3** *Let  $S$  be any surface.*

- (1)  $\text{PMap}(S)$  is residually finite if and only if  $S$  has finite genus.
- (2)  $\text{Map}(S)$  is residually finite if and only if  $S$  is of finite type.

Note that the finite-type cases are handled by Grossman [20]. Using the structure of  $\tau_w$ , we show that  $\text{PMap}(S)$  is residually finite whenever  $S$  has finite genus (Proposition 4.1). We then introduce the infinite-stranded braid group  $B_\infty$ , show that it fails to be residually finite (Corollary 4.3), and prove that it embeds in every finite-genus big mapping class group (Proposition 4.4). Thus,  $\text{Map}(S)$  is not residually finite when  $S$  is of infinite type and has finite genus. Proposition 4.6 shows that  $\text{PMap}_c(S)$ , the subgroup of  $\text{Map}(S)$  consisting of mapping classes with compact support, has no finite quotients when  $S$  has infinite genus. This fact is almost immediate from a theorem of Paris [31] and implies that when  $S$  has infinite genus,  $\text{Map}(S)$  and  $\text{PMap}(S)$  are not residually finite.

The reason  $\text{PMap}(S)$ , when  $S$  is a finite-genus surface, behaves differently than other big mapping class groups is the existence of forgetful homomorphisms to finite-type pure mapping class groups. We use these forgetful homomorphisms to build the topology  $\tau_w$  on  $\text{PMap}(S)$  mentioned above (see Section 3). Understanding basic topological properties of  $\tau_w$  is the key to characterizing Dehn twists and understanding the residual properties of  $\text{PMap}(S)$ . The construction of  $\tau_w$  naturally leads to an inverse limit of finite-type pure mapping class groups, which is the completion (as uniform spaces) of the associated big pure mapping class group. This inverse limit viewpoint is thoroughly discussed in Section 7.

As Dehn twists and the topology of  $\tau_w$  play a critical role in our understanding of  $\text{PMap}(S)$ , it is natural to understand which of these properties hold in the more standard compact-open topology. Equipping the group of homeomorphisms of a surface with the compact-open topology, we give  $\text{Map}(S)$  the corresponding quotient topology. We let  $\tau_q$  denote the restriction of this topology to  $\text{PMap}(S)$ . Note that if  $S$  is of finite

type, then  $\tau_q$  is the discrete topology. From the definition of  $\tau_w$  and the Dehn–Lickorish theorem, it follows that Dehn twists topologically generate  $(\text{PMap}(S), \tau_w)$ .

In the following theorem, we see that Dehn twists topologically generate  $(\text{PMap}(S), \tau_q)$  if  $S$  has finite genus; however, this can fail in the infinite-genus setting. Intuitively, a topological end of a surface is *accumulated by genus* if there is a sequence of handles converging to the end (see Section 2.2 for the precise definition). In order to build a topological generating set in the infinite-genus setting we introduce the notion of a *handle shift* in Section 6. Roughly speaking, a handle shift is a homeomorphism which applies a  $+1$  shift to a  $\mathbb{Z}$ -indexed collection of handles in the surface. This notion allows us to build a topological generating set for  $(\text{PMap}(S), \tau_q)$ :

**Theorem 4** *The set of Dehn twists topologically generate  $(\text{PMap}(S), \tau_q)$  if and only if  $S$  has at most one end accumulated by genus. If  $S$  has at least two ends accumulated by genus, then the set of Dehn twists together with the set of handle shifts topologically generate  $(\text{PMap}(S), \tau_q)$ .*

## 1.1 Outline

In Section 2 we give the necessary background, which focuses on the structure of the space of ends of an infinite-type surface.

In Section 3, we define the topology  $\tau_w$ , show it is Hausdorff (Lemma 3.4), and investigate the closure of the set of Dehn twists (Lemma 3.6), allowing us to characterize Dehn twists (Proposition 3.7). Section 4 is dedicated to investigating residual finiteness.

Given the results of Section 4, Section 5 focuses on finite-genus surfaces. In this section, we combine the characterization of Dehn twists from Section 3 and results regarding homomorphisms between finite-type mapping class groups to conclude the proof of Theorem 1 in Propositions 5.1 and 5.3.

In Section 6, we introduce the notion of a handle shift and prove Theorem 4. In Section 7, we discuss inverse limits of finite-type pure mapping class groups and finish the paper by showing the proper containment  $\tau_w \subset \tau_q$  in Section 7.1.

## Acknowledgements

The authors would like to thank Javier Aramayona, Ara Basmajian, Daryl Cooper, David Fernández-Bretón, Chris Leininger, Darren Long, and Scott Schneider for several helpful conversations. The authors also thank the referee for helpful comments.

Vlavis was supported in part by NSF RTG grant 1045119.

## 2 Background

### 2.1 Curves

A *simple closed curve* in a surface  $S$  is the image of a topological embedding  $\mathbb{S}^1 \hookrightarrow S$ . A simple closed curve is *trivial* if it is homotopic to a point; it is *peripheral* if it either is homotopic to a boundary component or bounds a once-punctured disk in  $S$ ; it is *essential* if it is neither trivial nor peripheral; it is *separating* if its complement is disconnected and *nonseparating* otherwise. If  $c$  is a simple closed curve in  $S$ , then we let  $[c]$  denote its isotopy class in  $S$ . Given two isotopy classes of simple closed curves  $[c]$  and  $[d]$ , their *geometric intersection number*, denoted by  $i([c], [d])$ , is defined to be

$$i([c], [d]) = \min\{|c' \cap d'| : [c'] = [c] \text{ and } [d'] = [d]\}.$$

When is it clear from context, we will conflate a simple closed curve with its isotopy class.

Recent work of Hernández, Morales and Valdez [22] assures us that, just as in the finite-type setting, simple closed curves play an essential role in the study of big mapping class groups. In particular, they extend the Alexander method to infinite-type surfaces [22, Theorem 1.1], yielding the following lemma:

**Lemma 2.1** [22, Corollary 1.2] *Suppose  $S$  is an infinite-type surface without boundary. If an element of  $\text{Map}^\pm(S)$  fixes the isotopy class of every simple closed curve, then it is trivial.*

We will need one final notion: if  $F \subset S$  is a subsurface, then we call  $F$  *essential* if it is connected and the inclusion  $F \hookrightarrow S$  induces a monomorphism  $\text{PMap}(F) \hookrightarrow \text{PMap}(S)$ , where a homeomorphism of  $F$  is extended to  $S$  by the identity.

### 2.2 Ends

In order to work with infinite-type surfaces, it is essential to understand their space of ends, which we now define.

**Definition 2.2** Let  $S$  be an infinite-type surface and fix an exhaustion  $\{K_i\}_{i \in \mathbb{N}}$  of  $S$  by compact subsurfaces, that is, each  $K_i$  is a compact surface of  $S$ ,  $K_i \subset K_{i+1}$ , and  $\bigcup_{i \in \mathbb{N}} K_i = S$ .

- An *end* of  $S$  is a sequence  $\{U_i\}_{i \in \mathbb{N}}$  such that each  $U_i$  is a connected component of  $S \setminus K_i$  and  $U_{i+1} \subset U_i$  for all  $i \in \mathbb{N}$ .

- Define  $U_i^*$  to be the set of all ends that contain  $U_i$ , where  $U_i$  is a connected component of  $S \setminus K_i$ . The *space of ends* of  $S$ , denoted by  $\text{Ends}(S)$ , consists — as a set — of the ends of  $S$ . It is equipped with the topology generated by sets of the form  $U_i^*$ .

Note that the construction of  $\text{Ends}(S)$  is independent of the choice of exhaustion.

The space  $\text{Ends}(S)$  is Hausdorff, totally disconnected, second countable, and compact (see [1, Chapter 1, Sections 36–37]); hence, it can be realized as a closed subset of the Cantor set. An end of  $S$  comes in two types:

- An end  $\{U_i\}$  is *accumulated by genus* if  $U_i$  has infinite genus for all  $i$ .
- An end  $\{U_i\}$  is *planar* if  $U_i$  is planar for all but finitely many  $i$ .

In addition, an end is *isolated* if it is an isolated point in the topology given above. We will use the word *puncture* to denote an isolated planar end.

Kerékjártó [36] showed that the homeomorphism type of an orientable infinite-type surface is determined by the quadruple

$$(g, b, \text{Ends}_\infty(S), \text{Ends}(S)),$$

where  $g \in \mathbb{N} \cup \{0, \infty\}$  is the genus of  $S$ ,  $b \in \mathbb{N}$  is the number of boundary components, and  $\text{Ends}_\infty(S) \subset \text{Ends}(S)$  is the subset of ends accumulated by genus (see Richards [34] for a proof).

An immediate consequence of the classification of infinite-type surfaces is:

**Lemma 2.3** (definition of  $\bar{S}$ ) *When  $S$  has finite genus, there exists a unique — up to homeomorphism — compact surface, which we label  $\bar{S}$ , containing  $S$  such that  $\bar{S} \setminus S$  is homeomorphic to  $\text{Ends}(S)$ .*

The surface  $\bar{S}$  given by Lemma 2.3 is the Freudenthal compactification of  $S$ . Note that the Freudenthal compactification exists for any surface, but is not a manifold in general.

Given the definition of  $\text{Ends}(S)$  it is clear that a homeomorphism of  $S$  induces a homeomorphism of  $\text{Ends}(S)$ . As any isotopy must fix the ends of  $S$ , we see that there is an induced action of  $\text{Map}(S)$  on  $\text{Ends}(S)$ . The *pure mapping class group*  $\text{PMap}(S)$  is the kernel of this action.

**Proposition 2.4** [8, Proposition I.1] *Let  $\mathbb{D}$  denote the unit disk and let  $\Pi$  be a totally disconnected compact subset contained in the interior of  $\mathbb{D}$ . Given a homeomorphism  $f: \Pi \rightarrow \Pi$  there exists a homeomorphism  $\hat{f}: \mathbb{D} \rightarrow \mathbb{D}$  and a regular neighborhood  $N$  of  $\partial\mathbb{D}$  such that  $\hat{f}|_{\Pi} = f$  and  $\hat{f}|_N = \text{id}$ .*

Observe that since a regular neighborhood of the boundary of  $\mathbb{D}$  is fixed pointwise, it follows that  $\hat{f}$  is orientation-preserving. This yields:

**Corollary 2.5** *If  $S$  has finite genus, then the induced homomorphism  $\text{Map}(S) \rightarrow \text{Homeo}(\text{Ends}(S))$  is surjective.*

### 3 The initial topology

For the entirety of this section, we require  $S$  to have finite genus and be of infinite type. In this setting, we will introduce a topology on  $\text{PMap}(S)$  that is suited to our algebraic study of the group. Define the set  $\Lambda(S)$  to be the set of finite subsets of  $\text{Ends}(S)$ , that is,

$$\Lambda(S) = \{\lambda \subset \text{Ends}(S) : |\lambda| < \infty\}.$$

When it is clear from context, we will simply write  $\Lambda$ . For each  $\lambda \in \Lambda$ , we define the surface  $S_{\lambda} = \bar{S} \setminus \lambda$ . The inclusion  $i_{\lambda}: S \hookrightarrow S_{\lambda}$  induces a homomorphism  $\varphi_{\lambda}: \text{PMap}(S) \rightarrow \text{PMap}(S_{\lambda})$ . This homomorphism is the forgetful homomorphism, where one forgets all ends of  $S$  not in  $\lambda$ .

The *initial topology*, denoted by  $\tau_w$ , on  $\text{PMap}(S)$  is the initial topology with respect to the family of maps  $\{\varphi_{\lambda}\}_{\lambda \in \Lambda}$ , that is, it is the coarsest topology such that  $\varphi_{\lambda}$  is continuous for each  $\lambda \in \Lambda$ , where we equip  $\text{PMap}(S_{\lambda})$  with the discrete topology. If at this point the reader is inclined to think about inverse limits, we discuss this viewpoint in Section 7.

For the rest of the section our goal is to present two key lemmas (Lemmas 3.4 and 3.6) regarding the algebraic structure of  $\text{PMap}(S)$ . Before starting, we need to understand the center of  $\text{PMap}(S)$ , which we denote by  $Z(\text{PMap}(S))$ . Recall that the pure mapping class group of any finite-type surface is generated by Dehn twists (see [17, Section 4.4.4]). Combining this fact with the definition of the initial topology above, we have the following lemma as an immediate corollary:

**Lemma 3.1** *( $\text{PMap}(S), \tau_w$ ) is topologically generated by Dehn twists.*



**Lemma 3.2** *An element of  $\text{PMap}(S)$  is central if and only if it fixes the isotopy class of every simple closed curve.*

**Proof** Let  $f \in \text{PMap}(S)$ . Recall that if  $c$  is a simple closed curve in  $S$ , then  $f \cdot T_c \cdot f^{-1} = T_{f(c)}$ . Now assume that  $f$  fixes the isotopy class of every simple closed curve. It follows that  $f \cdot T_c \cdot f^{-1} = T_{f(c)} = T_c$  for any simple closed curve  $c$ . Therefore,  $f$  commutes with every Dehn twist in  $\text{PMap}(S)$  and hence, by Lemma 3.1,  $f$  commutes with every element of a topological generating set for  $\text{PMap}(S)$ . It is an easy exercise to show that multiplication in  $(\text{PMap}(S), \tau_w)$  is continuous. Therefore, we can conclude that  $f \in Z(\text{PMap}(S))$ .

Now assume that  $f$  is central. It follows that  $T_c = f \cdot T_c \cdot f^{-1} = T_{f(c)}$  and hence  $f([c]) = [c]$  for every simple closed curve  $c$  in  $S$ .  $\square$

**Lemma 3.3**  *$Z(\text{PMap}(S))$  is generated by the Dehn twists about the boundary components of  $S$ . In particular, if  $S$  is borderless then  $\text{PMap}(S)$  has trivial center.*

**Proof** Let  $G < \text{PMap}(S)$  be the subgroup generated by the Dehn twists about the boundary components of  $S$ .

Let  $R$  be the surface obtained from  $S$  by gluing a once-punctured disk to each component of  $\partial S$ . Let  $\iota: S \hookrightarrow R$  be the associated inclusion and  $\iota_*: \text{PMap}(S) \rightarrow \text{PMap}(R)$  the induced homomorphism. By construction,  $\iota_*$  is surjective. To see this, note that  $\iota_*$  is continuous and every Dehn twist in  $\text{PMap}(R)$  is an image of a Dehn twist in  $\text{PMap}(S)$ .

As  $\iota_*$  is surjective, it follows that if  $f \in Z(\text{PMap}(S))$ , then  $\iota_*(f) \in Z(\text{PMap}(R))$ . By Lemma 3.2,  $\iota_*(f)$  fixes the isotopy class of every simple closed curve; therefore, as  $R$  is borderless, we can apply Lemma 2.1 to see that  $\iota_*(f)$  is trivial.

A direct replacement of the finite-type Alexander method with the infinite-type Alexander method [22] in the proof of [17, Theorem 3.18] implies  $G = \ker \iota_*$ ; in particular,  $f \in G$ . Therefore,  $Z(\text{PMap}(S)) \subset G$ .

Now, let  $f \in G$ . Then  $f$  fixes the isotopy class of every simple closed curve. Lemma 3.2 allows us to conclude that  $f$  is central and  $G \subset Z(\text{PMap}(S))$ .  $\square$

**Lemma 3.4**  *$\tau_w$  is Hausdorff.*

**Proof** To show the Hausdorff property, it is enough to show that any point can be separated from the identity with open sets. By the definition of  $\tau_w$  and continuity of left multiplication, it is enough to show that

$$\bigcap_{\lambda \in \Lambda} \ker \varphi_\lambda = \{\text{id}\}.$$

Let  $f \in \text{PMap}(S)$  be a nontrivial element. If  $f$  is central in  $\text{PMap}(S)$ , then by Lemma 3.3 it is a product of Dehn twists about boundary components and hence the same is true for  $\varphi_\lambda(f)$ . In particular,  $f$  is not an element of  $\ker \varphi_\lambda$  whenever  $|\lambda| > 1$ . Assuming  $f$  is not central, by Lemma 3.2, there exists an essential simple closed curve  $a$  in  $S$  such that  $f(a)$  is not homotopic to  $a$ . Choose an essential finite-type surface  $F \subset S$  so that  $\text{genus}(F) = \text{genus}(S)$  and both  $a$  and  $f(a)$  are contained in  $F$ . Let  $\lambda \in \Lambda$  be such that the intersection of  $\lambda$  with each component of  $\bar{S} \setminus F$  is nonempty. It follows that  $i_\lambda|_F$  is an embedding of  $F$  into  $S_\lambda$  as an incompressible surface; in particular,  $i_\lambda(f(a))$  and  $i_\lambda(a)$  are not homotopic in  $S_\lambda$ . It follows that  $\varphi_\lambda(f)$  does not fix the isotopy class of  $i_\lambda(a)$  in  $S_\lambda$ ; hence,  $f \notin \ker \varphi_\lambda$ .  $\square$

The proof of Lemma 3.4 tells us that  $\text{PMap}(S)$  inherits the residual properties of finite-type pure mapping class groups. In particular, Lemma 3.4 combined with the work of Grossman [20] implies that  $\text{PMap}(S)$  is residually finite. This is discussed in more depth in Section 4. In addition, it follows from the inverse limit construction in Section 7 and Lemma 3.4 that  $(\text{PMap}(S), \tau_w)$  is a topological group (see Proposition 7.1).

The goal of the next key lemma (Lemma 3.6) is to detect Dehn twists in  $\text{PMap}(S)$ . In the case of finite-type surfaces, the main idea behind Ivanov's results in [25] on the automorphisms of mapping class groups is to algebraically characterize Dehn twists. Given an automorphism preserving Dehn twists, one can then build a map on the collection of simple closed curves in order to determine the isotopy class of a homeomorphism. Building off the literature surrounding homomorphisms between mapping class groups, we take a similar approach in Section 5.

Given a simple closed curve  $a$  in  $S$  let  $[a]$  denote its isotopy class in  $S$  and let  $[a]_\lambda$  denote the isotopy class of  $i_\lambda(a)$  in  $S_\lambda$ . For the sake of the argument below, we record an obvious lemma:

**Lemma 3.5** For any  $\lambda \in \Lambda$ ,

$$i([a], [b]) \geq i([a]_\lambda, [b]_\lambda),$$

where  $a$  and  $b$  are any two simple closed curves in  $S$ .




Figure 1: The sequence of Dehn twists  $\{T_{a_n}\}_{n=2}^\infty$  in  $\text{PMap}(S)$ , where  $S = \mathbb{C} \setminus \mathbb{N}$ , converges to the identity in  $\tau_w$ , but does not converge in  $\tau_q$  to any homeomorphism.

Let  $\mathcal{D}$  denote the collection of Dehn twists in  $\text{PMap}(S)$  and let  $\mathcal{D}_\lambda$  denote the collection of Dehn twists in  $\text{PMap}(S_\lambda)$ . The subgroup of  $\text{PMap}(S)$  generated by  $\mathcal{D}$  is the group of compactly supported mapping classes, denoted by  $\text{PMap}_c(S)$ . We will use this notation throughout the article.

**Lemma 3.6** *The closure of  $\mathcal{D}$  in  $\tau_w$  is  $\overline{\mathcal{D}} = \mathcal{D} \cup \{\text{id}\}$ .*

We note that this statement also holds in  $\tau_q$ ; we give a proof in Section 6 (see Proposition 6.4). Additionally, the reader should be warned that the subtlety in the proof of the lemma lies in the fact that the stabilizer of an isotopy class of a curve is not open in  $\tau_w$ . This failure for a stabilizer of a curve to be open follows from the proof of Proposition 7.6 and is demonstrated by the example illustrated in Figure 1, which shows an example of a (nonintuitive) sequence of Dehn twists converging to the identity.

**Proof** Fix an exhaustion  $K_1 \subset K_2 \subset \dots$  of  $S$  by essential finite-type genus( $S$ )-subsurfaces. Observe, for later in the proof, that the direct limit of the groups  $\{\text{PMap}(K_n)\}_{n \in \mathbb{N}}$  is  $\text{PMap}_c(S)$ .

We first see that the identity is in the closure of  $\mathcal{D}$ . Recall that the sets of the form  $\ker \varphi_\lambda$  for  $\lambda \in \Lambda$  form a neighborhood basis for the identity in  $(\text{PMap}(S), \tau_w)$ . Given  $\lambda \in \Lambda$  we can find a clopen subset  $U$  of  $\text{Ends}(S)$  disjoint from  $\lambda$  and a disk  $D \subset \overline{S}$  such that  $U \subset D$ . The Dehn twist about  $\partial D$  is then an element in  $\mathcal{D} \cap \ker \varphi_\lambda$ . It follows that  $\mathcal{D}$  intersects every neighborhood of the identity and hence the identity is in  $\overline{\mathcal{D}}$ .

As  $\tau_w$  is not first countable when  $\text{Ends}(S)$  is uncountable (see Lemma 7.4), it is necessary for the remainder of the proof for us to work with nets as opposed to sequences; however, the reader will lose no intuition by replacing the word net with sequence.

Let  $I$  be a directed set and let  $\{T_{c_i}\}_{i \in I}$  be a net in  $\mathcal{D}$ , where  $T_{c_i}$  denotes the Dehn twist about the simple closed curve  $c_i$ . Assume the net converges to  $f \in (\text{PMap}(S), \tau_w)$ . Further assuming that  $f$  is not the identity, we will then show that  $f$  is a Dehn twist.

We first claim that  $f$  is an element of  $\text{PMap}_c(S)$ . Suppose not. Then, by Lemma 3.3,  $f$  is not in the center of  $\text{PMap}(S)$ , so there exists an essential simple closed curve  $a$  with  $[f(a)] \neq [a]$ . Choose  $N \in \mathbb{N}$  such that both  $a$  and  $f(a)$  can be homotoped into  $K_N$ . For each  $n \in \mathbb{N}$  with  $n > N$  we can find  $b_n$  homotopic into the complement of  $K_n$  such that  $[f(b_n)] \neq [b_n]$ . Fix  $n \in \mathbb{N}$  with  $n > N$ . Then there exists  $m \in \mathbb{N}$  with  $m > n$  and where  $b_n$  and  $f(b_n)$  are homotopic into  $K_m$ . Choose  $\lambda \in \Lambda$  such that  $\lambda$  intersects each component of  $\bar{S} \setminus K_m$ .

Given the choice of  $\lambda$ , we have  $[f(a)]_\lambda \neq [a]_\lambda$  and  $[f(b_n)]_\lambda \neq [b_n]_\lambda$ . It follows that  $\varphi_\lambda(f)$  is nontrivial and thus a Dehn twist, call it  $T$ , as our net is in  $\mathcal{D}$ . By construction,

$$T([a]_\lambda) = \varphi_\lambda(f)([a]_\lambda) \neq [a]_\lambda \quad \text{and} \quad T([b_n]_\lambda) = \varphi_\lambda(f)([b_n]_\lambda) \neq [b_n]_\lambda.$$

Choose a simple closed curve  $c$  in  $K_m$  such that  $T = T_{[c]_\lambda}$ . Then

- $i([c]_\lambda, [a]_\lambda) > 0$ ,
- $i([c]_\lambda, [b_n]_\lambda) > 0$ , and
- $i([a]_\lambda, [b_n]_\lambda) = 0$ ,

where the first two inequalities follow from the fact that  $T$  fixes neither  $[a]_\lambda$  nor  $[b_n]_\lambda$ . The last equality comes from the fact that  $a$  and  $b_n$  are disjoint combined with Lemma 3.5. A direct application of [17, Proposition 3.4] gives the inequality

$$i(T([a]_\lambda), [b_n]_\lambda) = i([c]_\lambda, [a]_\lambda)i([c]_\lambda, [b_n]_\lambda) > 0.$$

Another application of Lemma 3.5 yields

$$i([f(a)], [b_n]) \geq i([f(a)]_\lambda, [b_n]_\lambda) = i(T([a]_\lambda), [b_n]_\lambda) > 0.$$

As  $n$  was arbitrary, we see that  $f(a)$  must exit every compact set of  $S$ ; hence, it cannot be compact and  $f$  is not a homeomorphism.

It follows there exists  $n \in \mathbb{N}$  such that  $f \in \text{PMap}(K_n) < \text{PMap}_c(S)$ . Let  $m \in \mathbb{N}$  such that  $m > n$  and  $K_n$  is contained in the interior of  $K_m$ . Choose  $\lambda \in \Lambda$  such that  $\lambda$  intersects each component of  $\bar{S} \setminus K_m$  nontrivially. We have that  $\varphi_\lambda$  restricted to  $\text{PMap}(K_n)$  is injective and induced by the inclusion  $i_\lambda|_{K_n}: K_n \hookrightarrow \bar{S} \setminus \lambda$ ; in particular, as  $f \in \text{PMap}(K_n)$  and  $\varphi_\lambda(f)$  is a Dehn twist, we must have that  $f$  is a Dehn twist.  $\square$

As a direct consequence, we give a characterization of Dehn twists. For  $\lambda \in \Lambda$ , let  $\overline{\mathcal{D}}_\lambda$  be the union of the Dehn twists in  $\text{PMap}(S_\lambda)$  with the identity, so  $\overline{\mathcal{D}}_\lambda = \varphi_\lambda(\overline{\mathcal{D}})$ .

**Proposition 3.7**  $f \in \overline{\mathcal{D}}$  if and only if  $\varphi_\lambda(f) \in \overline{\mathcal{D}}_\lambda$  for every  $\lambda \in \Lambda$ .

**Proof** The forward direction is trivial. For the reverse, assume  $f \in \text{PMap}(S)$  such that  $\varphi_\lambda(f) \in \overline{\mathcal{D}}_\lambda$  for every  $\lambda \in \Lambda$ . For each  $\lambda \in \Lambda$  choose  $T_\lambda \in \overline{\mathcal{D}}$  such that  $\varphi_\lambda(T_\lambda) = \varphi_\lambda(f)$ . This implies that the net  $\{T_\lambda\}_{\lambda \in \Lambda}$  converges to  $f$ ; hence,  $f \in \overline{\mathcal{D}}$  by Lemma 3.6. □

## 4 Residual finiteness

A group is *residually finite* if the intersection of all of its finite-index proper normal subgroups is trivial. As noted earlier, the work of Grossman [20] tells us that mapping class groups of finite-type surfaces are residually finite. As this property is inherited by subgroups it follows that pure mapping class groups are also residually finite.

In this section, we will prove Theorem 3 in a series of propositions. We break the section into two subsections handling the finite- and infinite-genus cases separately.

### 4.1 Finite genus

As a start, we properly record the result mentioned in Section 3:

**Proposition 4.1** *If  $S$  has finite genus, then  $\text{PMap}(S)$  is residually finite.*

**Proof** Let  $f \in \text{PMap}(S)$  be an arbitrary nontrivial element. Then by Lemma 3.4 there exists  $\lambda \in \Lambda$  such that  $f \notin \ker \varphi_\lambda$ . Further, since  $\text{PMap}(S_\lambda)$  is residually finite [20] there exists a homomorphism  $\psi: \text{PMap}(S_\lambda) \rightarrow G$ , where  $G$  is a finite group and  $f \notin \ker(\psi \circ \varphi_\lambda)$ . □

In a slight change of perspective, we will focus on the full mapping class group  $\text{Map}(S)$  for the remainder of this subsection.

Let  $B_n$  denote the  $n$ -stranded braid group, which is also the mapping class group of the  $n$ -punctured disk. The classical presentation for  $B_n$  is

$$B_n = \left\langle \sigma_1, \dots, \sigma_{n-1} \mid \begin{array}{ll} \sigma_i \sigma_j = \sigma_j \sigma_i & \text{for } |i - j| \geq 2 \text{ and } 1 \leq i, j \leq n - 1, \\ \sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1} & \text{for } 1 \leq i \leq n - 2 \end{array} \right\rangle.$$

With this presentation we see that there are natural inclusions  $\iota_{n,m}: B_n \hookrightarrow B_m$  whenever  $n \leq m$ . In particular  $\langle B_n, \iota_{n,m} \rangle$  is a directed system of groups and we define  $B_\infty$  to be the direct limit. The presentation for  $B_\infty$  is immediate from that of  $B_n$ , namely,

$$B_\infty = \left\langle \{\sigma_i : i \in \mathbb{N}\} \mid \begin{array}{ll} \sigma_i \sigma_j = \sigma_j \sigma_i & \text{for } |i - j| \geq 2 \text{ and } i, j \in \mathbb{N}, \\ \sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1} & \text{for } i \in \mathbb{N} \end{array} \right\rangle.$$

We will proceed by proving that  $B_\infty$  is not residually finite and that it is a subgroup of every finite-genus big mapping class group.

**Proposition 4.2** *Every finitely generated quotient of  $B_\infty$  is cyclic.*

**Proof** Let  $f: B_\infty \rightarrow G$  be a surjective homomorphism to a finitely generated group. We can then find some  $N$  such that  $f|_{B_N}$  is surjective. For  $m > N$ ,  $\sigma_m$  commutes with every element of  $B_N$ ; hence,  $f(\sigma_m)$  is in the center of  $G$  for every  $m > N$ . In particular,  $f(\sigma_N)$  commutes with  $f(\sigma_{N+1})$ . From the relations, it is easy to see that if  $f(\sigma_i)$  commutes with  $f(\sigma_{i+1})$ , then  $f(\sigma_i) = f(\sigma_{i+1})$ . It now follows that  $G$  is generated by  $f(\sigma_1)$ . □

The abelianization of  $B_\infty$  is isomorphic to  $\mathbb{Z}$ ; in particular, from Proposition 4.2, it follows that every finite quotient of  $B_\infty$  factors through its abelianization. Thus the commutator subgroup is in the intersection of every proper finite-index normal subgroup of  $B_\infty$ . This shows:

**Corollary 4.3**  *$B_\infty$  is not residually finite.*

**Proposition 4.4** *Let  $S$  be an infinite-type surface. If  $S$  has finite genus, then  $B_\infty$  is a subgroup of  $\text{Map}(S)$ .*

In fact, Proposition 4.4 holds whenever  $S$  has infinitely many planar ends, but we will not use the full generality of the result in what follows.

**Proof** The proof splits into two cases:  $\text{Ends}(S)$  has either infinitely or finitely many isolated points. Let us start by assuming that  $\text{Ends}(S)$  has infinitely many isolated points. As  $\text{Ends}(S)$  is separable, the set of isolated ends must be countably infinite, so we can choose a labeling  $\{p_1, p_2, \dots\}$  of the isolated ends. For each  $n \in \mathbb{N}$  let  $D_n \subset S$  be an  $n$ -punctured disk satisfying

- (1)  $\text{Ends}(D_n) = \{p_1, \dots, p_n\}$ , and
- (2)  $D_n \subset D_{n+1}$ .

It follows that  $D_n$  is an essential subsurface of  $S$ ; in particular, the embedding  $D_n \hookrightarrow S$  induces an injection  $\text{Map}(D_n) \hookrightarrow \text{Map}(S)$ . Recall that  $\text{Map}(D_n) \cong B_n$ . By the universal property of direct limits, the inclusions  $D_n \hookrightarrow D_{n+1}$  induce an injective homomorphism  $B_\infty \hookrightarrow \text{Map}(S)$  as desired.

Now assume  $\text{Ends}(S)$  has a finite number of isolated points  $\{p_1, \dots, p_n\}$ . It follows that  $C = \text{Ends}(S) \setminus \{p_1, \dots, p_n\}$  is perfect. Let  $D \subset \bar{S}$  be a disk such that  $D \cap F = \emptyset$  and  $D \cap C = C'$ , where  $C'$  is a clopen subset of  $C$ . It follows from Brouwer's topological characterization of the Cantor set (see [27, Theorem 7.4]) that  $C'$  is homeomorphic to the Cantor set. The work of Funar and Kapoudjian [18, Section 7] implies  $B_\infty < \text{Map}(D \setminus C')$ ; thus,  $B_\infty < \text{Map}(S)$  as  $D \setminus C'$  is an essential subsurface.  $\square$

Combining Corollary 4.3 and Proposition 4.4, we have:

**Corollary 4.5** *If  $S$  is a finite-genus surface of infinite type, then  $\text{Map}(S)$  is not residually finite.*

## 4.2 Infinite genus

In this subsection we will prove the remaining pieces of Theorem 3 involving infinite-genus surfaces.

**Proposition 4.6** *If  $S$  has infinite genus, then  $\text{PMap}_c(S)$  has no finite quotients.*

This has an immediate corollary:

**Corollary 4.7**  *$\text{PMap}_c(S)$  is perfect and does not virtually surject onto  $\mathbb{Z}$ .*

Mapping class groups of finite-type surfaces are known to be perfect [32] (except for some low-complexity cases). Motivated by Kazhdan's property (T), Ivanov conjectured that mapping class groups of finite-type surfaces do not virtually surject onto  $\mathbb{Z}$  (see [28, Problem 2.11.A]). (Equivalently, this says  $H_1(G, \mathbb{Q}) = 0$  for any finite-index subgroup  $G$  of  $\text{Map}(S)$ .) Putman and Wieland [33] have proven stability conditions for this conjecture; it would be fascinating if the stability could manifest into a statement about big mapping class groups.

As a second corollary, we have our desired result in relation to Theorem 3:

**Corollary 4.8** *Neither  $\text{Map}(S)$  nor  $\text{PMap}(S)$  is residually finite when  $S$  has infinite genus.*

The heavy lifting in the proof of Proposition 4.6 is done by a theorem of Paris [31], which says that when  $S$  is a finite-type genus- $g$  surface, the minimal index of a proper subgroup of  $\text{PMap}(S)$  is bounded below by a linear function in  $g$ . (The work of Berrick, Gebhardt and Paris [9] gives the exact value as an exponential function in  $g$ .)

**Proof of Proposition 4.6** Let  $\{F_n\}_{n \in \mathbb{N}}$  be an exhaustion of  $S$  by essential finite-type surfaces. Then  $\text{genus}(F_n) \rightarrow \infty$  as  $n \rightarrow \infty$ . Let  $m_n$  denote the minimal index of a subgroup in  $\text{PMap}(F_n)$ . Then  $\lim_{n \rightarrow \infty} m_n = \infty$  [31].

Suppose  $N$  is a proper normal subgroup of  $\text{PMap}_c(S)$  with finite index  $k$ . As  $\text{PMap}_c(S)$  is the direct limit of  $\{\text{PMap}(F_n)\}_{n \in \mathbb{N}}$ , we have that there exists  $M \in \mathbb{N}$  such that  $\text{PMap}(F_m)$  is not a subgroup of  $N$  for all  $m > M$  (otherwise  $N$  would not be proper). It follows that for  $m > M$  the intersection  $N \cap \text{PMap}(F_m)$  is a proper normal subgroup of  $\text{PMap}(F_m)$  with index at most  $k$ . However, this implies that the sequence  $\{m_n\}_{n \in \mathbb{N}}$  is bounded, a contradiction.  $\square$

## 5 Morphisms

The goal of this section is to finish the proof of Theorem 1 with Propositions 5.1 and 5.3. We then deduce corollaries about the outer automorphism group. The tools in this section combine the results on the initial topology in Section 3 with the literature on homomorphisms between mapping class groups (namely, [4; 15]). By Proposition 4.1 and Corollary 4.8, the algebraic structure of  $\text{PMap}(S)$  detects whether  $S$  has finite or infinite genus. For the remainder of the section, we focus on finite-genus surfaces.

**Proposition 5.1** *Let  $S$  and  $S'$  be finite-genus surfaces. If the genus of  $S$  is at least 3 and strictly greater than that of  $S'$ , then there is no monomorphism from  $\text{PMap}(S)$  to  $\text{PMap}(S')$ .*

**Proof** Suppose  $S$  and  $S'$  are surfaces satisfying

$$\text{genus}(S) > \max\{3, \text{genus}(S')\}$$

and that  $\psi: \text{PMap}(S) \rightarrow \text{PMap}(S')$  is a homomorphism. Let  $F \subset S$  be an essential finite-type subsurface with  $\text{genus}(F) = \text{genus}(S)$  and  $\iota: \text{PMap}(F) \hookrightarrow \text{PMap}(S)$  the inclusion homomorphism. For each  $\lambda \in \Lambda(S')$ , we obtain a homomorphism  $\psi_\lambda: \text{PMap}(F) \rightarrow \text{PMap}(S'_\lambda)$  defined by the composition  $\psi_\lambda = \varphi_\lambda \circ \psi \circ \iota$ , where  $\varphi_\lambda$  is the forgetful map defined in Section 3. Because  $F$  and  $S'_\lambda$  are finite-type and  $\text{genus}(F) > \max\{3, \text{genus}(S'_\lambda)\}$ , it follows from [4, Proposition 7.1] that  $\psi_\lambda$  is trivial.



Further, as  $\lambda$  was arbitrary, we see that

$$\psi \circ \iota(\text{PMap}(F)) \subset \bigcap_{\lambda \in \Lambda(S')} \ker \varphi_\lambda,$$

implying that  $\iota(\text{PMap}(F)) \subset \ker \psi$  by Lemma 3.4; hence,  $\psi$  is not injective. □

Continuing with our progress towards Theorem 1, we can now assume that if  $\text{PMap}(S)$  and  $\text{PMap}(S')$  are isomorphic, then they have the same genus. We have one additional lemma before finishing the final proposition towards Theorem 1.

**Lemma 5.2** *Let  $S$  and  $S'$  be surfaces such that their genera are finite, equal, and at least 4. If  $\Psi: \text{PMap}(S) \rightarrow \text{PMap}(S')$  is an isomorphism, then  $\Psi$  preserves Dehn twists.*

**Proof** Let  $T_c$  denote the Dehn twist along a simple closed curve  $c$ . Choose an essential finite-type surface  $F \subset S$  such that  $c \subset F$  and  $\text{genus}(F) = \text{genus}(S)$ . The inclusion  $F \hookrightarrow S$  induces a homomorphism  $\iota: \text{PMap}(F) \rightarrow \text{PMap}(S)$ . For each  $\lambda \in \Lambda(S')$  we define  $\iota_\lambda: \text{PMap}(F) \rightarrow \text{PMap}(S'_\lambda)$  to be the composition  $\varphi_\lambda \circ \Psi \circ \iota$ , where again  $\varphi_\lambda$  is the forgetful map defined in Section 3. If  $\iota_\lambda$  is nontrivial, then [4, Theorem 1.1] implies  $\iota_\lambda$  is induced by an inclusion  $F \hookrightarrow S'_\lambda$  (note that this is where the lower bound on genus is required<sup>1</sup>).

In particular,  $\iota_\lambda$  sends Dehn twists in  $\text{PMap}(F)$  into  $\overline{\mathcal{D}}_\lambda$ . As  $T_c$  is the image of a Dehn twist in  $\text{PMap}(F)$  under  $\iota$ , we have that  $\varphi_\lambda \circ \Psi(T_c) \in \overline{\mathcal{D}}_\lambda$  for each  $\lambda \in \Lambda(S')$ . By Proposition 3.7,  $\Psi(T_c)$  is a Dehn twist. □

**Proposition 5.3** *Let  $S$  and  $S'$  be surfaces. If the genera of  $S$  and  $S'$  are finite, equal, and at least 4, then any isomorphism between  $\text{PMap}(S)$  and  $\text{PMap}(S')$  is induced by a homeomorphism.*

**Proof** Let  $\Psi: \text{PMap}(S) \rightarrow \text{PMap}(S')$  be an isomorphism. Observe that the centers  $Z(\text{PMap}(S))$  and  $Z(\text{PMap}(S'))$  of  $\text{PMap}(S)$  and  $\text{PMap}(S')$ , respectively, are isomorphic, and therefore,  $S$  and  $S'$  have the same number of boundary components by Lemma 3.3. By applying Lemma 5.2 to both  $\Psi$  and  $\Psi^{-1}$ , we see that  $\Psi$  restricts to an isomorphism from  $\text{PMap}_c(S)$  to  $\text{PMap}_c(S')$ .

---

<sup>1</sup>The statement of [4, Theorem 1.1] requires the genus to be at least 6; however, the remark immediately following the theorem statement in the article states that this requirement can be decreased to 4 under our assumptions.

Let  $F_1 \subset F_2 \subset \dots$  be an exhaustion of  $S$  by essential finite-type surfaces such that  $\text{genus}(F_n) = \text{genus}(S)$  and  $\partial S \subset \partial F_n$  for each  $n \in \mathbb{N}$ . Recall that essential surfaces are connected by definition. It follows that the direct limit of the groups  $\{\text{PMap}(F_n)\}_{n \in \mathbb{N}}$  with respect to the natural inclusions is  $\text{PMap}_c(S)$ .

Fix  $n \in \mathbb{N}$  and let  $d_1, \dots, d_m$  be a collection of essential simple closed curves in  $F_n$  such that the set  $\{T_{d_1}, \dots, T_{d_m}\}$  of Dehn twists generates  $\text{PMap}(F_n)$ . For each  $i \in \{1, \dots, m\}$  let  $d'_i$  be the simple closed curve in  $S'$  such that  $\Psi(T_{d_i}) = T_{d'_i}$ ; the existence of  $d'_i$  is guaranteed by Lemma 5.2. Let  $R'$  be a regular neighborhood of  $d'_1 \cup \dots \cup d'_m$ . It is possible that one or more boundary components of  $R'$  bounds either a disk or a once-punctured disk in  $S'$ . If this is the case, glue the associated disk or once-punctured disk onto these boundary components; we call the resulting surface  $R_n$ .

Observe that since any Dehn twist about a boundary component of  $S'$  can be written as a product of Dehn twists about curves contained in  $R_n$ , we can conclude that  $\partial S' \subset \partial R_n$ . We now claim that the surface  $R_n$  is connected: Indeed, observe that  $d_i$  and  $d_j$  are disjoint if and only if  $d'_i$  and  $d'_j$  are disjoint. Using that  $\Psi$  is an isomorphism, the above fact follows from the fact that  $T_{d_i}$  and  $T_{d_j}$  commute if and only if  $d_i$  and  $d_j$  are disjoint. It now follows that  $d'_1 \cup \dots \cup d'_m$  is connected as  $d_1 \cup \dots \cup d_m$  is connected; hence,  $R_n$  is connected.

As  $\text{genus}(F_n) = \text{genus}(S) = \text{genus}(S')$ , we have (again by [4, Proposition 7.1]) that  $\text{genus}(R_n) = \text{genus}(F_n)$ . Now if  $f \in \text{PMap}(S')$  is in the centralizer of  $\text{PMap}(R_n)$ , then  $f$  commutes with  $T_{d'_i}$  for each  $i \in \{1, \dots, m\}$ ; in particular,  $f$  is in the centralizer of  $\Psi(\text{PMap}(F_n))$ . Conversely, if  $f \in \text{PMap}(S')$  is in the centralizer of  $\Psi(\text{PMap}(F_n))$ , then again  $f$  commutes with the Dehn twist  $T_{d'_i}$  for every  $i \in \{1, \dots, m\}$ ; therefore, as the collection of curves  $\{d'_1, \dots, d'_m\}$  is filling in  $R_n$ , we can conclude that  $f$  is in the centralizer of  $\text{PMap}(R_n)$ . The above discussion also implies that an element in the centralizer of  $\text{PMap}(R_n)$  is supported in the complement of the interior of  $R_n$ ; therefore, by Lemma 3.3, the center of the centralizer of  $\text{PMap}(R_n)$  is generated by the boundary components of  $R_n$ . In particular, the rank of the center of the centralizer of  $\text{PMap}(R_n)$  is the number of boundary components of  $R_n$  (the same holds for  $F_n$  and  $\text{PMap}(F_n)$ ). Now the center of the centralizer of  $\text{PMap}(F_n)$  and that of  $\Psi(\text{PMap}(F_n))$  are isomorphic; it follows from the above that  $F_n$  and  $R_n$  have the same number of boundary components.

Let  $Z_n$  denote the center of  $\text{PMap}(F_n)$ . From the preceding discussion, we have established that the center of the centralizer of  $\text{PMap}(R_n)$  is the center of  $\text{PMap}(R_n)$ .



an automorphism of  $\text{PMap}_c(S)$  — in the special case where  $S$  has finite genus and  $\text{Ends}(S)$  is a Cantor set — preserves Dehn twists.

As pointed out in the introduction, Corollary 2 is an immediate consequence of Proposition 5.3 and Lemma 3.3, that is, for a borderless surface  $S$  we have

$$\text{Aut}(\text{PMap}(S)) = \text{Map}^\pm(S)$$

when  $4 \leq \text{genus}(S) < \infty$ . We point out an alternative proof: By Lemma 5.2, an automorphism permutes the Dehn twists and hence induces an automorphism of the curve graph (two curves are disjoint if and only if their associated Dehn twists commute). In an recent paper [23], it is shown that the automorphism group of the curve graph is the extended mapping class group. One can then follow a — by now — standard argument to obtain Corollary 2. (This is the route taken in the recent work [7] in which Question 1.1 is answered.)

As a corollary to Corollary 2, we see the topological structure of  $\text{Ends}(S)$  becoming apparent in the algebra of  $\text{PMap}(S)$ .

**Corollary 5.4** *If  $S$  is a borderless surface whose genus is finite and at least 4, then  $\text{Out}(\text{PMap}(S))$  is isomorphic to  $\text{Homeo}(\text{Ends}(S)) \times \mathbb{Z}/2\mathbb{Z}$ .*

**Proof** Corollary 2.5 tells us the homomorphism  $\text{Map}(S) \rightarrow \text{Homeo}(\text{Ends}(S))$  is surjective and by extension  $\text{Map}^\pm(S) \rightarrow \text{Homeo}(\text{Ends}(S))$  is also surjective. It follows that there exists an order-two orientation-reversing homeomorphism fixing all the ends of  $S$ ; let  $f \in \text{Map}^\pm(S)$  be such an element. It follows that

$$\text{Map}^\pm(S) = \text{Map}(S) \rtimes \mathbb{Z}/2\mathbb{Z},$$

where the  $\mathbb{Z}/2\mathbb{Z}$  factor is identified with  $\langle f \rangle$ . It easily follows from Corollary 2 and Corollary 2.5 that

$$\text{Out}(\text{PMap}(S)) = \text{Map}^\pm(S) / \text{PMap}(S) = \text{Homeo}(\text{Ends}(S)) \times \mathbb{Z}/2\mathbb{Z}. \quad \square$$

Given Corollary 5.4, it is natural to ask if the isomorphism type of  $\text{Out}(\text{PMap}(S))$  determines the homeomorphism type of  $\text{Ends}(S)$ . This is equivalent to asking: given a closed subset  $E$  of the Cantor set, does the isomorphism type of  $\text{Homeo}(E)$  determine the homeomorphism type of  $E$ ?

Note that there is an easy counterexample: if  $E = C \sqcup X$  where  $X$  is a one-point space and  $C$  is a Cantor space, then  $\text{Homeo}(C) \cong \text{Homeo}(E)$ . It was originally

conjectured by Monk [30] that this was the only counterexample; however, another counterexample was given by McKenzie [29, Theorem 6]. On the other hand, McKenzie [29, Theorem 5] showed the conjecture holds in a large subclass; namely, if the closure of the isolated points in  $E$  is clopen, then  $\text{Homeo}(E)$  determines  $E$ . (The work of Monk and McKenzie is in the category of boolean algebras; one can translate to the language above using Stone's representation theorem.)

In light of Theorem 1, it is natural to pose the following:

**Problem 5.5** Find correspondences between algebraic invariants of  $\text{PMap}(S)$  and topological invariants of  $S$ .

As an example, we highlight the work of Monk:

**Proposition 5.6** [30, Corollary 2.1 and Theorems 3 and 5] *Let  $E$  be a closed subset of the Cantor set with infinite cardinality.*

- (1)  *$\text{Homeo}(E)$  is simple if and only if  $E$  is homeomorphic to either a Cantor space or the disjoint union of a Cantor space and the one-point space.*
- (2) *If  $m \in \mathbb{N}$  with  $1 < m < \infty$ , then  $E$  is homeomorphic to  $C \sqcup \{p_1, \dots, p_m\}$  if and only if  $\text{Homeo}(E) \cong \text{Sym}_m \times \text{Homeo}(C)$ , where  $C$  is a Cantor space.*
- (3)  *$\text{Homeo}(E)$  has exactly two proper normal subgroups if and only if  $E$  is homeomorphic to the one-point compactification of  $\mathbb{N}$ .*

## 6 The compact-open topology

We begin by restating the definition of the mapping class group: Equip  $\text{Homeo}(S)$  with the compact-open topology. Then

$$\text{Map}(S) = \pi_0(\text{Homeo}^+(S, \partial S)),$$

where  $\text{Homeo}^+(S, \partial S)$  is the group of orientation-preserving homeomorphisms of  $S$  fixing  $\partial S$  pointwise. With this definition,  $\text{Map}(S)$  comes equipped with a natural topology, namely, the corresponding quotient topology, giving  $\text{Map}(S)$  the structure of a topological group. If  $S$  is a finite-type surface, then this is the discrete topology on  $\text{Map}(S)$  (one way to see this is as a consequence of the Alexander method [17, Proposition 2.8]). We set  $\tau_q$  to be the corresponding subspace topology on  $\text{PMap}(S)$ . It follows from Lemma 2.1 that the identity component of  $\text{Homeo}^+(S)$  is closed when  $S$  is borderless. This implies that the one-point sets in  $(\text{PMap}(S), \tau_q)$

are closed; it is a standard exercise to show that in a topological group this implies the group is Hausdorff. (A similar argument works for surfaces with boundary.)

Moreover, it is metrizable:

**Proposition 6.1**  $(\text{PMap}(S), \tau_q)$  is metrizable.

**Proof** As  $\text{Homeo}(S, \partial S)$  is first countable and  $\text{Map}^\pm(S)$  is the image under an open quotient map, it follows that  $(\text{PMap}(S), \tau_q)$  is first countable. By the Birkhoff–Kakutani theorem [10; 26], every first-countable Hausdorff topological group is metrizable.  $\square$

Recall that for a finite-type surface, a combination of the Dehn–Lickorish theorem and the Birman exact sequence (see [17, Chapter 4]) tells us that  $\text{PMap}(S)$  is generated by Dehn twists. In the infinite-type setting this of course cannot be true as  $\text{PMap}(S)$  is uncountable; however, building off the result in the finite-type setting we are able to obtain topological generating sets for  $\text{PMap}(S)$ . The remainder of this section is dedicated to proving Theorem 4.

Before we get to the proof of Theorem 4, we need to introduce a class of homeomorphisms contained in  $\text{PMap}(S)$ . Consider the surface  $\Sigma$  defined by taking the surface  $\mathbb{R} \times [0, 1]$ , removing the interior of each disk of radius  $\frac{1}{4}$  with center of the form  $(n, \frac{1}{2})$ , for each  $n \in \mathbb{Z}$ , and attaching a torus with one boundary component to the boundary of each such disk. Let  $\sigma$  be the homeomorphism of  $\Sigma$  that behaves like  $(x, y) \mapsto (x+1, y)$  on the interior of  $\Sigma$  and tapers to the identity in a neighborhood of  $\partial\Sigma$  (see Figure 2). One should think of  $\sigma$  as obtained by sliding the  $n^{\text{th}}$ -handle (in the disk with center  $(n, \frac{1}{2})$ ) horizontally until it reaches the position of the  $(n+1)^{\text{th}}$ -handle.

Given a surface  $S$  with at least two ends accumulated by genus, consider an embedding of  $\Sigma$  into  $S$  inducing an injection  $\text{Ends}(\Sigma) \hookrightarrow \text{Ends}(S)$ . The homeomorphism  $\sigma$  can then be extended to all of  $S$  by the identity. The isotopy class of the extension  $\sigma$  defines an element of  $\text{PMap}(S)$ ; we call such an element a *handle shift*.

In the proof of Proposition 6.2, we will make use of the notion of a pants decomposition: a *pants decomposition* of a surface  $S$  is a maximal collection  $\mathcal{P}$  of pairwise disjoint, pairwise nonisotopic, essential simple closed curves with the property that any simple closed curve on  $S$  intersects at most finitely curves in  $\mathcal{P}$ . The complement of the union of curves in a pants decomposition is a disjoint union of surfaces each of which is homeomorphic to a thrice-punctured sphere.

Towards Theorem 4 we prove the following proposition.




Figure 2: The surface  $\Sigma$ . Outside the dotted lines, the homeomorphism  $\sigma$  is the identity. The arcs  $\sigma(a)$  and  $b$  are homotopic inside  $\Sigma$ .

**Proposition 6.2** *The set of Dehn twists topologically generate  $(\text{PMap}(S), \tau_q)$  when  $S$  has at most one end accumulated by genus. If  $S$  has at least two ends accumulated by genus, then the set of Dehn twists together with the set of handle shifts topologically generate  $(\text{PMap}(S), \tau_q)$ .*

**Proof** Let  $\mathcal{D}$  denote the collection of Dehn twists in  $\text{PMap}(S)$ . Since Dehn twists generate finite-type pure mapping class groups and  $\text{PMap}_c(S)$  is a direct limit of finite-type pure mapping class groups, it follows that  $\langle \mathcal{D} \rangle = \text{PMap}_c(S)$ . In light of the finite-type results, we will assume  $S$  is of infinite type. When  $S$  has at least two ends accumulated by genus, let  $\text{PMap}_h(S)$  denote the subgroup of  $\text{PMap}(S)$  generated by Dehn twists and handle shifts.

Take  $f \in \text{PMap}(S)$  to be an arbitrary element. Abusing notation, we will conflate  $f$  with a representative homeomorphism. Fix a pants decomposition  $\mathcal{P}$  of  $S$ . Let  $\{P_1, P_2, \dots\}$  be the components of the complement of  $\bigcup_{\gamma \in \mathcal{P}} \gamma$  in  $S$ , labeled so that the surface  $R_n = \bigcup_{i=1}^n \bar{P}_i$  is connected, where  $\bar{P}_i$  is the closure of  $P_i$  in  $S$ . Choose a component, call it  $a$ , of  $\partial P_1$  and let  $F$  be a finite-type essential surface such that

- both  $a$  and  $f(a)$  are contained in  $F$ ,
- each component of  $\partial F$  is separating, and
- each component of  $\bar{S} \setminus F$  intersects  $\text{Ends}(S)$  nontrivially.

When  $S$  has at most one end accumulated by genus, we additionally require  $F$  to be such that

- at most one component of  $S \setminus F$  has positive genus.

By the classification of surfaces, the  $\text{PMap}(S)$ -orbit of  $a$ , up to isotopy, is determined by its partition of  $\text{Ends}(S)$ , its partition of  $\partial S$ , and the topological type of its complements. (This is true for any surface and any simple closed curve in the surface.) Observe that the construction of  $F$  guarantees that  $a$  and  $f(a)$  must induce the same partition of  $\partial F$  and  $\text{Ends}(F)$  as well.

First, if  $a$  is nonseparating, then both  $a$  and  $f(a)$  are nonseparating in  $F$ , so there exists  $g_a \in \text{PMap}(F) < \text{PMap}(S)$  such that  $g_a(a) = f(a)$ . In addition, we require that  $g_a$  fixes an orientation on  $a$ .

Now suppose  $a$  is separating. In the case where  $S$  has at most one end accumulated by genus, there is a component  $U$  of  $S \setminus a$  that has finite genus. Let  $W = U \cap F$  and  $V = f(U) \cap F$ . Note that since  $S$  has at most one end accumulated by genus,  $S \setminus F$  has at most one component with positive genus, and when  $S$  is a surface of finite genus,  $F$  can be chosen to have the same genus as  $S$ . It follows that  $\text{genus}(W) = \text{genus}(V)$ , and therefore, the components of  $F \setminus a$  and  $F \setminus f(a)$  have the same topological type. In particular, there exists  $g_a \in \text{PMap}(F) < \text{PMap}(S)$  such that  $g_a(a) = f(a)$ . In both of these cases,  $g_a$  is supported on a finite-type surface and hence is contained in  $\text{PMap}_c(S)$ .

Since  $g_a^{-1} \circ f$  fixes  $a$  and both  $f$  and  $g_a$  fix  $\text{Ends}(S)$  pointwise,  $g_a^{-1} \circ f$  restricts to a homeomorphism of each component of  $S \setminus a$ . Let  $b \neq a$  be another component of  $\partial P_1$ , and let  $S'$  be the component of  $S \setminus a$  containing  $b$ . We can then repeat the above process with  $S$  replaced by  $S'$  and  $f$  by  $g_a^{-1} \circ f|_{S'}$ . If  $P_1$  has a third boundary component, then we repeat this process with this final component. In either case, we have built a homeomorphism  $g_1 \in \text{PMap}_c(S)$  such that  $g_1(P_1) = f(P_1)$  and  $g_1^{-1} \circ f$  fixes each boundary component of  $P_1$ . Thus,  $f_1 = g_1^{-1} \circ f$  restricts to the identity on  $P_1 = R_1$ .

Let  $S_2$  be the component of  $S \setminus P_1$  containing  $P_2$ . Following the same reasoning as above and working in  $\text{PMap}(S_2)$ , we can find a product of Dehn twists in  $\text{PMap}(S_2)$ , call it  $g'_2$ , such that  $g'^{-1}_2 \circ f'_1$  is the identity when restricted to  $P_2$ , where  $f'_1$  is the restriction of  $f_1$  to  $S_2$ . Choose an element  $g''_2$  in the stabilizer of  $P_1$  in  $\text{PMap}_c(S)$  whose restriction to  $S_2$  agrees with  $g'_2$ . Defining  $g_2 = g_1 \circ g''_2 \in \text{PMap}_c(S)$  we have that  $g_2^{-1} \circ f$  restricts to the identity on  $R_2$  after composing with Dehn twists about the components of  $\partial P_1$  if necessary.

Continuing in this fashion, we construct  $g_n \in \text{PMap}_c(S)$  such that  $g_n^{-1} \circ f$  restricted to  $R_n$  is the identity. As  $R_n$  is an exhaustion of  $S$ , we see that  $g_n^{-1} \circ f$  converges to the identity; hence,  $g_n \rightarrow f$  as desired.



Now suppose that  $S$  has at least two ends accumulated by genus. Note that we may no longer be able to choose  $F$  to satisfy the fourth bullet point above. Let  $a$  be a separating curve with each component of  $S \setminus a$  being infinite genus and let  $U$  be a component of  $F \setminus a$ . It is possible that  $\text{genus}(V) \neq \text{genus}(W)$ , where  $W = U \cap F$  and  $V = f(U) \cap F$ . If this is the case, there exists a handle shift  $h \in \text{PMap}_h(S)$  such that  $h(a)$  induces the same partition of  $\partial F$  and  $\text{Ends}(F)$  as  $a$  and  $f(a)$ , where  $h(a) \subset F$ , and such that the genera of  $h(U) \cap F$  and  $V$  are equal. Thus, there exists  $g_a \in \text{PMap}(F) < \text{PMap}(S)$  such that  $g_a \circ h(a) = f(a)$ . Of course,  $g_a \circ h \in \text{PMap}_h(S)$ , so we can now repeat the argument above to produce  $k_n \in \text{PMap}_h(S)$  such that  $k_n \rightarrow f$ .  $\square$

The next proposition shows that introducing handle shifts is essential when  $S$  has at least two ends accumulated by genus.

**Proposition 6.3** *If  $S$  has at least two ends accumulated by genus, then  $\text{PMap}_c(S)$  is not dense in  $(\text{PMap}(S), \tau_q)$ .*

**Proof** We begin by describing a property of all elements of  $\text{PMap}_c(S)$ , and then show that handle shifts do not have this property and cannot be approximated by those elements that do. Choose a separating simple closed curve  $a$  such that each component of  $S \setminus a$  has infinite genus. Let  $f$  be an arbitrary element of  $\text{PMap}_c(S)$ . Let  $F$  be an essential finite-type subsurface of  $S$  so that  $f \in \text{PMap}(F) < \text{PMap}(S)$  and  $F$  satisfies the first three bullet points in the proof of Proposition 6.2. Let  $U$  be a component of  $F \setminus a$ . Then  $U$  and  $f(U)$  contain the same components of  $\partial F$ . Further,  $\text{genus}(U) = \text{genus}(f(U))$ , so either  $[f(a)] = [a]$  or  $i([a], [f(a)]) > 0$ .

Now, let  $b$  be a simple closed curve so that  $a$  and  $b$  cobound a genus-1 subsurface of  $S$ . It is not hard to see that there exists a handle shift  $h \in \text{PMap}(S)$  with  $h(a) = b$ . Thus, no element in  $\text{PMap}_c(S)$  can agree with  $h$  on the compact set  $a$ , since  $b = h(a)$  and  $a$  are distinct and disjoint; hence,  $\text{PMap}_c(S)$  is not dense in  $(\text{PMap}(S), \tau_q)$ .  $\square$

Propositions 6.2 and 6.3 together imply Theorem 4.

In Lemma 3.6, we gave the closure of the set of Dehn twists in  $\tau_w$ . In the finite-genus setting, the analogous result holds for  $\tau_q$  and is obtained by combining Lemma 3.6 and Proposition 7.6. Below we give a proof in general, regardless of whether the genus of  $S$  is finite or infinite:

**Proposition 6.4** *The closure of  $\mathcal{D}$  in  $\tau_q$  is  $\overline{\mathcal{D}} = \mathcal{D} \cup \{\text{id}\}$ .*

**Proof** Let  $K_1 \subset K_2 \subset \dots$  be an exhaustion of  $S$  by essential finite-type subsurfaces. We first see that the identity is in  $\overline{\mathcal{D}}$  (which is a repeated argument from the beginning of Lemma 3.6). Let  $b_n$  be a simple closed curve contained in the complement of  $K_n$ . Then the sequence  $\{T_{b_n}\}_{n \in \mathbb{N}} \subset \mathcal{D}$  converges to the identity in  $(\text{PMap}(S), \tau_q)$ .

Following the proof outline of Lemma 3.6, we want to show that if a sequence  $\{T_{a_i}\}_{i \in \mathbb{N}} \subset \mathcal{D}$  converges to a nonidentity element  $f \in (\text{PMap}(S), \tau_q)$ , then  $f$  is compactly supported.

Assume that  $f \notin \text{PMap}_c(S)$ . It follows that there exists a natural number  $N$  such that for every integer  $n$  with  $n \geq N$  there exists an integer  $i_n$  such that the curve  $a_{i_n}$ , up to isotopy, is contained in neither  $K_n$  nor its complement and intersects  $K_N$  nontrivially.

By possibly increasing  $i_n$ , we may assume that the curve  $a_{i_n}$  has nontrivial intersection with a single boundary component of  $\partial K_N$ , call it  $b$ . For each curve  $a_{i_n}$ , let  $b_n$  be a component of  $K_n$  such that  $i(a_{i_n}, b_n) > 0$ . As the compact-open topology agrees with the topology of compact convergence, there exists  $J \in \mathbb{N}$  such that, up to isotopy,  $f(b) = T_{a_j}(b)$  for all  $j > J$ . Apply [17, Proposition 3.4] to obtain

$$i(f(b), b_n) = i(T_{a_{i_n}}(b), b_n) = i(a_{i_n}, b)i(a_{i_n}, b_n) > 0,$$

whenever  $i_n > \max\{J, N\}$ . It follows that  $f(b)$  leaves every compact set; hence, it must not be compact and  $f$  cannot be a homeomorphism. Therefore,  $f \in \text{PMap}_c(S)$ .

We can now find  $m \in \mathbb{N}$  such that all but finitely many of the  $T_{a_i}$  are elements of  $\text{PMap}(K_m) < \text{PMap}_c(S)$ . The subspace topology on  $\text{PMap}(K_m)$  is discrete, so the sequence  $\{T_{a_i}\}_{i \in \mathbb{N}}$  is eventually constant; in particular,  $f$  is an element of  $\overline{\mathcal{D}}$ .  $\square$

Note that from the above proof we see that a sequence in  $\mathcal{D}$  converges to a Dehn twist in  $\tau_q$  if and only if it is eventually constant. This is not the case in  $\tau_w$ : Consider the curves  $a_n$  in Figure 1 from Section 3 with the modification that the curve  $a_n$  partitions off  $\{0, 1, n\}$  in  $\mathbb{N} \cup \{0\}$ . The sequence  $\{T_{a_n}\} \subset \mathcal{D}$  would then converge to  $T_{a_1}$  in  $(\text{PMap}(\mathbb{C} \setminus (\mathbb{N} \cup \{0\})), \tau_w)$ , but would not converge in  $\tau_q$ .

## 7 Inverse limits

For the entirety of this section,  $S$  will be a finite-genus surface. To simplify notation, for  $\lambda \in \Lambda(S)$ , we let

$$\begin{aligned} G_\lambda &= \text{PHomeo}^+(\overline{S}, \partial S, \lambda) \\ &= \{f \in \text{Homeo}^+(\overline{S}, \partial S) : f(p) = p \text{ for every } p \in \lambda\} \end{aligned}$$

and

$$G = \text{PHomeo}^+(\bar{S}, \partial S, \text{Ends}(S)) \\ = \{f \in \text{Homeo}^+(\bar{S}, \partial S) : f(p) = p \text{ for every } p \in \text{Ends}(S)\}.$$

Equip  $\text{Homeo}(\bar{S}, \partial S)$  with the compact-open topology and equip  $G$  and each  $G_\lambda$  with the subspace topology.

Observe that  $\text{Homeo}(S, \partial S)$  and  $\text{Homeo}(\bar{S}, \partial S, \text{Ends}(S))$  are isomorphic. Furthermore, it is clear that the compact-open topology on the latter group must be at least as fine as the former. However, the classical work of Arens [5, Theorem 3] tells us that the compact-open topology on the former group is the strongest admissible topology making it a topological group; hence, the two groups are topologically isomorphic. In other words, we may treat the ends of  $S$  as marked points.

The inclusion maps  $\iota_{\lambda,\mu}: G_\mu \hookrightarrow G_\lambda$ , whenever  $\lambda \subseteq \mu$ , make  $(\{G_\lambda\}, \{\iota_{\lambda,\mu}\})$  an inverse system. We can then set  $G_L$  to be the inverse limit and define  $\iota_\lambda: G_L \rightarrow G_\lambda$  to be the projections. The initial topology on  $G$  with respect to the collection  $\{\iota_\lambda\}$  gives  $G_L$  the structure of a topological group; this is the standard topology given to an inverse limit of topological groups. With this topology, we see that  $G_L$  is topologically isomorphic to the intersection of the  $G_\lambda$ , each endowed with the compact-open topology, so

$$G_L \cong \bigcap_{\lambda \in \Lambda} G_\lambda = G.$$

For the remainder of the section we will identify  $G$  and  $G_L$ .

As  $\text{PMap}(S) = \pi_0(G)$ , we want to investigate whether  $\text{PMap}(S)$  has the structure of an inverse limit. Similar to the above, the maps  $\iota_{\lambda,\mu}$  induce forgetful maps  $\varphi_{\lambda,\mu}: \text{PMap}(S_\mu) \rightarrow \text{PMap}(S_\lambda)$ .

This defines an inverse system  $(\{\text{PMap}(S_\lambda)\}, \{\varphi_{\lambda,\mu}\})$  and we can take the inverse limit

$$L(S) = \varprojlim_{\lambda \in \Lambda} \text{PMap}(S_\lambda).$$

Let  $\pi_\lambda: L(S) \rightarrow \text{PMap}(S_\lambda)$  be the projections defining  $L(S)$ . Giving  $\text{PMap}(S_\lambda)$  the discrete topology for each  $\lambda \in \Lambda$ , we equip  $L(S)$  with the initial topology with respect to  $\{\pi_\lambda\}_{\lambda \in \Lambda}$ . The universal property of inverse limits gives a continuous homomorphism  $\Phi: (\text{PMap}(S), \tau_w) \rightarrow L(S)$ .

**Proposition 7.1**  *$\Phi$  is a topological embedding. Further,  $\Phi$  is a topological isomorphism if and only if  $S$  is of finite type.*




Figure 3: The surface  $S = \mathbb{C} \setminus \mathbb{Z}$ . The sequence of Dehn twists  $\{T_{a_n}\}_{n \in \mathbb{N}}$  converges in  $L(S)$ , but not in  $(\text{PMap}(S), \tau_w)$ .

**Proof** Lemma 3.4 implies that  $\tau_w$  separates points and hence  $\Phi$  is injective. It is clear from the definition that  $\tau_w$  agrees with the subspace topology when  $\text{PMap}(S)$  is viewed as a subset of  $L(S)$ . Note that if  $S$  is of finite type, then  $L(S) = \text{PMap}(S)$ , so for the remainder we may assume that  $S$  is of infinite type.

We now show that  $\Phi$  fails to be onto. We will show this in the case  $S = \mathbb{C} \setminus \mathbb{Z}$ ; the general case can be ascertained by replacing each integer in  $\mathbb{Z}$  with a clopen set of  $\text{Ends}(S)$  in the general case.

For each  $n \in \mathbb{N}$ , let  $a_n$  be the boundary of a regular neighborhood of the line segment on the real axis connecting 1 and  $n + 1$  as in Figure 3. Let  $T_n \in \text{PMap}(S)$  denote the Dehn twist about  $a_n$ . For each  $\lambda \in \Lambda$ , the sequence  $\{\varphi_\lambda(T_n)\}$  is eventually constant; call this limit Dehn twist  $T_\lambda \in \text{PMap}(S_\lambda)$ . It follows that the sequence  $\{T_n\}_{n \in \mathbb{N}}$  converges to  $\bar{T} \in L(S)$ , where  $\bar{T}$  has coordinates  $\varphi_\lambda(\bar{T}) = T_\lambda$ .

We claim  $\bar{T}$  is not in  $\text{PMap}(S)$ ; we proceed by contradiction. Suppose  $\bar{T} \in \text{PMap}(S)$  and conflate  $\bar{T}$  with a representative homeomorphism. Let  $b_n$  be the boundary of the ball centered at the origin with radius  $n + \frac{1}{2}$ . Using Lemma 3.5 and a similar argument as in the proof of Lemma 3.6, we see that  $i(\bar{T}(b_1), b_n) > 0$ . It follows that  $\bar{T}(b_1)$  cannot be compact, a contradiction.  $\square$

We point out the contrast between the pure homeomorphism group of  $S$  being an inverse limit and the pure mapping class group failing to do so. It is natural to ask if the topologies  $\tau_w$  and  $\tau_q$  agree as the compact-open topology and the initial topologies agree on the level of homeomorphism groups. Proposition 7.6 tells us that this also fails to be the case (when  $S$  is of infinite type).

Now recall that a topological group is *Polish* if it is separable and completely metrizable. Such groups have many nice properties and are studied extensively in descriptive set

theory. Of particular interest is the work of Rosendal [35], which provides a theory of coarse geometry for Polish groups that are not locally compact.

**Proposition 7.2**  *$L(S)$  is Polish if and only if the cardinality of  $\text{Ends}(S)$  is countable.*

Before getting to the proof of Proposition 7.2, we separate out two lemmas that are interesting to us in their own right. Observe that by the definition of the initial topology and the Dehn–Lickorish theorem, the first lemma is immediate.

**Lemma 7.3**  *$L(S)$  is topologically generated by Dehn twists.*

**Lemma 7.4** *If the cardinality of  $\text{Ends}(S)$  is uncountable, then  $(\text{PMap}(S), \tau_w)$  is not first countable.*

The proof mimics the standard proof that an uncountable product of discrete topological spaces is not first countable.

**Proof** Suppose that  $\{U_n\}_{n \in \mathbb{N}}$  is a countable neighborhood basis of the identity in  $\text{PMap}(S)$ . For each  $n \in \mathbb{N}$  there exists a finite set  $\Lambda_n \subset \Lambda$  such that

$$V_n := \bigcap_{\lambda \in \Lambda_n} \ker \varphi_\lambda \subset U_n.$$

Therefore,  $\{V_n\}_{n \in \mathbb{N}}$  is a countable neighborhood basis for the identity.

Define  $C \subset \text{Ends}(S)$  such that  $\lambda \subset C$  if and only if there exists  $n \in \mathbb{N}$  with  $\lambda \in \Lambda_n$ . As  $C$  is countable, we can find  $\mu \in \Lambda$  such that  $\mu \cap C = \emptyset$  and  $|\mu| \geq 4$ . We claim that  $\ker \varphi_\mu$  is a neighborhood of the identity not containing any of the  $V_n$ . We proceed by contradiction: Suppose that  $V_n \subset \ker \varphi_\mu$ . Pick a simple closed curve  $c$  in  $S$  bounding a disk  $D$  in  $\bar{S}$  disjoint from each  $\lambda \in \Lambda_n$  and satisfying  $|D \cap \mu| = 2$ . The Dehn twist  $T_c$  about  $c$  satisfies  $T_c \in V_n$  and  $T_c \notin \ker \varphi_\mu$ . Thus, there is no countable neighborhood basis of the identity.  $\square$

**Proof of Proposition 7.2** If  $S$  is finite-type, then  $L(S)$  is equal to  $\text{PMap}(S)$  and has the discrete topology, so the result is immediate. Suppose now that  $\text{Ends}(S)$  is countably infinite. Let  $\{\lambda_n\}_{n \in \mathbb{N}} \subset \Lambda$  be cofinal, so that

$$L(S) = \varprojlim \text{PMap}(S_{\lambda_n}).$$

We can then define the metric  $d: L(S) \rightarrow \mathbb{R}$ : for any two elements  $f, g \in L(S)$ , set  $d(f, g) = 1$  if  $\pi_{\lambda_n}(f) \neq \pi_{\lambda_n}(g)$  for every  $n \in \mathbb{N}$ , otherwise set

$$d(f, g) = \min\{2^{-n} : \pi_{\lambda_n}(f) = \pi_{\lambda_n}(g)\}.$$

It is a routine exercise to check that  $d$  is in fact a complete metric and induces the initial topology on  $L(S)$ . Lemma 7.3 tells us that  $L(S)$  is separable; hence, it is Polish. To finish the proof, Lemma 7.4 tells us that  $L(S)$  is not metrizable if the cardinality of  $\text{Ends}(S)$  is uncountable.  $\square$

**Corollary 7.5** *If the cardinality of  $\text{Ends}(S)$  is countably infinite, then  $L(S)$  is homeomorphic to the Baire space  $\mathbb{N}^{\mathbb{N}}$ .*

(Another realization of the Baire space is the space of irrational numbers as a subspace of the real line.)

**Proof** First observe that  $L(S)$  is a subspace of a direct product of discrete spaces; hence, it has a basis of clopen sets, so it is zero-dimensional. Given that  $L(S)$  is zero-dimensional and Polish, to show that  $L(S)$  is homeomorphic to the Baire space it is enough to show that every compact set has empty interior; this utilizes the topological characterization of the Baire space by Alexandrov and Urysohn (see [27, Theorem 7.7]). As in the proof of Proposition 7.2, let  $\{\lambda_n\}_{n \in \mathbb{N}} \subset \Lambda$  be a cofinal subset. With this setup, the sets of the form  $\pi_n^{-1}(g)$ , where  $\pi_n = \pi_{\lambda_n}$  and  $g \in \text{PMap}(S_{\lambda_n})$ , give a basis for the topology of  $L(S)$ .

Let  $C \subset L(S)$  be compact. Assume that  $C$  has nonempty interior. Then there exists  $n \in \mathbb{N}$  and  $g \in \text{PMap}(S_{\lambda_n})$  such that  $\pi_n^{-1}(g) \subset C$ . As  $\pi_n^{-1}(g)$  is closed and  $C$  is compact, we must have that  $\pi_n^{-1}(g)$  is compact. But, then  $\pi_{n+1}(\pi_n^{-1}(g))$  is compact as  $\pi_{n+1}$  is continuous. This is a contradiction as  $\pi_{n+1}(\pi_n^{-1}(g))$  has infinite cardinality in a discrete space.  $\square$

Even when  $\text{Ends}(S)$  is uncountable,  $L(S)$ , as an inverse limit of discrete spaces, has the structure of a complete uniform space [24, Exercise IV.8]. Hence, in either case,  $L(S)$  can be viewed as the completion of  $\text{PMap}(S)$ .

## 7.1 Comparing topologies

Recall from Lemma 7.4 that if the cardinality of  $\text{Ends}(S)$  is uncountably infinite, then  $(\text{PMap}(S), \tau_w)$  is not metrizable, so, in this case,  $\tau_w \neq \tau_q$ ; moreover, we have:

**Proposition 7.6** *If  $S$  is of finite type, then  $\tau_w = \tau_q$ ; otherwise, there is a strict containment  $\tau_w \subset \tau_q$ .*

**Proof** We have already noted that when  $S$  is of finite type both  $\tau_q$  and  $\tau_w$  are discrete. Let us assume from now on that  $S$  is of infinite type. In the above notation, for  $\lambda \in \Lambda$ , let  $\psi_\lambda: G \rightarrow \text{PMap}(S_\lambda)$  be the homomorphism factoring through  $G_\lambda$ . As  $\psi_\lambda$  is continuous for each  $\lambda \in \Lambda$ , we have that the projection homomorphism  $q: G \rightarrow (\text{PMap}(S), \tau_w)$  is continuous. By the definition of  $\tau_q$ , this shows that  $\tau_w \subseteq \tau_q$ .

By the remark preceding the statement of Proposition 7.6, we may assume that the cardinality of  $\text{Ends}(S)$  is countable. To finish the proof, we will find a sequence that converges in  $\tau_w$  but not in  $\tau_q$ . Let  $K_1 \subset K_2 \subset \dots$  be an exhaustion of  $S$  by essential finite-type surfaces each of whose genus agrees with  $S$ . Choose a sequence of homeomorphisms  $\tilde{g}_n \in G$  such that  $\tilde{g}_n$  restricts to the identity on  $K_n$ , which implies that  $\tilde{g}_n \rightarrow 1$  in  $G$ . If  $g_n = q(\tilde{g}_n)$ , then  $g_n \rightarrow 1$  in  $\tau_q$ , hence in  $\tau_w$  as well.

For each  $\lambda \in \Lambda$ , let  $n_\lambda$  be the largest integer  $n$  such that  $\lambda$  intersects each component of the complement of  $K_n$  in  $\bar{S}$ . In addition, for each  $\lambda \in \Lambda$ , choose an isotopy class of an essential simple closed curve in  $S$ , denoted by  $c_\lambda$ , such that  $c_\lambda$  is trivial in  $S_\lambda$  and  $c_\lambda \cap K_{n_\lambda}$  fills  $K_{n_\lambda}$ . Let  $T_\lambda$  denote the Dehn twist about  $c_\lambda$  and define  $h_\lambda \in \text{PMap}(S)$  to be  $T_\lambda \circ g_{n_\lambda}$ .

The map  $\Lambda \rightarrow \text{PMap}(S)$  defined by  $\lambda \mapsto h_\lambda$  gives a net in  $\text{PMap}(S)$ . Now, the collection of sets of the form  $U_\mu = \ker \varphi_\mu$  for  $\mu \in \Lambda$  form a neighborhood subbasis for the identity in  $\tau_w$ . If  $\lambda \supset \mu$ , then we claim  $h_\lambda \in U_\mu$ . First, observe that  $K_{n_\mu} \subset K_{n_\lambda}$ , so that  $\tilde{g}_{n_\lambda}$  restricted to  $K_{n_\mu}$  is the identity. It follows that  $\varphi_\mu(g_{n_\lambda})$  is trivial. Also,

$$\varphi_\mu(T_\lambda) = \varphi_{\mu,\lambda} \circ \varphi_\lambda(T_\lambda)$$

is the identity as  $T_\lambda \in \ker \varphi_\lambda$  by definition. This shows that  $h_\lambda \in U_\mu$  for all  $\lambda \supset \mu$ . In particular, the net  $\{h_\lambda\}_{\lambda \in \Lambda}$  converges to the identity in  $\tau_w$ .

We claim that this net does not converge in  $\tau_q$ . Let

$$U = \{f \in \text{PMap}(S) : f = q(\tilde{f}), \text{ where } \tilde{f}(K_1) = K_1\}.$$

Then  $U$  is an open neighborhood of the identity in  $\tau_q$ . Indeed, let  $V$  be an open regular neighborhood of  $K_1$ . Then  $U$  is the image, under  $q$ , of

$$U(K_1, V) = \{\tilde{f} \in G : \tilde{f}(K_1) \subset V\},$$

which is a basis element for the compact-open topology on  $G$ . Further, as  $q$  is both a homomorphism and a quotient map, it is open; hence  $U$  is open. Now observe that  $h_\lambda \notin U$  for any  $\lambda$ ; this follows from the fact that given any simple closed curve  $c \subset K_1$ , we have  $h_\lambda(c) = T_\lambda(c)$ , which intersects  $\partial K_1$  nontrivially.  $\square$

## References

- [1] **L V Ahlfors, L Sario**, *Riemann surfaces*, Princeton Mathematical Series 26, Princeton Univ. Press (1960) MR
- [2] **J Aramayona, A Fossas, H Parlier**, *Arc and curve graphs for infinite-type surfaces*, Proc. Amer. Math. Soc. 145 (2017) 4995–5006 MR
- [3] **J Aramayona, L Funar**, *Asymptotic mapping class groups of closed surfaces punctured along Cantor sets*, preprint (2017) arXiv
- [4] **J Aramayona, J Souto**, *Homomorphisms between mapping class groups*, Geom. Topol. 16 (2012) 2285–2341 MR
- [5] **R Arens**, *Topologies for homeomorphism groups*, Amer. J. Math. 68 (1946) 593–610 MR
- [6] **J Bavard**, *Hyperbolicité du graphe des rayons et quasi-morphismes sur un gros groupe modulaire*, Geom. Topol. 20 (2016) 491–535 MR
- [7] **J Bavard, S Dowdall, K Rafi**, *Isomorphisms between big mapping class groups*, preprint (2017) arXiv
- [8] **F Béguin, S Crovisier, F Le Roux**, *Construction of curious minimal uniquely ergodic homeomorphisms on manifolds: the Denjoy–Rees technique*, Ann. Sci. École Norm. Sup. 40 (2007) 251–308 MR
- [9] **A J Berrick, V Gebhardt, L Paris**, *Finite index subgroups of mapping class groups*, Proc. Lond. Math. Soc. 108 (2014) 575–599 MR
- [10] **G Birkhoff**, *A note on topological groups*, Compositio Math. 3 (1936) 427–430 MR
- [11] **J S Birman, A Lubotzky, J McCarthy**, *Abelian and solvable subgroups of the mapping class groups*, Duke Math. J. 50 (1983) 1107–1120 MR
- [12] **D Calegari**, *Circular groups, planar groups, and the Euler class*, from “Proceedings of the Casson Fest” (C Gordon, Y Rieck, editors), Geom. Topol. Monogr. 7, Geom. Topol. Publ., Coventry (2004) 431–491 MR
- [13] **D Calegari**, *Mapping class groups: the next generation*, blog post (2014) Available at <https://tinyurl.com/geometry-and-the-imagination>
- [14] **J Cantwell, L Conlon**, *Endperiodic automorphisms of surfaces and foliations*, preprint (2010) arXiv



- [15] **F Castel**, *Geometric representations of the braid groups*, Astérisque 378, Soc. Math. France, Paris (2016) MR
- [16] **M G Durham, F Fanoni, N G Vlamis**, *Graphs of curves on infinite-type surfaces with mapping class group actions*, preprint (2016) arXiv
- [17] **B Farb, D Margalit**, *A primer on mapping class groups*, Princeton Mathematical Series 49, Princeton Univ. Press (2012) MR
- [18] **L Funar, C Kapoudjian**, *On a universal mapping class group of genus zero*, *Geom. Funct. Anal.* 14 (2004) 965–1012 MR
- [19] **L Funar, C Kapoudjian**, *The braided Ptolemy–Thompson group is finitely presented*, *Geom. Topol.* 12 (2008) 475–530 MR
- [20] **E K Grossman**, *On the residual finiteness of certain mapping class groups*, *J. London Math. Soc.* 9 (1974/75) 160–164 MR
- [21] **J L Harer**, *The virtual cohomological dimension of the mapping class group of an orientable surface*, *Invent. Math.* 84 (1986) 157–176 MR
- [22] **J H Hernández, I Morales, F Valdez**, *The Alexander method for infinite-type surfaces*, preprint (2017) arXiv
- [23] **J H Hernández, I Morales, F Valdez**, *Isomorphisms between curve graphs of infinite-type surfaces are geometric*, preprint (2017) arXiv
- [24] **J R Isbell**, *Uniform spaces*, Mathematical Surveys 12, Amer. Math. Soc., Providence, RI (1964) MR
- [25] **N V Ivanov**, *Automorphisms of Teichmüller modular groups*, from “Topology and geometry: Rohlin seminar” (O Y Viro, editor), *Lecture Notes in Math.* 1346, Springer (1988) 199–270 MR
- [26] **S Kakutani**, *Über die Metrisation der topologischen Gruppen*, *Proc. Imp. Acad.* 12 (1936) 82–84 MR
- [27] **A S Kechris**, *Classical descriptive set theory*, Graduate Texts in Mathematics 156, Springer (1995) MR
- [28] **R Kirby**, *Problems in low dimensional manifold theory*, *Proc. Sympos. Pure Math.* XXXII, Amer. Math. Soc., Providence, RI (1978) 273–312 MR
- [29] **R McKenzie**, *Automorphism groups of denumerable Boolean algebras*, *Canad. J. Math.* 29 (1977) 466–471 MR
- [30] **J D Monk**, *On the automorphism groups of denumerable Boolean algebras*, *Math. Ann.* 216 (1975) 5–10 MR
- [31] **L Paris**, *Small index subgroups of the mapping class group*, *J. Group Theory* 13 (2010) 613–618 MR
- [32] **J Powell**, *Two theorems on the mapping class group of a surface*, *Proc. Amer. Math. Soc.* 68 (1978) 347–350 MR

- [33] **A Putman, B Wieland**, *Abelian quotients of subgroups of the mappings class group and higher Prym representations*, J. Lond. Math. Soc. 88 (2013) 79–96 MR
- [34] **I Richards**, *On the classification of noncompact surfaces*, Trans. Amer. Math. Soc. 106 (1963) 259–269 MR
- [35] **C Rosendal**, *Coarse geometry of topological groups*, preprint (2017) Available at <https://tinyurl.com/Rosendal-Coarse-Geometry17>
- [36] **B von Kerékjártó**, *Vorlesungen über Topologie, I: Flächentopologie*, Grundle. Math. Wissen. 8, Springer (1923)

*Department of Mathematics, University of California, Santa Barbara  
Santa Barbara, CA, United States*

*Department of Mathematics, Queens College, CUNY  
Flushing, NY, United States*

`patel@math.ucsb.edu`, `nicholas.vlamis@qc.cuny.edu`

<http://www.patelp.com>, <http://math.nickvlamis.com>

Received: 1 December 2017      Revised: 23 April 2018