

Naturality of the contact invariant in monopole Floer homology under strong symplectic cobordisms

MARIANO ECHEVERRIA

The contact invariant is an element in the monopole Floer homology groups of an oriented closed three-manifold canonically associated to a given contact structure. A nonvanishing contact invariant implies that the original contact structure is tight, so understanding its behavior under symplectic cobordisms is of interest if one wants to further exploit this property.

By extending the gluing argument of Mrowka and Rollin to the case of a manifold with a cylindrical end, we will show that the contact invariant behaves naturally under a strong symplectic cobordism.

As quick applications of the naturality property, we give alternative proofs for the vanishing of the contact invariant in the case of an overtwisted contact structure, its nonvanishing in the case of strongly fillable contact structures and its vanishing in the reduced part of the monopole Floer homology group in the case of a planar contact structure. We also prove that a strong filling of a contact manifold which is an L -space must be negative definite.

57R17, 57R58

1. Stating the result and some applications	1796
2. Summary of the proof	1803
3. Setting up the equations	1811
4. Transversality and fiber products	1821
5. Stretching the neck	1832
6. Generalized gluing–excision theorem	1837
Appendix. Energy and compactness	1866
References	1872

1 Stating the result and some applications

Monopole Floer homology associates to a closed, oriented, connected three-manifold Y three abelian groups $\widetilde{\mathrm{HM}}_{\bullet}(Y)$, $\widehat{\mathrm{HM}}_{\bullet}(Y)$ and $\overline{\mathrm{HM}}_{\bullet}(Y)$, which are pronounced, respectively, “H M to”, “H M from” and “H M bar”. They admit a direct sum decomposition over spin-c structures of Y , in the sense that

$$\begin{aligned}\widetilde{\mathrm{HM}}_{\bullet}(Y) &= \bigoplus_{\mathfrak{s}} \widetilde{\mathrm{HM}}_{\bullet}(Y, \mathfrak{s}), \\ \widehat{\mathrm{HM}}_{\bullet}(Y) &= \bigoplus_{\mathfrak{s}} \widehat{\mathrm{HM}}_{\bullet}(Y, \mathfrak{s}), \\ \overline{\mathrm{HM}}_{\bullet}(Y) &= \bigoplus_{\mathfrak{s}} \overline{\mathrm{HM}}_{\bullet}(Y, \mathfrak{s}).\end{aligned}$$

In fact, the previous decomposition is finite; see the work of Kronheimer and Mrowka [22, Proposition 3.1.1]. The chain complexes whose homology are the previous groups are built using solutions of a perturbed version of the three-dimensional Seiberg–Witten equations, which are at the same time critical points of a perturbed Chern–Simons–Dirac functional [22, Section 4]. There are three different types of solutions (the boundary stable, boundary unstable and irreducible solutions) and each group uses two of the three types in their corresponding construction.

Now suppose that Y is equipped with a coorientable contact structure ξ compatible with the orientation of the manifold. In practice this means that there exists a globally defined one-form θ on Y for which $\xi = \ker \theta$ and $\theta \wedge d\theta$ is positive everywhere; see Geiges [15, Lemma 1.1.1]. As we will review in a moment, ξ determines a spin-c structure $\mathfrak{s}_{\xi}$ and one can exploit the additional structure provided by ξ in order to construct an element $c(\xi) \in \widetilde{\mathrm{HM}}_{\bullet}(-Y, \mathfrak{s}_{\xi})$ known as the contact invariant of (Y, ξ) .

It is important to observe that $c(\xi)$ belongs to the monopole Floer homology groups of the manifold $-Y$, that is, Y with the opposite orientation. This is because the contact invariant $c(\xi)$ should actually be regarded as a cohomology element $c(\xi) \in \widehat{\mathrm{HM}}^{\bullet}(Y, \mathfrak{s}_{\xi})$, and there is a natural isomorphism between the groups $\widehat{\mathrm{HM}}^{\bullet}(Y, \mathfrak{s}_{\xi})$ and $\widetilde{\mathrm{HM}}_{\bullet}(-Y, \mathfrak{s}_{\xi})$; see [22, Section 22.5]. However, we will work with the homology version of the contact invariant since most of the formulas in [22] are given explicitly for the homology groups.

Monopole Floer homology also has TFQT-like features, which concretely means that given a cobordism $W: Y \rightarrow Y'$ between two three-manifolds, there are group

homomorphisms between the corresponding homology groups,

$$\begin{aligned}\widetilde{\mathrm{HM}}_{\bullet}(W, \mathfrak{s}_W) &: \widetilde{\mathrm{HM}}_{\bullet}(Y, \mathfrak{s}_Y) \rightarrow \widetilde{\mathrm{HM}}_{\bullet}(Y', \mathfrak{s}_{Y'}), \\ \widehat{\mathrm{HM}}_{\bullet}(W, \mathfrak{s}_W) &: \widehat{\mathrm{HM}}_{\bullet}(Y, \mathfrak{s}_Y) \rightarrow \widehat{\mathrm{HM}}_{\bullet}(Y', \mathfrak{s}_{Y'}), \\ \overline{\mathrm{HM}}_{\bullet}(W, \mathfrak{s}_W) &: \overline{\mathrm{HM}}_{\bullet}(Y, \mathfrak{s}_Y) \rightarrow \overline{\mathrm{HM}}_{\bullet}(Y', \mathfrak{s}_{Y'}).\end{aligned}$$

Here $\mathfrak{s}_W$ denotes a spin-c structure which restricts in an appropriate sense to the given spin-c structures on Y and Y' . Just as in the contact case, if $(W, \omega): (Y, \xi) \rightarrow (Y', \xi')$ is equipped with a symplectic form ω , it determines a spin-c structure $\mathfrak{s}_\omega$, and so it makes sense to ask the *naturality question*, that is, whether or not

(1)
$$\widetilde{\mathrm{HM}}_{\bullet}(W^\dagger, \mathfrak{s}_\omega) \, c(\xi') \stackrel{?}{=} c(\xi),$$

where $W^\dagger: -Y' \rightarrow -Y$ denotes the cobordism turned “upside-down”. The main result of this work is that the answer to the previous question is positive in the case of a strong symplectic cobordism:

Theorem 1 *Let $(W, \omega): (Y, \xi) \rightarrow (Y', \xi')$ be a strong symplectic cobordism between two contact manifolds (Y, ξ) and (Y', ξ') . Then*

$$\widetilde{\mathrm{HM}}_{\bullet}(W^\dagger, \mathfrak{s}_\omega) \, c(\xi'; \mathbb{F}_2) = c(\xi; \mathbb{F}_2).$$

Here we have added an $\mathbb{F}_2$ to our notation of the contact invariant to emphasize that we are using the coefficient field $\mathbb{F} = \mathbb{Z}/2$ so that we can ignore orientation issues for the moduli spaces. However, throughout the paper we will drop the $\mathbb{F}_2$ from our notation for simplicity. Clearly one can also ask whether or not there is an analogous statement in the case of integer coefficients. Unfortunately, Theorem H in the work of Lin, Ruberman and Saveliev [31] shows that there is no canonical choice of sign in the definition of the contact invariant, so the best naturality statement one could hope for in this case is one given up to a sign.

At this point it is important to specify that our notion of a *strong symplectic cobordism* is that of a symplectic cobordism for which the symplectic form is given in collar neighborhoods of the concave *and* convex boundaries by symplectizations of the corresponding contact structures.

To give some context we should point out that this theorem appears stated as Theorem 2.4 in the work of Sivek [43], though the reference given is a paper by Mrowka and Rollin in preparation that was never published. Also, as will be discussed later in this paper, the “special” condition imposed on the cobordism in [43] and by Mrowka and Rollin in [35] can be removed.

One can also ask what is known in the twin versions of monopole Floer homology, namely, embedded contact homology and Heegaard Floer homology. It is not by any means obvious that the corresponding homology groups from Heegaard Floer and ECH are isomorphic to the ones coming from monopole Floer homology and the proof can be found in the work of Colin, Ghiggini and Honda [4; 5; 6], Kutluhan, Lee and Taubes [24; 25; 26; 27; 28] and Taubes [46; 47; 48; 49; 50]. Also, the corresponding contact invariants in each version are isomorphic to each other.

In Heegaard Floer homology naturality holds (for example) if (Y', ξ') is obtained from (Y, ξ) by Legendrian surgery along a Legendrian knot L ; see Lisca and Stipsicz [32, Theorem 2.3]. This is an interesting case because a 1-handle surgery, or a 2-handle surgery along a Legendrian knot K with framing -1 relative to the canonical framing, gives rise to a strong symplectic cobordism. On the ECH side the contact invariant is known to be well behaved with respect to weakly exact symplectic cobordisms; see Hutchings and Taubes [20, Remark 1.11]. Moreover, Michael Hutchings has communicated to the author that he can improve this result to the case of a strong symplectic cobordism, with the additional advantage that the contact manifolds can be disconnected; see Hutchings [18].

As we will see, the contact invariant with mod 2 coefficients is a useful tool for understanding contact structures and our naturality result is good enough to find properties of this invariant, though the properties we discuss in this work were previously proven by other means. Before we mention these applications, however, we will give some brief history that puts into perspective the construction of the contact invariant and why the following results were natural things to look for.

In [21], Kronheimer and Mrowka used the contact structure of Y to extend the definition of the Seiberg–Witten invariants to the case of a compact oriented four-manifold X bounding it.

More precisely, one considers the noncompact four-manifold $X^+ = X \cup_Y ([1, \infty) \times Y)$, where $[1, \infty) \times Y$ is given the structure of an almost Kähler cone using a symplectization ω of a contact form θ defining ξ . In particular, the symplectic form induced by θ determines a canonical spin-c structure $\mathfrak{s}_\omega$ on $[1, \infty) \times Y$, which we can think of as a complex vector bundle $S = S^+ \oplus S^-$ together with a Clifford multiplication $\rho: T^*([1, \infty) \times Y) \rightarrow \text{hom}_{\mathbb{C}}(S, S)$ satisfying certain conditions.

The canonical spin-c structure $\mathfrak{s}_\omega$ carries a canonical section Φ_0 of S^+ together with a canonical spin-c connection A_0 on the spinor bundle. Kronheimer and Mrowka

then study solutions of the Seiberg–Witten equations on X^+ which are asymptotic to (A_0, Φ_0) on the conical end. These solutions end up having uniform exponential decay with respect to the canonical configuration (A_0, Φ_0) (Proposition 3.15 in [21] or Propositions 5.7 and 5.10 in Zhang [53] for a similar situation), which means that the Seiberg–Witten equations on X^+ behave very similarly to how they would if the manifold were compact, more specifically, the moduli spaces of gauge-equivalence classes of such solutions are compact. This allows us, as in the closed manifold case, to define a map

$$\mathrm{SW}_{(X, \xi)}: \mathrm{Spin}^c(X, \xi) \rightarrow \mathbb{Z},$$

where $\mathrm{Spin}^c(X, \xi)$ denotes the set of isomorphism classes of relative spin-c structures on X that restrict to the spin-c structure $\mathfrak{s}_\xi$ on Y determined by the contact structure ξ . This map can be used to detect properties of contact structures on three-manifolds. For example, [21, Theorem 1.3] shows that for any closed three-manifold Y there are only finitely many homotopy classes of 2-plane fields which are realized as semifillable contact structures. In Section 1.3 of the same paper, Kronheimer and Mrowka mention as well that if (X, ξ) is a four-manifold with an overtwisted contact structure on its boundary, then $\mathrm{SW}_{(X, \xi)}$ vanishes identically.

The latter result is Corollary B in a different paper [35] by Mrowka and Rollin, where they analyzed how the map $\mathrm{SW}_{(X, \xi)}$ behaves under a symplectic cobordism $(W, \omega): (Y, \xi) \rightarrow (Y', \xi')$ which they called a *special symplectic cobordism* [35, page 4]. Theorem D in [35] shows that

$$(2) \quad \mathrm{SW}_{(X, \xi)} = \pm \mathrm{SW}_{(X \cup W, \xi')} \circ J,$$

where $J: \mathrm{Spin}^c(X, \xi) \rightarrow \mathrm{Spin}^c(X \cup W, \xi')$ is a canonical map that extends the spin-c structure of X across the cobordism W . With respect to $\mathbb{Z}/2\mathbb{Z}$ coefficients, the previous theorem can be interpreted as saying that the mod 2 Seiberg–Witten invariants are the same.

In order to detect more properties of the contact structure, we need to use the machinery of monopole Floer homology, whose canonical reference is the work of Kronheimer and Mrowka [22].

As first defined by Kronheimer, Mrowka, Ozsváth and Szabó [23, Section 6.3], one constructs the contact invariant $\mathfrak{c}(\xi) \in \widetilde{\mathrm{HM}}_\bullet(-Y, \mathfrak{s}_\xi)$ by studying the Seiberg–Witten equations on $(\mathbb{R}^+ \times -Y) \cup ([1, \infty) \times Y)$ which are asymptotic on the symplectic cone to the canonical configuration (A_0, Φ_0) mentioned before and asymptotic on

the half-cylinder to a solution of the (perturbed) three-dimensional Seiberg–Witten equations. We will give more details about this construction in the next section. However, it should be clear that based on the analogy with the numerical Seiberg–Witten invariants $\text{SW}_{(X,\xi)}$, one would expect the naturality property (our main result, [Theorem 1](#)) as well as the vanishing of the contact invariant for an overtwisted structure. It is the latter which we now indicate how to prove.

Corollary 2 *Let (Y, ξ) be an overtwisted contact three-manifold. Then the contact invariant of ξ vanishes, that is, $c(\xi) = 0$.*

Proof First we show that the 3–sphere S^3 admits an overtwisted structure ξ_{ot} for which $c(\xi_{\text{ot}}) = 0$. For this we will use the theorem of Eliashberg [[10](#), Theorem 1.6.1] on the existence of an overtwisted contact structure in every homotopy class of an oriented plane field, and the fact that the Floer groups of any three-manifold Y are graded by the set of homotopy classes of oriented plane fields [[22](#), Section 3.1].

Thanks to [[22](#), Proposition 3.3.1], which describes the Floer homology groups of S^3 , we can find a homotopy class of plane field $[\xi]$ for which $\widetilde{\text{HM}}_{[\xi]}(S^3) = 0$. Notice that in this case we are not specifying the spin-c structure because S^3 has only one up to isomorphism. By Eliashberg’s theorem we can choose an overtwisted structure ξ_{ot} in the homotopy class $[\xi]$. Now, $c(\xi_{\text{ot}})$ is supported in $\widetilde{\text{HM}}_{[\xi]}(-S^3) \simeq \widetilde{\text{HM}}_{[\xi]}(S^3) = 0$ and so it will automatically vanish, ie $c(\xi_{\text{ot}}) = 0$.

Now, if (Y, ξ) is an arbitrary overtwisted contact three-manifold, using a result of Etnyre and Honda [[12](#), Theorem 1.2], we can find a Stein cobordism $(W, \omega): (Y, \xi) \rightarrow (S^3, \xi_{\text{ot}})$. Such cobordisms are in fact strong cobordisms so we can conclude that

$$c(\xi) = \widetilde{\text{HM}}_{\bullet}(W^{\dagger}, \mathfrak{s}_{\omega}) c(\xi_{\text{ot}}) = \widetilde{\text{HM}}_{\bullet}(W^{\dagger}, \mathfrak{s}_{\omega})(0) = 0,$$

and therefore $c(\xi)$ vanishes. □

Remark 3 For a proof that does not use the naturality property, see Taubes [[45](#), Theorem 4.2]. The vanishing of the contact invariant for overtwisted contact structures is also known on the Heegaard Floer side; see Ozsváth and Szabó [[41](#), Theorem 1.4]. For a proof on the ECH side, see Hutchings’ blog [[19](#)]. In fact, in the case of ECH one can show that the contact invariant vanishes in the case of planar torsion; see Wendl [[51](#)]. The same is also true in the monopole Floer homology side thanks to our naturality result and Wendl [[52](#), Theorem 1].

Corollary 4 *Let (X, ω) be a strong filling of (Y, ξ) . Then the contact invariant of ξ is nonvanishing, that is, $c(\xi) \neq 0$.*

Proof By Darboux's theorem we can remove a standard small ball B of X to obtain a strong cobordism $(W, \omega): (S^3, \xi_{\text{tight}}) \rightarrow (Y, \xi)$. Naturality says that $c(\xi_{\text{tight}}) = \widetilde{\text{HM}}_{\bullet}(W^{\dagger}, \mathfrak{s}_{\omega}) c(\xi)$ but the left-hand side is nonvanishing and so we conclude that $c(\xi)$ is nonvanishing as well. $\square$

Remark 5 The Heegaard Floer version of this fact appears in the work of Ghiggini [16, Theorem 2.13]. That same paper contains an example of a weak filling where the contact invariant vanishes.

To explain the next corollary we do a quick review of some of the properties of the monopole Floer homology groups. Formally they behave like the ordinary homology groups $H_*(Z)$, $H_*(Z, A)$ and $H_*(A)$ for a pair of spaces in that they are related by a long exact sequence [22, Section 3.1]

$$(3) \quad \cdots \xrightarrow{i_*} \widetilde{\text{HM}}_{\bullet}(Y, \mathfrak{s}) \xrightarrow{j_*} \widehat{\text{HM}}_{\bullet}(Y, \mathfrak{s}) \xrightarrow{p_*} \overline{\text{HM}}_{\bullet}(Y, \mathfrak{s}) \xrightarrow{i_*} \widetilde{\text{HM}}_{\bullet}(Y, \mathfrak{s}) \xrightarrow{j_*} \cdots$$

An important subgroup of $\widehat{\text{HM}}_{\bullet}(Y, \mathfrak{s})$ is the image of $j_*: \widetilde{\text{HM}}_{\bullet}(Y, \mathfrak{s}) \rightarrow \widehat{\text{HM}}_{\bullet}(Y, \mathfrak{s})$, which is known as the *reduced Floer homology group* and denoted by $\text{HM}_{\bullet}(Y, \mathfrak{s})$, and in general it is of great interest to determine whether or not a particular element belongs to it. For example, if $j_* = 0$ we say that Y is an L -space in analogy with the terminology from Heegaard Floer [22, Section 42.6]. To relate this question to the naturality of the contact invariant, we need to use the fact that for a cobordism $(W^{\dagger}, \mathfrak{s}_W): (-Y', \mathfrak{s}_{Y'}) \rightarrow (-Y, \mathfrak{s}_Y)$ there is a commutative diagram,

$$(4) \quad \begin{array}{ccccccc} \cdots & \widetilde{\text{HM}}_{\bullet}(-Y', \mathfrak{s}_{Y'}) & \xrightarrow{j_*} & \widehat{\text{HM}}_{\bullet}(-Y', \mathfrak{s}_{Y'}) & \xrightarrow{p_*} & \overline{\text{HM}}_{\bullet}(-Y', \mathfrak{s}_{Y'}) & \xrightarrow{i_*} & \widetilde{\text{HM}}_{\bullet}(-Y', \mathfrak{s}_{Y'}) & \cdots \\ & \downarrow \widetilde{\text{HM}}_{\bullet}(W^{\dagger}, \mathfrak{s}_W) & & \downarrow \widehat{\text{HM}}_{\bullet}(W^{\dagger}, \mathfrak{s}_W) & & \downarrow \overline{\text{HM}}_{\bullet}(W^{\dagger}, \mathfrak{s}_W) & & \downarrow \widetilde{\text{HM}}_{\bullet}(W^{\dagger}, \mathfrak{s}_W) & \\ \cdots & \widetilde{\text{HM}}_{\bullet}(-Y, \mathfrak{s}_Y) & \xrightarrow{j_*} & \widehat{\text{HM}}_{\bullet}(-Y, \mathfrak{s}_Y) & \xrightarrow{p_*} & \overline{\text{HM}}_{\bullet}(-Y, \mathfrak{s}_Y) & \xrightarrow{i_*} & \widetilde{\text{HM}}_{\bullet}(-Y, \mathfrak{s}_Y) & \cdots \end{array}$$

Corollary 6 *Let (X, ω) be a strong filling of (Y', ξ') . Assume in addition that Y' is an L -space. Then X must be negative definite.*

Proof Suppose by contradiction that $b^+(X) \geq 1$. Remove a Darboux ball as before to obtain a cobordism $(W, \omega): (S^3, \xi_{\text{tight}}) \rightarrow (Y', \xi')$. By [22, Proposition 3.5.2], we have that $\overline{\text{HM}}_{\bullet}(W^{\dagger}, \mathfrak{s}_{\omega}) = 0$. By the commutative diagram and the fact that j_*

vanishes for Y' , we have that $c(\xi') \in \ker j_* = \text{im } i_*$. Hence $c(\xi') = i_*([\Psi'])$ for some $[\Psi'] \in \overline{\text{HM}}_\bullet(-Y', \mathfrak{s}_{\xi'})$ and the commutativity together with the naturality says that

$$0 = i_* \overline{\text{HM}}(W^\dagger, \mathfrak{s}_\omega)([\Psi']) = \widetilde{\text{HM}}_\bullet(W^\dagger, \mathfrak{s}_\omega) c(\xi') = c(\xi_{\text{tight}}),$$

which is a contradiction. $\square$

Remark 7 Corollary 6 appears as Theorem 1.4 in Ozsváth and Szabó [40].

Corollary 8 Suppose that (Y, ξ) is a planar contact manifold. Then $j_* c(\xi) = 0$ and in particular any strong filling of a planar contact manifold must be negative definite.

Proof Observe that the last statement is exactly the proof of the previous corollary, which only used the fact that $c(\xi) \in \ker j_*$. If (Y, ξ) is a planar contact manifold, Theorem 4 in Wendl [52] (and the remarks after it) shows that there is a strong symplectic cobordism $(W, \omega): (Y, \xi) \rightarrow (S^3, \xi_{\text{tight}})$. The result follows using the commutative diagram (4) and the fact that j_* vanishes on S^3 because it admits a metric of positive scalar curvature [22, Proposition 36.1.3]. $\square$

Remark 9 Theorem 1.2 in Ozsváth, Stipsicz and Szabó [39] shows that if the contact structure ξ on Y is compatible with a planar open book decomposition then its contact invariant vanishes when regarded as an element of the quotient group $\text{HF}_{\text{red}}(-Y, \mathfrak{s}_\xi)$. The second part of our corollary should be compared with Theorem 1.2 in Etnyre [11], where it is shown (among other things) that any symplectic filling of a planar contact manifold is negative definite.

The proof of the previous corollary can be extended to the case when Y' admits a metric with positive scalar curvature. First of all, it should be pointed out that this class of manifolds is not very large. Thanks to results of Schoen and Yau an orientable three-manifold with positive scalar curvature can always be obtained from a manifold with positive scalar curvature with $b_1 = 0$ by making a connected sum of a number of copies of $S^1 \times S^2$.

Corollary 10 Suppose that $(W, \omega): (Y, \xi) \rightarrow (Y', \xi')$ is a strong symplectic cobordism with Y' (hence $-Y'$) admitting a metric with positive scalar curvature. Then:

- (a) If $c_1(\mathfrak{s}_{\xi'})$ is not torsion, then the contact invariant $c(\xi')$ vanishes automatically and by naturality so will the contact invariant $c(\xi)$.
- (b) If $c_1(\mathfrak{s}_{\xi'})$ is torsion, then $j_* c(\xi') = 0$ and so by naturality $j_* c(\xi) = 0$. In particular, if there exists a strong cobordism $(W, \omega): (Y, \xi) \rightarrow (S^3, \xi_{\text{tight}})$ we must have that $j_* c(\xi) = 0$.

Proof Proposition 36.1.3 in [22] shows that j_* vanishes when $c_1(\mathfrak{s})$ is torsion and that the Floer groups are zero when $c_1(\mathfrak{s})$ is not torsion, from which the corollary follows immediately. $\square$

In the next section we will sketch the main argument in the proof of Theorem 1. It is our hope that this summary captures the essential ideas of the proof of our main theorem, since the remaining (and more technical) part of the paper will follow in large part the paper of Mrowka and Rollin [35], which is “required reading” for someone interested in understanding why the naturality theorem will be true.

Acknowledgements I would like to thank my advisor Thomas Mark for suggesting this problem to me and for all of his indispensable help and support. Also, I would like to thank Tomasz Mrowka for advising me with the gluing argument and other technical issues, Jianfeng Lin for many useful conversations and Boyu Zhang for explaining to me several aspects of his paper [53] and other discussions key for this paper. Finally, I would like to thank the referee for many helpful comments and corrections.

2 Summary of the proof

As stated before, we now give a brief summary of the main ideas involved in the proof of Theorem 1. In a nutshell, to show that $c(\xi)$ equals $\widetilde{\mathrm{HM}}_\bullet(W^\dagger, \mathfrak{s}_\omega) c(\xi')$, we will define an intermediate “hybrid” invariant $c(\xi', Y) \in \widetilde{\mathrm{HM}}_\bullet(-Y, \mathfrak{s}_\xi)$ which will work as a bridge between $c(\xi)$ and $\widetilde{\mathrm{HM}}_\bullet(W^\dagger, \mathfrak{s}_\omega) c(\xi')$. Namely, using a “stretching the neck” argument we will show that

$$\widetilde{\mathrm{HM}}_\bullet(W^\dagger, \mathfrak{s}_\omega) c(\xi') = c(\xi', Y),$$

while adapting the strategy of [35] (which as we will explain in a moment involves a “dilating the cone” argument), we will show that

$$c(\xi', Y) = c(\xi),$$

giving us the desired naturality result.

First we review the definition of the contact invariant, following [23, Section 6.2] (in that paper the contact invariant was denoted by $[\check{\psi}_{Y,\xi}]$ but we have decided to switch to the more standard notation used in Heegaard Floer homology). As mentioned in the introduction, given a contact manifold (Y, ξ) we construct the manifold

$$Z_{Y,\xi}^+ = (\mathbb{R}^+ \times -Y) \cup ([1, \infty) \times Y)$$

and study the Seiberg–Witten equations which are asymptotic to the canonical solution (A_0, Φ_0) on the conical end $[1, \infty) \times Y$ and to a critical point c of the three-dimensional Seiberg–Witten equations on the cylindrical end $\mathbb{R}^+ \times -Y$. To write the Seiberg–Witten equations a choice of spin- c structure needs to be made, and in this case the contact structure ξ determines a canonical spin- c structure $\mathfrak{s}$ on $Z_{Y,\xi}^+$, which we will describe later.

There is a gauge group action on such solutions and we define the moduli space $\mathcal{M}(Z_{Y,\xi}^+, \mathfrak{s}, [c])$ as the gauge-equivalence classes of the solutions to the Seiberg–Witten equations on $Z_{Y,\xi}^+$. As a matter of notation, $[\cdot]$ will represent the gauge-equivalence class of a configuration, so $[c]$ in this case denotes the gauge-equivalence class of the critical point c . The moduli space $\mathcal{M}(Z_{Y,\xi}^+, \mathfrak{s}, [c])$ is not equidimensional, in fact, it admits a partition into components of different topological type,

$$\mathcal{M}(Z_{Y,\xi}^+, \mathfrak{s}, [c]) = \bigcup_z \mathcal{M}_z(Z_{Y,\xi}^+, \mathfrak{s}, [c]),$$

where z indexes the different connected components of $\mathcal{M}(Z_{Y,\xi}^+, \mathfrak{s}, [c])$. We count points in the zero-dimensional moduli spaces (which will be compact, hence finite) and define

$$m_z(Z_{Y,\xi}^+, \mathfrak{s}, [c]) = \begin{cases} |\mathcal{M}_z(Z_{Y,\xi}^+, \mathfrak{s}, [c])| \bmod 2 & \text{if } \dim \mathcal{M}_z(Z_{Y,\xi}^+, \mathfrak{s}, [c]) = 0, \\ 0 & \text{otherwise.} \end{cases}$$

The contact invariant is then defined at the chain level as

(5)
$$c(\xi) = (c^o(\xi), c^s(\xi)) \in \check{C}_*(-Y, \mathfrak{s}_\xi) = \mathfrak{C}^o(-Y, \mathfrak{s}_\xi) \oplus \mathfrak{C}^s(-Y, \mathfrak{s}_\xi)$$

by

$$\begin{aligned} c^o(\xi) &= \sum_{[a] \in \mathfrak{C}^o(-Y, \mathfrak{s}_\xi)} \sum_z m_z(Z_{Y,\xi}^+, \mathfrak{s}, [a]) e_{[a]}, \\ c^s(\xi) &= \sum_{[a] \in \mathfrak{C}^s(-Y, \mathfrak{s}_\xi)} \sum_z m_z(Z_{Y,\xi}^+, \mathfrak{s}, [a]) e_{[a]}. \end{aligned}$$

In the above notation $\check{C}_*(-Y, \mathfrak{s}_\xi)$ is the free abelian group generated by the irreducible critical points $[a] \in \mathfrak{C}^o(-Y, \mathfrak{s}_\xi)$ and the boundary stable critical points $[a] \in \mathfrak{C}^s(-Y, \mathfrak{s}_\xi)$. Also, $e_{[a]}$ is a bookkeeping device for each critical point considered as a generator in the group. Lemma 6.6 in [23] then shows that $c(\xi)$ is a cycle, that is, it defines an element $c(\xi)$ of the monopole Floer homology group $\widetilde{\text{HM}}_\bullet(-Y, \mathfrak{s}_\xi)$.

Returning to the naturality question, suppose that we have a symplectic cobordism $(W, \omega): (Y, \xi) \rightarrow (Y', \xi')$ and we want to decide whether $\widetilde{\text{HM}}_\bullet(W^\dagger, \mathfrak{s}_\omega) c(\xi') = c(\xi)$.

Clearly this is equivalent to showing that at the *chain* level

$$\check{m}c(\xi') - c(\xi) \in \text{im } \check{\partial}_{-Y},$$

where $\check{\partial}_{-Y}: \check{C}_*(-Y, \mathfrak{s}_\xi) \rightarrow \check{C}_*(-Y, \mathfrak{s}_\xi)$ is the differential that generates $\widetilde{\text{HM}}_\bullet(-Y, \mathfrak{s}_\xi)$. Here $\check{m}$ is the chain map [22, Definition 25.3.3]

$$\check{m} = \begin{pmatrix} m_o^o & -m_o^u \bar{\partial}_u^s - \partial_o^u \bar{m}_u^s \\ m_s^o & \bar{m}_s^s - m_s^u \bar{\partial}_u^s - \partial_s^u \bar{m}_u^s \end{pmatrix}: \check{C}_\bullet(-Y', \mathfrak{s}_{\xi'}) \rightarrow \check{C}_\bullet(-Y, \mathfrak{s}_\xi).$$

To see what $\check{m}$ does, we will explain the meaning of m_s^o and $\bar{\partial}_u^s$, since the action of the remaining terms can be inferred easily from these two examples. The map m_s^o counts solutions on $W^\dagger: -Y' \rightarrow -Y$ with a half-cylinder attached on each end,

$$W_*^\dagger = (\mathbb{R}^- \times -Y') \cup W^\dagger \cup (\mathbb{R}^+ \times -Y),$$

which are asymptotic on $\mathbb{R}^- \times -Y'$ to an irreducible critical point $[a] \in \mathfrak{C}^o(-Y', \mathfrak{s}_{\xi'})$ and asymptotic on $\mathbb{R}^+ \times -Y$ to a boundary stable critical point $[b] \in \mathfrak{C}^s(-Y, \mathfrak{s}_\xi)$; see Figure 1.

Again, we obtain a moduli space $\mathcal{M}([a], W_*^\dagger, \mathfrak{s}_\omega, [b])$ and as before we can define

$$n_z([a], W_*^\dagger, \mathfrak{s}_\omega, [b]) = \begin{cases} |\mathcal{M}_z([a], W_*^\dagger, \mathfrak{s}_\omega, [b])| \bmod 2 & \text{if } \dim \mathcal{M}_z([a], W_*^\dagger, \mathfrak{s}_\omega, [b]) = 0, \\ 0 & \text{otherwise.} \end{cases}$$

On the other hand, the map $\bar{\partial}_u^s$ counts solutions on $\mathbb{R} \times -Y'$ which are asymptotic to a boundary stable critical point $[a] \in \mathfrak{C}^s(-Y', \mathfrak{s}_{\xi'})$ as $t \rightarrow -\infty$ and to a boundary unstable critical point $[b] \in \mathfrak{C}^u(-Y', \mathfrak{s}_{\xi'})$ as $t \rightarrow \infty$ (in our context a map like ∂_o^u would count solutions on $\mathbb{R} \times -Y$ instead). The bar indicates that we are only considering reducible solutions, ie solutions where the spinor vanishes identically. In the case of a cylinder there is a natural $\mathbb{R}$ action and the corresponding moduli space after we quotient out by this action is denoted by $\check{\mathcal{M}}([a], \mathfrak{s}_{\xi'}, [b])$ (the notation in [22] for this moduli space is $\check{M}_z([a], \mathfrak{s}_{\xi'}, [b])$). In this case we define

$$n_z([a], \mathfrak{s}_{\xi'}, [b]) = \begin{cases} |\check{\mathcal{M}}_z([a], \mathfrak{s}_{\xi'}, [b])| \bmod 2 & \text{if } \dim \check{\mathcal{M}}_z([a], \mathfrak{s}_{\xi'}, [b]) = 0, \\ 0 & \text{otherwise.} \end{cases}$$

From the formula one can see that $\check{m}c(\xi')$ has two terms, and since are working mod 2 we will write them without the signs to simplify the expression. The term corresponding to

$$m_o^o c^o(\xi') + m_o^u \bar{\partial}_u^s c^s(\xi') + \partial_o^u \bar{m}_u^s c^s(\xi')$$

is equivalent to

$$\begin{aligned}
 & \sum_{\substack{[a] \in \mathfrak{C}^o(-Y') \\ [c] \in \mathfrak{C}^o(-Y)}} \sum_{z_1, z_2} m_{z_1}(Z_{Y', \xi'}^+, s', [a]) n_{z_2}([a], W_*^\dagger, s_\omega, [c]) e_{[c]} \\
 & + \sum_{\substack{[a] \in \mathfrak{C}^s(-Y') \\ [b] \in \mathfrak{C}^u(-Y') \\ [c] \in \mathfrak{C}^o(-Y)}} \sum_{z_1, z_2, z_3} m_{z_1}(Z_{Y', \xi'}^+, s', [a]) \bar{n}_{z_2}([a], s_{\xi'}, [b]) n_{z_3}([b], W_*^\dagger, s_\omega, [c]) e_{[c]} \\
 & + \sum_{\substack{[a] \in \mathfrak{C}^s(-Y') \\ [b] \in \mathfrak{C}^u(-Y') \\ [c] \in \mathfrak{C}^o(-Y)}} \sum_{z_1, z_2, z_3} m_{z_1}(Z_{Y', \xi'}^+, s', [a]) \bar{n}_{z_2}([a], W_*^\dagger, s_\omega, [b]) n_{z_3}([b], s_\xi, [c]) e_{[c]}.
 \end{aligned}$$

Notice that if we fix a critical point $[c] \in \mathfrak{C}^o(-Y, s_\xi)$ we can consider the coefficient

$$\begin{aligned}
 (6) \quad & \sum_{[a] \in \mathfrak{C}^o(-Y')} \sum_{z_1, z_2} m_{z_1}(Z_{Y', \xi'}^+, s', [a]) n_{z_2}([a], W_*^\dagger, s_\omega, [c]) \\
 & + \sum_{\substack{[a] \in \mathfrak{C}^s(-Y') \\ [b] \in \mathfrak{C}^u(-Y')}} \sum_{z_1, z_2, z_3} m_{z_1}(Z_{Y', \xi'}^+, s', [a]) \bar{n}_{z_2}([a], s_{\xi'}, [b]) n_{z_3}([b], W_*^\dagger, s_\omega, [c]) \\
 & + \sum_{\substack{[a] \in \mathfrak{C}^s(-Y') \\ [b] \in \mathfrak{C}^u(-Y')}} \sum_{z_1, z_2, z_3} m_{z_1}(Z_{Y', \xi'}^+, s', [a]) \bar{n}_{z_2}([a], W_*^\dagger, s_\omega, [b]) n_{z_3}([b], s_\xi, [c]).
 \end{aligned}$$

Similarly, for each critical point $[c] \in \mathfrak{C}^s(-Y, s_\xi)$, the coefficient of $e_{[c]}$ in

$$m_s^o c^o(\xi') + \bar{m}_s^s c^s(\xi') + m_s^u \bar{\partial}_u^s c^s(\xi') + \partial_s^u \bar{m}_u^s c^s(\xi')$$

is given by

$$\begin{aligned}
 (7) \quad & \sum_{[a] \in \mathfrak{C}^o(-Y')} \sum_{z_1, z_2} m_{z_1}(Z_{Y', \xi'}^+, s', [a]) n_{z_2}([a], W_*^\dagger, s_\omega, [c]) \\
 & + \sum_{[a] \in \mathfrak{C}^s(-Y')} \sum_{z_1, z_2} m_z(Z_{Y', \xi'}^+, s', [a]) \bar{n}_{z_2}([a], W_*^\dagger, s_\omega, [c]) \\
 & + \sum_{\substack{[a] \in \mathfrak{C}^s(-Y') \\ [b] \in \mathfrak{C}^u(-Y')}} \sum_{z_1, z_2, z_3} m_{z_1}(Z_{Y', \xi'}^+, s', [a]) \bar{n}_{z_2}([a], s_{\xi'}, [b]) n_{z_3}([b], W_*^\dagger, s_\omega, [c]) \\
 & + \sum_{\substack{[a] \in \mathfrak{C}^s(-Y') \\ [b] \in \mathfrak{C}^u(-Y')}} \sum_{z_1, z_2, z_3} m_{z_1}(Z_{Y', \xi'}^+, s', [a]) \bar{n}_{z_2}([a], W_*^\dagger, s_\omega, [b]) n_{z_3}([b], s_\xi, [c]).
 \end{aligned}$$

Therefore, we want to show that up to a boundary term, $\sum_z m_z(Z_{Y', \xi'}^+, s, [c])$ is equal to (6) (if $[c]$ is irreducible) or (7) (if $[c]$ is boundary stable).




Figure 1: Manifold $W_*^\dagger$ with two cylindrical ends used to define the cobordism maps.

If there is any hope of showing the equality between these two quantities we need to find a geometric interpretation to the sums (6) and (7). In order to do this we will consider the Seiberg–Witten equations in a slightly more general scenario, one that combines the construction of the contact invariant with the cobordism. More precisely, we will study the Seiberg–Witten equations on

$$W_{\xi',Y}^+ = ([1,\infty) \times Y') \cup W^\dagger \cup (\mathbb{R}^+ \times -Y)$$

which are asymptotic on $[1,\infty) \times Y'$ to the canonical solution coming from the contact structure ξ' and asymptotic on $\mathbb{R}^+ \times -Y$ to a critical point $[c] \in \check{C}_*(-Y, \mathfrak{s}_\xi)$; see Figure 2. The moduli space of such solutions will naturally be denoted by $\mathcal{M}(W_{\xi',Y}^+, \mathfrak{s}_\omega, [c])$.

Thanks to the compactness arguments in [21; 22] and [53] (which guarantee uniform exponential decay along the conical end) we can proceed as before and define

$$m_z(W_{\xi',Y}^+, \mathfrak{s}_\omega, [c]) = \begin{cases} |\mathcal{M}_z(W_{\xi',Y}^+, \mathfrak{s}_\omega, [c])| \bmod 2 & \text{if } \dim \mathcal{M}_z(W_{\xi',Y}^+, \mathfrak{s}_\omega, [c]) = 0, \\ 0 & \text{otherwise.} \end{cases}$$




Figure 2: Manifold $W_{\xi',Y}^+$ used to define the “hybrid” invariant $c(\xi', Y)$.




Figure 3: $\widetilde{\mathrm{HM}}(W^\dagger, \mathfrak{s}_\omega) \, c(\xi') = c(\xi', Y)$ via a “stretching the neck” argument.

These numbers give rise to the hybrid invariant $c(\xi', Y)$ mentioned at the beginning of this section. In order to show the equality $\widetilde{\mathrm{HM}}_\bullet(W^\dagger, \mathfrak{s}_\omega) \, c(\xi') = c(\xi', Y)$ we must consider the parametrized moduli space

(8)
$$\bigcup_{L \in [0, \infty)} \{L\} \times \mathcal{M}(W_{\xi', Y}^+(L), \mathfrak{s}_\omega, [c]),$$

where $\mathcal{M}(W_{\xi', Y}^+(L), \mathfrak{s}_\omega, [c])$ denotes the moduli space of solutions to the Seiberg–Witten equations on the manifold

$$W_{\xi', Y}^+(L) = ([1, \infty) \times Y') \cup ([0, L] \times -Y') \cup W^\dagger \cup (\mathbb{R}^+ \times -Y).$$

The parametrized moduli space (8) is not compact; its compactification will be denoted by

(9)
$$\bigcup_{L \in [0, \infty]} \{L\} \times \mathcal{M}^+(W_{\xi', Y}^+(L), \mathfrak{s}_\omega, [c]),$$

where the definition of $\mathcal{M}^+(W_{\xi', Y}^+(\infty), \mathfrak{s}_\omega, [c])$ is given in (27). For now, it suffices to say that when we count the endpoints of all one-dimensional moduli spaces inside (9) we will get 0.

The count coming from the fiber over $L = 0$ will give the term $\sum_z m_z(W_{\xi', Y}^+, \mathfrak{s}_\omega, [c])$ corresponding to the hybrid invariant $c(\xi', Y)$ while the count coming from the fiber over $L = \infty$ will give the coefficients (6) and (7) of the image of $\check{m}c(\xi')$. Finally, the count coming from the other fibers will contribute a boundary term (see Proposition 32 for the precise statement). At the level of homology, this means that $\widetilde{\mathrm{HM}}_\bullet(W^\dagger, \mathfrak{s}_\omega) \, c(\xi') = c(\xi', Y)$ so at this point the naturality proof has been reduced to




Figure 4: Gluing technique for the “stretching the neck” argument.

showing that $c(\xi', Y) = c(\xi)$. Again, from the chain level perspective this means that up to boundary terms, for each critical point $[c]$ the numbers $\sum_z m_z(W_{\xi', Y}^+, \mathfrak{s}_\omega, [c])$ must equal $\sum_{z'} m_{z'}(Z_{Y, \xi}^+, \mathfrak{s}, [c])$.

If one were to replace the half-cylindrical end $\mathbb{R}^+ \times -Y$ with a compact piece X so that we could work with numbers instead of homology classes, the previous quantities would be the same by Theorem D in [35] (ie (2) in our paper). Therefore, it becomes clear at this point that what we need to do is adapt the Mrowka–Rollin theorem to the case in which we have a half-infinite cylinder.

Two things that change in this new setup are that certain inclusions of Sobolev spaces are no longer compact, and in order to achieve transversality (ie obtain unobstructed moduli spaces in the terminology of [35]) one must use the “abstract perturbations” defined by Kronheimer and Mrowka in [22]. In particular, these perturbations introduce new terms that do not appear in the usual linearizations of the Seiberg–Witten equations, so for the gluing argument we will employ one needs to check that the new contributions do not mess up the desired behavior of the linearized Seiberg–Witten equations. Namely, we will see that the contributions have leading terms which are quadratic in an appropriate sense. Had the leading term been linear, the gluing argument would not have worked.

Our gluing argument and the proof of [35, Theorem D] morally follow the same basic ideas as the other gluing arguments in gauge theory but as expected differ in the specific details (a few references include [22; 37; 42; 7; 34; 13; 14]). Perhaps the most common gluing argument in gauge theory is the one involving the “stretching the neck” operation on a closed oriented Riemannian four-manifold X which has a separating hypersurface Y inside it.




Figure 5: “Dilating the cone” argument used to show that $c(\xi', Y) = c(\xi)$.

Namely, one writes X as $X = X_1 \cup X_2$ and after choosing a metric which is cylindrical near Y one can stretch the metric along Y in order to have a cylinder $I_L \times Y$ of length L inserted between X_1 and X_2 as shown in Figure 4. The point is that as L increases, the Seiberg–Witten equations on $X_L = X_1 \cup (I_L \times Y) \cup X_2$ start behaving more like the solutions on the manifolds with cylindrical ends X_1^* and X_2^* . More precisely, one can start from solutions on X_1^* and X_2^* which agree on their respective ends in order to construct a presolution on X_L , that is, a configuration on X_L which is a solution to the Seiberg–Witten equations on X_L , except perhaps for a region supported on $I_L \times Y$. The main point of the gluing argument is that one can find an L_0 large enough that for all L bigger than L_0 we can obtain an actual solution to the Seiberg–Witten equations on X_L thanks to an application of the implicit function theorem for Banach spaces. In order for this to work it is imperative to have estimates that become independent of L .

Likewise, in our situation we want to take advantage of the fact that for a strong symplectic cobordism the symplectic structure is given near the boundary by the symplectization of the contact structure, so that in analogy with the cylindrical case we can perform a “dilating the cone” operation, where now the key parameter is a dilation parameter τ which determines the size of the cone C_τ determined by the symplectization of the contact structure near the boundary. As in the cylindrical case, the main idea is that once τ is sufficiently large, the moduli space of solutions to the Seiberg–Witten equations on the manifold in Figure 5 can be described in terms of the moduli space used to define the contact invariant of (Y, ξ) . Again, this will rely on an application of the implicit function theorem, which requires guaranteeing that certain estimates become independent of τ (once it becomes sufficiently large). As we will explain near the end of the paper this gluing theorem will establish that $c(\xi', Y) = c(\xi)$.

3 Setting up the equations

As explained in the previous section, we will analyze first the equations on $W_{\xi', Y}^+$. In particular, we begin by stating some basic geometric properties of the manifolds we are going to be working with.

Assumption 11 *Suppose we have a closed oriented three-manifold Y with contact structure ξ . We write $\xi = \ker \theta$ and choose the unique Riemannian metric g_θ such that [21, Section 2.3]:*

- *The contact form θ has unit length.*
- *$d\theta = 2 *_Y \theta$, where $*_Y$ is the Hodge star on (Y, g_θ) .*
- *If J is a choice of an almost complex structure on ξ then for any $v, w \in \xi$, $g_\theta(v, w) = d\theta(v, Jw)$.*

The contact structure ξ determines a canonical spin-c structure $\mathfrak{s}_\xi$: define the spinor bundle S as the rank-2 vector bundle $S = \underline{\mathbb{C}} \oplus \xi$, where $\underline{\mathbb{C}}$ is the trivial vector bundle and we are considering ξ as a complex line bundle. Moreover, there is a Clifford map $\rho_Y: TY \rightarrow \text{hom}(S, S)$ which identifies TY isometrically with the subbundle $\mathfrak{su}(S)$ of traceless, skew-adjoint endomorphisms equipped with the inner product $\frac{1}{2} \text{tr}(a^*b)$ [22, Section 1.1]. Using $(Y, g_\theta, \mathfrak{s}_\xi)$ we can write the configuration space on which the Seiberg–Witten equations are defined [22, Section 9.1]: for any integer or half-integer $k \geq 0$ define

$$\mathcal{C}_k(Y, \mathfrak{s}_\xi) = (B_{\text{ref}}, 0) + L_k^2(M; iT^*Y \oplus S) = \mathcal{A}_k(Y, \mathfrak{s}_\xi) \times L_k^2(Y; S),$$

where B_{ref} is a reference smooth connection on the spinor bundle S compatible with the Levi-Civita connection defined on TY and $\mathcal{A}_k(Y, \mathfrak{s}_\xi)$ denotes the (affine) space of spin-c connections of S with Sobolev regularity L_k^2 . We will always assume whenever needed that $k \geq 5$, but by elliptic regularity the constructions end up being independent of k because one can always find a smooth representative in each gauge-equivalence class of solutions to the Seiberg–Witten equations, so we will not dwell a lot on the actual value of k being used.

The gauge group $\mathcal{G}_{k+1}(Y)$ is

$$\mathcal{G}_{k+1}(Y) = \{u \in L_{k+1}^2(Y; \mathbb{C}) : |u| = 1 \text{ pointwise}\}.$$

It acts on the configuration space via

$$u \cdot (B, \Psi) = (B - u^{-1} du, u\Psi).$$

The action is not free at the *reducible configurations*, that is, the configurations $(B, 0)$ with the spinor component identically zero. The stabilizer at those configurations consists of the constant maps $u: Y \rightarrow S^1$ which we can identify with S^1 . To handle reducible configurations Kronheimer and Mrowka [22, Section 6.1] introduced the *blown-up configuration space*

$$\mathcal{C}_k^\sigma(Y, \mathfrak{s}_\xi) = \{(B, s, \phi) : \|\phi\|_{L^2(Y)}=1, s \geq 0\} = \mathcal{A}_k(Y, \mathfrak{s}_\xi) \times \mathbb{R}^{\geq} \times \mathbb{S}(L_k^2(Y; S)).$$

Here $\mathbb{S}(L_k^2(Y; S))$ denotes those elements ϕ in $L_k^2(Y; S)$ whose L^2 norm (not L_k^2 norm!) is equal to 1. In this case the gauge action is

$$u \cdot (B, s, \phi) = (B - u^{-1} du, s, u\phi),$$

and it is easy to check that the gauge group acts freely on this space. In fact, Lemma 9.1.1 in [22] shows that the space $\mathcal{C}_k^\sigma(Y, \mathfrak{s}_\xi)$ is naturally a Hilbert manifold with boundary and when $k \geq 1$, the space $\mathcal{G}_{k+1}(Y)$ is a Hilbert Lie group which acts smoothly and freely on $\mathcal{C}_k^\sigma(Y, \mathfrak{s}_\xi)$.

We are interested in triples (B, s, ϕ) which satisfy a perturbed version of the Seiberg–Witten equations. At this point the nature of the perturbations is not that important. For now it suffices to say that we will take them to be *strongly tame perturbations* [53, Definition 3.6]. As a technical point it is useful to note that the cylindrical functions constructed in [22, Section 11.1] are strongly tame perturbations, so the theorems from there which used this class of perturbations continue to work in this context. We will denote such a perturbation by $q_{Y, g_\theta, \mathfrak{s}_\xi}$. In general a strongly tame perturbation q can be regarded as a map $q: \mathcal{C}_k(Y, \mathfrak{s}_\xi) \rightarrow L_k^2(Y; iT^*Y \oplus S)$, where one thinks of the codomain as a copy of the tangent space $T_{(B, \psi)}\mathcal{C}_k(Y, \mathfrak{s}_\xi)$ for each configuration $(B, \psi) \in \mathcal{C}_k(Y, \mathfrak{s}_\xi)$. Since the codomain naturally splits one can write $q = (q^0, q^1)$ and in [22, Section 10.2] it is explained how q gives rise to a perturbation on the blown-up configuration space $q^\sigma = (q^0, \hat{q}^{1, \sigma})$ (notice that only the second component is modified).

The corresponding equations a triple (B, s, ϕ) satisfies are [22, Section 10.3]

(10)

$$\begin{aligned} \frac{1}{2} * F_{B^t} + s^2 \rho_Y^{-1} (\phi \phi^*)_0 + q_{Y, g_\theta, \mathfrak{s}_\xi}^0(B, s \phi) &= 0, \\ \Lambda_{q_{Y, g_\theta, \mathfrak{s}_\xi}}(B, s, \phi) s &= 0, \\ D_B \phi - \Lambda_{q_{Y, g_\theta, \mathfrak{s}_\xi}}(B, s, \phi) \phi + \tilde{q}_{Y, g_\theta, \mathfrak{s}_\xi}^1(B, s, \phi) &= 0, \end{aligned}$$

where:

- F_{B^t} denotes the curvature of the connection B^t on $\det(S)$.

- $(\phi\phi^*)_0$ denotes the trace-free part of the hermitian endomorphism $\phi\phi^*$, that is, $(\phi\phi^*)_0 = \phi\phi^* - \frac{1}{2}|\phi|^2 1_S$.
- D_B is the Dirac operator corresponding to the connection B .
- $\Lambda_{q_{Y,g_\theta,\mathfrak{s}_\xi}}(B, s, \phi) = \text{Re}\langle \phi, D_B\phi + \tilde{q}_{Y,g_\theta,\mathfrak{s}_\xi}^1(B, s, \phi) \rangle_{L^2(Y)}$ and $\tilde{q}^1(B, r, \psi) = \int_0^1 \mathcal{D}_{(B, sr\psi)} q^1(0, \psi) ds$ (here $\mathcal{D}$ denotes the linearization of the map q^1).

Using the equations (10) we can distinguish three types of solutions (or critical points) $\mathfrak{c} = (B, s, \phi)$ [23, Definition 4.4], the *irreducible critical point*, the *boundary stable reducible critical point* and the *boundary unstable reducible critical point*. What is important about this classification for us is that solutions of the four-dimensional Seiberg–Witten equations on $\mathbb{R} \times Y$ for which the spinor does not vanish identically can only be asymptotic as $t \rightarrow \infty$ to irreducible critical points or boundary stable reducible critical points. The gauge-equivalence class of any of these points will be denoted as $[c]$.

The triple $(Y, g_\theta, \mathfrak{s}_\xi)$ induces a spin-c structure on $(-Y, g_\theta)$ given by the same spinor bundle S_ξ and changing the Clifford multiplication from ρ_ξ to $-\rho_\xi$ [22, Section 22.5]. We will continue to denote this spin-c structure by $\mathfrak{s}_\xi$. Given this structure we can use the cylindrical metric and the spin-c structure induced by $-Y$ on the cylinder $\mathbb{R}^+ \times -Y$ [22, Section 4.3]. We use the perturbation $-q_{Y,g_\theta,\mathfrak{s}_\xi}$ on $-Y$.

Consider now the manifold

$$W_{\xi',Y}^+ = ([1, \infty) \times Y') \cup W^\dagger \cup (\mathbb{R}^+ \times -Y).$$

We will define the appropriate geometric structures needed on each piece together with the perturbations we will be using.

- On $\mathbb{R}^+ \times -Y$, we use the cylindrical metric and the canonical spin-c structure induced by $\mathfrak{s}_\xi$ on the cylinder. As explained in [22, Section 10.1], we have a four-dimensional perturbation

$$-\hat{q}_{Y,g_\theta,\mathfrak{s}_\xi}: \mathcal{C}_k(\mathbb{R}^+ \times -Y, \mathfrak{s}_\xi) \rightarrow L_k^2(\mathbb{R}^+ \times -Y; iT^*(\mathbb{R}^+ \times -Y) \oplus S)$$

on the half-cylinder $\mathbb{R}^+ \times -Y$, defined by restriction to each slice.

- On $W^\dagger$ we choose a metric g_W that is cylindrical in collar neighborhoods of the boundary components. To define the perturbation on $W^\dagger$ we follow [22, Section 24.1]. Since the Riemannian metric is cylindrical in the neighborhood of the boundary it contains on each boundary component an isometric copy of $I_1 \times -Y$ and $I_2 \times Y'$,

where $I_1 = (-C_1, 0]$ and $I_2 = (-C_2, 0]$. Since the argument is the same for both ends we will use generic notation. Let β be a cutoff function, equal to 1 near $t = 0$ and equal to 0 near $t = -C$. Let β_0 be a bump function with compact support in $(-C, 0)$, equal to one on a compact subset inside $(-C, 0)$, for example, the compact subset $[-\frac{1}{2}C, -\frac{1}{4}C]$. Choose another perturbation $\mathfrak{p}_0$ of the three-dimensional equations and consider the perturbation

$$\widehat{\mathfrak{p}}_W = \beta \widehat{\mathfrak{q}} + \beta_0 \widehat{\mathfrak{p}}_0.$$

It is useful to note that the reason why we use two perturbations is so that one can be varied when we use a transversality argument.

- On $[1, \infty) \times Y'$ we assume that the metric is cylindrical in a collar neighborhood $[1, C_K) \times Y'$ and on a complement of this neighborhood (like $N_K = [C_K + 1, \infty) \times Y'$ for instance) it is given by the metric

$$g_{K,\theta'} = dt \otimes dt + t^2 g_{\theta'}$$

with symplectic form

$$\omega_{\theta'} = \frac{1}{2} d(t^2 \theta').$$

Here K stands for Kähler, although in most cases the cone will not be a Kähler manifold (in fact this occurs only when (Y, ξ) is a Sasakian manifold [2]). The form is self-dual with respect to $g_{K,\theta'}$ and $|\omega_{\theta'}|_{g_{K,\theta'}} = \sqrt{2}$ pointwise. By [21, Lemma 2.1], on the symplectic cone we have a unit-length section Φ_0 associated to the canonical spinor bundle $S_{\omega_{\theta'}}$. For this section Φ_0 we have a corresponding connection A_0 such that $D_{A_0} \Phi_0 = 0$. Choose a smooth extension of (A_0, Φ_0) to all of $W_{\xi',Y}^+$ in such a way that (A_0, Φ_0) is translation invariant on the cylindrical end $\mathbb{R}^+ \times -Y$. Define

$$\mathfrak{p}_K = (-\tfrac{1}{2} \rho(F_{A_0'}^+) + (\Phi_0 \Phi_0^*)_0, 0),$$

and choose a cutoff function β_K which is supported on N_K and identically equal to 1 on $[C_K + 2, \infty)$. Choose also a cutoff function β_{N_K} which is supported on $[1, C_K) \times Y'$ and identically equal to 1 near the boundary $\partial([1, C_K) \times Y')$.

Our global perturbation will be

(11)
$$\begin{aligned} \mathfrak{p}_{W_{\xi',Y}^+} = & -\widehat{\mathfrak{q}}_{Y,g_{\theta},\mathfrak{s}_{\xi}} + (\beta \widehat{\mathfrak{q}}_{Y,g_{\theta},\mathfrak{s}_{\xi}} + \beta'_0 \widehat{\mathfrak{p}}_0) + (\beta'_0 \widehat{\mathfrak{p}}_0 + \beta' \widehat{\mathfrak{q}}_{Y',g_{\theta'},\mathfrak{s}_{\xi'}}) \\ & + (\beta_{N_K} \widehat{\mathfrak{q}}_{Y',g_{\theta'},\mathfrak{s}_{\xi'}} + \beta_K \mathfrak{p}_K), \end{aligned}$$

where β'_0 and β' are cutoff functions defined analogously for the other cylindrical neighborhood $I_2 \times Y'$.

In words the previous perturbation behaves as follows: If we start on the cylindrical end $\mathbb{R}^+ \times -Y$ we will see the translation-invariant perturbation $-\hat{q}_{Y,g_\theta,s_\xi}$. As we enter the cobordism through the boundary $-Y \subset W^\dagger$ (recall that $\partial W^\dagger = -Y \sqcup Y'$) this perturbation is modified into a combined perturbation $\beta \hat{q}_{Y,g_\theta,s_\xi} + \beta'_0 \hat{p}_0$, which is supported on a collar neighborhood of this end. After we exit this collar neighborhood we will see no perturbations until we reach again the collar neighborhood of the end $Y' \subset W^\dagger$, where the perturbation is $\beta'_0 \hat{p}'_0 + \beta' \hat{q}_{Y',g_{\theta'},s_{\xi'}}$. Finally, as we exit the cobordism we will see a perturbation identically equal to $\hat{q}_{Y',g_{\theta'},s_{\xi'}}$ for a small time until it becomes zero again and then it will eventually be changed into the perturbation identically equal to p_K . We will explain the reason why the perturbations were chosen in this way near the end of this section.

Now we must define the corresponding configuration space that we want to use in order to analyze the Seiberg–Witten equations. In general one needs to define the ordinary configuration space and its blow-up (see [22, Sections 13 and 24.2] for some motivation behind this construction). Because of the asymptotic condition we will impose, our solutions will always be irreducible so the gauge group action will be free without having to blow up the configuration space. Therefore, most of the time we will simply use the ordinary configuration space. However, if one wants to describe the compactification of the moduli spaces in terms of the space of broken trajectories then the blow-up model is more convenient so for completeness sake we will write the equations in the blow-up model (but we will switch to the ordinary configuration space when some computations become more transparent there).

We are interested in the configurations that solve the following perturbed version of the Seiberg–Witten equations:

(12)

$$\mathfrak{F}_p = \mathfrak{F} + p_{W_{\xi',Y}^+} = 0,$$

where the *unperturbed Seiberg–Witten map* is [22, Equation 4.12]

$$\mathfrak{F}(A, \Phi) = \left(\tfrac{1}{2}\rho(F_{A'}^+) - (\Phi\Phi^*)_0, D_A\Phi\right).$$

Both the perturbed and unperturbed maps are defined on elements of the following configuration space [53, Definition 3.5; 22, Definition 13.1]:

Definition 12 Define the *configuration space* (without blow-up) $\mathcal{C}_{k,\text{loc}}(W_{\xi',Y}^+, \mathfrak{s}_\omega)$ as follows. It will consist of pairs (A, Φ) such that:

- (1) A is a locally L^2_k spin-c connection for S and Φ is a locally L^2_k section of S^+ .
- (2) (A, Φ) is L^2_k close to the canonical solution on the conical end, that is,

$$A - A_0 \in L^2_k([1, \infty) \times Y', i T^*([1, \infty) \times Y')),$$
$$\Phi - \Phi_0 \in L^2_{k, A_0}([1, \infty) \times Y', S^+).$$

- Remark 13** (a) Recall that we chose an extension of A_0 to the cylindrical end in such a way that it was translation invariant so the condition that A is a locally L^2_k spin-c connection means that $A - A_0 \in L^2_{k, \text{loc}}(W^{+}_{\xi', Y}; i T^* W^{+}_{\xi', Y})$.
- (b) Notice that the second condition implies that Φ cannot be identically 0, that is, $\mathcal{C}_{k, \text{loc}}(W^{+}_{\xi', Y}, \mathfrak{s})$ contains no reducible configurations. In the notation of [22], we would write $\mathcal{C}_{k, \text{loc}}(W^{+}_{\xi', Y}, \mathfrak{s}) = \mathcal{C}^*_{k, \text{loc}}(W^{+}_{\xi', Y}, \mathfrak{s})$.
- (c) Because of the lack of a norm, the space $\mathcal{C}_{k, \text{loc}}(W^{+}_{\xi', Y}, \mathfrak{s})$ is not a Banach space unless we impose some asymptotic condition on the cylindrical end.

The *blown-up configuration space* $\mathcal{C}^{\sigma}_{k, \text{loc}}(W^{+}_{\xi', Y}, \mathfrak{s}_{\omega})$ is defined as follows:

Definition 14 If S denotes the spinor bundle, define the sphere $\mathbb{S}$ as the topological quotient of $L^2_{k, \text{loc}}(W^{+}_{\xi', Y}; S^+) \setminus 0$ by the action of $\mathbb{R}^+$ [22, Section 6.1]. The blown-up configuration space associated to $\mathcal{C}_{k, \text{loc}}(W^{+}_{\xi', Y}, \mathfrak{s}_{\omega})$ is

$$\mathcal{C}^{\sigma}_{k, \text{loc}}(W^{+}_{\xi', Y}, \mathfrak{s}_{\omega}) = \{(A, \mathbb{R}^+ \phi, \Phi) : \Phi \in \mathbb{R}^{\geq 0} \phi, \phi \in \mathbb{S} \text{ and } (A, \Phi) \in \mathcal{C}_{k, \text{loc}}(W^{+}_{\xi', Y}, \mathfrak{s}_{\omega})\}.$$

Just as its blown-down version, $\mathcal{C}^{\sigma}_{k, \text{loc}}(W^{+}_{\xi', Y}, \mathfrak{s}_{\omega})$ is not a Banach manifold, much less a Hilbert manifold, so we will not try to find useful slices on this space. These slices would have been “orthogonal” in some suitable sense to the gauge group action, which we will take to be

$$(13) \quad \mathcal{G}_{k+1}(W^{+}_{\xi', Y}) = \{u : W^{+}_{\xi', Y} \rightarrow \mathbb{C}^* : |u| = 1 \text{ and } 1 - u \in L^2_{k+1}([1, \infty) \times Y')\},$$

where the action of $u \in \mathcal{G}_{k+1}$ on a triple $(A, \mathbb{R}^+ \phi, \Phi) \in \mathcal{C}^{\sigma}_{k, \text{loc}}(W^{+}_{\xi', Y}, \mathfrak{s}_{\omega})$ is given by

$$(14) \quad u \cdot (A, \mathbb{R}^+ \phi, \Phi) = (A - u^{-1} du, \mathbb{R}^+(u\phi), u\Phi).$$

Using the Sobolev multiplication theorems on manifolds with bounded geometry [9, Chapter 1] it is not difficult to verify that $\mathcal{G}_{k+1}(W^{+}_{\xi', Y})$ is a Hilbert Lie group and that the previous formula indeed gives an action on the configuration space $\mathcal{C}_{k, \text{loc}}(W^{+}_{\xi', Y}, \mathfrak{s}_{\omega})$, that is:

Lemma 15 Suppose that $k \geq 4$. Then $\mathcal{G}_{k+1}(W_{\xi', Y}^+)$ is a Hilbert Lie group. Moreover, the action of $\mathcal{G}_{k+1}(W_{\xi', Y}^+)$ on $\mathcal{C}_{k, \text{loc}}(W_{\xi', Y}^+, \mathfrak{s}_\omega)$ is well defined in that:

- (i) If $(A, \Phi) \in \mathcal{C}_k(W_{\xi', Y}^+, \mathfrak{s}_\omega)$ and $u \in \mathcal{G}_{k+1}(W_{\xi', Y}^+)$ then $u \cdot (A, \Phi) \in \mathcal{C}_k(W_{\xi', Y}^+, \mathfrak{s}_\omega)$.
- (ii) Similarly, if $u \cdot (A, \Phi) = (\tilde{A}, \tilde{\Phi})$ for two configurations $(A, \Phi), (\tilde{A}, \tilde{\Phi}) \in \mathcal{C}_k(W_{\xi', Y}^+, \mathfrak{s}_\omega)$ and u is an $L_{k+1, \text{loc}}^2(W_{\xi', Y}^+)$ gauge transformation, then $1 - u \in L_{k+1}^2([1, \infty) \times Y')$.

Remark 16 A proof of this lemma can be found in the author's thesis [8, Lemma 17].

Therefore it makes sense to define

$$\mathcal{B}_{k, \text{loc}}^\sigma(W_{\xi', Y}^+, \mathfrak{s}_\omega) = \mathcal{C}_{k, \text{loc}}^\sigma(W_{\xi', Y}^+, \mathfrak{s}_\omega) / \mathcal{G}_{k+1}(W_{\xi', Y}^+).$$

Again, since the original space $\mathcal{C}_{k, \text{loc}}^\sigma(W_{\xi', Y}^+, \mathfrak{s}_\omega)$ is not a Banach manifold, we will not be interested in studying directly $\mathcal{B}_{k, \text{loc}}^\sigma(W_{\xi', Y}^+, \mathfrak{s}_\omega)$, although this is the space where the solutions to the Seiberg–Witten equations live.

To define the moduli space to the Seiberg–Witten equations, we need to introduce the τ model first. Let

$$[\mathfrak{c}] \in \mathfrak{C}^o(-Y, g_\theta, \mathfrak{s}_\xi, -\mathfrak{q}_{Y, g_\theta, \mathfrak{s}_\xi}) \cup \mathfrak{C}^s(-Y, g_\theta, \mathfrak{s}_\xi, -\mathfrak{q}_{Y, g_\theta, \mathfrak{s}_\xi})$$

be a critical point [22, Proposition 12.2.5] to the blown-up three-dimensional Seiberg–Witten equations on $-Y$ (10). Write $[\mathfrak{c}] = [(B, s, \phi)]$ and let $\mathfrak{c} = (B, s, \phi)$ be a smooth representative in $\mathcal{C}_k^\sigma(-Y, \mathfrak{s}_\xi)$. The critical point $\mathfrak{c}$ gives rise to a translation-invariant configuration $\gamma_{\mathfrak{c}}$ on the half-infinite cylinder $\mathbb{R}^+ \times -Y$.

Definition 17 Define on $\mathbb{R}^+ \times -Y$ the τ model $\mathcal{C}_{k, \text{loc}}^\tau(\mathbb{R}^+ \times -Y, \mathfrak{s}_\xi, \mathfrak{c})$ associated to $\mathfrak{c}$ as the space of triples [22, Section 13.3]

$$\gamma = (A, r(t), \phi(t)) \in \mathcal{A}_{k, \text{loc}}(\mathbb{R}^+ \times -Y, \mathfrak{s}_\xi) \times L_{k, \text{loc}}^2(\mathbb{R}^+; \mathbb{R}) \times L_{k, \text{loc}}^2(\mathbb{R}^+ \times -Y; S^+)$$

such that:

- (i) $\gamma - \gamma_{\mathfrak{c}} \in L_{k, \text{loc}}^2(iT^*(\mathbb{R}^+ \times -Y)) \times L_{k, \text{loc}}^2(\mathbb{R}^+; \mathbb{R}) \times L_{k, \text{loc}}^2(\mathbb{R}^+ \times -Y; S^+)$, ie γ is $L_{k, \text{loc}}^2$ close to $\gamma_{\mathfrak{c}}$.
- (ii) For all $t \in \mathbb{R}^+$, we have that $r(t) \geq 0$.
- (iii) For all $t \in \mathbb{R}^+$, we have that $\|\phi(t)\|_{L^2(-Y)} = 1$, ie on each slice the L^2 norm (not the L_k^2 norm) is 1.

There is a natural restriction of the gauge group $\mathcal{G}_{k+1}(W_{\xi',Y}^+)$ to $\mathbb{R}^+ \times -Y$ which acts on $\mathcal{C}_{k,\text{loc}}^\tau(\mathbb{R}^+ \times -Y, \mathfrak{s}_\xi, \mathfrak{c})$ via

$$u \cdot (A, r(t), \phi(t)) = (A - u^{-1} du, r(t), u\phi(t)).$$

The gauge-equivalence classes of configurations under this gauge group action will be denoted as

$$\mathcal{B}_{k,\text{loc}}^\tau(\mathbb{R}^+ \times -Y, \mathfrak{s}_\xi, [\mathfrak{c}]) = \mathcal{C}_{k,\text{loc}}^\tau(\mathbb{R}^+ \times -Y, \mathfrak{s}_\xi, \mathfrak{c})/\mathcal{G}_{k+1,\text{loc}}(\mathbb{R}^+ \times -Y).$$

We will also use the *unique continuation principle*, which will essentially allow us for the most part to avoid working with the blow-up model. The versions most convenient to us are [22, Propositions 7.1.4 and 10.8.1], which can still be used in our context because the perturbation $\mathfrak{p}_K$ used on the conical (symplectic) end involves no spinor component.

These imply that if a solution of the perturbed Dirac equation vanishes on a slice $\{t\} \times -Y$ of the cylindrical end $\mathbb{R}^+ \times -Y$, then it would have to vanish on the entire half-cylinder $\mathbb{R}^+ \times -Y$ and then on the entire four-manifold $W_{\xi',Y}^+$. However, since we will be interested in solutions which are asymptotic on the conical end to the spinor Φ_0 (which is nonvanishing), this cannot be the case so we can safely conclude that no such solutions will exist, that is, our spinor Φ will never vanish on an open set or a cylindrical slice. Thanks to this, the following definition makes sense (compare with [22, Definition 24.2.1]):

Definition 18 The moduli space $\mathcal{M}(W_{\xi',Y}^+, \mathfrak{s}_\omega, [\mathfrak{c}])$ for a critical point

$$[\mathfrak{c}] \in \mathfrak{C}^o(-Y, g_\theta, \mathfrak{s}_\xi, -\mathfrak{q}_{Y,g_\theta,\mathfrak{s}_\xi}) \cup \mathfrak{C}^s(-Y, g_\theta, \mathfrak{s}_\xi, -\mathfrak{q}_{Y,g_\theta,\mathfrak{s}_\xi})$$

consists of gauge-equivalence classes of triples

$$[A, \mathbb{R}^+ \phi, \Phi] \in \mathcal{B}_{k,\text{loc}}^\sigma(W_{\xi',Y}^+, \mathfrak{s}_\omega)$$

such that:

- (1) $(A, \mathbb{R}^+ \phi, \Phi) \in \mathcal{C}_{k,\text{loc}}^\sigma(W_{\xi',Y}^+, \mathfrak{s}_\omega)$ and (A, Φ) satisfies the perturbed Seiberg–Witten equations $\mathfrak{F}_\mathfrak{p}(A, \Phi) = 0$ on $W_{\xi',Y}^+$. Here $\mathfrak{p}$ refers to the perturbation explained before (12).
- (2) Because of the unique continuation principle, Φ cannot be identically zero on each of the cylindrical slices. Thus, for each t , we can define [22, Sections 6.2 and 13.1]

$$(r(t), \psi(t)) = \left(\|\check{\Phi}(t)\|_{L^2(-Y)}, \frac{\check{\Phi}(t)}{\|\check{\Phi}(t)\|_{L^2(-Y)}} \right).$$

Also, if we decompose the covariant derivative ∇_A in the $\frac{d}{dt}$ direction as

$$\nabla_{A,\frac{d}{dt}} = \frac{d}{dt} + a_t \otimes 1_S,$$

we require that $\gamma = (A, r(t), \psi(t))$ be an element of $C_{k,\text{loc}}^\tau(\mathbb{R}^+ \times -Y, \mathfrak{s}_\xi, \mathfrak{c})$ and that it solve the following Seiberg–Witten equations on the cylinder [22, Equation 10.9]:

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \check{A}^t &= -\frac{1}{2} *_Y F_{\check{A}^t} + da_t - r^2 \rho^{-1} (\psi \psi^*)_0 - \mathfrak{q}^0(\check{A}, r \psi), \\ \frac{d}{dt} r &= -\Lambda_{\mathfrak{q}}(\check{A}, r, \psi) r, \\ \frac{d}{dt} \psi &= -D_{\check{A}} \psi - a_t \psi - \tilde{\mathfrak{q}}^1(\check{A}, r, \psi) + \Lambda_{\mathfrak{q}}(\check{A}, r, \psi) \psi, \end{aligned}$$

where $\check{A}(t)$ denotes the restriction of A to the t slice. Moreover, we require that the gauge-equivalence class $[\gamma]$ of γ be asymptotic as $t \rightarrow \infty$ to $[\mathfrak{c}]$ in the sense of [22, Definition 13.1.1].

The moduli space $\mathcal{M}(W_{\xi',Y}^+, \mathfrak{s}_\omega, [\mathfrak{c}])$ is naturally a subset of $\mathcal{B}_{k,\text{loc}}^\sigma(W_{\xi',Y}^+, \mathfrak{s}_\omega)$. However, since the latter space is not in any natural way a Hilbert manifold we will use a fiber product description of $\mathcal{M}(W_{\xi',Y}^+, \mathfrak{s}_\omega, [\mathfrak{c}])$ instead [22, Lemmas 24.2.2 and 19.1.1]. The idea is that we can “break” the moduli space $\mathcal{M}(W_{\xi',Y}^+, \mathfrak{s}_\omega, [\mathfrak{c}])$ into three moduli spaces which we will show are Hilbert manifolds. These moduli spaces are the moduli space on the cobordism $\mathcal{M}(W^\dagger, \mathfrak{s}_\omega)$, the moduli space on the half-cylinder $\mathcal{M}^\tau(\mathbb{R}^+ \times -Y, \mathfrak{s}_\xi)$ and the moduli space on the conical end $\mathcal{M}([1, \infty) \times Y', \mathfrak{s}')$. We included a superscript τ for the second one to emphasize that its definition uses the τ model, which is described in Definition 17. The fiber product description will then also allows us to show that $\mathcal{M}(W_{\xi',Y}^+, \mathfrak{s}_\omega, [\mathfrak{c}])$ has a Hilbert manifold structure but in order to explain this we first begin with a lemma.

Lemma 19 *The moduli spaces $\mathcal{M}(W^\dagger, \mathfrak{s}_\omega)$ and $\mathcal{M}^\tau(\mathbb{R}^+ \times -Y, \mathfrak{s}_\xi)$ are Hilbert manifolds.*

Remark 20 Sections 4.1, 4.2 and 4.3 of the author’s thesis [8] give more details on the construction of these manifolds and the ideas behind the proof of this lemma.

Proof The piece corresponding to the moduli space $\mathcal{M}(W^\dagger, \mathfrak{s}_\omega)$ is described in [22, Proposition 24.3.1], where it is shown to be a Hilbert manifold.

Likewise, the moduli space $\mathcal{M}^\tau(\mathbb{R}^+ \times -Y, \mathfrak{s}_\xi)$ is described in [22, Theorem 14.4.2], where it is shown to be a Hilbert manifold (strictly speaking the authors analyzed an

entire cylinder $\mathbb{R} \times Y$ rather than a half-cylinder $\mathbb{R}^+ \times -Y$ but the analysis is essentially the same). □

We will start the next section showing that $\mathcal{M}([1, \infty) \times Y', \mathfrak{s}')$ is a Hilbert manifold, following the arguments in [22, Section 24]. At the end of the day, we obtain restriction (or trace) maps

$$\begin{aligned} R_\tau &\colon \mathcal{M}(\mathbb{R}^+ \times -Y, \mathfrak{s}_\xi, [\mathfrak{c}]) \rightarrow \mathcal{B}^\sigma_{k-\frac{1}{2}}(Y, \mathfrak{s}_\xi), \\ R^-_{\overline{W}} &\colon \mathcal{M}(W^\dagger, \mathfrak{s}_\omega) \rightarrow \mathcal{B}^\sigma_{k-\frac{1}{2}}(-Y, \mathfrak{s}_\xi), \\ R^+_{\overline{W}} &\colon \mathcal{M}(W^\dagger, \mathfrak{s}_\omega) \rightarrow \mathcal{B}^\sigma_{k-\frac{1}{2}}(Y', \mathfrak{s}_{\xi'}), \\ R_K &\colon \mathcal{M}([1, \infty) \times Y', \mathfrak{s}') \rightarrow \mathcal{B}^\sigma_{k-\frac{1}{2}}(-Y', \mathfrak{s}_\xi) \end{aligned}$$

given by restricting the (gauge-equivalence class of a) solution to the boundary of each of the corresponding manifolds. We should point out that there is an identification between $\mathcal{B}^\sigma_k(-Y, \mathfrak{s})$ and $\mathcal{B}^\sigma_k(Y, \mathfrak{s})$ [22, Section 22.5] and we can identify $\mathcal{M}(W_{\xi', Y}^+, \mathfrak{s}, [\mathfrak{c}])$ with the fiber product $\text{Fib}(R_\tau, R^-_{\overline{W}}, R^+_{\overline{W}}, R_K)$ given by

(15) $\{([\gamma_{\mathbb{R}^+ \times -Y}], [\gamma_W], [\gamma_{[1, \infty) \times Y'}]) \colon R_\tau[\gamma_{\mathbb{R}^+ \times -Y}] = R^-_{\overline{W}}[\gamma_W]$
 $\text{and } R^+_{\overline{W}}[\gamma_W] = R_K[\gamma_{[1, \infty) \times Y'}])\}.$

Now we can explain how to give $\mathcal{M}(W_{\xi', Y}^+, \mathfrak{s}, [\mathfrak{c}])$ a Hilbert manifold structure, which is stated precisely in Definition 26 and Theorem 27 of the next section.

For convenience write $R = (R_\tau, R^-_{\overline{W}}, R^+_{\overline{W}}, R_K)$ and suppose that

$$[\gamma] = ([\gamma_{\mathbb{R}^+ \times -Y}], [\gamma_W], [\gamma_{[1, \infty) \times Y'}]) \in \text{Fib}(R_\tau, R^-_{\overline{W}}, R^+_{\overline{W}}, R_K)$$

is such that the map R is transverse at $([\mathfrak{b}], [\mathfrak{b}'])$, where

$$[\mathfrak{b}] = R_\tau([\gamma_{\mathbb{R}^+ \times -Y}]) = R^-_{\overline{W}}([\gamma_W]) \quad \text{and} \quad [\mathfrak{b}'] = R^+_{\overline{W}}([\gamma_W]) = R_K([\gamma_{[1, \infty) \times Y'}]).$$

In other words, we want the linearized map $\mathcal{D}_{[\gamma]}R$ to be Fredholm and surjective. If this can be achieved, then near $[\gamma]$ the space $\text{Fib}(R_\tau, R^-_{\overline{W}}, R^+_{\overline{W}}, R_K)$ will have the structure of a smooth manifold of dimension $\dim \ker \mathcal{D}_{[\gamma]}R$. The Fredholm property is proven in Lemma 28 in the next section. The surjectivity of the map $\mathcal{D}_{[\gamma]}R$ may not be true for an arbitrary perturbation of the form described in (11) earlier; however, an application of Sard’s theorem shows that one can choose generic perturbations such that the surjectivity is achieved as well. In fact, achieving the surjectivity is essentially the same as the proof Kronheimer and Mrowka gave for the case of a manifold X^*

with cylindrical ends [22, Proposition 24.4.7]. By choosing a perturbation from this generic set, one can then guarantee that

$$\mathrm{Fib}(R_\tau, R_{\overline{W}}, R_W^+, R_K) = \mathcal{M}(W_{\xi', Y}^+, \mathfrak{s}_\omega, [c])$$

has the structure of a smooth manifold (possibly disconnected with components of different dimensions).

4 Transversality and fiber products

4.1 The moduli space on the conical end $\mathcal{M}([1, \infty) \times Y', \mathfrak{s}')$

We want to regard $\mathcal{M}([1, \infty) \times Y', \mathfrak{s}')$ as a Hilbert submanifold of $\mathcal{B}_k([1, \infty) \times Y', \mathfrak{s}')$, which will be a Hilbert manifold. Write for simplicity $K_{Y'} = [1, \infty) \times Y'$ and define

$$\mathcal{C}_k(K_{Y'}, \mathfrak{s}) = \{(A, \Phi) : A - A_0 \in L_k^2(i T^* K_{Y'}) \text{ and } \Phi - \Phi_0 \in L_{k, A_0}^2(S^+)\}.$$

We take the gauge group to be

$$\mathcal{G}_{k+1}(K_{Y'}) = \{u: K_{Y'} \rightarrow \mathbb{C} : |u| = 1, \ u \in L_{k+1, \mathrm{loc}}^2(K_{Y'}), \ 1 - u \in L_{k+1}^2(K_{Y'})\}.$$

Clearly $\mathcal{C}_k(K_{Y'}, \mathfrak{s}')$ will be a Hilbert manifold because of the L_k^2 asymptotic conditions. It is also easy to see that $\mathcal{G}_{k+1}(K_{Y'})$ will be a Hilbert Lie group. Thus, to show that

$$\mathcal{B}_k(K_{Y'}, \mathfrak{s}') = \mathcal{C}_k(K_{Y'}, \mathfrak{s}')/\mathcal{G}_{k+1}(K_{Y'})$$

is a Hilbert manifold we can use [22, Lemma 9.3.2], which we quote for convenience:

Lemma 21 *Suppose we have a Hilbert Lie group G acting smoothly and freely on a Hilbert manifold C with Hausdorff quotient. Suppose that at each $c \in C$, the map $d_0: T_c G \rightarrow T_c C$ (obtained from the derivative of the action) has closed range. Then the quotient C/G is also a Hilbert manifold.*

The Hilbert manifold structure is given as follows. If $S \subset C$ is any locally closed submanifold containing c satisfying

$$T_c C = \mathrm{im}(d_0) \oplus T_c S,$$

then the restriction of the quotient map $S \rightarrow C/G$ is a diffeomorphism from a neighborhood of c in S to a neighborhood of Gc in C/G . Therefore, we need to verify first that $\mathcal{B}_k(K_{Y'}, \mathfrak{s})$ is a Hausdorff space, which is the content of the next lemma.

Lemma 22 *The quotient space $\mathcal{B}_k(K_{Y'}, \mathfrak{s}')$ is Hausdorff.*

Proof We suppose that we have two gauge equivalent sequences $\gamma_n = (A_n, \Phi_n)$ and $\tilde{\gamma}_n = (\tilde{A}_n, \tilde{\Phi}_n)$ and want to show that the limits $\gamma = (A_\infty, \Phi_\infty)$ and $\tilde{\gamma} = (\tilde{A}_\infty, \tilde{\Phi}_\infty)$ they converge to are gauge equivalent as well. By an exhaustion argument and an application of [22, Proposition 9.3.1] we can find a gauge transformation u_∞ which establishes the gauge equivalence, ie $u_\infty \cdot \gamma = \tilde{\gamma}$. A priori we only have $u_\infty \in L^2_{k+1, \text{loc}}(K_{Y'})$ and to show that $1 - u_\infty \in L^2_{k+1}(K_{Y'})$ we can now apply condition (ii) of Lemma 15. $\square$

As is usually the case for Seiberg–Witten or Yang–Mills moduli spaces, we do not want any random slice to the gauge group action. Rather, we want to use the so-called Coulomb–Neumann slice [22, Section 9.3]. A tangent vector to $\mathcal{C}_k(K_{Y'}, \mathfrak{s}')$ at $\gamma = (A, \Phi)$ can be written as

$$(a, \Psi) \in L^2_k(K_{Y'}, iT^*K_{Y'}) \oplus L^2_{k, A_0}(K_{Y'}, S^+),$$

while the derivative of the gauge group action is

$$(16) \quad \begin{aligned} d_\gamma: L^2_{k+1}(K_{Y'}; i\mathbb{R}) &\rightarrow \mathcal{T}_k = L^2_k(K_{Y'}, iT^*K_{Y'}) \oplus L^2_{k, A_0}(K_{Y'}, S^+), \\ d_{(A, \Phi)}(\zeta) &= (-d\zeta, \zeta\Phi). \end{aligned}$$

We use the inner product

$$\langle (a_1, \Psi_1), (a_2, \Psi_2) \rangle_{L^2} = \int \langle a_1, a_2 \rangle + \text{Re} \langle \Psi_1, \Psi_2 \rangle$$

to define the formal adjoint of $d_{(A, \Phi)}$, which is given by [22, Lemma 9.3.3]

$$(17) \quad d_{(A, \Phi)}^*(a, \Psi) = -d^*a + i \text{Re} \langle i\Phi, \Psi \rangle.$$

To use Lemma 21 we just need to show that d_γ has closed range. To do this we will rely on [21, Theorem 3.3] and [53, Proposition 4.1].

First we need to define a map that we will soon use to show that $\mathcal{M}([1, \infty) \times Y', \mathfrak{s}')$ is a Hilbert manifold (for the proof of Lemma 24). The linearization of the unperturbed Seiberg–Witten map is [21, Equation 8]

$$\begin{aligned} \mathcal{D}_{(A, \Phi)} \mathfrak{F}: L^2_k(K_{Y'}, iT^*K_{Y'}) \oplus L^2_{k, A_0}(K_{Y'}, S^+) \\ \rightarrow L^2_{k-1}(K_{Y'}, i\mathfrak{su}(S^+)) \oplus L^2_{k-1, A_0}(K_{Y'}, S^-), \\ (a, \Psi) \mapsto (\rho(d^+a) - \{\Phi\Psi^* + \Psi\Phi^*\}_0, D_A\Psi + \rho(a)\Phi), \end{aligned}$$

where

$$\{\Phi\Psi^* + \Psi\Phi^*\}_0 = \Phi\Psi^* + \Psi\Phi^* - \frac{1}{2}\langle\Phi, \Psi\rangle - \frac{1}{2}\langle\Psi, \Phi\rangle = \Phi\Psi^* + \Psi\Phi^* - \operatorname{Re}\langle\Phi, \Psi\rangle.$$

Define the elliptic operator (in [53; 35; 21] this is the operator $\mathcal{D}$)

$$(18) \quad Q_{(A,\Phi)} = \mathcal{D}_{(A,\Phi)}\mathfrak{F} \oplus \mathbf{d}_{(A,\Phi)}^*$$

by

$$(a, \Psi) \mapsto (\rho(d^+a) - \{\Phi\Psi^* + \Psi\Phi^*\}_0, D_A\Psi + \rho(a)\Phi, -d^*a + i\operatorname{Re}\langle i\Phi, \Psi\rangle).$$

We also want a formula for the formal adjoint $Q_{(A,\Phi)}^*$: this is essentially Equation 24.10 in [22]. Modulo notational differences, we obtain

$$(19) \quad Q_{(A,\Phi)}^*(\eta, \psi, \vartheta) = ((d^+)^*\rho^*\eta + \rho^*(\psi\Phi^*) - d\vartheta, D_A^*\psi - \eta\Phi + \vartheta\Phi).$$

In particular, taking $\eta = 0$ and $\psi = 0$ one obtains

$$(20) \quad Q_{(A,\Phi)}^*(0, 0, \vartheta) = (-d\vartheta, \vartheta\Phi) = \mathbf{d}_{(A,\Phi)}(\vartheta).$$

Now we are finally ready to prove that $\mathcal{B}_k(K_{Y'}, \mathfrak{s})$ is Hausdorff.

Lemma 23 *Define at a configuration $\gamma = (A, \Phi)$ the subspaces*

$$\mathcal{K}_{k,\gamma} = \{(a, \Psi) : \mathbf{d}_{(A,\Phi)}^*(a, \Psi) = 0 \text{ and } \langle a|_{\partial K_{Y'}}, n \rangle = 0 \text{ at } \partial K_{Y'}\},$$

$$\mathcal{J}_{k,\gamma} = \operatorname{im} \mathbf{d}_\gamma.$$

As γ varies over $\mathcal{C}_k(K_{Y'}, \mathfrak{s})$, the subspaces $\mathcal{J}_{k,\gamma}$ and $\mathcal{K}_{k,\gamma}$ define complementary closed subbundles of $\mathcal{T}_{k,\gamma}$ and we have a smooth decomposition

$$(21) \quad TC_k(K_{Y'}, \mathfrak{s}) = \mathcal{J}_k \oplus \mathcal{K}_k.$$

Proof In order to show the smooth decomposition $TC_k(K_{Y'}, \mathfrak{s}) = \mathcal{J}_k \oplus \mathcal{K}_k$ we can follow the proof of [22, Proposition 9.3.4] and reduce this to the invertibility of the “Laplacian”

$$(22) \quad L_{k+1}^2(K_{Y'}, iT^*K_{Y'}) \ni \vartheta \mapsto \Delta\vartheta + |\Phi|^2\vartheta \in L_{k-1}^2(K_{Y'}, iT^*K_{Y'}).$$

This property can be proved using a parametrix argument (which is essentially the same as Lemma 28 and Theorem 38 in this paper): choosing a compact subset large enough for which $|\Phi|^2$ is not identically zero, one knows from [22, Proposition 9.3.4] that the operator (22) is invertible. On the other hand, [35, Lemma 2.3.2] says that on any four-manifold with conical end (like the manifold X^+ we just used), the operator (22)

is invertible. Notice that their lemma requires a solution to the Seiberg–Witten equations but this is only because in [35, Section 2.3], the authors were trying to find uniform bounds (independent of the solution used). At this stage this is not our concern so the proof they give near the end of that section can be adapted to any configuration. Therefore, we can splice these two inverses to get an approximate inverse to (22) on our domain of interest $K_{Y'}$. By choosing appropriate cutoff functions one can then guarantee that (22) will be invertible (again, the proof of Theorem 38 provides more details). Finally, notice that $\mathcal{J}_k$ is a closed subspace since it is the kernel of a continuous projection $T\mathcal{C}_k \rightarrow \mathcal{J}_k$. $\square$

Continuing with our analysis of our moduli space, we will now show:

Lemma 24 $\mathcal{M}(K_{Y'}, \mathfrak{s}')$ is a Hilbert submanifold of $\mathcal{B}_k(K_{Y'}, \mathfrak{s}')$.

Proof We seek an analogue of [22, Proposition 24.3.1]. The main point in the proof of that proposition was to show that the operator $Q_{(A, \Phi)}$ introduced in (18) is surjective. To show surjectivity, the idea in the book was to apply [22, Corollary 17.1.5]. We will not use directly the corollary but rather its proof.

(a) First one needs to check that

$$Q_{(A, \Phi)}: L_k^2(K_{Y'}; iT^*K_{Y'} \oplus S^+) \rightarrow L_{k-1}^2(K_{Y'}; i\mathfrak{su}(S^+) \oplus S^-)$$

has closed range. The proof in [22, Corollary 17.1.5] for the closed range property was based on [22, Theorem 17.1.3], which showed that on a compact manifold with boundary, the operator

$$(23) \quad Q_{(A, \Phi)} \oplus (\pi_0 \circ r): L_k^2(X; iT^*K_{Y'} \oplus S^+) \rightarrow L_{k-1}^2(X; i\mathfrak{su}(S^+) \oplus S^-)$$

is Fredholm, where X denotes the compact manifold with boundary and $\pi_0 \circ r$ denotes Atiyah–Patodi–Singer spectral conditions. If we can show that (23) continues to be Fredholm when X is replaced with $K_{Y'}$ we would thus be done. But this just follows from a parametrix argument, since we already know that on a manifold without boundary and with a symplectic end, the operator $Q_{(A, \Phi)}$ is Fredholm [21, Theorem 3.3], while on a compact manifold with boundary, $Q_{(A, \Phi)} \oplus (\pi_0 \circ r)$ is Fredholm. Again, this type of parametrix argument can be found in our proof of Lemma 30 for example.

(b) The next step is to show that $Q_{(A, \Phi)}^*$ has the property that every nonzero solution of $Q_{(A, \Phi)}^*v = 0$ for $v = (\eta, \psi, \vartheta)$ has nonzero restriction to the boundary $\partial K_{Y'}$.

Using (19), we can see that the equation $Q_{(A,\Phi)}^*(\eta, \psi, \vartheta) = 0$ becomes in the coordinates (a, Ψ) of $Q_{(A,\Phi)}$ (compare with [22, Equation 24.10])

$$(24) \quad \begin{aligned} (d^+)^* \rho^* \eta + \rho^* (\psi \Phi^*) - d\vartheta &= 0, \\ D_A^* \psi - \eta \Phi + \vartheta \Phi &= 0. \end{aligned}$$

As in [22, (24.15)], the equations in (24) have the shape

$$\frac{d}{dt}v + (L_0 + h(t))v = 0,$$

where L_0 is a self-adjoint elliptic operator on Y' and h is a time-dependent operator on Y' satisfying the conditions of the unique continuation lemma. Since v vanishes on the boundary, it vanishes on the collar too and therefore on the cone $K_{Y'}$.

With (a) and (b) the surjectivity of $Q_{(A,\Phi)}$ follows so the proof that the moduli space $\mathcal{M}([1, \infty) \times Y', \mathfrak{s}')$ is a Hilbert submanifold of $\mathcal{B}_k(K_{Y'}, \mathfrak{s}')$ follows from our initial remarks. $\square$

Remark 25 As in the other cases, we have a restriction map

$$R_K: \mathcal{M}([1, \infty) \times Y', \mathfrak{s}') \rightarrow \mathcal{B}_{k-\frac{1}{2}}^\sigma(-Y', \mathfrak{s}_{\xi'}).$$

4.2 Gluing the moduli spaces

Now that we know that each moduli space appearing in the fiber product description (15) is a Hilbert manifold, we need to show that their fiber product is a finite-dimensional manifold, possibly with components of different dimensions. As mentioned before, we have the restriction maps

$$\begin{aligned} R_\tau: \mathcal{M}(\mathbb{R}^+ \times -Y, \mathfrak{s}_\xi, [c]) &\rightarrow \mathcal{B}_{k-\frac{1}{2}}^\sigma(Y, \mathfrak{s}_\xi), \\ R_W^-: \mathcal{M}(W^\dagger, \mathfrak{s}_\omega) &\rightarrow \mathcal{B}_{k-\frac{1}{2}}^\sigma(-Y, \mathfrak{s}_\xi), \\ R_W^+: \mathcal{M}(W^\dagger, \mathfrak{s}_\omega) &\rightarrow \mathcal{B}_{k-\frac{1}{2}}^\sigma(Y', \mathfrak{s}_{\xi'}), \\ R_K: \mathcal{M}([1, \infty) \times Y', \mathfrak{s}') &\rightarrow \mathcal{B}_{k-\frac{1}{2}}^\sigma(-Y', \mathfrak{s}_{\xi'}). \end{aligned}$$

If we write as before an element $[\gamma] \in \mathcal{M}(W_{\xi', Y}^+, \mathfrak{s}_\omega, [c])$ as

$$\text{Fib}(R_\tau, R_W^-, R_W^+, R_K) \ni [\gamma] = ([\gamma]_{\mathbb{R}^+ \times -Y}, [\gamma]_W, [\gamma]_{[1, \infty) \times Y'})$$

and define

$$\mathcal{B}_{k-\frac{1}{2}}^\sigma(-Y, \mathfrak{s}_\xi) \ni \mathfrak{b} = R_W^-(\gamma_W) \quad \text{and} \quad \mathcal{B}_{k-\frac{1}{2}}^\sigma(Y', \mathfrak{s}_{\xi'}) \ni \mathfrak{b}' = R_W^+(\gamma_W),$$

then the derivatives of our restriction maps can be written as

$$\begin{aligned}\mathcal{D}R_{[\gamma_{\mathbb{R}+\times-Y}]}^{\tau} &: T_{[\gamma_{\mathbb{R}+\times-Y}]} \mathcal{M}^{\tau}(\mathbb{R}^{+} \times -Y, \mathfrak{s}_{\xi}, [c]) \rightarrow \mathcal{K}_{k-\frac{1}{2}, b}^{\sigma}(Y, \mathfrak{s}_{\xi}), \\ \mathcal{D}R_{\mathcal{W}, [\gamma_{\mathcal{W}}]}^{-} &: T_{[\gamma_{\mathcal{W}}]} \mathcal{M}(W^{\dagger}, \mathfrak{s}_{\omega}) \rightarrow \mathcal{K}_{k-\frac{1}{2}, b}^{\sigma}(-Y, \mathfrak{s}_{\xi}), \\ \mathcal{D}R_{\mathcal{W}, [\gamma_{\mathcal{W}}]}^{+} &: T_{[\gamma_{\mathcal{W}}]} \mathcal{M}(W^{\dagger}, \mathfrak{s}_{\omega}) \rightarrow \mathcal{K}_{k-\frac{1}{2}, b'}^{\sigma}(Y', \mathfrak{s}_{\xi'}), \\ \mathcal{D}R_{K, [\gamma_{[1, \infty) \times Y'}]} &: T \mathcal{M}_{[\gamma_{[1, \infty) \times Y'}]}([1, \infty) \times Y', \mathfrak{s}) \rightarrow \mathcal{K}_{k-\frac{1}{2}, b'}^{\sigma}(-Y', \mathfrak{s}_{\xi'}),\end{aligned}$$

where the right-hand side of each is the corresponding Coulomb slice at the configuration b or b' . The next definition is the analogue of [22, Definition 24.4.2]:

Definition 26 Let $[\gamma] \in \mathcal{M}(W_{\xi', Y}^{+}, \mathfrak{s}_{\omega}, [c])$ and let

$$\rho: \mathcal{M}(W_{\xi', Y}^{+}, \mathfrak{s}_{\omega}, [c]) \rightarrow \mathcal{M}^{\tau}(\mathbb{R}^{+} \times -Y, \mathfrak{s}_{\xi}, [c]) \times \mathcal{M}(W^{\dagger}, \mathfrak{s}_{\omega}) \times \mathcal{M}([1, \infty) \times Y', \mathfrak{s}')$$

be the restriction map. Write

$$\rho([\gamma]) = ([\gamma_1], [\gamma_2], [\gamma_3]) = ([\gamma_{\mathbb{R}+\times-Y}], [\gamma_{\mathcal{W}}], [\gamma_{[1, \infty) \times Y'}]) \in \text{Fib}(R_{\tau}, R_{\mathcal{W}}^{-}, R_{\mathcal{W}}^{+}, R_K)$$

and

$$\begin{aligned}[b] &= R_{\tau}([\gamma_{\mathbb{R}+\times-Y}]) = R_{\mathcal{W}}^{-}([\gamma_{\mathcal{W}}]) \in \mathcal{B}_{k-\frac{1}{2}}^{\sigma}(-Y, \mathfrak{s}_{\xi}), \\ [b'] &= R_{\mathcal{W}}^{+}([\gamma_{\mathcal{W}}]) = R_K([\gamma_{[1, \infty) \times Y'}]) \in \mathcal{B}_{k-\frac{1}{2}}^{\sigma}(Y', \mathfrak{s}_{\xi'}).\end{aligned}$$

We say that the moduli space $\mathcal{M}(W_{\xi', Y}^{+}, \mathfrak{s}_{\omega}, [c])$ is *regular* at $[\gamma]$ if the map

$$\begin{aligned}R = ((R_{\tau}, R_{\mathcal{W}}^{-}), (R_{\mathcal{W}}^{+}, R_K)): \text{Fib}(R_{\tau}, R_{\mathcal{W}}^{-}, R_{\mathcal{W}}^{+}, R_K) \\ \rightarrow \mathcal{B}_{k-\frac{1}{2}}^{\sigma}(-Y, \mathfrak{s}_{\xi}) \times \mathcal{B}_{k-\frac{1}{2}}^{\sigma}(Y', \mathfrak{s}_{\xi'})\end{aligned}$$

is transverse at $\rho[\gamma]$. That is, $(R_{\tau}, R_{\mathcal{W}}^{-})$ is transverse at $[b]$ while $(R_{\mathcal{W}}^{+}, R_K)$ is transverse at $[b']$.

Following the strategy in [22, Section 24.4], to show regularity what we really need is an analogue of [22, Lemma 24.4.1] (which is our next lemma). The other pieces used in [22] do not change so we can conclude the following transversality result (compare with [22, Proposition 24.4.7]):

Theorem 27 Let q_{-Y} and $q_{Y'}$ be fixed perturbations for $-Y$ and Y' respectively such that for all critical points $[a], [b] \in \mathcal{B}_{k-\frac{1}{2}}^{\sigma}(-Y, \mathfrak{s}_{\xi})$ and $[a'], [b'] \in \mathcal{B}_{k-\frac{1}{2}}^{\sigma}(Y', \mathfrak{s}_{\xi'})$, the moduli spaces of trajectories $\mathcal{M}([a], \mathbb{R} \times -Y, \mathfrak{s}_{\xi}, [b])$ and $\mathcal{M}([a'], \mathbb{R} \times Y', \mathfrak{s}_{\xi'}, [b'])$ are cut out transversely. Then there is a residual subset $\mathcal{P}_0$ of the large space of perturbations $\mathcal{P}(-Y, \mathfrak{s}_{\xi}) \times \mathcal{P}(Y', \mathfrak{s}_{\xi'})$ defined in [22, Section 11.6] for which the following holds:

if for any $(\mathfrak{p}_0, \mathfrak{p}'_0) \in \mathcal{P}_0 \subset \mathcal{P}(-Y, \mathfrak{s}_\xi) \times \mathcal{P}(Y', \mathfrak{s}_{\xi'})$ one forms the perturbation

$$\mathfrak{p}_{W_{\xi', Y}^+} = -\widehat{\mathfrak{q}}_{Y, g_\theta, \mathfrak{s}_\xi} + (\beta \widehat{\mathfrak{q}}_{Y, g_\theta, \mathfrak{s}_\xi} + \beta'_0 \widehat{\mathfrak{p}}_0) + (\beta'_0 \widehat{\mathfrak{p}}_0 + \beta' \widehat{\mathfrak{q}}_{Y', g_{\theta'}, \mathfrak{s}_{\xi'}}) + (\beta_{N_K} \widehat{\mathfrak{q}}_{Y', g_{\theta'}, \mathfrak{s}_{\xi'}} + \beta_K \mathfrak{p}_K)$$

described in (11), then the moduli space $\mathcal{M}(W_{\xi', Y}^+, \mathfrak{s}_\omega, [\mathfrak{c}], \mathfrak{p}_{W_{\xi', Y}^+})$ defined using the perturbation $\mathfrak{p}_{W_{\xi', Y}^+}$ is regular; in other words, we have transversality at $\rho[\gamma]$ for all $[\gamma] \in \mathcal{M}(W_{\xi', Y}^+, \mathfrak{s}_\omega, [\mathfrak{c}], \mathfrak{p}_{W_{\xi', Y}^+})$.

In particular, for any perturbation belonging to this residual set, the moduli space

$$\mathcal{M}(W_{\xi', Y}^+, \mathfrak{s}_\omega, [\mathfrak{c}], \mathfrak{p}_{W_{\xi', Y}^+})$$

will be a manifold whose components have dimensions equal to

$$\text{ind } \mathcal{D}_{\rho[\gamma]} R = \dim \ker \mathcal{D}_{\rho[\gamma]} R.$$

Again, the proof of this theorem is a consequence of the following lemma:

Lemma 28 *Let $[\gamma] \in \mathcal{M}(W_{\xi', Y}^+, \mathfrak{s}_\omega, [\mathfrak{c}])$. Then the sum of the derivatives*

$$(25) \quad \mathcal{D}_{\rho[\gamma]} R = (\mathcal{D}_{[\gamma_1]} R_\tau + \mathcal{D}_{[\gamma_2]} R_{\overline{W}}^-) \oplus (\mathcal{D}_{[\gamma_2]} R_{\overline{W}}^+ + \mathcal{D}_{[\gamma_3]} R_K)$$

mapping

$$\begin{aligned} T_{[\gamma_1]} \mathcal{M}^\tau(\mathbb{R}^+ \times -Y, \mathfrak{s}_\xi, [\mathfrak{c}]) \oplus T_{[\gamma_2]} \mathcal{M}(W^\dagger, \mathfrak{s}_\omega) \oplus T_{[\gamma_3]} \mathcal{M}([1, \infty) \times Y', \mathfrak{s}) \\ \rightarrow \mathcal{K}_{k-\frac{1}{2}, \mathfrak{b}}^\sigma(-Y) \oplus \mathcal{K}_{k-\frac{1}{2}, \mathfrak{b}'}(Y') \end{aligned}$$

is Fredholm.

Proof We will begin showing that the following maps are Fredholm and compact:

- (1) $\pi_{\mathfrak{b}} \circ \mathcal{D}_{[\gamma_1]} R_\tau$ is compact.
- (2) $(1 - \pi_{\mathfrak{b}}) \circ \mathcal{D}_{[\gamma_1]} R_\tau$ is Fredholm.
- (3) $(1 - \pi_{\mathfrak{b}'}) \circ \mathcal{D}_{[\gamma_2]} R_{\overline{W}}^+$ is compact.
- (4) $\pi_{\mathfrak{b}'} \circ \mathcal{D}_{[\gamma_2]} R_{\overline{W}}^+$ is Fredholm.
- (5) $(1 - \pi_{\mathfrak{b}}) \circ \mathcal{D}_{[\gamma_2]} R_{\overline{W}}^-$ is compact.
- (6) $\pi_{\mathfrak{b}} \circ \mathcal{D}_{[\gamma_2]} R_{\overline{W}}^-$ is Fredholm.
- (7) $\pi_{\mathfrak{b}'} \circ \mathcal{D}_{[\gamma_3]} R_K$ is compact.
- (8) $(1 - \pi_{\mathfrak{b}'}) \circ \mathcal{D}_{[\gamma_3]} R_K$ is Fredholm.

Here π_b and $\pi_{b'}$ are defined as follows [22, Sections 12.4 and 17.3]. We have a Hessian operator $\text{Hess}_q^\sigma: \mathcal{K}_k^\sigma \rightarrow \mathcal{K}_{k-1}^\sigma$ obtained by projecting $\mathcal{D}(\text{grad } \ell)^\sigma$ onto the subspace $\mathcal{K}_{k-1}^\sigma$. The spectrum of Hess_q^σ is real, discrete and with finite-dimensional generalized eigenspaces. If the operator is hyperbolic (that is, zero is not an eigenvalue) we have a spectral decomposition

$$\mathcal{K}_{k-\frac{1}{2},b}^\sigma = \mathcal{K}_b^+ \oplus \mathcal{K}_b^-,$$

where $\mathcal{K}_b^+$ is the closure of the span of the positive eigenspaces and $\mathcal{K}_b^-$ of the negative eigenspaces. In the nonhyperbolic case, we choose ϵ sufficiently small that there are no eigenvalues in $(0, \epsilon)$ and then define $\mathcal{K}_{k-\frac{1}{2},b}^\pm$ using the spectral decomposition of the operator $\text{Hess}_{q,b}^\sigma - \epsilon$. The effect is that the generalized 0 eigenspace belongs to $\mathcal{K}_b^-$.

Also, notice that the roles of the different operators are sometimes opposite because of the different orientations on the manifolds, namely,

$$\mathcal{K}_b^-(-Y) = \mathcal{K}_b^+(Y) \quad \text{and} \quad \mathcal{K}_{b'}^-(-Y') = \mathcal{K}_{b'}^+(Y').$$

We will now break the proof this lemma and the verification of (1)–(8) into three mini lemmas. □

Lemma 29 *Facts (1)–(6) stated in the proof of Lemma 28 hold.*

Proof By [22, Proposition 24.3.2], (3)–(6) are true (recall that in [22, Section 24.3] the boundary is a compact four-manifold that is allowed to be disconnected; in our case the boundary is simply $-Y \cup Y'$).

By the discussion in [22, Lemma 24.4.1], (1) and (2) are true. □

Now we turn to verifying (7) and (8). To explain what we need to do we will chase through some theorems of [22] and [30, Proposition 2.18, Lemma 3.17]. It is also useful to observe that (7) will be true because the proof in [22] is essentially a local argument near the boundary.

Lemma 30 *Facts (7) and (8) stated in the proof of Lemma 28 hold.*

Proof Assertions (7) and (8) are the “conical” versions of [22, Proposition 24.3.2]. The proof of that theorem relies on other results in [22] as follows: The proof refers to Theorem 17.3.2, which requires Proposition 17.2.6, which depends on Proposition 17.2.5. The latter uses essentially Theorem 17.1.3 and the only part that is not proven

explicitly is part (a), which depends on a parametrix argument (modeled on Proposition 14.2.1) of Theorem 17.1.4.

In a nutshell, we must do the following. Decompose $Q_{(A,\Phi)}$ as

$$\begin{aligned} Q_{(A,\Phi)} &= D_0 + K, \\ D_0(a, \Psi) &= (\rho(d^+a), D_{A_0}\Psi, -d^*a), \\ K(a, \Psi) &= (-\{\Phi\Psi^* + \Psi\Phi^*\}_0, \rho(A - A_0)\Psi + \rho(a)\Phi + i\operatorname{Re}\langle i\Phi, \Psi\rangle). \end{aligned}$$

On the collar of $\partial K_{Y'}$, we can write D_0 in the form

$$\frac{d}{dt} + L_0,$$

where $L_0: C^\infty(-Y'; E_0) \rightarrow C^\infty(-Y'; E_0)$ is a first-order, self-adjoint elliptic differential operator. We will not write the exact formulas for the domain and codomain since they would rather cumbersome. Rather we will denote the bundles involved by the letter E_0 when referring to the three-manifolds and by E for the four-manifolds just as [22] does.

If H_0^+ and H_0^- are the closures in $L^2_{\frac{1}{2}}(Y; E_0)$ of the spans of the eigenvectors belonging to positive and nonpositive eigenvalues of L_0 and

$$\Pi_0: L^2_{\frac{1}{2}}(Y; E_0) \rightarrow L^2_{\frac{1}{2}}(Y; E_0)$$

is the projection with image H_0^- and kernel H_0^+ , we need to show that the operator

$$Q_{(A,\Phi)} \oplus (\Pi_0 \circ r_{-Y'}): L^2_k(K_{Y'}; E) \rightarrow L^2_{k-1}(K_{Y'}; E) \oplus (H_0^- \cap L^2_{k-\frac{1}{2}})$$

is Fredholm. First, for notational purposes take the collar neighborhood of $\partial K_{Y'}$ to be $(-5, 0] \times -Y'$, where $\partial K_{Y'}$ has now been identified with $\{0\} \times -Y'$. Also write for simplicity

$$Q_{K_{Y'}} = Q_{(A,\Phi)}: L^2_k(K_{Y'}; E) \rightarrow L^2_{k-1}(K_{Y'}; E).$$

To show the Fredholm property mentioned above we will give a parametrix argument, which is essentially the same as the one used in [22, Proposition 14.2.1]. Namely, we modify the manifold $K_{Y'}$ in two different ways.

For the first modification we close up $K_{Y'}$ first by extending the collar neighborhood a little bit (to the left in Figure 6) and then finding a four-manifold X (dots on the left side of Figure 6) bounding Y' . For the second modification, we forget about the part of the cone $K_{Y'}$ which does not have a product structure; in other words, we take the




Figure 6: Closing up the cone $K_{Y'}$ into the manifold $X \cup K_{Y'}$. Simultaneously, we extend the product neighborhood $(-5, 0] \times -Y'$ of $K_{Y'}$ into a half-infinite cylinder $Z = (-\infty, 0] \times -Y'$.

collar neighborhood of $K_{Y'}$ and extend it into a half-infinite cylinder which extends indefinitely to the right in Figure 6. In particular, notice that we superimposed both modifications in our image to save some space but they do not interact with each other. Each modification provides a parametrix as follows.

Regarding the first modification, we can define the manifold $X^+ = X \cup \text{cylinder} \cup K_{Y'}$ and extend $\mathcal{Q}_{K_{Y'}}$ to an operator

$$\mathcal{Q}_{X^+}: L_k^2(X^+; E) \rightarrow L_{k-1}^2(X^+; E)$$

and by [21, Theorem 3.3] there is a parametrix (that is, $\mathcal{Q}_{X^+} P_{X^+} - I$ and $P_{X^+} \mathcal{Q}_{X^+} - I$ are compact operators), which we denote by

$$P_{X^+}: L_{k-1}^2(X^+; E) \rightarrow L_k^2(X^+; E).$$

Similarly, for the second modification we define the half-cylinder $Z = (-\infty, 0] \times -Y'$. By [22, Theorem 17.1.4], the operator

$$\mathcal{Q}_Z \oplus (\Pi_0 \circ r_{-Y'}): L_k^2(Z; E) \rightarrow L_{k-1}^2(Z; E) \oplus (H_0^- \cap L_{k-\frac{1}{2}}^2(-Y'; E_0))$$

has a parametrix

$$P_Z: L_{k-1}^2(Z; E) \oplus (H_0^- \cap L_{k-\frac{1}{2}}^2(-Y'; E_0)) \rightarrow L_k^2(Z; E).$$

Finally, to define the parametrix corresponding to $\mathcal{Q}_{K_{Y'}} \oplus (\Pi_0 \circ r_{-Y'})$, let $1 = \eta_1 + \eta_2$ be a partition of unity subordinate to a covering of $K_{Y'}$ by the open sets

$$U_1 = K_{Y'} \setminus ([-2, 0] \times -Y') \quad \text{and} \quad U_2 = (-3, 0] \times -Y'.$$

Let γ_1 be a function which is 1 on the support of η_1 and vanishes on $(-1, 0] \times -Y'$. Similarly, let γ_2 be 1 on the support of η_2 and vanishing outside $[-4, 0] \times Y'$. Define

$$P_{K_{Y'}}: L^2_{k-1}(K_{Y'}; E) \oplus (H_0^- \cap L^2_{k-\frac{1}{2}}) \rightarrow L^2_k(K_{Y'}; E),$$

$$e \mapsto \gamma_1 P_{X+}(\eta_1 e) + \gamma_2 P_Z(\eta_2 e).$$

Notice that thanks to how the supports of the functions were chosen, the function is actually well defined. A similar computation to that in [22, Proposition 14.2.1] shows that $P_{K_{Y'}}$ is a parametrix for $\mathcal{Q}_{K_{Y'}} \oplus (\Pi_0 \circ r_{-Y'})$. $\square$

Now we can finish the proof of Lemma 28.

Lemma 31 $\mathcal{D}_{\rho[\gamma]}R$ is a Fredholm map.

Proof Thanks to items (1)–(8) established at the beginning of the proof of Lemma 28 we can see that

$$\begin{aligned} \mathcal{D}_{[\gamma_1]}R_\tau + \mathcal{D}_{[\gamma_2]}R_{\bar{W}} \\ = \underbrace{(1 - \pi_b) \circ \mathcal{D}_{[\gamma_1]}R_\tau}_{\text{Fredholm}} + \underbrace{\pi_b \circ \mathcal{D}_{[\gamma_2]}R_{\bar{W}}}_{\text{Fredholm}} + \underbrace{\pi_b \circ \mathcal{D}_{[\gamma_1]}R_\tau}_{\text{compact}} + \underbrace{(1 - \pi_b) \circ \mathcal{D}_{[\gamma_2]}R_{\bar{W}}}_{\text{compact}}. \end{aligned}$$

Likewise,

$$\begin{aligned} \mathcal{D}_{[\gamma_3]}R_K + \mathcal{D}_{[\gamma_2]}R_W^+ \\ = \underbrace{(1 - \pi_{b'}) \circ \mathcal{D}_{[\gamma_3]}R_K}_{\text{Fredholm}} + \underbrace{\pi_{b'} \circ \mathcal{D}_{[\gamma_2]}R_W^+}_{\text{Fredholm}} + \underbrace{\pi_{b'} \circ \mathcal{D}_{[\gamma_3]}R_K}_{\text{compact}} + \underbrace{(1 - \pi_{b'}) \circ \mathcal{D}_{[\gamma_2]}R_W^+}_{\text{compact}}. \end{aligned}$$

Therefore,

$$(\mathcal{D}_{[\gamma_1]}R_\tau + \mathcal{D}_{[\gamma_2]}R_{\bar{W}}) \oplus (\mathcal{D}_{[\gamma_2]}R_W^+ + \mathcal{D}_{[\gamma_3]}R_K)$$

differs by the compact operator

$$(\pi_b \circ \mathcal{D}_{[\gamma_1]}R_\tau + (1 - \pi_b) \circ \mathcal{D}_{[\gamma_2]}R_{\bar{W}}) \oplus (\pi_{b'} \circ \mathcal{D}_{[\gamma_3]}R_K + (1 - \pi_{b'}) \circ \mathcal{D}_{[\gamma_2]}R_W^+)$$

from the direct sum of the Fredholm operators

$$((1 - \pi_b) \circ \mathcal{D}_{[\gamma_1]}R_\tau \oplus \pi_b \circ \mathcal{D}_{[\gamma_2]}R_{\bar{W}}) \oplus ((1 - \pi_{b'}) \circ \mathcal{D}_{[\gamma_3]}R_K \oplus \pi_{b'} \circ \mathcal{D}_{[\gamma_2]}R_W^+)$$

and so the result follows. $\square$

5 Stretching the neck

As promised when we explained our strategy for proving naturality, we will consider a parametrized moduli space following the ideas used in [23, Sections 4.9, 4.10 and 6.3] and [22, Sections 24.6, 26.1 and 27.4]. Thanks to the computations in [53, Sections 5.5 and 6], formally, our situation of cylinder + compact + cone behaves in the same way as if we were working in the context of cylinder + compact, which is where the theorems just mentioned strictly speaking apply.

Recall that we want to show that $\widetilde{\mathrm{HM}}_{\bullet}(W^{\dagger}, \mathfrak{s}_{\omega})\, c(\xi') = c(\xi', Y)$; in other words, at the chain level we must have

$$\check{m}c(\xi') - c(\xi', Y) \in \mathrm{im}\, \check{\partial}_{-Y}.$$

The strategy we spelled out consisted in attaching a cylinder of length L to $W^+_{\xi', Y}$ and studying the Seiberg–Witten equations on

$$W^+_{\xi', Y}(L) = ([1, \infty) \times Y') \cup ([0, L] \times -Y') \cup W^{\dagger} \cup (\mathbb{R}^+ \times -Y).$$

Equivalently, as explained in [22, Section 24.6], we can consider a family of metrics g_L and perturbations on $W^{\dagger}$, all of which are equal near Y' . For example, we can choose a fraction of the collar neighborhood near Y' and instead of using the product metric $dt \otimes dt + g_{Y'}$, we use a smoothed-out version of the metric $g_L = L^2 dt \otimes dt + g_{Y'}$, which agrees with the old metric outside this region. In any case, we obtain a parametrized configuration space

$$\mathfrak{M}_z(W^+_{\xi', Y}, \mathfrak{s}_{\omega}, [c]) = \bigcup_{L \in [0, \infty)} \{L\} \times \mathcal{M}_z(W^+_{\xi', Y}(L), \mathfrak{s}_{\omega}, [c])$$

which we can identify with a subset of $[0, \infty) \times \mathcal{B}^{\sigma}_{k, \mathrm{loc}}(W^+_{\xi', Y}, \mathfrak{s}_{\omega})$ as follows (see [22, remark before Definition 24.4.9] and [36, Section 2.3]):

For any $t \in [0, \infty)$ there is a unique automorphism $b_t\colon TW^+_{\xi', Y} \rightarrow TW^+_{\xi', Y}$ that is positive, symmetric with respect to g_0 and has the property that $g_0(u, v) = g_t(b_t(u), b_t(v))$. The map induced by b_t on orthonormal frames gives rise to a map of spinor bundles $\bar{b}_t\colon S^{\pm}_0 \rightarrow S^{\pm}_t$ associated to the metrics g_0 and g_t . This map is an isomorphism preserving the fiberwise length of spinors. The identification

$$[0, \infty) \times \mathcal{B}^{\sigma}_{k, \mathrm{loc}}(W^+_{\xi', Y}, \mathfrak{s}_{\omega}) \rightarrow \bigcup_{L \in [0, \infty)} \{L\} \times \mathcal{B}^{\sigma}_{k, \mathrm{loc}}(W^+_{\xi', Y}(L), \mathfrak{s}_{\omega})$$

is then given by

$$(26) \quad (L, A, \mathbb{R}^+ \phi, \Phi) \mapsto (L, A, \mathbb{R}^+ \bar{b}_L(\phi), \bar{b}_L(\Phi)).$$

Just as in [22, Proposition 26.1.3], the moduli space $\mathfrak{M}_z(W_{\xi', Y}^+, \mathfrak{s}_\omega, [c])$ is a smooth manifold with boundary. The boundary is the fiber over $L = 0$, that is, the original moduli space $\mathcal{M}_z(W_{\xi', Y}^+, \mathfrak{s}, [c])$. Each individual moduli space $\mathcal{M}_z(W_{\xi', Y}^+(L), \mathfrak{s}_\omega, [c])$ can be compactified into $\mathcal{M}_z^+(W_{\xi', Y}^+(L), \mathfrak{s}_\omega, [c])$ by adding broken trajectories as in [22, Definition 24.6.1]¹ and to compactify

$$\bigcup_{L \in [0, \infty)} \{L\} \times \mathcal{M}_z^+(W_{\xi', Y}^+(L), \mathfrak{s}, [c])$$

we add a fiber over $L = \infty$, which is denoted by $\mathcal{M}_z^+(W_{\xi', Y}^+(\infty), \mathfrak{s}, [c])$, where

$$(27) \quad W_{\xi', Y}^+(\infty) = (K_{Y'} \cup [\mathbb{R}^+ \times -Y']) \cup ([\mathbb{R}^- \times -Y'] \cup W^\dagger \cup [\mathbb{R}^+ \times -Y]).$$

An element in this space consists (at most) of a quadruple $([\gamma_{K'}], [\check{\gamma}_{Y'}], [\gamma_{W^\dagger}], [\check{\gamma}_Y])$ where:

- $[\gamma_{K'}] \in \mathcal{M}(Z_{Y', \xi'}^+, \mathfrak{s}', [\mathfrak{a}_{Y'}])$ is a solution on $[\mathbb{R}^+ \times -Y] \cup K_{Y'}$.
- $[\check{\gamma}_{Y'}] \in \check{\mathcal{M}}^+([\mathfrak{a}_{Y'}], \mathfrak{s}_{\xi'}, [\mathfrak{b}_{Y'}])$ is an unparametrized trajectory on the cylinder $\mathbb{R} \times -Y'$.
- $[\gamma_{W^\dagger}] \in \mathcal{M}([\mathfrak{b}_{Y'}], W_*^\dagger, \mathfrak{s}_\omega, [\mathfrak{b}_Y])$ is a solution on $W_*^\dagger$, that is, $W^\dagger$ with two cylindrical ends attached to it.
- $[\check{\gamma}_Y] \in \check{\mathcal{M}}^+([\mathfrak{b}_Y], \mathfrak{s}_\xi, [c])$ is an unparametrized trajectory on the cylinder $\mathbb{R} \times -Y$.

Just as in [22, Proposition 26.1.4], the space

$$\mathfrak{M}_z^+(W_{\xi', Y}^+, \mathfrak{s}, [c]) = \bigcup_{L \in [0, \infty]} \{L\} \times \mathcal{M}_z^+(W_{\xi', Y}^+(L), \mathfrak{s}, [c])$$

is compact and when it is of dimension one, the zero-dimensional strata over $L = \infty$ are of the following types (compare with [22, Proposition 26.1.6]):

- $\mathcal{M}_{Z_{Y', \xi'}^+} \times \mathcal{M}_{W_*^\dagger}$,
- $\mathcal{M}_{Z_{Y', \xi'}^+} \times \mathcal{M}_{W_*^\dagger} \times \check{\mathcal{M}}_{-Y}$,
- $\mathcal{M}_{Z_{Y', \xi'}^+} \times \check{\mathcal{M}}_{-Y'} \times \mathcal{M}_{W_*^\dagger}$.

¹More precisely, for us a broken trajectory asymptotic to $[c]$ consists of an element $[\gamma_0]$ in a moduli space $\mathcal{M}_{z_0}(W_{\xi', Y}^+(L), \mathfrak{s}_\omega, [c])$ and an unparametrized broken trajectory $[\check{\gamma}]$ in a moduli space $\check{\mathcal{M}}_z([\mathfrak{c}_0], \mathfrak{s}_\xi, [c])$.

Here $\check{\mathcal{M}}$ denotes an unparametrized moduli space. Also, in the last two cases the middle space denotes a boundary-obstructed moduli space, ie it denotes trajectories which connect a boundary stable point (as $t \rightarrow -\infty$) with a boundary unstable point (as $t \rightarrow \infty$).

The following theorem shows that up to a boundary term, $\sum_z m_z(W_{\xi',Y}^+, \mathfrak{s}_\omega, [c])$ equals one of the sums (6) or (7). It can be seen as the analogue of [23, Lemma 4.15] and [22, Proposition 24.6.10] (in fact, it was used implicitly in the proof of the pairing formula in [23, Proposition 6.8] and [53, Theorem 6.2]):

Proposition 32 *If $\mathfrak{M}_z(W_{\xi',Y}^+, \mathfrak{s}, [c])$ is zero-dimensional, then $\mathfrak{M}_z(W_{\xi',Y}^+, \mathfrak{s}, [c])$ is compact. If $\mathfrak{M}_z(W_{\xi',Y}^+, \mathfrak{s}_\omega, [c])$ is one-dimensional and contains irreducible trajectories, then the compactification $\mathfrak{M}_z^+(W_{\xi',Y}^+, \mathfrak{s}_\omega, [c])$ is a one-dimensional manifold whose boundary points are of the following types:*

- (1) *The fiber over $L = 0$, namely the space $\mathcal{M}_z(W_{\xi',Y}^+, \mathfrak{s}_\omega, [c])$.*
- (2) *The fiber over $L = \infty$, namely the three products described previously.*
- (3) *Products of the form*

$$\mathfrak{M}(W_{\xi',Y}^+, \mathfrak{s}_\omega, [b]) \times \check{\mathcal{M}}([b], \mathfrak{s}_\xi, [c])$$

or

$$\mathfrak{M}(W_{\xi',Y}^+, \mathfrak{s}_\omega, [a]) \times \check{\mathcal{M}}([a], \mathfrak{s}_\xi, [b]) \times \check{\mathcal{M}}([b], \mathfrak{s}_\xi, [c]),$$

where, in the latter case, the middle space is boundary obstructed.

In order to apply the proposition define $P = [0, \infty)$ and the numbers

$$m_z(W_{\xi',Y}^+, \mathfrak{s}_\omega, [a])_P = \begin{cases} |\mathfrak{M}_z(W_{\xi',Y}^+, \mathfrak{s}_\omega, [a])| \bmod 2 & \text{if } \dim \mathfrak{M}_z(W_{\xi',Y}^+, \mathfrak{s}, [a]) = 0, \\ 0 & \text{otherwise.} \end{cases}$$

Recall also that the differential on $\check{C}_\bullet(-Y, \mathfrak{s}_\xi) = \mathfrak{C}^o(-Y, \mathfrak{s}_\xi) \oplus \mathfrak{C}^s(-Y, \mathfrak{s}_\xi)$ is [22, Definition 22.1.3]

$$\check{\partial} = \begin{pmatrix} \partial_o^o & -\partial_o^u \bar{\partial}_u^s \\ \partial_s^o & \bar{\partial}_s^s - \partial_u^u \bar{\partial}_u^s \end{pmatrix}.$$

Suppose now that $\mathfrak{M}_z(W_{\xi',Y}^+, \mathfrak{s}_\omega, [c])$ is one-dimensional. We use the previous proposition to count the endpoints of $\mathfrak{M}_z^+(W_{\xi',Y}^+, \mathfrak{s}_\omega, [c])$ by making cases on $[c]$.

Case $[c] \in \mathfrak{C}^o(-Y, \mathfrak{s}_\xi)$ (irreducible critical point)

(1) The fiber over $L = 0$ gives the contributions

$$(28) \qquad \sum_z m_z(W_{\xi', Y}^+, \mathfrak{s}_\omega, [c]).$$

These numbers were used in the chain-level definition of $c(\xi', Y)$.

(2) The fiber over $L = \infty$ gives the contributions (6)

$$(29) \qquad \sum_{[a] \in \mathfrak{C}^o(-Y')} \sum_{z_1, z_2} m_{z_1}(Z_{Y', \xi'}^+, \mathfrak{s}', [a]) n_{z_2}([a], W_*^\dagger, \mathfrak{s}_\omega, [c]) \\ + \sum_{\substack{[a] \in \mathfrak{C}^s(-Y') \\ [b] \in \mathfrak{C}^u(-Y')}} \sum_{z_1, z_2, z_3} m_{z_1}(Z_{Y', \xi'}^+, \mathfrak{s}', [a]) \bar{n}_{z_2}([a], \mathfrak{s}_{\xi'}, [b]) n_{z_3}([b], W_*^\dagger, \mathfrak{s}_\omega, [c]) \\ + \sum_{\substack{[a] \in \mathfrak{C}^s(-Y') \\ [b] \in \mathfrak{C}^u(-Y)}} \sum_{z_1, z_2, z_3} m_{z_1}(Z_{Y', \xi'}^+, \mathfrak{s}', [a]) \bar{n}_{z_2}([a], W_*^\dagger, \mathfrak{s}_\omega, [b]) n_{z_3}([b], \mathfrak{s}_\xi, [c]).$$

These numbers were used in the chain-level definition of $\check{m}c(\xi')$.

(3) We obtain contributions of the form

$$(30) \qquad \sum_{[a] \in \mathfrak{C}^o(-Y)} \sum_{w_1, w_2} m_{w_1}(W_{\xi', Y}^+, \mathfrak{s}_\omega, [a])_P n_{w_2}([a], \mathfrak{s}_\xi, [c]) \\ + \sum_{\substack{[a] \in \mathfrak{C}^s(-Y) \\ [b] \in \mathfrak{C}^u(-Y')}} \sum_{w_1, w_2, w_3} m_{w_1}(W_{\xi', Y}^+, \mathfrak{s}_\omega, [a])_P \bar{n}_{w_2}([a], \mathfrak{s}_\xi, [b]) n_{w_3}([b], \mathfrak{s}_\xi, [c]).$$

These numbers will be used in a moment to define the boundary term.

By Proposition 32 the sum of (28), (29) and (30) corresponds to the number of points in the boundary of a one-dimensional compact manifold, hence it must equal 0.

Case $[c] \in \mathfrak{C}^s(-Y, \mathfrak{s}_\xi)$ (boundary stable critical point)

(1) The fiber over $L = 0$ gives the contributions

$$(31) \qquad \sum_z m_z(W_{\xi', Y}^+, \mathfrak{s}_\omega, [c]).$$

These numbers were used in the chain-level definition of $c(\xi', Y)$.

(2) The fiber over $L = \infty$ gives the contributions (6)

$$\begin{aligned}
 (32) \quad & \sum_{[a] \in \mathfrak{C}^o(-Y')} \sum_{z_1, z_2} m_{z_1}(Z_{Y', \xi', s'}^+, [a]) n_{z_2}([a], W_*^\dagger, s_\omega, [c]) \\
 & + \sum_{[a] \in \mathfrak{C}^s(-Y')} \sum_{z_1, z_2} m_z(Z_{Y', \xi', s'}^+, [a]) \bar{n}_{z_2}([a], W_*^\dagger, s_\omega, [c]) \\
 & + \sum_{\substack{[a] \in \mathfrak{C}^s(-Y') \\ [b] \in \mathfrak{C}^u(-Y')}} \sum_{z_1, z_2, z_3} m_{z_1}(Z_{Y', \xi', s'}^+, [a]) \bar{n}_{z_2}([a], s_\xi, [b]) n_{z_3}([b], W_*^\dagger, s_\omega, [c]) \\
 & + \sum_{\substack{[a] \in \mathfrak{C}^s(-Y') \\ [b] \in \mathfrak{C}^u(-Y)}} \sum_{z_1, z_2, z_3} m_{z_1}(Z_{Y', \xi', s'}^+, [a]) \bar{n}_{z_2}([a], W_*^\dagger, s_\omega, [b]) n_{z_3}([b], s_\xi, [c]).
 \end{aligned}$$

These numbers were used in the chain-level definition of $\check{m}c(\xi')$.

(3) We obtain contributions of the form

$$\begin{aligned}
 (33) \quad & \sum_{[a] \in \mathfrak{C}^o(-Y)} \sum_{w_1, w_2} m_{w_1}(W_{\xi', Y}^+, s_\omega, [a])_P n_{w_2}([a], s_\xi, [c]) \\
 & + \sum_{[a] \in \mathfrak{C}^s(-Y)} \sum_{w_1, w_2} m_{w_1}(W_{\xi', Y}^+, s_\omega, [a])_P \bar{n}_{w_2}([a], s_\xi, [c]) \\
 & + \sum_{\substack{[a] \in \mathfrak{C}^s(-Y) \\ [b] \in \mathfrak{C}^u(-Y)}} \sum_{w_1, w_2, w_3} m_{w_1}(W_{\xi', Y}^+, s_\omega, [a])_P \bar{n}_{w_2}([a], s_\xi, [b]) n_{w_3}([b], s_\xi, [c]).
 \end{aligned}$$

These numbers will be used in a moment to define the boundary term.

As before, the sum of (31), (32) and (33) equals 0.

Define the chain element $\psi \in \mathfrak{C}^o(-Y, s_\xi) \oplus \mathfrak{C}^s(-Y, s_\xi)$ by

$$\left(\sum_{[a] \in \mathfrak{C}^o(-Y)} \sum_{w_1} m_{w_1}(W_{\xi', Y}^+, s_\omega, [a])_P e_{[a]}, \sum_{[a] \in \mathfrak{C}^s(-Y)} \sum_{w_1} m_{w_1}(W_{\xi', Y}^+, s_\omega, [a])_P e_{[a]} \right).$$

It is not hard to see that

$$\check{\partial}\psi = \left(\sum_{[c] \in \mathfrak{C}^o(-Y)} C_o e_{[c]}, \sum_{[c] \in \mathfrak{C}^s(-Y)} C_s e_{[c]} \right),$$

where C_o equals (30) and C_s equals (33). In other words, we have the chain-level identity

$$\check{m}c(\xi') - c(\xi', Y) = \check{\partial}\psi,$$

which gives us the desired identity

$$\widetilde{\mathrm{HM}}_{\bullet}(W^{\dagger}, \mathfrak{s}_{\omega}) \mathfrak{c}(\xi') = \mathfrak{c}(\xi', Y),$$

concluding the first phase in the proof for the naturality of the contact invariant under strong symplectic cobordisms. Now we proceed to address the second part of the proof (as explained at the beginning of the paper). Namely, we will show that $\mathfrak{c}(\xi', Y)$ equals $\mathfrak{c}(\xi)$ by adapting Mrowka and Rollin's "dilating the cone" technique to the case of a manifold with cylindrical end.

6 Generalized gluing–excision theorem

6.1 Gluing and identifying spin-c structures

Before describing the modified gluing argument why will say very quickly why the “special” condition can be dropped for the symplectic cobordisms we are working with. More details can be found in [8, Section 6.1]. The definition Mrowka and Rollin used for a special symplectic cobordism, which appears near formula (1.1) of [35], we repeat for convenience:

Definition 33 A cobordism $(W, \omega): (Y, \xi) \rightarrow (Y', \xi')$ is said to be a *special symplectic cobordism* if:

- (1) With the symplectic orientation, $\partial W = -Y \sqcup Y'$ and ω is strictly positive on ξ and ξ' with their induced orientations.
- (2) The symplectic form is given in a collar neighborhood of the concave boundary by a symplectization of (Y, ξ) .
- (3) The map induced by the inclusion $i^*: H^1(W, Y'; \mathbb{Z}) \rightarrow H^1(Y; \mathbb{Z})$ is the zero map.

Notice that it is the last condition that makes the symplectic cobordism “special”. We want to work with strong cobordisms, which in particular means that the convex end is also given by a symplectization of (Y, ξ) and that the special condition does not appear. The reason why Mrowka and Rollin introduced this condition is that they were interested in guaranteeing the injectivity of a certain map

$$J: \mathrm{Spin}^c(X, \xi) \rightarrow \mathrm{Spin}^c(X \cup W, \xi'),$$

where X was a compact manifold with boundary a contact manifold (Y, ξ) and $\mathrm{Spin}^c(X, \xi)$ denotes the isomorphism classes of (relative) spin-c structures on X

whose restriction to Y induce the spin-c structure determined by ξ . A similar definition applies to $\text{Spin}^c(X \cup W, \xi')$, where now $X \cup W$ bounds (Y', ξ') . In the same way in which for a manifold without boundary Z the set of spin-c structures $\text{Spin}^c(Z)$ is an affine space over $H^2(Z; \mathbb{Z})$, the set $\text{Spin}^c(X, \xi)$ is an affine space over $H^2(X; Y; \mathbb{Z})$ (this is discussed in the first three pages of [21]).

We are interested in the situation when $X = \mathbb{R}^+ \times -Y$ and therefore the affine space $H^2(\mathbb{R}^+ \times -Y; Y; \mathbb{Z})$ reduces automatically to a singleton, so regardless of how the map

$$j: \text{Spin}^c(\mathbb{R}^+ \times -Y, \xi) \rightarrow \text{Spin}^c(\mathbb{R}^+ \times -Y \cup W, \xi')$$

is defined, it will automatically be injective. In fact, the definition of such a map is not difficult to give: as we already mentioned the contact structure ξ gives rise to a canonical spinor model S_ξ on Y (also on $-Y$ [22, Section 22.5]) and on $\mathbb{R}^+ \times -Y$ [22, Section 4.3].

This canonical spinor bundle model over $\mathbb{R}^+ \times -Y$ represents the (unique) isomorphism class $\mathfrak{s}(\mathbb{R}^+ \times -Y, \xi)$ of relative spin-c structure inside $\text{Spin}^c(\mathbb{R}^+ \times -Y, \xi)$. We define $j[\mathfrak{s}(\mathbb{R}^+ \times -Y, \xi)]$ by specifying a relative spin-c structure over $\mathbb{R}^+ \times -Y \cup W$ as follows.

Using the symplectic structure on W we have a canonical spinor bundle S_ω as well. This induces spinor bundles on ∂W as explained in [22, Section 4.5]. Since the symplectic structure is specified near the boundary by the corresponding contact structure because of the strong condition in our cobordism it is not difficult to identify in this way S_ξ with $S_\omega|_{-Y}$ and hence we produce a total spinor bundle over $(\mathbb{R}^+ \times -Y) \cup W$, which is representing $j[\mathfrak{s}(\mathbb{R}^+ \times -Y, \xi)]$ (more details can be found in the author's thesis cited before).

Another way to explain why the special condition was needed in the paper [35] is to say that X could have interesting topology, so there was an obstruction problem when trying to extend certain data defined on the complement of X (for example gauge transformations) to the entire manifold. However, in our case these obstructions disappear since we have replaced X with a half-cylinder.

6.2 Connected sum along Y

We will now adapt the gluing/excision theorem in [35] to our situation. More precisely we want an analogue of [35, Corollary 3.2.2]. The following construction is based on Sections 4.1 and 2.1.5 from that paper. There they proved a gluing result for a

class of manifolds with a so-called *AFAK end* Z , that is, an asymptotically flat almost Kähler end, the idea being that manifolds of this class behave nicely enough near the symplectic end that all the necessary analysis goes through. We recall the definition of an AFAK end [35, Definition 2.1.2].

Definition 34 (AFAK end manifold) An asymptotically flat almost Kähler end is a manifold Z which admits a decomposition $C_Z \cup_Y N$, where N is a not necessarily compact four-dimensional manifold, with contact boundary Y , endowed with a fixed contact form θ , and $C_Z = (0, T] \times Y$ for some $T > 0$.

In addition, Z is endowed with an almost Kähler structure (ω_Z, J_Z) and a proper function $\sigma_Z: Z \rightarrow (0, \infty)$ satisfying:

- (a) On $(0, T] \times Y \subset Z$ we have $\sigma_Z(t, y) = t$.
- (b) The almost Kähler structure on C_Z is that of an almost Kähler cone on (Y, θ) .
- (c) There is a constant $\kappa > 0$ such that the injectivity radius satisfies $\kappa \operatorname{inj}(x) > \sigma(x)$ for all $x \in Z$.
- (d) For each $x \in Z$, let e_x be the map $e_x: v \rightarrow \exp_x(\sigma_Z(x)v/\kappa)$, and γ_x be the metric on the unit ball in $T_x Z$ defined as $e_x^* \gamma_x / \sigma_Z(x)^2$. Then these metrics have bounded geometry in the sense that all covariant derivatives of the curvature are bounded by some constants independent of x .
- (e) For each $x \in Z$, let o_x denote the symplectic form $e_x^* \omega_Z / \sigma_Z(x)^2$ on the unit ball. Then o_x similarly approximates the translation-invariant symplectic form, with all its derivatives.
- (f) For all $\epsilon > 0$, the function $e^{-\epsilon \sigma_Z}$ is integrable on Z .
- (g) The map $H_c^1(N) \rightarrow H^1(Y)$ induced by the inclusion $Y = \partial N \subset N$, where $H_c^*(N)$ is the compactly supported de Rham cohomology, is identically 0.

The important things that we need to point out regarding this definition are that:

- The last condition (g) regarding the vanishing of the map between de Rham cohomologies mimics the special condition for a symplectic cobordism that we already discussed before. Therefore, in our context this condition is not needed.
- To our cobordism (W, ω) one can associate an AFAK end (Z, ω_Z) as explained in [35, Section 4.1]. We can simplify the construction there in our case because our cobordism is strong, so in fact we can exploit the fact that near the convex end ω is

also determined by a symplectization of the contact structure. We start by using a collar neighborhood $[T_0, T_1) \times Y$ of $Y \subset \partial W$ (with $T_0 > 1$) and a contact form θ such that the symplectic form ω near the concave end of that neighborhood is given by $\frac{1}{2} d(t^2\theta)$. We then glue a sharp cone on the boundary Y by extending the collar neighborhood into $(0, T_1) \times Y$ with its symplectic form. Likewise, we have a similar collar neighborhood near the convex end and we can therefore glue (after some reparametrizations) the half-infinite cone $[1, \infty) \times Y'$ with the symplectic form $\frac{1}{2} d(t'^2\theta')$, where t' denotes the time coordinate on $[1, \infty) \times Y'$. Therefore we take Z to be

$$Z = ((0, T_0) \times Y) \cup W \cup ([1, \infty) \times Y').$$

Moreover, we can find a “time coordinate” σ_Z on Z as described in Definition 34; ie properties (a)–(e) are satisfied (in fact, after reparametrization it can be taken to agree with the natural time coordinate on the third factor $[1, \infty) \times Y'$ of Z).

- Notice in particular that our symplectic form ω_Z has the property that it is exact except for a compact set (which is contained in W). Hence the class of manifolds we are using could be called *AFAKAE ends* (where AE stands for almost exact) but for convenience we will keep calling this manifold an AFAK end. After choosing a metric g_Z and almost complex structure J_Z on Z so that ω_Z is self-dual and of pointwise norm $\sqrt{2}$ the data $(Z, \omega_Z, J_Z, g_Z, \sigma_Z)$ will represent an AFAK end with the caveats mentioned above.

This is the class of manifolds to which the generalized excision/gluing theorem will apply, though the theorem will only be used for this particular Z . The idea will be to glue Z to the cylindrical end $\mathbb{R}^+ \times -Y$ using an operation that Mrowka and Rollin named *connected sum along Y* .

To be more precise, consider as before the symplectic cone $[1, \infty) \times Y$ for the contact form θ with metric

$$g_{K,\theta} = dt \otimes dt + t^2 g_\theta$$

and symplectic form

$$\omega_\theta = \frac{1}{2} d(t^2\theta).$$

Choose a number² $\tau > 1$ and identify an annulus $(1, \tau) \times Y$ in $[1, \infty) \times Y$ with an annulus $(1/\tau, 1) \times Y \subset Z$ using the dilation map

$$(1, \tau) \times Y \xrightarrow{\nu_\tau} (1/\tau, 1) \times Y, \quad (t, y) \mapsto (t/\tau, y).$$

²It goes without saying that this number is completely unrelated to the τ used in the τ -model of the configuration space.




Figure 7: Using the “connected sum along Y ” operation to obtain the family of manifolds M_τ .

Define M_τ as the union of $(\mathbb{R}^+ \times -Y) \cup [1, \tau) \times Y$ and $Z \cap \{\sigma_Z > 1/\tau\}$,

(34)
$$M_\tau = ((\mathbb{R}^+ \times -Y) \cup [1, \tau) \times Y) \cup (Z \cap \{\sigma_Z > 1/\tau\}),$$

glued along the previous annuli via the dilation map ν_τ .

In the figure, the gray regions represent the annuli that are identified and the dashed regions are the parts of the cone and Z that are taken off in the construction. We need to say how to redefine the geometric structures we had in place (metric, symplectic form, etc) so that they agree under the identification operation. The symplectic form can be taken as

$$\omega_{Z,\tau} = \tau^2 \omega_Z,$$

and the new “time coordinate” becomes

$$\sigma_{\tau,Z} = \tau \sigma_Z.$$

The metric is a dilation of the original metric, that is,

$$g_{\tau,Z} = \tau^2 g_Z.$$

In this way, with respect to $g_{\tau,Z}$, the form $\omega_{Z,\tau}$ is self-dual with norm $\sqrt{2}$. As usual, g_Z and ω_Z determine a compatible almost complex structure $J_{Z,\tau}$ which in fact is independent of τ , ie

$$J_{Z,\tau} = J_Z.$$

The natural Clifford multiplication is

$$\rho_{\tau,Z}(\eta) = \frac{\rho_Z(\eta)}{\tau} \quad \text{for } \eta \text{ a one-form,}$$

while the spinor bundle remains the same, ie

$$S_{\tau,Z} = S_Z.$$

We will specify the spin-c structure $\mathfrak{s}_\tau$ on M_τ as the isomorphism class of the following spinor bundle (S_τ, ρ_τ) :

- On $\mathbb{R}^+ \times -Y$ we use the spinor bundle $(S_{\mathbb{R}^+ \times -Y}, \rho_{\mathbb{R}^+ \times -Y})$ that the canonical spinor bundle (S_θ, ρ_θ) on Y induces on $\mathbb{R}^+ \times -Y$.
- Along the boundary, we identify $(S_{\mathbb{R}^+ \times -Y, \theta}, \rho_{\mathbb{R}^+ \times -Y, \theta})|_Y$ with $(S_{K_Y}, \rho_{K_Y})|_{\{1\} \times Y}$, where (S_{K_Y}, ρ_{K_Y}) denotes the canonical spinor bundle associated to the symplectic cone $K_Y = [1, \infty) \times Y$.
- Over $M_\tau \cap \{\sigma_{\tau,Z} < \tau\} = M_\tau \cap \{\sigma_Z < 1\} = M_\tau \cap \{(1, \tau) \times Y\}$ we use the spinor bundle (S_{K_Y}, ρ_{K_Y}) .
- Over $M_\tau \cap \{\sigma_{\tau,Z} > 1\} = M_\tau \cap \{\sigma_Z > 1/\tau\}$ we use the spinor bundle $(S_{\tau,Z}, \rho_{\tau,Z}) = (S_{\tau,Z}, \rho_Z/\tau)$.

To write the transition map from (S_{K_Y}, ρ_{K_Y}) to $(S_{\tau,Z}, \rho_{\tau,Z})$ over $M_\tau \cap \{1/\tau < \sigma_Z < 1\}$ observe that if e_Y^1, e_Y^2, e_Y^3 is a coframe at the slice $\{1\} \times Y \simeq Y$ then $dt, te_Y^1, te_Y^2, te_Y^3$ is a coframe on $(1, \tau) \times Y \subset K_Y$ while $\tau dt, \tau te_Y^1, \tau te_Y^2, \tau te_Y^3$ is a coframe on $\{1/\tau < \sigma_Z < 1\} \subset Z$. Therefore we can define $\tilde{\epsilon}_t^{01} = \frac{1}{\sqrt{2}}(dt - ite_Y^1)$, $\tilde{\epsilon}_t^{23} = \frac{1}{\sqrt{2}}t(e_Y^2 - ie_Y^3)$ and the identification map

$$\mathfrak{G}_\tau: S_{K_Y} \rightarrow S_{\tau,Z}, \quad \alpha_{K_Y} + \beta_{K_Y} \tilde{\epsilon}_t^{01} \wedge \tilde{\epsilon}_t^{23} \mapsto \alpha_{K_Y} + \tau^2 \beta_{K_Y} \tilde{\epsilon}_t^{01} \wedge \tilde{\epsilon}_t^{23}.$$

Remark 35 The construction in [35] required (in that paper's notation) the choice of an element $(\mathfrak{s}, h) \in \text{Spin}^c(M, \omega)$ [35, Section 2.1.7]. As we explained before, by using a half-infinite cylinder instead of a compact piece, all of our constructions can be done in a canonical way, which is why our description is simpler and we can drop the explicit reference to h .

Our (unperturbed) Seiberg–Witten map continues to be

$$\mathfrak{F}(A, \Phi) = \left(\frac{1}{2} \rho(F_{A'}^+) - (\Phi \Phi^*)_0, D_A \Phi \right).$$

To define the perturbations, write the half-infinite cylinder as

$$\mathbb{R}^+ \times -Y = ([0, 1] \times -Y) \cup ([1, \infty) \times -Y),$$

where $[0, 1] \times -Y$ is going to play the role of a trivial cobordism. By that we simply mean that the perturbations we use on $[0, 1] \times -Y$ are of the form $\hat{\mathfrak{p}} = \beta \hat{\mathfrak{q}} + \beta_0 \hat{\mathfrak{p}}_0$, where $\hat{\mathfrak{p}}$ coincides near $\{1\} \times -Y$ with a strongly tame perturbation $-\hat{\mathfrak{q}}_{Y,g_\theta,\mathfrak{s}_\xi}$ on $[1, \infty) \times -Y$ and near $\{0\} \times -Y$ it vanishes. On

$$Z_\tau = [1, \tau) \times Y \cup (Z \cap \{\sigma_Z > 1/\tau\})$$

consider the perturbation

$$\mathfrak{p}_{Z_\tau} = -\frac{1}{2}\rho_\tau(F_{A_{0,\tau}^+}^+) + (\Phi_{\tau,0}\Phi_{\tau,0}^*)_0,$$

where $(A_{0,\tau}^+, \Phi_{\tau,0})$ denotes the canonical solution. Again, similar to the perturbation $\mathfrak{p}_{\mathcal{W}_{\xi,Y}^+}$ defined in (11) we can produce a perturbation

(35)
$$\mathfrak{p}_{M_\tau} = -\hat{\mathfrak{q}}_{Y,g_\theta,\mathfrak{s}_\xi} + (\beta \hat{\mathfrak{q}}_{Y,g_\theta,\mathfrak{s}_\xi} + \beta'_0 \hat{\mathfrak{p}}_0) + \beta_K \mathfrak{p}_{Z_\tau}.$$

It is also useful to think of the manifold $Z_{Y,\xi}^+$ (where the contact invariant $\mathfrak{c}(\xi)$ of (Y, ξ) is defined) as the manifold M_τ obtained by taking “ $\tau = \infty$ ”. In other words, we will write

$$M_\infty \equiv Z_{Y,\xi}^+.$$

Notice that on this manifold we can also define a perturbation $\mathfrak{p}_{M_\infty}$ in exactly the same way as for $\mathfrak{p}_{M_\tau}$ (so it agrees with $-\hat{\mathfrak{q}}_{Y,g_\theta,\mathfrak{s}_\xi}$ on the half-cylinder $[1, \infty) \times -Y$, it agrees with $\mathfrak{p}_K$ on the cone $[1, \infty) \times Y$ and it is interpolated between these two perturbations on the finite cylinder $[0, 1] \times -Y$ through a perturbation $\beta \hat{\mathfrak{q}}_{Y,g_\theta,\mathfrak{s}_\xi} + \beta'_0 \hat{\mathfrak{p}}_0$).

Our previous transversality theorem, Theorem 27, now reads as follows:

Lemma 36 *For all critical points $[c] \in \mathfrak{C}^o(-Y, \mathfrak{s}_\xi) \oplus \mathfrak{C}^s(-Y, \mathfrak{s}_\xi)$ and each $0 < \tau \leq \infty$ there is a residual subset $\mathcal{P}_\tau$ of the large space of perturbations $\mathcal{P}(Y, \mathfrak{s}_\xi)$ such that for any $\mathfrak{p}_\tau \in \mathcal{P}_\tau$ the corresponding perturbation $\mathfrak{p}_{M_\tau}$ satisfies the property that all the moduli spaces $\mathcal{M}(M_\tau, \mathfrak{s}_\tau, [c], \mathfrak{p}_{M_\tau})$ are cut out transversely.*

When we study the properties of the gluing map it will become clear that we want to be able to choose a single perturbation $\mathfrak{p}_{\text{all}}$ such that when we plug it in the formula for $\mathfrak{p}_{M_\tau}$ it guarantees transversality *simultaneously* for all moduli spaces $\mathcal{M}(M_\tau, \mathfrak{s}_\tau, [c], \mathfrak{p}_{M_\tau})$. In other words, we would like to be able to choose a perturbation $\mathfrak{p}_{\text{all}} \in \bigcap_{0 < \tau \leq \infty} \mathcal{P}_\tau$. However, notice that without further restrictions $\bigcap_{0 < \tau \leq \infty} \mathcal{P}_\tau$ might be empty.

Fortunately, because we are ultimately interested in the case when τ is sufficiently large we can choose an increasing sequence τ_n with $\tau_n \rightarrow \infty$ and then use the

fact that the countable intersection of residual sets is residual [38, Theorem 1.4], so that $(\bigcap_n \mathcal{P}_{\tau_n}) \cap \mathcal{P}_\infty$ is residual as well. In particular this means that we can take $\mathfrak{p}_{\text{all}} \in (\bigcap_n \mathcal{P}_{\tau_n}) \cap \mathcal{P}_\infty$, which we will assume from now on.

Strictly speaking, since we will work with an additional family M'_τ obtained by using another connected sum operation with another AFAK end Z' we should really take $\mathfrak{p}_{\text{all}} \in (\bigcap_n \mathcal{P}_{\tau_n}) \cap \mathcal{P}_\infty \cap (\bigcap_n \mathcal{P}'_{\tau_n})$, where $\mathcal{P}'_{\tau_n}$ denotes the residual space of perturbations for the manifold M'_τ . However, for the proof of the gluing theorem we will end up taking $Z' = (0, \infty) \times Y$ (as in [35, Section 4.1]), in which case one can check that *all* the M'_τ end up coinciding with $Z^+_{Y,\xi} = M_\infty$. Hence, this point does not make much of a difference. Also, for notational convenience, we will keep writing the moduli spaces typically as $\mathcal{M}(M_\tau, \mathfrak{s}_\tau, [c])$ instead of $\mathcal{M}(M_{\tau_n}, \mathfrak{s}_{\tau_n}, [c])$.

6.3 Gluing map

Our main objective in this section is to adapt [35, Theorem 3.1.9] to our situation. First we need to define a pregluing map that allows us to compare solutions in the moduli spaces corresponding to the manifolds M_τ and M'_τ . This will then be promoted to an actual gluing map which basically says that once τ becomes sufficiently large the Seiberg–Witten solutions on M_τ are in bijective correspondence with the Seiberg–Witten solutions on M'_τ (the precise statement is Theorem 46).

As can be seen from Figure 7, one should think of the manifolds M_τ as being diffeomorphic versions of the manifold $W^{+}_{\xi',Y}$ described in Figure 2. The moduli space of Seiberg–Witten equations over each of the M_τ gives rise to a “ τ –hybrid” invariant $c(\xi', Y, \tau) \in \check{C}_*(-Y, \mathfrak{s}_\xi)$, but a standard deformation of metrics and perturbations argument which is explained at the end of the paper tells us that in fact they all define the same homology class $c(\xi', Y, \tau) = c(\xi', Y)$, where the right-hand side denotes our original “hybrid” invariant. On the other hand, when we take $Z' = (0, \infty) \times Y$, the resulting manifolds M'_τ agree with $Z^+_{Y,\xi}$ as mentioned at the end of the previous section. Therefore, from the moduli space of Seiberg–Witten equations over M'_τ we obtain the ordinary contact invariant $c(\xi)$ and then the our gluing argument will imply that these two invariants agree.

We write $(M_\tau, g_\tau, \omega_\tau, J_\tau, \sigma_\tau)$ and $(M'_\tau, g'_\tau, \omega'_\tau, J'_\tau, \sigma'_\tau)$ to make explicit the data required in our construction. Notice that on the domains $\{\sigma_\tau \leq \tau\} \subset M_\tau$ and $\{\sigma'_\tau \leq \tau\} \subset M'_\tau$ all the previous structures agree (including the spinor bundles and the canonical solutions). In fact, we can regard these regions as subsets of $Z^+_{Y,\xi}$.

Let (A, Φ) be a solution of the Seiberg–Witten equations on M_τ . We want to transport (A, Φ) into an approximate solution on M'_τ .

First we need to construct a spinor bundle $S'_{(A, \Phi)}$ associated to (A, Φ) on M'_τ . To be more precise, the isomorphism class of the spin-c structure is independent of the solution (A, Φ) that we use, but the particular model will depend on the solution since it will be used to define a transition function.

Lemma 48 in the [appendix](#) states that we can find a compact set C with the following significance: for every τ large enough and for every solution to the Seiberg–Witten equations on M_τ we have $|\alpha| \geq \frac{1}{2}$ on $M_\tau \setminus [(\mathbb{R}^+ \times -Y) \cup C]$ (recall that Φ is equal to (α, β) and is written in [\[35\]](#) instead as (β, γ)). We may write C as $C = [1, T] \times Y \subset Z_{Y, \xi}^+$, where T is large enough and independent of τ and the solution (A, Φ) . *From now on we will assume that τ is chosen so that it is larger than T .*

For $\tau > T$, we construct the spinor bundle $S'_{(A, \Phi)}$ on M'_τ as follows (in [\[35\]](#) this spinor bundle was called $S_{(A, \Phi)}$):

- (1) Over the region $M'_\tau \cap \{\sigma'_\tau \leq \tau\} \subset Z_{Y, \xi}^+$, we use the spinor bundle S_ξ determined by ξ . Over the region $M'_\tau \cap \{\sigma'_\tau \geq T\}$, we use the spinor bundle determined by the almost complex structure J'_τ , ie $S'_{J'_\tau}$. In other words,

$$S'_{(A, \Phi)} = \begin{cases} S_\xi & \text{over } M'_\tau \cap \{\sigma'_\tau \leq \tau\}, \\ S'_{J'_\tau} & \text{over } M'_\tau \cap \{\sigma'_\tau \geq T\}. \end{cases}$$

- (2) To specify what happens over the annulus $\{T \leq \sigma'_\tau \leq \tau\}$ define the map (gauge transformation)³

$$h_{(A, \Phi)}: M_\tau \setminus [(\mathbb{R}^+ \times -Y) \cup C] \rightarrow S^1, \quad h_{(A, \Phi)} = \frac{|\alpha|}{\alpha}.$$

Identifying both S_ξ and $S'_{J'_\tau}$ canonically with $\mathbb{C} \oplus \Lambda_2^+$, the transition map $S_\xi \rightarrow S'_{J'_\tau}$ becomes multiplication by $h_{(A, \Phi)}$.

Notice that if $u \in \mathcal{G}(M_\tau)$ then $h_{u \cdot (A, \Phi)} = u^{-1} h_{(A, \Phi)}$ and so we can easily build an isomorphism

$$u^\#: S'_{(A, \Phi)} \rightarrow S'_{u \cdot (A, \Phi)}.$$

Our next job is to construct a configuration on the spinor bundle $S'_{(A, \Phi)}$ over M'_τ . For this we recall a family of cutoff functions described in [\[35, Section 2.2.1\]](#). Let $\chi(t)$ be

³Here we do not use the notation $h'_{(A, \Phi)}$ that can be found in [\[35\]](#), since our isomorphism h is already canonical and so there is no need to distinguish $h'_{(A, \Phi)}$ from $h_{(A, \Phi)}$.




Figure 8: Defining the spinor bundle $S'_{(A,\Phi)}$ over M'_τ .

a smooth decreasing function such that

$$\chi(t) = \begin{cases} 0 & \text{if } t \geq 1, \\ 1 & \text{if } t \leq 0, \end{cases}$$

and define

$$\chi_\tau(t) = \chi\left(\frac{t-\tau}{N_0} + 1\right) = \begin{cases} 0 & \text{if } t \geq \tau, \\ 1 & \text{if } t \leq \tau - N_0, \end{cases}$$

where N_0 is a number that is fixed later to control the derivatives of χ_τ . With the help of this function define $S'_{(A,\Phi)}$ as follows:

- On the region $M'_\tau \cap \{\sigma'_\tau < \tau\}$ we can identify the structures on M_τ with those on M'_τ and so $(A, \Phi)|_{M_\tau \cap \{\sigma_\tau < \tau\}}$ defines a configuration on $S'_{(A,\Phi)}|_{M'_\tau \cap \{\sigma'_\tau \leq \tau\}}$.
- On the region $M'_\tau \cap \{\sigma'_\tau \geq T\}$ we can write Φ as a pair (α, β) and A as $A = A_{0,\tau} + a$ so if we regard $h_{(A,\Phi)}$ as a gauge transformation (the one that is used to transition on the annulus $M'_\tau \cap \{T \leq \sigma'_\tau \leq \tau\}$) then

$$\begin{aligned} h_{(A,\Phi)} \cdot (A, \Phi) &= h_{(A,\Phi)} \cdot (A_{0,\tau} + a, (\alpha, \beta)) \\ &= \left(A_{0,\tau} + a - \frac{\alpha}{|\alpha|} d\left(\frac{|\alpha|}{\alpha}\right), \left(|\alpha|, \frac{|\alpha|}{\alpha} \beta\right) \right) \\ &\equiv (A_{0,\tau} + \hat{a}, (\hat{\alpha}, \hat{\beta})). \end{aligned}$$

Notice that $\hat{\alpha}$ is a real function with $\hat{\alpha} \geq \frac{1}{2}$. Therefore we define, on $M'_\tau \cap \{\sigma'_\tau \geq T\}$,

$$(A, \Phi)^\# \equiv (A'_{0,\tau} + (\chi_\tau \circ \sigma'_\tau) \hat{a}, (\hat{\alpha}^{\chi_\tau \circ \sigma'_\tau}, (\chi_\tau \circ \sigma'_\tau) \hat{\beta})).$$

- On the end $\{\sigma'_\tau \geq \tau\}$ we set

$$(A, \Phi)^\# = (A'_{0,\tau}, \Phi'_{0,\tau}).$$

Since the construction is compatible with the gauge group action in the sense that

$$u^\# \cdot (A, \Phi)^\# = (u \cdot (A, \Phi))^\#,$$

we have constructed our *pregluing map*

$$(36) \quad \# : \mathcal{M}(M_\tau, \mathfrak{s}_\tau, [\mathfrak{c}]) \rightarrow (\mathcal{C}/\mathcal{G})(M'_\tau).$$

Lemma 51 of the [appendix](#) guarantees the following exponential decay estimate: there is a $\delta > 0$ and a T large enough that for every $N_0 \geq 1$, $k \in \mathbb{N}$, τ satisfying $\tau \geq T + N_0$ and solution (A, Φ) of the Seiberg–Witten equations on M_τ , we have that $(A, \Phi)^\#$ satisfies the Seiberg–Witten equations on $\{\sigma'_\tau \leq T\} \subset M'_\tau$ and

$$(37) \quad |\mathfrak{F}_{\mathfrak{p}_{M'_\tau}}(A, \Phi)^\#|_{C^k(g'_\tau, A^\#)} \leq c_k e^{-\delta \sigma_\tau}$$

on $\{\sigma'_\tau \geq T\} \subset M'_\tau$.

Our objective now is to modify the pregluing map $\# : \mathcal{M}(M_\tau, \mathfrak{s}_\tau, [\mathfrak{c}]) \rightarrow (\mathcal{C}/\mathcal{G})(M'_\tau)$ to obtain a *gluing map* [35, Theorem 3.1.9]

$$\mathfrak{G}_\tau : \mathcal{M}(M_\tau, \mathfrak{s}_\tau, [\mathfrak{c}]) \rightarrow \mathcal{M}(M'_\tau, \mathfrak{s}'_\tau, [\mathfrak{c}]).$$

We want to define $\mathfrak{G}_\tau$ at the level of configuration spaces in such a way that is gauge equivariant. Our proposal is that this map should decompose as

$$(38) \quad \mathfrak{G}_\tau(A, \Phi) = (A, \Phi)^\# + (\mathcal{D}_{(A, \Phi)^\#} \mathfrak{F}_{\mathfrak{p}_{M'_\tau}})^*(b', \psi'),$$

where $(b', \psi') \in L^2_{k,A}(i\mathfrak{su}(S_\tau^{++}) \oplus S_\tau^{--})$ is the quantity that needs to be determined.

Here $\mathcal{D}_{(A, \Phi)^\#} \mathfrak{F}_{\mathfrak{p}_{M'_\tau}}$ denotes the linearization of the perturbed Seiberg–Witten map $\mathfrak{F}_{\mathfrak{p}_{M'_\tau}}$. In the old days of Seiberg–Witten theory, where only the curvature equation was perturbed by some imaginary-valued self-dual two-form, this linearized map $\mathcal{D}_{(A, \Phi)^\#} \mathfrak{F}_{\mathfrak{p}_{M'_\tau}}$ would coincide with the linearization of the *unperturbed* Seiberg–Witten map $\mathcal{D}_{(A, \Phi)^\#} \mathfrak{F}$, since the perturbations were independent of the configuration $(A, \Phi)^\#$ being used. In fact, analyzing the formula (35), we can see that the discrepancy between these two maps is due to the (abstract) perturbations used on the cylindrical end, so to emphasize this point we may sometimes write $\mathcal{D}_{\mathfrak{q}, (A, \Phi)^\#} \mathfrak{F}$ instead of the more precise notation $\mathcal{D}_{(A, \Phi)^\#} \mathfrak{F}_{\mathfrak{p}_{M'_\tau}}$.

Recall that the perturbed Seiberg–Witten equation is

$$\mathfrak{F}_{\mathfrak{p}_{M'_\tau}}(A, \Phi) = \mathfrak{F}(A, \Phi) + \mathfrak{p}_{M'_\tau}(A, \Phi) = 0.$$

By definition, we want $\mathfrak{G}_\tau(A, \Phi)$ to solve the previous equation, which means that

$$\mathfrak{F} \mathfrak{G}_\tau(A, \Phi) + \mathfrak{p}_{M'_\tau} \mathfrak{G}_\tau(A, \Phi) = 0,$$

and implicitly we want to think of the previous equation as depending on (b', ψ') when we write $\mathfrak{G}_\tau(A, \Phi)$ in an explicit way as in (38). In order to write this equation in terms of (b', ψ') , one needs to perform many tedious calculations which are included in the author’s thesis [8, pages 88–96] but not here in order to simplify the exposition.

To explain the end result, it is good to compare our calculation with [35, Equation 3.2]. Following the notation in [35, Section 3], we define first the “perturbed Seiberg–Witten Laplacian”

(39)
$$\Delta_{2,q,(A,\Phi)^\#} = (\mathcal{D}_{(A,\Phi)^\#} \mathfrak{F}_q) \circ (\mathcal{D}_{(A,\Phi)^\#} \mathfrak{F}_q)^*$$

and the quadratic map

$$\mathcal{Q}(a, \phi) = -(\phi \phi^*)_0, \rho(a) \phi).$$

Finally, define the perturbation term

$$P(b', \psi') = \mathfrak{p}_{M'_\tau}(\mathfrak{G}(A, \Phi)) - \mathfrak{p}_{M'_\tau}(A, \Phi)^\# - (\mathcal{D}_{(A,\Phi)} \hat{q}) \circ (\mathcal{D}_{(A,\Phi)^\#} \mathfrak{F}_q)^*.$$

One can then show the following:

Theorem 37 *The configuration $\mathfrak{G}_q(A, \Phi) = (A, \Phi)^\# + (\mathcal{D}_{q,(A,\Phi)^\#} \mathfrak{F}_q)^*(b', \psi')$ is a solution to the perturbed Seiberg–Witten equations $\mathfrak{F}_{\mathfrak{p}_{M'_\tau}} \mathfrak{G}_q(A, \Phi) = 0$ if and only if*

(40)
$$\Delta_{2,q,(A,\Phi)^\#}(b', \psi') + \mathcal{Q} \circ (\mathcal{D}_{(A,\Phi)^\#} \mathfrak{F}_q)^*(b', \psi') + P(b', \psi') = -\mathfrak{F}_{\mathfrak{p}_{M'_\tau}}(A, \Phi)^\#.$$

Notice that the term $P(b', \psi')$ is a new term that does not appear in the usual linearization of the Seiberg–Witten equations. This appears solely because of the presence of the abstract perturbations used in [22]. To solve this equation we will need a sharp version of the contraction mapping theorem.

Namely, the basic idea is to define

$$V_q = \Delta_{2,q,(A,\Phi)^\#}(b', \psi').$$

Our intention is to show that $\Delta_{2,q,(A,\Phi)^\#}$ is invertible, so that if we define

$$S_{q,(A,\Phi)^\#}(V_q) \equiv -\mathcal{Q} \circ [(\mathcal{D}_{(A,\Phi)^\#} \mathfrak{F}_q)^*](\Delta_{2,q,(A,\Phi)^\#}^{-1} V_q) - P(\Delta_{2,q,(A,\Phi)^\#}^{-1} V_q),$$

the gluing equation (40) that we need to solve can be written as

$$V_q = S_{q,(A,\Phi)^\#}(V_q) - \mathfrak{F}_{\mathfrak{p}_{M'_\tau}}(A, \Phi)^\#.$$

The solution of this equation will be guaranteed once we show the hypotheses of [35, Proposition 2.3.5] are satisfied. Therefore, we will show first that $\Delta_{2,q,(A,\Phi)^\#}$ is indeed invertible.

6.4 Invertibility of $\Delta_{2,q,(A,\Phi)^\#}$

In this section we seek a version of [35, Proposition 3.1.2 and Corollary 3.1.6], namely, we want to show that:

Theorem 38 *For each $k \geq 0$ there exists a constant $c_k > 0$ such that for every τ large enough, every $N_0 \geq 1$ and every solution (A, Φ) of the Seiberg–Witten equations on M_τ belonging to the zero-dimensional strata of $\mathcal{M}(M_\tau, \mathfrak{s}_\tau, [c])$, the operator*

$$\begin{aligned} \Delta_{2,q,(A,\Phi)^\#}: L^2_{k+2,A^\#}(M'_\tau, g'_\tau) &\rightarrow L^2_{k,A^\#}(M'_\tau, g'_\tau), \\ (b', \psi') &\mapsto (\mathcal{D}_{q,(A,\Phi)^\#} \mathfrak{F}) \circ (\mathcal{D}_{q,(A,\Phi)^\#} \mathfrak{F})^*(b', \psi'), \end{aligned}$$

is an isomorphism, and moreover, its inverse $\Delta_{2,q,(A,\Phi)^\#}^{-1}$ satisfies for all (b', ψ')

$$c_k \|(b', \psi')\|_{L^2_{k+1}(g'_\tau, A^\#)} \geq \|\Delta_{2,q,(A,\Phi)^\#}^{-1}(b', \psi')\|_{L^2_{k+3}(g'_\tau, A^\#)}.$$

Before proceeding we make a few clarifications:

Remark 39 (a) The norms used for the gluing arguments are gauge-equivariant norms which depend on the configuration $(A, \Phi)^\#$ being used, as can be seen from our use of subscripts in the formulas for the Sobolev spaces.

(b) Our hypothesis regarding the fact that the solution $[(A, \Phi)]$ belongs to the zero-dimensional strata of the moduli space $\mathcal{M}(M_\tau, \mathfrak{s}_\tau, [c])$ has to do with the fact that we will need to find uniform bounds which we will depend (partly) on the norms of these solutions. Since we are using gauge-equivariant norms and the zero-dimensional moduli spaces $\mathcal{M}_0(M_\tau, \mathfrak{s}_\tau, [c])$ are compact, for a fixed τ there can only be finitely many terms to worry about. Clearly, a priori the bounds that we get still depend on the value of τ chosen, but we will see that a transversality argument will help us control these quantities in a way that is τ -independent. It should be pointed out that this assumption regarding the zero-dimensional strata is not that different from the hypothesis used in other gluing arguments. See for example [7, Theorem 4.17] (which uses a compactness assumption as well) or [22, Theorem 18.3.5] (which describes all small solutions of a moduli space).

(c) The strategy that we will use to prove the invertibility of $\Delta_{2,q,(A,\Phi)^\#}$ differs from the one employed in [35] mainly for the following reasons. In that work, the norm $\Delta_{2,q,(A,\Phi)^\#}$ was controlled by first controlling the norm of a different operator, $\square_{(A,\Phi)^\#} = Q_{q,(A,\Phi)} \circ Q_{q,(A,\Phi)}^*$ (defined in the proof of [35, Proposition 3.1.2]), and then relating the norms of these two operators through Equation (3.6) of that paper. However, these norms were only comparable because of that paper's Equation (3.7), which uses the fact that $D_{A^\#}\Phi^\#$ is almost zero. This was true in the case there because the usual Seiberg–Witten equations do not perturb the Dirac equation and since $(A, \Phi)^\#$ is very close to being a solution this means that $\Phi^\#$ is very close to being a harmonic spinor with respect to $D_{A^\#}$. However, the abstract perturbations q used in [22] do modify the Dirac equation, so any clear relationship between $\square_{(A,\Phi)^\#}$ and $D_{A^\#}$ is lost.

The proof of this theorem will follow a splicing argument similar to the one used in [33, Section 4.2.2] (or [7, Section 4.4]). Namely, we will separate the manifold M'_τ into two pieces (see Figure 8) and show the following:

Lemma 40 *We can find two operators*

$$\Delta_{2,q,(A,\Phi),\text{cyl}}^{-1}: L^2(M_\tau) \rightarrow L^2_{2,A}(M_\tau)$$

and

$$(41) \quad \widetilde{\Delta_{2,(A,\Phi),\text{end}}^{-1}}: L^2(N_\tau^+) \rightarrow L^2_{2,\tilde{A}}(N_\tau^+)$$

with the following properties:

(a) $\Delta_{2,q,(A,\Phi),\text{cyl}}^{-1}$ is the inverse to the Seiberg–Witten “Laplacian” defined as in (39),

$$\Delta_{2,q,(A,\Phi)} = (\mathcal{D}_{(A,\Phi)}\mathfrak{F}_q) \circ (\mathcal{D}_{(A,\Phi)}\mathfrak{F}_q)^*: L^2_{2,A}(M_\tau) \rightarrow L^2(M_\tau).$$

(b) Likewise, there is a Seiberg–Witten Laplacian

$$\Delta_{2,(A,\Phi)^\#,\text{end}} = (\mathcal{D}_{(A,\Phi)^\#}\mathfrak{F}) \circ (\mathcal{D}_{(A,\Phi)^\#}\mathfrak{F})^*$$

mapping

$$L^2_{2,A}(M'_\tau \setminus (\mathbb{R}^+ \times -Y)) \rightarrow L^2(M'_\tau \setminus (\mathbb{R}^+ \times -Y))$$

which can be extended to an invertible Seiberg–Witten Laplacian

$$\widetilde{\Delta_{2,(A,\Phi),\text{end}}} = (\widetilde{\mathcal{D}_{(A,\Phi)}}\mathfrak{F}_\eta) \circ (\widetilde{\mathcal{D}_{(A,\Phi)}}\mathfrak{F}_\eta)^*$$

mapping

$$L^2_{2,\tilde{A}}(N_\tau^+) \rightarrow L^2(N_\tau^+),$$

where N_τ^+ is a capped-off version of $M'_\tau \setminus (\mathbb{R}^+ \times -Y)$ defined in the proof.

Remark 41 As we will explain in the proof, $\widetilde{(A, \Phi)}$ denotes an extension of $(A, \Phi)^\#$ to the manifold N_τ^+ which agrees with $(A, \Phi)^\#$ on the region $Z_\tau \setminus ([1, \frac{1}{2}T] \times Y)$. Because of this we will write in the following lemmas $\Delta_{2, \widetilde{(A, \Phi)}, \text{end}}^{-1}$ as $\Delta_{2, (A, \Phi)^\#, \text{end}}^{-1}$.

Also, our construction of $\Delta_{2, \widetilde{(A, \Phi)}, \text{end}}$ is not very sharp since we are using this operator as a proxy for a parametrix argument that will be important in the next section.

Proof We will divide M'_τ into two regions, each of which is the natural location where one of the operators from the statement of the lemma can be considered:

• **The unperturbed region** $(\mathbb{R}^+ \times -Y) \cup [1, T] \times Y$ (construction of $\Delta_{2, q, (A, \Phi), \text{cyl}}^{-1}$) This refers to the region where $(A, \Phi)^\# = (A, \Phi)$, that is, the region where the solution (A, Φ) was not modified. Notice that this includes the cylinder $\mathbb{R}^+ \times -Y$ and the section of the cone $[1, T] \times Y$ and we can use the fact that the moduli space on (M_τ, g_τ) is regular to conclude that $Q_{q, (A, \Phi)}$ is surjective on M_τ [22, Definition 14.5.6]. Using that $Q_{q, (A, \Phi)} = \mathcal{D}_{(A, \Phi)} \mathfrak{F}_q \oplus \mathbf{d}_{(A, \Phi)}^*$ we conclude that $\mathcal{D}_{(A, \Phi)} \mathfrak{F}_q$ must be surjective as well. Now, since $Q_{q, (A, \Phi)}$ is a Fredholm operator we can easily see that $Q_{q, (A, \Phi)}^*$ must be injective. Moreover $Q_{q, (A, \Phi)}^* = (\mathcal{D}_{(A, \Phi)} \mathfrak{F}_q)^* \oplus \mathbf{d}_{(A, \Phi)}$ so $(\mathcal{D}_{(A, \Phi)} \mathfrak{F}_q)^*$ must be injective as well.

In particular, it is not difficult to check that because of this $\Delta_{2, q, (A, \Phi)}$ will be invertible as an operator on M_τ . To emphasize that we care about this operator when applied to sections supported on the unperturbed region (which contains the cylinder) we will write the inverse as a map

$$\Delta_{2, q, (A, \Phi), \text{cyl}}^{-1} \colon L^2(M_\tau) \rightarrow L^2_{2, A}(M_\tau).$$

• **The perturbed region** $M'_\tau \setminus (\mathbb{R}^+ \times -Y)$ (construction of $\Delta_{2, \widetilde{(A, \Phi)}, \text{end}}^{-1}$) This refers to the region where $(A, \Phi)^\#$ and (A, Φ) do not necessarily agree. Notice that this includes the region $[1, \tau] \times Y$ together with remaining piece of the AFAK end Z' . Moreover $\Delta_{2, q, (A, \Phi)^\#} = \Delta_{2, (A, \Phi)^\#}$ on this part of the manifold.

We now follow [35, Section 3.3], which uses a similar idea to what we are about to do, albeit for dealing with the orientability of the moduli spaces.

Recall that we can find an almost complex manifold N with boundary $\partial N = Y$ [17, Lemma 4.4]. We can glue this piece to $M'_\tau \setminus (\mathbb{R}^+ \times -Y)$ along the cone $[1, \tau] \times Y$ to obtain a manifold N_τ^+ without boundary and with an AFAK end (so these are precisely the class of manifolds that were studied in [17]).

In particular, since an almost complex structure now exists globally on N_τ^+ we can talk about the canonical pair $(A'_{0,\tau}, \Phi'_{0,\tau})$ globally, which we will denote as $\widetilde{(A'_{0,\tau}, \Phi'_{0,\tau})}$. This configuration is almost a solution to the Seiberg–Witten equations

$$\begin{aligned}\frac{1}{2}\rho(F_{A'}^+) - (\Phi\Phi^*)_0 &= \frac{1}{2}\rho(F_{\tilde{A}'_{0,\tau}}^+) - (\tilde{\Phi}'_{0,\tau}\tilde{\Phi}'_{0,\tau}{}^*)_0, \\ D_A\Phi &= 0\end{aligned}$$

on N_τ^+ . The only reason why we say almost is that on the almost complex manifold N we attached, the nondegenerate two-form ω_N induced by the metric and almost complex structure on N may not give rise to a symplectic form. In general, there is a torsion (Lee–Gauduchon) form η_N characterized by

$$d\omega_N = \omega_N \wedge \eta.$$

In particular, $\widetilde{(A'_{0,\tau}, \Phi'_{0,\tau})}$ will be a solution to the equations

$$\begin{aligned}\frac{1}{2}\rho(F_{A'}^+) - (\Phi\Phi^*)_0 &= \frac{1}{2}\rho(F_{\tilde{A}'_{0,\tau}}^+) - (\tilde{\Phi}'_{0,\tau}\tilde{\Phi}'_{0,\tau}{}^*)_0, \\ D_A\Phi &= \frac{1}{4}\rho(\eta)\Phi\end{aligned}$$

on N_τ^+ . Associated to these Seiberg–Witten equations there is a corresponding operator $\mathcal{Q}_{\eta,(\tilde{A}'_{0,\tau},\tilde{\Phi}'_{0,\tau})}$. The discussion in [35, Section 3.3.1] in fact shows that the moduli space of solutions is unobstructed at $\widetilde{(A'_{0,\tau}, \Phi'_{0,\tau})}$, that is, both of the operators $\mathcal{Q}_{\eta,(\tilde{A}'_{0,\tau},\tilde{\Phi}'_{0,\tau})}$ and $\mathcal{Q}_{\eta,(\tilde{A}'_{0,\tau},\tilde{\Phi}'_{0,\tau})}^*$ are invertible. Hence, as in the argument from the first part of this lemma we can conclude that the corresponding operator

$$\Delta_{2,\widetilde{(A'_{0,\tau}, \Phi'_{0,\tau})},\text{end}} = (\mathcal{D}_{\widetilde{(A'_{0,\tau}, \Phi'_{0,\tau})}}\mathfrak{F}_\eta) \circ (\mathcal{D}_{\widetilde{(A'_{0,\tau}, \Phi'_{0,\tau})}}\mathfrak{F}_\eta)^*: L^2_{2,\tilde{A}'_{0,\tau}}(N_\tau^+) \rightarrow L^2(N_\tau^+)$$

is invertible.

Now we explain what to do with the other configurations $(A, \Phi)^\#$. We define $\widetilde{(A, \Phi)}$ as an extension to N_τ^+ in such a way that:

- (a) $\widetilde{(A, \Phi)}$ agrees with $\widetilde{(A'_{0,\tau}, \Phi'_{0,\tau})}$ on the region $N \cup ([1, \frac{1}{4}T] \times Y)$.
- (b) $\widetilde{(A, \Phi)}$ agrees with $(A, \Phi)^\#$ on $N_\tau^+ \setminus (N([1, \frac{1}{2}T] \times Y)) = Z_\tau([1, \frac{1}{2}T] \times Y)$.
- (c) $\widetilde{(A, \Phi)}$ interpolates between $\widetilde{(A'_{0,\tau}, \Phi'_{0,\tau})}$ and $(A, \Phi)^\#$ on the remaining region $[\frac{1}{4}T, \frac{1}{2}T] \times Y$.

In particular, since $(A, \Phi)^\#$ also agrees with $\widetilde{(A'_{0,\tau}, \Phi'_{0,\tau})}$ whenever $t > \tau$, we find that $\widetilde{(A, \Phi)}$ differs from $\widetilde{(A'_{0,\tau}, \Phi'_{0,\tau})}$ at most on the region $[\frac{1}{4}T, \tau] \times Y$. Since being an invertible operator is an open condition, if we wanted to show that the corresponding

operator $\Delta_{2,(\tilde{A},\tilde{\Phi}),\text{end}}$ is invertible, it would suffice to show that it is close in the operator norm to $\Delta_{2,(\widetilde{A'_{0,\tau}},\widetilde{\Phi'_{0,\tau}}),\text{end}}$.

In fact, one can compare in a very explicit fashion both operators (we do something very similar in the next lemma for example hence we omit some of the details for now), so we just need to have chosen T from the beginning of our gluing construction in [Section 6.3](#) in such a way that this norm condition is satisfied, so that $\Delta_{2,(\tilde{A},\tilde{\Phi}),\text{end}}$ ends up being invertible (again, a similar argument appears in the next lemma). $\square$

Lemma 42 *For any solution (A, Φ) to the Seiberg–Witten equations on the manifold M_τ , the operator $\Delta_{2,q,(A,\Phi)^\#}$ is a Fredholm operator on M'_τ .*

Proof We will introduce some cutoff functions that will allow us to follow the usual splicing arguments: these will be denoted by η_{cyl} and η_{end} . They satisfy the following properties:

- $0 \leq \eta_{\text{cyl}}, \eta_{\text{end}} \leq 1$ and $\eta_{\text{cyl}}^2 + \eta_{\text{end}}^2 = 1$.
- η_{cyl} is supported on the unperturbed region. Moreover, $\eta_{\text{cyl}} \equiv 1$ on a small neighborhood of the region $\mathbb{R}^+ \times -Y \cup [1, \frac{1}{2}T] \times Y$. In particular, *the gradient of η_{cyl} is supported on the fixed region $[1, T] \times Y$.*
- η_{end} is supported on the perturbed region. Moreover, $\eta_{\text{end}} \equiv 1$ on a small neighborhood of $([T, \tau] \times Y) \cup \{Z' \cap \{\sigma_{Z'} > 1/\tau\}\}$. In particular, *the gradient of η_{end} is supported on the fixed region $[1, T] \times Y$.* Notice that also η_{end} will vanish in a small neighborhood of $\{\frac{1}{2}T\} \times Y$, since η_{cyl} is equal to 1 near that slice.
- For $\eta = \eta_{\text{cyl}}, \eta_{\text{end}}$ we have $|\nabla^n \eta| \leq (2/T)^n$.

Our proto-inverse will be the operator

$$\begin{aligned} & \tilde{\Delta}_{2,q,(A,\Phi)^\#}^{-1}: L^2(M'_\tau, g'_\tau) \rightarrow L^2_2(M'_\tau, g'_\tau), \\ (b', \psi') & \mapsto \eta_{\text{cyl}} \Delta_{2,q,(A,\Phi),\text{cyl}}^{-1}[\eta_{\text{cyl}}(b', \psi')] + \eta_{\text{end}} \Delta_{2,(A,\Phi)^\#, \text{end}}^{-1}[\eta_{\text{end}}(b', \psi')]. \end{aligned}$$

We will see that $\tilde{\Delta}_{2,q,(A,\Phi)^\#}^{-1}$ provides a parametrix for $\Delta_{2,q,(A,\Phi)^\#}$: this is because

$$\begin{aligned} & \Delta_{2,q,(A,\Phi)^\#}[\tilde{\Delta}_{2,q,(A,\Phi)^\#}^{-1}(b', \psi')] \\ &= \Delta_{2,q,(A,\Phi)^\#}[\eta_{\text{cyl}} \Delta_{2,q,(A,\Phi),\text{cyl}}^{-1}[\eta_{\text{cyl}}(b', \psi')] + \eta_{\text{end}} \Delta_{2,(A,\Phi)^\#, \text{end}}^{-1}[\eta_{\text{end}}(b', \psi')]], \end{aligned}$$

which equals

$$\begin{aligned} & \{ \mathcal{D}_{(A,\Phi)} \eta_{\text{cyl}}, \Delta_{2,q,(A,\Phi),\text{cyl}}^{-1} [\eta_{\text{cyl}}(b', \psi')] \} + \eta_{\text{cyl}}^2(b', \psi') \\ & + \{ \mathcal{D}_{(A,\Phi)} \eta_{\text{end}}, \Delta_{2,(A,\Phi)^\#, \text{end}}^{-1} [\eta_{\text{end}}(b', \psi')] \} + \eta_{\text{end}}^2(b', \psi') \\ & + \{ \mathcal{D}_{(A,\Phi)} \eta_{\text{cyl}}, \mathcal{D}_{(A,\Phi)} (\Delta_{2,q,(A,\Phi),\text{cyl}}^{-1} [\eta_{\text{cyl}}(b', \psi')]) \} \\ & + \{ \mathcal{D}_{(A,\Phi)} \eta_{\text{end}}, \mathcal{D}_{(A,\Phi)} (\Delta_{2,(A,\Phi)^\#, \text{end}}^{-1} [\eta_{\text{end}}(b', \psi')]) \} \\ & = \{ \mathcal{D}_{(A,\Phi)} \eta_{\text{cyl}}, \Delta_{2,q,(A,\Phi),\text{cyl}}^{-1} [\eta_{\text{cyl}}(b', \psi')] \} \\ & + \{ \mathcal{D}_{(A,\Phi)} \eta_{\text{end}}, \Delta_{2,(A,\Phi)^\#, \text{end}}^{-1} [\eta_{\text{end}}(b', \psi')] \} + (b', \psi') \\ & + \{ \mathcal{D}_{(A,\Phi)} \eta_{\text{cyl}}, \mathcal{D}_{(A,\Phi)} (\Delta_{2,q,(A,\Phi),\text{cyl}}^{-1} [\eta_{\text{cyl}}(b', \psi')]) \} \\ & + \{ \mathcal{D}_{(A,\Phi)} \eta_{\text{end}}, \mathcal{D}_{(A,\Phi)} (\Delta_{2,(A,\Phi)^\#, \text{end}}^{-1} [\eta_{\text{end}}(b', \psi')]) \}. \end{aligned}$$

Here the notation $\{ \cdot, \cdot \}$ is used to indicate a bilinear pointwise multiplication between some (higher order) derivatives of the cutoff functions and the elements in the domain. Also, the notation $\mathcal{D}_{(A,\Phi)} \eta_{\text{cyl}}$ means that this expression involves (higher order) derivatives of the perturbation (and a priori the configuration (A, Φ)) but their precise form is not important to us.

Notice that the first and last two terms are supported on the compact subset $[1, T] \times Y$, where $(A, \Phi)^\# = (A, \Phi)$. Also, we dropped the dependence on q for the derivatives $\mathcal{D}_{(A,\Phi)} \eta_\bullet$ since this perturbation affects only the cylindrical region. To analyze if there is any dependence of $\mathcal{D}_{(A,\Phi)} \eta_\bullet$ on (A, Φ) , we will study $\mathcal{D}_{(A,\Phi)} \eta_{\text{cyl}}$ since the other case is exactly the same. We need to compute

(42)
$$\Delta_{2,q,(A,\Phi)^\#}(\eta_{\text{cyl}}(b_{\text{cyl}}, \psi_{\text{cyl}})),$$

where we defined

$$(b_{\text{cyl}}, \psi_{\text{cyl}}) \equiv \Delta_{2,q,(A,\Phi),\text{cyl}}^{-1} [\eta_{\text{cyl}}(b', \psi')].$$

Notice that we may write

$$\Delta_{2,q,(A,\Phi)^\#} = (\mathcal{D}_{(A,\Phi)^\#} \mathfrak{F}_q) \circ (\mathcal{D}_{(A,\Phi)^\#} \mathfrak{F}_q)^* = (\mathcal{D}_{(A,\Phi)^\#} \mathfrak{F}) \circ (\mathcal{D}_{(A,\Phi)^\#} \mathfrak{F})^* = \Delta_{2,(A,\Phi)^\#},$$

since we are only interested in computing (42) on the region $[1, T] \times Y$, where η_{cyl} is not constant. The advantage of using this unperturbed Seiberg–Witten “Laplacian” is that we can give an explicit formula for it based on (18) and (19). We find that for

arbitrary (b', ψ') ,

$$\begin{aligned} & \Delta_{2, (A, \Phi)^\#}(b', \psi') \\ &= (\mathcal{D}_{(A, \Phi)^\#} \mathfrak{F})((d^+)^* \rho^* b' + \rho^*(\psi'(\Phi^\#)^*), D_{A^\#}^* \psi' - b' \Phi^\#) \\ &= \left(\rho(d^+[(d^+)^* \rho^* b' + \rho^*(\psi'(\Phi^\#)^*)]) \right. \\ & \quad \left. - \{ \Phi^\#(D_{A^\#}^* \psi' - b' \Phi^\#)^* + (D_{A^\#}^* \psi' - b' \Phi^\#)(\Phi^\#)^* \}_0, \right. \\ & \quad \left. D_{A^\#}(D_{A^\#}^* \psi' - b' \Phi^\#) + \rho((d^+)^* \rho^* b' + \rho^*(\psi'(\Phi^\#)^*)) \Phi^\# \right), \end{aligned}$$

and therefore $\Delta_{2, (A, \Phi)^\#}[(\eta_{\text{cyl}} b_{\text{cyl}}, \eta_{\text{cyl}} \psi_{\text{cyl}})]$ becomes

$$\begin{aligned} & \rho(d^+[(d^+)^* \rho^*(\eta_{\text{cyl}} b_{\text{cyl}}) + \rho^*(\eta_{\text{cyl}} \psi_{\text{cyl}}(\Phi^\#)^*)]) \\ & \quad - \{ \Phi^\#(D_{A^\#}^*(\eta_{\text{cyl}} \psi_{\text{cyl}}) - (\eta_{\text{cyl}} b_{\text{cyl}}) \Phi^\#)^* + (D_{A^\#}^*(\eta_{\text{cyl}} \psi_{\text{cyl}}) - (\eta_{\text{cyl}} b_{\text{cyl}}) \Phi^\#)(\Phi^\#)^* \}_0 \\ & \quad + \rho(d^+[(d^+)^* \rho^*(\eta_{\text{cyl}} b_{\text{cyl}}) + \rho^*((\eta_{\text{cyl}} \psi_{\text{cyl}})(\Phi^\#)^*)]) \\ & \quad - \{ \Phi^\#(D_{A^\#}^*(\eta_{\text{cyl}} \psi_{\text{cyl}}) - (\eta_{\text{cyl}} b_{\text{cyl}}) \Phi^\#)^* + (D_{A^\#}^*(\eta_{\text{cyl}} \psi_{\text{cyl}}) - (\eta_{\text{cyl}} b_{\text{cyl}}) \Phi^\#)(\Phi^\#)^* \}. \end{aligned}$$

Since the Dirac operator D satisfies the Leibniz rule [1, Proposition 3.38]

$$D(\eta\psi) = \rho(d\eta)\psi + \eta D\psi$$

the only derivatives of η_{cyl} that appear are those involving its exterior derivative, which is independent of the configuration $(A, \Phi)^\#$ that is used. A similar story is true for η_{end} . Since $\Delta_{2, q, (A, \Phi), \text{cyl}}^{-1}[\eta_{\text{cyl}}(b', \psi')]$ and $\Delta_{2, (A, \Phi)^\#, \text{end}}^{-1}[\eta_{\text{end}}(b', \psi')]$ are elements of $L_{2, A^\#}^2(M'_\tau)$, our previous discussion in fact tells us that the operator

$$K_{(A, \Phi)}: L^2([1, T] \times Y) \rightarrow L^2([1, T] \times Y)$$

that maps (b', ψ') to

$$\begin{aligned} & \{ \mathcal{D}\eta_{\text{cyl}}, \Delta_{2, q, (A, \Phi), \text{cyl}}^{-1}[\eta_{\text{cyl}}(b', \psi')] \} + \{ \mathcal{D}\eta_{\text{end}}, \Delta_{2, (A, \Phi)^\#, \text{end}}^{-1}[\eta_{\text{end}}(b', \psi')] \} \\ & \quad + \{ \mathcal{D}_{(A, \Phi)} \eta_{\text{cyl}}, \mathcal{D}_{(A, \Phi)}(\Delta_{2, q, (A, \Phi), \text{cyl}}^{-1}[\eta_{\text{cyl}}(b', \psi')]) \} \\ & \quad + \{ \mathcal{D}_{(A, \Phi)} \eta_{\text{end}}, \mathcal{D}_{(A, \Phi)}(\Delta_{2, (A, \Phi)^\#, \text{end}}^{-1}[\eta_{\text{end}}(b', \psi')]) \} \end{aligned}$$

can in fact be regarded as an operator

$$K_{(A, \Phi)}: L^2([1, T] \times Y) \rightarrow L^2([1, T] \times Y)$$

and using the compact inclusion $L^2([1, T] \times Y) \hookrightarrow L^2([1, T] \times Y)$ on a compact manifold we conclude that $K_{(A, \Phi)}$ is a compact operator. This provides the desired parametrix. $\square$

The next lemma addresses the bounds on the norms of this parametrix.

Lemma 43 *The norms of the parametrices can be chosen in a uniform way. That is, there is a constant C_T such that for all solutions (A, Φ) we have $\|K_{(A,\Phi)}\| \leq C_T/T$. In fact, one can find a constant C_∞ , independent of T , such that $\|K_{(A,\Phi)}\| \leq C_\infty/T$.*

Proof It is possible to show that the norms of the parametrices are uniform, that is, that there is a constant C_T such that for all solutions (A, Φ) we have $\|K_{(A,\Phi)}\| \leq C_T/T$. In fact, we will show something better, which is that one could have chosen a constant C_∞ which is independent of T ; in other words, $\|K_{(A,\Phi)}\| \leq C_\infty/T$.

Notice that a priori the only terms that may not seem controllable in terms of T are

(43)

$$\begin{aligned} &\{\mathcal{D}\eta_{\text{cyl}}, \Delta_{2,q,(A,\Phi),\text{cyl}}^{-1}[\eta_{\text{cyl}}(b', \psi')]\}, \\ &\{\mathcal{D}_{(A,\Phi)}\eta_{\text{cyl}}, \mathcal{D}_{(A,\Phi)}(\Delta_{2,q,(A,\Phi),\text{cyl}}^{-1}[\eta_{\text{cyl}}(b', \psi')])\}. \end{aligned}$$

We will discuss how to control the first term in (43) since the second is exactly the same. If we take a sequence of solutions (A_n, Φ_n) on M_{τ_n} then on $[1, T]$ it will converge strongly to a solution (A_∞, Φ_∞) on $Z_{Y,\xi}^+$ (this is due to Lemma 53 in the appendix, based on the compactness theorem of [35, Theorem 2.2.11]) and hence for all (b', ψ')

$$\{\mathcal{D}\eta_{\text{cyl}}, \Delta_{2,q,(A_n,\Phi_n),\text{cyl}}^{-1}[\eta_{\text{cyl}}(b', \psi')]\}$$

converges to

$$\{\mathcal{D}\eta_{\text{cyl}}, \Delta_{2,q,(A_\infty,\Phi_\infty),\text{cyl}}^{-1}[\eta_{\text{cyl}}(b', \psi')]\}.$$

It is clear then that it would be enough to have a uniform bound on the operator norms

$$\|\Delta_{2,q,(A_\infty,\Phi_\infty)}^{-1}\|_{L^2(Z_{Y,\xi}^+) \rightarrow L^2_{2,A_\infty}(Z_{Y,\xi}^+)}.$$

As we will make more explicit in the next proof, since we are taking a sequence of solutions (A_n, Φ_n) which belong to the zero-dimensional strata of the moduli spaces $\mathcal{M}(M_{\tau_n}, \mathfrak{s}_{\tau_n}, [\mathfrak{c}])$, the limiting solution (A_∞, Φ_∞) must belong to the zero-dimensional strata of $\mathcal{M}(Z_{Y,\xi}^+, \mathfrak{s}, [\mathfrak{c}])$, and since we are using gauge-equivariant norms, there are only finitely many values the previous operator norm can take (this is related to the second point in the remarks we made after stating the invertibility of the Laplacian). Therefore, we will have the uniform bound for the operator $K_{(A,\Phi)}$, that is, $\|K_{(A,\Phi)}\| \leq C_\infty/T$, where C_∞ is independent of τ , T and the solutions (A, Φ) used.

Therefore, there is no loss of generality in assuming that T was chosen from the beginning so that it would also satisfy the condition

$$\|K_{(A,\Phi)}\|_{L^2([1,T]\times Y) \rightarrow L^2_2([1,T]\times Y)} \leq \frac{C_\infty}{T} \leq \frac{1}{2}$$

for all the solutions of the Seiberg–Witten equations on M_τ . In particular, from the identity

$$\Delta_{2,q,(A,\Phi)^\#}[\tilde{\Delta}_{2,q,(A,\Phi)^\#}^{-1}(b', \psi')] = K_{(A,\Phi)}(b', \psi') + (b', \psi'),$$

we see that the operator norms satisfy

$$\|\Delta_{2,q,(A,\Phi)^\#} \tilde{\Delta}_{2,q,(A,\Phi)^\#}^{-1} - \text{Id}\|_{L^2_{2,A^\#}(M'_\tau, g'_\tau) \rightarrow L^2(M'_\tau, g_{\tau'})} \leq \frac{1}{2}.$$

In particular we conclude that each $\Delta_{2,q,(A,\Phi)^\#} \tilde{\Delta}_{2,q,(A,\Phi)^\#}^{-1}$ is invertible and they (and their inverses) are uniformly bounded since

$$\frac{1}{2} \leq \|\Delta_{2,q,(A,\Phi)^\#} \tilde{\Delta}_{2,q,(A,\Phi)^\#}^{-1}\|_{L^2_{2,A^\#}(M'_\tau, g'_\tau) \rightarrow L^2(M'_\tau, g_{\tau'})} \leq \frac{3}{2}.$$

Therefore the inverse of $\Delta_{2,q,(A,\Phi)^\#}$ is $\tilde{\Delta}_{2,q,(A,\Phi)^\#}^{-1}(\Delta_{2,q,(A,\Phi)^\#} \tilde{\Delta}_{2,q,(A,\Phi)^\#}^{-1})^{-1}$. $\square$

Returning to our proof of [Theorem 38](#), since $(\Delta_{2,q,(A,\Phi)^\#} \tilde{\Delta}_{2,q,(A,\Phi)^\#}^{-1})^{-1}$ is uniformly bounded we just need to check that $\tilde{\Delta}_{2,q,(A,\Phi)^\#}^{-1}$ is uniformly bounded to conclude that $\Delta_{2,q,(A,\Phi)^\#}$ is uniformly bounded (a similar argument would work to give uniform bounds on $\Delta_{2,q,(A,\Phi)^\#}^{-1}$). Looking at the definition of $\tilde{\Delta}_{2,q,(A,\Phi)^\#}^{-1}$ it becomes clear that it suffices to show that $\eta_{\text{cyl}} \Delta_{2,q,(A,\Phi)^\#}^{-1}[\eta_{\text{cyl}}(b', \psi')]$ is uniformly bounded.

Here we will use again the assumption we mentioned at the end of the previous proof. Namely, we are now assuming that the gauge-equivalence classes of our solutions $(A, \Phi) \in \mathcal{M}(M_\tau, \mathfrak{s}_\tau, [\mathfrak{c}])$ all belong to the zero-dimensional strata of the moduli spaces. Since the Laplacians are gauge equivariant in the sense that

$$\Delta_{2,q,u \cdot (A,\Phi)}[u \cdot (b, \psi)] = u \cdot \Delta_{2,q,(A,\Phi)}(b, \psi)$$

and we are using the gauge-equivariant norms $\|\cdot\|_{L^2_{k,A}}$, for each τ there are only finitely many gauge-equivalence classes we need to worry about, which immediately implies that for each τ we have a control on the Laplacians (and their inverses). Clearly we still need to see what happens as we vary τ .

Let K be a subset of $(\mathbb{R}^+ \times -Y) \cup ([1, \infty) \times Y)$ and use $\|\Delta_{2,q,(A,\Phi)}\|_{A,K}$ or $\|\Delta_{2,q,(A,\Phi)}^{-1}\|_{A,K}$ to denote the operator norms of $\Delta_{2,q,(A,\Phi)}$ and $\Delta_{2,q,(A,\Phi)}^{-1}$ when restricted to sections supported on K . Clearly if $K \subset K'$ then $\|\Delta_{2,q,(A,\Phi)}^{-1}\|_{A,K} \leq \|\Delta_{2,q,(A,\Phi)}^{-1}\|_{A,K'}$.

Now, recall that we are actually working with a sequence τ_n increasing to ∞ , so for each τ_n let $[(A_n, \Phi_n)] \in \mathcal{M}_0(M_{\tau_n}, \mathfrak{s}_{\tau_n}, [\mathfrak{c}])$ be a (gauge-equivalence class of) solution

belonging to the zero-dimensional strata. Notice that each compact subset $K \subset (\mathbb{R}^+ \times -Y) \cup ([1, \infty) \times Y)$ eventually belongs to all M_{τ_n} (once τ_n is sufficiently large) so the compactness theorem in this case says that we can choose representatives (A_n, Φ_n) which converge to a solution (A_∞, Φ_∞) which solves the equations on $Z_{\xi, Y}^+$ and this convergence is strong when restricted to the compact subset K . In particular, it is clear from this that

(44)

$$\|\Delta_{2,q,(A_n,\Phi_n)}^{-1}\|_{A_n,K} \rightarrow \|\Delta_{2,q,(A_\infty,\Phi_\infty)}^{-1}\|_{A_\infty,K}.$$

In fact, we must also have that the limiting solution (A_∞, Φ_∞) belongs to the zero-dimensional strata because the different strata are labeled by the index of the operator $Q_{q,(A,\Phi)}$ and this index can only decrease (this is how the broken trajectories appear). However, since the index of each element in the sequence was already zero, the index of the limiting configuration would need to be negative if it were to decrease but transversality rules this out, since we do not have negative-dimensional moduli spaces. Therefore the convergence is without broken trajectories, that is, $[(A_\infty, \Phi_\infty)] \in \mathcal{M}_0(Z_{Y,\xi}^+, \mathfrak{s}, [c])$. In particular, the fact that no energy is lost along the half-cylinder allows us to improve the convergence in (44) to (we will say more about this in a moment)

(45)

$$\|\Delta_{2,q,(A_n,\Phi_n)}^{-1}\|_{A_n,K_t} \rightarrow \|\Delta_{2,q,(A_\infty,\Phi_\infty)}^{-1}\|_{A_\infty,K_t},$$

where now $K_t = (\mathbb{R}^+ \times -Y) \cup ([1, t] \times Y)$ ($t > 1$ is arbitrary). In particular,

$$\|\Delta_{2,q,(A_\infty,\Phi_\infty)}^{-1}\|_{A_\infty,K_t} \leq \|\Delta_{2,q,(A_\infty,\Phi_\infty)}^{-1}\|_{A_\infty,Z_{Y,\xi}^+} \leq C,$$

where

$$C = \max\{\|\Delta_{2,q,(A_\infty,\Phi_\infty)}^{-1}\|_{A_\infty,Z_{Y,\xi}^+} : [(A_\infty, \Phi_\infty)] \in \mathcal{M}_0(Z_{Y,\xi}^+, \mathfrak{s}, [c])\}.$$

Since t and the sequence were both arbitrary this clearly gives us the uniform bound that we were after so we have proven [Theorem 38](#).

We will now say more about why the convergence (45) is true. For this we need to recall that thanks to the fiber product description of our moduli spaces, we can restrict each solution $[(A_n, \Phi_n)]$ to a solution on the cylindrical-end moduli space $\mathcal{M}(\mathbb{R}^+ \times -Y, \mathfrak{s}_\xi, [c])$, which we will denote as $[(A_n, \Phi_n)]_{\text{cyl}} \in \mathcal{M}(\mathbb{R}^+ \times -Y, \mathfrak{s}_\xi, [c])$.

Likewise, the limiting solution $[(A_\infty, \Phi_\infty)]$ can also be restricted to this moduli space so we have as well that $[(A_\infty, \Phi_\infty)]_{\text{cyl}} \in \mathcal{M}(\mathbb{R}^+ \times -Y, \mathfrak{s}_\xi, [c])$. When we described the configuration spaces at the beginning of the paper we used the topology of strong convergence on compact subsets $L_{k,\text{loc}}^2$ to define the moduli spaces. However, as explained in [\[22, Theorem 13.3.5\]](#), the same moduli space $\mathcal{M}(\mathbb{R}^+ \times -Y, \mathfrak{s}_\xi, [c])$ can

also be obtained if we had used the stronger topology of L^2_k convergence along the entire half-cylinder $\mathbb{R}^+ \times -Y$ (the authors of [22] really did this for the moduli space on the cylinder $\mathbb{R} \times Y$ but it does not affect our claim). Therefore, the convergence of $[(A_n, \Phi_n)]_{\text{cyl}}$ towards $[(A_\infty, \Phi_\infty)]_{\text{cyl}}$ can be regarded as a strong convergence with respect to the L^2_{k,A_c} norm, where A_c represents the translation-invariant connection associated to a smooth representative $\mathfrak{c}$ of the critical point $[c]$. In other words, we can choose representatives of $[(A_n, \Phi_n)]_{\text{cyl}}$ and $[(A_\infty, \Phi_\infty)]_{\text{cyl}}$ so that

$$A_n = A_c + a_n, \quad A_\infty = A_c + a_\infty, \quad \Phi_n = \Phi_c + \phi_n, \quad \Phi_\infty = \Phi_c + \phi_\infty,$$

where Φ_c is a translation-invariant representative of $\mathfrak{c}$ and we have that

$$\begin{aligned} \lim_{n \rightarrow \infty} \|A_n - A_\infty\|_{L^2_k(\mathbb{R}^+ \times -Y)} &= \lim_{n \rightarrow \infty} \|a_n - a_\infty\|_{L^2_k(\mathbb{R}^+ \times -Y)} = 0, \\ \lim_{n \rightarrow \infty} \|\Phi_n - \Phi_\infty\|_{L^2_{k,A_c}(\mathbb{R}^+ \times -Y)} &= \lim_{n \rightarrow \infty} \|\phi_n - \phi_\infty\|_{L^2_{k,A_c}(\mathbb{R}^+ \times -Y)} = 0. \end{aligned}$$

The norms $\|\cdot\|_{L^2_{k,A_n}}$ and $\|\cdot\|_{L^2_{k,A_c}}$ can now be compared thanks to the Sobolev multiplication theorems (since for example $\nabla_{A_n} \bullet = \nabla_{A_\infty} \bullet + (a_n - a_\infty) \otimes \bullet$ with similar formulas for the higher derivatives) and the previous limits make it clear that the operator norm convergence (44) on compact subsets K can be improved to the operator norm convergence (45) on sets of the form half-cylinder + compact.

In the next section we explain the properties of the gluing map one obtains using the invertibility of the Laplacian.

Remark 44 Many of the following arguments will have a similar structure to the one before. Namely, because we are taking solutions belonging to the zero-dimensional strata for an individual τ we will find a bound, but a priori this may depend on τ . However, as we take τ_n sufficiently large the bounds end up being controlled by a limiting case $\tau = \infty$, since we can invoke the strong convergence on the half-cylindrical end. Since the arguments are essentially the same in each case we will not repeat the strategy and will instead just say that it follows by similar arguments.

6.5 Definition and some properties of the gluing map

As explained before, if we write $V_q = \Delta_{2,q,(A,\Phi)^\#}(b', \psi')$ then solving the gluing equation (40) is equivalent to solving the equation

$$V_q = S_{q,(A,\Phi)^\#}(V_q) - \mathfrak{F}_{p_{M'_t}}(A, \Phi)^\#.$$

The solution of this equation requires an application of the contraction mapping theorem,

which requires us to first show that the map S_q is a uniform contraction in the following sense (this is an analogue of [35, Lemma 3.1.8]):

Theorem 45 *For every k large enough there exist constants $\alpha_k > 0$ and $\kappa_k \in (0, \frac{1}{2})$ such that for every τ large enough, every $N_0 \geq 1$ and every approximate solution of the Seiberg–Witten equations $(A, \Phi)^\#$ on M'_τ which comes from an actual solution (A, Φ) on M_τ whose gauge-equivalence class $[A, \Phi]$ belongs to the zero-dimensional strata of the moduli space $\mathcal{M}_0(M_\tau; \mathfrak{s}_\tau; [c])$, we have for all*

$$V_1, V_2 \in L^2_k(M'_\tau; i\mathfrak{su}(S^+) \oplus S^-, g'_\tau; A^\#),$$

if

$$\|V_1\|_{L^2_k(g'_\tau, A^\#)}, \|V_2\|_{L^2_k(g'_\tau, A^\#)} \leq \alpha_k,$$

then

$$\|S_{q, (A, \Phi)^\#}(V_2) - S_{q, (A, \Phi)^\#}(V_1)\|_{L^2_k(g'_\tau, A^\#)} \leq \kappa_k \|V_2 - V_1\|_{L^2_k(g'_\tau, A^\#)}.$$

Proof Recall that

$$S_{q, (A, \Phi)^\#}(V_q) = -\mathcal{Q} \circ [(\mathcal{D}_{(A, \Phi)^\#} \mathfrak{F}_q)^*](\Delta_{2, q, (A, \Phi)^\#}^{-1} V_q) - P(\Delta_{2, q, (A, \Phi)^\#}^{-1} V_q)$$

We will mention the main differences compared with the proof given in [35]. First of all, we need the bounds in [22, Proposition 11.4.1], which say that for $k \geq 2$

(46)
$$\|\mathcal{D}^l_{(A, \Phi)} \hat{q}\| \leq C(1 + \|a\|_{L^2_k(Z)})^{2k(l+1)}(1 + \|\Phi\|_{L^2_{k, A}(Z)})^{l+1}.$$

Here C is a constant independent of the configuration and in this theorem Z denotes a finite cylinder, while $A = A_{\text{ref}} + a \otimes 1$ for some reference configuration A_{ref} . Note that these bounds can be used on the half-cylinder $\mathbb{R}^+ \times -Y$ as well. Simply decompose it as

$$\mathbb{R}^+ \times -Y = \bigcup_{n \geq 0} \underbrace{[n, n+1] \times -Y}_{Z_n}.$$

If $\bullet$ denotes an element in the domain of $\mathcal{D}^l_{(A, \Phi)} \hat{q}$ then we have

$$\begin{aligned} \|\mathcal{D}^l_{(A, \Phi)} \hat{q}(\bullet)\|_{\mathbb{R}^+ \times -Y} &= \sum_{n=0}^\infty \|\mathcal{D}^l_{(A, \Phi)} \hat{q}(\bullet)\|_{Z_n} \\ &\leq C \sum_{n=0}^\infty (1 + \|a\|_{L^2_k(Z_n)})^{2k(l+1)} (1 + \|\Phi\|_{L^2_{k, A}(Z_n)})^{l+1} \|\bullet\|_{Z_n}, \end{aligned}$$

where in the last step we used the bounds coming from the operator norm (46). If we define

$$C_{n, (A, \Phi)} = (1 + \|a\|_{L^2_k(Z_n)})^{2k(l+1)} (1 + \|\Phi\|_{L^2_{k, A}(Z_n)})^{l+1},$$

then it is not too difficult to see that

$$C_{\max, (A, \Phi)} = \max_n C_{n, (A, \Phi)} < \infty.$$

One reason for this is that the previous quantities $C_{n, (A, \Phi)}$ do not differ too much from those for the translation-invariant solution $C_{n, (A_c, \Phi_c)}$, which are independent of n .

To compare $C_{n, (A, \Phi)}$ with $C_{n, (A_c, \Phi_c)}$, take our reference connection to be A_c and use the fact in this case $a = A - A_c$ can be chosen to be exponentially decaying as well as $\Phi - \Phi_c$, since (A, Φ) is asymptotic to (A_c, Φ_c) ; ([22, Section 13.5 and Proposition 13.6.1] or the [appendix](#)). Hence a term like

$$(1 + \|\Phi\|_{L^2_{k, A}(Z_n)})^{l+1}$$

can be bounded by

$$(1 + \|\Phi - \Phi_c\|_{L^2_{k, A}(Z_n)} + \|\Phi_c\|_{L^2_{k, A}(Z_n)})^{l+1},$$

where the term $\|\Phi - \Phi_c\|_{L^2_{k, A}(Z_n)}$ contributes less as n increases, say less than $1/n^2$ for sufficiently large n .

In any case we end up with

$$\|\mathcal{D}^l_{(A, \Phi)} \hat{q}(\bullet)\|_{\mathbb{R} + \times - Y} \leq C C_{\max, (A, \Phi)} \sum_{n=0}^{\infty} \|\bullet\|_{Z_n} = C C_{\max, (A, \Phi)} \|\bullet\|_{\mathbb{R} + \times - Y}.$$

Since $\bullet$ was arbitrary this says that each $\mathcal{D}^l_{(A, \Phi)} \hat{q}$ is a bounded operator on the half-cylinder. For each τ , we are only dealing with finitely many gauge-equivalence classes of solutions because of our assumption on the strata so the bounds are once again controlled for a fixed τ . By analogous arguments, one can find bounds which actually become independent of τ , so that $\|\mathcal{D}^l_{(A, \Phi)} \hat{q}\| \leq C_l$ for some constant C_l on the half-infinite cylinder.

The other ingredient is that the leading term of $P(\Delta^{-1}_{2, q, (A, \Phi)^{\#}} V_q)$ is quadratic in the following sense. To emphasize its dependence on V , we will write $P(\Delta^{-1}_{2, q, (A, \Phi)^{\#}} V_q)$ as

$$\begin{aligned} f(V) &= q((A, \Phi)^{\#} + (\mathcal{D}_{(A, \Phi)^{\#}} \mathfrak{F}_q)^* \Delta^{-1}_{2, q, (A, \Phi)^{\#}} V) - q(A, \Phi)^{\#} \\ &\quad - (\mathcal{D}_{(A, \Phi)^{\#}} q) \circ (\mathcal{D}_{(A, \Phi)^{\#}} \mathfrak{F}_q)^* \Delta^{-1}_{2, q, (A, \Phi)^{\#}} V. \end{aligned}$$

We want to compute $f'(V)$ and $f''(V)$, that is, the Banach space derivatives with respect to V . For this define the functions

$$\begin{aligned} f_1(V) &= q((A, \Phi)^{\#} + (\mathcal{D}_{(A, \Phi)^{\#}} \mathfrak{F}_q)^* \Delta^{-1}_{2, q, (A, \Phi)^{\#}} V) - q(A, \Phi)^{\#}, \\ f_2(V) &= (\mathcal{D}_{(A, \Phi)^{\#}} q) \circ (\mathcal{D}_{(A, \Phi)^{\#}} \mathfrak{F}_q)^* \Delta^{-1}_{2, q, (A, \Phi)^{\#}} V, \end{aligned}$$

so that $f(V) = f_1(V) - f_2(V)$. Since $f_2(V)$ is linear in V it is easy to determine that

$$f_2'(V) = (\mathcal{D}_{(A,\Phi)^\#} \mathfrak{q}) \circ (\mathcal{D}_{(A,\Phi)^\#} \mathfrak{F}_q)^* \Delta_{2,q,(A,\Phi)^\#}^{-1}(V).$$

Clearly f_2' is independent as a linear transformation of the “basepoint” (which is hidden in our notation) so we will have that $f_2^{(n)} = 0$ for $n \geq 2$. To compute the derivative of $f_1(V)$ think of the Taylor expansion of $\mathfrak{q}$ about $(A, \Phi)^\#$ (which plays the role of 0 in our affine space interpretation for the domain of $\mathfrak{q}$, so we can use [29, Corollary 4.4 in Chapter 1]). In this way

$$f_1(V) = (\mathcal{D}_{(A,\Phi)^\#} \mathfrak{q}) \circ ((\mathcal{D}_{(A,\Phi)^\#} \mathfrak{F}_q)^* \Delta_{2,q,(A,\Phi)^\#}^{-1} V + \frac{1}{2} (\mathcal{D}_{(A,\Phi)^\#}^2 \mathfrak{q}) V^{(2)} + \dots,$$

where $V^{(2)} = (V, V)$. Notice that the first term is exactly $f_2(V)$! Therefore

$$f_1'(V) = f_2(V) \quad \text{and} \quad f_2'' = (\mathcal{D}_{(A,\Phi)^\#}^2 \mathfrak{q}).$$

This means that the leading term for the Taylor expansion of $f(V)$ will be quadratic, that is

$$(47) \qquad f(V) = \frac{1}{2} (\mathcal{D}_{(A,\Phi)^\#}^2 \mathfrak{q}) V^{(2)} + \dots.$$

To see why this is important notice in the case of $-\mathcal{Q} \circ [(\mathcal{D}_{(A,\Phi)^\#} \mathfrak{F}_q)^*](\Delta_{2,q,(A,\Phi)^\#}^{-1} V_q)$ Mrowka and Rollin found a bound [35, after Equation 3.14] which can be adapted to our case to read

$$\begin{aligned} (48) \quad & \left\| \mathcal{Q} \circ [(\mathcal{D}_{(A,\Phi)^\#} \mathfrak{F}_q)^*](\Delta_{2,q,(A,\Phi)^\#}^{-1} V_2) \right. \\ & \left. - \mathcal{Q} \circ [(\mathcal{D}_{(A,\Phi)^\#} \mathfrak{F}_q)^*](\Delta_{2,q,(A,\Phi)^\#}^{-1} V_1) \right\|_{L_k^2(g'_\tau, A^\#)} \\ & \leq C'_k \|V_2 + V_1\|_{L_k^2(g_\tau, A^\#)} \|V_2 - V_1\|_{L_k^2(g'_\tau, A^\#)}, \end{aligned}$$

where C'_k , a constant which is independent of τ (once it is large enough), the approximate solution (A, Φ) and the constant $N_0 \geq 1$ are those used in the perturbations defining the connected sum along Y operation.

Since we are assuming that $\|V_1\|_{L_k^2}, \|V_2\|_{L_k^2} \leq \alpha_k$, we can use the triangle inequality to obtain that

$$\|V_2 + V_1\|_{L_k^2(g_\tau, A^\#)} \leq \|V_2\|_{L_k^2(g_\tau, A^\#)} + \|V_1\|_{L_k^2(g_\tau, A^\#)} \leq 2\alpha_k,$$

so the inequality (48) reads

$$\begin{aligned} & \left\| \mathcal{Q} \circ [(\mathcal{D}_{(A,\Phi)^\#} \mathfrak{F}_q)^*](\Delta_{2,q,(A,\Phi)^\#}^{-1} V_2) - \mathcal{Q} \circ [(\mathcal{D}_{(A,\Phi)^\#} \mathfrak{F}_q)^*](\Delta_{2,q,(A,\Phi)^\#}^{-1} V_1) \right\|_{L_k^2(g'_\tau, A^\#)} \\ & \leq 2\alpha_k C'_k \|V_2 - V_1\|_{L_k^2(g'_\tau, A^\#)}. \end{aligned}$$

Hence to make this contribution less than $\frac{1}{2}\kappa_k \|V_2 - V_1\|_{L_k^2(g'_\tau, A^\#)}$ we just need to take $\alpha_k < \kappa/(4C'_k)$.

Likewise, since

$$P(\Delta_{2,q,(A,\Phi)^\#}^{-1} V_2) - P(\Delta_{2,q,(A,\Phi)^\#}^{-1} V_1)$$

is the same as

$$f(V_2) - f(V_1)$$

and each has quadratic leading terms according to (47), the norm

$$\|P(\Delta_{2,q,(A,\Phi)^\#}^{-1} V_2) - P(\Delta_{2,q,(A,\Phi)^\#}^{-1} V_1)\|_{L_k^2(g'_\tau, A^\#)}$$

can now be bounded by an expression of the form

$$f(\alpha_k, C''_k) \|V_2 - V_1\|_{L_k^2(g'_\tau, A^\#)},$$

where $f(\alpha_k, C''_k)$ will be some expression in α_k whose particular details do not interest us and C''_k denotes constants that do not depend on τ or the solution used. In any case, the important thing is that we can again choose α_k so that $f(\alpha_k, C''_k) < \frac{1}{2}\kappa_k$ and so combining both inequalities the result follows. $\square$

At this point we can use the contraction mapping theorem [35, Proposition 2.3.5] to obtain our definition of the *gluing map* [35, Theorem 3.1.9]:

Theorem 46 *There exists constants $\alpha_k, c_k > 0$ such that for every τ large enough, every solution (A, Φ) of the Seiberg–Witten equations on M_τ whose gauge-equivalence class belongs to the zero-dimensional strata of the moduli space $\mathcal{M}(M_\tau; \mathfrak{s}_\tau; [c])$ and every constant $N_0 \geq 1$, there is a unique section (b', ψ') on M'_τ such that*

$$\mathfrak{G}_\tau(A, \Phi) = (A, \Phi)^\# + (\mathcal{D}_{q,(A,\Phi)^\#} \mathfrak{F}_q)^*(b', \psi')$$

is a solution of the Seiberg–Witten equations with $\|(b', \psi')\|_{L_{k+2}^2(g'_\tau, A^\#)} \leq \alpha_k$. Furthermore, the map is gauge equivariant and induces a map

$$\mathfrak{G}_\tau: \mathcal{M}_0(M_\tau; \mathfrak{s}_\tau; [c]) \rightarrow \mathcal{M}_0(M'_\tau; \mathfrak{s}'_\tau; [c]),$$

where $\mathcal{M}_0(M_\tau; \mathfrak{s}_\tau; [c])$ denotes the zero-dimensional strata of $\mathcal{M}(M_\tau; \mathfrak{s}_\tau; [c])$. Further,

$$(49) \quad \begin{aligned} \|(b', \psi')\|_{L_{k+2}^2(g'_\tau, A^\#)} &\leq c_k \|\mathfrak{F}_{p_{M'_\tau}}(A, \Phi)^\#\|_{L_k^2(g'_\tau, A^\#)}, \\ \|\mathfrak{G}_\tau(A, \Phi) - (A, \Phi)^\#\|_{L_{k+1}^2(g'_\tau, A^\#)} &\leq c_k \|\mathfrak{F}_{p_{M'_\tau}}(A, \Phi)^\#\|_{L_k^2(g'_\tau, A^\#)}. \end{aligned}$$

Moreover, $\mathfrak{G}_\tau$ is an injection and since the construction is reversible it is a bijection. Hence $\mathcal{M}_0(M_\tau; \mathfrak{s}_\tau; [c])$ and $\mathcal{M}_0(M'_\tau; \mathfrak{s}'_\tau; [c])$ have the same mod 2 cardinality.

Proof We need to verify that the gluing map preserves the dimensionality of the zero-dimensional strata. For this recall that if $[(A, \Phi)]$ belongs to $\mathcal{M}_0(M_\tau; \mathfrak{s}_\tau; [c])$, then the index of the operator

$$Q_{q,(A,\Phi)} = d^*_{(A,\Phi)} \oplus \mathcal{D}_{(A,\Phi)} \mathfrak{F}_{p_{M_\tau}}$$

is precisely the dimension of the stratum to which $[(A_n, \Phi_n)]$ belongs. Since the transversality condition already implied that $Q_{q,(A,\Phi)}$ was surjective we conclude that in fact $Q_{q,(A,\Phi)}$ is an invertible operator.

Now we use the same splicing procedure as in the case of finding the inverse for the Seiberg–Witten Laplacians $\Delta_{2,q,(A,\Phi)^\#}$. Namely, the operator $\eta_{\text{end}} Q_{(A'_{0,\tau}, \Phi'_\tau)} (\eta_{\text{end}} \cdot)$ associated to the canonical solution $(A_{0,\tau'}, \Phi'_\tau)$ on the AFAK end Z' will be invertible on a suitable domain using [35, Lemma 3.1.4]. Therefore, we can patch together $\eta_{\text{cyl}} Q_{q,(A,\Phi)}^{-1} (\eta_{\text{cyl}} \cdot)$ and $\eta_{\text{end}} Q_{(A'_{0,\tau'}, \Phi'_\tau)}^{-1} (\eta_{\text{end}} \cdot)$ to show that $Q_{q,(A,\Phi)^\#}$ will become invertible.

To compare $Q_{q,(A,\Phi)^\#}$ and $Q_{q,\mathfrak{G}_q(A,\Phi)}$ notice that inequality (37) and the bound in (49) allow us to conclude that the operator norms of $Q_{q,(A,\Phi)^\#}$ and $Q_{q,\mathfrak{G}_q(A,\Phi)}$ are very close to each other. Since being an invertible operator is an open condition it follows that $Q_{q,\mathfrak{G}_q(A,\Phi)}$ will have to be invertible as well.

Now we must address the injectivity of our map. It is essentially the same as the proof of [35, Corollary 3.2.2]. If the injectivity of the map is not true for τ large enough then we obtain a sequence $\tau_j \rightarrow \infty$ and solutions to the Seiberg–Witten equations (A_j, Φ_j) and $(\tilde{A}_j, \tilde{\Phi}_j)$ on M_{τ_j} such that for all j , $[A_j, \Phi_j] \neq [\tilde{A}_j, \tilde{\Phi}_j]$ while $[\mathfrak{G}_{\tau_j}(A_j, \Phi_j)] = [\mathfrak{G}_{\tau_j}(\tilde{A}_j, \tilde{\Phi}_j)]$. Moreover, after taking gauge transformations we can assume that the solutions have exponential decay and converge on every compact subset of $Z_{Y,\xi}^+$ to some solutions (A_∞, Φ_∞) and $(\tilde{A}_\infty, \tilde{\Phi}_\infty)$. Moreover, for all j we have $[A_j, \Phi_j] \neq [\tilde{A}_j, \tilde{\Phi}_j]$ as gauge-equivalence classes. We want to show that if $(A_\infty, \Phi_\infty) = (\tilde{A}_\infty, \tilde{\Phi}_\infty)$ then

$$(50) \qquad \| (A_j, \Phi_j) - (\tilde{A}_j, \tilde{\Phi}_j) \|_{L^2_{k+1}(g_\tau, A_j)} \rightarrow 0.$$

First of all, from (49) and (37) we already know that

$$(51) \qquad \| \mathfrak{G}_{\tau_j}(A_j, \Phi_j) - (A_j, \Phi_j)^\# \|_{L^2_{k+1}(g'_\tau, A^\#)} \rightarrow 0,$$

and hence $\mathfrak{G}_{\tau_j}(A_j, \Phi_j)$ converges on every compact subset towards (A_∞, Φ_∞) since (A_j, Φ_j) does. Similarly $\mathfrak{G}_{\tau_j}(\tilde{A}_j, \tilde{\Phi}_j)$ converges to $(\tilde{A}_\infty, \tilde{\Phi}_\infty)$. Because $\mathfrak{G}(A_j, \Phi_j)$ and $\mathfrak{G}(\tilde{A}_j, \tilde{\Phi}_j)$ are gauge equivalent for each j , the limits are also gauge equivalent.

Hence the limits of (A_j, Φ_j) and $(\tilde{A}_j, \tilde{\Phi}_j)$ are gauge equivalent. After making further gauge transformations, we can then assume that (A_j, Φ_j) and $(\tilde{A}_j, \tilde{\Phi}_j)$ converge toward the same limit (A_∞, Φ_∞) on $Z_{Y,\xi}^+$. In principle, this would be weak convergence along the cylindrical end $\mathbb{R}^+ \times Y$. However, by the discussion from before when we analyzed the restriction of a solution to the cylindrical moduli space $\mathcal{M}(\mathbb{R}^+ \times -Y, \mathfrak{s}_\xi, [\mathfrak{c}])$, we can actually assume that the convergence is strong along the entire cylindrical end; in other words, (A_j, Φ_j) and $(\tilde{A}_j, \tilde{\Phi}_j)$ are converging strongly towards (A_∞, Φ_∞) on the cylindrical end as well. This allows us to conclude that (50) is true.

Since we now have strong convergence along the cylinder, the estimates in [35] continue to hold in that we can find a “radius” r small enough (independent of τ) for which whenever there is j such that $\|(A_j, \Phi_j) - (\tilde{A}_j, \tilde{\Phi}_j)\|_{L^2_{k+1}(g_\tau, A_j)} < r$ then (A_j, Φ_j) and $(\tilde{A}_j, \tilde{\Phi}_j)$ are gauge equivalent (this is a much weaker version of [35, Proposition 3.2.1]). From (50) it is clear that such j will exist and hence we are done. $\square$

We have reached the proof of the naturality property for the contact invariant under strong symplectic cobordisms, Theorem 1. To see why $\widetilde{\text{HM}}_\bullet(W^\dagger, \mathfrak{s}_\omega) c(\xi') = c(\xi)$ recall that in Section 5 we showed that

$$\widetilde{\text{HM}}_\bullet(W^\dagger, \mathfrak{s}_\omega) c(\xi') = c(\xi', Y).$$

The gluing theorem we just proved was aimed at showing that

(52)

$$c(\xi', Y) = c(\xi).$$

To see why $c(\xi', Y) = c(\xi)$ we need to apply Theorem 46 to the case in which the second AFAK end is $Z' = (0, \infty) \times Y$. As explained before, it is not difficult to see that for this choice the corresponding manifolds M'_τ in fact all agree with each other in the sense that their metrics, spinor bundles, symplectic forms, etc are the same, and in fact coincide with the manifold $Z_{Y,\xi}^+$ used to define the contact invariant of (Y, ξ) . In particular, we have for all $\tau > 0$ that

$$|\mathcal{M}_0(M'_\tau, \mathfrak{s}', [\mathfrak{c}])| = |\mathcal{M}_0(Z_{Y,\xi}^+, \mathfrak{s}, [\mathfrak{c}])| \mod 2.$$

Now choose τ_{large} such that

$$|\mathcal{M}_0(M_\tau; \mathfrak{s}_{\tau_{\text{large}}}; [\mathfrak{c}])| = |\mathcal{M}_0(M'_\tau, \mathfrak{s}', [\mathfrak{c}])| = |\mathcal{M}_0(Z_{Y,\xi}^+, \mathfrak{s}, [\mathfrak{c}])| \mod 2.$$

If we think of using the numbers $|\mathcal{M}_0(M_\tau; \mathfrak{s}_{\tau_{\text{large}}}; [\mathfrak{c}])| \mod 2$ in order to define a chain-level element $c(\xi', Y, \tau_{\text{large}}) \in \check{C}_*(-Y, \mathfrak{s}_\xi)$ as in formula (5) (the τ -hybrid invariant we

discussed before), then the previous identity says that at the chain level

$$c(\xi', Y, \tau_{\text{large}}) = c(\xi),$$

which in particular gives the identity of homology classes

(53)

$$c(\xi', Y, \tau_{\text{large}}) = c(\xi).$$

Now, $c(\xi', Y, \tau_{\text{large}})$ is not the same chain-level element as the element $c(\xi', Y)$ we used during the initial sections of this paper. However, it is not difficult to see that we can use a one-parameter family of metrics $g(t)$ and perturbations $p_0(t)$ on $M_{\tau_{\text{large}}}$ (which is diffeomorphic to $W_{\xi', Y}^\dagger$) to go from one element to the other. Therefore, one can consider a parametrized moduli space and use the same argument as in [Section 5](#) to conclude that $c(\xi', Y, \tau_{\text{large}})$ and $c(\xi', Y)$ do define the same homology element in $\widetilde{\text{HM}}_\bullet(-Y, \mathfrak{s}_\xi)$; in other words

(54)

$$c(\xi', Y, \tau_{\text{large}}) = c(\xi', Y).$$

Combining the identities [\(52\)](#), [\(53\)](#) and [\(54\)](#) the naturality result follows; ie we have shown that for a strong symplectic cobordism $(W, \omega): (Y, \xi) \rightarrow (Y', \xi')$ one has

$$\widetilde{\text{HM}}_\bullet(W^\dagger, \mathfrak{s}_\omega) c(\xi') = c(\xi', Y).$$

Appendix Energy and compactness

Now we will briefly discuss the compactness arguments invoked during this paper. We will explain why the results in [\[53\]](#) allows us to extend the compactness results of [\[35\]](#) to the corresponding versions with cylindrical ends. First we will explain why the “dilating the cone” operation we have discussed in this paper can be regarded as a geometric way to implement Taubes’ perturbation for the Seiberg–Witten equations on symplectic manifolds [\[44, Equation 1.18\]](#).

We recall the perturbations used by Taubes. Let (X, ω) be a symplectic manifold. The analysis will be entirely local, so in fact X may be regarded as an open ball or a noncompact manifold if preferred. The positive part S_ω^+ of the spinor bundle determined by ω can be decomposed as

$$S_\omega^+ = \mathbb{C} \Phi_0 \oplus \langle \Phi_0 \rangle^\perp.$$

Here Φ_0 is the canonical spinor that appeared before in our paper: we can regard it as the constant function $\Phi_0: X \rightarrow \{1\} \subset \mathbb{C}$ identically equal to 1. The canonical

connection A_0 is then the unique spin-c connection for which $D_{A_0}\Phi_0 = 0$. The perturbed Seiberg–Witten equations on symplectic manifolds due to Taubes are

$$\text{SW}^r_{\text{Taubes}} \begin{cases} \frac{1}{2}\rho(F_A^+) - (\Phi\Phi^*)_0 = \frac{1}{2}\rho(F_{A_0}^+) - \frac{1}{4}ir\rho(\omega), \\ D_A\Phi = 0. \end{cases}$$

Here r is a parameter that eventually Taubes takes to be very large. For $r = 1$, the perturbations are cooked up in such a way that (A_0, Φ_0) solves $\text{SW}^1_{\text{Taubes}}$, since the Clifford identities say that $(\Phi_0\Phi_0^*)_0 = \frac{1}{4}i\rho(\omega)$. Moreover, $(A_0, \sqrt{r}\Phi_0)$ then solves $\text{SW}^r_{\text{Taubes}}$ for essentially the same reasons. At this point it is useful to note that one can also rescale the spinor as Taubes usually does. That is, one can write $\Phi = \sqrt{r}\phi$, so that $\text{SW}^r_{\text{Taubes}}$ becomes

(55)
$$\text{SW}^r_{\text{Taubes}} \begin{cases} \frac{1}{2}\rho(F_A^+) - r(\phi\phi^*)_0 = \frac{1}{2}\rho(F_{A_0}^+) - \frac{1}{4}ir\rho(\omega), \\ D_A\Phi = 0. \end{cases}$$

To see how to obtain $\text{SW}^r_{\text{Taubes}}$ from $\text{SW}^1_{\text{Taubes}}$,

$$\text{SW}^1_{\text{Taubes}} \begin{cases} \frac{1}{2}\rho(F_A^+) - (\Phi\Phi^*)_0 = \frac{1}{2}\rho(F_{A_0}^+) - \frac{1}{4}i\rho(\omega), \\ D_A\Phi = 0, \end{cases}$$

suppose that we dilate the metric. That is, for a parameter $\tau > 0$, we define

$$g_\tau = \tau^2 g,$$

where g was a metric compatible with ω , ie ω is a self-dual harmonic 2-form of the pointwise norm $\sqrt{2}$ with respect to g . Since τ is a constant then

$$\omega_\tau = \tau^2 \omega$$

will continue to be a symplectic form, now compatible with the metric g_τ . The spinor bundle will clearly not change with respect to this new metric; that is,

$$S_{\omega,\tau} = S_\omega,$$

while the Clifford map on one-forms is rescaled as

$$\rho_\tau = \frac{\rho}{\tau}.$$

On two-forms we find that $\rho_\tau = \rho/\tau^2$, so in particular $\rho_\tau(\omega_\tau) = \rho(\omega)$. A dilation is a very simple conformal change of metric, for which it is understood how the Dirac operator changes [3, Equation D.1]. In our case we find that

$$D_{A,g_\tau}\Phi = \tau^{-1}D_{A,g}\Phi,$$

so in particular being a harmonic spinor (ie $D_A\Phi = 0$) is a condition independent of the metric g_τ used, and the canonical connection will be preserved under the dilation, since A_0 satisfies the property $D_{A_0,g_\tau}\Phi_0 = 0$ regardless of the value of τ . In other words

$$A_{0,\tau} = A_0.$$

Moreover, the notion of self-duality is preserved under conformal changes of the metric, in particular, dilations, which means that the *Taubian geometric perturbations* of the Seiberg–Witten equations on the dilated symplectic manifolds (X, g_τ, ω_τ) ,

$$\text{SW}_{\text{Taubes}}^\tau \begin{cases} \frac{1}{2}\rho_\tau(F_A^+) - (\Phi\Phi^*)_0 = \frac{1}{2}\rho_\tau(F_{A_{0,\tau}}^+) - \frac{1}{4}i\rho_\tau(\omega_\tau), \\ D_{A,g_\tau}\Phi = 0, \end{cases}$$

can be rewritten in terms of the geometric structures on the original data (X, g, ω) as (recall that on two-forms, $\rho_\tau(F_A^+) = \rho(F_A^+)/\tau^2$)

(56)
$$\text{SW}_{\text{Taubes}}^\tau \begin{cases} \frac{1}{2}\rho(F_A^+) - \tau^2(\Phi\Phi^*)_0 = \frac{1}{2}\rho(F_{A_{0,\tau}}^+) - \frac{1}{4}i\tau^2\rho(\omega), \\ D_{A,g}\Phi = 0. \end{cases}$$

Setting $\tau = \sqrt{r}$, we see that $\text{SW}_{\text{Taubes}}^\tau$ is indistinguishable from $\text{SW}_{\text{Taubes}}^r$! Therefore, the results from papers which work with the perturbations $\text{SW}_{\text{Taubes}}^r$ can be translated immediately to our paper. In particular, the theorems from [53] are readily available to our situation. The only caveat is that [53] interprets the scaling of the spinors in a slightly different way. For example, in [53, Definition 3.5], the decay condition is written as $(\Phi - \sqrt{r}\Phi_0) \in L^2_{k,A_0}$, while in our definition of the configuration space, Definition 12, we write the decay condition as $(\Phi - \Phi_0) \in L^2_{k,A_0}$. In particular, when the results of [53] are translated into our context, there are additional factors of $\sqrt{r}$ one needs to remove.

The most important result proven from [53] for the purposes of our problem is the uniform bound on the symplectic energy for solutions of the Seiberg–Witten equations on manifolds with both symplectic and cylindrical ends. More precisely, when (A, Φ) is a configuration on M_τ we can restrict it to the symplectic region $M_\tau \setminus (\mathbb{R}^+ \times -Y)$ and consider the *symplectic energy*

(57)
$$E_\tau(A, \Phi) = \int_{M_\tau \setminus (\mathbb{R}^+ \times -Y)} \left[(1 - |\alpha|^2 - |\beta|^2)^2 + |F_a|^2 + |\widehat{\nabla}_A\Phi|^2 \right] \text{vol}^{g_\tau},$$

which is defined in [53, Section 5.3]. Here we decomposed Φ as (α, β) and A as $A = A_0 + a$. In (57), $\widehat{\nabla}_A$ refers to the *twisted Chern connection*, which can be defined as

$$\widehat{\nabla}_A\Phi = \nabla^C\Phi + a \otimes \Phi,$$

where ∇^C is the Chern connection the AFAK end structure determines (this is explained

in [35, Section 1.3.2]). Notice that in [53], $|\nabla_a \alpha|^2 + |\nabla'_A \beta|^2$ appears in the place of $|\widehat{\nabla}_A \Phi|^2$, but these two things in fact have the same meaning. The uniform bound on the symplectic energy E_τ that we need is the following.

Lemma 47 *Consider the sequence of manifolds M_τ defined in (34) (Section 6.2). There exists a constant κ such that for every τ large enough and every solution to the Seiberg–Witten equations (A, Φ) on M_τ asymptotic along the symplectic end to (A_0, Φ_0) and along the cylindrical end to a critical point $\mathfrak{c}$ of the three-dimensional Seiberg–Witten equations, we have that*

$$E_\tau(A, \Phi) \leq \kappa.$$

Proof Notice that if the cylindrical end is not present, then this result would follow from [35, Lemma 2.2.7]. In our case the perturbation term ϖ_τ which appears in the previously cited lemma of Mrowka and Rollin is not present, since this was only introduced for transversality purposes, which is not needed for the manifolds M_τ , thanks to the use of the abstract perturbations along the cylindrical end and the fiber product description of the moduli spaces.

As mentioned by Mrowka and Rollin, when their cylindrical end is not present, the fact that κ is independent of τ is a consequence of the uniform boundedness of the injectivity radius and curvature, as well as the other geometric quantities involved in the definition of these families of AFAK manifolds M_τ , as described in [35, Lemma 2.1.6]. In other words, when the cylindrical end is not present, finding a uniform control on the symplectic energy E_τ is a consequence of knowing that E_{τ_0} is bounded for some τ_0 sufficiently large (which was the content of [21, Lemma 3.17]) and then appealing to the uniform geometry of the manifolds M_τ .

Therefore, for our situation what we need to know is that in the presence of a cylindrical end it is still the case that one can find τ_0 large enough that $E_{\tau_0}(A, \Phi) \leq \kappa_0$ for all solutions (A, Φ) on M_{τ_0} and for some κ_0 independent of the Seiberg–Witten solution (A, Φ) on M_{τ_0} . In other words, we want the analogue of [21, Lemma 3.17] in the presence of a cylindrical end. The fact that κ_0 can be taken to be independent of τ_0 will follow again from the comments we just made in the previous paragraph regarding the uniform boundedness of the geometric quantities involved in the family of the AFAK manifolds. The existence of such a τ_0 now follows from [53, Proposition 5.12], given that we can reinterpret the equations on an arbitrary M_τ as equations on the fixed M_1 , at the cost of perturbing the curvature equation with factors of τ , as we explained at the beginning of the [appendix](#) when we discussed the Taubian perturbations (55) and (56). $\square$

Next we explain some of the auxiliary lemmas used in some of the proofs of our paper. The first one appeared in [Section 6.3](#), where the pregluing map was being constructed.

Lemma 48 *Consider the sequence of manifolds M_τ defined in [\(34\)](#) ([Section 6.2](#)). We are able to find a compact set $C = [1, T] \times Y$ contained in the symplectic region $M_\tau \setminus (\mathbb{R}^+ \times -Y)$ with the following significance: for every τ large enough and for every solution (A, Φ) to the Seiberg–Witten equations on M_τ asymptotic along the symplectic end to (A_0, Φ_0) and along the cylindrical end to a critical point $\mathfrak{c}$ of the three-dimensional Seiberg–Witten equations, we have $|\alpha| \geq \frac{1}{2}$ on $M_\tau \setminus [(\mathbb{R}^+ \times -Y) \cup C]$, where we wrote Φ as (α, β) along the symplectic end.*

Remark 49 Notice that τ will in fact not depend on the particular critical point $\mathfrak{c}$ we are asymptotic to.

Proof Observe that this corresponds to Lemma 2.2.8 in [\[35\]](#). The proof Mrowka and Rollin provided proceeds by contradiction, and eventually gives rise to a sequence of solutions to the Seiberg–Witten equations (A_j, Φ_j) on M_{τ_j} . However, they just care about the restrictions of these solutions to some balls $B(x_j, \sigma_{\tau_j}(x_j)/\kappa)$ centered at x_j of radii $\sigma_{\tau_j}(x_j)/\kappa$. The important feature of these balls is that they are all contained in the symplectic region $M_{\tau_j} \setminus (\mathbb{R}^+ \times -Y)$, so what is happening along the cylinder is of no importance, given that we already know the uniform control on the symplectic energies E_τ thanks to the previous lemma. Therefore, the proof they give in fact goes through in our setup, since on the symplectic region we are using the same perturbations as Mrowka and Rollin. □

Remark 50 The subsequence that Mrowka and Rollin obtained in the proof of [\[35, Lemma 2.2.8\]](#) can also be obtained from [\[54, Proposition 5.3\]](#), if in the notation of [\[54\]](#) we take the sequence $\{(M_n, g_n, p_n)\}$ to be the sequence of pointed balls $B(x_j, \sigma_{\tau_j}(x_j)/\kappa)$ that arise from the proof by contradiction.

The next lemma appeared right after the pregluing map [\(36\)](#) was defined.

Lemma 51 *Lemma 2.5.4 in [\[35\]](#) still holds. That is, there is a $\delta > 0$ and a T large enough that for every $N_0 \geq 1$, $k \in \mathbb{N}$, τ satisfying $\tau \geq T + N_0$ and solution (A, Φ) of the Seiberg–Witten equations on M_τ , we have that $(A, \Phi)^\#$ satisfies the Seiberg–Witten equations on $\{\sigma'_\tau \leq T\} \subset M'_\tau$ and*

(58)

$$|\mathfrak{F}_{p_{M'_\tau}}(A, \Phi)^\#|_{C^k(g'_\tau, A^\#)} \leq c_k e^{-\delta \sigma_\tau}$$

on $\{\sigma'_\tau \geq T\} \subset M'_\tau$.

Remark 52 Again, we are assuming that the solutions are asymptotic along the cylindrical end to some critical point c , but the constants of the lemma are insensitive to the particular critical point being used.

Proof Our proof is essentially the same as the proof of [35, Lemma 2.5.4]. The only thing we need to know is that we can find a gauge with uniform exponential decay, as in [35, Corollary 2.2.10]. In turn, the proof of that corollary was modeled on the proof of [21, Corollary 3.16]. This last corollary required knowing that the symplectic energy E_τ is uniformly bounded. But this is precisely the content of our Lemma 47. $\square$

The following lemma was used in the proof of Lemma 43.

Lemma 53 Let (A_n, Φ_n) be a sequence of solutions to the Seiberg–Witten equations on M_{τ_n} . Then, after making gauge transformations if necessary, we can assume that (A_n, Φ_n) converges weakly to a solution (A_∞, Φ_∞) on $Z_{Y,\xi}^+$. In particular, on every compact set K of $Z_{Y,\xi}^+$, which can be regarded as a subset of M_{τ_n} for all τ_n sufficiently large, we have strong convergence of (A_n, Φ_n) to (A_∞, Φ_∞) .

Therefore, if we take a sequence of solutions (A_n, Φ_n) on M_{τ_n} and restrict them to $[1, T] \times Y \subset M_{\tau_n}$, the solutions will converge (after any necessary gauge transformations) strongly to a solution (A_∞, Φ_∞) on $[1, T] \times Y \subset Z_{Y,\xi}^+$.

Proof Notice that when we restrict the solutions (A_n, Φ_n) to the cylindrical ends $\mathbb{R}^+ \times -Y$, which are the same for all M_{τ_n} , we already know that they will converge in the weak sense to some solution on $\mathbb{R}^+ \times -Y$, and hence strongly on compact subsets of $\mathbb{R}^+ \times -Y$ (after gauging if necessary). This compactness result is based on [22, Proposition 24.6.4], which only needs a bound on the topological energy $\mathcal{E}_q^{\text{top}}$ [22, Definition 24.6.3]. That the topological energy continues to be bounded in our situation is explained in [53, Remark 4 after the proof of Theorem 6.1].

Therefore, the only new thing to understand is why the convergence still holds when we restrict the solutions to the symplectic ends, since after that one can do a patching argument, as in the proof of [22, Proposition 24.6.4]. The convergence on the symplectic end follows the same argument as in the proof of the compactness theorem [35, Theorem 2.2.11], which is a consequence of being able to find gauge transformations along the symplectic end which give rise to a uniform exponential decay for (A_n, Φ_n) with respect to (A_0, Φ_0) . As we mentioned previously, in the proof of Lemma 51, this is a consequence of the uniform control on the symplectic energy E_{τ_n} , ie Lemma 47. $\square$

References

- [1] **N Berline, E Getzler, M Vergne**, *Heat kernels and Dirac operators*, corrected 1st edition, Grundle Math. Wissen. 298, Springer (2004) [MR](#)
- [2] **C P Boyer, K Galicki**, *Sasakian geometry*, Oxford Univ. Press (2008) [MR](#)
- [3] **B Charbonneau**, *Analytic aspects of periodic instantons*, PhD thesis, Massachusetts Institute of Technology (2004) [MR](#) Available at <https://dspace.mit.edu/handle/1721.1/26746>
- [4] **V Colin, P Ghiggini, K Honda**, *The equivalence of Heegaard Floer homology and embedded contact homology via open book decompositions, I*, preprint (2012) [arXiv](#)
- [5] **V Colin, P Ghiggini, K Honda**, *The equivalence of Heegaard Floer homology and embedded contact homology via open book decompositions, II*, preprint (2012) [arXiv](#)
- [6] **V Colin, P Ghiggini, K Honda**, *The equivalence of Heegaard Floer homology and embedded contact homology, III: From hat to plus*, preprint (2012) [arXiv](#)
- [7] **S K Donaldson**, *Floer homology groups in Yang–Mills theory*, Cambridge Tracts in Mathematics 147, Cambridge Univ. Press (2002) [MR](#)
- [8] **M Echeverria**, *Naturality of the contact invariant in monopole floer homology under strong symplectic cobordisms*, PhD thesis, University of Virginia (2019) [MR](#) Available at <https://doi.org/10.18130/v3-vykt-tc16>
- [9] **J Eichhorn**, *Gauge theory on open manifolds of bounded geometry*, Internat. J. Modern Phys. A 7 (1992) 3927–3977 [MR](#)
- [10] **Y Eliashberg**, *Classification of overtwisted contact structures on 3–manifolds*, Invent. Math. 98 (1989) 623–637 [MR](#)
- [11] **J B Etnyre**, *Planar open book decompositions and contact structures*, Int. Math. Res. Not. 2004 (2004) 4255–4267 [MR](#)
- [12] **J B Etnyre, K Honda**, *On symplectic cobordisms*, Math. Ann. 323 (2002) 31–39 [MR](#)
- [13] **D S Freed, K K Uhlenbeck**, *Instantons and four-manifolds*, 2nd edition, Mathematical Sciences Research Institute Publications 1, Springer (1991) [MR](#)
- [14] **K A Frøyshov**, *Compactness and gluing theory for monopoles*, Geom. Topol. Monogr. 15, Geom. Topol. Publ., Coventry (2008) [MR](#)
- [15] **H Geiges**, *An introduction to contact topology*, Cambridge Studies in Advanced Mathematics 109, Cambridge Univ. Press (2008) [MR](#)
- [16] **P Ghiggini**, *Ozsváth–Szabó invariants and fillability of contact structures*, Math. Z. 253 (2006) 159–175 [MR](#)
- [17] **R E Gompf**, *Handlebody construction of Stein surfaces*, Ann. of Math. 148 (1998) 619–693 [MR](#)
- [18] **M Hutchings**, *Embedded contact homology as a (symplectic) field theory*, in preparation

- [19] **M Hutchings**, *Strong cobordisms and the ECH contact invariant*, blog post (2012) Available at <https://tinyurl.com/floerhomology-blog-120507>
- [20] **M Hutchings**, **CH Taubes**, *Proof of the Arnold chord conjecture in three dimensions, II*, *Geom. Topol.* 17 (2013) 2601–2688 [MR](#)
- [21] **PB Kronheimer**, **TS Mrowka**, *Monopoles and contact structures*, *Invent. Math.* 130 (1997) 209–255 [MR](#)
- [22] **P Kronheimer**, **T Mrowka**, *Monopoles and three-manifolds*, *New Mathematical Monographs* 10, Cambridge Univ. Press (2007) [MR](#)
- [23] **P Kronheimer**, **T Mrowka**, **P Ozsváth**, **Z Szabó**, *Monopoles and lens space surgeries*, *Ann. of Math.* 165 (2007) 457–546 [MR](#)
- [24] **C Kutluhan**, **Y-J Lee**, **CH Taubes**, *HF = HM, I: Heegaard Floer homology and Seiberg–Witten Floer homology*, preprint (2011) [arXiv](#) To appear in *Geom. Topol.*
- [25] **C Kutluhan**, **Y-J Lee**, **CH Taubes**, *HF = HM, II: Reeb orbits and holomorphic curves for the ech/Heegaard–Floer correspondence*, preprint (2010) [arXiv](#) To appear in *Geom. Topol.*
- [26] **C Kutluhan**, **Y-J Lee**, **CH Taubes**, *HF = HM, III: Holomorphic curves and the differential for the ech/Heegaard Floer correspondence*, preprint (2010) [arXiv](#) To appear in *Geom. Topol.*
- [27] **C Kutluhan**, **Y-J Lee**, **CH Taubes**, *HF = HM, IV: The Seiberg–Witten Floer homology and ech correspondence*, preprint (2011) [arXiv](#) To appear in *Geom. Topol.*
- [28] **C Kutluhan**, **Y-J Lee**, **CH Taubes**, *HF = HM, V: Seiberg–Witten–Floer homology and handle addition*, preprint (2012) [arXiv](#) To appear in *Geom. Topol.*
- [29] **S Lang**, *Fundamentals of differential geometry*, *Graduate Texts in Mathematics* 191, Springer (1999) [MR](#)
- [30] **J Lin**, *The Seiberg–Witten equations on end-periodic manifolds and an obstruction to positive scalar curvature metrics*, *J. Topol.* 12 (2019) 328–371 [MR](#)
- [31] **J Lin**, **D Ruberman**, **N Saveliev**, *On the Frøyshov invariant and monopole Lefschetz number*, preprint (2018) [arXiv](#)
- [32] **P Lisca**, **A I Stipsicz**, *Seifert fibered contact three-manifolds via surgery*, *Algebr. Geom. Topol.* 4 (2004) 199–217 [MR](#)
- [33] **J Morgan**, **T Mrowka**, *On the gluing theorem for instantons on manifolds containing long cylinders*, unpublished manuscript
- [34] **JW Morgan**, **T Mrowka**, **D Ruberman**, *The L^2 -moduli space and a vanishing theorem for Donaldson polynomial invariants*, *International*, Cambridge, MA (1994) [MR](#)
- [35] **T Mrowka**, **Y Rollin**, *Legendrian knots and monopoles*, *Algebr. Geom. Topol.* 6 (2006) 1–69 [MR](#)

- [36] **T Mrowka, D Ruberman, N Saveliev**, *Seiberg–Witten equations, end-periodic Dirac operators, and a lift of Rohlin’s invariant*, J. Differential Geom. 88 (2011) 333–377 [MR](#)
- [37] **LI Nicolaescu**, *Notes on Seiberg–Witten theory*, Graduate Studies in Mathematics 28, Amer. Math. Soc., Providence, RI (2000) [MR](#)
- [38] **J C Oxtoby**, *Measure and category: a survey of the analogies between topological and measure spaces*, 2nd edition, Graduate Texts in Mathematics 2, Springer (1980) [MR](#)
- [39] **P Ozsváth, A Stipsicz, Z Szabó**, *Planar open books and Floer homology*, Int. Math. Res. Not. 2005 (2005) 3385–3401 [MR](#)
- [40] **P Ozsváth, Z Szabó**, *Holomorphic disks and genus bounds*, Geom. Topol. 8 (2004) 311–334 [MR](#)
- [41] **P Ozsváth, Z Szabó**, *Heegaard Floer homology and contact structures*, Duke Math. J. 129 (2005) 39–61 [MR](#)
- [42] **P Safari**, *Gluing Seiberg–Witten monopoles*, Comm. Anal. Geom. 13 (2005) 697–725 [MR](#)
- [43] **S Sivek**, *Monopole Floer homology and Legendrian knots*, Geom. Topol. 16 (2012) 751–779 [MR](#)
- [44] **CH Taubes**, *$SW \Rightarrow Gr$: from the Seiberg–Witten equations to pseudo-holomorphic curves*, J. Amer. Math. Soc. 9 (1996) 845–918 [MR](#)
- [45] **CH Taubes**, *The Seiberg–Witten equations and the Weinstein conjecture, II: More closed integral curves of the Reeb vector field*, Geom. Topol. 13 (2009) 1337–1417 [MR](#)
- [46] **CH Taubes**, *Embedded contact homology and Seiberg–Witten Floer cohomology, I*, Geom. Topol. 14 (2010) 2497–2581 [MR](#)
- [47] **CH Taubes**, *Embedded contact homology and Seiberg–Witten Floer cohomology, II*, Geom. Topol. 14 (2010) 2583–2720 [MR](#)
- [48] **CH Taubes**, *Embedded contact homology and Seiberg–Witten Floer cohomology, III*, Geom. Topol. 14 (2010) 2721–2817 [MR](#)
- [49] **CH Taubes**, *Embedded contact homology and Seiberg–Witten Floer cohomology, IV*, Geom. Topol. 14 (2010) 2819–2960 [MR](#)
- [50] **CH Taubes**, *Embedded contact homology and Seiberg–Witten Floer cohomology, V*, Geom. Topol. 14 (2010) 2961–3000 [MR](#)
- [51] **C Wendl**, *A hierarchy of local symplectic filling obstructions for contact 3–manifolds*, Duke Math. J. 162 (2013) 2197–2283 [MR](#)
- [52] **C Wendl**, *Non-exact symplectic cobordisms between contact 3–manifolds*, J. Differential Geom. 95 (2013) 121–182 [MR](#)
- [53] **B Zhang**, *A monopole Floer invariant for foliations without transverse invariant measure*, preprint (2016) [arXiv](#)

- [54] **R Zhang**, *Gauge theory and self-linking of Legendrian knots*, PhD thesis, Northeastern University (2016) [MR](#) Available at <https://search.proquest.com/docview/1786646026>

*Department of Mathematics, University of Virginia
Charlottesville, VA, United States*

*Current address: Department of Mathematics, Rutgers University
Piscataway, NJ, United States*

me3qr@virginia.edu

Received: 7 September 2018 Revised: 11 August 2019

