

Pillowcase covers: Counting Feynman-like graphs associated with quadratic differentials

ELISE GOUJARD
MARTIN MÖLLER

We prove the quasimodularity of generating functions for counting pillowcase covers, with and without Siegel–Veech weight. Similar to prior work on torus covers, the proof is based on analyzing decompositions of half-translation surfaces into horizontal cylinders. It provides an alternative proof of the quasimodularity results of Eskin and Okounkov and a practical method to compute area Siegel–Veech constants.

A main new technical tool is a quasipolynomiality result for 2–orbifold Hurwitz numbers with completed cycles.

11F11, 32G15; 14H30, 14N10, 30F30, 81T18

1. Introduction	2451
2. Counting covers of the pillow by global graphs	2455
3. Fermionic Fock space and its balanced subspace	2464
4. Hurwitz numbers and graph sums	2470
5. Constant coefficients of quasielliptic functions	2473
6. Quasimodularity of graph sums	2479
7. Quasipolynomiality of 2–orbifold double Hurwitz numbers	2484
8. Application to Siegel–Veech constants	2491
9. Example: $Q(2, 1, -1^3)$	2498
References	2508

1 Introduction

Mirror symmetry for elliptic curves can be phrased in its tropical version by stating that the Hurwitz number counting covers of elliptic curves can be computed as Feynman

integrals and that the corresponding generating functions are quasimodular forms; see Böhm, Bringmann, Buchholz and Markwig [3], Goujard and Möller [17], Dijkgraaf [9], Kaneko and Zagier [19] and Eskin and Okounkov [13]. A Feynman integral is physics-inspired terminology for an integral over a product of derivatives of propagators (ie Weierstrass \wp -functions), the form of the product being encoded by a (Feynman) graph.

The goal of this paper is to show that the mirror symmetry story has a complete analog in the scope of pillowcase covers, covers of the projective line with ramification profile given by the partition $(2, \dots, 2)$ over 3 points, with the profile $(\nu, 2, \dots, 2)$ over a special point (where the parts of the partition ν are all odd) and possibly a fixed finite number of even-order branch points elsewhere. (Section 2.1 provides an introduction to the relevant terminology for covers and Hurwitz spaces.) These pillowcase covers arise naturally in the volume computation for strata of quadratic differentials.

Our generalized Feynman graphs encode the decomposition into horizontal cylinders of these pillowcase covers. In this coding, the vertices correspond to the branch points, and the edges correspond to the cylinders. The generalized Feynman graphs have a special vertex 0 corresponding to the branch point with profile $(\nu, 2, \dots, 2)$ and, most important, come with an orientation of the half-edges, so that the edge contribution to the Feynman integrand is the Weierstrass \wp -function $\wp(z_i \pm z_j)$ evaluated at $z_i \pm z_j$, according to whether the half-edges are inconsistently or consistently oriented along the edge. The evaluation of those generalized Feynman integrals are quasimodular forms; see Section 5.1 for a definition that applies to a general cofinite Fuchsian group. In fact, the evaluations are now quasimodular only for the level subgroup $\Gamma_0(2)$, rather than for the full modular group $\mathrm{SL}(2, \mathbb{Z})$ in the case of torus covers. The argument is a rather straightforward generalization of the torus cover case; see Theorems 5.6 and 6.1, and compare to [17, Sections 5 and 6]. The two papers are intentionally parallel whenever possible, to facilitate comparison. In particular, both papers start with a correspondence theorem (Proposition 2.3) that can certainly be rephrased in terms of covers of tropical curves [17, Section 8; 3].

To arrive from there at our main goals, we need moreover a structure theorem for the algebra of shifted symmetric quasipolynomials and a polynomiality theorem for orbifold double Hurwitz numbers; see the end of the introduction. Altogether, we can first give another proof of the following theorem of Eskin and Okounkov [14]. We let $N^\circ(\Pi) = \sum N_d^\circ(\Pi)q^d$ be the generating series of the number $N_d^\circ(\Pi)$ of connected pillowcase covers of degree d with branching profile Π . We let $\ell(\Pi)$ be the number of parts of the partition Π , and $|\Pi|$ denotes its total cardinality.

Theorem 1.1 (= Corollary 7.4) *For any ramification profile Π the counting function $N^\circ(\Pi)$ for connected pillowcase covers of profile Π is a quasimodular form for the group $\Gamma_0(2)$ of mixed weight less than or equal to $|\Pi| + \ell(\Pi)$.*

The starting point of this paper was to obtain the following version for a weighted count, motivated by the computation of area Siegel–Veech constants (see Section 6 for a brief introduction, and Eskin, Kontsevich, Masur and Zorich [12; 11] for more background.).

Theorem 1.2 (= Corollary 8.2) *For any ramification profile Π and any odd integer $p \geq -1$, the generating series $c_p^\circ(\Pi)$ for counting connected pillowcase covers with p –Siegel–Veech weight is a quasimodular form for the group $\Gamma_0(2)$ of mixed weight at most $|\Pi| + \ell(\Pi) + p + 1$.*

To explain the use of this result, we compare the knowledge about strata of the moduli space of abelian differentials and quadratic differentials with respect to Masur–Veech volumes and Siegel–Veech constants at the time of writing. In the abelian case our understanding is nearly complete. Siegel–Veech constants can be computed recursively by computing ratios of Masur–Veech volumes of boundary strata [12]. These volumes can be computed efficiently by counting torus covers and closed formulas derived from this; see Eskin and Okounkov [13] and Chen, Möller and Zagier [6]. The volumes have an interpretation as intersection numbers of tautological classes — see Sauvaget [24] and Chen, Möller, Sauvaget and Zagier [5] — and the formulas are well-understood, so as to give large genus asymptotics in all detail; see [6; 5] and Aggarwal [1].

For the moduli space of quadratic differentials much less is known, except for strata of genus zero surfaces whose volumes are explicitly computable; see Athreya, Eskin and Zorich [2]. Siegel–Veech constants are also related to Masur–Veech volumes by a recursive procedure; see Masur and Zorich [21] and Goujard [15]. But these volumes are much harder to evaluate for higher genus, despite the work of Eskin and Okounkov [14], and some hints being given in Engel [10]. The behavior of the large genus asymptotics is conjectured in Delecroix, Goujard, Zograf and Zorich [7] for the sequence of principal strata. Only for the principal strata is an interpretation as an intersection number known [7].

In the current status of knowledge, Theorems 1.1 and 1.2 provide (besides structural insight) a somewhat reasonable practical way to compute volumes and Siegel–Veech

constants for strata of quadratic differentials by computing the coefficients for sufficiently many small d in order to determine the quasimodular form uniquely and then using the growth rate of the coefficients. This procedure is explained, along with the technical steps of the proof, in an example in Section 9. Algorithms that compute volumes and Siegel–Veech constants for quadratic differentials as efficiently as in the abelian case still have to be found.

Finally, we explain the “local” polynomiality results that are the intrinsic reason for quasimodularity. In the case of torus covers, double Hurwitz numbers (counting branched covers with a specified branching away from two special points) arise naturally by slicing the torus. These numbers are polynomials in the ramification orders at the two special points if one uses completed cycles at each slice; see eg Shadrin, Spitz and Zvonkine [26]. In other words, we look at a degeneration of the target elliptic curve so that there is a single branch point in each connected component of the nodal curve. Then the degeneration formula for Hurwitz covers expresses the corresponding Hurwitz numbers as a vacuum expectation of a product of local operators, hence the polynomiality. Together with a theorem that shows that certain graph sums with polynomial local contributions are quasimodular and a graph combination argument to pass to completed cycles p_ℓ , we obtained in [17] quasimodularity for torus covers.

In the case of pillowcase covers, slicing the pillowcase, some 2–orbifold Hurwitz numbers arise naturally at special slices. We show in Theorem 7.2 that these 2–orbifold Hurwitz numbers are quasipolynomiality (rather than just piecewise quasipolynomiality) if the 2–orbifold carries only products of the completed cycles \bar{p}_k in the algebra of shifted symmetric quasipolynomiality (see Section 3). Note that quasipolynomiality of 2–orbifold Hurwitz numbers fails even for the completed cycles p_ℓ . The quasipolynomiality is the cause of quasimodularity of the associated generating series for the subgroup $\Gamma_0(2)$ rather than the full group $\mathrm{SL}(2, \mathbb{Z})$.

Acknowledgements The authors are very grateful to Dmitri Zvonkine for suggesting the form of the one-sided pillowcase vertex operator. We thank Alex Eskin for sharing a manuscript of an old project with Andrei Okounkov that also discussed local surfaces and global graphs. We thank Philip Engel for many helpful comments that improved the exposition of the paper. Both authors acknowledge the hospitality of the *Max-Planck Institute for Mathematics* (MPIM, Bonn), where much of this work was done. Research of Goujard was partially supported by a public grant as part of the FMJH. Research of Möller is partially supported by the DFG-project MO 1884/1-1 and by the LOEWE-Schwerpunkt “Uniformisierte Strukturen in Arithmetik und Geometrie”.

2 Counting covers of the pillow by global graphs

The goal of this section is the basic correspondence theorem, Proposition 2.3, and its variants. It allows us to count covers of the pillow by counting graphs with various additional decorations. As in the abelian case, the correspondence theorem works only if we count coverings without unramified components. We thus start with standard remarks on the passage between the various ways of imposing connectivity in the counting problems.

2.1 Covers of the pillow and their Hurwitz tuples

We give a short introduction to Hurwitz spaces of covers of the pillow $B \cong \mathbb{C}P^1$ and recall some basic notions needed in the sequel. We provide the pillow with the flat metric that identifies B with two squares of side length $\frac{1}{2}$ glued back to back. We will denote the corners of the pillow by P_1, \dots, P_4 .

A *pillowcase cover* is a cover of degree $2d$ of B fully branched with d transpositions over three corners of the pillow, with all odd-order branching stocked together with transpositions over the remaining corner of the pillow, and with all other even-order branch points cyclic and located at arbitrary points different from the corners. We rephrase this condition in terms of ramification profile and introduce some notation used throughout the paper.

Let $\Pi = (\mu^{(1)}, \dots, \mu^{(n+4)})$ consist of the following types of partitions. We impose that $\mu^{(1)} = (v, 2^{d-|v|/2})$, where v is a partition of an even number into odd parts, that $\mu^{(2)} = \mu^{(3)} = \mu^{(4)} = (2^d)$ and finally that $\mu^{(i+4)} = (\mu_i, 1^{2d-\mu_i})$ with $\mu_i \geq 2$. We refer to both $\mu^{(i)}$ and to Π as a (local or global) *ramification profile* and we define g by the relation

$$\ell(\mu) + \ell(v) - |\mu| - \frac{1}{2}|v| = 2 - 2g,$$

where $\ell(\cdot)$ denotes the length of a partition and $|\cdot|$ the size of a partition. We write Π_\emptyset for the profile with $n = 4$ and $\mu^{(1)} = (2^d)$.

Let $H_d(\Pi)$ (or just H if the parameters are fixed) denote the n -dimensional *Hurwitz space* of degree $2d$, genus g coverings $p: X \rightarrow \mathbb{P}^1$ of a curve of genus zero with $n + 4$ branch points and ramification profile Π , ie we require that over the i^{th} branch point P_i there are $\ell(\mu^{(i)})$ ramification points, of ramification orders respectively given by the parts $\mu_j^{(i)}$ of the partition $\mu^{(i)}$.




Figure 1: Standard presentation of $\pi_1(\mathbb{P}^1 \setminus \{P_1, \dots, P_{n+4}\})$.

Let $\rho: \pi_1(\mathbb{P}^1 \setminus \{P_1, \dots, P_{n+4}\}, \mathbb{Z}) \rightarrow S_{2d}$ be the monodromy representation in the symmetric group of $2d$ elements associated with a covering in H . We use the convention that loops (and elements of the symmetric group) are composed from right to left. The elements $(\alpha_1, \alpha_2, \alpha_3, \alpha_4, \gamma_1, \dots, \gamma_n)$ as in the left picture of Figure 1 generate the fundamental group $\pi_1(\mathbb{P}^1 \setminus \{P_1, \dots, P_{n+4}\}, \mathbb{Z})$ with the relation

$$(1) \quad \alpha_1 \alpha_4 \gamma_1 \cdots \gamma_n = \alpha_2^{-1} \alpha_3^{-1}.$$

Given such a homomorphism ρ , we let $\alpha_i = \rho(\alpha_i)$ and $\gamma_i = \rho(\gamma_i)$ and call the tuple

$$(2) \quad h = (\alpha_1, \dots, \alpha_4, \gamma_1, \dots, \gamma_n) \in (S_{2d})^{n+4}$$

the *Hurwitz tuple* corresponding to ρ and the choice of generators. Conversely, a Hurwitz tuple as in (2) satisfying (1) and generating a transitive subgroup of S_{2d} defines a homomorphism ρ and thus a connected covering p . We denote by $\text{Hur}_d^0(\Pi)$ the set of all such Hurwitz tuples, the upper zero reflecting that we count connected coverings only. The set of all Hurwitz tuples (ie without the requirement of a transitive subgroup) is denoted by $\text{Hur}_d(\Pi)$. As an important technical intermediate notion we need covers *without unramified components*, ie covers $p: X \rightarrow \mathbb{P}^1$ that do not have a connected component X' such that $p|_{X'} = \pi_T \circ p'$ factors into an unramified covering p' and the torus double covering $\pi_T: E \rightarrow \mathbb{P}^1$ branched at P_1, \dots, P_4 . We let $\alpha_1^0 \in S_{2d}$ be the permutation with all transpositions of α_1 replaced by the identity. In terms of Hurwitz tuples, we define the tuples without unramified components equivalently as

$$\text{Hur}'_d(\Pi) = \{h \in \text{Hur}_d(\Pi) : \langle \alpha_1^0, \gamma_1, \dots, \gamma_n \rangle \text{ acts nontrivially on every } \langle h \rangle\text{-orbit}\}.$$

The corresponding countings of covers (as usual with weight $1/\text{Aut}(p)$) differ from the cardinalities of these sets of Hurwitz tuples by the simultaneous conjugation of the Hurwitz tuple, hence by a factor of $d!$. Consequently, we let

$$(3) \quad N_d(\Pi) = \frac{|\text{Hur}_d(\Pi)|}{d!}, \quad N'_d(\Pi) = \frac{|\text{Hur}'_d(\Pi)|}{d!}, \quad N_d^0(\Pi) = \frac{|\text{Hur}_d^0(\Pi)|}{d!},$$

and package these data into the generating series

$$(4) \quad N(\Pi) = \sum_{d=0}^{\infty} N_d(\Pi)q^d, \quad N'(\Pi) = \sum_{d=0}^{\infty} N'_d(\Pi)q^d, \quad N^0(\Pi) = \sum_{d=0}^{\infty} N_d^0(\Pi)q^d.$$

The connected components of a covering induce a partition of the branch points of α_1^0 and $\gamma_1, \dots, \gamma_n$. This implies that

$$(5) \quad N'(\Pi) = N(\Pi)/N(\Pi_{\emptyset}).$$

Similarly, the inclusion–exclusion expression for counting connected covers in terms of covers without unramified components carries over from the case of torus covers (eg [17, Proposition 2.1]).

2.2 Covers of the projective line with three marked points

We need coverings of the projective line branched over three points with two types of parametrizations. As in the case of torus coverings (see [17, Section 2.2] for more details and remarks on numbered vs unnumbered enumeration) we define

$$\text{Cov}(\mathbf{w}^-, \mathbf{w}^+, \mu) = \{(\pi: S \rightarrow \mathbb{P}^1, \sigma_0, \sigma_\infty) : \deg(\pi) = \sum w_i^+ = \sum w_i^-, \pi^{-1}(1) = [\mu], \pi^{-1}(0) = \mathbf{w}^-, \pi^{-1}(\infty) = \mathbf{w}^+\}$$

to be the set of coverings of \mathbb{P}^1 with fixed profile μ over 1, with profile $\mathbf{w}^- = (w_1^-, \dots, w_n^-)$ and $\mathbf{w}^+ = (w_1^+, \dots, w_n^+)$ over 0 and ∞ , respectively, and where σ_0 and σ_∞ are labelings of the branch points over 0 and ∞ . We usually consider \mathbf{w}^- and \mathbf{w}^+ as “input” and “output” tuples of variables. We denote by

$$(6) \quad A(\mathbf{w}^-, \mathbf{w}^+, \mu) = \sum_{\pi \in \text{Cov}(\mathbf{w}^-, \mathbf{w}^+, \mu)} \frac{1}{\text{Aut}(\pi)}$$

the automorphism-weighted count of these numbers and refer to this quantity as *triple Hurwitz numbers* (although some authors, eg [26], call them double Hurwitz numbers, referring to two sets \mathbf{w}^\pm of free variables).

The second type of covering has only one set of variables and a product of transpositions of the point at ∞ . That is, we define

$$\text{Cov}_2(\mathbf{w}, \nu) = \left\{ (\pi: S \rightarrow \mathbb{P}^1, \sigma_0) : \deg(\pi) = \sum w_i, \pi^{-1}(1) = [\nu, 2^{(\deg(\pi)-|\nu|)/2}], \right. \\ \left. \pi^{-1}(0) = \mathbf{w}, \pi^{-1}(\infty) = [2^{\deg(\pi)/2}] \right\}$$

to be the set of coverings with fixed profile over 1 and ∞ (but stabilized by transpositions rather than by adding ones as usual!) and with variable profile $\mathbf{w} = (w_1, \dots, w_k)$ over 0. Finally, we let

$$(7) \quad A_2(\mathbf{w}, \nu) = \sum_{\pi \in \text{Cov}_2(\mathbf{w}, \nu)} \frac{1}{\text{Aut}(\pi)}$$

and we refer to them as *simple Hurwitz numbers with 2-stabilization*.

As usual, all these notions have their respective variants for coverings without unramified components (decorated by a prime) and for connected coverings (decorated by an upper zero).

2.3 Global graphs and cylinder decompositions

We normalize the pillow to be the quotient orbifold $B = E_i / \pm$, where $E_i = \mathbb{C} / \mathbb{Z}[i]$ is the rectangular torus provided with the unique up to scale quadratic differential q_B such that $\pi_T^* q_B$ is holomorphic on E . The pillow comes with the distinguished points P_1, \dots, P_4 that are the images of $0, \frac{i}{2}, \frac{i+1}{2}, \frac{1}{2} \in E_i$, respectively. For the remaining branch points we usually use in the sequel the *branch point normalization* that the i^{th} branch point P_i is the π_T -image of a point with coordinates $z_i = x_i + \sqrt{-1} \varepsilon_i$ with $0 \leq \varepsilon_5 < \varepsilon_6 < \dots < \varepsilon_{n+4} < \frac{1}{2}$ and any $x_i \in [0, 1)$.

Let Σ be the set of zeros and poles of $p^* q_B$ on X . The horizontal foliation with respect to q_B on B and thus on every pillowcase cover $p: X \rightarrow B$ with respect to $p^* q_B$ is periodic. It induces a decomposition of X into horizontal cylinders as follows: X is a union of maximal horizontal cylinders, ie maximal collections of closed regular horizontal geodesics (that are homotopic on $X \setminus \Sigma$), which are glued together along graphs (in fact, ribbon graphs) of horizontal saddle connections joining singularities in Σ (see Figure 2). Note that the preimages of P_2, P_3 and P_4 and those preimages of P_1 with ramification index 2 correspond to regular points of the flat metric induced by $p^* q_B$ on X ; they are not taken into account in this cylinder decomposition (as well as the regular preimages of the other P_i 's). We encode this cylinder decomposition by a global graph, as follows.




Figure 2: A pillowcase cover, the global graph and the local surfaces.

Definition 2.1 The *global graph* Γ associated with the pillowcase covering surface $(X, q = p^*q_B)$ of ramification profile Π is a graph Γ with $n + 1$ vertices. The vertices correspond to the branch points that belong to $\pi(\Sigma)$, ie the points $P_1, P_5, P_6, \dots, P_{n+4}$ when ν and μ are nonempty. The edges correspond to maximal horizontal cylinders of X . An edge is incident on a vertex if the image of the boundary of the cylinder contains the corresponding branch point.

If ν is nonempty, we put a special label 0 on the vertex corresponding to P_1 .

Note that the vertex 0 is distinguished from other vertices, as it corresponds to several ramification points (all odd singularities of q). Consequently different cylinder boundaries may correspond to that same vertex.

We illustrate this using Figure 2, which gives a covering in the stratum $\mathcal{Q}(2, 1, -1^3)$; in other terms, it has a ramification profile given by $n = 1, \nu = (3, 1, 1, 1)$ and $\mu_5 = 2$ in the notation of Section 2.1. (For more background on strata of quadratic differentials, including the notation $\mathcal{Q}(2, 1, -1^3)$, see for example [27].) The small triangles (with different orientations) are the three simple poles; the diamond indicates the simple zero. These points map to the black square on the pillow. The white circle indicates the double zero, mapping the white circle on the pillow. This corresponds to the point P_5 while P_1, \dots, P_4 are the corners of the pillow.




Figure 3: Orientation of half-edges and height minimum a_e .

We call the *special layer* at height 0 (resp. $\frac{1}{2}$) a connected component of the preimage of the horizontal segment of height 0 joining P_1 and P_4 in B (resp. horizontal segment of height $\frac{1}{2}$ joining P_2 and P_3). We provide Γ with an *orientation of its half-edges* as follows. We provide the pillowcase without the special layers with one of the two choices of an orientation of the vertical direction, say the upward-pointing. We orient a half-edge at a vertex v outward-pointing if the orientation of the cylinder pointing towards the boundary representing the half-edge is consistent with the chosen global (“vertical”) orientation, and we orient the half-edge inward-pointing otherwise. In particular, all the half-edges starting at the vertex 0 (if it exists) are oriented outward-pointing. Recall that cylinders may cross the special layers at height 0 or $\frac{1}{2}$ any number of times. The two half-edges corresponding to a cylinder are oriented consistently (see Figure 3, leftmost and third arrow) if and only if the cylinder crosses the special layers an even number of times. We refer to this extra datum as an *orientation* $G \in \Gamma$.

To reconstruct a pillowcase covering from a global graph, we need, as in the case of torus coverings, two types of extra data that encode the geometry of the cylinders and the geometry of the local surfaces, respectively. The first extra datum is the cylinder geometry.

Each cylinder (corresponding to an edge e) has an integral positive width w_e and a real positive height h_e . The heights h_e are not arbitrary, but related to the position of the branch points. To define the space parametrizing possible heights, we need a finer classification of the set $E(\Gamma)$.

Definition 2.2 For a graph Γ with an orientation G we define $E^0(G) = E^0(\Gamma)$ the set of edges that are adjacent to the vertex 0. We denote by $E^+(G)$ the set of edges that are consistently oriented. We use a lower index ℓ to denote loop edges, ie edges linking a vertex to itself.

For all nonloop edges we denote by $v^+(e)$ (resp. $v^-(e)$) the label of the vertex whose height according to the branch point normalization is higher (resp. lower). We refer to

them as the “upper” (resp. “lower”) vertex of the edge e . For all inconsistently oriented loop edges we extend this notation to $v^+(e) = v^-(e) = v(e)$.

The boundaries of a cylinder map to horizontal trajectories in B emanating at branch points v^+ and v^- . Given an orientation of the half-edges, the heights must thus lie in the *height space*

$$(8) \quad \tilde{\mathbb{N}}^{E(G)} = \{(h_e)_{e \in E(G)} : h_e \in \mathbb{N}_{>0} \text{ if } e \in E_\ell^0(G) \cup E_\ell^+(G) \text{ and } h_e - \Delta(e) \in \mathbb{N}_{\geq a_e} \text{ if } e \in E(G) \setminus (E_\ell^0(G) \cup E_\ell^+(G))\},$$

where $\Delta(e) = \pm \varepsilon_{v^+(e)} \pm \varepsilon_{v^-(e)}$, where the sign in front of each ε is positive if and only if the edge at the corresponding vertex is incoming and with a_e depending on the orientation as indicated in Figure 3.

In the example of Figure 2, the global graph has two vertices, the special vertex 0 and another one that corresponds to the branch point P_5 . The picture is drawn using the coordinate $z_5 = 0.25 + 0.2i$ for P_5 , so $\varepsilon_5 = 0.2$. The edges e_1 and e_2 corresponding to cylinders C_1 and C_2 are consistently oriented, and there are no loops. We have $\Delta(e_1) = \varepsilon_5 - \varepsilon_0 = \varepsilon_5 = \Delta(e_2)$ and $\Delta(e_3) = -\varepsilon_5 - \varepsilon_0 = -\varepsilon_5$, and $a_1 = a_2 = 0$ and $a_3 = 1$. The height space is then given by the condition

$$h_1 \in \varepsilon_5 + \mathbb{N}_{\geq 0}, \quad h_2 \in \varepsilon_5 + \mathbb{N}_{\geq 0}, \quad h_3 \in -\varepsilon_5 + \mathbb{N}_{\geq 1} = \left(\frac{1}{2} - \varepsilon_5\right) + \mathbb{N}_{\geq 0},$$

and parametrizes the possible heights for the cylinders C_1 , C_2 and C_3 . In the picture, $h_1 = h_2 = 1.2$ and $h_3 = 0.8$.

We claim that the collection of heights of cylinders in a pillowcase covering belongs to the height space and that, conversely, each element in the height space can be realized by such a covering. The integrality of the heights corrected by $\Delta(e)$ follows directly from the branch point normalization and the conventions of half-edge markings. It remains to justify the lower bounds one for the corrected heights. This happens if and only if the cylinder has to go all the way up to the preimage of the height $\frac{1}{2}$ line and down again. Loops based at 0 and consistently oriented loops have this property. For the remaining loops, it depends on the orientation of the half-edges. Note that (second and fourth case in Figure 3) the lower bound a_e is independent of the choice. For nonloop edges the integer a_e encodes whether the cylinder has to go around the pillowcase. This completes the proof of the claim.

The last piece of local information for a cylinder is the *twist* $t_e \in \mathbb{Z} \cap [0, w_e - 1]$. The twist depends on the choice of a ramification point $P^-(e)$ and $P^+(e)$ in each of the two components adjacent to the cylinders and it is defined as the integer part of the

real part, $\lfloor \Re(\int_s \omega) \rfloor$, of the integral along the unique straight line joining $P^-(e)$ to $P^+(e)$ such that $t_e \in [0, w_e - 1]$. The exact values of the twist will hardly matter in the sequel. It is important to retain simply that there are w_e possibilities for the twist in a given cylinder.

2.4 The basic correspondence theorem

The second extra datum needed for the correspondence theorem is the local geometry at the vertices. We let X^0 be the complement of the core curves of the cylinders. We call the union of connected components of X^0 that carry the same label the *local surfaces* of (X, ω) . We label these local surfaces also by an integer in $\{0, 1, \dots, n\}$ according to the ramification point they carry. This labeling is well-defined, since p is a cover without unramified components. The restriction of the cover p to any local surface besides the one corresponding to the special vertex is metrically the pullback of a flat cylinder branched over one point, as in the case of torus coverings. We thus encode these local surfaces by elements in $\text{Cov}'(\mathbf{w}_v^-, \mathbf{w}_v^+, \mu_v)$ where \mathbf{w}_v^- and \mathbf{w}_v^+ are the widths of the incoming and outgoing edges. The restriction of the cover p to a neighborhood of the line at height 0 is precisely the type of cover parametrized by an element in $\text{Cov}_2(\mathbf{w}, \nu)$, with \mathbf{w} the tuple of widths of the outgoing edges.

For the following proposition we fix a ramification profile Π and let ν or μ_v be the component of the tuple Π that corresponds to the vertex v under the vertex marking conventions explained in Section 2.3.

Proposition 2.3 *There is a bijective correspondence between*

- (i) *flat surfaces (X, q) with a covering $p: X \rightarrow B$ of degree $2d$ and profile Π of the pillow B without unramified components and with $q = \pi^* q_B$, and*
- (ii) *isomorphism classes of tuples $(G, (w_e, h_e, t_e)_{e \in E(G)}, (\pi_v)_{v \in V(G)})$ consisting of*
 - *a global graph Γ with labeled vertices including a special vertex 0 if v is nonempty, without isolated vertices, together with an orientation $G \in \Gamma$ of the half-edges such that all half-edges emerging from the special vertex are outgoing;*
 - *a collection of real numbers $(w_e, h_e, t_e)_{e \in E(G)}$ representing the width, height and twist of the cylinder corresponding to e , where the widths w_e*

are integers, the tuple of heights $(h_e)_{e \in E(G)} \in \tilde{\mathbb{N}}^{E(G)}$ is in the height space, $t_e \in \mathbb{Z} \cap [0, w_e - 1]$ and these numbers satisfy

$$(9) \quad 2 \sum_{e \in E(G)} w_e h_e = 2d;$$

- a collection of \mathbb{P}^1 -coverings $(\pi_v)_{v \in V(G) \setminus \{0\}} \in \text{Cov}'(\mathbf{w}_v^-, \mathbf{w}_v^+, \mu_v)$ without unramified components, where \mathbf{w}_v^- is the tuple of widths at the incoming edges at v , \mathbf{w}_v^+ is the tuple of widths at the outgoing edges at v , and μ_v is the ramification profile given by the labels at the vertex v ; and
- a \mathbb{P}^1 -covering $\pi_0 \in \text{Cov}'_2(\mathbf{w}_0, \nu)$, where \mathbf{w}_0 is the tuple of widths at the outgoing edges at $v = 0$ and ν is the ramification profile given by the labels at the vertex $v = 0$,

up to the action of the group $\text{Aut}(\Gamma)$ of automorphisms of the labeled graph Γ .

Proof With the setup and the orientation of half-edges adapted to pillowcase coverings, the proof proceeds exactly as in the case of torus covers; see [17, Proposition 2.4]. \square

2.5 Variants of the correspondence theorem

For the proof of the main theorem we will also need variants of the correspondence theorem that arise from counting covers by graphs while declaring a subset of points P_i for $i \in S \subset \{5, \dots, n + 4\}$ to be part of the layer of the special vertex. For extreme cases $S = \emptyset$, we are back in the situation of the previous situation, while for $S = \{5, \dots, n + 4\}$ the global graph is tautologically just a single vertex with no edges, decorated by a local Hurwitz number which is just the global Hurwitz number we are interested in.

For the concrete statement, we start with the *branch point normalization*. We place the points P_i at $z_i = x_i + \sqrt{-1}\varepsilon_i$, where now $0 < \varepsilon_i < \kappa$ for all $i \in S$ and $\kappa < \varepsilon_i < \frac{1}{2}$ for all $i \in S^c = \{5, \dots, n + 4\} \setminus S$, and moreover, within these constraints, strictly increasing with i .

The *global graph* associated with $(X, q = p^*q_B)$ is now the following graph Γ_S with $n + 1 - |S|$ vertices. The special vertex 0 corresponds to the region $R = \{0 \leq \Im(z) \leq \kappa\}$ and the remaining vertices are indexed by S^c . Edges correspond the cylinders that are not entirely contained in a connected component of $p^{-1}(R)$. The notion of an *orientation* $G_S \in \Gamma_S$ carries over verbatim from the above discussion, and the same holds for the height space $\tilde{\mathbb{N}}^{E_S(G)}$, declaring $\varepsilon_i = 0$ for $i \in S$.

The simplification in the graph is accounted for by a more complex Hurwitz number at the special vertex. We extend the definition of Hurwitz numbers with 2–stabilization by

$$\begin{aligned} & \text{Cov}_2(\mathbf{w}, \{\mu_i\}_{i \in S}, \nu) \\ &= \{(\pi: S \rightarrow \mathbb{P}^1, \sigma_0) : \deg(\pi) = \sum w_i, \pi^{-1}(0) = \mathbf{w}, \pi^{-1}(1) = [\nu, 2^{(\deg(\pi)-|\nu|)/2}], \\ & \quad \pi^{-1}(a_i) = [\mu_i] \text{ for } i \in S, \pi^{-1}(\infty) = [2^{\deg(\pi)/2}]\} \end{aligned}$$

for some points $a_i \notin \{0, 1, \infty\}$, and set

$$(10) \quad A_2(\mathbf{w}, \{\mu_i\}_{i \in S}, \nu) = \sum_{\pi \in \text{Cov}_2(\mathbf{w}, \{\mu_i\}_{i \in S}, \nu)} \frac{1}{\text{Aut}(\pi)}.$$

The same proof as above now yields the following proposition:

Proposition 2.4 *There is a bijective correspondence between*

- (i) *flat surfaces (X, q) with a covering $p: X \rightarrow B$ of degree $2d$ and profile Π of the pillow B without unramified components and with $q = \pi^*q_B$, and*
- (ii) *isomorphism classes of tuples $(G_S, (w_e, h_e, t_e)_{e \in E(G_S)}, (\pi_v)_{v \in V(G_S)})$ as in Proposition 2.3, with $\pi_0 \in \text{Cov}'_2(\mathbf{w}_0, \nu)$ replaced by $\pi_0 \in \text{Cov}'_2(\mathbf{w}_0, \{\mu_i\}_{i \in S}, \nu)$, up to the action of the group $\text{Aut}(\Gamma)$ of automorphisms of the labeled graph Γ .*

3 Fermionic Fock space and its balanced subspace

In this section we briefly recall the necessary background material about fermionic Fock space and the balanced subspace for the evaluation of the w –brackets that compute the generating functions of pillowcase covers. (See also [23; 22; 14] for this formalism.) The new result here is Theorem 3.3, stating that the generalized shifted symmetric functions f_ℓ and g_ν are rich enough to generate the algebra $\bar{\Lambda}$.

Recall the definition $f_\mu(\lambda) = |\mu|! \chi^\lambda(\mu) / (z(\mu) \dim \chi^\lambda)$ of the normalized characters, where $z(\mu) = \prod_{m=1}^\infty m^{r_m(\mu)} \prod_{m=1}^\infty r_m(\mu)!$ denotes the order of the centralizer of the partition $\mu = 1^{r_1} 2^{r_2} 3^{r_3} \dots$. We also write f_ℓ for the special case that σ is a ℓ –cycle. The algebra of shifted symmetric polynomials is defined as $\Lambda^* = \varprojlim \Lambda^*(n)$, where $\Lambda^*(n)$ is the algebra of symmetric polynomials in the n variables $\lambda_1 - 1, \dots, \lambda_n - n$. The functions

$$(11) \quad P_\ell(\lambda) = \sum_{i=1}^\infty \left((\lambda_i - i + \frac{1}{2})^\ell - (-i + \frac{1}{2})^\ell \right) \quad \text{and} \quad P_\mu = \prod_i P_{\mu_i}$$

belong to Λ^* . We add constant terms corresponding to regularizations of these functions to obtain

$$(12) \quad p_\ell(\lambda) = P_\ell(\lambda) + (1 - 2^{-\ell})\zeta(-\ell).$$

Here $\ell! \beta_{\ell+1} = (1 - 2^{-\ell})\zeta(-\ell)$ with β_k defined by $B(z) := \frac{z}{2} / \sinh(\frac{z}{2}) = \sum_{k=0}^\infty \beta_k z^k$. Recall the first basic structure result.

Theorem 3.1 [20] *The algebra Λ^* is freely generated by all the p_ℓ (or equivalently by the P_ℓ) with $\ell \geq 1$. The functions f_μ belong to Λ^* . More precisely, as μ ranges over all partitions, these functions f_μ form a basis of Λ^* .*

Let $f: \mathbb{P} \rightarrow \mathbb{Q}$ be an arbitrary function on the set \mathbb{P} of all partitions. We define (following [14]) the w -brackets

$$(13) \quad \langle f \rangle_w = \frac{\sum_{\lambda \in \mathbb{P}} w(\lambda) f(\lambda) q^{|\lambda|}}{\sum_{\lambda \in \mathbb{P}} w(\lambda) q^{|\lambda|}} \in \mathbb{Q}[[q]],$$

where the difference with the q -brackets used to compute torus coverings is the weight function

$$(14) \quad \sqrt{w(\lambda)} = \frac{\dim(\lambda)}{|\lambda|!} f_{2, \dots, 2}(\lambda)^2, \quad w(\lambda) = \sqrt{w(\lambda)}^2.$$

The main reason for introducing w -brackets is the expression

$$(15) \quad N'(\Pi) = \langle g_\nu f_{\mu_5} \cdots f_{\mu_{n+4}} \rangle_w$$

for the Hurwitz numbers without unramified components. This follows directly from the classical Burnside formula (see [13, Section 2]).

The algebra Λ^* is enlarged to the algebra

$$(16) \quad \bar{\Lambda} = \mathbb{Q}[p_\ell, \bar{p}_k \mid k, \ell > 0]$$

of shifted symmetric quasipolynomials, where

$$(17) \quad \bar{p}_k(\lambda) = \sum_{i \geq 0} ((-1)^{\lambda_i - i + 1} (\lambda_i - i + \frac{1}{2})^\ell - (-1)^{-i + 1} (-i + \frac{1}{2})^\ell) + \gamma_k,$$

and where the constants γ_i are zero for i odd, $\gamma_0 = \frac{1}{2}$, $\gamma_2 = -\frac{1}{8}$, $\gamma_4 = \frac{5}{32}$ and in general are defined by the expansion $C(z) = 1/(e^{z/2} + e^{-z/2}) = \sum_{k \geq 0} \gamma_k z^k / k!$. We provide the algebra $\bar{\Lambda}$ with a grading by defining the generators to have

$$\text{wt}(p_k) = k + 1 \quad \text{and} \quad \text{wt}(\bar{p}_k) = k.$$

The main reason for introducing Λ^* is the following result. The reason for introducing g_ν will become clear by (15) in Section 4. In this section ν is always a partition consisting of an even number of odd parts.

Theorem 3.2 [14, Theorem 2] *There is a function $g_\nu \in \bar{\Lambda}$ of (mixed) weight less than or equal to $\frac{1}{2}|\nu|$ such that*

$$(18) \quad g_\nu(\lambda) = \frac{f_{(\nu, 2, 2, \dots)}(\lambda)}{f_{(2, 2, \dots)}(\lambda)} \quad \text{for } \lambda \text{ balanced.}$$

Our goal is the following converse, for which we define $\text{wt}(g_\nu) = \frac{1}{2}|\nu|$:

Theorem 3.3 *The elements g_ν generate $\bar{\Lambda}$ as a graded Λ^* -module, ie the subspace of $\bar{\Lambda}$ of weight less than or equal to n is generated by expressions hg_ν for $h \in \Lambda^*$ with $\text{wt}(h) + \text{wt}(g_\nu) \leq n$.*

Of course, the elements g_ν do not form a basis as there are many more g_ν than products of \bar{p}_k for a given weight. We recall the main steps of the proof of Theorem 3.2, since we need them for Theorem 3.3.

3.1 Fermionic Fock space

Let $\Lambda_0^{\infty/2}V$ be the *charge zero subspace of the half-infinite wedge or fermionic Fock space* over the countably infinite-dimensional vector space V . We denote the basis elements of V by underlined half-integers. An orthonormal basis of $\Lambda_0^{\infty/2}V$ is given by the elements

$$v_\lambda = \underline{\xi}_1 \wedge \underline{\xi}_2 \wedge \underline{\xi}_3 \wedge \dots, \quad \xi_i = \lambda_i - i + \frac{1}{2},$$

indexed by partitions $\lambda = (\lambda_1 \geq \lambda_2 \geq \dots)$. The basic operators on the half-infinite wedge is for any $k \in \mathbb{Z} + \frac{1}{2}$ the *creation operator* $\psi_k(v) = \underline{k} \wedge v$ and its adjoint ψ^* , the *annihilation operator*. For any function f on the real line we define the (unregularized) operators

$$\tilde{\mathcal{E}}_k[f] = \sum_{m \in \mathbb{Z} + \frac{1}{2}} f(m) : \psi_{m-k} \psi_m^* :,$$

where the colons denote the normally ordered product. We use the convention that x is the default variable on the real line. Consequently, if T is a term in x (typically a polynomial, or a character like $(-1)^x$ times a polynomial), we write $\tilde{\mathcal{E}}_k[T]$ as shorthand

for $\tilde{\mathcal{E}}_k[x \mapsto T(x)]$. The regularized operators for exponential arguments are defined by

$$\mathcal{E}_k[e^{zx}] := \mathcal{E}_k[x \mapsto e^{zx}] = \tilde{\mathcal{E}}_k[e^{zx}] + \delta_{0,k} \frac{B(z)}{z}$$

and in the presence of a character $(-1)^x$ by

$$\mathcal{E}_k[e^{(z+\pi i)x}] := \mathcal{E}_k[x \mapsto e^{(z+\pi i)x}] = \tilde{\mathcal{E}}_k[x \mapsto e^{(z+\pi i)x}] - i \delta_{k,0} C(z),$$

where

$$C(z) := \frac{1}{e^{z/2} + e^{-z/2}} = \sum_{k \geq 0} \gamma_k \frac{z^k}{k!}.$$

We frequently need three special cases of these operators. First,

$$\alpha_{-n} = \mathcal{E}_{-n}[1] = \sum_{m \in \mathbb{Z} + \frac{1}{2}} : \psi_{m+n} \psi_m^* :,$$

whose adjoint is denoted by $\alpha_n = \alpha_{-n}^*$. The *Murnaghan–Nakayama rule* says that

$$(19) \quad \prod_i \alpha_{-\mu_i} v_\emptyset = \sum_\lambda \chi^\lambda(\mu) v_\lambda.$$

Second, the expansion of the (regularized) formal power series

$$\mathcal{E}_0(z) := \mathcal{E}_0[x \mapsto e^{zx}] = \frac{1}{z} + \sum_{\ell \geq 1} \mathcal{P}_\ell \frac{z^\ell}{\ell!}$$

and the expansion of

$$i \mathcal{E}_0[x \mapsto e^{(z+\pi i)x}] = \sum_{\ell \geq 0} \frac{z^\ell}{\ell!} \bar{\mathcal{P}}_\ell$$

gives operators \mathcal{P}_ℓ and $\bar{\mathcal{P}}_k$ with the property

$$\mathcal{P}_\ell v_\lambda = p_\ell(\lambda) v_\lambda, \quad \bar{\mathcal{P}}_k v_\lambda = \bar{p}_k(\lambda) v_\lambda.$$

Finally, note that the unregularized $\mathcal{E}_0(z)$ -operator admits the useful formula

$$(20) \quad \tilde{\mathcal{E}}_0(z) = [y^0] \psi(e^z y) \psi^*(y),$$

where the interior expression can be checked by the commutator lemma for vertex operators (see [23, Lemma 7.1] or [18, Section 14]) to be

$$(21) \quad \begin{aligned} & \psi(xy) \psi^*(y) \\ &= \frac{1}{x^{1/2} - x^{-1/2}} \exp\left(\sum_{n>0} \frac{(xy)^n - y^n}{n} \alpha_{-n}\right) \exp\left(\sum_{n>0} \frac{y^{-n} - (xy)^{-n}}{n} \alpha_n\right). \end{aligned}$$

3.2 The balanced subspace

Note that the definition of the algebra Λ excludes the operator p_0 , the charge operator, since it is equal to zero on $\Lambda_0^{\infty/2}V$. Similarly, the definition of the algebra $\bar{\Lambda}$ excludes the operator

$$\bar{p}_0(\lambda) = \frac{1}{2} + \sum_{i \geq 0} ((-1)^{\lambda_i - i + 1} - (-1)^{-i + 1}).$$

A partition λ is called *balanced* if among the $\lambda_i - i + 1$ for $\lambda_i \geq 0$ there are as many odd as even numbers, ie if and only if $\bar{p}_0(\lambda) = \frac{1}{2}$. Every partition λ determines (by sorting the $\lambda_i - i + 1$ into even and odd) two partitions α and β , called the *2-quotients*, such that

$$\{\lambda_i - i + \frac{1}{2}\} = \{2(\alpha_i - i + \frac{1}{2}) + \bar{p}_0(\lambda)\} \cup \{2(\beta_i - i + \frac{1}{2}) - \bar{p}_0(\lambda)\}.$$

We let $\Lambda^{\text{bal}}V$ denote the *balanced subspace* of $\Lambda_0^{\infty/2}V$, ie the subspace spanned by the v_λ for λ balanced. It inherits from $\Lambda_0^{\infty/2}V$ the grading by eigenspace of the energy operator, ie $\Lambda^{\text{bal}}V = \bigoplus_{d \geq 0, \text{even}} \Lambda^{\text{bal}}V_d$. We use the shorthand notation

$$|[\rho; \bar{\rho}]\rangle = \frac{1}{\mathfrak{z}(\rho)\mathfrak{z}(\bar{\rho})} \prod_i \alpha_{-\rho_i} \prod_j \bar{\alpha}_{-\bar{\rho}_j} v_\emptyset$$

for the following reason [13]:

Proposition 3.4 For $\rho = (\rho_i)$ and $\bar{\rho} = (\bar{\rho}_i)$ running over all partitions with entries in $2\mathbb{Z}$, the elements $|[\rho; \bar{\rho}]\rangle$ form an orthogonal basis of $\Lambda^{\text{bal}}V$.

Proof of Theorem 3.2 By (19) the content of the theorem is that the orthogonal projection of $|[v, 2^{d-|v|/2}; \emptyset]\rangle$ to the balanced subspace $\Lambda^{\text{bal}}V$ is a linear combination of the projections of $|\prod_i \mathcal{P}_{\mu_i} \prod \bar{\mathcal{P}}_{\bar{\mu}_i} 2^d\rangle$ with $\mu = (\mu_i)_{i \geq 1}$ and $\bar{\mu} = (\bar{\mu}_i)_{i \geq 1}$ partitions with $\text{wt}(\mathcal{P}_\mu) + \text{wt}(\bar{\mathcal{P}}_{\bar{\mu}}) \leq \frac{1}{2}|v|$ with coefficients independent of d . For this purpose one calculates, using the commutation laws of the vertex operators, on the one hand that

$$(22) \quad \langle [\rho; \bar{\rho}] | [v, 2^{d-|v|/2}; \emptyset] \rangle = \frac{2^{\ell(v) - \ell(\bar{\rho})}}{2^{d-|v|/2} (d - |v|/2)! \mathfrak{z}(v) \mathfrak{z}(\bar{\rho})} C(v, \bar{\rho}) \quad \text{if } \rho = 2^{d-|v|/2},$$

where $C(v, \bar{\rho})$ is the number of ways to assemble the parts of $\bar{\rho}$ from the parts of v , and zero otherwise. In particular, $|v| = |\bar{\rho}|$ for (22) to be nonzero. The squared norms of the element $|[\rho; \bar{\rho}]\rangle$ for ρ and $\bar{\rho}$ having even parts only is equal to $1/\mathfrak{z}(\rho)\mathfrak{z}(\bar{\rho})$. In particular, the scalar product (22) divided by $\|[\rho; \bar{\rho}]\|^2$ is independent of d .

On the other hand, one computes using the commutation laws of the vertex operators that the brackets

$$(23) \quad D_{[\rho; \bar{\rho}], (\mu, \bar{\mu})}(d) = \left\langle [(\rho, 2^{(d-|\rho|)/2}); \bar{\rho}] \mid \prod_i \mathcal{P}_{\mu_i} \prod \bar{\mathcal{P}}_{\bar{\mu}_i} \mid 2^d \right\rangle / \|[(\rho, 2^{(d-|\rho|)/2}); \bar{\rho}]\|^2$$

for ρ an partition with only even parts of length different from two are nonzero only if

$$(24) \quad \Delta(\rho, \bar{\rho}, \mu, \bar{\mu}) := \text{wt}(\rho) + \text{wt}(\bar{\rho}) - \text{wt}(\mu) - \text{wt}(\bar{\mu}) \geq 0.$$

Since an additional factor \mathcal{P}_1 in (23) gives an additional factor $d - \frac{1}{24}$, we first consider $D_{[\rho; \bar{\rho}], (\mu, \bar{\mu})}(d)$ with μ without a part equal to one. Then, if (24) is attained, the scalar product $D_{[\rho; \bar{\rho}], (\mu, \bar{\mu})}(d)$ is nonzero if and only if $\mu = \frac{1}{2}\rho$ and $\bar{\mu} = \frac{1}{2}\bar{\rho}$. In general, $D_{[\rho; \bar{\rho}], (\mu, \bar{\mu})}(d)$ is a polynomial of degree $\frac{1}{2}\Delta(\rho, \bar{\rho}, \mu, \bar{\mu})$. This also implies that (for fixed weight of $[\rho; \bar{\rho}]$) the matrix $D_{[\rho; \bar{\rho}], (\mu, \bar{\mu})}(d)$ (with entries in $\mathbb{Q}[d]$) is an invertible matrix D , in fact block triangular.

Using these facts we can write

$$(25) \quad g_\nu = \sum_{|\bar{\rho}|=|\nu|} \frac{2^{\ell(\nu)-\ell(\bar{\rho})}}{\mathfrak{z}(\nu)} C(\nu, \bar{\rho}) \sum_{\mu, \bar{\mu}} (D^{-1})_{[\emptyset; \bar{\rho}], (\mu, \bar{\mu})} (p_1 + \frac{1}{24}) p_\mu \bar{p}_{\bar{\mu}},$$

where the sum is over all $(\mu, \bar{\mu})$ with $|\mu| + \ell(\mu) + |\bar{\mu}| \leq \frac{1}{2}|\nu|$. □

Lemma 3.5 For every fixed d the matrix $C(\nu, \bar{\rho})$, where ν is a partition of $2d$ consisting only of odd parts and $\bar{\rho}$ is a partition of $2d$ consisting only of even parts, has full rank equal to $\mathbb{P}(d)$.

Proof We order the rows $\bar{\rho}$ lexicographically and consider the submatrix with columns ν consisting of the partitions

$$\nu(\rho) = (\bar{\rho}_1 - 1, 1, \bar{\rho}_2 - 1, 1, \dots, \bar{\rho}_n - 1, 1)$$

formed from $\bar{\rho} = (\bar{\rho}_1 \geq \dots \geq \bar{\rho}_n)$. Since $C(\nu(\bar{\rho}), \bar{\rho}') \neq 0$ if and only if $\bar{\rho}' \leq \bar{\rho}$ lexicographically, the claim follows. □

Proof of Theorem 3.3 Since the matrix $D_{[\rho; \bar{\rho}], (\mu, \bar{\mu})}(d)$ is invertible, it suffices to prove by induction on the weight that the operators of contraction against

$$b_{\rho, \bar{\rho}} = \frac{1}{\|[(\rho, 2^{(d-|\rho|)/2}); \bar{\rho}]\|^2} \langle [(\rho, 2^{(d-|\rho|)/2}); \bar{\rho}] \mid$$

are in the Λ^* -module generated by the g_ν . For $\rho = \emptyset$ this follows from Lemma 3.5; in fact, those $b_{\emptyset, \bar{\rho}}$ can be spanned by g_ν with constant coefficients. For $\rho \neq \emptyset$ we use the expression of $b_{\rho, \bar{\rho}}$ as a linear combination of $p_\mu \bar{p}_\mu$. The terms with $\text{wt}(p_\mu \bar{p}_\mu) = \text{wt}(b_{\rho, \bar{\rho}})$ are either $p_{\rho/2} \bar{p}_{\bar{\rho}/2}$ or involve a factor of p_1^j for some $j > 0$ by the properties of the matrix $D_{[\rho; \bar{\rho}], (\mu, \bar{\mu})}(d)$. Consequently, these terms are generated by g_ν as a Λ^* -module by the induction hypothesis and the extra factor p_1^j does not alter this fact. For the terms with smaller weight the induction hypothesis applies directly. \square

4 Hurwitz numbers and graph sums

The goal of this section is to use the correspondence theorems to express any w -bracket in terms of auxiliary brackets that directly reflect the graph sums of the correspondence theorems. The precise form of the goal, Theorem 4.2, will involve in the auxiliary brackets only arguments for which the $A'(\cdot)$ -functions will later be proven to be polynomial.

We first define for any function F on partitions

$$(26) \quad A(\mathbf{w}^-, \mathbf{w}^+, F) = \frac{1}{\prod_i w_i^- \prod_i w_i^+} \sum_{|\lambda|=d} \chi^\lambda(\mathbf{w}^-) \chi^\lambda(\mathbf{w}^+) F(\lambda)$$

and we define the variant without unramified connected components, denoted by $A'(\mathbf{w}^-, \mathbf{w}^+, F)$, by the usual inclusion–exclusion formula (eg [17, equation (17) or (25)]). In our case $A'(\mathbf{w}^-, \mathbf{w}^+, F)$ equals $A^0(\mathbf{w}^-, \mathbf{w}^+, F)$, the connected variant, since there is only one ramification point. The reason for this definition is that on one hand the triple Hurwitz number introduced in (6) can be written using the Burnside lemma (see eg [17, Section 2]) as

$$A'(\mathbf{w}^-, \mathbf{w}^+, \mu) = A'(\mathbf{w}^-, \mathbf{w}^+, f_\mu).$$

On the other hand, we will use that the function with completed cycles argument

$$\bar{A}'(\mathbf{w}^-, \mathbf{w}^+, \mu) := A'\left(\mathbf{w}^-, \mathbf{w}^+, \frac{P_\mu}{\prod \mu_i}\right)$$

is a polynomial of even degree for $\mu = (\mu_1)$ being a partition consisting of a single part and for $\mu_1 + 1 - \ell(\mathbf{w}^-) - \ell(\mathbf{w}^+)$ even (see [26], rephrased as [17, Theorem 4.1]).

A new feature of pillowcase covers is the use of the one-variable analog

$$(27) \quad A_2(\mathbf{w}, F) = \frac{1}{\prod_i w_i} \sum_{|\lambda|=d} \sqrt{w(\lambda)} F(\lambda),$$

where the second variable has been replaced by the character for the fixed partition $(2, \dots, 2)$. We define the version without unramified connected components $A'_2(\mathbf{w}, F)$ by the usual inclusion–exclusion formula. In general it does not correspond to connected covers here. Again, the reason for this definition is two-fold. By the Burnside formula the simple Hurwitz numbers with 2–stabilization introduced in (7) and generalized in (10) can be written as

$$(28) \quad A'_2(\mathbf{w}, \{\mu_i\}_{i \in S}, \nu) = A'_2\left(\mathbf{w}, g_\nu \prod_{i \in S} f_{\mu_i}\right).$$

We study polynomiality properties of A'_2 for suitable $F = \prod \bar{p}_{k_i} \in \bar{\Lambda}$ in detail in Section 7.

Let Π be a profile as specified in Section 2.1. We decompose the Hurwitz number $N'(\Pi)$ according to the contribution of the global graphs, ie we write

$$N'(\Pi) = \frac{1}{|\text{Aut}(\Gamma)|} \sum_{\Gamma} N'(\Gamma, \Pi),$$

where the sum is over all (not necessarily connected) graphs Γ with $n = |\Pi|$ nonspecial labeled vertices and possibly a special vertex labeled 0 and where $\text{Aut}(\Gamma)$ are the automorphisms of the graph Γ that respect the vertex labeling. (Note that Γ has neither a labeling nor an orientation on the edges.) Following the results in the correspondence theorem we define an *admissible orientation* G of Γ (symbolically written as $G \in \Gamma$) to be an orientation of the half-edges of Γ such that all the half-edges at the special vertex 0 (if it exists) are outward-pointing. Now the following proposition is an immediate consequence of the correspondence theorem Proposition 2.3:

Proposition 4.1 *The contributions of individual labeled graphs to $N'(\Pi)$ can be expressed in terms of triple Hurwitz numbers as*

$$(29) \quad N'(\Gamma, \Pi) = \sum_{G \in \Gamma} N'(G, \Pi),$$

where

$$(30) \quad N'(G, \Pi) = \sum_{\substack{h \in \tilde{\mathbb{N}}^{E(G)} \\ w \in \mathbb{Z}_+^{E(G)}}} \prod_{e \in E(G)} w_e q^{h_e w_e} \cdot A'_2(\mathbf{w}_0, \nu) \prod_{v \in V(G) \setminus \{0\}} A'(\mathbf{w}_v^-, \mathbf{w}_v^+, \mu_\nu) \delta(v),$$

where $V(G)^* = V(G) \setminus \{0\}$ and where

$$(31) \quad \delta(v) = \delta \left(\sum_{i \in e_+(v)} w_i^+ - \sum_{i \in e_-(v)} w_i^- \right).$$

We formalize the type of expression appearing in the previous proposition by defining *auxiliary brackets*

$$(32) \quad \begin{aligned} [F_1, \dots, F_n; F_0] &= \sum_{\Gamma} [F_1, \dots, F_n; F_0]_{\Gamma}, \\ [F_1, \dots, F_n; F_0]_{\Gamma} &= \sum_{G \in \Gamma} [F_1, \dots, F_n; F_0]_G, \end{aligned}$$

where the sum is over all labeled graphs Γ with n vertices and over all admissible orientations, respectively, and where

$$\begin{aligned} [F_1, \dots, F_n; F_0]_G &= \sum_{\substack{h \in \tilde{N}^{E(G)} \\ w \in \mathbb{Z}_+^{E(G)}}} \prod_{i \in E(G)} w_i q^{h_i w_i} \cdot A'_2(\mathbf{w}_0, F_0) \prod_{v \in V(G)^*} A'(w_v^-, w_v^+, F_{\#v}) \delta(v). \end{aligned}$$

Here $\#v$ denotes the label of the vertex v . This notation is designed so that Proposition 4.1 can be restated as

$$(33) \quad \langle f_{\mu_1} \cdots f_{\mu_n} g_v \rangle_w = [f_{\mu_1}, \dots, f_{\mu_n}; g_v].$$

More generally, by verbatim the same proof, Proposition 2.4 can be restated as the generalization

$$(34) \quad \langle f_{\mu_1} \cdots f_{\mu_n} g_v \rangle_w = \left[\prod_{i \notin S} f_{\mu_i}; \prod_{i \in S} f_{\mu_i} g_v \right]$$

for any subset $S \subseteq \{1, \dots, n\}$.

We are now ready to formulate the goal of this section in detail.

Theorem 4.2 *The w -bracket of any element in $\bar{\Lambda}$ can be expressed as a finite linear combination of the auxiliary brackets, ie for every $\ell = (\ell_1, \dots, \ell_n)$ and every $\mathbf{k} = (\ell_1, \dots, k_m)$ there exist $c_{(\mathbf{t}, \mathbf{s})} \in \mathbb{Q}$ (depending on (ℓ, \mathbf{k})) such that*

$$(35) \quad \left\langle \prod_{j=1}^n p_{\ell_j} \prod_{i=1}^m \bar{p}_{k_i} \right\rangle_w = \sum_{(\mathbf{t}, \mathbf{s})} c_{(\mathbf{t}, \mathbf{s})} \left[p_{t_1}, \dots, p_{t_{\ell(\mathbf{t})}}; \prod_{i=1}^{\ell(\mathbf{s})} \bar{p}_{s_i} \right],$$

where the sum is over all (\mathbf{t}, \mathbf{s}) with $\sum_j (t_j + 1) + \sum_i s_i \leq \sum_j (\ell_j + 1) + \sum_i k_i$.

Proof The proof is by induction on the weight $w = \sum_{i=1}^m k_i$ of the \bar{p}_{k_i} -part of the bracket, the case of weight zero being trivial (no special vertex, ie as in the abelian case).

By Theorem 3.3 we can write the left-hand side of (35) as a linear combination of $\langle f_{\mu_1} \cdots f_{\mu_n} g_\nu \rangle_w$ with $\text{wt}(g_\nu) \leq w$. By (33) each such summand is equal to

$$(36) \quad [f_{\mu_1}, \dots, f_{\mu_n}; g_\nu] = \sum_{\mathbf{b}, \mathbf{a}} c(\mathbf{b}, \mathbf{a}) \left[f_{\mu_1}, \dots, f_{\mu_n}; \prod_{j \geq 1} p_{b_j} \prod_{i \geq 1} \bar{p}_{a_i} \right],$$

where the sum is over all partitions \mathbf{a} and \mathbf{b} , by Theorem 3.2. In the summands where \mathbf{b} is the empty partition, we replace f_{μ_i} by a linear combination of products of p_ℓ thanks to Theorem 3.1, and these contributions are of the required form of the right-hand side of (35). In all the summands with \mathbf{b} nonempty we use the converse base change of Theorem 3.1 to write the product of p_{b_j} as a linear combination of a product of f_{μ_j} . This is done using that p_k is a linear combination of f_k and products of p_l 's with $l < k$, and using induction on the weight. We can now use (34) from right to left to express all the terms as a sum of w -brackets with \bar{p}_{k_i} -part of weight $w - \sum_j b_j + 1$. Since \mathbf{b} is nonempty, we conclude thanks to the induction hypothesis. \square

5 Constant coefficients of quasielliptic functions

In this section we consider the constant coefficient (in z_1, \dots, z_n) of a function $f(\tau; z_1, \dots, z_n)$ that is quasielliptic, has a quasimodular transformation behavior and has poles at most at 2-torsion translates of the coordinate axes and diagonals. We show in Theorem 5.6 that this constant coefficient is indeed a quasimodular form for the subgroup $\Gamma(2)$ of $\text{SL}(2, \mathbb{Z})$.

5.1 Quasimodular forms

A *quasimodular form* for the cofinite Fuchsian group $\Gamma \subset \text{SL}(2, \mathbb{R})$ of weight k is a function $f: \mathbb{H} \rightarrow \mathbb{C}$ that is holomorphic on \mathbb{H} and the cusps of Γ and such that there exists an integer p and holomorphic functions $f_i: \mathbb{H} \rightarrow \mathbb{C}$ such that

$$(c\tau + d)^{-k} f\left(\frac{a\tau + b}{c\tau + d}\right) = \sum_{i=0}^p f_i(\tau) \left(\frac{c}{c\tau + d}\right)^i \quad \text{for all } \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma.$$

The smallest integer p with the above property is called the *depth* of the quasimodular form. By definition, quasimodular forms of depth zero are simply modular forms. The

basic examples of quasimodular forms are the Eisenstein series defined by

$$G_{2k}(\tau) = \frac{(2k - 1)!}{2(2\pi i)^{2k}} \sum_{(m,n) \in \mathbb{Z}^2 \setminus \{(0,0)\}} \frac{1}{(m + n\tau)^{2k}} = -\frac{B_{2k}}{4k} + \sum_{n=1}^{\infty} \sigma_{2k-1}(n)q^n.$$

Here B_l is the Bernoulli number, σ_l is the divisor sum function and $q = e^{2\pi i \tau}$. For $k \geq 2$ these are modular forms, while for $k = 1$ the Eisenstein series

$$G_2(\tau) = -\frac{1}{24} + \sum_{n=1}^{\infty} \sigma_1(n)q^n$$

is a quasimodular form of weight 2 and depth 1 for $SL(2, \mathbb{Z})$. By [19] we can write the ring of quasimodular forms for any group Γ with $Stab_{\infty}(\Gamma) = \pm \begin{pmatrix} 1 & \\ & 1 \end{pmatrix}$ as $QM(\Gamma) = \mathbb{C}[G_2] \otimes M(\Gamma)$ in terms of the ring of modular forms $M(\Gamma)$. Recall that $\Gamma(n)$ denotes the subgroup of $SL(2, \mathbb{Z})$ of matrices congruent to the identity mod n and that $\Gamma_0(n)$ denotes the subgroup of $SL(2, \mathbb{Z})$ congruent to $\begin{pmatrix} 1 & * \\ 0 & 1 \end{pmatrix}$ mod n . We will be mainly interested in the congruence groups $\Gamma_0(2)$ and

$$\Gamma(2) = \begin{pmatrix} 2 & 0 \\ 0 & 1 \end{pmatrix} \Gamma_0(4) \begin{pmatrix} 2 & 0 \\ 0 & 1 \end{pmatrix}^{-1} \subset \Gamma_0(2).$$

Since $M(\Gamma_0(2))$ is freely generated by $G_2^{\text{odd}}(\tau) = G_2(\tau) - 2G_2(2\tau)$ and $G_4(2\tau)$ by the transformation formula (eg [8, Proposition 4.2.1]) and the usual dimension formula for modular forms, we deduce that

$$(37) \quad QM(\Gamma_0(2)) \cong \mathbb{C}[G_2(\tau), G_2(2\tau), G_4(2\tau)]$$

is a polynomial ring. For $\Gamma_0(4)$ we restrict our attention to the subring of even-weight quasimodular forms. Since $M_{2*}(\Gamma_0(4))$ is freely generated by $G_2^{\text{odd}}(\tau)$ and $G_2^{\text{odd}}(2\tau)$ (again using the transformation formula together with the isomorphism $\Gamma_0(4) \cong \Gamma(2)$ given by conjugation with $\text{diag}(2, 1)$), we deduce that

$$(38) \quad QM(\Gamma(2)) \cong \mathbb{C}[G_2(\frac{1}{2}\tau), G_2(\tau), G_2(2\tau)].$$

We use the notation $q = e^{2\pi i \tau}$, hence $q^{1/2} = e^{\pi i \tau}$. Note that a typical element $QM(\Gamma(2))$ has a Fourier expansion in $q^{1/2}$. The following observation allows us to prove quasimodularity by the larger group $\Gamma_0(2)$:

Lemma 5.1 *For all $k \in \mathbb{N}$ the Eisenstein series $G_{2k}(\frac{1}{2}\tau)$, $G_{2k}(\tau)$ and $G_{2k}(2\tau)$ are quasimodular forms for $\Gamma(2)$. Moreover, any even-weight quasimodular form for $\Gamma(2)$ whose Fourier expansion is a series in q is in fact a quasimodular form for $\Gamma_0(2)$.*

Proof The second statement follows immediately from (37) and (38). □

5.2 Quasimodular forms as constant coefficients of quasielliptic functions

We are now ready to state the first main criterion for quasimodularity, involving the constant coefficients of some quasielliptic functions introduced below. We start with a general remark on the domains where the expansions are valid. Suppose that the meromorphic function $f(z_1, z_2, \dots, z_n; \tau)$ is periodic under $z_j \mapsto z_j + 1$ for each j and under $\tau \mapsto \tau + 1$. We can then write $f(z_1, z_2, \dots, z_n; \tau) = \bar{f}(\zeta_1, \dots, \zeta_n, q)$, where $\zeta_j = e^{2\pi iz_j}$ as above. For any permutation $\pi \in S_n$, we fix the domain

$$\Omega_\pi = |q|^{1/2} < |\zeta_{\pi(i)}| < |\zeta_{\pi(i+1)}| < 1 \quad \text{for all } i = 1, \dots, n-1.$$

On such a domain the *constant term with respect to all the ζ_i* is well-defined. It can be expressed as the integral

$$[\zeta_n^0, \dots, \zeta_1^0]_\pi \bar{f} = \frac{1}{(2\pi i)^n} \oint_{\gamma_n} \cdots \oint_{\gamma_1} f(z_1, \dots, z_n; \tau) dz_1 \cdots dz_n$$

along the integration paths

$$\gamma_j: [0, 1] \rightarrow \mathbb{C}, \quad t \mapsto iy_j + t,$$

where $0 \leq y_{\pi(1)} < y_{\pi(2)} < \cdots < y_{\pi(n)} < \frac{1}{2}$. We call these our *standard integration paths* for the permutation π . If the domain Ω_π is clear from the context, we also write $[\zeta^0]$ or $[\zeta_n^0, \dots, \zeta_1^0]$ as shorthand for the coefficient extraction $[\zeta_n^0, \dots, \zeta_1^0]_\pi$.

Let $\Delta = \Delta_\tau$ be the operator on meromorphic functions defined by

$$\Delta(f)(z) = f(z + \tau) - f(z).$$

A meromorphic function f is called *quasielliptic* (for the lattice $\mathbb{Z} + \tau\mathbb{Z}$) if $f(z + 1) = f(z)$ and if there exists some integer e such that $\Delta^e(f)$ is elliptic. The minimal such e is called the *order (of quasiellipticity)* of f .

We say that a meromorphic function $f: \mathbb{C}^n \times \mathbb{H} \rightarrow \mathbb{C}$ is quasielliptic if it is quasielliptic in each of the first n variables. For such a function we write $e = (e_1, \dots, e_n)$ for the tuple of orders of quasiellipticity in the n variables. Consequently, a quasielliptic function of order $(0, \dots, 0)$ is simply an elliptic function.

We write Δ_i for the operator Δ acting on the i^{th} variable. Note that these operators Δ_i commute. Let $T = \{0, \frac{1}{2}, \frac{1}{2}\tau, \frac{1}{2}(1 + \tau)\}$ be the set of 2-torsion points.

The functions we want to take constant coefficients of belong to the space in the following definition. It is similar to the quasielliptic quasimodular forms used in

[17, Definition 5.5]. The difference consists of allowing 2–torsion translates for the poles and requiring a modular transformation law for a smaller group. We do not decorate our new definition of $\mathcal{Q}_{n,e}^{(k)}$ by an extra symbol 2 to avoid overloading notation.

Definition 5.2 We define for $n \geq 0$, $k \geq 0$ and $e \geq 0$ the vector space of $\mathcal{Q}_{n,e}^{(k)}$ of quasielliptic quasimodular forms for $\Gamma(2)$ to be the space of meromorphic functions f on $\mathbb{C}^n \times \mathbb{H}$ in the variables $(z_1, \dots, z_n; \tau)$ that

- (i) have poles on \mathbb{C}^n at most at the $\mathbb{Z} + \tau\mathbb{Z}$ –translates of the diagonals $z_i = z_j$ and $z_i = -z_j$ and the 2–torsion points $z_i \in T$,
- (ii) are quasielliptic of order e , and
- (iii) are quasimodular of weight k for $\Gamma(2)$, ie f is holomorphic in τ on $\mathbb{H} \cup \{0, 1, \infty\}$ and there exists some $p \geq 0$ and functions $f_i(z_1, \dots, z_n; \tau)$ that are holomorphic in τ and meromorphic in the z_i such that

$$(c\tau + d)^{-k} f\left(\frac{z_1}{c\tau + d}, \dots, \frac{z_n}{c\tau + d}; \frac{a\tau + b}{c\tau + d}\right) = \sum_{i=0}^p f_i(z_1, \dots, z_n; \tau) \left(\frac{c}{c\tau + d}\right)^i$$

for all $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma(2)$.

Examples of such quasielliptic quasimodular forms will be constructed from the propagator, the shift $P(z; \tau) = (1/(2\pi i)^2)\wp(z; \tau) + 2G_2(\tau)$ of the Weierstrass \wp –function, and from the shift $Z(z; \tau) = -\zeta(z; \tau)/2\pi i + 2G_2(\tau)2\pi iz$ of the Weierstrass ζ –function. The reason for this shift, as well as the Fourier and Laurent series expansion of these functions, is summarized in [17, Section 5.2]. In particular we will need

$$P_{\text{even}}(z; \tau) = 2P(2z; 2\tau) \quad \text{and} \quad P_{\text{odd}}(z; \tau) = P(z; \tau) - P_{\text{even}}(z; \tau).$$

Proposition 5.3 The functions $P^{(k)}(z_i - a; \tau)$ for $a \in T$ and each of the functions $P^{(k)}(2z_i; \tau)$, $P^{(k)}(2z_i; 2\tau)$, $P_{\text{even}}^{(k)}(z_i; \tau)$, $P_{\text{odd}}^{(k)}(z_i; \tau)$, $P^{(k)}(z_i - z_j; \tau)$ and $P^{(k)}(z_i + z_j; \tau)$ belong to $\mathcal{Q}_{n,0}^{(k+2)}$.

The functions $Z(z_i - a; \tau)$ for $a \in T$ belong to $\mathcal{Q}_{n,e_i}^{(1)}$, where $e_i = (0, \dots, 0, 1, 0, \dots, 0)$, and the functions $Z(z_i - z_j; \tau)$ and $Z(z_i + z_j; \tau)$ belong to $\mathcal{Q}_{n,e_i + e_j}^{(1)}$.

Proof Since P is an elliptic meromorphic function with poles at $\mathbb{Z} + \tau\mathbb{Z}$, and quasimodular in the sense of (iii) of weight 2 for $\text{SL}(2, \mathbb{Z})$, or, more precisely,

$$(c\tau + d)^{-2} P\left(\frac{z}{c\tau + d}; \frac{a\tau + b}{c\tau + d}\right) = P(z; \tau) + \frac{c}{c\tau + d},$$

we deduce easily the result for the functions derived from P . In fact, the functions $P^{(k)}(z_i - \frac{1}{2}; \tau)$ (hence also $P^{(k)}(2z_i; 2\tau)$, $P_{\text{even}}^{(k)}(z_i; \tau)$ and $P_{\text{odd}}^{(k)}(z_i; \tau)$) are quasimodular for the bigger group $\Gamma_0(2)$. Moreover, the functions $P^{(k)}(2z_i; \tau)$ are quasimodular for the full group $SL(2, \mathbb{Z})$. We proceed similarly for the functions derived from Z , which is quasielliptic of order 1, quasimodular of weight one and depth one with $Z_1(z; \tau) = z$. □

Proposition 5.4 *The direct sum*

$$\mathcal{Q}_n = \bigoplus_{k \geq 0} \mathcal{Q}_n^{(k)}, \quad \text{where } \mathcal{Q}_n^{(k)} = \bigoplus_{e \geq 0} \mathcal{Q}_{n,e}^{(k)}$$

is a graded ring. The derivatives $\partial/\partial z_i$ map $\mathcal{Q}_n^{(k)}$ to $\mathcal{Q}_n^{(k+1)}$ for all $i = 1, \dots, n$ and the derivative $D_q = q \frac{\partial}{\partial q}$ maps $\mathcal{Q}_n^{(k)}$ to $\mathcal{Q}_n^{(k+2)}$.

For all $i, j \in \{1, \dots, n\}$ with $i \neq j$, the functions

$$(39) \quad L(z_i; \tau) = -\frac{1}{2} Z^2(z_i; \tau) + \frac{1}{2} P(z_i; \tau) - G_2(\tau) + \frac{1}{12}$$

as well as $L(2z_i; \tau)$, $L(2z_i; 2\tau)$, $L(z_i - z_j; \tau)$ and $L(z_i + z_j; \tau)$ belong to $\mathcal{Q}_n^{(0)} \oplus \mathcal{Q}_n^{(2)}$.

Proof The proof is similar to the $SL(2, \mathbb{Z})$ case (cf [17, Proposition 5.6]). □

From now on we omit the variable τ in the notation, if not necessary.

Proposition 5.5 *The vector space \mathcal{Q}_n is (additively) generated as a \mathcal{Q}_{n-1} -module by the functions $Z^e(z_n - a)$ and $Z^e(z_n - a)P^{(m)}(z_n - a)$ for $a \in T$ together with $Z^e(z_n + z_j)$ and $Z^e(z_n + z_j)P^{(m)}(z_n + z_j)$, $Z^e(z_n - z_j)$, $Z^e(z_n - z_j)P^{(m)}(z_n - z_j)$ for $j = 1, \dots, n - 1$ and for all $e \geq 0$ and $m \geq 0$.*

More precisely, if $f \in \mathcal{Q}_n^{(k)}$ then we can write

$$\begin{aligned} f(z_1, \dots, z_n) = & \sum_{a \in T} \left(\sum_{e,m} A_{e,m,j} Z^e(z_n - a) P^{(m)}(z_n - a) + \sum_e B_{a,e} Z^e(z_n - a) \right) \\ & + \sum_{e,m,i} C_{e,m,j} Z^e(z_n + z_i) P^{(m)}(z_n + z_i) + \sum_{e,i} D_{e,i} Z^e(z_n + z_i) \\ & + \sum_{e,m,i} E_{e,m,i} Z^e(z_n - z_i) P^{(m)}(z_n - z_i) + \sum_{e,i} F_{e,i} Z^e(z_n - z_i) + G \end{aligned}$$

with $A_{a,e,m}, C_{e,m,i}, E_{e,m,i} \in \mathcal{Q}_{n-1}^{(k-e-m+2)}$, $B_{a,e}, D_{e,i}, F_{e,i} \in \mathcal{Q}_{n-1}^{(k-e)}$ and $G \in \mathcal{Q}_{n-1}^{(k)}$.

Proof For every n we argue inductively on the order $e = \min_{j \geq 0} \{ \Delta_n^j(f) \text{ elliptic} \}$ of quasiellipticity with respect to the last variable. Suppose, without loss of generality, that $f \in \mathcal{Q}_n^{(k)}$ is homogeneous of weight k .

We first treat the case $e = 0$. We show that we can write

$$f = \sum_{a \in T} \left(\sum A_{m,a} P_a^{(m)} + B_a Z_a \right) + \sum C_{m,i} P_{i,+}^{(m)} + D_i Z_{i,+} + \sum E_{m,i} P_{i,-}^{(m)} + F_{i,j} Z_{i,j} + G$$

with $A_{m,a}, C_{m,i}, E_{m,i} \in \mathcal{Q}_{n-1}^{(k-m-2)}$, $B_a, D_i, F_{i,j} \in \mathcal{Q}_{n-1}^{(k-1)}$ and $G \in \mathcal{Q}_{n-1}^{(k)}$, where $P_{j,a}^{(m)} = P^{(m)}(z_n + a)$ for $a \in T$, $P_{i,-}^{(m)} = P^{(m)}(z_n - z_i)$ and $P_{i,+}^{(m)} = P^{(m)}(z_n + z_i)$, for all $m \geq 0$ and all $i = 1, \dots, n-1$, $Z_a = Z(z_n + a) - Z(z_n + z_{n-1}) + Z(z_{n-1})$ for $a \in T$, $Z_{i,+} = Z(z_n + z_i) - Z(z_n - z_i) - 2Z(z_i)$ for all $i = 1, \dots, n-1$ and $Z_{i,j} = Z(z_n - z_i) - Z(z_n - z_j) + Z(z_i - z_j)$ for all $1 \leq i < j \leq n-1$. By Proposition 5.3, these functions are clearly in $\mathcal{Q}_{n,0}$.

We proceed by induction on the pole orders, first along the divisors $z_n - a$, then along the divisors $z_n + z_i$, then along the divisors $z_n - z_i$. The rest of the proof is then totally similar to [17, Proposition 5.4]. The residue theorem ensures that we can eliminate the last poles with the functions $Z_{i,j}$ to end the procedure.

For the case $e > 0$, the proof is a straightforward adaptation of the proof of [17, Proposition 5.7]. □

Using this additive basis we can now prove the main result.

Theorem 5.6 *For any permutation π the constant term with respect to the domain Ω_π of a function in $\mathcal{Q}_n^{(k)}$ is a quasimodular form for $\Gamma(2)$ of mixed weight $\leq k$.*

Proof Again, the proof is exactly the same as in [17, Theorem 5.8]: we proceed by induction on the variables and show that the z_n -integrals of the additive generators are quasimodular forms of bounded mixed weight. For the generators involving products of Z and derivatives of P , we integrate by parts to decrease the order of quasiellipticity. The problem is then reduced to the computation of $[\zeta^0]Z^e(z - a)$. The last step of the proof is isolated in the statement of the next proposition. □

Proposition 5.7 *The constant coefficient $[\zeta^0]Z^e(z - a)$ for $a \in T$ is a quasimodular form for $SL(2, \mathbb{Z})$ of mixed weight less than or equal to e .*

Proof Since $Z(z)$ is 1-periodic, we clearly have $[\zeta^0]Z^e(z - \frac{1}{2}) = [\zeta^0]Z^e(z)$ and these coefficients are quasimodular forms for $SL(2, \mathbb{Z})$ of mixed weight less than or




Figure 4: Integration paths to evaluate $[\zeta^0]Z^e$.

equal to e , by [17, Proposition 5.9]. Similarly, $[\zeta^0]Z^e(z - \frac{1}{2}\tau - \frac{1}{2}) = [\zeta^0]Z^e(z - \frac{1}{2}\tau)$, so we just have to compute $[\zeta^0]Z^e(z - \frac{1}{2}\tau)$.

Using again the 1-periodicity of Z , we obtain, for all ℓ ,

$$(40) \quad [\zeta^0]Z^\ell(z - \frac{1}{2}\tau) = \int_{\gamma_1} Z^\ell(z - \frac{1}{2}\tau) dz = \int_{\gamma_2} Z^\ell(z) dz = \int_{\gamma_3} Z^\ell(-z) dz = [\zeta^0]Z^\ell(-z),$$

where the integration paths γ_1, γ_2 and γ_3 are as described in Figure 4. Since Z is odd, the constant coefficients $[\zeta^0]Z^\ell(z - \frac{1}{2}\tau)$ are then given by $(-1)^\ell [\zeta^0]Z^\ell$ (see [17] for some explicit values). □

Note that the proof of Theorem 5.6 provides an effective algorithm to compute constant coefficients of quasielliptic functions. Applications of this algorithm and explicit computations of quasimodular forms are detailed in Section 9.

6 Quasimodularity of graph sums

The goal of this section is to show the quasimodularity of graph sums of the form (41) below. The motivation for considering these sums will become apparent in comparison with the quasipolynomiality theorem in Section 7. We encourage the reader to look at Section 9 simultaneously with this one, we hope that all the notation will become transparent on the example that we treat in detail.

We will show quasimodularity for graphs that arise as global graphs as in Section 2.3 with the following extra decoration.

A *global graph with distinguished edges* is a graph with vertices with the labels $1, \dots, n$ and possibly a special vertex 0 and a subset $E^+(\Gamma)$ of $E(\Gamma)$ of distinguished edges such that no extremity of an edge in $E^+(\Gamma)$ is the vertex 0 . We let $V^*(\Gamma) = V(\Gamma) \setminus \{0\}$; we let $E^0(\Gamma)$ be the edges adjacent to 0 . Finally we define $E^*(\Gamma) = E(\Gamma) \setminus E^0(\Gamma)$ and $E^-(\Gamma) = E^*(\Gamma) \setminus E^+(\Gamma)$. An *admissible orientation* G of (Γ, E^+) is an orientation of the half-edges of Γ such that

- all half-edges adjacent to the vertex 0 are outgoing, and
- the orientations of the two half-edges are consistent on marked edges, and inconsistent on the other edges that are not adjacent to v_0 .

We write $G \in (\Gamma, E^+)$ for the specification of an admissible orientation. Obviously every admissible orientation G of Γ (in the sense of Section 4) is admissible for (Γ, E^+) for a uniquely determined subset $E^+ \subset E(\Gamma)$. We define the *set of parity conditions* to be $\text{PC}(\Gamma) = \{0, 1\}^{E^0(\Gamma)}$. It specifies a congruence class mod 2 for the width of each edge adjacent to the vertex 0 .

We consider here for fixed $\mathbf{m} = (m_1, \dots, m_{E(\Gamma)})$ and fixed $E^+(\Gamma)$ the graph sums

$$(41) \quad S(\Gamma, E^+, \mathbf{m}, \text{par}) = \sum_{G \in (\Gamma, E^+)} S(G, \mathbf{m}, \text{par})$$

over all admissible orientations G of the half-edges of Γ , where, for $\text{par} \in \text{PC}(\Gamma)$,

$$(42) \quad S(G, \mathbf{m}, \text{par}) = \sum_{\substack{h \in \tilde{\mathbb{N}}^{E(G)} \\ \mathbf{w}_* \in \mathbb{N}_{>0}^{E(G)^*}}} \sum_{\substack{\mathbf{w}_0 \in \mathbb{N}_{>0}^{E^0(\Gamma)} \\ \mathbf{w}_0 \cong \text{par} \pmod{2}}} \prod_{i \in E(G)} w_i^{m_i+1} q^{h_i w_i} \prod_{v \in V(G)^*} \delta(v).$$

Here $\tilde{\mathbb{N}}^{E(G)}$ is the height space introduced in (8) and $\delta(v)$ is as in (31). The goal of this section is to show the quasimodularity of these graph sums.

Theorem 6.1 *For a fixed tuple of nonnegative even integers $\mathbf{m} = (m_1, \dots, m_{|E(\Gamma)|})$ and for any $\text{par} \in \text{PC}(\Gamma)$, the graph sums $S(\Gamma, E^+, \mathbf{m}, \text{par})$ are quasimodular forms for the group $\Gamma_0(2)$ of mixed weight at most $k(\mathbf{m}) := \sum_i (m_i + 2)$.*

The following is a first simplification step for the computation. Recall the definition of the offsets a_e from Figure 3.

Lemma 6.2 *Replacing the height space $\tilde{\mathbb{N}}^{E(G)}$ by*

$$(43) \quad \hat{\mathbb{N}}^{E(G)} = \{(h'_e)_{e \in E(\Gamma)} : h'_e \in \mathbb{N}_{>0} \text{ if } e \in E_\ell^0(G) \cup E_\ell^+(G), h'_e \in \mathbb{N}_{\geq a_e} \text{ otherwise}\}$$

does not change the total sum $S(G, \mathbf{m}, \text{par})$.

Proof We apply the linear change of variables $h'_e = h_e - \Delta(e)$ with $\Delta(e)$ as in the line below (8). This maps $\tilde{\mathcal{N}}^{E(G)}$ bijectively onto $\hat{\mathcal{N}}^{E(G)}$. For notational convenience we set $\Delta(e) = 0$ for the remaining edges. Each summand of (42) for fixed $(w_1, \dots, w_{|E(G)|})$ is multiplied under the variable change by

$$q^{\sum_e \Delta(e)w_e} \prod_{v \in V(G)^*} \delta(v) = q^{\sum_v \varepsilon_v (\sum_{i \in e^-(v)} w_i - \sum_{i \in e^+(v)} w_i)} \prod_{v \in V(G)^*} \delta(v) = 1.$$

This implies the claim. □

6.1 The reduced graph

We will simplify the graph sums by isolating the contribution from the loop edges $E_\ell^0(G) \cup E_\ell^+(G)$. The *reduced graph* $\bar{\Gamma}$ is obtained from Γ by deleting those loops, ie the loops adjacent to the vertex v_0 and the loops among the distinguished edges.

Lemma 6.3 *The graph sums $S(\Gamma, \mathbf{m}, \text{par})$ factor as*

$$S(\Gamma, E^+, \mathbf{m}, \text{par}) = S_{\text{loops}}(\Gamma, \mathbf{m}, \text{par})S(\bar{\Gamma}, E^+, \mathbf{m}, \text{par}),$$

where

$$S_{\text{loops}}(\Gamma, \mathbf{m}, \text{par}) = \sum_{h \in \mathbb{N}_{>0}^{E_\ell^0(\Gamma) \cup E_\ell^+(\Gamma)}} \sum_{\mathbf{w} \in \mathbb{Z}_+^{E_\ell^+(\Gamma)}} \sum_{\substack{\mathbf{w}_0 \in \mathbb{N}_{>0}^{E_\ell^0(\Gamma)} \\ \mathbf{w}_0 \cong \text{par} \pmod{2}}} \prod_{i \in E_\ell^0(\Gamma) \cup E_\ell^+(\Gamma)} w_i^{m_i+1} q^{h_i w_i}.$$

Proof The length constraints $\delta(v)$ for $v \in V(G)^*$ are unchanged under removing a loop edge $e \in E^+(G)$ that contributes equally to both incoming and outgoing weight. The length parameters at the vertex 0 is always unconstrained. This proves the factorization we claim. □

Lemma 6.4 *If \mathbf{m} is even, $S_{\text{loops}}(\Gamma, \mathbf{m}, \text{par})$ is a quasimodular form for $\Gamma_0(2)$ of mixed weight $k(\mathbf{m})$.*

Proof The graph sum $S_{\text{loops}}(\Gamma, \mathbf{m}, \text{par})$ is a product of

$$S_m = \sum_{w,h=1}^\infty w^{m+1} q^{2wh} = G_{m+2}(q) - G_{m+2}(0),$$

$$S_{m,\text{even}} = \sum_{w,h=1}^\infty (2w)^{m+1} q^{(2w)h} = 2^{m+1}(G_{m+2}(q^2) - G_{m+2}(0)),$$

$$S_{m,\text{odd}} = \sum_{w,h=1}^{\infty} (2w-1)^{m+1} q^{(2w-1)h} = S_m - S_{m,\text{even}}.$$

All the right-hand sides are quasimodular forms for $\Gamma_0(2)$ by (37). □

6.2 Contour integrals

We now write the sum of the reduced graph as contour integral of suitable derivatives of the following variants of the propagator. Let $P_{\text{even}}(z; \tau) = 2P(2z; 2\tau)$ and $P_{\text{odd}}(z; \tau) = P(z; \tau) - P_{\text{even}}(z; \tau)$. We also use P_i for $i \in \mathbb{Z}/2$ to refer to these two functions. For a reduced graph $\bar{\Gamma}$, for \mathbf{m} even, and for a given parity condition $\text{par} \in \text{PC}(\Gamma)$, define

$$(44) \quad P_{\bar{\Gamma}, E^+, \mathbf{m}, \text{par}}(\mathbf{z}) = \prod_{i \in E^0(\bar{\Gamma})} P_{\text{par}_i}^{(m_i)}(z_{v_+(i)}) \cdot \prod_{i \in E^+(\bar{\Gamma})} P^{(m_i)}(z_{v_-(i)} - z_{v_+(i)}) \cdot \prod_{i \in E^-(\bar{\Gamma})} P^{(m_i)}(z_{v_-(i)} + z_{v_+(i)}),$$

where $v_+(i)$ and $v_-(i)$ are the two ends of the edge i , with v_- being the one of lower index (so for $i \in E^0(\bar{\Gamma})$ necessarily $v_-(i) = 0$). Note that since the graph is reduced, $v_+(i)$ and $v_-(i)$ can be the same, but only if $i \in E^-(\bar{\Gamma})$. Note that the variable z_0 does not appear in the expression $P_{\bar{\Gamma}, \mathbf{m}, \text{par}}(\mathbf{z})$ at all.

Proposition 6.5 *For a tuple of nonnegative even integers \mathbf{m}_1 and a parity condition par , we can express the graph sum as*

$$(45) \quad S(\bar{\Gamma}, E^+, \mathbf{m}, \text{par}) = [\zeta_n^0, \dots, \zeta_1^0] P_{\bar{\Gamma}, E^+, \mathbf{m}, \text{par}}(\mathbf{z}; \tau),$$

where the coefficient extraction is for the expansion on the domain $|q^{1/2}| < |\zeta_i| < |\zeta_{i+1}| < 1$ for all i .

Proof The proof is analogous to the proof of [17, Proposition 6.7] and we suggest to read the two proofs in parallel since we will not reproduce the bulky main formulas. We rather indicate the main changes. First note that in the domain specified above the inequalities

$$|q| < |\zeta_i \zeta_j| < 1 \quad \text{for all } i, j, \quad |q| < |q^{1/2}| < \left| \frac{\zeta_i}{\zeta_j} \right| < 1 \quad \text{for all } j > i$$

hold, and hence the Fourier expansions

$$\begin{aligned}
 P_{\text{even}}^{(m)}(z; \tau) &= \sum_{w \geq 1, \text{even}} w^{m+1} \left(\zeta^w \sum_{h \geq 0} q^{wh} + \zeta^{-w} \sum_{h \geq 1} q^{wh} \right), \\
 P_{\text{odd}}^{(m)}(z; \tau) &= \sum_{w \geq 1, \text{odd}} w^{m+1} \left(\zeta^w \sum_{h \geq 0} q^{wh} + \zeta^{-w} \sum_{h \geq 1} q^{wh} \right), \\
 (46) \quad P^{(m)}(z_i + z_j; \tau) &= \sum_{w \geq 1} w^{m+1} \left((\zeta_i \zeta_j)^w \sum_{h \geq 0} q^{wh} + (\zeta_i \zeta_j)^{-w} \sum_{h \geq 1} q^{wh} \right) \\
 &\hspace{25em} \text{for all } i, j, \\
 P^{(m)}(z_i - z_j; \tau) &= \sum_{w \geq 1} w^{m+1} \left(\left(\frac{\zeta_i}{\zeta_j} \right)^w \sum_{h \geq 0} q^{wh} + \left(\frac{\zeta_i}{\zeta_j} \right)^{-w} \sum_{h \geq 1} q^{wh} \right) \\
 &\hspace{25em} \text{for all } j > i
 \end{aligned}$$

are valid. For the proof we first consider the factors in (44) that involve the edges E_1 adjacent to the vertex 1, ie those involving the variable z_1 . The propagators P , P_{odd} or P_{even} in (44) are chosen so that the parity conditions for w_e specified in (42) hold. Each of the propagators in (46) has (for fixed (w, h)) two summands, which we consider as incoming (ζ -exponent $+w$) or outgoing (ζ -exponent $-w$). Consequently, expanding the product of propagators involving the edges in E_1 is a sum over all partitions $E_1 = J_1 \cup K_1$ of the incoming and outgoing terms. The integration with respect to z_1 forces that all contributions vanish except for those where the incoming w_e are equal to the outgoing w_e . This ensures the appearance of the factor $\delta(v_1)$ in (42). The proof proceeds by similarly considering the vertex 2 and expanding the propagator factors that involve the edges E_2 adjacent to this vertex but not already in E_1 , which produces a sum over all partitions $E_2 = J_2 \cup K_2$ according to whether the incoming or outgoing summand of the propagator has been taken.

The main difference to the abelian case is the consequence of orienting the half-edges. Suppose that e joins v_1 to v_2 . If $e \in E^+(\Gamma)$ then in all admissible orientations e is incoming at v_1 and outgoing at v_2 , or vice versa. If $e \in J_1$ is incoming at v_1 , we have to make sure that the propagator terms have ζ_2^{-w} , ie we have to use $P^{(m)}(z_1 - z_2)$. On the other hand, if $e \in E^-(\Gamma)$, then in all admissible orientations e is incoming or outgoing simultaneously at v_1 and v_2 . That is, ζ_1 and ζ_2 have to appear with the same w -exponent, whence the use of $P^{(m)}(z_1 + z_2)$. The reader can check that this orientation convention is also consistent for the special vertex 0 and that the range of the sums $h \geq 0$ versus $h \geq 1$ in (44) is consistent with the conditions of the height space that appear in the h -summation in (42). □

Proof of Theorem 6.1 This is now a direct consequence of Lemma 6.4 for the loop contribution, of Proposition 6.5 for the reduced graph and of Theorem 5.6 for quasimodularity (for $\Gamma(2)$) of contour integrals, combined with Lemma 5.1 to get quasimodularity for the bigger group $\Gamma_0(2)$. \square

7 Quasipolynomiality of 2-orbifold double Hurwitz numbers

The main result of this section is the quasipolynomiality of the simple Hurwitz numbers with 2-stabilization $A'_2(\mathbf{w}, F)$ in the case that F is a product of \bar{p}_k . The meaning of quasipolynomiality is that the restriction to a congruence class mod 2 in each variable is a polynomial. The crucial statement for the quasimodularity is that these polynomials are global, ie not piecewise polynomials depending on a chamber decomposition of the domain of \mathbf{w} . As a first application we combine this with the correspondence and quasimodularity theorems of the previous section to give in Corollary 7.4 another proof the Eskin–Okounkov theorem on the quasimodularity of the number of pillowcase covers.

7.1 The one-sided pillowcase operator

Our goal here is to write $A'_2(\mathbf{w}, F)$ in terms of vertex operators. For this purpose we define the *one-sided pillowcase operator*

$$\Gamma_{\sqrt{w}} = \exp\left(\sum_{i>0} \frac{\alpha_{-i}^2}{2i}\right).$$

Proposition 7.1 *The simple Hurwitz numbers with 2-stabilization can be expressed using the one-sided pillowcase operator as*

$$(47) \quad A_2(\mathbf{w}, F) = \frac{1}{\prod w_i} \left\langle 0 \left| \prod_{i=1}^{\ell(\mathbf{w})} \alpha_{w_i} \mathcal{F} \Gamma_{\sqrt{w}} \right| 0 \right\rangle,$$

where $\mathcal{F}v_\lambda = F(\lambda)v_\lambda$.

Proof We first observe that

$$(48) \quad \langle \Gamma_{\sqrt{w}} v_\emptyset, v_\lambda \rangle = \sum_{\nu} \chi^\lambda(2\nu) \prod_{i \geq 1} \frac{(1/2i)^{r_i(\nu)}}{r_i(\nu)!},$$

where 2ν is the partition obtained by repeating twice each row of the Young diagram of ν , ie if $\nu = 1^{r_1(\nu)} 2^{r_2(\nu)} \dots$ is written in terms of the multiplicities of the parts then

$2\nu = 1^{2r_1(\nu)}2^{2r_2(\nu)} \dots$. This observation follows from developing the exponential in $\Gamma_{\sqrt{w}}$ and the Murnaghan–Nakayama rule. It thus remains to show that

$$\sum_{\nu} \chi^{\lambda}(2\nu) \prod_{i=1}^{\nu_1} \frac{(1/2i)^{r_i(\nu)}}{r_i(\nu)!} = f_{2,2,\dots,2}(\lambda)^2 \frac{\dim \lambda}{|\lambda|!} = \sqrt{w(\lambda)}$$

or equivalently that

$$(49) \quad f_{2,\dots,2}(\lambda)^2 = \sum_{\nu} f_{2\nu}(\lambda) \cdot \prod_{i \geq 1} (i^{r_i} (2r_i - 1)!).$$

Let C_2 be the conjugacy class corresponding to the partition $(2, \dots, 2)$ and $C(2\nu)$ the conjugacy class of 2ν . Define

$$n_{C_2^2}^C = \#\{(a, b) \in C_2^2 \mid ab \in C\}$$

and let $e_C = \sum_{g \in C} [g]$ be central elements the group ring $\mathbb{Z}[G]$. Then $e_{C_2}e_{C_2} = \sum_C n_{C_2^2}^C e_C$. Note that e_C acts on any irreducible representation λ as multiplication by the scalar $f_C(\lambda)$. Consequently, the claim (49) is equivalent to

$$n_{C_2^2}^C = \begin{cases} \prod_i (i^{r_i} (2r_i - 1)!) & \text{if } C = C(2\nu), \\ 0 & \text{otherwise.} \end{cases}$$

To prove this, note first that the product $\tau = \rho\sigma$ of two permutations ρ and σ in C_2 is in $C(2\nu)$. This follows by recursively proving that $\sigma\tau^{-n}(P) = \tau^n(\sigma(P))$, i.e. the cycles starting at P and $\sigma(P)$ have the same length. To justify the combinatorial factor, assume first that all $r_i = 1$. In order to specify the factorization it is necessary and sufficient to specify for each i and for some point in an i -cycle its σ -image in the other i -cycle. The rest of the factorization is determined by the requirement of profile $(2, \dots, 2)$ and this initial choice. In the general case $r_i > 1$ we moreover have to match the $2r_i$ cycles of length i in pairs (which corresponds to the factor $(2r_i - 1)!$) and make a choice as above for each pair.

The proposition follows from (48) using again the Murnaghan–Nakayama rule. □

We next collect some rules to evaluate the right-hand side of (47). The standard commutation law between creation operators is

$$(50) \quad \alpha_n^a \alpha_{-n}^b = \sum_{k=0}^{\min(a,b)} \binom{a}{k} \frac{b!}{(b-k)!} n^k \alpha_{-n}^{b-k} \alpha_n^{a-k}.$$

Together with $[\alpha_m, \alpha_n] = 0$ for $m \neq -n$, this implies that we can deal with the $\alpha_{\pm n}$ for all n separately, ie

$$\left\langle \prod_n f_n(\alpha_n) g_n(\alpha_{-n}) v_\emptyset, v_\emptyset \right\rangle = \prod_n \langle f_n(\alpha_n) g_n(\alpha_{-n}) v_\emptyset, v_\emptyset \rangle,$$

where $f_n(\alpha_n) = \sum c_{j,n} \alpha_n^j$ and $g_n(\alpha_{-n}) = \sum c'_{j,n} \alpha_{-n}^j$. From the commutation relation

$$e^{c_n \alpha_{-n}} e^{d_n \alpha_n} = e^{-n c_n d_n} e^{d_n \alpha_n} e^{c_n \alpha_{-n}},$$

which follows directly from (50), we deduce successively the following relations involving the factors of $\Gamma_{\sqrt{w}}$. First

$$P := \prod_{i=1}^s (e^{c_{i,n} \alpha_{-n}} e^{d_{i,n} \alpha_n}) = e^{-n A_n} e^{D_n \alpha_n} e^{C_n \alpha_{-n}},$$

where $C_n = \sum_{i=1}^s c_{i,n}$, $D_n = \sum_{i=1}^s d_{i,n}$ and $A_n = \sum_{j=1}^s d_{j,n} \sum_{i=1}^j c_{i,n}$. Next, from

$$\begin{aligned} \langle 0 | e^{c_n \alpha_{-n}} e^{d_n \alpha_n} e^{\alpha_{-n}^2/2n} | 0 \rangle &= e^{n d_n^2/2}, \\ \langle 0 | \alpha_n e^{c_n \alpha_{-n}} e^{d_n \alpha_n} e^{\alpha_{-n}^2/2n} | 0 \rangle &= n(d_n + c_n) e^{n d_n^2/2}, \end{aligned}$$

we obtain

$$(51) \quad \begin{aligned} \langle 0 | P \cdot e^{\alpha_{-n}^2/2n} | 0 \rangle &= e^{n \tilde{A}_n} e^{n D_n^2/2}, \\ \langle 0 | \alpha_n \cdot P \cdot e^{\alpha_{-n}^2/2n} | 0 \rangle &= n(D_n + C_n) e^{n \tilde{A}_n} e^{n D_n^2/2}, \end{aligned}$$

where $\tilde{A}_n = D_n C_n - A_n = \sum_{j=2}^s d_{j,n} \sum_{i=1}^{j-1} c_{i,n}$, and, most generally,

$$(52) \quad \langle 0 | \alpha_n^\ell \cdot P \cdot e^{\alpha_{-n}^2/2n} | 0 \rangle = \sum_{i=0}^{\ell/2} \binom{\ell}{2i} \frac{(2i)!}{2^i (i)!} n^{\ell-i} (C_n + D_n)^{\ell-2i} e^{n \tilde{A}_n} e^{n D_n^2/2}.$$

7.2 Polynomiality

We can now evaluate the vertex operator expressions and prove the main result of this section.

Theorem 7.2 *The simple Hurwitz number with 2-stabilization $A'_2(\mathbf{w}, F)$ without unramified components is a quasipolynomial if F is a product of \bar{p}_k , ie for each coset $\mathbf{m} = (m_1, \dots, m_t) \in \{0, 1\}^n$ with $\sum m_i$ even there exists a polynomial $R_{F,\mathbf{m}} \in \mathbb{Q}[w_1, \dots, w_t]$ of degree $\text{wt}(F) - t$ such that*

$$A'_2(\mathbf{w}, F) = R_{F,\mathbf{m}}(\mathbf{w}) \quad \text{for all } \mathbf{w} \in 2\mathbb{N}^t + \mathbf{m}.$$

The proof relies on matching piecewise polynomial on sectors like $w_1 > w_2$ to form a global polynomial. The parity constraints to match the piecewise polynomials do not work out for elements in Λ^* , not even for $F = p_\ell$, if there is more than one boundary variable w_i . In fact, for any $w_1, w_2 \in \mathbb{N}$ with $w_1 + w_2$ even,

$$\frac{1}{5}A'_2((w_1, w_2), p_5) = \frac{7}{8}u^3 + \frac{13}{8}uv^2 - u,$$

where $u = \min(w_1, w_2)$ and $v = \max(w_1, w_2)$ is only piecewise polynomial, while, eg on the coset $\mathbf{m} = (0, 0)$,

$$\frac{1}{4}A'_2((2w_1, 2w_2), \bar{p}_4) = 10(2w_1)^2 + 10(2w_2)^2 - 3$$

is globally a polynomial.

The basic source of polynomiality is the following lemma, relevant for the case of $F = \bar{p}_k$:

Lemma 7.3 *For each $k \geq 1$ there is a polynomial Q_k of degree k such that, for $n \in \mathbb{N}$,*

$$Q_k(n) = [y^{-2n}][z^k]D, \quad \text{where } D(y, z) = \frac{1 + y^{-2}e^{-z}}{\sqrt{(1 - y^{-2})(1 - y^{-2}e^{-2z})}}.$$

Moreover, Q_k is even for k even and Q_k is odd for k odd, and $Q_k(0) = 0$ in both cases.

Proof We abbreviate $\partial_z = \partial/\partial z$. Since $[z^k] = k! \partial_z^k f|_{z=0}$, it suffices to write $\partial_z^k D(y, z)|_{z=0} = R_k(y^{-2})/(1 - y^{-2})^{k+1}$ for some polynomial $R_k(y^{-2})$ of degree $\leq k$ without constant coefficient. The relation $D(1/y, -z) = D(y, z)$ implies that R_k is palindromic, ie in the span of $y^{-s} + y^{-(2k-s)}$ for $s = 2, 4, \dots, 2k - 2$. Since by the binomial theorem

$$[y^{-2n}] \frac{1}{(1 - y^{-2})^{k+1}} = \frac{1}{k!} (n + k - 1) \cdots (n + 1)n$$

agrees with a polynomial of degree k for integers $n \geq 1 - k$, we obtain the polynomiality claim. The parity claim follows from R_k being palindromic. □

Proof of Theorem 7.2 We may shift \bar{p}_k by the regularization constant γ_k in order to use (20) and (21) and assume that $F = \prod_{j=1}^s (\bar{p}_{k_j} - \gamma_{k_j})$ for some k_j , not necessarily distinct. Proposition 7.1 now translates into

$$(53) \quad A_2(\mathbf{w}, F) = [y_1^0 \cdots y_s^0][z_1^{k_1} \cdots z_s^{k_s}] \prod_{n \geq 1} \frac{1}{n^{r_n(\mathbf{w})}} \langle 0 | \alpha_n^{r_n(\mathbf{w})} \Psi_F e^{\alpha_{-n}^2/2n} | 0 \rangle,$$

where $r_n(\mathbf{w})$ is the multiplicity of n in \mathbf{w} and where

$$\Psi_F = \prod_{j=1}^s \frac{1}{e^{z_j/2} + e^{-z_j/2}} \exp\left(\frac{y_j^n((-e^{z_j})^n - 1)}{n} \alpha_{-n}\right) \exp\left(\frac{y_j^{-n}(1 - (-e^{-z_j})^n)}{n} \alpha_n\right).$$

By (51), the first factor common to the evaluation of the brackets (53) for all n is $e^{nD_n^2/2}$, and it results in a product of

$$D_{[j]} := \exp\left(\sum_{n>0} \frac{y_j^{-2n}(1 - (-e^{-z_j})^n)^2}{2n}\right) = \frac{1 + y_j^{-2}e^{-z_j}}{\sqrt{(1 - y_j^{-2})(1 - y_j^{-2}e^{-2z_j})}}$$

and

$$D_{[ij]} := \frac{(1 + y_i y_j e^{-z_i})(1 + y_i y_j e^{-z_j})}{(1 - y_i y_j)(1 - y_i y_j e^{-z_i - z_j})}.$$

The second common factor $e^{n\tilde{A}_n}$ for all n results in a factor of

$$\tilde{A} := \prod_{j=2}^s \prod_{i=1}^{j-1} \frac{(1 + (y_j/y_i)e^{-z_i})(1 + (y_j/y_i)e^{z_j})}{(1 - y_j/y_i)(1 - (y_j/y_i)e^{z_j - z_i})}.$$

In order to built an arbitrary covering with boundary lengths \mathbf{w} from a covering without unramified components, we have to choose for each length n of the boundary components among the $\ell_n = r_n(\mathbf{w})$ an even number $2i$ of boundary components that are glued together in pairs to form cylinders, and the number of such gluings is $(2i)!/2^i (i)!$, the number of fixed-point-free involutions. That is, the combinatorial factor $\binom{\ell_n}{2i} (2i)!/2^i (i)!$ in front of the summand in (52) counts precisely these possibilities. Consequently, this formula implies that

$$(54) \quad A'_2(\mathbf{w}, F) = [y_1^0 \cdots y_s^0][z_1^{k_1} \cdots z_s^{k_s}] \left(\frac{\tilde{A} \prod_{i<j} D_{[ij]} \prod_j D_{[j]}}{\prod_j (e^{z_j/2} + e^{-z_j/2})} \prod_{n:r_n(\mathbf{w}) \geq 1} K_n^{r_n(\mathbf{w})} \right),$$

where $K_n = C_n + D_n$ as in the vertex operator manipulations above, ie

$$K_n = \sum_{j=1}^s \frac{y_j^n((-e^{z_j})^n - 1)}{n} + \frac{y_j^{-n}(1 - (-e^{-z_j})^n)}{n}.$$

The claim follows if we can show two statements: first that the expression (54) is piecewise polynomial and second that this expression with each K_n replaced by $\tilde{K}_n = nK_n$ is globally a polynomial. The factor $\prod_j (e^{z_j/2} + e^{-z_j/2})$ results just in a shift of z_j -degrees and will be ignored in the sequel. Note that we can write the last factor in (54) equivalently as $\prod_{n:r_n(\mathbf{w}) \geq 1} K_n^{r_n(\mathbf{w})} = \prod_{i=1}^t K_{w_i}$.

We start with the case $s = 1$, illustrating the main idea. Let t be the number of n with $r_n(\mathbf{w}) \geq 1$; say these are n_1, \dots, n_t . First, we want to show that

$$\mathbf{n} = (n_1, \dots, n_t) \mapsto [y_1^0] \left([z_1^{k-j}] \prod_{i=1}^t \tilde{K}_{n_i} \cdot [z_1^j] D_{[1]} \right)$$

is polynomial of degree k in the n_i for each $j \in [0, k]$ (and zero otherwise). To evaluate this, we can choose in each \tilde{K}_{n_i} -factor the $y_i^{n_i}$ -term or the $y_i^{-n_i}$ -term and then sum over the contributions of all choices. For each $\delta = (\delta_1, \dots, \delta_t) \in \{\pm 1\}^t$ we consider the linear form $f_\delta(n_1, \dots, n_t) = \sum_{i=1}^t \delta_i n_i$. We claim that already the sum of the contributions of f_δ and $f_{-\delta}$ is polynomial, ie that

$$\begin{aligned} \mathbf{n} \mapsto [z_1^{k-j}] \prod_{i=1}^t ((-1)^{m_i} e^{\delta_i n_i z_1} - 1) [y_1^{f_\delta(\mathbf{n})}] [z_1^j] D_{[1]} \\ + [z_1^{k-j}] \prod_{i=1}^t ((-1)^{m_i} e^{-\delta_i n_i z_1} - 1) [y_1^{f_{-\delta}(\mathbf{n})}] [z_1^j] D_{[1]} \end{aligned}$$

is the restriction of a polynomial to any collection of natural numbers $n_i \equiv m_i \pmod{2}$ is the fixed coset. If we write $f_\delta^+ = \max(0, f_\delta)$, the claim follows from the observation that $Q_j(\frac{1}{2} f_\delta^+(\mathbf{n})) + (-1)^j Q_j(\frac{1}{2} f_{-\delta}^+(\mathbf{n})) = Q_j(\frac{1}{2} f_\delta(\mathbf{n}))$ is globally a polynomial for \mathbf{n} in a fixed congruence class and for Q_j with the parity as in Lemma 7.3. The polynomiality for $s = 1$ and the \tilde{K}_n -version follow by summing up these expressions.

Second, we argue that $A_2'(\mathbf{w}, \bar{p}_k)$ without the additional factors n in \tilde{K}_n is a piecewise polynomial $k - t$, ie that the polynomial expression obtained previously using \tilde{K}_n is indeed divisible by n . The divisibility by n_i follows from adding the contribution of f_δ and $f_{\delta'}$, where δ' differs from δ precisely in the i^{th} digit, since $(n_i + n_j)^k + (n_i - n_j)^k$ is divisible by n_i independently of the parity of k .

For the general case $s \geq 1$ follows along the same lines. We first prove a generalization of Lemma 7.3, stating that for $\mathbf{k} = (k_1, \dots, k_s)$ there is a polynomial $Q_{\mathbf{k}}$ in s variables g_i of total degree $\sum k_i$ such that

$$Q_{\mathbf{k}}(g_1, \dots, g_s) = [y_1^{-2g_1} \dots y_s^{-2g_s}] [z_1^{k_1} \dots z_s^{k_s}] \left(\tilde{A} \prod_{i < j} D_{[ij]} \prod_j D_{[j]} \right)$$

if all $g_i \geq 0$. Moreover, this polynomial has the parity

$$(55) \quad Q_{\mathbf{k}}(g_1, \dots, -g_j, \dots, g_s) = (-1)^{k_j} Q_{\mathbf{k}}(g_1, \dots, -g_j, \dots, g_s).$$

To see this, we write

$$\begin{aligned} & \partial_{z_1}^{k_1} \cdots \partial_{z_s}^{k_s} \left(\tilde{A} \prod_{i < j} D_{[ij]} \prod_j D_{[j]} \right) \Big|_{z_1 = \cdots = z_s = 0} \\ &= \sum_{\ell, \mathbf{m}, \mathbf{n}} R_{\mathbf{k}, \ell, \mathbf{m}, \mathbf{n}}(y_1, \dots, y_s) \cdot \prod_i \frac{1}{(1 - y_i^2)^{\ell_i}} \cdot \prod_{i < j} \frac{1}{(1 - y_i y_j)^{m_{ij}}} \cdot \prod_{i < j} \frac{1}{(1 - y_j / y_i)^{n_{ij}}} \end{aligned}$$

for some polynomial $R_{\mathbf{k}, \ell, \mathbf{m}, \mathbf{n}}$ with the components of ℓ , \mathbf{m} and \mathbf{n} bounded in terms of \mathbf{k} . Using the binomial expansion of this expression we see that the coefficient $[y_1^{-2g_1} \cdots y_s^{-2g_s}]$ of this expression is a sum of polynomial expressions in nonnegative integers u_i , v_{ij} and w_{ij} over the bounded simplex defined by

$$u_i + \sum_j v_{ij} + \sum_{j < i} n_{ij} - \sum_{j > i} n_{ij} = g_i \quad \text{for } i = 1, \dots, s.$$

This sum is again a polynomial and this implies the polynomiality claim. Moreover, the argument $\tilde{A} \prod_{i < j} D_{[ij]} \prod_j D_{[j]}$ in the definition of $Q_{\mathbf{k}}$ is unchanged under the transformation $(y_i, z_i) \mapsto (1/y_i, -z_i)$ since each $D_{[i]}$ has this property and since this transformation swaps $D_{[ij]}$ with the (i, j) -factor of \tilde{A} . As in the case $s = 1$, this implies palindromic numerators and the parity statement (55).

We show similarly that for each tuple (j_1, \dots, j_s) separately the function

$$\mathbf{n} = (n_1, \dots, n_t) \mapsto [y_1^0] \left([z_1^{k_1 - j_1} \cdots z_s^{k_s - j_s}] \prod_{i=1}^t \tilde{K}_{n_i} \cdot [z_1^{j_1} \cdots z_s^{j_s}] \tilde{A} \prod_{i < j} D_{[ij]} \prod_j D_{[j]} \right)$$

is a polynomial of degree $\sum k_i$. Evaluation of $\prod_{i=1}^t \tilde{K}_{n_i}$ now leads to s linear forms $f_1(\mathbf{n}), \dots, f_s(\mathbf{n})$ with each n_i appearing in exactly one of the f_j , and with coefficient ± 1 . Given one such tuple, the contributions of $\pm f_1, \dots, \pm f_s$ add up to a global polynomial thanks to (55), as in the case $s = 1$. Finally, adding the contribution of f_1, \dots, f_s and the linear form with precisely the sign of n_i flipped gives the divisibility by n_i that was still left to prove. □

7.3 Quasimodularity of the number of pillowcase covers

Recall that we introduced in Section 2 the generating functions $N^0(\Pi)$ and $N^1(\Pi)$ of the number of pillowcase covers that are connected and without unramified components, respectively.

Corollary 7.4 For any ramification profile Π the counting function $N^0(\Pi)$ for connected pillowcase covers of profile Π is a quasimodular form for the group $\Gamma_0(2)$ of mixed weight less than or equal to $\text{wt}(\Pi) = \frac{1}{2}|v| + |\mu| + \ell(\mu)$.

Proof In (15) we recalled that $N'(\Pi)$ is the w -bracket of some element in $\bar{\Lambda}$ of weight $\text{wt}(\Pi)$. By Theorem 4.2 this series is thus a linear combination of auxiliary brackets with entries p_{ℓ_i} in the first arguments and a product of \bar{p}_{k_j} 's as last argument, where $\sum(l_i + 1) + \sum k_j \leq \text{wt}(\Pi)$. The classical polynomiality for triple Hurwitz numbers with p_k -arguments (summarized as [17, Theorem 4.1]) implies that the auxiliary function $A'(w^-, w^+, F)$ appearing in the definition (32) of auxiliary brackets is a polynomial of degree $\text{wt}(F) - \ell(w^-) - \ell(w^+)$. Similarly, Theorem 7.2 implies that $A'_2(\mathbf{w}, F)$ is a polynomial of degree $\text{wt}(F) - \ell(\mathbf{w})$. That is, the auxiliary bracket is the sum over all subsets $E^+(\Gamma)$ and all parity conditions $\text{par} \in \text{PC}(\Gamma)$ of graph sums of the form defined by (41) and (42) with $\sum m_i \leq \text{wt}(\Pi) - 2\ell(\mathbf{w})$. By Theorem 6.1, such a graph sum defines a quasimodular form of the weight as claimed. This gives the result for $N'(\Pi)$ and the claim for $N^0(\Pi)$ follows from inclusion–exclusion; see eg [17, Proposition 2.1]. □

8 Application to Siegel–Veech constants

In this section we show that counting pillowcase covers with certain weight functions also fall in the scope of the quasimodularity theorems and we prove Theorem 1.2.

Let $\lambda = (\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_k)$ be a partition. For $p \in \mathbb{Z}$ we define the p^{th} Siegel–Veech weight of λ to be $S_p(\lambda) = \sum_{j=1}^k \lambda_j^p$. With the conventions of Section 2, in particular the definition of Hurwitz tuples in (2), the core curves of the horizontal cylinders have the monodromies

$$\begin{aligned} \sigma_0 &= \alpha_1 \alpha_4 = \alpha_2 \alpha_3 (\gamma_1 \cdots \gamma_n)^{-1}, \\ \sigma_1 &= \alpha_1 \alpha_4 \gamma_1 = \alpha_2 \alpha_3 (\gamma_2 \cdots \gamma_n)^{-1}, \\ &\vdots \\ \sigma_n &= \alpha_1 \alpha_4 \gamma_1 \cdots \gamma_n = \alpha_2 \alpha_3. \end{aligned}$$

Motivated by the relation to area Siegel–Veech constants in Proposition 8.3 below, we define the Siegel–Veech weighted Hurwitz numbers of a Hurwitz tuple h to be

$$(56) \quad S_p(h) = \sum_{i=0}^n S_p(\sigma_i(h)).$$

Next, for $* \in \{', 0, \emptyset\}$ we package them into the generating series

$$(57) \quad c_p^*(d, \Pi) = \sum_{j=1}^{|\text{Hur}_d^*(\Pi)|} S_p(\alpha^{(j)}) \quad \text{and} \quad c_p^*(\Pi) = \sum_{d \geq 0} c_p^*(d, \Pi) q^d.$$

These series admit the following graph sum decomposition:

Proposition 8.1 *The generating series $c'_p(\Pi)$ can be expressed in terms of graph sums of triple Hurwitz numbers as*

$$c'_p(\Pi) = \sum_{\Gamma} \frac{1}{|\text{Aut}(\Gamma)|} c'_p(\Pi, \Gamma), \quad \text{where} \quad c'_p(\Pi, \Gamma) = \sum_{G \in \Gamma} c'_p(\Pi, G)$$

and where

$$c'_p(\Pi, G) = \sum_{\substack{h \in \tilde{\mathbb{N}}^{E(G)} \\ w \in \mathbb{Z}_+^{E(G)}}} \left(\sum_{e \in E(G)} h_e w_e^p \right) \prod_{e \in E(G)} w_e q^{h_e w_e} A'_2(\mathbf{w}_0, \nu) \cdot \prod_{v \in V(G)} A'(\mathbf{w}_v^-, \mathbf{w}_v^+, \mu_v) \delta(v).$$

Proof The definition (56) together with the definition of the Siegel–Veech weight is made such that a covering defined by a Hurwitz tuples is counted with the weight given by the sum over all horizontal cylinders C of $h(C)w(C)^p$. This results in the extra factor in the formula for $c'_p(\Pi, G)$ in comparison with the formula for $N'(\Pi, G)$ in (30). The whole formula is a direct consequence of the correspondence theorem, ie of Proposition 2.3. □

Corollary 8.2 *For any ramification profile Π and any odd $p \geq -1$ the generating series $c'_p(\Pi)$ for counting pillowcase covers without unramified components and with p –Siegel–Veech weight as well as the generating series $c_p^0(\Pi)$ for connected counting with p –Siegel–Veech weight are quasimodular forms for the group $\Gamma_0(2)$ of mixed weight $\leq \text{wt}(\Pi) + p + 1$.*

Roughly, this follows from the polynomiality of $A'(\cdot)$ and $A'_2(\cdot)$ (ie from Theorem 7.2) in a similar way as Corollary 7.4. The extra factor $\sum_{e \in E(G)} h_e w_e^p$ raises the degree of the polynomial by $p + 1$ and this results in the shifted weight. We explain the procedure in detail in Section 8.2.

8.1 Relation to area Siegel–Veech constants

Siegel–Veech constants measure the growth rates of the number of saddle connections or closed geodesics or equivalently embedded cylinders. Among the various possibilities

of weighting the count, the area weight is the most important due to its connection to the sum of Lyapunov exponents [11]. In detail,

$$c_{\text{area}}(X) = \lim_{L \rightarrow \infty} \frac{N_{\text{area}}(T, L)}{\pi L^2}, \quad \text{where } N_{\text{area}}(T, L) = \sum_{\substack{Z \subset X \text{ cylinder} \\ w(Z) \geq L}} \frac{\text{Area}(Z)}{\text{Area}(X)}.$$

is called the (area) Siegel–Veech constant of the flat surface X . These constants are interesting both for generic flat surfaces of a given singularity type and for pillowcase covers.

Proposition 8.3 *The area Siegel–Veech constant is related to Siegel–Veech weighted Hurwitz numbers by*

$$c_{\text{area}}(d, \Pi) = \frac{3}{\pi^2} \frac{c_{-1}^0(d, \Pi)}{N_d^0(\Pi)}.$$

In particular, knowing the numerator and the denominator of the right-hand side to be quasimodular forms, and thus knowing the asymptotic behavior of both $c_{-1}^0(d, \Pi)$ and $N_d^0(\Pi)$ as $d \rightarrow \infty$, allows us to compute the area Siegel–Veech constant of a generic surface with a given singularity type.

Proof The proof of [11, Theorem 4] or [6, Theorem 3.1] is easily adapted from torus covers to pillowcase covers. □

8.2 Quasimodularity of Siegel–Veech weighted graph sums

The goal of this section is to prove Corollary 8.2. This will follow from the following proposition. Recall from Section 6 the definition of the distinguished edges $E^+(\Gamma)$ and the parity conditions $\text{par} \in \text{PC}(\Gamma)$.

Proposition 8.4 *If $\mathbf{m} = (m_1, \dots, m_{|E(\Gamma)|})$ is a tuple of even integers, then for each $e_0 \in E(G)$ the graph sum $S_{e_0}^{\text{SV}}(\Gamma, E^+, \mathbf{m}, \text{par}) = \sum_{G \in (\Gamma, E^+)} S_{e_0}^{\text{SV}}(G, \mathbf{m}, \text{par})$, where*

$$S_{e_0}^{\text{SV}}(G, \mathbf{m}, \text{par}) = \sum_{\substack{h \in \mathbb{N}^{E(G)} \\ w \in \mathbb{N}_+^{E(G)^*}}} \sum_{\substack{w_0 \in \mathbb{N}_{>0}^{E^0(\Gamma)} \\ w_0 \cong \text{par} \pmod{2}}} \frac{h_{e_0}}{w_{e_0}} \prod_{e \in E(G)} w_i^{m_i+1} q^{h_i w_i} \prod_{v \in V(G)^*} \delta(v),$$

is a quasimodular form of mixed weight at most $k(\mathbf{m}) = \sum_i (m_i + 2)$.

Proof We may reduce to the reduced graph $\bar{\Gamma}$ by computing the loop contributions separately; compare Lemma 6.3 or rather [17, Lemma 7.5]. If the e_0 is not a loop, then the loop contributions are quasimodular by Lemma 6.4. If e_0 is a loop, we also need to take the extra factor h/w into account and note that $\sum_{w,h=1}^{\infty} hw^m q^{hw} = D_q S_m$ is quasimodular (for $m \geq 2$ even) and similarly for the odd and even variants appearing the proof of Lemma 6.4.

To deal with $\bar{\Gamma}$, we combine the construction of Section 6.2 and the Siegel–Veech weight in the proof of [17, Theorem 7.3]. More precisely, we define, if $e_0 \in E^*(\Gamma)$,

$$P_{\bar{\Gamma}, E^+, \mathbf{m}, \text{par}}^{\text{SV}}(\mathbf{z}) = \frac{D_q P^{(m_{e_0}-2)}(z_{v_1(e_0)} \pm z_{v_2(e_0)})}{P^{(m_{e_0})}(z_{v_1(e_0)} \pm z_{v_2(e_0)})} \cdot P_{\bar{\Gamma}, E^+, \mathbf{m}, \text{par}}(\mathbf{z})$$

if $m_{e_0} \geq 2$ and in the remaining case $m_{e_0} = 0$ we let

$$P_{\bar{\Gamma}, E^+, \mathbf{m}, \text{par}}^{\text{SV}}(\mathbf{z}) = \frac{L(z_{v_1(e_0)} \pm z_{v_2(e_0)})}{P(z_{v_1(e_0)} \pm z_{v_2(e_0)})} \cdot P_{\bar{\Gamma}, E^+, \mathbf{m}, \text{par}}(\mathbf{z})$$

with L as in (39). In both cases the sign is chosen according to $e_0 \in E^{\pm}(\Gamma)$. This definition replaces the factor corresponding to the edge e_0 in $P_{\bar{\Gamma}, E^+, \mathbf{m}, \text{par}}(\mathbf{z})$ by one with the extra factor h_{e_0}/w_{e_0} . This follows from the power series expansion of P and L given in [17, equation (41)]; compare also the proof of Theorem 7.3 in loc. cit. If $e_0 \in E^0(\Gamma)$, we define similarly

$$P_{\bar{\Gamma}, E^+, \mathbf{m}, \text{par}}^{\text{SV}}(\mathbf{z}) = \frac{D_q P_{\text{par}_{e_0}}^{(m_{e_0}-2)}(2z_{v_2(e_0)})}{P_{\text{par}_{e_0}}^{(m_{e_0})}(2z_{v_2(e_0)})} \cdot P_{\bar{\Gamma}, E^+, \mathbf{m}, \text{par}}(\mathbf{z})$$

or

$$P_{\bar{\Gamma}, E^+, \mathbf{m}, \text{par}}^{\text{SV}}(\mathbf{z}) = \frac{L_{\text{par}_{e_0}}(2z_{v_2(e_0)})}{P_{\text{par}_{e_0}}(2z_{v_2(e_0)})} \cdot P_{\bar{\Gamma}, E^+, \mathbf{m}, \text{par}}(\mathbf{z})$$

according to $m_{e_0} \geq 2$ or $m_{e_0} = 0$, respectively, where $L_{\text{even}}(z, \tau) = 2L(2z, 2\tau)$ and $L_{\text{odd}} = L - L_{\text{even}}$. With this modified prefactor, the same proof as in Proposition 6.5 shows that

$$(58) \quad S^{\text{SV}}(\bar{\Gamma}, E^+, \mathbf{m}, \text{par}) = [\zeta_n^0, \dots, \zeta_1^0] P_{\bar{\Gamma}, E^+, \mathbf{m}, \text{par}}^{\text{SV}}(\mathbf{z}; \tau).$$

Each of the factors in the definition of $P_{\bar{\Gamma}, E^+, \mathbf{m}, \text{par}}^{\text{SV}}(\mathbf{z})$ is a quasielliptic quasimodular form by Propositions 5.3 and 5.4. As in the case without Siegel–Veech weight, the claim follows from Theorem 5.6 and the upgrade, Lemma 5.1, to get quasimodularity by the group $\Gamma_0(2)$. □

Proof of Corollary 8.2 We want to apply Proposition 8.4. First, we pretend for the moment that the summation in Proposition 8.1 is over the normalized height space $\widehat{\mathbb{N}}^{E(G)}$ rather than over $\widetilde{\mathbb{N}}^{E(G)}$ and prove quasimodularity of the corresponding sum. Second, to reduce the graph sum expression for Siegel–Veech constants in Proposition 8.1 to those with polynomial entries, we have to mimic the argument leading to Theorem 4.2. Let

$$[F_1, \dots, F_n; F_0]^{p\text{-SV}} = \sum_{\Gamma} \sum_{G \in \Gamma} [F_1, \dots, F_n; F_0]_G^{p\text{-SV}}$$

and

$$\begin{aligned}
 & [F_1, \dots, F_n; F_0]_G^{p\text{-SV}} \\
 &= \sum_{\substack{h \in \widetilde{\mathbb{N}}^{E(G)} \\ w \in \mathbb{Z}_+^{E(G)}}} \left(\sum_{e \in E(G)} h_e w_e^p \right) \prod_{e \in E(G)} w_e q^{h_e w_e} A'_2(w_0, F_0) \prod_{v \in V(G)} A'(w_v^-, w_v^+, F_{\#_v}) \delta(v).
 \end{aligned}$$

Then Proposition 8.1 can be generalized using the correspondence theorem in the form Proposition 2.4 to

$$(59) \quad c'_p(\Pi) = \left[\prod_{i \notin S} f_{\mu_i}; \prod_{i \in S} f_{\mu_i} g_v \right]^{p\text{-SV}}$$

for any subset $S \subseteq \{1, \dots, n\}$. To prove quasimodularity we start with $S = \emptyset$ and decompose g_v as a linear combination of $P_{I,J} = \prod_{j \in J} p_{b_j} \prod_{i \in I} \bar{p}_{a_i}$ for some a_i and b_j . For the summands where $J = \emptyset$ we write the first argument of the bracket as a linear combination of products of p_k and use the polynomiality for triple Hurwitz numbers with p_k -argument and Theorem 7.2 to conclude thanks to Proposition 8.4.

To deal with the summands where $J \neq \emptyset$, we write $P_{I,J}$ as a linear combination of terms that are products of several f_k 's and one g_v . This is possible by Theorem 3.3. Since $J = \emptyset$, each term involves at least one f_k . For each term we now apply (59) twice to move the f_k -product to the first argument of the auxiliary bracket. By this procedure we have reduced the weight of the g_v in the second argument and we conclude by induction.

We have to justify the first simplification concerning height spaces. Note that the shift to $\widehat{\mathbb{N}}^{E(G)}$ does not change the q -exponents by Lemma 6.2, so we may focus on the Fourier coefficients. Since the expression for $c'_p(\Pi)$ in Proposition 8.1 involves a summation over all orientations, we may combine the contributions of G and the

reverse orientation $-G$, where the arrows of all edges except for those emanating from v_0 have been inverted. We thus obtain a prefactor of

$$\sum_{h \in \mathbb{N}_{\geq ae}} \frac{h - \Delta(e)}{w} + \sum_{h \in \mathbb{N}_{\geq (1-ae)}} \frac{h + \Delta(e)}{w} \quad \text{instead of} \quad \sum_{h \in \mathbb{N}_{\geq ae}} \frac{h}{w} + \sum_{h \in \mathbb{N}_{\geq (1-ae)}} \frac{h}{w}$$

when using $\widehat{\mathbb{N}}^{E(G)}$. The difference is only the term $h = 0$, ie without an h/w -prefactor, and we recursively know these graph sums to be quasimodular (in fact of smaller weight). Note that this argument also shows that the Siegel–Veech constant does not depend on the height parameters ε_i chosen on the base.

Finally, the quasimodularity for $c_p^0(\Pi)$ follows from the usual inclusion–exclusion formulas; see [6, Proposition 6.2] for the version with Siegel–Veech weight. \square

8.3 Siegel–Veech weight and representation theory

The reader familiar with [6] will recall that counting function for Hurwitz tuples, even with Siegel–Veech weight, can be expressed efficiently using the representation theory of the symmetric group. More precisely, for Π the profile of a torus cover,

$$(60) \quad N_d(\Pi) = \sum_{\lambda \in \mathcal{P}(d)} \prod_{i=1}^n f_{\mu_i}(\lambda) \quad \text{and} \quad c_p(d, \Pi) = \sum_{\lambda \in \mathcal{P}(d)} \prod_{i=1}^n f_{\mu_i}(\lambda) T_p(\lambda),$$

where $T_p(\lambda) = \sum_{\xi \in Y_\lambda} h(\xi)^{p-1}$ and where $h(\xi)$ is the hook-length of the cell ξ of the Young diagram Y_λ .

It would be very useful to have a similar formula for the Siegel–Veech weighted counting of pillowcase covers. We are only aware of the following much more complicated formula:

Proposition 8.5 *The number of all covers of degree d with profile Π counted with p -Siegel–Veech weight is*

$$c_p(d, \Pi) = \frac{1}{l(\mu) + 1} \sum_{k=0}^{l(\mu)} \sum_C S_p(C) \sum_{\lambda, \lambda'} \sqrt{w(\lambda)w(\lambda')} |C| \chi^\lambda(C) \chi^{\lambda'}(C) \cdot g_v(\lambda') \prod_{i=1}^k f_{\mu_i}(\lambda') \prod_{i=k+1}^{l(\mu)} f_{\mu_i}(\lambda).$$

In particular, we are not aware of an operator on Fock space whose q -trace computes the generating series with Siegel–Veech weight. Note that the \mathfrak{W} -operator

of [14, Theorem 4] has the property $\langle v_\lambda | \mathfrak{W} | v_\lambda \rangle = w(\lambda)$, but it is not true that $\langle v_\lambda | \mathfrak{W} | v_\nu \rangle = \sqrt{w(\lambda)w(\nu)}$ for $\lambda \neq \nu$. Finding a vertex operator with this property would be a way to use Proposition 8.5 to express Siegel–Veech weighted generating series as q -traces. Writing c_p as a vacuum expectation using the one-sided pillow-case operator $\Gamma_{\sqrt{w}}$ seems even more promising.

Note that the relation between $c_p(\Pi)$ and $c'_p(\Pi)$ is given by the formula

$$(61) \quad c_p(\Pi) = c'_p(\Pi)N() + N'(\Pi)c_p().$$

The proof of [6, Proposition 6.2] applies verbatim in our situation.

Proof The monodromy of the core curves of the cylinders of a Hurwitz tuple $h \in \text{Hur}_d(\Pi)$ is given by

$$\begin{aligned} \sigma_0 &= \alpha_1\alpha_4 = \alpha_2\alpha_3(\gamma_1 \cdots \gamma_n)^{-1}, \\ \sigma_1 &= \alpha_1\alpha_4\gamma_1 = \alpha_2\alpha_3(\gamma_2 \cdots \gamma_n)^{-1}, \\ &\vdots \\ \sigma_n &= \alpha_1\alpha_4\gamma_1 \cdots \gamma_n = \alpha_2\alpha_3. \end{aligned}$$

To count Hurwitz tuples with Siegel–Veech weight, say for the k^{th} cylinder, we split the defining equation as

$$(62) \quad \alpha_1\alpha_4\gamma_1, \dots, \gamma_k = c = \alpha_2\alpha_3\gamma_n^{-1} \cdots \gamma_{k+1}^{-1}$$

and count the solutions of each side separately. That is, we denote by $C_1, C_2, C_3, C_4, C, C_1^\mu, \dots, C_n^\mu$, respectively, the conjugacy classes of permutations of type

$$(v, 2^{|v|/2-d}), (2^d), (2^d), (2^d), (k^{c_k}), (\mu_1, 1^{d-\mu_1}), \dots, (\mu_n, 1^{d-\mu_n}).$$

We denote by $c_p(S_{2d}; C_1, C_4, C_1^\mu, \dots, C_k^\mu, C)$ the number of solutions of

$$(63) \quad \alpha_1\alpha_4\gamma_1, \dots, \gamma_k = c$$

with α_i of conjugacy class C_i and γ_i of conjugacy class C_i^ν , and c of conjugacy class C , counted with weight $S_p(c) = S_p(C)$. We claim that

$$\begin{aligned} c_p(S_{2d}; C_1, C_4, C_1^\mu, \dots, C_k^\mu, C) \\ = \frac{|C_1||C_4||C_1^\mu| \cdots |C_k^\mu||C|}{|G|} S_p(C) \cdot \sum_{\chi} \frac{\chi(C_1)\chi(C_4)\chi(C_1^\mu) \cdots \chi(C_k^\mu)\chi(C)}{\chi(1)^{k+1}}. \end{aligned}$$

To see this, we revisit the proof of the orthogonality relations; see [25, Theorem 7.2.1]. We introduce the class function

$$\phi(x) = S_p(C)1_{\{x \in C\}} = \sum_{\chi} c_{\chi} \chi$$

with

$$c_{\chi} = \int_G S_p(C)1_{\{x \in C\}} \bar{\chi}(x) dx = \frac{|C|}{|G|} S_p(C) \chi(C).$$

We have

$$\begin{aligned} I(\phi) &= \int_{G^{k+2}} \phi(t_1 \alpha_1 t_1^{-1} t_4 \alpha_4 t_4^{-1} s_1 \gamma_1 s_1^{-1} \cdots s_k \gamma_k s_k^{-1} y) dt_1 dt_4 ds_1 \cdots ds_k \\ &= \sum_{\chi} c_{\chi} \frac{\chi(C_1) \chi(C_4) \chi(C_1^{\mu}) \cdots \chi(C_k^{\mu}) \chi(y)}{\chi(1)^{k+2}} \end{aligned}$$

and the left-hand side is $I(\phi) = N_p(S_{2d}; C_1, C_4, C_1^{\mu}, \dots, C_k^{\mu}, C) / |G|^{k+2}$. Taking $y = 1$, we get the formula of the claim.

In the second step we count the solution of the right equality of (62) with weight $c_p(S_{2d}; C_1, C_4, C_1^{\mu}, \dots, C_k^{\mu}, C)$. The class function is now

$$\phi(x) = \frac{1}{|C|} c_p(S_{2d}; C_1, C_4, C_1^{\mu}, \dots, C_k^{\mu}, x) 1_{x \in C},$$

and its coefficients are

$$c_{\chi} = \frac{1}{|G|} N_p(S_{2d}; C_1, C_4, C_1^{\mu}, \dots, C_k^{\mu}, C) \bar{\chi}(C).$$

With the argument as above we conclude that the number of solutions counted with weight is

$$|C_2| |C_3| |C_n^{\mu}| \cdots |C_{k+1}^{\mu}| \sum_{\chi} c_{\chi} \frac{\chi(C_2) \chi(C_3) \chi(C_n^{\mu}) \cdots \chi(C_{k+1}^{\mu})}{\chi(1)^{n-k+1}}.$$

We sum now on all conjugacy classes C and use the definition of f_{μ} and g_{ν} to obtain the result. □

9 Example: $\mathcal{Q}(2, 1, -1^3)$

In this section, we treat the example of the stratum $\mathcal{Q}(2, 1, -1^3)$ from A to Z to illustrate all sections of the paper. This stratum is the lowest-dimensional example exhibiting all the relevant aspects.

Integer points in the stratum $\mathcal{Q}(2, 1, -1^3)$ correspond in our setting to covers of the pillow ramified over five points: the four corners P_1, \dots, P_4 and an additional point P_5 (see Figure 1), with the ramification profile $\Pi = (\mu^{(1)}, \mu^{(2)}, \mu^{(3)}, \mu^{(4)}, \mu^{(5)})$, where $\mu^{(1)} = (3, 1, 1, 1, 2^{d-3})$ over P_1 , $\mu^{(2)} = \mu^{(3)} = \mu^{(4)} = (2^d)$ over P_2, P_3 and P_4 , and $\mu^{(5)} = (2, 1^{2d-2})$ over P_5 . Here $2d$ is the degree of the cover. In this particular case, we cannot have covers with at least two ramified connected components, so $N'(\Pi) = N^0(\Pi)$.

9.1 Counting covers

By [14] and Theorem 1.1, the generating series $N^0(\Pi)$ is a quasimodular form for $\Gamma_0(2)$ of mixed weight less than or equal to 6. Computing the first coefficients of the series we get that¹

$$N^0(\Pi) = 360G_2(q^2)^3 - 360G_2(q)G_2(q^2)^2 + 72G_2(q)^2G_2(q^2) - 30G_4(q^2)G_2(q^2) - \frac{5}{4}G_4(q^2) + 3G_2(q)^2 + 15G_2(q^2)^2 - 15G_2(q)G_2(q^2).$$

Our goal is to retrieve this result by considering all graph contributions. Standard Hurwitz theory (see (27) and [14]) gives that

$$N^0(\Pi) = \frac{1}{N(\Pi_\emptyset)} \sum_{\lambda} g_{3,1,1,1}(\lambda) f_2(\lambda) w^2(\lambda) q^{|\lambda|/2},$$

where $g_{3,1,1,1}(\lambda) = f_{3,1,1,1,2,\dots,2}(\lambda) / f_{2,\dots,2}(\lambda)$ and $w(\lambda)$ is as in (14).

Following the strategy in Theorem 4.2 we need to express the graph sums with local contributions $g_{3,1,1,1}$ and f_2 by graph sums using the functions p_k and \bar{p}_k for which we have nice polynomiality results. We compute that

$$g_{(3,1,1,1)} = -\frac{1}{4}\bar{p}_1 p_1 + \frac{1}{108}\bar{p}_1^3 - \frac{1}{36}\bar{p}_2 \bar{p}_1 + \frac{3}{8}\bar{p}_1 + \frac{2}{27}\bar{p}_3, \quad f_2 = \frac{1}{2}p_2.$$

The definition

$$\bar{g}_{(3,1,1,1)} := \frac{1}{108}\bar{p}_1^3 - \frac{1}{36}\bar{p}_2 \bar{p}_1 + \frac{3}{8}\bar{p}_1 + \frac{2}{27}\bar{p}_3, \quad g_{(3,1,1,1)}^{\text{deg}} := -\frac{1}{4}\bar{p}_1$$

and the resulting decomposition

$$g_{(3,1,1,1)} = \bar{g}_{(3,1,1,1)} + g_{(3,1,1,1)}^{\text{deg}} p_1$$

¹We choose in this paper to consider series in q^d rather than as series in q^{2d} as in [14]. The integer d can be seen as the area of the cover while $2d$ is the degree of the cover.




Figure 5: Admissible graphs for $\bar{g}_{(3,1,1,1)} \cdot f_2$.

isolates the products of \bar{p}_i 's so we can use the polynomiality results of Section 7. Writing $\bar{g}_{3111} f_2 = \frac{1}{2} f_2 f_1 g_{11} + f_2 g_{3111}$ and $g_{3111}^{\text{deg}} p_1 f_2 = -\frac{1}{2} g_{11} f_2 f_2 + \frac{1}{48} g_{11} f_2$, we can make geometric sense of the formal decomposition, as counting degenerate covers, eg in the stratum $\mathcal{Q}(2, 0, -1^2)$. We determine the simple Hurwitz numbers with 2-stabilization A'_2 for products of \bar{p}_i 's, proven to be quasipolynomial in Theorem 7.2 by computing the first few terms. As a result,

$$A'_2((w), \bar{g}_{(3,1,1,1)}) = \frac{1}{24} w^2 + \frac{1}{3},$$

$$A'_2((w_1, w_2, w_3), \bar{g}_{(3,1,1,1)}) = \begin{cases} \frac{1}{2} & \text{if two } w_i \text{ are odd,} \\ \frac{3}{2} & \text{if all } w_i \text{ are even,} \end{cases}$$

$$A'_2((w), g_{(3,1,1,1)}^{\text{deg}}) = -\frac{1}{4}.$$

We recall that these polynomials are only defined for $\sum w_i$ even. Similarly, the double Hurwitz numbers A' for products of p_i 's are polynomials; in fact,

$$A'((w), (w), p_1) = 1, \quad A'((w_1), (w_2, w_3), f_2) = 1.$$

We have thus collected all local polynomials.

Now we glue the local surfaces together, encoding the gluings by the various possible global graphs. The contribution of $\bar{g}_{(3,1,1,1)} \cdot f_2$ is encoded in graphs with two vertices,




Figure 6: Admissible graphs for $\bar{g}_{(3,1,1,1)} \cdot f_2$.

one special vertex v_0 of valency 1 or 3, and another trivalent vertex v_1 since all other valencies result in a zero local polynomial and thus in a zero contribution. By convention, we represent v_0 as the bottom of the graph, marked with a cross. Disregarding orientations, we obtain three admissible graphs, shown in Figure 5.

Similarly, the contribution of $g_{(3,1,1,1)}^{\text{deg}} \cdot p_1 \cdot f_2$ is encoded in graphs with three vertices, one special vertex v_0 of valency 1, a vertex v_1 of valency 2 and a vertex v_2 of valency 3. The graphs are listed in Figure 6.

Next, we are supposed to sum over all possible orientations of these graphs. We sort the orientations by the subset of coherently oriented edges, ie by those distinguished with a $+$ in the notation of Section 6. Note that certain decorations give trivial contributions, such as the graph A with the loop decorated with $+$. In fact, considering all possible orientations of half-edges compatible with this decoration, we see that we get incompatible width conditions for the vertex v_1 . This implies that, integrating the corresponding propagator, we will get terms like $\delta(w_2 = w_1 + w_2)$ that are always trivial.

The next step is to associate to each decorated graph its propagator, and then to integrate this propagator (get its ζ^0 -coefficient) to obtain the contributions of each individual graph. In Tables 1 and 2 we consider only decorations with nontrivial contributions. For each vertex, we indicate the corresponding integration variable z_i (on the left of the vertex) and the corresponding local polynomial (on the right). For each decorated graph we give the associated propagator and the contribution to the volume. The contributions were computed using three independent methods: the reduction algorithm used in the proof of Theorem 5.6, the computation of the first terms of the q -series using graph sums (42), and, independently, using extraction of $[\zeta^0]$ coefficients, combined with a numerical test of quasimodularity (test of linear dependency with the basis of quasimodular forms).

We provide details of the method of Theorem 5.6 for the graphs A and D , to illustrate the algorithm in the proof. We denote by $T = \{0, \frac{1}{2}, \frac{1}{2}\tau, \frac{1}{2}(1 + \tau)\}$ the set of 2-torsion points and let $T^* = T \setminus \{0\}$. Note that, considering the residues at the poles at T , we have

$$P(2z; \tau) = \frac{1}{4} \sum_{a \in T} P(z - a; \tau).$$

Consequently, $P(2z)P(z)$ is an elliptic function with a pole of order 4 at 0, and poles of order 2 at T^* , whose residues are easily compensated by $\frac{1}{4} \frac{1}{6} P''(z)$ and $\frac{1}{4} P(z - a)P(a)$ for $a \in T^*$, respectively. Then, compensating for the remaining pole

graph Γ	$\frac{1}{ \text{Aut}(\Gamma) }$	propagator (P) contribution (C)
	$\frac{1}{2}$	<p>P: $(\frac{1}{24} P^{(2)}(z) + \frac{1}{3} P(z)) P(2z)$</p> <p>C: $\frac{7}{30} G_6 - \frac{8}{3} G_4 G_2 + \frac{5}{9} G_4 - \frac{4}{3} G_2^2$</p>
	$\frac{1}{2}$	<p>P: $\frac{1}{2} P_{\text{odd}}(z)^2 P_{\text{even}}(z)$</p> <p>C: $-\frac{1}{3}(16G_{22}G_{42} + 8G_2G_{42}) + \frac{1}{5}(-192G_{22}^3 + 608G_2G_{22}^2 - 384G_2^2G_{22} + 64G_2^3)$</p>
	$\frac{1}{6}$	<p>P: $\frac{3}{2} P_{\text{even}}(z)^3$</p> <p>C: $48G_{22}G_{42} - 16G_2G_{42} + \frac{1}{5}(-832G_{22}^3 + 768G_2G_{22}^2 - 384G_2^2G_{22} + 64G_2^3)$</p>
	$\frac{1}{4}$	<p>P: $\frac{1}{2} S_{0,\text{odd}} P_{\text{even}}(z) P(2z)$</p> <p>C: $(G_2 - 2G_{22} - \frac{1}{24})(\frac{5}{6} G_4 - 2G_2^2)$</p>
	$\frac{1}{4}$	<p>P: $\frac{3}{2} S_{0,\text{even}} P_{\text{even}}(z) P(2z)$</p> <p>C: $3(2G_{22} + \frac{1}{12})(\frac{5}{6} G_4 - 2G_2^2)$</p>

Table 1: Graphs for $\mathcal{Q}(2, 1, -1^3)$: contribution of $\bar{g}_{(3,1,1,1)} \cdot f_2$.

of order 2 at 0, we get

$$P(2z)P(z) = \frac{1}{24} P''(z) + G_2 P(z) + \frac{1}{4} \sum_{a \in T^*} P(a)(P(z) + P(z - a)) + \frac{5}{3} G_4 - 4G_2^2.$$

graph Γ	$\frac{1}{ \text{Aut}(\Gamma) }$	propagator (P) contribution (C)
	$\frac{1}{2}$	<p>P: $-\frac{1}{4}P(z_1)P(z_1-z_2)P(2z_2)$</p> <p>C: A-3B</p>
	$\frac{1}{2}$	<p>P: $-\frac{1}{4}P(z_1)P(z_1+z_2)P(2z_2)$</p> <p>C: A+B</p>
	1	<p>P: $-\frac{1}{4}P(z_2)P(z_1-z_2)P(z_1+z_2)$</p> <p>C: A+B</p>
	$\frac{1}{2}$	<p>P: $-\frac{1}{4}S_0P(z_2)P(2z_2)$</p> <p>C: $-\frac{1}{4}(G_2+\frac{1}{24})(-\frac{20}{3}G_{42}+80G_{22}^2-80G_2G_{22}+16G_2^2)$</p>

Table 2: Graphs for $Q(2, 1, -1^3)$: contribution of $g_{(3,1,1,1)}^{\text{deg}} \cdot p_1 \cdot f_2$.

Since P is the derivative of a 1-periodic function Z , the contour integral is then reduced to

$$[\zeta^0]P(2z)P(z) = \frac{5}{3}G_4 - 4G_2^2.$$

Proceeding similarly with the term $P(2z)P''(z)$, we get

$$P(2z)P''(z) = \frac{1}{80}P^{(4)}(z) + \frac{1}{2}G_2P''(z) + \frac{49}{2}G_4P(z) + \frac{1}{4} \sum_{a \in T^*} (P(a)P''(z) + P''(a)P(z-a)) + \frac{28}{5}G_6 - 64G_4G_2.$$

hence

$$[\zeta^0]P(2z)P''(z) = \frac{28}{5}G_6 - 64G_4G_2,$$

which gives the contribution of the graph A . The contributions of the graphs B and C are computed similarly using the decomposition

$$P(2z; 2\tau) = \frac{1}{4}(P(z; \tau) + P(z - \frac{1}{2}; \tau)).$$

Computing the contribution of the graph D_a we will see quasielliptic functions appearing. We first decompose $P(z_1 - z_2)P(2z_2)$ in the additive basis with respect to z_2 . For this purpose decompose as usual $P(2z_2)$ into the sum of the four contributions of the 2-torsion points. The term $P(z_1 - z_2)P(z_2)$ has a pole of order 2 at $z_2 = z_1$ (which is compensated by $P(z_1 - z_2)P(z_1)$), a pole of order 2 at $z_2 = 0$ (which is compensated by $P(z_1)P(z_2)$), and it also has a pole of order 1 at $z_2 = z_1$ (which is compensated by $Z(z_1 - z_2)P'(z_1)$), and a pole of order 1 at $z_2 = 0$ (finally compensated by $Z(z_2)P'(z_1)$). Proceeding similarly for the three other 2-torsion points, we get

$$P(z_1 - z_2)P(2z_2) = \frac{1}{4} \sum_{a \in T} ([P(z_2 - a) + P(z_2 - z_1)]P(z_1 - a) + [Z(z_2 - a) - Z(z_2 - z_1)]P'(z_1 - a)) + R(z_1),$$

where

$$R(z_1) = \frac{4}{3}P''(2z_1) - \frac{1}{4} \sum_{a \in T} (P'(z_1 - a)Z(z_1 - a)) - 4P(2z_1)G_2 + \frac{1}{6}G_2^2 - \frac{5}{3}G_4.$$

As a result,

$$[\zeta_2^0]P(z_1)P(z_1 - z_2)P(2z_2) = P(z_1)R(z_1).$$

We already showed how to treat terms like $P(z_1)P''(2z_1)$ and $P(z_1)P(2z_1)$, so we focus on $P(z_1)P'(z_1 - a)Z(z_1 - a)$ in the sequel. The product $S_2 = P(z_1)P'(z_1 - a)$ is an elliptic function, so, examining the pole orders, we get

$$S_2 = \begin{cases} \frac{1}{12}P^{(3)}(z_1) + 2P'(z_1)G_2 & \text{if } a = 0, \\ P''(a)[Z(z_1 - a) - Z(z_1)] + P(a)P'(z_1 - a) & \text{if } a = \frac{1}{2}, \\ P''(a)[Z(z_1 - a) - Z(z_1) + \frac{1}{2}] + P(a)P'(z_1 - a) & \text{if } a \in \{\frac{1}{2}\tau, \frac{1}{2}(1 + \tau)\}. \end{cases}$$

When multiplying the right-hand side by $Z(z_1 - a)$, the right-hand side belongs to the additive basis, except for the case $Z(z_1)Z(z_1 - a)$ for $a \in T^*$. But since

$$\Delta(Z(z_1)Z(z_1 - a)) = \frac{1}{2}\Delta(Z(z_1)^2 + Z(z_1 - a)^2),$$

the function $Z(z_1)Z(z_1 - a) - \frac{1}{2}(Z(z_1)^2 + Z(z_1 - a)^2)$ is elliptic with poles of order less than or equal to 2 at 0 and a , we get

$$Z(z_1)Z(z_1 - a) = \frac{1}{2}(Z(z_1)^2 + Z(z_1 - a)^2 - P(z_1) - P(z_1 - a)) + 3G_2 - \frac{1}{2}P(a) + \begin{cases} 0 & \text{if } a = \frac{1}{2}, \\ \frac{1}{2}[Z(z - a) - Z(a)] + \frac{1}{8} & \text{if } a = \frac{1}{2}\tau, \frac{1}{2}(1 + \tau), \end{cases}$$

and finally

$$[\zeta_1^0]Z(z_1)Z(z_1 - a) = G_2 - \frac{1}{2}P(a) + \begin{cases} \frac{1}{6} & \text{if } a = \frac{1}{2}, \\ -\frac{5}{24} & \text{if } a = \frac{1}{2}\tau, \frac{1}{2}(1 + \tau). \end{cases}$$

In total the contribution of graph D_a is a combination of the quasimodular forms

$$A = (-40G_{22} + 10G_2)G_{42} + 352G_{22}^3 - 408G_2G_{22}^2 + 144G_2^2G_{22} - 16G_2^3, \\ B = -\frac{5}{4}G_{42} + 12G_{22}^2 - 12G_2G_{22} + 3G_2^2,$$

as indicated in the table. Note that each contribution is a quasimodular form for $\Gamma(2)$, but also a series in q , so in fact it is a quasimodular form for $\Gamma_0(2)$ (as we remarked already in Lemma 5.1). The sum of all contributions in the table is finally the quasimodular form given at the beginning of this subsection.

9.2 Siegel–Veech weight

Everything is ready to compute the contributions of these graphs with Siegel–Veech weight. We have the same graphs with the same local polynomials; we just associate a slightly modified propagator to take care of the weight. The recipe to get this propagator from the old one is simple, for example if they aren't distinguished loops: for each edge of the graph replace the corresponding factor P by L and P^m by $D_q P^{m-2}$ if $m \geq 2$, and then sum over all edges of the graph (see Section 8.2 for a precise statement). The last step of integration is then similar to the previous case. We give the results in Table 3 (we do not copy the factors $1/|\text{Aut}(\Gamma)|$ which are the same; we group the graphs of same type). In the column contribution, lwt stands for lower-weight terms. In the compilation of the table we used

$$S_0^{\text{SV}} = \sum_{w,h=1}^{\infty} hq^{wh} = S_0 = G_2 + \frac{1}{24}, \quad S_{0,\text{even}}^{\text{SV}} = \sum_{w,h=1}^{\infty} hq^{(2w)h} = G_{22} + \frac{1}{24}$$

and $S_{0,\text{odd}}^{\text{SV}} = S_0^{\text{SV}} - S_{0,\text{even}}^{\text{SV}}$. Summing up, we get the generating series

$$c_{-1}^0(\Pi) = 245G_{22}^3 - 245G_2G_{22}^2 + 49G_2^2G_{22} - \frac{245}{12}G_2G_{42}G_{22} + \text{lwt},$$

graph	propagator (P) contribution (C)
A	$P: \frac{1}{2} \left[\left(\frac{1}{24} D_q P(z) + \frac{1}{3} L(z) \right) P(2z) + \left(\frac{1}{24} P^{(2)}(z) + \frac{1}{3} P(z) \right) L(2z) \right]$ $C: \frac{1}{2} \left[-\frac{880}{3} G_{22}^3 + 340 G_2 G_{22}^2 - 120 G_2^2 G_{22} + \frac{100}{3} G_{42} G_{22} + \frac{40}{3} G_2^3 + -\frac{25}{3} G_{42} G_2 \right] + \text{lwt}$
B	$P: \frac{1}{2} \cdot \frac{1}{2} (2L_{\text{odd}}(z) P_{\text{odd}}(z) P_{\text{even}}(z) + P_{\text{odd}}(z)^2 L_{\text{even}}(z)) + \frac{1}{6} \cdot \frac{3}{2} (3P_{\text{even}}(z)^2 L_{\text{even}}(z))$ $C: \frac{1}{2} \left[-\frac{560}{3} G_{22}^3 + 280 G_2 G_{22}^2 - 160 G_2^2 G_{22} + \frac{100}{3} G_{42} G_{22} + \frac{80}{3} G_2^3 - \frac{50}{3} G_{42} G_2 \right]$
C	$P: \frac{1}{4} \left[\left(\frac{1}{2} S_{0,\text{odd}} + \frac{3}{2} S_{0,\text{even}} \right) \cdot (L_{\text{even}}(z) P(2z) + P_{\text{even}}(z) L(2z)) \right. \\ \left. + \left(\frac{1}{2} S_{0,\text{odd}}^{\text{SV}} + \frac{3}{2} S_{0,\text{even}}^{\text{SV}} \right) \cdot P_{\text{even}}(z) P(2z) \right]$ $C: \frac{1}{4} \left[180 G_{22}^3 - 115 G_2 G_{22}^2 - 29 G_2^2 G_{22} - 15 G_{42} G_{22} + 13 G_2^3 - \frac{65}{12} G_{42} G_2 \right] + \text{lwt}$
D	$P: \frac{1}{2} \cdot \left(-\frac{1}{4} \right) \cdot (L(z_1) P(z_1 - z_2) P(2z_2) + P(z_1) L(z_1 - z_2) P(2z_2) \\ + P(z_1) P(z_1 - z_2) L(2z_2) + L(z_1) P(z_1 + z_2) P(2z_2) \\ + P(z_1) L(z_1 + z_2) P(2z_2) + P(z_1) P(z_1 + z_2) L(2z_2))$ $C: \frac{1}{2} \left[352 G_{22}^3 - 408 G_2 G_{22}^2 + 144 G_2^2 G_{22} - 40 G_{42} G_{22} - 16 G_2^3 + 10 G_{42} G_2 \right] + \text{lwt}$
E	$P: -\frac{1}{4} (L(z_1) P(z_1 - z_2) P(z_1 + z_2) + P(z_1) L(z_1 - z_2) P(z_1 + z_2) \\ + P(z_1) P(z_1 - z_2) L(z_1 + z_2))$ $C: 264 G_{22}^3 - 306 G_2 G_{22}^2 + 108 G_2^2 G_{22} - 30 G_{42} G_{22} - 12 G_2^3 + \frac{15}{2} G_{42} G_2 + \text{lwt}$
F	$P: \frac{1}{2} \cdot \left(-\frac{1}{4} \right) \cdot (P(z_2) P(2z_2) + L(z_2) P(2z_2) + P(z_2) L(2z_2))$ $C: \frac{1}{2} \left[-\frac{65}{2} G_2 G_{22}^2 + \frac{65}{2} G_2^2 G_{22} - \frac{13}{2} G_2^3 + \frac{65}{24} G_{42} G_2 \right] + \text{lwt}$

Table 3: Graphs for $\mathcal{Q}(2, 1, -1^3)$: Siegel–Veech contribution.

as can be checked using Proposition 8.5 and formula (61) (here $c'_p = c_p^0$). The lower-weight terms are weight 2 and 4 terms, as we can expect from the weight of $N^0(\Pi)$ (as L contains lower-weight terms). This series is proportional to $N^0(\Pi)$, since this stratum is nonvarying (see [4] for more explanation). We will see in the next section that the coefficient of proportionality is the Siegel–Veech constant of the stratum.

9.3 Contributions to volumes of strata and Siegel–Veech constants

Evaluation of volumes and Siegel–Veech constants of strata are closely related to the asymptotics of the generating series $N^0(\Pi)$ and $c_{-1}(\Pi)$ as q tends to 1 (or equivalently τ tends to 0), as stated in [14] (see also [16, Proposition 7]). These asymptotics can be easily obtained thanks to the quasimodularity property for Eisenstein series: the transformation $\tau \rightarrow -1/\tau$ relates the asymptotics as $\tau \rightarrow 0$ to the asymptotics as $\tau \rightarrow i\infty$.

We overview briefly the results of [6, Section 9] here and define two polynomials describing the growth of a quasimodular form (for $\Gamma_0(2)$) near $\tau = 0$, so at the same time the average growth of its Fourier coefficients.

We recall that $\text{QM}(\Gamma_0(2))$ is the space of even-weight quasimodular forms for $\Gamma_0(2)$, and it is generated as a polynomial ring by G_2, G_{22} and G_{42} ; see (37).

Definition 9.1 We define the map Ev as the unique algebra homomorphism from $\text{QM}(\Gamma_0(2))$ to $\mathbb{Q}[X]$ sending G_2 to $-\frac{1}{24}X - \frac{1}{2}$, G_{22} to $-\frac{1}{96}X - \frac{1}{4}$ and G_{42} to $\frac{1}{3840}X^2$.

Setting $h = -2\pi i \tau$, we define

$$\text{ev}[F](h) = \frac{1}{h^k} \text{Ev}[F]\left(-\frac{4\pi^2}{h}\right) \in \mathbb{Q}[\pi^2]\left[\frac{1}{h}\right]$$

for $F \in \text{QM}_{2k}(\Gamma_0(2))$ (weight $2k$ quasimodular form). This polynomial describes the growth of $F(\tau)$ near $\tau = 0$ directly (also for mixed-weight forms), as proved in the following proposition:

Proposition 9.2 For $F \in \text{QM}(\Gamma_0(2))$ we have

$$F(i\varepsilon) = \text{ev}[F](2\pi\varepsilon) + (\text{small}) \quad (\varepsilon \searrow 0),$$

where “small” means terms that tend exponentially quickly to 0.

Proof This is directly derived from the modularity properties $G_2(-1/\tau) = \tau^2 G_2(\tau) - \tau/4\pi i$ and $G_4(-1/\tau) = \tau^4 G_4(\tau)$. □

In Table 4 we give the h -evaluation of all individual graphs. Note that the h -evaluation of lower-weight terms is $O(1/h^4)$. In this table,

$$\text{vol} = \frac{2 \dim}{2^{\dim} \dim!} \cdot \lim_{h \rightarrow 0} (\text{ev}[N^0(\Pi)](h) \cdot h^{\dim})$$

is the volume of $\mathcal{Q}(2, 1, -1^3)$ (with respect to the Eskin–Okounkov convention; the volume for the Athreya–Eskin–Zorich convention has an additional factor $\frac{1}{2} \cdot 4^5 \cdot 3! = 3072$; see [16, Lemma 2] for discussion concerning volume normalizations), and $\dim = \dim_{\mathbb{C}} \mathcal{Q}(2, 1, -1^3) = 5$.

In the general case, the Siegel–Veech constant c_{area} is computed using

$$\frac{1}{3} \pi^2 c_{\text{area}} = \lim_{D \rightarrow \infty} \frac{\sum_{d=1}^D c_{-1}^0(d, \Pi)}{\sum_{d=1}^D N^0(d, \Pi)} = \frac{\lim_{h \rightarrow 0} (\text{ev}[c_{-1}^0(\Pi)](h) \cdot h^{\dim})}{\lim_{h \rightarrow 0} (\text{ev}[N^0(\Pi)](h) \cdot h^{\dim})},$$

since both limits are finite.

graph Γ	$\text{ev}[N^0(\Gamma, \Pi)](h)$	$\text{ev}[c_{-1}(\Gamma, \Pi)](h)$
A	$\frac{4}{45}\pi^4/h^5 + O(1/h^4)$	$O(1/h^4)$
B	$\frac{2}{15}\pi^4/h^5 + O(1/h^4)$	$\frac{5}{36}\pi^4/h^5 + O(1/h^4)$
C	$\frac{1}{9}\pi^4/h^5 + O(1/h^4)$	$\frac{11}{144}\pi^4/h^5 + O(1/h^4)$
D	$O(1/h^4)$	$O(1/h^4)$
E	$O(1/h^4)$	$O(1/h^4)$
F	$-\frac{1}{36}\pi^4/h^5 + O(1/h^4)$	$-\frac{13}{288}\pi^4/h^5 + O(1/h^4)$
total	$\text{vol} = \frac{1}{3072}\pi^4$	$\frac{1}{3}\pi^2 c_{\text{area}} = \frac{49}{72}$

Table 4: Graphs for $\mathcal{Q}(2, 1, -1^3)$: growth polynomials and contribution to the volume and the Siegel–Veech constant of the stratum.

In our example, the stratum is nonvarying, so the series $c_{-1}^0(\Pi)$ and $N^0(\Pi)$ are proportional (not the individual graph contributions though). The Siegel–Veech constant is just the proportionality factor. We get the constant

$$\frac{1}{3}\pi^2 c_{\text{area}}(\mathcal{Q}(2, 1, -1^3)) = \frac{49}{72},$$

in agreement with the value computed in [4].

References

- [1] **A Aggarwal**, *Large genus asymptotics for volumes of strata of abelian differentials*, J. Amer. Math. Soc. 33 (2020) 941–989 MR
- [2] **JS Athreya, A Eskin, A Zorich**, *Right-angled billiards and volumes of moduli spaces of quadratic differentials on $\mathbb{C}P^1$* , Ann. Sci. École Norm. Sup. 49 (2016) 1311–1386 MR
- [3] **J Böhm, K Bringmann, A Buchholz, H Markwig**, *Tropical mirror symmetry for elliptic curves*, J. Reine Angew. Math. 732 (2017) 211–246 MR Correction in 760 (2020), 163–164
- [4] **D Chen, M Möller**, *Nonvarying sums of Lyapunov exponents of abelian differentials in low genus*, Geom. Topol. 16 (2012) 2427–2479 MR
- [5] **D Chen, M Möller, A Sauvaget, D Zagier**, *Masur–Veech volumes and intersection theory on moduli spaces of abelian differentials*, Invent. Math. 222 (2020) 283–373 MR
- [6] **D Chen, M Möller, D Zagier**, *Quasimodularity and large genus limits of Siegel–Veech constants*, J. Amer. Math. Soc. 31 (2018) 1059–1163 MR

- [7] **V Delecroix, E Goujard, P Zograf, A Zorich**, *Masur–Veech volumes, frequencies of simple closed geodesics and intersection numbers of moduli spaces of curves*, preprint (2019) arXiv To appear in *Duke Math. J.*
- [8] **F Diamond, J Shurman**, *A first course in modular forms*, Grad. Texts in Math. 228, Springer (2005) MR
- [9] **R Dijkgraaf**, *Mirror symmetry and elliptic curves*, from “The moduli space of curves” (R Dijkgraaf, C Faber, G van der Geer, editors), Progr. Math. 129, Birkhäuser, Boston (1995) 149–163 MR
- [10] **P Engel**, *Hurwitz theory of elliptic orbifolds, I*, preprint (2017) arXiv To appear in *Geom. Topol.*
- [11] **A Eskin, M Kontsevich, A Zorich**, *Sum of Lyapunov exponents of the Hodge bundle with respect to the Teichmüller geodesic flow*, Publ. Math. Inst. Hautes Études Sci. 120 (2014) 207–333 MR
- [12] **A Eskin, H Masur, A Zorich**, *Moduli spaces of abelian differentials: the principal boundary, counting problems, and the Siegel–Veech constants*, Publ. Math. Inst. Hautes Études Sci. 97 (2003) 61–179 MR
- [13] **A Eskin, A Okounkov**, *Asymptotics of numbers of branched coverings of a torus and volumes of moduli spaces of holomorphic differentials*, Invent. Math. 145 (2001) 59–103 MR
- [14] **A Eskin, A Okounkov**, *Pillowcases and quasimodular forms*, from “Algebraic geometry and number theory” (V Ginzburg, editor), Progr. Math. 253, Birkhäuser, Boston (2006) 1–25 MR
- [15] **E Goujard**, *Siegel–Veech constants for strata of moduli spaces of quadratic differentials*, Geom. Funct. Anal. 25 (2015) 1440–1492 MR
- [16] **E Goujard**, *Volumes of strata of moduli spaces of quadratic differentials: getting explicit values*, Ann. Inst. Fourier (Grenoble) 66 (2016) 2203–2251 MR
- [17] **E Goujard, M Möller**, *Counting Feynman-like graphs: quasimodularity and Siegel–Veech weight*, J. Eur. Math. Soc. 22 (2020) 365–412 MR
- [18] **V G Kac**, *Infinite-dimensional Lie algebras*, 3rd edition, Cambridge Univ. Press (1990) MR
- [19] **M Kaneko, D Zagier**, *A generalized Jacobi theta function and quasimodular forms*, from “The moduli space of curves” (R Dijkgraaf, C Faber, G van der Geer, editors), Progr. Math. 129, Birkhäuser, Boston (1995) 165–172 MR
- [20] **S Kerov, G Olshanski**, *Polynomial functions on the set of Young diagrams*, C. R. Acad. Sci. Paris Sér. I Math. 319 (1994) 121–126 MR
- [21] **H Masur, A Zorich**, *Multiple saddle connections on flat surfaces and the principal boundary of the moduli spaces of quadratic differentials*, Geom. Funct. Anal. 18 (2008) 919–987 MR

- [22] **A Okounkov, R Pandharipande**, *Gromov–Witten theory, Hurwitz theory, and completed cycles*, Ann. of Math. 163 (2006) 517–560 MR
- [23] **R Ríos-Zertuche**, *An introduction to the half-infinite wedge*, from “Mexican mathematicians abroad: recent contributions” (N Bárcenas, F Galaz-García, M Moreno Rocha, editors), Contemp. Math. 657, Amer. Math. Soc., Providence, RI (2016) 197–237 MR
- [24] **A Sauvaget**, *Volumes and Siegel–Veech constants of $\mathcal{H}(2G - 2)$ and Hodge integrals*, Geom. Funct. Anal. 28 (2018) 1756–1779 MR
- [25] **J-P Serre**, *Topics in Galois theory*, Res. Notes in Math. 1, Jones & Bartlett, Boston (1992) MR
- [26] **S Shadrin, L Spitz, D Zvonkine**, *On double Hurwitz numbers with completed cycles*, J. Lond. Math. Soc. 86 (2012) 407–432 MR
- [27] **A Zorich**, *Flat surfaces*, from “Frontiers in number theory, physics, and geometry, I” (P Cartier, B Julia, P Moussa, P Vanhove, editors), Springer (2006) 437–583 MR

*Institut de Mathématiques de Bordeaux, Université de Bordeaux
Talence, France*

*Institut für Mathematik, Goethe-Universität Frankfurt
Frankfurt am Main, Germany*

elise.goujard@math.u-bordeaux.fr, moeller@math.uni-frankfurt.de

Received: 17 October 2018 Revised: 29 August 2019