

Topological spines of 4–manifolds

HEE JUNG KIM

DANIEL RUBERMAN

We show that infinitely many of the simply connected 4–manifolds constructed by Levine and Lidman that do not admit PL spines actually admit topological spines.

57N13; 57Q40, 57R67

1 Introduction

A remarkable recent paper of Levine and Lidman [21] gives examples X of smooth (equivalently, PL) simply connected 4–manifolds, homotopy equivalent to a 2–sphere, which do not admit PL spines. In other words, there is no PL (not necessarily locally flat) embedded sphere S in X that is a strong deformation retract of X . (See Cappell and Shaneson [6] and Matsumoto [24; 25] for results in higher dimensions and about spines for nonsimply connected manifolds.) In this note we show that an infinite family of the Levine–Lidman examples — those whose intersection form is $\langle 4 \rangle$ — admit a (tame) *topological spine*; this is a *locally* PL sphere that is a strong deformation retract of X .

Finding a spine for a compact manifold (especially in codimension 2) is typically phrased as a problem of homology surgery (see Cappell and Shaneson [5]), as one is looking for a suitable complement for a regular neighborhood of the spine. The complement would be a homology cobordism between the boundary of this regular neighborhood and the boundary of X . Indeed, the argument of Levine and Lidman is to find a 4–manifold X , homotopy equivalent to a 2–sphere, such that ∂X is not smoothly homology cobordant to integral surgery on any knot in the 3–sphere. In contrast, we show that for a family of their examples, ∂X is *topologically* homology cobordant to integral surgery on a knot in the 3–sphere. With some additional information about the fundamental group of the homology cobordism, this is sufficient to build the spine.

To state our main theorem, we use the notation of [21], where $\{W_p \mid p \in \mathbb{Z}\}$ denotes a family of oriented smooth 4–manifolds, each homotopy equivalent to a sphere, where a generator of $H_2(W_p)$ has self-intersection 4. For $p \notin \{-2, -1, 0\}$, the manifold W_p has no PL spine [21, Proposition 3.3].

Theorem 1.1 *For each p , the manifold W_p contains a tame topological spine.*

The paper [21] also includes examples of spineless 4-manifolds whose intersection forms are $\langle \pm k^2 \rangle$ for any nonzero k . It is natural to ask if these manifolds have topological spines; at present our method does not seem to extend to values of k greater than 2. We will comment on this further at the end of the paper. We remark that a recent paper of Hayden and Piccirillo [17] contains examples of smooth/PL manifolds homotopy equivalent to a 2-sphere of arbitrary self-intersection with no PL spine but which contain a topological spine.

Acknowledgements Thanks to Adam Levine and Tye Lidman for interesting correspondence, and to Patrick Orson for a useful discussion on some fine points of surgery theory. Ruberman was partially supported by NSF Grant DMS-1811111.

2 Preliminaries

First, we give the definition of a topological spine, following [26].


Definition 2.1 A spine of a compact topological manifold W with nonempty boundary is a tamely embedded complex $S \subset W$ such that S is a strong deformation retract of W and $S \hookrightarrow W$ is a simple homotopy equivalence. A PL spine in a PL manifold W is a spine that is a subcomplex.

In this paper, “spine” will always mean the topological version, unless specified that it is to be PL. We will be looking for a rather simple sort of spine in a manifold W homotopy equivalent to a 2-sphere, whose intersection form is $\langle n \rangle$ for some integer n . The complex S will be a 2-sphere that is locally flat, except near a single point, where it is a cone on a knot K . Such an S has a neighborhood $N(S)$ homeomorphic to $X_4(K)$, where $X_n(K)$ denotes the 4-manifold obtained by attaching an n -framed 2-handle to the 4-ball along K . We use the standard notation $S_n^3(K)$ for $\partial X_n(K)$.

We briefly recall the construction of Levine and Lidman. For any integer m , let Q_m denote the orientable (as a manifold) circle bundle over $\mathbb{R}P^2$ with twisted Euler class m . We assume throughout that $m = -4p - 3$ for an integer p ; note that m being odd implies that $H_1(Q_m) \cong \mathbb{Z}_4$. The fundamental group G_m of Q_m has order $4 \cdot |m|$ and in fact is a semidirect product

$$(1) \quad 1 \rightarrow \mathbb{Z}_{|m|} \rightarrow G_m \rightarrow \mathbb{Z}_4 \rightarrow 1$$

with a generator of \mathbb{Z}_4 acting on the subgroup $\mathbb{Z}_{|m|}$ by multiplication by -1 .

Figure 1: Circle bundle Q_{-4p-3} .

A number of surgery diagrams for Q_m are given in [21] to which we add (the well-known) variation in Figure 1; the box represents p full twists.

Note that for $p = 0, -1$ the component labeled J can be blown down to yield that Q_m is surgery on a knot in the 3-sphere. (It is not clear what happens for $p = -2$.) More generally, Q_m is $+4$ framed surgery on a knot in Y_p drawn in [21, Figure 2]. The manifold W_p of Levine–Lidman has boundary $M_p = Q_m \# -Y_p$, where Y_p is a certain Brieskorn homology sphere. The homology sphere summand is important to the construction of W_p but will not play much of a role for us.

Remark 2.2 We follow the convention of [21] by indexing some objects by p and others by m .

2.1 Twisted linking numbers

Suppose that M is an oriented 3-manifold, and that $p: \tilde{M} \rightarrow M$ is a regular covering space with covering group π . Suppose further that $H_1(\tilde{M}) = 0$. Suppose that L_1 and L_2 are nullhomotopic knots in M . Then we can consider linking numbers $\text{lk}_{\tilde{M}}$ between lifts of L_1 and L_2 . Fixing a particular lift \tilde{L}_1 and \tilde{L}_2 of each component, we can form a generating function

$$\tilde{\text{lk}}(L_1, L_2) = \sum_{g \in \pi} \text{lk}_{\tilde{M}}(\tilde{L}_1, \tilde{L}_2 \cdot g)g.$$

Note that by invariance of linking numbers under (orientation-preserving) homeomorphisms, the function $\widetilde{\text{lk}}$ determines linking numbers between arbitrary lifts of L_1 and L_2 . More precisely, $\text{lk}_{\widetilde{M}}(\widetilde{L}_1 \cdot h, \widetilde{L}_2 \cdot g) = \text{lk}_{\widetilde{M}}(\widetilde{L}_1, \widetilde{L}_2 \cdot gh^{-1})$.

If (L, ϕ) is a framed knot in M , then we define the self-linking $\widetilde{\text{slk}}(L; \phi)$ by choosing $L_1 = L$ and L_2 to be the push-off L^ϕ defined by the framing. The constant term in this generating function is the framing of \widetilde{L} . More precisely, the choice of lift \widetilde{L} determines a lift \widetilde{L}^ϕ , which is the push-off of \widetilde{L} by the lifted framing. Since L and \widetilde{L} are nullhomologous, the framings can be characterized by integers n and n' . Suppose that the covering group π is finite, of order d , and consider a chain \widetilde{C} in \widetilde{M} with boundary \widetilde{L} , projecting to a null-homology C for L . Then each intersection point of L^ϕ with C lifts to d intersection points in \widetilde{M} . Taking signs into account, we see that

$$dn = \sum_{g, h \in \pi} \text{lk}_{\widetilde{M}}(\widetilde{L} \cdot g, \widetilde{L}^\phi \cdot h) = d \left(\sum_{g \in \pi} \text{lk}_{\widetilde{M}}(\widetilde{L}, \widetilde{L}^\phi \cdot g) \right)$$

and so

$$(2) \quad n = n' + \left(\sum_{1 \neq g \in \pi} \text{lk}_{\widetilde{M}}(\widetilde{L}, \widetilde{L}^\phi \cdot g) \right) = n' + \left(\sum_{1 \neq g \in \pi} \text{lk}_{\widetilde{M}}(\widetilde{L}, \widetilde{L} \cdot g) \right).$$

Such linking numbers have been considered by many authors [9; 27; 28] mostly in the setting when π is a cyclic group, where there are good formulas (originating in Seifert's work [30] on branched covers) in terms of Seifert matrices. For more general covering groups π , we can also compute $\widetilde{\text{lk}}(L_1, L_2)$ in terms of data in M . This procedure seems to be well-known [15], so we will just summarize it.

Suppose that the covering $\widetilde{M} \rightarrow M$ corresponds to a homomorphism

$$r: \pi = \pi_1(M, x_0) \rightarrow \pi,$$

where x_0 is a basepoint. Note that the choice of lifts \widetilde{L}_i is equivalent to choosing a homotopy class β_i of arcs (modulo multiplication by elements of $\ker(r)$) from $x_0 \in M$ to each L_i . Write c_i for the endpoint of β_i on L_i . For each i , choose an immersion $\delta_i: D^2 \rightarrow M$ whose restriction to the boundary is L_i . Then to any intersection point x of L_1 and $\Delta_2 = \delta_2(D^2)$ we assign an element $g(x) \in \pi$ as follows. Note that Δ_2 is based, using β_2 . Then we define

$$g(x) = \beta_1 * \alpha_1 * \alpha_2^{-1} * \beta_2^{-1},$$

where α_1 is a path in L_1 from c_1 to x , and α_2 is the projection of a path in D^2 to a path in Δ_2 from c_2 to x .

Then by [15, Theorem 4.10] we have

$$(3) \quad \widetilde{\text{lk}}(L_1, L_2) = \sum_{x \in L_1 \cap \Delta_2} \text{sign}_x(L_1, \Delta_2) r(g(x)),$$

where $\text{sign}_x(L_1, \Delta_2)$ is the local intersection number at x .

From a minor variation on equation (3) we get the following recipe for varying the twisted linking number.

Lemma 2.3 *Let (L_1, L_2) be a link with both components nullhomotopic. Then for any element $q = \sum a_g g \in \mathbb{Z}[\pi]$, there is a knot $L_1(q)$ isotopic to L_1 such that $\widetilde{\text{lk}}(L_1(q), L_2) = \widetilde{\text{lk}}(L_1, L_2) + q$.*

Proof It clearly suffices to prove this for $q = \pm g$; the procedure is shown in Figure 2. In summary, send out a feeler from L_1 to L_2 following along the loop g (and the base paths for L_1 and L_2) and put a clasp, whose sign is determined by the sign in q . \square




Figure 2: Changing the twisted linking number.

If L is a knot, then we can insert a self-clasp of either sign along a loop $g \neq 1$ to get a new knot $L(g)$ such that the linking number $\text{lk}_{\widetilde{M}}(\widetilde{L}(g), \widetilde{L}(g) \cdot g)$ changes by ± 1 . If L has a framing ϕ , specified by a number n and we give $L(g)$ the same framing n , then by (2) the framing of $\widetilde{L}(g)$ has changed by ± 1 . More generally, we can do this for any finite collection of elements $g \neq 1$. Making use of (2) to compute the framing upstairs, we get the following.

Lemma 2.4 *For any nullhomotopic framed knot (L, ϕ) with framing n , and any*

$$q = \sum_{1 \neq g \in \pi} a_g g \in \mathbb{Z}[\pi],$$

there is a nullhomotopic framed knot $(L(q), \phi(q))$ with framing n such that

$$\widetilde{\text{slk}}(\widetilde{L}(q), \phi(q)) = \widetilde{\text{slk}}(\widetilde{L}, \phi) + \sum_{1 \neq g \in \pi} a_g.$$

2.2 Outline of the proof

The main step in the proof is to construct a knot k_p in S^3 so that $S_4^3(k_p)$ is topologically homology cobordant to $Q_m \# Y_p$. The homology cobordism will be glued onto $X_4(k_p)$ to give a topological 4-manifold with a spine; we will argue that the resulting manifold is homeomorphic to W_p , and hence W_p will have a topological spine. There are several remarks before we embark on the proof. The first is that in the topological category, Y_p is homology cobordant to S^3 , so we will instead find a homology cobordism between $S_4^3(k_p)$ and Q_m . At the end, we will have to reassemble everything to get the correct boundary for the manifold we construct.

A more substantial remark is that constructing a homology cobordism is an instance of *homology surgery* [5], since one is trying to create a manifold with a given homological rather than homotopy type. It is a standard observation, based on the Casson–Gordon invariants [7; 8], that homology surgery does not work well in dimension 4, even in the topological category; see for example [1] or [23]. So we will look to use something more like the usual surgery theory, which will produce a stronger conclusion. In particular, we will look for a homology cobordism V_m between $S_4^3(k_p)$ and Q_m with fundamental group G_m such that the induced cover of $S_4^3(k_p)$ is a homology sphere; in other words, V_m will be a $\mathbb{Z}[G_m]$ –homology cobordism. This is in a sense similar to applications [12] of surgery to knot slicing problems, where a homology surgery problem is solved by a judicious choice of fundamental group, so that Wall’s surgery theory can be used directly.

The construction of V_m has the following steps.

- (1) Find a knot K_p such that there is a map $f: S_4^3(K_p) \rightarrow Q_m$ that is a $\mathbb{Z}[G_m]$ –homology equivalence.
- (2) Construct a normal cobordism between f and the identity map from Q_m to itself.
- (3) Modify K_p , producing k_p and a normal cobordism such that the surgery obstruction in $L(\mathbb{Z}[G_m])$ is trivial.

The knots K_p in this outline are not the same as the knots K_p in [21], which play no role in the current work.

The second step turns out to be a straightforward deduction from known calculations, but the other two require a little more work. For instance, when $p = 1$, one can find a knot L (in fact the knot 12n0749) whose knot group maps to G_{-7} such that the induced 28-fold cover of $S^3_4(L)$ is a homology sphere. However, there is no degree-one map inducing this homology equivalence and in fact we couldn't find any knot in the tables (through 13 crossings, after which we gave up) with the desired properties for K_1 . A general construction for K_p is given in the next section.

3 Degree-one maps

In this section we carry out the first step in the construction outlined above. Unless otherwise specified, we write $m = -4p - 3$ for an arbitrary integer p . Recall [4] that a normal map $f: M \rightarrow X$ from a manifold M to a Poincaré complex X is a degree-one map with a choice of bundle map covering f from the stable normal bundle (in the appropriate category) of M to a lift of the Spivak normal fibration of X . We will deal with only the special case when both X and M are manifolds with trivial stable normal bundles, so any degree-one map is covered (not uniquely) by such a bundle map and can be considered as a normal map.

Proposition 3.1 *There is a knot K_p and a degree-one normal map $f: S^3_4(K_p) \rightarrow Q_m$ that is a $\mathbb{Z}[G_m]$ -homology equivalence.*

We will find the following terminology to be useful.

Definition 3.2 Let (J, K) be a link in S^3 , and let $\vec{\eta} \subset S^3 - (J \cup K)$ be a link.

- (a) $\vec{\eta}$ is a *mod K unlink* if its components bound disjoint discs in $S^3 - J$.
- (b) $\vec{\eta}$ is a *mod K unknotting link* if the linking number of every component η_i with J is 0 and if doing a series of right or left twists along the components of $\vec{\eta}$ unknots J .

The mod K terminology is intended to suggest that $\vec{\eta}$ has the desired property after we fill in K .

A very useful technique for constructing maps of degree one is the following, from [2, Proposition 3.2]; see also [13] for a converse result. Suppose M is a 3-manifold, and C a nullhomotopic curve. Then the result of Dehn surgery on C with any slope

maps to M with degree one. The proof of this fact in [2] is straightforward, but we offer the following 4-dimensional version of the proof, which seems perhaps a bit easier.

Lemma 3.3 *Let M be a 3-manifold, and let $\vec{C} = \{C_1, \dots, C_n\}$ be a framed link with all components nullhomotopic. If N is the result of surgery on \vec{C} (with the given framings) then there is a degree-one map from N to M .*

Note that this implies the stronger statement where surgery is replaced by Dehn surgery (or more properly, Dehn filling). The standard argument [29, Chapter 9.H] that a Dehn surgery can be done as a sequence of “honest” surgeries applies in an arbitrary 3-manifold. The auxiliary curves that are added in this process are nullhomotopic, and so one can apply Lemma 3.3.

Proof Let X be the result of adding 2-handles to $M \times I$ along $\vec{C} \times \{1\}$ with the given framings, so that $\partial X \cong N \cup M$. Since each component of \vec{C} is nullhomotopic, the identity map of M extends to a retraction from X to $M \times \{0\}$. Since N is homologous to M in X , the restriction of this map to N has degree one. \square

3.1 Proof of Proposition 3.1

We will assume $p \geq 0$ for simplicity; the case of negative p is similar with slightly different pictures and notation.

To apply this consider the curves η_1, \dots, η_p in Figure 3, on the right. We would like to do -1 -framed surgery along these curves to create a degree-one map; the problem is that they are not trivial in $\pi_1(Q_m)$. The curve η_0 will be used below in the proof of Lemma 3.4 to modify η_1, \dots, η_p ; the curve z will be used afterwards to change the fundamental group.

Note that $\vec{\eta} = \{\eta_1, \dots, \eta_p\}$ is a mod K unknotting link for J and that η_0 is an unknot mod K . It follows easily that there is a band sum of $\vec{\eta}$ with p copies of η_0 to make a link $\vec{\gamma} = \{\gamma_1, \dots, \gamma_p\}$ that is also a mod K unknotting link for J and such that the homotopy class of γ_i is $\eta_i^{-1} * \eta_0$. The calculation in the proof of Lemma 3.4 shows that $\eta_i^{-1} * \eta_0$ is trivial in $\pi_1(Q_m)$ for all i .

Note that we can find other mod K unknotting links for J by taking the band sum of components of $\vec{\gamma}$ with any mod K unknot; we need to know which homotopy classes contain such knots.




Figure 3: Surgery curves in Q_m : $p = 0$, left, and $p > 0$, right.

Lemma 3.4 Every element of $\pi_1(Q_m)$ is represented by a mod K unknot.

Proof We claim that $\pi_1(Q_m)$ is generated by the curves η_0 (with the indicated basepoint) and z . Both of these are mod K unknots, so any product of them is represented by a mod K unknot. Recalling that $m = -4p - 3$, it is known that Q_m can be represented by a surgery along (-1) -surgery on the $(2, 2p + 1)$ torus knot J and 0 -surgery on a curve grabbing twice K as in Figure 3 on the right, which is drawn for positive p . Let g_p be the fundamental group of the complement of the link (J, K) in S^3 . For $p = 0$, a Wirtinger presentation shows that g_0 is generated by the curves labeled x, y, z in Figure 3 on the left, with relations $y^{-1}x(yzy^{-1}z^{-1}) = 1$, $yzx^{-1}z^{-1} = 1$, and $yz^{-1}y^{-1}x^{-1}zx = 1$. Note that the complement of the link (J, K) for all the p is the same as for $p = 0$, since we do a twist around K , where the twist diffeomorphism from the complement of the link for $p = 0$ to that for arbitrary p takes $x \mapsto x$, $y \mapsto y$, and $z \mapsto (xy)^p z$. So, the fundamental group g_p is generated

by curves x, y, z_p , where z_p denotes $(xy)^p z$, with relations $y^{-1}x(yz_p y^{-1}z_p^{-1}) = 1$, $yz_p x^{-1}z_p^{-1} = 1$, and $yz_p^{-1}y^{-1}x^{-1}z_p x = 1$.

The fundamental group $\pi_1(Q_m)$ is obtained from g_p simply by adding the relations induced from surgery curves S_0, S_{-1} , which are parallel to K and J , with framings 0 and -1 , respectively. From Figure 3, we can read off a presentation for each surgery curve; starting near the dot on curve of J we get $1 = zy^{p+1}zy^{-2p-2}y^{-p} = zy^{p+1}zy^{-3p-2}$ for S_{-1} and obviously, the 0-surgery S_0 gives $xy = 1$.

So, we get a presentation for $\pi_1(Q_m)$ with generators x, y, z_p and relations

$$(4) \quad y^{-1}x(yz_p y^{-1}z_p^{-1}) = 1,$$

$$(5) \quad yz_p x^{-1}z_p^{-1} = 1,$$

$$(6) \quad yz_p^{-1}y^{-1}x^{-1}z_p x = 1,$$

$$(7) \quad xy = 1,$$

$$(8) \quad zy^{p+1}zy^{-3p-2} = 1.$$

Using (7), we get rid of the generator x and the relation (6) so that the presentation has generators y, z and relations

$$(9) \quad y^{-2}(yz y^{-1}z^{-1}) = 1, \quad yzyz^{-1} = 1 \quad \text{and} \quad zy^{p+1}zy^{-3p-2} = 1.$$

The first two relations in (9) are equivalent to $z^{-1}yz = y^{-1}$, which implies that $z^{-1}y^{p+1}z = y^{-p-1}$, so $y^{p+1}z = zy^{-p-1}$. Then the third relation is

$$1 = zy^{p+1}zy^{-3p-2} = z(zy^{-p-1})y^{-3p-2} = z^2y^{-4p-3}.$$

So, this reduction gives a presentation for $\pi_1(Q_m)$ as follows:

$$(10) \quad \langle y, z \mid z^2 = y^{4p+3}, z^{-1}yz = y^{-1} \rangle.$$

From Figure 3, we read off $\eta_0 = yzy^{-1}z^{-1}$ and $\eta_1 = y^2$, so the first relation in (9) implies that $1 = y^{-2}(yz y^{-1}z^{-1}) = \eta_1^{-1} * \eta_0 = y^{-2}\eta_0 = 1$, so $y^2 = \eta_0$. This makes the relation in (10) $z^2 = y^{4p+3} = y^{2(2p+1)} \cdot y = \eta_0^{2p+1} \cdot y$, so $\eta_0^{-(2p+1)}z^2 = y$, which shows that $\pi_1(Q_m)$ is generated by η_0 and z . This completes the proof. \square

Returning to the proof of Proposition 3.1, note that relation (7) implies that the curves η_1, \dots, η_p all represent the class $y^2 \in G_m = \pi_1(Q_m)$, as does η_0 . Let N_m be the result of (-1) framed surgery on each component of $\vec{\gamma}$. Then by the discussion above, there is a degree-one map from N_m to Q_m . By construction, after surgery on $\vec{\gamma}$,

we can blow down $\vec{\gamma}$ and J in succession. The result is $+4$ surgery on some new knot \hat{K}_m in the 3-sphere, which can in principle be drawn. It will be a little messy, and is about to get a lot messier so we do not draw it.

At this point we have achieved most of step (1), since $N_m = S_4^3(\hat{K}_m)$ maps with degree one to Q_m , and hence its fundamental group maps surjectively onto G_m and induces an isomorphism on integral homology. However, there is no reason that it should induce an isomorphism in twisted homology. For example, a direct calculation using SnapPy [10] and GAP [14] reveals that for $m = -7$, the homology of the induced 28-fold cover \tilde{N}_{-7} is quite large. We show how to modify $\vec{\gamma}$ (and hence \hat{K}_m) so as to kill the homology of \tilde{N}_m .

By construction, \tilde{N}_m is surgery on the preimage of γ in the universal cover of Q_m , which is the 3-sphere. As such, its homology is presented by the $p|G_m| \times p|G_m|$ matrix of linking numbers $\mathcal{L} = \text{lk}_{S^3}(\tilde{\gamma}_i \cdot g, \tilde{\gamma}_j' \cdot h)$, where γ_j' is the (-1) -pushoff of γ_j . The framing on $\tilde{\gamma}_j$ is determined by (2). Using Lemma 2.4 we can modify γ by self-clasping moves along elements of $\pi = G_m$ so that the first row of \mathcal{L} becomes the vector $(n', 0, \dots, 0)$. Since the framing on γ was -1 , (2) says that $n' = -1$ as well.

By Lemma 3.4, every element of G_m is represented by a mod K unknot, so we can do the self-clasping moves so that the modified version of $\vec{\gamma}$ remains a mod K unknotting link for J . It follows that the result of blowing down the modified $\vec{\gamma}$, followed by blowing down J , turns K into a knot K_p with all the desired properties.

Finally, since $S_4^3(K_p)$ and Q_m are both parallelizable, any degree-one map is covered by a map of stable normal bundles, and hence is a normal map. \square

Remark 3.5 This argument for altering the homology of a covering space was inspired by Jerry Levine's surgical construction of a knot with a given Alexander polynomial [22].

4 The surgery problem

First we show that the map constructed in the previous section gives rise to a normal map to $Q_m \times I$.

Proposition 4.1 Any degree-one normal map $f: Y \rightarrow Q_m$ that is a $\mathbb{Z}[G_m]$ -homology equivalence is normally cobordant to the identity map of Q_m .

Proof We make use of a version [20] of the surgery exact sequence for 3-manifolds, in which the usual structure set $\mathcal{S}(M)$ for homotopy equivalences to a manifold M^3 is replaced by the set of $\mathbb{Z}[\pi_1(M)]$ -homology equivalences. We are in a setting where we don't know (or care) if the homology equivalence is simple. Hence the relevant surgery groups are L^h groups, and the equivalence relation for the structure set is up to h -cobordism instead of homeomorphism or diffeomorphism. For our purposes, it doesn't matter whether we are working in the topological or smooth category; we will stick to TOP for convenience. Writing $\bar{\mathcal{S}}(M)$ for this modified structure set, there is an exact sequence

$$(11) \quad \bar{\mathcal{S}}(M) \xrightarrow{N} [M, G/\text{TOP}] \xrightarrow{\theta} L_3^h(\mathbb{Z}[\pi_1(M)]).$$

The first map is the normal invariant, while the second is the surgery obstruction. The maps in this sequence have been studied in some detail in [18], and we could base a proof of Proposition 4.1 on Theorem 2 of that paper. However, we give an alternate and perhaps slightly more direct argument.

By the well-known calculation [19] of the low-dimensional homotopy groups of G/TOP , we get that

$$[Q_m, G/\text{TOP}] = [Q_m, K(\mathbb{Z}_2, 1)] = H^1(Q_m; \mathbb{Z}_2) = \mathbb{Z}_2.$$

The map to \mathbb{Z}_2 is a codimension-one Arf invariant, and we claim that it vanishes. Because the projection $G_m \rightarrow \mathbb{Z}_4$ splits (G_m being a semidirect product as in (1)) the induced map $L_3(\mathbb{Z}[G_m]) \rightarrow L_3(\mathbb{Z}[\mathbb{Z}_4])$ is a split surjection. It is known [16] that $L_3(\mathbb{Z}[\mathbb{Z}_4]) \cong \mathbb{Z}_2$, with the nontrivial element being detected by a codimension-one Arf invariant. So in this case, the surgery obstruction map θ is an injection. Since the composition $\theta \circ N$ is trivial, and our given f is a $\mathbb{Z}[G_m]$ -homology equivalence, its normal invariant must be trivial. \square

4.1 Killing the surgery obstruction

At this point, we have a normal map $F: Z^4 \rightarrow Q_m \times I$, where

$$\partial Z = S_4^3(K_p) \sqcup -Q_m.$$

The restriction of F to Q_m is the identity, and the restriction to $S_4^3(K_p)$ is a $\mathbb{Z}[G_m]$ -homology equivalence. Hence there is a well-defined surgery obstruction $\theta(F) \in L_4(\mathbb{Z}[G_m])$ to doing surgery on Z to obtain a homotopy $Q_m \times I$. In principle,

the groups $L_4(\mathbb{Z}[G_m])$ are computable [32; 16] but in practice there are 2-torsion obstructions that are hard to evaluate.

Our approach is to alter the knot K_p one more time to kill the surgery obstruction. The main step is worth summarizing in a lemma; it applies both to L_4^s and L_4^h , either of which will be temporarily denoted L_4 . Recall [31, Chapter 5] that an element θ of $L_4(\mathbb{Z}[G])$ may be represented, possibly after stabilizations, by a free module A over $\mathbb{Z}[G]$ and a pair (λ, μ) . Here λ is a form on $A \times A$ that is Hermitian with respect to the involution induced by $g \rightarrow g^{-1}$ on $\mathbb{Z}[G]$ and μ is a quadratic form on A related to λ . Let B be a matrix for λ ; the image of B under the augmentation $\epsilon: \mathbb{Z}[G] \rightarrow \mathbb{Z}$ has a signature which we denote by $\text{sign}(\theta)$.

Lemma 4.2 *Suppose that K is a knot in S^3 , and that there is a 3-manifold Y along with a $\mathbb{Z}[G]$ -homology equivalence $f: S_n^3(K) \rightarrow Y$, where $G = \pi_1(Y)$. Then there is a knot K' and a topological normal cobordism (Z, F) from f to a $\mathbb{Z}[G]$ -homology equivalence $f': S_n^3(K') \rightarrow Y$ with surgery obstruction $\theta(Z, F) = \theta$. If $\text{sign}(\theta) \equiv 0 \pmod{16}$ then there is a smooth normal cobordism.*

Proof We remark first that any finite collection of elements of $\pi_1(S_n^3(K))$ may be represented by an unlink. For one can start with any link L in $S^3 - K$ representing those group elements, and then do crossing changes in S^3 , missing K , to make L into an unlink in $S^3 - K$.

Note that $\epsilon(\theta)$ is an even unimodular form over \mathbb{Z} . This follows from Theorem 5.2(iii) in [31], which depends on the fact that F is a normal map. A more direct argument is that since F is a normal map, Z is in fact a spin manifold, and hence its intersection form $\epsilon(\theta)$ is even.

By the previous remarks the signature of $\epsilon(\theta)$ is divisible by 8. Since there exist closed smooth spin manifolds of signature 16 and closed topological spin manifolds of index 8, we can change $\text{sign}(\theta)$ by multiples of 8 (resp. 16) in the topological (resp. smooth) categories without affecting the boundary. So in either category, our assumption on the signature means that we may assume that $\text{sign}(\theta) = 0$. Since $L_4(\mathbb{Z})$ is detected by the signature, this means that $\epsilon(\theta)$ is trivial in $L_4(\mathbb{Z})$.

As described above, λ is encoded as a matrix with respect to a basis for a free $\mathbb{Z}[G]$ -module A . Note that any integral change of basis for

$$A \otimes_{\mathbb{Z}[G]} \mathbb{Z}$$

lifts to a change of basis for A . Since $\epsilon(\theta) = 0 \in L_4(\mathbb{Z})$, it follows that there is a basis $\{e_i \mid i = 1, \dots, 2m\}$ such that $\epsilon(\lambda)$ is represented by a sum of hyperbolic forms

$$H = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

Let B be the matrix for λ with respect to this basis.

Now we follow the proof of Wall's realization theorem [31, Theorem 5.8], adding handles to $S_n^3(K) \times I$ along a framed link in $S_n^3(K) \times \{1\}$ to create the normal cobordism Z . More precisely, start with a 0-framed unlink \vec{C} in a ball $S_n^3(K)$ whose components C_i correspond to the basis vectors e_i . As in Lemma 2.3, for $i \neq j$ we introduce clasps between the circles C_i and C_j so that the twisted linking number (measured in $\mathbb{Z}[G]$) between those circles becomes B_{ij} . Similarly, introduce self-clasps into C_i along elements of $\pi_1(S_n^3(K))$ so that the self-intersection form will be given by μ . When handles are added along the components of the resulting 0-framed link, the intersection form of the resulting 4-manifold will be given by λ .

It follows that the upper boundary of Z , which results from doing surgery along \vec{C} , is n -framed surgery on the knot K' obtained by canceling all of the handles (viewed as attached to C in S^3). \square

Remark 4.3 In the case of interest, we let $Y = Q_m$, and apply Proposition 3.1 to obtain the knot K , and then Lemma 4.2 to change it into K' . In principle, one could trace through their proofs to determine the knots K and K' , respectively. In the special case $p = 1$, the group $L_4(\mathbb{Z}[G_{-7}]) \cong \mathbb{Z}^6$ is detected by multisignatures, which are computable. So perhaps in this case, the knot K' could be drawn.

5 Proof of Theorem 1.1

Putting together Propositions 3.1 and 4.1, we find a knot K_p , a $\mathbb{Z}[G_m]$ -homology equivalence $f: S_4^3(K_p) \rightarrow Q_m$, and a normal cobordism (Z, F) from f to the identity of Q_m . Making use of Lemma 4.2, we find another knot k_p , and a normal cobordism (Z', F') (possibly nonsmoothable) from f to a $\mathbb{Z}[G_m]$ -homology equivalence $f': S_4^3(k_p) \rightarrow Q_m$ such that the surgery obstruction $\theta(F')$ is the negative of $\theta(F)$. Gluing these two together gives a normal cobordism from f' to the identity of Q_m with trivial surgery obstruction.

Since $\pi_1(Q_m)$ is finite, Freedman's theorem [11] says that surgery on

$$Z \cup_{S^3_4(K_p)} Z'$$

can be done to get a homotopy equivalence from a new 4-manifold V_p to $Q_m \times I$.

Now recall that Levine–Lidman's manifold is denoted by W_p , and its boundary is $Q_m \# -Y_p$, where Y_p is a homology sphere. In the topological category, Y_p bounds a contractible manifold U , and we take the boundary connected sum of V_p with manifold U to get a homology cobordism between $S^3_4(k_p)$ and ∂W_p . By the d -invariant calculation in [21] this homology cobordism cannot be smooth, unless $p \in \{0, -1, -2\}$. Let

$$W'_p = X_4(k_p) \cup_{S^3_4(k_p)} (V_p \natural U),$$

and note that $\partial W'_p = Q_m \# -Y_p$.

By construction, W'_p has a topological spine, so Theorem 1.1 will be proved if we can identify it with W_p .

Claim W'_p is homeomorphic to W_p .

To prove the claim, we use the work of Boyer [3], which extends Freedman's classification of simply connected topological manifolds to the setting of manifolds with boundary.

By construction, the inclusion of $S^3_4(k_p)$ into V_p induces a surjection on the fundamental group. Since $X_4(k_p)$ is simply connected, it follows that W'_p is simply connected as well. Note further that W'_p is spin, since its intersection form is even. The classification for general boundaries is rather complicated, but in the present case the answer turns out to be quite simple. For an oriented 3-manifold M , Boyer constructs a map $c^t_L(M)$ from the set of homeomorphism classes of simply connected 4-manifolds with boundary M and intersection form L to a certain coset space $B^t_L(M)$ of the automorphisms of the linking form of M . When L is even, and M is a rational homology sphere, this map is an injection.

The fact that $H_1(Q_m \# -Y_p) \cong \mathbb{Z}_4$ implies [3, Proposition 0.6] that $B^t_L(Q_m \# -Y_p)$ has exactly one element. Hence W'_p is homeomorphic to W_p and the proof of Theorem 1.1 is complete. \square

Remark 5.1 It is reasonable to wonder if the other manifolds $W_{k,m}$ constructed in [21] have topological spines. These manifolds have intersection form $\langle k^2 \rangle$ for $k > 2$

and $|m| \gg 0$, and boundaries of the form $Q_{k,m} \# Y$. Here $Q_{k,m}$ is surgery on a link similar to the one in Figure 3 on the right, and Y is a homology sphere. So it seems likely that the portion of the argument that produces a knot and a degree homology equivalence on the result of surgery on that knot to $Q_{k,m}$ will still work. However, for our method to succeed, we need an honest surgery problem whose restriction to the boundary is a homology equivalence with coefficients in $\pi_1(Q_{k,m})$, and not just a homology surgery problem. The fundamental group of $Q_{k,m}$ is large and it is not clear what should be the fundamental group of a surgery problem that might be constructed to produce a homology cobordism.

References

- [1] **S Akbulut**, *A note on homology surgery and the Casson–Gordon invariant*, Math. Proc. Cambridge Philos. Soc. 85 (1979) 335–344 MR
- [2] **M Boileau, S Wang**, *Non-zero degree maps and surface bundles over S^1* , J. Differential Geom. 43 (1996) 789–806 MR
- [3] **S Boyer**, *Simply-connected 4–manifolds with a given boundary*, Trans. Amer. Math. Soc. 298 (1986) 331–357 MR
- [4] **W Browder**, *Surgery on simply-connected manifolds*, Ergeb. Math. Grenzgeb. 65, Springer (1972) MR
- [5] **S E Cappell, J L Shaneson**, *The codimension two placement problem and homology equivalent manifolds*, Ann. of Math. 99 (1974) 277–348 MR
- [6] **S E Cappell, J Shaneson**, *Totally spineless manifolds*, Illinois J. Math. 21 (1977) 231–239 MR
- [7] **A J Casson, C M Gordon**, *On slice knots in dimension three*, from “Algebraic and geometric topology, II” (R J Milgram, editor), Proc. Sympos. Pure Math. 32, Amer. Math. Soc., Providence, RI (1978) 39–53 MR
- [8] **A J Casson, C M Gordon**, *Cobordism of classical knots*, from “À la recherche de la topologie perdue” (L Guillou, A Marin, editors), Progr. Math. 62, Birkhäuser, Boston (1986) 181–199 MR
- [9] **J C Cha, K H Ko**, *Signature invariants of covering links*, Trans. Amer. Math. Soc. 358 (2006) 3399–3412 MR
- [10] **M Culler, N M Dunfield, M Goerner, J R Weeks**, *SnapPy, a computer program for studying the topology of 3–manifolds* Available at <http://snappy.computop.org>
- [11] **M H Freedman, F Quinn**, *Topology of 4–manifolds*, Princeton Math. Ser. 39, Princeton Univ. Press (1990) MR

- [12] **S Friedl, P Teichner**, *New topologically slice knots*, *Geom. Topol.* 9 (2005) 2129–2158
MR Correction in 10 (2006) 2501–2504
- [13] **S Gadgil**, *Degree-one maps, surgery and four-manifolds*, *Bull. Lond. Math. Soc.* 39 (2007) 419–424 MR
- [14] **The GAP Group**, *GAP: groups, algorithms, and programming*, software (2004) Version 4.4 Available at <http://www.gap-system.org>
- [15] **D L Goldsmith**, *A linking invariant of classical link concordance*, from “Knot theory” (J-C Hausmann, editor), *Lecture Notes in Math.* 685, Springer (1978) 135–170 MR
- [16] **I Hambleton, L R Taylor**, *A guide to the calculation of the surgery obstruction groups for finite groups*, from “Surveys on surgery theory, I” (S Cappell, A Ranicki, J Rosenberg, editors), *Ann. of Math. Stud.* 145, Princeton Univ. Press (2000) 225–274 MR
- [17] **K Hayden, L Piccirillo**, *The trace embedding lemma and spinelessness*, preprint (2019) arXiv To appear in *J. Differential Geom.*
- [18] **B Jahren, S Kwasik**, *Three-dimensional surgery theory, UNil-groups and the Borel conjecture*, *Topology* 42 (2003) 1353–1369 MR
- [19] **R C Kirby, L C Siebenmann**, *On the triangulation of manifolds and the Hauptvermutung*, *Bull. Amer. Math. Soc.* 75 (1969) 742–749 MR
- [20] **R C Kirby, L R Taylor**, *A survey of 4-manifolds through the eyes of surgery*, from “Surveys on surgery theory, II” (S Cappell, A Ranicki, J Rosenberg, editors), *Ann. of Math. Stud.* 149, Princeton Univ. Press (2001) 387–421 MR
- [21] **A S Levine, T Lidman**, *Simply connected, spineless 4-manifolds*, *Forum Math. Sigma* 7 (2019) art. id. e14 MR
- [22] **J Levine**, *A characterization of knot polynomials*, *Topology* 4 (1965) 135–141 MR
- [23] **J Levine, K E Orr**, *A survey of applications of surgery to knot and link theory*, from “Surveys on surgery theory, I” (S Cappell, A Ranicki, J Rosenberg, editors), *Ann. of Math. Stud.* 145, Princeton Univ. Press (2000) 345–364 MR
- [24] **Y Matsumoto**, *A 4-manifold which admits no spine*, *Bull. Amer. Math. Soc.* 81 (1975) 467–470 MR
- [25] **Y Matsumoto**, *Wild embeddings of piecewise linear manifolds in codimension two*, from “Geometric topology” (J C Cantrell, editor), Academic, New York (1979) 393–428 MR
- [26] **E K Pedersen**, *Spines of topological manifolds*, *Comment. Math. Helv.* 50 (1975) 41–44 MR
- [27] **M Powell**, *Twisted Blanchfield pairings and symmetric chain complexes*, *Q. J. Math.* 67 (2016) 715–742 MR

- [28] **J H Przytycki, A Yasuhara**, *Linking numbers in rational homology 3–spheres, cyclic branched covers and infinite cyclic covers*, Trans. Amer. Math. Soc. 356 (2004) 3669–3685 MR
- [29] **D Rolfsen**, *Knots and links*, Math. Lect. Ser. 7, Publish or Perish, Berkeley (1976) MR
- [30] **H Seifert**, *Die Verschlingungsinvarianten der zyklischen Knotenüberlagerungen*, Abh. Math. Sem. Univ. Hamburg 11 (1935) 84–101 MR
- [31] **C T C Wall**, *Surgery on compact manifolds*, Lond. Math. Soc. Monogr. 1, Academic, London (1970) MR
- [32] **C T C Wall**, *Classification of Hermitian forms, VI: Group rings*, Ann. of Math. 103 (1976) 1–80 MR

*Department of Mathematics, Western Washington University
Bellingham, WA, United States*

*Department of Mathematics, Brandeis University
Waltham, MA, United States*

heejungorama@gmail.com, ruberman@brandeis.edu

Received: 15 May 2019 Revised: 19 December 2019