

AG  
T

*Algebraic & Geometric  
Topology*

Volume 23 (2023)

**Suspension homotopy of 6-manifolds**

RUIZHI HUANG



# Suspension homotopy of 6–manifolds

RUIZHI HUANG

For a simply connected closed orientable manifold of dimension 6, we compute its homotopy decomposition after double suspension. This allows us to determine its  $K$ – and  $KO$ –groups easily. Moreover, in a special case we refine the decomposition to show the rigidity property of the manifold after double suspension.

[55P15](#), [55P40](#), [57R19](#); [55N15](#), [55P10](#)

## 1 Introduction

Let  $M$  be a closed orientable smooth manifold of dimension  $n$ . Numerous investigations have been made in geometric topology into the diffeomorphism or homeomorphism type of  $M$  in various cases. For instance, in the general case, Wall [30; 32] studied  $(s-1)$ –connected  $2s$ –manifolds and  $(s-1)$ –connected  $(2s+1)$ –manifolds. For concrete cases with specified dimension  $n$ , Barden [1] classified simply connected 5–manifolds, and Wall [31], Jupp [18] and Zhubr [35; 36] classified simply connected 6–manifolds. More recently, Kreck and Su [20] classified certain nonsimply connected 5–manifolds, while Crowley and Nordström [11] and Kreck [19] studied the classification problem of various kinds of 7–manifolds.

In the literature mentioned, the homotopy classification of  $M$  was usually carried out as a byproduct in terms of a system of invariants. However, it is almost impossible to extract nontrivial homotopy information of  $M$  directly from the classification. On the other hand, unstable homotopy theory is a powerful tool for studying the homotopy properties of manifolds. There have been several interesting investigations recently in this direction. For instance, Beben and Theriault [6] studied the loop decompositions of  $(s-1)$ –connected  $2s$ –manifolds, while Beben and Wu [8] and Huang and Theriault [17] studied the loop decompositions of the  $(s-1)$ –connected  $(2s+1)$ –manifolds. The homotopy groups of these manifolds were also investigated by Samik Basu and Somnath

Basu [2; 3] from a different point of view. Moreover, a theoretical method of loop decomposition was developed by Beben and Theriault [7], which is quite useful for studying the homotopy of manifolds. Additionally, the homotopy type of the suspension of a connected 4–manifold was determined by So and Theriault [28].

We study the homotopy of simply connected 6–manifolds. Let  $M$  be a simply connected closed orientable 6–manifold. By Poincaré duality and the universal coefficient theorem we have

$$(1) \quad H_*(M; \mathbb{Z}) = \begin{cases} \mathbb{Z}^{\oplus d} \oplus T & \text{if } * = 2, \\ \mathbb{Z}^{\oplus 2m} \oplus T & \text{if } * = 3, \\ \mathbb{Z}^{\oplus d} & \text{if } * = 4, \\ \mathbb{Z} & \text{if } * = 0, 6, \\ 0 & \text{otherwise,} \end{cases}$$

where  $m, d \geq 0$ , and  $T$  is a finitely generated abelian torsion group. Our first main theorem concerns the double suspension splitting of  $M$ . Let  $\Sigma X$  denote the suspension of any CW–complex  $X$ . Let  $P^n(T)$  be the Moore space such that the reduced cohomology  $\tilde{H}^*(P^n(T); \mathbb{Z})$  is isomorphic to  $T$  if  $* = n$  and 0 otherwise; see Neisendorfer [25].

**Theorem 1.1** *Suppose  $M$  is a simply connected closed orientable 6–manifold with homology of the form (1). Suppose that  $T$  has no 2 or 3–torsion. Then there is an integer  $c$  with  $0 \leq c \leq d$  determined by the cohomology ring of  $M$  such that:*

- If  $c = 0$ ,

$$\Sigma^2 M \simeq \Sigma W_0 \vee \bigvee_{j=1}^{d-1} (S^4 \vee S^6) \vee \bigvee_{i=1}^{2m} S^5 \vee P^6(T) \vee P^5(T),$$

where  $W_0 \simeq (S^3 \vee S^5) \cup e^7$ .

- If  $c = d$ ,

$$\Sigma^2 M \simeq \Sigma W_d \vee \bigvee_{i=1}^{d-1} \Sigma^2 \mathbb{C}P^2 \vee \bigvee_{i=1}^{2m} S^5 \vee P^6(T) \vee P^5(T),$$

where  $W_d \simeq \Sigma \mathbb{C}P^2 \cup e^7$ .

- If  $1 \leq c \leq d - 1$ ,

$$\Sigma^2 M \simeq \Sigma W_c \vee \bigvee_{i=1}^{c-1} \Sigma^2 \mathbb{C}P^2 \vee \bigvee_{j=1}^{d-c-1} (S^4 \vee S^6) \vee \bigvee_{i=1}^{2m} S^5 \vee P^6(T) \vee P^5(T),$$

where  $W_c \simeq (\Sigma \mathbb{C}P^2 \vee S^3 \vee S^5) \cup e^7$ .

**Theorem 1.1** classifies the homotopy type of  $\Sigma^2 M$  up to an indeterminate term  $\Sigma W_c$  for  $0 \leq c \leq d$ . The Steenrod square  $Sq^2: H^2(M; \mathbb{Z}/2\mathbb{Z}) \rightarrow H^4(M; \mathbb{Z}/2\mathbb{Z})$  determines  $c$ . Since we need only the suspension of  $W_c$ , the attaching map of the top cell of  $W_c$  can

be chosen so that it does not have a Whitehead product as a component. In general, the suspension of this attaching map depends on the manifold  $M$  itself; for instance, see [Theorem 1.3](#) below.

Nevertheless, [Theorem 1.1](#) is still useful, for instance, to calculate the  $K$ -group or the  $KO$ -group of  $M$  in [Corollary 1.2](#). In particular, when  $M$  is a Calabi–Yau threefold, it partially reproduces the result of Doran and Morgan [12, Corollary 1.10] about its  $K$ -group by a different method, and provides a new computation of its  $KO$ -group. Moreover, there are many examples of simply connected Calabi–Yau threefolds. For instance, based on Kreuzer and Skarke [21], Batyrev and Kreuzer [4] showed that there are exactly 473 800 760 families of simply connected Calabi–Yau threefolds corresponding to 4-dimensional reflexive polytopes.

**Corollary 1.2** *Let  $M$  be a manifold satisfying the conditions of [Theorem 1.1](#). Then the reduced  $K$ -group and  $KO$ -group of  $M$  are*

$$\tilde{K}(M) \cong \mathbb{Z}^{\oplus 2d+1} \oplus T \quad \text{and} \quad \tilde{KO}(M) \cong \bigoplus_d (\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}).$$

If we specialize to the case when  $d = 1$ , we can obtain a complete description of  $M$  after double suspension, based on the work of Yamaguchi [34] (also summarized and corrected by Baues [5]). In particular, Yamaguchi’s work [34] implies that a generator  $x \in H^2(M; \mathbb{Z}) \cong \mathbb{Z}$  has the property that  $x^2 = ky$  for some  $k \in \mathbb{Z}$  and a generator  $y \in H^4(M; \mathbb{Z})$  of infinite order. Let  $\eta_i^3 = \eta_{i+2} \circ \eta_{i+1} \circ \eta_i \in \pi_{i+3}(S^i)$  (see Toda [29]), where  $\eta_i \in \pi_{i+1}(S^i)$  is the Hopf element. Let  $V_3$  be the manifold that is the total space of the sphere bundle of the oriented  $\mathbb{R}^3$ -bundle over  $S^4$  determined by its first Pontryagin class  $p_1 = 12s_4$ , where  $s_4 \in H^4(S^4; \mathbb{Z})$  is a generator.

**Theorem 1.3** *Suppose  $M$  is a simply connected closed orientable 6-manifold with homology of the form (1) such that  $d = 1$ . Let  $x \in H^2(M; \mathbb{Z}) \cong \mathbb{Z}$  be a generator. Let  $k \in \mathbb{Z}$  be such that  $x^2 = ky$  for some generator  $y \in H^4(M; \mathbb{Z})$  of infinite order. Suppose  $T$  has no 2- or 3-torsion. Then:*

- If  $k$  is odd, then  $M$  is Spin. Moreover,

$$\Sigma^2 M \simeq \Sigma^2 \mathbb{C}P^3 \vee \bigvee_{i=1}^{2m} S^5 \vee P^6(T) \vee P^5(T)$$

when  $k \equiv \pm 1 \pmod{6}$ , while

$$\Sigma^2 M \simeq \Sigma^2 V_3 \vee \bigvee_{i=1}^{2m} S^5 \vee P^6(T) \vee P^5(T)$$

when  $k \equiv 3 \pmod{6}$ .

- If  $k$  is even and  $M$  is non-Spin,

$$\Sigma^2 M \simeq S^4 \vee \Sigma^4 \mathbb{C}P^2 \vee \bigvee_{i=1}^{2m} S^5 \vee P^6(T) \vee P^5(T).$$

- If  $k$  is even and  $M$  is Spin,

$$\Sigma^2 M \simeq (S^4 \cup_{\lambda \eta_4^3} e^8) \vee S^6 \vee \bigvee_{i=1}^{2m} S^5 \vee P^6(T) \vee P^5(T),$$

where  $\lambda \in \mathbb{Z}/2$  is determined by  $M$ .

It should be remarked that there is no indeterminacy in the term  $(S^4 \cup_{\lambda \eta_4^3} e^8)$  in the last decomposition of  $\Sigma^2 M$ . Indeed, the stable cube element  $\eta_n^3 \in \pi_{n+3}(S^n)$  (for  $n \geq 2$ ) is detected by the secondary operation  $\mathbb{T}$  (see Harper [13, Exercise 4.2.5]), and in our case the homotopy decomposition has to preserve the module structure induced by the cohomology operations. Moreover, it is clear that  $k \bmod 2$  and the spin condition of  $M$  are determined by the Steenrod square  $Sq^2$ . We will also see that  $\Sigma \mathbb{C}P^3$  and  $\Sigma V_3$  can be distinguished by the Steenrod power  $\mathcal{P}^1: H^3(\Sigma M; \mathbb{Z}/3\mathbb{Z}) \rightarrow H^7(\Sigma M; \mathbb{Z}/3\mathbb{Z})$ . Hence, we obtain the following rigidity result for manifolds of the type in [Theorem 1.3](#) after double suspension:

**Corollary 1.4** *Let  $M$  and  $M'$  be two manifolds of the type in [Theorem 1.3](#). Then  $\Sigma^2 M \simeq \Sigma^2 M'$  if and only if  $H^*(\Sigma^2 M; \mathbb{Z}) \cong H^*(\Sigma^2 M'; \mathbb{Z})$  as abelian groups, and  $H^*(\Sigma^2 M; \mathbb{Z}/p\mathbb{Z}) \cong H^*(\Sigma^2 M'; \mathbb{Z}/p\mathbb{Z})$  as  $\mathbb{Z}/2\mathbb{Z}\{Sq^2, \mathbb{T}\}$ -modules when  $p = 2$ , and as  $\mathbb{Z}/3\mathbb{Z}\{\mathcal{P}^1\}$ -modules when  $p = 3$ .*

The paper is organized as follows. In [Section 2](#) we reduce the decomposition problem of 6-manifolds to that of those whose third Betti numbers are zero. In [Section 3](#), we give a detailed procedure to decompose 6-manifolds after double suspension by the homology decomposition method. Sections 4 and 5 are devoted to proving [Theorems 1.1](#) and [1.3](#), respectively. In [Section 6](#), we compute some homotopy groups of odd primary Moore spaces used in [Section 3](#).

**Acknowledgements** Huang was supported by the National Natural Science Foundation of China (grants 11801544 and 11688101), and the Chen Jingrun Future Star Program of the AMSS. He would like to thank Professor Stephen Theriault for the international online lecture series *Loop space decompositions*, which stimulated his research interest in the homotopy of 6-manifolds.

## 2 Reducing to the case when $b_3(M) = 0$

The following well known splitting theorem for 6-manifolds was proved by Wall [31] in the smooth category, while Jupp [18] pointed out that the theorem holds in the topological category by the same argument.

**Theorem 2.1** [31, Theorem 1] *Suppose  $M$  is a simply connected closed orientable 6-manifold with third Betti number  $b_3(M) = 2m$ . Then there exists a 6-manifold  $M_1$  such that*

$$M \cong M_1 \# \underset{m}{\#}(S^3 \times S^3) \quad \text{and} \quad H_3(M_1; \mathbb{Q}) = 0.$$

**Corollary 2.2** *Let  $M$  and  $M_1$  be manifolds as in Theorem 2.1. Then*

$$\Sigma M \simeq \Sigma M_1 \vee \bigvee_{i=1}^m (S^4 \vee S^4).$$

**Proof** Let  $M'_1$  and  $M'$  be the 5-skeletons of  $M_1$  and  $M$ , respectively. It is known that  $S^3 \vee S^3$  is the 5-skeleton of  $S^3 \times S^3$  and  $\Sigma(S^3 \times S^3) \simeq \Sigma(S^3 \vee S^3) \vee S^7$ . In particular, there is a homotopy retraction  $r: \Sigma(S^3 \times S^3) \rightarrow \Sigma(S^3 \vee S^3)$ . For the connected sum  $M_1 \# (S^3 \times S^3)$ , there are the obvious pinch maps  $q_1: M_1 \# (S^3 \times S^3) \rightarrow M_1$  and  $q_2: M_1 \# (S^3 \times S^3) \rightarrow S^3 \times S^3$ . Consider the composition

$$\begin{aligned} \phi: \Sigma(M_1 \# (S^3 \times S^3)) &\xrightarrow{\mu'} \Sigma(M_1 \# (S^3 \times S^3)) \vee \Sigma(M_1 \# (S^3 \times S^3)) \\ &\xrightarrow{Eq_1 \vee (r \circ Eq_2)} \Sigma M_1 \vee \Sigma(S^3 \vee S^3), \end{aligned}$$

where  $\mu'$  is the standard comultiplication for a suspension, and  $E$  denotes the suspension of a map. It is easy to see that  $\phi$  induces an isomorphism on homology and so is a homotopy equivalence by Whitehead's theorem. Since

$$M' \simeq M'_1 \vee \bigvee_{i=1}^m (S^3 \vee S^3),$$

by repeating the above argument, we obtain the decomposition in the statement of the corollary. □

In Theorem 2.1 the connected summand  $M_1$  satisfies  $b_3(M_1) = 0$ , so by Corollary 2.2 it suffices to consider such 6-manifolds in the sequel.

### 3 Homology decomposition of $M$ after suitable suspensions

Let  $M$  be a simply connected closed orientable 6–manifold with  $b_3(M) = 0$ . By Poincaré duality and the universal coefficient theorem, we have

$$(2) \quad H_*(M; \mathbb{Z}) = \begin{cases} \mathbb{Z}^{\oplus d} \oplus T & \text{if } * = 2, \\ T & \text{if } * = 3, \\ \mathbb{Z}^{\oplus d} & \text{if } * = 4, \\ \mathbb{Z} & \text{if } * = 0, 6, \\ 0 & \text{otherwise,} \end{cases}$$

where  $d \geq 0$ , and  $T$  is a finitely generated abelian torsion group. We may write

$$(3) \quad T = \bigoplus_{k=1}^{\ell} \mathbb{Z} / p_k^{r_k} \mathbb{Z},$$

where each  $p_k$  is a prime and  $r_k \geq 1$ .

Instead of using a skeletal decomposition, we may apply a *homology decomposition* to study the cell structure of  $M$ . For any finitely generated abelian group  $A$ , let  $P^n(A)$  be the Moore space such that  $\tilde{H}^*(P^n(A); \mathbb{Z}) \cong A$  if  $* = n$  and 0 otherwise [25]. The information on the homotopy groups of  $P^n(T)$  used in this section will be proved in Section 6.

**Theorem 3.1** [14, Theorem 4H.3] *Let  $X$  be a simply connected CW–complex. Write  $H_i = H_i(X; \mathbb{Z})$ . Then there is a sequence of complexes  $X_{(i)}$  (for  $i \geq 2$ ) such that:*

- $H_j(X_{(i)}; \mathbb{Z}) \cong H_j(X; \mathbb{Z})$  for  $j \leq i$  and  $H_j(X_{(i)}; \mathbb{Z}) = 0$  for  $j > i$ .
- $X_{(2)} = P^3(H_2)$ , and  $X_{(i)}$  is defined by a homotopy cofibration

$$P^i(H_i) \xrightarrow{f_{i-1}} X_{(i-1)} \xrightarrow{l_{i-1}} X_{(i)},$$

where  $f_{i-1}$  induces a trivial homomorphism

$$f_{i-1*}: H_{i-1}(P^i(H_i); \mathbb{Z}) \rightarrow H_{i-1}(X_{(i-1)}; \mathbb{Z}).$$

- $X \simeq \text{hocolim}\{X_{(2)} \xrightarrow{l_2} \dots \xrightarrow{l_{i-2}} X_{(i-1)} \xrightarrow{l_{i-1}} X_{(i)} \xrightarrow{l_i} \dots\}$ .

From this theorem, it is clear that the homology decomposition is compatible with the suspension functor. That is, for  $X$  in Theorem 3.1 the sequence of the triples  $(\Sigma X_{(i)}, E f_i, E l_i)$  is a homology decomposition of  $\Sigma X$ .

### 3.1 Structure of $M_{(3)}$

By [Theorem 3.1](#) and (2), there is a homotopy cofibration

$$(4) \quad P^3(T) \xrightarrow{f_2} M_{(2)} \xrightarrow{\iota_2} M_{(3)}, \quad \text{where } M_{(2)} \simeq \bigvee_{i=1}^d S^2 \vee P^3(T).$$

Notice that from (3), we have  $P^n(T) \simeq \bigvee_{k=1}^{\ell} P^n(p_k^{r_k})$  by [23] or [25].

**Lemma 3.2** *The map  $f_2$  in (4) is nullhomotopic, and hence*

$$M_{(3)} \simeq \bigvee_{i=1}^d S^2 \vee P^3(T) \vee P^4(T).$$

**Proof** Since  $P^3(T) \simeq \bigvee_{k=1}^{\ell} P^3(p_k^{r_k})$ , there is the embedding  $j: \bigvee_{i=1}^{\ell} S^2 \rightarrow P^3(T)$  of the bottom cells. Consider the commutative diagram

$$\begin{array}{ccccc} \pi_2(\bigvee_{i=1}^{\ell} S^2) & \xrightarrow{j_*} & \pi_2(P^3(T)) & \xrightarrow{f_{2*}} & \pi_2(M_{(2)}) \\ \cong \downarrow \text{hur} & & \cong \downarrow \text{hur} & & \cong \downarrow \text{hur} \\ H_2(\bigvee_{i=1}^{\ell} S^2; \mathbb{Z}) & \xrightarrow{j_*} & H_2(P^3(T); \mathbb{Z}) & \xrightarrow{f_{2*}=0} & H_2(M_{(2)}; \mathbb{Z}) \end{array}$$

where the Hurewicz homomorphisms *hur* are isomorphisms by the Hurewicz theorem,  $f_{2*} = 0$  on homology by [Theorem 3.1](#), and both maps  $j_*$  are epimorphisms. In particular,  $f_{2*} \circ j_*$  is trivial on homotopy groups, and hence  $f_2 \circ j$  is nullhomotopic. Thus, combined with (4), we have the diagram of homotopy cofibrations

$$(5) \quad \begin{array}{ccccc} \bigvee_{i=1}^{\ell} S^2 & \longrightarrow & * & \longrightarrow & \bigvee_{i=1}^{\ell} S^3 \\ \downarrow \bigvee_{i=1}^{\ell} p_k^{r_k} & & \downarrow & & \downarrow c \\ \bigvee_{i=1}^{\ell} S^2 & \xrightarrow{0} & M_{(2)} & \longrightarrow & M_{(2)} \vee \bigvee_{i=1}^{\ell} S^3 \\ \downarrow j & & \parallel & & \downarrow \\ P^3(T) & \xrightarrow{f_2} & M_{(2)} & \xrightarrow{\iota_2} & M_{(3)} \end{array}$$

where  $p_k^{r_k}: S^n \rightarrow S^n$  is a map of degree  $p_k^{r_k}$ , and  $c$  is the induced map.

We claim that  $c$  can be chosen to be  $i_2 \circ (\bigvee_{i=1}^{\ell} p_k^{r_k})$  in (5), where

$$i_2: \bigvee_{i=1}^{\ell} S^3 \rightarrow M_{(2)} \vee \bigvee_{i=1}^{\ell} S^3$$

is the injection onto the sphere summands. Indeed, we may replace (5) by a strictly commutative diagram up to homeomorphism. Start from the upper left square of (5). It



becomes strictly commutative once we replace the 0 map by the constant map. On the one hand, extending the new square one step to the right by taking the mapping cone, we obtain  $\tilde{c}: \bigvee_{i=1}^{\ell} S^3 \rightarrow M_{(2)} \vee \bigvee_{i=1}^{\ell} S^3$ , which is exactly equal to  $i_2 \circ (\bigvee_{i=1}^{\ell} p_k^{rk})$ , and the upper right square automatically commutes. On the other hand, by the homotopy extension property of a cofibration, we may replace  $f_2: P^3(T) \rightarrow M_{(2)}$  by a map  $\tilde{f}_2$  such that  $\tilde{f}_2 \simeq f_2$  and the lower left square strictly commutes. It follows that the mapping cone of  $\tilde{f}_2$  is homotopy equivalent to  $M_{(3)}$  and is homeomorphic to the mapping cone of  $i_2 \circ (\bigvee_{i=1}^{\ell} p_k^{rk})$ . Hence we can choose  $c = i_2 \circ (\bigvee_{i=1}^{\ell} p_k^{rk})$  in (5).

Now from (5) it follows that

$$M_{(3)} \simeq M_{(2)} \vee P^4(T) \simeq \bigvee_{i=1}^d S^2 \vee P^3(T) \vee P^4(T), \quad \square$$

The following corollary follows from Lemma 3.2 and will be used in Lemma 3.6.

**Corollary 3.3** *The homotopy cofiber of the obvious inclusion*

$$j: P^3(T) \vee P^4(T) \rightarrow M_{(3)} \rightarrow M$$

is a Poincaré duality complex  $V$  with cell structure

$$V = \bigvee_{i=1}^d S^2 \cup e_{(1)}^4 \cup e_{(2)}^4 \cdots \cup e_{(d)}^4 \cup e^6.$$

Moreover, by [31, Theorem 8]  $V$  is homotopy equivalent to a closed smooth manifold.

### 3.2 Structure of $M_{(5)}$

By Theorem 3.1 and Lemma 3.2, there is a homotopy cofibration

$$(6) \quad \bigvee_{i=1}^d S^3 \xrightarrow{f_3} M_{(3)} \xrightarrow{i_3} M_{(4)} = M_{(5)}, \quad \text{where } M_{(3)} \simeq \bigvee_{i=1}^d S^2 \vee P^3(T) \vee P^4(T).$$

We need to study the map

$$f_3: \bigvee_{i=1}^d S^3 \rightarrow \bigvee_{i=1}^d S^2 \vee P^3(T) \vee P^4(T).$$

Let  $i_3: P^4(T) \rightarrow \bigvee_{i=1}^d S^2 \vee P^3(T) \vee P^4(T)$  be the inclusion. Define the complex  $Y$  by the homotopy cofibration

$$P^4(T) \xrightarrow{i_3 \circ i_3} M_{(4)} = M_{(5)} \rightarrow Y.$$

**Lemma 3.4** The map  $f_3$  in (6) factors as

$$f_3: \bigvee_{i=1}^d S^3 \xrightarrow{f'_3} \bigvee_{i=1}^d S^2 \vee P^3(T) \xrightarrow{i_1 \vee i_2} \bigvee_{i=1}^d S^2 \vee P^3(T) \vee P^4(T)$$

for some  $f'_3$ , where  $i_1$  and  $i_2$  are inclusions. Moreover, there is a homotopy cofibration

$$(7) \quad \bigvee_{i=1}^d S^3 \xrightarrow{f'_3} \bigvee_{i=1}^d S^2 \vee P^3(T) \xrightarrow{\iota'_3} Y,$$

and

$$M_{(5)} \simeq Y \vee P^4(T).$$

**Proof** First, there is the diagram of homotopy cofibrations

$$\begin{array}{ccccc} * & \longrightarrow & P^4(T) & \xlongequal{\quad} & P^4(T) \\ \downarrow & & \downarrow i_3 & & \downarrow \iota_3 \circ i_3 \\ \bigvee_{i=1}^d S^3 & \xrightarrow{f_3} & M_{(3)} & \xrightarrow{\iota_3} & M_{(5)} \\ \parallel & & \downarrow q_{1,2} & & \downarrow \\ \bigvee_{i=1}^d S^3 & \xrightarrow{f'_3} & \bigvee_{i=1}^d S^2 \vee P^3(T) & \xrightarrow{\iota'_3} & Y \end{array}$$

where  $q_{1,2}$  is the obvious projection,  $\iota'_3$  is induced from  $\iota_3$ , and  $f'_3 := q_{1,2} \circ f_3$ . The diagram immediately implies that (7) is a homotopy cofibration.

Let  $q_3: \bigvee_{i=1}^d S^2 \vee P^3(T) \vee P^4(T) \rightarrow P^4(T)$  be the canonical projection. By Theorem 3.1,  $f_{3*}: H_3(\bigvee_{i=1}^d S^3; \mathbb{Z}) \rightarrow H_3(\bigvee_{i=1}^d S^2 \vee P^3(T) \vee P^4(T); \mathbb{Z})$  is trivial. In particular,  $q_{3*} \circ f_{3*} = 0$ . Then, by the Hurewicz Theorem,  $q_3 \circ f_3$  is nullhomotopic. Further, by the Hilton–Milnor Theorem (see §XI.6 of [33], for instance),  $\pi_3(S^2 \vee P^3(T) \vee P^4(T)) \cong \pi_3(S^2 \vee P^3(T)) \oplus \pi_3(P^4(T))$ , and hence

$$[\bigvee_{i=1}^d S^3, S^2 \vee P^3(T) \vee P^4(T)] \cong [\bigvee_{i=1}^d S^3, S^2 \vee P^3(T)] \oplus [\bigvee_{i=1}^d S^3, P^4(T)].$$

Under this isomorphism, the homotopy class of  $f_3$  corresponds to  $[f'_3] + [q_3 \circ f_3]$ . However since we already showed that  $[q_3 \circ f_3] = 0$ , we have  $f_3 \simeq (i_1 \vee i_2) \circ f'_3$ , so  $M_{(5)} \simeq Y \vee P^4(T)$  as required.  $\square$

### 3.3 Structure of $\Sigma M_{(5)}$

From this point, we may need extra conditions on the torsion group  $T$ . First recall that we already showed that by Lemma 3.4 there is a homotopy cofibration

$$(8) \quad \bigvee_{i=1}^d S^3 \xrightarrow{f'_3} \bigvee_{i=1}^d S^2 \vee P^3(T) \xrightarrow{\iota'_3} Y, \quad \text{where } M_{(5)} \simeq Y \vee P^4(T).$$

Let

$$q_1: \bigvee_{i=1}^d S^2 \vee P^3(T) \rightarrow \bigvee_{i=1}^d S^2$$

be the canonical projection, let  $f_3'' := q_1 \circ f_3' = q_1 \circ q_{1,2} \circ f_3$ , and also recall that  $P^n(T) \simeq \bigvee_{k=1}^{\ell} P^n(P_k^{r_k})$ . Let us suppose  $p_k \geq 3$  for each  $k$  from now on.

**Lemma 3.5** *Suppose  $T$  has no 2-torsion. Then there is a homotopy equivalence*

$$\Sigma M_{(5)} \simeq \Sigma X \vee P^5(T) \vee P^4(T),$$

where  $X$  is the homotopy cofiber of the map  $f_3'': \bigvee_{i=1}^d S^3 \rightarrow \bigvee_{i=1}^d S^2$ .

**Proof** There is the diagram of homotopy cofibrations

$$\begin{array}{ccccc}
 * & \longrightarrow & P^3(T) & \xlongequal{\quad} & P^3(T) \\
 \downarrow & & \downarrow i_2 & & \downarrow \iota_3' \circ i_2 \\
 \bigvee_{i=1}^d S^3 & \xrightarrow{f_3'} & \bigvee_{i=1}^d S^2 \vee P^3(T) & \xrightarrow{\iota_3'} & Y \\
 \parallel & & \downarrow q_1 & & \downarrow \\
 \bigvee_{i=1}^d S^3 & \xrightarrow{f_3''} & \bigvee_{i=1}^d S^2 & \longrightarrow & X
 \end{array}$$

where  $i_2$  is the canonical inclusion. Since  $\pi_4(P^4(P^r)) = 0$  for odd  $p$  by Lemma 6.3, we have  $\Sigma Y \simeq \Sigma X \vee P^4(T)$ . The lemma then follows from (8). □

### 3.4 Structure of $\Sigma^2 M$

Recall, when  $T$  has no 2-torsion, by Lemma 3.5 there is a homotopy cofibration

$$(9) \quad S^5 \xrightarrow{f_5} M_{(5)} \xrightarrow{\iota_5} M, \quad \text{where } \Sigma M_{(5)} \simeq \Sigma X \vee P^5(T) \vee P^4(T).$$

Further, by Corollary 3.3 we have the homotopy cofibration

$$S^5 \rightarrow X \rightarrow V,$$

where  $X$  is defined in Lemma 3.5 without restriction on  $T$ , and  $V$  is a closed smooth manifold. We now further suppose  $T$  has no 3-torsion.

**Lemma 3.6** *Suppose  $T$  has no 2- or 3-torsion. Then*

$$\Sigma^2 M \simeq \Sigma^2 V \vee P^6(T) \vee P^5(T).$$

**Proof** By the Hilton–Milnor theorem, we may write the suspension of  $f_5$  as

$$Ef_5 = g_5^{(1)} + g_5^{(2)} + g_5^{(3)} + \theta: S^6 \rightarrow \Sigma M_{(5)} \simeq \Sigma X \vee P^5(T) \vee P^4(T),$$

for some  $\theta$ , where  $E\theta = 0$ ,  $g_5^{(i)} = q_i \circ Ef_5$ , and  $q_i$  is the canonical projection of  $\Sigma X \vee P^5(T) \vee P^4(T)$  onto its  $i^{\text{th}}$  summand. Then by Lemma 6.4  $g_5^{(2)} = 0$ , and  $Eg_5^{(3)} = 0$  by Lemma 6.6. It follows that  $E^2 f_5 = Eg_5^{(1)}$ . Furthermore, there is the diagram of homotopy cofibrations

$$\begin{array}{ccccc}
 * & \longrightarrow & P^4(T) \vee P^3(T) & \simeq & P^4(T) \vee P^3(T) \\
 \downarrow & & \downarrow j_5 & & \downarrow j \\
 S^5 & \xrightarrow{f_5} & M_{(5)} & \xrightarrow{\iota_5} & M \\
 \parallel & & \downarrow \pi_5 & & \downarrow \pi \\
 S^5 & \longrightarrow & X & \longrightarrow & V
 \end{array}$$

where the homotopy cofibration in the last column is defined in Corollary 3.3 by using Lemma 3.2, and similarly the homotopy cofibration in the middle column can be also defined by using Lemma 3.2. Then it is clear that  $g_5^{(1)} \simeq E(\pi_5 \circ f_5)$  and the lemma follows. □

### 4 Proof of Theorem 1.1 and Corollary 1.2

In Lemma 3.6 we established the double suspension splitting of  $M$  when  $b_3(M) = 0$  and are now left to consider the homotopy type of  $V$  after suspension. Recall that  $V$  is a Poincaré duality complex of dimension 6, and its 5-skeleton  $V_5 = X$  is the homotopy cofiber of the map  $f_3'': \bigvee_{i=1}^d S^3 \rightarrow \bigvee_{i=1}^d S^2$  by Lemma 3.5. The following lemma, as a special case of [16, Lemma 6.1], determines the suspension homotopy type of  $X$ .

**Lemma 4.1** [16, Lemma 6.1] *There is a homotopy equivalence*

$$\Sigma X \simeq \bigvee_{i=1}^c \Sigma \mathbb{C}P^2 \vee \bigvee_{j=1}^{d-c} (S^3 \vee S^5)$$

for some  $0 \leq c \leq d$ .

We may apply the method in [16, Section 3] to decompose  $\Sigma^2 V$ , in the same way that we used it to prove [16, Lemmas 6.4 and 6.6].

**Lemma 4.2** Suppose  $\Sigma X$  decomposes as in Lemma 4.1.

- If  $c = 0$ ,

$$\Sigma^2 V \simeq \Sigma W_0 \vee \bigvee_{j=1}^{d-1} (S^4 \vee S^6),$$

where  $W_0 \simeq (S^3 \vee S^5) \cup e^7$ .

- If  $c = d$ ,

$$\Sigma^2 V \simeq \Sigma W_d \vee \bigvee_{i=1}^{d-1} \Sigma^2 \mathbb{C}P^2,$$

where  $W_d \simeq \Sigma \mathbb{C}P^2 \cup e^7$ .

- If  $1 \leq c \leq d - 1$ ,

$$\Sigma^2 V \simeq \Sigma W_c \vee \bigvee_{i=1}^{c-1} \Sigma^2 \mathbb{C}P^2 \vee \bigvee_{j=1}^{d-c-1} (S^4 \vee S^6),$$

where  $W_c \simeq (\Sigma \mathbb{C}P^2 \vee S^3 \vee S^5) \cup e^7$ .

**Proof** Since the proof is similar to those of [16, Lemmas 6.4 and 6.6], we only sketch it. The interested reader can find the details in [16, Section 3]. Using Lemma 4.1, let  $g: S^6 \rightarrow \Sigma X \simeq \bigvee_{i=1}^c \Sigma \mathbb{C}P^2 \vee \bigvee_{j=1}^{d-c} (S^3 \vee S^5)$  be the attaching map of the top cell of  $\Sigma V$ . To apply the method in [16, Section 3], we only need information about the homotopy groups  $\pi_6(\Sigma \mathbb{C}P^2) \cong \mathbb{Z}/6\mathbb{Z}$  by [22, Proposition 8.2(i)],  $\pi_6(S^3) \cong \mathbb{Z}/12\mathbb{Z}$  and  $\pi_6(S^5) \cong \mathbb{Z}/2$ , which are all finite cyclic groups. Then we can represent the attaching map  $Eg$  of the top cell of  $\Sigma^2 V$  by a matrix  $B$ , and apply [16, Lemma 3.1] to transform  $B$  to a simpler matrix  $C$ . The new matrix representation  $C$  of the attaching map, corresponding to a base change of  $\Sigma X$  through a self homotopy equivalence, will give the desired decomposition.  $\square$

**Proof of Theorem 1.1** First, by Theorem 2.1 and Corollary 2.2, we have

$$\Sigma M \simeq \Sigma M_1 \vee \bigvee_{i=1}^m (S^4 \vee S^4),$$

where  $M_1$  is a closed 6-manifold with homology of the form (2). In particular  $b_3(M_1) = 0$ . Hence, by Lemmas 3.6 and 4.2, we have that if  $1 \leq c \leq d - 1$  then

$$\Sigma^2 M \simeq \Sigma W_c \vee \bigvee_{i=1}^{c-1} \Sigma^2 \mathbb{C}P^2 \vee \bigvee_{j=1}^{d-c-1} (S^4 \vee S^6) \vee P^6(T) \vee P^5(T) \vee \bigvee_{i=1}^{2m} S^5,$$

where  $W_c \simeq (\Sigma \mathbb{C}P^2 \vee S^3 \vee S^5) \cup e^7$ . The decompositions for the cases when  $c = 0$  or  $c = d$  can be obtained similarly. Finally, notice that  $c$  records the number of the nontrivial Steenrod square  $Sq^2: H^2(\Sigma^2 M; \mathbb{Z}/2\mathbb{Z}) \rightarrow H^4(\Sigma^2 M; \mathbb{Z}/2\mathbb{Z})$ , which is preserved by the decomposition and the suspension operator. Since  $Sq^2$  is the cup square on the elements of  $H^2(M; \mathbb{Z}/2\mathbb{Z})$ , this completes the proof of Theorem 1.1.  $\square$

|                       |              |   |                            |              |                          |                          |   |              |   |   |   |
|-----------------------|--------------|---|----------------------------|--------------|--------------------------|--------------------------|---|--------------|---|---|---|
| $i \bmod 2$           | 0            | 1 | $j \bmod 8$                | 0            | 1                        | 2                        | 3 | 4            | 5 | 6 | 7 |
| $\tilde{K}^{-i}(S^0)$ | $\mathbb{Z}$ | 0 | $\widetilde{KO}^{-j}(S^0)$ | $\mathbb{Z}$ | $\mathbb{Z}/2\mathbb{Z}$ | $\mathbb{Z}/2\mathbb{Z}$ | 0 | $\mathbb{Z}$ | 0 | 0 | 0 |

Table 1:  $\tilde{K}^{-i}(S^0)$  and  $\widetilde{KO}^{-j}(S^0)$ .

To prove [Corollary 1.2](#) we need Bott periodicity, which is described by [Table 1](#), from which we can easily calculate the following, where only  $\widetilde{KO}^2(P^5(T)) = 0$  requires that  $T$  has no 2-torsion:

**Lemma 4.3** *Let  $W_c$  be the complex defined in [Lemma 4.2](#) for  $0 \leq c \leq d$ .*

- $\tilde{K}(P^5(T)) \cong T$  and  $\tilde{K}(P^6(T)) = 0$ .
- $\widetilde{KO}^2(P^5(T)) = \widetilde{KO}^2(P^6(T)) = 0$ .
- $\widetilde{KO}^1(\Sigma\mathbb{C}P^2) \cong \widetilde{KO}^1(S^3 \vee S^5) \cong \mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}$ .
- $\widetilde{KO}^1(\Sigma^2\mathbb{C}P^2) \cong \widetilde{KO}^1(S^4 \vee S^6) = 0$ .
- $\widetilde{KO}^1(W_0) \cong \widetilde{KO}^1(W_d) \cong \mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}$  and  $\widetilde{KO}^1(W_c) \cong \bigoplus_2(\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z})$ .

**Proof of [Corollary 1.2](#)** We only compute the  $\widetilde{KO}$ -group of  $M$  when  $1 \leq c \leq d - 1$ , as the other cases can be computed similarly. By [Theorem 1.1](#) and [Lemma 4.3](#), we have

$$\begin{aligned}
 \widetilde{KO}(M) &\cong \widetilde{KO}^2(\Sigma^2 M) \cong \widetilde{KO}^2(\Sigma W_c) \oplus \bigoplus_{i=1}^{c-1} \widetilde{KO}^2(\Sigma^2\mathbb{C}P^2) \oplus \bigoplus_{j=1}^{d-c-1} \widetilde{KO}^2(S^4 \vee S^6) \\
 &\quad \oplus \bigoplus_{j=1}^{2m} \widetilde{KO}^2(S^5) \oplus \widetilde{KO}^2(P^6(T)) \oplus \widetilde{KO}^2(P^5(T)) \\
 &\cong \bigoplus_2(\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}) \oplus \bigoplus_{c-1}(\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}) \oplus \bigoplus_{d-c-1}(\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}) \\
 &= \bigoplus_d(\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}). \quad \square
 \end{aligned}$$

## 5 The case when $H^2(M; \mathbb{Z}) \cong \mathbb{Z}$

By [Lemma 3.6](#), we may consider the torsion free case first. In [\[34\]](#), Yamaguchi classified the homotopy types of CW-complexes of the form  $V \simeq S^2 \cup e^4 \cup e^6$ . Specializing to the case when  $V$  is a manifold, we can summarize the necessary result in the following theorem (see [\[5, Section 1\]](#)):

**Theorem 5.1** [34, Corollary 4.6, Lemmas 2.6 and 4.3] *Let  $V \simeq S^2 \cup_{k\eta_2} e^4 \cup_b e^6$  be a closed smooth manifold, where  $k\eta_2$  with  $k \in \mathbb{Z}$  and  $b$  are the attaching maps of the cells  $e^4$  and  $e^6$ , respectively. Then the top attaching map  $b$  is determined by a generator  $b_W \in \pi_5(S^2 \cup_{k\eta_2} e^4)$  of infinite order, the second Stiefel–Whitney class  $\omega_2(V) \in H^2(V; \mathbb{Z}/2)$  of  $V$ , and an indeterminacy term  $b' \in \mathbb{Z}/2$  which depends on the following three cases:*

- *If  $k$  is odd, then  $V$  is Spin, the homotopy type of  $V$  is uniquely determined by  $k$ , and  $b = b_W$ .*
- *If  $k$  is even and  $V$  is non-Spin, then the homotopy type of  $V$  is uniquely determined by  $k$ , and  $b = b_W + \tilde{\eta}_4$  with  $\tilde{\eta}_4$  representing the generator of a  $\mathbb{Z}/2$  summand determined by  $\omega_2(V)$ .*
- *If  $k$  is even and  $V$  is Spin, then  $V$  has precisely two homotopy types depending on the value of  $b' \in \mathbb{Z}/2$ , and  $b = b_W + b'$ .*

**Remark 5.2** In Theorem 5.1,  $b_W$ , as a generator of the  $\mathbb{Z}$ –summand of  $\pi_5(S^2 \cup_{k\eta_2} e^4)$ , is indeed a relative Whitehead product when  $k \neq 0$  by [34, Lemma 2.6]. It is possible that the suspension  $Eb_W$  is not nullhomotopic. The class  $\tilde{\eta}_4$  is derived from the homotopy class of

$$S^5 \xrightarrow{b} S^2 \cup_{k\eta_2} e^4 \xrightarrow{q} S^4,$$

where  $q$  is the quotient map onto the 4–cell of  $S^2 \cup_{k\eta_2} e^4$  (see [5, Section 1]). The class  $b'$  is from a class of  $\pi_5(S^2) \cong \mathbb{Z}/2\{\eta_2^3\}$  by [34, Lemma 2.6] or [5, Section 1]. Also, as pointed out in Mathematical Reviews [26], the original theorem of [34] was misstated, but is corrected here and in [5, Section 1] as well.

Thanks to Theorem 5.1 and Remark 5.2, we can describe the suspension homotopy type of  $V$ . Recall that  $\pi_6(\Sigma\mathbb{C}P^2) \cong \mathbb{Z}/6\mathbb{Z}\{E\pi_2\}$  [22, Theorem 8.2(i)], where  $\pi_2: S^5 \rightarrow \mathbb{C}P^2$  is the Hopf map with the cofibre  $\mathbb{C}P^3$ , and  $E$  is the suspension of a map.

**Proposition 5.3** *Let  $V \simeq S^2 \cup_{k\eta_2} e^4 \cup_b e^6$  be a closed smooth manifold.*

- *If  $k$  is odd, then  $V$  is Spin and*

$$\Sigma V \simeq \Sigma\mathbb{C}P^2 \cup_{k'E\pi_2} e^7,$$

where  $k' = 1$  or  $3$  is such that  $k' \equiv \pm k \pmod{6}$ .

- If  $k$  is even and  $V$  is non-Spin then

$$\Sigma V \simeq S^3 \vee \Sigma^3 \mathbb{C}P^2.$$

- If  $k$  is even and  $V$  is Spin then

$$\Sigma V \simeq (S^3 \cup_{b'\eta_3} e^7) \vee S^5,$$

where  $b' \in \mathbb{Z}/2$  is from [Theorem 5.1](#).

**Proof** When  $k$  is even the decompositions follow immediately from [Theorem 5.1](#) and [Remark 5.2](#). When  $k$  is odd,  $V$  is Spin and  $\Sigma V \simeq \Sigma \mathbb{C}P^2 \cup_{Eb_W} e^7$  by [Theorem 5.1](#). Also notice that  $\Sigma \mathbb{C}P^2 \cup_{Eb_W} e^7 \simeq \Sigma \mathbb{C}P^2 \cup_{-Eb_W} e^7$ . Hence, to prove the statement in the proposition it suffices to show that the suspension map

$$E: \pi_5(S^2 \cup_{k\eta_2} e^4) \rightarrow \pi_6(\Sigma \mathbb{C}P^2)$$

sends the generator  $b_W$  to  $kE\pi_2$  up to sign.

For this purpose, start with the diagram of homotopy cofibrations

$$(10) \quad \begin{array}{ccccc} S^3 & \xrightarrow{k\eta_2} & S^2 & \longrightarrow & S^2 \cup_{k\eta_2} e^4 \\ \downarrow k & & \parallel & & \downarrow r \\ S^3 & \xrightarrow{\eta_2} & S^2 & \longrightarrow & \mathbb{C}P^2 \\ \downarrow & & \downarrow & & \downarrow \\ P^4(k) & \longrightarrow & * & \longrightarrow & P^5(k) \end{array}$$

which defines the map  $r$ . Then there is the diagram of homotopy fibrations

$$(11) \quad \begin{array}{ccccccc} S^1 & \longrightarrow & Z & \longrightarrow & S^2 \cup_{k\eta_2} e^4 & \xrightarrow{f_x} & K(\mathbb{Z}, 2) \\ \parallel & & \downarrow \tilde{r} & & \downarrow r & & \parallel \\ S^1 & \longrightarrow & S^5 & \xrightarrow{\pi_2} & \mathbb{C}P^2 & \xrightarrow{f_c} & K(\mathbb{Z}, 2) \end{array}$$

where  $f_c$  and  $f_x$  represent the generators  $c \in H^2(\mathbb{C}P^2; \mathbb{Z})$  and  $x \in H^2(S^2 \cup_{k\eta_2} e^4; \mathbb{Z})$ , respectively, and  $Z$  is the homotopy fibre of  $f_x$  mapping to  $S^5$  by the induced map  $\tilde{r}$ . By analyzing the Serre spectral sequences of the homotopy fibrations in (11), it can be shown that  $Z \simeq P^4(k) \cup e^5$  and  $\tilde{r}^*: H^5(S^5; \mathbb{Z}) \rightarrow H^5(Z; \mathbb{Z})$  is of degree  $k$ . Since by [Lemma 6.3](#)  $\pi_4(P^4(k)) = 0$  when  $k$  is odd, we see that  $Z \simeq P^4(k) \vee S^5$ , and then  $\tilde{r}_*$  is of degree  $k$  on homology. Moreover, by the naturality of the Hurewicz homomorphism



and Lemma 6.4, it is easy to see that  $\tilde{r}_*: \pi_5(Z) \cong \mathbb{Z} \rightarrow \pi_5(S^5)$  is of degree  $k$ . It follows that  $r_*: \pi_5(S^2 \cup_{k\eta_2} e^4) \cong \mathbb{Z} \rightarrow \pi_5(\mathbb{C}P^2) \cong \mathbb{Z}$  is of degree  $k$  by (11).

Now the naturality of suspension map induces the commutative diagram

$$(12) \quad \begin{array}{ccc} \pi_5(S^2 \cup_{k\eta_2} e^4) & \xrightarrow{r_*} & \pi_5(\mathbb{C}P^2) \\ \downarrow E & & \downarrow E \\ \pi_6(\Sigma\mathbb{C}P^2) & \xrightarrow{Er_*} & \pi_6(\Sigma\mathbb{C}P^2) \end{array}$$

where  $E: \pi_5(\mathbb{C}P^2) \cong \mathbb{Z} \rightarrow \pi_6(\Sigma\mathbb{C}P^2) \cong \mathbb{Z}/6\mathbb{Z}$  is surjective by [22, Theorem 8.2(i)]. We have shown that  $r_*$  in (12) is of degree  $k$ . On the other hand, from the last column of (10) we have the homotopy cofibration

$$\Sigma\mathbb{C}P^2 \xrightarrow{Er} \Sigma\mathbb{C}P^2 \rightarrow P^6(k).$$

Applying the Blakers–Massey theorem [9], we obtain the exact sequence

$$(13) \quad \pi_6(\Sigma\mathbb{C}P^2) \xrightarrow{Er_*} \pi_6(\Sigma\mathbb{C}P^2) \rightarrow \pi_6(P^6(k)).$$

Since  $\pi_6(P^6(k)) = 0$  by Lemma 6.3 and  $\pi_6(\Sigma\mathbb{C}P^2) \cong \mathbb{Z}/6\mathbb{Z}\{E\pi_2\}$ , we see that  $Er_*$  is an isomorphism from (13). Then by (12),  $E: \pi_5(S^2 \cup_{k\eta_2} e^4) \rightarrow \pi_6(\Sigma\mathbb{C}P^2)$  sends the generator  $b_W$  to  $kE\pi_2$  up to sign. This proves the statement in the case when  $k$  is odd, and we have completed the proof of the proposition.  $\square$

Now we can prove Theorem 1.3 and Corollary 1.4.

**Proof of Theorem 1.3** First, by Theorem 2.1, Corollary 2.2 and Lemma 3.6, we have

$$\begin{aligned} \Sigma^2 M &\simeq \Sigma^2 M_1 \vee \bigvee_{i=1}^m (S^5 \vee S^5) \\ &\simeq \Sigma^2 V \vee P^6(T) \vee P^5(T) \vee \bigvee_{i=1}^m (S^5 \vee S^5), \end{aligned}$$

where  $M_1$  is a closed 6–manifold with homology of the form (2) such that  $b_3(M_1) = 0$  and  $d = 1$ . Moreover, by Corollary 3.3 and the assumption on the ring structure of  $H^*(M; \mathbb{Z})$ , we have  $V \simeq S^2 \cup_{k\eta_2} e^4 \cup_b e^6$  for some attaching map  $b$ . Let  $\lambda = b'$  in Theorem 5.1. The theorem for the two cases when  $k$  is even then follows immediately from Proposition 5.3. For the case when  $k$  is odd, recall that there is the fibre bundle [15, Section 1.1]

$$S^2 \rightarrow \mathbb{C}P^3 \xrightarrow{\sigma} S^4,$$

with its first Pontryagin class  $p_1 = 4s_4$  where  $s_4 \in H^4(S^4; \mathbb{Z})$  is a generator. Taking the pullback of this bundle with the self-map of  $S^4$  of degree 3, we obtain the 6-manifold  $V_3$  with bundle projection  $\sigma_3$  onto  $S^4$  in the following diagram of  $S^2$ -bundles:

$$\begin{array}{ccccc} S^2 & \longrightarrow & V_3 & \xrightarrow{\sigma_3} & S^4 \\ \parallel & & \downarrow & & \downarrow 3 \\ S^2 & \longrightarrow & \mathbb{C}P^3 & \xrightarrow{\sigma} & S^4 \end{array}$$

From this diagram it is easy to see that the first Pontryagin class of  $\sigma_3$  is  $12s_4$  as required, and  $x^2 = 3y$ , where by abuse of notation  $x, y \in H^*(V_3; \mathbb{Z})$  are two generators such that  $\deg(x) = 2$  and  $\deg(y) = 4$ . Hence by Proposition 5.3,  $\Sigma V \simeq \Sigma \mathbb{C}P^3$  when  $k \equiv \pm 1 \pmod 6$  and  $\Sigma V \simeq \Sigma V_3$  when  $k \equiv 3 \pmod 6$ , and then the two decompositions when  $k$  is odd in the theorem follow.  $\square$

**Proof of Corollary 1.4** As discussed before Corollary 1.4, the number  $k \pmod 2$  and the spin condition of  $M$  are determined by the Steenrod square  $Sq^2$ . Since the attaching maps of the top cells of  $\Sigma \mathbb{C}P^3$  and  $\Sigma V_3$  are  $E\pi_2$  of order 6 and  $3E\pi_2$  of order 2, respectively, by Proposition 5.3, after localization at 3 we can consider the Steenrod power  $\mathcal{P}^1 : H^3(\Sigma M; \mathbb{Z}/3\mathbb{Z}) \rightarrow H^7(\Sigma M; \mathbb{Z}/3\mathbb{Z})$ . Then since  $\Sigma V_3 \simeq_{(3)} S^3 \vee S^5 \vee S^7$ ,  $\mathcal{P}^1$  acts trivially on its cohomology. In contrast,  $\Sigma \mathbb{C}P^3 \simeq_{(3)} S^3 \cup_{\alpha_1} e^7 \vee S^5$  with  $\alpha_1$  an element detected by  $\mathcal{P}^1$  [13, Section 1.5.5]. Hence,  $\Sigma \mathbb{C}P^3$  and  $\Sigma V_3$  can be distinguished by  $\mathcal{P}^1$ . Moreover, the stable cube element  $\eta_n^3 \in \pi_{n+3}(S^n)$  (for  $n \geq 2$ ) is detected by the secondary operation  $\mathbb{T}$  [13, Exercise 4.2.5]. Therefore the two cases of  $S^4 \cup_{\lambda \eta_4^3} e^8$  depending on  $\lambda \in \mathbb{Z}$  can be distinguished by  $\mathbb{T}$ . From the above discussions on cohomology operations, we can prove the corollary easily by the decompositions in Theorem 1.3.  $\square$

## 6 Some computations on homotopy groups of odd primary Moore spaces

In this section, we work out the homotopy groups of Moore spaces used in Section 3. Consider the Moore space  $P^{2n+1}(p^r)$  with  $n \geq 1$ ,  $p \geq 3$  and  $r \geq 1$ . We have the homotopy fibration

$$(14) \quad F^{2n+1}\{p^r\} \rightarrow P^{2n+1}(p^r) \xrightarrow{q} S^{2n+1},$$

where  $q$  is the pinch map of the bottom cell. Cohen, Moore and Neisendorfer proved the following the famous decomposition theorem:

**Theorem 6.1** [10; 24] *Let  $p$  be an odd prime. Then there is a  $p$ -local homotopy equivalence*

$$\Omega F^{2n+1}\{p^r\} \simeq_{(p)} S^{2n-1} \times \prod_{k=1}^{\infty} S^{2p^k n-1}\{p^{r+1}\} \times \Omega \Sigma \bigvee_{\alpha} P^{n_{\alpha}}(p^r),$$

where  $S^i\{p^r\}$  is the homotopy fibre of the degree map  $p^r : S^i \rightarrow S^i$ , and  $\bigvee_{\alpha} P^{n_{\alpha}}(p^r)$  is an infinite bouquet of mod  $p^r$  Moore spaces, with only finitely many Moore spaces in each dimension and the least value of  $n_{\alpha}$  being  $4n - 1$ .

We also need the following classical result:

**Lemma 6.2** [25, Proposition 6.2.2] *Let  $p$  be an odd prime. Then there is a homotopy equivalence*

$$P^m(p^r) \wedge P^n(p^r) \simeq P^{m+n}(p^r) \vee P^{m+n-1}(p^r).$$

**Lemma 6.3** [27; 28] *Let  $p$  be an odd prime. Then there are isomorphisms*

$$\pi_3(P^3(p^r)) = \mathbb{Z}/p^r\mathbb{Z} \quad \text{and} \quad \pi_n(P^n(p^r)) = 0$$

for  $n \geq 4$ .

**Proof** The cases when  $n = 3$  and  $4$  were already proved in [28, Lemma 2.1] and [27, Lemma 3.3], respectively, while the remaining cases follow immediately from the Freudenthal suspension theorem. □

**Lemma 6.4** *Let  $p$  be an odd prime. Then there is an isomorphism*

$$\pi_{n+1}(P^n(p^r)) = 0$$

for  $n \geq 3$ .

**Proof** The isomorphism  $\pi_4(P^3(p^r)) = 0$  was shown in [27, Lemma 3.3]. Let us consider  $\pi_5(P^4(p^r))$ . By the classical EHP-sequence [33, Chapter XII, Theorem 2.2], there is an exact sequence

$$0 = \pi_4(P^3(p^r)) \rightarrow \pi_5(P^4(p^r)) \xrightarrow{H} \pi_5(P^4(p^r) \wedge P^3(p^r)) \xrightarrow{P} \pi_3(P^3(p^r)) \rightarrow \pi_4(P^4(p^r)) = 0.$$

By Lemma 6.2,

$$\pi_5(P^4(p^r) \wedge P^3(p^r)) \cong \pi_5(P^6(p^r) \vee P^7(p^r)) \cong \mathbb{Z}/p^r\mathbb{Z}.$$

Hence, by Lemma 6.3 and the above exact sequence,  $P$  is an isomorphism, implying that  $\pi_5(P^4(p^r)) = 0$ . The remaining cases follow immediately from the Freudenthal suspension theorem, and this completes the proof of the lemma.  $\square$

In the remaining two lemmas, we exclude the case when  $p = 3$ .

**Lemma 6.5** *Let  $p \geq 5$ . Then there is an isomorphism*

$$\pi_{n+2}(P^n(p^r)) = 0$$

for  $n \geq 6$ .

**Proof** By the Freudenthal suspension theorem, it suffices to show  $\pi_9(P^7(p^r)) = 0$ . To do this, let us compute  $\pi_9(F^7\{p^r\})$  first. By Theorem 6.1,

$$\pi_9(F^7\{p^r\}) \cong \pi_8(\Omega F^7\{p^r\}) \cong \pi_8(S^5)_{(p)}.$$

Since  $\pi_8(S^5) \cong \mathbb{Z}/24\mathbb{Z}$  and  $p \geq 5$ , we have  $\pi_9(F^7\{p^r\}) = 0$ . Now from the exact sequence of homotopy groups of the homotopy fibration (14) (for  $n = 3$ )

$$0 = \pi_9(F^7\{p^r\}) \rightarrow \pi_9(P^7(p^r)) \rightarrow \pi_9(S^7)_{(p)} = 0,$$

we see that  $\pi_9(P^7(p^r)) = 0$ .  $\square$

**Lemma 6.6** *Let  $p \geq 5$ . The suspension morphism*

$$E: \pi_6(P^4(p^r)) \rightarrow \pi_7(P^5(p^r)) \cong \mathbb{Z}/p^r\mathbb{Z}$$

is trivial.

**Proof** On the one hand there is the EHP-sequence of  $P^4(p^r)$

$$\pi_6(P^4(p^r)) \xrightarrow{E} \pi_7(P^5(p^r)) \xrightarrow{H} \pi_7(P^5(p^r) \wedge P^4(p^r)) \rightarrow \pi_5(P^4(p^r)) = 0,$$

where  $\pi_5(P^4(p^r)) = 0$  by Lemma 6.4, and

$$\pi_7(P^5(p^r) \wedge P^4(p^r)) \cong \pi_7(P^8(p^r) \vee P^9(p^r)) \cong \mathbb{Z}/p^r\mathbb{Z}$$

by Lemma 6.2. It follows that

$$(15) \quad \pi_7(P^5(p^r))/\text{Im}(E) \cong \mathbb{Z}/p^r\mathbb{Z}.$$

On the other hand there is the EHP-sequence of  $P^5(p^r)$

$$\pi_9(P^6(p^r) \wedge P^5(p^r)) \xrightarrow{P} \pi_7(P^5(p^r)) \rightarrow \pi_8(P^6(p^r)) = 0,$$

where  $\pi_8(P^6(p^r)) = 0$  by Lemma 6.5, and

$$\pi_9(P^6(P^r) \wedge P^5(p^r)) \cong \pi_9(P^{10}(P^r) \vee P^{11}(p^r)) \cong \mathbb{Z}/p^r\mathbb{Z}$$

by Lemma 6.2. It follows that

$$(16) \quad \pi_7(P^5(p^r)) \cong \mathbb{Z}/p^r\mathbb{Z}/\text{Ker}(P).$$

By (15) and (16), we see that  $\pi_7(P^5(p^r)) \cong \mathbb{Z}/p^r\mathbb{Z}$ , and  $\text{Im}(E) = \text{Ker}(P) = 0$ .  $\square$

## References

- [1] **D Barden**, *Simply connected five-manifolds*, Ann. of Math. 82 (1965) 365–385 [MR](#) [Zbl](#)
- [2] **S Basu**, *The homotopy type of the loops on  $(n-1)$ -connected  $(2n+1)$ -manifolds*, from “Algebraic topology and related topics” (M Singh, Y Song, J Wu, editors), Springer (2019) 1–25 [MR](#) [Zbl](#)
- [3] **S Basu, S Basu**, *Homotopy groups of highly connected manifolds*, Adv. Math. 337 (2018) 363–416 [MR](#) [Zbl](#)
- [4] **V Batyrev, M Kreuzer**, *Integral cohomology and mirror symmetry for Calabi–Yau 3-folds*, from “Mirror symmetry, V” (N Yui, S-T Yau, J D Lewis, editors), AMS/IP Stud. Adv. Math. 38, Amer. Math. Soc., Providence, RI (2006) 255–270 [MR](#) [Zbl](#)
- [5] **H-J Baues**, *The degree of maps between certain 6-manifolds*, Compositio Math. 110 (1998) 51–64 [MR](#) [Zbl](#)
- [6] **P Beben, S Theriault**, *The loop space homotopy type of simply-connected four-manifolds and their generalizations*, Adv. Math. 262 (2014) 213–238 [MR](#) [Zbl](#)
- [7] **P Beben, S Theriault**, *Homotopy groups of highly connected Poincaré duality complexes*, Doc. Math. 27 (2022) 183–211 [MR](#) [Zbl](#)
- [8] **P Beben, J Wu**, *The homotopy type of a Poincaré duality complex after looping*, Proc. Edinb. Math. Soc. 58 (2015) 581–616 [MR](#) [Zbl](#)
- [9] **A L Blakers, W S Massey**, *The homotopy groups of a triad, II*, Ann. of Math. 55 (1952) 192–201 [MR](#) [Zbl](#)
- [10] **FR Cohen, J C Moore, J A Neisendorfer**, *Torsion in homotopy groups*, Ann. of Math. 109 (1979) 121–168 [MR](#) [Zbl](#)
- [11] **D Crowley, J Nordström**, *The classification of 2-connected 7-manifolds*, Proc. Lond. Math. Soc. 119 (2019) 1–54 [MR](#) [Zbl](#)
- [12] **CF Doran, J W Morgan**, *Algebraic topology of Calabi–Yau threefolds in toric varieties*, Geom. Topol. 11 (2007) 597–642 [MR](#) [Zbl](#)

- [13] **J R Harper**, *Secondary cohomology operations*, Graduate Studies in Mathematics 49, Amer. Math. Soc., Providence, RI (2002) [MR](#) [Zbl](#)
- [14] **A Hatcher**, *Algebraic topology*, Cambridge Univ. Press (2002) [MR](#) [Zbl](#)
- [15] **F Hirzebruch, T Berger, R Jung**, *Manifolds and modular forms*, Aspects of Mathematics E20, Vieweg, Brunswick (1992) [MR](#) [Zbl](#)
- [16] **R Huang**, *Homotopy of gauge groups over high-dimensional manifolds*, Proc. Roy. Soc. Edinburgh Sect. A 152 (2022) 182–208 [MR](#) [Zbl](#)
- [17] **R Huang, S Theriault**, *Loop space decompositions of  $(2n-2)$ -connected  $(4n-1)$ -dimensional Poincaré duality complexes*, Res. Math. Sci. 9 (2022) art.id.53 [MR](#) [Zbl](#)
- [18] **P E Jupp**, *Classification of certain 6-manifolds*, Proc. Cambridge Philos. Soc. 73 (1973) 293–300 [MR](#) [Zbl](#)
- [19] **M Kreck**, *On the classification of 1-connected 7-manifolds with torsion free second homology*, J. Topol. 11 (2018) 720–738 [MR](#) [Zbl](#)
- [20] **M Kreck, Y Su**, *On 5-manifolds with free fundamental group and simple boundary links in  $S^5$* , Geom. Topol. 21 (2017) 2989–3008 [MR](#) [Zbl](#)
- [21] **M Kreuzer, H Skarke**, *Complete classification of reflexive polyhedra in four dimensions*, Adv. Theor. Math. Phys. 4 (2000) 1209–1230 [MR](#) [Zbl](#)
- [22] **J Mukai**, *The  $S^1$ -transfer map and homotopy groups of suspended complex projective spaces*, Math. J. Okayama Univ. 24 (1982) 179–200 [MR](#) [Zbl](#)
- [23] **J Neisendorfer**, *Primary homotopy theory*, Mem. Amer. Math. Soc. 232, Amer. Math. Soc., Providence, RI (1980) [MR](#) [Zbl](#)
- [24] **J A Neisendorfer**, *3-Primary exponents*, Math. Proc. Cambridge Philos. Soc. 90 (1981) 63–83 [MR](#) [Zbl](#)
- [25] **J Neisendorfer**, *Algebraic methods in unstable homotopy theory*, New Mathematical Monographs 12, Cambridge Univ. Press (2010) [MR](#) [Zbl](#)
- [26] **S Oka**, review of [34], *Mathematical Reviews* (1982) Available at <http://mathscinet.ams.org/mathscinet-getitem?mr=0672527>
- [27] **T So**, *Homotopy types of gauge groups over non-simplyconnected closed 4-manifolds*, Glasg. Math. J. 61 (2019) 349–371 [MR](#) [Zbl](#)
- [28] **T So, S Theriault**, *The suspension of a 4-manifold and its applications*, preprint (2019) [arXiv 1909.11129](https://arxiv.org/abs/1909.11129)
- [29] **H Toda**, *Composition methods in homotopy groups of spheres*, Annals of Mathematics Studies 49, Princeton Univ. Press (1962) [MR](#) [Zbl](#)
- [30] **C T C Wall**, *Classification of  $(n-1)$ -connected  $2n$ -manifolds*, Ann. of Math. 75 (1962) 163–189 [MR](#) [Zbl](#)

- [31] **C T C Wall**, *Classification problems in differential topology, V: On certain 6–manifolds*, Invent. Math. 1 (1966) 355–374 [MR](#) [Zbl](#)
- [32] **C T C Wall**, *Classification problems in differential topology, VI: Classification of  $(s-1)$ –connected  $(2s+1)$ –manifolds*, Topology 6 (1967) 273–296 [MR](#) [Zbl](#)
- [33] **G W Whitehead**, *Elements of homotopy theory*, Graduate Texts in Mathematics 61, Springer (1978) [MR](#) [Zbl](#)
- [34] **K Yamaguchi**, *On the homotopy type of CW complexes with the form  $S^2 \cup e^4 \cup e^6$* , Kodai Math. J. 5 (1982) 303–312 [MR](#) [Zbl](#)
- [35] **A V Zhubr**, *Classification of simply connected six-dimensional spinor manifolds*, Izv. Akad. Nauk SSSR Ser. Mat. 39 (1975) 839–859 [MR](#) [Zbl](#) In Russian; translated in *Math. USSR-Izv.* 9 (1975) 793–812
- [36] **A V Zhubr**, *Closed simply connected six-dimensional manifolds: proofs of classification theorems*, Algebra i Analiz 12 (2000) 126–230 [MR](#) [Zbl](#) In Russian; translated in *St. Petersburg Math. J.* 12 (2001) 605–680

*Institute of Mathematics, Chinese Academy of Sciences  
Beijing, China*

[huangrz@amss.ac.cn](mailto:huangrz@amss.ac.cn)

<https://sites.google.com/site/hrzsea/>

Received: 15 April 2021      Revised: 27 September 2021

# ALGEBRAIC & GEOMETRIC TOPOLOGY

[msp.org/agt](http://msp.org/agt)

## EDITORS

### PRINCIPAL ACADEMIC EDITORS

John Etnyre  
[etnyre@math.gatech.edu](mailto:etnyre@math.gatech.edu)  
Georgia Institute of Technology

Kathryn Hess  
[kathryn.hess@epfl.ch](mailto:kathryn.hess@epfl.ch)  
École Polytechnique Fédérale de Lausanne

### BOARD OF EDITORS

|                        |  |                     |  |
|------------------------|--|---------------------|--|
| Matthew Ando           | University of Illinois<br><a href="mailto:mando@math.uiuc.edu">mando@math.uiuc.edu</a>                                   | Tsuyoshi Kobayashi  | Nara Women's University<br><a href="mailto:tsuyoshi09@gmail.com">tsuyoshi09@gmail.com</a>              |
| Julie Bergner          | University of Virginia<br><a href="mailto:jeb2md@eservices.virginia.edu">jeb2md@eservices.virginia.edu</a>               | Christine Lescop    | Université Joseph Fourier<br><a href="mailto:lescop@ujf-grenoble.fr">lescop@ujf-grenoble.fr</a>        |
| Joan Birman            | Columbia University<br><a href="mailto:jb@math.columbia.edu">jb@math.columbia.edu</a>                                    | Robert Lipshitz     | University of Oregon<br><a href="mailto:lipshitz@uoregon.edu">lipshitz@uoregon.edu</a>                 |
| Steven Boyer           | Université du Québec à Montréal<br><a href="mailto:cohf@math.rochester.edu">cohf@math.rochester.edu</a>                  | Norihiro Minami     | Nagoya Institute of Technology<br><a href="mailto:nori@nitech.ac.jp">nori@nitech.ac.jp</a>             |
| Indira Chatterji       | CNRS & Université Côte d'Azur (Nice)<br><a href="mailto:indira.chatterji@math.cnrs.fr">indira.chatterji@math.cnrs.fr</a> | Andrés Navas Flores | Universidad de Santiago de Chile<br><a href="mailto:andres.navas@usach.cl">andres.navas@usach.cl</a>   |
| Fred Cohen             | University of Rochester<br><a href="mailto:cohf@math.rochester.edu">cohf@math.rochester.edu</a>                          | Thomas Nikolaus     | University of Münster<br><a href="mailto:nikolaus@uni-muenster.de">nikolaus@uni-muenster.de</a>        |
| Alexander Dranishnikov | University of Florida<br><a href="mailto:dranish@math.ufl.edu">dranish@math.ufl.edu</a>                                  | Robert Oliver       | Université Paris-Nord<br><a href="mailto:bobol@math.univ-paris13.fr">bobol@math.univ-paris13.fr</a>    |
| Tobias Ekholm          | Uppsala University, Sweden<br><a href="mailto:tobias.ekholm@math.uu.se">tobias.ekholm@math.uu.se</a>                     | Luis Paris          | Université de Bourgogne<br><a href="mailto:lparis@u-bourgogne.fr">lparis@u-bourgogne.fr</a>            |
| Mario Eudave-Muñoz     | Univ. Nacional Autónoma de México<br><a href="mailto:mario@matem.unam.mx">mario@matem.unam.mx</a>                        | Jérôme Scherer      | École Polytech. Féd. de Lausanne<br><a href="mailto:jerome.scherer@epfl.ch">jerome.scherer@epfl.ch</a> |
| David Futер            | Temple University<br><a href="mailto:dfuter@temple.edu">dfuter@temple.edu</a>  | Peter Scott         | University of Michigan<br><a href="mailto:pscott@umich.edu">pscott@umich.edu</a>                       |
| Soren Galatius         | Stanford University<br><a href="mailto:galatius@math.stanford.edu">galatius@math.stanford.edu</a>                        | Zoltán Szabó        | Princeton University<br><a href="mailto:szabo@math.princeton.edu">szabo@math.princeton.edu</a>         |
| John Greenlees         | University of Warwick<br><a href="mailto:john.greenlees@warwick.ac.uk">john.greenlees@warwick.ac.uk</a>                  | Ulrike Tillmann     | Oxford University<br><a href="mailto:tillmann@maths.ox.ac.uk">tillmann@maths.ox.ac.uk</a>              |
| J. Elisenda Grigsby    | Boston College<br><a href="mailto:grigsbyj@bc.edu">grigsbyj@bc.edu</a>   | Maggy Tomova        | University of Iowa<br><a href="mailto:maggy-tomova@uiowa.edu">maggy-tomova@uiowa.edu</a>               |
| Ian Hambleton          | McMaster University<br><a href="mailto:ian@math.mcmaster.ca">ian@math.mcmaster.ca</a>                                    | Chris Wendl         | Humboldt-Universität zu Berlin<br><a href="mailto:wendl@math.hu-berlin.de">wendl@math.hu-berlin.de</a> |
| Hans-Werner Henn       | Université Louis Pasteur<br><a href="mailto:henn@math.u-strasbg.fr">henn@math.u-strasbg.fr</a>                           | Daniel T. Wise      | McGill University, Canada<br><a href="mailto:daniel.wise@mcgill.ca">daniel.wise@mcgill.ca</a>          |
| Daniel Isaksen         | Wayne State University<br><a href="mailto:isaksen@math.wayne.edu">isaksen@math.wayne.edu</a>                             |                     |  |

---

See inside back cover or [msp.org/agt](http://msp.org/agt) for submission instructions.


The subscription price for 2023 is US \$650/year for the electronic version, and \$940/year (+ \$70, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Algebraic & Geometric Topology is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [Current Mathematical Publications](#) and the [Science Citation Index](#).

Algebraic & Geometric Topology (ISSN 1472-2747 printed, 1472-2739 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840.

---

AGT peer review and production are managed by EditFlow<sup>®</sup> from MSP.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2023 Mathematical Sciences Publishers



# ALGEBRAIC & GEOMETRIC TOPOLOGY

Volume 23

Issue 1 (pages 1–508)

2023

---

|   |     |
|---|-----|
| Proper 2–equivalences between infinite ended finitely presented groups                      | 1   |
| MANUEL CÁRDENAS, FRANCISCO FERNÁNDEZ LASHERAS, ANTONIO QUINTERO and RANJA ROY               |     |
| Groups acting on CAT(0) cube complexes with uniform exponential growth                      | 13  |
| RADHIKA GUPTA, KASIA JANKIEWICZ and THOMAS NG   |     |
| Admissible replacements for simplicial monoidal model categories                            | 43  |
| HALDUN ÖZGÜR BAYINDIR and BORIS CHORNY  |     |
| The bridge number of arborescent links with many twigs                                      | 75  |
| SEBASTIAN BAADER, RYAN BLAIR, ALEXANDRA KJUCHUKOVA and FILIP MISEV                          |     |
| $G_\infty$ –ring spectra and Moore spectra for $\beta$ –rings                               | 87  |
| MICHAEL STAHLHAUER  |     |
| External Spanier–Whitehead duality and homology representation theorems for diagram spaces  | 155 |
| MALTE LACKMANN  |     |
| On the surjectivity of the $\mathrm{tmf}$ –Hurewicz image of $A_1$                          | 217 |
| VIET-CUONG PHAM   |     |
| The handlebody group and the images of the second Johnson homomorphism                      | 243 |
| QUENTIN FAES  |     |
| Recognition of connective commutative algebra spectra through an idempotent quasiadjunction | 295 |
| RENATO VASCONCELLOS VIEIRA  |     |
| $SL_2$ quantum trace in quantum Teichmüller theory via writhe                               | 339 |
| HYUN KYU KIM, THANG T Q LÊ and MIRI SON   |     |
| Constraints on families of smooth 4–manifolds from $\mathrm{Pin}^-(2)$ –monopole            | 419 |
| HOKUTO KONNO and NOBUHIRO NAKAMURA  |     |
| Suspension homotopy of 6–manifolds  | 439 |
| RUIZHI HUANG  |     |
| Time-periodic solutions of Hamiltonian PDEs using pseudoholomorphic curves                  | 461 |
| OLIVER FABERT and NIEK LAMOREE  |     |