

AG
T

*Algebraic & Geometric
Topology*

Volume 23 (2023)

Uniform foliations with Reeb components

JOAQUÍN LEMA



Uniform foliations with Reeb components

JOAQUÍN LEMA

A foliation on a compact manifold is uniform if each pair of leaves of the induced foliation on the universal cover are at finite Hausdorff distance from each other. We study uniform foliations with Reeb components. We give examples of such foliations on a family of closed 3–manifolds with infinite fundamental group. Furthermore, we prove some results concerning the behavior of a uniform foliation with Reeb components on general 3–manifolds.

57R30

1 Introduction

Consider a foliation \mathcal{F} on a compact Riemannian 3–manifold M . This foliation lifts to a foliation $\tilde{\mathcal{F}}$ on the universal cover \tilde{M} . We will say that \mathcal{F} is *uniform* if any pair of leaves of $\tilde{\mathcal{F}}$ are at finite Hausdorff distance from each other.

A lot can be said about a uniform foliation if we further assume \mathcal{F} to be Reebless; see for example Fenley and Potrie [6] and Thurston [17]. In this paper, we will focus on the opposite case. More precisely, we will study the following question posed by Fenley and Potrie [6, Question 1]:

Question 1.1 If \mathcal{F} is uniform in M with infinite fundamental group, does it follow that \mathcal{F} is also Reebless?

Our first result will be to give a negative answer to the question (see Section 4):

Theorem 1.2 For every $l, m \in \mathbb{N}$ and every choice M_1, \dots, M_l of 3–manifolds with finite fundamental group, there exists a uniform foliation with Reeb components on $M = (\#_{i=1}^m S^1 \times S^2) \# (\#_{i=1}^l M_i)$.

One can arrange the foliations given by Theorem 1.2 to be C^∞ –smooth; see Remark 4.3. We will also give an example of uniform foliation with Reeb components on the solid torus, which is trivial on the boundary. However, these 3–manifolds are “small” in the

sense that they are the only ones not admitting an immersed essential closed surface of genus $g \geq 1$; see [Proposition 2.5](#). In this paper, we will say that an immersed surface is *essential* if the immersion induces an injective morphism from the fundamental group of the surface to $\pi_1(M)$.

We can say the following for the rest of the 3–manifolds:

Theorem 1.3 *Let \mathcal{F} be a uniform foliation on a compact 3–manifold M admitting an immersion $i: \Sigma \rightarrow M$ from a closed surface Σ of genus $g \geq 1$ such that $i_*: \pi_1(\Sigma) \rightarrow \pi_1(M)$ is injective. Then $i(\Sigma)$ must intersect the set of Reeb components.*

The idea of the proof is to put the immersion in general position with respect to the foliation, and then apply Poincaré–Bendixson theory on the induced foliation on $\tilde{\Sigma} \cong \mathbb{R}^2$.

[Theorem 1.3](#) and some remarks in [Section 3](#) motivate us to modify [Question 1.1](#): does an *irreducible* 3–manifold with infinite fundamental group admit a uniform foliation with Reeb components?

Acknowledgements I am profoundly grateful to my advisor Rafael Potrie for his patience and suggestions during every step of my thesis. This work could not be done without him. I would also like to thank Sergio Fenley for his comments on earlier versions of this paper, which motivated the statement of [Theorem 1.3](#).

The author was supported by a CAP grant for Master’s students at UdelaR.

2 Preliminaries

2.1 Foliations on 3–manifolds

We will work on a compact 3–manifold M endowed with a $C^{\infty,0+}$ codimension-one foliation \mathcal{F} . From now on, by foliation on a 3–manifold we refer to a foliation of codimension one. We will assume some familiarity with foliation theory; the reader may find a comprehensive treatment of the subject in [\[3; 4\]](#) and [\[9; 10\]](#).

A Reeb component is a foliation of the solid torus such that the boundary is a leaf, and there is a circle’s worth of planar leaves in the interior spiraling towards the boundary torus. We will also call some quotient of this foliation a Reeb component. These are crucial in the study of foliations on 3–manifolds by Novikov’s celebrated theorem [\[11\]](#).

It says (among other things) that if \mathcal{F} is orientable and transversely orientable, then it is Reebless if and only if every leaf L is essential (as defined in the introduction). In this sense, for Reebless foliations, the topology of a leaf is tied to the topology of the manifold.

Considering this fact, it is natural to ask if there exists some property of a foliation \mathcal{F} with Reeb components “reading” the topology of the 3–manifold. This does not seem plausible, as every 3–manifold admits a foliation with Reeb components. Furthermore, Thurston shows in [16] that every plane field on M is homotopic to the tangent space of a foliation. His construction is local in nature, so we do not care about the global topology of M . The Reeb components play a key role because holes can easily be filled using them. See [4, Section 8.5] for a detailed treatment of this construction.

However, we believe that if a compact 3–manifold M is “big enough”, then it does not admit a *uniform* foliation with Reeb components; see Section 3. In particular, Lemma 3.1 would tell us that every foliation on these manifolds must have leaves lifting to unbounded sets on the universal cover. This would be interesting because not much is known about the behavior of a foliation with Reeb components on a general 3–manifold.

Suppose that M is equipped with a Riemannian metric, and let us denote the universal cover of M by $p: \tilde{M} \rightarrow M$, equipping \tilde{M} with the pullback metric.

Definition 2.1 Let \mathcal{F} be a foliation on a compact manifold M , and denote the lifted foliation on \tilde{M} by $\tilde{\mathcal{F}}$. We will say that \mathcal{F} is *uniform* if any pair of leaves of $\tilde{\mathcal{F}}$ are at finite Hausdorff distance from each other.

This notion was introduced by Thurston in [17], who further required \mathcal{F} to be Reebless; see also [2, Section 9.3]. We stick to the definition given by S Fenley and R Potrie in [6].

2.2 Turbulization

To prove Theorem 1.2, we will rely on a method for modifying foliations along a transverse curve known as *turbulization*.

Suppose that a 3–manifold M is endowed with an oriented and cooriented foliation \mathcal{F} admitting an embedded closed curve τ transverse to the foliation. Let $\overline{N(\tau)}$ be the closure of an embedded tubular neighborhood of τ . Then the orientability and coorientability of \mathcal{F} implies that if $N(\tau)$ is small enough, $\mathcal{F}|_{\overline{N(\tau)}}$ is homeomorphic to the product foliation by disks on the solid torus $\overline{N(\tau)}$.




Figure 1: Left: some leaves of the foliation \mathcal{T} . Right: the function λ controlling the vector field normal to $\ker \omega$.

Fix an identification of $\overline{N(\tau)}$ with $D^2 \times S^1$ sending the leaves of $\mathcal{F}|_{\overline{N(\tau)}}$ to $D^2 \times \{\cdot\}$. We will denote points in $D^2 \times S^1$ using cylindrical coordinates, by which we mean that (r, θ, z) represents a point whose projection to D^2 has polar coordinates (r, θ) and which projects to $z \in S^1$. Notice that the plane field tangent to the foliation by disks is given by the kernel of the 1-form $\alpha_0 = dz$.

Now we will construct a new foliation \mathcal{T} on $D^2 \times S^1$ with a Reeb component on the interior and coinciding with the foliation by disks on a neighborhood of the boundary torus. This will give us a new foliation \mathcal{F}' on M , defined as \mathcal{F} in the complement of $\overline{N(\tau)}$ and as \mathcal{T} on $\overline{N(\tau)}$.

Take $\lambda: [0, 1] \rightarrow [-\frac{\pi}{2}, \frac{\pi}{2}]$ to be a smooth function which is a strictly increasing bijection restricted to the interval $[0, \frac{3}{4}]$, and satisfies $\lambda(\frac{2}{3}) = 0$ and $\lambda|_{[3/4, 1]} = \frac{\pi}{2}$; see Figure 1, right. Now define the 1-form

$$\omega_{(r,\theta,z)} = \cos(\lambda(r)) dr + \sin(\lambda(r)) dz.$$

This is well-defined on $D^2 \times S^1$ if $\lambda^{(k)}(0) = 0$ for every $k \geq 1$. Using Frobenius' theorem, one can check that $\pi = \ker \omega$ is an integrable plane field. The foliation tangent to π is the desired foliation \mathcal{T} depicted at the left of Figure 1.


If we are careful with the choice of λ , it can be shown that the resulting foliation \mathcal{F}' on M is as regular as \mathcal{F} . The reader can find a detailed description of this construction in [3, Example 3.3.11].

2.3 General position

Let \mathcal{F} be a foliation on a 3-manifold M and $i: \Sigma \rightarrow M$ an immersion. We can always take a small perturbation of i in order to suppose that $i(\Sigma)$ is “as transverse as possible” to the foliation.

Definition 2.2 Let Σ be a closed surface and let $i : \Sigma \rightarrow M$ be an immersion into a 3-manifold M endowed with a foliation \mathcal{F} . We will say that *the immersion is in general position with respect to \mathcal{F}* if the following happens:

- (1) Except at a finite set of points $\{p_1, \dots, p_l\} \subset \Sigma$, i is transverse to \mathcal{F} .
- (2) The points $\{i(p_1), \dots, i(p_l)\}$ lie on distinct leaves of \mathcal{F} .
- (3) For every p_k and every submersion ϕ defined on a neighborhood around $i(p_k)$ locally defining the foliation¹ sending $i(p_k)$ to zero, there exists a neighborhood U_k of p_k such that $\phi \circ i|_{U_k}$ is topologically conjugated to f_c or f_s , where $f_c(x, y) = x^2 + y^2$ and $f_s(x, y) = x^2 - y^2$. By this we mean that there exists a homeomorphism between U_k and some neighborhood U'_k of $0 \in \mathbb{R}^2$ such that the diagram



commutes, where σ may be c or s .

Geometrically the last condition says that the tangencies between Σ and the foliation look like a critical point of a Morse function on Σ (up to homeomorphism); see Figure 2.

The following theorem dates back to Haefliger when the foliation \mathcal{F} is sufficiently regular. It was generalized to the case of C^0 foliations by Solodov in [14].

Theorem 2.3 *Let $i : \Sigma \rightarrow M$ be an immersion of a closed surface Σ on a 3-manifold M endowed with a oriented and cooriented foliation \mathcal{F} . Then for every $\varepsilon > 0$, there exists an immersion $j : \Sigma \rightarrow M$ in general position with respect to \mathcal{F} and ε -close to i in the C^0 -topology.*

Suppose that \mathcal{F} is oriented and cooriented and $i : \Sigma \rightarrow M$ is an immersion in general position with respect to \mathcal{F} . The first condition of Definition 2.2 tells us that \mathcal{F} induces a foliation on $\Sigma \setminus \{p_1, \dots, p_l\}$ whose leaves are intersections of leaves of \mathcal{F} and $i(\Sigma)$. The third condition gives us a model neighborhood around a singularity; more precisely, they look like saddles in the case of f_s , or centers in the case of f_c ; see Figure 2.

¹By this we mean a map $\phi : U \rightarrow \mathbb{R}$ such that the preimages of the regular values are disks contained in a leaf.




Figure 2: Bottom: tangencies of a surface in general position (in yellow) and the foliation (in green). Top: the induced singular foliation on the surface. The orientation and coorientation of \mathcal{F} induces an orientation on the singular foliation.

These local models give us what is called a *singular foliation with Morse singularities* on Σ . The orientation of \mathcal{F} induces an orientation on this singular foliation (this is an orientation outside its singular points).

The second condition tells us that a separatrix cannot join two distinct singularities; see below for a definition of separatrix.

2.4 Singular foliations of Morse type

In this section, we will fix some notation that we are going to use throughout this text. Let Σ be a surface endowed with an oriented singular foliation with Morse singularities \mathcal{G} . Suppose that p is some nonsingular point; we will denote the leaf passing through p by L_p . If L_p is noncompact (ie not a circle) then p separates L_p into two components L_p^+ and L_p^- , where L_p^+ is composed of points greater than p with respect to the order imposed by the orientation, and L_p^- of points smaller than p .

If L is some noncompact leaf, we will say that the ω -limit of the leaf is the set $\omega(L) = \bigcap_{p \in L} \overline{L_p^+}$. Analogously we define the α -limit of the leaf as $\alpha(L) = \bigcap_{p \in L} \overline{L_p^-}$. A leaf L is a *separatrix* if the α - or ω -limit of L is a singularity.

Let C be a union of singularities and separatrices S_i such that the α -limit and ω -limit of S_i are singularities in C . This set defines a directed graph with one vertex for each

singularity in C and one edge for each S_i , with the orientation induced by the foliation. We will say that C is a *closed graph* if there exists a closed path on the graph which travels through every edge only once (respecting the orientation).

We will end up studying Morse-type singular foliations on \mathbb{R}^2 . In this context, we can apply the classical Poincaré–Bendixson theorem (see for instance [12, Theorem 1.8]):

Theorem 2.4 (Poincaré–Bendixson) *Let \mathcal{G} be a Morse type singular foliation on \mathbb{R}^2 and L a leaf with compact closure. Then $\alpha(L)$ and $\omega(L)$ can be a saddle singularity, a closed leaf or a closed graph.*

2.5 Essential immersions

Let M be some closed 3–manifold. We will say that a 2–sided embedded closed surface Σ is *compressible* if there exists some embedded disk D on M such that $D \cap \Sigma = \partial D$ and ∂D is not homotopically trivial in Σ . We will call D a *compressing disk* for Σ . A 2–sided embedded surface Σ of genus $g \geq 1$ is *incompressible* if there are no compressing disks.

If a 2–sided embedded surface Σ is compressible, we can do surgery on a compressing disk D in order to obtain a simpler surface [8, Section 1.2]. This operation preserves the homology class $[\Sigma] \in H_2(M, \mathbb{Q})$. Doing finitely many surgeries on Σ , we obtain an embedded 2–sided surface, whose connected components are incompressible surfaces or spheres. The sum of the homology class of the connected components is the homology class of Σ .

The disk theorem [8, Theorem 3.1] tells us that a 2–sided embedded surface Σ is incompressible if and only if the embedding $i : \Sigma \rightarrow M$ induces an injective morphism $i_* : \pi_1(\Sigma) \rightarrow \pi_1(M)$, ie the surface is essentially embedded.

Thanks to the virtual Haken conjecture proved by Agol in [1], we can say exactly which closed 3–manifolds admit essentially embedded closed surfaces of genus $g \geq 1$:

Proposition 2.5 *A closed 3–manifold M admits an essentially immersed surface Σ of genus $g \geq 1$ if and only if some factor of the prime decomposition of M is irreducible with an infinite fundamental group.*

Proof The “if” part is a direct consequence of the virtual Haken conjecture. Suppose P is a factor of the prime decomposition, which is irreducible with an infinite fundamental

group. Then it has a finite cover \hat{P} which admits an embedded incompressible surface of genus $g \geq 1$. Projecting this surface to P and avoiding the balls of the connected sums, we obtain an essential immersion on M .

To see the “only if” part, it is enough to show that the fundamental group of the remaining 3–manifolds does not admit a subgroup isomorphic to the fundamental group of a closed surface. No factor of the prime decomposition of M is irreducible with an infinite fundamental group. Therefore, factors can be irreducible with a finite fundamental group or $S^1 \times S^2$ (prime but not irreducible). The fundamental group of such a 3–manifold must be a free product of infinite cyclic and finite groups.

If the fundamental group of a closed surface Γ were isomorphic to a subgroup of such a group, then the Kurosh subgroup theorem [13, Section 1.5.5] tells us that Γ is isomorphic to a free product of infinite cyclic groups and finite groups. However, the fundamental group of a closed surface is freely indecomposable; this means that it is not isomorphic to a free product of two nontrivial groups (this follows from Stallings’ end theorem [15, Section 4.A.6]). Therefore it would be isomorphic to a cyclic group or a finite group. Both options are impossible. \square

We remark that the last proof shows that the only closed 3–manifolds which do not admit an essential immersion are those where we can use [Theorem 1.2](#).

3 Some remarks

We begin with some general facts about uniform foliations *with Reeb components* on 3–manifolds. We will assume that M is equipped with some Riemannian metric; this induces a metric on the universal cover $p: \tilde{M} \rightarrow M$.

The following was observed by Fenley and Potrie in [6]; we include the proof for completeness.

Lemma 3.1 *Let \mathcal{F} be a uniform foliation with Reeb components on a compact manifold M . Then every lift of a leaf to the universal cover \tilde{M} has compact closure. In particular, the inclusion of a leaf $i: L \rightarrow M$ induces a morphism $i_*: \pi_1(L) \rightarrow \pi_1(M)$ with finite image.*

Proof It is enough to show that a boundary leaf of some Reeb component lifts to a compact leaf because any pair of leaves of the foliation $\tilde{\mathcal{F}}$ are at finite Hausdorff

distance from each other. So assume for the sake of contradiction that the lift of a boundary leaf of some Reeb component R is noncompact. Let $q \in R$ and $\gamma \subset R$ be a loop at q homotopic to the core of the Reeb component. The homotopy class of this curve is nontrivial; otherwise, R would lift to a compact set. Choose some lift of γ starting at $\tilde{q} \in p^{-1}(q)$, we claim that $d_{\tilde{M}}(\tilde{q}, \gamma^n(\tilde{q})) \rightarrow_n \infty$.

One way to see this is to use that the map $\alpha \in \pi_1(M) \rightarrow \alpha(\tilde{p})$ is a quasi-isometry (by the Milnor–Švarc lemma). This fact implies that the distance between a lift of a plane on the interior of the Reeb component and $\gamma^n(\tilde{q})$ goes to infinity with n , contradicting the uniform condition.

To see that the inclusion of a leaf $i: L \rightarrow M$ induces a morphism with finite image, suppose for the sake of contradiction that $\#(i_*\pi_1(L)) = \infty$. Then the cardinality of the stabilizer of a connected component \tilde{L} of $p^{-1}(L)$ is infinite. In particular, the orbit of every $p \in \tilde{L}$ is infinite. But the action is proper, so \tilde{L} escapes every compact set, contradicting the last paragraph. \square

This lemma motivates us to look for counterexamples in foliations such that every leaf of $\tilde{\mathcal{F}}$ has compact closure. The following gives us a criterion for a foliation to verify this condition.

Lemma 3.2 *Let \mathcal{F} be a foliation on a compact manifold M . Suppose that there exist compact leaves A_1, \dots, A_k such that every connected component of the complement of $\bigcup_{i=1}^k p^{-1}(A_i)$ has compact closure. Then \mathcal{F} is uniform.*

Proof Take a leaf L of $\tilde{\mathcal{F}}$ different from a lift of some A_i . Then some connected component of the complement of $\bigcup_{i=1}^k p^{-1}(A_i)$ contains L because the boundaries of these regions are composed of leaves. Therefore, the leaf has compact closure because each one of these regions is compact. \square

This remark motivated our first counterexamples to [Question 1.1](#); see [Constructions 4.2](#) and [4.10](#). However, the reader may notice that the existence of these compact leaves imposes conditions on the topology of the 3-manifold M . [Lemma 3.1](#) allows us to use compression disks on compact leaves until we get spheres on M . These spheres must bound some topology in order for us to use [Lemma 3.2](#).

The following is a consequence of this idea.

Lemma 3.3 *Let \mathcal{F} be an oriented and cooriented uniform foliation with Reeb components on a closed, irreducible 3–manifold M with an infinite fundamental group. Then every compact leaf is a torus, bounding a solid torus on one of its sides.*

Proof The inclusion i of a compact leaf L induces the zero morphism $i_*: \pi_1(L) \rightarrow \pi_1(M)$, because the fundamental group of a closed, irreducible 3–manifold is torsion-free. We can compress the surface L using the disk theorem until we get a union of spheres S . The compression operation preserves the class $[L] \in H_2(M; \mathbb{Q})$. However, S is composed of spheres and M is irreducible, therefore $[L] = [S] = 0$.

This implies that the leaf L cannot admit a closed transversal τ , because if it did then the intersection product between $[\tau] \in H_1(M; \mathbb{Q})$ and $[L] \in H_2(M; \mathbb{Q})$ would be nontrivial and $[L] \neq 0$. A theorem of Goodman (see [3, Theorem 6.3.5]) implies that L must be a torus. To see that it bounds a solid torus, do surgery on L with some compressing disk to get a sphere. This sphere bounds a ball on one side, so we obtain a solid torus bounded by L by reversing the compression operation. \square

Lemmas 3.1 and 3.3 paint a strange picture. For instance, if the universal cover is \mathbb{R}^3 then the foliation fills all the space, but every compact leaf cannot bound topology. An exceptional minimal set is not much worse because they must have compact closure. On the other hand, our main theorem, Theorem 1.3, tells us that the set of Reeb components must intersect every essentially immersed surface. Figure 3 shows a possible picture for the set of Reeb components on a uniform foliation on \mathbb{T}^3 .

Question 3.4 *If \mathcal{F} is uniform in an irreducible 3–manifold M with an infinite fundamental group, does it follow that \mathcal{F} is also Reebless?*

4 Constructions

We begin with an explicit counterexample to Question 1.1 on $S^1 \times S^2$. To do so, consider the characterization of $S^1 \times S^2$ given by the following remark.

Remark 4.1 The manifold $S^1 \times S^2$ is homeomorphic to M , which is constructed as follows. Start with the solid torus $S^1 \times D^2$ and a circle K , which is the boundary of an embedded disk D . Identify a tubular neighborhood of the knot K with $S^1 \times D^2$ in such a way that the longitude $S^1 \times \{\cdot\}$ bounds an embedded disk in the complement of the neighborhood. Drilling this tubular neighborhood, we obtain a compact 3–manifold N with two boundary components homeomorphic to $S^1 \times S^1$. Identifying both boundary components via the identity map, we obtain the closed 3–manifold M .




Figure 3: Can we have a uniform foliation on \mathbb{T}^3 so that the Reeb components are the ones depicted in this image?

To prove this, notice that $S^1 \times S^2$ is homeomorphic to the quotient of $\mathbb{R}^3 \setminus \{0\}$ induced by the properly discontinuous \mathbb{Z} -action generated by the map $f(x) = \frac{1}{2}x$. If S^2 is the standard sphere, the set bounded by S^2 and $f(S^2)$ is a fundamental domain. Now consider a torus S resulting from S^2 by adding a small handle. Notice that $f(S)$ is contained in the solid torus bounded by S and containing 0. Furthermore, the set N bounded by S and $f(S)$ is also a fundamental domain for the action. Therefore, $S^1 \times S^2$ is homeomorphic to N , where we identify both boundary components via the map f . This identification coincides with the identification defining M .

We will be referring to a curve K on a 3-manifold which is the boundary of an embedded disk as an *unknot*. We will always equip an unknot with a frame such that the longitude of the tubular neighborhood is the boundary of some embedded disk in the complement of the neighborhood.

The following construction is motivated by [Lemma 3.2](#).

Construction 4.2 We will construct a uniform foliation with Reeb components \mathcal{F} on $S^1 \times S^2$ which admits a transversal unknot.

Thanks to [Remark 4.1](#), it is enough to find a foliation tangential to the boundary of the manifold N obtained from $S^1 \times D^2$ by drilling a tubular neighborhood of an unknot.




Figure 4: The foliation on N before turbulizing along τ .

To see this, notice that $S^1 \times S^2$ is obtained from N by identifying the boundary components. The foliation on N projects to a foliation of $S^1 \times S^2$ with a toric leaf S coming from the boundary components of N . We saw in [Remark 4.1](#) that the lift of S to the universal cover $\mathbb{R}^3 \setminus \{0\}$ bounds a compact fundamental domain. Therefore this foliation falls under the hypothesis of [Lemma 3.2](#). In particular, it is uniform.

We may construct a foliation tangential to the boundary on N as follows: let D^2 be the unit disk on \mathbb{R}^2 and $A \subset D^2$ be the annuli with inner and outer radii $\frac{1}{2}$ and 1, respectively. Equip A with a Reeb component and consider the product foliation induced on $S^1 \times A \subset S^1 \times D^2$. We can complete this foliation to $S^1 \times D^2$ by adding a Reeb component on $S^1 \times (D^2 \setminus A)$. Notice that the core of a Reeb annulus is transversal to the foliation; this induces a transversal τ to the foliation on the solid torus bounding a disk. Turbulizing along τ and drilling the newly generated Reeb component, we obtain the desired foliation on N ; see [Figure 4](#).

Remark 4.3 The constructed foliation can be made C^∞ . We just have to choose the Reeb annulus and the turbulization process in such a way that the resulting foliation on N has C^∞ -trivial holonomy on the boundary; see [[3](#), Definition 3.4.1]. The reader may check that each construction on this section can be made C^∞ . We will not pay attention to this.

Now we describe a procedure that will allow us to start with a pair of foliated 3-manifolds M and N and construct an explicit foliation on its connected sum. Furthermore, if we look at M and N with balls removed inside $M \# N$, the foliation on the connected sum coincides with the original pair of foliations outside the boundary of the drilled balls. The idea comes from the following remark.

Remark 4.4 Let M_1 and M_2 be a pair of 3-manifolds. For $i = 1, 2$, consider an unknot K_i on each M_i contained in a certain 3-ball $B_i \subset M_i$ — for example, a tubular neighborhood of the spanned disk. Drilling a tubular neighborhood of K_i contained on B_i from M_i , we obtain N_i . Identifying this tubular neighborhood with $S^1 \times D^2$ via the standard framing, we get an identification between the boundary component of N_i coming from K_i and $S^1 \times S^1$.

We claim that if we start from $N_1 \cup N_2$ and we identify the boundaries of the drilled tubular neighborhoods of K_1 and K_2 via the map $f: S^1 \times S^1 \rightarrow S^1 \times S^1$ defined as $f(x, y) = (y, x)$, we obtain a 3-manifold M homeomorphic to the connected sum $M_1 \# M_2$. To prove this claim, notice that if we start with the balls B_i , drill an unknot K_i from each of them, and sew the resulting sets along the boundary tori according to the map f , then we obtain S^3 with a pair of 3-balls removed, ie $S^2 \times [0, 1]$. Therefore $M \cong M_1 \# S^3 \# M_2$, as desired.

Lemma 4.5 *Let M_1 and M_2 be compact 3-manifolds endowed with foliations \mathcal{F}_1 and \mathcal{F}_2 . Suppose that there exists an unknot K_i in M_i , transverse to \mathcal{F}_i and contained in a ball B_i . Then there exists a foliation \mathcal{F} on $M_1 \# M_2$ admitting a transversal unknot and coinciding with \mathcal{F}_i on $M_i \setminus B_i$. Furthermore if \mathcal{F}_1 and \mathcal{F}_2 are uniform, then \mathcal{F} is also uniform.*

Proof We start by turbulizing \mathcal{F}_i along the transversal K_i . This process can be done by letting the foliation \mathcal{F}_i fixed outside the ball B_i . Drilling the newly generated Reeb component, we obtain a 3-manifold N_i equipped with a foliation tangential to the component S_i of ∂N_i coming from the Reeb component. Identifying $S_i \subset \partial M_i$ as in Remark 4.4, we obtain the desired foliation \mathcal{F} on $M_1 \# M_2$. Notice that the leaf obtained by identifying S_i is a torus S such that the inclusion $i: S \rightarrow M_1 \# M_2$ induces the zero morphism on the fundamental group. Near this leaf, there are a lot of transversal unknots resulting from the turbulization process.

Now we will see that this construction preserves the uniform condition. Notice that as \mathcal{F}_i admits a homotopically trivial transversal, it must have Reeb components by Novikov's theorem; see, for instance, [4, Theorem 9.1.4]. Therefore, Lemma 3.1 tells us that every leaf of the induced foliation $\tilde{\mathcal{F}}_i$ on the universal cover \tilde{M}_i has compact closure.

Let N_i be as above and consider the cover $q_i: \hat{N}_i \rightarrow N_i$ defined as \tilde{M}_i with the solid tori bounded by lifts of S_i drilled, and the projection $q_i = p_i|_{p_i^{-1}(N_i)}$, where $p_i: \tilde{M}_i \rightarrow M_i$

is the universal cover of M_i . We claim that if $p: \tilde{M} \rightarrow M_1 \# M_2$ is the universal cover of the connected sum, then each connected component of $p^{-1}(N_i)$ is homeomorphic to \hat{N}_i (notice that each N_i is naturally embedded in $M_1 \# M_2$).

To see this, it suffices to notice that if $j_i: N_i \subset M_i \hookrightarrow M_1 \# M_2$ is the inclusion on the connected sum, then $\ker(j_i)_*$ is isomorphic to $\pi_1(\hat{N}_i)$, by Galois correspondence. By Van Kampen's theorem $\pi_1(N_i) \cong \pi_1(M_i) \# \langle m_i \rangle$, where m_i is the homotopy class of a meridian of the drilled solid torus on M_i . The morphism $(j_i)_*$ sends each word γ of $\pi_1(N_i)$ to the word obtained by deleting all appearances of $\langle m_i \rangle$ from γ , because $i_*: \pi_1(S) \rightarrow \pi_1(M_1 \# M_2)$ is zero. Therefore, $\ker(j_i)_*$ is the normalizer of $\langle m_i \rangle$, which is exactly the image of $\pi_1(\hat{N}_i) \subset \pi_1(N_i)$.

To conclude the lemma, notice that every leaf apart from S is either contained in N_1 or N_2 . In any case, the lift of each of those leaves cannot leave a connected component of $p^{-1}(N_i)$ because its boundary is composed of leaves. The restricted foliation on this connected component is the foliation on $\hat{N}_i \subset \tilde{M}_i$. Therefore every leaf has compact closure, and the foliation is uniform. \square

Lemma 4.5 and **Construction 4.2** gives us the following.

Corollary 4.6 *Given $k \in \mathbb{N}$, there exists a uniform foliation \mathcal{F} with Reeb components on $\#_{i=1}^k S^1 \times S^2$. Furthermore, this foliation admits a transverse unknot.*

Every foliation on a compact 3-manifold with a finite fundamental group is uniform because the universal cover is compact. Furthermore, it has Reeb components by Novikov's theorem. We will see that we can construct such foliations in a way that we fall under the hypothesis of **Lemma 4.5**.

For the next construction, we will use the notion of *open book decomposition*. An open book decomposition on a closed 3-manifold M is a pair (B, π) , where B is an oriented link on M (called the *binding*) and $\pi: M \setminus B \rightarrow S^1$ is a fibration. We ask the fibers $\pi^{-1}(\cdot)$ to be the interior of compact surfaces whose boundary is B ; these are called the *pages* of the open book. There is a classical theorem (due to Alexander) saying that every closed 3-manifold has an open book decomposition; see [5] for a nice introduction to these objects.

Construction 4.7 We will construct a (uniform) foliation \mathcal{F} on every closed 3-manifold M with a finite fundamental group, such that there exists a torus transverse to \mathcal{F} which bounds a solid torus T . In particular, every meridian of ∂T is an unknot.

To see this, we will briefly recall the argument which shows that every closed 3-manifold admits a foliation. Choose an open book decomposition (B, π) on M . Let T be a tubular neighborhood of B and identify every connected component of T with $S^1 \times D^2$. With this identification, define a tubular neighborhood $T' \subset T$ whose connected components are $S^1 \times D'^2 \subset S^1 \times D^2$, where D'^2 is an open disk properly contained on D^2 .

On $M \setminus T'$ we have a foliation transverse to the boundary defined by the fibers of the fibration $\pi: M \setminus B \rightarrow S^1$. Spinning this foliation along the boundary and filling the boundary with a Reeb component (see for instance [2, Example 4.10]), we obtain a foliation on M . Notice that ∂T is composed of a torus transversal to the constructed foliation, bounding a solid torus.

Construction 4.7 and Corollary 4.6 immediately give us Theorem 1.2.

Corollary 4.8 *Given $k \in \mathbb{N}$ and a choice M_1, \dots, M_l of 3-manifolds with finite fundamental group, there exists a uniform foliation with Reeb components on*

$$\left(\#_{i=1}^k S^1 \times S^2 \right) \# \left(\#_{i=1}^l M_i \right).$$

We will finish this section by sketching an example of a uniform foliation on the solid torus inducing the trivial foliation by meridians on the boundary. To do so, we will use the following well-known lemma. A visual proof can be found in [7, Section 5.3].

Lemma 4.9 *Identify \mathbb{T}^3 as the quotient $\mathbb{R}^3/\mathbb{Z}^3$, and let T_x, T_y and T_z be disjoint tubular neighborhoods of curves on \mathbb{T}^3 which are projections of lines parallel to the x -, y - and z -axes, respectively. Then, if L is the Borromean link on S^3 and N is a tubular neighborhood of L , there exists a homeomorphism*

$$h: \mathbb{T}^3 \setminus (T_x \cup T_y \cup T_z) \rightarrow S^3 \setminus N.$$

Furthermore, the homeomorphism takes a meridian of T_x, T_y and T_z to a longitude of a component of the Borromean ring, ie a curve on a component of N bounding a disk on S^3 .

Construction 4.10 We will construct a uniform foliation with Reeb components on the solid torus $S^1 \times D^2$, inducing a foliation by closed curves of meridional slope on the boundary.




Figure 5: Left: a foliation \mathcal{F}_1 on \mathbb{T}^3 transverse to curves parallel to the x -, y - and z -axes. Right: the image of a compact leaf via h on the complement of the Borromean rings.

Start with \mathbb{T}^3 endowed with a foliation \mathcal{F}_1 , as depicted in Figure 5. This foliation is transversal to three closed curves c_x , c_y and c_z lifting to \mathbb{R}^3 as lines parallel to the x -, y - and z -axes, respectively. Take tubular neighborhoods T_x , T_y and T_z of c_x , c_y and c_z , respectively, where the foliation restricts to a trivial foliation by disks.

By Lemma 4.9, there is a homeomorphism $h: \mathbb{T}^3 \setminus (T_x \cup T_y \cup T_z) \rightarrow S^3 \setminus N$, with N a tubular neighborhood of the Borromean link. Notice that $S^3 \setminus N$ is homeomorphic to a solid torus, with the tubular neighborhood of two linked unknots drilled as to the right of Figure 5. To see this, look at the complement of the tubular neighborhood of one link component.

The homeomorphism h sends the foliation on \mathbb{T}^3 to a foliation \mathcal{F}_2 on the solid torus minus the link. This foliation is transverse to the boundary: it induces on the boundary of the solid torus the trivial foliation by meridians and the trivial foliation by longitudes around the link components. Spiraling around the link in the interior of the solid torus and then adding Reeb components, we obtain a foliation \mathcal{F} on the solid torus, which has the desired behavior on the boundary.

We claim that \mathcal{F} is uniform. To see this, take a compact leaf L of \mathcal{F}_1 in \mathbb{T}^3 intersecting only the curve c_z . Without loss of generality, $h(\partial T_z)$ is identified with the boundary of our solid torus with a link drilled. The leaf L does not separate $\mathbb{T}^3 \setminus (T_x \cup T_y \cup T_z)$. Therefore, its image $h(L)$ is a leaf of the foliation \mathcal{F} which does not separate the solid torus. A nonseparating surface on the solid torus must have a nontrivial intersection

number with the curves $S^1 \times \{\cdot\}$. This implies that when we lift the foliation to the universal cover, the lifts of this compact leaf bound a fundamental domain for $S^1 \times D^2$. Explicitly, this leaf on the solid torus looks like the yellow surface at the right of [Figure 5](#); see the proof of [Lemma 4.9](#) in [7, Section 5.3]. We conclude the uniform condition using [Lemma 3.2](#).

5 Proof of Theorem 1.3

In this section, M will be a compact 3-manifold endowed with a uniform foliation \mathcal{F} with Reeb components. We equip M with a Riemannian metric; this induces on the universal cover $p: \tilde{M} \rightarrow M$ the pullback metric. Lifting the foliation \mathcal{F} to \tilde{M} we obtain a foliation $\tilde{\mathcal{F}}$. By [Lemma 3.1](#), every leaf of this foliation has compact closure.

We will prove [Theorem 1.3](#) by contradiction. Suppose that there exists an essential immersion $i: \Sigma \rightarrow M$ from a closed surface Σ of genus $g \geq 1$ such that $i(\Sigma)$ does not intersect \mathcal{R} , the union of the Reeb components. Up to taking a finite cover, we can assume that \mathcal{F} is oriented and cooriented. Using [Theorem 2.3](#) we can assume that i is in general position with respect to the foliation. This is because there are immersions C^0 -close to i which are in general position, therefore if $i(\Sigma)$ is at a finite distance from \mathcal{R} , a sufficiently small perturbation of i will also be disjoint from \mathcal{R} . The foliation \mathcal{F} induces on Σ a singular foliation of Morse type, which will be denoted by \mathcal{G} .

Denote by $q: \mathbb{R}^2 \cong \tilde{\Sigma} \rightarrow \Sigma$ the universal cover of Σ . Lifting the immersion we obtain a map $\tilde{i}: \mathbb{R}^2 \cong \tilde{\Sigma} \rightarrow \tilde{M}$. As the immersion is essential, \tilde{i} is equivariant with respect to the actions of $\pi_1(\Sigma)$ on \mathbb{R}^2 and \tilde{M} . More precisely, for every $\gamma \in \pi_1(\Sigma)$ we have $\tilde{i} \circ \gamma = i_*(\gamma) \circ \tilde{i}$. The immersed plane $\tilde{i}(\mathbb{R}^2)$ is also in general position with respect to $\tilde{\mathcal{F}}$. In fact the induced singular foliation on \mathbb{R}^2 is exactly the lift of \mathcal{G} to the universal cover; we will denote it by $\tilde{\mathcal{G}}$. Notice that every leaf of $\tilde{\mathcal{G}}$ has compact closure because every leaf of $\tilde{\mathcal{F}}$ has compact closure.

We can classify every closed graph of $\tilde{\mathcal{G}}$:

Remark 5.1 Let S be a separatrix of $\tilde{\mathcal{G}}$ whose α -limit and ω -limit are singularities. Then $\alpha(S) = \omega(S)$. To see this, let $\{x\}$ and $\{y\}$ be the α - and ω -limit, respectively. The separatrices of \mathcal{G} cannot joint two distinct singularities by the second item in the definition of immersions in general position; therefore $y = \gamma x$ for some $\gamma \in \pi_1(\Sigma)$.




Figure 6: The possible configurations for the saddles in $\tilde{\mathcal{G}}$.

If γ is not the identity, then $q(S \cup \{x\} \cup \{\gamma x\})$ is a homotopically nontrivial curve. As the immersion is essential, this curve is sent to a homotopically nontrivial curve on M contained on certain leaf of \mathcal{F} . This is a contradiction by [Lemma 3.1](#); therefore $x = y$.

We will call a closed leaf or closed graph of $\tilde{\mathcal{G}}$ which is maximal with respect to inclusion (ie which is not strictly contained on another closed graph) a *generalized closed leaf*. The last remark tells us that a separatrix limits on at most one singularity. Therefore, if S is a separatrix of a generalized closed leaf limiting on a singularity x , then $S \cup \{x\}$ bounds a disk (by the Jordan curve theorem). This leaves us with exactly two possible configurations for a generalized closed leaf containing a singularity. Either the disk bounded by one separatrix contains the disk bounded by the other, or both disks are disjoint; both configurations are depicted in [Figure 6](#). These disks project homeomorphically to Σ , because their boundaries project homeomorphically.

Suppose that $D \subset \mathbb{R}^2$ is a disk bounded by a separatrix S of a generalized closed leaf. Then there is a well-defined notion of holonomy transport on the side of D . More precisely, take a half-open transversal τ of a point of S , with $\tau \subset D$. Start with a point x in τ and look at the next point of intersection between the leaf passing through x and τ (with respect to the orientation on \mathcal{G}). This defines a map from τ to itself. Choosing finitely many foliated neighborhoods of S and a model neighborhood of its corresponding singularity, one can see that this map is well-defined up to taking a smaller half-open transversal. Analogously, we can define the holonomy on the unbounded side of a generalized closed leaf.

Lemma 5.2 *Let \mathcal{F} be an oriented and transversely oriented uniform foliation with Reeb components on a compact 3-manifold M , and let $i: \Sigma \rightarrow M$ be an essential immersion of a surface Σ of genus ≥ 1 in general position with respect to \mathcal{F} . Suppose we further assume that $i(\Sigma)$ is disjoint from the set of Reeb components. Then every leaf of the induced singular foliation \mathcal{G} on Σ is either compact or a separatrix such that its α - and ω -limits coincide.*

Proof First, we prove that no generalized closed leaf of $\tilde{\mathcal{G}}$ has holonomy on any of its sides. Let C be a generalized closed leaf. If we had nontrivial holonomy on a certain side of C , then the curve $q(C) \subset \Sigma$ would be sent via i to a homotopically nontrivial curve on a leaf of \mathcal{F} . To see this, if $q(C)$ were sent to a homotopically trivial curve on a leaf of \mathcal{F} then it would have trivial holonomy on \mathcal{F} , and therefore trivial holonomy on $\tilde{\mathcal{G}}$.

Now we can use standard Haefliger-type arguments: by [Remark 5.1](#), $q(C)$ bounds one or two disks which are sent via i to immersed disks on M . The curve $i(q(C))$ is not homotopically trivial on its corresponding leaf of \mathcal{F} ; therefore, the boundary of at least one of the immersed disks is not homotopically trivial. This implies the existence of a vanishing cycle on at least one of the immersed disks; see for instance [\[4, Lemmas 9.2.2 and 9.2.4\]](#). However, Novikov's theorem says that if a leaf admits a vanishing cycle, then this leaf is the boundary of a Reeb component; see [\[4, Theorem 9.4.1\]](#). Therefore, $i(\Sigma)$ intersects the set of Reeb components, which is a contradiction.

Now we prove that either every leaf of \mathcal{G} is compact, or a separatrix limiting on a single singularity. It is enough to see this on $\tilde{\mathcal{G}}$. Suppose that L is a separatrix of $\tilde{\mathcal{G}}$ with $\alpha(S) \neq \omega(S)$, or a noncompact leaf. In any case, $\omega(L)$ or $\alpha(L)$ must be a generalized closed leaf or a closed leaf, by [Remark 5.1](#) and the Poincaré–Bendixson theorem. This generalized closed leaf must have nontrivial holonomy on one of its sides, which is a contradiction by the last paragraphs. \square

It is not hard to see that a singular foliation \mathcal{G} on Σ without holonomy must have a generalized closed leaf which is homotopically nontrivial:

Lemma 5.3 *Let Σ be a closed Riemannian surface of genus $g \geq 1$ and \mathcal{G} an oriented singular foliation of Morse type. Furthermore, suppose that every leaf is compact or a separatrix with the same α - and ω -limit. Then some leaf of the lifted singular foliation $\tilde{\mathcal{G}}$ on $\mathbb{R}^2 \cong \tilde{\Sigma}$ is unbounded.*

Proof Assume for the sake of contradiction that every leaf of $\tilde{\mathcal{G}}$ is bounded, and therefore has compact closure. Let \mathcal{C} be the set of generalized closed leaves of $\tilde{\mathcal{G}}$, and define the function $A: \mathcal{C} \rightarrow \mathbb{R}_{\geq 0}$ such that $A(C)$ is the area of the bounded component of the complement of $C \in \mathcal{C}$. Each connected component of the bounded component of an element of \mathcal{C} is a disk that projects homeomorphically to Σ , therefore its area cannot be greater than the area of Σ . This shows that the function A is bounded.

We claim that we can achieve the supremum $\sup_{C \in \mathcal{C}} A(C)$. Choose $x_n \in \mathbb{R}^2$ such that there exists a generalized closed leaf C_n with $x_n \in C_n$ and $A(C_n) \rightarrow \sup_{C \in \mathcal{C}} A(C)$ an increasing sequence. As the surface Σ is compact, up to translating $\{x_n\}$ by $\pi_1(\Sigma)$ we can assume that the sequence $\{x_n\}_{n \in \mathbb{N}}$ converges to $x \in \mathbb{R}^2$.

Let C be the generalized closed leaf passing through x ; we claim that $A(C)$ is the desired supremum. To see this, notice that if $m > n$ then $A(C_n) < A(C_m)$ and therefore, C_n is contained on the bounded component of C_m . This implies that for n sufficiently big, C_n is contained on the bounded component of C . Otherwise, there would exist some C_k bounding a region containing C_n with $n > k$, which is a contradiction.

Now choose C as in the last paragraph. This leaf cannot have holonomy because every leaf is compact. Therefore we can find a foliated neighborhood of C composed of closed orbits. In particular, some closed leaf bounding C exists, which contradicts the fact that C bounds the largest area. □

Therefore, [Lemma 5.2](#) tells us that the induced foliation on the surface Σ satisfies the hypothesis of [Lemma 5.3](#). However, this lemma contradicts the uniformity of the foliation \mathcal{F} . This contradiction concludes the proof of [Theorem 1.3](#).

Remark 5.4 Suppose that \mathcal{F} is an oriented and transversely oriented uniform foliation with Reeb components on a compact 3-manifold M , and let $i: \Sigma \rightarrow M$ be an essential immersion in general position as before. We just proved that $i(\Sigma)$ must intersect the




Figure 7: Turbulizing around the attracting/repelling fixed points we obtain a uniform foliation on \mathbb{T}^2 .

set of Reeb components. Thus, the induced singular foliation on Σ is uniform and necessarily contains vanishing cycles.

There exist singular foliations on surfaces satisfying these conditions; see for example [Figure 7.2](#). It is not obvious how the singular foliation induced on Σ would help in answering [Question 3.4](#). We believe that an extra argument is needed.

References

- [1] **I Agol**, *The virtual Haken conjecture*, Doc. Math. 18 (2013) 1045–1087 [MR](#) [Zbl](#)
- [2] **D Calegari**, *Foliations and the geometry of 3–manifolds*, Oxford Univ. Press (2007) [MR](#) [Zbl](#)
- [3] **A Candel**, **L Conlon**, *Foliations, I*, Graduate Studies in Mathematics 23, Amer. Math. Soc., Providence, RI (2000) [MR](#) [Zbl](#)
- [4] **A Candel**, **L Conlon**, *Foliations, II*, Graduate Studies in Mathematics 60, Amer. Math. Soc., Providence, RI (2003) [MR](#) [Zbl](#)
- [5] **JB Etnyre**, *Lectures on open book decompositions and contact structures*, from “Floer homology, gauge theory, and low-dimensional topology” (D A Ellwood, P S Ozsváth, A I Stipsicz, Z Szabó, editors), Clay Math. Proc. 5, Amer. Math. Soc., Providence, RI (2006) 103–141 [MR](#) [Zbl](#)
- [6] **SR Fenley**, **R Potrie**, *Minimality of the action on the universal circle of uniform foliations*, Groups Geom. Dyn. 15 (2021) 1489–1521 [MR](#) [Zbl](#)
- [7] **D Hardorp**, *All compact orientable three dimensional manifolds admit total foliations*, Mem. Amer. Math. Soc. 233, Amer. Math. Soc., Providence, RI (1980) [MR](#) [Zbl](#)
- [8] **A E Hatcher**, *Notes on basic 3–manifold topology*, course notes (2007) Available at <http://pi.math.cornell.edu/~hatcher/3M/3Mfds.pdf>
- [9] **G Hector**, **U Hirsch**, *Introduction to the geometry of foliations, A: Foliations on compact surfaces, fundamentals for arbitrary codimension, and holonomy*, Aspects of Mathematics 1, Vieweg & Sohn, Braunschweig (1981) [MR](#) [Zbl](#)
- [10] **G Hector**, **U Hirsch**, *Introduction to the geometry of foliations, B: Foliations of codimension one*, 2nd edition, Aspects of Mathematics E3, Vieweg & Sohn, Braunschweig (1987) [MR](#) [Zbl](#)
- [11] **SP Novikov**, *The topology of foliations*, Trudy Moskov. Mat. Obšč. 14 (1965) 248–278 [MR](#) [Zbl](#) In Russian; translated in Trans. Mosc. Math. Soc. 14 (1965) 268–304
- [12] **J Palis, Jr**, **W de Melo**, *Geometric theory of dynamical systems: an introduction*, Springer (1982) [MR](#) [Zbl](#)

²We thank Sergio Fenley for pointing out this example.

- [13] **J-P Serre**, *Trees*, Springer (1980) [MR](#) [Zbl](#)
- [14] **V V Solodov**, *Components of topological foliations*, *Mat. Sb.* 119(161) (1982) 340–354, 447 [MR](#) [Zbl](#) In Russian; translated in *Math. USSR Sb.* 47 (1984) 329–343
- [15] **J Stallings**, *Group theory and three-dimensional manifolds*, Yale Mathematical Monographs 4, Yale Univ. Press, New Haven, CT (1971) [MR](#) [Zbl](#)
- [16] **W P Thurston**, *Existence of codimension-one foliations*, *Ann. of Math.* 104 (1976) 249–268 [MR](#) [Zbl](#)
- [17] **W P Thurston**, *Three-manifolds, foliations and circles, I*, preprint (1997) [arXiv math/9712268](#)

Centro de Matemática, Udelar
Montevideo, Uruguay

joalema@cmat.edu.uy

Received: 8 December 2021 Revised: 25 April 2022

ALGEBRAIC & GEOMETRIC TOPOLOGY

msp.org/agt

EDITORS

PRINCIPAL ACADEMIC EDITORS

John Etnyre
etnyre@math.gatech.edu
Georgia Institute of Technology

Kathryn Hess
kathryn.hess@epfl.ch
École Polytechnique Fédérale de Lausanne

BOARD OF EDITORS

Julie Bergner	University of Virginia jeb2md@eservices.virginia.edu	Robert Lipshitz	University of Oregon lipshitz@uoregon.edu
Steven Boyer	Université du Québec à Montréal cohf@math.rochester.edu	Norihiko Minami	Nagoya Institute of Technology nori@nitech.ac.jp
Tara E Brendle	University of Glasgow tara.brendle@glasgow.ac.uk	Andrés Navas	Universidad de Santiago de Chile andres.navas@usach.cl
Indira Chatterji	CNRS & Univ. Côte d'Azur (Nice) indira.chatterji@math.cnrs.fr	Thomas Nikolaus	University of Münster nikolaus@uni-muenster.de
Alexander Dranishnikov	University of Florida dranish@math.ufl.edu	Robert Oliver	Université Paris 13 bobol@math.univ-paris13.fr
Tobias Ekholm	Uppsala University, Sweden tobias.ekholm@math.uu.se	Jessica S Purcell	Monash University jessica.purcell@monash.edu
Mario Eudave-Muñoz	Univ. Nacional Autónoma de México mario@matem.unam.mx	Birgit Richter	Universität Hamburg birgit.richter@uni-hamburg.de
David Futer	Temple University dfuter@temple.edu	Jérôme Scherer	École Polytech. Féd. de Lausanne jerome.scherer@epfl.ch
John Greenlees	University of Warwick john.greenlees@warwick.ac.uk	Vesna Stojanoska	Univ. of Illinois at Urbana-Champaign vesna@illinois.edu
Ian Hambleton	McMaster University ian@math.mcmaster.ca	Zoltán Szabó	Princeton University szabo@math.princeton.edu
Matthew Hedden	Michigan State University mhedden@math.msu.edu	Maggy Tomova	University of Iowa maggy-tomova@uiowa.edu
Hans-Werner Henn	Université Louis Pasteur henn@math.u-strasbg.fr	Nathalie Wahl	University of Copenhagen wahl@math.ku.dk
Daniel Isaksen	Wayne State University isaksen@math.wayne.edu	Chris Wendl	Humboldt-Universität zu Berlin wendl@math.hu-berlin.de
Thomas Koberda	University of Virginia thomas.koberda@virginia.edu	Daniel T Wise	McGill University, Canada daniel.wise@mcgill.ca
Christine Lescop	Université Joseph Fourier lescop@ujf-grenoble.fr		


See inside back cover or msp.org/agt for submission instructions.

The subscription price for 2023 is US \$650/year for the electronic version, and \$940/year (+ \$70, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Algebraic & Geometric Topology is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [Current Mathematical Publications](#) and the [Science Citation Index](#).

Algebraic & Geometric Topology (ISSN 1472-2747 printed, 1472-2739 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840.

AGT peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2023 Mathematical Sciences Publishers

ALGEBRAIC & GEOMETRIC TOPOLOGY

Volume 23

Issue 9 (pages 3909–4400)

2023

Two-dimensional extended homotopy field theories	3909
KÜRŞAT SÖZER	
Efficient multisections of odd-dimensional tori	3997
THOMAS KINDRED	
Bigrading the symplectic Khovanov cohomology	4057
ZHECHI CHENG	
Fibrations of 3–manifolds and asymptotic translation length in the arc complex	4087
BALÁZS STRENNER	
A uniformizable spherical CR structure on a two-cusped hyperbolic 3–manifold	4143
YUEPING JIANG, JIEYAN WANG and BAOHUA XIE	
A connection between cut locus, Thom space and Morse–Bott functions	4185
SOMNATH BASU and SACHCHIDANAND PRASAD	
Staircase symmetries in Hirzebruch surfaces	4235
NICKI MAGILL and DUSA MCDUFF	
Geometric triangulations of a family of hyperbolic 3–braids	4309
BARBARA NIMERSHIEM	
Beta families arising from a v_2^9 self-map on $S/(3, v_1^8)$	4349
EVA BELMONT and KATSUMI SHIMOMURA	
Uniform foliations with Reeb components	4379
JOAQUÍN LEMA	