

AG
T

*Algebraic & Geometric
Topology*

Volume 24 (2024)

Equivariant Seiberg–Witten–Floer cohomology

DAVID BARAGLIA
PEDRAM HEKMATI



Equivariant Seiberg–Witten–Floer cohomology

DAVID BARAGLIA

PEDRAM HEKMATI

We develop an equivariant version of Seiberg–Witten–Floer cohomology for finite group actions on rational homology 3–spheres. Our construction is based on an equivariant version of the Seiberg–Witten–Floer stable homotopy type, as constructed by Manolescu. We use these equivariant cohomology groups to define a series of d –invariants $d_{G,c}(Y, \mathfrak{s})$ which are indexed by the group cohomology of G . These invariants satisfy a Frøyshov-type inequality under equivariant cobordisms. Lastly, we consider a variety of applications of these d –invariants: concordance invariants of knots via branched covers, obstructions to extending group actions over bounding 4–manifolds, Nielsen realisation problems for 4–manifolds with boundary and obstructions to equivariant embeddings of 3–manifolds in 4–manifolds.

57K31; 57K10, 57K41

1 Introduction

In this paper we develop an equivariant version of Seiberg–Witten–Floer cohomology for rational homology 3–spheres equipped with the action of a finite group. Our approach is modelled on the construction of a Seiberg–Witten–Floer stable homotopy type due to Manolescu [49], which we now briefly recall. Let Y be a rational homology 3–sphere and \mathfrak{s} a spin^c –structure on Y . Given a metric g on Y , the construction of [49] yields an S^1 –equivariant stable homotopy type $SWF(Y, \mathfrak{s}, g)$. The Seiberg–Witten–Floer cohomology of (Y, \mathfrak{s}) is then given (up to a degree shift) by the S^1 –equivariant cohomology of $SWF(Y, \mathfrak{s}, g)$:

$$HSW^*(Y, \mathfrak{s}) = \tilde{H}_{S^1}^{*+2n(Y, \mathfrak{s}, g)}(SWF(Y, \mathfrak{s}, g)),$$

where $n(Y, \mathfrak{s}, g)$ is a rational number given by a certain combination of eta invariants.

The stable homotopy type $SWF(Y, \mathfrak{s}, g)$ depends on the choice of metric, but only up to a suspension. Given two metrics, g_0 and g_1 , one obtains a canonical homotopy equivalence

$$(1-1) \quad SWF(Y, \mathfrak{s}, g_1) \cong \Sigma^{SF(\{D_s\})\mathbb{C}} SWF(Y, \mathfrak{s}, g_0),$$

where $SF(\{D_s\})$ denotes the spectral flow for the family of Dirac operators $\{D_s\}$ determined by a path of metrics $\{g_s\}$ from g_0 to g_1 . The rational numbers $n(Y, \mathfrak{s}, g)$ are defined in such a way that they split the spectral flow in the sense that

$$(1-2) \quad SF(\{D_s\}) = n(Y, \mathfrak{s}, g_1) - n(Y, \mathfrak{s}, g_0).$$

Hence we obtain a canonical isomorphism

$$\tilde{H}_{S^1}^{*+2n(Y, \mathfrak{s}, g_1)}(SWF(Y, \mathfrak{s}, g_1)) \cong \tilde{H}_{S^1}^{*+2n(Y, \mathfrak{s}, g_0)}(SWF(Y, \mathfrak{s}, g_0)).$$

This shows that the Seiberg–Witten–Floer cohomology $HSW^*(Y, \mathfrak{s})$ does not depend on the choice of metric g .

By working in an appropriately defined S^1 –equivariant Spanier–Whitehead category in which suspension by fractional amounts of \mathbb{C} is allowed, Manolescu defined the Seiberg–Witten–Floer homotopy type of (Y, \mathfrak{s}) to be

$$SW(Y, \mathfrak{s}) = \Sigma^{-n(Y, \mathfrak{s}, g)} \mathbb{C} SWF(Y, \mathfrak{s}, g).$$

This is independent of the choice of g by (1-1) and (1-2).

Now suppose that a finite group G acts on Y by orientation-preserving diffeomorphisms which preserve the isomorphism class of \mathfrak{s} . Let g be a G –invariant metric on Y . Lifting the action of G to the associated spinor bundle determines an S^1 extension

$$1 \rightarrow S^1 \rightarrow G_{\mathfrak{s}} \rightarrow G \rightarrow 1.$$

Manolescu’s construction of the stable homotopy type $SWF(Y, \mathfrak{s}, g)$ can be carried out $G_{\mathfrak{s}}$ –equivariantly, so that $SWF(Y, \mathfrak{s}, g)$ may be promoted to a $G_{\mathfrak{s}}$ –equivariant stable homotopy type. This is analogous to the construction in [50] of the $\text{Pin}(2)$ –equivariant Seiberg–Witten–Floer stable homotopy type of (Y, \mathfrak{s}) where \mathfrak{s} is a spin–structure on Y . The main difference is that in our construction, the additional symmetries that comprise the group $G_{\mathfrak{s}}$ come from symmetries of Y rather than internal symmetries of the Seiberg–Witten equations.

We define the G –equivariant Seiberg–Witten–Floer cohomology of (Y, \mathfrak{s}) to be

$$HSW_G^*(Y, \mathfrak{s}) = \tilde{H}_{G_{\mathfrak{s}}}^{*+2n(Y, \mathfrak{s}, g)}(SWF(Y, \mathfrak{s}, g)).$$

The right-hand side is independent of the choice of metric g by much the same argument as in the S^1 –equivariant case.

We make some remarks concerning this construction.

- (1) Throughout this paper we have chosen to work with cohomology instead of homology. This is simply a matter of preference and we could just as well work with Seiberg–Witten–Floer homology groups.
- (2) Instead of Borel equivariant cohomology, we could take co-Borel cohomology or Tate cohomology, which correspond to the different versions of Heegaard Floer cohomology; see Lidman and Manolescu [45, Corollary 1.2.4].
- (3) In a similar fashion we can also define the G –equivariant Seiberg–Witten–Floer K –theory

$$KSW_G^*(Y, \mathfrak{s}) = \tilde{K}_{G_{\mathfrak{s}}}^{*+2n(Y, \mathfrak{s}, g)}(SWF(Y, \mathfrak{s}, g)).$$

More generally we could use any generalised equivariant cohomology theory in which the Thom isomorphism holds.

(4) We have not attempted to construct a metric independent $G_{\mathfrak{s}}$ -equivariant stable homotopy type. To do this one would need to split the equivariant spectral flow $SF_{G_{\mathfrak{s}}}(\{D_{\mathfrak{s}}\})$ in the same way that $n(Y, \mathfrak{s}, g)$ splits the nonequivariant spectral flow, as in (1-2).

1.1 Main results

Throughout we work with cohomology with coefficients in a field \mathbb{F} . To avoid the necessity of local systems we assume that either $\mathbb{F} = \mathbb{Z}/2\mathbb{Z}$, or that the order of G is odd (see Section 3.1). We now outline the main properties of equivariant Seiberg–Witten–Floer cohomology.

Module structure $HSW_G^*(Y, \mathfrak{s})$ is a graded module over $H_{G_{\mathfrak{s}}}^*$ (where for a group K we write H_K^* for $H_K^*(pt)$). In particular if $G_{\mathfrak{s}}$ is the trivial extension then $HSW_G^*(Y, \mathfrak{s})$ is a graded module over $H_G^*[U]$, where $\deg(U) = 2$.

Theorem 1.1 (spectral sequence) *There is a spectral sequence $E_r^{p,q}$ abutting to $HSW_G^*(Y, \mathfrak{s})$ whose second page is given by*

$$E_2^{p,q} = H^p(BG; HSW^q(Y, \mathfrak{s})).$$

Theorem 1.2 (localisation) *Suppose the extension $G_{\mathfrak{s}}$ is trivial and choose a trivialisation $G_{\mathfrak{s}} \cong S^1 \times G$. Then $H_{G_{\mathfrak{s}}}^* \cong H_G^*[U]$ and the localisation $U^{-1}HSW_G^*(Y, \mathfrak{s})$ of $HSW_G^*(Y, \mathfrak{s})$ with respect to U is a free $H_G^*[U, U^{-1}]$ -module of rank 1.*

L-spaces We say that Y is an L -space with respect to \mathfrak{s} and \mathbb{F} if $HSW^*(Y, \mathfrak{s})$ is isomorphic to a free $\mathbb{F}[U]$ -module of rank 1.

Theorem 1.3 *Suppose that $G_{\mathfrak{s}}$ is a split extension. If Y is an L -space with respect to \mathfrak{s} and \mathbb{F} , then the spectral sequence given in Theorem 1.1 degenerates at E_2 . Moreover,*

$$HSW_G^*(Y, \mathfrak{s}) \cong HSW^*(Y, \mathfrak{s}) \otimes_{\mathbb{F}} H_G^*.$$

Correction terms Suppose that $G_{\mathfrak{s}}$ is a split extension. For each nonzero $c \in H_G^*$ we obtain an invariant

$$d_{G,c}(Y, \mathfrak{s}) \in \mathbb{Q}$$

which may be thought of as a generalisation to the equivariant setting of the d -invariant $d(Y, \mathfrak{s})$. We also set $d_{G,0}(Y, \mathfrak{s}) = -\infty$.

Theorem 1.4 *The equivariant d -invariants satisfy the following properties:*

- (1) $d_{G,1}(Y, \mathfrak{s}) \geq d(Y, \mathfrak{s})$, where 1 is the generator of $H_G^0(pt)$;
- (2) $d_{G,c_1+c_2}(Y, \mathfrak{s}) \leq \max\{d_{G,c_1}(Y, \mathfrak{s}), d_{G,c_2}(Y, \mathfrak{s})\}$;
- (3) $d_{G,c_1c_2}(Y, \mathfrak{s}) \leq \min\{d_{G,c_1}(Y, \mathfrak{s}), d_{G,c_2}(Y, \mathfrak{s})\}$;
- (4) $d_{G,c_1}(Y, \mathfrak{s}) + d_{G,c_2}(\bar{Y}, \mathfrak{s}) \geq 0$ whenever $c_1c_2 \neq 0$;

- (5) if Y is an L -space with respect to \mathfrak{s} and \mathbb{F} , then $d_{G,c}(Y, \mathfrak{s}) = d(Y, \mathfrak{s})$ for all $c \neq 0$;
- (6) $d_{G,c}(Y, \mathfrak{s})$ is invariant under equivariant rational homology cobordism.

We find it convenient to also define corresponding equivariant δ -invariants by setting

$$\delta_{G,c}(Y, \mathfrak{s}) = \frac{1}{2}d_{G,c}(Y, \mathfrak{s}).$$

Our primary motivation for considering the equivariant d -invariants is that they are necessary for the formulation of our equivariant generalisation of Frøyshov’s inequality described below.

Cobordism maps Suppose that (W, \mathfrak{s}) is a G -equivariant cobordism from (Y_1, \mathfrak{s}_1) to (Y_2, \mathfrak{s}_2) (see Section 4.3 for the precise statement). Then W induces a morphism of graded H_G^* -modules

$$SW_G(W, \mathfrak{s}): HSW_G^*(Y_2, \mathfrak{s}_2) \rightarrow HSW_G^{*+b_+(W)-2\delta(W,\mathfrak{s})}(Y_1, \mathfrak{s}_1)$$

where $\delta(W, \mathfrak{s}) = \frac{1}{8}(c_1(\mathfrak{s})^2 - \sigma(W))$.

Theorem 1.5 (equivariant Frøyshov inequality) *Let W be a smooth, compact, oriented 4-manifold with boundary and with $b_1(W) = 0$. Suppose that G acts smoothly on W preserving the orientation and a spin^c -structure \mathfrak{s} . Suppose that the extension $G_{\mathfrak{s}}$ is trivial. Suppose each component of ∂W is a rational homology 3-sphere and that G sends each component of ∂W to itself. Let $e \in H_G^{b_+(W)}$ be the image in $H_G^*(\text{pt}; \mathbb{F})$ of the Euler class of any G -invariant maximal positive definite subspace of $H^2(W; \mathbb{R})$. Let $c \in H_G^*$ and suppose that $ce \neq 0$.*

- (1) If $\partial W = Y$ is connected, then

$$\delta(W, \mathfrak{s}) \leq \delta_{G,c}(Y, \mathfrak{s}|_Y) \quad \text{and} \quad \delta_{G,ce}(\bar{Y}, \mathfrak{s}|_Y) \leq \delta(\bar{W}, \mathfrak{s}).$$

- (2) If $\partial W = \bar{Y}_1 \cup Y_2$ has two connected components, then

$$\delta_{G,ce}(Y_1, \mathfrak{s}|_{Y_1}) + \delta(W, \mathfrak{s}) \leq \delta_{G,c}(Y_2, \mathfrak{s}|_{Y_2}).$$

Knot concordance invariants Let K be a knot in S^3 and let $Y = \Sigma_2(K)$ be the double cover of S^3 branched over K . Then Y has an action of $G = \mathbb{Z}_2$ generated by the covering involution. Further, Y has a spin^c -structure \mathfrak{t}_0 uniquely determined by the condition that it arises from a spin -structure. Set $\mathbb{F} = \mathbb{Z}_2$. Then $H_G^* \cong \mathbb{F}[Q]$, where $\deg(Q) = 1$. For each $j \geq 0$, we define an invariant $\delta_j(K) \in \mathbb{Z}$ by setting

$$\delta_j(K) = 4\delta_{\mathbb{Z}_2, Q^j}(\Sigma_2(K), \mathfrak{t}_0).$$

Let $\sigma(K)$ and $g_4(K)$ denote the signature and smooth 4-genus of K .

Theorem 1.6 *The invariants $\delta_j(K)$ have the following properties:*

- (1) $\delta_j(K)$ is a knot concordance invariant;
- (2) $\delta_0(K) \geq \delta(K)$, where $\delta(K)$ is the Manolescu–Owens invariant [51];
- (3) $\delta_{j+1}(K) \leq \delta_j(K)$ for all $j \geq 0$;

- (4) $\delta_j(K) \geq -\frac{1}{2}\sigma(K)$ for all $j \geq 0$ and $\delta_j(K) = -\frac{1}{2}\sigma(K)$ for $j \geq g_4(K) - \frac{1}{2}\sigma(K)$;
- (5) $\delta_j(-K) \geq \frac{1}{2}\sigma(K)$ for all $j \geq 0$ and $\delta_j(-K) = \frac{1}{2}\sigma(K)$ for $j \geq g_4(K) + \frac{1}{2}\sigma(K)$;
- (6) if $\Sigma_2(K)$ is an L -space, then $\delta_j(K) = \delta(K) = -\frac{1}{2}\sigma(K)$ and $\delta_j(-K) = \delta(-K) = \frac{1}{2}\sigma(K)$ for all $j \geq 0$.

In particular, if K is quasialternating, then $\Sigma_2(K)$ is an L -space; see Ozsváth and Szabó [57]. So we recover the main result of Lisca and Owens [47] that $\delta(K) = -\frac{1}{2}\sigma(K)$ for quasialternating knots.

The concordance invariants $\delta_j(K)$ can also be used to strengthen the inequality $g_4(K) \geq \frac{1}{2}|\sigma(K)|$; see Murasugi [52].

Theorem 1.7 For a knot K , let $j_+(K)$ be the smallest positive integer such that $\delta_j(K) = -\frac{1}{2}\sigma(K)$ and $j_-(K)$ the smallest positive integer such that $\delta_j(-K) = \frac{1}{2}\sigma(K)$. Then

$$g_4(K) \geq \max\left\{-\frac{1}{2}\sigma(K) + j_-(K), \frac{1}{2}\sigma(K) + j_+(K)\right\}.$$

Corollary 1.8 If $\delta(K) > -\frac{1}{2}\sigma(K)$ and $\sigma(K) \geq 0$, then

$$g_4(K) \geq \frac{1}{2}|\sigma(K)| + 1$$

Proof If $\delta(K) > -\frac{1}{2}\sigma(K)$, then $\delta_0(K) \geq \delta(K) > -\frac{1}{2}\sigma(K)$, and thus $j_+(K) \geq 1$. Hence

$$g_4(K) \geq \frac{1}{2}\sigma(K) + 1 = \frac{1}{2}|\sigma(K)| + 1. \quad \square$$

One can obtain even more knot concordance invariants by considering higher order cyclic branched covers; see Remark 6.7.

1.2 Applications

We outline here some of the applications of equivariant Seiberg–Witten–Floer cohomology. These are considered in more detail in Section 7.

1.2.1 Nonextendable actions (Section 7.2) Let Y be a rational homology 3–sphere equipped with an orientation-preserving action of G and let W be a smooth 4–manifold which bounds Y . The equivariant d -invariants give obstructions to extending the action of G over W .

Example 1.9 The Brieskorn homology sphere $Y = \Sigma(p, q, r)$ where p, q and r are pairwise coprime is the branched cyclic p -fold cover of the torus knot $T_{q,r}$. Let $\tau : Y \rightarrow Y$ be a generator of the \mathbb{Z}_p -action determined by this covering. For certain values of p, q and r it can be shown that Y bounds a contractible 4–manifold. For example, $\Sigma(2, 3, 13)$ bounds a contractible 4–manifold; see Akbulut and Kirby [2]. It can be shown that τ is smoothly isotopic to the identity; hence it follows that τ can be extended as a diffeomorphism over any 4–manifold bounded by Y . On the other hand we show in Proposition 7.2

that if p is prime then $\delta_{\mathbb{Z}_p,1}(Y, \mathfrak{s}) = -\lambda(Y)$ is minus the Casson invariant of Y (where \mathfrak{s} is the unique spin^c -structure on Y), which is nonzero. We further show that the nonvanishing of $\delta_{\mathbb{Z}_p,1}(Y, \mathfrak{s})$ implies that τ cannot be extended as a smooth \mathbb{Z}_p -action to any contractible 4-manifold bounded by Y . This partially recovers the nonextendability results of Anvari and Hambleton [6; 7] for Brieskorn homology 3-spheres bounded by contractible 4-manifolds.

On the other hand, our nonextendability result also holds in situations not covered by Anvari and Hambleton. Suppose now that $Y = \Sigma(p, q, r)$ bounds a *rational* homology 4-ball W . For example, Fintushel and Stern showed that $\Sigma(2, 3, 7)$ bounds a rational homology 4-ball, although it does not bound an integral homology 4-ball [26]. More examples can be found in Akbulut and Larson [4] and Şavk [21]. We show in Section 7.2 that the nonvanishing of $\delta_{\mathbb{Z}_p,1}(\Sigma(p, q, r))$, where p is prime, implies that the \mathbb{Z}_p -action cannot be extended to any rational homology 4-ball W bounded by Y , provided that p does not divide the order of $H^2(W; \mathbb{Z})$.

1.2.2 Realisation problems (Section 7.3) Let W be a smooth 4-manifold with boundary an integral homology sphere Y . Suppose that a finite group G acts on $H^2(W; \mathbb{Z})$ preserving the intersection form. We say that the action of G on $H^2(W; \mathbb{Z})$ can be realised by diffeomorphisms if there is a smooth orientation-preserving action of G on W inducing the given action on $H^2(W; \mathbb{Z})$. The equivariant d -invariants give obstructions to realising such actions by diffeomorphism. This extends the nonrealisation results of the first author [10; 11] for closed 4-manifolds to the case of 4-manifolds with nonempty boundary.

Example 1.10 Suppose that $b_1(W) = 0$ and that $H^2(W; \mathbb{Z})$ has no 2-torsion and even intersection form. Suppose that Y is an L -space. Suppose that an action of $G = \mathbb{Z}_p$ on $H^2(W; \mathbb{Z})$ is given, where p is prime and that the subspace of $H^2(W; \mathbb{R})$ fixed by G is negative definite. If $\frac{1}{8}\sigma(W) < -\delta(Y, \mathfrak{s})$ (where \mathfrak{s} is the unique spin^c -structure on Y) then the action of \mathbb{Z}_p on $H^2(W; \mathbb{Z})$ is not realisable by a smooth \mathbb{Z}_p -action on W . Note that we are not making any assumptions about the action of \mathbb{Z}_p on the boundary.

1.2.3 Equivariant embeddings of 3-manifolds in 4-manifolds (Section 7.4) Let Y be a rational homology 3-sphere equipped with an orientation-preserving action of G . By an equivariant embedding of Y into a 4-manifold X , we mean an embedding $Y \rightarrow X$ such that the action of G on Y extends over X .

Example 1.11 Let $Y = \Sigma(2, 2s - 1, 2s + 1)$ where s is odd, equipped with the involution τ obtained from viewing Y as the branched double cover $\Sigma_2(T_{2s-1, 2s+1})$. Then Y embeds in S^4 ; see Budney and Burton [13, Theorem 2.13]. On the other hand, $\delta_j(Y, \mathfrak{s}) \neq 0$ for some j . We will show that the nonvanishing of this invariant implies that Y cannot be equivariantly embedded in S^4 .

It is known that every 3-manifold Y embeds in the connected sum $\#^n(S^2 \times S^2)$ of n copies of $S^2 \times S^2$ for some sufficiently large n [1, Theorem 2.1]. Aceto, Golla and Larson define the *embedding number* $\varepsilon(Y)$ of Y to be the smallest n for which Y embeds in $\#^n(S^2 \times S^2)$. Here we consider an equivariant version of the embedding number. To obtain interesting results we need to make an assumption on the kinds of group actions allowed.

Definition 1.12 Let $G = \mathbb{Z}_p = \langle \tau \rangle$ where p is a prime number. We say that a smooth, orientation-preserving action of G on $X = \#^n(S^2 \times S^2)$ is *admissible* if $H^2(X; \mathbb{Z})^\tau = 0$, where

$$H^2(X; \mathbb{Z})^\tau = \{x \in H^2(X; \mathbb{Z}) \mid \tau(x) = x\}.$$

We define the *equivariant embedding number* $\varepsilon(Y, \tau)$ of (Y, τ) to be the smallest n for which Y embeds equivariantly in $\#^n(S^2 \times S^2)$ for some admissible \mathbb{Z}_p -action on $\#^n(S^2 \times S^2)$, if such an embedding exists. We set $\varepsilon(Y, \tau) = \infty$ if there is no such embedding.

Example 1.13 Let $Y = \Sigma(2, 3, 6n + 1) = \Sigma_2(T_{3, 6n+1})$ and equip Y with the covering involution τ . We show that

$$2n \leq \varepsilon(\Sigma(2, 3, 6n + 1), \tau) \leq 12n.$$

Suppose that n is odd. Then from [1, Proposition 3.5], the (nonequivariant) embedding number of $\Sigma(2, 3, 6n + 1)$ is 10. In particular, we see that $\varepsilon(\Sigma(2, 3, 6n + 1), \tau) > \varepsilon(\Sigma(2, 3, 6n + 1))$ for all odd $n > 5$. We also show that

$$\varepsilon(\Sigma(2, 3, 7), \tau) = 12,$$

whereas $\varepsilon(\Sigma(2, 3, 7)) = 10$.

1.3 Comparison with other works

In [30], Hendricks, Lipshitz and Sarkar introduce equivariant versions of several types of Floer homology, mostly focusing on the case that the group is \mathbb{Z}_2 . In particular they define a \mathbb{Z}_2 -equivariant version of HF^- , which is a module over $H_{S^1 \times \mathbb{Z}_2}^*(pt; \mathbb{Z}_2) = \mathbb{Z}_2[U, Q]$. This construction shares many similarities with the equivariant Seiberg–Witten–Floer cohomology constructed in this paper, such as a localisation isomorphism and a spectral sequence relating the equivariant and ordinary Floer homologies. In fact, it seems reasonable to conjecture that our constructions are isomorphic.

In [5], Alfieri, Kang and Stipsicz consider a \mathbb{Z}_2 -equivariant Heegaard Floer homology $HFB^-(K)$ for a branched double cover $Y = \Sigma_2(K)$ of a knot K , constructed in a manner similar to involutive Heegaard Floer homology — see Hendricks and Manolescu [31] — except that the involution arises from the covering involution on Y . These groups are modules over the ring $\mathbb{Z}_2[U, Q]/(Q^2)$. From this group they obtain knot concordance invariants $\bar{\delta}(K), \underline{\delta}(K)$. A similar approach was taken by Dai, Hedden and Mallick [22] to obtain ι -complexes — see Hendricks, Manolescu and Zemke [32, Definition 8.1] — associated to involutions on Y . Since $\mathbb{Z}_2[U, Q]/(Q^2) = H_{S^1 \times \mathbb{Z}_2}^*(pt; \mathbb{Z}_2)$, we suspect that the group $HFB^-(K)$ may be isomorphic to the \mathbb{Z} -equivariant Seiberg–Witten–Floer homology of $\Sigma_2(K)$.

In [46, Remark 3.1], Lidman and Manolescu define equivariant Seiberg–Witten–Floer homology in the special case that G acts freely on Y . Their construction coincides with ours in such cases.

1.4 Structure of the paper

In Section 2 we recall the construction of Seiberg–Witten–Floer spectra using finite-dimensional approximation and the Conley index. In Section 3 we extend this construction to the G -equivariant setting,

arriving at the construction of the G -equivariant Seiberg–Witten–Floer cohomology in Section 3.4. In the remainder of Section 3 we introduce the equivariant d -invariants and establish their basic properties. Section 4 is concerned with the behaviour of equivariant Seiberg–Witten–Floer cohomology and the d -invariants under equivariant cobordism. In Section 5 we specialise to the case that G is a cyclic group of prime order. In Section 6 we consider the case of branched double covers of knots with their natural involution to obtain knot concordance invariants. Finally in Section 7 we carry out some explicit computations of d -invariants and consider various applications.

2 Seiberg–Witten–Floer spectra

2.1 Seiberg–Witten trajectories

Throughout we let Y be a rational homology 3-sphere, ie Y is a compact, oriented, smooth 3-manifold with $b_1(Y) = 0$. References for the material in this section are [45; 49].

Let g be a Riemannian metric on Y and let \mathfrak{s} be a spin^c -structure with associated spinor bundle S . Let $\rho: TY \rightarrow \text{End}(S)$ denote Clifford multiplication, satisfying $\rho(v)\rho(w) + \rho(w)\rho(v) = -2g(v, w)$. The spinor bundle S is equipped with a Hermitian metric $\langle \cdot, \cdot \rangle$ which we take to be antilinear in the first variable. Let $\mathfrak{su}(S)$ be the Lie algebra bundle of trace-free skew-adjoint endomorphisms of S and $\mathfrak{sl}(S)$ the Lie algebra bundle of trace-free endomorphisms of S . Then ρ induces an isomorphism $\rho: TY \rightarrow \mathfrak{su}(S)$ which extends by complexification to an isomorphism $\rho: TY_{\mathbb{C}} \rightarrow \mathfrak{sl}(S)$ satisfying $\rho(\bar{v}) = -\rho(v)^*$. Using the metric g to identify TY and T^*Y we will also view ρ as a map $\rho: T^*Y \rightarrow \mathfrak{su}(S)$. We extend ρ to 2-forms by the rule $\rho(v \wedge w) = \frac{1}{2}[\rho(v), \rho(w)]$. It follows that $\rho(\lambda) = -\rho(*\lambda)$ for any 2-form λ . Define a Hermitian inner product on $\mathfrak{su}(S)$ by $\langle a, b \rangle = \frac{1}{2} \text{tr}(a^*b)$. Then for any tangent vectors u and v , we have $\langle \rho(u), \rho(v) \rangle = g(\bar{u}, v)$. Define a map

$$\tau: S \times S \rightarrow T^*Y_{\mathbb{C}}$$

by setting

$$\tau(\phi, \psi) = \rho^{-1}(\phi \otimes \psi^*)_0,$$

where $(\phi \otimes \psi^*)_0$ is the trace-free part of $\phi \otimes \psi^*$. That is, if ξ is any spinor, then

$$(\phi \otimes \psi^*)(\xi) = \phi \langle \psi, \xi \rangle - \frac{1}{2} \langle \psi, \phi \rangle \xi.$$

Then it follows that

$$\tau(\psi, \phi) = -\overline{\tau(\phi, \psi)}, \quad \tau(a\psi, b\phi) = a\bar{b}\tau(\psi, \phi).$$

In particular, $\tau(\phi, \phi)$ is imaginary and $\tau(c\phi, c\phi) = |c|^2\tau(\phi, \phi)$. We also have the identity

$$\langle \tau(\phi, \phi), v \rangle = \frac{1}{2} \langle \phi, \rho(v)\phi \rangle$$

for all spinors ϕ and vectors v .

Let $L = \det(S)$ be the determinant line bundle of S . Let $\Gamma(L)$ denote the space of $U(1)$ –connections on L , which is an affine space over $i\Omega^1(Y)$. We will write such a connection as $2A$. Then $2A$ determines a $\text{spin}^c(3)$ –connection on S whose $u(1)$ part is A and whose $\text{spin}(3)$ part is the Levi-Civita connection. Abusing terminology, we will refer to A as a spin^c –connection.

Given a spin^c –connection A , we let D_A denote the associated Dirac operator on S . Fix a reference spin^c –connection A_0 . Then we may write $A = A_0 + a$ for some $a \in i\Omega^1(Y)$. It follows that

$$D_A(\psi) = D_{A_0+a}(\psi) = D_{A_0}(\psi) + \rho(a)\psi.$$

Since $b_1(Y) = 0$, it follows that L admits a flat connection. We will assume that A_0 defines a flat connection on L .

We define the configuration space of Y to be

$$C(Y) = \Gamma(L) \times \Gamma(S).$$

$C(Y)$ depends on g and \mathfrak{s} but we omit this from the notation. $C(Y)$ is an affine space modelled on $i\Omega^1(Y) \oplus \Gamma(S)$. In particular, the tangent space $T_{(A,\phi)}C(Y)$ to any point $(A, \phi) \in C(Y)$ can naturally be identified with $i\Omega^1(Y) \oplus \Gamma(S)$. There is a natural metric on $i\Omega^1(Y) \oplus \Gamma(S)$, the L^2 –metric

$$\langle (a_1, \phi_1), (a_2, \phi_2) \rangle_{L^2} = -\int_Y a_1 \wedge *a_2 + \int_Y \text{Re}\langle \phi_1, \phi_2 \rangle \, d\text{vol}_Y.$$

This defines a (constant) Riemannian metric on $C(Y)$. We will need to work with Sobolev completions. Given a flat reference spin^c –connection A_0 , Sobolev norms are defined using A_0 and g . Fix an integer $k \geq 4$. Later we will work with the L^2_{k+1} –completion of $C(Y)$ and L^2_{k+2} –gauge transformations.

Having fixed a reference connection A_0 , we identify $C(Y)$ with $i\Omega^1(Y) \oplus \Gamma(S)$. Thus an element $(A, \phi) \in C(Y)$ will be identified with $(a, \phi) \in i\Omega^1(Y) \oplus \Gamma(S)$, where $A = A_0 + a$. To simplify notation, we will write D_a in place of D_{A_0+a} .

The Chern–Simons–Dirac functional $\mathcal{L}: C(Y) \rightarrow \mathbb{R}$ (with respect to A_0) is defined as

$$\mathcal{L}(a, \phi) = \frac{1}{2} \left(\int_Y \langle \phi, D_a \phi \rangle \, d\text{vol}_Y - \int_Y a \wedge da \right).$$

The gauge group $\mathcal{G} = \mathcal{C}^\infty(Y, S^1)$ acts on $C(Y)$ by

$$u \cdot (a, \phi) = (a - u^{-1} du, u \cdot \phi).$$

Observe that $D_{a-u^{-1}du}(u\phi) = uD_a\phi$, so

$$\langle u\phi, D_{a-u^{-1}du}(u\phi) \rangle = \langle u\phi, uD_a\phi \rangle = \langle \phi, D_a\phi \rangle.$$

It follows that \mathcal{L} is gauge invariant and we can regard \mathcal{L} as a function on the quotient space $C(Y)/\mathcal{G}$. The goal of Seiberg–Witten–Floer theory is to construct some sensible notion of Morse homology of \mathcal{L} on

$C(Y)/\mathcal{G}$. Consider the formal L^2 -gradient of \mathcal{L} ; that is, the function $\text{grad}(\mathcal{L}): C(Y) \rightarrow i\Omega^1(Y) \oplus \Gamma(S)$ such that

$$\langle \text{grad}(\mathcal{L})(a, \phi), (a', \phi') \rangle_{L^2} = \left. \frac{d}{dt} \right|_{t=0} (\mathcal{L}(a, \phi) + t(a', \phi'))$$

for all $(a, \phi) \in C(Y)$ and all $(a', \phi') \in i\Omega^1(Y) \oplus \Gamma(S)$. A short calculation gives

$$\text{grad}(\mathcal{L}) = (*da + \tau(\phi, \phi), D_a\phi).$$

A critical point of \mathcal{L} is a point where $\text{grad}(\mathcal{L})$ vanishes. So (a, ϕ) is a critical point if and only if it satisfies

$$*da + \tau(\phi, \phi) = 0, \quad D_a\phi = 0.$$

These are the 3-dimensional Seiberg–Witten equations.

A trajectory for the downwards gradient flow is a differentiable map $x: \mathbb{R} \rightarrow L^2_{k+1}(C(Y))$ such that

$$\frac{d}{dt}x(t) = -\text{grad}(\mathcal{L})(x(t)).$$

If $x(t) = (a(t), \phi(t)) \in L^2_{k+1}(Y, iT^*Y \oplus S)$, then

$$\frac{d}{dt}a(t) = -*da(t) - \tau(\phi(t), \phi(t)), \quad \frac{d}{dt}\phi(t) = -D_{a(t)}\phi(t).$$

A key observation is that such trajectories can be reinterpreted as solutions of the 4-dimensional Seiberg–Witten equations on the cylinder $X = \mathbb{R} \times Y$.

Definition 2.1 A Seiberg–Witten trajectory $x(t) = (a(t), \phi(t))$ is said to be of *finite type* if both $\mathcal{L}(x(t))$ and $\|\phi(t)\|_{\mathcal{G}^0}$ are bounded functions of t .

2.2 Restriction to the global Coulomb slice

Define the *global Coulomb slice* (with respect to A_0) to be the subspace

$$V = \text{Ker}(d^*) \oplus \Gamma(S) \subset C(Y).$$

Given $(a, \phi) \in C(Y)$, there exists an element of V which is gauge equivalent to (a, ϕ) , namely

$$(a - df, e^f\phi)$$

where $d^*(a - df) = 0$, so $\Delta f = d^*(a)$. If we impose the condition $\int_Y f d\text{vol}_Y = 0$, then there is a unique solution to these equations given by $f = Gd^*a$, where G is the Green's operator for the Laplacian $\Delta = dd^*$ on functions.

We have a globally defined map $\Pi: C(Y) \rightarrow V$, called the *global Coulomb projection*,

$$\Pi(a, \phi) = (a - df, e^f\phi)$$

where $\Delta f = d^*(a)$ and $\int_Y f d\text{vol} = 0$.

Restricting to the global Coulomb slice V uses up all of the gauge symmetry except for the S^1 subgroup of constant gauge transformations. Instead of working on $C(Y)$ with full gauge symmetry, we work on V with S^1 symmetry.

As $b_1(Y) = 0$, every map $u: Y \rightarrow S^1$ can be written as $u = e^f$ for some $f: Y \rightarrow i\mathbb{R}$. Moreover, f is unique up to addition of an integer multiple of $2\pi i$. We define \mathcal{G}_0 to be the subgroup of gauge transformations of the form $u = e^f$ for some $f: Y \rightarrow i\mathbb{R}$ with $\int_Y f \, d\text{vol} = 0$. It is easy to see that $\mathcal{G} = \mathcal{G}_0 \times S^1$.

We have that \mathcal{G}_0 acts freely on $C(Y)$ and the quotient space can be identified with V . This determines a metric \tilde{g} on V as follows. Take the restriction of the L^2 -metric on $C(Y)$ to the subbundle of the tangent bundle orthogonal to the gauge orbits. This construction is \mathcal{G}_0 -invariant and descends to a metric \tilde{g} on V .

The Chern–Simons–Dirac functional \mathcal{L} is gauge invariant; hence the gradient $\text{grad}(\mathcal{L})$ is orthogonal to the gauge orbits. It follows that the projection of $\text{grad}(\mathcal{L})$ to V coincides with taking the gradient of $\mathcal{L}|_V$ with respect to \tilde{g} . So the trajectories of $\text{grad}(\mathcal{L})$ on $C(Y)$ project to the trajectories of $\mathcal{L}|_V$, where the gradient of $\mathcal{L}|_V$ is taken using the metric \tilde{g} . Thus the trajectories on V have the form

$$\frac{d}{dt}(a(t), \phi(t)) = (- * da - \tau(\phi, \phi), -D_a\phi) - (-df, f\phi)$$

for a function $f: Y \rightarrow i\mathbb{R}$. The function f is uniquely determined by the conditions that $\int_Y f \, d\text{vol}_Y = 0$ and that $*da + \tau(\phi, \phi) - df$ is in the kernel of d^* . Hence $df = (1 - \pi)\tau(\phi, \phi)$, where π denotes the L^2 orthogonal projection to $\text{Ker}(d^*)$. We have that

$$\frac{d}{dt}(a(t), \phi(t)) = (- * da - \pi\tau(\phi, \phi), -D_a\phi - f\phi) = -(l + c)(a, \phi),$$

where

$$l(a, \phi) = (*da, D\phi)$$

is the linear part and

$$c(a, \phi) = (\pi\tau(\phi, \phi), \rho(a)\phi + f\phi)$$

is given by the nonlinear terms.

Let χ denote the gradient of $\mathcal{L}|_V$ with respect to \tilde{g} . Then $\chi = l + c$ extends to a map

$$\chi = l + c: V_{k+1} \rightarrow V_k,$$

where V_k denotes the L^2_k -Sobolev completion of V . The map l is a linear Fredholm operator. Using Sobolev multiplication and an estimate on the unique solution to $df = (1 - \pi)\tau(\phi, \phi)$, $\int_Y f \, d\text{vol}_Y = 0$, it follows that c viewed as a map $V_{k+1} \rightarrow V_{k+1}$ is continuous. Hence $c: V_{k+1} \rightarrow V_k$ is compact. The flow lines of χ on V will be called Seiberg–Witten trajectories in the Coulomb gauge. We say that such a trajectory $x(t) = (a(t), \phi(t))$ is of finite type if $\mathcal{L}(x(t))$ and $\|\phi(t)\|_{\mathcal{G}_0}$ are bounded independent of t . Clearly the finite type Seiberg–Witten trajectories in the Coulomb gauge are precisely the projection to V of the Seiberg–Witten trajectories in $C(Y)$ of finite type.

2.3 Finite-dimensional approximation

Let V_λ^μ denote the direct sum of all eigenspaces of l in the range $(\lambda, \mu]$ and let \tilde{p}_λ^μ be the L^2 -orthogonal projection from V to V_λ^μ . Note that V_λ^μ is a finite-dimensional subspace of V . For technical reasons we replace the projections \tilde{p}_λ^μ with smoothed out versions

$$p_\lambda^\mu = \int_0^1 \rho(\theta) \tilde{p}_{\lambda+\theta}^{\mu-\theta} d\theta$$

where $\rho: \mathbb{R} \rightarrow \mathbb{R}$ is smooth, nonnegative, nonzero precisely on $(0, 1)$ and $\int_{\mathbb{R}} \rho(\theta) d\theta = 1$. This is to make p_λ^μ vary continuously with μ and λ . The reason for doing this is to show that the Conley index is independent of the choices of μ and λ , up to a suspension. This is achieved by continuously increasing or decreasing μ and λ to get a continuous family of flows and using homotopy invariance of the Conley index under continuous deformation of the flow.

Consider the gradient flow equation

$$\frac{d}{dt}x(t) = -(l + p_\lambda^\mu c)x(t),$$

where $x: \mathbb{R} \rightarrow V_\lambda^\mu$. We call this an *approximate Seiberg–Witten trajectory*.

Let $B(R)$ denote the open ball of radius R in $L_{k+1}^2(V)$. Using the a priori estimates for the Seiberg–Witten equations, it can be shown that there exists an $R > 0$ such that all the finite type trajectories of $l + c$ are in $B(R)$ [49, Proposition 1]. This boundedness property does not necessarily hold for approximate trajectories, since the crucial estimates that hold for the Seiberg–Witten equations do not apply to the approximate trajectories. However, we have the following result which acts as a kind of substitute:

Proposition 2.2 [49, Proposition 3] *For any $-\lambda$ and μ sufficiently large, if an approximate trajectory $x: \mathbb{R} \rightarrow L_{k+1}^2(V_\lambda^\mu)$ satisfies $x(t) \in \overline{B(2R)}$ for all t , then in fact $x(t) \in B(R)$ for all t .*

This result will allow us to construct the Seiberg–Witten–Floer homotopy type of (Y, \mathfrak{s}) using Conley indices.

2.4 The Conley index

Suppose we have a 1-parameter group $\{\varphi_t\}$ of diffeomorphisms of an n -dimensional manifold M (not necessarily compact). The example to keep in mind is the gradient flow of a Morse function. Given a compact subset $N \subseteq M$, the invariant set of N is

$$\text{Inv}(N, \varphi) = \{x \in N \mid \varphi_t(x) \in N \text{ for all } t \in \mathbb{R}\}.$$

A compact subset $N \subseteq M$ is called an *isolating neighbourhood* if $\text{Inv}(N, \varphi) \subseteq \text{int } N$. An *isolated invariant set* is a subset $S \subseteq M$ such that $S = \text{Inv}(N, \varphi)$ for some isolating neighbourhood. Note that S must be compact since it is a closed subset of N and N is required to be compact.

Definition 2.3 Let S be an isolated invariant set. An *index pair* (N, L) for S is a pair of compact sets $L \subseteq N \subseteq M$ such that

- $\text{Inv}(N - L, \varphi) = S \subseteq \text{int}(N - L)$;
- L is an exit set for N , that is, for all $x \in N$, if there exists $t > 0$ such that $\varphi_t(x)$ is not in N , then there exists τ with $0 \leq \tau < t$ with $\varphi_\tau(x) \in L$;
- L is positively invariant in N , that is, if $x \in L$ and $t > 0$ and such that $\varphi_s(x) \in N$ for all $0 \leq s \leq t$, then $\varphi_s(x) \in L$ for all $0 \leq s \leq t$.

Any isolated invariant set S admits an index pair (N, L) . The *Conley index* of S is the based homotopy type

$$I(S) = (N/L, [L]).$$

The Conley index is independent of the choice of index pair (N, L) in a strong way. Namely for any two pairs (N_1, L_1) and (N_2, L_2) , there is a canonical homotopy equivalence $N_1/L_1 \cong N_2/L_2$. The composition of two such canonical homotopy equivalences $N_1/L_1 \cong N_2/L_2$ and $N_2/L_2 \cong N_3/L_3$ coincides up to homotopy with the canonical homotopy equivalence $N_1/L_1 \cong N_3/L_3$ (one says that the collection of Conley indices N/L forms a *connected simple system*). By abuse of terminology, if (N, L) is an index pair for S we say that $I = N/L$ is “the” Conley index of S .

Example 2.4 Consider a Morse function with critical point of index p , say

$$f(x_1, \dots, x_n) = \frac{1}{2}(-x_1^2 - \dots - x_p^2 + x_{p+1}^2 + \dots + x_n^2).$$

The negative gradient of f using the Euclidean metric is

$$-\text{grad}(f)(x) = (x_1, \dots, x_p, -x_{p+1}, \dots, -x_n).$$

It follows that the downwards gradient flow is given by

$$\varphi_t(x) = (e^t x_1, \dots, e^t x_p, e^{-t} x_{p+1}, \dots, e^{-t} x_n).$$

Let $S = \{0\}$ be the critical point. This is an isolated invariant set. In fact, the only invariant point of φ is the origin, so we could take $N = D^p \times D^{n-p}$ as an isolating neighbourhood (where D^j is the closed j -dimensional unit disc). Then $L = S^{p-1} \times D^{n-p}$ is an exit set for N . It is easy to see that (N, L) satisfies the condition for an index pair for S . The Conley index is $I(S) = D^p \times D^{n-p} / (S^{p-1} \times D^{n-p})$, which is homotopy equivalent to S^p , a p -dimensional sphere.

Example 2.5 If M is a compact manifold and φ is a Morse–Smale gradient flow on M , then the set S of all critical points and all flow lines between them is an isolated invariant set. The reduced homology of $I(S)$ is known to be isomorphic to the homology of M .

On the other hand, if M is noncompact, then we cannot take S to be all critical points of M and all flow lines starting or terminating at a critical point, because there could be flow lines going off to $-\infty$ or coming in from $+\infty$ and then S would not be compact.

We also need the equivariant Conley index. Let G be a compact Lie group acting smoothly on M , preserving a flow φ and an isolated invariant set S . It turns out that one can find a G -invariant index pair (N, L) for S and one can define the G -equivariant Conley index to be the pointed G -equivariant homotopy type

$$I_G(S) = (N/L, [L]).$$

It can be shown that this is well defined, up to G -equivariant homotopy equivalence. Moreover $I_G(S)$ has the based homotopy type of a finite G -CW complex.

Example 2.6 Consider again the example of the Morse function on \mathbb{R}^n given by

$$f(x_1, \dots, x_n) = \frac{1}{2}(-x_1^2 - \dots - x_p^2 + x_{p+1}^2 + \dots + x_n^2).$$

Now suppose that G is a compact Lie group which acts linearly on \mathbb{R}^n preserving f . Note that f defines an $O(n-p, p)$ -structure on \mathbb{R}^n and the fact that G preserves f just means that the action of G on \mathbb{R}^n factors through a homomorphism $G \rightarrow O(n-p, p)$. As G is compact, we may as well assume (after a linear change of coordinates) that G maps to the maximal compact subgroup $O(n-p) \times O(p)$. So we can decompose \mathbb{R}^n as

$$\mathbb{R}^n = V_+ \oplus V_-,$$

where V_+ and V_- are real orthogonal representations of G of dimensions $n-p$ and p respectively. Once again, take $S = \{0\}$ as our isolated invariant set. As our Conley index, we can take $N = D(V_-) \times D(V_+)$ and $L = S(V_-) \times D(V_+)$; hence

$$I_G(S) = D(V_-) \times D(V_+) / S(V_-) \times D(V_+) \cong D(V_-) / S(V_-) \cong (V_-)^+,$$

where $(V_-)^+$ is the one-point compactification of V_- . We see that the action of G on the Conley index is determined by the representation of G on the subspace of the tangent space at the critical point in the direction of the negative eigenvalues of the Hessian of f .

Let G and H be compact Lie groups and suppose that $G \times H$ acts smoothly on M . Suppose that $\{\varphi_t\}$ is a $G \times H$ -invariant flow. Then G acts smoothly on the submanifold M^H and the restriction of $\{\varphi_t\}$ defines a G -invariant flow on M^H . In such a situation we can consider the relation between $G \times H$ -equivariant Conley indices for the flow on M and G -equivariant Conley indices for the restriction of the flow to M^H .

Proposition 2.7 *Let $G \times H$ act smoothly on M , preserving a flow $\{\varphi_t\}$, and let (N, L) be a $G \times H$ -equivariant index pair for an isolated invariant set $S = \text{Inv}(A)$. Then (N^H, L^H) is a G -equivariant index pair for the isolated invariant set $S^H = \text{Inv}(A^H)$. Moreover, $(N/L)^H \cong N^H/L^H$.*

Proof First note that A^H is compact because A is compact. Moreover,

$$\text{Inv}(A^H) = \{a \in A^H \mid \varphi_t(a) \in A^H \text{ for all } t\} = \text{Inv}(A) \cap A^H = S \cap A^H = S^H.$$

Let M be a topological space and let $P, Q \subseteq M$ be subspaces. Give Q the induced topology. Then

$$\text{int}_M(P) \cap Q = \left(\bigcup_{U \subseteq P \text{ open in } M} U \right) \cap Q = \bigcup_{U \subseteq P \text{ open in } M} (U \cap Q) \subseteq \text{int}_Q(P \cap Q).$$

Applying this to $P = A$ and $Q = M^H$, we get

$$\text{int}_M(A) \cap M^H \subseteq \text{int}_{M^H}(A^H).$$

Then since $S \subseteq \text{int}_M(A)$ by the assumption that A is an isolating neighbourhood, it follows that

$$S^H \subseteq \text{int}_M(A) \cap M^H \subseteq \text{int}_{M^H}(A^H).$$

So A^H is an isolating neighbourhood in M^H for S^H .

Now let (N, L) be an index pair for S . So N and L are compact and $L \subseteq N$. This implies that N^H and L^H are compact and $L^H \subseteq N^H$. Next, since

$$\text{Inv}(N - L) = S \subseteq \text{int}_M(N - L),$$

it follows that

$$\begin{aligned} \text{Inv}(N^H - L^H) = S^H &= S \cap M^H \subseteq \text{int}_M(N - L) \cap M^H \\ &\subseteq \text{int}_{M^H}((N - L) \cap M^H) = \text{int}_{M^H}(N^H - L^H). \end{aligned}$$

We verify that L^H is an exit set for N^H . Let $x \in N^H$ and suppose $\varphi_t(x) \notin N^H$ for some $t > 0$. Then it follows that $\varphi_t(x) \notin N$, for if $\varphi_t(x) \in N$, then it would imply that $\varphi_t(x) \in N \cap M^H = N^H$, since φ_t preserves M^H . But L is an exit set for N , so there exists $\tau \in [0, t)$ with $\varphi_\tau(x) \in L$. It follows that $\varphi_\tau(x) \in L \cap M^H = L^H$. Hence L^H is an exit set for N^H .

We check that L^H is positively invariant in N^H . Suppose $x \in L^H$ and there exists a $t > 0$ for which $\varphi_s(x) \in N^H$ for all $s \in [0, t]$. Then since L is positively invariant in N , it follows that $\varphi_s(x) \in L$ for all $s \in [0, t]$. Hence $\varphi_s(x) \in L \cap M^H = L^H$ for all $s \in [0, t]$.

We have verified that (N^H, L^H) is an index pair for S^H . Moreover it is straightforward to check that $(N/L)^H = N^H/L^H$. □

2.5 Equivariant Spanier–Whitehead category

In this section we recall the construction of the category \mathfrak{C} from [49], which is an S^1 –equivariant version of the Spanier–Whitehead category. In Section 3.3 we will modify this construction to accommodate a finite group action on Y .

We work with pointed topological spaces with a basepoint-preserving action of S^1 . The objects of \mathfrak{C} are triples (X, m, n) , where X is a pointed topological space with S^1 –action, and $m, n \in \mathbb{Z}$.¹ We further

¹In [49] n is allowed to take on rational values. This is needed to construct a Seiberg–Witten–Floer spectrum which does not depend on the choice of metric. For our purposes it suffices to consider only integral values of n .

require that X has the S^1 -homotopy type of an S^1 -CW complex, which holds for Conley indices on manifolds. The set of morphisms between two objects (X, m, n) to (X', m', n') will be denoted by $\{(X, m, n), (X', m', n')\}_{S^1}$ and is defined to be

$$\{(X, m, n), (X', m', n')\}_{S^1} = \operatorname{colim}[(\mathbb{R}^k \oplus \mathbb{C}^l)^+ \wedge X, (\mathbb{R}^{k+m-m'} \oplus \mathbb{C}^{l+n-n'})^+ \wedge X']_{S^1},$$

where $[\cdot, \cdot]_{S^1}$ denotes the set of S^1 -equivariant homotopy classes and the colimit is taken over all k and l such that $k \geq m' - m$ and $l \geq n' - n$. The maps that define the colimit are given by suspensions where we smash on the left and for any topological space Z , we let Z^+ denote the one-point compactification with its obvious basepoint.

Any pointed space X with S^1 -action defines an object of \mathfrak{C} , namely $(X, 0, 0)$. We often simply write this as X . For any finite-dimensional representation E of S^1 , we let Σ^E denote the reduced suspension operation

$$\Sigma^E X = E^+ \wedge X.$$

This operation extends to \mathfrak{C} by taking $\Sigma^E(X, m, n) = (\Sigma^E X, m, n)$. We are mainly interested in the case that E is a real vector space with trivial S^1 -action, or E is a complex vector space with S^1 acting by scalar multiplication. If E is a real vector space with trivial action, then one finds that

$$\Sigma^E(X, m, n) \cong (X, m - \dim_{\mathbb{R}}(E), n).$$

The isomorphism depends on a choice of isomorphism $E \cong \mathbb{R}^{\dim_{\mathbb{R}}(E)}$. Up to homotopy there are two choices since $\operatorname{GL}(E, \mathbb{R})$ has two components. If E is a complex vector space and S^1 acts by scalar multiplication, then

$$\Sigma^E(X, m, n) \cong (X, m, n - \dim_{\mathbb{C}}(E)).$$

The isomorphism is unique up to homotopy as $\operatorname{GL}(E, \mathbb{C})$ is connected. We can define desuspension by a real vector space E with trivial S^1 -action as

$$\Sigma^{-E}(X, m, n) = ((E)^+ \wedge X, m + 2 \dim_{\mathbb{R}}(E), n).$$

Then $\Sigma^{-E} \Sigma^E Z \cong Z$ by an isomorphism which is canonical up to homotopy. We can define desuspension by a complex vector space E with S^1 acting by scalar multiplication by

$$\Sigma^{-E}(X, m, n) = (X, m, n + \dim_{\mathbb{C}}(E)).$$

Then $\Sigma^{-E} \Sigma^E Z \cong Z$ by an isomorphism which is canonical up to homotopy.

For $Z = (X, m, n) \in \mathfrak{C}$, we define the reduced equivariant cohomology of Z to be

$$\tilde{H}_{S^1}^j(Z) = \tilde{H}_{S^1}^{j+m+2n}(X).$$

The cohomology is well defined as a consequence of the Thom isomorphism.

2.6 Seiberg–Witten–Floer cohomology

Consider as before a rational homology 3–sphere Y and a spin^c –structure \mathfrak{s} . Let $R > 0$, λ and μ be as in Proposition 2.2. We want to take the Conley index of the set of all critical points in $B(R)$ and flow lines between them which lie in $B(R)$ for all time for the approximate Seiberg–Witten flow $l + p_\lambda^\mu c$. The problem is that there could be trajectories that go to infinity in a finite amount of time. Hence we do not have a flow $\{\varphi_t\}$ in the sense of a 1–parameter group of diffeomorphisms. To get around this issue, let u_λ^μ be a compactly supported smooth cutoff function which is identically 1 on $B(3R)$. For consistency purposes we assume that $u_\lambda^\mu = u_{\lambda'}^{\mu'}|_{V_\lambda^\mu}$ for $\lambda' \leq \lambda$ and $\mu' \geq \mu$. One way of doing this is to take $u_\lambda^\mu(v) = \rho(\|v\|)$, where ρ is smooth, compactly supported and $\rho(t) = 1$ for $t < 3$.

For each μ and λ , the vector field $u_\lambda^\mu(l + p_\lambda^\mu c)$ is compactly supported, so it generates a well-defined flow $\varphi_{\lambda,t}^\mu$ on V_λ^μ . Since $u_\lambda^\mu = 1$ on $\overline{B(2R)}$, Proposition 2.2 still applies to the trajectories of $u_\lambda^\mu(l + p_\lambda^\mu c)$. It follows that

$$\text{Inv}(V_\lambda^\mu \cap \overline{B(2R)}) = S_\lambda^\mu,$$

where S_λ^μ is the set of critical points and flow lines between critical points for the approximate Seiberg–Witten flow $l + p_\lambda^\mu c$ which lie in $B(R)$. Therefore S_λ^μ is an isolated invariant set. Moreover, S^1 preserves the approximate flow; hence we may take the S^1 –equivariant Conley index

$$I_\lambda^\mu = I_{S^1}(S_\lambda^\mu).$$

This is an S^1 –equivariant homotopy type. However it is not quite an invariant of (Y, \mathfrak{s}) because it depends on the choice of metric g as well as the values of λ , μ and R . Note that it is independent of the choice of u_λ^μ because of the assumption that $u_\lambda^\mu = 1$ on $B(3R)$. To get a genuine invariant we must understand how I_λ^μ changes as we vary these parameters.

Let λ^* and μ^* satisfy Proposition 2.2. Suppose that $\lambda' \leq \lambda \leq \lambda^*$ and $\mu' \geq \mu \geq \mu^*$. We wish to compare the Conley indices I_λ^μ , $I_\lambda^{\mu'}$ and $I_{\lambda'}^{\mu'}$. In other words, what happens if we increase either μ or $-\lambda$, staying in the range where μ and $-\lambda$ are sufficiently large.

We use the following invariance property of the Conley index: Suppose we have a family $\{\varphi_t(s)\}$ of flows depending continuously on $s \in [0, 1]$. Suppose that a fixed compact set A is an isolating neighbourhood for all $s \in [0, 1]$ and let $S(s) = \text{Inv}(A, \varphi_t(s))$. Then $I(S_0, \varphi_t(0)) \cong I(S_1, \varphi_t(1))$ by a canonical homotopy equivalence.

Consider increasing μ to μ' . The finite energy trajectories of $l + p_\lambda^\mu c p_\lambda^\mu$ in $V_\lambda^{\mu'}$ must actually lie in V_λ^μ . Therefore $A = \overline{B(2R)} \cap V_\lambda^{\mu'}$ is an isolating neighbourhood for S_λ^μ in $V_\lambda^{\mu'}$. Let $\mu(s) = (1 - s)\mu + s\mu'$ for $s \in [0, 1]$ and let $\tilde{\varphi}_\lambda^{\mu(s)}$ denote the flow of $u_\lambda^{\mu(s)}(l + p_\lambda^{\mu(s)} c p_\lambda^{\mu(s)})$ on $V_\lambda^{\mu(s)}$. Then for each $s \in [0, 1]$, A is an isolating neighbourhood for $S_\lambda^{\mu(s)}$ in $V_\lambda^{\mu(s)}$ with respect to the flow $\tilde{\varphi}_\lambda^{\mu(s)}$. Hence

$$\text{Inv}(\tilde{\varphi}_\lambda^\mu, A) \cong \text{Inv}(\tilde{\varphi}_\lambda^{\mu'}, A) = I_\lambda^{\mu'}.$$

But $\tilde{\varphi}^\mu$ is easily seen to be homotopic to the product of the flow of $u_\lambda^\mu(l + p_\lambda^\mu c)$ on V_λ^μ with a linear flow on W generated by $l|_W$, where W is the orthogonal complement of V_λ^μ in $V_\lambda^{\mu'}$. The Conley index of a product of flows is just the smash product of Conley indices. Combined with Example 2.6, we see that

$$\text{Inv}(\tilde{\varphi}_\lambda^\mu, A) \cong I_\lambda^\mu \wedge W_-^+,$$

where W_- is the part of W spanned by negative eigenvalues of l . But W is contained in the positive eigenvalues of l , so $W_- = 0$ and hence

$$I_\lambda^{\mu'} \cong I_\lambda^\mu.$$

Now consider decreasing λ to λ' . An identical argument to the one above gives

$$I_{\lambda'}^\mu \cong I_\lambda^\mu \wedge W_-^+,$$

where W is the orthogonal complement of V_λ^μ in $V_{\lambda'}^\mu$. In this case W is spanned by negative eigenspaces of l , so $W_- = W = V_{\lambda'}^\lambda$ and

$$I_{\lambda'}^\mu \cong I_\lambda^\mu \wedge (V_{\lambda'}^\lambda)^+.$$

This implies that

$$\Sigma^{-V_\lambda^0} I_\lambda^\mu$$

does not depend on the values of μ and λ (provided μ and $-\lambda$ are sufficiently large).

Definition 2.8 Given (Y, \mathfrak{s}) and a metric g , we set

$$SWF(Y, \mathfrak{s}, g) = \Sigma^{-V_\lambda^0(g)} I_\lambda^\mu(g)$$

for suitably chosen μ, λ and R .

We have established that the homotopy type of $SWF(Y, \mathfrak{s}, g)$ does not depend on the choices of μ and λ , or more precisely, any two choices of μ and λ are related by a canonical homotopy equivalence. One also checks that it does not depend on the choice of R . So up to homotopy, $SWF(Y, \mathfrak{s}, g)$ depends only on Y, \mathfrak{s} and g .

Next we consider varying the metric g . Consider a smooth homotopy g_s for $s \in [0, 1]$ joining two metrics g_0 and g_1 , which is constant near $s = 0$. Assuming that the g_s are all sufficiently close to each other in a suitable topology, we can arrange that there exists R, μ^* and λ^* such that Proposition 2.2 is true for all $s \in [0, 1]$ and all μ and λ with $\mu \geq \mu^*$ and $\lambda \leq \lambda^*$. This suffices, as compactness of $[0, 1]$ implies that any smooth path g_s can be broken up into finitely many subpaths over which this assumption holds.

We assume that there exists some $\lambda < \lambda^*$ and $\mu > \mu^*$ such that λ and μ are not eigenvalues of l_s for any $s \in [0, 1]$. This property will hold for all sufficiently small paths. The spaces $(V_\lambda^\mu)_s$ then form a smooth vector bundle over $[0, 1]$. We can trivialise this vector bundle and identify all these spaces with a single V_λ^μ . Further, we assume that $B(R)_{s_1} \subset B(2R)_{s_2}$ for each $s_1, s_2 \in [0, 1]$. Here we think of the balls as subsets of the same space V_λ^μ . Once again, this property will hold for all small enough paths. Then

$$\bigcap_{s \in [0, 1]} \overline{B(2R)_s}$$

is a compact isolating neighbourhood for S_λ^μ in any metric g_s with the flow $(\varphi_\lambda^\mu)_s$. The Conley index will be independent of s and hence

$$(I_\lambda^\mu)_0 \cong (I_\lambda^\mu)_1.$$

However, we do not have that $\Sigma^{-(V_\lambda^0)_0}(I_\lambda^\mu)_0$ equals $\Sigma^{-(V_\lambda^0)_1}(I_\lambda^\mu)_1$. The reason is that some eigenvalues in (λ, μ) may change sign. On the other hand, any eigenvalue greater than μ or less than λ cannot change sign, by our assumption that μ and λ are not eigenvalues of l_s for any $s \in [0, 1]$. Hence the difference between $(V_\lambda^0)_0$ and $(V_\lambda^0)_1$ is given in terms of the spectral flow of the family of operators $\{l_s\}$ for $s \in [0, 1]$.

The operator l can be split into real and complex components. The real part has no spectral flow, so we only need to consider the complex part, which is the Dirac operator D_s . The spectral flow can be expressed using the Atiyah–Patodi–Singer (APS) index theorem on the cylinder $X = [0, 1] \times Y$; see [8; 9]. Let \hat{g} be the metric on X given by g_s in the vertical direction and $(ds)^2$ in the horizontal direction. Let S_s denote the spinor bundle associated to (s, g_s) . The bundles S_s can all be identified with $S = S_0$, but with varying Clifford multiplication. The spin^c -structure s lifts to a spin^c -structure on X . Let S^\pm denote the spinor bundles of this spin^c -structure. Then S^\pm can be identified with the pullback of S to X . Suppose for each s we have chosen a flat reference connection A_s . Since we have identified S_s with S for all s , we get an induced identification of $L_s = \det(S_s)$ with $L = L_0$. Then $A_s = A_0 + i\alpha_s$ for some closed real 1-form α_s . The path of spin^c -connections $\{A_s\}$ fit together to form a spin^c -connection \hat{A} on the determinant line L pulled back to X . Let \hat{D} be the Dirac operator determined by \hat{g} and \hat{A} . Then $\hat{D}(\psi) = \partial_s \psi + D_s \psi$. After a possible reparametrisation we can assume that (g_s, A_s) is constant near the boundary. Applying the APS index theorem to the Dirac operator \hat{D} on the cylinder $[0, 1] \times Y$, one can write the spectral flow $SF(\{D_s\})$ as

$$SF(\{D_s\}) = \frac{1}{2}(\eta(D_1) - k(D_1)) - \frac{1}{2}(\eta(D_0) - k(D_0)) + \int_{[0,1] \times Y} \left(-\frac{1}{24}p_1(\hat{g}) + \frac{1}{8}c_1(\hat{A})^2\right),$$

where $\eta(D)$ is the eta invariant of D , $k(D) = \dim_{\mathbb{C}}(\text{Ker}(D))$, $p_1(\hat{g})$ is the first Pontryagin form of \hat{g} , and $c_1(\hat{A})$ is the Chern form $(i/2\pi)F_{2\hat{A}}$, where $F_{2\hat{A}}$ is the curvature of the induced connection $2\hat{A}$ on L . Now since α_s is closed for each s , we get $F_{2\hat{A}} = ds \wedge 2i\partial_s \alpha_s$ and hence $c_1(\hat{A})^2 = 0$. So

$$(2-1) \quad SF(\{D_s\}) = \frac{1}{2}(\eta(D_1) - k(D_1)) - \frac{1}{2}(\eta(D_0) - k(D_0)) - \frac{1}{24} \int_{[0,1] \times Y} p_1(\hat{g}).$$

Let $\eta_{\text{sign}}(g_s)$ denote the eta invariant of the signature operator on Y defined by g_s . Then from the APS index theorem for the signature operator together with the fact that the signature operator has no spectral flow, we find

$$(2-2) \quad \eta_{\text{sign}}(g_1) - \eta_{\text{sign}}(g_0) = \frac{1}{3} \int_{[0,1] \times Y} p_1(\hat{g}).$$

Combining (2-1) and (2-2), we see that

$$SF(\{D_s\}) = \frac{1}{2}(\eta(D_1) - k(D_1)) - \frac{1}{2}(\eta(D_0) - k(D_0)) - \frac{1}{8}(\eta_{\text{sign}}(g_1) - \eta_{\text{sign}}(g_0)),$$

and hence

$$SF(\{D_s\}) = n(Y, \mathfrak{s}, g_0) - n(Y, \mathfrak{s}, g_1),$$

where we have defined

$$n(Y, \mathfrak{s}, g) = \frac{1}{2}\eta(D) - \frac{1}{2}k(D) - \frac{1}{8}\eta_{\text{sign}}(g).$$

We will show that $n(Y, \mathfrak{s}, g)$ is a rational number. Let (W, \mathfrak{s}_W) be a spin^c 4-manifold bounding (Y, \mathfrak{s}) . This always exists because $\Omega_3^{\text{spin}^c} = 0$. Extend the Dirac operator D on Y to a Dirac operator \widehat{D} in the same way as we did for the cylinder $[0, 1] \times Y$. The APS index theorem for the Dirac operator and signature operator on W combined give

$$\text{ind}_{\text{APS}}(\widehat{D}) = \frac{1}{8}(c_1(\mathfrak{s}_W)^2 - \sigma(W)) + \frac{1}{2}(\eta_{\text{dir}} - k) - \frac{1}{8}\eta_{\text{sign}}$$

and thus

$$(2-3) \quad n(Y, \mathfrak{s}, g) = \text{ind}_{\text{APS}}(\widehat{D}) - \delta(W, \mathfrak{s}),$$

where we set

$$\delta(W, \mathfrak{s}) = \frac{1}{8}(c_1(\mathfrak{s}_W)^2 - \sigma(W)).$$

This shows that $n(Y, \mathfrak{s}, g)$ is a rational number since $\text{ind}_{\text{APS}}(\widehat{D})$ is an integer and $\delta(W, \mathfrak{s})$ is a rational number.

Definition 2.9 The *Seiberg–Witten–Floer cohomology* of (Y, \mathfrak{s}, g) is defined as

$$HSW^j(Y, \mathfrak{s}) = \widetilde{H}_{S^1}^{j+2n(Y, \mathfrak{s}, g)}(SWF(Y, \mathfrak{s}, g)),$$

where $j \in \mathbb{Q}$ and as usual the coefficient group \mathbb{F} has been omitted from the notation.

Below we will show that $HSW^*(Y, \mathfrak{s})$ is independent of the choice of metric g (and other auxiliary choices); hence it is a well defined topological invariant of the pair (Y, \mathfrak{s}) .

Notice that because of the grading shift by $2n(Y, \mathfrak{s}, g)$ the cohomology groups $HSW^*(Y, \mathfrak{s})$ are concentrated in rational degrees. It was shown by Lidman and Manolescu [45] that $HSW^*(Y, \mathfrak{s})$ is isomorphic to the Seiberg–Witten monopole Floer cohomology as defined by Kronheimer and Mrowka [39]. Together with the equivalence of monopole Floer homology and Heegaard Floer homology due to the work of Kutluhan, Lee and Taubes [40; 41; 42; 43; 44], Colin, Ghiggini and Honda [16; 17; 15] and Taubes [60; 61; 62; 63; 64], we have isomorphisms

$$HSW^*(Y, \mathfrak{s}) \cong HF_{-*}^-(\overline{Y}, \mathfrak{s}) \cong HF_+^*(Y, \mathfrak{s}),$$

where HF_{-*}^- denotes the minus version of Heegaard Floer homology and HF_+^* denotes the plus version of Heegaard Floer *cohomology* (taken with respect to the same coefficient group \mathbb{F}). Here we use a grading convention for HF^- such that $HF^-(S^3)$ starts in degree 0. Through the work of [20; 33; 58], the isomorphism is known to preserve the absolute gradings. Using co-Borel, Tate or nonequivariant

cohomologies gives similar isomorphisms to the other versions of Heegaard Floer homology; see [45, Corollary 1.2.4] for the precise statement.

If we have two metrics g_0 and g_1 , then the spectral flow of a path joining them satisfies

$$SF(\{D_s\}) = n(Y, \mathfrak{s}, g_1) - n(Y, \mathfrak{s}, g_0).$$

On the other hand, from the definition of spectral flow,

$$SF(\{D_s\}) = \dim(V_\lambda^0(g_0)) - \dim(V_\lambda^0(g_1)).$$

It follows that

$$SWF(Y, \mathfrak{s}, g_1) \cong \Sigma^{SF(\{D_s\})} SWF(Y, \mathfrak{s}, g_0),$$

and hence

$$\tilde{H}_{S^1}^{j+2SF(\{D_s\})}(SWF(Y, \mathfrak{s}, g_1)) \cong \tilde{H}_{S^1}^j(SWF(Y, \mathfrak{s}, g_0))$$

by the Thom isomorphism. Replacing j by $j + 2n(Y, \mathfrak{s}, g_0)$, we have

$$\tilde{H}_{S^1}^{j+2n(Y, \mathfrak{s}, g_1)}(SWF(Y, \mathfrak{s}, g_1)) \cong \tilde{H}_{S^1}^{j+2n(Y, \mathfrak{s}, g_0)}(SWF(Y, \mathfrak{s}, g_0)).$$

Hence the Seiberg–Witten–Floer cohomology $\tilde{H}_{S^1}^{j+2n(Y, \mathfrak{s}, g)}(SWF(Y, \mathfrak{s}, g))$ is independent of the metric. The above isomorphism is canonical in the sense that it does not depend on the choice of path from g_0 to g_1 . This follows from the fact that the space of all metrics on Y is contractible, so any two paths with the same endpoints are homotopic.

2.7 Duality

Definition 2.10 Let V be a finite-dimensional representation of a compact Lie group G . Two pointed, finite G -CW complexes X and X' are equivariantly V -dual if there exists a G -map

$$\varepsilon: X \wedge X' \rightarrow V^+$$

such that for any subgroup $H \subseteq G$, the fixed-point map

$$\varepsilon^H: X^H \wedge (X')^H \rightarrow (V^H)^+$$

induces a nonequivariant duality between X^H and $(X')^H$, in the sense of nonequivariant Spanier–Whitehead duality.

Consider the Conley index I_λ^μ associated to (Y, \mathfrak{s}, g) for suitably chosen R , μ and λ . One finds that reversing orientation of Y has the effect of reversing the Chern–Simons–Dirac flow. From [19], it follows that $I_\lambda^\mu(Y)$ and $I_\mu^\lambda(\bar{Y})$ are V_λ^μ -dual, so there exists a duality map

$$\varepsilon: I_\lambda^\mu(Y) \wedge I_\mu^\lambda(\bar{Y}) \rightarrow (V_\lambda^\mu)^+.$$

Notice that $\dim(V_\lambda^0(Y)) + \dim(V_{-\mu}^0(\bar{Y})) - 2k(D) = \dim(V_\lambda^\mu(Y))$, where $k(D)$ is the dimension of the kernel of D . Desuspending, we obtain a duality map

$$\varepsilon: SWF(Y, \mathfrak{s}, g) \wedge SWF(\bar{Y}, \mathfrak{s}, g) \rightarrow S^{-k(D)\mathbb{C}}.$$

We also have

$$(2-4) \quad n(Y, \mathfrak{s}, g) + n(\bar{Y}, \mathfrak{s}, g) = -k(D).$$

2.8 Fixed points

Definition 2.11 Let $s \geq 0$ be an integer. We say that a finite pointed S^1 -CW complex X is of type SWF at level s if

- the S^1 -fixed-point set X^{S^1} is homotopy equivalent to the sphere $(\mathbb{R}^s)^+$;
- the action of S^1 is free on the complement $X - X^{S^1}$.

Proposition 2.12 Given (Y, \mathfrak{s}, g) , let R, μ and λ be as in Proposition 2.2. Then $\overline{B(2R)} \cap V_\lambda^\mu$ is an isolating neighbourhood for $S_\lambda^\mu = \text{Inv}(\overline{B(2R)} \cap V_\lambda^\mu)$. Let $I_\lambda^\mu = I_{S^1}(S_\lambda^\mu)$ be the Conley index. Then I_λ^μ is of type SWF at level $s = \dim(V_\lambda^0(\mathbb{R}))$, where $V_\lambda^0(\mathbb{R})$ denotes the S^1 -invariant part of V_λ^0 .

Proof Let (N, L) be an index pair for S_λ^μ so that $I_\lambda^\mu = N/L$. Then by Proposition 2.7, $(I_\lambda^\mu)^{S^1} = N^{S^1}/L^{S^1}$ is the Conley index of $(S_\lambda^\mu)^{S^1}$. Further, we have that $\overline{B(2R)} \cap V_\lambda^\mu(\mathbb{R})$ is an isolating neighbourhood for $(S_\lambda^\mu)^{S^1}$, where $V_\lambda^\mu(\mathbb{R})$ denotes the S^1 -invariant part of V_λ^μ . It is easy to see that $c = 0$ on $V_\lambda^\mu(\mathbb{R})$, where c is the nonlinear part of the Seiberg–Witten flow. Thus the restriction of the approximate Seiberg–Witten flow $u_\lambda^\mu(l + p_\lambda^\mu c)$ to $V_\lambda^\mu(\mathbb{R})$ is the flow $u_\lambda^\mu l$. Restricted to $B(3R)$ this is just the linear flow associated to l . The real part of l has zero kernel, because $b_1(Y) = 0$. It follows that the Conley index of $(S_\lambda^\mu)^{S^1}$ is the Conley index of $\{0\}$ in $V_\lambda^\mu(\mathbb{R})$ with respect to the linear flow of l . This is $(V_\lambda^0(\mathbb{R}))^+$. Thus we have shown that the S^1 -fixed-point set of I_λ^μ is homotopy equivalent to $(V_\lambda^0(\mathbb{R}))^+$. Furthermore, S^1 acts freely on $V_\lambda^\mu - V_\lambda^\mu(\mathbb{R})$; hence S^1 acts freely on $N - N^{S^1}$ and therefore also on $(I_\lambda^\mu) - (I_\lambda^\mu)^{S^1}$. \square

Using the identities

$$(2-5) \quad (\mathbb{R}^+ \wedge X)^{S^1} = \mathbb{R}^+ \wedge X^{S^1}, \quad (\mathbb{C}^+ \wedge X)^{S^1} = X^{S^1},$$

we see that

- if X is of type SWF at level s , then $\mathbb{R}^+ \wedge X$ is of type SWF at level $s + 1$;
- if X is of type SWF at level s , then $\mathbb{C}^+ \wedge X$ is of type SWF at level s .

Now let $Z = (X, m, n)$ belong to the equivariant Spanier–Whitehead category \mathcal{C} . We say that Z is of type SWF at level s if X is of type SWF of level $s + m$. The above remarks shows that this is a well-defined notion.

We have shown that the Conley index I_λ^μ is of type SWF at level $s = \dim(V_\lambda^0(\mathbb{R}))$, where

$$V_\lambda^0 = V_\lambda^0(\mathbb{R}) \oplus V_\lambda^0(\mathbb{C})$$

denotes the decomposition of V_λ^0 into copies of \mathbb{R} and \mathbb{C} . Now we recall that

$$SWF(Y, \mathfrak{s}, g) = \Sigma^{-V_\lambda^0} I_\lambda^\mu.$$

It follows that $SWF(Y, \mathfrak{s}, g)$ is of type SWF at level 0.

Let X be a space of type SWF at level s . Let $\iota: X^{S^1} \rightarrow X$ denote the inclusion of the fixed-point set. Using the localisation theorem in equivariant cohomology [23, III (3.8)], it follows that the pullback map $\iota^*: \tilde{H}_{S^1}^*(X) \rightarrow \tilde{H}_{S^1}^*(X^{S^1})$ is not identically zero. Therefore, we may define the d -invariant $d(X)$ of X by

$$d(X) = \min\{j \mid x \in \text{Im}(\iota^*) \text{ for some } x \in \tilde{H}_{S^1}^j(X^{S^1}) \text{ with } x \neq 0\}.$$

Note that $d(X)$ could potentially depend on the choice of coefficient group, so we may write the invariant as $d(X; \mathbb{F})$ if we wish to indicate the dependence on \mathbb{F} .

We also define the δ -invariant of X by $\delta(X) = \frac{1}{2}d(X)$. Using (2-5) and the Thom isomorphism, one finds

$$d(\mathbb{R}^+ \wedge X) = d(X) + 1, \quad d(\mathbb{C}^+ \wedge X) = d(X) + 2.$$

Now if $Z = (X, m, n)$ is of type SWF, we define the d -invariant $d(Z)$ of Z to be

$$d(Z) = d(X) - m - 2n \in \mathbb{Z}.$$

From [45, Corollary 1.2.3], it follows that the d -invariant $d(Y, \mathfrak{s})$ as defined by Heegaard Floer homology (with coefficient group \mathbb{F}) is given in terms of $SWF(Y, \mathfrak{s}, g)$ by

$$d(Y, \mathfrak{s}) = d(SWF(Y, \mathfrak{s}, g)) - 2n(Y, \mathfrak{s}, g).$$

For notational convenience we also define $\delta(Y, \mathfrak{s}) = \frac{1}{2}d(Y, \mathfrak{s})$.

3 Equivariant Seiberg–Witten–Floer cohomology

3.1 Assumption on G and \mathbb{F}

Throughout this paper we will assume that one of the two following conditions hold:

- (1) G is an arbitrary finite group and $\mathbb{F} = \mathbb{Z}/2\mathbb{Z}$, or
- (2) \mathbb{F} is an arbitrary field and the order of G is odd.

Condition (1) ensures that we do not need to concern ourselves with questions of orientability. Condition (2) ensures that any S^1 -central extension \tilde{G} acts orientation-preservingly on all of its finite-dimensional

representations. Hence under either condition, the Thom isomorphism holds without requiring local coefficients,

$$\tilde{H}_{\tilde{G}}^*(X) \cong \tilde{H}_{\tilde{G}}^{*+\dim_{\mathbb{R}}(V)}(V^+ \wedge X).$$

Here X is any \tilde{G} -space and V is any finite-dimensional representation of \tilde{G} .

3.2 Lifting G -actions

Recall that Y denotes a rational homology 3-sphere. Suppose that a finite group G acts on Y by orientation-preserving diffeomorphisms and suppose G preserves the isomorphism class of a spin^c -structure \mathfrak{s} . We will construct a G -equivariant version of the Seiberg–Witten–Floer cohomology of (Y, \mathfrak{s}) .

Choose a G -invariant metric g on Y and a reference spin^c -connection A_0 such that the connection on the determinant line L is flat. Let $g \in G$ and choose a lift $\hat{g}: S \rightarrow S$ of g to the spinor bundle S , which is possible since G preserves the isomorphism class of \mathfrak{s} . Then $\hat{g}^{-1}A_0\hat{g} = A_0 + a$ for some $a \in i\Omega^1(Y)$. Since A_0 and $\hat{g}^{-1}A_0\hat{g}$ are flat, we must have $da = 0$. Moreover, $b_1(Y) = 0$ implies that $a = df$ for some $f: Y \rightarrow i\mathbb{R}$. Setting $\tilde{g} = e^{-f}\hat{g}$, it follows that \tilde{g} is a lift of g which preserves A_0 . Any other lift of g that preserves A_0 is of the form $c\tilde{g}$ with $c \in U(1)$ a constant. Let $G_{\mathfrak{s}}$ denote the set of all possible lifts of elements of G which preserve A_0 . Then $G_{\mathfrak{s}}$ is a group and we have a central extension

$$1 \rightarrow S^1 \rightarrow G_{\mathfrak{s}} \rightarrow G \rightarrow 1.$$

Now we carry out the construction of the Conley index of a finite-dimensional approximation of the Chern–Simons–Dirac flow $G_{\mathfrak{s}}$ -equivariantly, instead of just S^1 -equivariantly.

3.3 $G_{\mathfrak{s}}$ -equivariant Spanier–Whitehead category

In this section \tilde{G} denotes any S^1 central extension of G . We will construct a category $\mathfrak{C}(\tilde{G})$, the \tilde{G} -equivariant version of \mathfrak{C} .

Recall from Section 2.5 that the category \mathfrak{C} was constructed so that there exists a desuspension functor Σ^{-V} for any real vector space V with trivial S^1 -action or any complex vector space where S^1 acts by scalar multiplication. We now construct a category $\mathfrak{C}(\tilde{G})$ in which we can desuspend by real representations of \tilde{G} , where S^1 acts trivially, and by complex representations, where S^1 acts by scalar multiplication. We are lead to consider the following two types of finite-dimensional representations of \tilde{G} :

Type (1) V is a real representation of \tilde{G} and S^1 acts trivially.

Type (2) V is a complex representation \tilde{G} and S^1 acts on V by scalar multiplication.

Type (1) representations correspond canonically to real representations of G .

Type (2) representations correspond to projective unitary representations of G such that the pullback to G of the central extension $S^1 \rightarrow U(n) \rightarrow \text{PU}(n)$ gives an extension isomorphic to \tilde{G} . If \tilde{G} is split,

then type (2) representations are in bijection with complex representations of G . However, the bijection depends on a choice of splitting of \tilde{G} .

To define stable homotopy groups, we need to consider suspensions with explicitly chosen representations. In other words, we need to work at the level of representations and not just isomorphism classes. Let V_1, \dots, V_p be a complete set of irreducible representations of type (1), and W_1, \dots, W_q a complete set of irreducible representations of type (2). Any representation of type (1) is isomorphic to a direct sum of copies of V_1, \dots, V_p and likewise any representation of type (2) is a direct sum of copies of W_1, \dots, W_q .

If $m = (m_1, \dots, m_p), m' = (m'_1, \dots, m'_p) \in \mathbb{Z}^p$, we say $m \geq m'$ if $m_i \geq m'_i$ for each i . If $m \in \mathbb{Z}^p$ satisfies $m \geq 0$, then we set

$$V(m) = V_1^{\oplus m_1} \oplus \dots \oplus V_p^{\oplus m_p}.$$

Similarly, if $n = (n_1, \dots, n_q) \in \mathbb{Z}^q$ satisfies $n \geq 0$, then we set

$$W(n) = W_1^{\oplus n_1} \oplus \dots \oplus W_q^{\oplus n_q}.$$

The category $\mathfrak{C}(\tilde{G})$ has as objects triples (X, m, n) , where

- X is a pointed topological space with a basepoint-preserving \tilde{G} -action and the homotopy type of a \tilde{G} -CW complex;
- $m \in \mathbb{Z}^p$;
- $n \in \mathbb{Z}^q$.

Let (X, m, n) and (X', m', n') be two objects of $\mathfrak{C}(\tilde{G})$. The set of morphisms from (X, m, n) to (X', m', n') , denoted by $\{(X, m, n), (X', m', n')\}^{\tilde{G}}$, is defined to be

$$\operatorname{colim}_{k,l} [(V(k))^+ \wedge (W(l))^+ \wedge X, (V(k+m-m'))^+ \wedge (W(l+n-n'))^+ \wedge X']^{\tilde{G}}.$$

The colimit is taken over all $k \in \mathbb{N}^p$ and $l \in \mathbb{N}^q$ such that $k \geq m' - m$ and $l \geq n' - n$. The maps that define the colimit are given by suspensions where we smash on the left.

Let Y be any pointed \tilde{G} -space. We obtain a functor $Y \wedge : \mathfrak{C}(\tilde{G}) \rightarrow \mathfrak{C}(\tilde{G})$ which is defined on objects by $Y \wedge (X, m, n) = (Y \wedge X, m, n)$ and on morphisms in the evident way. In particular, if V is any finite-dimensional representation of \tilde{G} , we define the reduced suspension

$$\Sigma^V Z = V^+ \wedge Z.$$

We define desuspension by a representation V of type (1) as

$$\Sigma^{-V} (X, m, n) = ((V)^+ \wedge X, m + 2[V], n),$$

where $[V] = (v_1, \dots, v_p)$ and v_i is the multiplicity of V_i in V . Then $\Sigma^{-V} \Sigma^V Z \cong Z$, where the isomorphism is canonical up to homotopy. For any representation W of type (2) we define

$$\Sigma^{-W} (X, m, n) = (X, m, n + [W]),$$

where $[W] = (w_1, \dots, w_q)$ and w_i is the multiplicity of W_i in W . We have that $\Sigma^{-W} \Sigma^W Z \cong Z$ by an isomorphism which is canonical up to homotopy. In fact, such an isomorphism is induced by a choice of isomorphism $W \cong W([W])$. But for any pair of isomorphic complex representations, the space of isomorphism is connected (by Schur’s lemma it is a torsor for a product of complex general linear groups). Therefore the isomorphism $W \cong W([W])$ is unique up to homotopy.

For $Z = (X, m, n) \in \mathfrak{C}(\tilde{G})$, we define the reduced equivariant cohomology of Z to be

$$\tilde{H}_{\tilde{G}}^j(Z) = \tilde{H}_{\tilde{G}}^{j+|m|+2|n|}(X),$$

where $|m|$ and $|n|$ are defined as

$$|m| = \sum_{i=1}^p m_i \dim_{\mathbb{R}}(V_i) \quad \text{for } m = (m_1, \dots, m_p), \quad |n| = \sum_{i=1}^q n_i \dim_{\mathbb{C}}(W_i) \quad \text{for } n = (n_1, \dots, n_q).$$

The cohomology is well defined as a consequence of the Thom isomorphism.

3.4 G -equivariant Seiberg–Witten–Floer cohomology

Let Y be a rational homology 3–sphere and G a finite group acting on Y preserving the isomorphism class of a spin^c–structure \mathfrak{s} . Let $G_{\mathfrak{s}}$ be the S^1 –central extension of G obtained by lifting G to the spinor bundle corresponding to \mathfrak{s} . We repeat the construction of the Conley index $I_{\lambda}^{\mu}(g)$ from Section 2.6, except that now we carry out the construction $G_{\mathfrak{s}}$ –equivariantly. Restricting to the subgroup $S^1 \subseteq G_{\mathfrak{s}}$, $I_{\lambda}^{\mu}(g)$ agrees with the S^1 –equivariant Conley index as previously constructed.

We need to understand how $I_{\lambda}^{\mu}(g)$ depends on μ , λ , the choice of G –invariant metric g , and the constant R . As in the S^1 case, first consider variations of μ and λ . Carrying out a similar argument but $G_{\mathfrak{s}}$ –equivariantly, we see that $I_{\lambda}^{\mu}(g)$ simply changes by suspension. Analogous to the nonequivariant case we define

$$SWF(Y, \mathfrak{s}, g) = \Sigma^{-V_{\lambda}^0(g)} I_{\lambda}^{\mu}(g) \in \mathfrak{C}(G_{\mathfrak{s}}),$$

where $V_{\lambda}^0(g)$ is defined as before, but now carries a $G_{\mathfrak{s}}$ –action. Note that $V_{\lambda}^0(g)$ is the sum of a representation of type (1) and a representation of type (2), so the desuspension $\Sigma^{-V_{\lambda}^0(g)}$ is defined. Then up to canonical isomorphisms $SWF(Y, \mathfrak{s}, g)$ depends only on the triple (Y, \mathfrak{s}, g) .

We consider the dependence of $SWF(Y, \mathfrak{s}, g)$ on the metric g . The argument is much the same as before except done $G_{\mathfrak{s}}$ –equivariantly. Let g_0 and g_1 be two G –invariant metrics. The space of such metrics is contractible, so we may choose a path $\{g_s\}$ from g_0 to g_1 . Then as in the nonequivariant case, the signature operator has no spectral flow and we have

$$SWF(Y, \mathfrak{s}, g_1) = \Sigma^{SF_{G_{\mathfrak{s}}}(\{D_s\})} SWF(Y, \mathfrak{s}, g_0),$$

where now $SF_{G_{\mathfrak{s}}}(\{D_s\})$ is the equivariant spectral flow of $\{D_s\}$. Thus $SF_{G_{\mathfrak{s}}}(\{D_s\})$ is to be understood as a virtual representation of $G_{\mathfrak{s}}$ [25, Section 2]. Since the S^1 subgroup of $G_{\mathfrak{s}}$ acts by scalar multiplication

on spinors, it follows that $SF_{G_\mathfrak{s}}(\{D_\mathfrak{s}\})$ is a type (2) virtual representation. From the Thom isomorphism and the fact that the underlying rank of $SF_{G_\mathfrak{s}}(\{D_\mathfrak{s}\})$ is $SF(\{D_\mathfrak{s}\}) = n(Y, \mathfrak{s}, g_1) - n(Y, \mathfrak{s}, g_0)$, we obtain a canonical isomorphism

$$\tilde{H}_{G_\mathfrak{s}}^{j+2n(Y,\mathfrak{s},g_1)}(SWF(Y, \mathfrak{s}, g_1)) \cong \tilde{H}_{G_\mathfrak{s}}^{j+2n(Y,\mathfrak{s},g_0)}(SWF(Y, \mathfrak{s}, g_0)).$$

This motivates the following definition:

Definition 3.1 The G –equivariant Seiberg–Witten–Floer cohomology of (Y, \mathfrak{s}, g) is defined as

$$HSW_G^j(Y, \mathfrak{s}) = \tilde{H}_{G_\mathfrak{s}}^{j+2n(Y,\mathfrak{s},g)}(SWF(Y, \mathfrak{s}, g)).$$

By the argument above, the $HSW_G^*(Y, \mathfrak{s})$ depends only on (Y, \mathfrak{s}) and the G –action.

For a group K we write H_K^* for $H_K^*(\text{pt})$. Since $HSW^*(Y, \mathfrak{s})$ is defined using equivariant cohomology, it is a graded module over the ring $H_{S^1}^* = \mathbb{F}[U]$, where $\deg(U) = 2$. Similarly $HSW_G^*(Y, \mathfrak{s})$ is a graded module over $H_{G_\mathfrak{s}}^*$. Restricting from $G_\mathfrak{s}$ to S^1 , we obtain forgetful maps

$$HSW_G^*(Y, \mathfrak{s}) \rightarrow HSW^*(Y, \mathfrak{s}), \quad H_{G_\mathfrak{s}}^* \rightarrow H_{S^1}^*$$

compatible with the module structures.

Observe that since S^1 is the identity component of $G_\mathfrak{s}$, the action of $G_\mathfrak{s}$ on $HSW^*(Y, \mathfrak{s})$ descends to an action of G . So we may regard $HSW^*(Y, \mathfrak{s})$ as a G –module.

Theorem 3.2 There is a spectral sequence $E_r^{p,q}$ abutting to $HSW_G^*(Y, \mathfrak{s})$ whose second page is given by

$$E_2^{p,q} = H^p(BG; HSW^q(Y, \mathfrak{s})).$$

Proof For a $G_\mathfrak{s}$ –space M , let $M_{G_\mathfrak{s}}$ denote the Borel model for the $G_\mathfrak{s}$ –action and M_{S^1} the Borel model for the S^1 –action obtained by restriction. The composition $M_{G_\mathfrak{s}} \rightarrow BG_\mathfrak{s} \rightarrow BG$ is a fibration with fibre M_{S^1} . Applying the Leray–Serre spectral sequence, we get a spectral sequence which abuts to $\tilde{H}_{G_\mathfrak{s}}^*(M)$ and has $E_2^{p,q} = H^p(BG; \tilde{H}_{S^1}^q(M))$. More generally if M is the formal desuspension of a $G_\mathfrak{s}$ –space, then via an application of the Thom isomorphism a similar spectral sequence exists. Applying this to $HSW_G^*(Y, \mathfrak{s})$ gives the theorem. \square

Definition 3.3 Let Y be a rational homology 3–sphere and \mathfrak{s} a spin^c –structure. We say that Y is an L –space (with respect to \mathfrak{s} and \mathbb{F}) if the action of U on $HSW^*(Y, \mathfrak{s})$ is injective. Equivalently $HSW^*(Y, \mathfrak{s})$ is a free $\mathbb{F}[U]$ –module of rank 1.

Remark 3.4 The usual definition of an L –space is that $HF_{\text{red}}^+(Y, \mathfrak{s}) = 0$ for all spin^c –structures and where the coefficient group is \mathbb{Z} . From the universal coefficient theorem it follows that an L –space in this sense is an L –space with respect to any spin^c –structure \mathfrak{s} and any coefficient group \mathbb{F} .

Suppose that the extension $G_\mathfrak{s}$ is split. A choice of splitting induces an isomorphism $G_\mathfrak{s} \cong S^1 \times G$ and an isomorphism $H_{G_\mathfrak{s}}^* \cong H_G^*[U]$. We stress that these isomorphisms depend on the choice of splitting.

Theorem 3.5 Suppose that $G_{\mathfrak{s}}$ is a split extension. If Y is an L -space (with respect to \mathfrak{s} and \mathbb{F}), then the spectral sequence given in Theorem 3.2 degenerates at E_2 . Moreover,

$$HSW_G^*(Y, \mathfrak{s}) \cong HSW^*(Y, \mathfrak{s}) \otimes_{\mathbb{F}} H_G^* \cong H_G^*[U]\theta,$$

where θ has degree $d(Y, \mathfrak{s})$.

Proof If Y is an L -space (with respect to \mathfrak{s} and \mathbb{F}) then

$$HSW^*(Y, \mathfrak{s}) \cong \mathbb{F}[U]\theta,$$

where θ has degree $d(Y, \mathfrak{s})$. We claim that G acts trivially on $HSW^*(Y, \mathfrak{s})$. This can be seen as follows. First, since $HSW^*(Y, \mathfrak{s})$ is up to a degree shift the S^1 -equivariant cohomology of the Conley index $I = I_{\lambda}^{\mu}$, it suffices to prove the result for I . Let $\iota: I^{S^1} \rightarrow I$ be the inclusion of the S^1 fixed-point set. Since Y is an L -space, U acts injectively on $HSW^*(Y, \mathfrak{s})$. Together with the localisation theorem in equivariant cohomology, this implies that ι^* is injective. Hence it suffices to show that G acts trivially on $\tilde{H}_{S^1}^*(I^{S^1})$. But I^{S^1} has the homotopy type of a sphere, so if v is a generator of $\tilde{H}_{S^1}^*(I^{S^1})$ and $g \in G$, then $g^*(v) = \pm v$ according to whether or not g acts orientation-preservingly. Our assumptions on G and \mathbb{F} (see Section 3.1) ensures that $g^*(v) = v$ for all $g \in G$. This proves the claim.

Letting $E_r^{p,q}$ denote the spectral sequence for $HSW_G^*(Y, \mathfrak{s})$, it follows easily that

$$E_2^{p,q} \cong H^*(BG; \mathbb{F}[U]\theta) \cong H_G^*[U]\theta \cong HSW^*(Y, \mathfrak{s}) \otimes_{\mathbb{F}} H_G^*.$$

It remains to show that the differentials d_2, d_3, \dots are all zero. In fact since θ has the lowest q -degree of any term in $E_2^{p,q}$, it follows that $d_j(\theta) = 0$ for all $j \geq 2$. Then since the differentials commute with the $H_G^* \cong H_G^*[U]$ -module structure, it follows that d_2, d_3, \dots all vanish. \square

3.5 Spaces of type G -SWF

We introduce a G -equivariant analogue of spaces of type SWF. We then define a G -equivariant analogue of the d -invariant.

Let \tilde{G} be an extension of G by S^1 . If \tilde{G} acts on a space X , then we get an induced action of $G = \tilde{G}/S^1$ on the fixed-point set X^{S^1} . We write $\bar{G} = S^1 \times G$ for the trivial extension of G .

Definition 3.6 Let $s \geq 0$ be an integer. We say that a finite pointed \tilde{G} -CW complex X is of type G -SWF at level s if

- the S^1 -fixed-point set X^{S^1} is G -homotopy equivalent to a sphere $(V)^+$, where V is a real representation of G of dimension s ;
- the action of S^1 is free on the complement $X - X^{S^1}$.

More generally, let V be a finite-dimensional representation which is the direct sum of representations of type (1) and (2). An equivariant spectrum $Z = \Sigma^{-V} X \in \mathfrak{C}(\tilde{G})$ is said to be of type G -SWF at level s if X is G -SWF at level $s + \dim(V^{S^1})$.

Assume that \tilde{G} is split and choose a splitting $\tilde{G} \cong \bar{G}$. Let X be a space of type G –SWF at level s . Let $\iota: X^{S^1} \rightarrow X$ denote the inclusion of the fixed-point set. Recall that $H_{S^1}^* \cong \mathbb{F}[U]$, where $\deg(U) = 2$. Similarly $H_{\bar{G}}^* \cong H_G^*[U]$. The localisation theorem in equivariant cohomology implies that

$$\iota^*: U^{-1} \tilde{H}_{\bar{G}}^*(X) \rightarrow U^{-1} \tilde{H}_{\bar{G}}^*(X^{S^1})$$

is an isomorphism. Note $X^{S^1} \cong (V)^+$, where V is s –dimensional, so

$$\tilde{H}_{\bar{G}}^*(X^{S^1}) \cong H_G^*[U]\tau,$$

where $\deg(\tau) = s$. Therefore it also follows that

$$U^{-1} \tilde{H}_{\bar{G}}^*(X^{S^1}) \cong H_G^*[U, U^{-1}]\tau.$$

Then for each $c \in H_G^*$, it follows that there exists an $x \in \tilde{H}_{\bar{G}}^*(X)$ for which $\iota^*(x) = cU^k\tau$, for some $k \geq 0$. Set $\Lambda_G(X) = \tilde{H}_{\bar{G}}^*(X^{S^1})$. Then $\Lambda_G(X)$ is a free $H_G^*[U]$ –module of rank 1 and $\iota: X^{S^1} \rightarrow X$ induces a map

$$\iota^*: \tilde{H}_{\bar{G}}^*(X) \rightarrow \Lambda_G(X)$$

of $H_G^*[U]$ –modules. Introduce a filtration

$$\Lambda_G(X) = F_0 \supseteq F_1 \supseteq F_2 \supseteq \dots$$

on $\Lambda_G(X)$ by setting

$$F_j = H_G^{*\geq j} \Lambda_G(X),$$

where $H_G^{*\geq j} = \bigoplus_{k \geq j} H_G^k$. This is the filtration induced by the fibration

$$X^{S^1} \times_{\bar{G}} B\bar{G} \rightarrow BG.$$

Let τ denote the generator of $\Lambda_G(X)$. Then for $j \geq 0$ we have obvious identifications

$$F_j/F_{j+1} \cong H_G^j[U]\tau.$$

Now let c be a nonzero element in H_G^* of degree $|c| = \deg(c)$. By the discussion above we know that $cU^k\tau$ is in the image of ι^* for some $k \geq 0$. Hence we may define:

Definition 3.7 Let c be a nonzero element in H_G^* of degree $|c| = \deg(c)$. We define $d_{G,c}(X) \in \mathbb{Z}$ by

$$d_{G,c}(X) = \min\{2k + s \mid \iota^*(x) \in F_{|c|} \text{ and } \iota^*(x) = cU^k\tau \text{ mod } F_{|c|+1} \text{ for some } x \in \tilde{H}_{\bar{G}}^{s+2k+|c|}(X)\}.$$

For convenience we set $d_{G,0}(X) = -\infty$. Then if c is an element of H_G^* , we write $c = c_0 + c_1 + \dots + c_r$, where $c_i \in H_G^i$ and set

$$d_{G,c}(X) = \max\{d_{G,c_0}(X), \dots, d_{G,c_r}(X)\}.$$

Note that $d_{G,ac}(X) = d_{G,c}(X)$ for any $a \in \mathbb{F}^*$.

In concrete terms, the condition that $\iota^*(x) \in F_{|c|}$ and $\iota^*(x) = cU^k \tau \pmod{F_{|c|+1}}$ means that $\iota^*(x)$ is of the form

$$\iota^*(x) = cU^k \tau + c_1U^{k-1} \tau + \dots + c_rU^{k-r} \tau$$

for some $r \geq 0$ and some $c_1, \dots, c_r \in H_G^{*\geq(|c|+1)}$.

Remark 3.8 Let X be a space of type G -SWF. The definition of $d_{G,c}(X)$ does not depend on a choice of splitting of $S^1 \rightarrow \tilde{G} \rightarrow G$. Indeed, two splittings differ by a homomorphism $\phi: G \rightarrow S^1$. Let $\alpha = \phi^*(U) \in H_G^2$. The change of splitting acts on $H_G^*[U]$ by sending U to $U + \alpha$. Then since $(U + \alpha)^k = U^k + \dots$, where \dots denotes terms involving lower powers of U , it follows that $d_{G,c}(X)$ does not depend on the choice of splitting of \tilde{G} .

Proposition 3.9 Let X be a space of type G -SWF for the trivial extension. Then for all $c_1, c_2 \in H_G^*$,

$$d_{G,c_1+c_2}(X) \leq \max\{d_{G,c_1}(X), d_{G,c_2}(X)\}, \quad d_{G,c_1c_2}(X) \leq \min\{d_{G,c_1}(X), d_{G,c_2}(X)\}.$$

Proof Let s be the level of X . First consider the case that c_1 and c_2 are homogeneous, that is, $c_1 \in H_G^{|c_1|}$ and $c_2 \in H_G^{|c_2|}$ for some $|c_1|$ and $|c_2|$. Then by Definition 3.7, there exist $x_1 \in \tilde{H}_G^{d_{G,c_1}(X)+|c_1|}(X)$ and $x_2 \in \tilde{H}_G^{d_{G,c_2}(X)+|c_2|}(X)$ such that

$$\iota^*(x_1) = c_1U^{k_1} \tau + \dots, \quad \iota^*(x_2) = c_2U^{k_2} \tau + \dots,$$

where \dots denotes terms that are in the next stage of the filtration and $k_i = \frac{1}{2}(d_{G,c_i}(X) - s)$ for $i = 1, 2$. Note that if c_1 or c_2 are zero then we take x_1 or x_2 to be zero.

If $|c_1| \neq |c_2|$, then by Definition 3.7, we have $d_{G,c_1+c_2}(X) = \max\{d_{G,c_1}(X), d_{G,c_2}(X)\}$. Now suppose that $|c_1| = |c_2|$. Let $k = \max\{k_1, k_2\}$ and set $x = U^{k-k_1}x_1 + U^{k-k_2}x_2 \in \tilde{H}_G^{2k+s+|c_1|}(X)$. Then

$$\iota^*(x) = (c_1 + c_2)U^k \tau + \dots$$

and hence, from the definition of $d_{G,c_1+c_2}(X)$,

$$d_{G,c_1+c_2}(X) \leq 2k + s = \max\{2k_1 + s, 2k_2 + s\} = \max\{d_{G,c_1}(X), d_{G,c_2}(X)\}.$$

Next we observe that $c_2x_1 \in \tilde{H}_G^{d_{G,c_1}(X)+|c_1|+|c_2|}(X)$ and

$$\iota^*(c_2x_1) = (c_1c_2)U^{k_1} \tau,$$

and so it follows that $d_{G,c_1c_2}(X) \leq d_{G,c_1}(X)$. Exchanging the roles of x_1, x_2 and c_1, c_2 , we similarly find that $d_{G,c_1c_2}(X) \leq d_{G,c_2}(X)$; hence

$$d_{G,c_1c_2}(X) \leq \min\{d_{G,c_1}(X), d_{G,c_2}(X)\}.$$

Now suppose that c_1 and c_2 are not necessarily homogeneous. We may write $c_1 = a_0 + a_1 + \dots + a_r$ and $c_2 = b_0 + b_1 + \dots + b_r$, for some $r \geq 0$, where $a_i, b_i \in H_G^i$. By Definition 3.7,

$$d_{G,c_1}(X) = \max_i \{d_{G,a_i}(X)\}, \quad d_{G,c_2}(X) = \max_i \{d_{G,b_i}(X)\}.$$

Then since $c_1 + c_2 = (a_0 + b_0) + (a_1 + b_1) + \cdots + (a_r + b_r)$,

$$\begin{aligned} d_{G,c_1+c_2}(X) &= \max_i \{d_{G,a_i+b_i}(X)\} \\ &\leq \max_i \{\max\{d_{G,a_i}(X), d_{G,b_i}(X)\}\} \\ &= \max\{\max_i \{d_{G,a_i}(X)\}, \max_i \{d_{G,b_i}(X)\}\} \\ &= \max\{d_{G,c_1}(X), d_{G,c_2}(X)\}. \end{aligned}$$

Next, we have $c_1 c_2 = \sum_{i,j} a_i b_j$ and hence

$$d_{G,c_1 c_2}(X) \leq \max_{i,j} \{d_{G,a_i b_j}(X)\} \leq \max_{i,j} \{d_{G,a_i}(X)\} = \max_i \{d_{G,a_i}(X)\} = d_{G,c_1}(X),$$

where we used $d_{G,a_i b_j}(X) \leq d_{G,a_i}(X)$. Similarly we get $d_{G,c_1 c_2}(X) \leq d_{G,c_2}(X)$, and hence

$$d_{G,c_1 c_2}(X) \leq \min\{d_{G,c_1}(X), d_{G,c_2}(X)\}. \quad \square$$

Recall that the ordinary (nonequivariant) d -invariant of X , $d(X)$, is defined by

$$d(X) = \min\{j \mid \iota^*(x) \neq 0 \text{ for some } x \in \tilde{H}_{S^1}^j(X)\}.$$

It is not hard to see that $d(X) = d_{\{e\},1}(X)$, where $\{e\}$ denotes the trivial group and 1 is the generator of $H^0(\text{pt})$.

Proposition 3.10 *Let X be a space of type G -SWF for the trivial extension. Then*

$$d_{G,1}(X) \geq d(X).$$

Proof By the definition of $d_{G,1}(X)$, there exists $x \in \tilde{H}_{\bar{G}}^{d_{G,1}(X)}(X)$ such that $\iota^*(x) = U^k \tau + \cdots$, where $k = \frac{1}{2}(d_{G,1}(X) - s)$ and s is the level of X . Let $y \in \tilde{H}_{S^1}^{d_{G,1}(X)}(X)$ be the image of x under the map induced by $S^1 \rightarrow \bar{G}$. Then it follows that $\iota^*(y) = U^k \tau \in \tilde{H}_{S^1}^{d_{G,1}(X)}(X^{S^1})$. In particular, $\iota^*(y) \neq 0$, and hence $d_{G,1}(X) \geq d(X)$ by the definition of $d(X)$. \square

Let S^1 act trivially on \mathbb{R} and act by scalar multiplication on \mathbb{C} . Let V be a real representation of G . Then $V_{\mathbb{R}} = \mathbb{R} \otimes_{\mathbb{R}} V$ and $V_{\mathbb{C}} = \mathbb{C} \otimes_{\mathbb{R}} V$ may be regarded as representations of $\bar{G} = S^1 \times G$, where S^1 acts on the first factor and G on the second.

Proposition 3.11 *Let X be a space of type G -SWF for the trivial extension and let V be a finite-dimensional representation of G of type (1) or (2), as in Section 3.3. Then for any $c \in H_G^*$,*

$$d_{G,c}(V^+ \wedge X) = d_{G,c}(X) + \dim_{\mathbb{R}}(V).$$

Proof This result follows easily from the Thom isomorphism, together with the fact that in the type (2) case, the \bar{G} -equivariant Euler class of V has the form

$$e_{\bar{G}}(V) = U^{\dim(V)} + c_{G,1}(V)U^{\dim(V)-1} + \cdots + c_{G,\dim(V)}(V),$$

where $c_{G,j}(V) \in H_G^{2j}$ denotes the j^{th} G -equivariant Chern class of V . \square

If $Z = \Sigma^{-V} X \in \mathfrak{C}(\bar{G})$ is an equivariant spectrum of type G -SWF, we define the $d_{G,c}$ -invariant $d_{G,c}(Z)$ of Z to be

$$d_{G,c}(Z) = d_{G,c}(X) - \dim_{\mathbb{R}}(V).$$

This is well defined by Proposition 3.11. We also define a corresponding δ -invariant by setting $\delta_{G,c}(Z) = \frac{1}{2}d_{G,c}(Z)$.

Definition 3.12 Let X and Y be spaces of type G -SWF for the trivial extension of G , where X has level s and Y has level t . Let $f : X \rightarrow Y$ be an $S^1 \times G$ -equivariant map. Consider the restriction

$$f^{S^1} : X^{S^1} \rightarrow Y^{S^1}$$

of f to the fixed-point set. Note that $\tilde{H}_G^*(X^{S^1})$ is a free H_G^* -module starting in degree s . Let $\tau_{X^{S^1}}$ denote a generator. Then $\tau_{X^{S^1}}$ is unique up to an element of \mathbb{F}^* . Similarly $\tilde{H}_G^*(Y^{S^1})$ is a free H_G^* -module starting in degree t and we let $\tau_{Y^{S^1}}$ denote a generator. Then there exists a uniquely determined $\mu \in H_G^{t-s}$ such that

$$(f^{S^1})^*(\tau_{Y^{S^1}}) = \mu \tau_{X^{S^1}}.$$

We call $\mu = \text{deg}(f^{S^1})$ the *degree* of f^{S^1} . If we choose different generators for $\tilde{H}_G^*(X^{S^1})$ or $\tilde{H}_G^*(Y^{S^1})$, then $\text{deg}(f^{S^1})$ changes by an element of \mathbb{F}^* ; hence $\text{deg}(f^{S^1})$ is well defined up to multiplication by elements of \mathbb{F}^* . If $t < s$, then $\text{deg}(f^{S^1}) = 0$.

Note that suspension does not change the degree of f^{S^1} . Hence we can more generally speak of the degree of f^{S^1} when f is a stable map between spectra of type G -SWF.

Proposition 3.13 Let $f : X \rightarrow Y$ be a \bar{G} -equivariant map of spaces of type G -SWF for the trivial extension, where X has level s and Y has level t . Let $\mu = \text{deg}(f^{S^1}) \in H_G^{t-s}$ be the degree of f^{S^1} . Then for any nonzero $c \in H_G^*$,

$$d_{G,c\mu}(X) - s \leq d_{G,c}(Y) - t.$$

Proof We prove the result when $c \in H_G^{|c|}$ is homogeneous. The general case follows easily from this. The inclusion of the fixed-point sets gives a commutative diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \iota \uparrow & & \uparrow \iota \\ X^{S^1} & \xrightarrow{f^{S^1}} & Y^{S^1} \end{array}$$

Consider the induced commutative diagram in equivariant cohomology

$$\begin{array}{ccc} \tilde{H}_G^*(Y) & \xrightarrow{f^*} & \tilde{H}_G^*(X) \\ \downarrow \iota^* & & \downarrow \iota^* \\ \tilde{H}_G^*(Y^{S^1}) & \xrightarrow{(f^{S^1})^*} & \tilde{H}_G^*(X^{S^1}) \end{array}$$

From the definition of $d_{G,c}(Y)$, there exists some $x \in \tilde{H}_{\bar{G}}^{d_{G,c}(Y)+|c|}(Y)$ such that

$$\iota^*(x) = cU^k \tau_{Y^{S^1}} + \dots,$$

where $k = \frac{1}{2}(d_{G,c}(Y) - t)$. Then by commutativity of the diagram,

$$\iota^*(f^*(x)) = (f^{S^1})^*(\iota^*(x)) = (f^{S^1})^*(cU^k \tau_{Y^{S^1}} + \dots) = c\mu U^k \tau_{X^{S^1}} + \dots.$$

It follows that

$$d_{G,c\mu}(X) \leq d_{G,c}(Y) + |c| - |c\mu| = d_{G,c}(Y) - |\mu| = d_{G,c}(Y) - t + s.$$

Hence

$$d_{G,c\mu}(X) - s \leq d_{G,c}(Y) - t. \quad \square$$

3.6 Alternative characterisation of $d_{G,c}$

In this section we will give an alternative characterisation of $d_{G,c}$ which does not directly refer to ι^* and is sometimes more convenient for computations.

Let X be a space of type G –SWF for the trivial extension \bar{G} . Set $\Lambda_G^* = \tilde{H}_{\bar{G}}^*(X^{S^1})$. The inclusion of the fixed points $\iota: X^{S^1} \rightarrow X$ induces a map $\iota^*: \tilde{H}_{\bar{G}}^*(X) \rightarrow \Lambda_G^*$. Recall that Λ_G^* is a free $H_G^*[U]$ module of rank 1. Let τ denote a generator of Λ_G^* , so $\Lambda_G^* \cong H_G^*[U]\tau$. Recall that we have a filtration F_j on Λ_G^* given by $F_j = H_G^{*\geq j}\Lambda_G^*$. Similarly, there is a filtration on $\tilde{H}_{\bar{G}}^*(X)$ which comes from the spectral sequence for equivariant cohomology. We will denote this filtration by \mathcal{F}_j . Then $\iota^*(\mathcal{F}_j) \subseteq F_j$ because the inclusion ι induces a map between spectral sequences.

Let $c \in H_G^*$ be a nonzero element of degree $|c|$. Recall that the invariant $d_{G,c}(X)$ is defined by

$$d_{G,c}(X) = \min\{i \mid \iota^*(x) = cU^k \tau \text{ mod } F_{|c|+1} \text{ for some } x \in \tilde{H}_{\bar{G}}^i(X) \text{ and } k \geq 0\} - |c|.$$

The localisation theorem in equivariant cohomology implies that upon localising with respect to U , ι^* becomes an isomorphism

$$\iota^*: U^{-1}\tilde{H}_{\bar{G}}^*(X) \rightarrow U^{-1}\Lambda_G \cong H_G^*[U, U^{-1}]\tau.$$

In particular, there exists an element $\theta \in \tilde{H}_{\bar{G}}^{2k+\text{deg}(\tau)}(X)$ such that $\iota^*(\theta) = U^l \tau$ for some $l \geq 0$. Fix a choice of such a θ . The localisation isomorphism implies that $\iota^*(x) = 0$ if and only if $U^k x = 0$ for some $k \geq 0$.

Proposition 3.14 *Let $c \in H_G^*$ be a nonzero element of degree $|c|$. Then*

$$d_{G,c}(X) = \min\{i \mid U^n x = cU^k \theta \text{ mod } \mathcal{F}_{|c|+1} \text{ for some } x \in \tilde{H}_{\bar{G}}^i(X) \text{ and } n, k \geq 0\} - |c|.$$

Proof Let

$$a_{G,c}(X) = \min\{i \mid U^n x = cU^k \theta \pmod{\mathcal{F}_{|c|+1}} \text{ for some } x \in \tilde{H}_{\bar{G}}^i(X) \text{ and } n, k \geq 0\} - |c|.$$

Then we need to show that $d_{G,c}(X) = a_{G,c}(X)$. Suppose $x \in \tilde{H}_{\bar{G}}^{a_{G,c}(X)+|c|}(X)$ satisfies $U^n x = cU^k \theta \pmod{\mathcal{F}_{|c|+1}}$ for some $n, k \geq 0$. Then

$$U^n \iota^*(x) = cU^k \iota^*(\theta) = cU^{k+l} \tau \pmod{F_{|c|+1}}.$$

Since U is injective on Λ_G we must have $k+l \geq n$ and we can cancel U^n from both sides to get

$$\iota^*(x) = cU^{k+l-n} \tau \pmod{F_{|c|+1}}.$$

Hence $d_{G,c}(X) \leq \deg(x) - |c| = a_{G,c}(X)$. Conversely, let $x \in \tilde{H}_{\bar{G}}^{d_{G,c}(X)+|c|}(X)$ satisfy $\iota^*(x) = cU^k \tau \pmod{F_{|c|+1}}$ for some $k \geq 0$. Then

$$\iota^*(x) = cU^k \tau + c_1 U^{k-1} \tau + c_2 U^{k-2} \tau + \cdots + c_k \tau,$$

where $c_i \in H_G^{|c|+2i}$. Since $\iota^*(\theta) = U^l \tau$, it follows that

$$\iota^*(U^l x) = \iota^*(cU^k \theta + c_1 U^{k-1} \theta + \cdots + c_k \theta).$$

Next recall that ι^* is an isomorphism after localising with respect to U . Hence if $\iota^*(y_1) = \iota^*(y_2)$, then $U^n y_1 = U^n y_2$ for some $n \geq 0$ and we have

$$U^{n+l} x = cU^{n+k} \theta + c_1 U^{n+k-1} \theta + \cdots + c_k U^n \theta = cU^{n+k} \theta \pmod{\mathcal{F}_{|c|+1}}.$$

From the definition of $a_{G,c}(X)$, it follows that $a_{G,c}(X) \leq \deg(x) - |c| = d_{G,c}(X)$. We have shown $d_{G,c}(X) \leq a_{G,c}(X)$ and $a_{G,c}(X) \leq d_{G,c}(X)$; hence $d_{G,c}(X) = a_{G,c}(X)$. \square

3.7 Equivariant d -invariants for rational homology 3-spheres

We return to the setting that Y is a rational homology 3-sphere, G is a finite group acting on Y preserving the isomorphism class of a spin^c -structure \mathfrak{s} . Choose a G -invariant metric g and let $G_{\mathfrak{s}}$ be the S^1 -central extension of G obtained by lifting G to the spinor bundle corresponding to \mathfrak{s} . Now suppose that $G_{\mathfrak{s}}$ is a trivial extension; hence $G_{\mathfrak{s}} \cong \bar{G}$. From the construction of the Conley index, one finds that $SWF(Y, \mathfrak{s}, g)$ is of type G -SWF at level 0.

Definition 3.15 Let G act on Y and let \mathfrak{s} be a G -invariant spin^c -structure. Suppose that the corresponding S^1 -extension $G_{\mathfrak{s}}$ is trivial and choose an isomorphism of extensions $G_{\mathfrak{s}} \cong \bar{G}$. For any $c \in H_G^*$ we define the invariant $d_{G,c}(Y, \mathfrak{s})$ by

$$d_{G,c}(Y, \mathfrak{s}) = d_{G,c}(SWF(Y, \mathfrak{s}, g)) - 2n(Y, \mathfrak{s}, g).$$

We also set $\delta_{G,c}(Y, \mathfrak{s}) = \frac{1}{2} d_{G,c}(Y, \mathfrak{s})$.

The definition of $d_{G,c}(Y, \mathfrak{s})$ does not depend on the choice of isomorphism $G_{\mathfrak{s}} \cong \overline{G}$ by Remark 3.8. The definition also does not depend on the choice of metric g as a consequence of the relation

$$SWF(Y, \mathfrak{s}, g_1) = \Sigma^{SF_{G_{\mathfrak{s}}(\{D, \mathfrak{s}\})}} SWF(Y, \mathfrak{s}, g_0)$$

and the Thom isomorphism.

We only define the invariants $d_{G,c}(Y, \mathfrak{s})$ in the case that $G_{\mathfrak{s}}$ is a trivial extension. This is because the definition of $d_{G,c}(Y, \mathfrak{s})$ uses localisation by U , but $U \in H_{S^1}^*$ does not necessarily extend to a class in $H_{G_{\mathfrak{s}}}^*$, unless $G_{\mathfrak{s}}$ is a trivial extension.

The inclusion $\iota: (V_{\lambda}^0(\mathbb{R}))^+ \rightarrow I_{\lambda}^{\mu}$ of the S^1 –fixed points of the Conley index desuspends to a map $\iota: \Sigma^{-V_{\lambda}^0(\mathbb{C})} S^0 \rightarrow SWF(Y, \mathfrak{s}, g)$; hence we get a homomorphism

$$\iota^*: HSW_G^*(Y, \mathfrak{s}) \rightarrow \Lambda_G^*(Y, \mathfrak{s}),$$

where we have set $\Lambda_G^*(Y, \mathfrak{s}) = \tilde{H}_{\overline{G}}^{*+2n(Y, \mathfrak{s}, g)}(\Sigma^{-V_{\lambda}^0(\mathbb{C})} S^0)$. This is a free $H_G^*[U]$ –module and we let τ denote a generator. As in Section 3.5 we filter $\Lambda_G^*(Y, \mathfrak{s})$ by setting $F_j = H_G^{*\geq j} \Lambda_G^*(Y, \mathfrak{s})$. The construction of ι^* and $\Lambda_G^*(Y, \mathfrak{s})$ depend on the choice of metric g , but the construction for any two metrics are related by a canonical homomorphism. The d –invariants of (Y, \mathfrak{s}) are given by

$$d_{G,c}(Y, \mathfrak{s}) = \min\{2k + j \mid \iota^*(x) \in F_{|c|} \text{ and } \iota^*(x) = cU^k \tau \text{ mod } F_{|c|+1} \text{ for some } x \in SWF_G^{j+2k+|c|}(Y, \mathfrak{s})\}.$$

Recall that $d(\overline{Y}, \mathfrak{s}) = -d(Y, \mathfrak{s})$. On the other hand, the behaviour of the invariants $d_{G,c}(Y, \mathfrak{s})$ under orientation reversal is not so straightforward. For example, it follows from Proposition 3.10 that

$$(3-1) \quad -d_{G,1}(\overline{Y}, \mathfrak{s}) \leq d(Y, \mathfrak{s}) \leq d_{G,1}(Y, \mathfrak{s}).$$

In particular, $d_{G,1}(\overline{Y}, \mathfrak{s}) = -d_{G,1}(Y, \mathfrak{s})$ can only occur if $d_{G,1}(Y, \mathfrak{s}) = d(Y, \mathfrak{s})$ and $d_{G,1}(\overline{Y}, \mathfrak{s}) = d(\overline{Y}, \mathfrak{s})$. From (3-1), we also get that

$$d_{G,1}(Y, \mathfrak{s}) + d_{G,1}(\overline{Y}, \mathfrak{s}) \geq 0.$$

We will show in Theorem 4.4 that the invariants $d_{G,c}$ satisfy a stronger positivity condition.

Proposition 3.16 *Let G act on Y and let \mathfrak{s} be a G –invariant spin^c –structure. Suppose that the corresponding extension $G_{\mathfrak{s}}$ is trivial. If Y is an L –space (with respect to \mathfrak{s} and \mathbb{F}), then for all nonzero $c \in H_G^*$,*

$$d_{G,c}(Y, \mathfrak{s}) = d(Y, \mathfrak{s}).$$

Proof If Y is an L –space (with respect to \mathfrak{s} and \mathbb{F}) then

$$HSW^*(Y, \mathfrak{s}) \cong \mathbb{F}[U]\theta,$$

where θ has degree $d(Y, \mathfrak{s})$. From Theorem 3.5, there exists a class $\hat{\theta} \in HSW_G^*(Y, \mathfrak{s})$ which maps to θ under the forgetful map $HSW_G^*(Y, \mathfrak{s}) \rightarrow HSW^*(Y, \mathfrak{s})$ and we have that $HSW_G^*(Y, \mathfrak{s})$ is a free $H_G^*[U]$ –module generated by $\hat{\theta}$. We must also have that $\iota^*(\hat{\theta}) = U^k \tau \text{ mod } F_1$ for some $k \geq 0$, where τ is a

generator of $\Lambda_G(Y, \mathfrak{s})$. This holds because F_1 is the kernel of the forgetful map $\Lambda_G(Y, \mathfrak{s}) \rightarrow \Lambda_{\{1\}}(Y, \mathfrak{s})$. So for any nonzero $c \in H_G^j$ we have that $\iota^*(c\hat{\theta}) = c\iota^*(\hat{\theta}) = cU^k\tau \bmod F_{1+|c|}$; hence

$$d_{G,c}(Y, \mathfrak{s}) \leq \deg(c\hat{\theta}) - j = \deg(\hat{\tau}) = d(Y, \mathfrak{s}).$$

That is, $d_{G,c}(Y, \mathfrak{s}) \leq d(Y, \mathfrak{s})$ for all nonzero homogeneous c . On the other hand it is clear that there is no class of lower degree in $HSW_G^*(Y, \mathfrak{s})$ which maps under ι^* to a class of the form $cU^{k'}\tau \bmod F_{1+|c|}$. Hence $d_{G,c}(Y, \mathfrak{s}) = d(Y, \mathfrak{s})$ for all nonzero homogeneous c . Clearly the result extends to all nonzero c . \square

4 Behaviour under cobordisms

We show that equivariant cobordisms of rational homology 3–spheres induce maps on equivariant Seiberg–Witten–Floer cohomology. We follow the construction of Manolescu [49], incorporating the corrections due to Khandhawit [37]. Since our construction is a straightforward extension of that of Manolescu and Khandhawit, differing only in the replacement of S^1 by the larger group $G_{\mathfrak{s}}$, we will be brief.

4.1 Finite-dimensional approximation

Let W be a compact, oriented smooth 4–manifold with boundary $Y = \partial W$ a disjoint union of rational homology spheres $Y = \bigcup_j Y_j$. Assume further that $b_1(W) = 0$ and that W is connected. If \mathfrak{s} is a spin^c –structure on W , then the restriction of \mathfrak{s} to Y determines a spin^c –structure $\mathfrak{s}|_Y$ on Y . Since the boundary of W is a union of rational homology 3–spheres, we have $H^2(W, \partial W; \mathbb{R}) \cong H^2(W; \mathbb{R})$ and by Poincaré–Lefschetz duality we obtain a nondegenerate intersection form on $H^2(W; \mathbb{R})$. Given a metric g on W which is isometric to a product metric in a collar neighbourhood of ∂W , we let $H^+(W)$ denote the space of self-dual L^2 –harmonic 2–forms on the cylindrical end manifold \widehat{W} obtained from W by attaching half-infinite cylinders $[0, \infty) \times Y$ to W . It follows from [8, Proposition 4.9] that the natural map $H^+(W) \rightarrow H^2(W; \mathbb{R})$ is injective and identifies $H^+(W)$ with a maximal positive definite subspace of $H^2(W; \mathbb{R})$.

Suppose now that G acts smoothly and orientation-preservingly on W and that this action sends each connected component of ∂W to itself. Hence by restriction G acts on each Y_i by orientation-preserving diffeomorphisms. Assume further that G preserves the isomorphism class of a spin^c –structure \mathfrak{s} on W . Set $\mathfrak{s}_i = \mathfrak{s}|_{Y_i}$. Then the action of G on Y_i preserves \mathfrak{s}_i . Similar to Section 3.2 we obtain an S^1 –extension $G_{\mathfrak{s}}$ of G . Restricting to Y_i , we obtain an isomorphism of extensions $G_{\mathfrak{s}} \cong G_{\mathfrak{s}_i}$. Hence if $G_{\mathfrak{s}}$ is split, then it follows that each of the extensions $G_{\mathfrak{s}_i}$ is also split. Moreover a splitting of $G_{\mathfrak{s}}$ determines corresponding splittings of each $G_{\mathfrak{s}_i}$.

Choose a G –invariant metric g on W which is isometric to a product $(-\epsilon, 0] \times Y$ in some equivariant collar neighbourhood of Y (see [34, Theorem 3.5] for existence of equivariant collar neighbourhoods). To see that such a metric exists, first choose a G –invariant metric g_Y on Y . Then choose an arbitrary

metric g' on W which equals $(dt)^2 + g_Y$ in some equivariant collar neighbourhood $(-\epsilon, 0] \times Y$. Then let g be obtained from g' by averaging over G . Let S^\pm denote the spinor bundles on W corresponding to \mathfrak{s} . We note here that under these assumptions G preserves the subspace $H^+(W)$ of $H^2(W; \mathbb{R})$ defined by g . Let $\Omega_g^1(W)$ denote the space of 1-forms on W in double Coulomb gauge with respect to Y [37, Definition 1]. This space is easily seen to be preserved by the action of G on 1-forms. The double Coulomb gauge condition ensures that if $a \in \Omega_g^1(W)$ and $\phi \in \Gamma(S^+)$, then $(a, \phi)|_{Y_j}$ lies in the global Coulomb slice corresponding to Y_j . Let us temporarily assume that $Y = \partial W$ is connected. Let \hat{A} be a spin^c-connection on W such that in a collar neighbourhood of Y it equals the pullback of A_0 . Using the same argument as in Section 3.2, we can assume that \hat{A} is $G_{\mathfrak{s}}$ -invariant. Then using \hat{A} as a reference connection, we obtain a map which may be thought of as the Seiberg–Witten equations on W together with boundary conditions,

$$SW^\mu : i\Omega_g^1(W) \oplus \Gamma(S^+) \rightarrow i\Omega_+^2(W) \oplus \Gamma(S^-) \oplus V_{-\infty}^\mu, \\ (a, \phi) \mapsto (F_{\hat{A}+a}^+ - \sigma(\phi, \phi), D_{\hat{A}+a}(\phi), p^\mu(a, \phi)|_Y),$$

where p^μ is the orthogonal projection from V to $V_{-\infty}^\mu$. Taking a finite-dimensional approximation as described in [37; 49], one obtains a map

$$\Psi_{\mu, \lambda, U, U'} : (U')^+ \rightarrow (U^+) \wedge I_\lambda^\mu,$$

where $U' \subset i\Omega_g^1(W) \oplus \Gamma(S^+)$ and $U \subset i\Omega_+^2(W) \oplus \Gamma(S^-)$ are finite-dimensional G -invariant subspaces which satisfy

$$(4-1) \quad U \oplus V_\lambda^0 \oplus \text{Ker}(L^0) \cong U' \oplus \text{Coker}(L^0)$$

and L^0 is a Fredholm linear operator defined in [49, Section 9]. Since $SWF(Y, \mathfrak{s}, g) = \Sigma^{-V_\lambda^0} I_\lambda^\mu$, we can rewrite the map $\Psi_{\mu, \lambda, U, U'}$ as

$$\Psi_{\mu, \lambda, U, U'} : (U')^+ \rightarrow (U)^+ \wedge (V_\lambda^0)^+ \wedge SWF(Y, \mathfrak{s}, g).$$

Taking the smash product with $\text{Ker}(L^0)$ and using (4-1), we see that $\Psi_{\mu, \lambda, U, U'}$ is stably equivalent to a map

$$f : \text{Ker}(L^0)^+ \rightarrow \text{Coker}(L^0)^+ \wedge SWF(Y, \mathfrak{s}, g).$$

The real part of L^0 has zero kernel and cokernel isomorphic to $H^+(W)$. The complex part of L^0 can be identified with the Dirac operator $D_{\hat{A}}$ with Atiyah–Patodi–Singer (APS) boundary conditions. Thus

$$\text{Ker}(L^0) \cong \text{Ker}_{\text{APS}}(D_{\hat{A}}^\pm), \quad \text{Coker}(L^0) \cong H^+(W) \oplus \text{Coker}_{\text{APS}}(D_{\hat{A}}^\pm),$$

where $\text{Ker}_{\text{APS}}(D_{\hat{A}}^\pm)$ and $\text{Coker}_{\text{APS}}(D_{\hat{A}}^\pm)$ denote the kernel and cokernel of $D_{\hat{A}}^\pm$ with APS boundary conditions. Hence we obtain a $G_{\mathfrak{s}}$ -equivariant map

$$f : \text{Ker}_{\text{APS}}(D_{\hat{A}}^\pm)^+ \rightarrow (H^+(W))^+ \wedge \text{Coker}_{\text{APS}}(D_{\hat{A}}^\pm)^+ \wedge SWF(Y, \mathfrak{s}, g).$$

Note that f is only a map in the stable sense; that is, f is a morphism in the category $\mathcal{C}(G_{\mathfrak{s}})$.

Recall that the S^1 -fixed-point set of I_λ^μ is $V_\lambda^0(\mathbb{R})^+$. The inclusion $(V_\lambda^0(\mathbb{R})) \rightarrow I_\lambda^\mu$ of the S^1 -fixed points desuspends to a map $\iota: S^0 \rightarrow SWF(Y, \mathfrak{s}, g)$. By restricting to S^1 -fixed points we obtain a commutative diagram

$$\begin{array}{ccc} \text{Ker}_{\text{APS}}(D_A^+)^+ & \xrightarrow{f} & (H^+(W))^+ \wedge \text{Coker}_{\text{APS}}(D_A^+)^+ \wedge SWF(Y, \mathfrak{s}, g) \\ \uparrow & & \uparrow \\ S^0 & \xrightarrow{f^{S^1}} & (H^+(W))^+ \end{array}$$

Using that the Seiberg–Witten equations reduce to linear equations on the S^1 -fixed-point set, one finds that $f^{S^1}: S^0 \rightarrow (H^+(W))^+$ is the obvious map given by the one-point compactification of the inclusion $\{0\} \rightarrow H^+(W)$. Thus according to Definition 3.12, f^{S^1} has degree equal to $e(H^+(W))$, the image of the equivariant Euler class of $H^+(W)$ in $H_G^*(\text{pt}; \mathbb{F})$. For instance, if $\mathbb{F} = \mathbb{Z}/2\mathbb{Z}$ then $e(H^+(W))$ is the $b_+(W)$ th equivariant Stiefel–Whitney class. We will refer to $e(H^+(W))$ as the \mathbb{F} -Euler class of $H^+(W)$.

So far we have restricted to the case that the boundary ∂W is connected. More generally, if $\partial W = \bigcup_j Y_j$ is a union of rational homology 3–spheres then much the same construction applies. The Conley index I_λ^μ is now given by the smash product of the Conley indices of each component; hence f is now a map of the form

$$f: \text{Ker}_{\text{APS}}(D_A^+)^+ \rightarrow (H^+(W))^+ \wedge \text{Coker}_{\text{APS}}(D_A^+)^+ \wedge \bigwedge_j SWF(Y_j, \mathfrak{s}_j, g_j).$$

We still have that the degree of f^{S^1} is $e(H^+(W))$.

4.2 Equivariant Frøyshov inequality

In this section we prove an equivariant generalisation of Frøyshov’s inequality [28].

Theorem 4.1 *Let W be a smooth, compact, oriented 4–manifold with boundary and with $b_1(W) = 0$. Suppose that G acts smoothly on W preserving the orientation and a spin^c -structure \mathfrak{s} . Suppose that the extension $G_\mathfrak{s}$ is trivial. Suppose each component of ∂W is a rational homology 3–sphere and that G sends each component of ∂W to itself. Let $e \in H_G^{b_+(W)}$ be the \mathbb{F} -Euler class of any G -invariant maximal positive definite subspace of $H^2(W; \mathbb{R})$. Let $c \in H_G^*$ and suppose that $ce \neq 0$.*

(1) *If $\partial W = Y$ is connected, then*

$$\delta(W, \mathfrak{s}) \leq \delta_{G,c}(Y, \mathfrak{s}|_Y) \quad \text{and} \quad \delta_{G,ce}(\bar{Y}, \mathfrak{s}|_Y) \leq \delta(\bar{W}, \mathfrak{s}),$$

where we have defined

$$\delta(W, \mathfrak{s}) = \frac{1}{8}(c_1(\mathfrak{s})^2 - \sigma(W)).$$

(2) *If $\partial W = \bar{Y}_1 \cup Y_2$ has two connected components, then*

$$\delta_{G,ce}(Y_1, \mathfrak{s}|_{Y_1}) + \delta(W, \mathfrak{s}) \leq \delta_{G,c}(Y_2, \mathfrak{s}|_{Y_2}).$$

Proof We will give the proof in the case that W is connected. The general case follows easily from this by applying the theorem to each component of W . To simplify notation we will write \mathfrak{s} instead of $\mathfrak{s}|_Y$ and write g instead of $g|_Y$. In case (1), $\partial W = Y$ is connected.

As in Section 4.1, choosing suitable metrics and reference connections we obtain a stable map

$$f : \text{Ker}_{\text{APS}}(D_A^+) \rightarrow (H^+(W))^+ \wedge \text{Coker}_{\text{APS}}(D_A^+) \wedge SWF(Y, \mathfrak{s}, g)$$

such that the degree of f^{S^1} is e . Applying Proposition 3.13 to f , we obtain

$$2 \dim_{\mathbb{C}}(\text{Ker}_{\text{APS}}(D_A^+)) + b_+(W) \leq b_+(W) + 2 \dim_{\mathbb{C}}(\text{Coker}_{\text{APS}}(D_A^+)) + d_c(Y, \mathfrak{s}) + 2n(Y, \mathfrak{s}, g)$$

which simplifies to

$$\text{ind}_{\text{APS}}(D_A^+) \leq \frac{1}{2}d_c(Y, \mathfrak{s}) + n(Y, \mathfrak{s}, g) = \delta_c(Y, \mathfrak{s}) + n(Y, \mathfrak{s}, g).$$

Combined with (2-3) we get $\delta(W, \mathfrak{s}) \leq \delta_c(Y, \mathfrak{s})$.

Next recall from Section 2.7 the duality map

$$\varepsilon : SWF(Y, \mathfrak{s}, g) \wedge SWF(\bar{Y}, \mathfrak{s}, g) \rightarrow S^{-k(D)\mathbb{C}}$$

where $k(D) = \dim_{\mathbb{C}}(\text{Ker}(D))$. By the definition of equivariant duality,

$$\varepsilon^{S^1} : SWF(Y, \mathfrak{s}, g)^{S^1} \wedge SWF(\bar{Y}, \mathfrak{s}, g)^{S^1} \rightarrow S^0$$

is a nonequivariant duality. It follows that ε^{S^1} has degree 1. Taking the map f , suspending by $SWF(\bar{Y}, \mathfrak{s}, g)$ and composing with ε , we obtain a stable map

$$h : \text{Ker}_{\text{APS}}(D_A^+) \wedge SWF(\bar{Y}, \mathfrak{s}, g) \rightarrow (H^+(W))^+ \wedge \text{Coker}_{\text{APS}}(D_A^+) \wedge S^{-k(D)\mathbb{C}}$$

such that the degree of h^{S^1} is e . Applying Proposition 3.13 to h , we obtain

$$\begin{aligned} 2 \dim_{\mathbb{C}}(\text{Ker}_{\text{APS}}(D_A^+)) + d_{G,ce}(\bar{Y}, \mathfrak{s}, g) + 2n(\bar{Y}, \mathfrak{s}, g) + b_+(W) \\ \leq b_+(W) + 2 \dim_{\mathbb{C}}(\text{Coker}_{\text{APS}}(D_A^+)) - 2k(D) \end{aligned}$$

which simplifies to

$$\text{ind}_{\text{APS}}(D_A^+) + \delta_{G,ce}(\bar{Y}, \mathfrak{s}, g) + n(\bar{Y}, \mathfrak{s}, g) \leq -k(D).$$

Using (2-3) and (2-4), we obtain $\delta(W, \mathfrak{s}) \leq -\delta_{G,ce}(\bar{Y}, \mathfrak{s})$, or equivalently $\delta_{G,ce}(\bar{Y}, \mathfrak{s}|_Y) \leq \delta(\bar{W}, \mathfrak{s})$.

The proof of case (2) is similar. We start with the map

$$f : \text{Ker}_{\text{APS}}(D_A^+) \rightarrow (H^+(W))^+ \wedge \text{Coker}_{\text{APS}}(D_A^+) \wedge SWF(\bar{Y}_1, \mathfrak{s}, g) \wedge SWF(Y_2, \mathfrak{s}, g).$$

Suspending by $SWF(Y_1, \mathfrak{s}, g)$ and applying the duality map corresponding to Y_1 we obtain a map

$$h : \text{Ker}_{\text{APS}}(D_A^+) \wedge SWF(Y_1, \mathfrak{s}, g) \rightarrow (H^+(W))^+ \wedge \text{Coker}_{\text{APS}}(D_A^+) \wedge SWF(Y_2, \mathfrak{s}, g) \wedge S^{-k(D_1)\mathbb{C}},$$

where $k(D_1)$ is the dimension of the kernel of the Dirac operator on Y_1 . Applying Proposition 3.13 to this map and simplifying, we obtain the inequality $\delta_{G,ce}(Y_1, \mathfrak{s}|_{Y_1}) + \delta(W, \mathfrak{s}) \leq \delta_{G,c}(Y_2, \mathfrak{s}|_{Y_2})$. \square

Definition 4.2 Let (Y_1, \mathfrak{s}_1) and (Y_2, \mathfrak{s}_2) be rational homology 3–spheres equipped with spin^c –structures. Suppose that G acts orientation-preservingly on Y_1 and Y_2 and preserves the spin^c –structures \mathfrak{s}_1 and \mathfrak{s}_2 . A G –equivariant rational homology cobordism from (Y_1, \mathfrak{s}_1) to (Y_2, \mathfrak{s}_2) is a rational homology cobordism W from Y_1 to Y_2 such that the G –action and spin^c –structure $\mathfrak{s}_1 \cup \mathfrak{s}_2$ on ∂W extend over W . We say that (Y_1, \mathfrak{s}_1) and (Y_2, \mathfrak{s}_2) are G –equivariantly rational homology cobordant if there exists a G –equivariant rational homology cobordism from (Y_1, \mathfrak{s}_1) to (Y_2, \mathfrak{s}_2) .

Similarly we define the notion of a G –equivariant integral homology cobordism and say that two integral homology 3–spheres Y_1 and Y_2 on which G acts are G –equivariantly integral homology cobordant if there is a G –equivariant integral homology cobordism from Y_1 to Y_2 . Note that since Y_1 and Y_2 are integral homology 3–spheres, they have unique spin^c –structures which are automatically G –invariant and any G –equivariant integral homology cobordism from Y_1 to Y_2 has a unique spin^c –structure which restricts on the boundary to the unique spin^c –structures on Y_1 and Y_2 .

Corollary 4.3 *The G –equivariant δ –invariants $\delta_{G,c}(Y, \mathfrak{s})$ are invariant under G –equivariant rational homology cobordism; that is, if (W, \mathfrak{s}) is a G –equivariant rational homology cobordism from (Y_1, \mathfrak{s}_1) to (Y_2, \mathfrak{s}_2) and if the extensions $G_{\mathfrak{s}_1}$ and $G_{\mathfrak{s}_2}$ are trivial, then $\delta_{G,c}(Y_1, \mathfrak{s}_1) = \delta_{G,c}(Y_2, \mathfrak{s}_2)$ for all $c \in H_G^*$.*

Proof Since W is a rational homology cobordism, we have $H^2(W; \mathbb{R}) = 0$. So $\delta(W, \mathfrak{s}) = 0$ and $e = e(H^+(W)) = 1$. Therefore Theorem 4.1 gives

$$\delta_{G,c}(Y_1, \mathfrak{s}_1) \leq \delta_{G,c}(Y_2, \mathfrak{s}_2).$$

Similarly, viewing \overline{W} as a G –equivariant rational homology cobordism from Y_2 to Y_1 , we get

$$\delta_{G,c}(Y_2, \mathfrak{s}_2) \leq \delta_{G,c}(Y_1, \mathfrak{s}_1).$$

Hence $\delta_{G,c}(Y_1, \mathfrak{s}_1) = \delta_{G,c}(Y_2, \mathfrak{s}_2)$. □

Theorem 4.4 *Let Y be a rational homology 3–sphere, G a finite group acting on Y preserving orientation and the isomorphism class of a spin^c –structure \mathfrak{s} and suppose that $G_{\mathfrak{s}}$ is a trivial extension. Then for any $c_1, c_2 \in H_G^*$ with $c_1 c_2 \neq 0$,*

$$\delta_{c_1}(Y) + \delta_{c_2}(\overline{Y}) \geq 0.$$

Proof The proof is similar to that of Theorem 4.1. Let $W = [0, 1] \times Y$ be the trivial cobordism from Y to itself. Choosing suitable metrics and reference connections we obtain a stable map

$$f : \text{Ker}_{\text{APS}}(D_{\hat{A}}^{\dagger})^+ \rightarrow \text{Coker}_{\text{APS}}(D_{\hat{A}}^{\dagger})^+ \wedge \text{SWF}(\overline{Y}, \mathfrak{s}, g) \wedge \text{SWF}(Y, \mathfrak{s}, g).$$

Note that $H^+(W) = \{0\}$ and hence $e(H^+(W)) = 1$. Applying Proposition 3.13 to this map we see that for any $c_1, c_2 \in H_G^*$ with $c_1 c_2 \neq 0$,

$$\text{ind}_{\text{APS}}(D_{\hat{A}}^{\dagger}) \leq \delta_{G, c_1 c_2}(\text{SWF}(\overline{Y}, \mathfrak{s}, g) \wedge \text{SWF}(Y, \mathfrak{s}, g)).$$

From the definition of the δ -invariant it is clear that

$$\delta_{G,c_1c_2}(SWF(\bar{Y}, \mathfrak{s}, g) \wedge SWF(Y, \mathfrak{s}, g)) \leq \delta_{G,c_1}(SWF(\bar{Y}, \mathfrak{s}, g)) + \delta_{G,c_2}(SWF(Y, \mathfrak{s}, g)),$$

and hence

$$\text{ind}_{\text{APS}}(D_{\hat{A}}^+) \leq \delta_{G,c_1}(\bar{Y}, \mathfrak{s}) + \delta_{G,c_2}(Y, \mathfrak{s}) + n(\bar{Y}, \mathfrak{s}, g) + n(Y, \mathfrak{s}, g).$$

On the other hand, for $W = [0, 1] \times Y$, equation (2-3) reduces to

$$\text{ind}_{\text{APS}}(D_{\hat{A}}^+) = n(\bar{Y}, \mathfrak{s}, g) + n(Y, \mathfrak{s}, g).$$

Hence we obtain $0 \leq \delta_{G,c_1}(\bar{Y}, \mathfrak{s}) + \delta_{G,c_2}(Y, \mathfrak{s})$. □

4.3 Induced cobordism maps

In this section we show that equivariant cobordisms induce maps on equivariant Seiberg–Witten–Floer cohomology.

Theorem 4.5 *Let W be a smooth, compact, oriented 4–manifold with boundary and with $b_1(W) = 0$. Suppose that G acts smoothly on W preserving the orientation and a spin^c -structure \mathfrak{s} . Suppose that $\partial W = \bar{Y}_1 \cup Y_2$ where Y_1 and Y_2 are rational homology 3–spheres and set $\mathfrak{s}_i = \mathfrak{s}|_{Y_i}$. Suppose G sends Y_i to itself. Then there is a morphism of graded $H_{G_{\mathfrak{s}_i}}^*$ -modules*

$$SW_G(W, \mathfrak{s}): HSW_G^*(Y_2, \mathfrak{s}_2) \rightarrow HSW_G^{*+b_+(W)-2\delta(W,\mathfrak{s})}(Y_1, \mathfrak{s}_1)$$

such that the diagram

$$\begin{array}{ccc} HSW_G^*(Y_2, \mathfrak{s}_2) & \xrightarrow{SW_G(W,\mathfrak{s})} & HSW_G^{*+b_+(W)-2\delta(W,\mathfrak{s})}(Y_1, \mathfrak{s}_1) \\ \downarrow & & \downarrow \\ HSW^*(Y_2, \mathfrak{s}_2) & \xrightarrow{SW(W,\mathfrak{s})} & HSW^{*+b_+(W)-2\delta(W,\mathfrak{s})}(Y_1, \mathfrak{s}_1) \end{array}$$

commutes, where the vertical arrows are the forgetful maps to nonequivariant Seiberg–Witten–Floer cohomology and $SW(W, \mathfrak{s})$ is the morphism of Seiberg–Witten–Floer cohomology groups induced by (W, \mathfrak{s}) .

Proof We give the proof in the case W is connected. The general case follows by a similar argument. As in the proof of Theorem 4.1, choosing suitable metrics and reference connections, we obtain a stable map

$$h: S^{\text{ind}_{\text{APS}}(D_{\hat{A}}^+)} \wedge SWF(Y_1, \mathfrak{s}_1, g_1) \rightarrow (H^+(W))^+ \wedge SWF(Y_2, \mathfrak{s}_2, g_2) \wedge S^{-k(D_1)\mathbb{C}},$$

where $k(D_1)$ is the dimension of the kernel of the Dirac operator on Y_1 . The induced map in equivariant cohomology takes the form

$$h^*: \tilde{H}_{G_s}^j((H^+(W))^+ \wedge SWF(Y_2, \mathfrak{s}_2, g_2) \wedge S^{-k(D_1)\mathbb{C}}) \rightarrow \tilde{H}_{G_s}^j(S^{\text{ind}_{\text{APS}}(D_{\hat{A}}^+)} \wedge SWF(Y_1, \mathfrak{s}_1, g_1)).$$

Using the Thom isomorphism, this is equivalent to

$$h^*: \tilde{H}_{G_s}^{j-b_+(W)+2k(D_1)}(SWF(Y_2, \mathfrak{s}_2, g_2)) \rightarrow \tilde{H}_{G_s}^{j-2\text{ind}_{\text{APS}}(D_A^+)}(SWF(Y_1, \mathfrak{s}_1, g_1)).$$

By (2-3), $\text{ind}_{\text{APS}} = \delta(W, \mathfrak{s}) + n(Y_2, \mathfrak{s}_2, g_2) + n(\bar{Y}_1, \mathfrak{s}_1, g_1)$ and by (2-4), $n(Y_1, \mathfrak{s}_1, g_1) + n(\bar{Y}_1, \mathfrak{s}_1, g_1) = -k(D_1)$. Replacing j by $j + b_+(W) - 2k(D_1) + 2n(Y_2, \mathfrak{s}_2, g_2)$, we see that h^* takes the form

$$h^*: \tilde{H}_{G_s}^{j+2n(Y_2, \mathfrak{s}_2, g_2)}(SWF(Y_2, \mathfrak{s}_2, g_2)) \rightarrow \tilde{H}_{G_s}^{j+b_+(W)-2\delta(W, \mathfrak{s})+2n(Y_1, \mathfrak{s}_1, g_1)}(SWF(Y_1, \mathfrak{s}_1, g_1)).$$

Then since $HSW_G^*(Y_i, \mathfrak{s}_i) = \tilde{H}_{G_s}^{*+2n(Y_i, \mathfrak{s}_i, g_i)}(SWF(Y_i, \mathfrak{s}_i, g_i))$, we see that h^* is equivalent to a map

$$SW_G(W, \mathfrak{s}): HSW_G^*(Y_2, \mathfrak{s}_2) \rightarrow HSW_G^{*+b_+(W)-2\delta(W, \mathfrak{s})}(Y_1, \mathfrak{s}_1).$$

Since this is a map of equivariant cohomologies induced by an equivariant map of spaces, it follows that $SW_G(W, \mathfrak{s})$ is a morphism of graded $H_{G_s}^*$ -modules. Restricting to the subgroup $S^1 \rightarrow G_s$, we obtain the commutative diagram in the statement of the theorem. \square

5 The case $G = \mathbb{Z}_p$

In this section we specialise to the case $G = \mathbb{Z}_p$ and $\mathbb{F} = \mathbb{Z}_p$, where p is a prime number. Then for $p = 2$ we have $H_G^* \cong \mathbb{F}[Q]$, where $\deg(Q) = 1$, and if p is odd we have $H_G^* \cong \mathbb{F}[R, S]/(R^2)$, where $\deg(R) = 1$ and $\deg(S) = 2$. Suppose $G = \langle \tau \rangle$ acts smoothly and orientation-preservingly on a rational homology 3–sphere Y , preserving a spin^c –structure \mathfrak{s} . The action of G is equivalent to giving an orientation-preserving diffeomorphism $\tau: Y \rightarrow Y$ such that $\tau^p = \text{id}$ and $\tau^*(\mathfrak{s}) = \mathfrak{s}$. Choose a lift $\tau' \in G_s$ of τ . Then $(\tau')^p = \zeta$ for some $\zeta \in S^1$. Replacing τ' by $\tilde{\tau} = \zeta^{-1/p} \tau'$, where $\zeta^{1/p}$ is a p^{th} root of ζ , we see that $\tilde{\tau}^p = \text{id}$. Hence G_s is a trivial extension.

5.1 δ –invariants

Definition 5.1 If $p = 2$, then for any integer $j \geq 0$, we define $d_j(Y, \mathfrak{s}, \tau, 2) = d_{\mathbb{Z}_2, Q^j}(Y, \mathfrak{s})$. If p is odd, then for any integer $j \geq 0$, we define $d_j(Y, \mathfrak{s}, \tau, p) = d_{\mathbb{Z}_p, S^j}(Y, \mathfrak{s})$. We also set $\delta_j(Y, \mathfrak{s}, \tau, p) = \frac{1}{2}d_j(Y, \mathfrak{s}, \tau, p)$. When p and τ are understood we will omit them from the notation and simply write $d_j(Y, \mathfrak{s})$ and $\delta_j(Y, \mathfrak{s})$.

In the case p is odd, one may also consider the invariants $d_{\mathbb{Z}_p, RS^j}(Y, \mathfrak{s})$. For simplicity we will not consider these invariants.

Theorem 5.2 *We have the following properties:*

- (1) $\delta_0(Y, \mathfrak{s}) \geq \delta(Y, \mathfrak{s})$;
- (2) $\delta_{j+1}(Y, \mathfrak{s}) \leq \delta_j(Y, \mathfrak{s})$ for all $j \geq 0$;
- (3) the sequence $\{\delta_j(Y, \mathfrak{s})\}_{j \geq 0}$ is eventually constant;

- (4) $\delta_j(Y, \mathfrak{s}) + \delta_j(\bar{Y}, \mathfrak{s}) \geq 0$ for all $j \geq 0$;
- (5) if Y is an L -space, then $\delta_j(Y, \mathfrak{s}) = \delta(Y, \mathfrak{s})$ for all $j \geq 0$.

Proof Item (1) is a restatement of Proposition 3.10. Item (2) follows from Proposition 3.9, taking $c_1 = Q^j$ and $c_2 = Q$ in the case $p = 2$, and $c_1 = S^j$ and $c_2 = S$ in the case p is odd. Item (4) is a special case of Theorem 4.4. For (3), first note that the difference $\delta_j(Y, \mathfrak{s}) - \delta_{j+1}(Y, \mathfrak{s})$ is always an integer because $\delta_{G,c}(Y, \mathfrak{s}) + n(Y, \mathfrak{s}, g) \in \mathbb{Z}$ for any metric g . From (2) and (4) and the fact that $n(Y, \mathfrak{s}, g) + n(\bar{Y}, \mathfrak{s}, g) \in \mathbb{Z}$, it follows that $\delta_j(Y, \mathfrak{s}) + \delta_j(\bar{Y}, \mathfrak{s})$ is a nonnegative, decreasing, integer-valued function. Hence the value of $\delta_j(Y, \mathfrak{s}) + \delta_j(\bar{Y}, \mathfrak{s})$ must eventually be constant. Using (2) again, it follows that $\delta_j(Y, \mathfrak{s})$ and $\delta_j(\bar{Y}, \mathfrak{s})$ are eventually constant. Item (5) is a restatement of Proposition 3.16. \square

Next, we specialise Theorem 4.1 to the case $G = \mathbb{Z}_p$.

Theorem 5.3 *Let W be a smooth, compact, oriented 4–manifold with boundary and with $b_1(W) = 0$. Suppose that $\tau : W \rightarrow W$ is an orientation-preserving diffeomorphism of order p and \mathfrak{s} is a spin^c –structure preserved by τ . Suppose each component of ∂W is a rational homology 3–sphere and that τ sends each component of ∂W to itself. Suppose that the subspace of $H^2(W; \mathbb{R})$ fixed by τ is negative definite. Then for all $j \geq 0$:*

- (1) If $\partial W = Y$ is connected, then

$$\delta(W, \mathfrak{s}) \leq \delta_j(Y, \mathfrak{s}|_Y) \quad \text{and} \quad \delta(\bar{W}, \mathfrak{s}) \geq \begin{cases} \delta_{j+b_+(W)}(\bar{Y}, \mathfrak{s}|_Y) & \text{if } p = 2, \\ \delta_{j+b_+(W)/2}(\bar{Y}, \mathfrak{s}|_Y) & \text{if } p \text{ is odd.} \end{cases}$$

- (2) If $\partial W = \bar{Y}_1 \cup Y_2$ has two connected components, then

$$\delta_j(Y_2, \mathfrak{s}|_{Y_2}) \geq \begin{cases} \delta_{j+b_+(W)}(Y_1, \mathfrak{s}|_{Y_1}) + \delta(W, \mathfrak{s}) & \text{if } p = 2, \\ \delta_{j+b_+(W)/2}(Y_1, \mathfrak{s}|_{Y_1}) + \delta(W, \mathfrak{s}) & \text{if } p \text{ is odd.} \end{cases}$$

Proof Let $H^+(W)$ denote a τ –invariant maximal positive definite subspace of $H^2(W; \mathbb{R})$ (which always exists because $G = \langle \tau \rangle$ is finite) and let e denote the image of the Euler class of $H^+(W)$ in $H_{\mathbb{Z}_p}^*$. To deduce the result from Theorem 4.1, we just need to check that $eQ^j \neq 0$ for all $j \geq 0$ if $p = 2$, and $eS^j \neq 0$ for all $j \geq 0$ if p is odd.

In the case $p = 2$, e is the top Stiefel–Whitney class of $H^+(W)$, which is easily seen to be $Q^{b_+(W)}$ because our assumption that the subspace of $H^2(W; \mathbb{R})$ fixed by τ is negative definite implies that τ acts as -1 on $H^+(W)$. Then clearly $eQ^j \neq 0$ for all $j \geq 0$.

Now suppose p is odd. Let L_i be the complex 1–dimensional representation on which τ acts as multiplication by ζ^i , $\zeta = e^{2\pi i/p}$. Any finite-dimensional real representation of G is the direct sum of a trivial representation and copies of the underlying real representations of the L_i for $1 \leq i \leq p-1$. The hypothesis that the subspace of $H^2(W; \mathbb{R})$ fixed by τ is negative definite means that as a representation of G , $H^+(W)$ contains no trivial summand. Hence $H^+(W)$ admits a complex structure such that

$H^+(W) \cong \bigoplus_{i=1}^{p-1} L_i^{m_i}$ for some integers $m_i \geq 0$. The Euler class of $H^+(W)$ is equal to its top Chern class. Under the map $H^2(\mathbb{Z}_p; \mathbb{Z}) \rightarrow H^2(\mathbb{Z}_p; \mathbb{Z}_p)$ one finds that $c_1(L_i)$ gets sent to iS . Hence

$$e = \prod_{i=1}^{p-1} (iS)^{m_i},$$

from which it is clear that $eS^j \neq 0$ for all $j \geq 0$. □

Remark 5.4 Suppose that p is odd. Then as in the proof of Theorem 5.3, $H^+(V)$ admits a complex structure. So if p is odd and the assumptions of Theorem 5.3 hold, then $b_+(W)$ must be even.

To keep notation simple, we will henceforth set $b'_\pm(W) = b_\pm(W)$ if $p = 2$ and $b'_\pm(W) = \frac{1}{2}b_\pm(W)$ if p is odd. Then (1) and (2) of Theorem 5.3 can be written more uniformly as

$$\delta(W, \mathfrak{s}) \leq \delta_j(Y, \mathfrak{s}|_Y) \quad \text{and} \quad \delta_{j+b'_+(W)}(\bar{Y}, \mathfrak{s}|_Y) \leq \delta(\bar{W}, \mathfrak{s}),$$

and

$$\delta_{j+b'_+(W)}(Y_1, \mathfrak{s}|_{Y_1}) + \delta(W, \mathfrak{s}) \leq \delta_j(Y_2, \mathfrak{s}|_{Y_2}).$$

Corollary 5.5 *Let W be a smooth, compact, oriented 4–manifold with boundary and with $b_1(W) = 0$. Suppose that $\tau : W \rightarrow W$ is an orientation-preserving diffeomorphism of order p and \mathfrak{s} is a spin^c –structure preserved by τ . Suppose that $Y = \partial W$ is a rational homology 3–sphere. Suppose that the subspace of $H^2(W; \mathbb{R})$ fixed by τ is zero. Then*

- (1) $\delta_j(Y, \mathfrak{s}|_Y) \geq -\frac{1}{8}\sigma(W)$ for all $j \geq 0$ and $\delta_j(Y, \mathfrak{s}|_Y) = -\frac{1}{8}\sigma(W)$ for $j \geq b'_-(W)$;
- (2) $\delta_j(\bar{Y}, \mathfrak{s}|_Y) \geq \frac{1}{8}\sigma(W)$ for all $j \geq 0$ and $\delta_j(\bar{Y}, \mathfrak{s}|_Y) = \frac{1}{8}\sigma(W)$ for $j \geq b'_+(W)$.

Proof It suffices to prove (1) since (2) follows by reversing orientation on W and Y . Since \mathbb{Z}_p preserves \mathfrak{s} , it follows that the image of $c_1(\mathfrak{s})$ in real cohomology lies in the subspace of $H^2(W; \mathbb{R})$ fixed by \mathbb{Z}_p . By assumption this space is zero; hence $c_1(\mathfrak{s}) = 0$ in real cohomology and hence $c_1(\mathfrak{s})^2 = 0$. So $\delta(W, \mathfrak{s}) = -\frac{1}{8}\sigma(W)$. Then from Theorem 5.3(1), we get $\delta_j(Y, \mathfrak{s}) \geq -\frac{1}{8}\sigma(W)$ for all $j \geq 0$. Reversing orientation on W and Y and applying Theorem 5.3(1), we also get that $\delta_{j+b'_-(W)}(Y, \mathfrak{s}) \leq -\frac{1}{8}\sigma(W)$ for all $j \geq 0$, or equivalently, $\delta_j(Y, \mathfrak{s}) \leq -\frac{1}{8}\sigma(W)$ for all $j \geq b'_-(W)$. Combining inequalities, we see that $\delta_j(Y, \mathfrak{s}) = -\frac{1}{8}\sigma(W)$ for $j \geq b'_-(W)$. □

5.2 Some algebraic results

In this section we collect some algebraic results which will be useful for computing δ invariants.

Let Y be a rational homology 3–sphere, $\tau : Y \rightarrow Y$ an orientation-preserving diffeomorphism of prime order p and \mathfrak{s} a spin^c –structure preserved by τ . Take $G = \mathbb{Z}_p = \langle \tau \rangle$ and $\mathbb{F} = \mathbb{Z}_p$. Let $\{E_r^{p,q}, d_r\}_{r \geq 2}$ denote the spectral sequence relating equivariant and nonequivariant Seiberg–Witten–Floer cohomologies. Then

$$E_2^{p,q} = H^p(\mathbb{Z}_p, HSW^q(Y, \mathfrak{s})),$$

where \mathbb{Z}_p acts on $HSW^q(Y, \mathfrak{s})$ via the action induced by τ . To simplify notation we will write H^q for $HSW^q(Y, \mathfrak{s})$ and d for $d(Y, \mathfrak{s})$. So $E_2^{p,q} = H^p(\mathbb{Z}_p, H^q)$. For fixed q , H^q is a finite-dimensional representation of \mathbb{Z}_p over \mathbb{F} . Moreover, for all sufficiently large k ,

$$(5-1) \quad H^{d+2k} = \mathbb{F}, \quad H^{d+2k+1} = 0.$$

Recall that H_G^* is isomorphic to $\mathbb{F}[Q]$ for $p = 2$ and to $\mathbb{F}[R, S]/(R^2)$ for odd p . In the case $p = 2$ we will set $S = Q^2$, so in all cases $S \in H_G^2$.

Lemma 5.6 *If V is a finite-dimensional representation of \mathbb{Z}_p over $\mathbb{F} = \mathbb{Z}_p$, then*

$$S: H^i(\mathbb{Z}_p; V) \rightarrow H^{i+1}(\mathbb{Z}_p; V)$$

is surjective for all $i \geq 0$ and an isomorphism for all $i \geq 1$. Furthermore, $\dim_{\mathbb{F}}(H^i(\mathbb{Z}_p; V)) \leq \dim_{\mathbb{F}}(V)$.

Proof Since \mathbb{Z}_p acts freely on S^1 , it follows from [12, page 114] that there is an element $\nu \in H^2(\mathbb{Z}_p; \mathbb{Z})$ (independent of V) such that the cup product $\nu: H^i(\mathbb{Z}_p; V) \rightarrow H^{i+2}(\mathbb{Z}_p; V)$ is an isomorphism for $i > 0$ and surjective for $i = 0$. Since V is a representation of \mathbb{Z}_p over \mathbb{F} , the same statement holds if we replace ν by its image in $H^2(\mathbb{Z}_p; \mathbb{F})$, which must have the form aS for some $a \in \mathbb{F}$. Moreover, $a \neq 0$ follows by considering the case that $V = \mathbb{Z}_p$ is the trivial representation. Hence the cup product $S: H^i(\mathbb{Z}_p; V) \rightarrow H^{i+2}(\mathbb{Z}_p; V)$ is an isomorphism for $i > 0$ and surjective for $i = 0$. We have by induction that $\dim_{\mathbb{F}}(H^i(\mathbb{Z}_p; V)) \leq \dim_{\mathbb{F}}(H^0(\mathbb{Z}_p; V))$ if i is even and $\dim_{\mathbb{F}}(H^i(\mathbb{Z}_p; V)) \leq \dim_{\mathbb{F}}(H^1(\mathbb{Z}_p; V))$ if i is odd. Then since $H^0(\mathbb{Z}_p; V)$ and $H^1(\mathbb{Z}_p; V)$ can both be expressed as certain subquotients of V , it follows that $\dim_{\mathbb{F}}(H^i(\mathbb{Z}_p; V)) \leq \dim_{\mathbb{F}}(V)$ for all i . \square

Lemma 5.7 *For each $r \geq 2$, the map $S: E_r^{p,q} \rightarrow E_r^{p+2,q}$ is surjective for all $p \geq 0$ and an isomorphism for all $p \geq r - 1$.*

Proof Recall that $E_2^{p,q} = H^p(\mathbb{Z}_p, H^q)$. Hence $S: E_2^{p,q} \rightarrow E_2^{p+2,q}$ is surjective for all p and an isomorphism for all $p \geq 1$, by Lemma 5.6. This proves the case $r = 2$. Now we proceed by induction. Let $r > 2$ and suppose that $S: E_{r-1}^{p,q} \rightarrow E_{r-1}^{p+2,q}$ is surjective for all $p \geq 0$ and an isomorphism for all $p \geq r - 1$. Let $x \in E_r^{p+2,q}$. Then $x = [y]$ for some $y \in E_{r-1}^{p+2,q}$ with $d_{r-1}(y) = 0$. By the inductive hypothesis $y = Sz$ for some $z \in E_{r-1}^{p,q}$. Then $Sd_{r-1}(z) = d_{r-1}(Sz) = d_{r-1}(y) = 0$. That is, $Sd_{r-1}(z) = 0$. However, $d_{r-1}(z) \in E_{r-1}^{p+r-1,q+2-r}$ and $p + r - 1 \geq r - 2$, so $S: E_{r-1}^{p+r-1,q+2-r} \rightarrow E_{r-1}^{p+r+1,q+2-r}$ is an isomorphism by the inductive hypothesis. Hence $Sd_{r-1}(z) = 0$ implies that $d_{r-1}(z) = 0$. So z defines a class $w = [z] \in E_r^{p,q}$. Then $Sw = [Sz] = [y] = x$. Hence $S: E_r^{p,q} \rightarrow E_r^{p+2,q}$ is surjective for all $p \geq 0$.

Now suppose that $p \geq r - 1$ and consider $x \in E_r^{p,q}$ satisfying $Sx = 0$. Write $x = [y]$ for some $y \in E_{r-1}^{p,q}$ satisfying $d_{r-1}(y) = 0$. Then $0 = Sx = S[y] = [Sy]$. Hence $Sy = d_{r-1}(z)$ for some $z \in E_{r-1}^{p-r+3,q+r-2}$. By the inductive hypothesis and since $p - r + 3 \geq (r - 1) - r + 3 = 2$, we have that $z = Sw$ for some $w \in E_{r-1}^{p-r+1,q+r-2}$. Hence $Sy = d_{r-1}(z) = d_{r-1}(Sw) = Sd_{r-1}(w)$. By the inductive hypothesis,

$S: E_{r-1}^{p,q} \rightarrow E_{r-1}^{p+2,q}$ is injective; hence $y = d_{r-1}(w)$ and $x = [y] = [d_{r-1}(w)] = 0$. So $S: E_r^{p,q} \rightarrow E_r^{p+2,q}$ is injective for $p \geq r - 1$. □

From the above lemma, we see that $E_r^{p,q}$ does not depend on p , provided $p \geq r - 1$. Let M_r^q denote $E_r^{p,q}$ for $p \geq r - 1$. Moreover, since the differentials $\{d_r\}$ for the spectral sequence $E_r^{p,q}$ commute with S , they induce differentials $d_r: M_r^q \rightarrow M_r^{q+1-r}$ for which M_{r+1} is the cohomology of $d_r: M_r \rightarrow M_r$. Thus M_{r+1} is a subquotient of M_r .

For any module V over $\mathbb{F}[U]$, we define

$$V_{\text{red}} = \{x \in V \mid U^k x = 0 \text{ for some } k \geq 0\} \quad \text{and} \quad V^\infty = V/V_{\text{red}}.$$

Lemma 5.8 *For each $r \geq 2$, the image of the differential d_r is contained in $(E_r^{*,*})_{\text{red}}$.*

Proof By (5-1) there exists a k_0 such that $H^{d+2k} = \mathbb{F}$ and $H^{d+2k+1} = 0$ for all $k \geq k_0$. Hence the action of τ is trivial in these degrees and we have

$$E_2^{p,d+2k} = \mathbb{F}, \quad E_2^{p,d+2k+1} = 0$$

for all $k \geq k_0$. Since $SWF(Y, \mathfrak{s}, g)$ is a space of type \mathbb{Z}_p -SWF, the localisation theorem in equivariant cohomology implies that there exists a $k_1 \geq k_0$ such that the generator $x \in E_2^{0,d+2k_1} = \mathbb{F}$ satisfies $d_r(x) = 0$ for all $r \geq 2$. Then if $y \in E_2^{p,q}$ with $q \geq d + 2k_1$, it follows that y is of the form $y = cU^a x$ for some $a \geq 0$, where $c \in H_G^p$. Hence $d_r(y) = 0$ for all $r \geq 2$. Now let $y \in E_r^{p,q}$ where p and q are arbitrary. Then there exists some $a \geq 0$ such that $q + 2a \geq d + 2k_1$; hence $U^a d_r(y) = d_r(U^a y) = 0$. Therefore $d_r(y) \in (E_r^{*,*})_{\text{red}}$. □

Recall that H^∞ is a free $\mathbb{F}[U]$ -module of rank 1 with generator in degree d . Hence we may write $H^\infty = \mathbb{F}[U]\theta$ where $\deg(\theta) = d$. Next, observe that $E_2^{0,*}$ is the τ -invariant part of H^* , hence may be regarded as an $\mathbb{F}[U]$ -submodule of H^* . Similarly, since $E_{r+1}^{0,*}$ is the kernel of d_r restricted to $E_r^{0,*}$, it follows that $E_{r+1}^{0,*}$ can be identified with an $\mathbb{F}[U]$ -submodule of $E_r^{0,*}$. Hence $\{E_r^{0,*}\}$ may be regarded as a decreasing sequence of $\mathbb{F}[U]$ -submodules of H^* . Let S_r denote the image of $E_r^{0,*}$ under the quotient map $H^* \rightarrow H^\infty = H^*/H_{\text{red}}$. The localisation theorem in equivariant cohomology implies that S_r is nonzero and that the sequence S_r eventually stabilises. Then since S_r is a nonzero graded submodule of $H^\infty = \mathbb{F}[U]\theta$, it follows that $S_r = \mathbb{F}[U]U^{m_r}\theta$ for some nonnegative integer m_r . Note also that the sequence $\{m_r\}$ is increasing and is eventually constant.

Lemma 5.9 *For each $r \geq 2$,*

$$m_{r+1} - m_r \leq \dim_{\mathbb{F}}((M_r)_{\text{red}}) - \dim_{\mathbb{F}}((M_{r+1})_{\text{red}}).$$

Proof The classes $U^{j+m_r}\theta$ with $0 \leq j < m_{r+1} - m_r$ form a basis for S_r/S_{r+1} . Choose a lift $x_r \in E_r^{0,d+2m_r}$ of $U^{m_r}\theta \in S_r$. Then $d_r(U^j x_r) \neq 0$ for $0 \leq j < m_{r+1} - m_r$, for if $d_r(U^j x_r) = 0$ for

some $0 \leq j < m_{r+1} - m_r$, then we would have $U^{j+m_r}\theta \in S_{r+1}$. Observe that $d_r(U^j x_r) \in E_r^{r,*}$. By Lemma 5.7 and the definition of M_r , we see that $d_r(U^j x_r)$ can be identified with a nonzero element of M_r . Moreover, $d_r(U^j x_r) \in (M_r)_{\text{red}}$, by Lemma 5.8. Now since the $d_r(U^j x_r)$ are nonzero and have distinct degrees, they span a subspace of $(M_r)_{\text{red}}$ of dimension $m_{r+1} - m_r$. Furthermore, this subspace lies in the image of d_r ; hence $m_{r+1} - m_r \leq \dim_{\mathbb{F}}((M_r)_{\text{red}}) - \dim_{\mathbb{F}}((M_{r+1})_{\text{red}})$. \square

Proposition 5.10 *Suppose that τ acts trivially on $HSW^*(Y, \mathfrak{s})$. Then*

$$\delta_1(Y, \mathfrak{s}) - \delta(Y, \mathfrak{s}) \leq \dim_{\mathbb{F}}(HSW_{\text{red}}(Y, \mathfrak{s})).$$

Proof Recall that $d = d(Y, \mathfrak{s})$. Hence $\delta(Y, \mathfrak{s}) = \frac{1}{2}d$. From the definition of the invariant $\delta_1(Y, \mathfrak{s})$, it follows that for all sufficiently large r ,

$$\delta_1(Y, \mathfrak{s}) = m_r + \delta(Y, \mathfrak{s}).$$

By Lemma 5.9, for each $r \geq 2$,

$$m_{r+1} - m_r \leq \dim_{\mathbb{F}}((M_r)_{\text{red}}) - \dim_{\mathbb{F}}((M_{r+1})_{\text{red}}),$$

and summing from 2 to $r - 1$, we get

$$m_r - m_2 \leq \dim_{\mathbb{F}}((M_2)_{\text{red}}).$$

However since τ acts trivially on $HSW^*(Y, \mathfrak{s})$, we have that $E_2^{p,*} = HSW^*(Y, \mathfrak{s})$ for all $p \geq 0$. Hence $m_2 = 0$, $M_2 = HSW^*(Y, \mathfrak{s})$ and $(M_2)_{\text{red}} = HSW_{\text{red}}(Y, \mathfrak{s})$. Taking r sufficiently large,

$$\delta_1(Y, \mathfrak{s}) - \delta(Y, \mathfrak{s}) = m_r = m_r - m_2 \leq \dim_{\mathbb{F}}(HSW_{\text{red}}). \quad \square$$

6 Branched double covers of knots

6.1 Concordance invariants

Let $K \subset S^3$ be a knot in S^3 . Let $Y = \Sigma_2(K)$ be the branched double cover of S^3 , branched over K . Let $\pi : Y \rightarrow S^3$ denote the covering map. One finds that $b_1(Y) = 0$. Manolescu and Owens [51] define a knot invariant

$$\delta(K) = 2d(\Sigma_2(K), \mathfrak{t}_0) = 4\delta(\Sigma_2(K), \mathfrak{t}_0),$$

where \mathfrak{t}_0 is the spin^c -structure induced from the unique spin -structure on $\Sigma_2(K)$ (see [51, Section 2] for an explanation of this). It is shown in [51] that $\delta(K)$ is always integer-valued and descends to a surjective group homomorphism $\delta : \mathcal{C} \rightarrow \mathbb{Z}$, where \mathcal{C} is the smooth concordance group of knots in S^3 .

The covering involution on Y determines an action of $G = \mathbb{Z}_2$ on Y preserving \mathfrak{t}_0 (by uniqueness of the underlying spin -structure). Hence, for each $j \geq 0$, we may define the knot invariant

$$\delta_j(K) = 2d_j(\Sigma_2(K), \mathfrak{t}_0) = 4\delta_j(\Sigma_2(K), \mathfrak{t}_0).$$

Since $d_j(\Sigma_2(K), \mathfrak{t}_0) - d(\Sigma_2(K), \mathfrak{t}_0) \in 2\mathbb{Z}$, it follows that $\delta_j(K) - \delta(K) \in 4\mathbb{Z}$. Then, since $\delta(K)$ is integer-valued, it follows that the $\delta_j(K)$ are also integer-valued and moreover $\delta_j(K) = \delta(K) \pmod{4}$.

Proposition 6.1 *For each $j \geq 0$, $\delta_j(K)$ depends only on the concordance class of K ; hence δ_j descends to a concordance invariant $\delta_j: \mathcal{C} \rightarrow \mathbb{Z}$.*

Proof For an oriented knot K , recall that $-K$ denotes the knot obtained by reversing orientation on S^3 and K . It follows that $\Sigma_2(-K) = \overline{\Sigma_2(K)}$. A concordance of oriented knots K_1 and K_2 is a smooth embedding of $\Sigma = [0, 1] \times S^1$ in $[0, 1] \times S^3$ having boundary $-K_1 \cup K_2$. Taking the double cover of $[0, 1] \times S^3$ branched along Σ gives a \mathbb{Z}_2 -equivariant cobordism W from $\Sigma_2(K_1)$ to $\Sigma_2(K_2)$. From the calculations in [36, Section 3], one sees that W is a rational homology cobordism. We claim that W is spin. To see this, choose a smoothly embedded surface Σ in D^4 whose boundary is K_1 . Let W' be the double cover of $D^4 \cup [0, 1] \times S^3 \cong D^4$ branched over $\Sigma \cup [0, 1] \times S^1$. From [35] we see that W' is spin. Since W is embedded in W' , it follows that W is spin as well. Any spin-structure \mathfrak{t} on W will restrict on each component of the boundary to the unique spin-structure on the branched double cover $\Sigma_2(K_i)$. The result now follows by applying Corollary 4.3 to (W, \mathfrak{t}) . \square

We note that the δ_j are *not* group homomorphisms.

Let $\sigma(K)$ denote the signature of K and $g_4(K)$ the smooth 4-genus. Set $\sigma'(K) = -\frac{1}{2}\sigma(K)$. We also define $b_+(K) = g_4(K) - \sigma'(K)$ and $b_-(K) = g_4(K) + \sigma'(K)$.

Proposition 6.2 *The knot concordance invariants δ_j have the properties*

- (1) $\delta_0(K) \geq \delta(K)$;
- (2) $\delta_{j+1}(K) \leq \delta_j(K)$ for all $j \geq 0$;
- (3) $\delta_j(K) \geq \sigma'(K)$ for all $j \geq 0$ and $\delta_j(K) = \sigma'(K)$ for $j \geq b_-(K)$;
- (4) $\delta_j(-K) \geq -\sigma'(K)$ for all $j \geq 0$ and $\delta_j(-K) = -\sigma'(K)$ for $j \geq b_+(K)$;
- (5) if $\Sigma_2(K)$ is an L -space, then $\delta_j(K) = \delta(K)$ and $\delta_j(-K) = \delta(-K)$ for all $j \geq 0$.

Proof Items (1), (2) and (5) follow from (1), (2) and (5) of Theorem 5.2. For (3) and (4), choose a smooth embedded surface $\Sigma \subset D^4$ in the 4-ball of genus $g_4(K)$ which bounds K . Let W be the double cover of D^4 branched along Σ . From [35] it follows that W is spin. Let \mathfrak{t} be any spin-structure on W . Then $\mathfrak{t}|_{\Sigma_2(K)} = \mathfrak{t}_0$ by uniqueness of \mathfrak{t}_0 . Next, observe that $H^2(W; \mathbb{R})^{\mathbb{Z}_2} = H^2(D^4; \mathbb{R}) = 0$. Then (3) and (4) follow by applying Corollary 5.5 to (W, \mathfrak{t}) . \square

Corollary 6.3 *If K is a knot such that $\Sigma_2(K)$ is an L -space, then $\delta(K) = \sigma'(K)$.*

Proof This follows by (3) and (5) of Proposition 6.2 \square

Remark 6.4 In particular, if K is quasialternating, then $\Sigma_2(K)$ is an L -space [57]. This recovers the main result of [47] that $\delta(K) = \sigma'(K)$ for quasialternating knots.

Theorem 6.5 For a knot K , let $j_+(K)$ be the smallest positive integer such that $\delta_j(K) = \sigma'(K)$ and $j_-(K)$ the smallest positive integer such that $\delta_j(-K) = -\sigma'(K)$. Then

$$g_4(K) \geq \max\{\sigma'(K) + j_-(K), -\sigma'(K) + j_+(K)\}.$$

Remark 6.6 Observe that the right-hand side of this inequality is at least $\frac{1}{2}|\sigma(K)|$. Hence we have obtained a strengthening of the well-known inequality $g_4(K) \geq \frac{1}{2}|\sigma(K)|$ [52].

Proof From Proposition 6.2 we have that $\delta_j(K) = \sigma'(K)$ for $j \geq g_4(K) + \sigma'(K)$ and $\delta_j(-K) = -\sigma'(K)$ for $j \geq g_4(K) - \sigma'(K)$. Hence $j_+(K) \leq g_4(K) + \sigma'(K)$ and $j_-(K) \leq g_4(K) - \sigma'(K)$. \square

Remark 6.7 In this section we have used branched double covers $\Sigma_2(K)$ of knots equipped with their natural \mathbb{Z}_2 -action to obtain a sequence of concordance invariants. Similarly, for any odd prime p we may consider the cyclic branched cover $\Sigma_p(K)$ with its natural \mathbb{Z}_p -action. Once again there is a canonical spin^c -structure \mathfrak{t}_0 [29] and so we may define a sequence of invariants

$$\delta_{(p),j}(K) = 2d_{\mathbb{Z}_p, S^j}(\Sigma_p(K), \mathfrak{t}_0)$$

depending on a prime p and an integer $j \geq 0$. By similar arguments to the $p = 2$ case one finds that these are integer-valued knot concordance invariants of K .

7 Computations and applications

7.1 Brieskorn homology spheres

Let p, q and r be pairwise coprime positive integers and let $Y = \Sigma(p, q, r)$ be the corresponding Brieskorn integral homology 3-sphere. Then Y has a unique spin^c -structure and so when speaking of the Floer homology of Y we omit the mention of the spin^c -structure.

Recall that $\Sigma(p, q, r)$ can be realised as the p -fold cyclic cover of S^3 branched along the torus knot $T_{q,r}$. In particular, this construction defines an action of \mathbb{Z}_p on Y . Let $\tau: Y \rightarrow Y$ denote the generator of this action. Recall that $\Sigma(p, q, r)$ is obtained by taking the link of the singularity

$$\{(x, y, z) \in \mathbb{C}^3 \mid x^p + y^q + z^r = 0\}.$$

Then τ is given by $(x, y, z) \mapsto (e^{2\pi i/p}x, y, z)$. This map is isotopic to the identity through the homotopy $(x, y, z) \mapsto (e^{2\pi i qrt}x, e^{2\pi i prt}y, e^{2\pi i pqt}z)$ for $t \in [0, (qr)^*]$, where $0 < (qr)^* < p$ denotes the multiplicative inverse of $qr \bmod p$. It follows that τ acts trivially on $HF^+(Y)$.

Henceforth we will assume that p is a prime number. Set $\mathbb{F} = \mathbb{Z}_p$ and recall that $H_{\mathbb{Z}_p}^* \cong \mathbb{F}[Q]$ where $\deg(Q) = 1$ if $p = 2$, and $H_{\mathbb{Z}_p}^* \cong \mathbb{F}[R, S]/(R^2)$ where $\deg(R) = 1$ and $\deg(S) = 2$ if p is odd. Let \mathfrak{s} denote the unique spin^c -structure on Y . As in Section 5, we let $\delta_j(Y, \mathfrak{s}, \tau, p)$ denote

$\delta_{\mathbb{Z}_p, Q^p}(Y, \mathfrak{s})$ for $p = 2$ or $\delta_{\mathbb{Z}_p, S^p}(Y, \mathfrak{s})$ for odd p . We will further abbreviate this to $\delta_j(Y, \tau)$. When $p = 2$, $\delta_j(Y, \tau) = \frac{1}{4}\delta_j(T_{q,r})$, where $\delta_j(K)$ denotes the knot concordance invariant introduced in Section 6.1. More generally, $\delta_j(Y, \tau) = \frac{1}{4}\delta_{(p),j}(T_{q,r})$, where $\delta_{(p),j}(K)$ is the knot concordance invariant defined in Remark 6.7.

Example 7.1 Let $(p, q, r) = (2, 3, 5)$. Then $Y = \Sigma(2, 3, 5)$ is the Poincaré homology 3–sphere. Since $\Sigma(2, 3, 5)$ has spherical geometry, it is an L –space [56, Proposition 2.3]. Therefore

$$\delta_j(T_{3,5}) = \delta(T_{3,5}) = \sigma'(T_{3,5}) = 4 \quad \text{for all } j \geq 0.$$

The property of being an L –space does not depend on the choice of orientation, so we also have

$$\delta_j(-T_{3,5}) = \delta(-T_{3,5}) = -4 \quad \text{for all } j \geq 0.$$

The same argument applied to $p = 3$ or 5 gives

$$\delta_{(3),j}(T_{2,5}) = \delta_{(5),j}(T_{2,3}) = 4 \quad \text{for all } j \geq 0$$

and

$$\delta_{(3),j}(-T_{2,5}) = \delta_{(5),j}(-T_{2,3}) = -4 \quad \text{for all } j \geq 0.$$

Proposition 7.2 Let p, q and r be positive, pairwise coprime integers and assume that p is prime. Then $\delta_j(\Sigma(p, q, r), \tau) = -\lambda(\Sigma(p, q, r))$ for all $j \geq 0$, where $\lambda(\Sigma(p, q, r))$ is the Casson invariant of $\Sigma(p, q, r)$. Furthermore,

$$\delta_{(p),j}(T_{q,r}) = -\frac{1}{2} \sum_{j=1}^{p-1} \sigma_{j/p}(T_{q,r}) \quad \text{for all } j \geq 0,$$

where $\sigma_\alpha(K)$ is the Tristram–Levine signature of K .

Proof Recall that $Y = \Sigma(p, q, r)$ is the boundary of a negative definite plumbing [53] whose plumbing graph has only one bad vertex in the terminology of [55]. Then it follows from [55, Corollary 1.4] that $HF^+(\bar{Y})$ is concentrated in even degrees. Consequently, $HF_{\text{red}}^+(Y)$ is concentrated in odd degrees. (Note that [55] uses \mathbb{Z} coefficients, but it is shown there that $HF_{\text{red}}^+(Y; \mathbb{Z})$ has no torsion and hence by the universal coefficient theorem, [55, Corollary 1.4] also holds for \mathbb{Z}_p coefficients.) Therefore,

$$(7-1) \quad \chi(HF_{\text{red}}^+(Y)) = \dim_{\mathbb{F}}(HF_{\text{red,even}}^+(Y)) - \dim_{\mathbb{F}}(HF_{\text{red,odd}}^+(Y)) = -\dim_{\mathbb{F}}(HF_{\text{red}}^+(Y)).$$

By [54, Theorem 1.3], $\chi(HF_{\text{red}}^+(Y))$ is related to the Casson invariant $\lambda(Y)$ via the formula

$$(7-2) \quad \chi(HF_{\text{red}}^+(Y)) = \lambda(Y) + \delta(Y).$$

Hence

$$(7-3) \quad \dim_{\mathbb{F}}(HF_{\text{red}}^+(Y)) = -\lambda(Y) - \delta(Y).$$

Moreover, from [18; 27],

$$\lambda(\Sigma(p, q, r)) = \frac{1}{8} \sum_{j=1}^{p-1} \sigma_{j/p}(T_{q,r}) = \frac{1}{8} \sigma(M(p, q, r)),$$

where $M(p, q, r)$ is the Milnor fibre

$$M(p, q, r) = \{(x, y, z) \in \mathbb{C}^3 \mid x^p + y^q + z^r = \delta\} \cap D^6$$

(where δ is a sufficiently small nonzero complex number). Recall that $M(p, q, r)$ is a compact smooth 4–manifold with boundary diffeomorphic to $\Sigma(p, q, r)$. Moreover, $M(p, q, r)$ has the homotopy type of a wedge of 2–spheres, so $b_1(M(p, q, r)) = 0$. Further, $M(p, q, r)$ is a p –fold cyclic cover of D^4 branched along a surface bounding $T_{q,r}$. Hence the action of $\mathbb{Z}_p = \langle \tau \rangle$ on Y extends to $M(p, q, r)$. From [29, Lemma 2.1] it follows that there is a \mathbb{Z}_p invariant spin–structure \mathfrak{t}_0 on $M(p, q, r)$. Since $M(p, q, r)$ is a cyclic p –fold cover of D^4 , it follows that the subspace of $H^2(M(p, q, r); \mathbb{R})$ fixed by τ is zero. Hence Corollary 5.5 may be applied, giving

$$\delta_j(Y, \tau) \geq -\frac{1}{8} \sigma(M(p, q, r)) = -\lambda(Y) \quad \text{for all } j \geq 0.$$

Since τ acts trivially on $HF^+(Y)$, Proposition 5.10 implies that

$$\delta_0(Y, \tau) - \delta(Y) \leq \dim_{\mathbb{F}}(HF_{\text{red}}^+(Y)) = -\lambda(Y) - \delta(Y).$$

Hence $\delta_0(Y, \tau) \leq -\lambda(Y)$. On the other hand, $\delta_0(Y, \tau) \geq \delta_j(Y, \tau) \geq -\lambda(Y)$ for any $j \geq 0$. Hence $\delta_j(Y, \tau) = -\lambda(Y)$ for all $j \geq 0$. Therefore we also have

$$\delta_{(p),j}(T_{q,r}) = 4\delta_j(Y, \tau) = -4\lambda(Y) = -\frac{1}{2} \sum_{j=1}^{p-1} \sigma_{j/p}(T_{q,r})$$

for all $j \geq 0$. □

The above result shows that the values of $\delta_{(p),j}(T_{q,r})$ do not depend on j . In contrast, the values of $\delta_{(p),j}(-T_{q,r})$ usually do depend on j , as the following propositions illustrate.

Proposition 7.3 *Let $(a, b) = (3, 6n - 1)$ for $n \geq 1$. Then*

$$\delta(-T_{3,6n-1}) = -4, \quad \sigma'(-T_{3,6n-1}) = -4n$$

and

$$\delta_j(-T_{3,6n-1}) = \begin{cases} -4\left(\lfloor \frac{1}{2}j \rfloor + 1\right) & \text{if } 0 \leq j \leq 2n - 3, \\ -4n & \text{if } j \geq 2n - 2. \end{cases}$$

Proof The case $n = 1$ is already covered in Example 7.1, so we assume $n \geq 2$. Set $Y_{a,b} = \Sigma_2(T_{a,b}) = \Sigma(2, a, b)$ and let τ be the covering involution. Then $\delta_j(-T_{3,6n-1}) = 4\delta_j(\bar{Y}_{3,6n-1})$. From the computations in [54, Section 8] we find that $d(\bar{Y}_{3,6n-1}) = -2$, $SWF_{\text{red}}^*(\bar{Y}_{3,6n-1}) = (\mathbb{F}_{-2})^{n-1}$, where the subscript indicates degree. To simplify notation we let $V = SWF_{\text{red}}^*(\bar{Y}_{3,6n-1}) = (\mathbb{F}_{-2})^{n-1}$. Then

$$E_2^{*,*} \cong \mathbb{F}[U, Q]\theta \oplus V[Q],$$

where the bidegree is given as follows: θ and all elements of V have bidegree $(0, -2)$, U has bidegree $(0, 2)$ and Q has bidegree $(1, 0)$. Then $E_2^{p,q} = 0$ for $q < -2$. It follows that all the differentials in the spectral sequence are zero on θ and on V , since d_r sends $E_r^{p,-2}$ to $E_r^{p+r,-1-r}$ and $-1-r < -2$ for $r \geq 2$. Hence d_r is zero on all of E_r and $E_\infty^{*,*} \cong E_2^{*,*}$. Let \mathcal{F}_j denote the filtration on $HSW_{\mathbb{Z}_2}^*(\bar{Y}_{3,6n-1})$ corresponding to the spectral sequence, so $\mathcal{F}_j/\mathcal{F}_{j+1} \cong E_\infty^{j,*}$. In particular, $\mathcal{F}_1/\mathcal{F}_2 \cong \mathbb{F}[U]\theta \oplus V$. Choose lifts of θ and V to \mathcal{F}_1 . We lift $U^j\theta$ by taking the lift of θ and applying U^j . Hence we obtain a short exact sequence of $\mathbb{F}[U, Q]$ -modules

$$0 \rightarrow \mathbb{F}[U] \oplus V \rightarrow HSW_{\mathbb{Z}_2}^*(\bar{Y}_{3,6n-1}) \rightarrow \mathcal{F}_2 \rightarrow 0.$$

Next, for each $j \geq 0$, Q induces an isomorphism $Q: \mathcal{F}_j/\mathcal{F}_{j+1} \rightarrow \mathcal{F}_{j+1}/\mathcal{F}_{j+2}$; hence by applying Q repeatedly to $\mathbb{F}[U]\theta \oplus V$, we obtain a splitting of the filtration $\{\mathcal{F}_j\}$ as $\mathbb{F}[Q]$ -modules. The splittings give an isomorphism of $\mathbb{F}[Q]$ -modules

$$HSW_{\mathbb{Z}_2}^*(\bar{Y}_{3,6n-1}) \cong \mathbb{F}[U, Q]\theta \oplus V[Q].$$

However, this is not necessarily an isomorphism of $\mathbb{F}[U, Q]$ -modules. Under this isomorphism, U corresponds to an endomorphism of the form

$$\hat{U} = U_2 + QU_1 + Q^2U_0 + Q^3U_{-1} + \dots,$$

where $U_j: HSW^*(\bar{Y}_{3,6n-1}) \rightarrow HSW^{*+j}(\bar{Y}_{3,6n-1})$ and $U_2 = U$. Since $HSW^*(\bar{Y}_{3,6n-1})$ is concentrated in even degrees, $U_j = 0$ for odd j . Moreover, our construction is such that $U_j\theta = 0$ for $j \neq 2$. It follows that $U_j = 0$ for $j < 0$, as V is concentrated in a single degree. So we get

$$\hat{U} = U + Q^2U_0$$

for some $U_0: V \rightarrow HSW^0(\bar{Y}_{3,6n-1})$.

To simplify notation set $d_j = d_j(\bar{Y}_{3,6n-1})$. Using Proposition 3.14 we obtain the following characterisation of d_j :

$$d_j = \min\{i \mid \hat{U}^r x = U^m Q^j \theta \pmod{Q^{j+1}} \text{ for some } x \in HSW_{\mathbb{Z}_2}^i(\bar{Y}_{3,6n-1}) \text{ and } r, m \geq 0\} - j.$$

Recall that $\delta_j(-T_{3,6n-1}) = \sigma'(-T_{3,6n-1}) = -4n$ for sufficiently large j . Hence $d_j = -2n$ for sufficiently large j . Choose such a j . From the above characterisation of d_j there exists $x \in HSW_{\mathbb{Z}_2}^{j-2n}(\bar{Y}_{3,6n-1})$ such that $\hat{U}^r x = U^m Q^j \theta + \dots$ where \dots denotes terms of higher order in Q . We have that $x = Q^a y$ for some $a \leq j$. Then $\hat{U}^r Q^a y = U^m Q^j \theta + \dots$. Since Q is injective we may cancel, giving $\hat{U}^r y = U^m Q^{j-a} \theta + \dots$. If $a = j$, then $\hat{U}^r y = U^m \theta + \dots$. But $\hat{U} = U + Q^2U_0$, so $\hat{U}^r y = U^r y + \dots$; hence $U^r y = U^m \theta + \dots$. From the definition of the usual d -invariant we must have $\deg(y) \geq d(\bar{Y}_{3,6n-1}) = -2$. Hence $j - 2n = \deg(x) = a + \deg(y) = j - 2$, which is a contradiction since we have assumed that $n > 1$. It follows that $a < j$. We must have $y \in V$ for if $y = U^b \theta \pmod{V}$, then we would have $\hat{U}^r x = U^{r+b} Q^a \theta + \dots$, which contradicts $\hat{U}^r x = U^m Q^j \theta + \dots$ as $a < j$. Therefore $y \in V$. In particular $\deg(y) = -2$ and $j - 2n = \deg(x) = a - 2$.

Let b be the smallest positive integer such that $U_0^b y \notin V$. Such a b exists since

$$\hat{U}^r y = (U + Q^2 U_0)^r y = U^m Q^{j-a} \theta + \dots$$

and U is zero on V . Then it follows that $r \geq b$ and

$$\hat{U}^r y = (U + Q^2 U_0)^r y = U^{r-b} Q^{2b} (U_0^b y) + \dots = U^m Q^{j-a} \theta + \dots .$$

Hence $2b = j - a$. So we have shown that $j = a + 2b$ and $j - 2n = a - 2$. Hence $b = n - 1$. But since $\dim_{\mathbb{F}}(V) = n - 1 = b$, it follows that there exists a $v \in V$ such that $v, U_0 v, U_0^2 v, \dots, U_0^{n-2} v$ is a basis for V and $U_0^{n-1} v = \theta \bmod V$. Now it is straightforward to see that the sequence $\{d_j\}$ must have the form $-2, -2, -4, -4, -6, -6, \dots$, for $j \leq 2n - 3$ and $d_j = -2n$ for $j \geq 2n - 2$. \square

Proposition 7.4 *Let $(a, b) = (3, 6n + 1)$ for $n \geq 1$. Then*

$$\delta(-T_{3,6n+1}) = 0, \quad \sigma'(-T_{3,6n+1}) = -4n$$

and

$$\delta_j(-T_{3,6n+1}) = \begin{cases} -4 \lfloor \frac{1}{2} j \rfloor & \text{if } 0 \leq j \leq 2n - 1, \\ -4n & \text{if } j \geq 2n. \end{cases}$$

Proof By [54, Section 8], $d(\bar{Y}_{3,6n+1}) = 0$, $SWF_{\text{red}}^*(\bar{Y}_{3,6n+1}) = (\mathbb{F}_0)^n$ and $\sigma'(-T_{3,6n+1}) = 4n$. From here essentially the same argument as in Proposition 7.3 gives the result. \square

Remark 7.5 We can use Theorem 6.5 and the computations in Propositions 7.3 and 7.4 to obtain a lower bound for the 4–genus. From Proposition 7.3, we see that $\sigma'(T_{3,6n-1}) = 4n$ and $j_-(T_{3,6n-1}) = 2n - 2$; hence $g_4(T_{3,6n-1}) \geq 2n - 2 + 4n = 6n - 2$. On the other hand, from the positive solution to the Milnor conjecture [38], we know that $g_4(T_{a,b}) = \frac{1}{2}(a - 1)(b - 1)$. In particular, $g_4(T_{3,6n-1}) = 6n - 2$. Hence the above estimate for $g_4(T_{3,6n-1})$ is actually sharp.

Similarly, by Proposition 7.4, $\sigma'(T_{3,6n+1}) = 4n$ and $j_-(T_{3,6n+1}) = 2n$. So we obtain an estimate $g_4(T_{3,6n+1}) \geq 6n$. Once again, this estimate is sharp since $g_4(T_{3,6n+1}) = \frac{1}{2}(3 - 1)(6n + 1 - 1) = 6n$.

7.2 Nonextendable actions

Let Y be a rational homology 3–sphere equipped with an orientation-preserving action of G . Let W be a smooth 4–manifold with boundary Y . In this section we are concerned with the question of whether the G –action can be extended to W . In particular we are interested in finding obstructions to such an extension.

Proposition 7.6 *Let Y be an integral homology 3–sphere and \mathfrak{s} the unique spin^c –structure on Y . Let G act orientation-preservingly on Y and suppose the extension $G_{\mathfrak{s}}$ is trivial. Suppose that Y is the boundary of a contractible 4–manifold W . If the action of G extends over W then $\delta_{G,c}(Y, \mathfrak{s}) = \delta_{G,c}(\bar{Y}, \mathfrak{s}) = 0$ for every nonzero $c \in H_G^*$.*

Proof Suppose that the G -action extends to W . Since W is contractible, there is a unique spin^c -structure \mathfrak{t} on W . By uniqueness it is G -invariant and $\mathfrak{t}|_Y = \mathfrak{s}$. Theorem 4.1 gives $\delta_{G,c}(Y, \mathfrak{s}) \leq 0$ and $\delta_{G,c}(Y, \mathfrak{s}) \geq 0$; hence $\delta_{G,c}(Y, \mathfrak{s}) = 0$. Reversing orientations, we also find that $\delta_{G,c}(\bar{Y}, \mathfrak{s}) = 0$. \square

Example 7.7 Akbulut–Kirby constructed examples of contractible 4-manifolds bounding integral homology spheres, in particular $\Sigma(2, 5, 7)$, $\Sigma(3, 4, 5)$ and $\Sigma(2, 3, 13)$ bound contractible 4-manifolds [2, Theorem 2]. Further examples were given by Casson and Harer, in particular $\Sigma(2, 2s - 1, 2s + 1)$ for odd s bounds a contractible 4-manifold [14].

Now let $Y = \Sigma(2, 3, 13)$ and let τ be the involution obtained by viewing Y as the branched double cover $\Sigma_2(T_{3,13})$. Then $\delta_2(\bar{Y}) = -1$ by Proposition 7.4. Then it follows from Proposition 7.6 that τ does not extend to an involution on any contractible 4-manifold W bounded by Y . On the other hand, since τ is isotopic to the identity, τ does extend to a diffeomorphism on W .

Similarly if we let $Y = \Sigma(2, 2s - 1, 2s + 1) = \Sigma_2(T_{2s-1, 2s+1})$, where s is odd, and let τ be the covering involution, then Y bounds a contractible 4-manifold W but τ does not extend to an involution on W because $\delta_j(Y) = -\frac{1}{8}\sigma(T_{2s-1, 2s+1}) = \frac{1}{2}(s^2 - 1) \neq 0$ for all $j \geq 0$.

More generally, let $Y = \Sigma(p, q, r)$ where p, q and r are pairwise coprime positive integers. Assume that p is prime and let $\mathbb{Z}_p = \langle \tau \rangle$ act on Y by realising Y as the p -fold cyclic branched cover $\Sigma_p(T_{q,r})$. Then $\delta_0(Y, \tau) = -\lambda(\Sigma(p, q, r))$. From [59, Chapter 19], it can be seen that $\lambda(\Sigma(p, q, r)) < 0$ and hence the \mathbb{Z}_p -action on $Y = \Sigma(p, q, r)$ is nonextendable over contractible 4-manifolds bounded by Y . We have thus recovered a special case of the nonextendability results of Anvari and Hambleton [6; 7].

If we relax the condition that W is contractible to being a rational homology 4-ball, then we get a similar result, except that we have to make an assumption on the order of $H^2(W; \mathbb{Z})$.

Proposition 7.8 *Let Y be an integral homology 3-sphere and \mathfrak{s} the unique spin^c -structure on Y . Let $G = \mathbb{Z}_p$ for a prime p act orientation-preservingly on Y , and suppose that the extension $G_{\mathfrak{s}}$ is trivial. Suppose that Y is the boundary of a compact, oriented, smooth rational homology 4-ball W and assume that p does not divide the order of $H^2(W; \mathbb{Z})$. If the action of G extends over W then $\delta_{G,c}(Y, \mathfrak{s}) = \delta_{G,c}(\bar{Y}, \mathfrak{s}) = 0$ for every nonzero $c \in H_G^*$.*

Proof The set of spin^c -structures on W has cardinality $|H^2(W; \mathbb{Z})|$ and $G = \mathbb{Z}_p$ acts on this set. By assumption, p does not divide this number and hence there must exist a spin^c -structure \mathfrak{t} whose stabiliser group is not trivial. Since p is prime, this means \mathfrak{t} is fixed by all of G . From here, the rest of the proof is the same as for Proposition 7.6. \square

Example 7.9 Let $Y = \Sigma(p, q, r)$ where p, q and r are relatively prime and assume that p is prime. Let \mathbb{Z}_p act on Y as described in Section 7.1. Recall from Proposition 7.2 that $\delta_{\mathbb{Z}_p, 1}(Y, \mathfrak{s}) = -\lambda(\Sigma(p, q, r))$. As in Example 7.7, $\lambda(\Sigma(p, q, r)) < 0$ and hence $\delta_{\mathbb{Z}_p, 1}(Y, \mathfrak{s}) > 0$.

Therefore, by Proposition 7.8, if W is a compact, oriented, smooth rational homology 4–ball bounded by Y and if p does not divide the order of $H^2(W; \mathbb{Z})$, then the action of G does not extend over W . Thus we have obtained a partial extension of the results of Anvari–Hambleton to the case of rational homology 4–balls.

Fintushel and Stern [26] showed that $\Sigma(2, 3, 7)$ bounds a rational homology 4–ball, although it does not bound an integral 4–ball. Akbulut and Larson [4] showed that, for n odd, $\Sigma(2, 4n + 1, 12n + 5)$ and $\Sigma(3, 3n + 1, 12n + 5)$ bound rational 4–balls but not integral 4–balls. More examples, $\Sigma(2, 4n + 3, 12n + 7)$ and $\Sigma(3, 3n + 2, 12n + 7)$ for even n , were constructed by Şavk [21]. Taking $p = 2$ or 3 , the above Brieskorn spheres admit \mathbb{Z}_p –actions with nonzero delta invariants, as in Example 7.7. Hence the \mathbb{Z}_p –action does not extend to any oriented rational homology 4–ball W with boundary Y , provided the order of $H^2(W; \mathbb{Z})$ is coprime to p . However, it does not seem straightforward to determine whether the above examples are bounded by rational 4–balls satisfying this coprimality condition.

Proposition 7.10 *Let Y be an integral homology 3–sphere and \mathfrak{s} the unique spin^c –structure on Y . Let G act orientation-preservingly on Y and suppose that the extension $G_{\mathfrak{s}}$ is trivial. Suppose that Y is the boundary of a smooth, compact, oriented 4–manifold with $b_1(W) = 0$ and suppose that $H^2(W; \mathbb{Z})$ has no 2–torsion.*

- *If $H^2(W; \mathbb{R})$ is positive definite and $\delta_{G,1}(Y, \mathfrak{s}) > 0$, then the G –action on Y cannot be extended to a smooth G –action on W acting trivially on $H^2(W; \mathbb{Z})$.*
- *If $H^2(W; \mathbb{R})$ is negative definite and $\delta_{G,c}(Y, \mathfrak{s}) < 0$ for some $c \in H_G^*$, then the G –action on Y cannot be extended to a smooth G –action on W acting trivially on $H^2(W; \mathbb{Z})$.*

Proof Suppose the G –action on Y extends to a smooth G –action on W acting trivially on $H^2(W; \mathbb{Z})$. Since $H^2(W; \mathbb{Z})$ has no 2–torsion, a spin^c –structure \mathfrak{t} on W is determined uniquely by $c_1(\mathfrak{t})$. Since G acts trivially on $H^2(W; \mathbb{Z})$, it follows that G preserves every spin^c –structure. Furthermore, $\mathfrak{t}|_Y = \mathfrak{s}$ for any spin^c –structure on W by uniqueness of \mathfrak{t} .

If $H^2(W; \mathbb{R})$ is negative definite, then Theorem 4.1 may be applied to any spin^c –structure \mathfrak{t} on W , giving

$$\delta(W, \mathfrak{t}) \leq \delta_{G,c}(Y, \mathfrak{s})$$

for all \mathfrak{t} and all $c \in H_G^*$. Since Y is an integral homology sphere, the intersection form on the $H^2(W; \mathbb{Z})/\text{torsion}$ is unimodular. By the main theorem of [24], there exists a spin^c –structure \mathfrak{t} such that $\delta(W, \mathfrak{t}) \geq 0$. Hence $\delta_{G,c}(Y, \mathfrak{s}) \geq 0$. The proof in the case that $H^2(W; \mathbb{R})$ is positive definite is similarly obtained. \square

Example 7.11 Consider again $Y = \Sigma(p, q, r)$ with the same \mathbb{Z}_p –action. Recall from Proposition 7.2 that $\delta_{\mathbb{Z}_p,1}(Y, \mathfrak{s}) = -\lambda(\Sigma(p, q, r))$. As in Example 7.9, $\delta_{\mathbb{Z}_p,1}(Y, \mathfrak{s}) > 0$. So by Proposition 7.10, the action of \mathbb{Z}_p on Y cannot be extended to any smooth, compact, oriented 4–manifold W such that $b_1(W) = 0$, $H^2(W; \mathbb{Z})$ has no 2–torsion and with \mathbb{Z}_p acting trivially on $H^2(W; \mathbb{Z})$.

7.3 Realisation problems

In this section we are concerned with the following realisation problem. Let W be a smooth 4–manifold with boundary an integral homology sphere Y . Suppose that a finite group G acts on $H^2(W; \mathbb{Z})$ preserving the intersection form. We say that the action of G on $H^2(W; \mathbb{Z})$ can be realised by diffeomorphisms if there is a smooth orientation-preserving action of G on W inducing the given action on $H^2(W; \mathbb{Z})$.

For simplicity we will assume that $G = \mathbb{Z}_p$ for a prime p so that all extensions $G_{\mathfrak{s}}$ are trivial.

Proposition 7.12 *Let W be a smooth, compact, oriented 4–manifold with $b_1(W) = 0$ and with boundary $Y = \partial W$ an L –space integral homology sphere. Suppose that an action of $G = \mathbb{Z}_p$ on $H^2(W; \mathbb{Z})$ is given and suppose that $H^2(W; \mathbb{Z})$ has no 2–torsion. Suppose that the subspace of $H^2(W; \mathbb{R})$ fixed by G is negative definite. If the action of G on $H^2(W; \mathbb{Z})$ can be realised by diffeomorphisms, then*

$$\delta(W, \mathfrak{s}) \leq \delta(Y, \mathfrak{s}|_Y)$$

for every spin^c –structure \mathfrak{s} on W for which $c_1(\mathfrak{s})$ is invariant.

Proof This is essentially a special case of Theorem 5.3. Note that since $H^2(W; \mathbb{Z})$ is assumed to have no 2–torsion, any spin^c –structure \mathfrak{s} for which $c_1(\mathfrak{s})$ is invariant is preserved by G . So if G is realisable by diffeomorphisms, then Theorem 5.3 gives $\delta(W, \mathfrak{s}) \leq \delta_{G,1}(Y, \mathfrak{s}|_Y)$. But we have assumed that Y is an L –space, so $\delta_{G,1}(Y, \mathfrak{s}|_Y) = \delta(Y, \mathfrak{s}|_Y)$. \square

Example 7.13 We consider a specialisation of Proposition 7.12 as follows. Take $G = \mathbb{Z}_p$. Assume Y is an L –space integral homology 3–sphere and let \mathfrak{s} be the unique spin^c –structure. Suppose that W is a smooth, compact, oriented 4–manifold with $b_1(W) = 0$ and with boundary Y . Suppose that the intersection form on $H^2(W; \mathbb{Z})$ is even and that $H^2(W; \mathbb{Z})$ has no 2–torsion. Then W is spin and it has a unique spin–structure \mathfrak{t} . By uniqueness, the restriction of \mathfrak{t} to the boundary equals \mathfrak{s} . Suppose that an action of $G = \mathbb{Z}_p$ on $H^2(W; \mathbb{Z})$ is given and that the subspace of $H^2(W; \mathbb{R})$ fixed by G is negative definite. Then applying Proposition 7.12 to (W, \mathfrak{t}) , we find that $\delta(W, \mathfrak{t}) = -\frac{1}{8}\sigma(W) \leq \delta(Y, \mathfrak{s})$. Therefore, if $\frac{1}{8}\sigma(W) < -\delta(Y, \mathfrak{s})$ then the action of \mathbb{Z}_p on $H^2(W; \mathbb{Z})$ is not realisable by a smooth \mathbb{Z}_p –action on W . For example, if $W = K3 \# W_0$ is the connected sum of a $K3$ surface with W_0 , the negative definite plumbing of the E_8 graph, then $\partial W = Y = \Sigma(2, 3, 5)$ is the Poincaré homology 3–sphere which is an L –space. Then W satisfies all the above conditions and $\frac{1}{8}\sigma(W) = -3 < \delta(Y, \mathfrak{s}) = -1$. Hence for any prime p , any \mathbb{Z}_p –action on $H^2(W; \mathbb{Z})$ such that the invariant subspace of $H^2(W; \mathbb{R})$ is negative definite cannot be realised by a smooth \mathbb{Z}_p –action on W .

Corollary 7.14 *Let W be a smooth, compact, oriented 4–manifold with $b_1(W) = 0$ and with boundary $Y = \partial W$ an L –space integral homology sphere. Suppose that W is spin and that $H^2(W; \mathbb{Z})$ has no 2–torsion. If there is a smooth involution on W which acts as -1 on $H^2(W; \mathbb{R})$, then $\delta(Y, \mathfrak{s}) = -\frac{1}{8}\sigma(W)$.*

Proof Since W is spin, there is a spin^c –structure \mathfrak{s} for which $c_1(\mathfrak{s}) = 0$. Proposition 7.12 then implies that $-\frac{1}{8}\sigma(W) \leq \delta(Y, \mathfrak{s})$. The same argument applied to \overline{W} gives $\frac{1}{8}\sigma(W) \leq \delta(Y, \mathfrak{s})$. \square

7.4 Equivariant embeddings of 3–manifolds in 4–manifolds

Let Y be a rational homology 3–sphere equipped with an orientation-preserving action of G . By an equivariant embedding of Y into a 4–manifold X , we mean an embedding $Y \rightarrow X$ such that the action of G on Y extends over X . We consider some existence and nonexistence results for equivariant embeddings.

Proposition 7.15 *Suppose that Y is an integral homology 3–sphere. Let \mathfrak{s} be the unique spin^c –structure on Y and assume that $G_{\mathfrak{s}}$ is a trivial extension. If Y can be equivariantly embedded in S^4 , then $\delta_{G,c}(Y, \mathfrak{s}) = \delta_{G,c}(\bar{Y}, \mathfrak{s}) = 0$ for every nonzero $c \in H_G^*$.*

Proof If Y embeds equivariantly in S^4 , then we obtain an equivariant decomposition $S^4 = W_+ \cup_Y W_-$. Mayer–Vietoris and Poincaré–Lefschetz imply that W_{\pm} are integral homology 4–balls, hence are contractible by Whitehead’s theorem. The result now follows from Proposition 7.6. \square

Example 7.16 Let $Y = \Sigma(2, 3, 13)$, equipped with the involution τ obtained from viewing Y as the branched double cover $\Sigma_2(T_{3,13})$. Then Y embeds in S^4 [13, Theorem 2.13]. On the other hand, $\delta_2(\bar{Y}, \mathfrak{s}) = -1$ by Proposition 7.4. Hence Y cannot be equivariantly embedded in S^4 .

It is known that every 3–manifold Y embeds in the connected sum $\#^n(S^2 \times S^2)$ of n copies of $S^2 \times S^2$ for some sufficiently large n [1, Theorem 2.1]. Aceto, Golla and Larson define the *embedding number* $\varepsilon(Y)$ of Y to be the smallest n for which Y embeds in $\#^n(S^2 \times S^2)$. Here we consider an equivariant version of the embedding number. To obtain interesting results we need to make an assumption on the kinds of group actions allowed.

Definition 7.17 Let $G = \mathbb{Z}_p = \langle \tau \rangle$, where p is a prime number. We say that a smooth, orientation-preserving action of G on $X = \#^n(S^2 \times S^2)$ is *admissible* if $H^2(X; \mathbb{Z})^{\tau} = 0$, where

$$H^2(X; \mathbb{Z})^{\tau} = \{x \in H^2(X; \mathbb{Z}) \mid \tau(x) = x\}.$$

One way of constructing admissible actions is as follows. Let X be the p –fold cyclic cover of S^4 , branched over an unknotted embedded surface $\Sigma \subset S^4$ of genus g . Then X is diffeomorphic to $\#^{g(p-1)}(S^2 \times S^2)$ [3, Corollary 4.3] and the action of \mathbb{Z}_p on X as a cyclic branched cover is admissible — as can be seen from the proof of Theorem 9.3 in [11].

Let $\mathbb{Z}_p = \langle \tau \rangle$ act on a rational homology 3–sphere Y . We define the *equivariant embedding number* $\varepsilon(Y, \tau)$ of (Y, τ) to be the smallest n for which Y embeds equivariantly in $\#^n(S^2 \times S^2)$ for some admissible \mathbb{Z}_p –action on $\#^n(S^2 \times S^2)$, if such an embedding exists. We set $\varepsilon(Y, \tau) = \infty$ if there is no such embedding.

Recall that the *double slice genus* [48, Section 5] $g_{\text{ds}}(K)$ of a knot K in S^3 is defined as the minimal genus of an unknotted compact oriented surface S embedded in S^4 whose intersection with the equator S^3 is K . From the definition, it follows that $2g_4(K) \leq g_{\text{ds}}(K) \leq 2g_3(K)$, where $g_3(K)$ is the 3–genus of K .

Proposition 7.18 *Let $Y = \Sigma_2(K)$ be the branched double cover of a knot K and let τ be the covering involution on Y . Then $\varepsilon(Y, \tau) \leq g_{\text{ds}}(K)$.*

Proof Let S be an unknotted embedded surface in S^4 of genus $g_{\text{ds}}(K)$ intersecting the equator in K . Let W be the double cover of S^4 branched along S . Then by [3, Corollary 4.3], W is diffeomorphic to $\#^{g_{\text{ds}}(K)}(S^2 \times S^2)$. The covering involution on W is admissible since W is a branched double cover of S^4 . Clearly $Y = \Sigma_2(K) = \partial W$ embeds equivariantly in W and so $\varepsilon(Y, \tau) \leq g_{\text{ds}}(K)$. \square

Proposition 7.19 *Let $\mathbb{Z}_p = \langle \tau \rangle$ act orientation-preservingly on an integral homology 3–sphere Y . Let $j(Y, \tau)$ be the smallest positive integer such that $\delta_j(Y, \mathfrak{s}, \tau, p) + \delta_j(\bar{Y}, \mathfrak{s}, \tau, p) = 0$, or $j(Y, \tau) = \infty$ if no such j exists. Here \mathfrak{s} is the unique spin^c–structure on Y . Then $\varepsilon(Y, \tau) \geq j(Y, \tau)$ if $p = 2$ and $\varepsilon(Y, \tau) \geq 2j(Y, \tau)$ if p is odd.*

Proof To simplify notation we write $\delta_j(Y)$ for $\delta_j(Y, \mathfrak{s}, \tau, p)$. Suppose that Y embeds equivariantly in $X = \#^n(S^2 \times S^2)$, for an admissible action of τ . Then we obtain an equivariant splitting $X = X_+ \cup_Y X_-$. Let \mathfrak{t} be the unique spin–structure on X . By uniqueness, \mathfrak{t} is τ –invariant and $\mathfrak{t}|_Y = \mathfrak{s}$. Corollary 5.5 applied to X_+ gives $\delta_j(Y) = -\frac{1}{8}\sigma(X_+)$ for $j \geq b'_-(X_+)$ and $\delta_j(\bar{Y}) = \frac{1}{8}\sigma(X_+)$ for $j \geq b'_+(X_+)$. Let $n' = n$ if $p = 2$ or $\frac{1}{2}n$ if p is odd. Since $b'_\pm(X_+) \leq b'_\pm(X) = n'$, we see that $\delta_j(Y) + \delta_j(\bar{Y}) = 0$ for $j \geq n'$. Hence $j(Y, \tau) \leq n'$. Therefore $\varepsilon(Y, \tau) \geq j(Y, \tau)$ if $p = 2$ and $\varepsilon(Y, \tau) \geq 2j(Y, \tau)$ if p is odd. \square

Example 7.20 Let $Y = \Sigma(2, 3, 6n + 1) = \Sigma_2(T_{3, 6n+1})$ and equip Y with the covering involution τ . Then $g_3(T_{3, 6n+1}) = 6n$; hence $\varepsilon(Y, \tau) \leq 12n$, by Proposition 7.18. By Proposition 7.4, we see that $j(Y, \tau) = 2n$ and so $\varepsilon(Y, \tau) \geq 2n$. So we have an estimate on the equivariant embedding number of the form

$$2n \leq \varepsilon(\Sigma(2, 3, 6n + 1), \tau) \leq 12n.$$

Suppose that n is odd. Then by [1, Proposition 3.5], the (nonequivariant) embedding number of $\Sigma(2, 3, 6n+1)$ is given by $\varepsilon(\Sigma(2, 3, 6n+1)) = 10$. In particular, $\varepsilon(\Sigma(2, 3, 6n+1), \tau) > \varepsilon(\Sigma(2, 3, 6n+1))$ for all odd $n > 5$. Also, since we obviously have $\varepsilon(Y, \tau) \geq \varepsilon(Y)$, we see that

$$10 \leq \varepsilon(\Sigma(2, 3, 7), \tau) \leq 12.$$

In fact, we will now prove that $\varepsilon(\Sigma(2, 3, 7), \tau) = 12$. Suppose that $Y = \Sigma(2, 3, 7)$ embeds equivariantly in $X = \#^n(S^2 \times S^2)$ for some admissible involution, where $n \leq 12$. Then we obtain an equivariant splitting $X = X_+ \cup_Y X_-$. Since Y is an integral homology sphere, the intersection forms on X_\pm are unimodular. They are also even, since X is spin. Moreover the Rochlin invariant of Y is 1. So the intersection forms of X_\pm must contain at least one E_8 or $-E_8$ summand. Proposition 7.2 implies that $\delta_j(Y) = 1$ for all $j \geq 0$ and Proposition 7.4 implies that $\delta_j(\bar{Y}) = 0$ for $j = 0, 1$ and $\delta_j(\bar{Y}) = -1$ for $j \geq 2$. Since $n \leq 12$, Corollary 5.5 applied to X_\pm then implies that the intersection form of X_+ must be of the form $\alpha H \oplus (-E_8)$ for some $\alpha \geq 2$ (where H is the hyperbolic lattice) and similarly the intersection form

of X_- must be of the form $\alpha'H \oplus (E_8)$ for some $\alpha' \geq 2$. The intersection form of X is then $(\alpha + \alpha' + 8)H$ and so $n = \alpha + \alpha' + 8 \geq 2 + 2 + 8 = 12$. This proves that

$$\varepsilon(\Sigma(2, 3, 7), \tau) = 12.$$

Example 7.21 Let $Y = \Sigma(2, 3, 5) = \Sigma_2(T_{3,5})$ and equip Y with the covering involution τ . Then $g_3(T_{3,5}) = 4$; hence $\varepsilon(Y, \tau) \leq 8$, by Proposition 7.18. On the other hand, $\varepsilon(Y, \tau) \geq \varepsilon(Y)$ and $\varepsilon(Y) = 8$ by [1, Proposition 3.4], so $\varepsilon(\Sigma(2, 3, 5), \tau) = 8$.

References

- [1] **P Aceto, M Golla, K Larson**, *Embedding 3–manifolds in spin 4–manifolds*, J. Topol. 10 (2017) 301–323 MR Zbl
- [2] **S Akbulut, R Kirby**, *Mazur manifolds*, Michigan Math. J. 26 (1979) 259–284 MR Zbl
- [3] **S Akbulut, R Kirby**, *Branched covers of surfaces in 4–manifolds*, Math. Ann. 252 (1979/80) 111–131 MR Zbl
- [4] **S Akbulut, K Larson**, *Brieskorn spheres bounding rational balls*, Proc. Amer. Math. Soc. 146 (2018) 1817–1824 MR Zbl
- [5] **A Alfieri, S Kang, A I Stipsicz**, *Connected Floer homology of covering involutions*, Math. Ann. 377 (2020) 1427–1452 MR Zbl
- [6] **N Anvari, I Hambleton**, *Cyclic group actions on contractible 4–manifolds*, Geom. Topol. 20 (2016) 1127–1155 MR Zbl
- [7] **N Anvari, I Hambleton**, *Cyclic branched coverings of Brieskorn spheres bounding acyclic 4–manifolds*, Glasg. Math. J. 63 (2021) 400–413 MR Zbl
- [8] **M F Atiyah, V K Patodi, I M Singer**, *Spectral asymmetry and Riemannian geometry, I*, Math. Proc. Cambridge Philos. Soc. 77 (1975) 43–69 MR Zbl
- [9] **M F Atiyah, V K Patodi, I M Singer**, *Spectral asymmetry and Riemannian geometry, III*, Math. Proc. Cambridge Philos. Soc. 79 (1976) 71–99 MR Zbl
- [10] **D Baraglia**, *Obstructions to smooth group actions on 4–manifolds from families Seiberg–Witten theory*, Adv. Math. 354 (2019) art. id. 106730 MR Zbl
- [11] **D Baraglia**, *Constraints on families of smooth 4–manifolds from Bauer–Furuta invariants*, Algebr. Geom. Topol. 21 (2021) 317–349 MR Zbl
- [12] **K S Brown**, *Cohomology of groups*, Graduate Texts in Math. 87, Springer (1982) MR Zbl
- [13] **R Budney, B A Burton**, *Embeddings of 3–manifolds in S^4 from the point of view of the 11–tetrahedron census*, Exp. Math. 31 (2022) 988–1013 MR Zbl
- [14] **A J Casson, J L Harer**, *Some homology lens spaces which bound rational homology balls*, Pacific J. Math. 96 (1981) 23–36 MR Zbl
- [15] **V Colin, P Ghiggini, K Honda**, *The equivalence of Heegaard Floer homology and embedded contact homology, III: From hat to plus*, preprint (2012) arXiv 1208.1526

- [16] **V Colin, P Ghiggini, K Honda**, *The equivalence of Heegaard Floer homology and embedded contact homology via open book decompositions, I*, preprint (2012) arXiv 1208.1074
- [17] **V Colin, P Ghiggini, K Honda**, *The equivalence of Heegaard Floer homology and embedded contact homology via open book decompositions, II*, preprint (2012) arXiv 1208.1077
- [18] **O Collin, N Saveliev**, *A geometric proof of the Fintushel–Stern formula*, Adv. Math. 147 (1999) 304–314 MR Zbl
- [19] **O Cornea**, *Homotopical dynamics: Suspension and duality*, Ergodic Theory Dynam. Systems 20 (2000) 379–391 MR Zbl
- [20] **D Cristofaro-Gardiner**, *The absolute gradings on embedded contact homology and Seiberg–Witten Floer cohomology*, Algebr. Geom. Topol. 13 (2013) 2239–2260 MR Zbl
- [21] **O Şavk**, *More Brieskorn spheres bounding rational balls*, Topology Appl. 286 (2020) art. id. 107400 MR Zbl
- [22] **I Dai, M Hedden, A Mallick**, *Corks, involutions, and Heegaard Floer homology*, preprint (2020) arXiv 2002.02326v2
- [23] **T tom Dieck**, *Transformation groups*, De Gruyter Studies in Mathematics 8, de Gruyter, Berlin (1987) MR Zbl
- [24] **N D Elkies**, *A characterization of the \mathbb{Z}^n lattice*, Math. Res. Lett. 2 (1995) 321–326 MR Zbl
- [25] **H Fang**, *Equivariant spectral flow and a Lefschetz theorem on odd-dimensional Spin manifolds*, Pacific J. Math. 220 (2005) 299–312 MR Zbl
- [26] **R Fintushel, R J Stern**, *A μ -invariant one homology 3-sphere that bounds an orientable rational ball*, from “Four-manifold theory” (C Gordon, editor), Contemp. Math. 35, Amer. Math. Soc., Providence, RI (1984) 265–268 MR Zbl
- [27] **R Fintushel, R J Stern**, *Instanton homology of Seifert fibred homology three spheres*, Proc. London Math. Soc. 61 (1990) 109–137 MR Zbl
- [28] **K A Frøyshov**, *Monopole Floer homology for rational homology 3-spheres*, Duke Math. J. 155 (2010) 519–576 MR Zbl
- [29] **J E Grigsby, D Ruberman, S Strle**, *Knot concordance and Heegaard Floer homology invariants in branched covers*, Geom. Topol. 12 (2008) 2249–2275 MR Zbl
- [30] **K Hendricks, R Lipshitz, S Sarkar**, *A flexible construction of equivariant Floer homology and applications*, J. Topol. 9 (2016) 1153–1236 MR Zbl
- [31] **K Hendricks, C Manolescu**, *Involutive Heegaard Floer homology*, Duke Math. J. 166 (2017) 1211–1299 MR Zbl
- [32] **K Hendricks, C Manolescu, I Zemke**, *A connected sum formula for involutive Heegaard Floer homology*, Selecta Math. 24 (2018) 1183–1245 MR Zbl
- [33] **Y Huang, V G B Ramos**, *An absolute grading on Heegaard Floer homology by homotopy classes of oriented 2-plane fields*, J. Symplectic Geom. 15 (2017) 51–90 MR Zbl
- [34] **M Kankaanrinta**, *Equivariant collaring, tubular neighbourhood and gluing theorems for proper Lie group actions*, Algebr. Geom. Topol. 7 (2007) 1–27 MR Zbl
- [35] **L H Kauffman**, *Branched coverings, open books and knot periodicity*, Topology 13 (1974) 143–160 MR Zbl

- [36] **L H Kauffman, L R Taylor**, *Signature of links*, Trans. Amer. Math. Soc. 216 (1976) 351–365 MR Zbl
- [37] **T Khandhawit**, *A new gauge slice for the relative Bauer–Furuta invariants*, Geom. Topol. 19 (2015) 1631–1655 MR Zbl
- [38] **P B Kronheimer, T S Mrowka**, *Gauge theory for embedded surfaces, I*, Topology 32 (1993) 773–826 MR Zbl
- [39] **P Kronheimer, T Mrowka**, *Monopoles and three-manifolds*, New Mathematical Monographs 10, Cambridge Univ. Press (2007) MR Zbl
- [40] **Ç Kutluhan, Y-J Lee, C H Taubes**, *HF = HM, I: Heegaard Floer homology and Seiberg–Witten Floer homology*, Geom. Topol. 24 (2020) 2829–2854 MR Zbl
- [41] **Ç Kutluhan, Y-J Lee, C H Taubes**, *HF = HM, II: Reeb orbits and holomorphic curves for the ech/Heegaard Floer correspondence*, Geom. Topol. 24 (2020) 2855–3012 MR Zbl
- [42] **Ç Kutluhan, Y-J Lee, C H Taubes**, *HF = HM, III: Holomorphic curves and the differential for the ech/Heegaard Floer correspondence*, Geom. Topol. 24 (2020) 3013–3218 MR Zbl
- [43] **Ç Kutluhan, Y-J Lee, C Taubes**, *HF = HM, IV: The Seiberg–Witten Floer homology and ech correspondence*, Geom. Topol. 24 (2020) 3219–3469 MR Zbl
- [44] **Ç Kutluhan, Y-J Lee, C H Taubes**, *HF = HM, V: Seiberg–Witten Floer homology and handle additions*, Geom. Topol. 24 (2020) 3471–3748 MR Zbl
- [45] **T Lidman, C Manolescu**, *The equivalence of two Seiberg–Witten Floer homologies*, Astérisque 399, Soc. Math. France, Paris (2018) MR Zbl
- [46] **T Lidman, C Manolescu**, *Floer homology and covering spaces*, Geom. Topol. 22 (2018) 2817–2838 MR Zbl
- [47] **P Lisca, B Owens**, *Signatures, Heegaard Floer correction terms and quasi-alternating links*, Proc. Amer. Math. Soc. 143 (2015) 907–914 MR Zbl
- [48] **C Livingston, J Meier**, *Doubly slice knots with low crossing number*, New York J. Math. 21 (2015) 1007–1026 MR Zbl
- [49] **C Manolescu**, *Seiberg–Witten–Floer stable homotopy type of three-manifolds with $b_1 = 0$* , Geom. Topol. 7 (2003) 889–932 MR Zbl
- [50] **C Manolescu**, *Pin(2)–equivariant Seiberg–Witten Floer homology and the triangulation conjecture*, J. Amer. Math. Soc. 29 (2016) 147–176 MR Zbl
- [51] **C Manolescu, B Owens**, *A concordance invariant from the Floer homology of double branched covers*, Int. Math. Res. Not. 2007 (2007) art. id. rnm077 MR Zbl
- [52] **K Murasugi**, *On a certain numerical invariant of link types*, Trans. Amer. Math. Soc. 117 (1965) 387–422 MR Zbl
- [53] **W D Neumann, F Raymond**, *Seifert manifolds, plumbing, μ –invariant and orientation reversing maps*, from “Algebraic and geometric topology” (K C Millett, editor), Lecture Notes in Math. 664, Springer (1978) 163–196 MR Zbl
- [54] **P Ozsváth, Z Szabó**, *Absolutely graded Floer homologies and intersection forms for four-manifolds with boundary*, Adv. Math. 173 (2003) 179–261 MR Zbl
- [55] **P Ozsváth, Z Szabó**, *On the Floer homology of plumbed three-manifolds*, Geom. Topol. 7 (2003) 185–224 MR Zbl

- [56] **P Ozsváth, Z Szabó**, *On knot Floer homology and lens space surgeries*, *Topology* 44 (2005) 1281–1300 MR Zbl
- [57] **P Ozsváth, Z Szabó**, *On the Heegaard Floer homology of branched double-covers*, *Adv. Math.* 194 (2005) 1–33 MR Zbl
- [58] **V G B Ramos**, *Absolute gradings on ECH and Heegaard Floer homology*, *Quantum Topol.* 9 (2018) 207–228 MR Zbl
- [59] **N Saveliev**, *Lectures on the topology of 3-manifolds: an introduction to the Casson invariant*, de Gruyter, Berlin (1999) MR Zbl
- [60] **C H Taubes**, *Embedded contact homology and Seiberg–Witten Floer cohomology, I*, *Geom. Topol.* 14 (2010) 2497–2581 MR Zbl
- [61] **C H Taubes**, *Embedded contact homology and Seiberg–Witten Floer cohomology, II*, *Geom. Topol.* 14 (2010) 2583–2720 MR Zbl
- [62] **C H Taubes**, *Embedded contact homology and Seiberg–Witten Floer cohomology, III*, *Geom. Topol.* 14 (2010) 2721–2817 MR Zbl
- [63] **C H Taubes**, *Embedded contact homology and Seiberg–Witten Floer cohomology, IV*, *Geom. Topol.* 14 (2010) 2819–2960 MR Zbl
- [64] **C H Taubes**, *Embedded contact homology and Seiberg–Witten Floer cohomology, V*, *Geom. Topol.* 14 (2010) 2961–3000 MR Zbl

*School of Mathematical Sciences, The University of Adelaide
Adelaide, SA, Australia*

*Department of Mathematics, The University of Auckland
Auckland, New Zealand*

david.baraglia@adelaide.edu.au, p.hekmati@auckland.ac.nz

Received: 13 January 2022 Revised: 27 July 2022

ALGEBRAIC & GEOMETRIC TOPOLOGY

msp.org/agt

EDITORS

PRINCIPAL ACADEMIC EDITORS

John Etnyre
etnyre@math.gatech.edu
Georgia Institute of Technology

Kathryn Hess
kathryn.hess@epfl.ch
École Polytechnique Fédérale de Lausanne

BOARD OF EDITORS

Julie Bergner	University of Virginia jeb2md@eservices.virginia.edu	Robert Lipshitz	University of Oregon lipshitz@uoregon.edu
Steven Boyer	Université du Québec à Montréal cohf@math.rochester.edu	Norihiko Minami	Nagoya Institute of Technology nori@nitech.ac.jp
Tara E Brendle	University of Glasgow tara.brendle@glasgow.ac.uk	Andrés Navas	Universidad de Santiago de Chile andres.navas@usach.cl
Indira Chatterji	CNRS & Univ. Côte d'Azur (Nice) indira.chatterji@math.cnrs.fr	Thomas Nikolaus	University of Münster nikolaus@uni-muenster.de
Alexander Dranishnikov	University of Florida dranish@math.ufl.edu	Robert Oliver	Université Paris 13 bobol@math.univ-paris13.fr
Tobias Ekholm	Uppsala University, Sweden tobias.ekholm@math.uu.se	Jessica S Purcell	Monash University jessica.purcell@monash.edu
Mario Eudave-Muñoz	Univ. Nacional Autónoma de México mario@matem.unam.mx	Birgit Richter	Universität Hamburg birgit.richter@uni-hamburg.de
David Futer	Temple University dfuter@temple.edu	Jérôme Scherer	École Polytech. Féd. de Lausanne jerome.scherer@epfl.ch
John Greenlees	University of Warwick john.greenlees@warwick.ac.uk	Vesna Stojanoska	Univ. of Illinois at Urbana-Champaign vesna@illinois.edu
Ian Hambleton	McMaster University ian@math.mcmaster.ca	Zoltán Szabó	Princeton University szabo@math.princeton.edu
Matthew Hedden	Michigan State University mhedden@math.msu.edu	Maggy Tomova	University of Iowa maggy-tomova@uiowa.edu
Hans-Werner Henn	Université Louis Pasteur henn@math.u-strasbg.fr	Nathalie Wahl	University of Copenhagen wahl@math.ku.dk
Daniel Isaksen	Wayne State University isaksen@math.wayne.edu	Chris Wendl	Humboldt-Universität zu Berlin wendl@math.hu-berlin.de
Thomas Koberda	University of Virginia thomas.koberda@virginia.edu	Daniel T Wise	McGill University, Canada daniel.wise@mcgill.ca
Christine Lescop	Université Joseph Fourier lescop@ujf-grenoble.fr		


See inside back cover or msp.org/agt for submission instructions.

The subscription price for 2024 is US \$705/year for the electronic version, and \$1040/year (+\$70, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Algebraic & Geometric Topology is indexed by Mathematical Reviews, Zentralblatt MATH, Current Mathematical Publications and the Science Citation Index.

Algebraic & Geometric Topology (ISSN 1472-2747 printed, 1472-2739 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840.

AGT peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<https://msp.org/>

© 2024 Mathematical Sciences Publishers

ALGEBRAIC & GEOMETRIC TOPOLOGY

Volume 24 Issue 1 (pages 1–594) 2024

Chow–Witt rings of Grassmannians	1
MATTHIAS WENDT	
Higher chromatic Thom spectra via unstable homotopy theory	49
SANATH K DEVALAPURKAR	
The deformation space of nonorientable hyperbolic 3–manifolds	109
JUAN LUIS DURÁN BATALLA and JOAN PORTI	
Realization of Lie algebras and classifying spaces of crossed modules	141
YVES FÉLIX and DANIEL TANRÉ	
Knot Floer homology, link Floer homology and link detection	159
FRASER BINNS and GAGE MARTIN	
Models for knot spaces and Atiyah duality	183
SYUNJI MORIYA	
Automorphismes du groupe des automorphismes d’un groupe de Coxeter universel	251
YASSINE GUERCH	
The $RO(C_4)$ cohomology of the infinite real projective space	277
NICK GEORGAKOPOULOS	
Annular Khovanov homology and augmented links	325
HONGJIAN YANG	
Smith ideals of operadic algebras in monoidal model categories	341
DAVID WHITE and DONALD YAU	
The persistent topology of optimal transport based metric thickenings	393
HENRY ADAMS, FACUNDO MÉMOLI, MICHAEL MOY and QINGSONG WANG	
A generalization of moment-angle manifolds with noncontractible orbit spaces	449
LI YU	
Equivariant Seiberg–Witten–Floer cohomology	493
DAVID BARAGLIA and PEDRAM HEKMATI	
Constructions stemming from nonseparating planar graphs and their Colin de Verdière invariant	555
ANDREI PAVELESCU and ELENA PAVELESCU	
Census L –space knots are braid positive, except for one that is not	569
KENNETH L BAKER and MARC KEGEL	
Branched covers and rational homology balls	587
CHARLES LIVINGSTON	