

AG
T

*Algebraic & Geometric
Topology*

Volume 24 (2024)

Projective modules and the homotopy classification of (G, n) -complexes

JOHN NICHOLSON



Projective modules and the homotopy classification of (G, n) -complexes

JOHN NICHOLSON

A (G, n) -complex is an n -dimensional CW-complex with fundamental group G and whose universal cover is $(n-1)$ -connected. If G has periodic cohomology then, for appropriate n , we show that there is a one-to-one correspondence between the homotopy types of finite (G, n) -complexes and the orbits of the stable class of a certain projective $\mathbb{Z}G$ -module under the action of $\text{Aut}(G)$. We develop techniques to compute this action explicitly and use this to give an example where the action is nontrivial.

[55P15](#); [20C05](#), [55U15](#), [57K20](#)

1 Introduction

For a group G and $n \geq 2$, a (G, n) -complex is a connected n -dimensional CW-complex X for which $\pi_1(X) \cong G$ and \tilde{X} is $(n-1)$ -connected. Equivalently, it is the n -skeleton of a $K(G, 1)$. For example, a finite $(G, 2)$ -complex is equivalently a finite 2-complex X with $\pi_1(X) \cong G$. An example of a finite $(G, 3)$ -complex is a closed 3-manifold M with $\pi_1(M) \cong G$ finite. Given a group G and $n \geq 2$, a finite (G, n) -complex exists if and only if G has type F_n in the sense of Wall [1965].

Let $\text{HT}(G, n)$ be the set of homotopy types of finite (G, n) -complexes, which can be viewed as a graph with edges between each X and $X \vee S^n$. It is well known that $\text{HT}(G, n)$ is a tree [Whitehead 1939], i.e. a connected acyclic graph, and has a grading coming from $(-1)^n \chi(X)$ which takes a minimum value $\chi_{\min}(G, n)$. The problem of determining the structure of $\text{HT}(G, n)$ as a tree has a long history which dates back to Cockcroft and Swan [1961] and Dyer and Sieradski [1973; 1975].

In the case of finite abelian groups, the structure of $\text{HT}(G, n)$ has been classified through a series of articles by Metzler [1976], Sieradski and Dyer [1979], Browning [1979] and Linnell [1993]. However, much less is known for nonabelian groups and an important class of examples are the groups with k -periodic cohomology, i.e. finite groups for which the Tate cohomology groups satisfy $\hat{H}^i(G; \mathbb{Z}) \cong \hat{H}^{i+k}(G; \mathbb{Z})$ for all $i \in \mathbb{Z}$. For example, if G is finite and n is even, then it was shown by Browning [1978] that $\chi(X) = \chi(Y)$ implies $X \vee S^n \simeq Y \vee S^n$ (see also [Hambleton and Kreck 1993]). However, when n is odd, this is known only when G does not have k -periodic cohomology for $k \mid n + 1$ (see Question 7.4).

The aim of this article is to make new progress towards the classification over groups with periodic cohomology, building upon work of Dyer [1976] and Johnson [2003].

1.1 Main results

Let $\text{PHT}(G, n)$ be the tree of polarised homotopy types of finite (G, n) -complexes, i.e. the homotopy types of pairs (X, ρ) where $\rho: \pi_1(X) \cong G$.

Let G be a finite group and let $C(\mathbb{Z}G)$ denote the projective class group, i.e. the equivalence classes of finitely generated projective $\mathbb{Z}G$ -modules where $P \sim Q$ if $P \oplus \mathbb{Z}G^i \cong Q \oplus \mathbb{Z}G^j$ for some i and j . Note that a class $[P] \in C(\mathbb{Z}G)$ can be viewed as the set of (nonzero) projective $\mathbb{Z}G$ -modules P_0 for which $P_0 \sim P$, and this has the structure of a graded tree with edges between each P_0 and $P_0 \oplus \mathbb{Z}G$. Let $T_G \leq C(\mathbb{Z}G)$ denote the Swan subgroup (see [Section 3.2](#)). If G has k -periodic cohomology, then the *Swan finiteness obstruction* is an element $\sigma_k(G) \in C(\mathbb{Z}G)/T_G$ which vanishes if and only if there exists a finite CW-complex X with $\pi_1(X) \cong G$ and $\tilde{X} \simeq S^{k-1}$.

Recall that a finitely presented group G has the *D2 property* if every cohomologically 2-dimensional finite complex X with $\pi_1(X) \cong G$ is homotopy equivalent to a finite 2-complex.

Theorem A *Let G have k -periodic cohomology and let $n = ik$ or $ik - 2$ for some $i \geq 1$. Then there is an injective map of graded trees*

$$\Psi: \text{PHT}(G, n) \rightarrow [P_{(G,n)}]$$

for any projective $\mathbb{Z}G$ -module $P_{(G,n)}$ with $\sigma_{ik}(G) = [P_{(G,n)}] \in C(\mathbb{Z}G)/T_G$. Furthermore, Ψ is bijective if and only if $n \geq 3$ or if $n = 2$ and G has the D2 property.

Remark 1.1 (a) If G satisfies the Eichler condition, then $[P_{(G,n)}]$ has *cancellation* in the sense that $P_1 \oplus \mathbb{Z}G \cong P_2 \oplus \mathbb{Z}G$ implies $P_1 \cong P_2$ for all $P_1, P_2 \in [P_{(G,n)}]$ (see [\[Jacobinski 1968\]](#)). This implies that $\text{PHT}(G, n)$ and $\text{HT}(G, n)$ have cancellation in the sense that $X \vee S^n \simeq Y \vee S^n$ implies $X \simeq Y$, and recovers the main result of Dyer [\[1976\]](#).

(b) An equivalent statement appeared in [\[Johnson 2003\]](#) in the case $n = 2$, though the proof contained a small gap which was patched up in [\[Nicholson 2021b\]](#) using a theorem of Browning [\[1978\]](#).

Our proof is based on the work of Hambleton and Kreck [\[1993\]](#) and is independent of [\[Browning 1978; Johnson 2003\]](#). After establishing preliminaries in [Sections 2 and 3](#), we will prove general cancellation theorems for chain complexes of projective modules in [Section 4](#). This suffices to prove [Theorem A](#) due to the correspondence between $\text{PHT}(G, n)$ and the tree of algebraic n -complexes (see [Proposition 5.1](#)). In [Theorem 5.3](#), we give a detailed version of [Theorem A](#) which contains an explicit description of the map Ψ .

We then use of this description of Ψ to determine the induced action of $\text{Aut}(G)$ on $[P_{(G,n)}]$ via the bijection $\text{HT}(G, n) \cong \text{PHT}(G, n)/\text{Aut}(G)$. To state the induced action, consider the following two operations for M a (left) projective $\mathbb{Z}G$ -module:

- (1) If $\theta \in \text{Aut}(G)$, then let M_θ be the $\mathbb{Z}G$ -module whose abelian group is that of M but with action $g \cdot x = \theta(g)x$ for $g \in G$ and $x \in M$ (see [Lemma 6.1](#)).

- (2) If r represents a class in $(\mathbb{Z}/|G|)^\times$ and $I \subseteq \mathbb{Z}G$ is the augmentation ideal, then (I, r) is a projective $\mathbb{Z}G$ -module. The tensor product $(I, r) \otimes M$ is a projective $\mathbb{Z}G$ -module since (I, r) is a two-sided ideal (see [Lemma 4.15](#)).

In [Section 6](#), we will prove the following which is the main result of this article. Note that every projective $\mathbb{Z}G$ -module has the form $P \oplus \mathbb{Z}G^r$ where P has rank one and $r \geq 0$ (see [Section 3.1](#)).

Theorem B *Let G have k -periodic cohomology and let $n = ik$ or $ik - 2$ for some $i \geq 1$. Then Ψ induces an injective map of graded trees*

$$\bar{\Psi}: \text{HT}(G, n) \rightarrow [P_{(G,n)}]/\text{Aut}(G),$$

where the action by $\theta \in \text{Aut}(G)$ is given by

$$\theta: P \oplus \mathbb{Z}G^r \mapsto ((I, \psi_k(\theta)^i) \otimes P_\theta) \oplus \mathbb{Z}G^r,$$

where P has rank one, for some map $\psi_k: \text{Aut}(G) \rightarrow (\mathbb{Z}/|G|)^\times$ which depends only on G and k . Furthermore, $\bar{\Psi}$ is bijective if and only if $n \geq 3$ or if $n = 2$ and G has the D2 property.

This reduces the problem of determining when cancellation occurs in the homotopy trees to the purely algebraic problem of determining cancellation for $[P]$ and $[P]/\text{Aut}(G)$ which will be dealt with in [\[Nicholson 2020\]](#).

1.2 Computing the action of $\text{Aut}(G)$

After proving [Theorems A and B](#), the remainder of this article will be devoted to exploring the action of $\text{Aut}(G)$ on $[P_{(G,n)}]$. This includes establishing some general theory in preparation for the more detailed computations in [\[Nicholson 2020\]](#).

First, and perhaps somewhat surprisingly, we could find no example where the $\text{Aut}(G)$ -action described in [Theorem B](#) does not take the form of the simpler action $P \mapsto P_\theta$. In all examples computed, we had $(I, \psi_k(\theta)) \cong \mathbb{Z}G$ which implies that $(I, \psi_k(\theta)^i) \cong \mathbb{Z}G$. If $P \oplus \mathbb{Z}G^r \in [P_{(G,n)}]$ where P has rank one, then this implies that $\theta(P) \cong P_\theta \oplus \mathbb{Z}G^r \cong (P \oplus \mathbb{Z}G^r)_\theta$. In particular, $\theta(P) \cong P_\theta$ for all $P \in [P_{(G,n)}]$. We therefore ask the following:

Question 7.3 *Does there exist G with k -periodic cohomology and $\theta \in \text{Aut}(G)$ for which $(I, \psi_k(\theta))$ is not free?*

There are two approaches to finding examples where $(I, \psi_k(\theta))$ is not free. The first is to find an example where $(I, \psi_k(\theta))$ is not even stably free. It was shown by Dyer [\[1976, page 276\]](#) and Davis [\[1983\]](#) that $(I, \psi_k(\theta))$ is stably free when $\sigma_k(G) = 0$. Davis [\[1983, page 488\]](#) asked whether this also holds when $\sigma_k(G) \neq 0$. The second approach is to find an example where $(I, \psi_k(\theta))$ is stably free but not free. This is likely to be difficult since the general question of whether (I, r) can be stably free but not free is still open and dates back to Wall's problems list [\[1979b, Problem A4\]](#).

In [Section 8](#), we develop a general method to compute the action $P \mapsto P_\theta$. We will then use this to give the following example where the action is nontrivial. Let Q_{4n} denote the quaternion group of order $4n$, which has 4–periodic cohomology. Since $\sigma_4(Q_{4n}) = 0$, we can take $[P_{(Q_{4n}, 2)}] = [\mathbb{Z}Q_{4n}] = \bigcup_{r \geq 1} \text{SF}_r(\mathbb{Z}Q_{4n})$ where $\text{SF}_r(\mathbb{Z}Q_{4n})$ is the set of stably free $\mathbb{Z}Q_{4n}$ –modules of rank $r \geq 1$. As above, let $\theta \in \text{Aut}(Q_{4n})$ act on $[\mathbb{Z}Q_{4n}]$ by $\theta: P \mapsto (I, \psi_4(\theta)^i) \otimes P_\theta$ for some $i \geq 1$. We show:

Theorem C $\text{Aut}(Q_{24})$ acts nontrivially on $[\mathbb{Z}Q_{24}]$. More specifically, we have $|\text{SF}_1(\mathbb{Z}Q_{24})| = 3$ and $|\text{SF}_1(\mathbb{Z}Q_{24})/\text{Aut}(Q_{24})| = 2$.

This is in contrast to the case Q_{4n} for $2 \leq n \leq 5$, where $|\text{SF}_1(\mathbb{Z}Q_{4n})| = 1$, and the case Q_{28} , where $|\text{SF}_1(\mathbb{Z}Q_{28})| = |\text{SF}_1(\mathbb{Z}Q_{28})/\text{Aut}(Q_{28})| = 2$ (see [Table 1](#)).

1.3 Overview of the wider project

This article is the first of a two-part series (followed by [\[Nicholson 2020\]](#)) in which we explore the classification of finite (G, n) –complexes over groups with periodic cohomology. These results are motivated by the following.

Wall’s D2 problem for groups with 4–periodic cohomology In the language above, the D2 problem asks whether every finitely presented group G has the D2 property. This dates back to Wall’s paper on finiteness conditions [\[1965\]](#) and is currently open. The case where G has 4–periodic cohomology was proposed to contain a counterexample to the D2 problem [\[Cohen 1977\]](#), and has since been studied extensively. In this case, Johnson [\[2003\]](#) proved [Theorem A](#) when $n = 2$ and, using results of Swan [\[1983\]](#), he established the D2 property for many new groups. In [\[Nicholson 2021a; 2021b\]](#), we extended these results and determined when $\text{PHT}(G, 2)$ has cancellation.

In the case where $\text{PHT}(G, 2)$ has noncancellation, the D2 property has only been proven for Q_{28} (see [\[Mannan and Popiel 2021; Nicholson 2021b\]](#)). This motivated [Theorem B](#) in the case $n = 2$ since one imagines it might be easier to prove that $\bar{\Psi}$ is bijective rather than Ψ . The question of when $\text{HT}(G, 2)$ has cancellation is answered in [\[Nicholson 2020, Theorem A\]](#).

Stable and unstable classification of manifolds If X is a finite (G, n) –complex, then there exists an embedding $i: X \hookrightarrow \mathbb{R}^{2n+1}$. The boundary of a smooth regular neighbourhood of i gives a smooth closed $2n$ –manifold $M(X)$. If X is determined up to simple homotopy, then $M(X)$ is well defined up to s –cobordism which coincides with homeomorphism in the case where G is finite by work of Freedman. Furthermore, $M(X \vee S^n) \cong M(X) \# (S^n \times S^n)$. This can be found in [\[Bokor et al. 2021, Section 5\]](#).

Kreck and Schafer [\[1984\]](#) used this to construct smooth closed $4n$ –manifolds M_1 and M_2 for every $n \geq 1$ such that $M_1 \# (S^{2n} \times S^{2n}) \cong M_2 \# (S^{2n} \times S^{2n})$ are diffeomorphic but $M_1 \not\cong M_2$. Their examples have the form $M(X_i)$ where the $X_i \in \text{HT}(G, n)$ are the noncancellation examples for G abelian found

by Metzler, Sieradski and Dyer [Metzler 1976; Sieradski 1977; Sieradski and Dyer 1979]. Recently, Conway, Crowley, Powell and Sixt constructed examples of both simply connected M_i [Conway et al. 2023] and infinitely many M_i [Conway et al. 2021] for all $n \geq 2$. However, the examples of Kreck and Schafer remain the only known examples in dimension 4. In classifying $\text{HT}(G, n)$ when G has periodic cohomology, we hope to create a second family of examples both in dimension 4 and in higher dimensions.

Conventions

All rings R will be assumed to have a multiplicative identity and all R -modules will be assumed to be finitely generated left R -modules.

Recall that groups with periodic cohomology are necessarily finite. For most of this article, we will therefore restrict to the case where G is a finite group. However, we will briefly consider finitely presented groups more generally at the start of Sections 5 and 6.

Acknowledgements

I would like to thank FE A Johnson for his guidance and many interesting conversations. I would also like to thank Ian Hambleton, Jonathan Hillman and Wajid Mannan for helpful comments and references. Finally, I would like to thank the referee for their careful reading and suggestions which improved the exposition of this article. This work was supported by the UK Engineering and Physical Sciences Research Council (EPSRC) grant EP/N509577/1 as well as the Heilbronn Institute for Mathematical Research.

2 Extensions of modules

Let R be a ring. Recall our convention that all R -modules are assumed to be finitely generated left R -modules. For R -modules A and B , define $\text{Ext}_R^n(A, B)$ to be the set of exact sequences

$$E = (0 \rightarrow B \xrightarrow{i} E_{n-1} \xrightarrow{\partial_{n-1}} E_{n-2} \xrightarrow{\partial_{n-2}} \cdots \xrightarrow{\partial_2} E_1 \xrightarrow{\partial_1} E_0 \xrightarrow{\varepsilon} A \rightarrow 0)$$

for R -modules E_i considered up to congruence, i.e. the equivalence relation generated by *elementary congruences* which are chain maps of the form

$$\begin{array}{c} E \\ \downarrow \varphi \\ E' \end{array} = \left(\begin{array}{cccccccc} 0 & \longrightarrow & B & \longrightarrow & E_{n-1} & \longrightarrow & \cdots & \longrightarrow & E_0 & \longrightarrow & A & \longrightarrow & 0 \\ & & \downarrow \text{id} & & \downarrow \varphi_{n-1} & & & & \downarrow \varphi_0 & & \downarrow \text{id} & & \\ 0 & \longrightarrow & B & \longrightarrow & E'_{n-1} & \longrightarrow & \cdots & \longrightarrow & E'_0 & \longrightarrow & A & \longrightarrow & 0 \end{array} \right)$$

That is, two extensions E and E' are *congruent* if there exists extensions $E^{(i)}$ for $0 \leq i \leq n$ such that $E = E^{(0)}$, $E' = E^{(n)}$ and, for $i \leq n-1$, there exists an elementary congruence of the form $\varphi: E^{(i)} \rightarrow E^{(i+1)}$ or $\varphi: E^{(i+1)} \rightarrow E^{(i)}$.

We write extensions in $\text{Ext}_R^n(A, B)$ as $E = (E_*, \partial_*)$ where the maps $i: B \rightarrow E_{n-1}$ and $\varepsilon: E_0 \rightarrow A$ are understood. We will often write $\partial_i = \partial_i^E$, $i = i_E$ and $\varepsilon = \varepsilon_E$ when the need arises to distinguish different extensions.

This is an abelian group under Baer sum, and coincides with the usual definition of $\text{Ext}_R^n(A, B)$ [Weibel 1994, Section 3.4]. We will assume familiarity with the basic operations on extensions such as pullback, pushout and the Yoneda product [Johnson 2003, Section 24].

Worth emphasising however is the operation of stabilisation. If $E = (E_*, \partial_*) \in \text{Ext}_R^n(A, B)$, then define the *stabilised complex* $E \oplus R \in \text{Ext}_R^n(A, B \oplus R)$ by

$$E \oplus R = (0 \rightarrow B \oplus R \xrightarrow{\begin{pmatrix} i & 0 \\ 0 & 1 \end{pmatrix}} E_{n-1} \oplus R \xrightarrow{\begin{pmatrix} \partial_{n-1} \\ 0 \end{pmatrix}} E_{n-2} \rightarrow \cdots \rightarrow E_0 \rightarrow A \rightarrow 0).$$

This gives a well-defined map of abelian groups

$$-\oplus R: \text{Ext}_R^n(A, B) \rightarrow \text{Ext}_R^n(A, B \oplus R).$$

Let $\text{Proj}_R^n(A, B)$ denote the subset of $\text{Ext}_R^n(A, B)$ consisting of extensions (P_*, ∂_*) with the P_i projective. This is closed under Baer sum, and so is a subgroup, and is also preserved by pullbacks, pushouts, the Yoneda product and stabilisation. The following is a consequence of the cocycle description of Ext [Wall 1979a, Lemma 1.1].

Lemma 2.1 (shifting) *If A, B, C and D are R -modules, $E \in \text{Proj}_R^k(B, C)$ and $k, n, m \geq 1$, then the Yoneda product induces bijections*

$$-\circ E: \text{Ext}_R^n(C, D) \rightarrow \text{Ext}_R^{n+k}(B, D), \quad E \circ -: \text{Ext}_R^m(A, B) \rightarrow \text{Ext}_R^{m+k}(A, C).$$

This can be viewed as a sort of cancellation theorem for extensions up to congruence in the sense that $F \circ E \cong F' \circ E$ or $E \circ F \cong E \circ F'$ implies that $F \cong F'$.

A simple consequence of this is the following lemma. This can be interpreted as a kind of duality theorem for projective extensions.

Lemma 2.2 (duality) *If A, B and C are R -modules, $F \in \text{Proj}_R^k(A, C)$ and $k > n \geq 1$, then there are bijections*

$$\begin{aligned} \Psi_F: \text{Proj}_R^n(A, B) &\rightarrow \text{Proj}_R^{k-n}(B, C), & E &\mapsto (-\circ E)^{-1}(F), \\ \Psi_F^{-1}: \text{Proj}_R^{k-n}(B, C) &\rightarrow \text{Proj}_R^n(A, B), & E' &\mapsto (E' \circ -)^{-1}(F). \end{aligned}$$

We now turn our attention to an equivalence relation on $\text{Ext}_R^n(A, B)$ which is weaker than congruence. For R -modules A and B , and $E, E' \in \text{Ext}_R^n(A, B)$, a chain map $\varphi: E \rightarrow E'$ is said to be a *chain homotopy equivalence* if the restriction to the unaugmented chain complexes $\varphi: (E_*, \partial_*)_{0 \leq * < n} \rightarrow (E'_*, \partial'_*)_{0 \leq * < n}$ is a chain homotopy equivalence.

If $E, E' \in \text{Proj}_R^n(A, B)$ then, since a chain map between projective chain complexes is a chain homotopy equivalence if and only if it is a homology equivalence [Johnson 2003, Theorem 46.6], a chain homotopy equivalence $\varphi: E \rightarrow E'$ can equivalently be defined as a chain map of the form

$$\begin{array}{c} E \\ \downarrow \varphi \\ E' \end{array} = \begin{pmatrix} 0 \longrightarrow B \longrightarrow P_{n-1} \longrightarrow \cdots \longrightarrow P_0 \longrightarrow A \longrightarrow 0 \\ \phantom{P_{n-1}} \\ \phantom{P_{n-1}} \\ \phantom{P_{n-1}} \end{pmatrix}$$

where φ_A and φ_B are R -module isomorphisms. When convenient, we will often abbreviate this to $\varphi = (\varphi_B, \varphi_{n-1}, \dots, \varphi_0, \varphi_A)$. It follows easily that a congruence is a chain homotopy equivalence. We define $\text{hProj}_R^n(A, B)$ to be set of equivalence classes in $\text{Proj}_R^n(A, B)$ up to chain homotopy equivalences, which is an abelian group under Baer sum.

For special choices of modules, the shifting lemma and the duality lemma also hold for chain homotopy equivalences. We define \mathbb{Z} to be the R -module with underlying abelian group \mathbb{Z} and trivial R -action, i.e. $r \cdot n = n$ for all $r \in R$ and $n \in \mathbb{Z}$.

Lemma 2.3 (shifting) *If A and B are R -modules, $F \in \text{Proj}_R^k(\mathbb{Z}, \mathbb{Z})$ and $n, m, k \geq 1$, then the Yoneda product induces bijections*

$$-\circ F: \text{hProj}_R^n(\mathbb{Z}, A) \rightarrow \text{hProj}_R^{n+k}(\mathbb{Z}, A), \quad F \circ -: \text{hProj}_R^m(B, \mathbb{Z}) \rightarrow \text{hProj}_R^{m+k}(B, \mathbb{Z}).$$

Proof First note that $-\circ F$ induces maps on the chain homotopy classes by extending the map to $\pm \text{id}$ on F . This is necessarily surjective. To see that it is injective, suppose that there is a chain homotopy equivalence $\varphi: E_1 \circ F \rightarrow E_2 \circ F$. By considering $-\varphi$ if necessary, we can assume that $\varphi_{\mathbb{Z}} = \text{id}$, so

$$E_2 \circ F \cong (\varphi_A)_*(E_1 \circ F) = (\varphi_A)_*(E_1) \circ F.$$

By Lemma 2.1, this implies that $E_2 \cong (\varphi_A)_*(E_1)$ and so $E_1 \simeq E_2$ as required. \square

The proof of the duality lemma in this setting is similar and so will be omitted.

Lemma 2.4 (duality) *If A is an R -module, $F \in \text{Proj}_R^k(\mathbb{Z}, \mathbb{Z})$ and $k > n \geq 1$, then there are bijections*

$$\begin{aligned} \Psi_F: \text{hProj}_R^n(\mathbb{Z}, A) &\rightarrow \text{hProj}_R^{k-n}(A, \mathbb{Z}), & E &\mapsto (-\circ E)^{-1}(F), \\ \Psi_F^{-1}: \text{hProj}_R^{k-n}(A, \mathbb{Z}) &\rightarrow \text{hProj}_R^n(\mathbb{Z}, A), & E' &\mapsto (E' \circ -)^{-1}(F). \end{aligned}$$

We now specialise to the case where the underlying abelian group of R is finitely generated and torsion-free, and where R is a ring with involution, i.e. a ring with an antiautomorphism $r \mapsto \bar{r}$ such that $\bar{\bar{r}} = r$ for all $r \in R$. For example, for a finite group G , the group ring $\mathbb{Z}G$ has underlying abelian group $\mathbb{Z}^{|G|}$ and involution $\sum_{i=1}^n n_i g_i \mapsto \sum_{i=1}^n n_i g_i^{-1}$ where $n_i \in \mathbb{Z}$ and $g_i \in G$. Using this involution, any right R -module A can be viewed as a left R -module under the action $r \cdot x = x \cdot \bar{r}$ for $r \in R$ and $x \in A$. If A is a left R -module, then $A^* = \text{Hom}_R(A, R)$ is a right R -module under the action $(\varphi \cdot r)(x) = \varphi(x)r$ for $\varphi \in A^*$ and $r \in R$. We will view A^* as a left R -module using the involution on R .

Note that $(\cdot)^*$ can be viewed as a functor of R -modules: if $f: A_1 \rightarrow A_2$ is a map of R -modules, we can define $f^*: A_2^* \rightarrow A_1^*$ by $\varphi \mapsto \varphi \circ f$. For $E = (P_*, \partial_*) \in \text{Proj}_R^n(A, B)$, define the *dual extension* by

$$E^* = (0 \rightarrow A^* \xrightarrow{\varepsilon^*} P_0^* \xrightarrow{\partial_1^*} P_1^* \xrightarrow{\partial_2^*} \dots \xrightarrow{\partial_{n-2}^*} P_{n-2}^* \xrightarrow{\partial_{n-1}^*} P_{n-1}^* \xrightarrow{i^*} B^* \rightarrow 0).$$

The dual of a projective module is projective since $P \oplus Q \cong R^n$ implies that $P^* \oplus Q^* \cong (R^n)^* \cong R^n$. In particular, the P_i^* are projective R -modules.

Whilst E^* is not exact in general, it is true under mild assumptions on the modules involved. We say that an R -module A is an R -lattice if its underlying abelian group is finitely generated and torsion-free. For example, if P is a (finitely generated) projective R -module, then P is an R -lattice. This follows from the fact that $P \leq R^n$ is an R -submodule for some n and so its underlying abelian group is a subgroup of \mathbb{Z}^m where $m = n \cdot \text{rank}_{\mathbb{Z}}(R)$.

Recall that the *evaluation map* is the map $e_A: A \rightarrow A^{**}$, defined by $x \mapsto (f \mapsto f(x))$. We say an R -module is *reflexive* if e_A is an R -module isomorphism.

Lemma 2.5 *If A is an R -lattice, then A is reflexive.*

Remark 2.6 Since projective R -modules are R -lattices, this implies that they are reflexive. We note that this is true for arbitrary rings R , not just rings with involution whose underlying abelian group is finitely generated and torsion free.

This follows by noting that, if $A \cong_{\text{Ab}} \mathbb{Z}^k$, then the R -module structure is determined by a map $\rho_A: R \rightarrow M_k(\mathbb{Z})$. It can be shown that $\rho_{A^*}(r) = \rho_A(\bar{r})^T$ using the induced identification $A^* \cong_{\text{Ab}} \mathbb{Z}^k$, from which the claim follows.

It follows easily from this that the reflexivity property of R -lattices also holds on the level of extensions.

Lemma 2.7 (reflexivity) *If A and B are R -lattices and $n \geq 1$, then dualising gives an isomorphism of abelian groups*

$$*: \text{hProj}_R^n(A, B) \rightarrow \text{hProj}_R^n(B^*, A^*).$$

*If $E \in \text{Proj}_R^n(A, B)$, then there is a chain homotopy equivalence $e: E \rightarrow E^{**}$ induced by the evaluation maps.*

This has the following useful consequence which, in the language of [Johnson 2003, Theorem 28.5], says that projective R -modules are *injective relative to the class of R -lattices*.

Lemma 2.8 *Suppose A, B and E are R -lattices such that $(E, -) \in \text{Ext}_R^1(A, B)$ and P is a projective R -module. Then, for any map $f: B \rightarrow P$, there exists $\tilde{f}: E \rightarrow P$ such that $\tilde{f} \circ i = f$, i.e.*

$$\begin{array}{ccccccc} 0 & \longrightarrow & B & \xrightarrow{i} & E & \xrightarrow{\varepsilon} & A \longrightarrow 0 \\ & & \downarrow f & & \swarrow \tilde{f} & & \\ & & P & & & & \end{array}$$

We conclude this section by discussing an important invariant of projective extensions. Let $P(R)$ denote the R -module isomorphism classes of (finitely generated) projective R -modules and define the *projective class group* $C(R)$ as the quotient of $P(R)$ by the stable isomorphisms, where $P, Q \in P(R)$ are *stably isomorphic*, written $[P] = [Q]$, if $P \oplus R^i \cong Q \oplus R^j$ for some $i, j \geq 0$. This forms a group under direct sum and coincides with the Grothendieck group of the monoid $P(R)$.

For a projective extension

$$E = (0 \rightarrow B \xrightarrow{i} P_{n-1} \xrightarrow{\partial_{n-1}} P_{n-2} \xrightarrow{\partial_{n-2}} \dots \xrightarrow{\partial_2} P_1 \xrightarrow{\partial_1} P_0 \xrightarrow{\varepsilon} A \rightarrow 0),$$

we define the *Euler class* $e(E) = \sum_{i=0}^{n-1} (-1)^i [P_i] \in C(R)$. This is known to be a congruence invariant [Wall 1979a, Lemma 1.3]. In fact, more is true:

Lemma 2.9 *If A and B are R -modules, the Euler class defines a map*

$$e: \text{hProj}_R^n(A, B) \rightarrow C(R),$$

i.e. e is a chain homotopy invariant.

Proof Suppose $E_1, E_2 \in \text{Proj}_R^n(A, B)$ and that $\varphi: E_1 \rightarrow E_2$ is a chain homotopy equivalence. Then $E_2 \cong (\varphi_A)^*((\varphi_B)_*(E_1))$ and, since e is a congruence invariant, $e(E_2) = e((\varphi_A)^*((\varphi_B)_*(E_1)))$. Since pushout and pullback by automorphisms can be made to not affect the isomorphism classes of the modules in the extension, this implies that $e((\varphi_A)^*((\varphi_B)_*(E_1))) = e(E_1)$ and so e is a chain homotopy invariant. \square

The following tells us how the Euler class interacts with the Yoneda product.

Lemma 2.10 *Let A, B and C be R -modules. If $E \in \text{hProj}_R^n(A, B)$ and $F \in \text{hProj}_R^m(B, C)$, then*

$$e(F \circ E) = e(E) + (-1)^n e(F).$$

Proof Let $E = (P_*, \partial_*)_{* = 0}^{n-1}$ and let $F = (P_{*+n}, \partial_{*+n})_{* = 0}^{m-1}$. Then $F \circ E = (P_*, \partial_*)_{* = 0}^{n+m-1}$ and

$$e(F \circ E) = \sum_{i=0}^{n+m-1} (-1)^i [P_i] = \sum_{i=0}^{n-1} (-1)^i [P_i] + \sum_{i=0}^{m-1} (-1)^{i+n} [P_{i+n}] = e(E) + (-1)^n e(F). \quad \square$$

For a class $\chi \in C(R)$, we define $\text{Proj}_R^n(A, B; \chi)$ to be the subset of $\text{Proj}_R^n(A, B)$ consisting of those extensions with $e(E) = \chi$, and we can define $\text{hProj}_R^n(A, B; \chi)$ similarly as a subset of $\text{hProj}_R^n(A, B)$.

We have the following nice interpretations for the extensions $E \in \text{Proj}_R^n(A, B)$ with $e(E) = 0$. This follows easily by repeatedly forming the direct sum with length two extensions $P \xrightarrow{\cong} P$ for various $P \in P(R)$.

Lemma 2.11 *If A and B are R -modules and $n \geq 2$, then every congruence class in $\text{Proj}_R^n(A, B; 0)$ has a representative E of the form $E = (F_*, \partial_*)$ with the F_i free.*

This fails in the case $n = 1$, where it is not possible to form the direct sum with length two extensions $R \xrightarrow{\cong} R$ without altering the chain homotopy type. In fact, for a projective extension

$$E = (0 \rightarrow B \rightarrow P \rightarrow A \rightarrow 0),$$

we can define the *unstable Euler class* $\hat{e}(E) = P \in P(R)$.

Lemma 2.12 *If A and B are R -modules, the unstable Euler class defines a map*

$$\hat{e}: \text{hProj}_R^1(A, B) \rightarrow P(R).$$

Proof For $E_1 = (P_1, -)$, $E_2 = (P_2, -) \in \text{Proj}_R^1(A, B)$, recall that a chain map $\varphi: E_1 \rightarrow E_2$ is a chain homotopy equivalence if it induces a chain homotopy equivalence between the length one chain complexes P_1 and P_2 , i.e. if the restriction $\varphi|_{P_1}: P_1 \rightarrow P_2$ is an isomorphism. \square

3 Projective $\mathbb{Z}G$ -modules and the Swan finiteness obstruction

Throughout this section, we will let G be a finite group. The results of the previous section apply in the case $R = \mathbb{Z}G$ since $\mathbb{Z}G$ is a ring with involution which is finitely generated and torsion-free as an abelian group. The aim of this section will be to recall some of the special features of projective modules over $\mathbb{Z}G$ and to introduce the Swan finiteness obstruction.

3.1 Preliminaries on projective $\mathbb{Z}G$ -modules

We will now summarise the main special properties of (finitely generated) projective $\mathbb{Z}G$ -modules in the case where G is finite.

The first was shown by Swan [1960a, Theorem A].

Proposition 3.1 *Let P be a projective $\mathbb{Z}G$ -module. Then there is a projective ideal $I \subseteq \mathbb{Z}G$ such that $P \cong I \oplus \mathbb{Z}G^r$ for some $r \geq 0$.*

For a prime p , let \mathbb{Z}_p denote the p -adic integers and let $\mathbb{Z}_{(p)} = \{a/b \mid a, b \in \mathbb{Z}, p \nmid b\} \leq \mathbb{Q}$ denote the localisation at p . The next property that projective modules over $\mathbb{Z}G$ have is that they are locally free in the following sense (see [Swan 1980, Section 2] for further discussion).

Proposition 3.2 *Let P be a projective $\mathbb{Z}G$ -module. There exists $n \geq 0$ such that*

- (i) $P \otimes \mathbb{Z}_{(p)} \cong \mathbb{Z}_{(p)}G^n$ are isomorphic as $\mathbb{Z}_{(p)}G$ -modules,
- (ii) $P \otimes \mathbb{Q} \cong \mathbb{Q}G^n$ are isomorphic as $\mathbb{Q}G$ -modules,
- (iii) $P \otimes \mathbb{Z}_p \cong \mathbb{Z}_pG^n$ are isomorphic as \mathbb{Z}_pG -modules,
- (iv) $P \otimes \mathbb{Q}_p \cong \mathbb{Q}_pG^n$ are isomorphic as \mathbb{Q}_pG -modules.

Proof Items (ii) and (iv) each follow from [Swan 1970, Theorem 4.2]. Given this, (i) and (iii) now follow from [Swan 1970, Theorem 2.21]. \square

We define the *rank* of P , denoted by $\text{rank}(P)$, to be the $n \geq 0$ in the proposition above. For example, if $I \subseteq \mathbb{Z}G$ is a nonzero projective ideal, then it can be shown that $\text{rank}(I) = 1$; see [Swan 1960a, Section 7].

Let $P(\mathbb{Z}G)$ denote the set of $\mathbb{Z}G$ -module isomorphism classes of nonzero projective $\mathbb{Z}G$ -modules. This is a monoid under direct sum. Since $\text{rank}(P \oplus Q) = \text{rank}(P) + \text{rank}(Q)$ for all $P, Q \in P(\mathbb{Z}G)$, there is a surjective homomorphism of monoids

$$\text{rank}: P(\mathbb{Z}G) \rightarrow \mathbb{Z}, \quad P \mapsto \text{rank}(P).$$

Note that $\text{rank}(P) = 0$ if and only if $P = 0$. That is, if P is a nonzero projective $\mathbb{Z}G$ -module, then $\text{rank}(P) \geq 1$. This has the following consequence.

Corollary 3.3 *Let P be a nonzero projective $\mathbb{Z}G$ -module. Then there exists a surjection $\varphi: P \rightarrow \mathbb{Z}$.*

Proof Let $n = \text{rank}(P) \geq 1$ and consider the composition

$$P \xrightarrow{x \mapsto x \otimes 1} P \otimes \mathbb{Q} \xrightarrow{\cong} \mathbb{Q}G^n \xrightarrow{\pi_1} \mathbb{Q}G \xrightarrow{\varepsilon} \mathbb{Q}$$

where π_1 is projection onto the first coordinate and ε is the augmentation map. Since P is finitely generated, the image of the composition is a finitely generated subgroup of \mathbb{Q} and so is isomorphic to \mathbb{Z} . This gives the required surjection. \square

3.2 Swan modules

We will now define Swan modules which are a special type of projective module first introduced in [Swan 1960b, Section 6]. Let $\varepsilon: \mathbb{Z}G \rightarrow \mathbb{Z}$ denote the augmentation map and let $I = \text{Ker}(\varepsilon) \subseteq \mathbb{Z}G$ denote the augmentation ideal. For any $r \in \mathbb{Z}$ coprime to $|G|$, the ideal $(I, r) \subseteq \mathbb{Z}G$ is projective and depends only on $r \bmod |G|$ up to $\mathbb{Z}G$ -isomorphism [Swan 1960b]. Since (I, r) is a nonzero ideal, it has rank one as a projective $\mathbb{Z}G$ -module by the remarks in Section 3.1.

The modules (I, r) are known as *Swan modules* and the map

$$S: (\mathbb{Z}/|G|)^\times \rightarrow C(\mathbb{Z}G)$$

given by $r \mapsto [(I, r)]$ is known as the *Swan map*. This is a well-defined group homomorphism [Swan 1960b], and we define the *Swan subgroup* to be $T_G = \text{Im}(S) \leq C(\mathbb{Z}G)$.

Whilst we will not make explicit use of it in this article, we will briefly mention the closely related ideal $(N, r) \subseteq \mathbb{Z}G$ where $N = \sum_{g \in G} g$ denotes the group norm. Many authors take the (N, r) to be Swan modules instead of the ideals (I, r) . In fact, the two notions are equivalent, as the following proposition shows.

Proposition 3.4 *If G is a finite group and $r \in (\mathbb{Z}/|G|)^\times$, then $(I, r) \cong (N, r^{-1})$.*

This is presumably well known, but we will include a detailed proof here since we are not aware that one is currently available in the literature.

Proof By the uniqueness of pullbacks, it will suffice to prove that both (I, r) and (N, r^{-1}) arise as pullbacks of the map $r : \mathbb{Z} \rightarrow \mathbb{Z}/|G|$ which sends $1 \mapsto r$, and the map $\varepsilon : \mathbb{Z}G/(N) \rightarrow \mathbb{Z}/|G|$ which sends $x + (N) \mapsto \varepsilon(x) + |G|$.

First let $i : I \hookrightarrow (I, r)$ denote inclusion, let $\varphi : (I, r) \rightarrow \mathbb{Z}G/(N)$ and let $q : \mathbb{Z}G \rightarrow \mathbb{Z}G/(N)$ denote the quotient map. Then there is a diagram

$$\begin{array}{ccccccc}
 0 & \longrightarrow & I & \xrightarrow{i} & (I, r) & \xrightarrow{(1/r)\varepsilon} & \mathbb{Z} & \longrightarrow & 0 \\
 & & \downarrow \text{id} & & \downarrow q & & \downarrow r & & \\
 0 & \longrightarrow & I & \xrightarrow{j} & \mathbb{Z}G/(N) & \xrightarrow{\varepsilon} & \mathbb{Z}/|G| & \longrightarrow & 0
 \end{array}$$

where q and $(1/r)\varepsilon$ denote the restrictions of these maps to $(I, r) \subseteq \mathbb{Z}G$ and $j = q \circ i$. It can be checked that the diagram commutes and that the rows are exact, and so the right hand square is a pullback.

Now let $s \in \mathbb{Z}$ be such that $s = r^{-1} \in (\mathbb{Z}/|G|)^\times$, so that $(N, r^{-1}) \cong (N, s)$. Define $f : (N, s) \rightarrow \mathbb{Z}G/(N)$ by sending $Nx + sy \mapsto y$. Then consider the diagram

$$\begin{array}{ccccccc}
 0 & \longrightarrow & I & \xrightarrow{s} & (N, s) & \xrightarrow{\varepsilon} & \mathbb{Z} & \longrightarrow & 0 \\
 & & \downarrow \text{id} & & \downarrow f & & \downarrow r & & \\
 0 & \longrightarrow & I & \xrightarrow{j} & \mathbb{Z}G/(N) & \xrightarrow{\varepsilon} & \mathbb{Z}/|G| & \longrightarrow & 0
 \end{array}$$

Similarly, it can be checked that this commutes and that the rows are exact. □

3.3 Projective extensions

We will now consider the classification of extensions $\text{Proj}_{\mathbb{Z}G}^n(\mathbb{Z}, A)$ for a fixed $\mathbb{Z}G$ -module A . The following can be found in [Johnson 2003, Proposition 34.2] and shows that any two elements of $\text{Proj}_{\mathbb{Z}G}^n(\mathbb{Z}, A)$ are related by pullbacks. Note that this isomorphism depends on the choice of E and so only exists when $\text{Proj}_{\mathbb{Z}G}^n(\mathbb{Z}, A)$ is nonempty.

Proposition 3.5 *Let A be a $\mathbb{Z}G$ -module and $n \geq 1$. Then, for any $E \in \text{Proj}_{\mathbb{Z}G}^n(\mathbb{Z}, A)$, there is a bijection*

$$(m_\bullet)^* : (\mathbb{Z}/|G|)^\times \rightarrow \text{Proj}_{\mathbb{Z}G}^n(\mathbb{Z}, A)$$

given by $r \mapsto (m_r)^(E)$, where $m_r : \mathbb{Z} \rightarrow \mathbb{Z}$ denotes multiplication by r .*

Remark 3.6 This corresponds to the fact that extensions with fixed ends are determined by their k -invariants; see, for example, [Johnson 2003, Chapter 6].

Let e denote the stable Euler class as defined in Section 2. The next result computes the image of projective extensions under the stable Euler class.

Proposition 3.7 Let e denote the stable Euler class. Let $n \geq 1$ and let A be a $\mathbb{Z}G$ -module such that there exists $E \in \text{Proj}_{\mathbb{Z}G}^n(\mathbb{Z}, A)$. If $e(E) = [P]$, then

$$e(\text{Proj}_{\mathbb{Z}G}^n(\mathbb{Z}, A)) = [P] + T_G \subseteq C(\mathbb{Z}G).$$

Proof This was proven in [Swan 1960b, Lemmas 7.3 and 7.4] in the case $A = \mathbb{Z}$, and the proof for arbitrary A is analogous. We will outline the steps here for the convenience of the reader.

The first step is to show that, for any $E, E' \in \text{Proj}_{\mathbb{Z}G}^n(\mathbb{Z}, A)$, we have $e(E') - e(E) \in T_G$. By applying Schanuel's lemma (see [Swan 1960b, Proposition 1.1]) to the duals $E^*, (E')^* \in \text{Proj}_{\mathbb{Z}G}^n(A^*, \mathbb{Z})$, we get an isomorphism $\mathbb{Z} \oplus e(E^*) \cong \mathbb{Z} \oplus e((E')^*)$ and so $e((E')^*) - e(E^*) \in T_G$ by [Swan 1960b, Lemma 6.2]. Since $e(E^*) = e(E)^*$ and projective $\mathbb{Z}G$ -modules are reflexive, dualising gives that $e(E') - e(E) \in T_G$.

The second step is to show that, given $E \in \text{Proj}_{\mathbb{Z}G}^n(\mathbb{Z}, A)$, there exists $E' \in \text{Proj}_{\mathbb{Z}G}^n(\mathbb{Z}, A)$ such that $e(E') - e(E) = [(I, r)]$. This can be constructed in the same way as in [Swan 1960b, Lemma 7.4]. That is, using [Swan 1960b, Remark 2.1]. \square

3.4 The Swan finiteness obstruction

We will now specialise further to the case $A = \mathbb{Z}$. Recall that a finite group G is said to have k -periodic cohomology if there is an isomorphism of abelian groups $\hat{H}^i(G; \mathbb{Z}) \cong \hat{H}^{i+k}(G; \mathbb{Z})$ for all $i \in \mathbb{Z}$.

Remark 3.8 Many authors define finite groups with periodic cohomology by the a priori stronger condition that there exists a class $u \in \hat{H}^k(G; \mathbb{Z})$ such that cup product induces an isomorphism

$$u \cup -: \hat{H}^i(G; \mathbb{Z}) \rightarrow \hat{H}^{i+k}(G; \mathbb{Z})$$

for all $i \in \mathbb{Z}$. These definitions are equivalent since, if $\hat{H}^i(G; \mathbb{Z}) \cong \hat{H}^{i+k}(G; \mathbb{Z})$ for all $i \in \mathbb{Z}$, then $\hat{H}^k(G; \mathbb{Z}) \cong \hat{H}^0(G; \mathbb{Z}) \cong \mathbb{Z}/|G|$ which implies that the condition above holds by [Brown 1982, VI.9.1].

The following can be extracted from [Cartan and Eilenberg 1956, Chapter XII].

Proposition 3.9 Let G be a finite group. Then G has k -periodic cohomology if and only if $\text{Proj}_{\mathbb{Z}G}^k(\mathbb{Z}, \mathbb{Z})$ is nonempty.

If G has k -periodic cohomology then, since $\text{Proj}_{\mathbb{Z}G}^k(\mathbb{Z}, \mathbb{Z})$ is nonempty, Proposition 3.7 implies that there exists $P \in P(\mathbb{Z}G)$ for which

$$e(\text{Proj}_{\mathbb{Z}G}^k(\mathbb{Z}, \mathbb{Z})) = [P] + T_G \subseteq C(\mathbb{Z}G)$$

where $P(\mathbb{Z}G)$ denotes the set of nonzero projective $\mathbb{Z}G$ -modules. We can then quotient by T_G to get a unique class in $C(\mathbb{Z}G)/T_G$ which depends only on G and k . The Swan finiteness obstruction is defined as

$$\sigma_k(G) = [P] \in C(\mathbb{Z}G)/T_G.$$

Recall that a group G has *free period* k if there exists $E = (F_*, \partial_*) \in \text{Proj}_{\mathbb{Z}G}^k(\mathbb{Z}, \mathbb{Z})$ with the F_i free. The following is [Swan 1960b, Proposition 5.1].

Proposition 3.10 *Let G have k -periodic cohomology. Then $\sigma_k(G) = 0$ if and only if G has free period k .*

Remark 3.11 By a construction of Milnor, this is equivalent to the existence of a finite CW-complex X with $\pi_1(X) \cong G$ and $\tilde{X} \simeq S^{k-1}$ [Swan 1960b, Proposition 3.1]. Examples of groups with $\sigma_k(G) \neq 0$ were found by Milgram [1985].

We will conclude this section by giving a constraint on the projective $\mathbb{Z}G$ -modules P which can arise as a representative of the Swan finiteness obstruction.

We would like to compare $[P]$ and $[P^*]$ when $\sigma_k(G) = [P] + T_G$. This is difficult for general projectives since there exists finite groups G and projectives P for which $[P^*] \neq \pm[P]$, even in $C(\mathbb{Z}G)/T_G$. For example, we can take $G = \mathbb{Z}/37^2$ [Curtis and Reiner 1987, Theorem 50.56]. However, in our situation, we have the following.

Proposition 3.12 *If G has k -periodic cohomology, and $\sigma_k(G) = [P] + T_G$, then*

$$[P] = -[P^*] \in C(\mathbb{Z}G)/T_G.$$

Proof By Proposition 3.7, there exists $E \in \text{Proj}_{\mathbb{Z}G}^k(\mathbb{Z}, \mathbb{Z})$ with $e(E) = [P]$ and, by forming the direct sum with length two extensions $\mathbb{Z}G \xrightarrow{\cong} \mathbb{Z}G$, we can assume that

$$E \cong (0 \rightarrow \mathbb{Z} \xrightarrow{i} P \xrightarrow{\partial_{k-1}} F_{k-2} \xrightarrow{\partial_{k-2}} \dots \xrightarrow{\partial_1} F_0 \xrightarrow{\varepsilon} \mathbb{Z} \rightarrow 0)$$

for some F_i free. Dualising then gives that

$$E^* \cong (0 \rightarrow \mathbb{Z} \xrightarrow{\varepsilon^*} F_0 \xrightarrow{\partial_1^*} \dots \xrightarrow{\partial_{k-2}^*} F_{k-2} \xrightarrow{\partial_{k-1}^*} P^* \xrightarrow{i^*} \mathbb{Z} \rightarrow 0)$$

and, since k is necessarily even [Cartan and Eilenberg 1956, page 261], Schanuel’s lemma implies that

$$\mathbb{Z} \oplus P \oplus P^* \oplus F \cong \mathbb{Z} \oplus F'$$

for some F and F' free. By [Swan 1960b, Lemma 6.2], we then get that $[P \oplus P^*] \in T_G$. □

Remark 3.13 For a finite group G , the standard involution on $C(\mathbb{Z}G)$ is given by $[P] \mapsto -[P^*]$; see [Curtis and Reiner 1987, Section 50E]. This turns $C(\mathbb{Z}G)$ into a $\mathbb{Z}C_2$ -module where the C_2 -action is given by the involution. This additional structure has proven to be a useful for computing class groups [Curtis and Reiner 1987, page 284]. Note that T_G is fixed by this involution. This follows from the fact that $(I, r)^* \cong (N, r) \cong (I, r^{-1})$ by [Swan 1983, Lemma 17.1] and Proposition 3.4 respectively. Hence the involution induces an involution on $C(\mathbb{Z}G)/T_G$ and so endows it with a natural $\mathbb{Z}C_2$ -module structure. With respect to this action, Proposition 3.12 says that $\sigma_k(G) \in (C(\mathbb{Z}G)/T_G)^{C_2}$.

4 Classification of projective chain complexes

We would now like to consider more generally the classification of projective extensions over $\mathbb{Z}G$ with only one fixed end. Throughout this section, G will denote a finite group. For $n \geq 0$, a *projective n -complex* $E = (P_*, \partial_*)$ over $\mathbb{Z}G$ is a chain complex consisting of an exact sequence

$$E = (P_n \xrightarrow{\partial_n} P_{n-1} \xrightarrow{\partial_{n-1}} \dots \xrightarrow{\partial_1} P_0)$$

where $H_0(P_*) \cong \mathbb{Z}$ and the P_i are (finitely generated) projective $\mathbb{Z}G$ -modules. An *algebraic n -complex* is a projective n -complex such that the P_i are free.

Let $\text{Proj}(G, n)$ denote the set of chain homotopy types of projective n -complexes over $\mathbb{Z}G$, which is a graded graph with edges between each $E = (P_*, \partial_*)$ and

$$E \oplus \mathbb{Z}G = (P_n \oplus \mathbb{Z}G \xrightarrow{(\partial_n, 0)} P_{n-1} \xrightarrow{\partial_{n-1}} \dots \xrightarrow{\partial_1} P_0).$$

Similarly, let $\text{Alg}(G, n)$ denote the set of chain homotopy types of algebraic n -complexes over $\mathbb{Z}G$, which is also a graded graph under stabilisation. By extending the projective n -complex by $\text{Ker}(\partial_n)$, it is easy to see that there is a bijection

$$\text{Proj}(G, n) \cong \coprod_{A \in \text{Mod}(\mathbb{Z}G)} \text{hProj}_{\mathbb{Z}G}^{n+1}(\mathbb{Z}, A).$$

By abuse of notation, we will assume they are the same, i.e. that an extension $E \in \text{Proj}(G, n)$ lies in $\text{hProj}_{\mathbb{Z}G}^{n+1}(\mathbb{Z}, A)$ for some A . For a class $\chi \in C(\mathbb{Z}G)$, let $\text{Proj}(G, n; \chi)$ denote the subset of projective extensions E with $e(E) = \chi$. Note that $\text{Alg}(G, n) \cong \text{Proj}(G, n; 0)$ for $n \geq 2$.

4.1 General classification of projective n -complexes

The following is well known; see [Mannan 2007, Theorem 1.1; Hambleton et al. 2013, Proof of Lemma 8.12].

Theorem 4.1 *If $n \geq 0$ and $\chi \in C(\mathbb{Z}G)$, then $\text{Proj}(G, n; \chi)$ is a graded tree, i.e. if $E, E' \in \text{Proj}(G, n)$ have $e(E) = e(E')$, then $E \oplus \mathbb{Z}G^i \simeq E' \oplus \mathbb{Z}G^j$ for some $i, j \geq 0$.*

We will now prove a cancellation theorem for projective n -complexes. Our proof will be modelled on Hambleton and Kreck's proof [1993, Theorem B] that, if X and Y are finite 2-complexes with finite fundamental group such that $X \simeq X_0 \vee S^2$ and $X \vee S^2 \simeq Y \vee S^2$, then $X \simeq Y$. This idea was applied to algebraic 2-complexes in [Hambleton 2019].

If A is a $\mathbb{Z}G$ -module, then $x \in A$ is *unimodular* if there exists a map $f: A \rightarrow \mathbb{Z}G$ such that $f(x) = 1$. Let $\text{Um}(A) \subseteq A$ denote the set of unimodular elements in A .

Lemma 4.2 *Let A and B be $\mathbb{Z}G$ -modules. Then:*

- (i) *If $\varphi: A \rightarrow B$ is an isomorphism, then $\varphi(\text{Um}(A)) = \text{Um}(B)$.*
- (ii) *$(0, 1) \in \text{Um}(A \oplus \mathbb{Z}G)$, i.e. if $\varphi: A \oplus \mathbb{Z}G \rightarrow B$ is an isomorphism, then $\varphi(0, 1) \in \text{Um}(B)$.*

Suppose a $\mathbb{Z}G$ -module A has a splitting $A = A_1 \oplus A_2 \oplus \cdots \oplus A_n$. Then a map $f: A_i \rightarrow A_j$ can be viewed as an endomorphism of A by extending it to vanish everywhere else. Write $\text{GL}(A)$ for the group of automorphisms of A and define

$$E(A_i, A_j) = \langle 1 + f, 1 + g \mid f: A_i \rightarrow A_j, g: A_j \rightarrow A_i \rangle \leq \text{GL}(A)$$

to be the subgroup of *elementary automorphisms* for $i \neq j$, where $1: A \rightarrow A$ denotes the identity map.

The main result we will use is the following, which can be proven by combining [Hambleton and Kreck 1993, Corollary 1.12 and Lemma 1.16]. Let $\mathbb{Z}_{(p)} = \{a/b \mid a, b \in \mathbb{Z}, p \nmid b\} \leq \mathbb{Q}$ denote the localisation at a prime p and $A_{(p)} = A \otimes \mathbb{Z}_{(p)}$.

Theorem 4.3 *Suppose A is a $\mathbb{Z}G$ -module for which $\mathbb{Z}_{(p)} \oplus A_{(p)}$ is a free $\mathbb{Z}_{(p)}G$ -module for all but finitely many primes p . If $F_1, F_2 \cong \mathbb{Z}G$, then*

$$\mathcal{G} = \langle E(F_1, A \oplus F_2), E(F_2, A \oplus F_1) \rangle \leq \text{GL}(A \oplus F_1 \oplus F_2)$$

acts transitively on $\text{Um}(A \oplus F_1 \oplus F_2)$.

We will now establish criteria for which the above conditions hold for a $\mathbb{Z}G$ -module A . First recall that, by an extension of Maschke’s theorem of representations, the group ring RG is semisimple whenever R is a commutative ring such that $|G| \in R^\times$. This is the case when $R = \mathbb{Z}_{(p)}$ for p a prime not dividing $|G|$. This has the following consequence.

Lemma 4.4 *Let $n \geq 1$ be odd, let p be a prime not dividing $|G|$ and let A be a $\mathbb{Z}G$ -module for which $\text{Proj}_{\mathbb{Z}G}^n(\mathbb{Z}, A)$ is nonempty. Then $\mathbb{Z}_{(p)} \oplus A_{(p)}$ is a free $\mathbb{Z}_{(p)}G$ -module.*

Proof Let $E = (P_*, \partial_*) \in \text{Proj}_{\mathbb{Z}G}^n(\mathbb{Z}, A)$. Recall that localisation is an exact functor (since, for example, $\mathbb{Z}_{(p)}$ is a flat module). Hence we obtain $E_{(p)} = ((P_*)_{(p)}, \partial_*) \in \text{Proj}_{\mathbb{Z}_{(p)}G}^n(\mathbb{Z}_{(p)}, A_{(p)})$ where the ∂_* are the induced maps. By the extension of Maschke’s theorem mentioned above, $\mathbb{Z}_{(p)}G$ is semisimple and so the exact sequence $E_{(p)}$ splits completely. This implies that there is an isomorphism of $\mathbb{Z}_{(p)}G$ -modules

$$\mathbb{Z}_{(p)} \oplus A_{(p)} \oplus \bigoplus_{i \text{ odd}} (P_i)_{(p)} \cong \bigoplus_{i \text{ even}} (P_i)_{(p)}.$$

By Proposition 3.2, the $(P_i)_{(p)}$ are all free $\mathbb{Z}_{(p)}G$ -modules. It follows that $\mathbb{Z}_{(p)} \oplus A_{(p)}$ is a stably free $\mathbb{Z}_{(p)}G$ -module. Since $\mathbb{Z}_{(p)}G$ is semisimple, this implies that $\mathbb{Z}_{(p)} \oplus A_{(p)}$ is a free $\mathbb{Z}_{(p)}G$ -module. \square

Note that the fact that $\text{GL}(A \oplus \mathbb{Z}G^2)$ acts transitively on $\text{Um}(A \oplus \mathbb{Z}G^2)$ already implies the following cancellation theorem for modules.

Corollary 4.5 *Suppose A is a $\mathbb{Z}G$ -module, $A \cong A_0 \oplus \mathbb{Z}G$ and $\mathbb{Z}_{(p)} \oplus (A_0)_{(p)}$ is a free $\mathbb{Z}_{(p)}G$ -module for all but finitely many primes p . Then $A \oplus \mathbb{Z}G \cong A' \oplus \mathbb{Z}G$ implies $A \cong A'$.*

Proof Let $\psi: A \oplus \mathbb{Z}G \rightarrow A' \oplus \mathbb{Z}G$ be an isomorphism and let $x = \psi^{-1}(0, 1) \in \text{Um}(A \oplus \mathbb{Z}G)$. Since $A = A_0 \oplus \mathbb{Z}G$, Theorem 4.3 implies that $\text{GL}(A \oplus \mathbb{Z}G)$ acts transitively on $\text{Um}(A \oplus \mathbb{Z}G)$ and so there is

an isomorphism $\varphi: A \oplus \mathbb{Z}G \rightarrow A \oplus \mathbb{Z}G$ such that $\varphi(0, 1) = x$. Hence $\psi \circ \varphi: A \oplus \mathbb{Z}G \rightarrow A' \oplus \mathbb{Z}G$ has $(\psi \circ \varphi)(0, 1) = (0, 1)$ and so induces an isomorphism $(\psi \circ \varphi)|_A: A \rightarrow A' \oplus \mathbb{Z}G / \text{Im}(0 \oplus \mathbb{Z}G) \cong A'$. \square

We will upgrade the above argument from modules to projective n -complexes. The existence of a well-understood subgroup $\mathcal{G} \leq \text{GL}(A \oplus \mathbb{Z}G^2)$ which acts transitively on $\text{Um}(A \oplus \mathbb{Z}G^2)$ is important since we need only show that elements in \mathcal{G} can be extended to chain homotopy equivalences on the short exact sequences.

Theorem 4.6 *Let $n \geq 0$ be even and let $E, E' \in \text{Proj}(G, n)$. If $E \simeq E_0 \oplus \mathbb{Z}G$ and $E \oplus \mathbb{Z}G \simeq E' \oplus \mathbb{Z}G$, then $E \simeq E'$.*

Proof Let $E_0 \in \text{hProj}_{\mathbb{Z}G}^{n+1}(\mathbb{Z}, A_0)$, $E = (P_*, \partial_*) \in \text{hProj}_{\mathbb{Z}G}^{n+1}(\mathbb{Z}, A)$ and $E' = (P'_*, \partial'_*) \in \text{hProj}_{\mathbb{Z}G}^{n+1}(\mathbb{Z}, A')$. If $\psi: E \oplus \mathbb{Z}G \rightarrow E' \oplus \mathbb{Z}G$ denotes the given chain homotopy equivalence in $\text{hProj}_{\mathbb{Z}G}^{n+1}(\mathbb{Z}, A_0 \oplus \mathbb{Z}G^2)$ and $\psi_A: A_0 \oplus \mathbb{Z}G^2 \rightarrow A' \oplus \mathbb{Z}G$ is the induced map on the left, consider $x = \psi_A^{-1}(0, 1) \in \text{Um}(A_0 \oplus \mathbb{Z}G^2)$.

We now claim that there exists a self chain homotopy equivalence $\varphi: E \oplus \mathbb{Z}G \rightarrow E \oplus \mathbb{Z}G$ such that the induced map $\varphi_A: A \oplus \mathbb{Z}G \rightarrow A \oplus \mathbb{Z}G$ has $\varphi_A(0, 1) = x$.

Let $F_1, F_2 \cong \mathbb{Z}G$ be such that $A = A_0 \oplus F_1$ and $A \oplus \mathbb{Z}G = A_0 \oplus F_1 \oplus F_2$. Since $\text{Proj}_{\mathbb{Z}G}^{n+1}(\mathbb{Z}, A_0)$ is nonempty and $n + 1$ is odd, we can combine [Theorem 4.3](#) and [Lemma 4.4](#) to get that there exists $\varphi_A \in \mathcal{G} = \langle E(F_1, A_0 \oplus F_2), E(F_2, A_0 \oplus F_1) \rangle \leq \text{GL}(A_0 \oplus F_1 \oplus F_2)$ such that $\varphi_A(0, 0, 1) = x$. We claim that φ_A can be extended to a chain homotopy equivalence $\varphi: E \oplus \mathbb{Z}G \rightarrow E \oplus \mathbb{Z}G$.

First recall that $E(F_2, A_0 \oplus F_1) = E(F_2, A) \leq \text{GL}(A \oplus F_2)$ is generated by elements of the form $\begin{pmatrix} 1 & 0 \\ f & 1 \end{pmatrix}$ for $f: A \rightarrow F_2$ and $\begin{pmatrix} 1 & g \\ 0 & 1 \end{pmatrix}$ for $g: F_2 \rightarrow A$.

If $i: A \hookrightarrow P$, then there exists $\tilde{f}: P \rightarrow \mathbb{Z}G$ such that $\tilde{f} \circ i = f$ by [Lemma 2.8](#). It is straightforward to verify that the following diagrams commute, and so are chain homotopy equivalences:

$$\begin{aligned} E \oplus \mathbb{Z}G & \downarrow \varphi_1 \\ E \oplus \mathbb{Z}G & \\ E \oplus \mathbb{Z}G & \downarrow \varphi_2 \\ E \oplus \mathbb{Z}G & \end{aligned} = \begin{pmatrix} 0 \longrightarrow A \oplus \mathbb{Z}G \xrightarrow{\begin{pmatrix} i & 0 \\ 0 & 1 \end{pmatrix}} P_n \oplus \mathbb{Z}G \xrightarrow{(\partial_n, 0)} P_{n-1} \xrightarrow{\partial_{n-1}} \dots \xrightarrow{\partial_1} P_0 \longrightarrow 0 \\ \downarrow \begin{pmatrix} 1 & 0 \\ f & 1 \end{pmatrix} & \downarrow \begin{pmatrix} 1 & 0 \\ \tilde{f} & 1 \end{pmatrix} & \downarrow \text{id}_{P_{n-1}} & \downarrow \text{id}_{P_0} \\ 0 \longrightarrow A \oplus \mathbb{Z}G \xrightarrow{\begin{pmatrix} i & 0 \\ 0 & 1 \end{pmatrix}} P_n \oplus \mathbb{Z}G \xrightarrow{(\partial_n, 0)} P_{n-1} \xrightarrow{\partial_{n-1}} \dots \xrightarrow{\partial_1} P_0 \longrightarrow 0 \end{pmatrix}$$

$$\begin{pmatrix} 0 \longrightarrow A \oplus \mathbb{Z}G \xrightarrow{\begin{pmatrix} i & 0 \\ 0 & 1 \end{pmatrix}} P_n \oplus \mathbb{Z}G \xrightarrow{(\partial_n, 0)} P_{n-1} \xrightarrow{\partial_{n-1}} \dots \xrightarrow{\partial_1} P_0 \longrightarrow 0 \\ \downarrow \begin{pmatrix} 1 & g \\ 0 & 1 \end{pmatrix} & \downarrow \begin{pmatrix} 1 & i \circ g \\ 0 & 1 \end{pmatrix} & \downarrow \text{id}_{P_{n-1}} & \downarrow \text{id}_{P_0} \\ 0 \longrightarrow A \oplus \mathbb{Z}G \xrightarrow{\begin{pmatrix} i & 0 \\ 0 & 1 \end{pmatrix}} P_n \oplus \mathbb{Z}G \xrightarrow{(\partial_n, 0)} P_{n-1} \xrightarrow{\partial_{n-1}} \dots \xrightarrow{\partial_1} P_0 \longrightarrow 0 \end{pmatrix}$$

Similarly, we can show that the generators of $E(F_1, A_0 \oplus F_2)$ extend to chain homotopy equivalences. Hence, by writing $\varphi_A \in \mathcal{G}$ as the composition of maps of this form, we can get a chain homotopy equivalence $\varphi: E \oplus \mathbb{Z}G \rightarrow E \oplus \mathbb{Z}G$ by taking the composition of equivalences on each of the generators.

Now consider the map

$$\psi \circ \varphi = (\psi_A \circ \varphi_A, \psi_P \circ \varphi_P, \text{id}, \dots, \text{id}): E \oplus \mathbb{Z}G \rightarrow E' \oplus \mathbb{Z}G.$$




Figure 1: A graded tree which is a fork.

Since $(\psi_A \circ \varphi_A)(0, 1) = (0, 1)$, it must have the form $\psi_A \circ \varphi_A = \begin{pmatrix} \phi_A & 0 \\ 0 & 1 \end{pmatrix}$ since it is an isomorphism. By commutativity, $(\psi_P \circ \varphi_P)(0, 1) = (0, 1)$ and so similarly $\psi_P \circ \varphi_P = \begin{pmatrix} \phi_P & 0 \\ 0 & 1 \end{pmatrix}$ for some $\phi_P: P \rightarrow P'$. We are now done by noting that the triple $(\phi_A, \phi_P, \text{id}, \dots, \text{id})$ defines a chain homotopy equivalence $E \simeq E'$. □

We say that a graded tree is a *fork* if it has a single vertex at each nonminimal grade and a finite set of vertices at the minimal grade.

Corollary 4.7 *If $n \geq 0$ is even, G is a finite group and $\chi \in C(\mathbb{Z}G)$, then $\text{Proj}(G, n; \chi)$ is a fork. In particular, $\text{Alg}(G, n)$ is a fork for $n \geq 2$ even.*

This recovers the even-dimensional case of a result of Browning [1978, Theorem 5.4]. This fails in odd dimensions, i.e. there are examples of finite groups G for which $\text{Alg}(G, n)$ is not a fork for some n odd [Dyer 1979].

4.2 Projective 0-complexes and the unstable Euler class

We now consider the case $n = 0$. Recall that $P(\mathbb{Z}G)$ denotes the set of $\mathbb{Z}G$ -module isomorphism classes of (finitely generated) nonzero projective $\mathbb{Z}G$ -modules. This is a graded graph with edges between each P and $P \oplus \mathbb{Z}G$.

Note that a projective 0-complex has the form

$$E = (0 \rightarrow A \xrightarrow{i} P \xrightarrow{\varepsilon} \mathbb{Z} \rightarrow 0),$$

and so consists of a nonzero projective module $P \in P(\mathbb{Z}G)$ as well as the additional data (A, i, ε) . If \hat{e} is the unstable Euler class, then $\hat{e}: \text{Proj}(G, 0) \rightarrow P(\mathbb{Z}G)$ is a map of graded graphs since

$$\hat{e}(E \oplus \mathbb{Z}G) \cong \hat{e}(E) \oplus \mathbb{Z}G.$$

We will now show the following:

Theorem 4.8 *If G is a finite group, then the unstable Euler class gives an isomorphism of graded graphs*

$$\hat{e}: \text{Proj}(G, 0) \rightarrow P(\mathbb{Z}G).$$

Remark 4.9 Such a statement is implicit in the proof of [Johnson 2003, Theorem IV, Theorem 57.4], though the argument there contains an error and can only be used to recover the statement above in the case of projective modules of rank one. This, however, suffices since one can instead rely on the cancellation theorems of Hambleton and Kreck [1993, Theorem B] or Browning [1978, Theorem 5.4] at that stage in the proof.

Proof To see that \hat{e} is surjective, let $P \in P(\mathbb{Z}G)$. By Corollary 3.3, there is a surjection $\varphi: P \rightarrow \mathbb{Z}$ and this defines an extension $E = (P, -) \in \text{hProj}_{\mathbb{Z}G}^1(\mathbb{Z}, \text{Ker}(\varphi))$ which has $\hat{e}(E) = P$.

We will now show injectivity. First let $E = (P, -) \in \text{hProj}_{\mathbb{Z}G}^1(\mathbb{Z}, A)$ and let $E' = (P, -) \in \text{hProj}_{\mathbb{Z}G}^1(\mathbb{Z}, A')$. We will begin by considering the case where P has rank one. To show that $E \simeq E'$, it suffices to find isomorphisms $\varphi_A: A \rightarrow A'$ and $\varphi_{\mathbb{Z}}: \mathbb{Z} \rightarrow \mathbb{Z}$ such that the following diagram commutes:

$$\begin{array}{c} E \\ \downarrow \varphi \\ E' \end{array} = \begin{pmatrix} 0 \longrightarrow A \xrightarrow{i} P \xrightarrow{\varepsilon} \mathbb{Z} \longrightarrow 0 \\ \phantom{\xrightarrow{i}} \phantom{\xrightarrow{\varepsilon}} \phantom{\mathbb{Z}} \\ \phantom{\xrightarrow{i}} \phantom{\xrightarrow{\varepsilon}} \phantom{\mathbb{Z}} \\ \phantom{\xrightarrow{i}} \phantom{\xrightarrow{\varepsilon}} \phantom{\mathbb{Z}} \\ \phantom{\xrightarrow{i}} \phantom{\xrightarrow{\varepsilon}} \phantom{\mathbb{Z}} \\ 0 \longrightarrow A' \xrightarrow{i'} P \xrightarrow{\varepsilon'} \mathbb{Z} \longrightarrow 0 \end{pmatrix}$$

Consider the maps $\bar{\varepsilon} = \varepsilon \otimes \mathbb{Q}, \bar{\varepsilon}' = \varepsilon' \otimes \mathbb{Q}: P \otimes \mathbb{Q} \cong \mathbb{Q}G \rightarrow \mathbb{Q}$. Since \mathbb{Q} has trivial G -action, each map is determined by the fact that $\bar{\varepsilon}(g) = \bar{\varepsilon}'(g) = 0$ for all $g \in G$ and $\bar{\varepsilon}(1) = \bar{\varepsilon}'(1) = x_i$ for some $x_i \in \mathbb{Q}^\times$. Hence $\text{Ker}(\bar{\varepsilon}) = \text{Ker}(\bar{\varepsilon}')$ and so $(\varepsilon' \circ i) \otimes \mathbb{Q} = 0$. Since A is a $\mathbb{Z}G$ lattice, this implies that $\varepsilon' \circ i = 0$ and so we can define maps φ_A and $\varphi_{\mathbb{Z}}$ as above. Now $\varphi_{\mathbb{Z}}$ is necessarily surjective and so an isomorphism. Hence φ_A is an isomorphism by the five lemma, and so $E \simeq E'$.

Now suppose E and E' are as above but with $\text{rank}(P) \geq 2$. By Proposition 3.1, this implies that there exists P_0 of rank one such that $P \cong P_0 \oplus \mathbb{Z}G^i$ for some $i \geq 1$. Since \hat{e} is surjective, there exists $E_0 = (P_0, -) \in \text{hProj}_{\mathbb{Z}G}^1(\mathbb{Z}, A_0)$ for some A_0 . By Theorem 4.1, there exists $j \geq 0$ for which $E_0 \oplus \mathbb{Z}G^{i+j} \simeq E \oplus \mathbb{Z}G^j \simeq E' \oplus \mathbb{Z}G^j$. Since $i \geq 1$, Theorem 4.6 then implies $E_0 \oplus \mathbb{Z}G^i \simeq E \simeq E'$. \square

For use in later sections, it will be necessary to further refine the isomorphism given by \hat{e} . Consider following two decompositions (where \cong denotes bijection):

$$\text{Proj}(G, 0) \cong \coprod_{\chi \in C(\mathbb{Z}G)} \text{Proj}(G, 0; \chi) \cong \coprod_{A \in \text{Mod}(\mathbb{Z}G)} \text{hProj}_{\mathbb{Z}G}^1(\mathbb{Z}, A).$$

We will begin by determining the image of $\text{Proj}(G, 0; \chi)$ under \hat{e} . This is immediate from Theorem 4.8 and the definition of $\text{hProj}_{\mathbb{Z}G}^1(\mathbb{Z}, A; \chi)$. For convenience, we will write $\chi = [P]$ for some $P \in P(\mathbb{Z}G)$.

Proposition 4.10 *Let $P \in P(\mathbb{Z}G)$. Then there is an isomorphism of graded trees*

$$\hat{e}: \text{Proj}(G, 0; [P]) \rightarrow [P].$$

We will next determine the image of $\text{Proj}_{\mathbb{Z}G}^1(\mathbb{Z}, A)$ under \hat{e} for A a fixed $\mathbb{Z}G$ -module such that $\text{Proj}_{\mathbb{Z}G}^1(\mathbb{Z}, A)$ is nonempty. Recall that, if $E \in \text{Proj}_{\mathbb{Z}G}^1(\mathbb{Z}, A)$, then [Proposition 3.5](#) implies that there is a bijection

$$(m_\bullet)^*: (\mathbb{Z}/|G|)^\times \rightarrow \text{Proj}_{\mathbb{Z}G}^1(\mathbb{Z}, A)$$

given by $r \mapsto (m_r)^*(E)$, where $m_r: \mathbb{Z} \rightarrow \mathbb{Z}$ denotes multiplication by r .

If M is a (left) $\mathbb{Z}G$ -module and $r \in (\mathbb{Z}/|G|)^\times$, then the tensor product $(I, r) \otimes M$ can be considered as a (left) $\mathbb{Z}G$ -module since (I, r) is a two-sided ideal. This allows us to find an explicit form for pullbacks of extensions. We will begin with the following special case.

Lemma 4.11 *Let A be a $\mathbb{Z}G$ -module and suppose $E = (P, -) \in \text{Proj}_{\mathbb{Z}G}^1(\mathbb{Z}, A)$ where $\text{rank}(P) = 1$. Then, for any $r \in (\mathbb{Z}/|G|)^\times$, there are maps \bar{i} and \bar{e} such that*

$$(m_r)^*(E) \cong (0 \rightarrow A \xrightarrow{\bar{i}} (I, r) \otimes P \xrightarrow{\bar{e}} \mathbb{Z} \rightarrow 0).$$

Proof Let $E = (P, -) \in \text{Proj}_{\mathbb{Z}G}^1(\mathbb{Z}, A)$ and note that we have the diagrams

$$\begin{array}{ccc} (I, r) & \xrightarrow{(1/r)\varepsilon} & \mathbb{Z} \\ \downarrow i & & \downarrow r \\ \mathbb{Z}G & \xrightarrow{\varepsilon} & \mathbb{Z} \end{array} \quad \begin{array}{ccc} (I, r) \otimes P & \xrightarrow{(1/r)\varepsilon \otimes 1} & \mathbb{Z} \otimes P \\ \downarrow i \otimes 1 & & \downarrow r \otimes 1 \\ \mathbb{Z}G \otimes P & \xrightarrow{\varepsilon \otimes 1} & \mathbb{Z} \otimes P \end{array}$$

where $i: (I, r) \hookrightarrow \mathbb{Z}G$ is inclusion. It can be checked directly that the first diagram is a pullback, and this implies that the second diagram is a pullback since P is projective and so flat. Since $\text{rank}(P) = 1$, we can choose identifications $\mathbb{Z}G \otimes P \cong P$ and $\mathbb{Z} \otimes P \cong \mathbb{Z}$ for which $\varepsilon \otimes 1$ corresponds to ε^E . We now have a map $(\text{id}_A, \varphi, m_r): E' \rightarrow E$ where $E' = ((I, r) \otimes P, -)$. Hence $E' \cong (m_r)^*(E)$ by uniqueness of pullbacks. □

We can now upgrade this to the general case using [Theorem 4.8](#).

Lemma 4.12 *Let A be a $\mathbb{Z}G$ -module and suppose $E = (P, -) \in \text{Proj}_{\mathbb{Z}G}^1(\mathbb{Z}, A)$.*

- (i) *There exists a projective $\mathbb{Z}G$ -module P_0 with $\text{rank}(P_0) = 1$ and $k \geq 0$ such that $P \cong P_0 \oplus \mathbb{Z}G^k$ and*

$$E \cong (0 \rightarrow A \xrightarrow{i_0} P_0 \oplus \mathbb{Z}G^k \xrightarrow{(\varepsilon_0, 0)} \mathbb{Z} \rightarrow 0)$$

for some maps i_0 and $\varepsilon_0: P_0 \rightarrow \mathbb{Z}$.

- (ii) *With P_0, i_0 and ε_0 as above,*

$$(m_r)^*(E) \cong (0 \rightarrow A \xrightarrow{\bar{i}_0} ((I, r) \otimes P_0) \oplus \mathbb{Z}G^k \xrightarrow{(\bar{\varepsilon}_0, 0)} \mathbb{Z} \rightarrow 0)$$

for some maps \bar{i}_0 and $\bar{\varepsilon}_0: (I, r) \otimes P_0 \rightarrow \mathbb{Z}$.

Proof (i) Since $P \twoheadrightarrow \mathbb{Z}$, we know that P is nonzero. Hence, by [Proposition 3.1](#), there exists a projective $\mathbb{Z}G$ -module P_0 with $\text{rank}(P_0) = 1$ and $k \geq 0$ such that $P \cong P_0 \oplus \mathbb{Z}G^k$. Since \hat{e} is an

isomorphism of graded trees, there exists $E_0 \in \text{Proj}_{\mathbb{Z}G}^1(\mathbb{Z}, A_0)$ for some $\mathbb{Z}G$ -module A_0 such that $E \cong E_0 \oplus \mathbb{Z}G^k$. Write

$$E_0 = (0 \rightarrow A_0 \xrightarrow{i'_0} P_0 \xrightarrow{\varepsilon_0} \mathbb{Z} \rightarrow 0)$$

for some i'_0 and ε_0 . The result follows by forming $E_0 \oplus \mathbb{Z}G^k$.

- (ii) The result follows by noting that $(m_r)^*(E_0 \oplus \mathbb{Z}G^k) \cong (m_r)^*(E_0) \oplus \mathbb{Z}G^k$ and evaluating $(m_r)^*(E_0)$ using Lemma 4.11. □

Remark 4.13 The proof of (i) also implies that $A \cong A_0 \oplus \mathbb{Z}G^k$.

This implies the following. This is the analogue of Proposition 3.7 which established the corresponding result for the stable Euler class e .

Proposition 4.14 Let A be a $\mathbb{Z}G$ -module and suppose $E = (P, -) \in \text{Proj}_{\mathbb{Z}G}^1(\mathbb{Z}, A)$. Then

$$\hat{e}(\text{Proj}_{\mathbb{Z}G}^1(\mathbb{Z}, A)) = \{((I, r) \otimes P_0) \oplus \mathbb{Z}G^k \mid r \in (\mathbb{Z}/|G|)^\times\} \subseteq P(\mathbb{Z}G)$$

where P_0 is any rank one projective $\mathbb{Z}G$ -module such that $P \cong P_0 \oplus \mathbb{Z}G^k$ for $k \geq 0$.

For completeness, as well as for later use, we will note the following which is a consequence of [Fröhlich et al. 1974, Remark 1.30]. This shows that Propositions 3.7 and 4.14 agree in the case $n = 1$.

Lemma 4.15 Let P be a projective $\mathbb{Z}G$ -module with $\text{rank}(P) = 1$ and let $r \in (\mathbb{Z}/|G|)^\times$. Then

$$[(I, r) \otimes P] = [(I, r)] + [P] \in C(\mathbb{Z}G).$$

5 Polarised homotopy classification of (G, n) -complexes

Recall that, for a group G , a G -polarised space is a pair (X, ρ_X) where X is a topological space and $\rho_X: \pi_1(X, *) \rightarrow G$ is a given isomorphism. We say that two G -polarised spaces (X, ρ_X) and (Y, ρ_Y) are polarised homotopy equivalent if there exists a homotopy equivalence $h: X \rightarrow Y$ such that $\rho_X = \rho_Y \circ \pi_1(h)$.

Let $\text{PHT}(G, n)$ denote the set of polarised homotopy types of finite (G, n) -complexes over G . This is a graded graph with edges between each (X, ρ_X) and $(X \vee S^2, (\rho_X)^+)$ where $(\rho_X)^+$ is induced by ρ_X and the collapse map $X \vee S^2 \rightarrow X$.

If X is a finite CW-complex, then the cellular chain complex $C_*(\tilde{X})$ can be viewed as a chain complex of $\mathbb{Z}[\pi_1(X)]$ -modules under the monodromy action. We can use a G -polarisation $\rho: \pi_1(X) \rightarrow G$ to get a chain complex of $\mathbb{Z}G$ -modules $C_*(\tilde{X}, \rho)$ which is the same as $C_*(\tilde{X})$ as a chain complex of abelian groups but with action $g \cdot x = \rho^{-1}(g)x$ for all $g \in G$ and $x \in C_i(\tilde{X})$ for some $i \geq 0$.

The following is a mild generalisation of [Nicholson 2021b, Theorem 1.1]:

Proposition 5.1 *Let G be a finitely presented group and let $n \geq 2$. Then there is an injective map of graded trees*

$$\tilde{C}_* : \text{PHT}(G, n) \rightarrow \text{Alg}(G, n)$$

induced by the map $(X, \rho) \mapsto C_(\tilde{X}, \rho)$. Furthermore:*

- (i) *If $n \geq 3$, then \tilde{C}_* is bijective.*
- (ii) *If $n = 2$, then \tilde{C}_* is bijective if and only if G has the D2 property.*

Remark 5.2 (a) Even if G does not satisfy the D2 property, Proposition 5.1 can be replaced with an isomorphism $\tilde{C}_* : \text{D2}(G) \rightarrow \text{Alg}(G, 2)$ where $\text{D2}(G)$ denotes the polarised homotopy tree of D2-complexes over G [Nicholson 2021b, Theorems 1.1].

(b) This is often vacuous in the case $n \geq 3$ since $\text{PHT}(G, n)$ and $\text{Alg}(G, n)$ are often empty. More specifically, $\text{PHT}(G, n)$ is nonempty if and only if G is of type F_n . $\text{Alg}(G, n)$ is nonempty if and only if G has type FP_n (see [Bieri 1976]), and it is well known that $F_n \iff \text{FP}_n$ for finitely presented groups. This situation arises since there exist finitely presented groups which are not of type F_n for $n \geq 3$ [Stallings 1963].

(c) This fails in general for nonfinitely presented groups. In particular, for each $n \geq 2$, Bestvina and Brady [1997] constructed a nonfinitely presented group G of type FP_n . Here $\text{PHT}(G, n)$ is empty and $\text{Alg}(G, n)$ is nonempty and so \tilde{C}_* is not bijective.

We will now use the results from the previous section to study projective n -complexes over groups with periodic cohomology. By Proposition 5.1, this will lead to a proof of the following more detailed version of Theorem A. Note that, if X is a finite (G, n) -complex, then

$$\pi_n(X) \cong H_n(\tilde{X}) \cong \text{Ker}(\partial_n : C_n(\tilde{X}) \rightarrow C_{n-1}(\tilde{X}))$$

are isomorphisms of $\mathbb{Z}G$ -modules.

Theorem 5.3 *Let G have k -periodic cohomology, let $n = ik$ or $ik - 2$ for some $i \geq 1$ and let $P_{(G,n)}$ be a projective $\mathbb{Z}G$ -module with $\sigma_{ik}(G) = [P_{(G,n)}] \in C(\mathbb{Z}G)/T_G$. Let $F \in \text{Proj}_{\mathbb{Z}G}^{ik}(\mathbb{Z}, \mathbb{Z})$ be such that $e(F) = [P_{(G,n)}]$. Then there is an injective map of graded trees*

$$\Psi : \text{PHT}(G, n) \rightarrow [P_{(G,n)}],$$

defined as follows:

- (i) *If $n = ik - 2$, then $\Psi : X \mapsto P$, where P is the unique projective $\mathbb{Z}G$ -module for which*

$$(0 \rightarrow \mathbb{Z} \xrightarrow{\alpha} P^* \xrightarrow{\beta} \pi_n(X) \rightarrow 0) \circ C_*(\tilde{X}) \simeq F$$

for some α and β .

(ii) If $n = ik$, then $\Psi: X \mapsto P$, where P is the unique projective $\mathbb{Z}G$ -module for which

$$C_*(\tilde{X}) \simeq (0 \rightarrow \pi_n(X) \xrightarrow{\alpha} P \xrightarrow{\beta} \mathbb{Z} \rightarrow 0) \circ F$$

for some α and β .

Furthermore, Ψ is bijective if and only if $n \geq 3$ or $n = 2$ and G has the D2 property.

Remark 5.4 The definition of $P_{(G,n)}$ depends on G , n and k . Note that n and k determine i except when $k = 2$ where $n = ik = (i + 1)k - 2$. However, in this case there is no ambiguity since G is cyclic [Swan 1965, Lemma 5.2] and so $\sigma_{2i}(G) = 0$ for all i .

First note that, when G has periodic cohomology, we get the following two relations between projective complexes of different dimensions.

Lemma 5.5 Suppose G has k -periodic cohomology and let $F \in \text{Proj}_{\mathbb{Z}G}^k(\mathbb{Z}, \mathbb{Z})$. If $n \geq 0$, then we have isomorphisms of graded graphs

$$- \circ F: \text{Proj}(G, n) \rightarrow \text{Proj}(G, n + k), \quad * \circ \Psi_F: \text{Proj}(G, n) \rightarrow \text{Proj}(G, k - (n + 2)),$$

where $n + 2 \leq k$ in the second case.

Proof The first isomorphism is immediate from the shifting lemma. The second isomorphism consists of the compositions

$$\text{hProj}_{\mathbb{Z}G}^{n+1}(\mathbb{Z}, A) \xrightarrow{\Psi_F} \text{hProj}_{\mathbb{Z}G}^{k-n-1}(A, \mathbb{Z}) \xrightarrow{*} \text{hProj}_{\mathbb{Z}G}^{k-n-1}(\mathbb{Z}, A^*)$$

for all $\mathbb{Z}G$ -modules A . These are bijections by the duality and reflexivity lemmas.

To see that the image of the full map is $\text{Proj}(G, k - (n + 2))$ note that, if B is such that $\text{hProj}_{\mathbb{Z}G}^{k-n-2}(\mathbb{Z}, B)$ is nonzero, then B is a $\mathbb{Z}G$ -lattice since it is a submodule of a free module. By Lemma 2.5, $B^{**} \cong B$ and so there is an isomorphism $* \circ \Psi_F: \text{hProj}_{\mathbb{Z}G}^{n+1}(\mathbb{Z}, B^*) \rightarrow \text{hProj}_{\mathbb{Z}G}^{k-n-1}(\mathbb{Z}, B)$. \square

Remark 5.6 Furthermore, if $E \in \text{Proj}(G, n)$ has $\chi = e(E)$, then it is easy to see that $e(E \circ F) = e(F) + \chi$ since k is even and $e((\psi_F(E))^*) = e(F)^* - \chi^*$.

The proof of Theorem 5.3 will now consist of applying Lemma 5.5 in the case $k \mid n$ or $n + 2$ and then composing with the isomorphism from Theorem 4.8.

We will need the following result of Wall [1979a, Corollary 12.6].

Proposition 5.7 If G has k -periodic cohomology, then

$$2\sigma_k(G) = 0 \in C(\mathbb{Z}G)/T_G.$$

By iterating extensions using the Yoneda product, it can be shown that $n\sigma_k(G) = \sigma_{nk}(G)$ and so this theorem is equivalent to showing that $\sigma_{2k}(G) = 0$, i.e. that the obstruction vanishes whenever k is not the minimal period.

Theorem 5.8 *If G has k -periodic cohomology and $\sigma_k(G) = [P_{(G,n)}] + T_G$ for some $P_{(G,n)} \in P(\mathbb{Z}G)$, then there exists $F \in \text{Proj}_{\mathbb{Z}G}^k(\mathbb{Z}, \mathbb{Z})$ such that there are isomorphisms of graded trees*

$$\text{Alg}(G, k) \xrightarrow{(-\circ F)^{-1}} \text{Proj}(G, 0; [P_{(G,n)}]) \xrightarrow{\hat{e}} [P_{(G,n)}].$$

Proof By Proposition 5.7, $\sigma_k(G) = [P_{(G,n)}] + T_G = -[P_{(G,n)}] + T_G$ and so there exists $F \in \text{Proj}_{\mathbb{Z}G}^k(\mathbb{Z}, \mathbb{Z})$ with $e(F) = -[P_{(G,n)}]$ by Proposition 3.7.

If $E \in \text{Alg}(G, k)$, then $e(E) = 0$ and so $e((-\circ F)^{-1}) = -(-1)^k e(F)$ by Lemma 2.10. Since k is even, this is equal to $[P_{(G,n)}]$. Hence the map $(-\circ F)^{-1}$ is as described. By Lemma 5.5, we get that $(-\circ F)^{-1}$ is an isomorphism.

That \hat{e} is an isomorphism follows from Proposition 4.10. □

Theorem 5.9 *If G has k -periodic cohomology and $\sigma_k(G) = [P_{(G,n)}] + T_G$ for some $P_{(G,n)} \in P(\mathbb{Z}G)$, then there exists $F \in \text{Proj}_{\mathbb{Z}G}^k(\mathbb{Z}, \mathbb{Z})$ such that there are isomorphisms of graded trees*

$$\text{Alg}(G, k - 2) \xrightarrow{* \circ \Psi_F} \text{Proj}(G, 0; [P_{(G,n)}]) \xrightarrow{\hat{e}} [P_{(G,n)}].$$

Proof By Proposition 3.12, we have that $\sigma_k(G) = [P_{(G,n)}] + T_G = -[P_{(G,n)}^*] + T_G$ and so there exists $F \in \text{Proj}_{\mathbb{Z}G}^k(\mathbb{Z}, \mathbb{Z})$ with $e(F) = -[P_{(G,n)}^*]$ by Proposition 3.7.

If $E \in \text{Alg}(G, k - 2)$, then $e(\Psi_F(E)) = -e(F)$ by Lemma 2.10. This implies that

$$e((*\circ \Psi_F)(E)) = -e(F)^* = [P_{(G,n)}]$$

and so the map $* \circ \Psi_F$ is as described. By Lemma 5.5, $* \circ \Psi_F$ is an isomorphism.

That \hat{e} is an isomorphism follows from Proposition 4.10, as in the previous theorem. □

Proof of Theorem 5.3 If G has k -periodic cohomology, then it also has ik -periodic cohomology for any $i \geq 1$. Hence, by swapping k for ik , we can assume $i = 1$. By combining Theorems 5.8 and 5.9 with Proposition 5.1, we obtain injective maps of graded trees $\Psi: \text{PHT}(G, n) \rightarrow [P_{(G,n)}]$ for $n = k$ or $k - 2$, which are bijective as required. It remains to show that, in each case, Ψ has the form given in (i) and (ii).

If $n = k - 2$, then $(*\circ \Psi_F)(C_*(\tilde{X})) \simeq (0 \rightarrow A \rightarrow P \rightarrow \mathbb{Z} \rightarrow 0)$ for some A and some $P \in [P_{(G,n)}]$. By Lemma 2.7, $\Psi_F(C_*(\tilde{X})) \simeq (0 \rightarrow \mathbb{Z} \rightarrow P^* \rightarrow A^* \rightarrow 0)$. Hence $(0 \rightarrow \mathbb{Z} \rightarrow P^* \rightarrow A^* \rightarrow 0) \circ C_*(\tilde{X}) \simeq F$ and $A^* \cong \pi_n(X)$.

If $n = k$, then $(-\circ F)^{-1}(C_*(\tilde{X})) \simeq (0 \rightarrow A \rightarrow P \rightarrow \mathbb{Z} \rightarrow 0)$ for some A and some $P \in [P_{(G,n)}]$. Hence $C_*(\tilde{X}) \simeq (0 \rightarrow A \rightarrow P \rightarrow \mathbb{Z} \rightarrow 0) \circ F$ and $A \cong \pi_n(X)$. □

This completes the proof of Theorem A.

6 Homotopy classification of (G, n) -complexes

For a finitely presented group G , an automorphism $\theta \in \text{Aut}(G)$ acts on $\text{PHT}(G, n)$ by $(X, \rho) \mapsto (X, \theta \circ \rho)$. It is straightforward to see that

$$\text{HT}(G, n) \cong \text{PHT}(G, n) / \text{Aut}(G)$$

and the goal of this chapter will be to determine the induced action of $\text{Aut}(G)$ on $[P_{(G,n)}]$ under the isomorphism $\text{PHT}(G, n) \cong [P_{(G,n)}]$ obtained in [Theorem 5.3](#).

6.1 Preliminaries on the action of $\text{Aut}(G)$

We begin by defining natural actions of $\text{Aut}(G)$ on $\mathbb{Z}G$ -modules and chain complexes of $\mathbb{Z}G$ -modules. First, for a $\mathbb{Z}G$ -module A and $\theta \in \text{Aut}(G)$, let A_θ denote the $\mathbb{Z}G$ -module whose underlying abelian group is that of A and whose action is $g \cdot x = \theta(g)x$ where $g \in G$, $x \in A$. This action has the following basic properties:

Lemma 6.1 *Let $\theta \in \text{Aut}(G)$.*

(i) *There is a $\mathbb{Z}G$ -module isomorphism*

$$i_\theta: \mathbb{Z}G \rightarrow \mathbb{Z}G_\theta, \quad \sum_{g \in G} a_i g_i \mapsto \sum_{g \in G} a_i \theta(g_i).$$

(ii) *If $A, B \in \text{Mod}(\mathbb{Z}G)$, then $(A \oplus B)_\theta \cong A_\theta \oplus B_\theta$.*

(iii) *If $P \in P(\mathbb{Z}G)$, then $P_\theta \in P(\mathbb{Z}G)$.*

We can extend the action to chain complexes as follows. If A and B are $\mathbb{Z}G$ -modules and

$$E = (E_*, \partial_*) \in \text{Ext}_{\mathbb{Z}G}^n(A, B),$$

then we define $E_\theta \in \text{Ext}_{\mathbb{Z}G}^n(A_\theta, B_\theta)$ by

$$E_\theta = (0 \rightarrow B_\theta \xrightarrow{\partial_n} (E_{n-1})_\theta \xrightarrow{\partial_{n-1}} (E_{n-2})_\theta \rightarrow \cdots \rightarrow (E_1)_\theta \xrightarrow{\partial_1} (E_0)_\theta \xrightarrow{\partial_0} A_\theta \rightarrow 0).$$

It is easy to see that this is well defined up to chain homotopy and, by the lemma above, it preserves projective extensions and so also induces a map on $\text{hProj}_{\mathbb{Z}G}^n(A, B)$. The following is immediate from the definition of $\tilde{C}_*(X, \rho)$.

Lemma 6.2 *If $E \in \text{Alg}(G, n)$, then the induced action of $\theta \in \text{Aut}(G)$ on E is given by $\theta \cdot E = E_\theta$, i.e. if $E = \tilde{C}_*(X, \rho)$, then $E_\theta = \tilde{C}_*(X, \theta \circ \rho)$.*

We now establish a few basic properties of this action which we will use later in this section. From now on, we will specialise to the case where G is a finite group. First, we note that the action commutes with dualising.

Lemma 6.3 *If A and B are $\mathbb{Z}G$ -lattices, $E \in \text{Proj}_{\mathbb{Z}G}^n(A, B)$ for $n \geq 1$ and $\theta \in \text{Aut}(G)$, then*

$$(E_\theta)^* \cong (E^*)_\theta.$$

Proof We begin by proving the corresponding statement for modules, i.e. that, if A is a $\mathbb{Z}G$ -lattice, then $(A_\theta)^* \cong (A^*)_\theta$. Let $A \cong_{\text{Ab}} \mathbb{Z}^k$, so that the $\mathbb{Z}G$ -module structure is determined by an integral representation $\rho_A: G \rightarrow \text{GL}_k(\mathbb{Z})$. As remarked earlier, $\rho_{A^*}(g) = \rho_A(g^{-1})^T$ and it is easy to see that $\rho_{A_\theta} = \rho_A \circ \theta$. Therefore $(A_\theta)^* \cong (A^*)_\theta$ follows by noting that

$$\rho_{(A_\theta)^*}(g) = \rho_{A_\theta}(g^{-1})^T = \rho_A(\theta(g^{-1}))^T = \rho_A(\theta(g)^{-1})^T$$

and

$$\rho_{(A^*)_\theta}(g) = \rho_{A^*}(\theta(g)) = \rho_A(\theta(g)^{-1})^T.$$

The result for extensions now follows immediately since θ only affects the underlying modules and not the maps between them. □

In light of this, for $\mathbb{Z}G$ -lattices A and B and $E \in \text{Proj}_{\mathbb{Z}G}^n(A, B)$, it now makes sense to write A_θ^* and E_θ^* . Note that the action also commutes with pushouts.

Lemma 6.4 *If $\theta \in \text{Aut}(G)$, $f: B_1 \rightarrow B_2$ is a $\mathbb{Z}G$ -module homomorphism and $E \in \text{Ext}_{\mathbb{Z}G}^n(A, B_1)$, then*

$$f_*(E_\theta) \cong (f_*(E))_\theta.$$

6.2 Proof of Theorem B

In the case where $A = B = \mathbb{Z}$, we can consider this as an action on $\text{Proj}_{\mathbb{Z}G}^n(\mathbb{Z}, \mathbb{Z})$ by using the identification $\mathbb{Z}_\theta \cong \mathbb{Z}$.

Lemma 6.5 *If G has k -periodic cohomology, then there exists a unique map $\psi_k: \text{Aut}(G) \rightarrow (\mathbb{Z}/|G|)^\times$ such that, for every $F \in \text{Proj}_{\mathbb{Z}G}^k(\mathbb{Z}, \mathbb{Z})$ and $\theta \in \text{Aut}(G)$,*

$$F_\theta \cong (m_{\psi_k(\theta)})_*(F).$$

Proof Fix an extension $F_0 \in \text{Proj}_{\mathbb{Z}G}^k(\mathbb{Z}, \mathbb{Z})$. By dualising and then applying [Proposition 3.5](#), it follows that every extension in $\text{Proj}_{\mathbb{Z}G}^k(\mathbb{Z}, \mathbb{Z})$ is of the form $(m_r)_*(F_0)$ for some $r \in (\mathbb{Z}/|G|)^\times$. For $\theta \in \text{Aut}(G)$, define $\psi_k(\theta) = r \in (\mathbb{Z}/|G|)^\times$ for any $r \in (\mathbb{Z}/|G|)^\times$ such that $(F_0)_\theta \cong (m_r)_*(F_0)$.

If $F \in \text{Proj}_{\mathbb{Z}G}^k(\mathbb{Z}, \mathbb{Z})$, then $F \cong (m_r)_*(F_0)$ for a unique $r \in (\mathbb{Z}/|G|)^\times$. By [Lemma 6.4](#), we now have that

$$\begin{aligned} F_\theta &\cong ((m_r)_*(F_0))_\theta \cong (m_r)_*((F_0)_\theta) \cong (m_r)_*((m_{\psi_n(\theta)})_*(F_0)) \\ &\cong (m_{\psi_n(\theta)})_*((m_r)_*(F_0)) \cong (m_{\psi_n(\theta)})_*(F). \end{aligned}$$

□

Lemma 6.6 *If $E, E' \in \text{Proj}_{\mathbb{Z}G}^k(\mathbb{Z}, \mathbb{Z})$ and $r \in \mathbb{Z}$ is coprime to $|G|$, then*

$$E \circ (m_r)_*(E') \cong (m_r)_*(E) \circ E'.$$

Proof Consider the pushout map $\nu: E' \rightarrow (m_r)_*(E')$. Since this induces m_r on the left copy of \mathbb{Z} , we can extend it to a map $\tilde{\nu}: E \circ E' \rightarrow E \circ (m_r)_*(E')$ which induces multiplication by $r \in \mathbb{Z} \subseteq \mathbb{Z}G$ on every module in E , i.e.

$$\begin{array}{c} E \circ E' \\ \downarrow \tilde{\nu} \\ E \circ (m_r)_*(E') \end{array} = \left(\begin{array}{cccccccccccc} 0 & \rightarrow & \mathbb{Z} & \xrightarrow{i} & P_{k-1} & \xrightarrow{\partial_{k-1}} & \cdots & \xrightarrow{\partial_1} & P_0 & \xrightarrow{i' \circ \varepsilon} & P'_{k-1} & \xrightarrow{\partial'_{k-1}} & \cdots & \xrightarrow{\partial'_1} & P'_0 & \xrightarrow{\varepsilon'} & \mathbb{Z} & \rightarrow & 0 \\ & & \downarrow r & & \downarrow r & & & & \downarrow r & & \downarrow \nu_{k-1} & & & & \downarrow \nu_0 & & \downarrow 1 & & \\ 0 & \rightarrow & \mathbb{Z} & \xrightarrow{i} & P_{k-1} & \xrightarrow{\partial_{k-1}} & \cdots & \xrightarrow{\partial_1} & P_0 & \xrightarrow{i' \circ \varepsilon} & P'_{k-1} & \xrightarrow{\partial'_{k-1}} & \cdots & \xrightarrow{\partial'_1} & P'_0 & \xrightarrow{\varepsilon'} & \mathbb{Z} & \rightarrow & 0 \end{array} \right)$$

By the uniqueness of pushouts, this implies that $E \circ (m_r)_*(E') \cong (m_r)_*(E \circ E') = (m_r)_*(E) \circ E'$ as required. □

Note that, if G has k -periodic cohomology and $k \mid n$, then it also has n -periodic cohomology and so ψ_n can still be defined using Lemma 6.5. The above lemma now allows us to give the following relation between ψ_k and ψ_n for $k \mid n$.

Lemma 6.7 *If G has k -periodic cohomology, $i \geq 1$ and $\theta \in \text{Aut}(G)$, then*

$$\psi_{ik}(\theta) = \psi_k(\theta)^i.$$

Proof For $F \in \text{Proj}_{\mathbb{Z}G}^k(\mathbb{Z}, \mathbb{Z})$ and $F^i \in \text{Proj}_{\mathbb{Z}G}^{ik}(\mathbb{Z}, \mathbb{Z})$, Lemma 6.5 implies that $F_\theta \cong (m_{\psi_k(\theta)})_*(F)$ and $(F^i)_\theta \cong (m_{\psi_{ik}(\theta)})_*(F^i)$. Since $(F^i)_\theta \cong (F_\theta)^i$, this implies that $(m_{\psi_{ik}(\theta)})_*(F^i) \cong ((m_{\psi_k(\theta)})_*(F))^i$.

By repeated application of Lemma 6.6,

$$(m_{\psi_{ik}(\theta)})_*(F^i) \cong ((m_{\psi_k(\theta)})_*(F))^i \cong (m_{\psi_k(\theta)})_*^i(F^i) \cong (m_{\psi_k(\theta)^i})_*(F^i)$$

and so $\psi_{ik}(\theta) \cong \psi_k(\theta)^i \pmod{|G|}$ by the extension of Proposition 3.5 to arbitrary extensions via the shifting lemma. □

In order to prove Theorem B, it suffices to check what the action of $\text{Aut}(G)$ corresponds to under the isomorphisms described in Section 5. Similarly to Section 5, it will suffice to consider the cases where $k = n$ or $n + 2$.

Theorem 6.8 *Suppose that G has k -periodic cohomology and $\sigma_k(G) = [P_{(G,n)}] + T_G$ for some $P_{(G,n)} \in P(\mathbb{Z}G)$. If $F \in \text{Proj}_{\mathbb{Z}G}^k(\mathbb{Z}, \mathbb{Z})$ is such that $e(F) = -[P_{(G,n)}]$, then*

$$\begin{aligned} \text{hProj}_{\mathbb{Z}G}^{k+1}(\mathbb{Z}, A; 0) &\xrightarrow{(-\circ F)^{-1}} \text{hProj}_{\mathbb{Z}G}^1(\mathbb{Z}, A; [P_{(G,n)}]) \xrightarrow{-\hat{e}} [P_{(G,n)}], \\ E &\mapsto E' \mapsto P \oplus \mathbb{Z}G^r, \\ E_\theta &\mapsto (m_{\psi_k(\theta)})_*((E')_\theta) \mapsto ((I, \psi_k(\theta)) \otimes P_\theta) \oplus \mathbb{Z}G^r, \end{aligned}$$

where P is a rank one projective $\mathbb{Z}G$ -module and $r \geq 0$.

Proof For the first map, it suffices to check that $(\psi_k(\theta))^*((E')_\theta) \circ F \simeq E_\theta$. Since $E' \circ F = E$, we have $(E')_\theta \circ F_\theta \simeq E_\theta$. By Lemma 6.5, $F_\theta \cong (m_{\psi_n(\theta)})_*(F)$ and so

$$E_\theta \simeq (E')_\theta \circ (m_{\psi_n(\theta)})_*(F) = (m_{\psi_n(\theta)})^*((E')_\theta) \circ F.$$

The form for the second map follows directly from Lemma 4.12. □

Theorem 6.9 Suppose that G has k -periodic cohomology and $\sigma_k(G) = [P_{(G,n)}] + T_G$ for some $P_{(G,n)} \in P(\mathbb{Z}G)$. If $F \in \text{Proj}_{\mathbb{Z}G}^k(\mathbb{Z}, \mathbb{Z})$ is such that $e(F) = -[P_{(G,n)}^*]$, then

$$\begin{aligned} \text{hProj}_{\mathbb{Z}G}^{k-1}(\mathbb{Z}, A; 0) &\xrightarrow{\Psi_F} \text{hProj}_{\mathbb{Z}G}^1(A, \mathbb{Z}; [P_{(G,n)}^*]) \xrightarrow{*} \text{hProj}_{\mathbb{Z}G}^1(\mathbb{Z}, A^*; [P_{(G,n)}]) \xrightarrow{\hat{e}} [P_{(G,n)}], \\ E &\mapsto E' \mapsto (E')^* \mapsto P \oplus \mathbb{Z}G^r, \end{aligned}$$

$$E_\theta \mapsto (m_{\psi_k(\theta)^{-1}})_*((E')_\theta) \mapsto (m_{\psi_k(\theta)})^*((E')_\theta^*) \mapsto ((I, \psi_k(\theta)) \otimes P_\theta) \oplus \mathbb{Z}G^r,$$

where P is a rank one projective $\mathbb{Z}G$ -module and $r \geq 0$.

Proof For this first map, it suffices to check that $(m_{\psi_k(\theta)^{-1}})_*((E')_\theta) \circ E_\theta \simeq F$. Since $E' \circ E \simeq F$, we have $(E')_\theta \circ E_\theta \simeq F_\theta$. By Lemma 6.5, $F_\theta \cong (m_{\psi_n(\theta)})_*(F)$ and so

$$F \simeq (m_{\psi_k(\theta)^{-1}})_*((E')_\theta \circ E_\theta) \simeq (m_{\psi_k(\theta)^{-1}})_*((E')_\theta) \circ E_\theta.$$

For the second map, it is easy to see that pushouts dualise to pullbacks in the other direction, i.e. if $E_0 = (m_{\psi_k(\theta)^{-1}})_*((E')_\theta)$, then $(m_{\psi_k(\theta)^{-1}})_*(E_0^*) \simeq (E')_\theta^*$ and so $E_0^* \simeq (m_{\psi_k(\theta)})^*((E')_\theta^*)$. The form for the third map follows directly from Lemma 4.12. □

If G has k -periodic cohomology and $n = ik$ or $ik - 2$ for some $i \geq 1$, then the above shows that the induced action of $\theta \in \text{Aut}(G)$ on $[P_{(G,n)}]$ is given by $P \oplus \mathbb{Z}G^r \mapsto ((I, \psi_{ik}(\theta)) \otimes P_\theta) \oplus \mathbb{Z}G^r$ where P has rank one and $r \geq 0$. Furthermore, $\psi_{ik}(\theta) = \psi_k(\theta)^i$ by Lemma 6.7.

This completes the proof of Theorem B except for a possible discrepancy in the case where $k = 2$ and i is not determined by the fact that $n = ik$ or $ik - 2$ (see Remark 5.4). However, in this case, G is cyclic and so $(I, r) \cong \mathbb{Z}G$ for all $r \in (\mathbb{Z}/|G|)^\times$ by [Swan 1960b, Corollary 6.1]. Hence $(I, \psi_k(\theta)^i) \cong \mathbb{Z}G$ is independent of i .

7 Stably free Swan modules and (G, n) -complexes

Before computing the action of $\text{Aut}(G)$ on $[P_{(G,n)}]$, we will pause to consider the role of Swan modules in the classification of (G, n) -complexes. We begin by considering the map

$$\psi_k : \text{Aut}(G) \rightarrow (\mathbb{Z}/|G|)^\times$$

where G has k -periodic cohomology.

If $\theta \in \text{Aut}(G)$, then the action $E \mapsto E_\theta$ induces an action of $\text{Aut}(G)$ on $H^k(G; \mathbb{Z}) = \text{Ext}_{\mathbb{Z}G}^k(\mathbb{Z}, \mathbb{Z})$. This agrees with the usual action coming from the alternate definition of $H^k(-; \mathbb{Z})$ as a functor on groups

[Cartan and Eilenberg 1956, Chapter XII]. This implies that $\text{Im}(\psi_k) = \text{Aut}_k(G)$ which is defined in [Dyer 1976, Section 8]. We will now give several examples of maps $\psi_k: \text{Aut}(G) \rightarrow (\mathbb{Z}/|G|)^\times$.

Cyclic If $C_n = \langle x \mid x^n = 1 \rangle$ is the cyclic group of order n , then

$$\text{Aut}(C_n) = \{\theta_i: x \mapsto x^i \mid i \in (\mathbb{Z}/n)^\times\}$$

and $\psi_2: \text{Aut}(C_n) \rightarrow (\mathbb{Z}/n)^\times$ sends $\theta_i \mapsto i$ by [Swan 1960b, Proposition 8.1]. This is surjective and so recovers the classical results $T_{C_n} = 1$.

Dihedral If $D_{4n+2} = \langle x, y \mid x^{2n+1} = y^2 = 1, yxy^{-1} = x^{-1} \rangle$ is the dihedral group of order $4n + 2$, then

$$\text{Aut}(D_{4n+2}) = \{\theta_{i,j}: x \mapsto x^i, y \mapsto x^j y \mid i \in (\mathbb{Z}/(2n+1))^\times, j \in \mathbb{Z}/(2n+1)\}$$

and $\psi_4: \text{Aut}(D_{4n+2}) \rightarrow (\mathbb{Z}/(4n+2))^\times$ sends $\theta_{i,j} \mapsto i^2$ by the discussion in [Johnson 2002, Section 5]. Since $(\mathbb{Z}/(4n+2))^\times = \pm((\mathbb{Z}/(4n+2))^\times)^2$, this recovers the result $T_{D_{4n+2}} = 1$.

Quaternionic Let $Q_{4n} = \langle x, y \mid x^n = y^2, yxy^{-1} = x^{-1} \rangle$ is the quaternion group of order $4n$. For $n = 2$, it is shown in [Swan 1960b, Proposition 8.3] that $\psi_4: \text{Aut}(Q_8) \rightarrow (\mathbb{Z}/8)^\times$ sends $\theta \mapsto 1$ for all $\theta \in \text{Aut}(G)$. For $n \geq 3$,

$$\text{Aut}(Q_{4n}) = \{\theta_{i,j}: x \mapsto x^i, y \mapsto x^j y \mid i \in (\mathbb{Z}/2n)^\times, j \in \mathbb{Z}/2n\}$$

and $\psi_4: \text{Aut}(Q_{4n}) \rightarrow (\mathbb{Z}/4n)^\times$ sends $\theta_{i,j} \mapsto i^2$ by, for example, [Golasiński and Gonçalves 2004, Proposition 1.1].

The following was noted by Davis [1983] and Dyer [1976, Note (b)]. It would be interesting to know, as was asked by Davis, whether this holds in the case $\sigma_k(G) \neq 0$.

Proposition 7.1 *If G has free period k , then $S \circ \psi_k = 0$, i.e. $(I, \psi_k(\theta))$ is stably free for all $\theta \in \text{Aut}(G)$.*

Proof Note that Theorems 6.8 and 6.9 each show that $[P] = [(I, \psi_k(\theta)) \otimes P_\theta]$ for all $P \in P(\mathbb{Z}G)$ of rank one such that $\sigma_k(G) = [P] + T_G$. By Lemma 4.15, the composition

$$\text{Aut}(G) \xrightarrow{\psi_k} (\mathbb{Z}/|G|)^\times \xrightarrow{S} T_G \leq C(\mathbb{Z}G)$$

is given by $S \circ \psi_k: \theta \mapsto [P] - [P_\theta]$ which is well defined since θ gives a well-defined action on $C(\mathbb{Z}G)$. By Lemma 6.1, $(\mathbb{Z}G)_\theta \cong \mathbb{Z}G$ and so the composition is trivial in the case where $\sigma_k(G) = 0$. \square

We say that a finite group G has *weak cancellation* if every stably free Swan module is free. The following was asked by Dyer [1976, page 266] and later appeared in Wall's problems list [1979b, Problem A4].

Question 7.2 *Does there exist G with periodic cohomology and $r \in (\mathbb{Z}/|G|)^\times$ such that (I, r) is stably free but not free?*

This is equivalent to asking whether every group with periodic cohomology has weak cancellation and is still open, even for arbitrary finite groups. There are two important consequences that a negative answer to [Question 7.2](#) would have.

First, recall the following question from the introduction. Note that, if $(I, \psi_k(\theta))$ is free, then the action described in [Theorem B](#) has the simpler form $P \mapsto P_\theta$.

Question 7.3 *Does there exist G with k -periodic cohomology and $\theta \in \text{Aut}(G)$ for which $(I, \psi_k(\theta))$ is not free?*

It follows from [Proposition 7.1](#) that, if G has free period k and has weak cancellation, then $(I, \psi_k(\theta)) \cong \mathbb{Z}G$ for all $\theta \in \text{Aut}(G)$. In particular, if [Question 7.2](#) has a negative answer, then the only groups for which the action in [Theorem B](#) might not have the form $P \mapsto P_\theta$ are the groups with $\sigma_k(G) \neq 0$.

Second, consider the following:

Question 7.4 *Let $n \geq 2$, let G be finite and let X and Y be finite (G, n) -complexes with $\chi(X) = \chi(Y)$. Then $X \vee rS^n \simeq Y \vee rS^n$ for some r . Does $r = 1$ always work?*

This is equivalent to asking whether $\text{HT}(G, n)$ is a fork when G is finite. The case where n is even was proven by Browning [1978], and also follows by combining [Corollary 4.7](#) and [Proposition 5.1](#). When n is odd, this is known to hold provided G does not have k -periodic cohomology for any $k \mid n + 1$. If G has k -periodic cohomology for $k \mid n + 1$, then this holds provided G has weak cancellation (see [Dyer 1976, pages 276–277]). In particular, if [Question 7.2](#) has a negative answer, then [Question 7.4](#) has an affirmative answer. Note that the corresponding question for infinite groups is also still open (see [Nicholson 2021c, Problem B2]).

8 Milnor squares and the classification of projective modules

Given the observations in the previous section, the primary obstacle to computing sufficiently interesting examples of $\text{HT}(G, n)$ and $\text{PHT}(G, n)$ for our groups is the classification of projective $\mathbb{Z}G$ -modules.

One method to classify projective R -modules over a ring R is to relate this to the classification of projective modules over simpler rings using Milnor squares. In this section, we will present a refinement of the basic theory of Milnor squares which will also allow us to determine how a ring automorphism $\alpha \in \text{Aut}(R)$ acts on the class of projective R -modules. We will then apply these methods in [Section 9](#).

Suppose R and S are rings and $f: R \rightarrow S$ is a ring homomorphism. We can use this to turn S into an (S, R) -bimodule, with right-multiplication by $r \in R$ given by $x \cdot r = xf(r)$ for any $x \in S$. If M is an R -module, we can define the *extension of scalars* of M by f as the tensor product

$$f_{\#}(M) = S \otimes_R M$$

since S as a right R -module and M as a left R -module, and we consider this as a left S -module where left multiplication by $s \in S$ is given by $s \cdot (x \otimes m) = (sx) \otimes m$ for any $x \in S$ and $m \in M$. This comes equipped with maps of abelian groups

$$f_*: M \rightarrow f_*(M)$$

sending $m \mapsto 1 \otimes m$, and defines a covariant functor from R -modules to S -modules [Curtis and Reiner 1981, page 227]. It has the following basic properties which follow from the standard properties of tensor products such as associativity [Mac Lane 1963, page 145].

Lemma 8.1 *Let $f: R \rightarrow S$ and $g: S \rightarrow T$ be ring homomorphisms and let M and N be R -modules. Then*

- (i) $f_*(M \oplus N) \cong f_*(M) \oplus f_*(N)$,
- (ii) $f_*(R) \cong S$,
- (iii) $(g \circ f)_*(M) \cong (g_* \circ f_*)(M)$.

If $P(R)$ denotes the set of isomorphism classes of projective R -modules, then the first two properties show that f_* induces a map $f_*: P(R) \rightarrow P(S)$ which restricts to each stable class.

Recall that, if R, R_1, R_2 and R_0 are rings, then a pullback diagram

$$\mathcal{R} = \begin{array}{ccc} R & \xrightarrow{i_2} & R_2 \\ \downarrow i_1 & & \downarrow j_2 \\ R_1 & \xrightarrow{j_1} & R_0 \end{array}$$

is a Milnor square if either j_1 or j_2 are surjective. If $P_1 \in P(R_1)$ and $P_2 \in P(R_2)$ are such that there is an R_0 -module isomorphism $h: (j_1)_*(P_1) \rightarrow (j_2)_*(P_2)$, then define

$$M(P_1, P_2, h) = \{(x, y) \in P_1 \times P_2 \mid h((j_1)_*(x)) = (j_2)_*(y)\} \leq P_1 \times P_2,$$

which is an R -module where multiplication by $r \in R$ is given by $r \cdot (x, y) = ((i_1)_*(r)x, (i_2)_*(r)y)$. It was shown by Milnor that $M(P_1, P_2, h)$ is projective [Milnor 1971, Theorem 2.1]. Let $\text{Aut}_R(P)$ denote the set of R -module automorphisms of an R -module P . The main result on Milnor squares is as follows. This is a consequence of the results in [Milnor 1971, Section 2] and the precise statement can be found in [Swan 1980, Proposition 4.1].

Theorem 8.2 *Suppose \mathcal{R} is a Milnor square and $P_i \in P(R_i)$ for $i = 0, 1, 2$ are such that*

$$P_0 \cong (j_1)_*(P_1) \cong (j_2)_*(P_2)$$

as R_0 -modules. Then there is a one-to-one correspondence

$$\text{Aut}_{R_1}(P_1) \backslash \text{Aut}_{R_0}(P_0) / \text{Aut}_{R_2}(P_2) \leftrightarrow \{P \in P(R) \mid (i_1)_*(P) \cong P_1, (i_2)_*(P) \cong P_2\}$$

given by sending a coset $[h]$ to $M(P_1, P_2, h)$ for any representative h .

Now suppose $\alpha \in \text{Aut}(R)$. If M is an R -module, define M_α as the R -module whose abelian group is that of M but whose R -action is given by $r \cdot m = \alpha(r)m$ for $r \in R$ and $m \in M$. For example, if $R = \mathbb{Z}G$, then $\theta \in \text{Aut}(G)$ induces a map $\theta \in \text{Aut}(\mathbb{Z}G)$ and M_θ coincides with the definition given earlier.

This is a special case of restriction of scalars, but can also be viewed as a part of extension of scalars as follows.

Lemma 8.3 *Let R be a ring and let $\alpha \in \text{Aut}(R)$. If M is an R -module, then there is an isomorphism of R -modules*

$$\psi: M_\alpha \rightarrow (\alpha^{-1})_\#(M)$$

given by $m \mapsto 1 \otimes m$.

From this, it is clear that this action has basic properties which are analogous to Lemma 6.1. The following is then immediate by combining Lemmas 8.1 and 8.3.

Corollary 8.4 *Suppose $f: R \rightarrow S$ is a ring homomorphism and $\alpha \in \text{Aut}(R)$ and $\beta \in \text{Aut}(S)$ are such that $f \circ \alpha = \beta \circ f$. If M is an R -module, then*

$$f_\#(M_\alpha) \cong f_\#(M)_\beta.$$

We can turn the set of Milnor squares into a category with morphisms defined as follows. If \mathcal{R} and \mathcal{R}' are Milnor squares, then a morphism is a quadruple

$$\hat{\alpha} = (\alpha, \alpha_1, \alpha_2, \alpha_0): \mathcal{R} \rightarrow \mathcal{R}'$$

where $\alpha: R \rightarrow R'$ and $\alpha_i: R_i \rightarrow R'_i$ such that there is a commutative diagram

$$\begin{array}{ccccc}
 R & \xrightarrow{\quad} & R_2 & & \\
 \downarrow & \searrow \alpha & \downarrow & \searrow \alpha_2 & \\
 & & R & \xrightarrow{\quad} & R_2 \\
 & & \downarrow & & \downarrow \\
 R_1 & \xrightarrow{\quad} & R_0 & & \\
 \downarrow & \searrow \alpha_1 & \downarrow & \searrow \alpha_0 & \\
 & & R_1 & \xrightarrow{\quad} & R_0
 \end{array}$$

Let $\text{Aut}(\mathcal{R})$ denote the set of automorphisms of a Milnor square \mathcal{R} , i.e. the set of isomorphisms $\hat{\alpha}: \mathcal{R} \rightarrow \mathcal{R}$.

Lemma 8.5 *Let \mathcal{R} be a Milnor square and let $P_1 \in P(R_1)$ and $P_2 \in P(R_2)$ be such that there is an R_0 -module isomorphism $h: (j_1)_\#(P_1) \rightarrow (j_2)_\#(P_2)$. If $\hat{\alpha} = (\alpha, \alpha_1, \alpha_2, \alpha_0) \in \text{Aut}(\mathcal{R})$, then*

$$M(P_1, P_2, h)_\alpha \cong M((P_1)_{\alpha_1}, (P_2)_{\alpha_2}, h)$$

where, on the right, we view h as a map $h: (j_1)_\#(P_1)_{\alpha_0} \rightarrow (j_2)_\#(P_2)_{\alpha_0}$.

Proof Let $P = M(P_1, P_2, h)$ so that, by [Theorem 8.2](#), $(i_1)_\#(P) \cong P_1$ and $(i_2)_\#(P) \cong P_2$. It is easy to see directly that the natural map

$$M((i_1)_\#(P), (i_2)_\#(P), h) \rightarrow M((i_1)_\#(P_\alpha), (i_2)_\#(P_\alpha), h)$$

is an isomorphism. We are then done by applying [Corollary 8.4](#). □

This has the following simplification when P_1 and P_2 are free of rank one. Here we will use the identification $\text{Aut}_{R_0}(R_0) \cong R_0^\times$ which sends $h: R_0 \rightarrow R_0$ to $h(1) \in R_0^\times$.

Lemma 8.6 *Let \mathcal{R} be a Milnor square and let $u \in R_0^\times$. If $\hat{\alpha} = (\alpha, \alpha_1, \alpha_2, \alpha_0) \in \text{Aut}(\mathcal{R})$, then*

$$M(R_1, R_2, u)_\alpha \cong M(R_1, R_2, \alpha_0^{-1}(u)).$$

Proof Fix identifications $\psi_i: (j_i)_\#(R_i) \rightarrow R_0$ and let $h: (j_1)_\#(R_1) \rightarrow (j_1)_\#(R_1)$ be such that

$$(\psi_2 \circ h \circ \psi_1^{-1})(1) = u \in R_0^\times.$$

By [Lemma 8.5](#),

$$M(R_1, R_2, h)_\alpha \cong M((R_1)_{\alpha_1}, (R_2)_{\alpha_2}, h)$$

where $h: ((j_1)_\#(R_1))_{\alpha_0} \rightarrow ((j_1)_\#(R_1))_{\alpha_0}$ coincides with h as a map of abelian groups. For $i = 0, 1, 2$, let $c_i: R_i \rightarrow (R_i)_{\alpha_i}$ be the isomorphism which sends $1 \mapsto 1$. Then it is easy to see that

$$\begin{array}{ccc} (j_i)_\#(R_i) & \xrightarrow{1 \otimes c_i} & ((j_i)_\#((R_i)_{\alpha_i})) & \xrightarrow{f} & ((j_i)_\#(R_i))_{\alpha_0} \\ \downarrow \psi_i & & & & \downarrow \psi_i \\ R_0 & \xrightarrow{c_0} & & & (R_0)_{\alpha_0} \end{array}$$

commutes for $i = 1, 2$, where $f: (j_i)_\#((R_i)_{\alpha_i}) \rightarrow ((j_i)_\#(R_i))_{\alpha_0}$ is the isomorphism coming from [Corollary 8.4](#). Using the isomorphisms c_i for $i = 1, 2$, we get that

$$M((R_1)_{\alpha_1}, (R_2)_{\alpha_2}, h) \cong M(R_1, R_2, h_0)$$

where $h_0: (j_1)_\#(R_1) \rightarrow (j_2)_\#(R_2)$ induces $h: ((j_1)_\#(R_1))_{\alpha_0} \rightarrow ((j_1)_\#(R_1))_{\alpha_0}$ via $f \circ (1 \otimes c_i)$. Let $u_0 = (\psi_2 \circ h_0 \circ \psi_1^{-1})(1) \in R_0^\times$. Then, since the above diagram commutes, we get the commutative diagram

$$\begin{array}{ccc} R_0 & \xrightarrow{\psi_2 \circ h_0 \circ \psi_1^{-1}} & R_0 & & 1 & \longrightarrow & u_0 \\ \downarrow c_0 & & \downarrow c_0 & & \downarrow & & \downarrow \\ (R_0)_{\alpha_0} & \xrightarrow{\psi_2 \circ h_0 \circ \psi_1^{-1}} & (R_0)_{\alpha_0} & & 1 & \longrightarrow & \alpha_0(u_0) \end{array}$$

which implies that $u = \alpha_0(u_0)$ and so $u_0 = \alpha_0^{-1}(u)$, as required. □

If \mathcal{R} is a Milnor square, we say that $\alpha \in \text{Aut}(R)$ extends across \mathcal{R} if there exists $\hat{\alpha} = (\alpha, \alpha_1, \alpha_2, \alpha_0) \in \text{Aut}(\mathcal{R})$. The following gives conditions under which this induced map is unique.

Lemma 8.7 *Let \mathcal{R} be a pullback square with all maps surjective. If $\alpha \in \text{Aut}(R)$ extends across \mathcal{R} , then it does so uniquely. That is, there exist unique maps α_1, α_2 and α_0 for which $\hat{\alpha} = (\alpha, \alpha_1, \alpha_2, \alpha_0) \in \text{Aut}(\mathcal{R})$.*

Proof This follows from the simple observation that, if $f: R \twoheadrightarrow S$ is a surjective ring homomorphism and $\alpha: R \rightarrow R$ and $\beta_1, \beta_2: S \rightarrow S$ are ring homomorphisms such that $f \circ \alpha = \beta_i \circ f$ for $i = 1, 2$, then $\beta_1 = \beta_2$. To see this, note that the conditions imply that $(\beta_1 - \beta_2) \circ f = 0$ and so $\beta_1 = \beta_2$ on $\text{Im}(f)$. Since f is surjective, $\text{Im}(f) = S$ and so $\beta_1 = \beta_2$. \square

We conclude this section with the following result which is a consequence of [Theorem 8.2](#) and [Lemmas 8.6](#) and [8.7](#).

Proposition 8.8 *Let \mathcal{R} be a pullback square with all maps surjective and such that every $\alpha \in \text{Aut}(R)$ extends across \mathcal{R} . Then there is a one-to-one correspondence*

$$R_1^\times \backslash (R_0^\times / \text{Aut}(R)) / R_2^\times \leftrightarrow \{P \in P(R) : (i_1)_\#(P) \cong R_1, (i_2)_\#(P) \cong R_2\} / \text{Aut}(R)$$

where $\alpha \in \text{Aut}(R)$ acts on R_0^\times by sending $r \mapsto \alpha_0^{-1}(r)$ for $r \in R_0^\times$ and where $\alpha_0 \in \text{Aut}(R_0)$ is the unique automorphism such that $\hat{\alpha} = (\alpha, \alpha_1, \alpha_2, \alpha_0) \in \text{Aut}(\mathcal{R})$.

9 Example: quaternion groups

The aim of this section is to illustrate how [Theorems A](#) and [B](#) can be combined with the known techniques to classify projective $\mathbb{Z}G$ -modules to obtain a detailed classification of finite (G, n) -complexes up to homotopy equivalence.

For $k \geq 2$, recall that the quaternion group of order $4k$ has presentation

$$Q_{4k} = \langle x, y \mid x^k = y^2, yxy^{-1} = x^{-1} \rangle.$$

It is a finite 3-manifold group and so has free period 4. For $n \geq 2$ even, [Theorem A](#) and [Proposition 5.1](#) imply that $\text{PHT}(Q_{4k}, n) \cong [\mathbb{Z}Q_{4k}] = \bigcup_{r \geq 1} \text{SF}_r(\mathbb{Z}Q_{4k})$ where $\text{SF}_r(\mathbb{Z}Q_{4k})$ is the set of stably free $\mathbb{Z}Q_{4k}$ -modules of rank $r \geq 1$.

Since stably free $\mathbb{Z}G$ -modules of rank ≥ 2 are free for G finite [[Swan 1960a](#)] (or since $\text{PHT}(G, n)$ is a fork by [Corollary 4.7](#)), it remains to compute $\text{SF}_1(\mathbb{Z}Q_{4k})$. This was completed by Swan [[1983](#), [Theorem III](#)] for $k \leq 9$. For $k \leq 7$, he showed that $|\text{SF}_1(\mathbb{Z}Q_{4k})| = 1$ for $2 \leq k \leq 5$, $|\text{SF}_1(\mathbb{Z}Q_{24})| = 3$ and $|\text{SF}_1(\mathbb{Z}Q_{28})| = 2$. It also follows from his classification that $\mathbb{Z}Q_{4k}$ has weak cancellation in all these cases and so the action of $\theta \in \text{Aut}(Q_{4k})$ on $[\mathbb{Z}Q_{4k}]$ sends $P \mapsto P_\theta$ (see [Section 7](#)).

In the case Q_{28} , the action of $\text{Aut}(Q_{28})$ on $[\mathbb{Z}Q_{28}]$ is trivial since $(\mathbb{Z}Q_{28})_\theta \cong \mathbb{Z}Q_{28}$ for all $\theta \in \text{Aut}(Q_{28})$ and so this must also hold for the nonfree stably free module also. The main result of this section will be to compute the action in the case Q_{24} .

Theorem 9.1 *$\text{Aut}(Q_{24})$ acts nontrivially on $[\mathbb{Z}Q_{24}]$. More specifically, we have $|\text{SF}_1(\mathbb{Z}Q_{24})| = 3$ and $|\text{SF}_1(\mathbb{Z}Q_{24})/\text{Aut}(Q_{24})| = 2$.*

G	Q_8	Q_{12}	Q_{16}	Q_{20}	Q_{24}	Q_{28}
$\text{PHT}(G, n)$	•	•	•	•	•••	••
$\text{HT}(G, n)$	•	•	•	•	••	••

Table 1: Minimal complexes for any n even with $n \neq 2$.

All of this is summarised in Table 1, which gives the structure of $\text{PHT}(G, n)$ and $\text{HT}(G, n)$ when $n \neq 2$ is even. These graded trees are both forks by Corollary 4.7 and each dot represents a finite (G, n) -complex at the minimal level.

Remark 9.2 This also holds in the case $n = 2$ provided G has the D2 property. This holds trivially in the cases Q_8, Q_{12}, Q_{16} and Q_{20} , and is otherwise only known to be true in the case Q_{28} by [Nicholson 2021b, Theorem 7.7] using the presentation of Mannan and Popiel [2021].

We will now proceed to the proof of Theorem 9.1. First let x and y be generators for Q_{24} in the presentation given above. Let $\Lambda = \mathbb{Z}Q_{24}/(x^6 + 1)$ and note that the quotient map $f: \mathbb{Z}Q_{24} \twoheadrightarrow \Lambda$ induces a map

$$f_{\#}: \text{SF}_1(\mathbb{Z}Q_{24}) \rightarrow \text{SF}_1(\Lambda)$$

by Lemma 8.1. This is a bijection by the proof of [Swan 1983, Theorem 11.14].

Now note that the factorisation $x^6 + 1 = (x^2 + 1)(x^4 - x^2 + 1)$ implies that the ideals $I = (x^2 + 1)$ and $J = (x^4 - x^2 + 1)$ have $I \cap J = (x^6 + 1)$ and $I + J = (3, x^2 + 1)$. It follows from [Curtis and Reiner 1987, Example 42.3] that we have a pullback diagram

$$\begin{array}{ccc} \Lambda & \longrightarrow & \mathbb{Z}Q_{24}/(x^4 - x^2 + 1) \\ \downarrow & & \downarrow \\ \mathbb{Z}Q_{24}/(x^2 + 1) & \longrightarrow & \mathbb{F}_3Q_{24}/(x^2 + 1) \end{array}$$

which is a Milnor square since all maps are surjective.

For a field \mathbb{F} , let $\mathbb{H}_{\mathbb{F}} = \mathbb{F}[i, j]$ denote the quaternions over \mathbb{F} and we define $\mathbb{H}_{\mathbb{Z}} = \mathbb{Z}[i, j]$ and $\mathbb{Z}[\zeta_{12}, j]$ to be subrings of $\mathbb{H}_{\mathbb{R}}$, where $\zeta_{12} = e^{2\pi i/12}$ is the 12th root of unity in the i direction. It is straightforward to check that there are isomorphisms of rings

$$\begin{aligned} \phi_1: \mathbb{H}_{\mathbb{Z}} &\rightarrow \mathbb{Z}Q_{24}/(x^2 + 1), & i &\mapsto x, j \mapsto y \\ \phi_2: \mathbb{Z}[\zeta_{12}, j] &\rightarrow \mathbb{Z}Q_{24}/(x^4 - x^2 + 1), & \zeta_{12} &\mapsto x, j \mapsto y. \end{aligned}$$

Using this, we can rewrite the Milnor square above as

$$\mathcal{R} = \begin{array}{ccc} \Lambda & \xrightarrow{i_2} & \mathbb{Z}[\zeta_{12}, j] \\ \downarrow i_1 & & \downarrow j_2 \\ \mathbb{H}_{\mathbb{Z}} & \xrightarrow{j_1} & \mathbb{H}_{\mathbb{F}_3} \end{array} \quad \begin{array}{ccc} x, y & \longmapsto & \zeta_{12}, j \\ \downarrow & & \downarrow \\ i, j & \longmapsto & i, j \end{array}$$

By [Swan 1983, Lemma 8.14], the induced map $(i_2)_*: C(\Lambda) \rightarrow C(\mathbb{Z}[\zeta_{12}, j])$ is an isomorphism. It also follows from [Swan 1983, page 84] that the rings $\mathbb{H}_{\mathbb{Z}}$ and $\mathbb{Z}[\zeta_{12}, j]$ both have stably free cancellation, i.e. that every stably free module is free. It follows easily that

$$SF_1(\Lambda) = \{P \in P(\Lambda) : (i_1)_\#(P) \cong \mathbb{H}_{\mathbb{Z}}, (i_2)_\#(P) \cong \mathbb{Z}[\zeta_{12}, j]\}.$$

In particular, by combining with Theorem 8.2, we get that there is a bijection

$$SF_1(\Lambda) \leftrightarrow \mathbb{H}_{\mathbb{Z}}^\times \backslash \mathbb{H}_{\mathbb{F}_3}^\times / \mathbb{Z}[\zeta_{12}, j]^\times.$$

Lemma 9.3 $\mathbb{H}_{\mathbb{Z}}^\times \backslash \mathbb{H}_{\mathbb{F}_3}^\times / \mathbb{Z}[\zeta_{12}, j]^\times = \{[1], [1 + j], [1 + k]\}.$

Proof If $N : \mathbb{H}_{\mathbb{F}_3} \rightarrow \mathbb{F}_3$ is the norm, then $\mathbb{H}_{\mathbb{F}_3}^\times = N^{-1}(\pm 1)$. Now note that $\mathbb{H}_{\mathbb{Z}}^\times = \{\pm 1, \pm i, \pm j, \pm k\}$, and it is easy to check that

$$\mathbb{H}_{\mathbb{Z}}^\times \backslash \mathbb{H}_{\mathbb{F}_3}^\times = \{[1], [1 + i], [1 + j], [1 + k], [1 + i + j + k], [1 - i - j - k]\}.$$

By [Magurn et al. 1983, Lemma 7.5(b)], $\mathbb{Z}[\zeta_{12}, j]^\times = \mathbb{Z}[\zeta_{12}]^\times \cdot \langle j \rangle$ and so it remains to determine

$$\text{Im}(\mathbb{Z}[\zeta_{12}, j]^\times \rightarrow \mathbb{H}_{\mathbb{Z}}^\times \backslash \mathbb{H}_{\mathbb{F}_3}^\times) = \text{Im}(\mathbb{Z}[\zeta_{12}]^\times \rightarrow \mathbb{H}_{\mathbb{Z}}^\times \backslash \mathbb{H}_{\mathbb{F}_3}^\times) \subseteq \{[1], [1 + i]\},$$

where the last inclusion follows since $\zeta_{12} \mapsto i$ and $\mathbb{H}_{\mathbb{Z}}^\times \backslash \langle 1, i \rangle = \{[1], [1 + i]\}.$

Consider the n^{th} cyclotomic polynomial

$$\Phi_n(x) = \prod_{k \in \mathbb{Z}_n^\times} (x - \zeta_n^k).$$

It is well known, and can be shown using Möbius inversion, that $\Phi_n(1) = 1$ if n is not a prime power. In particular, $\Phi_{12}(1) = 1$ and this implies that $1 - \zeta_{12} \in \mathbb{Z}[\zeta_{12}]^\times$. Hence

$$[1 + i] = [1 - i] \in \text{Im}(\mathbb{Z}[\zeta_{12}]^\times \rightarrow \mathbb{H}_{\mathbb{Z}}^\times \backslash \mathbb{H}_{\mathbb{F}_3}^\times).$$

The result then follows since

$$j(1 + i + j + k)(1 + i) = 1 + k, \quad -j(1 - i - j - k)(1 + i) = 1 + j$$

implies that $[1 + j] = [1 - i - j - k]$ and $[1 + k] = [1 + i + j + k]$ in $\mathbb{H}_{\mathbb{Z}}^\times \backslash \mathbb{H}_{\mathbb{F}_3}^\times / \mathbb{Z}[\zeta_{12}, j]^\times$. □

This implies that $|SF_1(\mathbb{Z}Q_{24})| = 3$, which recovers the result of Swan. In order to determine the action of $\text{Aut}(Q_{24})$ on $SF_1(\mathbb{Z}Q_{24})$, first recall from Section 7 that

$$\text{Aut}(Q_{24}) = \{\theta_{a,b} : x \mapsto x^a, y \mapsto x^b y \mid a \in (\mathbb{Z}/12)^\times, b \in \mathbb{Z}/12\}.$$

If \mathcal{R} denotes the Milnor square defined above, then the following is easy to check.

Lemma 9.4 *If $a \in (\mathbb{Z}/12)^\times$ and $b \in \mathbb{Z}/12$, then $\theta_{a,b} \in \text{Aut}(Q_{24})$ extends to a Milnor square automorphism*

$$\hat{\theta}_{a,b} = (\theta'_{a,b}, \theta^1_{a,b}, \theta^2_{a,b}, \bar{\theta}_{a,b}) \in \text{Aut}(\mathcal{R})$$

where, for $a = 2a_0 + 1$, the maps are defined as follows:

(i) $\theta'_{a,b} \in \text{Aut}(\mathbb{Z}Q_{24}/(x^6 + 1))$ is given by $x \mapsto x^a$ and $y \mapsto x^b y$.

(ii) $\theta^1_{a,b} \in \text{Aut}(\mathbb{H}_{\mathbb{Z}})$ and $\bar{\theta}_{a,b} \in \text{Aut}(\mathbb{H}_{\mathbb{F}_3})$ are each given by

$$i \mapsto i^a = (-1)^{a_0} i, \quad j \mapsto j^b = \begin{cases} (-1)^{b_0} j & \text{if } b = 2b_0 + 1, \\ (-1)^{b_0} k & \text{if } b = 2b_0. \end{cases}$$

(iii) $\theta^2_{a,b} \in \text{Aut}(\mathbb{Z}[\zeta_{12}, j])$ is given by $\zeta_{12} \mapsto \zeta_{12}^a$ and $j \mapsto \zeta_{12}^b j$.

Since \mathcal{R} is a pullback square with all maps surjective, we can now apply [Proposition 8.8](#). By combining with [Lemma 9.3](#), this implies that there is a bijection

$$\text{SF}_1(\mathbb{Z}Q_{24})/\text{Aut}(Q_{24}) \leftrightarrow \{[1], [1 + j], [1 + k]\}/\text{Aut}(Q_{24})$$

where $\theta_{a,b} \in \text{Aut}(Q_{24})$ acts on the double cosets via the action described in [Lemma 9.4](#). In particular,

$$\bar{\theta}_{a,b}([1 + j]) = \begin{cases} [1 + (-1)^{b_0} j] = [1 + j] & \text{if } b = 2b_0 + 1, \\ [1 + (-1)^{b_0} k] = [1 + k] & \text{if } b = 2b_0, \end{cases}$$

and so $\bar{\theta}_{a,b}$ acts nontrivially when b is even. Hence $|\text{SF}_1(\mathbb{Z}Q_{24})/\text{Aut}(Q_{24})| = 2$. This completes the proof of [Theorem 9.1](#).

References

- [Bestvina and Brady 1997] **M Bestvina, N Brady**, *Morse theory and finiteness properties of groups*, Invent. Math. 129 (1997) 445–470 [MR](#) [Zbl](#)
- [Bieri 1976] **R Bieri**, *Homological dimension of discrete groups*, Queen Mary College, London (1976) [MR](#) [Zbl](#)
- [Bokor et al. 2021] **I Bokor, D Crowley, S Friedl, F Hebestreit, D Kasproski, M Land, J Nicholson**, *Connected sum decompositions of high-dimensional manifolds*, from “2019–20 MATRIX annals” (DR Wood, J de Gier, CE Praeger, T Tao, editors), MATRIX Book Ser. 4, Springer (2021) 5–30 [MR](#) [Zbl](#)
- [Brown 1982] **KS Brown**, *Cohomology of groups*, Grad. Texts in Math. 87, Springer (1982) [MR](#) [Zbl](#)
- [Browning 1978] **WJ Browning**, *Homotopy types of certain finite CW-complexes with finite fundamental group*, PhD thesis, Cornell University (1978) Available at <https://www.proquest.com/docview/302945517>
- [Browning 1979] **WJ Browning**, *Finite CW-complexes of cohomological dimension 2 with finite abelian π_1* , unpublished (1979)
- [Cartan and Eilenberg 1956] **H Cartan, S Eilenberg**, *Homological algebra*, Princeton Landmarks Math. Phys. 19, Princeton Univ. Press (1956) [MR](#) [Zbl](#)
- [Cockcroft and Swan 1961] **WH Cockcroft, RG Swan**, *On the homotopy type of certain two-dimensional complexes*, Proc. Lond. Math. Soc. 11 (1961) 194–202 [MR](#) [Zbl](#)

- [Cohen 1977] **J M Cohen**, *Complexes dominated by a 2-complex*, *Topology* 16 (1977) 409–415 [MR](#) [Zbl](#)
- [Conway et al. 2021] **A Conway, D Crowley, M Powell, J Sixt**, *Stably diffeomorphic manifolds and modified surgery obstructions*, preprint (2021) [arXiv 2109.05632](#)
- [Conway et al. 2023] **A Conway, D Crowley, M Powell, J Sixt**, *Simply connected manifolds with large homotopy stable classes*, *J. Aust. Math. Soc.* 115 (2023) 172–203 [MR](#) [Zbl](#)
- [Curtis and Reiner 1981] **C W Curtis, I Reiner**, *Methods of representation theory, I: With applications to finite groups and orders*, Wiley, New York (1981) [MR](#) [Zbl](#)
- [Curtis and Reiner 1987] **C W Curtis, I Reiner**, *Methods of representation theory, II: With applications to finite groups and orders*, Wiley, New York (1987) [MR](#) [Zbl](#)
- [Davis 1983] **J F Davis**, *A remark on: ‘Homotopy equivalences and free modules’ [Topology 21 (1982) 91–99] by S Plotnick*, *Topology* 22 (1983) 487–488 [MR](#) [Zbl](#)
- [Dyer 1976] **M N Dyer**, *Homotopy classification of (π, m) -complexes*, *J. Pure Appl. Algebra* 7 (1976) 249–282 [MR](#) [Zbl](#)
- [Dyer 1979] **M N Dyer**, *Nonminimal roots in homotopy trees*, *Pacific J. Math.* 80 (1979) 371–380 [MR](#) [Zbl](#)
- [Dyer and Sieradski 1973] **M N Dyer, A J Sieradski**, *A homotopy classification of 2-complexes with finite cyclic fundamental group*, *Bull. Amer. Math. Soc.* 79 (1973) 75–77 [MR](#) [Zbl](#)
- [Dyer and Sieradski 1975] **M N Dyer, A J Sieradski**, *Trees of homotopy types of 2-dimensional CW complexes, II*, *Trans. Amer. Math. Soc.* 205 (1975) 115–125 [MR](#) [Zbl](#)
- [Fröhlich et al. 1974] **A Fröhlich, I Reiner, S Ullom**, *Class groups and Picard groups of orders*, *Proc. Lond. Math. Soc.* 29 (1974) 405–434 [MR](#) [Zbl](#)
- [Golasiński and Gonçalves 2004] **M Golasiński, D L Gonçalves**, *Spherical space forms: homotopy types and self-equivalences*, from “Categorical decomposition techniques in algebraic topology” (G Arone, J Hubbuck, R Levi, M Weiss, editors), *Progr. Math.* 215, Birkhäuser, Basel (2004) 153–165 [MR](#) [Zbl](#)
- [Hambleton 2019] **I Hambleton**, *Two remarks on Wall’s D_2 problem*, *Math. Proc. Cambridge Philos. Soc.* 167 (2019) 361–368 [MR](#) [Zbl](#)
- [Hambleton and Kreck 1993] **I Hambleton, M Kreck**, *Cancellation of lattices and finite two-complexes*, *J. Reine Angew. Math.* 442 (1993) 91–109 [MR](#) [Zbl](#)
- [Hambleton et al. 2013] **I Hambleton, S Pamuk, E Yalçın**, *Equivariant CW-complexes and the orbit category*, *Comment. Math. Helv.* 88 (2013) 369–425 [MR](#) [Zbl](#)
- [Jacobinski 1968] **H Jacobinski**, *Genera and decompositions of lattices over orders*, *Acta Math.* 121 (1968) 1–29 [MR](#) [Zbl](#)
- [Johnson 2002] **F E A Johnson**, *Explicit homotopy equivalences in dimension two*, *Math. Proc. Cambridge Philos. Soc.* 133 (2002) 411–430 [MR](#) [Zbl](#)
- [Johnson 2003] **F E A Johnson**, *Stable modules and the $D(2)$ -problem*, *Lond. Math. Soc. Lect. Note Ser.* 301, Cambridge Univ. Press (2003) [MR](#) [Zbl](#)
- [Kreck and Schafer 1984] **M Kreck, J A Schafer**, *Classification and stable classification of manifolds: some examples*, *Comment. Math. Helv.* 59 (1984) 12–38 [MR](#) [Zbl](#)
- [Linnell 1993] **P A Linnell**, *Minimal free resolutions and (G, n) -complexes for finite abelian groups*, *Proc. Lond. Math. Soc.* 66 (1993) 303–326 [MR](#) [Zbl](#)

- [Mac Lane 1963] **S Mac Lane**, *Homology*, Grundle. Math. Wissen. 114, Springer (1963) [MR](#) [Zbl](#)
- [Magurn et al. 1983] **B Magurn**, **R Oliver**, **L Vaserstein**, *Units in Whitehead groups of finite groups*, J. Algebra 84 (1983) 324–360 [MR](#) [Zbl](#)
- [Mannan 2007] **W H Mannan**, *The $D(2)$ property for D_8* , Algebr. Geom. Topol. 7 (2007) 517–528 [MR](#) [Zbl](#)
- [Mannan and Popiel 2021] **W H Mannan**, **T Popiel**, *An exotic presentation of Q_{28}* , Algebr. Geom. Topol. 21 (2021) 2065–2084 [MR](#) [Zbl](#)
- [Metzler 1976] **W Metzler**, *Über den Homotopietyp zweidimensionaler CW-Komplexe und Elementartransformationen bei Darstellungen von Gruppen durch Erzeugende und definierende Relationen*, J. Reine Angew. Math. 285 (1976) 7–23 [MR](#) [Zbl](#)
- [Milgram 1985] **R J Milgram**, *Evaluating the Swan finiteness obstruction for periodic groups*, from “Algebraic and geometric topology” (A Ranicki, N Levitt, F Quinn, editors), Lecture Notes in Math. 1126, Springer (1985) 127–158 [MR](#) [Zbl](#)
- [Milnor 1971] **J Milnor**, *Introduction to algebraic K -theory*, Ann. of Math. Ser. 72, Princeton Univ. Press (1971) [MR](#) [Zbl](#)
- [Nicholson 2020] **J Nicholson**, *Cancellation for (G, n) -complexes and the Swan finiteness obstruction*, preprint (2020) [arXiv 2005.01664](#) To appear in Int. Math. Res. Not.
- [Nicholson 2021a] **J Nicholson**, *A cancellation theorem for modules over integral group rings*, Math. Proc. Cambridge Philos. Soc. 171 (2021) 317–327 [MR](#) [Zbl](#)
- [Nicholson 2021b] **J Nicholson**, *On CW-complexes over groups with periodic cohomology*, Trans. Amer. Math. Soc. 374 (2021) 6531–6557 [MR](#) [Zbl](#)
- [Nicholson 2021c] **J Nicholson**, *Stably free modules and the unstable classification of 2-complexes*, preprint (2021) [arXiv 2108.02220](#)
- [Sieradski 1977] **A J Sieradski**, *A semigroup of simple homotopy types*, Math. Z. 153 (1977) 135–148 [MR](#) [Zbl](#)
- [Sieradski and Dyer 1979] **A J Sieradski**, **M N Dyer**, *Distinguishing arithmetic for certain stably isomorphic modules*, J. Pure Appl. Algebra 15 (1979) 199–217 [MR](#) [Zbl](#)
- [Stallings 1963] **J Stallings**, *A finitely presented group whose 3-dimensional integral homology is not finitely generated*, Amer. J. Math. 85 (1963) 541–543 [MR](#) [Zbl](#)
- [Swan 1960a] **R G Swan**, *Induced representations and projective modules*, Ann. of Math. 71 (1960) 552–578 [MR](#) [Zbl](#)
- [Swan 1960b] **R G Swan**, *Periodic resolutions for finite groups*, Ann. of Math. 72 (1960) 267–291 [MR](#) [Zbl](#)
- [Swan 1965] **R G Swan**, *Minimal resolutions for finite groups*, Topology 4 (1965) 193–208 [MR](#) [Zbl](#)
- [Swan 1970] **R G Swan**, *K -theory of finite groups and orders*, Lect. Notes in Math. 149, Springer (1970) [MR](#) [Zbl](#)
- [Swan 1980] **R G Swan**, *Strong approximation and locally free modules*, from “Ring theory and algebra, III” (B R McDonald, editor), Lect. Notes Pure Appl. Math. 55, Dekker, New York (1980) 153–223 [MR](#) [Zbl](#)
- [Swan 1983] **R G Swan**, *Projective modules over binary polyhedral groups*, J. Reine Angew. Math. 342 (1983) 66–172 [MR](#) [Zbl](#)
- [Wall 1965] **C T C Wall**, *Finiteness conditions for CW-complexes*, Ann. of Math. 81 (1965) 56–69 [MR](#) [Zbl](#)
- [Wall 1979a] **C T C Wall**, *Periodic projective resolutions*, Proc. Lond. Math. Soc. 39 (1979) 509–553 [MR](#) [Zbl](#)

- [Wall 1979b] *List of problems*, from “Homological group theory” (C T C Wall, editor), Lond. Math. Soc. Lect. Note Ser. 36, Cambridge Univ. Press (1979) 369–394 [MR](#) [Zbl](#)
- [Weibel 1994] **C A Weibel**, *An introduction to homological algebra*, Cambridge Stud. Adv. Math. 38, Cambridge Univ. Press (1994) [MR](#) [Zbl](#)
- [Whitehead 1939] **J H C Whitehead**, *Simplicial spaces, nuclei and m -groups*, Proc. Lond. Math. Soc. 45 (1939) 243–327 [MR](#) [Zbl](#)

School of Mathematics and Statistics, University of Glasgow
Glasgow, United Kingdom

john.nicholson@glasgow.ac.uk

Received: 19 July 2022 Revised: 4 April 2023

ALGEBRAIC & GEOMETRIC TOPOLOGY

msp.org/agt

EDITORS

PRINCIPAL ACADEMIC EDITORS

John Etnyre
etnyre@math.gatech.edu
Georgia Institute of Technology

Kathryn Hess
kathryn.hess@epfl.ch
École Polytechnique Fédérale de Lausanne

BOARD OF EDITORS

Julie Bergner	University of Virginia jeb2md@eservices.virginia.edu	Robert Lipshitz	University of Oregon lipshitz@uoregon.edu
Steven Boyer	Université du Québec à Montréal cohf@math.rochester.edu	Norihiko Minami	Yamato University minami.norihiko@yamato-u.ac.jp
Tara E Brendle	University of Glasgow tara.brendle@glasgow.ac.uk	Andrés Navas	Universidad de Santiago de Chile andres.navas@usach.cl
Indira Chatterji	CNRS & Univ. Côte d'Azur (Nice) indira.chatterji@math.cnrs.fr	Thomas Nikolaus	University of Münster nikolaus@uni-muenster.de
Alexander Dranishnikov	University of Florida dranish@math.ufl.edu	Robert Oliver	Université Paris 13 bobol@math.univ-paris13.fr
Tobias Ekholm	Uppsala University, Sweden tobias.ekholm@math.uu.se	Jessica S Purcell	Monash University jessica.purcell@monash.edu
Mario Eudave-Muñoz	Univ. Nacional Autónoma de México mario@matem.unam.mx	Birgit Richter	Universität Hamburg birgit.richter@uni-hamburg.de
David Futer	Temple University dfuter@temple.edu	Jérôme Scherer	École Polytech. Féd. de Lausanne jerome.scherer@epfl.ch
John Greenlees	University of Warwick john.greenlees@warwick.ac.uk	Vesna Stojanoska	Univ. of Illinois at Urbana-Champaign vesna@illinois.edu
Ian Hambleton	McMaster University ian@math.mcmaster.ca	Zoltán Szabó	Princeton University szabo@math.princeton.edu
Matthew Hedden	Michigan State University mhedden@math.msu.edu	Maggy Tomova	University of Iowa maggy-tomova@uiowa.edu
Hans-Werner Henn	Université Louis Pasteur henn@math.u-strasbg.fr	Nathalie Wahl	University of Copenhagen wahl@math.ku.dk
Daniel Isaksen	Wayne State University isaksen@math.wayne.edu	Chris Wendl	Humboldt-Universität zu Berlin wendl@math.hu-berlin.de
Thomas Koberda	University of Virginia thomas.koberda@virginia.edu	Daniel T Wise	McGill University, Canada daniel.wise@mcgill.ca
Christine Lescop	Université Joseph Fourier lescop@ujf-grenoble.fr		

See inside back cover or msp.org/agt for submission instructions.

The subscription price for 2024 is US \$705/year for the electronic version, and \$1040/year (+\$70, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Algebraic & Geometric Topology is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [Current Mathematical Publications](#) and the [Science Citation Index](#).

Algebraic & Geometric Topology (ISSN 1472-2747 printed, 1472-2739 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840.

AGT peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**

nonprofit scientific publishing

<https://msp.org/>

© 2024 Mathematical Sciences Publishers

ALGEBRAIC & GEOMETRIC TOPOLOGY

Volume 24 Issue 4 (pages 1809–2387) 2024

Möbius structures, quasimetrics and completeness	1809
MERLIN INCERTI-MEDICI	
$\mathbb{Z}/p \times \mathbb{Z}/p$ actions on $S^n \times S^n$	1841
JIM FOWLER and COURTNEY THATCHER	
\mathbb{Z}_k -stratifolds	1863
ANDRÉS ÁNGEL, CARLOS SEGOVIA and ARLEY FERNANDO TORRES	
Relative systoles in hyperelliptic translation surfaces	1903
CORENTIN BOISSY and SLAVYANA GENINSKA	
Smooth singular complexes and diffeological principal bundles	1913
HIROSHI KIHARA	
Natural symmetries of secondary Hochschild homology	1953
DAVID AYALA, JOHN FRANCIS and ADAM HOWARD	
The shape of the filling-systole subspace in surface moduli space and critical points of the systole function	2011
YUE GAO	
Moduli spaces of geometric graphs	2039
MARA BELOTTI, ANTONIO LERARIO and ANDREW NEWMAN	
Classical shadows of stated skein representations at roots of unity	2091
JULIEN KORINMAN and ALEXANDRE QUESNEY	
Commensurators of thin normal subgroups and abelian quotients	2149
THOMAS KOBERDA and MAHAN MJ	
Pushouts of Dwyer maps are $(\infty, 1)$ -categorical	2171
PHILIP HACKNEY, VIKTORIYA OZORNOVA, EMILY RIEHL and MARTINA ROVELLI	
A variant of a Dwyer–Kan theorem for model categories	2185
BORIS CHORNY and DAVID WHITE	
Integral generalized equivariant cohomologies of weighted Grassmann orbifolds	2209
KOUSHIK BRAHMA and SOUMEN SARKAR	
Projective modules and the homotopy classification of (G, n) -complexes	2245
JOHN NICHOLSON	
Realization of Lie algebras of derivations and moduli spaces of some rational homotopy types	2285
YVES FÉLIX, MARIO FUENTES and ANICETO MURILLO	
On the positivity of twisted L^2 -torsion for 3-manifolds	2307
JIANRU DUAN	
An algebraic C_2 -equivariant Bézout theorem	2331
STEVEN R COSTENOBLE, THOMAS HUDSON and SEAN TILSON	
Topologically isotopic and smoothly inequivalent 2-spheres in simply connected 4-manifolds whose complement has a prescribed fundamental group	2351
RAFAEL TORRES	
Remarks on symplectic circle actions, torsion and loops	2367
MARCELO S ATALLAH	
Correction to the article Hopf ring structure on the mod p cohomology of symmetric groups	2385
LORENZO GUERRA	