

AG
T

*Algebraic & Geometric
Topology*

Volume 24 (2024)

Remarks on symplectic circle actions, torsion and loops

MARCELO S ATALLAH



Remarks on symplectic circle actions, torsion and loops

MARCELO S ATALLAH

We study loops of symplectic diffeomorphisms of closed symplectic manifolds. Our main result, which is valid for a large class of symplectic manifolds, shows that the flux of a symplectic loop vanishes whenever its orbits are contractible. As a consequence, we obtain a new vanishing result for the flux group and new instances where the presence of a fixed point of a symplectic circle action is a sufficient condition for it to be Hamiltonian. We also obtain applications to symplectic torsion; more precisely, nontrivial elements of $\text{Symp}_0(M, \omega)$ that have finite order.

53D22, 53D40, 57R17; 57S15, 57S17

1 Introduction

1.1 The flux group and the e -homomorphism

Let (M, ω) be a closed symplectic manifold and denote by $\widetilde{\text{Symp}}_0(M, \omega)$ the universal cover of the identity component $\text{Symp}_0(M, \omega)$ of the group of symplectic diffeomorphisms. The *flux homomorphism*

$$\widetilde{\text{Flux}}: \widetilde{\text{Symp}}_0(M, \omega) \rightarrow H^1(M; \mathbb{R})$$

is defined by assigning to each class $\tilde{\psi} \in \widetilde{\text{Symp}}_0(M, \omega)$ a cohomology class

$$\widetilde{\text{Flux}}(\tilde{\psi}) = \int_0^1 [l_{X^t} \omega] dt,$$

where X^t is the time-dependent vector field induced by a symplectic path $\{\psi_t\}$ representing $\tilde{\psi}$. In particular, if γ is a 1-cycle in M , we have that

$$(1) \quad \langle \widetilde{\text{Flux}}(\{\psi_t\}), [\gamma] \rangle = \int_{[0,1] \times \mathbb{R}/\mathbb{Z}} \alpha^* \omega,$$

where $\alpha: [0, 1] \times \mathbb{R}/\mathbb{Z} \rightarrow M$ is given by setting $\alpha(t, s) = \psi_t(\gamma(s))$. We shall often denote the flux of $\tilde{\psi}$ by $\widetilde{\text{Flux}}(\{\psi_t\})$. The *flux group* Γ is the image of the restriction of the flux homomorphism to $\pi_1(\text{Symp}_0(M, \omega)) \subset \widetilde{\text{Symp}}_0(M, \omega)$. The $\widetilde{\text{Flux}}$ map descends to a homomorphism

$$\text{Flux}: \text{Symp}_0(M, \omega) \rightarrow H^1(M; \mathbb{R}) / \Gamma,$$

whose kernel was shown by Banyaga [2] to be equal to $\text{Ham}(M, \omega)$. In particular, we have the exact sequence

$$1 \rightarrow \text{Ham}(M, \omega) \rightarrow \text{Symp}_0(M, \omega) \rightarrow H^1(M; \mathbb{R}) / \Gamma \rightarrow 1,$$

which implies that $\text{Ham}(M, \omega)$ coincides with $\text{Symp}_0(M, \omega)$ if and only if $H^1(M; \mathbb{R})$ vanishes. For a basepoint $x_0 \in M$, denote by

$$\text{ev}: \pi_1(\text{Symp}_0(M, \omega), \text{id}) \rightarrow \pi_1(M, x_0)$$

the *evaluation homomorphism* given by setting $\text{ev}(\{\psi_t\}) = \{\psi_t(x_0)\}$. The image of ev lies in the center of $\pi_1(M)$; see Polterovich [25]. Therefore, since M is connected, the evaluation maps for different choices of basepoints $x_0 \in M$ are identified. Hence, we write ev without making specific reference to x_0 . Observe that the evaluation map factors through the flux group Γ yielding the following commutative diagram:¹

$$(2) \quad \begin{array}{ccc} \pi_1(\text{Symp}_0(M, \omega)) & & \\ \widetilde{\text{Flux}} \downarrow & \searrow \text{ev} & \\ \Gamma & \xrightarrow{e} & \pi_1(M) \end{array}$$

To see that the e -homomorphism is well-defined note that if

$$\widetilde{\text{Flux}}(\{\phi_t\}) = \widetilde{\text{Flux}}(\{\psi_t\}),$$

then the loop $\{\phi_t \circ \psi_t^{-1}\}$ of symplectic diffeomorphisms can be homotoped to a Hamiltonian loop, which is known to have contractible orbits; see McDuff [16]. An important consequence of diagram (2) is that whenever e is injective, a symplectic loop with contractible orbits can be homotoped to a Hamiltonian loop. It is not hard to see that this is the case for (M, ω) symplectically aspherical: $[\omega]$ vanishes on the image $H_2^S(M; \mathbb{Z})$ of the Hurewicz map $\pi_2(M) \rightarrow H_2(M; \mathbb{Z})$. Indeed, if a symplectic loop $\{\psi_t\}$ has trivial evaluation and γ is any 1-cycle in M , then the torus $\alpha(T^2)$, where $\alpha(t, s) = \psi_t(\gamma(s))$, will have the symplectic area of a sphere. Equation (1) then implies that the flux of the loop $\{\psi_t\}$ vanishes.

The injectivity of the e -homomorphism has important consequences in the theory of symplectic circle actions discussed in Section 1.2. Furthermore, it allows one to deduce the vanishing of the Flux group in cases where the evaluation map is trivial. In particular, we have the following:

Proposition 1.1 *Let (M, ω) be a closed symplectic manifold with injective e -homomorphism, and suppose further that either $\chi(M) \neq 0$ or that $\pi_1(M)$ has finite center. Then the flux group Γ is trivial.*

Proof When $\chi(M) \neq 0$ we have that ev is trivial, a fact that is true even in the more general setting of loops of diffeomorphisms on a closed manifold; see Lê and Ono [12]. Therefore, Γ is trivial by the injectivity of e . Now, suppose that $\pi_1(M)$ has finite center and let $\{\psi_t\}$ be a symplectic loop. Then, there exists some positive integer k such that $\text{ev}(\{\psi_t\})^k = 1$. Thus, we have

$$e(\widetilde{\text{Flux}}(\{\psi_t^k\})) = \text{ev}(\{\psi_t^k\}) = \text{ev}(\{\psi_t\})^k = 1.$$

We conclude, by the injectivity of e and the fact that $\widetilde{\text{Flux}}$ is a homomorphism to a torsion-free group, that the loop $\{\psi_t\}$ has no flux. □

¹The fact that ev factors through the flux group was pointed out to me by Egor Shelukhin.

Remark In the case where (M, ω) is symplectically aspherical and the Euler characteristic is nonzero, Proposition 1.1 can be obtained from [12, Proposition 1.7, Corollary 4.2]. Furthermore, the fact that the flux group vanishes was established in Lalonde, McDuff and Polterovich [11] and Polterovich [24]. We also note that in Kędra, Kotschick and Morita [9], the implication of Proposition 1.1 is proven under the weak Lefschetz assumption.

Our main result provides more sufficient conditions for the injectivity of the e -homomorphism.

Definition 1.2 Let (M, ω) be a closed symplectic manifold of dimension $2n$. We say that it satisfies condition (\star) if at least one of the following is true:

- **Symplectically aspherical** The cohomology class $[\omega]$ vanishes on the image $H_2^S(M; \mathbb{Z})$ of the Hurewicz map $\pi_2(M) \rightarrow H_2(M; \mathbb{Z})$.
- **Spherically monotone** There exists a constant $\lambda \in \mathbb{R} \setminus \{0\}$ such that

$$[\omega]|_{\pi_2(M)} = \lambda \cdot c_1(M)|_{\pi_2(M)},$$

where $c_1(M)$ denotes the first Chern class associated with (M, ω) . We say *positive* (resp. *negative*) *spherically monotone* when $\lambda > 0$ (resp. $\lambda < 0$).

- **Weak Lefschetz property** The multiplication map

$$[\omega]^{n-1} : H^1(M; \mathbb{R}) \rightarrow H^{2n-1}(M; \mathbb{R})$$

is injective (hence an isomorphism).

Remark (examples satisfying (\star)) The standard symplectic torus $(T^{2n}, dp \wedge dq)$ and any closed oriented surface Σ_g of genus $g \geq 1$ with the standard area form are symplectically aspherical. Symplectic products of the form $(M \times N, \omega_M \oplus \omega_N)$, where (M, ω_M) is positive (resp. negative) homologically monotone and (N, ω_N) is aspherical, are positive (resp. negative) spherically monotone but not homologically monotone. In particular, $\mathbb{C}P^n \times T^{2m}$ with symplectic form $\omega_{FS} \oplus dp \wedge dq$ is positive spherically monotone. Hypersurfaces of $\mathbb{C}P^n$ defined by setting $z_0^m + \dots + z_n^m = 0$, with $m > n + 1$, are negative homologically monotone; see McDuff and Salamon [18]. Finally, all closed Kähler manifolds satisfy the weak Lefschetz property.

Theorem 1.3 Let (M, ω) be a closed symplectic manifold satisfying (\star) . Then the e -homomorphism is injective.

When the weak Lefschetz property is satisfied the injectivity of the e -homomorphism follows from classical arguments that were known at least as early as the work of Ono in [20]; see also McDuff and Salamon [19] and Lalonde, McDuff and Polterovich [11]. Indeed, if $\{\psi_t\}$ is a symplectic loop inducing a symplectic vector field X_t , then the homology classes of its orbits are Poincaré dual to the class

$$\frac{1}{\text{Vol}(M)} \left[\widetilde{\text{Flux}}(\{\psi_t\}) \wedge \frac{\omega^{n-1}}{(n-1)!} \right] \in H^{2n-1}(M; \mathbb{R}).$$

Therefore, by the injectivity of $[\omega]^{n-1}$, we have that $\widetilde{\text{Flux}}(\{\psi_t\}) = 0$ whenever $\text{ev}(\{\psi_t\}) = 1$, since every orbit of $\{\psi_t\}$ is homologically trivial.

1.1.1 Outline of the proof of Theorem 1.3 When (M, ω) is spherically monotone we provide two proofs of the injectivity of the e -homomorphism, which are detailed in Section 3. The first proof relies on looking at this problem from the perspective of Floer–Novikov theory developed by Lê and Ono [30]; see also Ono [22; 23]. Floer–Novikov cohomology is a natural generalization of Hamiltonian Floer cohomology in the sense that to a symplectic path $\{\psi_t\}$ based at identity with nondegenerate endpoint $\psi = \psi_1$, we can associate a Floer-type cohomology group $\text{HFN}^*(\{\psi_t\}; J)$ that, up to a natural isomorphism, depends only on the flux of $\{\psi_t\}$. The proof has two key steps. First, using ideas in [23] we obtain an isomorphism

$$\text{HFN}(\{\psi_t\}; J) \cong H^*(M; \mathbb{Q}) \otimes \Lambda_\omega$$

for a symplectic path $\{\psi_t\}$ with $\widetilde{\text{Flux}}(\{\psi_t\}) = [\theta] \in \ker e$. Next, we show that when (M, ω) is spherically monotone, the Floer–Novikov cohomology of a symplectic path $\{\psi_t\}$ is isomorphic to the Morse–Novikov cohomology $\text{HN}^*(M, \theta)$ of its flux. A simple rank comparison then shows that this is only possible when $\widetilde{\text{Flux}}(\{\psi_t\}) = 0$, which concludes the argument.

The second proof follows from a result of McDuff [14, Theorem 1], from which the triviality of Γ in the homologically monotone setting follows. Let $\{\psi_t\}$ be a symplectic loop. For a loop γ , set $\alpha(t, s) = \psi_t(\gamma(s))$ as before. McDuff’s result implies that the 2-cycle A_γ represented by the torus $\text{im}(\alpha)$ satisfies

$$(3) \quad \langle c_1(M), A_\gamma \rangle = 0.$$

If $\{\psi_t\}$ has trivial evaluation, then $\text{im}(\alpha)$ can be represented by a sphere. Thus, we obtain

$$\langle \widetilde{\text{Flux}}(\{\psi_t\}), [\gamma] \rangle = \langle [\omega], A_\gamma \rangle = \lambda \cdot \langle c_1(M), A_\gamma \rangle = 0,$$

where $\lambda \neq 0$ is the monotonicity constant. Since γ is arbitrary, we conclude that $\widetilde{\text{Flux}}(\{\psi_t\}) = 0$. While this proof is easier, it heavily relies on McDuff’s result, which was proven using highly nontrivial topological arguments. The first proof, on the other hand, is symplectic in nature.

1.2 Symplectic circle actions

Let $S^1 = \mathbb{R}/\mathbb{Z}$ be the standard circle group. Let (M, ω) be a symplectic manifold equipped with a smooth S^1 -action generated by a vector field X . The contraction of the symplectic form with the vector field X defines a 1-form $\iota_X \omega$ on M . The circle action is called *symplectic* whenever $\iota_X \omega$ is closed, and *Hamiltonian* if it is also exact. Knowing that the action is Hamiltonian has several advantages. For example, one can use a primitive H of $\iota_X \omega$, referred to as a *moment map*, to obtain a symplectic quotient of M at a regular value of H . This procedure is used, in particular, to reduce the dimension of the phase-space associated to problems arising in classical mechanics that have a circular symmetry.

If $H^1(M; \mathbb{R}) = 0$, it is clear that every symplectic circle action is Hamiltonian. Otherwise, it becomes substantially more difficult to determine whether $\iota_X \omega$ is an exact 1-form. Finding sufficient and necessary conditions assuring that a symplectic circle action is Hamiltonian has been a subject of interest at least as early as the work of T. Frankel in the late 1950s showing the following:

Theorem 1.4 (Frankel [6]) *A symplectic circle action on a closed Kähler manifold is Hamiltonian if and only if it has fixed points.*

This result was later generalized by Ono [20] to closed Lefschetz manifolds. The condition of having fixed points was shown to be sufficient to guarantee the exactness of the circle action in a few other instances. McDuff [15] proved it when M has dimension four, while Ono [21] and Ginzburg [7] showed it in the symplectically aspherical case. Tolman and Weitsman [29] proved that it remains true for semifree circle actions with isolated fixed points. Finally, Lupton and Oprea [13] and McDuff [14] showed that every symplectic circle action is Hamiltonian when (M, ω) is homologically monotone, i.e. the cohomology class $[\omega]$ is a nonzero multiple of the first Chern class $c_1(M)$ — a fact which is not true in the more general spherically monotone setting. Indeed, one can consider the symplectic circle action on $T^2 \times S^2$ (with the product symplectic form of the standard symplectic structures) given by rotation in the first factor and identity on the second.

On the other hand, McDuff constructed in [15] a non-Hamiltonian circle action with fixed tori on a closed 6-dimensional Calabi–Yau symplectic manifold (see Cho and Kim [3]), showing that the condition $M^{S^1} \neq \emptyset$ alone is not sufficient to guarantee the exactness of $\iota_X \omega$ for general closed symplectic manifolds. McDuff and Salamon then asked if every symplectic circle action with isolated fixed points on a closed symplectic manifold is Hamiltonian. This question was answered in the negative by Tolman in [28]; see also Jang and Tolman [8]. It remains unclear when such examples can exist, or from another viewpoint, how large the class is of closed symplectic manifolds for which the presence of fixed points is equivalent to the exactness of the circle action. Nonetheless, the e -homomorphism gives a partial characterization of this class. In particular, we have the following:

Proposition 1.5 *Let (M, ω) be a closed symplectic manifold such that the e -homomorphism is injective. Then a symplectic circle action is Hamiltonian if and only if it has fixed points.*

Proof If a symplectic circle action has a fixed point, then the symplectic loop $\{\psi_t\}$ induced by it has trivial evaluation. The injectivity of the e -homomorphism implies that $[\iota_X \omega] = \widetilde{\text{Flux}}(\{\psi_t\}) = 0$. Here, X is the time-independent vector field generating the action. \square

In view of Theorem 1.3, we obtain the following.

Theorem 1.6 *Let (M, ω) be a closed spherically monotone symplectic manifold. Then, a symplectic circle action is Hamiltonian if and only if it has a fixed point.*

A result of McDuff [17] showed that every closed symplectic manifold that admits a Hamiltonian circle action is uniruled in the Gromov–Witten sense. In particular, they are geometrically uniruled: for each ω -compatible almost complex structure J and each point $x \in M$, there is a nonconstant J -holomorphic sphere u such that $x \in \text{im}(u)$. Symplectic manifolds that are symplectically Calabi–Yau are not geometrically uniruled, neither are (spherically) negative monotone ones. By considering this fact in addition to Theorem 1.6 we obtain the following corollary.

Corollary 1.7 *Let the circle act symplectically and nontrivially on a closed symplectic manifold (M, ω) such that*

$$c_1(M)|_{\pi_2(M)} = \lambda \cdot [\omega]|_{\pi_2(M)}$$

for $\lambda \in \mathbb{R}$. Then:

- (i) *If $\lambda > 0$, the action is Hamiltonian if and only if it has fixed points.*
- (ii) *If $\lambda < 0$, the action is non-Hamiltonian, and has no fixed points.*
- (iii) *If $\lambda = 0$ the action is non-Hamiltonian.*

Remark When $\lambda \neq 0$ there are examples of non-Hamiltonian symplectic circle actions. In particular, consider the symplectic product $\mathbb{T}^2 \times M$ of the standard symplectic torus with any closed spherically monotone symplectic manifold, and the symplectic circle action given by $t \cdot (x, y, p) = (x + t, y, p)$ for $(x, y) \in T^2$ and $p \in M$.

Another closely related question was raised by McDuff and Salamon [19]. They asked if there exists a symplectic free circle action whose orbits are contractible. Kotschick [10] proved this to be the case for all symplectic manifolds of dimension four, even if the action is only assumed to be smooth. Furthermore, Kotschick produced examples of symplectic free circle actions with contractible orbits in every even dimension greater than or equal to six. As a corollary of Theorem 1.3 and the argument in the proof of Proposition 1.5, we obtain the following.

Theorem 1.8 *Let (M, ω) be a closed symplectic manifold satisfying (\star) . Then every free symplectic circle action must have noncontractible orbits.*

1.3 Applications to symplectic torsion

The injectivity of the e -homomorphism, when combined with results in Atallah and Shelukhin [1], provides applications to questions about the existence of finite subgroups of $\text{Symp}_0(M, \omega)$. In [1], it was shown that if (M, ω) is positive spherically monotone, then the existence of a nontrivial finite subgroup of $\text{Ham}(M, \omega)$ implies that (M, ω) is geometrically uniruled. Furthermore, it was shown that if (M, ω) is negative spherically monotone, then there are no nontrivial finite subgroups of $\text{Ham}(M, \omega)$. Therefore, as a corollary of Proposition 1.1 and Theorem 1.3 we obtain the following.

Theorem 1.9 *Let (M, ω) be a closed symplectic manifold such that*

$$c_1(M)|_{\pi_2(M)} = \lambda \cdot [\omega]|_{\pi_2(M)}$$

for $\lambda \in \mathbb{R} \setminus \{0\}$. Further, suppose that either $\chi(M) \neq 0$ or $\pi_1(M)$ has finite center. Then:

- (i) *If $\lambda > 0$, then the existence of a nontrivial finite subgroup of $\text{Symp}_0(M, \omega)$ implies that (M, ω) is geometrically uniruled.*
- (ii) *If $\lambda < 0$, then all finite subgroups of $\text{Symp}_0(M, \omega)$ are trivial.*

Remark The symplectic product of $(\mathbb{C}P^n, \omega_{\text{FS}})$ with any closed oriented surface Σ_g of genus $g \geq 2$ with the standard area form satisfies condition (i) of Theorem 1.9. More generally, any symplectic product of the form $(M \times N, \omega_M \oplus \omega_N)$, where (M, ω_M) is positive spherically monotone with $\chi(M) \neq 0$ and (N, ω_N) is symplectically aspherical with $\chi(N) \neq 0$. Furthermore, following Dimca [4], the Euler characteristic of the degree m hypersurface $X_m \subset \mathbb{C}P^n$ defined by setting $z_0^m + \dots + z_n^m = 0$ is given by

$$\chi(X_m) = \frac{1}{m}((1-m)^{n+1} - 1) + n + 1.$$

Therefore, $\chi(X_m) \neq 0$ when $m > n + 1$. Hence, X_m , which is negative monotone, satisfies condition (ii) of Theorem 1.9.

The injectivity of the e -homomorphism also gives information about the presence of fixed points of a nontrivial symplectic diffeomorphism $\psi \in \text{Symp}_0(M, \omega)$ of finite order. The following definition given by Polterovich in [24] naturally fits into this context.

Definition 1.10 A fixed point x of a symplectic diffeomorphism $\psi \in \text{Symp}_0(M, \omega)$ is of *contractible type* if there exists a symplectic path $\{\psi_t\}$ based at the identity with $\psi_1 = \psi$ such that the loop $\{\psi_t(x)\}$ is contractible in M .

The presence of a fixed point of contractible type of a nontrivial $\psi \in \text{Symp}_0(M, \omega)$ of finite order implies that it must be Hamiltonian. Indeed, we have the following:

Proposition 1.11 *Let (M, ω) be a closed symplectic manifold such that the e -homomorphism is injective. Further suppose that $\psi \in \text{Symp}_0(M, \omega)$ is nontrivial of finite order, ie $\psi^d = \text{id}$ for some integer $d > 1$. Then ψ is Hamiltonian if and only if it admits a fixed point of contractible type.*

Proof Suppose $x \in \text{Fix}(x)$ is of contractible type. Let $\{\psi_t\}$ be a symplectic path with $\psi_1 = \psi$ such that $\{\psi_t(x)\}$ is contractible in M . Note that $\{\psi_t^d(x)\}$ is a symplectic loop. Then,

$$e(\widetilde{\text{Flux}}(\{\psi_t^d\})) = [\{\psi_t^d(x)\}] = [\{\psi_t(x)\}]^d = 1.$$

Hence, by the injectivity of e and the fact that $\widetilde{\text{Flux}}$ is a homomorphism to a torsion-free group, we have that $\widetilde{\text{Flux}}(\{\psi_t\}) = 0$. In particular, ψ is Hamiltonian. □

Corollary 1.12 *Let (M, ω) be a closed symplectic manifold satisfying (\star) such that $\chi(M) \neq 0$. Then all nontrivial symplectic diffeomorphisms $\psi \in \text{Symp}_0(M, \omega)$ of finite order are Hamiltonian.*

We note that any orientation-preserving diffeomorphism of a closed manifold, which is of finite order and has finitely many fixed points, must have the number of fixed points equal to the Lefschetz number of the diffeomorphism. Therefore, when the Euler characteristic of (M, ω) is zero and $\psi \in \text{Symp}_0(M, \omega)$ is torsion, it either has no fixed points or nonisolated ones. In the latter case, there is no a priori reason for the existence of a fixed point of contractible type. However, in the case of the standard symplectic torus, Polterovich [24] showed that any fixed point of a symplectic diffeomorphism $\psi \in \text{Symp}_0(\mathbb{T}^{2n}, dp \wedge dq)$ is of contractible type. We can adapt that argument to prove the following:

Corollary 1.13 *Let (M, ω) be a simply connected closed symplectic manifold satisfying (\star) . Then every non-Hamiltonian $\psi \in \text{Symp}_0(\mathbb{T}^{2n} \times M, dp \wedge dq \oplus \omega)$ of finite order has no fixed points.*

Proof Let $(x, p) \in \mathbb{T}^{2n} \times M$ be a fixed point of ψ , and $\{\psi_t\}$ a symplectic path such that $\psi_0 = \text{id}$ and $\psi_1 = \psi$. Consider the lift

$$\tilde{\psi}_t: \mathbb{R}^{2n} \times M \rightarrow \mathbb{R}^{2n} \times M$$

of ψ_t to the universal cover of $\mathbb{T}^{2n} \times M$ and pick $(\tilde{x}, p) \in \pi^{-1}(\{(x, p)\})$. Then, $\tilde{\psi}(\tilde{x}, p) = (\tilde{x} + a, p)$ for some $a \in \mathbb{Z}^{2n}$. We can then define a symplectic flow $f_t \times \text{id}_M$ on $\mathbb{T}^{2n} \times M$ by setting

$$f_t(y) = y - t \cdot a \pmod{1}.$$

Note that its lift to the universal cover is given by

$$(\tilde{f}_t \times \text{id}_M)(y, q) = (y - t \cdot a, q).$$

Therefore, by setting $\varphi_t = f_t \circ \psi_t$, we obtain a symplectic path based at the identity with $\varphi_1 = \psi$, and whose lift $\tilde{\varphi}_t$ satisfies $\tilde{\varphi}(\tilde{x}, p) = (\tilde{x}, p)$. Consequently, the loop $\{\varphi_t(x, p)\}$ is contractible in $\mathbb{T}^{2n} \times M$. The corollary then follows by noting that $\mathbb{T}^{2n} \times M$ satisfies condition (\star) and by Proposition 1.11. \square

2 Preliminaries

2.1 Floer–Novikov cohomology

In Section 2.1.2 we review the construction of Floer–Novikov cohomology after Ono [23]. This is a cohomological version of the construction introduced by Lê and Ono [30] with a slightly smaller coefficient ring. In Section 2.1.3 we review a variant of Floer–Novikov cohomology introduced in Ono [22] which enables the comparison between symplectic paths with different flux. We refer to Lê and Ono [12] for an in-depth discussion of the variants of Floer–Novikov cohomology and the relations among them. We also briefly recall the definition of classical Morse–Novikov cohomology and outline a few important properties it satisfies. For further details on Morse–Novikov cohomology, see [30; 26; 5].

2.1.1 Morse–Novikov cohomology Let M be a closed smooth manifold and let θ be a closed 1–form on M . Fix a ground field \mathbb{K} . For our purposes it will be sufficient to consider the case $\mathbb{K} = \mathbb{Q}$. Denote by $I_\theta: \pi_1(M) \rightarrow \mathbb{R}$ the homomorphism given by integrating θ over a representative loop; that is,

$$I_\theta([\gamma]) = \int_\gamma \theta.$$

In particular, I_θ depends only on the cohomology class $[\theta]$. Denote by $\pi: \bar{M}^\theta \rightarrow M$ the covering space of M corresponding to $\ker I_\theta \subset \pi_1(M)$. It is the minimal abelian covering on which $\pi^*\theta$ is exact. The covering transformation group is given by $G_\theta = \pi_1(M)/\ker I_\theta$. We denote by Λ_θ the completion of the group ring of G_θ with respect to the filtration induced by I_θ ; that is,

$$\Lambda_\theta = \left\{ \sum_i a_i g_i \mid a_i \in \mathbb{K} \text{ and } g_i \in G_\theta \text{ satisfy condition (A)} \right\},$$

where condition (A) is:

(A) For each $c \in \mathbb{R}$, the set $\{i \mid a_i \neq 0, I_\theta(g_i) < c\}$ is finite.

The fact that \mathbb{K} is a field implies that so is Λ_θ . Let \bar{f} be a choice of primitive for $\pi^*\theta$. Then, to each $\tilde{x} \in \text{Crit}(\bar{f})$ there corresponds a zero $x = \pi(\tilde{x})$ of θ . A 1–form θ is said to be Morse if all the critical points of \bar{f} are nondegenerate. The Morse–Novikov cochain complex in degree k with coefficients in \mathbb{K} is defined as

$$\text{CN}^k(M, \theta) = \left\{ \sum_i a_i \tilde{x}_i \mid a_i \in \mathbb{K} \text{ and } \tilde{x}_i \in \text{Crit}(\bar{f}), \text{ where } \text{index}(\tilde{x}_i) = k, \text{ satisfy condition (B)} \right\},$$

where condition (B) is:

(B) For each $c \in \mathbb{R}$ the set $\{i \mid a_i \neq 0, \bar{f}(\tilde{x}_i) < c\}$ is finite.

Note that $\text{CN}^*(M, \theta)$ is finitely generated over Λ_θ . For a choice of Riemannian metric g on M , the coboundary operator δ is defined by counting bounded gradient trajectories of \bar{f} with respect to the pullback metric π^*g that are emerging from a critical point $\tilde{x} \in \text{Crit}(\bar{f})$ and converging to critical points $\tilde{y} \in \text{Crit}(\bar{f})$ such that $\text{index}(\tilde{y}) - \text{index}(\tilde{x}) = 1$. We may assume that the gradient flow is of Morse–Smale type. The Morse–Novikov cohomology of θ is defined as $\text{HN}^*(M, \theta) = H^*((\text{CN}(M, \theta), \delta))$, and is a finitely generated Λ_θ –module. The resulting cohomology is independent of the choice of Riemannian metric for which the flow is of Morse–Smale type. Furthermore, cohomologous 1–forms have canonically isomorphic Morse–Novikov cohomologies. That is, if $[\theta_1] = [\theta_2]$, then $\text{HN}^*(M, \theta_1)$ is isomorphic to $\text{HN}^*(M, \theta_2)$ as graded vector spaces over $\Lambda_{\theta_1} = \Lambda_{\theta_2}$. More generally, we have the following.

Proposition 2.1 (Lê and Ono [30, Theorem C.2]) *Suppose that θ_1 and θ_2 are closed 1–forms such that $\ker I_{\theta_1} \subset \ker I_{\theta_2}$. Then, for each degree k ,*

$$\dim_{\Lambda_{\theta_1}} \text{HN}^k(M, \theta_1) \leq \dim_{\Lambda_{\theta_2}} \text{HN}^k(M, \theta_2).$$

In addition, we have the following useful proposition, which distinguishes Morse–Novikov cohomology from usual Morse cohomology.

Proposition 2.2 (Ono [23, Proposition 4.12]) *Let θ be a closed 1–form such that $[\theta] \neq 0$. Then*

$$\dim_{\Lambda_\theta} \text{HN}^k(M, \theta) = 0 \quad \text{for } k = 0 \text{ or } k = \dim(M).$$

In particular, when θ is not exact, we have that

$$(4) \quad \dim_{\Lambda_\theta} \text{HN}(M, \theta) < \dim_{\mathbb{Q}} \text{H}(M).$$

2.1.2 Floer–Novikov cohomology Let (M, ω) be a closed symplectic manifold, and $\{\psi_t\}$ a symplectic path based at the identity with endpoint $\psi_1 = \psi$. The deformation lemma in [30] implies that we can suppose, without loss of generality, that there exists a 1–periodic smooth family of smooth functions $F_t \in C^\infty(M)$, with $F_{t+1} = F_t$, such that

$$(5) \quad \iota_{X^t} \omega = \theta + dF_t,$$

where X^t is the time–dependent vector field induced by the symplectic isotopy $\{\psi_t\}$ and θ is a closed 1–form representing $\widetilde{\text{Flux}}(\{\psi_t\})$. When θ is exact, equation (5) becomes the usual Hamiltonian equation for $H_t = F_t + f$, where $df = \theta$. Similarly, we have a formal closed 1–form on the contractible component $\mathcal{L}M$ of the loop space of M . Indeed, for a loop $x \in \mathcal{L}M$ and $v \in T_x \mathcal{L}M = \Gamma(x^* TM)$, we define

$$a_{\{\psi_t\}}(v) = \int_0^1 \omega(v(t), x'(t) - X^t(x(t))) dt = \int_0^1 \omega(v(t), x'(t)) dt + \int_0^1 (\theta + dF_t)(v(t)) dt.$$

The idea is to find a suitable cover on which $a_{\{\psi_t\}}$ is exact, and then to define Floer–Novikov cohomology as an analog of Morse theory for a primitive \mathcal{A}_{ψ_t} . Consider the homomorphisms $\mathcal{I}_\theta, \mathcal{I}_\omega, \mathcal{I}_{c_1} : \pi_1(\mathcal{L}M) \rightarrow \mathbb{R}$ defined as

$$\mathcal{I}_\theta = I_\theta \circ \text{Ev}_*, \quad \mathcal{I}_\omega(\{x_s\}) = \int_{C(\{x_s\})} \omega, \quad \mathcal{I}_{c_1}(\{x_s\}) = \langle c_1(M), [C(\{x_s\})] \rangle,$$

for a loop $\{x_s\}_{s \in [0,1]}$ in $\mathcal{L}M$. Here, $\text{Ev} : \mathcal{L}M \rightarrow M$ is given by evaluation at $t = 0$, and $C(\{x_s\})$ is the 2–cycle represented by

$$\mathbb{S}^1 \times \mathbb{S}^1 \ni (s, t) \mapsto x_s(t) \in M.$$

We denote by $\widetilde{\mathcal{L}M} \rightarrow \mathcal{L}M$ the covering space of $\mathcal{L}M$ corresponding to

$$\ker(\mathcal{I}_\omega + \mathcal{I}_\theta) \cap \ker \mathcal{I}_{c_1} \subset \pi_1(\mathcal{L}M).$$

A description of $\widetilde{\mathcal{L}M}$ can be given in the following manner. Choose a primitive \bar{f} of $\pi^* \theta$ on \bar{M}^θ , and consider pairs (\tilde{x}, u) composed of a loop $\tilde{x} \in \mathcal{L}\bar{M}^\theta$ and a capping $u : \mathbb{D} \rightarrow M$ such that $u|_{\partial\mathbb{D}} = \pi \circ \tilde{x}$. Define the equivalence relation \sim given by $(\tilde{x}, u) \sim (\tilde{y}, w)$ if and only if

$$\pi \circ \tilde{x} = \pi \circ \tilde{y}, \quad \int_u \omega + \bar{f}(\tilde{x}(0)) = \int_w \omega + \bar{f}(\tilde{y}(0)), \quad \langle c_1(M), u \# (-w) \rangle = 0.$$

Here, $u\#(-w)$ corresponds to the sphere obtained by gluing the two disks along their common boundaries with the orientation of w reversed. Each element in $\tilde{\mathcal{L}}M$ corresponds to an equivalence class $[\tilde{x}, u]$ of such a pair. With this description, the covering map $\Pi: \tilde{\mathcal{L}}M \rightarrow \mathcal{L}M$ is given by $[\tilde{x}, u] \mapsto \pi \circ \tilde{x} = x$. A choice of primitive for $\Pi^*a_{\{\psi_t\}}$ is given by

$$\mathcal{A}_{\{\psi_t\}}([\tilde{x}, u]) = \int_0^1 (\bar{f} + F_t \circ \pi)(\tilde{x}(t)) dt + \int_u \omega,$$

where F_t is as in equation (5). The critical points $\mathcal{P}(\{\psi_t\})$ of $\mathcal{A}_{\{\psi_t\}}$ are the lifts to $\tilde{\mathcal{L}}M$ of the fixed points $x \in \text{Fix}(\psi)$ of ψ such that $[\{\psi_t(x)\}] = 1$ in $\pi_1(M)$. To a critical point $[\tilde{x}, u]$ we assign an index $\text{CZ}([\tilde{x}, u])$ given by the Conley–Zehnder index of $x = \pi \circ \tilde{x}$ with respect to the trivialization x^*TM , which extends to u^*TM . Note that the covering transformation group of $\mathcal{L}M$ is given by

$$G_{\theta,\omega} = \frac{\pi_1(\mathcal{L}M)}{\ker(\mathcal{F}_\omega + \mathcal{F}_\theta) \cap \ker \mathcal{F}_{c_1}}.$$

Let $\Lambda_{\theta,\omega}$ be Novikov ring given by the completion of the group ring of $G_{\theta,\omega}$ with respect to the filtration induced by $\mathcal{F}_\omega + \mathcal{F}_\theta$; that is,

$$\Lambda_{\theta,\omega} = \left\{ \sum_i a_i g_i \mid a_i \in \mathbb{Q} \text{ and } g_i \in G_{\theta,\omega} \text{ satisfy condition (A')} \right\},$$

where condition (A') is:

(A') For each $c \in \mathbb{R}$ the set $\{i \mid a_i \neq 0, (\mathcal{F}_\omega + \mathcal{F}_\theta)(g_i) < c\}$ is finite.

Suppose that ψ is nondegenerate and let $J = \{J_t\}$ be a family of ω -compatible almost complex structures. The Floer–Novikov cochain complex is defined as

$$\text{CFN}^k(\{\psi_t\}; J) = \left\{ \sum_i a_i [\tilde{x}_i, u_i] \mid a_i \in \mathbb{Q} \text{ and } [\tilde{x}_i, u_i] \in \mathcal{P}(\{\psi_t\}) \text{ satisfy condition (B')} \right\},$$

where condition (B') is:

(B') For each $c \in \mathbb{R}$ the set $\{i \mid a_i \neq 0, \mathcal{A}_{\{\psi_t\}}([\tilde{x}_i, u_i]) < c\}$ is finite and $\text{CZ}([\tilde{x}_i, u_i]) = k$ for all i .

The graded \mathbb{Q} -vector space $\text{CFN}^*(\{\psi_t\}; J)$ is endowed with the Floer–Novikov coboundary operator δ_{FN} , which is defined as the signed count of isolated solutions (modulo the \mathbb{R} -action) of the asymptotic boundary value problem on maps $u: \mathbb{R} \times S^1 \rightarrow M$ defined by the gradient of $\mathcal{A}_{\{\psi_t\}}$; see [30; 23]. In other words, the coboundary operator counts the finite-energy solutions to the Floer equation

$$\frac{\partial u}{\partial s} + J_t(u) \left(\frac{\partial u}{\partial t} - X_t(u) \right) = 0, \quad \lim_{s \rightarrow \pm\infty} \tilde{u}(s, t) = \tilde{x}^\pm(t),$$

for some lift $\tilde{u}: \mathbb{R} \times S^1 \rightarrow M$ such that $\text{CZ}([\tilde{x}^+, u^+]) - \text{CZ}([\tilde{x}^-, u^-]) = 1$ and $[\tilde{x}^+, u^- \# u] = [\tilde{x}^+, u^+]$. The Floer–Novikov cohomology is defined as

$$\text{HFN}^*(\{\psi_t\}; J) = H^*(\text{CFN}(\{\psi_t\}; J), \delta_{\text{FN}}).$$

It can be shown to be independent of the choice of J . In addition, it only depends on the flux of the symplectic path.

Theorem 2.3 (Lê and Ono [30, Theorem 4.3]) *Suppose that $\{\psi_t^{(1)}\}$ and $\{\psi_t^{(2)}\}$ are symplectic paths with nondegenerate endpoints. Suppose that $\{\psi_t^{(1)} \circ (\psi_t^{(2)})^{-1}\}$ has zero flux. Then,*

$$\text{HFN}^*(\{\psi_t^{(1)}\}; J^{(1)}) \cong \text{HFN}^*(\{\psi_t^{(2)}\}; J^{(2)})$$

as $\Lambda_{\theta, \omega}$ -modules. Here, $[\theta]$ is the flux of both paths.

Remark When $\{\varphi_H^t\}$ is a Hamiltonian isotopy, $\overline{M}^\theta = M$ and $(\text{CFN}^*(\{\psi\}; J), \delta)$ reduces to the usual Floer cochain complex $(\text{CF}(H; J), \delta)$ associated with the Hamiltonian function H . Therefore, the PSS isomorphism yields

$$(6) \quad \text{HFN}^*(\{\varphi_H^t\}; J) = \text{HF}^*(H; J) \cong \text{H}^{*+n}(M; \mathbb{Q}) \otimes_{\mathbb{Q}} \Lambda_{\omega}.$$

The main reason for the choice of Novikov ring in this construction is so that we have the following two statements.

Theorem 2.4 (Ono [23, Theorem 4.10]) *Let $\{\psi_t^{(1)}\}$ and $\{\psi_t^{(2)}\}$ be symplectic paths with nondegenerate endpoints $\psi_1^{(1)} = \psi_1^{(2)} = \psi$. Suppose that the symplectic loop $\{\psi_t^{(1)} \circ (\psi_t^{(2)})^{-1}\}$ has trivial evaluation, ie $\text{ev}(\{\psi_t^{(1)} \circ (\psi_t^{(2)})^{-1}\}) = 1$. Then, we have a ring isomorphism $\Psi: \Lambda_{\theta_1, \omega} \xrightarrow{\cong} \Lambda_{\theta_2, \omega}$, and*

$$\text{HFN}^*(\{\psi_t^{(1)}\}; J^{(1)}) \cong \text{HFN}^*(\{\psi_t^{(2)}\}; J^{(2)})$$

as $\Lambda_{\theta_1, \omega}$ -modules, where $[\theta_i]$ corresponds to the flux of $\{\psi_t^{(i)}\}$ for $i = 1, 2$. The module action of $\Lambda_{\theta_1, \omega}$ on $\text{HFN}^*(\{\psi_t^{(2)}\}; J^{(2)})$ is the one induced by Ψ .

Proof Let $\xi_t = \psi_t^{(1)} \circ (\psi_t^{(2)})^{-1}$. In [23, Theorem 4.10] the conclusion of Theorem 2.4 was proven under the assumption that the map $\Psi: \{\gamma(t)\} \mapsto \{\xi_t(\gamma(t))\}$ preserves the component \mathcal{LM} of the free loop space consisting of contractible loops. Therefore, Theorem 2.4 follows from the observation that Ψ preserves \mathcal{LM} whenever $\text{ev}(\{\xi_t\}) = 1$. As detailed in [23, Section 4], it is the map $\Psi: \mathcal{LM} \rightarrow \mathcal{LM}$ that induces the isomorphism $\text{HFN}^*(\{\psi_t^{(1)}\}; J^{(1)}) \cong \text{HFN}^*(\{\psi_t^{(2)}\}; J^{(2)})$. \square

Theorem 2.5 (Ono [23, Theorem 3.12]) *Let $\{\psi_t\}$ be a symplectic path based at identity with sufficiently small flux $[\theta]$. Then,*

$$\text{HFN}^*(\{\psi_t\}; J) \cong \text{HN}^{*+n}(M, \theta) \otimes_{\Lambda_{\theta}} \Lambda_{\theta, \omega}.$$

Remark The Floer–Novikov cohomology $\text{HFN}^*(\{\psi_t\}; J)$ of a symplectic path $\{\psi_t\}$ is not always isomorphic to the Morse–Novikov cohomology $\text{HN}(M, \theta)$ of its flux $[\theta]$; this has also been observed in Seidel [27]. This can be seen by studying the examples in Jang and Tolman [8], Tolman [28] and McDuff [15] of non-Hamiltonian symplectic circle actions with fixed points. Indeed, in these cases we

have symplectic paths with nontrivial flux and trivial evaluation. In particular, if we suppose that such an isomorphism exists, Theorem 2.4 together with Theorem 2.5 would imply that $\dim_{\Lambda_\theta} \text{HN}(M, \theta) = \dim_{\mathbb{Q}} H(M; \mathbb{Q})$, which is in contradiction to Proposition 2.2. This also shows that the e -homomorphism is not injective in general.

2.1.3 A variant of Floer–Novikov cohomology for changing flux In this section we recall a variant, introduced in [22], of the Floer–Novikov cohomology construction presented in Section 2.1.2, which allows the comparison between symplectic paths that have different flux.

Let (M, ω) be a closed symplectic manifold. Suppose $\{\psi_t\}$ is a symplectic path with endpoint $\psi_1 = \psi$ and flux $[\theta]$. Let $p: \tilde{M} \rightarrow M$ be an abelian covering space of M such that $p^*\theta$ is exact; \bar{M}^θ is the smallest choice of such a covering space. Let $\{\tilde{F}_t\}$ be a smooth family of smooth functions on \tilde{M} such that $d\tilde{F}_t = p^*\theta$. Just as before, we would like to make a choice of covering space of $\mathcal{L}M$ on which the pullback of $a_{\{\psi_t\}}$ is exact. We denote by $P: \tilde{\mathcal{L}}\tilde{M} \rightarrow \mathcal{L}M$ the covering space of $\mathcal{L}M$ associated with

$$\ker \mathcal{F}_\omega \cap \ker \mathcal{F}_{c_1} \cap \text{Ev}_*^{-1}(p_*(\pi_1(\tilde{M}))) \subset \pi_1(\mathcal{L}M).$$

This covering can be seen as the space of pairs (\tilde{x}, u) , $u|_{\partial\mathbb{D}} = \pi \circ \tilde{x}$, under the equivalence relation defined by $(\tilde{x}, u) \sim (\tilde{y}, w)$ if and only if

$$\tilde{x} = \tilde{y}, \quad \int_u \omega = \int_w \omega, \quad \langle c_1(M), u \# (-w) \rangle = 0.$$

On $\tilde{\mathcal{L}}\tilde{M}$, the pullback $P^*a_{\psi_t}$ is exact and a choice of primitive is given by

$$\tilde{\mathcal{A}}_{\{\psi_t\}}([\tilde{x}, u]) = \int_0^1 \tilde{F}_t(\tilde{x}(t)) dt + \int_u \omega.$$

Similarly, the critical points $\tilde{\mathcal{P}}(\{\psi_t\})$ of $\tilde{\mathcal{A}}_{\{\psi_t\}}$ are lifts to $\tilde{\mathcal{L}}\tilde{M}$ of the fixed points $x \in \text{Fix}(\psi)$ of ψ that satisfy $[\{\psi_t(x)\}] = 1 \in \pi_1(M)$. Just as in Section 2.1.2, to each critical point $[\tilde{x}, u]$ we assign a Conley–Zehnder type index. The covering transformation group of $\tilde{\mathcal{L}}\tilde{M}$ is given by

$$\tilde{G}_{\theta, \omega} = \frac{\pi_1(\mathcal{L}M)}{\ker \mathcal{F}_\omega \cap \ker \mathcal{F}_{c_1} \cap \text{Ev}_*^{-1}(p_*(\pi_1(\tilde{M})))},$$

and we denote by $\tilde{\Lambda}_{\theta, \omega}$ the Novikov ring given by the completion of its group ring with respect to the filtration induced by $\mathcal{F}_\omega + \mathcal{F}_\theta$; that is,

$$\tilde{\Lambda}_{\theta, \omega} = \left\{ \sum_i a_i g_i \mid a_i \in \mathbb{Q} \text{ and } g_i \in \tilde{G}_{\theta, \omega} \text{ satisfy condition (A')} \right\}.$$

Suppose that ψ is nondegenerate and let $J = \{J_t\}$ be a family of ω -compatible almost complex structures. The Floer–Novikov cochain complex is defined as

$$\text{CFN}^k(\{\psi_t\}, \tilde{M}; J) = \left\{ \sum_i a_i [\tilde{x}_i, u_i] \mid a_i \in \mathbb{Q} \text{ and } [\tilde{x}_i, u_i] \in \tilde{\mathcal{P}}(\{\psi_t\}) \text{ satisfy condition (B')} \right\}.$$

The coboundary operator $\tilde{\delta}_{FN}$ is defined by the same formula as in Section 2.1.2. The Floer–Novikov homology $\text{HFN}_*(\{\psi_t\}, \tilde{M}; J)$ associated with the covering space \tilde{M} is defined as the homology of $(\text{CFN}(\{\psi_t\}, \tilde{M}; J), \tilde{\delta}_{FN})$. It is independent of the choice of almost complex structure J and depends only on the flux of the symplectic path. The following theorem allows the comparison between the ranks of the Floer–Novikov cohomology of symplectic paths with flux lying in the kernel of p^* .

Proposition 2.6 (Ono [22, Proposition 4.8]) *Let $\{\psi_t^{(1)}\}$ and $\{\psi_t^{(2)}\}$ be symplectic paths such that $\widetilde{\text{Flux}}(\{\psi_t^{(i)}\}) \in \ker\{p^*: H^1(M; \mathbb{R}) \rightarrow H^1(\tilde{M}; \mathbb{R})\}$ for $i = 1, 2$. Then,*

$$\text{rank}_{\tilde{\Lambda}_{\theta_1, \omega}} \text{HFN}^*(\{\psi_t^{(1)}\}, \tilde{M}; J^{(1)}) = \text{rank}_{\tilde{\Lambda}_{\theta_2, \omega}} \text{HFN}^*(\{\psi_t^{(2)}\}, \tilde{M}; J^{(2)}),$$

where $[\theta_i]$ corresponds to the flux of $\{\psi_t^{(i)}\}$ for $i = 1, 2$.

3 Proof of Theorem 1.3

3.1 Proof using Floer–Novikov theory

When (M, ω) is either symplectically aspherical or satisfies the weak Lefschetz property, we have presented proofs based on classical arguments in Section 1.1. We shall, therefore, consider the spherically monotone case. Let $\{\psi_t\}$ be a symplectic loop with

$$\widetilde{\text{Flux}}(\{\psi_t\}) = [\theta] \in \Gamma.$$

Observe that spherical monotonicity implies that

$$(7) \quad \ker(\mathcal{F}_\omega + \mathcal{F}_\theta) \cap \ker \mathcal{F}_{c_1} = \ker \mathcal{F}_\omega \cap \ker \mathcal{F}_\theta \cap \ker \mathcal{F}_{c_1}.$$

Set $\tilde{M} = \overline{M}^\theta$ in the construction of the variant of Floer–Novikov homology defined in Section 2.1.3. In this case, note that $\text{Ev}_*^{-1}(p_*(\pi_1(\overline{M}^\theta))) = \ker \mathcal{F}_\theta$. Indeed, this is a consequence of $\pi_1(\overline{M}^\theta) = \ker I_\theta$, which follows from the defining property of \overline{M}^θ , and of the equality $\mathcal{F}_\theta = I_\theta \circ \text{Ev}_*$. These observations allow one to deduce that for any symplectic path $\{\xi_t\}$ with flux $[\theta]$, we have $\tilde{\Lambda}_{\theta, \omega} = \Lambda_{\theta, \omega}$, and

$$(8) \quad (\text{CFN}^*(\{\xi_t\}, \overline{M}^\theta; J), \tilde{\delta}_{FN}) = (\text{CFN}^*(\{\xi_t\}; J), \delta_{FN})$$

by definition (we reiterate that monotonicity is used in an important way; see the remark below).

Remark In general, $\text{HFN}_*(\{\psi_t\}, \tilde{M}; J)$ is different from $\text{HFN}_*(\{\psi_t\}; J)$ even in the case $\tilde{M} = \overline{M}^\theta$. Nonetheless, whenever $\ker \mathcal{F}_{c_1} \subset \ker \mathcal{F}_\omega$ we have that equation (8) holds and, arguing as in the preceding paragraph, that

$$\text{HFN}_*(\{\psi_t\}, \overline{M}^\theta; J) = \text{HFN}_*(\{\psi_t\}; J).$$

This holds for (M, ω) spherically (positive or negative) monotone or symplectically aspherical. We refer to [12] for more details on the relationship between these cohomology theories.

Now, further suppose that $\{\psi_t\} \in \ker \text{ev}$. Let $\{\varphi_t\}$ be a Hamiltonian path with endpoint φ , which is generated by a nondegenerate Hamiltonian H , and set $\psi'_t = \psi_t \circ \varphi_t$. Then $\{\psi'_t\}$ and $\{\varphi_t\}$ are two symplectic paths with nondegenerate endpoints $\psi' = \varphi$. Since $\text{ev}(\{\psi_t\}) = 1$, Theorem 2.4 implies that $\Lambda_{\theta,\omega} \cong \Lambda_\omega$ and that

$$(9) \quad \text{HFN}^*(\{\psi'_t\}, J') \cong \text{HFN}^*(\{\varphi_t\}; J)$$

as $\Lambda_{\theta,\omega}$ -modules. Since φ_t is Hamiltonian, we have that

$$(10) \quad \text{HFN}^*(\{\varphi_t\}; J) \cong \text{HF}^*(H; J) \cong \text{H}^*(M; \mathbb{Q}) \otimes_{\mathbb{Q}} \Lambda_\omega.$$

For $\varepsilon > 0$, let $\{\psi_t^{\varepsilon\theta}\}_{t \in [0,1]}$ be the symplectic path induced by the symplectic vector field $X_{\varepsilon\theta}$ defined by $\iota_{X_{\varepsilon\theta}} \omega = \varepsilon\theta$. Then, for all $\varepsilon > 0$, we have that

$$\widetilde{\text{Flux}}(\{\psi_t^{\varepsilon\theta}\}) = \varepsilon[\theta] \in \ker\{\pi^* : \text{H}^1(M; \mathbb{R}) \rightarrow \text{H}^1(\overline{M}^\theta; \mathbb{R})\}.$$

Therefore, Proposition 2.6 implies that

$$(11) \quad \text{rank}_{\widetilde{\Lambda}_{\theta,\omega}} \text{HFN}^*(\{\psi'_t\}, \overline{M}^\theta; J) = \text{rank}_{\widetilde{\Lambda}_{\varepsilon\theta,\omega}} \text{HFN}^*(\{\psi_t^{\varepsilon\theta}\}, \overline{M}^\theta; J).$$

Finally, equations (8)–(11), Theorem 2.5 and Proposition 2.1 imply that for $\varepsilon > 0$ sufficiently small,

$$\begin{aligned} \text{rank}_{\Lambda_\omega} \text{H}^*(M; \mathbb{Q}) \otimes_{\mathbb{Q}} \Lambda_\omega &= \text{rank}_{\Lambda_\omega} \text{HFN}^*(\{\varphi_t\}; J) = \text{rank}_{\Lambda_{\theta,\omega}} \text{HFN}^*(\{\psi'_t\}, J') \\ &= \text{rank}_{\widetilde{\Lambda}_{\theta,\omega}} \text{HFN}^*(\{\psi'_t\}, \overline{M}^\theta; J) = \text{rank}_{\widetilde{\Lambda}_{\varepsilon\theta,\omega}} \text{HFN}^*(\{\psi_t^{\varepsilon\theta}\}, \overline{M}^\theta; J) \\ &= \text{rank}_{\Lambda_{\varepsilon\theta,\omega}} \text{HFN}^*(\{\psi_t^{\varepsilon\theta}\}; J) = \text{rank}_{\Lambda_{\varepsilon\theta,\omega}} \text{HN}^*(M, \varepsilon\theta) \otimes_{\Lambda_{\varepsilon\theta}} \Lambda_{\varepsilon\theta,\omega} \\ &= \text{rank}_{\Lambda_{\theta,\omega}} \text{HN}^*(M, \theta) \otimes_{\Lambda_\theta} \Lambda_{\theta,\omega}. \end{aligned}$$

Hence, θ must be exact. Indeed, if $[\theta] \neq 0$, then by Proposition 2.2

$$\begin{aligned} \text{rank}_{\Lambda_\omega} \text{H}^*(M; \mathbb{Q}) \otimes_{\mathbb{Q}} \Lambda_\omega &= \dim_{\mathbb{Q}} \text{H}^*(M; \mathbb{Q}) \\ &> \dim_{\Lambda_\theta} \text{HN}^{*+n}(M, \theta) = \text{rank}_{\Lambda_{\theta,\omega}} \text{HN}^{*+n}(M, \theta) \otimes_{\Lambda_\theta} \Lambda_{\theta,\omega}, \end{aligned}$$

in contradiction with the above equalities. This concludes the proof of Theorem 1.3.

3.2 A proof using a result of McDuff

It follows from McDuff [14, Theorem 1] that if $\{\psi_t\}$ is a symplectic loop, then \mathcal{F}_{c_1} vanishes on the elements of $\pi_1(\mathcal{L}M)$ that are represented by $\{\psi_t(\gamma(s))\}_{s,t \in [0,1]}$ for a loop $\gamma \in \mathcal{L}M$. If $\{\psi_t\}$ has trivial evaluation, then $\{\psi_t(\gamma(s))\}_{s,t \in [0,1]}$ determines a homotopy class $A_\gamma \in \pi_2(M)$ with $\langle c_1(M), A_\gamma \rangle = 0$. If (M, ω) is spherically monotone, for every 1-cycle in M given by a loop γ we have that

$$\langle \widetilde{\text{Flux}}(\{\psi_t\}), [\gamma] \rangle = \langle [\omega], A_\gamma \rangle = \lambda \langle c_1(M), A_\gamma \rangle = 0.$$

Therefore, $\widetilde{\text{Flux}}(\{\psi_t\}) = 0$, which yields once again the conclusion of Theorem 1.3. In addition, note that if $\{\psi_t\}$ has trivial evaluation and flux $[\theta] \neq 0$ we can produce a homotopy class $\alpha \in \pi_1(\mathcal{L}M)$ such that $\alpha \in \ker(\mathcal{F}_\omega + \mathcal{F}_\theta) \cap \ker \mathcal{F}_{c_1}$ while $\mathcal{F}_\theta(\alpha) = -\mathcal{F}_\omega(\alpha) \neq 0$. Indeed the class represented by the loop-of-loops $\alpha(s, t) = \psi_t(\gamma(s))$ satisfies these properties. We are then able to conclude the following.

Corollary 3.1 *If the e -homomorphism is not injective, then*

$$\ker \mathcal{F}_\omega \cap \ker \mathcal{F}_\theta \cap \ker \mathcal{F}_{c_1} \subsetneq \ker(\mathcal{F}_\omega + \mathcal{F}_\theta) \cap \ker \mathcal{F}_{c_1}$$

for all $[\theta] \in \ker e$.

This shows, in hindsight, why the equality of Novikov rings in the spherically monotone setting yielded a proof of injectivity of the e -homomorphism.

Acknowledgements

This work is part of the author's PhD thesis carried out at the Université de Montréal under the supervision of Egor Shelukhin, whom we thank for the inspiring interactions and countless suggestions. We thank Dustin Connery-Grigg for the suggestion to re-examine McDuff's result [14], which yielded an alternative proof of the main theorem. This research was partially supported by Fondation Courtois, by the ISM's excellence scholarship, and by the J Armand Bombardier excellence scholarship. This material is based upon work supported by the National Science Foundation under grant DMS-1928930, while the author was in residence at the Simons Laufer Mathematical Sciences Institute (previously known as MSRI), Berkeley, California during the Fall 2022 semester.

References

- [1] **M S Atallah, E Shelukhin**, *Hamiltonian no-torsion*, *Geom. Topol.* 27 (2023) 2833–2897 MR Zbl
- [2] **A Banyaga**, *Sur la structure du groupe des difféomorphismes qui préservent une forme symplectique*, *Comment. Math. Helv.* 53 (1978) 174–227 MR Zbl
- [3] **Y Cho, M K Kim**, *An example of circle actions on symplectic Calabi–Yau manifolds with non-empty fixed points*, preprint (2013) arXiv 1304.0540
- [4] **A Dimca**, *Singularities and topology of hypersurfaces*, Springer (1992) MR Zbl
- [5] **M Farber**, *Topology of closed one-forms*, *Math. Surv. Monogr.* 108, Amer. Math. Soc., Providence, RI (2004) MR Zbl
- [6] **T Frankel**, *Fixed points and torsion on Kähler manifolds*, *Ann. of Math.* 70 (1959) 1–8 MR Zbl
- [7] **V L Ginzburg**, *Some remarks on symplectic actions of compact groups*, *Math. Z.* 210 (1992) 625–640 MR Zbl
- [8] **D Jang, S Tolman**, *Non-Hamiltonian actions with fewer isolated fixed points*, *Int. Math. Res. Not.* 2023 (2023) 6045–6077 MR Zbl
- [9] **J Kędra, D Kotschick, S Morita**, *Crossed flux homomorphisms and vanishing theorems for flux groups*, *Geom. Funct. Anal.* 16 (2006) 1246–1273 MR Zbl
- [10] **D Kotschick**, *Free circle actions with contractible orbits on symplectic manifolds*, *Math. Z.* 252 (2006) 19–25 MR Zbl
- [11] **F Lalonde, D McDuff, L Polterovich**, *On the flux conjectures*, from “Geometry, topology, and dynamics” (F Lalonde, editor), *CRM Proc. Lecture Notes* 15, Amer. Math. Soc., Providence, RI (1998) 69–85 MR Zbl

- [12] **H V Lê, K Ono**, *Floer–Novikov cohomology and symplectic fixed points, revisited*, Proc. Int. Geom. Cent. 13 (2020) 89–114 MR Zbl
- [13] **G Lupton, J Oprea**, *Cohomologically symplectic spaces: toral actions and the Gottlieb group*, Trans. Amer. Math. Soc. 347 (1995) 261–288 MR Zbl
- [14] **D McDuff**, *Symplectic diffeomorphisms and the flux homomorphism*, Invent. Math. 77 (1984) 353–366 MR Zbl
- [15] **D McDuff**, *The moment map for circle actions on symplectic manifolds*, J. Geom. Phys. 5 (1988) 149–160 MR Zbl
- [16] **D McDuff**, *A survey of the topological properties of symplectomorphism groups*, from “Topology, geometry and quantum field theory” (U Tillmann, editor), Lond. Math. Soc. Lect. Note Ser. 308, Cambridge Univ. Press (2004) 173–193 MR Zbl
- [17] **D McDuff**, *Hamiltonian S^1 –manifolds are uniruled*, Duke Math. J. 146 (2009) 449–507 MR Zbl
- [18] **D McDuff, D Salamon**, *J–holomorphic curves and symplectic topology*, Amer. Math. Soc. Colloq. Publ. 52, Amer. Math. Soc., Providence, RI (2004) MR Zbl
- [19] **D McDuff, D Salamon**, *Introduction to symplectic topology*, 3rd edition, Oxford Univ. Press (2017) MR Zbl
- [20] **K Ono**, *Equivariant projective imbedding theorem for symplectic manifolds*, J. Fac. Sci. Univ. Tokyo Sect. IA Math. 35 (1988) 381–392 MR Zbl
- [21] **K Ono**, *Obstruction to circle group actions preserving symplectic structure*, Hokkaido Math. J. 21 (1992) 99–102 MR Zbl
- [22] **K Ono**, *Floer–Novikov cohomology and symplectic fixed points*, J. Symplectic Geom. 3 (2005) 545–563 MR Zbl
- [23] **K Ono**, *Floer–Novikov cohomology and the flux conjecture*, Geom. Funct. Anal. 16 (2006) 981–1020 MR Zbl
- [24] **L Polterovich**, *Growth of maps, distortion in groups and symplectic geometry*, Invent. Math. 150 (2002) 655–686 MR Zbl
- [25] **L Polterovich**, *Floer homology, dynamics and groups*, from “Morse theoretic methods in nonlinear analysis and in symplectic topology” (P Biran, O Cornea, F Lalonde, editors), NATO Sci. Ser. II Math. Phys. Chem. 217, Springer (2006) 417–438 MR Zbl
- [26] **L Polterovich, D Rosen**, *Function theory on symplectic manifolds*, CRM Monogr. Ser. 34, Amer. Math. Soc., Providence, RI (2014) MR Zbl
- [27] **P Seidel**, *Braids and symplectic four-manifolds with abelian fundamental group*, Turkish J. Math. 26 (2002) 93–100 MR Zbl
- [28] **S Tolman**, *Non-Hamiltonian actions with isolated fixed points*, Invent. Math. 210 (2017) 877–910 MR Zbl
- [29] **S Tolman, J Weitsman**, *On semifree symplectic circle actions with isolated fixed points*, Topology 39 (2000) 299–309 MR Zbl
- [30] **LH Vân, K Ono**, *Symplectic fixed points, the Calabi invariant and Novikov homology*, Topology 34 (1995) 155–176 MR Zbl

*Department of Mathematics and Statistics, University of Montreal
Montreal QC, Canada*

marcelo.sarkis.atallah@umontreal.ca

Received: 20 December 2022 Revised: 14 May 2023

ALGEBRAIC & GEOMETRIC TOPOLOGY

msp.org/agt

EDITORS

PRINCIPAL ACADEMIC EDITORS

John Etnyre
etnyre@math.gatech.edu
Georgia Institute of Technology

Kathryn Hess
kathryn.hess@epfl.ch
École Polytechnique Fédérale de Lausanne

BOARD OF EDITORS

Julie Bergner	University of Virginia jeb2md@eservices.virginia.edu	Robert Lipshitz	University of Oregon lipshitz@uoregon.edu
Steven Boyer	Université du Québec à Montréal cohf@math.rochester.edu	Norihiko Minami	Yamato University minami.norihiko@yamato-u.ac.jp
Tara E Brendle	University of Glasgow tara.brendle@glasgow.ac.uk	Andrés Navas	Universidad de Santiago de Chile andres.navas@usach.cl
Indira Chatterji	CNRS & Univ. Côte d'Azur (Nice) indira.chatterji@math.cnrs.fr	Thomas Nikolaus	University of Münster nikolaus@uni-muenster.de
Alexander Dranishnikov	University of Florida dranish@math.ufl.edu	Robert Oliver	Université Paris 13 bobol@math.univ-paris13.fr
Tobias Ekholm	Uppsala University, Sweden tobias.ekholm@math.uu.se	Jessica S Purcell	Monash University jessica.purcell@monash.edu
Mario Eudave-Muñoz	Univ. Nacional Autónoma de México mario@matem.unam.mx	Birgit Richter	Universität Hamburg birgit.richter@uni-hamburg.de
David Futer	Temple University dfuter@temple.edu	Jérôme Scherer	École Polytech. Féd. de Lausanne jerome.scherer@epfl.ch
John Greenlees	University of Warwick john.greenlees@warwick.ac.uk	Vesna Stojanoska	Univ. of Illinois at Urbana-Champaign vesna@illinois.edu
Ian Hambleton	McMaster University ian@math.mcmaster.ca	Zoltán Szabó	Princeton University szabo@math.princeton.edu
Matthew Hedden	Michigan State University mhedden@math.msu.edu	Maggy Tomova	University of Iowa maggy-tomova@uiowa.edu
Hans-Werner Henn	Université Louis Pasteur henn@math.u-strasbg.fr	Nathalie Wahl	University of Copenhagen wahl@math.ku.dk
Daniel Isaksen	Wayne State University isaksen@math.wayne.edu	Chris Wendl	Humboldt-Universität zu Berlin wendl@math.hu-berlin.de
Thomas Koberda	University of Virginia thomas.koberda@virginia.edu	Daniel T Wise	McGill University, Canada daniel.wise@mcgill.ca
Christine Lescop	Université Joseph Fourier lescop@ujf-grenoble.fr		


See inside back cover or msp.org/agt for submission instructions.

The subscription price for 2024 is US \$705/year for the electronic version, and \$1040/year (+\$70, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Algebraic & Geometric Topology is indexed by Mathematical Reviews, Zentralblatt MATH, Current Mathematical Publications and the Science Citation Index.

Algebraic & Geometric Topology (ISSN 1472-2747 printed, 1472-2739 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840.

AGT peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<https://msp.org/>

© 2024 Mathematical Sciences Publishers

ALGEBRAIC & GEOMETRIC TOPOLOGY

Volume 24 Issue 4 (pages 1809–2387) 2024

Möbius structures, quasimetrics and completeness	1809
MERLIN INCERTI-MEDICI	
$\mathbb{Z}/p \times \mathbb{Z}/p$ actions on $S^n \times S^n$	1841
JIM FOWLER and COURTNEY THATCHER	
\mathbb{Z}_k -stratifolds	1863
ANDRÉS ÁNGEL, CARLOS SEGOVIA and ARLEY FERNANDO TORRES	
Relative systoles in hyperelliptic translation surfaces	1903
CORENTIN BOISSY and SLAVYANA GENINSKA	
Smooth singular complexes and diffeological principal bundles	1913
HIROSHI KIHARA	
Natural symmetries of secondary Hochschild homology	1953
DAVID AYALA, JOHN FRANCIS and ADAM HOWARD	
The shape of the filling-systole subspace in surface moduli space and critical points of the systole function	2011
YUE GAO	
Moduli spaces of geometric graphs	2039
MARA BELOTTI, ANTONIO LERARIO and ANDREW NEWMAN	
Classical shadows of stated skein representations at roots of unity	2091
JULIEN KORINMAN and ALEXANDRE QUESNEY	
Commensurators of thin normal subgroups and abelian quotients	2149
THOMAS KOBERDA and MAHAN MJ	
Pushouts of Dwyer maps are $(\infty, 1)$ -categorical	2171
PHILIP HACKNEY, VIKTORIYA OZORNOVA, EMILY RIEHL and MARTINA ROVELLI	
A variant of a Dwyer–Kan theorem for model categories	2185
BORIS CHORNY and DAVID WHITE	
Integral generalized equivariant cohomologies of weighted Grassmann orbifolds	2209
KOUSHIK BRAHMA and SOUMEN SARKAR	
Projective modules and the homotopy classification of (G, n) -complexes	2245
JOHN NICHOLSON	
Realization of Lie algebras of derivations and moduli spaces of some rational homotopy types	2285
YVES FÉLIX, MARIO FUENTES and ANICETO MURILLO	
On the positivity of twisted L^2 -torsion for 3-manifolds	2307
JIANRU DUAN	
An algebraic C_2 -equivariant Bézout theorem	2331
STEVEN R COSTENOBLE, THOMAS HUDSON and SEAN TILSON	
Topologically isotopic and smoothly inequivalent 2-spheres in simply connected 4-manifolds whose complement has a prescribed fundamental group	2351
RAFAEL TORRES	
Remarks on symplectic circle actions, torsion and loops	2367
MARCELO S ATALLAH	
Correction to the article Hopf ring structure on the mod p cohomology of symmetric groups	2385
LORENZO GUERRA	