

AG
T

*Algebraic & Geometric
Topology*

Volume 24 (2024)

Correction to the article
Hopf ring structure on the mod p cohomology of symmetric groups

LORENZO GUERRA



Correction to the article Hopf ring structure on the mod p cohomology of symmetric groups

LORENZO GUERRA

We correct Propositions 5.4, 5.5 and 5.6 of the author's previous article (*Algebr. Geom. Topol.* 17 (2017) 957–982).

20J06

In our previous paper [1], the formulas describing the action of the mod p Steenrod algebra on Hopf ring generators are incorrect. This is not caused by a flaw in the general argument, but by our misstatement of Hung and Minh's theorem on mod p modular invariants. In this note, we refer to our original article for notation and conventions.

We begin by providing the accurate statement of Hung and Minh's theorem. Although these two authors decided to compute coefficients explicitly, we believe that, for our purpose, their result is best restated using the total Steenrod class $\mathcal{P}^* = \sum_{r \geq 0} \mathcal{P}^r$.

Hung and Minh's theorem [2, Theorem 4.1, page 42] *In $[\mathbb{Z}_p[y_1, \dots, y_n] \otimes \Lambda(x_1, \dots, x_n)]^{\mathrm{Gl}_n(\mathbb{Z}_p)}$, the algebra of modular invariants for $\mathrm{Gl}_n(\mathbb{Z}_p)$, for all $0 \leq s < n$ and $0 \leq s_1 < \dots < s_k < n$,*

$$\begin{aligned} \mathcal{P}^*(d_{s,n-s}) &= (d_{0,n} + \dots + d_{s,n-s})(d_{0,n} + \dots + d_{n-1,1} + 1)^{p-1} - (d_{0,n} + \dots + d_{s-1,n-s+1})^p, \\ \mathcal{P}^*(R_{n,s_1,\dots,s_k}) &= \sum_{\substack{0 \leq t_1 < t_2 < \dots < t_{k+1} \leq n \\ s_{i-1} < t_i \leq s_i}} \sum_{i=1}^{k+1} [(-1)^{k+1-i} R_{n,t_1,\dots,\hat{t}_i,\dots,t_{k+1}} d_{t_i,n-t_i}] (d_{0,n} + \dots + d_{n-1,1} + 1)^{p-2}. \end{aligned}$$

In the expression above, we let, by convention, $s_{-1} = -1$, $s_{k+1} = n$, $R_{n,t_1,\dots,t_k,n} = 0$, and $d_{n,0} = 1$.

We now provide the corrected formulas for the Steenrod powers of generators.

Corrected version of Proposition 5.4 Let $0 \leq k < n$. Let $\mathrm{Outgrowth}(\gamma_{n-k,p^k})$ be the set of full-width Hopf monomials $x \in H^*(\Sigma_{p^n}; \mathbb{Z}_p)$ of type C with $\mathrm{effsc}(x) \geq n - k$ and $\mathrm{ht}(x) \leq p$. Then, for all $0 \leq r \leq p^n - p^k$,

$$\mathcal{P}^r(\gamma_{n-k,p^k}) = \sum_{\substack{x \in \mathrm{Outgrowth}(\gamma_{n-k,p^k}) \\ \deg(x) = 2(p^n - p^k + r(p-1))}} c_{n,k,x} x,$$

where $c_{n,k,x}$ is a scalar coefficient calculated as follows.

If $x = \prod_{i=1}^n \gamma_{i,p^{n-i}}^{e_i}$ ($e_i \geq 0$) is a gathered block, then

$$c_{n,k,x} = (-1)^{n-k+\sum_{i=1}^n ie_i} \frac{1}{p!} \frac{p!}{\prod_{i=0}^n e_i!} \left(\sum_{i=n-k}^n e_i \right),$$

where we let by convention $e_0 = p - \sum_{i=1}^n e_i$ and we reduce $c_{n,k,x} \bmod p$ after calculating its expression in \mathbb{Z} first. If x is not a gathered block, we can write x as a nonzero scalar multiple of a transfer product $b_1 \odot \cdots \odot b_r$, where each b_j is a column of width equal to a power p^{l_j} of p . Then, we put $c_{n,k,x} = \prod_{j=1}^r c_{l_j, l_j-n+k, b_j}$.

Corrected version of Proposition 5.5 • Let $1 \leq j \leq k$. Let $\text{Outgrowth}(\alpha_{j,k})$ be the set of full-width Hopf monomials $x \in H^*(\Sigma_{p^k}; \mathbb{Z}_p)$ of type A with $\text{effsc}(x) \geq k$ and $\text{ht}(x) \leq p$. Then, for all $0 \leq r < p^k - p^{k-j}$,

$$\mathcal{P}^r(\alpha_{j,k}) = \sum_{\substack{x \in \text{Outgrowth}(\alpha_{j,k}) \\ \deg(x) = 2p^k - 2p^{k-j} - 1 + 2r(p-1)}} c'_{k,j,x} x,$$

where $c'_{k,j,x}$ is a scalar coefficient calculated as follows.

If $x = \alpha_{t,k} \prod_{i=1}^k \gamma_{i,p^{k-i}}^{e_i}$ with $e_i \geq 0$ and $1 \leq t \leq k$ is a gathered block, then

$$c'_{k,j,x} = \begin{cases} (-1)^{j+t+\sum_{i=1}^k ie_i} ((p-1)!/\prod_{i=0}^k e_i!) (\sum_{l=j}^k e_l) & \text{if } 1 \leq t < j, \\ -(-1)^{j+t+\sum_{i=1}^k ie_i} ((p-1)!/\prod_{i=0}^k e_i!) (\sum_{l=0}^{j-1} e_l) & \text{if } j \leq t \leq k, \end{cases}$$

where we let by convention $e_0 = p - 1 - \sum_{i=1}^n e_i$. All Hopf monomials $x \in \text{Outgrowth}(\alpha_{j,k})$ have this form.

• Let $1 \leq i < j \leq k$. Let $\text{Outgrowth}(\beta_{i,j,p^{k-j}})$ be the set of full-width Hopf monomials $x \in H^*(\Sigma_{p^k}; \mathbb{Z}_p)$ of type B with $\text{effsc}(x) \geq j$ and $\text{ht}(x) \leq p$. Then, for all $0 \leq r \leq p^k - p^{k-j} - p^{k-i}$,

$$\mathcal{P}^r(\beta_{i,j,p^{k-j}}) = \sum_{\substack{x \in \text{Outgrowth}(\beta_{i,j,p^{k-j}}) \\ \deg(x) = 2(p^k - p^{k-j} - p^{k-i} + r(p-1))}} c''_{k,i,j,x} x,$$

where $c''_{k,i,j,x}$ is a scalar coefficient calculated as follows.

If $x = \beta_{t,u,p^{k-u}} \prod_{i=1}^k \gamma_{i,p^{k-i}}^{e_i}$ with $e_i \geq 0$ and $1 \leq t < u \leq k$ is a gathered block, then

$$c''_{k,i,j,x} = \begin{cases} -(-1)^{i+j+t+u+\sum_{m=1}^k me_m} ((p-1)!/\prod_{m=0}^k e_m!) (\sum_{l=j}^k e_l) & \text{if } 1 \leq t < i \leq u < j \leq k, \\ (-1)^{i+j+t+u+\sum_{m=1}^k me_m} ((p-1)!/\prod_{m=0}^k e_m!) (\sum_{l=i}^{j-1} e_l) & \text{if } 1 \leq t < i < j \leq u \leq k, \\ -(-1)^{i+j+t+u+\sum_{m=1}^k me_m} ((p-1)!/\prod_{m=0}^k e_m!) (\sum_{l=0}^{i-1} e_l) & \text{if } i \leq t < j \leq u \leq k, \\ 0 & \text{otherwise,} \end{cases}$$

where we let by convention $e_0 = p - 1 - \sum_{m=1}^k e_m$. If x is not a gathered block, we can write x as a nonzero scalar multiple of a transfer product $b_1 \odot \cdots \odot b_r$, where each b_m is a column of width equal to a power p^{l_m} of p . Then, we put $c''_{k,i,j,x} = \prod_{m=1}^r c''_{l_m,i,j,b_m}$.

Corrected version of Proposition 5.6 *The following formulas hold:*

- $\beta(\alpha_{j,k}) = \gamma_{k,1}$ if $j = k$ and is equal to 0 otherwise.
- $\beta(\beta_{i,j,p^k}) = \alpha_{i,j} \odot \beta_{i,j,p^k-1}$.
- $\beta(\gamma_{j,p^k}) = 0$.

The proof of these propositions is essentially unchanged from the author's original article, except that we use the correct statement of Hung and Mihن's theorem to compute coefficients.

References

- [1] **L Guerra**, *Hopf ring structure on the mod p cohomology of symmetric groups*, Algebr. Geom. Topol. 17 (2017) 957–982 MR Zbl
- [2] **N H V Hung, P A Minh**, *The action of the mod p Steenrod operations on the modular invariants of linear groups*, Vietnam J. Math. 23 (1995) 39–56 MR Zbl

Dipartimento di Matematica, Università di Roma Tor Vergata
Rome, Italy

guerra@axp.mat.uniroma2.it

Received: 21 December 2023

ALGEBRAIC & GEOMETRIC TOPOLOGY

msp.org/agt

EDITORS

PRINCIPAL ACADEMIC EDITORS

John Etnyre etnyre@math.gatech.edu Georgia Institute of Technology	Kathryn Hess kathryn.hess@epfl.ch École Polytechnique Fédérale de Lausanne
--------------------------------------------------------------------------	----------------------------------------------------------------------------------

BOARD OF EDITORS

Julie Bergner	University of Virginia jeb2mld@eservices.virginia.edu	Robert Lipshitz	University of Oregon lipshitz@uoregon.edu
Steven Boyer	Université du Québec à Montréal cohf@math.rochester.edu	Norihiko Minami	Yamato University minami.norihiko@yamato-u.ac.jp
Tara E Brendle	University of Glasgow tara.brendle@glasgow.ac.uk	Andrés Navas	Universidad de Santiago de Chile andres.navas@usach.cl
Indira Chatterji	CNRS & Univ. Côte d'Azur (Nice) indira.chatterji@math.cnrs.fr	Thomas Nikolaus	University of Münster nikolaus@uni-muenster.de
Alexander Dranishnikov	University of Florida dranish@math.ufl.edu	Robert Oliver	Université Paris 13 bobol@math.univ-paris13.fr
Tobias Ekholm	Uppsala University, Sweden tobias.ekholm@math.uu.se	Jessica S Purcell	Monash University jessica.purcell@monash.edu
Mario Eudave-Muñoz	Univ. Nacional Autónoma de México mario@matem.unam.mx	Birgit Richter	Universität Hamburg birgit.richter@uni-hamburg.de
David Futer	Temple University dfuter@temple.edu	Jérôme Scherer	École Polytech. Féd. de Lausanne jerome.scherer@epfl.ch
John Greenlees	University of Warwick john.greenlees@warwick.ac.uk	Vesna Stojanoska	Univ. of Illinois at Urbana-Champaign vesna@illinois.edu
Ian Hambleton	McMaster University ian@math.mcmaster.ca	Zoltán Szabó	Princeton University szabo@math.princeton.edu
Matthew Hedden	Michigan State University mhedden@math.msu.edu	Maggy Tomova	University of Iowa maggy.tomova@uiowa.edu
Hans-Werner Henn	Université Louis Pasteur henn@math.u-strasbg.fr	Natalie Wahl	University of Copenhagen wahl@math.ku.dk
Daniel Isaksen	Wayne State University isaksen@math.wayne.edu	Chris Wendl	Humboldt-Universität zu Berlin wendl@math.hu-berlin.de
Thomas Koberda	University of Virginia thomas.koberda@virginia.edu	Daniel T Wise	McGill University, Canada daniel.wise@mcgill.ca
Christine Lescop	Université Joseph Fourier lescop@ujf-grenoble.fr		

See inside back cover or msp.org/agt for submission instructions.

The subscription price for 2024 is US \$705/year for the electronic version, and \$1040/year (+\$70, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Algebraic & Geometric Topology is indexed by Mathematical Reviews, Zentralblatt MATH, Current Mathematical Publications and the Science Citation Index.

Algebraic & Geometric Topology (ISSN 1472-2747 printed, 1472-2739 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840.

AGT peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY
 mathematical sciences publishers

nonprofit scientific publishing

<https://msp.org/>

© 2024 Mathematical Sciences Publishers

ALGEBRAIC & GEOMETRIC TOPOLOGY

Volume 24 Issue 4 (pages 1809–2387) 2024

Möbius structures, quasimetrics and completeness MERLIN INCERTI-MEDICI	1809
$\mathbb{Z}/p \times \mathbb{Z}/p$ actions on $S^n \times S^n$ JIM FOWLER and COURTNEY THATCHER	1841
\mathbb{Z}_k -stratifolds ANDRÉS ÁNGEL, CARLOS SEGOVIA and ARLEY FERNANDO TORRES	1863
Relative systoles in hyperelliptic translation surfaces CORENTIN BOISSY and SLAVYANA GENINSKA	1903
Smooth singular complexes and diffeological principal bundles HIROSHI KIHARA	1913
Natural symmetries of secondary Hochschild homology DAVID AYALA, JOHN FRANCIS and ADAM HOWARD	1953
The shape of the filling-systole subspace in surface moduli space and critical points of the systole function YUE GAO	2011
Moduli spaces of geometric graphs MARA BELOTTI, ANTONIO LERARIO and ANDREW NEWMAN	2039
Classical shadows of stated skein representations at roots of unity JULIEN KORINMAN and ALEXANDRE QUESNEY	2091
Commensurators of thin normal subgroups and abelian quotients THOMAS KOBERDA and MAHAN MJ	2149
Pushouts of Dwyer maps are $(\infty, 1)$ -categorical PHILIP HACKNEY, VIKTORIYA OZORNOVA, EMILY RIEHL and MARTINA ROVELLI	2171
A variant of a Dwyer–Kan theorem for model categories BORIS CHORNY and DAVID WHITE	2185
Integral generalized equivariant cohomologies of weighted Grassmann orbifolds KOUSHIK BRAHMA and SOUMEN SARKAR	2209
Projective modules and the homotopy classification of (G, n) -complexes JOHN NICHOLSON	2245
Realization of Lie algebras of derivations and moduli spaces of some rational homotopy types YVES FÉLIX, MARIO FUENTES and ANICETO MURILLO	2285
On the positivity of twisted L^2 -torsion for 3-manifolds JIANRU DUAN	2307
An algebraic C_2 -equivariant Bézout theorem STEVEN R COSTENOBLE, THOMAS HUDSON and SEAN TILSON	2331
Topologically isotopic and smoothly inequivalent 2-spheres in simply connected 4-manifolds whose complement has a prescribed fundamental group RAFAEL TORRES	2351
Remarks on symplectic circle actions, torsion and loops MARCELO S ATALLAH	2367
Correction to the article Hopf ring structure on the mod p cohomology of symmetric groups LORENZO GUERRA	2385