

AG  
T

*Algebraic & Geometric  
Topology*

Volume 24 (2024)

**Algebraic generators of the skein algebra of a surface**

RAMANUJAN SANTHAROUBANE





# Algebraic generators of the skein algebra of a surface

RAMANUJAN SANTHAROUBANE

Let  $\Sigma$  be a surface with negative Euler characteristic, genus at least one and at most one boundary component. We prove that the Kauffman bracket skein algebra of  $\Sigma$  over the field of rational functions can be algebraically generated by a finite number of simple closed curves that are naturally associated to certain generators of the mapping class group of  $\Sigma$ . The action of the mapping class group on the skein algebra gives canonical relations between these generators. From this, we conjecture a presentation for a skein algebra of  $\Sigma$ .

57K31

## 1 Introduction

### 1.1 Main results

This paper is focused on finding algebraic generators of the Kauffman bracket skein algebra of a surface. Throughout this paper, we will refer to the Kauffman bracket skein algebra simply as the skein algebra. Let  $\Sigma$  be a compact oriented connected surface of genus at least one and with at most one boundary component. Moreover, we will suppose that  $\Sigma$  has negative Euler characteristic. We denote by  $\mathcal{S}(\Sigma, \mathbb{Q}(A))$  the skein module of  $\Sigma \times [0, 1]$  with coefficients in the field of rational function  $\mathbb{Q}(A)$  and by  $S(\Sigma)$  the skein module over  $\mathbb{Z}[A^{\pm 1}]$ . The module  $\mathcal{S}(\Sigma, \mathbb{Q}(A))$  is equipped with a natural product given by stacking banded links. For  $\gamma$  is a simple closed curve on  $\Sigma$ , we write  $\gamma$  for the element  $\gamma \times [\frac{1}{2}, \frac{2}{3}]$  in  $\mathcal{S}(\Sigma, \mathbb{Q}(A))$  and we denote by  $t_\gamma$  the Dehn twist along  $\gamma$ .

**Theorem 1.1** *Let  $\{\gamma_j\}_{j \in I}$  be a finite set of nonseparating simple closed curves such that*

- (1) *for any  $i, j \in I$ , the curves  $\gamma_i$  and  $\gamma_j$  intersect at most once;*
- (2) *the set  $\{t_{\gamma_j}\}_{j \in I}$  generates the mapping class group of  $\Sigma$ .*

*Then  $\{\gamma_j\}_{j \in I}$  generates  $\mathcal{S}(\Sigma, \mathbb{Q}(A))$  as a  $\mathbb{Q}(A)$ -algebra. Moreover,  $\{\gamma_j\}_{j \in I}$  generates*

$$S(\Sigma) \otimes \mathbb{Z} \left[ A^{\pm 1}, \frac{1}{A^2 - A^{-2}} \right]$$

*as a  $\mathbb{Z}[A^{\pm 1}, 1/(A^2 - A^{-2})]$ -algebra.*

We recall that the mapping class group of  $\Sigma$  is  $\pi_0(\text{Homeo}^+(\Sigma, \partial\Sigma))$ . We will now give an interpretation of some relations that should hold for the generators in the previous theorem. Let us fix  $\{\gamma_j\}_{j \in I}$  a set of simple closed curves on  $\Sigma$  satisfying the hypothesis of Theorem 1.1. Let  $\mathbb{Q}(A)\langle I \rangle$  be the free

noncommutative  $\mathbb{Q}(A)$ -algebra generated by  $\{X_j\}_{j \in I}$ . The theorem says that there exists a surjective algebra homomorphism

$$(1) \quad \Psi: \mathbb{Q}(A)\langle I \rangle \rightarrow \mathcal{S}(\Sigma, \mathbb{Q}(A))$$

defined by

$$\Psi(X_j) = \frac{\gamma_j}{A^2 - A^{-2}} \quad \text{for all } j \in I.$$

Now for  $j \in I$  and  $\epsilon \in \{-1, 1\}$ , let  $T_j^\epsilon: \mathbb{Q}(A)\langle I \rangle \rightarrow \mathbb{Q}(A)\langle I \rangle$  be the homomorphism of  $\mathbb{Q}(A)$ -algebra defined by

$$(2) \quad T_j^\epsilon(X_k) = \begin{cases} X_k & \text{if } \iota(\gamma_j, \gamma_k) = 0, \\ \epsilon(A^\epsilon X_j X_k - A^{-\epsilon} X_k X_j) & \text{if } \iota(\gamma_j, \gamma_k) = 1. \end{cases}$$

Here  $\iota$  is the geometric intersection of simple closed curves. With the convention that a Dehn twist always turn to the right with respect to the orientation of the surface, we can check that  $\Psi$  exchanges the actions of the  $T_j$  and the  $t_j$  in the sense that

$$\Psi(T_j^\epsilon X) = t_{\gamma_j}^\epsilon(\Psi(X)) \quad \text{for all } X \in \mathbb{Q}(A)\langle I \rangle, j \in I, \text{ and } \epsilon \in \{-1, 1\}.$$

Let  $\Gamma(\Sigma)$  be the mapping class group of  $\Sigma$  and let  $\bar{\Gamma}(\Sigma)$  be the group  $\Gamma(\Sigma)$  modulo its center. Suppose that  $I = \{1, \dots, N\}$  and let us denote each  $t_{\gamma_j}$  simply by  $t_j$ . Note that the map  $t_j^\epsilon \mapsto T_j^\epsilon$  does not extend to an action of  $\bar{\Gamma}(\Sigma)$  on  $\mathbb{Q}(A)\langle I \rangle$ . Indeed  $\mathbb{Q}(A)\langle I \rangle$  is a free noncommutative algebra and the relations in  $\bar{\Gamma}(\Sigma)$  satisfied by the Dehn twists  $\{t_j^{\pm 1}\}_{j \in J}$  have no reason to be satisfied by the operators  $\{T_j^{\pm 1}\}_{j \in J}$ . Hence the relations between the  $\{t_j^{\pm 1}\}_{j \in J}$  give relations between the generators  $\{\gamma_j\}_{j \in J}$  in  $\mathcal{S}(\Sigma, \mathbb{Q}(A))$ . Suppose that  $\bar{\Gamma}(\Sigma)$  has the following presentation with respect to the generators  $\{t_j\}_{j \in I}$ :

$$\bar{\Gamma}(\Sigma) = \langle t_1, \dots, t_N \mid R_1(t_1, \dots, t_N) = \dots = R_K(t_1, \dots, t_N) = 1 \rangle$$

where  $K$  is an integer and the  $R_k(t_1, \dots, t_N)$  are some words in  $\{t_j^{\pm 1}\}_{j \in I}$ . Let  $\mathcal{R}$  be the bi-ideal of  $\mathbb{Q}(A)\langle I \rangle$  generated by the elements

$$\begin{aligned} R_k(T_1, \dots, T_N)X_i - X_i & \quad \text{for } 1 \leq i \leq N \text{ and } 1 \leq k \leq K, \\ T_j T_j^{-1} X_i - X_i & \quad \text{for } 1 \leq i, j \leq N, \\ X_i X_j - X_j X_i & \quad \text{for } \iota(\gamma_i, \gamma_j) = 0. \end{aligned}$$

We define

$$(3) \quad \mathcal{A}(\Gamma(\Sigma)) = \frac{\mathbb{Q}(A)\langle I \rangle}{\mathcal{R}},$$

which is a quotient of  $\mathbb{Q}(A)\langle I \rangle$  on which the actions of the  $T_j^{\pm 1}$  extend to a canonical action of  $\bar{\Gamma}(\Sigma)$ . A direct consequence of Theorem 1.1 is the following:

**Corollary 1.2** *The canonical map*

$$\mathcal{A}(\Gamma(\Sigma)) \rightarrow \mathcal{S}(\Sigma, \mathbb{Q}(A))$$

*is surjective.*

**Conjecture 1.3** *There exists a presentation of  $\overline{\Gamma}(\Sigma)$  for which  $\mathcal{A}(\Gamma(\Sigma))$  is isomorphic to  $\mathcal{S}(\Sigma, \mathbb{Q}(A))$  as a noncommutative  $\mathbb{Q}(A)$ -algebra.*

## 1.2 Notes and references

Bullock [1999] was the first to find algebraic generators of the skein algebra of a surface. His generators are over  $\mathbb{Z}[A^{\pm 1}]$  and not over  $\mathbb{Q}(A)$ . The number of his generators is exponential in the genus of the surface whereas here we have a linear number (by choosing the right generators of  $\Gamma(\Sigma)$ ).

It was shown in [Przytycki and Sikora 2000] that each  $\mathcal{S}(\Sigma)$  has a generating set of cardinality which is cubic in the genus of the surface.

Finite generation was also proved by Abdiel and Frohman [2017, Theorem 3.7]. Frohman and Kania-Bartoszyńska [2018] studied the skein algebra when  $A$  is evaluated at a root of unity. They proved that it is generated over its center by a pair of subalgebras from pants decomposition. Their generators have some similarities with the one in the current paper.

Presentations of skein algebras of surfaces are only known in genus zero and one. Bullock and Przytycki [2000] found such a presentation for the one-holed torus, the four-holed sphere and two-holed torus. They related some of these algebras to nonstandard deformations of Lie algebras.

When  $A$  is specialized to  $-1$ , it was shown by Bullock [1997] and Przytycki and Sikora [2000] that the skein algebra of a surface is isomorphic to the ring of algebraic functions of the  $\mathrm{SL}(2, \mathbb{C})$  character variety of the surface. Moreover, for  $A = \sqrt{-1}$ , Marché [2011] gave an homological interpretation of the skein algebra of the surface. Note that the map  $\Psi$  defined in (1) is not defined if  $A$  is specialized to a 4<sup>th</sup> primitive root of unity. It is possible to see that if we specialize  $A$  at a 4<sup>th</sup> root of unity in the algebra  $\mathcal{A}(\Gamma(\Sigma))$  we find something different from the algebras studied by Bullock, Marché, Przytycki and Sikora.

Humphries generators [1979] and Lickorish generators [1964] are examples of generators of the mapping class groups satisfying the hypothesis of Theorem 1.1. Moreover, presentations for both of these generating sets are known; we refer to book of Farb and Margalit [2012] for more details.

We consider  $\Gamma(\Sigma)$  quotiented by its center because the center of the mapping class group acts trivially on the skein algebra of the surface.

## Acknowledgements

I want to thank C Frohman, J Marché and G Masbaum for helpful conversations.

## 2 Quick review of the skein algebra

For any compact oriented manifold  $M$  (maybe with boundary), we denote by  $\mathcal{S}(M)$  the Kauffman bracket skein module with coefficients in  $\mathbb{Z}[A^{\pm 1}]$ . We recall that it is the free  $\mathbb{Z}[A^{\pm 1}]$ -module generated by the




Figure 1: Kauffman triple.

set of isotopy classes of banded links in the interior of  $M$  quotiented by the following so-called skein relations. First

$$(4) \quad L_x = AL_\infty + A^{-1}L_0$$

where  $L_x, L_0$  and  $L_\infty$  are any three banded links in  $M$  which are the same outside a small 3–ball but differ inside as in Figure 1. In this case, the triple  $L_x, L_0, L_\infty$  is called a Kauffman triple. The second relation satisfied in  $\mathcal{S}(M)$  is

$$L \cup D = -(A^2 + A^{-2})L$$

where  $L$  is any link in  $M$  and  $D$  is a trivial banded knot. We define  $\mathcal{S}(M, \mathbb{Q}(A))$  to be the  $\mathbb{Q}(A)$ –vector space  $\mathcal{S}(M) \otimes \mathbb{Q}(A)$  where the tensor product is made over  $\mathbb{Z}[A^{\pm 1}]$ .

Let  $\Sigma$  be a compact oriented connected surface; we denote by  $\mathcal{S}(\Sigma, \mathbb{Q}(A))$  the space  $\mathcal{S}(\Sigma \times [0, 1], \mathbb{Q}(A))$ . Stacking banded links on top of each other gives  $\mathcal{S}(\Sigma, \mathbb{Q}(A))$  the structure of a  $\mathbb{Q}(A)$ –algebra.

A multiloop is a disjoint union of non-null-homotopic simple close curve inside  $\Sigma$ . For  $\gamma \subset \Sigma$  a multiloop we write  $\gamma$  for the banded link  $\gamma \times [\frac{1}{2}, \frac{2}{3}]$  in  $\mathcal{S}(\Sigma, \mathbb{Q}(A))$  and we will still call this banded link a multiloop. A well-known theorem is the following:

**Theorem 2.1** *The set of isotopy classes of multiloops on  $\Sigma$  is a basis of the  $\mathbb{Q}(A)$ –vector space  $\mathcal{S}(\Sigma, \mathbb{Q}(A))$ .*

In particular this theorem clearly implies that simple closed curves generate  $\mathcal{S}(\Sigma, \mathbb{Q}(A))$  as an algebra. Recall that  $\Gamma(\Sigma) = \pi_0(\text{Homeo}^+(\Sigma, \partial\Sigma))$  acts canonically on  $\mathcal{S}(\Sigma, \mathbb{Q}(A))$ . If  $\gamma \subset \Sigma$  is a simple closed curve, we denote by  $t_\gamma$  the Dehn twist along  $\gamma$ . We use the turn right convention for  $t_\gamma$ . More precisely, let  $\mathcal{N} \subset \Sigma$  be an annulus neighborhood of  $\gamma$ , we can identify  $\mathcal{N}$  with  $S^1 \times [0, 1]$  using an orientation preserving homeomorphism. Outside  $\mathcal{N}$ , the map  $t_\gamma$  is defined to be the identity and on  $\mathcal{N}$  it is given by the map  $(e^{i\theta}, t) \mapsto (e^{i(\theta-2\pi t)}, t)$ . Figure 2 helps to visualize this definition.

The following lemma can be obtained by applying the skein relation (4).




Figure 2

**Lemma 2.2** Let  $\alpha$  and  $\beta$  be two simple close curves intersecting once. Then in  $\mathcal{S}(\Sigma, \mathbb{Q}(A))$ ,

$$t_\alpha^\epsilon(\beta) = \frac{A^\epsilon \alpha \beta - A^{-\epsilon} \beta \alpha}{\epsilon(A^2 - A^{-2})}.$$

### 3 Proof of Theorem 1.1

Let  $\Gamma(\Sigma)$  be the mapping class group of  $\Sigma$ . Let  $\{\gamma_j\}_{j \in I}$  be a set of simple closed curves satisfying the hypothesis of the Theorem 1.1 and let  $\mathcal{B}$  be the subalgebra of  $\mathcal{S}(\Sigma, \mathbb{Q}(A))$  generated by  $\{\gamma_j\}_{j \in I}$ .

**Lemma 3.1**  $\mathcal{B}$  is stable by the action of  $\Gamma(\Sigma)$ .

**Proof** Since  $\{\gamma_j\}_{j \in I}$  generates  $\Gamma(\Sigma)$ , it enough to prove that for any  $j, k \in I$  we have  $t_{\gamma_j}^{\pm 1}(\gamma_k) \in \mathcal{B}$ . If  $\gamma_j$  does not intersect  $\gamma_k$  then  $t_{\gamma_j}^{\pm 1}(\gamma_k) = \gamma_k \in \mathcal{B}$ . Now if  $\gamma_j$  intersects  $\gamma_k$  once then, by Lemma 2.2,

$$t_{\gamma_j}^{\pm 1}(\gamma_k) = \frac{A^{\pm 1} \gamma_j \gamma_k - A^{\mp 1} \gamma_k \gamma_j}{\pm(A^2 - A^{-2})} \in \mathcal{B}. \quad \square$$

**Lemma 3.2** If  $\mathcal{C}_0$  denotes the set of nonseparating simple closed curves then  $\mathcal{C}_0 \subset \mathcal{B}$ .

**Proof** Let  $\gamma \in \mathcal{C}_0$  and  $\gamma_0 \in \{\gamma_j\}_{j \in I}$ ; there exists  $\phi \in \Gamma(\Sigma)$  such that  $\phi(\gamma_0) = \gamma$ . Since  $\gamma_0$  belongs to  $\mathcal{B}$  which is stable by the action of  $\Gamma(\Sigma)$  (see the previous lemma), we have  $\gamma \in \mathcal{B}$ . □

**Lemma 3.3** If  $\mathcal{C}_1$  denotes the set of separating simple closed curves then  $\mathcal{C}_1 \subset \mathcal{B}$ .

**Proof** Suppose that the genus of  $\Sigma$  is  $g \geq 1$ . Let  $\delta_1, \dots, \delta_g$  be the curves in Figure 3, where  $\delta_g$  is trivial when  $\Sigma$  does not have boundary. Let  $j \in \{1, \dots, g\}$  and let  $z_j$  and  $z'_j$  be the two nonseparating curves in the torus with two boundary components defined by  $\delta_j, \delta_{j-1}$ , as shown in Figure 4.

By applying the skein relations, we have

$$z'_j z_j = A^2 x_j x'_j + A^{-2} y_j y'_j + \delta_j + \delta_{j-1}$$

where  $\delta_0 = -A^2 - A^{-2}$  and  $x_j, x'_j, y_j$  and  $y'_j$  are nonseparating curves. By Lemma 3.2,  $z'_j, z_j, x_j, x'_j, y_j$  and  $y'_j$  are in  $\mathcal{B}$ , so by an induction on  $j$  we can prove that for all  $1 \leq j \leq g$  we have  $\delta_j \in \mathcal{B}$ .

Now if  $\gamma$  is a separating curve, there exists  $\phi \in \Gamma(\Sigma)$  and  $j_0$  such that  $\gamma = \phi(\delta_{j_0})$ . Since  $\mathcal{B}$  is stable by the action of  $\Gamma(\Sigma)$ , we have  $\gamma \in \mathcal{B}$ . □




Figure 3




Figure 4

**Proof of Theorem 1.1**  $\mathcal{S}(\Sigma, \mathbb{Q}(A))$  is algebraically generated by simple closed curves. Hence, combining Lemmas 3.2 and 3.3, we can conclude the proof. Moreover, we can still conclude by Lemmas 3.1, 3.2 and 3.3 that the  $\mathbb{Z}[A^{\pm 1}, 1/(A^2 - A^{-2})]$ -algebra generated by  $\{\gamma_j\}_{j \in I}$  is

$$S(\Sigma) \otimes \mathbb{Z}\left[A^{\pm 1}, \frac{1}{A^2 - A^{-2}}\right]. \quad \square$$

### 4 Interpretation of the relations in the skein algebra

Let  $\{\gamma_j\}_{j \in I}$  be a set of simple closed curves on  $\Sigma$  satisfying the hypothesis of Theorem 1.1. Recall that  $\mathbb{Q}(A)\langle I \rangle$  is the free noncommutative  $\mathbb{Q}(A)$ -algebra generated by  $\{X_j\}_{j \in I}$ .

**Definition 4.1** For  $X, Y \in \mathbb{Q}(A)\langle I \rangle$  we define  $[X, Y]_A := AXY - A^{-1}YX$ .

Recall that the maps  $\{T_j^\epsilon\}_{j \in I}$  are defined by (2) in the introduction. Recall also that given a presentation of  $\bar{\Gamma}(\Sigma)$  with respect to the generating set  $\{t_{\gamma_j}\}_{j \in I}$ , we defined  $\mathcal{A}(\Gamma(\Sigma))$  via (3) (see the introduction). By definition, any relation satisfied by the  $\{t_{\gamma_j}\}_{j \in I}$  (which may not appear in the given presentation) gives some relation in  $\mathcal{A}(\Gamma(\Sigma))$ . Let us focus on the relations

$$(5) \quad T_j^{-1}T_i^{-1}T_j^{-1}T_iT_jT_iX_a - X_a = 0 \quad \text{for } \iota(\gamma_i, \gamma_j) = 1 \text{ and } a \in I,$$

$$(6) \quad T_jT_j^{-1}X_i - X_i = 0 \quad \text{for } i, j \in I,$$

$$(7) \quad X_iX_j - X_jX_i = 0 \quad \text{for } \iota(\gamma_i, \gamma_j) = 0.$$

Note that these relations hold but are not a complete set of relations in  $\mathcal{A}(\Gamma(\Sigma))$ . The first relation comes from the braid relations in the mapping class group.

**Proposition 4.2** In  $\mathcal{A}(\Gamma(\Sigma))$ , the relations (5), (6) and (7) are equivalent to

$$(8) \quad [[X_j, X_i]_A, X_j]_A = X_i \quad \text{for } \iota(\gamma_i, \gamma_j) = 1,$$

$$(9) \quad X_iX_j - X_jX_i = 0 \quad \text{for } \iota(\gamma_i, \gamma_j) = 0.$$



**Proof** Let  $i, j \in I$ . Note that if  $\iota(\gamma_i, \gamma_j) = 0$ , the relation (6) is empty and if  $\iota(\gamma_i, \gamma_j) = 1$ , this relation gives  $[[X_j, X_i]_A, X_j]_A = X_i$ .

Let  $i, j \in I$  such that  $\iota(\gamma_i, \gamma_j) = 1$ . Because of (6), the relation (5) can be rewritten as

$$T_i T_j T_i X_k = T_j T_i T_j X_k$$

for all  $k \in I$ . It is easy to check that this relation is implied by (8) and (9).  $\square$

**Remark 4.3** We did not include  $T_j^{-1} T_j X_i = X_i$  in the relations defining  $\mathcal{A}(\Gamma(\Sigma))$  because they give  $[X_j, [X_i, X_j]_A]_A = X_i$  for  $\iota(\gamma_i, \gamma_j) = 1$  which is the same as  $[[X_j, X_i]_A, X_j]_A = X_i$ .

#### 4.1 The case of the one-holed torus

Let  $\Sigma$  be a surface of genus one with one boundary component. Its mapping class group is the braid group  $B_3$  whose presentation is  $\langle t_1, t_2 \mid t_1 t_2 t_1 = t_2 t_1 t_2 \rangle$ . Here  $t_1$  is Dehn twist along the canonical meridian of  $\Sigma$  and  $t_2$  is the Dehn twist around the longitude of  $\Sigma$ . Note that these two curves satisfy the hypothesis of Theorem 1.1. The center of this group is the group generated by  $(t_1 t_2 t_1)^2$  and  $\bar{\Gamma}(\Sigma)$  is  $\text{PSL}_2(\mathbb{Z})$  with presentation

$$\bar{\Gamma}(\Sigma) = \langle t_1, t_2 \mid t_1 t_2 t_1 = t_2 t_1 t_2, (t_1 t_2 t_1)^2 = 1 \rangle.$$

In this case  $\mathcal{A}(\Gamma(\Sigma))$  is a noncommutative algebra generated by  $X_1$  and  $X_2$ . Because of Proposition 4.2, the only relations between  $X_1$  and  $X_2$  are

$$[X_1, [X_2, X_1]_A]_A = X_2, \quad [X_2, [X_1, X_2]_A]_A = X_1, \quad (T_1 T_2 T_1)^2 X_1 = X_1, \quad (T_1 T_2 T_1)^2 X_2 = X_2.$$

It is easy to check that the two last relations are implied by the two first one. Therefore,

$$\mathcal{A}(\Gamma(\Sigma)) = \langle X_1, X_2 \mid [X_1, [X_2, X_1]_A]_A = X_2, [X_2, [X_1, X_2]_A]_A = X_1 \rangle.$$

From [Bullock 1999, Theorem 2.1] the skein module of the one-holed torus is isomorphic to  $\mathcal{A}(\Gamma(\Sigma))$ . Therefore Conjecture 1.3 holds for the one-holed torus.

## References

- [Abdiel and Frohman 2017] **N Abdiel, C Frohman**, *The localized skein algebra is Frobenius*, *Algebr. Geom. Topol.* 17 (2017) 3341–3373 MR Zbl
- [Bullock 1997] **D Bullock**, *Rings of  $\text{SL}_2(\mathbb{C})$ -characters and the Kauffman bracket skein module*, *Comment. Math. Helv.* 72 (1997) 521–542 MR Zbl
- [Bullock 1999] **D Bullock**, *A finite set of generators for the Kauffman bracket skein algebra*, *Math. Z.* 231 (1999) 91–101 MR Zbl
- [Bullock and Przytycki 2000] **D Bullock, JH Przytycki**, *Multiplicative structure of Kauffman bracket skein module quantizations*, *Proc. Amer. Math. Soc.* 128 (2000) 923–931 MR Zbl

- [Farb and Margalit 2012] **B Farb, D Margalit**, *A primer on mapping class groups*, Princeton Math. Ser. 49, Princeton Univ. Press (2012) MR Zbl
- [Frohman and Kania-Bartoszyńska 2018] **C Frohman, J Kania-Bartoszyńska**, *The structure of the Kauffman bracket skein algebra at roots of unity*, Math. Z. 289 (2018) 889–920 MR Zbl
- [Humphries 1979] **S P Humphries**, *Generators for the mapping class group*, from “Topology of low-dimensional manifolds” (R A Fenn, editor), Lecture Notes in Math. 722, Springer (1979) 44–47 MR Zbl
- [Lickorish 1964] **W B R Lickorish**, *A finite set of generators for the homeotopy group of a 2–manifold*, Proc. Cambridge Philos. Soc. 60 (1964) 769–778 MR Zbl
- [Marché 2011] **J Marché**, *The Kauffman skein algebra of a surface at  $\sqrt{-1}$* , Math. Ann. 351 (2011) 347–364 MR Zbl
- [Przytycki and Sikora 2000] **J H Przytycki, A S Sikora**, *On skein algebras and  $Sl_2(\mathbb{C})$ –character varieties*, Topology 39 (2000) 115–148 MR Zbl

Laboratoire Mathématiques d’Orsay, CNRS, Université Paris-Saclay  
Orsay, France

Current address: Department of Mathematics, University of Virginia  
Charlottesville, VA, United States

ramanujan.santharoubane@gmail.com

Received: 4 March 2022      Revised: 21 November 2022

# ALGEBRAIC & GEOMETRIC TOPOLOGY

[msp.org/agt](https://msp.org/agt)

## EDITORS

### PRINCIPAL ACADEMIC EDITORS

John Etnyre  
etnyre@math.gatech.edu  
Georgia Institute of Technology

Kathryn Hess  
kathryn.hess@epfl.ch  
École Polytechnique Fédérale de Lausanne

### BOARD OF EDITORS

Julie Bergner	University of Virginia jeb2md@eservices.virginia.edu	Robert Lipshitz	University of Oregon lipshitz@uoregon.edu
Steven Boyer	Université du Québec à Montréal cohf@math.rochester.edu	Norihiko Minami	Yamato University minami.norihiko@yamato-u.ac.jp
Tara E Brendle	University of Glasgow tara.brendle@glasgow.ac.uk	Andrés Navas	Universidad de Santiago de Chile andres.navas@usach.cl
Indira Chatterji	CNRS & Univ. Côte d'Azur (Nice) indira.chatterji@math.cnrs.fr	Thomas Nikolaus	University of Münster nikolaus@uni-muenster.de
Alexander Dranishnikov	University of Florida dranish@math.ufl.edu	Robert Oliver	Université Paris 13 bobol@math.univ-paris13.fr
Tobias Ekholm	Uppsala University, Sweden tobias.ekholm@math.uu.se	Jessica S Purcell	Monash University jessica.purcell@monash.edu
Mario Eudave-Muñoz	Univ. Nacional Autónoma de México mario@matem.unam.mx	Birgit Richter	Universität Hamburg birgit.richter@uni-hamburg.de
David Futer	Temple University dfuter@temple.edu	Jérôme Scherer	École Polytech. Féd. de Lausanne jerome.scherer@epfl.ch
John Greenlees	University of Warwick john.greenlees@warwick.ac.uk	Vesna Stojanoska	Univ. of Illinois at Urbana-Champaign vesna@illinois.edu
Ian Hambleton	McMaster University ian@math.mcmaster.ca	Zoltán Szabó	Princeton University szabo@math.princeton.edu
Matthew Hedden	Michigan State University mhedden@math.msu.edu	Maggy Tomova	University of Iowa maggy-tomova@uiowa.edu
Hans-Werner Henn	Université Louis Pasteur henn@math.u-strasbg.fr	Nathalie Wahl	University of Copenhagen wahl@math.ku.dk
Daniel Isaksen	Wayne State University isaksen@math.wayne.edu	Chris Wendl	Humboldt-Universität zu Berlin wendl@math.hu-berlin.de
Thomas Koberda	University of Virginia thomas.koberda@virginia.edu	Daniel T Wise	McGill University, Canada daniel.wise@mcgill.ca
Christine Lescop	Université Joseph Fourier lescop@ujf-grenoble.fr		

---

See inside back cover or [msp.org/agt](https://msp.org/agt) for submission instructions.


The subscription price for 2024 is US \$705/year for the electronic version, and \$1040/year (+\$70, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Algebraic & Geometric Topology is indexed by Mathematical Reviews, Zentralblatt MATH, Current Mathematical Publications and the Science Citation Index.

Algebraic & Geometric Topology (ISSN 1472-2747 printed, 1472-2739 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840.

---

AGT peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<https://msp.org/>

© 2024 Mathematical Sciences Publishers

# ALGEBRAIC & GEOMETRIC TOPOLOGY

Volume 24 Issue 5 (pages 2389–2970) 2024

---

Formal contact categories	2389
BENJAMIN COOPER	
Comparison of period coordinates and Teichmüller distances	2451
IAN FRANKEL	
Topological Hochschild homology of truncated Brown–Peterson spectra, I	2509
GABRIEL ANGELINI-KNOLL, DOMINIC LEON CULVER and EVA HÖNING	
Points of quantum $SL_n$ coming from quantum snakes	2537
DANIEL C DOUGLAS	
Algebraic generators of the skein algebra of a surface	2571
RAMANUJAN SANTHAROUBANE	
Bundle transfer of $L$ -homology orientation classes for singular spaces	2579
MARKUS BANAGL	
A reduction of the string bracket to the loop product	2619
KATSUHIKO KURIBAYASHI, TAKAHITO NAITO, SHUN WAKATSUKI and TOSHIHIRO YAMAGUCHI	
Asymptotic dimensions of the arc graphs and disk graphs	2655
KOJI FUJIWARA and SAUL SCHLEIMER	
Representation stability for homotopy automorphisms	2673
ERIK LINDELL and BASHAR SALEH	
The strong Haken theorem via sphere complexes	2707
SEBASTIAN HENSEL and JENNIFER SCHULTENS	
What are GT-shadows?	2721
VASILY A DOLGUSHEV, KHANH Q LE and AIDAN A LORENZ	
A simple proof of the Crowell–Murasugi theorem	2779
THOMAS KINDRED	
The Burau representation and shapes of polyhedra	2787
ETHAN DLUGIE	
Turning vector bundles	2807
DIARNUID CROWLEY, CSABA NAGY, BLAKE SIMS and HUIJUN YANG	
Rigidification of cubical quasicategories	2851
PIERRE-LOUIS CURIEN, MURIEL LIVERNET and GABRIEL SAADIA	
Tautological characteristic classes, I	2889
JAN DYMARA and TADEUSZ JANUSZKIEWICZ	
Homotopy types of suspended 4-manifolds	2933
PENGCHENG LI	
The braid indices of the reverse parallel links of alternating knots	2957
YUANAN DIAO and HUGH MORTON	