

AG
T

*Algebraic & Geometric
Topology*

Volume 24 (2024)

A reduction of the string bracket to the loop product

KATSUHIKO KURIBAYASHI

TAKAHITO NAITO

SHUN WAKATSUKI

TOSHIHIRO YAMAGUCHI



A reduction of the string bracket to the loop product

KATSUHIKO KURIBAYASHI

TAKAHITO NAITO

SHUN WAKATSUKI

TOSHIHIRO YAMAGUCHI

The negative cyclic homology for a differential graded algebra over the rational field has a quotient of the Hochschild homology as a direct summand if the S -action is trivial. With this fact, we show that the string bracket in the sense of Chas and Sullivan is reduced to the loop product followed by the BV operator on the loop homology provided the given manifold is *BV-exact*. The reduction is indeed derived from the equivalence between the BV-exactness and the triviality of the S -action. Moreover, it is proved that a Lie bracket on the loop cohomology of the classifying space of a connected compact Lie group possesses the same reduction. By using these results, we consider the nontriviality of string brackets. We also show that a simply connected space with positive weights is BV-exact. Furthermore, the *higher BV-exactness* is discussed featuring the cobar-type Eilenberg–Moore spectral sequence.

55P35, 55P50, 55T20

1. Introduction	2620
2. String brackets described in terms of the Hochschild homology	2624
3. Preliminaries	2630
4. Proofs of assertions	2634
5. The string brackets for formal spaces	2639
6. Computation of the string bracket for a nonformal space	2641
7. The cobar-type EMSS and r -BV-exactness	2645
Appendix A. A non-BV-exact space	2649
Appendix B. Connes' B -map in the Gysin exact sequence	2650
List of symbols	2651
References	2652

1 Introduction

Let LM be the free loop space, namely, the space of continuous maps from the circle S^1 to a space M with compact-open topology. The rotation on the domain space S^1 of LM induces an S^1 -action on LM . Then we have the S^1 -equivariant homology $H_*^{S^1}(LM) = H_*(ES^1 \times_{S^1} LM)$ for a space M . The *string bracket* is a Lie bracket on the S^1 -equivariant homology of the free loop space LM of an orientable closed manifold M , which is introduced by Chas and Sullivan in [8]. The bracket is defined by using the loop product on the loop homology $H_*(LM)$ and maps in the Gysin exact sequence of the S^1 -principal bundle

$$(1.1) \quad S^1 \rightarrow ES^1 \times LM \rightarrow ES^1 \times_{S^1} LM.$$

In particular, the Batalin–Vilkovisky (BV) identity of the BV operator on the loop homology induces the Jacobi identity for the string bracket; see the proof of [8, Theorem 6.1].

As for computations of the string brackets, Basu [2] and Félix, Thomas and Vigué-Poirrier [19] have determined explicitly the rational string bracket of the product of spheres. For a simply connected closed manifold M whose rational cohomology is generated by a single element, the rational string bracket is trivial though the rational loop product of M is highly nontrivial; see [2, Theorem 3.4] and [19, Section 5.2, Example 1]. On the other hand, a result due to Tabing [43] shows that the integral string bracket of the sphere is nontrivial.

The loop homology of the classifying space BG of a connected compact Lie group G in the sense of Chataur and Menichi [9] admits the BV algebra structure; see also [27, Theorem C.1]. Therefore, the same argument as that about manifolds allows us to deduce that the string cohomology of BG is endowed with a graded Lie algebra structure; see Proposition 2.3 and Chen, Eshmatov and Liu [11, Theorem 1.1].

The aim of this article is to investigate general methods for computing the rational string brackets for a manifold and the classifying space of a connected compact Lie group. The key strategy is to use Jones' isomorphisms

$$H^*(LM; \mathbb{Q}) \cong \mathrm{HH}_*(A_{\mathrm{PL}}(M)) \quad \text{and} \quad H_{S^1}^*(LM; \mathbb{Q}) \cong \mathrm{HC}_*^-(A_{\mathrm{PL}}(M)),$$

where $A_{\mathrm{PL}}(M)$ is the polynomial de Rham algebra over \mathbb{Q} of a simply connected space M and the right-hand sides of the isomorphisms denote the Hochschild homology and the negative cyclic homology of the complex, respectively; see Section 3 for more details. Furthermore, the decomposition theorem of the negative cyclic homology and the cyclic homology (additive K-theory in the sense of Feigin and Tsygan [20]) in Vigué-Poirrier [46] and Kuribayashi and Yamaguchi [29] is applied in the computation; see Theorem 2.15. It turns out that for a simply connected closed manifold M , the rational string bracket for M is reduced to the loop product of M followed by the BV operator provided the manifold possesses the exactness of the operator; see Definition 2.9.

Assertion 1.2 *Let M be a simply connected closed manifold. Suppose further that M is **BV-exact**. Then the string bracket in the string homology $H_*^{S^1}(LM; \mathbb{Q})$ is regarded as the **loop bracket** in the loop homology $H_*(LM; \mathbb{Q})$ up to isomorphism, and hence the string bracket is determined by the Gerstenhaber bracket in the Hochschild cohomology of the polynomial de Rham algebra $A_{\text{PL}}(M)$ of M .*

The detail is described in Corollary 2.16. In particular, the nilpotency of the string bracket is equivalent to that of the Gerstenhaber bracket. We stress that the Gerstenhaber algebra in Assertion 1.2 is considered with the Lie model for M without using the loop product; see Félix, Menichi and Thomas [16]. It is worth mentioning that the BV-exactness, which is introduced to consider the reduction of the string brackets, is a new homotopy invariant deeply related to other traditional rational homotopy invariants for spaces. We discuss and summarize this topic in Assertion 1.3 below and several paragraphs before the assertion.

Félix, Thomas and Vigué-Poirrier [19] have given an explicit description of the rational string bracket of M with its Sullivan model. On the other hand, our method for computing the string bracket is formulated with the loop product and the BV operator on the loop homology. Moreover, the BV-exactness is also described in terms of the loop homology. Therefore, it is possible to make a computation of the dual to the string bracket on the equivariant homology $H_*^{S^1}(LM; \mathbb{Q})$ by considering *only* behavior of the BV operator on the loop homology $H_*(LM; \mathbb{Q})$; see Remark 2.14 for more details. This is an advantage of our result.

In the case of the classifying space, the same strategy as above is applicable in the computation of the string bracket. In fact, for the classifying space BG of *every* compact connected Lie group G , the rational string bracket for BG is described as the BV operator followed by the dual loop coproduct; see Theorems 2.7(i) and 2.8(i). As for general properties of the string brackets, the theorems allow us to deduce that the Lie bracket on the string cohomology $H_{S^1}^*(LBG; \mathbb{Q})$ is highly nontrivial even if $\text{rank } G = 1$; see Proposition 5.2. Moreover, Propositions 5.3 and 5.4 assert that the loop homology endowed with the string bracket of a simply connected Lie group G is nilpotent if and only if $\text{rank } G = 1$.

The notion of a *Gorenstein space* due to Félix, Halperin and Thomas [14] enables us to deal with a manifold and the classifying space of a Lie group simultaneously. As a consequence, with the influence of string topology on Gorenstein spaces (see Félix and Thomas [18]), we have Theorems 2.7, 2.8 and 2.15 mentioned above.

We moreover propose a method for computing the string bracket of a non-BV-exact space M . To this end, we introduce a bracket on the cobar-type Eilenberg–Moore spectral sequence (EMSS) converging to $H_{S^1}^*(LM; \mathbb{Q})$ which is compatible with the string bracket of the target; see Theorem 7.7. Moreover, the EMSS carries a decomposition compatible with the Hodge decomposition of the target; see Remark 7.3. While there is no computational example obtained by applying the spectral sequence, in future work, it is expected that the EMSS is applicable in computing the string bracket explicitly; see Section 1.1 problems. As described above, the BV-exactness is a key to computing string brackets on Gorenstein spaces. Moreover, it is worthwhile mentioning that the BV-exactness for a space M is equivalent to the triviality

of the S -action in Connes' exact sequence; see Theorem 2.11. In fact, the new invariant is only described in terms of the Hochschild homology while the S -action is defined on the negative cyclic homology. A deep consideration due to Vigué-Poirrier in [45; 46] shows that the S -action on the negative cyclic homology is trivial if M is formal. Thus we see that the class of BV-exact spaces contains that of formal spaces; see Corollary 2.13.


With historical perspectives, we comment on relationships among notions of p -universality in Mimura, O'Neill and Toda [36], *positive weights* in Body and Douglas [4], the BV-exactness and its variants; see Definition 2.20 for positive weights.

By definition, simply connected spaces X and Y are said to be p -equivalent if there is a map $f : X \rightarrow Y$ which induces $H^*(X; \mathbb{Z}/p) \cong H^*(Y; \mathbb{Z}/p)$, where p is a prime or zero and $\mathbb{Z}/0 = \mathbb{Q}$. In [41], Serre raised the so-called symmetry question whether the existence of a p -equivalence $X \rightarrow Y$ implies the existence of a p -equivalence in the reverse direction $Y \rightarrow X$. However, in general, the p -equivalence does not satisfy the symmetricity.

Mimura, O'Neil and Toda [36] defined the notion of a p -universal space and proved that in the full subcategory of p -universal spaces of the category of simply connected spaces whose homotopy types are those of finite CW complexes, the p -equivalence is indeed an equivalence relation. We observe that the p -universality does not depend on p or 0; see [36, Proposition 2.9]. Afterward, Body and Douglas [4] defined the concept of positive weights for Sullivan minimal models. Scheerer's result [40, Theorem 2], in turn, yields that the two notions of p -universality and positive weights are equivalent.

By using the EMSS mentioned above, we also introduce the notion of r -BV-exactness; see Definition 7.11. The r -BV-exactness for a simply connected space M is equivalent to the collapsing at the E_{r+1} -term of the EMSS for M ; see Corollary 7.5. The decomposition of the EMSS allows us to deduce that the notion of BV-exactness is indeed equivalent to that of 1-BV-exactness; see Theorem 7.10. Thus r -BV-exactness is regarded as a higher version of BV-exactness. We summarize important relationships among invariants mentioned above.

Assertion 1.3 *The following implications concerning rational homotopy invariants hold for a simply connected space X :*



Here the reduced cohomology $\tilde{H}_{\mathfrak{S}^1}^*(LX; \mathbb{Q})$ is the cokernel of the map $H_{\mathfrak{S}^1}^*(*; \mathbb{Q}) \rightarrow H_{\mathfrak{S}^1}^*(LX; \mathbb{Q})$ induced by the trivial map, and the S -action on $\tilde{H}_{\mathfrak{S}^1}^*(LX; \mathbb{Q})$ is defined by the multiplication of the generator of $\tilde{H}^*(BS^1; \mathbb{Q})$ with the map induced by the projection q of the fibration

$$LX \rightarrow ES^1 \times_{S^1} LX \xrightarrow{q} BS^1.$$

Observe that the equivalence $(*)$ holds if X has the homotopy type of a finite CW complex.

As mentioned above, a simply connected space admitting positive weights is BV-exact. Proposition 6.1 gives an example of a nonformal BV-exact manifold. Moreover, we obtain an elliptic and non-BV-exact space in Appendix A.

This manuscript is organized as follows. In Section 2, our results are stated in detail. In Section 3, we recall the Hochschild homology, the cyclic homology and Connes' exact sequences. Moreover, the Gorenstein space in the sense of Félix, Halperin and Thomas [14] is also recalled. Section 4 provides the proofs of our results described in Section 2. Section 5 discusses the nilpotency of the string homology of a Lie group and the classifying space of a Lie group. In Section 6, the BV-exactness for a nonformal manifold of dimension 11 is considered. Thanks to the reduction for computing the bracket described in Section 2, we determine explicitly the dual string bracket for the manifold; see Theorem 6.6. We believe that the result gives the first example which computes the string bracket of a non formal space. Section 7 considers the cobar-type Eilenberg–Moore spectral sequence (EMSS) for computing string brackets of non-BV-exact manifolds.

In Appendix A, we obtain an example of an elliptic and non-BV-exact space. Appendix B describes the Gysin exact sequence associated with the principal bundle $S^1 \rightarrow ES^1 \times LM \xrightarrow{p} ES^1 \times_{S^1} LM$ for a simply connected space M in terms of Sullivan models; see Whitehead [52, (5.12) Theorem] for the exact sequence.

Finally, on page 2651 there is a list of symbols used repeatedly in this article.

1.1 Problems

We propose questions and problems on topics in this article.

- (P1) If a space is BV-exact, then does it admit positive weights?
- (P2) For each $r > 1$, is there an r -BV-exact space which is not $(r-1)$ -BV-exact?
- (P3) Is a space r -BV-exact for some $r < \infty$?
- (P4) By making use of the EMSS in Section 7, compute explicitly the string brackets of a non-BV-exact manifold.

2 String brackets described in terms of the Hochschild homology

While the underlying field in Proposition 2.3 below is of *arbitrary* characteristic, other results described in this section hold for a field of characteristic zero.

Let \mathbb{K} be a field and denote the singular homology and cohomology with coefficients in \mathbb{K} by $H_*(-)$ and $H^*(-)$, respectively. For an orientable closed manifold M of dimension d , the Chas and Sullivan loop product \bullet on the shifted homology $\mathbb{H}_*(LM) := H_{*+d}(LM)$ is unital, associative and graded commutative; see [8]. Consider the principal bundle $S^1 \rightarrow ES^1 \times LM \xrightarrow{p} ES^1 \times_{S^1} LM$. The bundle gives rise to the homology Gysin sequence

$$\cdots \rightarrow \mathbb{H}_{*-d}(LM) \xrightarrow{p_*} H_*^{S^1}(LM) \xrightarrow{c} H_{*-2}^{S^1}(LM) \xrightarrow{M} \mathbb{H}_{*-d-1}(LM) \rightarrow \cdots$$

The *string bracket* $[\cdot, \cdot]$ on $H_*^{S^1}(LM)$ is defined by

$$(2.1) \quad [a, b] := (-1)^{|a|-d} p_*(M(a) \bullet M(b)) \quad \text{for } a, b \in H_*^{S^1}(LM).$$

The bracket is of degree $2-d$ and gives a Lie algebra structure to the equivariant homology of LM .

Let G be a connected compact Lie group of dimension d . We write $\mathbb{H}^*(LBG) := H^{*+d}(LBG)$ and $\mathcal{H}^*(LBG) := H_{S^1}^{*+d+1}(LBG)$. With this notation, the cohomology Gysin sequence associated with the principal bundle $S^1 \rightarrow ES^1 \times LBG \xrightarrow{p} ES^1 \times_{S^1} LBG$ induces an exact sequence of the form

$$\cdots \rightarrow \mathcal{H}^{*-2}(LBG) \xrightarrow{S} \mathcal{H}^*(LBG) \xrightarrow{\pi := p^*} \mathbb{H}^{*+1}(LBG) \xrightarrow{\beta} \mathcal{H}^{*-1}(LBG) \rightarrow \cdots$$

Chataur and Menichi [9] have proved that there exists an associative and graded commutative multiplication \odot on $\mathbb{H}^*(LBG)$ which is induced by the dual loop coproduct with an appropriate sign; see [27, Corollary B.3] and also Section 3. Then the *dual string cobracket* $[\cdot, \cdot]$ on $\mathcal{H}^*(LBG)$ is defined by

$$(2.2) \quad [x, y] := (-1)^{\|x\|} \beta(\pi(x) \odot \pi(y)) \quad \text{for } x, y \in \mathcal{H}^*(LBG).$$

Here the notation $\|x\|$ means the degree of x as an element in the shifted cohomology.

Proposition 2.3 *Let G be a connected compact Lie group of dimension d and \mathbb{K} a field of arbitrary characteristic. Then the dual string cobracket gives $\mathcal{H}^*(LBG)$ a graded Lie algebra structure.*

Remark 2.4 Proposition 2.3 is a particular case of [9, Theorem 65] and [11, Theorem 1.1]. The result [9, Theorem 65] shows the Lie algebra structure on a homological conformal field theory. The result [11, Theorem 1.1] describes a gravity algebra structure on the negative cyclic homology of a mixed complex; see [21] for a gravity algebra. We give an elementary proof of this proposition by taking care of sign convention in Section 4.

We relate the string brackets (ie the string bracket (2.1) and the dual string cobracket (2.2)) above to the Hochschild homology and the cyclic homology. Let Ω be a connected differential graded algebra (DGA) over a field \mathbb{K} of arbitrary characteristic. A DGA Ω is called a *cochain algebra* if the differential

is of degree +1. If the differential of a DGA Ω decreases degree by one, we call the DGA Ω a *chain algebra*. Let Ω be a chain algebra, which is nonpositive; that is, $\Omega = \bigoplus_{i \leq 0} \Omega_i$. We recall Connes' exact sequences [30, Theorem 2.2.1 and Proposition 5.1.5] for the Hochschild homology, cyclic homology and the negative cyclic homology of Ω , which are of the form

$$\begin{aligned}
 (2.5) \quad & \dots \rightarrow \mathrm{HH}_{n+1}(\Omega) \xrightarrow{I} \mathrm{HC}_{n+1}(\Omega) \xrightarrow{S'} \mathrm{HC}_{n-1}(\Omega) \xrightarrow{B_{\mathrm{HH}}} \mathrm{HH}_n(\Omega) \rightarrow \dots, \\
 & \dots \rightarrow \mathrm{HC}_{n+2}^-(\Omega) \xrightarrow{S=\times u} \mathrm{HC}_n^-(\Omega) \xrightarrow{\pi} \mathrm{HH}_n(\Omega) \xrightarrow{\beta} \mathrm{HC}_{n+1}^-(\Omega) \rightarrow \dots, \\
 & \dots \rightarrow \mathrm{HC}_{n+1}^-(\Omega) \xrightarrow{\times u} \mathrm{HC}_{n-1}^{\mathrm{per}}(\Omega) \xrightarrow{\tilde{\pi}} \mathrm{HC}_{n-1}(\Omega) \xrightarrow{B_{\mathrm{HC}}} \mathrm{HC}_n^-(\Omega) \rightarrow \dots.
 \end{aligned}$$

Here S denotes the S -action and the maps B_{HH} , β and B_{HC} are induced by Connes' B -map B ; see Section 3.1 for more details. The reduced versions of the Hochschild homology and the negative cyclic homology of Ω are denoted by $\widetilde{\mathrm{HH}}_*(\Omega)$ and $\widetilde{\mathrm{HC}}_*(\Omega)$, respectively; see Section 3.1.

Remark 2.6 Following Jones [26], we define the Hochschild homology and the cyclic homology for a chain algebra but not a cochain algebra. For a cochain algebra Ω , we define a chain algebra $\Omega_{\#}$ by $(\Omega_{\#})_{-i} = \Omega^i$ for i . Thus, for a nonnegative cochain algebra \mathcal{M} , we have a nonpositive chain algebra $\mathcal{M}_{\#}$. The Hochschild homology and the negative cyclic homology of \mathcal{M} are defined by $\mathrm{HH}_*(\mathcal{M}_{\#})$ and $\mathrm{HC}_*(\mathcal{M}_{\#})$, respectively. By abuse of notation, we may write $\mathrm{HH}_*(\mathcal{M})$ and $\mathrm{HC}_*(\mathcal{M})$ for $\mathrm{HH}(\mathcal{M}_{\#})$ and $\mathrm{HC}(\mathcal{M}_{\#})$, respectively.

The constructions of the string brackets above are generalized with *Gorenstein spaces*. An orientable manifold and the classifying space of a connected Lie group are typical examples of Gorenstein spaces; see Section 3 for the definition and fundamental properties of a Gorenstein space. For a Gorenstein space M of dimension d , we define a comultiplication \bullet^{\vee} and a multiplication \odot on the cohomology $H^*(LM; \mathbb{K})$, which are called the *dual loop product* and the *dual loop coproduct*, respectively; see Section 3. Therefore, by using the formulae (2.1) and (2.2) above, we have the string bracket and the dual string cobracket for a Gorenstein space M with $\bullet := (\bullet^{\vee})^{\vee}$ and \odot , respectively; see Theorems 2.7 and 2.8 below for more details. We do not know the string brackets satisfy the Jacobi identity for general Gorenstein spaces. However, as seen in Theorem 2.8, these constructions indeed give generalizations of brackets (2.1) on manifolds and (2.2) on classifying spaces.

The following theorem asserts that the dual to the string bracket in the sense of Chas and Sullivan for a manifold is the dual loop product followed by the BV operator. Moreover, we see that the string bracket in Proposition 2.3 is described as the BV operator followed by the dual loop coproduct.

In the rest of this section, we further assume that \mathbb{K} is a field of characteristic zero and a DGA Ω is locally finite; that is the homology $H_i(\Omega)$ is finite-dimensional for each $i \leq 0$.

Theorem 2.7 *Let M be a simply connected Gorenstein space and Ω the chain algebra $A_{\text{PL}}(M)_{\#} \otimes_{\mathbb{Q}} \mathbb{K}$. Suppose that the S -action on the reduced negative cyclic homology $\widetilde{\text{HC}}_*^-(\Omega)$ is trivial. Then:*

(i) *There is a commutative diagram*

$$\begin{array}{ccc}
 ((\widetilde{\text{HH}}_*(\Omega)/\text{Im } \Delta) \oplus \mathbb{K}[u])^{\otimes 2} & \xrightarrow[\cong]{\Xi \otimes \Xi} & \text{HC}_*^-(\Omega)^{\otimes 2} \\
 \Delta \otimes \Delta \downarrow & & \pi \otimes \pi \downarrow \\
 \text{HH}_*(\Omega)^{\otimes 2} & & \text{HH}_*(\Omega)^{\otimes 2} \\
 \odot \downarrow & & \odot \downarrow \\
 \text{HH}_*(\Omega) & & \text{HH}_*(\Omega) \\
 \text{Cokernel} \downarrow & & \beta \downarrow \\
 (\widetilde{\text{HH}}_*(\Omega)/\text{Im } \Delta) \oplus \mathbb{K}[u] & \xrightarrow[\cong]{\Xi} & \text{HC}_*^-(\Omega)
 \end{array}$$

Here $\Delta = B_{\text{HH}} \circ I : \text{HH}_*(\Omega) \rightarrow \text{HH}_*(\Omega)$ is the “BV operator”, \odot is the product described in Section 3.3, Cokernel is defined by (projection on the cokernel, 0), and the horizontal isomorphism Ξ is defined by the composite

$$(\widetilde{\text{HH}}_*(\Omega)/\text{Im } \Delta) \oplus \mathbb{K}[u] \xrightarrow[\cong]{I} \widetilde{\text{HC}}_*(\Omega) \oplus \mathbb{K}[u] \xrightarrow[\cong]{B_{\text{HC}}} \widetilde{\text{HC}}_*^-(\Omega) \oplus \mathbb{K}[u] \xrightarrow[\cong]{\text{sp}} \text{HC}_*^-(\Omega),$$

with the map sp in Remark 3.1 below.

(ii) *There is a commutative diagram*

$$\begin{array}{ccc}
 ((\widetilde{\text{HH}}_*(\Omega)/\text{Im } \Delta) \oplus \mathbb{K}[u])^{\otimes 2} & \xrightarrow[\cong]{\Xi \otimes \Xi} & \text{HC}_*^-(\Omega)^{\otimes 2} \\
 \text{Cokernel} \otimes \text{Cokernel} \uparrow & & \beta \otimes \beta \uparrow \\
 \text{HH}_*(\Omega)^{\otimes 2} & & \text{HH}_*(\Omega)^{\otimes 2} \\
 \bullet \vee \uparrow & & \bullet \vee \uparrow \\
 \text{HH}_*(\Omega) & & \text{HH}_*(\Omega) \\
 \Delta \uparrow & & \pi \uparrow \\
 (\widetilde{\text{HH}}_*(\Omega)/\text{Im } \Delta) \oplus \mathbb{K}[u] & \xrightarrow[\cong]{\Xi} & \text{HC}_*^-(\Omega).
 \end{array}$$

Here $\Delta = B_{\text{HH}} \circ I$ is the BV operator of the BV algebra $\text{HH}_*(\Omega)$, and the horizontal isomorphism Ξ is the one defined in (i).

We call the right-hand vertical composites in Theorem 2.7(i) and (ii) the *dual string cobracket* and the *dual string bracket*, respectively.

Note that the condition on the S -action can be replaced with BV-exactness; see Definition 2.9 and Remark 2.14 for details. It is also worth mentioning that the composite $B_{\text{HH}} \circ I$ is nothing but the cohomological Batalin–Vilkovisky (BV) operator Δ on the Hochschild homology of a DGA Ω if Ω is the polynomial de Rham algebra of a manifold or the classifying space of a connected Lie group. By abuse of terminology, we may call $B_{\text{HH}} \circ I$ the BV operator in general.

As mentioned above, under the isomorphisms Θ_1 and Θ_2 due to Jones [26, Theorem A], the loop cohomology $H^*(LM)$ and the string cohomology $H_{S^1}^*(LM)$ are identified with the Hochschild homology and the negative cyclic homology of $A_{PL}(M)$, respectively. Thus, we have:

Theorem 2.8 (i) *The dual string cobracket for BG described in Proposition 2.3 coincides with that in Theorem 2.7(i) up to isomorphisms Θ_1 and Θ_2 .*

(ii) *Let M be a simply connected closed manifold of dimension d . Then the dual $[\cdot, \cdot]^\vee$ to the string bracket in the sense of Chas and Sullivan on M coincides with the dual string bracket in Theorem 2.7(ii) up to isomorphisms Θ_1 and Θ_2 .*

In view of [27, Theorem 4.1], Theorem 2.7(i) and Theorem 2.8(i) allow us to compute the dual string cobracket on $H_{S^1}^*(LBG; \mathbb{K})$ explicitly if \mathbb{K} is a field of characteristic zero; see Section 5. We observe that the classifying space BG is formal and then the S -action is trivial; see Corollary 2.13 below.

Moreover, by dualizing Theorem 2.7(ii) and Theorem 2.8(ii), we have Theorem 2.15, described below, for computing the string bracket of a manifold. It turns out that, in the rational case, the original string bracket can be formulated as the loop product followed by the BV operator on the loop homology. Before describing our main result concerning a manifold, we need a notion of the Batalin–Vilkovisky exactness.

Definition 2.9 A DGA Ω is *Batalin–Vilkovisky exact (BV-exact)* if $\text{Im } \tilde{B} = \text{Ker } \tilde{B}$, where the reduced operator $\tilde{B}: \widetilde{HH}_*(\Omega) \rightarrow \widetilde{HH}_*(\Omega)$, is a restriction of Connes’ B -operator $B := \pi \circ \beta: HH_*(\Omega) \rightarrow HH_*(\Omega)$. We say that a simply connected space M is *BV-exact* if the polynomial de Rham algebra $A_{PL}(M)$ of M is.

Remark 2.10 Let M be a simply connected closed manifold. The result [17, Proposition 2] implies that the dual of the BV operator $\Delta': H_*(LM) \rightarrow H_{*+1}(LM)$ is identified with the operator B in Definition 2.9 under the isomorphism Θ_1 mentioned above. Then, it follows that a manifold M is BV-exact if and only if $\text{Im } \tilde{\Delta}' = \text{Ker } \tilde{\Delta}'$ for the reduced BV operator $\tilde{\Delta}': \tilde{H}_*(LM) \rightarrow \tilde{H}_{*+1}(LM)$.

Theorem 2.11 *A simply connected DGA Ω is BV-exact if and only if the reduced S -action on $\widetilde{HC}_*(\Omega)$ is trivial.*

We refer the reader to Theorem 7.10 for a generalization of the result. An important example with trivial reduced S -action is given by the following proposition due to Vigué-Poirrier.

Proposition 2.12 [46, Proposition 5] *If a simply connected DGA Ω is formal, then the reduced S -action on $\widetilde{HC}_*(\Omega)$ is trivial.*

By combining Theorem 2.11 and Proposition 2.12, we have:

Corollary 2.13 *If a simply connected DGA Ω is formal, then it is BV-exact. As a consequence, a simply connected manifold whose rational cohomology is generated by a single element and the classifying space of a compact connected Lie group are BV-exact.*

We also have a generalization of the corollary; see Theorem 2.21.

Remark 2.14 It follows from Theorem 2.11 that the condition on the S -action in Theorems 2.7 and 2.8 may be replaced with the BV-exactness. This implies that the string brackets are determined exactly with the loop (co)products and the BV operator on the Hochschild homology of a DGA Ω without dealing with the cyclic homology of Ω itself provided Ω is BV-exact. There is an isomorphism

$$\tilde{\Delta} : \widetilde{\text{HH}}_*(\Omega) / \text{Im } \tilde{\Delta} = \widetilde{\text{HH}}_*(\Omega) / \text{Ker } \tilde{\Delta} \xrightarrow{\cong} \text{Im } \tilde{\Delta} = \text{Ker } \tilde{\Delta}.$$

Dualizing Theorems 2.7(ii) and 2.8(ii), we have:

Theorem 2.15 *Let M be a simply connected closed manifold and \mathbb{K} a field of characteristic zero. Assume further that M is BV-exact. Then there exists a commutative diagram*

$$\begin{array}{ccccc} H_*^{S^1}(LM; \mathbb{K})^{\otimes 2} & \xrightarrow[\cong]{\Phi \otimes \Phi} & (\text{Ker } \tilde{\Delta}' \oplus \mathbb{K}[u])^{\otimes 2} & \xrightarrow{\text{inc} \oplus 0} & H_*(LM; \mathbb{K})^{\otimes 2} \\ \downarrow \text{the string bracket} & & & & \downarrow \text{the loop product} \bullet \\ H_*^{S^1}(LM; \mathbb{K}) & \xrightarrow[\cong]{\Phi} & (\text{Ker } \tilde{\Delta}' \oplus \mathbb{K}[u]) & \xleftarrow{\Delta'} & H_*(LM; \mathbb{K}) \end{array}$$

Here $\tilde{\Delta}' : \tilde{H}_*(LM; \mathbb{K}) \rightarrow \tilde{H}_{*+1}(LM; \mathbb{K})$ denotes the reduced BV operator on the homology, and Φ is the dual of the composite of the isomorphisms Θ_2 and Ξ described in Theorem 2.7.

The shifted homology $\mathbb{H}_*(LM) := H_{*+d}(LM)$ for an orientable closed manifold M of dimension d admits a BV algebra structure with the loop product \bullet and the BV operator Δ' ; see [8]. It turns out that the homology is endowed with a Gerstenhaber algebra structure whose Lie bracket (loop bracket) $\{ , \}$ is given by

$$\{a, b\} = (-1)^{|a|}(\Delta'(a \bullet b) - (\Delta'a) \bullet b - (-1)^{|a|}a \bullet (\Delta'b)) \quad \text{for } a, b \in \mathbb{H}_*(LM).$$

If a and b are in the kernel of Δ' , then $\{a, b\} = (-1)^{|a|}\Delta'(a \bullet b)$. Therefore, by virtue of Theorem 2.15, we have:

Corollary 2.16 *Under the same assumption and notations as in Theorem 2.15, the rational string bracket of the loop space LM is regarded as a restriction of the loop bracket up to the isomorphism Φ .*

Remark 2.17 (i) Proposition 2.12 implies that Theorems 2.7, 2.8 and 2.15 are applicable to a formal simply connected closed manifold.

(ii) It follows from [10, Theorem 8.5] that the loop homology of an orientable closed manifold admits a gravity algebra structure extending the Lie algebra structure on the string homology. Theorem 2.15 may enable us to determine a gravity algebra structure on the string homology of a BV-exact manifold M ; see Example 5.6.

Remark 2.18 In general, the cyclic homology (additive K–theory [20]) for a DGA does *not* appear as the singular homology of any topological space because the homology is of \mathbb{Z} –grading. We stress that, however, the cyclic homology is used to investigate the string brackets for a manifold and the classifying space of a Lie group. In fact, the horizontal isomorphism Ξ in Theorem 2.7 factors through the cyclic homology of $A_{\text{PL}}(M)_{\#}$.

Remark 2.19 By using the description of the dual loop product Dlp in [28, Theorem 2.3] and Theorem 2.7, we may relate the dual of the string bracket to the cup product on $H^*(LM; \mathbb{K})$ for a manifold M . In fact, the isomorphism Ξ in Theorem 2.7 is a morphism of algebras if the S –action is trivial; see [29, Theorem 2.5]. We observe that the additive K–theory $K^+(\Omega) := \text{HC}_{*-1}(\Omega)$ for a chain algebra Ω is a graded algebra with the Loday–Quillen $*$ –product in [31]; see [29, Proposition 1.1].

We relate the BV-exactness to a more familiar rational homotopy invariant.

Definition 2.20 A simply connected space X admits *positive weights* if the Sullivan minimal model $(\wedge V, d)$ for X has a direct sum decomposition $V = \bigoplus_{i>0} V_{(i)}$ satisfying $d(V_{(i)}) \subset (\wedge V)_{(i)}$. A nonzero element in $V_{(i)}$ is said to have *weight* i , and the weight on V is extended in a multiplicative way to $\wedge V$. For $x \in (\wedge V)_{(i)}$, its weight is written by $\text{wt}(x) = i$.

Many spaces admit positive weights.

- (1) The Sullivan minimal model $\mathcal{M}(X)$ of a formal space X is given by the bigraded model $(\wedge V, d)$ of its cohomology algebra $H^*(X; \mathbb{Q})$ [24, Section 3], whose lower degree is given by $dV_p \subset (\wedge V)_{p-1}$ for $p > 0$ and $dV_0 = 0$. Then the space X admits positive weights defined by $\text{wt}(v) := |v| + p$ for $v \in V_p$.
- (2) If a space X has a two stage Sullivan minimal model $\mathcal{M}(X) = (\wedge(V_0 \oplus V_1), d)$ with $dV_0 = 0$ and $dV_1 \subset \wedge V_0$, then X admits positive weights defined by $\text{wt}(v) := |v| + i$ for $v \in V_i$. For example, a homogeneous space is such a space even if it is not formal; see also Section 6 for such a manifold.
- (3) It is known that smooth complex algebraic varieties admit positive weights coming from its mixed Hodge structure [38]. In the paper, the Sullivan minimal models are discussed over \mathbb{C} , but admitting positive weights is reduced to that over \mathbb{Q} ; see [5, Theorem 2.7].

Theorem 2.21 A simply connected space X admitting positive weights is BV-exact.

A simply connected space does not necessarily admit positive weights. In fact, there exist a four cell complex [37, Section 4] and elliptic spaces [1, Section 5] not admitting positive weights; see also Appendix A. It is worth mentioning that every finite group is realized as the group of self-homotopy equivalences of a rationalized elliptic space which does not admit positive weights; see [13].

3 Preliminaries

In this section, we recall the Hochschild homology and the cyclic homology together with relationships between them and the loop homology.

3.1 Hochschild and cyclic homology

In this section we recall the definitions of the Hochschild chain complex and the cyclic bar complex in [22] and [23]. Let Ω be a connected commutative DGA over a field \mathbb{K} of arbitrary characteristic endowed with a differential d of degree -1 . We call a DGA Ω *nonpositive* if $\Omega = \bigoplus_{i \leq 0} \Omega_i$. In what follows, it is assumed that a DGA is nonpositively graded algebra with the properties above unless otherwise stated. The degree of a homogeneous element x of a graded algebra is denoted by $|x|$.

First we recall the Hochschild chain complex together with the Connes' B -operator. Write $\bar{\Omega} = \Omega/\mathbb{K}$ and $C(\Omega) = \sum_{k=0}^{\infty} \Omega \otimes \bar{\Omega}^{\otimes k}$. We define \mathbb{K} -linear maps $b, B: C(\Omega) \rightarrow C(\Omega)$ of degrees -1 and 1 by

$$\begin{aligned}
 b(w_0, \dots, w_k) &= \\
 & - \sum_{i=0}^k (-1)^{\epsilon_{i-1}} (w_0, \dots, w_{i-1}, dw_i, w_{i+1}, \dots, w_k) \\
 & \quad - \sum_{i=0}^{k-1} (-1)^{\epsilon_i} (w_0, \dots, w_{i-1}, w_i w_{i+1}, w_{i+2}, \dots, w_k) + (-1)^{(|w_i|-1)\epsilon_{k-1}} (w_k w_0, \dots, w_{k-1}), \\
 B(w_0, \dots, w_k) &= \sum_{i=0}^k (-1)^{(\epsilon_{i-1}+1)(\epsilon_k-\epsilon_{i-1})} (1, w_i, \dots, w_k, w_0, \dots, w_{i-1}).
 \end{aligned}$$

Here $\deg(w_0, \dots, w_k) = |w_0| + \dots + |w_k| + k$ for $(w_0, \dots, w_k) \in C(\Omega)$, $\epsilon_i = |w_0| + \dots + |w_i| - i$ and $|u| = -2$. Note that the formulae $bB + Bb = 0$ and $b^2 = B^2 = 0$ hold. The chain complex $(C(\Omega), b)$ is called the *Hochschild chain complex*. The *Hochschild homology* $\mathrm{HH}_*(\Omega)$ and the *reduced Hochschild homology* $\widetilde{\mathrm{HH}}_*(\Omega)$ are the homologies of the complexes $(C(\Omega), b)$ and $(C(\Omega)/\mathbb{K}, b)$, respectively.

The *cyclic bar complex* is the complex $(C(\Omega)[u^{-1}], b + uB)$, where b and B are regarded as $\mathbb{K}[u^{-1}]$ -linear maps extending b and B on $C(\Omega)$. Its homology is denoted by $\mathrm{HC}_*(\Omega)$ and called the *cyclic homology*. The *negative cyclic homology* $\mathrm{HC}_*^-(\Omega)$, the *reduced negative cyclic homology* $\widetilde{\mathrm{HC}}_*^-(\Omega)$ and the *periodic cyclic homology* $\mathrm{HC}_*^{\mathrm{per}}(\Omega)$ of a DGA Ω are defined as the homologies of the complexes $(C(\Omega)[[u]], b + uB)$, $((C(\Omega)/\mathbb{K})[[u]], b + uB)$ and $(C(\Omega)[[u, u^{-1}]], b + uB)$, respectively. Since a DGA in our case has negative degree, the power series algebra $C(\Omega)[[u]]$ coincides with the polynomial algebra $C(\Omega)[u]$; similarly, $(C(\Omega)/\mathbb{K})[[u]] = (C(\Omega)/\mathbb{K})[u]$ and $C(\Omega)[[u, u^{-1}]] = C(\Omega)[u, u^{-1}]$.

We recall Connes' exact sequences (2.5). The projection of the cyclic complex onto itself gives rise to the map S' . More precisely, we have $S'(\sum_{i \geq 0} x_i u^{-i}) = \sum_{i \geq 0} x_{i+1} u^{-i}$. Observe that the cyclic homology $\mathrm{HC}_*(\Omega)$ and the negative cyclic homology $\mathrm{HC}_*^-(\Omega)$ are $\mathbb{K}[u]$ -modules, where $|u| = -2$. The multiplication $S = \times u: \mathrm{HC}_{n+2}^-(\Omega) \rightarrow \mathrm{HC}_n^-(\Omega)$ is called the S -action on the negative cyclic homology.

For the connecting homomorphism β in Connes' exact sequence (2.5), we see that $\beta([a_0]) = [B(a_0)]$. Moreover, we have $B_{\text{HH}}([\sum_{i \geq 0} x_i u^{-i}]) = [B(x_0)]$ and $B_{\text{HC}}([\sum_{i \geq 0} x_i u^{-i}]) = [B(x_0)]$.

Remark 3.1 Under the same notation as above, the unit and augmentation of Ω yield a split exact sequence of $\mathbb{K}[u]$ -modules of the form $0 \rightarrow C(\mathbb{K})[u] \rightarrow C(\Omega)[u] \rightarrow (C(\Omega)/\mathbb{K})[u] \rightarrow 0$. Then the splitting map $s': \widetilde{\text{HC}}_*^-(\Omega) \rightarrow \text{HC}_*^-(\Omega)$ gives rise to an isomorphism $\text{sp}: \widetilde{\text{HC}}_*^-(\Omega) \oplus \mathbb{K}[u] \xrightarrow{\cong} \text{HC}_*^-(\Omega)$. We observe that $C(\mathbb{K})[u] = \mathbb{K}[u] = \text{HC}_*^-(\mathbb{K})$.

3.2 Sullivan minimal models

Let $\mathcal{M}(Z) = (\wedge V, d)$ be the Sullivan minimal model of a nilpotent CW complex Z of finite type [15]. It is a free \mathbb{Q} -commutative DGA with a \mathbb{Q} -graded vector space $V = \bigoplus_{i \geq 1} V^i$, where $\dim V^i < \infty$, and a decomposable differential in the sense that $d(V^i) \subset (\wedge^+ V \cdot \wedge^+ V)^{i+1}$ and $d \circ d = 0$. Here $\wedge^+ V$ denotes the ideal of $\wedge V$ generated by elements of positive degree. Observe that $\mathcal{M}(Z)$ determines the rational homotopy type of Z ; that is, the spatial realization $\|\mathcal{M}(Z)\|$ is homotopy equivalent to Z_0 , the rationalization of Z . In particular, we see that

$$V^n \cong \text{Hom}(\pi_n(Z), \mathbb{Q}) \quad \text{and} \quad H^*(\wedge V, d) \cong H^*(Z; \mathbb{Q}),$$

the second isomorphism being of graded algebras. A space X is said to be *formal* if there exists a quasi-isomorphism $\rho: \mathcal{M}(X) \rightarrow (H^*(X; \mathbb{Q}), 0)$ of DGA's. We refer the reader to [15] for more details.

In what follows, let \mathbb{K} be a field of characteristic zero unless otherwise specifically mentioned. Let \mathcal{M} be a free DGA $(\wedge V, d)$ with $V = \bigoplus_{i > 1} V^i$ over \mathbb{K} . We denote by (\mathcal{L}, δ, s) the double complex defined in [7]. Namely, $\mathcal{L} = \wedge(V \oplus \bar{V})$, s is the unique derivation of degree -1 defined by $s(v) = \bar{v}$, $s(\bar{v}) = 0$ and δ is the unique derivation of degree $+1$ which satisfies $\delta|_V = d$ and $\delta s + s\delta = 0$. Here \bar{V} is the suspension of V ; that is, $\bar{V}^n = V^{n+1}$. By [7, Theorem 2.4(i)], we see that the map $\Theta: C(\mathcal{M}) \rightarrow \mathcal{L}$ defined by $\Theta(a_0, a_1, \dots, a_p) = (1/p!)a_0 s(a_1) \cdots s(a_p)$ is a chain map between the double complexes $(C(\mathcal{M}), b, B)$ and (\mathcal{L}, δ, s) . Moreover, it follows from [7, Theorem 2.4(ii)–(iii)] that the map Θ induces isomorphisms $H(\Theta): \text{HH}_*(\mathcal{M}) = H_*(C(\mathcal{M}), b) \cong H_*(\mathcal{L}, \delta)$ and $H(\Theta \otimes 1): \text{HC}_*^-(\mathcal{M}) = H_*(C(\mathcal{M})[u], b + uB) \cong H_*(\mathcal{L}[u], \delta + u \cdot s)$.

Remark 3.2 As mentioned in Section 3.1, the connecting homomorphism β in Connes' exact sequence (2.5) is given by $\beta([a_0]) = [B(a_0)]$. It follows that $\beta([a_0]) = [s(a_0)]$ up to the isomorphism $H(\Theta)$; see again [7, Theorem 2.4(i)].

Let X be a simply connected space of finite type and LX the free loop space of X . Then the Sullivan minimal model of LX over \mathbb{K} , $\mathcal{M}(LX)$, is given by (\mathcal{L}, δ) (see [48]), and the Sullivan minimal model of the orbit space $ES^1 \times_{S^1} LX$, $\mathcal{M}(ES^1 \times_{S^1} LX)$, is given by $(\mathcal{E}, D) := (\mathcal{L}[u], \delta + u \cdot s)$ (see [47, Theorem A]). Thus we have isomorphisms $\text{HH}_*(\mathcal{M}(X)) \cong H^{-*}(LX; \mathbb{K})$ and $\text{HC}_*^-(\mathcal{M}(X)) \cong H^{-*}(ES^1 \times_{S^1} LX; \mathbb{K})$ by composing Θ_1 and Θ_2 with $H(\Theta)$ and $H(\Theta \otimes 1)$, respectively.

3.3 Loop product and coproduct on Gorenstein spaces

In order to introduce uniformly the loop product due to Chas and Sullivan and the dual loop coproduct due to Chataur and Menichi, we recall the notion of a Gorenstein DGA introduced by Félix, Halperin and Thomas in [14].

Let A be an augmented DGA over \mathbb{K} . We call A a *Gorenstein algebra* of dimension d if

$$\dim \text{Ext}_A^*(\mathbb{K}, A) = \begin{cases} 0 & \text{if } * \neq d, \\ 1 & \text{if } * = d. \end{cases}$$

Here Ext is defined by using semifree resolutions; see [14, Appendix] for details. A path-connected space M is called a *Gorenstein space* of dimension d if the polynomial de Rham algebra $A_{\text{PL}}(M)$ is a Gorenstein algebra of dimension d .

The result [14, Theorem 3.1] implies that a simply connected Poincaré duality space, for example a simply connected closed orientable manifold of dimension d , is a Gorenstein space of dimension d . It follows from [14, Proposition 3.2] that the classifying space BG of a connected compact Lie group G is also a Gorenstein space of dimension $-\dim G$. The following result due to Félix and Thomas is a key to defining the loop product and the loop coproduct on the loop homology of a Gorenstein space.

Theorem 3.3 [20, Theorem 12] *Let M be a simply connected Gorenstein space of dimension d whose cohomology with coefficients in \mathbb{Q} is of finite type. Then, for any integer k ,*

$$\text{Ext}_{A_{\text{PL}}(M^n)}^k(A_{\text{PL}}(M), A_{\text{PL}}(M^n)) \cong H^{k-(n-1)d}(M; \mathbb{Q}),$$

where $A_{\text{PL}}(M)$ is considered an $A_{\text{PL}}(M^n)$ -module via the diagonal map $\text{Diag}: M \rightarrow M^n$.

For a Gorenstein space M as in Theorem 3.3, let $D(\text{Mod-}A_{\text{PL}}(M^n))$ be the derived category of right $A_{\text{PL}}(M^n)$ -modules. In the category, we define $\text{Diag}^!$ by the map which corresponds to a generator of the one-dimensional vector space $H^0(M; \mathbb{Q})$ under the isomorphism $\text{Ext}_{A_{\text{PL}}(M^n)}^{(n-1)d}(A_{\text{PL}}(M), A_{\text{PL}}(M^n)) \cong H^0(M)$. Moreover, for a homotopy fiber square

$$\begin{array}{ccc} E' & \xrightarrow{q} & E \\ p' \downarrow & & \downarrow p \\ M & \xrightarrow{\text{Diag}} & M^n \end{array}$$

there exists a unique map $q^!$ in $\text{Ext}_{A_{\text{PL}}(E)}^{(n-1)d}(A_{\text{PL}}(E'), A_{\text{PL}}(E))$ which fits into the commutative diagram in $D(\text{Mod-}A_{\text{PL}}(M^n))$

$$\begin{array}{ccc} A_{\text{PL}}^*(E') & \xrightarrow{q^!} & A_{\text{PL}}^{*+(n-1)d}(E) \\ (p')^* \uparrow & & \uparrow p^* \\ A_{\text{PL}}^*(M) & \xrightarrow{\text{Diag}^!} & A_{\text{PL}}^{*+(n-1)d}(M^n) \end{array}$$

The result follows from the same proof as that of [18, Theorems 1 and 2].

We recall the definition of the loop product on a simply connected Gorenstein space M . Consider the diagram

$$(3.4) \quad \begin{array}{ccccc} LM & \xleftarrow{\text{Comp}} & LM \times_M LM & \xrightarrow{q} & LM \times LM \\ & & \downarrow & & \downarrow (\text{ev}_0, \text{ev}_0) \\ & & M & \xrightarrow{\text{Diag}} & M \times M \end{array}$$

where the right-hand square is the pull-back of the evaluation map $(\text{ev}_0, \text{ev}_0)$ defined by $\text{ev}_0(\gamma) = \gamma(0)$ along the diagonal map Diag , and Comp denotes the concatenation of loops. By definition, the composite

$$q^! \circ (\text{Comp})^* : A_{\text{PL}}(LM) \rightarrow A_{\text{PL}}(LM \times_M LM) \rightarrow A_{\text{PL}}(LM \times LM)$$

induces Dlp , the dual to the loop product on $H^*(LM; \mathbb{Q})$; see [18, Introduction].

We define a product \bullet on $\mathbb{H}_*(LM) := H_{*+d}(LM)$, which is called the *loop product*, by

$$a \bullet b = (-1)^{d(|a|+d)} ((\text{Dlp})^\vee)(a \otimes b)$$

for a and $b \in \mathbb{H}_*(LM)$; see [12, Proposition 4] and [44, Definition 3.2].

In order to recall the loop coproduct for a Gorenstein space M , we consider the commutative diagram

$$\begin{array}{ccccc} LM \times LM & \xleftarrow{q} & LM \times_M LM & \xrightarrow{\text{Comp}} & LM \\ & & \downarrow & & \downarrow l \\ & & M & \xrightarrow{\text{Diag}} & M \times M \end{array}$$

where $l : LM \rightarrow M \times M$ is a map defined by $l(\gamma) = (\gamma(0), \gamma(\frac{1}{2}))$. By definition, the composite

$$\text{Comp}^! \circ q^* : A_{\text{PL}}(LM \times LM) \rightarrow A_{\text{PL}}(LM \times_M LM) \rightarrow A_{\text{PL}}(LM)$$

induces the dual to the loop coproduct Dlcp on $H^*(LM)$. We define a product \odot on the shifted cohomology $\mathbb{H}^*(LM) = H^{*-d}(LM)$, called the *dual loop coproduct*, by

$$a \odot b = (-1)^{d(d-|a|)} \text{Dlcp}(a \otimes b) \quad \text{for } a \otimes b \in H^*(LM) \otimes H^*(LM).$$

Remark 3.5 The product \bullet on $\mathbb{H}_*(LM)$ is associative and graded commutative if M is a simply connected Poincaré duality space; see [28, Proposition 2.7]. So is the product \odot on $\mathbb{H}^*(LM)$ if M is the classifying space BG of a connected Lie group G ; see [9] and [27, Theorem B.1]. Moreover, so are both of \bullet and \odot if M is a Gorenstein space with $\dim(\bigoplus_n \pi_n(M) \otimes \mathbb{Q}) < \infty$; see [39, Theorem 1.1] and [50, Theorem 1.5].

Remark 3.6 By the same fashion as above, a Gorenstein space is defined on an arbitrary field \mathbb{K} . Then Theorem 3.3 remains true after replacing $A_{\text{PL}}(X)$ with the singular cochain algebra of X with coefficients in \mathbb{K} . That is the original assertion in [18]. Moreover, the constructions of the loop product and the loop coproduct are applicable to the Gorenstein space M ; that is, those products are defined on the singular cohomology of LM with coefficient in \mathbb{K} ; see [18]. However, we only use such an algebra defined on a field of characteristic zero for our purpose.

We conclude this section with the definition of a BV algebra. In the next section, the notion plays an important role in defining the dual string cobracket of the classifying space of a Lie group.

Definition 3.7 A graded algebra (\mathbb{H}^*, \odot) equipped with an operator Δ on \mathbb{H}^* of degree -1 is a BV algebra if $\Delta \circ \Delta = 0$ and the Batalin–Vilkovisky identity holds; that is, for any elements a, b and c in \mathbb{H}^* ,

$$\begin{aligned} \Delta(a \odot b \odot c) &= \Delta(a \odot b) \odot c + (-1)^{\|a\|} a \odot \Delta(b \odot c) + (-1)^{\|b\|\|a\|+\|b\|} b \odot \Delta(a \odot c) \\ &\quad - \Delta(a) \odot b \odot c - (-1)^{\|a\|} a \odot \Delta(b) \odot c - (-1)^{\|a\|+\|b\|} a \odot b \odot \Delta(c), \end{aligned}$$

where $\|\alpha\|$ stands for the degree of an element α in \mathbb{H}^* .

4 Proofs of assertions

The strategy of the proof of Proposition 2.3 is exactly that of [8, Theorem 6.2]. In order to make the sign computation more clear in our setting, we give the proof.

Proof of Proposition 2.3 It is readily seen that the dual string cobracket satisfies skew-symmetry since the multiplication m is commutative. Indeed, we have

$$[y, x] = (-1)^{\|y\|} \beta(\pi(y) \odot \pi(x)) = (-1)^{\|x\|(\|y\|+1)+1} \beta(\pi(x) \odot \pi(y)) = -(-1)^{\|x\|\|y\|} [x, y].$$

Let $\Delta: \mathbb{H}^*(LBG) \rightarrow \mathbb{H}^{*-1}(LBG)$ be the cohomological BV operator stated in [27, Appendix E]. Observe that Δ coincides with the composite $\pi\beta$. It follows from [27, Corollary C.3] that the triple $(\mathbb{H}^*(LBG), \odot, \Delta)$ is a BV algebra; hence the bracket

$$\{a, b\} := (-1)^{\|a\|} \Delta(a \odot b) - (-1)^{\|a\|} \Delta(a) \odot b - a \odot \Delta(b)$$

satisfies the Poisson identity

$$(4.1) \quad \{a, b \odot c\} = \{a, b\} \odot c + (-1)^{(\|a\|-1)\|b\|} b \odot \{a, c\}.$$

In the case where $a = \pi(x)$, $b = \pi(y)$ and $c = \pi(z)$, applying β to (4.1) we see that $\beta\{\pi(x), \pi(y) \odot \pi(z)\}$ coincides with

$$\beta(\{\pi(x), \pi(y)\} \odot \pi(z) + (-1)^{(\|\pi(x)\|-1)\|\pi(y)\|} \pi(y) \odot \{\pi(x), \pi(z)\}).$$

Since $\Delta\pi = 0$ and $\beta\Delta = 0$, it follows that

$$\begin{aligned} \{\pi(x), \pi(y)\} &= (-1)^{\|x\|-1} \Delta(\pi(x) \odot \pi(y)) = -\pi[x, y], \\ \beta\{\pi(x), \pi(y) \odot \pi(z)\} &= -(-1)^{\|y\|} \beta(\pi(x) \odot \pi[y, z]) = -(-1)^{\|x\|+\|y\|} [x, [y, z]]. \end{aligned}$$

Therefore, by combining the formulae, we see that

$$\begin{aligned} -(-1)^{\|x\|+\|y\|} [x, [y, z]] &= -\beta(\pi[x, y] \odot \pi(z)) - (-1)^{(\|\pi(x)\|-1)\|\pi(y)\|} \beta(\pi(y) \odot \pi[x, z]) \\ &= -(-1)^{\|x\|+\|y\|} [[x, y], z] - (-1)^{(\|\pi(x)\|-1)\|\pi(y)\|+\|y\|} [y, [x, z]] \\ &= (-1)^{\|x\|+\|y\|+(\|x\|+\|y\|)\|z\|} [z, [x, y]] + (-1)^{\|x\|(\|y\|+\|z\|)+\|x\|+\|y\|} [y, [z, x]]. \end{aligned}$$

Multiplying the both sides of the above equality by $(-1)^{\|x\|+\|y\|+1+\|x\|\|z\|}$, we have

$$(-1)^{\|x\|\|z\|}[x, [y, z]] = -(-1)^{\|y\|\|z\|}[z, [x, y]] - (-1)^{\|x\|\|y\|}[y, [z, x]],$$

which is indeed the Jacobi identity. This completes the proof. \square

Proof of Theorem 2.7 We will first prove (i). Recall the homomorphisms $B_{\text{HC}}: \text{HC}_{n-1}(\Omega) \rightarrow \text{HC}_n^-(\Omega)$ and $B_{\text{HH}}: \text{HC}_{n-1}(\Omega) \rightarrow \text{HH}_n(\Omega)$ in Connes' exact sequence in Section 3, which are defined by $B_{\text{HC}}(\sum_{i \geq 0} x_i u^{-i}) = Bx_0$ and $B_{\text{HH}}(\sum_{i \geq 0} x_i u^{-i}) = Bx_0$. The result [29, Theorem 2.5(i)] implies that B is an isomorphism. By assumption, the S -action is trivial. Then, by [29, Theorems 2.5(ii)–(iii)], the map I is an isomorphism. By a direct calculation, we see that $\pi \circ \Xi = B_{\text{HH}} \circ I$ and $\Xi \circ \text{Cokernel} = \beta$. The same consideration as above enables us to obtain the result (ii). \square

Proof of Theorem 2.8 The assertions (i) and (ii) follow from [26, Theorem A]; see also [10, Theorem 8.3]. In fact, the dual of the homology Gysin exact sequence for the fibration $S^1 \rightarrow ES^1 \times LM \rightarrow ES^1 \times_{S^1} LM$ is identified with the Connes exact sequence under isomorphisms Θ_1 and Θ_2 mentioned in the sentence before Theorem 2.8; see [6, Theorem B] and Appendix B for a description of the Gysin sequence in terms of rational models. With those isomorphisms, we compare the dual to string bracket for a manifold and the dual string cobracket for BG with the dual string bracket and the dual string cobracket in Theorem 2.7, respectively.

To this end, we recall that a simply connected closed manifold M of dimension d is a Gorenstein space of dimension d . Moreover, the classifying space BG of a connected compact Lie group G is a Gorenstein space of dimension $d = -\dim G$; see [14]. Thus the result [18, Theorem A] and observations in [18, pages 419–420] yield that the dual loop product \bullet^\vee for the manifold M and the dual loop coproduct \odot for the classifying space BG are nothing but the dual to the loop product and the dual to the loop coproduct, respectively. It turns out that the bracket on $H_*(LM; \mathbb{K})$ for the manifold M and the dual string cobracket on $H^*(LBG; \mathbb{K})$ coincide with the original string brackets (2.1) and (2.2), respectively. Thus, we have the results. \square

Proof of Theorem 2.15 Let Ω be the DGA $\Omega = A_{\text{PL}}(M)_\# \otimes_{\mathbb{Q}} \mathbb{K}$ for M . We observe that the dual of the BV operator $\Delta': H_*(LM; \mathbb{K}) \rightarrow H_*(LM; \mathbb{K})$ on the homology is regarded as the BV operator $\Delta: \text{HH}_*(\Omega) \rightarrow \text{HH}_*(\Omega)$ in Theorem 2.7; see Remark 2.10.

Let $\widetilde{\text{HH}}_*$ denote the reduced Hochschild homology $\widetilde{\text{HH}}_*(\Omega)$. Dualizing the reduced BV operator $\widetilde{\Delta}': \widetilde{H}_*(LM) \rightarrow \widetilde{H}_*(LM)$, we have an exact sequence (*):

$$\widetilde{\text{HH}}_* \xrightarrow{\widetilde{\Delta}^\vee} \widetilde{\text{HH}}_* \xrightarrow{\pi} \widetilde{\text{HH}}_*/\text{Im } \widetilde{\Delta}^\vee \rightarrow 0.$$

Observe that $\widetilde{\Delta}^\vee = B_{\text{HH}} \circ I = \Delta$. By considering the dual exact sequence of (*), we see that π gives rise to the isomorphism $\pi^\vee: \text{Ker } \widetilde{\Delta}' = \text{Ker}(\Delta^\vee) \xrightarrow{\cong} (\widetilde{\text{HH}}_*/\text{Im } \Delta)^\vee$. Theorem 2.8(ii) yields the result. \square

In the rest of the section, we prove Theorems 2.11 and 2.21. First we prove the “if” part of Theorem 2.11.

Proof of the “if” part of Theorem 2.11 Let Ω be a simply connected DGA such that the reduced S -action on $\widetilde{HC}_*^-(\Omega)$ is trivial. Consider the reduced version of Connes’ exact sequence

$$\cdots \rightarrow \widetilde{HC}_{n+2}^-(\Omega) \xrightarrow{S=0} \widetilde{HC}_n^-(\Omega) \xrightarrow{\pi} \widetilde{HH}_n(\Omega) \xrightarrow{\beta} \widetilde{HC}_{n+1}^-(\Omega) \rightarrow \cdots,$$

which splits into a short exact sequence

$$0 \rightarrow \widetilde{HC}_n^-(\Omega) \xrightarrow{\pi} \widetilde{HH}_n(\Omega) \xrightarrow{\beta} \widetilde{HC}_{n+1}^-(\Omega) \rightarrow 0.$$

By definition, there is a decomposition $\widetilde{B} = \pi \circ \beta: \widetilde{HH}_*(\Omega) \rightarrow \widetilde{HH}_*(\Omega)$ and hence the above short exact sequence implies $\text{Ker } \widetilde{B} = \text{Ker } \beta = \text{Im } \pi = \text{Im } \widetilde{B}$. \square

In order to prove the “only if” part of Theorem 2.11, we recall the notion of the proper exactness of a sequence of complexes defined in [42].

Definition 4.2 Let $M_1 \rightarrow M_2 \rightarrow M_3$ be a sequence of complexes and chain maps (of arbitrary degrees).

- (i) The sequence is H -exact at M_2 if the sequence of cohomology $H(M_1) \rightarrow H(M_2) \rightarrow H(M_3)$ is exact.
- (ii) The sequence is Z -exact at M_2 if the sequence of modules of cycles $Z(M_1) \rightarrow Z(M_2) \rightarrow Z(M_3)$ is exact.
- (iii) The sequence is *proper exact* (at M_2) if the sequence is exact (as a sequence of underlying graded modules), H -exact and Z -exact [42].
- (iv) The sequence is *weakly proper exact* at M_2 if the sequence is exact and H -exact.

The following lemma is useful to prove the proper exactness from the weak proper exactness of a given sequence.

Lemma 4.3 Let $M_0 \xrightarrow{f_0} M_1 \xrightarrow{f_1} M_2 \xrightarrow{f_2} M_3 \xrightarrow{f_3} M_4$ be a sequence of complexes which is proper exact at M_2 and weakly proper exact at M_1 and M_3 . Then it is proper exact also at M_3 .

Proof For simplicity, we assume that the degrees of the chain maps are zero. We show that $\text{Ker } Z(f_3) \subset \text{Im } Z(f_2)$. For any x_3 in $\text{Ker } Z(f_3)$, there exists an element $y_2 \in M_2$ such that $f_2(y_2) = x_3$ by the exactness at M_3 . By the proper exactness at M_2 , we see that $dy_2 = f_1(y_1)$ for some $y_1 \in Z(M_1)$. Since $H(f_1)[y_1] = [dy_2] = 0$, it follows from the H -exactness at M_1 that $y_1 - f_0 y_0 = dz$ for some $[y_0] \in H(M_0)$ and $z \in M_1$. It is readily seen that $f_2(y_2 - f_1 z) = f_2 y_2 = x_3$ and $d(y_2 - f_1 z) = 0$. We have the result. \square

It is proved that the weak proper exactness for a long sequence yields the proper exactness.

Proposition 4.4 A weakly proper exact sequence $0 \rightarrow M_0 \rightarrow M_1 \rightarrow M_2 \rightarrow \dots$ starting from 0 is always proper exact.

Proof Since the sequence $0 \rightarrow 0 \rightarrow 0 \rightarrow M_0 \rightarrow M_1$ is weakly proper exact at M_0 and proper exact at 0, it follows from Lemma 4.3 that the sequence is proper exact at M_0 . Similarly, the sequence $0 \rightarrow 0 \rightarrow M_0 \rightarrow M_1 \rightarrow M_2$ gives proper exactness at M_1 . By repeating this argument, we can prove the proper exactness at M_n for all n . \square

Remark 4.5 By the same argument as in the proof above, we can also prove the dual of Proposition 4.4, which asserts that a weakly proper exact sequence ending with 0 is always proper exact.

Next we give a key lemma for proving Theorem 2.11.

Lemma 4.6 Let $M_0 \rightarrow M_1 \xrightarrow{f_1} M_2 \xrightarrow{f_2} M_3$ be a proper exact sequence. Then one has

$$\text{Im } d \cap \text{Ker } f_2 = d(\text{Ker } f_2).$$

Proof The Z -exactness at M_2 and the H -exactness at M_2 give the result. The details are left to the reader. \square

Note that the consequence in Lemma 4.6 is equivalent to the exactness of the sequence of modules of coboundaries.

Now we begin the proof of the “only if” part of Theorem 2.11. Let $(\wedge V, d)$ be a Sullivan model of the DGA Ω with $V = V^{\geq 2}$. Define $(\tilde{\mathcal{L}}, \delta) = (\wedge^+(V \oplus \bar{V}), \delta)$ and $(\tilde{\mathcal{E}}, D) = (\wedge u \otimes \tilde{\mathcal{L}}, D)$; see Section 3. Then $(\tilde{\mathcal{L}}, \delta)$ and $(\tilde{\mathcal{E}}, D)$ are chain models for the reduced Hochschild homology and the reduced negative cyclic homology of Ω , respectively. Let $\tilde{s}: \tilde{\mathcal{L}} \rightarrow \tilde{\mathcal{L}}$ be the derivation defined by $\tilde{s}(v) = \bar{v}$ and $\tilde{s}(\bar{v}) = 0$ for $v \in V$. Now we have a direct sum decomposition $(\tilde{\mathcal{L}}, \delta) = \bigoplus_n (\tilde{\mathcal{L}}^{(n)}, \delta)$ of complexes, where $\tilde{\mathcal{L}}^{(n)} = \tilde{\mathcal{L}} \cap (\wedge V \otimes \wedge^n \bar{V})$. Then \tilde{s} decomposes into a sequence $0 \rightarrow \tilde{\mathcal{L}}^{(0)} \rightarrow \tilde{\mathcal{L}}^{(1)} \rightarrow \tilde{\mathcal{L}}^{(2)} \rightarrow \dots$ of complexes.

Lemma 4.7 The sequence $0 \rightarrow \tilde{\mathcal{L}}^{(0)} \rightarrow \tilde{\mathcal{L}}^{(1)} \rightarrow \tilde{\mathcal{L}}^{(2)} \rightarrow \dots$ is exact; that is, $\text{Ker } \tilde{s} = \text{Im } \tilde{s}$ in $\tilde{\mathcal{L}}$.

Proof Take a basis $\{v_\lambda\}_\lambda$ of V . Then we have $(\mathcal{L}, s) \cong \bigotimes_\lambda (\wedge(v_\lambda, \bar{v}_\lambda), s)$ and hence $H(\mathcal{L}, s) \cong \mathbb{Q}$, which is equivalent to $H(\tilde{\mathcal{L}}, \tilde{s}) \cong 0$. \square

Remark 4.8 The operator $\tilde{B}: \widetilde{\text{HH}}_*(\Omega) \rightarrow \widetilde{\text{HH}}_*(\Omega)$ is nothing but the homomorphism $H(\tilde{s})$ up to the isomorphism $H(\Theta)$. This follows from the definition of the map B in Section 3 and Remark 3.2.

Now we recall a result of Vigué-Poirrier which gives a description of the cyclic homology in terms of $\tilde{\mathcal{L}}$. Here we give a proof for the convenience of the reader.

Lemma 4.9 [45, Lemma 2] *The canonical inclusion $\Phi: (\text{Ker } \tilde{s}, d) \rightarrow (\tilde{\mathcal{E}}, D)$ is a quasi-isomorphism.*

Proof Define bounded double complexes $\{K^{p,q}\}$ and $\{\tilde{\mathcal{E}}^{p,q}\}$ by $K^{p,0} = (\text{Ker } \tilde{s})^p$ and $K^{p,q} = 0$ for $q \neq 0$, and $\tilde{\mathcal{E}}^{p,q} = \wedge^q u \otimes \tilde{\mathcal{L}}^{p-q}$. Then their total chain complexes are $(\text{Ker } \tilde{s}, \delta)$ and $(\tilde{\mathcal{E}}, D)$, respectively, and the inclusion Φ gives rise to a morphism of double complexes. Now consider the filtration with respect to p . By Lemma 4.7, we have $E_1^{p,0} K = E_1^{p,0} \tilde{\mathcal{E}} = (\text{Ker } \tilde{s})^p$ and $E_1^{p,q} K = E_1^{p,q} \tilde{\mathcal{E}} = 0$ for $q \neq 0$. Hence $E_1 \Phi$ is an isomorphism and so is $H\Phi$ by the convergence of the spectral sequences. \square

Now we describe the S -action $S = u \times (-): H(\tilde{\mathcal{E}}) \rightarrow H(\tilde{\mathcal{E}})$ in terms of $\text{Ker } \tilde{s}$. By Lemma 4.7, we have an exact sequence $0 \rightarrow \text{Ker } \tilde{s} \rightarrow \tilde{\mathcal{L}} \xrightarrow{\tilde{s}} \text{Ker } \tilde{s} \rightarrow 0$, and its connecting homomorphism $c: H(\text{Ker } \tilde{s}) \rightarrow H(\text{Ker } \tilde{s})$ is given by $c([\tilde{s}\alpha]) = [\delta\alpha]$. Note that any element in $H(\text{Ker } \tilde{s})$ can be written as $[\tilde{s}\alpha]$ for some $\alpha \in \tilde{\mathcal{L}}$ with $\delta\tilde{s}\alpha = 0$, since $\text{Ker } \tilde{s} = \text{Im } \tilde{s}$ by Lemma 4.7. By a straightforward computation, we have:

Lemma 4.10 *The map c coincides with S through $H\Phi$ up to sign, ie $S \circ H\Phi = -H\Phi \circ c$.*

We are ready to prove the “only if” part of Theorem 2.11.

Proof of the “only if” part of Theorem 2.11 By Lemmas 4.9 and 4.10, in order to prove the assertion it suffices to show that the connecting homomorphism c is trivial. To this end, we show that $[\delta\alpha] = 0$ in $H(\text{Ker } \tilde{s})$ for any $\alpha \in \tilde{\mathcal{L}}$ with $\delta\tilde{s}\alpha = 0$; see the argument before Lemma 4.10. Remark 4.8 yields that the BV-exactness of the DGA Ω is equivalent to the condition that the sequence $(*)$,

$$0 \rightarrow \tilde{\mathcal{L}}^{(0)} \rightarrow \tilde{\mathcal{L}}^{(1)} \rightarrow \tilde{\mathcal{L}}^{(2)} \rightarrow \dots,$$

is weakly proper exact. Thus, by Proposition 4.4, we see that the sequence $(*)$ is proper exact. Moreover, Lemma 4.6 implies that $\text{Ker } \tilde{s} \cap \text{Im } \delta = \delta(\text{Ker } \tilde{s})$. Therefore, it follows that $\delta\alpha \in \text{Ker } \tilde{s} \cap \text{Im } \delta = \delta(\text{Ker } \tilde{s})$ for any $\alpha \in \tilde{\mathcal{L}}$ with $\delta\tilde{s}\alpha = 0$. We have the result. \square

We conclude this section proving Theorem 2.21. The proof is given by slightly modifying the proof of [46, Proposition 5].

Proof of Theorem 2.21 Recall that $(\tilde{\mathcal{L}}, \delta) = (\wedge^+(V \oplus \bar{V}), \delta)$ is a model of the Hochschild complex. For a derivation $\theta: \wedge V \rightarrow \wedge V$ of degree 0 with $\theta d = d\theta$ and $\theta(V) \subset \wedge^+ V$, define derivations $L_\theta, e_\theta: \tilde{\mathcal{L}} \rightarrow \tilde{\mathcal{L}}$ by $L_\theta(v) = \theta v$, $L_\theta(\bar{v}) = \tilde{s}\theta v$, $e_\theta(v) = 0$ and $e_\theta(\bar{v}) = \theta v$. Then, as derivations on $\tilde{\mathcal{L}}$, we have $[L_\theta, \tilde{s}] = [L_\theta, \delta] = [e_\theta, \delta] = 0$ and $[e_\theta, \tilde{s}] = L_\theta$. Hence L_θ induces $H(L_\theta): H(\text{Ker } \tilde{s}) \rightarrow H(\text{Ker } \tilde{s})$ and it follows that $H(L_\theta) \circ c = 0: H(\text{Ker } \tilde{s}) \rightarrow H(\text{Ker } \tilde{s})$ by a straightforward computation from the above equations.

Now we let θ be the derivation defined by $\theta(x) = \text{wt}(x)x$ for weight-homogeneous elements $x \in \wedge V$. Then for any weight-homogeneous element $\alpha \in H(\text{Ker } \tilde{s})$, we have $0 = H(L_\theta) \circ c(\alpha) = \text{wt}(\alpha)c(\alpha)$, where the weight on $\tilde{\mathcal{L}}$ is defined as an extension of that on $\wedge V$ with $\text{wt}(\bar{v}) = \text{wt}(v)$ for $v \in V$. By the positivity of the weight, we have $c(\alpha) = 0$ and hence $c = 0$. Therefore, Lemmas 4.9 and 4.10 imply the triviality of the reduced S -action, which is equivalent to the BV-exactness by Theorem 2.11. \square

5 The string brackets for formal spaces

In this section, we consider string brackets for formal spaces as an application of Theorem 2.7.

5.1 Dual string cobrackets for classifying spaces

We begin by considering the string bracket for the classifying space of a connected Lie group of rank one.

Example 5.1 The result [27, Theorem 4.1] enables us to compute the dual loop coproduct on the loop cohomology $\mathbb{H}^*(LBG; \mathbb{Q})$ for every compact connected Lie group G . Thus, in particular, by Theorem 2.7, we determine explicitly the Lie algebra structure of $\mathcal{H}^*(LBSU(2)) := H_{S^1}^{*+3+1}(LBSU(2); \mathbb{Q})$ endowed with the dual string cobracket. In fact, we see that

$$\begin{aligned} \mathcal{H}^* &:= \mathcal{H}^*(LBSU(2)) \cong (\widehat{HH}_*(\Omega)/\text{Im } \Delta)_{-* -3 -2} \oplus (\mathbb{Q}[u])_{-* -3 -1} \\ &\cong \mathbb{Q}\{x, x^2, \dots, x^n, \dots\} \oplus \mathbb{Q}\{1, u, u^2, \dots, u^k, \dots\} \end{aligned}$$

as vector spaces, where Ω denotes the Sullivan minimal model for $SU(2)$. Observe that $|x^n| = 4n - 5$ and $|1| = -4$ for x^n and $1 \in \mathcal{H}^*(LBSU(2))$. The formula in [27, Theorem 4.1] for the loop product \odot yields that $\Delta(x^n) \odot 1 = nx^{n-1}$, $\Delta(x^n) \odot \Delta(x^m) = \pm nm \Delta(x)x^{n+m-2}$ and $\Delta(1) = 0$ in $\mathbb{H}^*(LBSU(2))$. Therefore, we see that $[1, 1] = 0$, $[x^n, x^m] = 0$ for $m, n \geq 1$, $[u^l, \alpha] = 0$ for every $\alpha \in \mathcal{H}^*$, $l \geq 1$ and $[x^n, 1] = -nx^{n-1}$ for $n \geq 1$.

Next we consider the dual string cobracket for the classifying space of G with arbitrary rank.

Proposition 5.2 For each n , the n -fold dual string cobracket $[\mathcal{H}, [\mathcal{H}, \dots, [\mathcal{H}, \mathcal{H}] \dots]]$ is nontrivial on $\mathcal{H}^* := H_{S^1}^{*+\dim G+1}(LBG; \mathbb{K})$.

Proof For the case $\text{rank } G = 1$, Example 5.1 implies the result. Assume that $N := \text{rank } G \geq 2$. Recall the result [27, Theorem 4.3], which asserts that the loop cohomology $\mathbb{H}^*(LBG) := H^{*+\dim G}(LBG)$ is isomorphic to the tensor product of algebra

$$H^*(BG) \otimes H_{-*}(G) = \mathbb{K}[y_1, \dots, y_N] \otimes \wedge(x_1^\vee, \dots, x_N^\vee)$$

equipped with the BV operator Δ given by $\Delta(x_i^\vee x_j^\vee) = \Delta(y_i y_j) = \Delta(x_i^\vee) = \Delta(y_i) = 0$ and

$$\Delta(y_i x_j^\vee) = \begin{cases} 0 & \text{if } i \neq j, \\ 1 & \text{if } i = j. \end{cases}$$

Thus, an induction argument with the BV identity enables us to deduce that

$$\Delta(y_1^{k_1} \cdots y_N^{k_N} x_{i_1}^\vee \cdots x_{i_s}^\vee) = \sum_{1 \leq j \leq s} (-1)^{d_j} k_{i_j} y_1^{k_1} \cdots y_{i_j}^{k_{i_j}-1} \cdots y_N^{k_N} x_{i_1}^\vee \cdots \widehat{x_{i_j}^\vee} \cdots x_{i_s}^\vee,$$

where $\widehat{}$ denotes omission and $d_j = |x_{i_1}^\vee| + \cdots + |x_{i_{j-1}}^\vee|$. Therefore, it follows that

$$\Delta(y_2 x_2^\vee x_1^\vee) \odot \Delta(y_1^l x_1^\vee x_2^\vee) = x_1^\vee \odot l y_1^{l-1} x_2^\vee = l y_1^{l-1} x_1^\vee x_2^\vee.$$

Moreover, we see that $\Delta(y_1^{l-1}x_1^\vee x_2^\vee) \neq 0$ for $l \geq 2$. Then the element $y_1^{l-1}x_1^\vee x_2^\vee$ is not in $\text{Im } \Delta$. Observe that $\Delta^2 = 0$. We consider an n -fold bracket of the form

$$\alpha := [y_2 x_2^\vee x_1^\vee, [y_2 x_2^\vee x_1^\vee, \dots, [y_2 x_2^\vee x_1^\vee, y_1^l x_1^\vee x_2^\vee] \dots]] \quad \text{for } l > n.$$

It turns out that

$$\alpha = l(l-1) \cdots (l-(n-1)) y_1^{l-n} x_1^\vee x_2^\vee \neq 0$$

in the codomain $(\widetilde{\text{HH}}_*(\Omega)/\text{Im } \widetilde{\Delta}) \oplus \mathbb{K}[u]$ of the dual string cobracket. Theorem 2.7(i) allows us to obtain the result. □

5.2 String brackets for manifolds

As an application of Theorem 2.8 (or Theorem 2.15), we give another proof of the first half of the result [2, Theorem 3.4] due to Basu and [19, Example 5.2] due to Félix, Thomas and Vigué-Poirrier.

Proposition 5.3 *For a simply connected closed manifold M such that $H^*(M; \mathbb{Q})$ is generated by a single element, the string bracket is trivial.*

Proof The result [17, Theorem 1] implies that the loop homology of M is isomorphic to the Hochschild cohomology of $A_{\text{PL}}(M)$ endowed with the BV algebra structure due to Menichi [33]. We observe that M is formal. Therefore, Theorem 2.15 and explicit computations in [34, Theorem 16] and [53, Main Theorem] yield the result. In fact, for elements α_1 and α_2 in $\text{Im } \widetilde{\Delta} = \text{Ker } \widetilde{\Delta}$, we have $\Delta(\alpha_1 \bullet \alpha_2) = 0$; see Theorem 2.11 and Remark 2.10. In particular, we observe the case where $H^*(M) \cong H^*(S^n)$ with n odd. Then the generator a_{-n} of the loop homology $\mathbb{H}_*(LM) := H_{*+n}(LM)$ with odd degree is in $H_0(LM)$. Then the generator a_{-n} is not in $\text{Ker } \widetilde{\Delta}$; see Theorem 2.15. □

The result [35, Theorem 39] due to Menichi gives an explicit form of the BV operator on the rational loop homology of a connected compact Lie group. We can also apply the result in our computation. In particular, the behavior of the string bracket as seen in Proposition 5.3 changes drastically in case of a Lie group with rank greater than one.

Proposition 5.4 (cf [19, Example 5.2]) *Let G be a simply connected Lie group with rank greater than one. The Lie algebra $\mathcal{H}_* = H_{*-\dim G+2}^{\mathcal{S}^1}(LG; \mathbb{Q})$ endowed with the string bracket is nonnilpotent. More precisely, for any n , the n -fold bracket $[\mathcal{H}, [\mathcal{H}, \dots, [\mathcal{H}, \mathcal{H}] \dots]]$ is nontrivial.*

Proof We first observe that a simply connected Lie group is formal. Indecomposable elements x_1, \dots, x_N in $\mathbb{H}_*(G)$ are in the reduced homology $\widetilde{H}_*(LG)$ because $N := \text{rank } G > 1$. Thus, it follows from [35, Theorem 39] and [25, Theorem 1] that x_i and $(s^{-1}x_j)^k$ are in $\text{Ker } \widetilde{\Delta}$. Moreover, there exists a nontrivial n -fold string bracket. For example, for $k > n$, we see that on \mathcal{H}_* ,

$$[x_j, [x_j, \dots, [x_j, (s^{-1}x_j)^k] \dots]] = \pm k(k-1) \cdots (k-(n-1))(s^{-1}x_j)^{k-n} \neq 0.$$

This follows from the explicit formula of the BV operator in [35, Theorem 39] and Theorem 2.15. Observe that x_j is in $\text{Ker } \widetilde{\Delta}$ if $\text{rank } G > 1$. We have the result. □

5.3 Gravity algebras

The *gravity algebra* with higher Lie brackets was introduced by Getzler [21]. We consider a gravity algebra structure which appears on the string homology of a manifold and the classifying space of a Lie group; see, for example, [10, Definition 8.1] for the definition of the gravity algebra.

Example 5.5 The result [11, Theorem 1.1] due to Chen, Eshmatov and Liu shows that the negative cyclic homology of a DGA Ω admits a gravity algebra structure if the Hochschild homology of Ω has a BV algebra structure. The higher Lie bracket $[\dots,]: (\text{HC}_*^-(\Omega))^{\otimes n} \rightarrow \text{HC}_*^-(\Omega)$ is defined for $n \geq 2$ by

$$[x_1, \dots, x_n] = (-1)^{(n-1)|x_1| + (n-2)|x_2| + \dots + |x_{n-1}|} \beta(\pi(x_1) \odot \pi(x_2) \odot \dots \odot \pi(x_n)),$$

where \odot denotes the dual loop coproduct on the Hochschild homology.

Let G be a connected Lie group. We see that all higher Lie brackets are nontrivial for the classifying space BG . For the case where $\text{rank } G \geq 2$, it follows from Theorem 2.7 that

$$[y_i x_i^\vee, \dots, y_i x_i^\vee, y_2 x_2^\vee x_1^\vee, y_1^l x_1^\vee x_2^\vee] = \pm 1 \odot \dots \odot 1 \odot \Delta(y_2 x_2^\vee x_1^\vee) \odot \Delta(y_1^l x_1^\vee x_2^\vee) = l y_1^{l-1} x_1^\vee x_2^\vee \neq 0$$

with the same notation as in the proof of Proposition 5.2. Suppose that $\text{rank } G = 1$. Then, with the same notation as in Example 5.1, we see that $[x^2, \dots, x^2] = \pm \text{Coker}(\Delta(x^2) \odot \dots \odot \Delta(x^2) \odot \Delta(x^2) \odot 1) = \pm \text{Coker}(\Delta(x^2) \odot \dots \odot \Delta(x^2) \odot 2x) = \pm 2^{n-1} x^{n-1} \neq 0$ for the higher Lie bracket of rank n .

Example 5.6 In [10], Chen has proved that the string homology of an orientable closed manifolds admits a gravity algebra structure extending the Lie algebra structure; see [10, Theorem 8.5] for more details. Let G be a simply connected Lie group. We see that all higher Lie brackets in the string homology of G are nontrivial if and only if $\text{rank } G > 1$. In fact, in case of $\text{rank } G > 1$, by applying Theorem 2.15 to the higher Lie bracket of G , we have $[x_j, s^{-1}x_j, \dots, s^{-1}x_j] = \pm k(s^{-1}x_j)^{k-1}$ in $H_*^{S^1}(LG; \mathbb{Q})$ with the same notation as in Proposition 5.4. If $\text{rank } G = 1$, the only generator x_1 of odd degree is not in $\text{Ker } \tilde{\Delta}$ and then all higher Lie brackets are trivial; see the computation in the proof of Proposition 5.4.

6 Computation of the string bracket for a nonformal space

In this section, we consider the string bracket of a nonformal and BV-exact manifold. We begin recalling a nonformal manifold in [19, 6.4 Example].

Let $UTS^6 \rightarrow S^6$ be the unit tangent bundle over S^6 . Then, we have a simply connected 11-dimensional manifold M which fits in the pullback diagram

$$\begin{array}{ccc} M & \longrightarrow & UTS^6 \\ \downarrow & & \downarrow p \\ S^3 \times S^3 & \xrightarrow{f} & S^6 \end{array}$$

where $f : S^3 \times S^3 \rightarrow S^6$ is a smooth map homotopic to the map defined by collapsing the 3–skeleton into a point. Since the Euler class of the unit tangent bundle mentioned above is nontrivial, it follows that the minimal model of M has the form $\mathcal{M} = (\wedge(x, y, z), d)$, where $d(x) = 0 = d(y)$, $d(z) = xy$, $|x| = |y| = 3$ and $|z| = 5$. It is readily seen that M is nonformal since the Massey product $\langle x, x, y \rangle$ does not vanish; see [24, page 277]. Moreover, we have:

Proposition 6.1 *The 11–dimensional manifold M is BV-exact.*

Proposition 6.1 is proved by computing the Hochschild homology explicitly. To this end, we recall the minimal model \mathcal{M} for M mentioned above. The Hochschild homology of \mathcal{M} is the homology of the Sullivan algebra $\mathcal{L} = (\wedge(x, y, z, \bar{x}, \bar{y}, \bar{z}), d)$, where $d(\bar{x}) = 0 = d(\bar{y})$, $d(\bar{z}) = -\bar{x}y + x\bar{y}$; see Section 3. To compute $H(\mathcal{L})$, we define its subcomplex \mathcal{L}' by $\wedge(x, y, \bar{x}, \bar{y}, \bar{z})$. By a simple calculation, we have the following lemma.

Lemma 6.2 *The set $\{\bar{x}^p \bar{y}^q, x\bar{x}^p \bar{y}^q, y\bar{y}^q, xy\bar{z}^r \mid p, q, r \geq 0\}$ forms a basis of $H(\mathcal{L}')$.*

Next we compute $H(\mathcal{L})$ by comparing with $H(\mathcal{L}')$ and \mathcal{L}/\mathcal{L}' .

Proposition 6.3 *The following set forms a basis of the Hochschild homology $H(\mathcal{L})$:*

$$\{\bar{x}^p \bar{y}^q, x\bar{x}^p \bar{y}^q, y\bar{y}^q, xy\bar{z}^{r+1}, xz\bar{x}^p \bar{y}^q, yz\bar{y}^q, xyz\bar{z}^r, z\bar{x}^{p+1} \bar{y}^q - x\bar{x}^p \bar{y}^q \bar{z}, z\bar{y}^{q+1} - y\bar{y}^q \bar{z}\},$$

where p, q and r run over all nonnegative integers.

Proof Since there is an isomorphism of complexes $(\mathcal{L}/\mathcal{L}', d) \cong (\mathbb{Q}\{z\}, 0) \otimes (\mathcal{L}', d)$, Lemma 6.2 implies that the set $\{z\bar{x}^p \bar{y}^q, xz\bar{x}^p \bar{y}^q, yz\bar{y}^q, xyz\bar{z}^r \mid p, q, r \geq 0\}$ forms a basis of $H(\mathcal{L}/\mathcal{L}')$. Consider the long exact sequence associated with the short exact sequence $0 \rightarrow \mathcal{L}' \rightarrow \mathcal{L} \rightarrow \mathcal{L}/\mathcal{L}' \rightarrow 0$. The connecting homomorphism $H(\mathcal{L}/\mathcal{L}') \rightarrow H(\mathcal{L})$ sends z to xy and the other basis elements to zero. Hence each basis element of $H(\mathcal{L}')$ or $H(\mathcal{L}/\mathcal{L}')$ corresponds to a basis element of $H(\mathcal{L})$, except for z and xy . By lifting basis elements of $H(\mathcal{L}/\mathcal{L}')$ to cocycles in \mathcal{L} , we get the above basis. \square

Proof of Proposition 6.1 Let $\tilde{\mathcal{L}}$ be the reduced complex $\wedge^+(x, y, z, \bar{x}, \bar{y}, \bar{z})$. Recall that the reduced operation \tilde{B} in Definition 2.9 is modeled by the map $H\tilde{s}: H(\tilde{\mathcal{L}}) \rightarrow H(\tilde{\mathcal{L}})$ induced by the derivation $\tilde{s}: \tilde{\mathcal{L}} \rightarrow \tilde{\mathcal{L}}, v \mapsto \bar{v}$ for $v = x, y, z$; see Remark 3.2.

By using the basis given in Proposition 6.3, we see that

$$\begin{aligned} H\tilde{s}(x\bar{x}^p \bar{y}^q) &= \bar{x}^{p+1} \bar{y}^q, & H\tilde{s}(y\bar{y}^q) &= \bar{y}^{q+1}, & H\tilde{s}(xz\bar{x}^p \bar{y}^q) &= z\bar{x}^{p+1} \bar{y}^q - x\bar{x}^p \bar{y}^q \bar{z}, \\ H\tilde{s}(yz\bar{y}^q) &= z\bar{y}^{q+1} - y\bar{y}^q \bar{z}, & H\tilde{s}(xyz\bar{z}^r) &= \frac{r+2}{r+1} xy\bar{z}^{r+1}. \end{aligned}$$

This proves $\text{Ker } H\tilde{s} = \text{Im } H\tilde{s}$. \square

Remark 6.4 A program [51] on a personal computer for computing the homology of a DGA helps us in proving Proposition 6.3. In fact, the computer calculation shows the basis in the proposition, while our proof is by hand.

Remark 6.5 In the minimal model \mathcal{M} , we define weights of x , y and z by 1, 1 and 2, respectively. Then, it is readily seen that the model \mathcal{M} for the manifold M admits positive weights. Therefore, Theorem 2.21 enables us to conclude that M is BV-exact. However, the explicit generators of the Hochschild homology of \mathcal{M} represented in Proposition 6.3 are used in the computation below of the string bracket of M . We adhere to the proof of Proposition 6.1.

The negative cyclic homology of \mathcal{M} is isomorphic to the homology of $\mathcal{C} = (\wedge(u) \otimes \wedge(x, y, z, \bar{x}, \bar{y}, \bar{z}), D)$; see Section 3. Here, the differential D is given by

$$D(x) = u\bar{x}, \quad D(y) = u\bar{y}, \quad D(z) = xy + u\bar{z}, \quad D(\bar{x}) = D(\bar{y}) = 0, \quad D(\bar{z}) = -\bar{x}y + x\bar{y}.$$

Then the morphism β in Theorem 2.7 is induced by the derivation $s: \mathcal{L} \rightarrow \mathcal{C}$.

It follows from the BV-exactness of the manifold M that $\mathrm{HC}_*^-(\mathcal{M})$ decomposes into a direct sum $\mathbb{Q}[u] \oplus \mathrm{Im} \tilde{\beta}$, where $\tilde{\beta}$ is a morphism induced by the map $\tilde{s}: \tilde{\mathcal{L}} \rightarrow \tilde{\mathcal{C}}$ on the reduced complexes. Hence, by applying $\tilde{\beta}$ to the basis except for 1 in Proposition 6.3, we see that $\mathrm{Im} \tilde{\beta}$ is spanned by the homology classes

$$\zeta_{p,q} := \frac{1}{p!q!} \bar{x}^p \bar{y}^q, \quad \eta_{p,q} := \begin{cases} \frac{1}{p!q!} (z\bar{x}^p \bar{y}^q - x\bar{x}^{p-1} \bar{y}^q \bar{z}) & \text{if } p \neq 0, \\ \frac{1}{q!} (z\bar{y}^q - y\bar{y}^{q-1} \bar{z}) & \text{if } p = 0, \end{cases} \quad \theta_r := \frac{r+1}{r} xy\bar{z}^r,$$

for $p, q \geq 0, r \geq 1$ with $(p, q) \neq (0, 0)$. We also put $\zeta_{0,0} = 1$ for convenience. Denote by Dsb the dual string bracket $[\cdot, \cdot]^{\vee}$ over \mathbb{Q} stated in Theorem 2.8.

Theorem 6.6 For the dual string bracket Dsb over \mathbb{Q} of the 11-dimensional manifold M , one has

$$\begin{aligned} \mathrm{Dsb}(\zeta_{p,q}) &= \sum_{i=0}^{p+1} \sum_{j=0}^{q+1} \{i(q+1) - j(p+1)\} (\zeta_{i,j} \otimes \eta_{p+1-i, q+1-j} - \eta_{p+1-i, q+1-j} \otimes \zeta_{i,j}), \\ \mathrm{Dsb}(\eta_{p,q}) &= \theta_2 \otimes \zeta_{p,q} - \zeta_{p,q} \otimes \theta_2 - \sum_{i=0}^{p+1} \sum_{j=0}^{q+1} \{i(q+1) - j(p+1)\} \eta_{i,j} \otimes \eta_{p+1-i, q+1-j}, \\ \mathrm{Dsb}(\theta_r) &= 0. \end{aligned}$$

Proof We first compute the dual loop product Dlp by the rational model described in [19]. Let $\mathcal{M} = \wedge V$ be the minimal model for M , $\mathcal{P} = (\wedge V)^{\otimes 2} \otimes \wedge \bar{V}$ the Sullivan model for the free path space stated in [15, Section 15] and $\varepsilon_{\mathcal{P}}: \mathcal{P} \rightarrow \wedge V$ the $(\wedge V)^{\otimes 2}$ -semifree resolution of $\wedge V$ which is given by the multiplication of $\wedge V$ and the canonical augmentation of $\wedge \bar{V}$.

By virtue of [19, Lemma 1], we see that a DGA morphism $\mathcal{P} \rightarrow \mathcal{P} \otimes_{\wedge V} \mathcal{P}$ defined by $v_1 \otimes v_2 \mapsto v_1 \otimes 1 \otimes v_2$, $\bar{x} \mapsto 1 \otimes \bar{x} + \bar{x} \otimes 1$, $\bar{y} \mapsto 1 \otimes \bar{y} + \bar{y} \otimes 1$, $\bar{z} \mapsto 1 \otimes \bar{z} + \bar{z} \otimes 1 - \frac{1}{2}\bar{x} \otimes \bar{y} + \frac{1}{2}\bar{y} \otimes \bar{x}$ for $v_i \in V$ is a Sullivan representative for the composition of free paths. This induces a Sullivan representative $\mathcal{M}_{\mathrm{Comp}}: \mathcal{L} \rightarrow \mathcal{L} \otimes_{\wedge V} \mathcal{L}$ for Comp in (3.4) which has formulae

$$\begin{aligned} \mathcal{M}_{\mathrm{Comp}}(v) &= v, & \mathcal{M}_{\mathrm{Comp}}(\bar{x}) &= 1 \otimes \bar{x} + \bar{x} \otimes 1, \\ \mathcal{M}_{\mathrm{Comp}}(\bar{y}) &= 1 \otimes \bar{y} + \bar{y} \otimes 1, & \mathcal{M}_{\mathrm{Comp}}(\bar{z}) &= 1 \otimes \bar{z} + \bar{z} \otimes 1 - \frac{1}{2}\bar{x} \otimes \bar{y} + \frac{1}{2}\bar{y} \otimes \bar{x}, \end{aligned}$$

where $v \in V$. Recall the morphism $\varepsilon_{\mathcal{P}} \otimes 1: \mathcal{P} \otimes_{(\wedge V)^{\otimes 2}} \mathcal{L}^{\otimes 2} \rightarrow \wedge V \otimes_{(\wedge V)^{\otimes 2}} \mathcal{L}^{\otimes 2}$ appeared in the model for Dlp. A section σ of the morphism $\varepsilon_{\mathcal{P}} \otimes 1$ is given by

$$\begin{aligned} \sigma(v) &= v \otimes 1, & \sigma(\bar{x} \otimes 1) &= 1 \otimes (\bar{x} \otimes 1), & \sigma(1 \otimes \bar{x}) &= 1 \otimes (1 \otimes \bar{x}), & \sigma(\bar{y} \otimes 1) &= 1 \otimes (\bar{y} \otimes 1), \\ \sigma(1 \otimes \bar{y}) &= 1 \otimes (1 \otimes \bar{y}), & \sigma(\bar{z} \otimes 1) &= 1 \otimes (\bar{z} \otimes 1), & \sigma(1 \otimes \bar{z}) &= 1 \otimes (1 \otimes \bar{z}) - \bar{x} \otimes (1 \otimes \bar{y}) + \bar{y} \otimes (1 \otimes \bar{x}). \end{aligned}$$

Define a $(\wedge V)^{\otimes 2}$ -morphism $\text{Diag}^!: \mathcal{P} \rightarrow (\wedge V)^{\otimes 2}$ of degree 11 by

$$\text{Diag}^!(1) = (-x \otimes 1 + 1 \otimes x)(-y \otimes 1 + 1 \otimes y)(-z \otimes 1 + 1 \otimes z), \quad \text{Diag}^!|_{\wedge + \bar{V}} \equiv 0,$$

which gives a representative of a nonzero element in $\text{Ext}_{(\wedge V)^{\otimes 2}}^{11}(\wedge V, \wedge V)$; see [49, Section 5] for the detail about a construction of the shriek map $\text{Diag}^!$. Then, the result [19, Theorem A] yields that the composite

$$\mathcal{L} \xrightarrow{\mathcal{M}_{\text{Comp}}} \mathcal{L} \otimes_{\wedge V} \mathcal{L} \cong \wedge V \otimes_{(\wedge V)^{\otimes 2}} \mathcal{L}^{\otimes 2} \xrightarrow{\sigma} \mathcal{P} \otimes_{(\wedge V)^{\otimes 2}} \mathcal{L}^{\otimes 2} \xrightarrow{\text{Diag}^! \otimes 1} \mathcal{L}^{\otimes 2}$$

induces the dual loop product Dlp on homology. This rational model and a straightforward computation enable us to compute Dlp explicitly. In fact, we have

$$\begin{aligned} \text{Dlp}(\bar{x}^p \bar{y}^q) &= xyz \otimes \bar{x}^p \bar{y}^q + \bar{x}^p \bar{y}^q \otimes xyz \\ &\quad + \sum_{i=0}^p \sum_{j=0}^q \binom{p}{i} \binom{q}{j} (-x \bar{x}^i \bar{y}^j \otimes yz \bar{x}^{p-i} \bar{y}^{q-j} + y \bar{x}^i \bar{y}^j \otimes xz \bar{x}^{p-i} \bar{y}^{q-j} \\ &\quad \quad \quad - xz \bar{x}^i \bar{y}^j \otimes y \bar{x}^{p-i} \bar{y}^{q-j} + yz \bar{x}^i \bar{y}^j \otimes x \bar{x}^{p-i} \bar{y}^{q-j}), \end{aligned}$$

$$\begin{aligned} \text{Dlp}(z \bar{x}^p \bar{y}^q - x \bar{x}^{p-1} \bar{y}^q \bar{z}) &= xyz \otimes (z \bar{x}^p \bar{y}^q - x \bar{x}^{p-1} \bar{y}^q \bar{z}) + (z \bar{x}^p \bar{y}^q - x \bar{x}^{p-1} \bar{y}^q \bar{z}) \otimes xyz \\ &\quad + xyz \bar{z} \otimes x \bar{x}^{p-1} \bar{y}^q + x \bar{x}^{p-1} \bar{y}^q \otimes xyz \bar{z} + xy \bar{z} \otimes xz \bar{x}^{p-1} \bar{y}^q - xz \bar{x}^{p-1} \bar{y}^q \otimes xy \bar{z} \\ &\quad - \sum_{i=0}^p \sum_{j=0}^q \binom{p}{i} \binom{q}{j} (xz \bar{x}^i \bar{y}^j \otimes yz \bar{x}^{p-i} \bar{y}^{q-j} - yz \bar{x}^i \bar{y}^j \otimes xz \bar{x}^{p-i} \bar{y}^{q-j}), \end{aligned}$$

$$\begin{aligned} \text{Dlp}(z \bar{y}^q - y \bar{y}^{q-1} \bar{z}) &= -xyz \otimes (z \bar{y}^q - y \bar{y}^{q-1} \bar{z}) - (z \bar{y}^q - y \bar{y}^{q-1} \bar{z}) \otimes xyz + xyz \bar{z} \otimes y \bar{y}^{q-1} + y \bar{y}^{q-1} \otimes xyz \bar{z} \\ &\quad + xy \bar{z} \otimes yz \bar{y}^{q-1} - yz \bar{y}^{q-1} \otimes xy \bar{z} - \sum_{j=0}^q \binom{q}{j} (xz \bar{y}^j \otimes yz \bar{y}^{q-j} - yz \bar{y}^j \otimes xz \bar{y}^{q-j}), \end{aligned}$$

$$\text{Dlp}(xy \bar{z}^r) = \sum_{i=0}^r \binom{r}{i} (-xyz \bar{z}^i \otimes xy \bar{z}^{r-i} + xy \bar{z}^i \otimes xyz \bar{z}^{r-i}).$$

It follows from Theorem 2.8(ii) that

$$\begin{aligned} \text{Dsb}(\zeta_{p,q}) &= \frac{1}{p!q!} (\beta \otimes \beta) \bullet^{\vee} (\bar{x}^p \bar{y}^q), & \text{Dsb}(\eta_{p,q}) &= \frac{1}{p!q!} (\beta \otimes \beta) \bullet^{\vee} (z \bar{x}^p \bar{y}^q - x \bar{x}^{p-1} \bar{y}^q \bar{z}), \\ \text{Dsb}(\eta_{0,q}) &= \frac{1}{q!} (\beta \otimes \beta) \bullet^{\vee} (z \bar{y}^q - y \bar{y}^{q-1} \bar{z}), & \text{Dsb}(\theta_r) &= \frac{r+1}{r} (\beta \otimes \beta) \bullet^{\vee} (xy \bar{z}^r). \end{aligned}$$

Therefore, by these formulae and the computations of Dlp above, we have the result. \square

7 The cobar-type EMSS and r -BV-exactness

Let X be a simply connected space. We define a cobracket on the cobar-type Eilenberg–Moore spectral sequence converging to the rational equivariant cohomology of the free loop space LX , which is compatible with the dual to the string bracket in the sense of Chas and Sullivan [8] if X is a simply connected closed manifold.

We begin by recalling the spectral sequence associated with a filtered complex (A, F, d) . Consider the submodules $Z_r^{p,q}$ and $B_r^{p,q}$ defined by

$$(7.1) \quad Z_r^{p,q} := F^p A^{p+q} \cap d^{-1}(F^{p-r} A^{p+q+1}) \quad \text{and} \quad B_r^{p,q} := F^p A^{p+q} \cap d(F^{p-r} A^{p+q-1}).$$

With the submodules of A , we have a spectral sequence $\{E_r, d_r\}$ whose E_r -term is defined by $E_r^{p,q} := Z_r^{p,q} / (Z_{r-1}^{p+1,q-1} + B_{r-1}^{p,q})$; see [32, Proof of Theorem 2.6].

We use the same notation as that in Section 2. In particular, for a cochain algebra A , we define a chain algebra $A_\#$ by $(A_\#)_{-i} = A^i$ for i . The converse is also considered; that is, for a chain algebra Ω , we have a cochain algebra $\Omega^\#$ defined by $((\Omega)^\#)^i = \Omega_{-i}$ for i ; see Remark 2.6.

Let $(\wedge V, d)$ be a Sullivan model of a simply connected commutative cochain algebra A . Define $(\mathcal{L}, \delta) = (\wedge(V \oplus \bar{V}), \delta)$ and $(\mathcal{E}, D) = (\wedge u \otimes \mathcal{L}, D)$; see Section 3. Then complexes (\mathcal{L}, δ) and (\mathcal{E}, D) compute the Hochschild homology and the negative cyclic homology of $A_\#$, respectively. Thus we have the cobar-type Eilenberg–Moore spectral sequence (the EMSS for short) $\{E_r^{*,*}, d_r\}$ converging to $\text{HC}_*^-(A) := (\text{HC}_*^-(A_\#))^\#$ as an algebra with

$$E_2^{*,*} \cong \text{Cotor}_{\wedge(t)}^{*,*}(\text{HH}_*(A), \mathbb{Q})$$

as a bigraded algebra, where $|t| = 1$ and the $\wedge(t)$ -comodule structure on the Hochschild homology $\text{HH}_*(A) := (\text{HH}_*(A_\#))^\#$ is induced by the derivation s in the cyclic complex (\mathcal{E}, D) . In fact, the $\wedge(t)$ -comodule structure $\nabla: \mathcal{L} \rightarrow \mathcal{L} \otimes \wedge(t)$ on (\mathcal{L}, δ) is given by $\nabla(\alpha) = \gamma(\alpha) \otimes t + \alpha \otimes 1$, where $\gamma(\alpha) = (-1)^{|\alpha|} s(\alpha)$. A map assigning the element au^n in the cyclic complex \mathcal{E} to an element $a[t|\cdots|t]$ in the n^{th} cobar complex gives rise to an isomorphism of complexes. As a consequence, we have isomorphisms

$$\text{Cotor}_{\wedge(t)}^*(\mathcal{L}, \mathbb{Q}) \cong H(\mathcal{E}, D) \cong \text{HC}_*^-(A_\#)^\#.$$

Remark 7.2 The isomorphisms above allow us to work in the category of $\wedge(t)$ -comodule in order to investigate the negative cyclic homology of a DGA.

We observe that, by construction, there is an isomorphism $E_1^{0,*} \cong \text{HH}_*(A)$. In particular, when we choose the polynomial de Rham algebra $A_{\text{PL}}(M)$ for a simply connected space M as the DGA A , the spectral sequence converges to the S^1 -equivariant cohomology $\text{HC}_*^-(A) \cong H_{S^1}^*(LM; \mathbb{Q})$, with

$$E_2^{*,*} \cong \text{Cotor}_{H^*(S^1; \mathbb{Q})}^{*,*}(H^*(LM; \mathbb{Q}), \mathbb{Q}).$$

One has a direct sum decomposition $(\tilde{\mathcal{L}}, \delta) = \bigoplus_n (\tilde{\mathcal{L}}^{(n)}, \delta)$ of complexes, where $\tilde{\mathcal{L}}^{(n)} = \tilde{\mathcal{L}} \cap (\wedge^n V \otimes \wedge^n \bar{V})$ with $\mathcal{L} = \tilde{\mathcal{L}} \oplus \mathbb{Q}$. Then the reduced derivation \tilde{s} decomposes $(\tilde{\mathcal{L}}, \delta)$ into a sequence $0 \rightarrow \tilde{\mathcal{L}}^{(0)} \rightarrow \tilde{\mathcal{L}}^{(1)} \rightarrow \tilde{\mathcal{L}}^{(2)} \rightarrow \dots$ of complexes. Thus it follows that the EMSS $\{E_r^{*,*}, d_r\}$ is decomposed as

$$\{E_r^{*,*}, d_r\} = \bigoplus_{N \in \mathbb{Z}} \{({}_N E_r^{*,*}, d_r)\} \oplus \{\mathbb{Q}[u], 0\},$$

where $\text{bideg } u = (1, 1)$, each spectral sequence $\{({}_N E_r^{*,*}, d_r)\}$ for $N \geq 0$ is constructed by the double complex

$$({}_N)\mathcal{K}: 0 \rightarrow \tilde{\mathcal{L}}^{(N)} \rightarrow \tilde{\mathcal{L}}^{(N+1)} \otimes \mathbb{Q}\{u\} \rightarrow \tilde{\mathcal{L}}^{(N+2)} \otimes \mathbb{Q}\{u^2\} \rightarrow \dots$$

and for $N < 0$, the spectral sequence $\{({}_N E_r^{*,*}, d_r)\}$ is obtained by the double complex

$$({}_N)\mathcal{K}: 0 \rightarrow 0 \rightarrow \dots \rightarrow 0 \rightarrow \tilde{\mathcal{L}}^{(0)} \otimes \mathbb{Q}\{u^{-N}\} \rightarrow \tilde{\mathcal{L}}^{(1)} \otimes \mathbb{Q}\{u^{-N+1}\} \rightarrow \dots .$$

Here, the double complex $({}_N)\mathcal{K}$ is regarded as a filtered complex associated with the horizontal degrees. Thus, in the spectral sequence $\{({}_N E_r^{*,*}, d_r)\}$ for $N < 0$, we have $({}_N E_r^{i,*} = 0$ for $i < -N$. We observe that each spectral sequence $\{({}_N E_r^{*,*}, d_r)\}$ converges to the target as an algebra.

Remark 7.3 The direct sum of the targets of the spectral sequences $\{({}_N E_r^{*,*}, d_r)\}$ is nothing but the Hodge decomposition of $\text{HC}_*^-(A)$; that is, we have $\widetilde{\text{HC}}_*^-(A) = \bigoplus_{N \geq 0} H(\mathcal{K}(N))$; see [6, Section 2]. If A is the polynomial de Rham algebra $A_{\text{PL}}(X)$ for a simply connected space X , then the direct summands in the Hodge decomposition are identified with the eigenspaces of the Adams operation on $\tilde{H}_{S^1}^*(LX; \mathbb{Q})$; see [6, Theorem 3.2] for the identification. We refer the reader to [30, 4.5.4] for the operation. The result [3, Theorem 1.1] shows that the string bracket respects the Hodge decomposition in some sense. Thus, we are also interested in computations of string brackets, as described in Section 1.1, together with the consideration of the Hodge decomposition.

Proposition 7.4 *If the spectral sequence $\{({}_0 E_r, d_r)\}$ collapses at E_r -term, then so does $\{({}_N E_r, d_r)\}$ for each integer N , and then $\text{Tot } E_r \cong H_{S^1}^*(LX)$ as a vector space.*

Thus, it is readily seen that the collapsing of the EMSS is governed by that of the zeroth spectral sequence.

Corollary 7.5 *The spectral sequence $\{({}_0 E_r, d_r)\}$ collapses at the E_r -term if and only if so does $\{E_r^{*,*}, d_r\}$.*

Lemma 7.6 *For $l \geq r - 1$ and $N \in \mathbb{Z}$, we have $({}_0 E_r^{l+N, *+N} \cong ({}_N E_r^{l,*}$.*

Proof For $N < 0$, the multiplication $u^{-N} \times: ({}_0 E_r^{*,*} \rightarrow ({}_N E_r^{*-N, * - N}$ gives an isomorphism. Assume that $N \geq 0$. By definition, we see that

$$\begin{aligned} ({}_N E_r^{l,*} &= ({}_N Z_r^{l,*} / (({}_N Z_{r-1}^{l+1, * - 1} + ({}_N B_{r-1}^{l,*}), \\ ({}_0 E_r^{l+N, *+N} &= ({}_0 Z_r^{l+N, *+N} / (({}_0 Z_{r-1}^{l+N+1, *+N-1} + ({}_0 B_{r-1}^{l+N, *+N}), \end{aligned}$$

where $(N)Z$ and $(N)B$ denote the subcomplexes of $(N)\mathcal{H}$ defined in (7.1) for the filtered complex $A = (N)\mathcal{H}$. Moreover, we have $(N)Z_r^{l,*} \cong (0)Z_r^{l+N,*+N}$ and $(N)Z_{r-1}^{l+1,*-1} \cong (0)Z_{r-1}^{l+N+1,*+N-1}$. Since $l \geq r - 1$, it follows that $(N)B_{r-1}^{l,*} \cong (0)B_{r-1}^{l+N,*+N}$. Then the multiplication $u^N \times: (N)E_r^{l,*} \cong (0)E_r^{l+N,*+N}$ is an isomorphism. \square

Proof of Proposition 7.4 Lemma 4.7 yields that the spectral sequence $\{(0)E_r^{*,*}, d_r\}$ converges to 0, the trivial module. The assumption and Lemma 7.6 imply that $(N)E_r^{l,*} = 0$ for $l \geq r - 1$ and N . \square

Theorem 7.7 Let M be a simply connected closed manifold, and A the polynomial de Rham algebra $A_{\text{PL}}(M)$ of M . Then the map $[\cdot, \cdot]_r^{\vee}: E_r^{p,*} \rightarrow (E_r^{*,*} \otimes E_r^{*,*})^{p,*+d-2}$ defined by $[\cdot, \cdot]_r^{\vee} \equiv 0$ for $p > 0$ and for $p = 0$, the composite

$$E_r^{0,*} = \text{Ker } d_{r-1} \xrightarrow{i} \text{HH}_*(A) \xrightarrow{\bullet^{\vee}} \text{HH}_*(A)^{\otimes 2} \xrightarrow{\Delta \otimes \Delta} E_r^{0,*} \otimes E_r^{0,*}$$

gives rise to a cobracket on the spectral sequence, where i denotes the inclusion. That is, it is compatible with the differentials and $H([\cdot, \cdot]_r^{\vee}) = [\cdot, \cdot]_{r+1}^{\vee}$. Moreover, the cobracket $[\cdot, \cdot]_{\infty}^{\vee}$ is compatible with the dual to the string bracket on $H_*^{S^1}(LM)$ at the E_{∞} -term in the sense that the composite

$$H_{S^1}^*(LM) \xrightarrow{\pi} E_{\infty}^{0,*} \xrightarrow{i} \text{HH}_*(A) \xrightarrow{\bullet^{\vee}} \text{HH}_*(A)^{\otimes 2} \xrightarrow{\Delta \otimes \Delta} E_{\infty}^{0,*} \otimes E_{\infty}^{0,*}$$

coincides with the dual to the string bracket modulo $F^1 H_{S^1}^*(LM)$. Here π is the projection and $\{F^l H_{S^1}^*(LM)\}_{l \geq 0}$ is the decreasing filtration associated with the spectral sequence.

Proof By dimensional reasons, it is readily seen that $(d_r \otimes 1 \pm 1 \otimes d_r) \circ [\cdot, \cdot]_r^{\vee} = 0 = [\cdot, \cdot]_r^{\vee} \circ d_r$ for $p > 0$. Moreover, we see that every element in the image of Δ in $E_r^{0,*}$ is a permanent cocycle. In fact, for $w \in \text{Im } \Delta$, we have $Dw = (\delta + us)w = 0$. Then, it follows that $(d_r \otimes 1 \pm 1 \otimes d_r) \circ [\cdot, \cdot]_r^{\vee} = 0 = [\cdot, \cdot]_r^{\vee} \circ d_r$ in $E_r^{0,*}$. By the definition of the cobrackets, we have $H([\cdot, \cdot]_r^{\vee}) = [\cdot, \cdot]_{r+1}^{\vee}$. In fact, the left-hand side is the restriction of $[\cdot, \cdot]_r^{\vee}$ in the nontrivial case.

Consider the compatibility of the cobracket at the E_{∞} -term. We have a commutative diagram

$$\begin{array}{ccccccc} \text{HC}_*^-(A) & \xrightarrow{\pi} & \text{HH}_*(A) & \xrightarrow{\bullet^{\vee}} & \text{HH}_*(A)^{\otimes 2} & \xrightarrow{\beta \otimes \beta} & \text{HC}_*^-(A)^{\otimes 2} \\ \cong \downarrow & & \parallel & & & \nearrow \beta \otimes \beta & \downarrow \text{pr} \\ \text{HC}_*^-(A)/F^1 \oplus F^1/F^2 \oplus \dots \oplus F^* & & & & & & (\text{HC}_*^-(A)/F^1)^{\otimes 2} \\ \cong \downarrow & & & & & & \downarrow \cong \\ \bigoplus_{p+q=*} E_{\infty}^{p,q} & \xrightarrow{\text{pr}} & E_{\infty}^{0,*} = \text{Ker } d_{*+1} & \xrightarrow{i} & \text{HH}_*(A) & \xrightarrow{\bullet^{\vee}} & \text{HH}_*(A)^{\otimes 2} & \xrightarrow{\Delta \otimes \Delta} & E_{\infty}^{0,*} \otimes E_{\infty}^{0,*} \end{array}$$

where $\{F^l\}_{l \geq 0}$ denotes the decreasing filtration of $\text{HC}_*^-(A) \cong H_{S^1}^*(LM)$ associated with the spectral sequence. In fact, the commutativity of the left-hand side square and the right-hand side triangle follows from the construction of the spectral sequence; see for example [32, Proof of Theorem 2.6]. Theorem 2.8(ii) implies the upper sequence is the dual of the string bracket. We have the result. \square

Proposition 7.8 For each N , the S -action $u \times$ on (\mathcal{E}, D) gives rise to a map

$$S: \{(N)E_r^{*,*}, d_r\} \rightarrow \{(N-1)E_r^{*+1,*+1}, d_r\}$$

on the spectral sequence which is compatible with the S -action on the negative cyclic homology $\mathrm{HC}_*^-(A)$.

Proof The S -action on (\mathcal{E}, D) gives rise to a map $(N)\mathcal{H} \rightarrow (N-1)\mathcal{H}$ which increases the filtration degree by $+1$ and is compatible with the differential. Then the map induces the action S on the spectral sequence. \square

Lemma 7.9 Suppose that the E_r -term $(0)E_r^{p,q}$ in $\{(0)E_r^{*,*}, d_r\}$ is trivial for any (p, q) . Then the $(r-1)$ times S -action $S^{r-1}: \widetilde{\mathrm{HC}}_*^-(A) \rightarrow \widetilde{\mathrm{HC}}_*^-(A)$ is trivial.

Proof Let x be an element in $\widetilde{\mathrm{HC}}_*^-(A)$. Then x is in $\widetilde{\mathrm{HC}}_*^{-(n)}(A)$ for some $n \geq 0$ and then it is represented by an element α in $(n)E_\infty^{t,*}$ for some $t \geq 0$. Thus the element $S^{r-1}x$ is represented by $S^{r-1}\alpha \in (n-(r-1))E_\infty^{t+(r-1),*}$. By assumption, it follows from Lemma 7.6 that $(N)E_r^{l,*} = 0$ for $l \geq r-1$ and $N \in \mathbb{Z}$. This implies that $S^{r-1}\alpha = 0$ in the E_∞ -term and that there is no extension problem; that is, $S^{r-1}x = S^{r-1}\alpha = 0$ in $\mathrm{HC}_*^{-(n-(r-1))}(A) \subset \mathrm{HC}_*^-(A)$. This completes the proof. \square

Moreover, we have:

Theorem 7.10 The E_r -term $(0)E_r^{p,q}$ in $\{(0)E_r^{*,*}, d_r\}$ is trivial for any (p, q) if and only if the $(r-1)$ times S -action S^{r-1} on $\widetilde{\mathrm{HC}}_*^-(A)$ is.

Proof The “only if” part follows from Lemma 7.9. To prove the “if” part, we assume that S^{r-1} is trivial on $\widetilde{\mathrm{HC}}_*^-(A)$. Take any element $x = x_p \otimes u^p + x_{p+1} \otimes u^{p+1} + \cdots \in (0)Z_r^{p,*}$, where $x_i \in \widetilde{\mathcal{L}}^{(i)}$ is zero for sufficiently large i . By the definition of $(0)Z_r^{p,*}$, the total differential increases the filtration degree of x by r , ie we have $dx_p = 0$ and $\tilde{s}x_i + dx_{i+1} = 0$ for $p \leq i \leq p+r-2$. Now we have an element $[\tilde{s}x_{p+r-1}] \in H(\mathrm{Ker} \tilde{s})$ and the above equation implies that $[dx_{p+1}] = S^{r-1}[sx_{p+r-1}] = 0 \in H(\mathrm{Ker} \tilde{s})$ by Lemma 4.10 and the assumption of triviality of S^{r-1} . By Lemma 4.7, we see that $\mathrm{Ker} \tilde{s} = \mathrm{Im} \tilde{s}$. Thus, there is an element $v_p \in \widetilde{\mathcal{L}}^{(p)}$ with $d\tilde{s}v_p = dx_{p+1}$. By using these elements, we define $y = (x_{p+1} - \tilde{s}v_p) \otimes u^{p+1} + x_{p+2} \otimes u^{p+2} + x_{p+3} \otimes u^{p+3} + \cdots \in (0)Z_{r-1}^{p+1,*}$. Then we can show $x - y = x_p \otimes u^p + \tilde{s}v_p \otimes u^{p+1} \in (0)B_{r-1}^{p,*}$ by the same argument as above. It follows that $x = y + (x - y) \in (0)Z_{r-1}^{p+1,*} + (0)B_{r-1}^{p,*}$ and hence $[x] = 0 \in (0)Z_r^{p,*} / (0)Z_{r-1}^{p+1,*} + (0)B_{r-1}^{p,*} = (0)E_r^{p,*}$. Since x is an arbitrary element of $(0)Z_r^{p,*}$, this proves the “if” part. \square

The BV-exactness of a space is equivalent to the condition that the E_2 -term of the spectral sequence $\{(0)E_r^{*,*}, d_r\}$ is trivial. Then Theorem 7.10 gives another proof of Theorem 2.11. This consideration allows us to propose a *higher version* of the BV-exactness.

Definition 7.11 A simply connected space X is r -BV-exact if the E_{r+1} -term $(0)E_{r+1}^{p,q}$ in the spectral sequence $\{(0)E_r^{*,*}, d_r\}$ associated with X is trivial for any (p, q) .

Indeed, there exists a non-BV-exact space in the class of rational elliptic spaces; see Appendix A below. While we are interested in the hierarchy of rational spaces defined by the r -BV-exactness as seen in Section 1.1, we do not pursue the topic in this manuscript.

Appendix A A non-BV-exact space

We describe an example of a non-BV-exact space. Let $(\wedge V, d)$ be the minimal model

$$(\wedge(x_1, x_2, y_1, y_2, y_3, z), d)$$

of an elliptic space X of dimension 228, given in [1, Example 5.2]. The degrees are given by $|x_1| = 10$, $|x_2| = 12$, $|y_1| = 41$, $|y_2| = 43$, $|y_3| = 45$ and $|z| = 119$. The differential is as follows:

$$\begin{aligned} dx_1 &= 0, & dy_1 &= x_1^3 x_2, \\ dx_2 &= 0, & dy_2 &= x_1^2 x_2^2, & dz &= x_2(y_1 x_2 - x_1 y_2)(y_2 x_2 - x_1 y_3) + x_1^{12} + x_2^{10}. \\ & & dy_3 &= x_1 x_2^3, \end{aligned}$$

Note that X does not admit positive weights. Indeed, let $\text{wt}(x_1) = i$ and $\text{wt}(x_2) = j$. Then $\text{wt}(y_1) = 3i + j$, $\text{wt}(y_2) = 2i + 2j$ and $\text{wt}(y_3) = i + 3j$. By dz , we have the equations $5i + 6j = 12i = 10j$ induced from $\text{wt}(x_2(y_1 x_2 - x_1 y_2)(y_2 x_2 - x_1 y_3)) = \text{wt}(x_1^{12}) = \text{wt}(x_2^{10})$. Thus we obtain $i = j = 0$.

Let $\omega = x_1^{14} y_2 y_3 - x_1^{13} x_2 y_1 y_3 + x_1^{12} x_2^2 y_1 y_2$ be the representing cocycle of the fundamental class of the manifold, which is considered as an element of $\wedge^+ V = \tilde{\mathcal{L}}^{(0)} \subset \tilde{\mathcal{L}}$. Then $[\omega] \notin \text{Im}(H(\tilde{s}): 0 \rightarrow H(\tilde{\mathcal{L}}^{(0)})) = 0$. On the other hand, we have $[\omega] \in \text{Ker}(H(\tilde{s}): H(\tilde{\mathcal{L}}^{(0)}) \rightarrow H(\tilde{\mathcal{L}}^{(1)}))$, since $\tilde{s}(\omega) = \delta(\alpha)$ for the element α defined by

$$\begin{aligned} \alpha &= -1380x_1^{11}x_2^6\bar{y}_3 - 5290x_1^{11}x_2^5y_3\bar{x}_2 - 114x_1^{10}y_1y_2\bar{y}_2 + 114x_1^{10}y_1y_3\bar{y}_1 \\ &\quad - 114x_1^9x_2y_1y_2\bar{y}_1 + \frac{93}{2}x_1^2y_2y_3\bar{z} + x_1^2y_2z\bar{y}_3 - x_1^2y_3z\bar{y}_2 + 114x_1x_2^7y_2y_3\bar{y}_3 \\ &\quad - \frac{93}{2}x_1x_2y_1y_3\bar{z} - x_1x_2y_1z\bar{y}_3 - 114x_1x_2y_2z\bar{y}_2 + 115x_1x_2y_3z\bar{y}_1 + 113x_1y_1y_3z\bar{x}_2 \\ &\quad + 572x_1y_2y_3z\bar{x}_1 + 115x_2^9\bar{z} - 114x_2^8y_1y_3\bar{y}_3 + 114x_2^8y_2y_3\bar{y}_2 + 1150x_2^8z\bar{x}_2 \\ &\quad + \frac{93}{2}x_2^2y_1y_2\bar{z} + 115x_2^2y_1z\bar{y}_2 - 115x_2^2y_2z\bar{y}_1 - 340x_2y_1y_2z\bar{x}_2 - 229x_2y_1y_3z\bar{x}_1. \end{aligned}$$

Hence we have $\text{Im } H(\tilde{s}) \subsetneq \text{Ker } H(\tilde{s})$, ie $(\wedge V, d)$ is not BV-exact. Note that we have found the element α by using the program [51] mentioned in Remark 6.4, while the equality is also checked by hand.

Finally we consider the differentials in the spectral sequence $\{({}_0)E_r, d_r\}$ defined in Section 7. Since $d_1[\omega] = H(\tilde{s})[\omega] = 0$, the cocycle ω defines an element $[\omega] \in ({}_0)E_2$, where $({}_0)E_2$ is considered as a subquotient of $({}_0)E_1 = H(\tilde{\mathcal{L}}, \delta)$. Then the equality $\tilde{s}(\omega) = \delta(\alpha)$ enables us to compute $d_2[\omega] = [\tilde{s}\alpha] \neq 0 \in ({}_0)E_2$, where the nontriviality is proved by using the program [51]. Thus this Sullivan algebra gives an example such that $d_2 \neq 0$ on $({}_0)E_2$. Note that it is currently unknown whether $({}_0)E_3 = 0$ (ie 2-BV-exact) or not.

Appendix B Connes’ B–map in the Gysin exact sequence

In this section, by giving precisely a rational model for the *integration over the fiber* $\beta: H^{*+1}(LX; \mathbb{Q}) \rightarrow H_{S^1}^*(LX; \mathbb{Q})$, we describe the Gysin exact sequence of the S^1 –principal bundle

$$S^1 \rightarrow ES^1 \times LX \xrightarrow{p} ES^1 \times_{S^1} LX$$

in terms of Sullivan models for LX and $ES^1 \times_{S^1} LX$. As a consequence, the Gysin sequence is identified with Connes’ exact sequence under the isomorphisms $HH_*(\mathcal{M}(X)) \cong H^{-*}(LX; \mathbb{Q})$ and $HC_*^-(\mathcal{M}(X)) \cong H^{-*}(ES^1 \times_{S^1} LX; \mathbb{Q})$ described in Section 3.

The cohomology Gysin sequence associated with the bundle has the form

$$\dots \rightarrow H_{S^1}^{*-2}(LX) \xrightarrow{S} H_{S^1}^*(LX) \xrightarrow{p^*} H^*(LX) \xrightarrow{\beta} H_{S^1}^{*-1}(LX) \rightarrow \dots,$$

in which S is defined by the cup product with the Euler class $q^*(u)$, where u is the generator of $H^2(BS^1)$. The S^1 –principal bundle above fits in the pullback diagram

$$\begin{array}{ccccc} & & S^1 & \xlongequal{\quad} & S^1 \\ & & \downarrow & & \downarrow \\ LX & \longrightarrow & ES^1 \times LX & \longrightarrow & ES^1 \\ \parallel & & \downarrow p & & \downarrow \\ LX & \longrightarrow & ES^1 \times_{S^1} LX & \xrightarrow{q} & BS^1 \end{array}$$

in which the lower sequence is the fiber bundle associated with the universal bundle $ES^1 \rightarrow BS^1$.

We recall the Sullivan models \mathcal{L} and \mathcal{E} defined in Section 3. In the model \mathcal{E} for $ES^1 \times_{S^1} LX$, we write u for the Euler class $q^*(u)$. Thus the map S in the Gysin sequence is regarded as the multiplication by u in the models, namely the S –action in Connes’ exact sequence. In order to obtain rational models for p and β , we here consider the relative Sullivan algebra $\mathcal{L}^\wedge := (\mathcal{E} \otimes \wedge(e), \hat{\delta})$ with base \mathcal{E} , where $\hat{\delta}(e) = u$ and $|e| = 1$.

Lemma B.1 *The canonical projection $\rho: \mathcal{L}^\wedge \rightarrow \mathcal{L}$ is a homotopy equivalence.*

Proof We define a DGA morphism $\iota: \mathcal{L} \rightarrow \mathcal{L}^\wedge$ by $\iota(\alpha) = \alpha + (-1)^{|\alpha|}s(\alpha)e$ for $\alpha \in \mathcal{L}$, where s is the derivation on \mathcal{L} stated in Section 3. Then, we have $\rho \circ \iota = 1$ by definition. Moreover, a homotopy $\mathcal{L}^\wedge \rightarrow \mathcal{L}^\wedge \otimes \wedge(t, dt)$ defined by $e \mapsto et, u \mapsto ut - e dt$ and $\alpha \mapsto \alpha + (-1)^{|\alpha|}s(\alpha)e(1 - t)$ for $\alpha \in \mathcal{E}$ implies that $\iota \circ \rho$ is homotopic to 1. This completes the proof. □

Proposition B.2 *The derivation $s: \mathcal{L} \rightarrow \mathcal{E}$ is a rational model for β .*

Proof From Lemma B.1, the inclusion $\mathcal{E} \hookrightarrow \mathcal{L}^\wedge$ is a rational model for the principal bundle

$$p: ES^1 \times LM \rightarrow ES^1 \times_{S^1} LM.$$

Since β is the fiber integration associated with the principal bundle, it is modeled by a map $\int_{\mathcal{E}}: \mathcal{L}^\wedge \rightarrow \mathcal{E}$ defined by $\int_{\mathcal{E}}(\alpha_0 + \alpha_1 e) = \alpha_1$ for $\alpha_i \in \mathcal{E}$. Therefore, the result follows since the composite $\int_{\mathcal{E}} \iota$ coincides with the derivation s . □

List of symbols

symbol	meaning	page
•	the loop product	2633
\odot	the dual loop coproduct	2633
$[\ ,]$	the string bracket, dual string cobracket	2624
Δ	the BV operator on the Hochschild homology of a differential graded algebra	2626
Δ'	the BV operator on the homology of LM	2627
$HH_*(\Omega)$	the Hochschild homology of a DGA Ω	2630
$\widetilde{HH}_*(\Omega)$	the reduced Hochschild homology, $HH_*(\Omega) \cong \widetilde{HH}_*(\Omega) \oplus \mathbb{K}$	2630
$HC_*^-(\Omega)$	the negative cyclic homology of a DGA Ω	2630
$\widetilde{HC}_*^-(\Omega)$	the reduced negative cyclic homology, $HC_*^-(\Omega) \cong \widetilde{HC}_*^-(\Omega) \oplus \mathbb{K}[u]$	2630
S	the S -action on the negative cyclic homology	2630
$\mathcal{L}, (\mathcal{L}, \delta)$	the Sullivan minimal model for the free loop space LM (and the Hochschild homology)	2631
$\mathcal{E}, (\mathcal{E}, D)$	the Sullivan minimal model for the Borel construction $ES^1 \times_{S^1} LM$ (and the negative cyclic homology)	2631
$(\widetilde{\mathcal{L}}, \delta)$	the reduced version of (\mathcal{L}, δ)	2637
$(\widetilde{\mathcal{L}}^{(n)}, \delta)$	a direct summand of $(\widetilde{\mathcal{L}}, \delta) = \bigoplus_n (\widetilde{\mathcal{L}}^{(n)}, \delta)$	2637
s	a derivation on \mathcal{L} , which is a chain model of Δ	2631

Acknowledgements The authors thank Jean-Claude Thomas and Luc Menichi for comments on the first draft of this manuscript. The authors are grateful to the referee for a very careful reading of the manuscript, valuable suggestions and comments.

Kuribayashi was partially supported by a Grant-in-Aid for Scientific Research (B) 21H00982 from Japan Society for the Promotion of Science. Naito was supported by JSPS KAKENHI grant JP18K13403. Wakatsuki was supported by JSPS KAKENHI grant 20J00404. Yamaguchi was partially supported by JSPS KAKENHI grant 20K03591.

References

- [1] **M Arkowitz, G Lupton**, *Rational obstruction theory and rational homotopy sets*, *Math. Z.* 235 (2000) 525–539 MR Zbl
- [2] **S Basu**, *Transversal string topology & invariants of manifolds*, PhD thesis, State University of New York at Stony Brook (2011) MR Available at <https://www.proquest.com/docview/898888965>
- [3] **Y Berest, A C Ramadoss, Y Zhang**, *Hodge decomposition of string topology*, *Forum Math. Sigma* 9 (2021) art. id. e33 MR Zbl
- [4] **R Body, R Douglas**, *Unique factorization of rational homotopy types*, *Pacific J. Math.* 90 (1980) 21–26 MR Zbl
- [5] **R Body, M Mimura, H Shiga, D Sullivan**, *p -universal spaces and rational homotopy types*, *Comment. Math. Helv.* 73 (1998) 427–442 MR Zbl
- [6] **D Burghlelea, Z Fiedorowicz, W Gajda**, *Adams operations in Hochschild and cyclic homology of de Rham algebra and free loop spaces, K -Theory* 4 (1991) 269–287 MR Zbl
- [7] **D Burghlelea, M Vigué-Poirrier**, *Cyclic homology of commutative algebras, I*, from “Algebraic topology: rational homotopy” (Y Félix, editor), *Lecture Notes in Math.* 1318, Springer (1988) 51–72 MR Zbl
- [8] **M Chas, D Sullivan**, *String topology*, preprint (1999) arXiv math/9911159
- [9] **D Chataur, L Menichi**, *String topology of classifying spaces*, *J. Reine Angew. Math.* 669 (2012) 1–45 MR Zbl
- [10] **X Chen**, *An algebraic chain model of string topology*, *Trans. Amer. Math. Soc.* 364 (2012) 2749–2781 MR Zbl
- [11] **X Chen, F Eshmatov, L Liu**, *Gravity algebra structure on the negative cyclic homology of Calabi–Yau algebras*, *J. Geom. Phys.* 147 (2020) 103522, 19 MR Zbl
- [12] **RL Cohen, JDS Jones, J Yan**, *The loop homology algebra of spheres and projective spaces*, from “Categorical decomposition techniques in algebraic topology” (G Arone, J Hubbuck, R Levi, M Weiss, editors), *Progr. Math.* 215, Birkhäuser, Basel (2004) 77–92 MR Zbl
- [13] **C Costoya, A Viruel**, *Every finite group is the group of self-homotopy equivalences of an elliptic space*, *Acta Math.* 213 (2014) 49–62 MR Zbl
- [14] **Y Félix, S Halperin, J-C Thomas**, *Gorenstein spaces*, *Adv. in Math.* 71 (1988) 92–112 MR Zbl
- [15] **Y Félix, S Halperin, J-C Thomas**, *Rational homotopy theory*, *Graduate Texts in Math.* 205, Springer (2001) MR Zbl
- [16] **Y Félix, L Menichi, J-C Thomas**, *Gerstenhaber duality in Hochschild cohomology*, *J. Pure Appl. Algebra* 199 (2005) 43–59 MR Zbl
- [17] **Y Félix, J-C Thomas**, *Rational BV -algebra in string topology*, *Bull. Soc. Math. France* 136 (2008) 311–327 MR Zbl
- [18] **Y Félix, J-C Thomas**, *String topology on Gorenstein spaces*, *Math. Ann.* 345 (2009) 417–452 MR Zbl
- [19] **Y Félix, J-C Thomas, M Vigué-Poirrier**, *Rational string topology*, *J. Eur. Math. Soc.* 9 (2007) 123–156 MR Zbl
- [20] **BL Feigin, BL Tsygan**, *Additive K -theory*, from “ K -theory, arithmetic and geometry” (Y I Manin, editor), *Lecture Notes in Math.* 1289, Springer (1987) 67–209 MR Zbl

- [21] **E Getzler**, *Two-dimensional topological gravity and equivariant cohomology*, *Comm. Math. Phys.* 163 (1994) 473–489 MR Zbl
- [22] **E Getzler, JDS Jones**, *A_∞ -algebras and the cyclic bar complex*, *Illinois J. Math.* 34 (1990) 256–283 MR Zbl
- [23] **E Getzler, JDS Jones, S Petrack**, *Differential forms on loop spaces and the cyclic bar complex*, *Topology* 30 (1991) 339–371 MR Zbl
- [24] **S Halperin, J Stasheff**, *Obstructions to homotopy equivalences*, *Adv. in Math.* 32 (1979) 233–279 MR Zbl
- [25] **RA Hepworth**, *String topology for Lie groups*, *J. Topol.* 3 (2010) 424–442 MR Zbl
- [26] **JDS Jones**, *Cyclic homology and equivariant homology*, *Invent. Math.* 87 (1987) 403–423 MR Zbl
- [27] **K Kuribayashi, L Menichi**, *The Batalin–Vilkovisky algebra in the string topology of classifying spaces*, *Canad. J. Math.* 71 (2019) 843–889 MR Zbl
- [28] **K Kuribayashi, L Menichi, T Naito**, *Derived string topology and the Eilenberg–Moore spectral sequence*, *Israel J. Math.* 209 (2015) 745–802 MR Zbl
- [29] **K Kuribayashi, T Yamaguchi**, *On additive K -theory with the Loday–Quillen $*$ -product*, *Math. Scand.* 87 (2000) 5–21 MR Zbl
- [30] **J-L Loday**, *Cyclic homology*, *Grundlehren Math. Wissen.* 301, Springer (1992) MR Zbl
- [31] **J-L Loday, D Quillen**, *Cyclic homology and the Lie algebra homology of matrices*, *Comment. Math. Helv.* 59 (1984) 569–591 MR Zbl
- [32] **J McCleary**, *A user’s guide to spectral sequences*, 2nd edition, *Cambridge Studies in Advanced Mathematics* 58, Cambridge Univ. Press (2001) MR Zbl
- [33] **L Menichi**, *Batalin–Vilkovisky algebra structures on Hochschild cohomology*, *Bull. Soc. Math. France* 137 (2009) 277–295 MR Zbl
- [34] **L Menichi**, *String topology for spheres*, *Comment. Math. Helv.* 84 (2009) 135–157 MR Zbl
- [35] **L Menichi**, *A Batalin–Vilkovisky algebra morphism from double loop spaces to free loops*, *Trans. Amer. Math. Soc.* 363 (2011) 4443–4462 MR Zbl
- [36] **M Mimura, RC O’Neill, H Toda**, *On p -equivalence in the sense of Serre*, *Jpn. J. Math.* 40 (1971) 1–10 MR Zbl
- [37] **M Mimura, H Toda**, *On p -equivalences and p -universal spaces*, *Comment. Math. Helv.* 46 (1971) 87–97 MR Zbl
- [38] **JW Morgan**, *The algebraic topology of smooth algebraic varieties*, *Inst. Hautes Études Sci. Publ. Math.* 48 (1978) 137–204 MR Zbl
- [39] **T Naito**, *String operations on rational Gorenstein spaces*, preprint (2013) arXiv 1301.1785
- [40] **H Scheerer**, *Fibrewise P -universal nilpotent fibrations*, *Proc. Roy. Soc. Edinburgh Sect. A* 98 (1984) 89–104 MR Zbl
- [41] **J-P Serre**, *Groupes d’homotopie et classes de groupes abéliens*, *Ann. of Math.* 58 (1953) 258–294 MR Zbl
- [42] **L Smith**, *Homological algebra and the Eilenberg–Moore spectral sequence*, *Trans. Amer. Math. Soc.* 129 (1967) 58–93 MR Zbl

- [43] **F Y Tabing**, *String homology and Lie algebra structures*, PhD thesis, University of California, Santa Cruz (2015) MR arXiv 1610.03933
- [44] **H Tamanoi**, *Cap products in string topology*, *Algebr. Geom. Topol.* 9 (2009) 1201–1224 MR Zbl
- [45] **M Vigué-Poirrier**, *Sur l’algèbre de cohomologie cyclique d’un espace 1–connexe applications à la géométrie des variétés*, *Illinois J. Math.* 32 (1988) 40–52 MR Zbl
- [46] **M Vigué-Poirrier**, *Homologie cyclique des espaces formels*, *J. Pure Appl. Algebra* 91 (1994) 347–354 MR Zbl
- [47] **M Vigué-Poirrier, D Burghlea**, *A model for cyclic homology and algebraic K–theory of 1–connected topological spaces*, *J. Differential Geom.* 22 (1985) 243–253 MR Zbl
- [48] **M Vigué-Poirrier, D Sullivan**, *The homology theory of the closed geodesic problem*, *J. Differential Geometry* 11 (1976) 633–644 MR Zbl
- [49] **S Wakatsuki**, *Description and triviality of the loop products and coproducts for rational Gorenstein spaces*, preprint (2016) arXiv 1612.03563
- [50] **S Wakatsuki**, *Coproducts in brane topology*, *Algebr. Geom. Topol.* 19 (2019) 2961–2988 MR Zbl
- [51] **S Wakatsuki**, *Kohomology: cohomology calculator for Sullivan algebras* (2024) Available at <https://github.com/shwaka/kohomology/index.html>
- [52] **G W Whitehead**, *Elements of homotopy theory*, *Graduate Texts in Math.* 61, Springer (1978) MR Zbl
- [53] **T Yang**, *A Batalin–Vilkovisky algebra structure on the Hochschild cohomology of truncated polynomials*, *Topology Appl.* 160 (2013) 1633–1651 MR Zbl

*Department of Mathematical Sciences, Faculty of Science, Shinshu University
Matsumoto, Nagano, Japan*

*Nippon Institute of Technology
Gakuendai, Miyashiro-machi, Minamisaitama-gun, Saitama, Japan*

*Department of Mathematical Sciences, Faculty of Science, Shinshu University
Matsumoto, Nagano, Japan*

*Faculty of Education, Kochi University
Akebono-cho, Kochi, Japan*

kuri@math.shinshu-u.ac.jp, naito.takahito@nit.ac.jp,
shun.wakatsuki@math.nagoya-u.ac.jp, tyamag@kochi-u.ac.jp

Received: 12 May 2022 Revised: 11 May 2023

ALGEBRAIC & GEOMETRIC TOPOLOGY

msp.org/agt

EDITORS

PRINCIPAL ACADEMIC EDITORS

John Etnyre
etnyre@math.gatech.edu
Georgia Institute of Technology

Kathryn Hess
kathryn.hess@epfl.ch
École Polytechnique Fédérale de Lausanne

BOARD OF EDITORS

Julie Bergner	University of Virginia jeb2md@eservices.virginia.edu	Robert Lipshitz	University of Oregon lipshitz@uoregon.edu
Steven Boyer	Université du Québec à Montréal cohf@math.rochester.edu	Norihiko Minami	Yamato University minami.norihiko@yamato-u.ac.jp
Tara E Brendle	University of Glasgow tara.brendle@glasgow.ac.uk	Andrés Navas	Universidad de Santiago de Chile andres.navas@usach.cl
Indira Chatterji	CNRS & Univ. Côte d'Azur (Nice) indira.chatterji@math.cnrs.fr	Thomas Nikolaus	University of Münster nikolaus@uni-muenster.de
Alexander Dranishnikov	University of Florida dranish@math.ufl.edu	Robert Oliver	Université Paris 13 bobol@math.univ-paris13.fr
Tobias Ekholm	Uppsala University, Sweden tobias.ekholm@math.uu.se	Jessica S Purcell	Monash University jessica.purcell@monash.edu
Mario Eudave-Muñoz	Univ. Nacional Autónoma de México mario@matem.unam.mx	Birgit Richter	Universität Hamburg birgit.richter@uni-hamburg.de
David Futер	Temple University dfuter@temple.edu	Jérôme Scherer	École Polytech. Féd. de Lausanne jerome.scherer@epfl.ch
John Greenlees	University of Warwick john.greenlees@warwick.ac.uk	Vesna Stojanoska	Univ. of Illinois at Urbana-Champaign vesna@illinois.edu
Ian Hambleton	McMaster University ian@math.mcmaster.ca	Zoltán Szabó	Princeton University szabo@math.princeton.edu
Matthew Hedden	Michigan State University mhedden@math.msu.edu	Maggy Tomova	University of Iowa maggy-tomova@uiowa.edu
Hans-Werner Henn	Université Louis Pasteur henn@math.u-strasbg.fr	Nathalie Wahl	University of Copenhagen wahl@math.ku.dk
Daniel Isaksen	Wayne State University isaksen@math.wayne.edu	Chris Wendl	Humboldt-Universität zu Berlin wendl@math.hu-berlin.de
Thomas Koberda	University of Virginia thomas.koberda@virginia.edu	Daniel T Wise	McGill University, Canada daniel.wise@mcgill.ca
Christine Lescop	Université Joseph Fourier lescop@ujf-grenoble.fr		


See inside back cover or msp.org/agt for submission instructions.

The subscription price for 2024 is US \$705/year for the electronic version, and \$1040/year (+\$70, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Algebraic & Geometric Topology is indexed by Mathematical Reviews, Zentralblatt MATH, Current Mathematical Publications and the Science Citation Index.

Algebraic & Geometric Topology (ISSN 1472-2747 printed, 1472-2739 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840.

AGT peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<https://msp.org/>

© 2024 Mathematical Sciences Publishers

ALGEBRAIC & GEOMETRIC TOPOLOGY

Volume 24 Issue 5 (pages 2389–2970) 2024

Formal contact categories	2389
BENJAMIN COOPER	
Comparison of period coordinates and Teichmüller distances	2451
IAN FRANKEL	
Topological Hochschild homology of truncated Brown–Peterson spectra, I	2509
GABRIEL ANGELINI-KNOLL, DOMINIC LEON CULVER and EVA HÖNING	
Points of quantum SL_n coming from quantum snakes	2537
DANIEL C DOUGLAS	
Algebraic generators of the skein algebra of a surface	2571
RAMANUJAN SANTHAROUBANE	
Bundle transfer of L -homology orientation classes for singular spaces	2579
MARKUS BANAGL	
A reduction of the string bracket to the loop product	2619
KATSUHIKO KURIBAYASHI, TAKAHITO NAITO, SHUN WAKATSUKI and TOSHIHIRO YAMAGUCHI	
Asymptotic dimensions of the arc graphs and disk graphs	2655
KOJI FUJIWARA and SAUL SCHLEIMER	
Representation stability for homotopy automorphisms	2673
ERIK LINDELL and BASHAR SALEH	
The strong Haken theorem via sphere complexes	2707
SEBASTIAN HENSEL and JENNIFER SCHULTENS	
What are GT-shadows?	2721
VASILY A DOLGUSHEV, KHANH Q LE and AIDAN A LORENZ	
A simple proof of the Crowell–Murasugi theorem	2779
THOMAS KINDRED	
The Burau representation and shapes of polyhedra	2787
ETHAN DLUGIE	
Turning vector bundles	2807
DIARMUID CROWLEY, CSABA NAGY, BLAKE SIMS and HUIJUN YANG	
Rigidification of cubical quasicategories	2851
PIERRE-LOUIS CURIEN, MURIEL LIVERNET and GABRIEL SAADIA	
Tautological characteristic classes, I	2889
JAN DYMARA and TADEUSZ JANUSZKIEWICZ	
Homotopy types of suspended 4-manifolds	2933
PENGCHENG LI	
The braid indices of the reverse parallel links of alternating knots	2957
YUANAN DIAO and HUGH MORTON	