

AG  
T

*Algebraic & Geometric  
Topology*

Volume 24 (2024)

**Strongly shortcut spaces**

NIMA HODA





# Strongly shortcut spaces

NIMA HODA

We define the strong shortcut property for rough geodesic metric spaces, generalizing the notion of strongly shortcut graphs. We show that the strong shortcut property is a rough similarity invariant. We give several new characterizations of the strong shortcut property, including an asymptotic cone characterization. We use this characterization to prove that asymptotically  $CAT(0)$  spaces are strongly shortcut. We prove that if a group acts metrically properly and coboundedly on a strongly shortcut rough geodesic metric space then it has a strongly shortcut Cayley graph and so is a strongly shortcut group. Thus we show that  $CAT(0)$  groups are strongly shortcut.

To prove these results, we use several intermediate results which we believe may be of independent interest, including what we call the circle tightening lemma and the fine Milnor–Schwarz lemma. The circle tightening lemma describes how one may obtain a quasi-isometric embedding of a circle by performing surgery on a rough Lipschitz map from a circle that sends antipodal pairs of points far enough apart. The fine Milnor–Schwarz lemma is a refinement of the Milnor–Schwarz lemma that gives finer control on the multiplicative constant of the quasi-isometry from a group to a space it acts on.

20F65, 20F67, 51F30

1. Introduction	3291
2. Basic notions and definitions	3296
3. Characterizing the strong shortcut property	3298
4. The circle tightening lemma	3305
5. The fine Milnor–Schwarz lemma	3321
6. Asymptotically $CAT(0)$ spaces	3323
References	3324

## 1 Introduction

The study of interactions between nonpositive curvature and infinite group theory have a long history dating back to the work of Max Dehn on fundamental groups of surfaces in the early 20<sup>th</sup> century. These

ideas have been developed in a variety of directions since that time and have become particularly relevant in recent decades with the emergence of geometric group theory. Various theories of nonpositively curved groups have been developed: small cancellation groups, CAT(0) groups, cubulated groups, systolic groups, quadric groups, etc. However, while the case of *negatively* curved groups has been satisfactorily unified by the seminal work of Gromov [1987] on hyperbolic groups, to this date there is no satisfactory general notion of a nonpositively curved group.

Strongly shortcut graphs were introduced in earlier work [Hoda 2022] as graphs satisfying a weak notion of nonpositive curvature. They were shown to unify a broad family of graphs of interest in geometric group theory and metric graph theory including hyperbolic graphs, standard Cayley graphs of finitely generated Coxeter groups and 1–skeletons of finite-dimensional CAT(0) cube complexes, systolic complexes and quadric complexes [Haettel et al. 2023; Hoda 2022]. They are finitely presented and have polynomial isoperimetric functions and so have decidable word problem [Hoda 2022]. Strongly shortcut groups are defined as those groups admitting a proper and cocompact action on a strongly shortcut graph [Hoda 2022]. They include a wide family of groups satisfying various nonpositive curvature conditions, including hyperbolic groups, Coxeter groups, cocompactly cubulated groups, systolic groups, quadric groups, finitely presented small cancellation groups, Helly groups, hierarchically hyperbolic groups and even the discrete Heisenberg groups [Haettel et al. 2023; Hoda 2022; Le Donne and Paddeu 2023].

A graph  $\Gamma$  is *strongly shortcut* if, for some  $K > 1$ , there is a bound on the lengths of cycles  $\alpha: S \rightarrow \Gamma$  for which  $d_X(\alpha(p), \alpha(\bar{p})) \geq (1/K) \cdot (|S|/2)$  for every antipodal pair of points  $p, \bar{p} \in S$ . By [Hoda 2022], a graph  $\Gamma$  is *strongly shortcut* if and only if, for some  $K > 1$ , there is a bound on the lengths of the  $K$ –bi-Lipschitz embedded cycles of  $\Gamma$ . A result of Papasoglu [1996, page 793] implies that strongly shortcut groups have simply connected asymptotic cones. By another result of Papasoglu [1996, page 805], this implies that strongly shortcut groups have linear isodiametric functions and, by a result of Riley [2003, Theorem C], this implies that strongly shortcut groups have linear filling length functions.

In this paper, we introduce a generalization of this notion to rough geodesic metric spaces. A metric space  $X$  is *R–rough geodesic* if, for every  $x_1, x_2 \in X$ , there exists a function  $f: [0, \ell] \rightarrow X$  such that  $f(0) = x_1$ ,  $f(\ell) = x_2$ ,  $\ell = d(x_1, x_2)$  and

$$|s - t| - R \leq d(f(s), f(t)) \leq |s - t| + R$$

for any  $s$  and  $t$  in the interval  $[0, \ell]$ . This is the same as  $X$  being  $(1, R)$ –quasigeodesic. The special case  $R = 0$  is that of geodesic metric spaces. An  $R$ –rough geodesic metric space  $X$  is *strongly shortcut* if, for some  $K > 1$ , there is a bound on the lengths of Riemannian circles  $S$  for which there exists an  $R$ –rough 1–Lipschitz map  $\alpha: S \rightarrow X$  that satisfies  $d_X(\alpha(p), \alpha(\bar{p})) \geq (1/K) \cdot (|S|/2)$  for every antipodal pair of points  $p, \bar{p} \in S$ . Such a map  $\alpha$  is called a  $1/K$ –*almost isometric R–circle*. We give several characterizations of the strong shortcut property, we show that a group acting metrically properly and coboundedly on a strongly shortcut rough geodesic metric space is a strongly shortcut group and we prove a few other results that may be of independent interest in metric geometry and geometric group

theory. The results of this paper are applied in several upcoming papers [Haettel et al. 2023; Hoda and Krishna M S 2023; Le Donne and Paddeu 2023].

Below is a summary of our main results.

Theorem A gives several conditions that are equivalent to the strong shortcut property for rough geodesic metric spaces. Of particular note is condition (5) which expresses the strong shortcut property in terms of asymptotic cones. Conditions (2) and (3) generalize [Hoda 2022, Proposition 3.5], which expresses the strong shortcut property for graphs in terms of bi-Lipschitz cycles. These two generalizations have their own advantages: for geodesic metric spaces (ie when  $R = 0$ ) condition (2) expresses the strong shortcut property purely in terms of bi-Lipschitz maps from circles, whereas condition (3) avoids the dependence on  $R$ . Condition (4) expresses the strong shortcut property in terms of nonapproximability of certain finite metric spaces at large scale. Conditions (4) and (5) also make sense for general metric spaces and we prove that they are equivalent for general metric spaces (see Theorem 3.6). Thus one may consider condition (4) of Theorem A as a definition of the strong shortcut property for general metric spaces.

**Theorem A** (Theorem 3.8) *Let  $X$  be an  $R$ -rough geodesic metric space. The following conditions are equivalent:*

- (1)  $X$  is strongly shortcut.
- (2) There exists an  $L > 1$  such that there is a bound on the lengths of the  $(L, 4R)$ -quasi-isometric embeddings of Riemannian circles in  $X$ .
- (3) There exists an  $L > 1$  such that for every  $C \geq 0$  there is a bound on the lengths of the  $(L, C)$ -quasi-isometric embeddings of Riemannian circles in  $X$ .
- (4) For some  $L > 1$  and some  $n \in \mathbb{N}$ , there is a bound on the  $\lambda > 0$  for which there exists an  $L$ -bi-Lipschitz embedding of  $\lambda S_n^0$  in  $X$ , where  $S_n^0$  is the vertex set of the cycle graph  $S_n$  of length  $n$  and  $\lambda S_n^0$  is  $S_n^0$  with the metric scaled by  $\lambda$ .
- (5) No asymptotic cone of  $X$  contains an isometric copy of the Riemannian circle of unit length.

The main difficulty in proving Theorem A is in the implication (2)  $\implies$  (1). This is because a  $1/K$ -almost isometric  $R$ -circle in  $X$  does not need to be an  $(L, 4R)$ -quasi-isometric embedding for any  $L > 1$ ; while the almost isometric condition only concerns pairs of antipodal points, the quasi-isometry condition concerns all pairs of points. The idea of the proof is that given a  $1/K$ -almost isometric  $R$ -circle  $\alpha$  with  $K > 1$  sufficiently close to 1, we can perform surgery on  $\alpha$  in order to obtain an  $(L, 4R)$ -quasi-isometric embedding where  $L$  depends on  $K$  in such a way that if  $K \rightarrow 1$  then  $L \rightarrow 1$  also. The contrapositive  $\neg(1) \implies \neg(2)$  then readily follows since any family of arbitrarily long  $R$ -circles with almost isometric constant  $K$  approaching 1 could then be surgered to produce a family of quasi-isometric embeddings with the multiplicative constant  $L$  tending to 1. The circle surgery result, which we call the circle tightening lemma, is stated in slightly simplified form in Theorem G below and expressed more formally in Lemma 4.5.

**Theorem B** (Theorems 3.5 and 3.8) *Let  $\Gamma$  be a graph. Then  $\Gamma$  is strongly shortcut as a graph if and only if  $\Gamma$  is strongly shortcut as a geodesic metric space.*

Theorem C gives several conditions that are equivalent to the strong shortcut property for groups. Condition (3) reduces the property to the existence of a strongly shortcut Cayley graph. The proof is a direct application of the fine Milnor–Schwarz lemma (Theorem H) and stability of the strong shortcut property under scaling and quasi-isometric perturbation of the metric (Theorem F).

**Theorem C** (Corollary 5.5) *Let  $G$  be a group. The following conditions are equivalent.*

- (1)  $G$  is strongly shortcut.
- (2)  $G$  acts metrically properly and coboundedly on a strongly shortcut rough geodesic metric space.
- (3)  $G$  has a finite generating set  $S$  for which the Cayley graph of  $(G, S)$  is strongly shortcut.

Asymptotically CAT(0) spaces and groups were first introduced and studied by Kar [2011]. A metric space  $X$  is *asymptotically* CAT(0) if every asymptotic cone of  $X$  is CAT(0). A group is *asymptotically* CAT(0) if it acts properly and cocompactly on an asymptotically CAT(0) proper geodesic metric space. Examples of asymptotically CAT(0) spaces include CAT(0) spaces, Gromov-hyperbolic spaces and  $\widehat{\text{SL}(2, \mathbb{R})}$  with the Sasaki metric [Kar 2011].

**Theorem D** (Theorems 6.1 and 6.2) *Asymptotically CAT(0) rough geodesic metric spaces are strongly shortcut. Consequently, (asymptotically) CAT(0) groups are strongly shortcut.*

Theorem E shows that the strong shortcut condition is preserved under taking asymptotic cones. This was suggested as a desirable property for a general notion of nonpositive curvature in Gromov [1993, Section 6.E].

**Theorem E** (Corollary 3.9) *Let  $X$  be an  $R$ -rough geodesic metric space. If  $X$  is strongly shortcut then every asymptotic cone of  $X$  is strongly shortcut.*

Theorem F below has several consequences. In addition to showing that the strong shortcut property descends to isometric subspaces and is a rough similarity invariant, it implies that for a given strongly shortcut space, a sufficiently small bi-Lipschitz distortion of the metric preserves the strong shortcut property. This is another property which is discussed in Gromov [1993, Section 6.E].

**Theorem F** (Corollary 3.10) *Let  $X$  be a strongly shortcut rough geodesic metric space. Then there exists an  $L_X > 1$  such that whenever  $Y$  is a rough geodesic metric space and  $f: Y \rightarrow X$  is an  $(L_X, C)$ -quasi-isometric embedding up to scaling, with  $C > 0$ , then  $Y$  is also strongly shortcut. In particular, the strong shortcut property is a rough similarity invariant of rough geodesic metric spaces.*

In fact, Theorem F holds for general metric spaces with condition (4) of Theorem A in place of the strong shortcut property (see Proposition 3.4). It should be noted that the strong shortcut property is not




Figure 1: The closed outer path in black is a 1-Lipschitz embedding  $\alpha$  of a Riemannian circle  $S$ . This embedding  $\alpha$  has poor bi-Lipschitz constant but only because it badly distorts distances between relatively nearby pairs of points of  $S$  (pairs contained in the subpaths  $\alpha|_{Q_i}$ ). If we consider only antipodal pairs of points of  $S$  then the distortion of their distances under  $\alpha$  is much less than in these worst cases. In other words  $\alpha$  has low distortion when viewed *globally*. The circle tightening lemma tells us that if the global distortion of  $\alpha$  is low enough then we can perform surgery on  $\alpha$ , replacing distorted subpaths of arbitrarily low total relative length with efficient alternatives (the red paths in the figure) in order to obtain an arbitrarily good bi-Lipschitz constant.

a quasi-isometry invariant so one cannot hope to remove the dependence on  $X$  of the quasi-isometry constant  $L_X$ . See Section 3.1 for an example.

The following result, which we call the circle tightening lemma, states that a map from a circle that satisfies a rough Lipschitz upper bound and that, on antipodes, satisfies a bi-Lipschitz lower bound can be upgraded through surgery to a rough bi-Lipschitz map. See Figure 1. The circle tightening lemma is essential in the proof of Theorem A. We believe it may be of independent interest. Here we express a slightly simplified version of the circle tightening lemma. For the formal statement, please see Lemma 4.5.

**Theorem G** (circle tightening lemma, Lemma 4.5) *Let  $X$  be an  $R$ -rough geodesic metric space with  $R \geq 0$ , let  $L > 1$  and let  $\varepsilon > 0$ . There exists a  $K > 1$  such that if  $\alpha : S \rightarrow X$  is a sufficiently long  $R$ -rough 1-Lipschitz map from a Riemannian circle  $S$  satisfying*

$$d_X(\alpha(p), \alpha(\bar{p})) \geq \frac{1}{K} d_S(p, \bar{p})$$

*for every antipodal pair  $p, \bar{p} \in S$  then there exists a countable collection  $\{Q_i\}_i$  of pairwise disjoint closed segments in  $S$  of total length  $\sum_i |Q_i| < \varepsilon|S|$  such that shortening the  $Q_i$  and replacing the  $\alpha|_{Q_i} : Q_i \rightarrow X$  we can obtain from  $\alpha$  an  $(L, 4R)$ -quasi-isometric embedding of a circle.*

Note that in the statement of the circle tightening lemma, the rough geodesicity constant  $R$  may be equal to 0 in which case the result is about 1-Lipschitz maps and  $L$ -bi-Lipschitz maps in a geodesic metric space.

We call the following refinement of the Milnor–Schwarz lemma the fine Milnor–Schwarz lemma. It is used in the proof of Theorem C. It essentially says that if a group  $G$  acts metrically properly and coboundedly on a rough geodesic space  $X$  then, up to scaling, the group  $G$  has word metrics that are quasi-isometric to  $X$  with multiplicative constant arbitrarily close to 1. We believe it may be of independent interest.

**Theorem H** (fine Milnor–Schwarz lemma, Lemma 5.2, Remark 5.1) *Let  $(X, d)$  be a rough geodesic metric space. Let  $G$  be a group acting metrically properly and coboundedly on  $X$  by isometries. Fix  $x_0 \in X$ . For  $t > 0$  let  $S_t$  be the finite set defined by*

$$S_t = \{g \in G \mid d(x_0, gx_0) \leq t\}$$

*and consider the word metric  $d_{S_t}$  defined by  $S_t$ . (For those  $t$  where  $S_t$  does not generate  $G$ , we allow  $d_{S_t}$  to take the value  $\infty$ ). Let  $K_t$  be the infimum of all  $K > 1$  for which*

$$(G, td_{S_t}) \rightarrow X, \quad g \mapsto g \cdot x_0,$$

*is a  $(K, C_K)$ –quasi-isometry for some  $C_K \geq 0$ . Then  $K_t \rightarrow 1$  as  $t \rightarrow \infty$ .*

## Structure of the paper

In Section 2 we introduce basic notions that will be used throughout the paper. In Section 3 prove various characterizations of the strong shortcut property and prove that it is a rough similarity invariant. In Section 4 we state and prove the circle tightening lemma. In Section 5 we state and prove the fine Milnor–Schwarz lemma. In Section 6, we apply the results of the previous sections to prove that asymptotically CAT(0) groups are strongly shortcut.

## Acknowledgements

The author would like to thank Pierre Pansu for some valuable discussions about asymptotic cones of strongly shortcut graphs. This work was partially supported by the ERC grant GroIsRan and an NSERC postdoctoral fellowship.

## 2 Basic notions and definitions

Let  $X$  and  $Y$  be metric spaces, let  $S$  be a set and let  $R$  be a nonnegative real. A function  $f: S \rightarrow Y$  is  $R$ –roughly onto if every  $y \in Y$  is at distance at most  $R$  from some point in  $f(S) \subset Y$ . An  $R$ –rough isometric embedding from  $X$  to  $Y$  is a function  $f: X \rightarrow Y$  such that

$$d(x_1, x_2) - R \leq d(f(x_1), f(x_2)) \leq d(x_1, x_2) + R$$

for all  $x_1, x_2 \in X$ . An  $R$ –rough isometric embedding is the same as a  $(1, R)$ –quasi-isometric embedding. An  $R$ –rough isometric embedding  $f: X \rightarrow Y$  is an  $R$ –rough isometry if it is roughly onto. An  $R$ –rough isometry is the same as a  $(1, R)$ –quasi-isometry.

An  $R$ -rough geodesic in  $X$  from  $x_1$  to  $x_2$  is an  $R$ -rough isometric embedding  $f$  from the interval  $[0, \ell] \subset \mathbb{R}$  to  $X$  with  $\ell = d(x_1, x_2)$  such that  $f(0) = x_1$  and  $f(\ell) = x_2$ . An  $R$ -rough geodesic is the same as a  $(1, R)$ -quasigeodesic. A metric space  $(X, d)$  is  $R$ -rough geodesic if every pair of points in  $X$  is joined by an  $R$ -rough geodesic.<sup>1</sup> Note that rough geodesicity implies weak geodesicity, as used by Kasparov and Skandalis [1994; 2003] and others [Lafforgue 2002; Mineyev and Yu 2002]. A natural question is whether or not every rough geodesic space can be thickened, in the sense of Gromov [1993, Section 1.B] to a geodesic metric space.

Let  $X$  and  $Y$  be metric spaces, let  $R \geq 0$  and let  $K \geq 1$ . An  $R$ -rough  $K$ -Lipschitz map from  $Y$  to  $X$  is a function  $\alpha: Y \rightarrow X$  such that

$$d(\alpha(p), \alpha(q)) \leq Kd(p, q) + R$$

for all  $p, q \in Y$ . An  $R$ -path in  $X$  is an  $R$ -rough 1-Lipschitz map  $\alpha: P \rightarrow X$  from an interval  $P \subset \mathbb{R}$ . An  $R$ -circle in  $X$  is an  $R$ -rough 1-Lipschitz map  $\alpha: S \rightarrow X$  from a Riemannian circle  $S$ . We use the notation  $|F|$  to denote the length of  $F$ , where  $F$  is an interval, a Riemannian circle or a finite union of closed segments in an interval or in a Riemannian circle.

**Remark 2.1** The concatenation of two  $R$ -paths need not be an  $R$ -path. However, if  $\alpha_1: P_1 \rightarrow X$  and  $\alpha_2: P_2 \rightarrow X$  are a pair of concatenable  $R$ -paths and  $\gamma: [0, R] \rightarrow X$  is the constant path at the point of concatenation then the concatenation  $\alpha_1\gamma\alpha_2$  is an  $R$ -path.

An  $R$ -circle  $\alpha: S \rightarrow X$  is  $1/K$ -almost isometric, for some  $K > 1$ , if

$$d(\alpha(p), \alpha(\bar{p})) \geq \frac{1}{K} \cdot \frac{|S|}{2}$$

for every antipodal pair of points  $p, \bar{p} \in S$ .

**Definition 2.2** An  $R$ -rough geodesic metric space  $X$  is *strongly shortcut* if, for some  $K > 1$ , there is a bound on the lengths of the  $1/K$ -almost isometric  $R$ -cycles of  $X$ .

**Remark 2.3** By Theorem 3.8, the apparent dependence on  $R$  in Definition 2.2 is not essential. That is, if  $X$  is an  $R$ -rough geodesic metric space and  $R' > R$  then  $X$  is strongly shortcut if and only if it is strongly shortcut when viewed as an  $R'$ -rough geodesic metric space.

We view graphs as geodesic metric spaces with each edge isometric to a unit interval. For a graph  $\Gamma$ , we use the notation  $\Gamma^0$  to denote the vertex set of  $\Gamma$  with its subspace metric. The *cycle graph*  $S_n$  of length  $n$  is the graph isometric to a Riemannian circle of length  $n$ . A *cycle* in a graph  $\Gamma$  is a combinatorial

<sup>1</sup>We include the condition  $\ell = d(x_1, x_2)$  in the definition of an  $R$ -rough geodesic  $f$  only for convenience. If we do not assume it, then we can recover it up to slightly increasing the rough geodesicity constant to  $R' = (1 + \sqrt{2})R$ . Indeed, the remaining conditions on  $f$  imply that  $|\ell - d(x_1, x_2)| \leq R$  and it can be shown that either  $d(x_1, x_2) \leq R'$  (in which case any function  $[0, d(x_1, x_2)] \rightarrow \{x_1, x_2\}$  is an  $R'$ -rough isometric embedding) or the composition of  $f$  with the orientation preserving linear bijection  $[0, d(x_1, x_2)] \rightarrow [0, \ell]$  results in an  $R'$ -rough isometric embedding.

map  $S_n \rightarrow \Gamma$  from some cycle graph  $S_n$  to  $\Gamma$ . A *path graph* is a graph isometric to a real interval. A *combinatorial path* in a graph  $\Gamma$  is a combinatorial map  $P \rightarrow \Gamma$  from a path graph  $P$ .

Note that if  $\alpha: S_n \rightarrow \Gamma$  is a cycle in a graph  $\Gamma$  then  $\alpha$  is a 1-Lipschitz map from a Riemannian circle to a geodesic metric space or, in the language we have established above, an  $R$ -circle in an  $R$ -rough geodesic metric space where  $R = 0$ .

**Definition 2.4** A graph  $\Gamma$  is *strongly shortcut as a graph* if, for some  $K > 1$ , there is a bound on the lengths of the  $1/K$ -almost isometric cycles of  $\Gamma$ .

**Remark 2.5** By Theorems 3.5 and 3.8 and Corollary 3.10, the following conditions are equivalent for a graph  $\Gamma$ .

- (1)  $\Gamma$  is strongly shortcut as a graph.
- (2)  $\Gamma$  is strongly shortcut as a geodesic metric space.
- (3)  $\Gamma^0$  is strongly shortcut as a rough geodesic metric space.

If  $X$  is a metric space and  $\lambda > 0$  then we write  $\lambda X$  to denote the metric space obtained from  $X$  by scaling the metric by  $\lambda$ .

### 3 Characterizing the strong shortcut property

In this section we will give various characterizations of the strong shortcut property.

**Lemma 3.1** Let  $\alpha: S \rightarrow X$  be a  $1/K$ -almost isometric  $R$ -circle in a metric space  $X$ . Then

$$d(\alpha(p), \alpha(q)) \geq d(p, q) - \frac{K-1}{K} \cdot \frac{|S|}{2} - 2R$$

for all  $p, q \in S$ .

**Proof** Let  $p', q' \in S \setminus \{p, q\}$  be antipodal and suppose that a geodesic segment of  $S$  visits  $p', p, q$  and  $q'$ , in that order. Then

$$\begin{aligned} \frac{1}{K} \cdot \frac{|S|}{2} &\leq d(\alpha(p'), \alpha(q')) \\ &\leq d(\alpha(p'), \alpha(p)) + d(\alpha(p), \alpha(q)) + d(\alpha(q), \alpha(q')) \\ &\leq d(p', p) + R + d(\alpha(p), \alpha(q)) + d(q, q') + R \\ &= d(p', p) + d(q, q') + d(\alpha(p), \alpha(q)) + 2R \\ &= \frac{|S|}{2} - d(p, q) + d(\alpha(p), \alpha(q)) + 2R, \end{aligned}$$

from which we can obtain the desired inequality. □

The following definition is very useful because it applies to general metric spaces.

**Definition 3.2** A metric space  $X$  approximates  $n$ -gons if for every  $K > 1$  and every  $n \in \mathbb{N}$  there exist  $K$ -bi-Lipschitz embeddings of  $\lambda S_n^0$  in  $X$  for arbitrarily large  $\lambda > 0$ .

We will see that in the case of a graph or a rough geodesic metric space nonapproximation of  $n$ -gons is equivalent to the strong shortcut property. Thus it would make sense to define the strong shortcut property for general metric spaces as nonapproximability of  $n$ -gons.

**Definition 3.3** Let  $X$  and  $Y$  be metric spaces. A function  $f : Y \rightarrow X$  is a  $(K, C)$ -quasi-isometry up to scaling if there exists a  $\lambda > 0$  such that  $f$  is a  $(K, C)$ -quasi-isometry when viewed as a function from  $\lambda Y$  to  $X$ . A function  $f : Y \rightarrow X$  is a rough similarity if, for some  $C > 0$ , the function  $f$  is a  $(1, C)$ -quasi-isometry up to scaling. A property  $\mathcal{P}$  of metric spaces is a rough similarity invariant if whenever  $X$  satisfies  $\mathcal{P}$  and  $f : Y \rightarrow X$  is a rough similarity then  $Y$  also satisfies  $\mathcal{P}$ . A property  $\mathcal{P}$  of metric spaces is a rough approximability invariant if, for any metric space  $X$  satisfying  $\mathcal{P}$ , there exists an  $L_X > 1$  such that whenever  $C > 0$  and  $f : Y \rightarrow X$  is an  $(L_X, C)$ -quasi-isometric embedding up to scaling, then  $Y$  also satisfies  $\mathcal{P}$ .

**Proposition 3.4** Nonapproximability of  $n$ -gons is a rough approximability invariant of metric spaces. In particular, nonapproximability of  $n$ -gons is a rough similarity invariant of metric spaces.

**Proof** Let  $X$  be a metric space that does not approximate  $n$ -gons. Then there is a  $K > 1$ , an  $n \in \mathbb{N}$  and a  $\Lambda > 0$  such that any  $K$ -bi-Lipschitz embedding of  $\lambda S_n^0$  in  $X$  satisfies  $\lambda < \Lambda$ .

Let  $Y$  be a metric space, let  $t > 0$ , let  $L \in (1, K)$  and let  $f : tY \rightarrow X$  be an  $(L, C)$ -quasi-isometric embedding. Let  $K' \in (1, K/L)$  and let  $\lambda' > CLK'/t$ . We will show that there is a bound on the  $\lambda'$  for which there exists a  $K'$ -bi-Lipschitz embedding  $\alpha : \lambda' S_n^0 \rightarrow Y$ . Viewing such an  $\alpha$  as map from  $t\lambda' S_n^0$  to  $tY$  the composition  $f \circ \alpha : t\lambda' S_n^0 \rightarrow X$  is an  $(LK', C)$ -quasi-isometric embedding. But the minimum distance between distinct points in  $t\lambda' S_n^0$  is  $t\lambda'$  and so one can show that  $f \circ \alpha$  is a  $(t\lambda' LK' + C)/(t\lambda' - LK'C)$ -bi-Lipschitz embedding from  $t\lambda' S_n^0$ . But

$$\frac{t\lambda' LK' + C}{t\lambda' - LK'C} \rightarrow LK' < K$$

as  $\lambda' \rightarrow \infty$  so there is a  $\Lambda_0$  such that if  $\lambda' \geq \Lambda_0$  then  $(t\lambda' LK' + C)/(t\lambda' - LK'C) < K$ . So if we had  $\lambda' \geq \Lambda' = \max\{\Lambda_0, \Lambda/t\}$  then  $f \circ \alpha$  would be a  $K$ -bi-Lipschitz embedding of  $\lambda S_n^0$  in  $X$  with  $\lambda = t\lambda' \geq \Lambda$ , which would be a contradiction. Thus  $\Lambda'$  bounds the  $\lambda'$  for which there exists a  $K'$ -bi-Lipschitz embedding  $\alpha : \lambda' S_n^0 \rightarrow Y$ , as required.  $\square$

**Theorem 3.5** Let  $\Gamma$  be a graph. Then the following conditions are equivalent.

- (1)  $\Gamma$  is not strongly shortcut as a graph.
- (2)  $\Gamma$  approximates  $n$ -gons.

**Proof** (1)  $\implies$  (2) Let  $K' > 1$  and let  $\alpha : S_{n'} \rightarrow \Gamma$  be a  $1/K'$ -almost isometric cycle. Let  $n \in \mathbb{N}$  and subdivide  $S_{n'}$  into  $n$  segments of equal length, ignoring the original graph structure on  $S_{n'}$ . Let  $Y \subset S_{n'}$  be

the set of endpoints of the segments. Then  $Y$  is isometric to  $\lambda S_n$  for  $\lambda = n'/n$ . Let  $\alpha'$  be the composition of the inclusion  $Y \hookrightarrow S_{n'}$  with  $\alpha$ . Let  $p, q \in Y$  be distinct. Then  $d(p, q) \geq |S_{n'}|/n$  and, by Lemma 3.1,

$$d(\alpha'(p), \alpha'(q)) \geq d(p, q) - \frac{K' - 1}{K'} \cdot \frac{|S_{n'}|}{2} \geq d(p, q) - \frac{K' - 1}{K'} \cdot \frac{nd(p, q)}{2} = \left(1 - \frac{n(K' - 1)}{2K'}\right)d(p, q),$$

but  $d(\alpha'(p), \alpha'(q)) \leq d(p, q)$  so, when  $K'$  is small enough that  $n(K' - 1)/2K' < 1$ , the map  $\alpha'$  is  $K$ -bi-Lipschitz for

$$K = \left(1 - \frac{n(K' - 1)}{2K'}\right)^{-1}.$$

Thus, given an arbitrary  $n \in \mathbb{N}$  and an  $\alpha'$  as above with  $K'$  small enough, we can obtain a  $K$ -bi-Lipschitz embedding of  $\lambda S_n$  in  $\Gamma$  with

$$K = \left(1 - \frac{n(K' - 1)}{2K'}\right)^{-1}, \quad \lambda = \frac{n'}{n}.$$

Since  $\Gamma$  is not strongly shortcut, there exist  $\alpha$  as above with  $K' > 1$  arbitrarily close to 1 and with  $n'$  arbitrarily large. But  $K \rightarrow 1$  as  $K' \rightarrow 1$  and  $\lambda \rightarrow \infty$  as  $n' \rightarrow \infty$  so we have  $K$ -bi-Lipschitz embeddings of  $\lambda S_n$  with  $K$  arbitrarily close to 1 and  $\lambda$  arbitrarily large.

(2)  $\implies$  (1) Let  $n \in \mathbb{N}$ , let  $K > 1$ , let  $\lambda > K$  and let  $\alpha: \lambda S_n^0 \rightarrow \Gamma$  be a  $K$ -bi-Lipschitz embedding. There is a retraction  $r: \Gamma \rightarrow \Gamma^0$  such that  $r$  is a  $(1, 1)$ -quasi-isometry. Then the composition  $r \circ \alpha$  is a  $(K, 1)$ -quasi-isometric embedding. But distinct points in  $\lambda S_n^0$  are at distance at least  $\lambda$  and, since  $K < \lambda$ , this implies that  $r \circ \alpha$  is  $L$ -bi-Lipschitz, where  $L = (K\lambda + 1)/(\lambda - K)$ . View  $S_n$  as the Cayley graph of  $\mathbb{Z}/n\mathbb{Z}$  with generating set  $\{1\}$  and, for  $i \in \mathbb{Z}/n\mathbb{Z}$ , let  $v_i$  be the vertex of  $S_n$  corresponding to  $i$ . Then, for each  $i$ , we have  $d(r \circ \alpha(v_i), r \circ \alpha(v_{i+1})) \leq \lfloor L\lambda \rfloor$  so there is a combinatorial path  $\gamma_i: P_i \rightarrow \Gamma$  of length  $m_i \in \{\lfloor L\lambda \rfloor - 1, \lfloor L\lambda \rfloor\}$  from  $r \circ \alpha(v_i)$  to  $r \circ \alpha(v_{i+1})$ . For each  $i$ , identify the endpoint of  $P_i$  with the initial point of  $P_{i+1}$  to obtain a cycle  $\gamma: S_m \rightarrow \Gamma$  with  $m = \sum_{i=1}^n m_i$ . Then  $\alpha$  factors through  $\gamma$  via the embedding that sends  $v_i$  to the initial point of  $P_i \subset S_m$ . So, viewing  $S_n^0$  as a subset of  $S_m^0$  via this embedding, we have  $r \circ \alpha(v_i) = \gamma(v_i)$ , for each  $i$ . Let  $x \in S_m$  and let  $v_i$  minimize  $d(x, v_i)$ . Then  $d(x, v_i) \leq \frac{1}{2}\lfloor L\lambda \rfloor$  and if  $\bar{x}$  is the antipode of  $x$  and  $\bar{i} = i + \lfloor \frac{n}{2} \rfloor$  then  $d(\bar{x}, v_{\bar{i}}) \leq \lfloor L\lambda \rfloor + \frac{n}{2}$ , so

$$\begin{aligned} d(\gamma(x), \gamma(\bar{x})) &\geq d(\gamma(v_i), \gamma(v_{\bar{i}})) - d(\gamma(x), \gamma(v_i)) - d(\gamma(\bar{x}), \gamma(v_{\bar{i}})) \\ &\geq d(\gamma(v_i), \gamma(v_{\bar{i}})) - d(x, v_i) - d(\bar{x}, v_{\bar{i}}) \\ &\geq d(\gamma(v_i), \gamma(v_{\bar{i}})) - \frac{1}{2}\lfloor L\lambda \rfloor - \lfloor L\lambda \rfloor - \frac{1}{2}n \\ &= d(r \circ \alpha(v_i), r \circ \alpha(v_{\bar{i}})) - \frac{3}{2}\lfloor L\lambda \rfloor - \frac{1}{2}n \\ &\geq \frac{1}{L}d_{\lambda S_n^0}(v_i, v_{\bar{i}}) - \frac{3}{2}\lfloor L\lambda \rfloor - \frac{1}{2}n \\ &= \frac{\lambda}{L}\lfloor \frac{1}{2}n \rfloor - \frac{3}{2}\lfloor L\lambda \rfloor - \frac{1}{2}n \\ &\geq \frac{\lambda}{L}\left(\frac{1}{2}(n - 1)\right) - \frac{3}{2}L\lambda - \frac{1}{2}n = \frac{1}{m}\left(\frac{\lambda(n - 1)}{L} - 3L\lambda - n\right) \cdot \frac{1}{2}|S_m|. \end{aligned}$$

Since  $m \leq nL\lambda$ , the above computation implies that

$$d(\gamma(x), \gamma(\bar{x})) \geq \left( \frac{n-1}{nL^2} - \frac{3}{n} - \frac{1}{L\lambda} \right) \frac{|S_m|}{2}$$

so, given  $\alpha$  as above, we can obtain a  $1/K'$ -almost isometric cycle in  $\Gamma$  of length  $m$ , where

$$\frac{1}{K'} = \left( \frac{n-1}{nL^2} - \frac{3}{n} - \frac{1}{L\lambda} \right).$$

We need only show there exist  $\alpha$  for which  $1/K'$  is arbitrarily close to 1 and  $m$  is arbitrarily large. By hypothesis, there exist  $\alpha$  for which  $K = (n+1)/n$  and  $\lambda > n$ , for arbitrary  $n \in \mathbb{N}$ . But then, as  $n \rightarrow \infty$ , we have  $m \geq n(\lfloor L\lambda \rfloor - 1) \rightarrow \infty$  and  $L = (K\lambda + 1)/(\lambda - K) \rightarrow K$  so

$$\frac{1}{K'} = \left( \frac{n-1}{nL^2} - \frac{3}{n} - \frac{1}{L\lambda} \right) \rightarrow 1. \quad \square$$

Let  $X$  be a metric space. Let  $\mathcal{U}$  be a nonprincipal ultrafilter on  $\mathbb{N}$ . Let  $(b^{(m)})_{m \in \mathbb{N}}$  be a sequence in  $X$ . Let  $(s^{(m)})_{m \in \mathbb{N}}$  be a sequence of positive reals such that  $s^{(m)} \rightarrow \infty$  as  $m \rightarrow \infty$ . Consider the set

$$\mathfrak{X}' = \left\{ (x_m)_{m \in \mathbb{N}} \mid \left( \frac{d(x_m, b^{(m)})}{s^{(m)}} \right)_{m \in \mathbb{N}} \text{ is bounded} \right\}$$

of sequences in  $X$  that are bounded with respect to the basepoint sequence  $(b^{(m)})_m$  and the scaling sequence  $(s^{(m)})_m$ . For  $(x_m)_m, (x'_m)_m \in \mathfrak{X}'$ ,

$$\bar{d}((x_m)_m, (x'_m)_m) = \lim_{\mathcal{U}} \frac{d(x_m, x'_m)}{s^{(m)}}$$

defines a pseudometric on  $\mathfrak{X}'$ . The asymptotic cone  $\mathfrak{X}$  of  $X$  with respect to the nonprincipal ultrafilter  $\mathcal{U}$ , the basepoint sequence  $(b^{(m)})_m$  and the scaling sequence  $(s^{(m)})_m$  is the metric space obtained from  $\mathfrak{X}'$  and  $\bar{d}$  by identifying  $(x_m)_m$  and  $(x'_m)_m$  whenever  $\bar{d}((x_m)_m, (x'_m)_m) = 0$ .

Note that Theorem 3.6 and Corollary 3.7 apply to general metric spaces and not just rough geodesic metric spaces.

**Theorem 3.6** *Let  $X$  be a metric space. Then the following conditions are equivalent.*

- (1) *There is an asymptotic cone of  $X$  that contains an isometric copy of the Riemannian circle of unit length.*
- (2)  *$X$  approximates  $n$ -gons.*

**Proof** (1)  $\implies$  (2) Suppose  $S \subset \mathcal{X}$  is a subspace isometric to the Riemannian circle of unit length in the asymptotic cone  $\mathcal{X}$  of  $X$  with respect to a nonprincipal ultrafilter  $\mathcal{U}$ , a basepoint sequence  $(b^{(m)})_m$  and a scaling sequence  $(s^{(m)})_m$ . Take any  $n \in \mathbb{N}$ , any  $K > 1$  and any  $\Lambda > 0$ . We will construct a  $K$ -bi-Lipschitz map  $\alpha: \lambda S_n^0 \rightarrow X$  with  $\lambda \geq \Lambda$ . Subdivide  $S$  into  $n$  segments of equal length and let  $S^0$  denote the set of endpoints of the segments. For each  $\varepsilon > 0$  and each  $p, q \in S^0$  represented by  $(p_m)_m$  and  $(q_m)_m$ , there is an  $A_\varepsilon^{p,q} \in \mathcal{U}$  such that

$$d(p, q) - \varepsilon \leq \frac{d(p_m, q_m)}{s^{(m)}} \leq d(p, q) + \varepsilon$$

for all  $m \in A_\varepsilon^{p,q}$ . There are finitely many pairs  $p, q \in S^0$  so  $A_\varepsilon = \bigcap_{p,q} A_\varepsilon^{p,q} \in \mathcal{U}$ . Then, for any distinct  $p, q \in S^0$ ,

$$\left(1 - \frac{\varepsilon}{d(p,q)}\right) s^{(m)} d(p,q) \leq d(p_m, q_m) \leq \left(1 + \frac{\varepsilon}{d(p,q)}\right) s^{(m)} d(p,q)$$

for all  $m \in A_\varepsilon$ . But  $d(p, q) \geq \frac{1}{n}$  and so

$$(1 - n\varepsilon) s^{(m)} d(p, q) \leq d(p_m, q_m) \leq (1 + n\varepsilon) s^{(m)} d(p, q)$$

for all  $m \in A_\varepsilon$ . So if  $n\varepsilon < 1$  then, for  $m \in A_\varepsilon$ , the map

$$\alpha_m : s^{(m)} S^0 \rightarrow X, \quad p \mapsto p_m,$$

is bi-Lipschitz with bi-Lipschitz constant

$$\max\left\{1 + n\varepsilon, \frac{1}{1 - n\varepsilon}\right\} = \frac{1}{1 - n\varepsilon}.$$

The space  $s^{(m)} S^0$  is isometric to  $(s^{(m)}/n) S_n^0$  so if we chose  $\varepsilon$  small enough that  $n\varepsilon < 1$  and  $1/(1 - n\varepsilon) < K$  and we take  $m \in A_\varepsilon$  large enough that  $s^{(m)}/n \geq \Lambda$  then we can take  $\alpha = \alpha_m$ .

(2)  $\implies$  (1) For  $m \in \mathbb{N}$ , there exists a  $(m + 1)/m$ -bi-Lipschitz map  $\alpha_m : \lambda_m S_{2^m}^0 \rightarrow X$  with  $\lambda_m \geq m$ . Metrize the group  $(1/2^m)\mathbb{Z} = \{k/2^m \mid k \in \mathbb{Z}\} \subset \mathbb{R}$  with the subspace metric and metrize the quotient group  $(1/2^m)\mathbb{Z}/\mathbb{Z}$  with the quotient metric. Then  $(1/2^m)S_{2^m}^0$  is isometric to  $(1/2^m)\mathbb{Z}/\mathbb{Z}$ . Via this isometry we identify the vertex set  $S_{2^m}^0$  with the elements of  $(1/2^m)\mathbb{Z}/\mathbb{Z}$ . Thus we view  $(1/2^m)S_{2^m}^0$  as a metric subspace of the Riemannian circle of unit length  $S = \mathbb{R}/\mathbb{Z}$ . By this identification, the union  $S_{\mathbb{D}} = \bigcup_{m \in \mathbb{N}} S_{2^m}^0 \subset S$  is the *dyadic circle*  $\mathbb{Z}[\frac{1}{2}]/\mathbb{Z}$ . The dyadic circle  $S_{\mathbb{D}}$  is dense in  $S$ . Thus, since asymptotic cones are complete metric spaces [Dru\u0219u and Kapovich 2018, Proposition 10.70], it will suffice to isometrically embed  $S_{\mathbb{D}}$  into an asymptotic cone of  $X$ .

View  $\alpha_m$  as an  $(m + 1)/m$ -bi-Lipschitz map from  $(1/2^m)S_{2^m}^0$  to  $(1/\lambda_m 2^m)X$ . Set  $b^{(m)} = \alpha_m(0)$  and set  $s^{(m)} = \lambda_m 2^m$ . Every nonzero element of  $S_{\mathbb{D}}$  can be uniquely represented as  $k/2^\ell$  with  $k$  odd and satisfying  $0 \leq k < 2^\ell$ . For any such representation  $k/2^\ell$  and any  $m \in \mathbb{N}$ , set

$$x_{k/2^\ell}^{(m)} = \begin{cases} b^{(m)} & \text{if } m < \ell, \\ \alpha_m(k/2^\ell) & \text{if } m \geq \ell, \end{cases}$$

and set  $x_0^{(m)} = b^{(m)}$ . Then, for any nonprincipal ultrafilter  $\mathcal{U}$ , the expression  $p \mapsto (x_p^{(m)})_m$  defines an isometric embedding of  $S_{\mathbb{D}}$  into the asymptotic cone  $\mathcal{X}$  of  $X$  with respect to  $\mathcal{U}$ , the basepoint sequence  $(b^{(m)})_m$  and the scaling sequence  $S^{(m)}$ . Indeed, for every  $p, q \in S_{\mathbb{D}}$ ,

$$\frac{d(x_p^{(m)}, x_q^{(m)})}{s^{(m)}} = \frac{d(\alpha_m(p), \alpha_m(q))}{\lambda_m 2^m} \leq \frac{(m + 1)/m \cdot \lambda_m d_{S_{2^m}}(p, q)}{\lambda_m 2^m} = \frac{m + 1}{m} \cdot d_{S_{\mathbb{D}}}(p, q)$$

and

$$\frac{d(x_p^{(m)}, x_q^{(m)})}{s^{(m)}} = \frac{d(\alpha_m(p), \alpha_m(q))}{\lambda_m 2^m} \geq \frac{m/(m + 1) \cdot \lambda_m d_{S_{2^m}}(p, q)}{\lambda_m 2^m} = \frac{m}{m + 1} \cdot d_{S_{\mathbb{D}}}(p, q)$$

whenever  $m$  is large enough. □

**Corollary 3.7** *Let  $X$  be a metric space and let  $\mathfrak{X}$  be an asymptotic cone of  $X$ . Suppose that  $X$  does not approximate  $n$ -gons. Then  $\mathfrak{X}$  does not approximate  $n$ -gons.*

**Proof** By Theorem 3.6, it suffices to show that any asymptotic cone  $\mathfrak{X}'$  of  $\mathfrak{X}$  does not contain an isometric copy of a Riemannian circle of unit length. But  $\mathfrak{X}'$  is isometric to an asymptotic cone of  $X$  [Druţu and Kapovich 2018, Corollary 10.80] so does not contain an isometric copy of a Riemannian circle of unit length by Theorem 3.6.  $\square$

**Theorem 3.8** *Let  $X$  be an  $R$ -rough geodesic metric space. The following conditions are equivalent.*

- (1)  $X$  is not strongly shortcut.
- (2) For every  $L > 1$  there exist  $(L, 4R)$ -quasi-isometric embeddings of arbitrarily long Riemannian circles in  $X$ .
- (3) For every  $L > 1$  there is a  $C \geq 0$  such that there exist  $(L, C)$ -quasi-isometric embeddings of arbitrarily long Riemannian circles in  $X$ .
- (4)  $X$  approximates  $n$ -gons.
- (5) There is an asymptotic cone of  $X$  that contains an isometric copy of the Riemannian circle of unit length.

**Proof** Conditions (4) and (5) are equivalent for general metric spaces, by Theorem 3.6. So it will suffice to prove the equivalence of conditions (1), (2), (3) and (4).

(1)  $\implies$  (2) Let  $N = 2$ , let  $L > 1$  be arbitrary, let  $K > 1$  be small enough to satisfy Lemma 4.5 and let  $\alpha: S \rightarrow X$  be a  $1/K$ -almost isometric  $R$ -circle in  $X$  with  $|S|$  arbitrarily larger than the  $M$  from Lemma 4.5. Then the limit  $R$ -circle  $\alpha_\infty: S_\infty \rightarrow X$  given by Lemma 4.5 is an  $(L, 4R)$ -quasi-isometric embedding of a Riemannian circle of length at least  $\frac{1}{2}|S|$ .

(2)  $\implies$  (3) This is immediate.

(3)  $\implies$  (4) Let  $\alpha: S \rightarrow X$  be an  $(L, C)$ -quasi-isometric embedding of a Riemannian circle. Let  $n \in \mathbb{N}$ , subdivide  $S$  into  $n$  segments of equal length and let  $Y$  be the set of endpoints of the segments. Then  $Y$  is isometric to  $(|S|/n)S_n$  and

$$\left(\frac{1}{L} - \frac{nC}{|S|}\right)d(p, q) \leq d(\alpha(p), \alpha(q)) \leq \left(L + \frac{nC}{|S|}\right)d(p, q)$$

for distinct  $p, q \in Y$ . By hypothesis, there exist arbitrarily long  $\alpha$  with  $L$  arbitrarily close to 1 and so  $\alpha|_Y$  is a  $K$ -bi-Lipschitz embedding of  $\lambda S_n$  for  $\lambda = |S|/n$  arbitrarily large and

$$K = \max\left\{\left(\frac{1}{L} - \frac{nC}{|S|}\right)^{-1}, L + \frac{nC}{|S|}\right\}$$

arbitrarily close to 1.

(4)  $\implies$  (1) Let  $n \in \mathbb{N}$ , let  $L > 1$ , let  $\lambda > 0$  and let  $\alpha: \lambda S_n^0 \rightarrow X$  be an  $L$ -bi-Lipschitz embedding. View  $S_n$  as the Cayley graph of  $\mathbb{Z}/n\mathbb{Z}$  with generating set  $\{1\}$  and, for  $i \in \mathbb{Z}/n\mathbb{Z}$ , let  $v_i$  be the vertex of  $S_n$  corresponding to  $i$ . Then, for each  $i$ , we have  $d(\alpha(v_i), \alpha(v_{i+1})) \leq L\lambda$  so, by scaling an  $R$ -rough

geodesic, there is an  $R$ -path  $\gamma'_i: P_i \rightarrow X$  of length  $|P_i| = L\lambda$  from  $\alpha(v_i)$  to  $\alpha(v_{i+1})$ . Let  $\gamma_i$  be the concatenation  $c_i\gamma'_i$  where  $c_i: [0, R] \rightarrow X$  is the constant path of length  $R$  at  $\alpha(v_i)$ . For each  $i$ , identify the endpoint of  $P_i$  with the initial point of  $P_{i+1}$  to obtain an  $R$ -circle  $\gamma: S \rightarrow X$  with  $|S| = n(L\lambda + R)$ . Then  $\alpha$  factors through  $\gamma$  via the embedding that sends  $v_i$  to the initial point of  $P_i \subset S$ . So, viewing  $S_n^0$  as a subset of  $S$  via this embedding, we have  $\alpha(v_i) = \gamma(v_i)$  for each  $i$ . Let  $x \in S$  and let  $v_i$  minimize  $d(x, v_i)$ . Then  $d(x, v_i) \leq \frac{1}{2}(L\lambda + R)$  and if  $\bar{x}$  is the antipode of  $x$  and  $\bar{i} = i + \lfloor \frac{n}{2} \rfloor$  then  $d(\bar{x}, v_{\bar{i}}) \leq L\lambda + R$ , so

$$\begin{aligned} d(\gamma(x), \gamma(\bar{x})) &\geq d(\gamma(v_i), \gamma(v_{\bar{i}})) - d(\gamma(x), \gamma(v_i)) - d(\gamma(\bar{x}), \gamma(v_{\bar{i}})) \\ &\geq d(\gamma(v_i), \gamma(v_{\bar{i}})) - d(x, v_i) - R - d(\bar{x}, v_{\bar{i}}) - R \\ &\geq d(\gamma(v_i), \gamma(v_{\bar{i}})) - \frac{1}{2}(L\lambda + R) - (L\lambda + R) - 2R \\ &= d(\alpha(v_i), \alpha(v_{\bar{i}})) - \frac{1}{2}(3L\lambda + 7R) \\ &\geq \frac{1}{L}d_{\lambda S_n^0}(v_i, v_{\bar{i}}) - \frac{1}{2}(3L\lambda + 7R) \\ &= \frac{\lambda}{L} \lfloor \frac{1}{2}n \rfloor - \frac{1}{2}(3L\lambda + 7R) \\ &\geq \frac{\lambda}{L}(\frac{1}{2}(n-1)) - \frac{1}{2}(3L\lambda + 7R) \\ &= \frac{1}{|S|} \left( \frac{\lambda(n-1)}{L} - 3L\lambda - 7R \right) \cdot \frac{1}{2}|S| \\ &= \frac{1}{n(L\lambda + R)} \left( \frac{\lambda(n-1)}{L} - 3L\lambda - 7R \right) \cdot \frac{1}{2}|S| \\ &= \left( \frac{\lambda n - \lambda}{L^2\lambda n + nLR} - \frac{3L\lambda}{L\lambda n + nR} - \frac{7R}{L\lambda n + nR} \right) \cdot \frac{1}{2}|S|. \end{aligned}$$

So, given  $\alpha$  as above, we can obtain a  $1/K$ -almost isometric  $R$ -circle in  $X$  of length  $|S| = n(L\lambda + R)$ , where

$$\frac{1}{K} = \left( \frac{\lambda n - \lambda}{L^2\lambda n + nLR} - \frac{3L\lambda}{L\lambda n + nR} - \frac{7R}{L\lambda n + nR} \right).$$

We need only show there exist  $\alpha$  for which  $1/K$  is arbitrarily close to 1 and  $|S|$  is arbitrarily large. By hypothesis, there exist  $\alpha$  for which  $L = (n + 1)/n$  and  $\lambda > n$ , for arbitrary  $n \in \mathbb{N}$ . But then, as  $n \rightarrow \infty$ , we have  $|S| = n(L\lambda + R) \rightarrow \infty$  and

$$\frac{1}{K} = \left( \frac{\lambda n - \lambda}{L^2\lambda n + nLR} - \frac{3L\lambda}{L\lambda n + nR} - \frac{7R}{L\lambda n + nR} \right) \rightarrow 1. \quad \square$$

**Corollary 3.9** *Let  $X$  be a metric space. If  $X$  is strongly shortcut then every asymptotic cone of  $X$  is strongly shortcut.*

**Proof** This follows immediately from Theorem 3.8 and Corollary 3.7. □

**Corollary 3.10** *The strong shortcut property is a rough approximability invariant of rough geodesic metric spaces. In particular, the strong shortcut property is a rough similarity invariant of rough geodesic metric spaces.*

**Proof** This follows immediately from Theorem 3.8 and Proposition 3.4. □




Figure 2: Continuing the pattern, one obtains an infinite graph that is strongly shortcut because it is the 1–skeleton of a finite-dimensional CAT(0) cube complex. Subdividing the *interior* edges of each  $n \times n$  grid results in a quasi-isometric graph that is not strongly shortcut.

### 3.1 Instability under quasi-isometries

In light of Corollary 3.10, we should point out that the strong shortcut property is not a quasi-isometry invariant. The 1–skeleton of an  $n \times n$  grid of squares is strongly shortcut but subdividing its *interior* edges causes its boundary cycle to become isometrically embedded. We can construct a strongly shortcut graph  $\Gamma$  that contains isometric copies of 1–skeletons of larger and larger  $n \times n$  grids. See Figure 2. Subdividing the interior edges of each  $n \times n$  grid of  $\Gamma$  does not change the quasi-isometry type but results in a graph that is not strongly shortcut because it contains arbitrarily long isometrically embedded cycles.

## 4 The circle tightening lemma

The circle tightening lemma describes how one may perform surgery on an almost isometric  $R$ –circle to obtain a quasi-isometrically embedded  $R$ –circle, assuming the various constants are chosen appropriately. A version of this lemma first appeared implicitly in the proof of a proposition in an earlier work [Hoda 2022, Proposition 3.5] where it applied only to graphs. Here we state and prove a generalization to (rough) geodesic metric spaces.

### 4.1 Tightening sequence for a Riemannian circle

Let  $S$  be a Riemannian circle. A *tightening sequence* for  $S$  is a sequence of intervals and Riemannian circles  $(P_i)_i$ , a sequence of Riemannian circles  $(S_i)_i$  and sequences of maps

$$S = S_0 \leftrightarrow P_0 \rightarrow S_1 \leftrightarrow P_1 \rightarrow S_2 \leftrightarrow P_2 \rightarrow \dots$$

such that, for each  $i$ , either

- (1) (a)  $P_i \hookrightarrow S_i$  and  $P_i \rightarrow S_{i+1}$  are continuous paths of unit speed,
- (b)  $P_i \rightarrow S_{i+1}$  is injective on the interior  $P_i^\circ$  of  $P_i$ ,
- (c)  $|P_i| \geq \frac{1}{2}|S|$ , and
- (d)  $|S_{i+1}| < |S_i|$ ; or
- (2) (a)  $P_i^\circ = P_i = S_i = S_{i+1}$ , and
- (b)  $P_i \hookrightarrow S_i$  and  $P_i \rightarrow S_{i+1}$  are identity maps.




Figure 3: In a circle tightening sequence, the circle  $S_{i+1}$  is either equal to  $S_i$  or is obtained from  $S_i$  by replacing some geodesic segment  $Q_i$  of  $S_i$  with a shorter segment  $\bar{Q}_i$ , possibly of zero length.

See Figure 3. Then, for each  $i$ , the circle  $S_{i+1}$  is obtained from  $S_i$  by replacing  $Q_i = S \setminus P_i^o$  with  $\bar{Q}_i$ , where either  $Q_i = \bar{Q}_i = \emptyset$  or  $Q_i$  and  $\bar{Q}_i$  are intervals with  $|\bar{Q}_i| < |Q_i|$ . So we also have a commutative diagram of 1-Lipschitz maps






Figure 4: A circle tightening sequence that is disjoint up to 4 but not disjoint up to 5. The outer circle is the initial circle  $S = S_0$ . For  $i \geq 0$ , the circle  $S_{i+1}$  is obtained from  $S_i$  by replacing the geodesic segment  $Q_i \subset S_i$  (indicated by perpendicular markings) with a shorter sequence  $\bar{Q}_i$ . The segment  $Q_4$  (drawn in cyan) is the first replaced segment that cannot be viewed as a subspace of  $S$  since it is not contained in  $P_{0,4}^o$ , which can be viewed as  $S \setminus \bigcup_{i=0}^3 Q_i$ .

where each  $Q_i \rightarrow \bar{Q}_i$  is affine. We call the  $Q_i$  the *tightened segments* of the tightening sequence. We let  $\pi^{(i)}$  denote the composition  $\pi_{i-1} \circ \pi_{i-2} \circ \dots \circ \pi_0$ . A tightening sequence is *eventually constant* if  $S_i = S_{i+1}$  for all large enough  $i$ .

Let  $P_{0,j}^\circ$  be the limit of the diagram



in the category of topological spaces and continuous maps. Concretely, we have  $P_{0,0}^\circ = S_0$ ,  $P_{0,1}^\circ = P_0^\circ$  and  $P_{0,j}^\circ = P_{0,j-1}^\circ \cap P_{j-1}^\circ$  where the intersection is taken in  $S_{j-1}$ . Thus we have the commutative diagram



We can think of  $P_{0,j}^\circ$  as the original points of  $S$  that are not replaced until at least step  $j$  of the “construction” of the  $S_i$ , where the  $j^{\text{th}}$  step of the construction refers to the operation of replacing  $Q_j$  with  $\bar{Q}_j$  in order to obtain  $S_{j+1}$  from  $S_j$ .

We say that a tightening sequence is *disjoint up to  $j$*  if  $Q_i \subset P_{0,i}^\circ$  for all  $i < j$ , where  $P_{0,i}^\circ$  is viewed as a subspace of  $S_i$  via the embedding  $P_{0,i}^\circ \hookrightarrow S_i$ . See Figure 4. We say that a tightening sequence is *completely disjoint* if it is disjoint up to  $j$  for every  $j$ .

If a tightening sequence is disjoint up to  $j$ , for  $i < j$ , we have  $Q_i \sqcup P_{0,i+1}^\circ = P_{0,i}^\circ \hookrightarrow S$ . So, for  $i < j$ , we may think of the  $Q_i$  as disjoint subspaces of  $S$  with  $S \setminus \bigcup_{i=0}^{j-1} Q_i = P_{0,j}^\circ$  in  $S$ . Since  $S_j$  is obtained from  $S$  by replacing  $Q_i$  with  $\bar{Q}_i$ , for each  $i < j$ , we see then that the  $\bar{Q}_i$ , with  $i < j$ , embed disjointly in  $S_j$  with  $P_{0,j}^\circ = S_j \setminus \bigcup_{i=0}^{j-1} \bar{Q}_i$  in  $S_j$ .

If a tightening sequence is completely disjoint then the  $Q_i$  all embed disjointly in  $S$  and the complement of their union in  $S$  is  $P_{0,\infty}^\circ = \bigcap_{j=1}^\infty P_{0,j}^\circ$ .

**Lemma 4.1** Consider a tightening sequence for a Riemannian circle  $S$  with the same notation as above. If the tightening sequence is completely disjoint and the sum  $\sum_{i=0}^\infty |Q_i|$  of the tightened segment lengths is strictly less than  $|S|$  then

$$\delta: S \times S \rightarrow \mathbb{R}_{\geq 0}, \quad (x, y) \mapsto \lim_{i \rightarrow \infty} d_{S_i}(\pi^{(i)}(x), \pi^{(i)}(y)),$$

defines a pseudometric on  $S$  such that the induced metric quotient  $S_\infty$  of  $(S, \delta)$ , called the **limit Riemannian circle** of the tightening sequence, is a Riemannian circle of length  $\lim_{i \rightarrow \infty} |S_i|$ .

**Proof** The  $\pi^{(i)}$  are isomorphisms on fundamental group so, for each  $i$ , we have a commuting diagram

$$\begin{array}{ccc} \mathbb{R} & \xrightarrow{\tilde{\pi}_i} & \mathbb{R} \\ \downarrow & & \downarrow \\ S_i & \xrightarrow{\pi_i} & S_{i+1} \end{array}$$

where  $\mathbb{R} \rightarrow S_i$  and  $\mathbb{R} \rightarrow S_{i+1}$  are the quotient maps from  $(\mathbb{R}, +)$  by the subgroups  $|S_i|\mathbb{Z}$  and  $|S_{i+1}|\mathbb{Z}$ , respectively. Then, since  $\pi_i$  is 1-Lipschitz, so is  $\tilde{\pi}_i$ . Without loss of generality, the map  $\tilde{\pi}_i$  sends 0 to 0 and preserves order, in the sense that  $s \leq r$  implies  $\tilde{\pi}_i(s) \leq \tilde{\pi}_i(r)$ . Let  $\tilde{\pi}^{(i)}: \mathbb{R} \rightarrow \mathbb{R}$  be the composition  $\tilde{\pi}_0 \circ \tilde{\pi}_1 \circ \dots \circ \tilde{\pi}_{i-1}$  so that the diagram

$$\begin{array}{ccc} \mathbb{R} & \xrightarrow{\tilde{\pi}^{(i)}} & \mathbb{R} \\ \downarrow & & \downarrow \\ S & \xrightarrow{\pi^{(i)}} & S_i \end{array}$$

commutes and satisfies the same properties as the previous diagram. Then for  $r \in \mathbb{R}$ , the sequence  $(\pi^{(i)}(r))_i$  is either nonnegative and nonincreasing or nonpositive and nondecreasing. In either case the limit exists so we can define a limit function

$$\pi^{(\infty)}: \mathbb{R} \rightarrow \mathbb{R}, \quad r \mapsto \lim_{i \rightarrow \infty} \pi^{(i)}(r),$$

which is also 1-Lipschitz, sends 0 to 0 and preserves order.

By assumption  $\sum_{i=1}^{\infty} |Q_i| < |S|$  so  $\pi^{(i)}(|S|) = |S_i| \geq |S| - \sum_{i=1}^{\infty} |Q_i| > 0$  and so

$$\pi^{(\infty)}(|S|) = \lim_{i \rightarrow \infty} |S_i| > 0.$$

For  $r \in \mathbb{R}$ , we have  $\pi^{(i)}(r + |S|) = \pi^{(i)}(r) + |S_i|$  so

$$\pi^{(\infty)}(r + |S|) = \pi^{(\infty)}(r) + \lim_{i \rightarrow \infty} |S_i|,$$

which implies that if  $\mathbb{R} \rightarrow \bar{S}$  is the quotient map of  $(\mathbb{R}, +)$  with kernel  $(\lim_{i \rightarrow \infty} |S_i|)\mathbb{Z}$  then the map

$$\pi^{(\infty)}: S \rightarrow S_{\infty}, \quad r + |S|\mathbb{Z} \mapsto \pi^{(\infty)}(r) + \left(\lim_{i \rightarrow \infty} |S_i|\right)\mathbb{Z},$$

is well defined and makes the diagram

$$\begin{array}{ccc} \mathbb{R} & \xrightarrow{\tilde{\pi}^{(\infty)}} & \mathbb{R} \\ \downarrow & & \downarrow \\ S & \xrightarrow{\pi^{(\infty)}} & \bar{S} \end{array}$$

commute.

Then  $|\bar{S}| = \lim_{i \rightarrow \infty} |S_i|$  and, for  $r + |S|\mathbb{Z}$  and  $s + |S|\mathbb{Z}$  in  $S$ ,

$$\begin{aligned} d_{\bar{S}}(\pi^{(\infty)}(r + |S|\mathbb{Z}), \pi^{(\infty)}(s + |S|\mathbb{Z})) &= d_{\bar{S}}(\pi^{(\infty)}(r) + |\bar{S}|\mathbb{Z}, \pi^{(\infty)}(s) + |\bar{S}|\mathbb{Z}) \\ &= \min_{k \in \mathbb{Z}} |\pi^{(\infty)}(r) - \pi^{(\infty)}(s) + |\bar{S}|k| \end{aligned}$$


$$\begin{aligned}
 &= \min_{k \in I} |\pi^{(\infty)}(r) - \pi^{(\infty)}(s) + |\bar{S}|k| \\
 &= \min_{k \in I} \lim_{i \rightarrow \infty} |\pi^{(i)}(r) - \pi^{(i)}(s) + |\bar{S}_i|k| \\
 &= \lim_{i \rightarrow \infty} \min_{k \in I} |\pi^{(i)}(r) - \pi^{(i)}(s) + |\bar{S}_i|k| \\
 &= \lim_{i \rightarrow \infty} \min_{k \in \mathbb{Z}} |\pi^{(i)}(r) - \pi^{(i)}(s) + |\bar{S}_i|k| \\
 &= \lim_{i \rightarrow \infty} d_{S_i}(\pi^{(i)}(r) + |S_i|\mathbb{Z}, \pi^{(i)}(s) + |S_i|\mathbb{Z}) \\
 &= \lim_{i \rightarrow \infty} d_{S_i}(\pi^{(i)}(r + |S|\mathbb{Z}), \pi^{(i)}(s + |S|\mathbb{Z})),
 \end{aligned}$$

where  $I = \{k \in \mathbb{Z} \mid |k| \leq \lceil |r - s|/|\bar{S}| \rceil\}$ . So the pseudometric on  $S$  pulled back from  $\pi^{(\infty)}$  is  $\delta$ . Then, since  $\pi^{(\infty)}$  is surjective, this implies that  $\bar{S}$  is  $S_\infty$ , the induced metric quotient of  $(S, \delta)$  and  $\pi^\infty$  is the quotient map. □

**Remark 4.2** If a completely disjoint tightening sequence of a Riemannian circle is eventually constant then, for large enough  $i$ , the limit Riemannian circle  $S_\infty$  is isometric to the  $i^{\text{th}}$  Riemannian circle of the sequence  $S_i$ .

### 4.2 Tightening sequence for an $R$ -circle


Let  $R \geq 0$  and let  $X$  be an  $R$ -rough geodesic metric space. Let  $\alpha: S \rightarrow X$  be an  $R$ -circle. A *tightening sequence* for  $\alpha$  is a sequence of intervals and Riemannian circles  $(P_i)_i$ , a sequence of Riemannian circles  $(S_i)_i$  and sequences of maps as in the commutative diagram



such that each  $\alpha_i$  is an  $R$ -circle and the sequence of maps

$$S = S_0 \leftrightarrow P_0 \rightarrow S_1 \leftrightarrow P_1 \rightarrow S_2 \leftrightarrow P_2 \rightarrow \dots$$

is a tightening sequence for  $S$ . Then, by the discussion of Section 4.1, we have a diagram



where the bounded planar regions are commuting triangles and squares.

**Lemma 4.3** Consider a tightening sequence for an  $R$ -circle  $\alpha : S \rightarrow X$  in an  $R$ -rough geodesic metric space  $X$ , with the same notation as above. If the tightening sequence is completely disjoint and the sum  $\sum_{i=0}^{\infty} |Q_i|$  of the tightened segment lengths is strictly less than  $|S|$  then

$$\alpha_{\infty} : S_{\infty} \rightarrow X, \quad x \mapsto \lim_{i \rightarrow \infty} \alpha_i \circ \pi^{(i)} \circ \sigma(x),$$

defines an  $R$ -circle, called a **limit  $R$ -circle** of the tightening sequence, where  $S_{\infty}$  is the limit Riemannian circle of the tightening sequence and  $\sigma : S_{\infty} \rightarrow S$  is a section of the quotient map  $S \rightarrow S_{\infty}$ .

**Proof** Since the tightening sequence is completely disjoint, we may think of the  $Q_i$  as a collection of disjoint segments in  $S$ . For  $x \in S$  either  $x \notin \bigcup_{i=0}^{\infty} Q_i$  and  $(\alpha_i \circ \pi^{(i)}(x))_{i=0}^{\infty}$  is a constant sequence or  $x \in Q_j$  for some  $j$  and the tail sequence  $(\alpha_i \circ \pi^{(i)}(x))_{i=j+1}^{\infty}$  is constant. In either case, the limit  $\lim_{i \rightarrow \infty} \alpha_i \circ \pi^{(i)}(x)$  exists so we have a function  $\alpha'_{\infty} : S \rightarrow X$  given by  $\alpha'_{\infty}(x) = \lim_{i \rightarrow \infty} \alpha_i \circ \pi^{(i)}(x)$ .

Let  $S_{\infty}$  be the limit Riemannian circle given by Lemma 4.1. So  $S_{\infty}$  is the induced metric quotient of  $(S, \delta)$  where  $\delta$  is the pseudometric give by  $\delta(x, y) = \lim_{i \rightarrow \infty} d_{S_i}(\pi^{(i)}(x), \pi^{(i)}(y))$ . Since each  $\alpha_i$  is an  $R$ -circle, for  $x, y \in S$ ,

$$d_X(\alpha_i \circ \pi^{(i)}(x), \alpha_i \circ \pi^{(i)}(y)) \leq d_{S_i}(\pi^{(i)}(x), \pi^{(i)}(y)) + R$$

for all  $i$ , and thus

$$d_X(\alpha'_{\infty}(x), \alpha'_{\infty}(y)) \leq \delta(x, y) + R$$

by taking limits as  $i \rightarrow \infty$ .

Then, for  $x, y \in S_{\infty}$ ,

$$d_X(\alpha_{\infty}(x), \alpha_{\infty}(y)) = d_X(\alpha'_{\infty}(\sigma(x)), \alpha'_{\infty}(\sigma(y))) \leq \delta(\sigma(x), \sigma(y)) + R = d_{S_{\infty}}(x, y) + R. \quad \square$$

**Remark 4.4** If a completely disjoint tightening sequence for an  $R$ -circle  $\alpha : S \rightarrow X$  is eventually constant then, for large enough  $i$ , the limit  $R$ -circle  $\alpha_{\infty} : S_{\infty} \rightarrow X$  is isometric over  $X$  to the  $i^{\text{th}}$   $R$ -circle of the sequence  $\alpha_i$ . This means that there is an isometry  $S_{\infty} \rightarrow S$  such that diagram

$$\begin{array}{ccc} S_{\infty} & \longrightarrow & S_i \\ & \searrow \alpha_{\infty} & \downarrow \alpha_i \\ & & X \end{array}$$

commutes.

We are ready now to state the circle tightening lemma.

**Lemma 4.5** (circle tightening lemma) Let  $N > 1$ , let  $L > 1$ , let  $K > 1$  be small enough (depending on  $N$  and  $L$ ), let  $R \geq 0$ , let  $M > 0$  be large enough (depending on  $N, L, K$  and  $R$ ) and let  $C \geq 4R$ .

Let  $\alpha : S \rightarrow X$  be an  $R$ -circle in an  $R$ -rough geodesic metric space. If  $\alpha$  is  $1/K$ -almost isometric and  $|S| > M$  then  $\alpha$  has a completely disjoint tightening sequence such that the total length  $\sum_{i=0}^{\infty} |Q_i|$  of the tightened segments is at most  $|S|/N$  and the limit  $R$ -circle  $\alpha_{\infty}$  is an  $(L, C)$ -quasi-isometric embedding. If, additionally,  $C > 0$  then such a tightening sequence exists that is eventually constant.

Lemma 4.5 is a consequence of Claims 4.7, 4.8, 4.11 and 4.14 and the strict inequalities of Claim 4.12 below, but to understand these claims we need to first define greedy tightening sequences and prove some properties about them.

### 4.3 Greedy tightening sequences

In order to prove Lemma 4.5 we will need to describe a tightening sequence that is constructed inductively by greedily choosing segments to tighten. Let  $\alpha: S \rightarrow X$  be an  $R$ -circle in an  $R$ -rough geodesic metric space  $X$ , with  $R \geq 0$ . Let  $C \geq 4R$  and let  $L > 1$ . We will inductively define a tightening sequence for  $\alpha$  with the same notation as in the previous sections. Suppose we have  $\alpha_i: S_i \rightarrow X$ . If  $\alpha_i$  is an  $(L, C)$ -quasi-isometric embedding then we extend the sequence as follows.

- (1) We set  $P_i^\circ = P_i = S_i = S_{i+1}$  and  $Q_i = \bar{Q}_i = \emptyset$ .
- (2) We let  $P_i \hookrightarrow S_i$  and  $P_i \rightarrow S_{i+1}$  be identity maps.
- (3) We let  $\alpha_{i+1} = \alpha_i$ .

Otherwise, the set

$$J_i = \left\{ (p, q) \in S_i \times S_i \mid d_X(\alpha_i(p), \alpha_i(q)) < \frac{1}{L}d_{S_i}(p, q) - C \right\}$$

is nonempty and  $d_{S_i}(p, q) > LC \geq 0$  for any  $(p, q) \in J_i$  and so  $s_i = \sup\{d_{S_i}(p, q) \mid (p, q) \in J_i\} > 0$ . By compactness of  $S$ , there is a sequence  $(p_i^{(n)}, q_i^{(n)})_n$  in  $J_i$  that converges to some  $(p_i, q_i) \in S \times S$  with  $d_{S_i}(p_i, q_i) = s_i$  as  $n \rightarrow \infty$ . Then

$$\begin{aligned} d_X(\alpha_i(p_i), \alpha(q_i)) &\leq d_X(\alpha_i(p_i), \alpha(p_i^{(n)})) + d_X(\alpha_i(p_i^{(n)}), \alpha(q_i^{(n)})) + d_X(\alpha_i(q_i^{(n)}), \alpha(q_i)) \\ &< d_{S_i}(p_i, p_i^{(n)}) + R + \frac{1}{L}d_{S_i}(p_i^{(n)}, q_i^{(n)}) - C + d_{S_i}(q_i^{(n)}, q_i) + R \\ &\rightarrow \frac{1}{L}d_{S_i}(p_i, q_i) - C + 2R \end{aligned}$$

as  $n \rightarrow \infty$ . So

$$(*) \quad d_X(\alpha_i(p_i), \alpha(q_i)) \leq \frac{1}{L}d_{S_i}(p_i, q_i) - C + 2R,$$

$$(\dagger) \quad d_{S_i}(p_i, q_i) > 0 \quad \text{and} \quad d_{S_i}(p_i, q_i) \geq L(C - 2R)$$

hold. Let  $Q_i$  be a geodesic segment of  $S_i$  between  $p_i$  and  $q_i$ . In the case where  $p_i$  and  $q_i$  are antipodal in  $S_i$ , there are two geodesic segments between  $p_i$  and  $q_i$ ; in this case we let  $Q_i$  be the geodesic segment whose intersection with  $P_{0,i}^\circ$  has greatest total length. Let  $P_i^\circ$  be the complement of  $Q_i$  and let  $P_i$  be the closure of  $P_i^\circ$ . Let  $\gamma'_i: \bar{Q}'_i \rightarrow X$  be an  $R$ -rough geodesic from  $\alpha_i(p_i)$  to  $\alpha_i(q_i)$ . For  $x \in X$ , let  $c_x: [0, R] \rightarrow X$  denote the constant path of length  $R$  at  $x$ . Let  $\gamma_i: \bar{Q}_i \rightarrow X$  be the concatenation  $c_{\alpha_i(p_i)}\gamma'_i c_{\alpha_i(q_i)}$ . We have

$$0 \leq |\bar{Q}_i| - 2R = d_X(\alpha_i(p_i), \alpha_i(q_i)) \leq \frac{1}{L}|Q_i| - C + 2R$$

and so, since  $C \geq 4R$ ,

$$(\ddagger) \quad |\bar{Q}_i| \leq \frac{1}{L}|Q_i|.$$

We obtain  $\alpha_{i+1}: S_{i+1} \rightarrow X$  from  $\alpha_i|_{P_i}$  and  $\gamma_i$  by identifying the corresponding endpoints of  $\bar{Q}_i$  with  $p_i$  and  $q_i$  in  $P_i$ . Then, by consideration of Remark 2.1, the map  $\alpha_{i+1}$  is an  $R$ -circle.

**Remark 4.6** The inequalities

$$d_X(\alpha_{i+1}(p_i), \alpha_{i+1}(x)) \leq d_{S_{i+1}}(p_i, x), \quad d_X(\alpha_{i+1}(q_i), \alpha_{i+1}(x)) \leq d_{S_{i+1}}(q_i, x)$$

hold for any  $x \in \bar{Q}_i$ .

This completes the description of our inductive construction. Any tightening sequence for  $\alpha$  obtained in this way is called an  $(L, C)$ -greedy tightening sequence for  $\alpha$ . The importance of this construction for us is evident from the following claim.

**Claim 4.7** Let  $L > 1$ , let  $K > 1$ , let  $C \geq 4R$  and consider an  $(L, C)$ -greedy tightening sequence for a  $1/K$ -almost isometric  $R$ -circle  $\alpha: S \rightarrow X$  in an  $R$ -rough geodesic metric space  $X$ . If the tightening sequence is completely disjoint and the sum  $\sum_{i=0}^{\infty} |Q_i|$  of the tightened segment lengths is strictly less than  $|S|$  then any limiting  $R$ -circle  $\alpha_{\infty}: S_{\infty} \rightarrow X$  is an  $(L, C)$ -quasi-isometric embedding.

**Proof** Let  $S_{\infty}$  be the limit Riemannian circle given by Lemma 4.1. So  $S_{\infty}$  is the induced metric quotient of  $(S, \delta)$  where  $\delta$  is the pseudometric given by  $\delta(x, y) = \lim_{i \rightarrow \infty} d_{S_i}(\pi^{(i)}(x), \pi^{(i)}(y))$ . Let  $\alpha_{\infty}: S_{\infty} \rightarrow X$  be a limit  $R$ -circle as in Lemma 4.3. So  $\alpha_{\infty}$  is an  $R$ -circle defined by  $\alpha_{\infty}(x) = \lim_{i \rightarrow \infty} \alpha_i \circ \pi^{(i)} \circ \sigma(x)$ , where  $\sigma: S_{\infty} \rightarrow S$  is a section of the quotient map  $S \rightarrow S_{\infty}$ .

If  $\alpha_{\infty}$  is not an  $(L, C)$ -quasi-isometric embedding then, since  $R \leq C$ ,

$$d_X(\alpha_{\infty}(x), \alpha_{\infty}(y)) < \frac{1}{L}d_{S_{\infty}}(x, y) - C$$

for some  $x, y \in S_{\infty}$ , which then must be distinct. But

$$d_{S_{\infty}}(x, y) = \lim_{i \rightarrow \infty} d_{S_i}(\pi^{(i)}(\sigma(x)), \pi^{(i)}(\sigma(y))),$$

$$d_X(\alpha_{\infty}(x), \alpha_{\infty}(y)) = \lim_{i \rightarrow \infty} d_X(\alpha_i \circ \pi^{(i)}(\sigma(x)), \alpha_i \circ \pi^{(i)}(\sigma(y))),$$

where, by complete disjointness,  $(\pi^{(i)}(\sigma(x)))_i$  and  $(\pi^{(i)}(\sigma(y)))_i$  are eventually constant. So, for all large enough  $j$ ,

$$d_X(\alpha_j \circ \pi^{(j)}(\sigma(x)), \alpha_j \circ \pi^{(j)}(\sigma(y))) = \lim_{i \rightarrow \infty} d_X(\alpha_i \circ \pi^{(i)}(\sigma(x)), \alpha_i \circ \pi^{(i)}(\sigma(y)))$$

but also, for all large enough  $j$ ,

$$\lim_{i \rightarrow \infty} d_X(\alpha_i \circ \pi^{(i)}(\sigma(x)), \alpha_i \circ \pi^{(i)}(\sigma(y))) < \frac{1}{L}d_{S_j}(\pi^{(j)}(\sigma(x)), \pi^{(j)}(\sigma(y))) - C$$

so, for all large enough  $j$ ,

$$d_X(\alpha_j(\pi^{(j)} \circ \sigma(x)), \alpha_j(\pi^{(j)} \circ \sigma(y))) < \frac{1}{L}d_{S_j}(\pi^{(j)} \circ \sigma(x), \pi^{(j)} \circ \sigma(y)) - C$$

which implies that  $(\pi^{(j)} \circ \sigma(x), \pi^{(j)} \circ \sigma(y)) \in J_j$ , for all large enough  $j$ . But then, for all large enough  $j$ ,

$$d_{S_j}(\pi^{(j)} \circ \sigma(x), \pi^{(j)} \circ \sigma(y)) \leq s_j = |Q_j|$$

with  $\lim_{j \rightarrow \infty} |Q_j| = 0$  so

$$d_{S_\infty}(x, y) = \lim_{j \rightarrow \infty} d_{S_j}(\pi^{(j)} \circ \sigma(x), \pi^{(j)} \circ \sigma(y)) = 0$$

which is a contradiction. □

#### 4.4 Eventual constantness and greedy tightening sequences

Consider a greedy tightening sequence with notation as in Section 4.3. Note that if  $S_i = S_{i+1}$  for some  $i$  then  $S_i = S_{i+1} = S_{i+2} = \dots$ , so the tightening sequence is eventually constant. Moreover, for any  $i$  for which  $S_i \neq S_{i+1}$ , we have

$$\begin{aligned} |S_i| - |S_{i+1}| &= |Q_i| - |\bar{Q}_i| \geq |Q_i| - \frac{1}{L}|Q_i| = \left(1 - \frac{1}{L}\right)|Q_i| \\ &\geq \left(1 - \frac{1}{L}\right)L(C - 2R) = (L - 1)(C - 2R) \end{aligned}$$

by (†) and (‡). If  $R > 0$  then, since  $C \geq 4R$ , we have  $C > 2R > 0$ . If  $R = 0$  then  $C > 2R$  is equivalent to  $C > 0$ . Thus, if  $C > 0$  then  $|S_i| - |S_{i+1}| \geq (L - 1)(C - 2R) > 0$ . This implies the following claim.

**Claim 4.8** *If  $R \geq 0$ ,  $L > 1$  and  $C > 0$  then any  $(L, C)$ -greedy tightening sequence for an  $R$ -circle in an  $R$ -rough geodesic space is eventually constant.*

#### 4.5 Disjointness and greedy tightening sequences

Consider a greedy tightening sequence with notation as in Section 4.3. Assume that  $\alpha$  is  $1/K$ -almost isometric and that the tightening sequence is disjoint up to  $j$ . Recall that, by the discussion in Section 4.1, we may think of the  $Q_i$ , with  $i < j$ , as disjoint subspaces of  $S$ .

If  $i < j$  and  $Q_i \neq \emptyset$  then, by Lemma 3.1 and (\*),

$$|Q_i| - \frac{K-1}{K} \cdot \frac{|S|}{2} - 2R \leq d_X(\alpha(p_i), \alpha(q_i)) \leq \frac{1}{L}|Q_i| - C + 2R,$$

but  $C \geq 4R$  so

$$|Q_i| - \frac{K-1}{K} \cdot \frac{|S|}{2} \leq \frac{1}{L}|Q_i|$$

for all  $i < j$ . Hence, we have established the following claim.

**Claim 4.9** *If an  $(L, C)$ -greedy tightening sequence for a  $1/K$ -almost isometric  $R$ -circle is disjoint up to  $j$  then*

$$|Q_i| \leq \left( \frac{K-1}{K} \cdot \frac{L}{L-1} \right) \frac{|S|}{2}$$

for any  $i < j$ , where  $Q_i$  is the  $i^{\text{th}}$  replaced segment of the tightening sequence.

By this claim, we can find a pair of points  $p, q$  in the closure of  $S \setminus (\bigcup_{i=1}^{j-1} Q_i)$  at distance

$$d_S(p, q) \geq \frac{|S|}{2} - \frac{K-1}{K} \cdot \frac{L}{L-1} \cdot \frac{|S|}{4}.$$

Let  $A_1$  and  $A_2$  be the two segments of  $S$  between  $p$  and  $q$ . If  $I_1 = \{i < j \mid Q_i \subseteq A_1\}$  then, since  $\alpha_j$  is an  $R$ -circle,

$$\begin{aligned} d_X(\alpha(p), \alpha(q)) &\leq d_{S_j}(p, q) + R \\ &\leq |A_1| - \sum_{i \in I_1} |Q_i| + \sum_{i \in I_1} |\bar{Q}_i| + R \\ &\leq |A_1| - \sum_{i \in I_1} |Q_i| + \frac{1}{L} \sum_{i \in I_1} |Q_i| + R \\ &= |A_1| - \frac{L-1}{L} \sum_{i \in I_1} |Q_i| + R \end{aligned}$$

where the last inequality follows by  $(\ddagger)$ . Corresponding relations for  $A_2$  and  $I_2 = \{i < j \mid Q_i \subseteq A_2\}$  also hold, and so, by Lemma 3.1,

$$\begin{aligned} |S| - \frac{L-1}{L} \sum_{i < j} |Q_i| + 2R &\geq 2d_X(\alpha(p), \alpha(q)) \geq 2d_S(p, q) - \frac{K-1}{K} \cdot |S| - 4R \\ &\geq |S| - \left( \frac{K-1}{K} \cdot \frac{L}{L-1} \right) \frac{|S|}{2} - \frac{K-1}{K} \cdot |S| - 4R, \end{aligned}$$

which establishes the following claim.

**Claim 4.10** *If an  $(L, C)$ -greedy tightening sequence for a  $1/K$ -almost isometric  $R$ -circle is disjoint up to  $j$  then*

$$\sum_{i < j} |Q_i| \leq \left( \frac{K-1}{K} \cdot \frac{L(3L-2)}{2(L-1)^2} \right) |S| + \frac{6LR}{L-1}$$

where  $Q_i$  is the  $i^{\text{th}}$  replaced segment of the tightening sequence.

Claim 4.10 implies that

$$\sum_{i < j} |Q_i| \leq \left( \frac{K-1}{K} \cdot \frac{L(3L-2)}{2(L-1)^2} + \frac{6LR}{|S|(L-1)} \right) |S|,$$

so if

$$\frac{K-1}{K} \cdot \frac{L(3L-2)}{2(L-1)^2} < \frac{1}{N}$$

then, if  $|S| > M$  for some  $M$  depending only on  $K, L, R$  and  $N$ , then

$$\frac{K-1}{K} \cdot \frac{L(3L-2)}{2(L-1)^2} + \frac{6LR}{|S|(L-1)} < \frac{1}{N}$$

and so  $\sum_{i < j} |Q_i| < |S|/N$ . Since

$$\frac{K-1}{K} \cdot \frac{L(3L-2)}{2(L-1)^2} < \frac{1}{N}$$

is equivalent to

$$K < \frac{NL(3L - 2)}{(3N - 2)L^2 - (2N - 4)L - 2},$$

we have established the following claim.

**Claim 4.11** *Let  $N > 1$ , let  $L > 1$ , let  $R \geq 0$  and let  $K > 1$  satisfy*

$$K < \frac{NL(3L - 2)}{(3N - 2)L^2 - (2N - 4)L - 2}.$$

*Then there exists an  $M > 0$  such that if an  $(L, C)$ -greedy tightening sequence for a  $1/K$ -almost isometric  $R$ -circle  $\alpha: S \rightarrow X$  of length  $|S| > M$  is disjoint up to  $j$  then*

$$\sum_{i < j} |Q_i| < \frac{|S|}{N}$$

*where the  $Q_i$  are the tightened segments of the tightening sequence.*

The following claim about rational functions has a short and elementary proof. We will make use of it below.

**Claim 4.12** *The inequalities*

$$1 < \frac{L(9L^2 - 3L - 4)}{7L^3 + 3L^2 - 10L + 2}, \quad \frac{L(9L^2 - 3L - 4)}{7L^3 + 3L^2 - 10L + 2} \leq \frac{L(5L - 4)}{3L^2 - 2},$$

$$1 < \frac{NL(3L - 2)}{(3N - 2)L^2 - (2N - 4)L - 2}, \quad \frac{L(9L^2 - 3L - 4)}{7L^3 + 3L^2 - 10L + 2} \leq \frac{L(7L - 6)}{5L^2 - 2L - 2}$$

*hold for any  $L > 1$  and  $N > 1$ .*

The next claim is essential in proving disjointness of greedy tightening sequences.

**Claim 4.13** *Let  $K > 1$ ,  $L > 1$  and  $R \geq 0$ . There exists an  $M > 0$  such that if  $X$  is an  $R$ -rough geodesic metric space,  $\alpha: S \rightarrow X$  is a  $1/K$ -almost isometric  $R$ -circle with  $|S| > M$  and*

$$K < \frac{L(9L^2 - 3L - 4)}{7L^3 + 3L^2 - 10L + 2},$$

*and  $C \geq 4R$  then any  $(L, C)$ -greedy tightening sequence for  $\alpha$  that is disjoint up to  $j$  satisfies the following statement. With notation as above, if  $(p, q) \in J_j$  and  $Q$  is a geodesic segment from  $p$  to  $q$  in  $S_j$ , then  $Q \subset P_{0,j}^\circ$ , where we view  $P_{0,j}^\circ$  as a subspace of  $S_j$  via the embedding  $P_{0,j}^\circ \hookrightarrow S_j$ .*

**Proof** First we will show that  $Q$  is not contained in  $\bar{Q}_i$  for any  $i < j$ . Recall that  $\bar{Q}_i$  is the concatenation  $A\bar{Q}'_iB$  where  $\alpha_j|_{\bar{Q}'_i}: \bar{Q}'_i \rightarrow X$  is an  $R$ -rough geodesic and  $\alpha_j$  is constant on  $A$  and  $B$ , each of which is

isometric to  $[0, R]$ . By  $(\dagger)$ , we have  $|Q| \geq (C - 2R)L \geq (4R - 2R)L \geq 2R$  and  $|Q| > 0$  so we cannot have  $Q \subseteq A$  or  $Q \subseteq B$ . We also cannot have  $\bar{Q}'_i \subseteq Q \subseteq \bar{Q}_i$  since then

$$|\bar{Q}_i| - 2R = d_X(\alpha_i(p_i), \alpha_i(q_i)) = d_X(\alpha_j(p), \alpha_j(q)) < \frac{1}{L}|Q| - C \leq \frac{1}{L}|\bar{Q}_i| - C < |\bar{Q}_i| - C,$$

which contradicts  $C \geq 4R$ . So, if  $Q \subseteq \bar{Q}_i$  then some endpoint of  $Q$  is contained in  $\bar{Q}'_i$ . But this implies that  $\alpha_j|_Q: Q \rightarrow X$  is a  $2R$ -rough geodesic and so, by  $(*)$  and  $(\dagger)$ ,

$$|Q| - 2R \leq d_X(\alpha_j(p), \alpha_j(q)) < \frac{1}{L}|Q| - C < |Q| - C,$$

which, again, contradicts  $C \geq 4R$ . Thus we see that  $Q$  is not contained in  $\bar{Q}_i$  for an  $i < j$ . Hence  $Q$  intersects  $P_{0,j}^\circ$  nontrivially in  $S_j$ .

See Figure 5. We will define a segment  $\bar{A}_p \subset S_j$  containing  $p$  and a corresponding segment  $A_p \subset S$ . (Note that  $\bar{A}_p$  does not denote the closure of  $A_p$  here.) If  $p$  is contained in the interior of  $\bar{Q}_i$  for some  $i < j$  then let  $\bar{A}_p = \bar{Q}_i$  and let  $A_p = Q_i$ . Otherwise, let  $A_p = \bar{A}_p = \{p\}$ . Define  $A_q$  and  $\bar{A}_q$  similarly for  $q$ . A priori, it is possible that  $\bar{A}_p = \bar{A}_q$ . Let  $Q^-$  be obtained from  $Q$  by subtracting the interiors of  $\bar{A}_p$  and  $\bar{A}_q$  and let  $Q^+ \rightarrow S_j$  extend  $Q \hookrightarrow S_j$  so as to include a full copy of  $\bar{A}_p$  and a full copy of  $\bar{A}_q$ . Let  $Q_0^- \subset S$  be obtained from  $Q^- \subset S_j$  by replacing any  $\bar{Q}_i \subset Q^-$  with  $Q_i \subset S$ , for  $i < j$ . Let  $Q_0^+ \rightarrow S$  be obtained from  $Q^+ \rightarrow S_j$  by replacing any  $\bar{Q}_i \hookrightarrow S_j$ , where  $i < j$ , with  $Q_i \hookrightarrow S$ .

Let  $p^+$  and  $q^+$  be the images of the endpoints of  $Q_0^+$  in  $S$ , with  $p^+$  the endpoint corresponding to  $p$  and  $q^+$  the endpoint corresponding to  $q$ . Let  $p^-$  and  $q^-$  be the endpoints of  $Q_0^-$  in  $S$ , with  $p^-$  the endpoint corresponding to  $p$  and  $q^-$  the endpoint corresponding to  $q$ . Then we have

$$\begin{aligned} d_X(\alpha(p^+), \alpha(q^+)) &\leq d_X(\alpha(p^+), \alpha_j(p)) + d_X(\alpha_j(p), \alpha_j(q)) + d_X(\alpha_j(q), \alpha(q^+)) \\ &\leq d_{\bar{A}_p}(p^+, p) + d_X(\alpha_j(p), \alpha_j(q)) + d_{\bar{A}_q}(q, q^+) \\ &< d_{\bar{A}_p}(p^+, p) + \frac{1}{L}|Q| - C + d_{\bar{A}_q}(q, q^+) \\ &= d_{\bar{A}_p}(p^+, p) + \frac{1}{L}(d_{\bar{A}_p}(p, p^-) + |Q^-| + d_{\bar{A}_q}(q^-, q)) + d_{\bar{A}_q}(q, q^+) - C \\ &\leq d_{\bar{A}_p}(p^+, p) + d_{\bar{A}_p}(p, p^-) + \frac{1}{L}|Q^-| + d_{\bar{A}_q}(q^-, q) + d_{\bar{A}_q}(q, q^+) - C \\ &= |\bar{A}_p| + \frac{1}{L}|Q^-| + |\bar{A}_q| - C \\ &\leq |\bar{A}_p| + \frac{1}{L}|Q_0^-| + |\bar{A}_q| - C \\ &\leq \frac{1}{L}|A_p| + \frac{1}{L}|Q_0^-| + \frac{1}{L}|A_q| - C = \frac{1}{L}|Q_0^+| - C \end{aligned}$$

where the second inequality follows from Remark 4.6 and the last inequality follows from  $(\ddagger)$ . By assumption,  $Q$  nontrivially intersects at least one  $\bar{Q}_i$ , with  $i < j$ . Let  $m$  be minimal such that  $Q$  nontrivially intersects  $\bar{Q}_m$ . Then, since  $Q$  intersects  $P_{0,j}^\circ$  nontrivially, the image of  $Q_0^+ \rightarrow S_m$  must strictly contain  $Q_m$ .




Figure 5: In the proof of Claim 4.13, from a geodesic segment  $Q$  of  $S_j$  (indicated by a green outline) we obtain  $Q^{-1}$  by removing the interiors of any  $\bar{Q}_i$  that are partially contained in  $Q$ . We obtain  $Q^+ \rightarrow S_j$  by extending the inclusion  $Q \hookrightarrow S_j$  to include full copies of any  $\bar{Q}_i$  that are partially contained in  $Q$ . From  $Q^- \hookrightarrow S_j$  and  $Q^+ \rightarrow S_j$  we obtain  $Q_0^- \hookrightarrow S$  and  $Q_0^+ \rightarrow S$  in  $S = S_0$  by replacing any  $\bar{Q}_i \hookrightarrow S_j$  with  $Q_i \hookrightarrow S$ . The  $\bar{Q}_i$  with  $i < j$  are drawn in red in  $S_j$ . The  $Q_i$  with  $i < j$  are drawn with perpendicular markings in  $S$ .

So, if  $Q_0^+ \rightarrow S_m$  was the inclusion of a geodesic segment, we would have  $(p^+, q^+) \in J_m$ , which would contradict  $d_{S_m}(p_m, q_m) = s_m$ . Thus  $|Q_0^+| > \frac{1}{2}|S_m|$ . But then

$$\begin{aligned} |Q_0^+| &> \frac{|S_m|}{2} \geq \frac{|S|}{2} - \frac{1}{2} \sum_{i < m} |Q_i| \geq \frac{|S|}{2} - \left( \frac{K-1}{K} \cdot \frac{L(3L-2)}{2(L-1)^2} \right) \frac{|S|}{2} - \frac{3LR}{L-1} \\ &= \left( 1 - \frac{K-1}{K} \cdot \frac{L(3L-2)}{2(L-1)^2} - \frac{6LR}{|S|(L-1)} \right) \frac{|S|}{2} \end{aligned}$$

by Claim 4.10 while

$$|Q_0| \leq \left( \frac{K-1}{K} \cdot \frac{L}{L-1} \right) \frac{|S|}{2}$$

by Claim 4.9. So  $|Q_0^+| \leq |Q_0|$  would imply

$$1 - \frac{K-1}{K} \cdot \frac{L(3L-2)}{2(L-1)^2} - \frac{6LR}{|S|(L-1)} < \frac{K-1}{K} \cdot \frac{L}{L-1}$$

which is equivalent to the inequality

$$(\S) \quad 1 < \frac{K-1}{K} \cdot \frac{L(5L-4)}{2(L-1)^2} + \frac{6LR}{|S|(L-1)}.$$

By hypothesis and Claim 4.12, we have  $K < L(5L-4)/(3L^2-2)$  which is equivalent to

$$1 > \frac{K-1}{K} \cdot \frac{L(5L-4)}{2(L-1)^2}$$

so if  $|S| > M'$  for some  $M'$  depending only on  $K, L$  and  $R$  then we would have

$$1 > \frac{K-1}{K} \cdot \frac{L(5L-4)}{2(L-1)^2} + \frac{6LR}{|S|(L-1)}$$

and this would contradict (§). Hence, assuming  $|S|$  is greater than this  $M'$ , we have  $|Q_0^+| > |Q_0|$ .

Then if  $Q_0^+ \rightarrow S$  were the inclusion of a geodesic segment, we would have  $(p^+, q^+) \in J_0$  and this would contradict  $d_S(p_0, q_0) = s_0$ . Thus  $|Q_0^+| > \frac{1}{2}|S|$ . On the other hand

$$\begin{aligned}
|Q_0^+| &= |Q_0^-| + |A_p| + |A_q| \\
&\leq |Q^-| + \sum_{i < j} (|Q_i| - |\bar{Q}_i|) + |A_p| + |A_q| \\
&\leq \frac{|S_j|}{2} + \sum_{i < j} (|Q_i| - |\bar{Q}_i|) + |A_p| + |A_q| \\
&= \frac{|S|}{2} + \frac{1}{2} \sum_{i < j} (|Q_i| - |\bar{Q}_i|) + |A_p| + |A_q| \\
&\leq \frac{|S|}{2} + \frac{1}{2} \sum_{i < j} |Q_i| + |A_p| + |A_q| \\
&\leq \frac{|S|}{2} + \left( \frac{K-1}{K} \cdot \frac{L(3L-2)}{4(L-1)^2} \right) |S| + \frac{3LR}{L-1} + \left( \frac{K-1}{K} \cdot \frac{L}{L-1} \right) |S| \\
&= \frac{|S|}{2} + \frac{K-1}{K} \cdot \frac{L}{L-1} \cdot \left( \frac{3L-2}{4(L-1)} + 1 \right) |S| + \frac{3LR}{L-1} \\
&= \frac{|S|}{2} + \frac{K-1}{K} \cdot \frac{L}{L-1} \cdot \frac{7L-6}{4(L-1)} \cdot |S| + \frac{3LR}{L-1} \\
&= \left( \frac{1}{2} + \frac{K-1}{K} \cdot \frac{L(7L-6)}{4(L-1)^2} + \frac{3LR}{|S|(L-1)} \right) |S|,
\end{aligned}$$

where the last inequality follows from Claims 4.9 and 4.10. By hypothesis and Claim 4.12,

$$K < \frac{L(7L-6)}{5L^2-2L-2},$$

which is equivalent to

$$\frac{1}{2} + \frac{K-1}{K} \cdot \frac{L(7L-6)}{4(L-1)^2} < 1.$$

Thus, if  $|S| > M''$  for some  $M''$  depending only on  $K$ ,  $L$  and  $R$  then

$$\frac{1}{2} + \frac{K-1}{K} \cdot \frac{L(7L-6)}{4(L-1)^2} + \frac{3LR}{|S|(L-1)} < 1$$

and so  $|Q_0^+| < |S|$  so that  $Q_0^+$  embeds in  $S$ . In this case, the endpoints  $p^+$  and  $q^+$  of  $Q_0^+$  in  $S$  are at distance

$$d_S(p^+, q^+) \geq \left( \frac{1}{2} - \frac{K-1}{K} \cdot \frac{L(7L-6)}{4(L-1)^2} - \frac{3LR}{|S|(L-1)} \right) |S|$$

but we also have

$$d_X(\alpha(p^+), \alpha(q^+)) < \frac{1}{L}|Q_0^+| - C \leq \frac{1}{L} \left( \frac{1}{2} + \frac{K-1}{K} \cdot \frac{L(7L-6)}{4(L-1)^2} + \frac{3LR}{|S|(L-1)} \right) |S| - C$$

which, by Lemma 3.1 and  $C \geq 4R$ , implies

$$\frac{1}{L} \left( \frac{1}{2} + \frac{K-1}{K} \cdot \frac{L(7L-6)}{4(L-1)^2} + \frac{3LR}{|S|(L-1)} \right) |S| > \left( \frac{1}{2} - \frac{K-1}{K} \cdot \frac{L(7L-6)}{4(L-1)^2} - \frac{3LR}{|S|(L-1)} \right) |S| - \frac{K-1}{K} \cdot \frac{|S|}{2}$$

which is equivalent to the inequality

$$(¶) \quad \frac{K-1}{K} \cdot \frac{9L^2 - 3L - 4}{2(L-1)^2} + \frac{6R(L+1)}{|S|(L-1)} > \frac{L-1}{L}.$$

By hypothesis,

$$K < \frac{L(9L^2 - 3L - 4)}{7L^3 + 3L^2 - 10L + 2}$$

which is equivalent to

$$\frac{K-1}{K} \cdot \frac{9L^2 - 3L - 4}{2(L-1)^2} < \frac{L-1}{L}$$

so if  $|S| > M'''$  for some  $M'''$  depending only on  $K, L$  and  $R$  then we would have

$$\frac{K-1}{K} \cdot \frac{9L^2 - 3L - 4}{2(L-1)^2} + \frac{6R(L+1)}{|S|(L-1)} < \frac{L-1}{L}$$

which contradicts (¶).

Therefore, if  $|S| > M = \max\{M', M'', M'''\}$ , which depends only on  $K, L$  and  $R$  then assuming the existence of a  $j$  for which  $Q \not\subset P_{0,j}^\circ$  leads us to a contradiction.  $\square$

**Claim 4.14** *Let  $K > 1, L > 1$  and  $R \geq 0$ . There exists an  $M > 0$  such that if  $X$  is an  $R$ -rough geodesic metric space,  $\alpha: S \rightarrow X$  is a  $1/K$ -almost isometric  $R$ -circle with  $|S| > M$  and*

$$K < \frac{L(9L^2 - 3L - 4)}{7L^3 + 3L^2 - 10L + 2},$$

*and  $C \geq 4R$  then any  $(L, C)$ -greedy tightening sequence for  $\alpha$  is completely disjoint.*

**Proof** Consider an  $(L, C)$ -greedy tightening sequence for  $\alpha$  with notation as above. For the sake of finding a contradiction, suppose  $j \geq 1$  is the least integer with  $Q_j \not\subset P_{0,j}^\circ$ . As above we view the  $\bar{Q}_i$  with  $i < j$  as disjoint segments of  $S_j$  with  $S_j \setminus \bigcup_{i=0}^{j-1} \bar{Q}_m = P_{0,j}^\circ$ .

Since  $Q_j \not\subset P_{0,j}^\circ$ , we have  $\bar{Q}_m \cap Q_j \neq \emptyset$ , for some  $m < j$ . Recall that  $(p_j, q_j)$  is the limit of a sequence  $(p_j^{(n)}, q_j^{(n)})_n$  in  $J_j$ . For each  $n$ , let  $Q_j^{(n)}$  be a geodesic segment between  $p_j^{(n)}$  and  $q_j^{(n)}$  in  $S_j$ . By Claim 4.13,  $Q_j^{(n)} \subset P_{0,j}^\circ$ , so  $\bar{Q}_m \cap Q_j \subset \{p_m, q_m\} \cap \{p_j, q_j\}$ . Without loss of generality, we may assume  $q_m = p_j$ . See Figure 6.

Since  $Q_j^{(n)} \subset P_{0,j}^\circ$ , we may think of the  $Q_j^{(n)}$  as segments of  $S_m$ , by the embedding  $P_{0,j}^\circ \hookrightarrow S_m$ . Each  $Q_j^{(n)}$  is a geodesic segment in  $S_m$  since the complementary segment of  $Q_j^{(n)}$  in  $S_m$  is even longer than the complementary segment of  $Q_j^{(n)}$  in  $S_j$ . Thus  $(p_j^{(n)}, q_j^{(n)})_n$  is a sequence in  $J_m$ . For each  $n$ , let  $Q_m^{(n)}$  be a segment between  $p_m^{(n)}$  and  $q_m^{(n)}$  such that  $(Q_m^{(n)})_n$  converges to  $Q_m$  in Hausdorff distance. The




Figure 6: A greedy tightening sequence that is disjoint up to  $j$  but not up to  $j + 1$  with  $Q_j$  intersecting the prior  $Q_i$  only at endpoints, as in the proof of Claim 4.14.

circular orders on the triples  $(p_m^{(n)}, q_m^{(n)}, q_j^{(n)})$  and  $(p_m^{(n)}, p_j^{(n)}, q_j^{(n)})$  are eventually constant and equal. Let  $(A^{(n)})_n$  be a sequence of segments in  $S_m$  from  $p_m^{(n)}$  to  $q_j^{(n)}$  such that  $A^{(n)}$  eventually contains  $q_m^{(n)}$  or, equivalently, eventually contains  $p_j^{(n)}$ .

Let  $\varepsilon > 0$  satisfy  $\varepsilon < \frac{1}{3}|Q_j|$  and  $\varepsilon \leq LR/(L + 1)$ . Then, for  $n$  large enough,

$$|A^{(n)}| \geq |Q_m^{(n)}| + |Q_j^{(n)}| - d_{S_m}(q_m^{(n)}, p_j^{(n)}) > |Q_m| - \varepsilon + |Q_j| - \varepsilon - \varepsilon = |Q_m| + |Q_j| - 3\varepsilon > |Q_m|$$

and

$$\begin{aligned} d_X(\alpha_m(p_m^{(n)}), \alpha_m(q_j^{(n)})) &\leq d_X(\alpha_m(p_m^{(n)}), \alpha_m(q_m^{(n)})) + d_X(\alpha_m(q_m^{(n)}), \alpha_m(p_j^{(n)})) + d_X(\alpha_m(p_j^{(n)}), \alpha_m(q_j^{(n)})) \\ &< \frac{1}{L}d_{S_m}(p_m^{(n)}, q_m^{(n)}) - C + d_{S_m}(q_m^{(n)}, p_j^{(n)}) + R + \frac{1}{L}d_{S_m}(p_j^{(n)}, q_j^{(n)}) - C \\ &< \frac{1}{L}|Q_m| + \varepsilon - C + \varepsilon + R + \frac{1}{L}|Q_j| + \varepsilon - C \\ &= \frac{1}{L}(|Q_m| + |Q_j|) - 2C + R + 3\varepsilon \\ &< \frac{1}{L}(|A^{(n)}| + 3\varepsilon) - 2C + R + 3\varepsilon \\ &= \frac{1}{L}|A^{(n)}| - 2C + R + \frac{3(L + 1)}{L} \cdot \varepsilon \\ &\leq \frac{1}{L}|A^{(n)}| - C \end{aligned}$$

since  $C \geq 4R$ . So, if  $A^{(n)}$  is a geodesic segment for arbitrarily large  $n$  then, for some  $n$ , we would have  $(p_m^{(n)}, q_j^{(n)}) \in J_m$  and  $d_{S_m}(p_m^{(n)}, q_j^{(n)}) > |Q_m| = s_m$ , which is a contradiction. Thus eventually

$$|Q_m^{(n)}| + |Q_j^{(n)}| = |A^{(n)}| > \frac{1}{2}|S_m|$$

and so  $|Q_m| + |Q_j| \geq \frac{1}{2}S_m$ .

Then, by Claim 4.9,

$$\frac{S_m}{2} \leq 2 \left( \frac{K-1}{K} \cdot \frac{L}{L-1} \right) \frac{|S|}{2}$$

while

$$\begin{aligned} \frac{|S_m|}{2} &\geq \frac{|S|}{2} - \frac{1}{2} \sum_{i < m} |Q_i| \geq \frac{|S|}{2} - \left( \frac{K-1}{K} \cdot \frac{L(3L-2)}{2(L-1)^2} \right) \frac{|S|}{2} - \frac{3LR}{L-1} \\ &= \left( 1 - \frac{K-1}{K} \cdot \frac{L(3L-2)}{2(L-1)^2} - \frac{6LR}{|S|(L-1)} \right) \frac{|S|}{2} \end{aligned}$$

by Claim 4.10. Combining these we obtain

$$\frac{K-1}{K} \cdot \frac{L(7L-6)}{2(L-1)^2} + \frac{6LR}{|S|(L-1)} \geq 1$$

but, by hypothesis and Claim 4.12,

$$K < \frac{L(7L-6)}{5L^2-2L-2},$$

which is equivalent to

$$\frac{K-1}{K} \cdot \frac{L(7L-6)}{2(L-1)^2} < 1,$$

so, if  $|S|$  is large enough (depending only on  $K$ ,  $L$  and  $R$ ) then we have a contradiction.  $\square$

## 5 The fine Milnor–Schwarz lemma

The fine Milnor–Schwarz lemma is a refinement of the Milnor–Schwarz lemma that gives finer control on the multiplicative constant of the quasi-isometry. In this section we will state and prove this version of the Milnor–Schwarz lemma. As a consequence we will prove that every strongly shortcut group has a strongly shortcut Cayley graph. A corresponding statement should hold for any rough approximability invariant of metric spaces.

Let  $(X, d)$  be a rough geodesic metric space. Let  $G$  be a group acting coboundedly on  $X$  by isometries. Let

$$S_{x_0,t} = \{g \in G \mid d(x_0, gx_0) \leq t\}$$

for  $x_0 \in X$  and  $t \in \mathbb{R}_{\geq 0}$ .

**Remark 5.1** If the action of  $G$  is metrically proper then the  $S_{x_0,t}$  are finite.

Let  $\Gamma_{x_0,t}$  be the graph with vertex set  $G$  and with an edge of length  $t$  between  $g$  and  $g'$  whenever  $g' = gs$  for some  $s \in S_{x_0,t}$ . So when  $S_{x_0,t}$  generates  $G$ , the graph  $\Gamma_{x_0,t}$  is the Cayley graph of  $G$  for the generating set  $S_{x_0,t}$  with edges scaled by  $t$ . Let  $d_{x_0,t}$  be the graph metric on  $\Gamma_{x_0,t}$  where we set  $d_{x_0,t}(g, h) = \infty$  when  $g$  and  $h$  are in different components of  $\Gamma_{x_0,t}$ . Then, when  $S_{x_0,t}$  generates  $G$ , the metric  $d_{x_0,t}$  is the word metric on  $G$  for the generating set  $S_{x_0,t}$  scaled by  $t$ . Let  $f_{x_0,t}: (G, d_{x_0,t}) \rightarrow (X, d)$  be defined

by  $f_{x_0,t}(g) = gx_0$ . Let  $K_{x_0,t}$  be the infimum of all  $K \geq 1$  for which there exists some  $C_K > 0$  such that  $f_{x_0,t}$  is a  $(K, C_K)$ -quasi-isometry. Note that if  $f_{x_0,t}$  is not a quasi-isometry (eg if  $S_{x_0,t}$  does not generate  $G$ ) then  $K_{x_0,t} = \infty$ .

**Lemma 5.2** (fine Milnor–Schwarz lemma) *Let  $(X, d)$  be a rough geodesic metric space. Let  $G$  be a group acting coboundedly on  $X$  by isometries. Let  $x_0 \in X$  and let  $K_{x_0,t}$  be defined as above for  $t \in \mathbb{R}_{\geq 0}$ . Then  $K_{x_0,t} \rightarrow 1$  as  $t \rightarrow \infty$ .*

**Proof** Let  $g \in G$ . We will prove that  $d(f_{x_0,t}(1), f_{x_0,t}(g)) \leq d_{x_0,t}(1, g)$ . If  $d_{x_0,t}(1, g) = \infty$  there is nothing to show and so  $d_{x_0,t}(1, g) = Mt$  for some  $M \in \mathbb{N}_{\geq 0}$  and there is a combinatorial path defined by

$$1 = g_0, g_1, g_2, \dots, g_M = g$$

in  $\Gamma_{x_0,t}$ . The  $g_{i-1}^{-1}g_i$  are contained in  $S_{x_0,t}$  and so, by the triangle inequality,

$$\begin{aligned} d(f_{x_0,t}(1), f_{x_0,t}(g)) &= d(x_0, gx_0) \\ &\leq d(x_0, g_1x_0) + d(g_1x_0, g_2x_0) + \dots + d(g_{M-1}x_0, gx_0) \\ &= d(x_0, g_1x_0) + d(x_0, g_1^{-1}g_2x_0) + \dots + d(x_0, g_{M-1}^{-1}gx_0) \\ &\leq Mt \\ &= d_{x_0,t}(1, g). \end{aligned}$$

We now establish a lower bound on  $D = d(f_{x_0,t}(1), f_{x_0,t}(g))$ . Since  $G$  acts coboundedly on  $X$ , the orbit  $Gx_0$  is a quasi-onto subspace of  $X$ . Hence  $Gx_0$  is roughly isometric to  $X$  and so  $Gx_0$  is also a rough geodesic metric space. Let  $R$  be the rough geodesicity constant of  $Gx_0$ . Let  $\alpha: [0, D] \rightarrow Gx_0$  be an  $R$ -rough geodesic from  $f_{x_0,t}(1) = x_0$  to  $f_{x_0,t}(g) = gx_0$ . Assume that  $t > R$  and subdivide  $[0, D]$  into at most  $\lceil D/(t - R) \rceil$  segments of length at most  $t - R$ . Let  $0 = a_0 < a_1 < a_2 < \dots < a_M = D$  be the endpoints of the segments and let  $g_i x_0 = \alpha(a_i)$ , for each  $i$ , with  $g_0 = 1$  and  $g_M = g$ . Then

$$d(x_0, g_i^{-1}g_{i+1}x_0) = d(g_i x_0, g_{i+1}x_0) = d(\alpha(a_i), \alpha(a_{i+1})) \leq |a_i - a_{i+1}| + R \leq t$$

for each  $i$ . Thus  $g_i^{-1}g_{i+1} \in S_{x_0,t}$ , for each  $i$ , and so

$$1 = g_0, g_1, g_2, \dots, g_{M-1}, g_M = g$$

defines a combinatorial path in  $\Gamma_{x_0,t}$ . Hence

$$\begin{aligned} d_{x_0,t}(1, g) &\leq tM \leq t \left\lceil \frac{D}{t - R} \right\rceil \\ &= t \left\lceil \frac{d(f_{x_0,t}(1), f_{x_0,t}(g))}{t - R} \right\rceil \\ &\leq t \left( \frac{d(f_{x_0,t}(1), f_{x_0,t}(g))}{t - R} + 1 \right) \\ &= \frac{t}{t - R} d(f_{x_0,t}(1), f_{x_0,t}(g)) + t \end{aligned}$$

and so we have  $((t - R)/t)d_{x_0,t}(1, g) - (t - R) \leq d(f_{x_0,t}(1), f_{x_0,t}(g))$ .

For  $g, g' \in G$ ,

$$d(f_{x_0,t}(g), f_{x_0,t}(g')) = d(gx_0, g'x_0) = d(x_0, g^{-1}g'x_0) = d(f_{x_0,t}(1), f_{x_0,t}(g^{-1}g'))$$

and  $d_{x_0,t}(g, g') = d_{x_0,t}(1, g^{-1}g')$ , so

$$\frac{t-R}{t}d_{x_0,t}(g, g') - (t-R) \leq d(f_{x_0,t}(g), f_{x_0,t}(g')) \leq d_{x_0,t}(g, g');$$

hence  $1 \leq K_{x_0,t} \leq (t-R)/t$  when  $t > R$ . This implies that  $K_{x_0,t} \rightarrow 1$  as  $t \rightarrow \infty$ . □

**Corollary 5.3** (Milnor–Schwarz lemma) *If the  $G$ -action on  $X$  is metrically proper then  $G$  is finitely generated and, for any finite generating set  $S$ , the map*

$$f_S : (G, d_S) \rightarrow X, \quad g \mapsto gx_0,$$

*is a quasi-isometry, where  $d_S$  is the word metric for the generating set  $S$ .*

**Proof** By Lemma 5.2, if  $t$  is large enough then  $K_{x_0,t} < \infty$ . Thus  $S_{x_0,t}$  generates  $G$ . By Remark 5.1, the generating set  $S_{x_0,t}$  is finite so  $G$  is finitely generated. Since  $K_{x_0,t} < \infty$  and since scaling the metric  $d_S$  by a factor of  $t$  preserves the quasi-isometry type, the map  $f_S$  is a quasi-isometry for  $S = S_{x_0,t}$ . But the identity map on  $G$  is a bi-Lipschitz equivalence from  $(G, d_{S'})$  to  $(G, d_{S''})$  where  $S'$  and  $S''$  are any two finite generating sets and  $d_{S'}$  and  $d_{S''}$  are the corresponding word metrics. Thus  $f_S$  is a quasi-isometry for any generating set  $S$ . □

**Corollary 5.4** *Let  $G$  be a group. If  $G$  acts metrically properly and coboundedly on a strongly shortcut rough geodesic metric space  $X$  then  $G$  has a finite generating set  $S$  for which the Cayley graph of  $(G, S)$  is strongly shortcut. In particular, the group  $G$  is strongly shortcut.*

**Proof** By Corollary 3.10, there exists an  $L_X > 1$  such that whenever  $C > 0$  and  $Y$  is a rough geodesic metric space and  $f : Y \rightarrow X$  is an  $(L_X, C)$ -quasi-isometry up to scaling, then  $Y$  is strongly shortcut. But, by Lemma 5.2, there is a Cayley graph  $\Gamma$  of  $G$ , a  $C > 0$  and an  $(L_X, C)$ -quasi-isometry up to scaling  $f : \Gamma \rightarrow X$ . So  $\Gamma$  is strongly shortcut as a rough geodesic metric space. Then, by Remark 2.5, the Cayley graph  $\Gamma$  is strongly shortcut as a graph. □

**Corollary 5.5** *Let  $G$  be a group. The following conditions are equivalent:*

- (1)  $G$  is strongly shortcut.
- (2)  $G$  acts metrically properly and coboundedly on a strongly shortcut rough geodesic metric space.
- (3)  $G$  has a finite generating set  $S$  for which the Cayley graph of  $(G, S)$  is strongly shortcut.

## 6 Asymptotically CAT(0) spaces

In this section we will apply the characterizations of Section 3 to prove that asymptotically CAT(0) rough geodesic metric spaces are strongly shortcut. By the results of Section 5, this will imply that asymptotically CAT(0) groups are strongly shortcut.

Asymptotically CAT(0) spaces and groups were first introduced and studied by Kar [2011]. A metric space  $X$  is *asymptotically* CAT(0) if every asymptotic cone of  $X$  is CAT(0). A group is *asymptotically* CAT(0) if it acts properly and cocompactly on an asymptotically CAT(0) proper geodesic metric space. (Note that the condition that the action be on a proper metric space does not make this definition more restrictive than the definition given by Kar [2011] since the definition of proper action in Kar [2011, page 77] seems to be that of Bridson and Haefliger [1999, Section I.8.2] and any geodesic metric space admitting a cocompact action that is proper by this more restricted definition is a proper metric space.) For an introduction to CAT(0) geodesic metric spaces, see Bridson and Haefliger [1999].

**Theorem 6.1** *Asymptotically CAT(0) rough geodesic metric spaces are strongly shortcut.*

**Proof** By uniqueness of geodesics in CAT(0) geodesic metric spaces, there is no isometric copy of a Riemannian circle in the asymptotic cone of an asymptotically CAT(0) metric space  $X$ . So, by Theorem 3.8, any asymptotically CAT(0) rough geodesic metric space is strongly shortcut.  $\square$

**Theorem 6.2** *Asymptotically CAT(0) groups are strongly shortcut.*

**Proof** A proper and cocompact action on a proper metric space is metrically proper and cobounded. So, by Theorem 6.1, any asymptotically CAT(0) group  $G$  acts metrically properly and coboundedly on a strongly shortcut geodesic metric space. Thus, by Corollary 5.5, the group  $G$  is strongly shortcut.  $\square$

## References

- [Bridson and Haefliger 1999] **MR Bridson, A Haefliger**, *Metric spaces of non-positive curvature*, Grundle. Math. Wissen. 319, Springer (1999) MR Zbl
- [Druţu and Kapovich 2018] **C Druţu, M Kapovich**, *Geometric group theory*, Amer. Math. Soc. Colloq. Publ. 63, Amer. Math. Soc., Providence, RI (2018) MR Zbl
- [Gromov 1987] **M Gromov**, *Hyperbolic groups*, from “Essays in group theory” (S M Gersten, editor), Math. Sci. Res. Inst. Publ. 8, Springer (1987) 75–263 MR Zbl
- [Gromov 1993] **M Gromov**, *Asymptotic invariants of infinite groups*, from “Geometric group theory, II” (G A Niblo, M A Roller, editors), Lond. Math. Soc. Lect. Note Ser. 182, Cambridge Univ. Press (1993) 1–295 MR Zbl
- [Haettel et al. 2023] **T Haettel, N Hoda, H Petyt**, *Coarse injectivity, hierarchical hyperbolicity and semihyperbolicity*, *Geom. Topol.* 27 (2023) 1587–1633 MR Zbl
- [Hoda 2022] **N Hoda**, *Shortcut graphs and groups*, *Trans. Amer. Math. Soc.* 375 (2022) 2417–2458 MR Zbl
- [Hoda and Krishna MS 2023] **N Hoda, S Krishna MS**, *Relatively hyperbolic groups with strongly shortcut parabolics are strongly shortcut*, *Math. Proc. Cambridge Philos. Soc.* 175 (2023) 367–380 MR Zbl
- [Kar 2011] **A Kar**, *Asymptotically CAT(0) groups*, *Publ. Mat.* 55 (2011) 67–91 MR Zbl
- [Kasparov and Skandalis 1994] **G Kasparov, G Skandalis**, *Groupes “boliques” et conjecture de Novikov*, *C. R. Acad. Sci. Paris Sér. I Math.* 319 (1994) 815–820 MR Zbl

- [Kasparov and Skandalis 2003] **G Kasparov, G Skandalis**, *Groups acting properly on “bolic” spaces and the Novikov conjecture*, Ann. of Math. 158 (2003) 165–206 MR Zbl
- [Lafforgue 2002] **V Lafforgue**, *K–théorie bivariante pour les algèbres de Banach et conjecture de Baum–Connes*, Invent. Math. 149 (2002) 1–95 MR Zbl
- [Le Donne and Paddeu 2023] **E Le Donne, N Paddeu**, *Escape from compact sets of normal curves in Carnot groups*, preprint (2023) arXiv 2304.03205
- [Mineyev and Yu 2002] **I Mineyev, G Yu**, *The Baum–Connes conjecture for hyperbolic groups*, Invent. Math. 149 (2002) 97–122 MR Zbl
- [Papasoglu 1996] **P Papasoglu**, *On the asymptotic cone of groups satisfying a quadratic isoperimetric inequality*, J. Differential Geom. 44 (1996) 789–806 MR Zbl
- [Riley 2003] **T R Riley**, *Higher connectedness of asymptotic cones*, Topology 42 (2003) 1289–1352 MR Zbl

*Department of Mathematics, Cornell University  
Ithaca, NY, United States*

nima.hoda@mail.mcgill.ca

Received: 16 October 2020      Revised: 14 November 2022



# ALGEBRAIC & GEOMETRIC TOPOLOGY

msp.org/agt

## EDITORS

### PRINCIPAL ACADEMIC EDITORS

John Etnyre  
etnyre@math.gatech.edu  
Georgia Institute of Technology

Kathryn Hess  
kathryn.hess@epfl.ch  
École Polytechnique Fédérale de Lausanne

### BOARD OF EDITORS

Julie Bergner	University of Virginia jeb2md@eservices.virginia.edu	Christine Lescop	Université Joseph Fourier lescop@ujf-grenoble.fr
Steven Boyer	Université du Québec à Montréal cohf@math.rochester.edu	Robert Lipshitz	University of Oregon lipshitz@uoregon.edu
Tara E Brendle	University of Glasgow tara.brendle@glasgow.ac.uk	Norihiko Minami	Yamato University minami.norihiko@yamato-u.ac.jp
Indira Chatterji	CNRS & Univ. Côte d'Azur (Nice) indira.chatterji@math.cnrs.fr	Andrés Navas	Universidad de Santiago de Chile andres.navas@usach.cl
Alexander Dranishnikov	University of Florida dranish@math.ufl.edu	Robert Oliver	Université Paris 13 bobol@math.univ-paris13.fr
Tobias Ekholm	Uppsala University, Sweden tobias.ekholm@math.uu.se	Jessica S Purcell	Monash University jessica.purcell@monash.edu
Mario Eudave-Muñoz	Univ. Nacional Autónoma de México mario@matem.unam.mx	Birgit Richter	Universität Hamburg birgit.richter@uni-hamburg.de
David Futер	Temple University dfuter@temple.edu	Jérôme Scherer	École Polytech. Féd. de Lausanne jerome.scherer@epfl.ch
John Greenlees	University of Warwick john.greenlees@warwick.ac.uk	Vesna Stojanoska	Univ. of Illinois at Urbana-Champaign vesna@illinois.edu
Ian Hambleton	McMaster University ian@math.mcmaster.ca	Zoltán Szabó	Princeton University szabo@math.princeton.edu
Matthew Hedden	Michigan State University mhedden@math.msu.edu	Maggy Tomova	University of Iowa maggy-tomova@uiowa.edu
Hans-Werner Henn	Université Louis Pasteur henn@math.u-strasbg.fr	Chris Wendl	Humboldt-Universität zu Berlin wendl@math.hu-berlin.de
Daniel Isaksen	Wayne State University isaksen@math.wayne.edu	Daniel T Wise	McGill University, Canada daniel.wise@mcgill.ca
Thomas Koberda	University of Virginia thomas.koberda@virginia.edu	Lior Yanovski	Hebrew University of Jerusalem lior.yanovski@gmail.com
Markus Land	LMU München markus.land@math.lmu.de		

---

See inside back cover or [msp.org/agt](https://msp.org/agt) for submission instructions.

---

The subscription price for 2024 is US \$705/year for the electronic version, and \$1040/year (+\$70, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Algebraic & Geometric Topology is indexed by Mathematical Reviews, Zentralblatt MATH, Current Mathematical Publications and the Science Citation Index.


---

Algebraic & Geometric Topology (ISSN 1472-2747 printed, 1472-2739 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840.

---

AGT peer review and production are managed by EditFlow<sup>®</sup> from MSP.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<https://msp.org/>

© 2024 Mathematical Sciences Publishers

# ALGEBRAIC & GEOMETRIC TOPOLOGY

Volume 24 Issue 6 (pages 2971–3570) 2024

---

Definition of the cord algebra of knots using Morse theory	2971
ANDREAS PETRAK	
An analogue of Milnor's invariants for knots in 3-manifolds	3043
MIRIAM KUZBARY	
Wall-crossing from Lagrangian cobordisms	3069
JEFF HICKS	
Foliated open books	3139
JOAN E LICATA and VERA VÉRTESI	
Algebraic and Giroux torsion in higher-dimensional contact manifolds	3199
AGUSTIN MORENO	
Locally equivalent Floer complexes and unoriented link cobordisms	3235
ALBERTO CAVALLO	
Strongly shortcut spaces	3291
NIMA HODA	
Extendable periodic automorphisms of closed surfaces over the 3-sphere	3327
CHAO WANG and WEIBIAO WANG	
Bounding the Kirby–Thompson invariant of spun knots	3363
ROMÁN ARANDA, PUTTIPONG PONGTANAPAIAN, SCOTT A TAYLOR and SUIXIN (CINDY) ZHANG	
Dynamics of veering triangulations: infinitesimal components of their flow graphs and applications	3401
IAN AGOL and CHI CHEUK TSANG	
L-spaces, taut foliations and the Whitehead link	3455
DIEGO SANTORO	
Horizontal decompositions, I	3503
PAOLO LISCA and ANDREA PARMA	
The homology of a Temperley–Lieb algebra on an odd number of strands	3527
ROBIN J SROKA	
Hyperbolic homology 3-spheres from drum polyhedra	3543
RAQUEL DÍAZ and JOSÉ L ESTÉVEZ	