

AG
T

*Algebraic & Geometric
Topology*

Volume 24 (2024)

**On products of beta and gamma elements
in the homotopy of the first Smith–Toda spectrum**

KATSUMI SHIMOMURA
MAO-NO-SUKE SHIMOMURA

On products of beta and gamma elements in the homotopy of the first Smith–Toda spectrum

KATSUMI SHIMOMURA
MAO-NO-SUKE SHIMOMURA

We determine the first cohomology of the monochromatic comodule M_2^1 at an odd prime, and apply the results to show nontrivialities of some products of beta and gamma elements in the homotopy groups of the Smith–Toda spectrum $V(1)$. The cohomology for a prime greater than three was previously determined by the first author. Here we verify them and determine the cohomology at the prime 3 by elementary calculation. The cohomology will be a stepping stone for computing the cohomology of the monochromatic comodule M_0^3 , which we hope to determine for a long time.

[55Q45](#); [55Q51](#), [55T15](#)

1 Introduction

Let p be an odd prime number and $\mathcal{S}_{(p)}$ denote the stable homotopy category of p –local spectra. Let $S \in \mathcal{S}_{(p)}$ denote the sphere spectrum. Then the mod p Moore spectrum M and the first Smith–Toda spectrum $V(1)$ are given by the cofiber sequences

$$(1.1) \quad S \xrightarrow{p} S \xrightarrow{i} M \xrightarrow{j} \Sigma S \quad \text{and} \quad \Sigma^q M \xrightarrow{\alpha} M \xrightarrow{i_1} V(1) \xrightarrow{j_1} \Sigma^{q+1} M.$$

Here $p \in \pi_0(S) \cong \mathbb{Z}_{(p)}$, and $\alpha \in [M, M]_q$ denotes the Adams map. Hereafter we put

$$q = 2p - 2 \in \mathbb{Z}.$$

To study the homotopy groups $\pi_*(X)$ of a spectrum X , we adopt the Adams–Novikov spectral sequence

$$(1.2) \quad E_2^{s,t}(X) = H^{s,t} \text{BP}_*(X) \Rightarrow \pi_{t-s}(X).$$

Hereafter we abbreviate as

$$H^{s,t} M = \text{Ext}_{\text{BP}_*(\text{BP})}^{s,t}(\text{BP}_*, M)$$

for a $\text{BP}_*(\text{BP})$ –comodule M over the Hopf algebroid

$$(1.3) \quad (\text{BP}_*, \text{BP}_*(\text{BP})) = (\mathbb{Z}_{(p)}[v_1, v_2, \dots], \text{BP}_*[t_1, t_2, \dots])$$

based on the Brown–Peterson spectrum $\text{BP} \in \mathcal{S}_{(p)}$. We note that the v_i are Hazewinkel’s generators and the degrees of v_i and t_i are $|v_i| = 2p^i - 2 = |t_i|$; see Miller, Ravenel and Wilson [2, (1.1)].

Let

$$(1.4) \quad I_n = (p, v_1, \dots, v_{n-1}) \quad \text{and} \quad J_j = (p, v_1, v_2^j)$$

for $v_0 = p$ denote the invariant ideals of BP_* . Since $\mathrm{BP}_*(\alpha) = v_1$, the cofiber sequences (1.1) induce the short exact sequences

$$(1.5) \quad 0 \rightarrow \mathrm{BP}_* \xrightarrow{p} \mathrm{BP}_* \xrightarrow{i_*} \mathrm{BP}_*/I_1 \rightarrow 0 \quad \text{and} \quad 0 \rightarrow \mathrm{BP}_*/I_1 \xrightarrow{v_1} \mathrm{BP}_*/I_1 \xrightarrow{(i_1)_*} \mathrm{BP}_*/I_2 \rightarrow 0$$

along with the isomorphisms

$$\mathrm{BP}_*(S) = \mathrm{BP}_*, \quad \mathrm{BP}_*(M) = \mathrm{BP}_*/I_1 \quad \text{and} \quad \mathrm{BP}_*(V(1)) = \mathrm{BP}_*/I_2.$$

Furthermore, we have a short exact sequence

$$(1.6) \quad 0 \rightarrow \mathrm{BP}_*/I_2 \xrightarrow{v_2^j} \mathrm{BP}_*/I_2 \xrightarrow{\bar{i}_j} \mathrm{BP}_*/J_j \rightarrow 0$$

for $j \geq 1$. We denote by

$$\delta_0: H^s \mathrm{BP}_*/I_1 \rightarrow H^{s+1} \mathrm{BP}_*, \quad \delta_1: H^s \mathrm{BP}_*/I_2 \rightarrow H^{s+1} \mathrm{BP}_*/I_1, \quad \bar{\delta}_j: H^s \mathrm{BP}_*/J_j \rightarrow H^{s+1} \mathrm{BP}_*/I_2$$

the connecting homomorphisms associated to the short exact sequences (1.5) and (1.6). We define the Greek letter elements by

$$\begin{aligned} \bar{\beta}'_s &= \delta_1(v_2^s) \in E_2^1(M) = H^1 \mathrm{BP}_*/I_1 & \text{for } v_2^s \in H^0 \mathrm{BP}_*/I_2, \\ \bar{\beta}_s &= \delta_0 \delta_1(v_2^s) \in E_2^2(S) = H^2 \mathrm{BP}_* & \text{for } v_2^s \in H^0 \mathrm{BP}_*/I_2, \\ \bar{\gamma}''_{s/j} &= \bar{\delta}_j(v_3^s) \in E_2^1(V(1)) = H^1 \mathrm{BP}_*/I_2 & \text{for } v_3^s \in H^0 \mathrm{BP}_*/J_j, \end{aligned}$$

and $\bar{\gamma}''_s = \bar{\gamma}''_{s/1} \in E_2^1(V(1))$. We notice that $1 \leq j \leq p^n$ if $p^n \mid s$, so that $v_3^s \in H^0 \mathrm{BP}_*/J_j$.

Let \mathbb{Z} and \mathbb{N} denote the set of all integers and its subset consisting of all nonnegative integers, respectively. We denote by $\mathbb{Z}^{(p)} (= \mathbb{Z} \setminus p\mathbb{Z})$ and $\mathbb{N}^{(p)} (= \mathbb{N} \setminus p\mathbb{N})$ the sets of the integers prime to p , and decompose $\mathbb{Z}^{(p)}$ into the three summands

$$\mathbb{Z}^{(p)} = \mathbb{Z}_0 \coprod \mathbb{Z}_1 \coprod \mathbb{Z}_2,$$

for

$$(1.7) \quad \begin{aligned} \mathbb{Z}_0 &= \{s \in \mathbb{Z}^{(p)} : p \nmid (s+1)\}, & \mathbb{Z}_1 &= \{s \in \mathbb{Z}^{(p)} : p^2 \mid (s+1)\}, \\ \mathbb{Z}_2 &= \{s \in \mathbb{Z}^{(p)} : p \mid (s+1) \text{ and } p^2 \nmid (s+1)\}. \end{aligned}$$

We consider subsets of \mathbb{N} :

$$\begin{aligned} 2\mathbb{N}_{>0} &= \{s \in \mathbb{N} : s \text{ is even} \geq 2\}, & \mathbb{N}_1 &= \{s \in \mathbb{N}^{(p)} : p^2 \nmid (s+p+1) \text{ or } p^3 \mid (s+p+1)\}, \\ \overline{2\mathbb{N}} &= \{s \in \mathbb{N} : s \text{ is odd}\}, & \mathbb{N}_2 &= \{s \in \mathbb{N}^{(p)} : p \nmid (s+2) \text{ or } p^3 \mid (s+2)(s+2+p)\}. \end{aligned}$$

Furthermore, we put $\mathbb{Z}_i^+ = \mathbb{Z}_i \cap \mathbb{N}$ for $i = 0, 1, 2$. We introduce the subsets U_1 , U'_1 , U_2 and U'_2 of $\mathbb{N}^{(p)} \times \mathbb{N}$ given by

$$\begin{aligned} U_1 &= (\mathbb{N}^{(p)} \times 2\mathbb{N}) \cup (\mathbb{Z}_0^+ \times \mathbb{N}), \\ U'_1 &= (\mathbb{N}^{(3)} \times \{0\}) \cup (\mathbb{N}_1 \times 2\mathbb{N}_{>0}) \cup ((\mathbb{Z}_0^+ \cap \mathbb{N}_2) \times \mathbb{N}) \cup (\mathbb{Z}_0^+ \times \{1\}), \\ U_2 &= (\mathbb{N}_1 \times 2\mathbb{N}) \cup (((\mathbb{Z}_0^+ \cap \mathbb{N}_2) \cup \mathbb{Z}_1^+) \times \mathbb{N}) \cup (\mathbb{N}^{(p)} \times \{1\}), \\ U'_2 &= (\mathbb{N}_1 \times \{0\}) \cup (\mathbb{N}^{(3)} \times (\{1\} \cup 2\mathbb{N}_{>0})) \cup ((\mathbb{Z}_0^+ \cup \mathbb{Z}_1^+) \times \mathbb{N}). \end{aligned}$$

Our main result is the following:

Theorem 1.8 *Let p be an odd prime. In the Adams–Novikov E_2 -term for computing $\pi_*(V(1))$, $\bar{\beta}_1$ and $\bar{\beta}_2$ act on the gamma elements $\bar{\gamma}''_{sp^r/j}$ (for $(s, r) \in \mathbb{N}^{(p)} \times \mathbb{N}$ and $1 \leq j \leq p^r$) by*

$$\begin{aligned}\bar{\gamma}''_{sp^r/j} \bar{\beta}_1 &\neq 0 \quad \text{for } (s, r) \in U_1 \text{ if } p \geq 5, \text{ and for } (s, r) \in U'_1 \text{ if } p = 3, \\ \bar{\gamma}''_{sp^r/j} \bar{\beta}_2 &\neq 0 \quad \text{for } (s, r) \in U_2 \text{ if } p \geq 5, \text{ and for } (s, r) \in U'_2 \text{ if } p = 3,\end{aligned}$$

in $E_2^3(V(1))$.

There is a way to define $\gamma''_{sp^r/j}$ for $j \leq a_r$ (a_r is defined in (2.7)) so that $v_2^{j-1} \gamma''_{sp^r/j} = \gamma''_{sp^r}$, and the theorem holds for such extended gamma elements. Also $\bar{\beta}_s \equiv \binom{s}{2} v_2^{s-2} \bar{\beta}_2 + s(2-s) v_2^{s-1} \bar{\beta}_1 \pmod{I_2}$ (see Oka and Shimomura [5, Lemma 4.4]), and so

$$\bar{\gamma}''_{sp^r/j} \bar{\beta}_t = \binom{t}{2} \bar{\gamma}''_{sp^r/j-t+2} \bar{\beta}_2 + t(2-t) \bar{\gamma}''_{sp^r/j-t+1} \bar{\beta}_1.$$

Thus Theorem 1.8 implies nontriviality of the products of $\bar{\gamma}''_{sp^r/j}$ and $\bar{\beta}_t$.

The Adams–Novikov differential d_r is 0 if $q \nmid (r-1)$ by the sparseness of the spectral sequence (1.2). This shows that the products in the theorem are not in the image of any differentials d_r . It is well known that the elements $\bar{\beta}_1$ and $\bar{\beta}_2$ converge to the homotopy elements $\beta_1, \beta_2 \in \pi_*(S)$, respectively, in the spectral sequence (1.2) for $X = S$.

Corollary 1.9 *Let p be an odd prime. If $\bar{\gamma}''_{sp^r/j} \in E_2^1(V(1))$ is a permanent cycle detecting $\gamma''_{sp^r/j} \in \pi_*(V(1))$, then $\gamma''_{sp^r/j} \beta_i \neq 0$ for $i = 1, 2$ in the homotopy groups $\pi_*(V(1))$ for (s, r) given in Theorem 1.8.*

Toda [12, Theorem 1] and Oka [4, Theorem 4.2] showed that γ''_s and $\gamma''_{sp/2}$ are permanent cycles for $p \geq 7$.

Corollary 1.10 *Let $p \geq 7$, and r and s be integers with $(s, r) \in \mathbb{N}^{(p)} \times \mathbb{N}$. Then, in $\pi_*(V(1))$,*

$$\gamma''_{sp^r/j} \beta_1 \neq 0 \quad \text{if } r \text{ is even or } p \nmid (s+1),$$

$$\gamma''_{sp^{2r}/j} \beta_2 \neq 0 \quad \text{if } p^2 \nmid (s+p+1) \text{ or } p^3 \mid (s+p+1),$$

$$\gamma''_{sp^{2r+1}/j} \beta_2 \neq 0 \quad \text{for } r \geq 1 \text{ if } p \nmid (s+1)(s+2), p^2 \mid (s+1) \text{ or } p^3 \mid (s+2)(s+2+p),$$

and $\gamma''_{sp/j} \beta_2 \neq 0$, where $j = 1, 2$.

Theorem 1.8 follows from Theorem 2.9, which states the structure of the first cohomology of the monochromatic comodule M_2^1 . The cohomology $H^1 M_2^1$ was determined by the first author [10] based on the computation in [9] at a prime ≥ 5 . Here we determine the cohomology based on elementary calculation at an odd prime. The generators are explicitly given so that we can use the result easily in further computation. This result will be a stepping stone for determining the long-desired cohomology $H^* M_0^3$.

This paper is organized as follows: In Section 2, we state the main result, Theorem 2.9, which gives the structure of $H^1 M_2^1$. In Section 3, we prove Theorems 2.9 and 1.8 assuming Lemma 3.4, whose proof will be given in the Section 6. Section 4 is devoted to introducing some formulas, cochains and relations for the following sections. We refine the elements $x_{3,i}$ given by Miller, Ravenel and Wilson [2, (5.11)] to define x_i , which induce the cochains $y_{s,i}, y'_{s,i} \in \Omega^1 E(3)_*$ in Section 5.

Acknowledgments The authors would like to express their gratitude to the referee for the careful reading of the manuscript and useful suggestions.

2 The structure of $H^1 M_2^1$

In this section, we state the structure of $H^1 M_2^1$ for an odd prime p . The structure was given in [10] for primes $p \geq 5$.

We begin with defining the monochromatic $\mathrm{BP}_*(\mathrm{BP})$ -comodules N_n^s and M_n^s inductively by

$$N_n^0 = \mathrm{BP}_*/I_n, \quad M_n^s = v_{s+n}^{-1} N_n^s,$$

for the ideal I_n in (1.4) and the short exact sequence

$$(2.1) \quad 0 \rightarrow N_n^s \xrightarrow{\iota_n^s} M_n^s \xrightarrow{\kappa_n^s} N_n^{s+1} \rightarrow 0$$

see [2, Section 3.A]. Since BP_* is a $\mathrm{BP}_*(\mathrm{BP})$ -comodule with structure map η_R , the right unit map of the Hopf algebroid $\mathrm{BP}_*(\mathrm{BP})$, these monochromatic comodules have the structure maps induced from η_R .

Let $E(3)$ denote the third Johnson–Wilson spectrum, which yields a Hopf algebroid

$$(E(3)_*, E(3)_*(E(3))) = (\mathbb{Z}_{(p)}[v_1, v_2, v_3, v_3^{-1}], E(3)_* \otimes_{\mathrm{BP}_*} \mathrm{BP}_*(\mathrm{BP}) \otimes_{\mathrm{BP}_*} E(3)_*).$$

Its structure maps are induced from the Hopf algebroid $(\mathrm{BP}_*, \mathrm{BP}_*(\mathrm{BP}))$ in (1.3). Since we have the Miller–Ravenel change of rings theorem

$$H^* M = \mathrm{Ext}_{\mathrm{BP}_*(\mathrm{BP})}^*(\mathrm{BP}_*, M) \cong \mathrm{Ext}_{E(3)_*(E(3))}^*(E(3)_*, E(3)_* \otimes_{\mathrm{BP}_*} M)$$

for a v_3 -local $\mathrm{BP}_*(\mathrm{BP})$ -comodule M [1, Theorem 3.10], we denote the cohomology of an $E(3)_*(E(3))$ -comodule M also by

$$H^s M = \mathrm{Ext}_{E(3)_*(E(3))}^s(E(3)_*, M).$$

By virtue of the change of rings theorem, we denote simply by M_n^s the $E(3)_*(E(3))$ -comodule $E(3)_* \otimes_{\mathrm{BP}_*} M_n^s$. We consider the Ext group as the cohomology group of the cobar complex

$$(2.2) \quad \Omega^s M = M \otimes_{E(3)_*} E(3)_*(E(3)) \otimes_{E(3)_*} \cdots \otimes_{E(3)_*} E(3)_*(E(3))$$

for s factors of $E(3)_*(E(3))$, with well-known differentials $d_r: \Omega^r M \rightarrow \Omega^{r+1} M$; see (4.1).

The cohomology $H^t M_n^s$ of the monochromatic comodules with $s+n=3$ are determined in the following cases (see [8, Theorems 6.3.12 and 6.3.14; 2, Theorem 5.10]):

$$(2.3) \quad \begin{aligned} H^0 M_3^0 &= K(3)_*, \\ H^1 M_3^0 &= K(3)_* \{h_0, h_1, h_2, \zeta_3\}, \\ H^2 M_3^0 &= K(3)_* \{g_i, k_i, b_i, h_i \zeta_3 : i \in \mathbb{Z}/3\}, \\ H^0 M_2^1 &= K(2)_*/k(2)_* \oplus \bigoplus_{i \geq 0, s \in \mathbb{Z}^{(p)}} k(2)_*/(v_2^{a_i}) \{x_i^s / v_2^{a_i}\}. \end{aligned}$$

Indeed, we read off $H^s M_3^0 = K(3)_* \otimes H^s S(3)$ from [8, Proposition 6.2.1], where $S(3)$ is the Hopf algebra defined in [8, Section 6.2]. The cohomology groups $H^* M_3^0$ and $H^0 M_1^2$ for $p \geq 5$ are also determined by Ravenel [8, Theorem 6.3.34] and Nakai [3], respectively. Here

$$k(2)_* = \mathbb{Z}/p[v_2], \quad K(2)_* = \mathbb{Z}/p[v_2, v_2^{-1}] \quad \text{and} \quad K(3)_* = \mathbb{Z}/p[v_3, v_3^{-1}],$$

where $K(3)_* = E(3)_*/I_3 = M_3^0$. The elements $x_i (= x_{3,i})$ are introduced in [2, (5.11)] and are such that $x_i \equiv v_3^{p^i} \pmod{I_3}$ (see Lemma 5.1), and the generators h_i, ζ_3, g_i, k_i and b_i are defined by cocycles in the cobar complex $\Omega^* E(3)_*/I_3$ as follows:

$$(2.4) \quad h_i = [t_1^{p^i}], \quad \zeta_3 = [Z], \quad g_i = [G_i], \quad k_i = [K_i] \quad \text{and} \quad b_i = [b_{1,i}].$$

Hereafter $[x]$ denotes the cohomology class represented by a cocycle x , and the representatives in (2.4) are defined by

$$(2.5) \quad \begin{aligned} Z &= -v_3^{-1} ct_3 + v_3^{-p} t_3^p + v_3^{-p^2} t_3^{p^2} - v_3^{-p} t_1^p t_2^{p^2}, & G_i &= t_1^{p^i} \otimes t_2^{p^i} + \frac{1}{2} t_1^{2p^i} \otimes t_1^{p^{i+1}}, \\ K_i &= t_2^{p^i} \otimes t_1^{p^{i+1}} + \frac{1}{2} t_1^{p^i} \otimes t_1^{2p^{i+1}}, & b_{1,i} &= \sum_{k=1}^{p-1} \frac{1}{p} \binom{p}{k} t_1^{kp^i} \otimes t_1^{(p-k)p^i}. \end{aligned}$$

Here ct_3 is the Hopf conjugation of t_3 (see Lemma 4.3). We notice that G_i, K_i and $b_{1,i}$ are also cocycles of $\Omega^* E(3)_*/I_2$, and of $\Omega^* \text{BP}_*/I_2$ in [2, (1.9)].

Remark 2.6 The generators g_i and k_i in (2.3) are given by the Massey products $\langle h_i, h_{i+1}, h_i \rangle$ and $\langle h_{i+1}, h_{i+1}, h_i \rangle$, respectively, in [8, Theorem 6.3.4]. These are represented by cocycles

$$G_i'' = t_2^{p^i} \otimes t_1^{p^{i+1}} + t_1^{p^i} \otimes ct_2^{p^i}$$

and K_i' in (4.20) in the cobar complex $\Omega^* E(3)_*/I_2$, since these Massey products have no indeterminacy. By (4.21), K_i' is homologous to K_i . Also $d_1(t_1^{p^i} t_2^{p^i}) = -2G_i - G_i''$, and G_i'' is homologous to $-2G_i$. Since p is odd, we may replace generators g_i and k_i by $[G_i]$ and $[K_i]$, and set as in (2.4).

We introduce integers $e(n), a_n, j_{s,n}$ and $j'_{s,n}$ for integers $n(\geq 0)$ and s by

$$(2.7) \quad \begin{aligned} e(n) &= \frac{p^n - 1}{p - 1} \quad \text{for } n \geq 0, \\ a_n &= \begin{cases} 1 & \text{for } n = 0, \\ p^n + (p^{n-1} - 1)/(p + 1) & \text{for odd } n \geq 1, \\ p^n + p(p^{n-2} - 1)/(p + 1) & \text{for even } n \geq 2, \end{cases} \\ j_{s,n} &= \begin{cases} 2 & \text{for } s \in \mathbb{Z}_0 \text{ and } n = 0, \\ 2p^2 - p + 1 & \text{for } s \in \mathbb{Z}_0 \text{ and } n = 2, \\ 2a_n + 1 & \text{for } s \in \mathbb{Z}_0, \text{ even } n \geq 4, \\ a_{n+2} - a_{n+1} & \text{for } s \in \mathbb{Z}_1 \text{ and even } n \geq 0, \\ p + 1 & \text{for } s \in \mathbb{Z}^{(p)} \text{ and } n = 1, \\ e(3)p^{n-2} - p + 1 & \text{for } s \in \mathbb{Z}^{(p)} \text{ and odd } n \geq 3, \end{cases} \end{aligned}$$

$$j'_{s,0} = \begin{cases} 2 & \text{for } p \nmid s(s-1), \\ 2p & \text{for } s = tp + 1 \text{ and } p \nmid t(t-1), \\ p^2 + 1 & \text{for } s = tp^2 + 1 \text{ and } p \nmid t, \\ a_n + p & \text{for } s = tp^n + 1 \text{ with } n > 2 \text{ and } p \nmid t, \\ a_n + 1 & \text{for } s = tp^n + e(n) \text{ with even } n \geq 2 \text{ and } p \nmid (t-1), \\ a_n + 2 & \text{for } s = tp^n + e(n) \text{ with odd } n > 2 \text{ and } p \nmid (t-1), \end{cases}$$

$$j'_{s,n} = \begin{cases} 2p & \text{for } s \in \mathbb{Z}_0 \text{ and } n = 1, \\ 2pa_{n-1} + p & \text{for } s \in \mathbb{Z}_0 \text{ and odd } n \geq 3, \\ pa_{n+1} - pa_n & \text{for } s \in \mathbb{Z}_1 \text{ and odd } n \geq 1, \\ p^2 + p & \text{for } s \in \mathbb{Z}^{(p)} \text{ and } n = 2, \\ e(3)p^{n-2} - 1 + \bar{1} & \text{for } s \in \mathbb{Z}^{(p)} \text{ and even } n \geq 4. \end{cases}$$

Here $\bar{1} = 0$ if $p \geq 5$ and 1 if $p = 3$, the \mathbb{Z}_i are the subsets of the integers \mathbb{Z} defined in (1.7), and the integers a_n are $a_{3,n}$ in [2, (5.13)]. We note that

$$(2.8) \quad a_n + a_{n-1} = e(3)p^{n-2} - 1 \quad \text{for } n \geq 2 \quad \text{and} \quad p^n + a_{n-2} - p^{n-3} = a_n \quad \text{for } n \geq 3.$$

Theorem 2.9 *Let p be an odd prime. Then $H^1 M_2^1$ is the direct sum of the $k(2)_*$ -module $B_\infty = K(2)_*/k(2)_*\{h_0, h_1, \tilde{\xi}_2, \xi_3\}$ and the $k(2)_*$ -cyclic modules generated by*

$$\begin{aligned} &(\xi_3)_{sp^n/a_n} \quad \text{for } (s, n) \in \mathbb{Z}^{(p)} \times \mathbb{N}, \\ &(h_0)_{sp^n/j_{s,n}} \quad \text{for } (s, n) \in ((\mathbb{Z}_0 \cup \mathbb{Z}_1) \times 2\mathbb{N}) \cup (\mathbb{Z}^{(p)} \times \overline{2\mathbb{N}}), \\ &(h_1)_{sp^n/j'_{s,n}} \quad \text{for } (s, n) \in ((\mathbb{Z}_0 \cup \mathbb{Z}_1) \times \overline{2\mathbb{N}}) \cup ((\mathbb{Z}^{(p)} \times 2\mathbb{N}) \setminus \{(1, 0)\}), \\ &(h_2)_{tp-1/p-1} \quad \text{for } t \in \mathbb{Z}. \end{aligned}$$

There is a little difference between the cases for $p \geq 5$ and $p = 3$. In the theorem, $\tilde{\xi}_2 (= (h_1)_1)$ denotes the homology class of z in (4.18) (see also (3.8)), and the generators $(\xi)_{s/j}$ for $\xi = [X]$ in $H^1 M_3^0$ denote

$$(\xi)_{s/j} = [v_3^s X / v_2^j + *]$$

for a cocycle $v_3^s X / v_2^j + *$ of the cobar complex $\Omega^1 M_2^1$ with an element $*$ killed by v_2^{j-1} . The element v_2 acts on $(\xi)_{s/j}$ by

$$(2.10) \quad v_2(\xi)_{s/j} = (\xi)_{s/j-1} \quad \text{and} \quad v_2(\xi)_{s/1} = 0,$$

and so $(\xi)_{s/j}$ generates a cyclic $k(2)_*$ -module isomorphic to $k(2)_*/(v_2^j)$:

$$k(2)_*\{(\xi)_{s/j}\} \cong k(2)_*/(v_2^j).$$

3 Proofs of Theorems 2.9 and 1.8

In this section, we assume Lemma 3.4, which will be verified by a routine calculation in Section 6.

3.1 Proof of Theorem 2.9

For the monochromatic comodules defined in Section 2, we have a short exact sequence

$$(3.1) \quad 0 \rightarrow M_3^0 \xrightarrow{\eta} M_2^1 \xrightarrow{v_2} M_2^1 \rightarrow 0,$$

where $\eta(x) = x/v_2$ (see [2, (3.10)]), which induces the long exact sequence

$$(3.2) \quad \cdots \rightarrow H^0 M_2^1 \xrightarrow{\delta_0} H^1 M_3^0 \xrightarrow{\eta_*} H^1 M_2^1 \xrightarrow{v_2} H^1 M_2^1 \xrightarrow{\delta_1} H^2 M_3^0 \rightarrow \cdots.$$

From [2, (5.18)], we read off the following:

Proposition 3.3 *The cokernel of $\delta_0: H^0 M_2^1 \rightarrow H^1 M_3^0$ is a \mathbb{Z}/p -module generated by $(h_0)_0, (h_1)_0$,*

$$\begin{aligned} (h_0)_{sp^{2k}} & \text{ for } s \in \mathbb{Z}_0 \cup \mathbb{Z}_1, & (h_0)_{tp^{2k+1}} & \text{ for } t \in \mathbb{Z}^{(p)}, & (h_1)_{tp^{2k}} & \text{ for } t \in \mathbb{Z}^{(p)}, \\ (h_1)_{sp^{2k+1}} & \text{ for } s \in \mathbb{Z}_0 \cup \mathbb{Z}_1, & (h_2)_{tp^{-1}} & \text{ for } t \in \mathbb{Z}, & (\xi_3)_t & \text{ for } t \in \mathbb{Z}, \end{aligned}$$

for $k \geq 0$. Here \mathbb{Z}_i is a subset of \mathbb{Z} given in (1.7), and $(\xi)_s = v_3^s \xi$ for $\xi \in \{h_i, \xi_3 : i \in \mathbb{Z}/3\}$.

Let $(x)_s \in \Omega^1 E(3)_*$ denote a cochain satisfying

$$(x)_s \equiv v_3^s x \pmod{I_3}.$$

Lemma 3.4 *The following cochains exist in $\Omega^1 E(3)_*/I_2$:*

(1) $(t_1)_{sp^{2k}}$ and $(t_1^p)_{sp^{2k+1}}$ for $s \in \mathbb{Z}_0$ such that

$$d_1((t_1)_{sp^{2k}}) \equiv \begin{cases} s(s+1)v_2^2 v_3^{s-1-p} G_2 & k=0, \\ s(s+1)v_2^2 p^{2-p+1} v_3^{sp^2-2p} G_1 & k=1, \\ -3s(s+1)v_2^{2a_{2k}} v_3^{(sp-2)p^{2k-1}} K_0 & k \geq 2, p \geq 5, \\ -2s(s+1)v_2^{2a_{2k+1}} v_3^{3^{2k-1}(3s-2)} (b_{1,0} + t_1^p \otimes Z') & k \geq 2, p=3, \end{cases}$$

$$d_1((t_1^p)_{sp^{2k+1}}) \equiv \begin{cases} s(s+1)v_2^{2p} v_3^{sp^2-2p} G_0 & k=0, \\ s(s+1)v_2^{2pa_{2k}+p} v_3^{(sp-2)p^{2k}} b_{1,1} & k \geq 1, \end{cases}$$

(2) $(t_1)_{sp^{2k}}$ and $(t_1^p)_{sp^{2k+1}}$ for $s = tp^2 - 1 \in \mathbb{Z}_1$ such that

$$d_1((t_1)_{sp^{2k}}) \equiv v_2^{a_{2k}+2-a_{2k+1}} v_3^{(tp-1)p^{2k+1}} b_{1,0}, \quad d_1((t_1^p)_{sp^{2k+1}}) \equiv v_2^{pa_{2k}+2-pa_{2k+1}} v_3^{(tp-1)p^{2k+2}} b_{1,1},$$

for $k \geq 0$,

(3) $(t_1)_{sp^{2k+1}}$ and $(t_1^p)_{sp^{2k}}$ for $s \in \mathbb{Z}^{(p)}$ such that

$$d_1((t_1^p)_{tp^{k+1}}) \equiv \begin{cases} t(t-1)v_2^{2p} v_3^{tp^{-1}} G_0 & k=1, \\ -tv_2^{p^2+1} v_3^{(tp-1)p} G_1 & k=2, \\ -2tv_2^{a_k+p} v_3^{(tp-1)p^{k-1}} G_0 & \text{odd } k \geq 3, \\ 2tv_2^{a_k+p} v_3^{(tp-1)p^{k-1}} K_0 & \text{even } k \geq 4, \end{cases}$$

$$d_1((t_1^p)_{tp^k+e(k)}) \equiv \begin{cases} -(t-1)v_2^{a_k+1} v_3^{tp^k+pe(k-2)} G_1 & \text{even } k \geq 2, \\ -(t-1)v_2^{a_k+2} v_3^{tp^k+pe(k-2)} b_{1,1} & \text{odd } k \geq 3, \end{cases}$$

$$d_1((t_1^p)_{sp^{2k}}) \equiv \begin{cases} s(s-1)v_2^2 v_3^{s-2} K_1 & k=0, \\ -sv_2^{p^2+p} v_3^{sp^2-p-1} K_0 & k=1, \\ -3sv_2^{e(3)p^{2k-2}-1} v_3^{(sp^2-p-1)p^{2k-2}} K_0 & p \geq 5, k \geq 2, \\ -sv_2^{3^{2k-2}e(3)} v_3^{(9s-4)3^{2k-2}} (b_{1,0} + Z' \otimes t_1^p) & p=3, k \geq 2, \end{cases}$$

$$d_1((t_1)_{sp^{2k+1}}) \equiv \begin{cases} -sv_2^{p+1} v_3^{(s-2)p} K_2 & k=0, \\ sv_2^{e(3)p^{2k-1}-p+1} v_3^{(sp^2-p-1)p^{2k-1}} b_{1,1} & k \geq 1, \end{cases}$$

$$(4) \quad (t_1^{p^2})_{tp-1} \text{ such that } d_1((t_1^{p^2})_{tp-1}) \equiv v_2^{p-1} v_3^{tp-p} b_{1,2}.$$

Here G_i , K_i and $b_{1,i}$ are the cocycles of $\Omega^2 E(3)_*/I_2$ in (2.5), Z' is an element in Lemma 5.1, and $x \equiv v_2^a y$ denotes the congruence modulo J_{a+1} .

Let $d_1((x)_t) \equiv v_2^j y \bmod J_{j+1}$ be a congruence in Lemma 3.4. Then $\delta_1([x]_{t/j}) = [y]$ for the connecting homomorphism δ_1 in (3.2). Here $[x]_{t/j} = [(x)_t/v_2^j] \in H^1 M_2^1$ denotes the cohomology class of the cocycle $(x)_t/v_2^j$ of $\Omega^1 M_2^1$. Thus the cochains in Lemma 3.4 give rise to elements $(h_0)_{sp^r/j_{s,r}}$ and $(h_1)_{sp^r/j'_{s,r}}$ of $H^1 M_2^1$ as well as their δ_1 -images. Furthermore, we have elements

$$(\zeta_3)_{tp^n/a_n} = x_n^t \zeta_3 / v_2^{a_n} \in H^1 M_2^1$$

for the elements $x_n (= x_{3,n})$ introduced in [2, (5.11)] (see Lemma 5.1) with

$$(3.5) \quad \delta_1((\zeta_3)_{tp^n/a_n}) = \begin{cases} (h_2 \zeta_3)_{t-1} & n=0, \\ (h_0 \zeta_3)_{(tp-1)p^{n-1}} & n \text{ is odd}, \\ (h_1 \zeta_3)_{(tp-1)p^{n-1}} & n \text{ is even } \geq 2, \end{cases}$$

by [2, (5.18)] (or Lemma 5.1). As a $k(2)_*$ -module, $K(2)_*/k(2)_*\{\xi\} = \mathbb{Z}/p\{(\xi)_{0/j} : j \geq 1\}$ with $v_2(\xi)_{0/j} = (\xi)_{0/j-1}$ and $v_2(\xi)_{0/1} = 0$; see (2.10).

Let B be the $k(2)_*$ -module of the theorem. Each direct summand of B is a submodule of $H^1 M_2^1$, which defines a $k(2)_*$ -module map $f: B \rightarrow H^1 M_2^1$. Furthermore, assigning $(\xi)_{s/1} \in B$ to the generator $(\xi)_s$ of the cokernel of δ_0 , we have a homomorphism $\bar{\eta}_*: H^1 M_3^0 \rightarrow B$ by Proposition 3.3. These homomorphisms fit in the commutative diagram

$$\begin{array}{ccccccccc} H^0 M_2^1 & \xrightarrow{\delta_0} & H^1 M_3^0 & \xrightarrow{\bar{\eta}_*} & B & \xrightarrow{v_2} & B & \xrightarrow{\delta'_1} & H^2 M_3^0 \\ \parallel & & \parallel & & \downarrow f & & \downarrow f & & \parallel \\ H^0 M_2^1 & \xrightarrow{\delta_0} & H^1 M_3^0 & \xrightarrow{\eta_*} & H^1 M_2^1 & \xrightarrow{v_2} & H^1 M_2^1 & \xrightarrow{\delta_1} & H^2 M_3^0 \end{array}$$

where we define δ'_1 by $\delta_1 f$. It suffices to show that the upper sequence is exact by [2, Remark 3.11]. By the definition of B , the subsequence $H^0 M_2^1 \xrightarrow{\delta_0} H^1 M_3^0 \xrightarrow{\bar{\eta}_*} B \xrightarrow{v_2} B$ is exact and the composite $B \xrightarrow{v_2} B \xrightarrow{\delta'_1} H^2 M_3^0$ is zero.

Suppose that the δ'_1 -images of the generators are linearly independent, and take $\xi \in \text{Ker } \delta'_1$ to be a homogeneous element. Then

$$\xi = \sum_k c_k \xi_k \quad \text{for generators } \xi_k \text{ of } B \text{ and scalars } c_k \in k(2)_* \quad \text{and} \quad 0 = \delta'_1(\xi) = \sum_k \bar{c}_k \delta'_1(\xi_k)$$

for the image \bar{c}_k of c_k under the projection $k(2)_* \rightarrow \mathbb{Z}/p$ sending v_2 to zero. Since the $\delta'_1(\xi_k)$ are linearly independent we see $\bar{c}_k = 0$, and so we have $c'_k \in k(2)_*$ such that $c_k = v_2 c'_k$. Therefore

$$\xi = \sum_k v_2 c'_k \xi_k \in \text{Im } v_2,$$

and we see that the upper sequence of the above diagram is exact if the δ'_1 -images of the generators are linearly independent.

The δ'_1 -image is a \mathbb{Z}/p -submodule of $H^2 M_3^0$ in (2.3) with generators of the form $(\rho)_s$ for $\rho \in \{g_i, k_i, b_i, h_i \zeta_3 : i \in \mathbb{Z}/3\}$ by Lemma 3.4 and (3.5). Moreover, Lemma 3.4 and (3.5) show that the δ'_1 -image of each generator ξ_k has only one summand of form $(\rho)_s$,

$$(h_0 \zeta_3)_{(tp-1)p^{2n}}, \quad (h_1 \zeta_3)_{(tp-1)p^{2n-1}}, \quad (h_2 \zeta_3)_{t-1}, \quad (g_2)_{s-1-p}, \\ (k_1)_{s-2}, \quad (k_2)_{(s-2)p}, \quad (b_0)_{(tp-1)p^{2n+1}} \text{ for } p \geq 5, \quad (b_2)_{tp-p},$$

except for

g_0	$(g_0)_{sp-2}$	$(g_0)_{(tp-1)p^{2n}}$		
g_1	$(g_1)_{(sp-2)p}$	$(g_1)_{(tp-1)p}$	$(g_1)_{tp^{2n}+pe(2n-2)}$	
k_0	$(k_0)_{(sp-2)p^{2n-1}}$	$(k_0)_{(tp-1)p^{2n-1}}$	$(k_0)_{(sp^2-p-1)p^{2n}}$	$(p \geq 5)$
k_0	$(k_0)_{3^{2n-1}(3t-1)}$	$(k_0)_{9s-4}$		$(p = 3)$
b_0	$(b_0)_{3^{2n-1}(3s-2)}$	$(b_0)_{3^{2n+1}(3t-1)}$	$(b_0)_{3^{2n-2}(9s-4)}$	$(p = 3)$
b_1	$(b_1)_{(sp-2)p^{2n}}$	$(b_1)_{(tp-1)p^{2n+2}}$	$(b_1)_{tp^{2n+1}+pe(2n-1)}$	$(b_1)_{(sp^2-p-1)p^{2n-1}}$

These show that the δ'_1 -images $\delta'_1(\xi_k)$ for the generators ξ_k of B are different from each other, and so they are linearly independent. \square

3.2 Proof of Theorem 1.8

Let $\delta_2^0: H^* N_2^1 \rightarrow H^{*+1} N_2^0$ be the connecting homomorphism associated to the short exact sequence (2.1), and consider the diagram

$$\begin{array}{ccccc} H^2 M_2^0 & \xrightarrow{(\kappa_2^0)_*} & H^2 N_2^1 & \xrightarrow{\delta_2^0} & H^3 N_2^0 = E_2^3(V(1)) \\ & & \downarrow \iota_2^1 & & \\ H^1 M_2^1 & \xrightarrow{\delta_1} & H^2 M_3^0 & \xrightarrow{\eta_*} & H^2 M_2^1 \end{array}$$

of exact sequences for δ_1 in (3.2). The connecting homomorphism $\bar{\delta}_j$ associated to (1.6) is factorized into the composite $\bar{\delta}_j: H^s \text{BP}_*/J_j \xrightarrow{\iota_j} H^s N_2^1 \xrightarrow{\delta_2^0} H^{s+1} N_2^0$ for the homomorphism $\hat{\iota}_j$ given by $\hat{\iota}_j(x) = x/v_2^j$. It follows that

$$(3.6) \quad \bar{\gamma}_{sp^r/j}'' = \delta_2^0(v_3^{sp^r}/v_2^j) \in H^1 N_2^0 = E_2^1(V(1)) \quad \text{for } v_3^{sp^r}/v_2^j \in H^0 N_2^1.$$

Since δ_2^0 is a $k(2)_*$ -module map, we have

$$(3.7) \quad v_2^{j-1} \bar{\gamma}_{sp^r/j}'' = v_2^{j-1} \delta_2^0(v_3^{sp^r}/v_2^j) = \delta_2^0(v_2^{j-1} v_3^{sp^r}/v_2^j) = \delta_2^0(v_3^{sp^r}/v_2) = \bar{\gamma}_{sp^r}''.$$

It is well known that

$$\bar{\beta}_1 = -b_0 = [-b_{1,0}] \quad \text{and} \quad \bar{\beta}_2 = 2k_0 = [2K_0] \in H^2 N_3^0$$

for the cocycles $b_{1,0}$ and K_0 in (2.5); see [5, Lemma 4.4]. This defines elements $v_3^{sp^r} \bar{\beta}_i/v_2 \in H^2 N_2^1$ for $i = 1, 2$, and

$$\delta_2^0(v_3^{sp^r} \bar{\beta}_i/v_2) = \gamma_{sp^r}'' \bar{\beta}_i \in E_2^3(V(1)) \quad (\text{by (3.6)}).$$

We also see that for $v_3^{sp^r} \bar{\beta}_i \in H^2 M_3^0$,

$$\eta_*(v_3^{sp^r} \bar{\beta}_i) = \iota_2^1(v_3^{sp^r} \bar{\beta}_i/v_2) \in H^2 M_2^1.$$

From Lemma 3.4, the elements $v_3^{sp^r} \bar{\beta}_1 = -(b_0)_{sp^r}$ and $v_3^{sp^r} \bar{\beta}_2 = 2(k_0)_{sp^r} \in H^2 M_3^0$ may be in the image of δ_1 if

$$p \geq 5 \quad \text{and} \quad (s, r) \in (\mathbb{Z}_1^+ \cup \mathbb{Z}_2^+) \times \overline{2\mathbb{N}}$$

or

$$p = 3 \quad \text{and} \quad (s, r) \in (\overline{\mathbb{N}}_1 \times 2\mathbb{N}_{>0}) \cup ((\mathbb{Z}_1^+ \cup \mathbb{Z}_2^+) \times \overline{2\mathbb{N}}) \cup (\overline{\mathbb{N}}_2 \times \overline{2\mathbb{N}}_{>1}),$$

and if

$$p \geq 5 \quad \text{and} \quad (s, r) \in (\overline{\mathbb{N}}_1 \times 2\mathbb{N}) \cup (\mathbb{Z}_2^+ \times \overline{2\mathbb{N}}_{>1}) \cup (\overline{\mathbb{N}}_2 \times \overline{2\mathbb{N}}_{>1})$$

or

$$p = 3 \quad \text{and} \quad (s, r) \in (\overline{\mathbb{N}}_1 \times \{0\}) \cup (\mathbb{Z}_2^+ \times \overline{2\mathbb{N}}_{>1}),$$

respectively. Here $\overline{\mathbb{N}}_i = \mathbb{N}^{(p)} \setminus \mathbb{N}_i$ for $i = 1, 2$. Therefore, if a pair (s, r) satisfies the condition of the theorem, then the element $v_3^{sp^r} \bar{\beta}_i$ is not in the image of δ_1 , and survives to $\iota_2^1(v_3^{sp^r} \bar{\beta}_i/v_2)$ under the homomorphism η_* . Thus $v_3^{sp^r} \bar{\beta}_i/v_2 \neq 0 \in H^2 N_2^1$ under the conditions.

Ravenel determined in [8, Theorem 6.3.24; 7, Theorem 3.2] that

$$(3.8) \quad H^2 M_2^0 = \begin{cases} K(2)_* \{h_0 \tilde{\zeta}_2, h_1 \tilde{\zeta}_2, b_0, b_1, \xi\} & p = 3, \\ K(2)_* \{h_0 \tilde{\zeta}_2, h_1 \tilde{\zeta}_2, g_0, g_1\} & p \geq 5, \end{cases}$$

where $\tilde{\zeta}_2 = v_2^{p+1} \zeta_2 = [-z]$ for ζ_2 in [2, Proposition 3.18] and z in (4.18). This shows that the elements $v_3^{sp^r} \bar{\beta}_i/v_2$ for $i = 1, 2$ are not in the image of $(\kappa_2^0)_*$, and hence survive to $\gamma_{sp^r}'' \bar{\beta}_i \in E_2^3(V(1))$. Moreover, $\gamma_{sp^r/j}'' \bar{\beta}_i \neq 0 \in E_2^3(V(1))$ if $v_2^{j-1} \gamma_{sp^r/j}'' \bar{\beta}_i = \gamma_{sp^r}'' \bar{\beta}_i$ is not zero, where the equality follows by (3.7). \square

4 Some cochains in the cobar complex $\Omega^* E(3)_*$

In the rest of this paper, we consider $E(3)_*(E(3))$ -comodules whose structure maps are induced from the right unit map $\eta_R: E(3)_* \rightarrow E(3)_*(E(3))$. We consider the cobar complex $\Omega^* M$ of a comodule M in (2.2), whose differentials are given by

$$(4.1) \quad d_0(v) = \eta_R(v) - v \in \Omega^1 E(3)_*, \quad \text{and} \quad d_1(x) = 1 \otimes x - \Delta(x) + x \otimes 1 \in \Omega^2 E(3)_*$$

for $v \in \Omega^0 E(3)_* = E(3)_*$ and $x \in \Omega^1 E(3)_* = E(3)_*(E(3))$. For the differentials d_0 and d_1 , we have relations (see [11, (2.3.2)])

$$\begin{aligned}
 (4.2) \quad & d_0(vv') = vd_0(v') + d_0(v)\eta_R(v'), \\
 & d_1(vx) = d_0(v) \otimes x + vd_1(x), \\
 & d_1(xy) = -x \otimes y - y \otimes x + d_1(x)\Delta y + (x \otimes 1 + 1 \otimes x)d_1(y), \\
 & d_1(x\eta_R(v)) = d_1(x)(1 \otimes \eta_R(v)) - x \otimes d_0(v),
 \end{aligned}$$

for $v, v' \in E(3)_*$ and $x, y \in E(3)_*(E(3))$. A formula for the Hopf conjugation $c: \mathrm{BP}_*(\mathrm{BP}) \rightarrow \mathrm{BP}_*(\mathrm{BP})$ is given in [6, (3)], and immediately implies the following:

Lemma 4.3 *The Hopf conjugation $c: E(3)_*(E(3)) \rightarrow E(3)_*(E(3))$ acts as*

$$ct_1 = -t_1, \quad ct_2 = t_1^{p+1} - t_2 \quad \text{and} \quad ct_3 \equiv t_2 t_1^{p^2} - t_1 c t_2^p - t_3 \pmod{I_2}.$$

For the right unit $\eta_R: \mathrm{BP}_* \rightarrow \mathrm{BP}_*(\mathrm{BP})$, we have a well-known formula (see [6, (11)])

$$(4.4) \quad \eta_R(v_n) \equiv v_n + v_{n-1} t_1^{p^{n-1}} - v_{n-1}^p t_1 \pmod{I_{n-1}}.$$

A routine calculation using (4.1) and (4.4) shows the following:

Lemma 4.5 *Put $\sigma_n = \sum_{k=0}^{n-1} v_2^{p^{2k} a_{2n-2k-1} - p^{2k+1}} v_3^{p^{2k}} \in E(3)_*$. Then*

$$d_0(\sigma_n) \equiv v_2^{p^{2n-2}} t_1^{p^{2n}} - v_2^{a_{2n-1}} t_1 \pmod{I_2}.$$

In $E(3)_*(E(3))$ we have $\eta_R(v_4) = 0 = \eta_R(v_5)$, which give rise to relations

$$(4.6) \quad v_3 t_1^{p^3} \equiv t_1 \eta_R(v_3)^p - v_2 t_2^{p^2} + v_2^p t_2 \quad \text{and} \quad v_3 t_2^{p^3} \equiv t_2 \eta_R(v_3)^{p^2} - v_2 t_3^{p^2} - v_2 w^p + v_2^p t_3 \pmod{I_2}$$

(see [6, (12) and (16); 8, Corollary 4.3.21]), where $w \in E(3)_*(E(3)) (= w_1(v_3, v_2 t_1^{p^2}, -v_2^p t_1)$ in [8, Corollary 4.3.21]) is an element defined by

$$(4.7) \quad pw = v_3^p + v_2^p t_1^{p^3} - v_2^{p^2} t_1^p + y^p - \eta_R(v_3)^p$$

for $y \in (p, v_1)$ in $\eta_R(v_3) = v_3 + v_2 t_1^{p^2} - v_2^p t_1 + y$; see (4.4).

The diagonal $\Delta: E(3)_*(E(3)) \rightarrow E(3)_*(E(3)) \otimes_{E(3)_*} E(3)_*(E(3))$ of the Hopf algebroid $E(3)_*(E(3))$ acts on the elements t_i and ct_i by

$$\begin{aligned}
 (4.8) \quad & \Delta(t_1) = t_1 \otimes 1 + 1 \otimes t_1, \\
 & \Delta(t_2) \equiv t_2 \otimes 1 + t_1 \otimes t_1^p + 1 \otimes t_2 - v_1 b_{1,0} \pmod{(p, v_1^2)}, \\
 & \Delta(t_3) \equiv t_3 \otimes 1 + t_2 \otimes t_1^{p^2} + t_1 \otimes t_2^p + 1 \otimes t_3 - v_2 b_{1,1} \pmod{I_2}, \\
 & \Delta(t_4) \equiv t_4 \otimes 1 + t_3 \otimes t_1^{p^3} + t_2 \otimes t_2^{p^2} + t_1 \otimes t_3^p + 1 \otimes t_4 - v_3 b_{1,2} \pmod{I_3},
 \end{aligned}$$

(see [6, Theorem 8; 8, Corollary 4.3.15]) and so

$$\begin{aligned}
 d_1(ct_2) &\equiv -t_1^p \otimes t_1, \\
 (4.9) \quad d_1(ct_3) &\equiv ct_2^p \otimes t_1 + t_1^{p^2} \otimes ct_2 - v_2 b_{1,1} \pmod{I_2}, \\
 d_1(ct_4) &\equiv t_1^{p^3} \otimes ct_3 - ct_2^{p^2} \otimes ct_2 + ct_3^p \otimes t_1 - v_3 b_{1,2} \pmod{I_3},
 \end{aligned}$$

since $\Delta(cx) = (c \otimes c)T\Delta(x)$ for the switching map T given by $T(x \otimes y) = y \otimes x$, where $b_{1,k}$ is the cocycle in (2.5).

The fact $d_1(t_1^{p^{k+1}}) \equiv -pb_{1,k} \pmod{p^2}$ implies not only that the cochain $b_{1,k} \in \Omega^2 E(3)_*/(p)$ is a cocycle, but also the following lemma:

Lemma 4.10 *The cochain w in (4.7) satisfies*

$$w \equiv -v_2 v_3^{p-1} t_1^{p^2} \pmod{J_2} \quad \text{and} \quad d_1(w) \equiv -v_2^p b_{1,2} + v_2^{p^2} b_{1,0} \pmod{I_2}.$$

Corollary 4.11 *Put $W_n = \sum_{i=0}^{n-1} v_2^{p^{2i}} a_{2n-2i-p^{2i+2}} w^{p^{2i}}$. Then*

$$d_1(W_n) \equiv -v_2^{p^{2n-1}} b_{1,2n} + v_2^{a_{2n}} b_{1,0} \pmod{I_2}.$$

We generalize the relations (4.6) and obtain the following proposition from [8, (4.3.1) and Lemma 4.3.11]; [6, Theorem 1] (see [9, Proposition 2.1]):

Proposition 4.12 *There exist elements T_n for $n \geq 0$ satisfying $T_n \equiv t_n^p \pmod{I_3}$ and*

$$v_2^{p^{k+1}} t_{k+1} + t_k \eta_R(v_3)^{p^k} \equiv v_1 T_{k+2} + v_2 T_{k+1}^p + v_3 T_k^{p^2} \pmod{(p, v_1^2)}$$

for $k \geq 0$. In particular, $T_0 = 1$, $T_1 \equiv t_1^p$, $T_2 \equiv t_2^p$ and $T_3 \equiv t_3^p + w \pmod{I_2}$.

Proof We begin by recalling some notation from [8, Section 4.3]. For a sequence $J = (j_1, j_2, \dots, j_m)$ of positive integers we set $|J| = m$ and $\|J\| = \sum_{i=1}^m j_i$, and an element $v_J \in E(3)_*$ is defined recursively by $v_{(j,J)} = v_j v_J^{p^j}$. Let $w_k(S)$ for a set S be symmetric polynomials of degree p^n such that $w_0(S) = \sum_{x \in S} x$ and $\sum_{x \in S} x^{p^n} = \sum_{k=0}^n p^k w_k(S)^{p^{n-k}}$. We then define sets S_n out of a set $S = \{a_{i,j}\}$ recursively by

$$S_n = \{a_{i,j} : i + j = n\} \cup \bigcup_{|J| > 0} \{v_J w_{|J|}(S_{n-\|J\|})^{p^{\|J\| - |J|}}\}.$$

By [8, (4.3.1) and Lemma 4.3.11],

$$(4.13) \quad w_0(C_n) \equiv \sum_{i+j=n}^F t_i \eta_R(v_j)^{p^i} \equiv \sum_{i+j=n}^F v_i t_j^{p^i} \equiv w_0(D_n) \pmod{(p)}$$

for the sets

$$C = \{t_i \eta_R(v_j)^{p^i}\} \quad \text{and} \quad D = \{v_i t_j^{p^i}\}.$$

In $E(3)_*(E(3))$, put

$$w(S_n) = \sum_J v_J^p w_{|J|+1}(S_{n-\|J\|})^{p^{\|J\| - |J|}} \quad \text{and} \quad T_n = t_n^p - w(C_n) + w(D_n).$$

Then the proposition follows from (4.13) and the congruences

$$\begin{aligned} w_0(C_n) &\equiv v_2^{p^{n-2}} t_{n-2} + t_{n-3} \eta_R(v_3)^{p^{n-3}} + v_1 w(C_{n-1}) + v_2 w(C_{n-2})^p + v_3 w(C_{n-3})^{p^2}, \\ w_0(D_n) &\equiv v_1 t_{n-1}^p + v_2 t_{n-2}^{p^2} + v_3 t_{n-3}^{p^3} + v_1 w(D_{n-1}) + v_2 w(D_{n-2})^p + v_3 w(D_{n-3})^{p^2}, \end{aligned}$$

seen by the relation

$$v_{(k,J)} w_{|(k,J)|} (S_{n-\|(k,J)\|})^{p^{\|(k,J)\| - |(k,J)|}} = v_k v_J^{p^k} w_{|J|+1} (S_{n-k-\|J\|})^{p^{\|J\| - |J| + k - 1}}. \quad \square$$

Lemma 4.14 For $n \geq 0$,

$$\eta_R(v_2^{p-1} v_3^{e(n)}) \equiv \sum_{i=0}^n (-1)^{n-i} v_2^{p^{i+1}e(n-i)+p-1} v_3^{e(i)} t_{n-i}^{p^i} - v_2^p w_n^p + v_1 v_2^{p-2} w_{n+1} \pmod{(p, v_1^2)}.$$

Here

$$(4.15) \quad w_n = \sum_{i=1}^n (-1)^i v_2^{e(i-1)} T_i \eta_R(v_3^{p^{i-1}e(n-i)}).$$

Proof In this proof, every congruence is considered modulo (p, v_1^2) . By Proposition 4.12, $t_k \eta_R(v_3^{p^k}) \equiv \tilde{T}_k - v_2^{p^{k+1}} t_{k+1}$ for $\tilde{T}_k = v_1 T_{k+2} + v_2 T_{k+1}^p + v_3 T_k^{p^2}$, which implies inductively

$$t_1 \eta_R(v_3^{pe(n)}) \equiv - \sum_{i=1}^n (-1)^i v_2^{p^2e(i-1)} \tilde{T}_i \eta_R(v_3^{p^{i+1}e(n-i)}) + (-1)^n v_2^{p^2e(n)} t_{n+1},$$

and hence

$$(4.16) \quad \begin{aligned} t_1 \eta_R(v_3^{pe(n)}) &\equiv -v_1 v_2^{-p-1} w_{n+2} + v_2^{1-p} w_{n+1}^p - v_3 w_n^{p^2} + (-1)^n v_2^{p^2e(n)} t_{n+1} \\ &\quad - v_1 v_2^{-p-1} (t_1^p \eta_R(v_3) - v_2 t_2^p) \eta_R(v_3^{pe(n)}) + v_2^{1-p} t_1^{p^2} \eta_R(v_3^{pe(n)}). \end{aligned}$$

Now we prove the lemma by induction. For $n = 0$, it follows from the facts $\eta_R(v_2) \equiv v_2 + v_1 t_1^p$ by (4.4) and $w_1 = -t_1^p$.

Assuming the case for n , we obtain the case for $n + 1$ from (4.16) and

$$\begin{aligned} \eta_R(v_2^{p-1} v_3^{e(n+1)}) &\equiv v_2^{-p^2+2p-1} v_3 \eta_R(v_2^{p-1} v_3^{e(n)})^p + v_2^{p-1} (v_2 t_1^{p^2} + v_1 t_2^p) \eta_R(v_3^{pe(n)}) \\ &\quad - v_2^{2p-1} t_1 \eta_R(v_3^{pe(n)}) - v_1 v_2^{p-2} t_1^p \eta_R(v_3^{e(n+1)}), \end{aligned}$$

given by $\eta_R(v_2^{p-1} v_3) \equiv v_2^{p-1} (v_3 + v_2 t_1^{p^2} - v_2^p t_1 + v_1 t_2^p) - v_1 v_2^{p-2} t_1^p \eta_R(v_3)$. Here $\eta_R(v_3)$ is given in [2, (5.7)]. \square

Evaluate the congruence in Lemma 4.14 under d_1 , and compare the v_1 -multiples. Then we deduce the following corollary; see [9, Proposition 2.3]. Indeed, if $v_1 v_2^{p-2} d_1(w_{n+1}) \equiv A + v_1 B \pmod{(p, v_1^2)}$ for some A and B involving no v_1 , then $A \equiv 0 \pmod{(p, v_1^2)}$ and $v_2^{p-2} d_1(w_{n+1}) \equiv B \pmod{I_2}$.

Corollary 4.17 For the elements w_n in (4.15),

$$d_1(w_{n+1}) \equiv - \sum_{i=0}^{n-1} (-1)^{n-i} v_2^{p^{i+1}e(n-i)} w_{i+1} \otimes t_{n-i}^{p^i} - (-1)^n v_2^{e(n+1)} b_n \pmod{I_2}.$$

Here b_n is an element in $d_1(t_n) \equiv a_n + v_1 b_n \pmod{(p, v_1^2)}$ for a_n and b_n involving no v_1 . In particular, $b_2 = b_{1,0}$ by (4.8).

We have the cocycle z in $\Omega^1 E(3)_*/I_2$ given by

$$(4.18) \quad z = v_3 t_1^p + v_2 c t_2^p - v_2^p t_2 = t_1^p \eta_R(v_3) - v_2 t_2^p + v_2^p c t_2 = -w_2 + v_2^p c t_2,$$

which represents the element $-v_2^{p+1} \zeta_2 \in H^1 M_2^0$; see [2, Proposition 3.18(c)] and (3.8). In particular,

$$(4.19) \quad t_1^p \eta_R(v_3) \equiv z + v_2 t_2^p - v_2^p c t_2 \pmod{I_2}.$$

We further have cocycles G'_i and $K'_i \in \Omega^2 E(3)_*/I_2$ for $i \in \{0, 1, 2\}$ defined by

$$(4.20) \quad G'_i = c t_2^{p^i} \otimes t_1^{p^i} + \frac{1}{2} t_1^{p^{i+1}} \otimes t_1^{2p^i} \quad \text{and} \quad K'_i = t_1^{p^{i+1}} \otimes c t_2^{p^i} + \frac{1}{2} t_1^{2p^{i+1}} \otimes t_1^{p^i},$$

which are homologous to G_i and K_i in (2.5), respectively. Indeed,

$$(4.21) \quad d_1(g_i) \equiv G'_i - G_i \quad \text{and} \quad d_1(\xi_i) \equiv K'_i - K_i \pmod{I_2},$$

for $i \in \{0, 1, 2\}$, and for $g_i, \xi_i \in \Omega^1 E(3)_*$ given by

$$(4.22) \quad g_i = t_1^{p^i} t_2^{p^i} - \frac{1}{2} t_1^{p^{i+1}+2p^i} \quad \text{and} \quad \xi_i = t_1^{p^{i+1}} t_2^{p^i} - \frac{1}{2} t_1^{2p^{i+1}+p^i}.$$

We also have a similar relation

$$(4.23) \quad d_1(t_1^p t_2) \equiv -(t_1^p \otimes t_2 + c t_2 \otimes t_1^p) - 2K_0 \pmod{I_2}.$$

Lemma 4.24 In $\Omega^1 E(3)_*$, put

$$\omega_1 = \eta_R(v_3) t_2 - v_2 t_3 + v_2^p t_1 t_2, \quad \omega_2 = \frac{1}{2} \eta_R(v_3) t_1^{2p} - v_2^p \xi_0 \quad \text{and} \quad \tilde{\omega}_2 = -w_3 - v_2^{pe(2)} t_1^p t_2.$$

Then, modulo I_2 ,

$$d_1(\omega_1) \equiv -t_1 \otimes z - v_2^p b_{1,1} - 2v_2^p G_0, \quad d_1(\omega_2) \equiv -t_1^p \otimes z - v_2 G_1 + v_2^p K_0,$$

$$d_1(\tilde{\omega}_2) \equiv v_2^{p^2} z \otimes t_1^p + 2v_2^{p^2+p} K_0 + v_2^{e(3)} b_{1,0}.$$

Proof In this proof we consider congruences modulo I_2 . A routine calculation shows the congruence for $d_1(\omega_1)$:

$$d_1(\eta_R(v_3) t_2) \equiv -t_1 \otimes (z + \underbrace{v_2 t_2^p}_a + \underbrace{v_2^p t_2 - v_2^p t_1^{p+1}}_c) - t_2 \otimes (\underbrace{v_2 t_1^p}_b - \underbrace{v_2^p t_1}_d), \quad (\text{by (4.2) and (4.19)})$$

$$d_1(-v_2 t_3) \equiv v_2 (\underbrace{t_1 \otimes t_2^p}_a + \underbrace{t_2 \otimes t_1^{p^2}}_b - v_2 b_{1,1}), \quad (\text{by (4.8)})$$

$$d_1(v_2^p t_1 t_2) \equiv -v_2^p (\underbrace{t_1 \otimes t_2}_a + \underbrace{t_2 \otimes t_1}_a + \underbrace{t_1^2 \otimes t_1^p}_c + \underbrace{t_1 \otimes t_1^{p+1}}_c). \quad (\text{by (4.8) and (4.2)})$$

Here the underlined terms with the same label cancel each other and the wavy underlined terms make $-2v_2^p G_0$.

For $d_1(\omega_2)$, we calculate

$$d_1\left(\frac{1}{2}\eta_R(v_3)t_1^{2p}\right) \equiv -t_1^p \otimes (z + \underbrace{v_2 t_2^p}_G - \underbrace{v_2^p c t_2}_{K'}) - \underbrace{\frac{1}{2}v_2 t_1^{2p} \otimes t_1^{p^2}}_G + \underbrace{\frac{1}{2}v_2^p t_1^{2p} \otimes t_1}_{K'}. \quad (\text{by (4.2) and (4.19)})$$

Add $d_1(-v_2^p \mathfrak{k}_0)$, and we obtain the desired congruence by (4.21).

We verify $d_1(\tilde{\omega}_2)$ by

$$d_1(w_3) \equiv -v_2^{pe(2)} w_1 \otimes t_2 + v_2^{p^2} w_2 \otimes t_1^p - v_2^{e(3)} b_{1,0} \quad (\text{by Corollary 4.17})$$

$$\equiv -\frac{v_2^{pe(2)}(-t_1^p) \otimes t_2}{a} + v_2^{p^2}(-z + \frac{v_2^p c t_2}{b}) \otimes t_1^p - v_2^{e(3)} b_{1,0}, \quad (\text{by (4.18) and (4.15)})$$

$$d_1(v_2^{pe(2)} t_1^p t_2) \equiv -v_2^{pe(2)} \left(\frac{(t_1^p \otimes t_2 + c t_2 \otimes t_1^p)}{a} + 2K_0 \right). \quad (\text{by (4.23)}) \quad \square$$

5 The elements x_i and deriving elements y_i and y'_i

In [2, (5.11)], Miller, Ravenel and Wilson introduced elements $x_{3,i} \in v_3^{-1}\text{BP}_*$. We refine them, and define the elements $x_i \in E(3)_*$ by

$$\begin{aligned} x_i &= v_3^{p^i} \quad \text{for } i = 0, 1, 2, \\ x_3 &= x_2^p - v_2^{p^3-1} v_3^{(p-1)p^2+1}, \\ x_4 &= x_3^p - v_2^{e(2)p^3-p-1} v_3^{(p^2-e(2))p^2+p+1}, \\ x_{2k+1} &= x_{2k}^p - v_2^{pa_{2k}-1} x_{2k-1}^{(p-1)p} v_3 - v_2^{e(3)p^{2k-1}-e(3)} v_3^{(p^2-e(2))p^{2k-1}+p+1}, \\ x_{2k+2} &= x_{2k+1}^p - 2v_2^{e(3)p^{2k}-e(3)} v_3^{(p^2-e(2))p^{2k}+p+1}, \end{aligned}$$

for $k \geq 2$.

Lemma 5.1 (see [9, Proposition 3.1]) *In $\Omega^1 E(3)_*$, we have*

$$\begin{aligned} d_0(x_0) &\equiv v_2 t_1^{p^2} - v_2^p t_1 \quad \text{mod } I_2, \\ d_0(x_1) &\equiv v_2^p v_3^{p-1} t_1 - v_2^{p+1} v_3^{-1} t_2^{p^2} \quad \text{mod } J_{2p}, \\ d_0(x_i) &\equiv v_2^{a_i} (x_{i-1}^{p-1} t_1^{p^{\varepsilon_i}} + B_i) \quad \text{mod } J_{e(3)p^{i-2}} \quad \text{for } i \geq 2. \end{aligned}$$

Here $\varepsilon_i = \frac{1}{2}(1 + (-1)^i)$, and the B_i are given by

i	2	3	$2k$	$2k+1$
B_i	$-v_2^p v_3^{c(2)} t_2$	$v_2^{p^2-p} v_3^{c(3)} (z - v_2^p t_1^{p+1})$	$v_2^{a_{2k}-1-p} v_3^{c(2k)} (z - v_2^p t_2)$	$v_2^{a_{2k}-p} v_3^{c(2k+1)} (2z - v_2^p c t_2)$

for $c(k) = (p^2 - p - 1)p^{k-2}$. For $i \geq 4$, add $v_2^{a_{i-1}+1} v_3^{c(i)} Z'$ to B_i if we consider the congruence modulo $J_{e(3)p^{i-2}+1}$. Here Z' is a cocycle homologous to aZ for some $a \in \mathbb{Z}/p$.

Proof This follows from a routine calculation: For $i \leq 2$, it follows from (4.4) and from (4.6).

We obtain $d_0(x_3)$ from (4.19) and

$$d_0(v_3^{(p-1)p^2+1}) \equiv v_3^{(p-1)p^2} (v_2 t_1^{p^2} - v_2^p t_1) - v_2^{a_2} v_3^{(p-1)p^2-p} (t_1^p \eta_R(v_3) - v_2^p t_2) \pmod{J_{e(3)}}$$

by (4.2), (4.4) and the congruence on $d_0(x_2)$. We note that $\eta_R(v_3^{p+1}) = v_3^{p+1} + v_2 z^p - v_2^{p^2} z$ by [2, (3.20)], and obtain

$$\begin{aligned} d_0(v_3^{(p^2-e(2))p^2+p+1}) \\ \equiv v_3^{(p^2-e(2))p^2} (v_2 z^p - v_2^p z) - v_2^{a_2} v_3^{(p^2-e(2))p^2-p} t_1^p (v_3^{p+1} + v_2 z^p) + v_2^{p^2+p} v_3^{(p^2-e(2))p^2} t_2 \pmod{J_{e(3)}}. \end{aligned}$$

The congruence on $d_0(x_4)$ follows from this and the congruence on $d_0(x_3)$, together with the definition of the element x_3 .

Inductively suppose that

$$d_0(x_{2k}) \equiv v_2^{a_{2k}} x_{2k-1}^{p-1} t_1^p + v_2^{e(3)p^{2k-2}-e(2)} v_3^{(p^2-e(2))p^{2k-2}} (z - v_2^p t_2) \pmod{J_{e(3)}p^{2k-2}}.$$

Then we calculate

$$\begin{aligned} d_0(x_{2k}^p) &\equiv \frac{v_2^{pa_{2k}} x_{2k-1}^{(p-1)p} t_1^{p^2}}{a} + v_2^{e(3)p^{2k-1}-e(2)p} v_3^{(p^2-e(2))p^{2k-1}} \left(\frac{z^p}{b} - \frac{v_2^{p^2} t_2^p}{c} \right) \\ d_0(-v_2^{pa_{2k}-1} x_{2k-1}^{(p-1)p} v_3) \\ &\equiv -v_2^{pa_{2k}-1} x_{2k-1}^{(p-1)p} \frac{(v_2 t_1^{p^2} - v_2^p t_1)}{a} + v_2^{e(3)p^{2k-1}-p-1} x_{2k-1}^{p^2-p-1} (z + \frac{v_2 t_2^p}{c} - v_2^p c t_2), \end{aligned}$$

where the second congruence follows by (4.2) and (4.19), and

$$d_0(-v_2^{e(3)p^{2k-1}-e(3)} v_3^{(p^2-e(2))p^{2k-1}+p+1}) \equiv -v_2^{e(3)p^{2k-1}-e(3)} v_3^{(p^2-e(2))p^{2k-1}} \left(\frac{v_2 z^p}{b} - v_2^{p^2} z \right).$$

Therefore

$$d_0(x_{2k+1}) \equiv v_2^{pa_{2k}+p-1} x_{2k-1}^{(p-1)p} t_1^p + v_2^{e(3)p^{2k-1}-e(2)} v_3^{(p^2-e(2))p^{2k-1}} (2z - v_2^p c t_2).$$

Also

$$\begin{aligned} d_0(x_{2k+1}^p) &\equiv v_2^{pa_{2k+1}} x_{2k-1}^{(p-1)p^2} t_1^p + v_2^{e(3)p^{2k}-e(2)p} v_3^{(p^2-e(2))p^{2k}} (2z^p - v_2^{p^2} c t_2^p) \\ &\equiv v_2^{pa_{2k+1}} (x_{2k+1}^{p-1} - v_2^{pa_{2k}-1} x_{2k-1}^{(p^2-p-1)p} v_3) t_1^p + v_2^{e(3)p^{2k}-e(2)p} v_3^{(p^2-e(2))p^{2k}} (2z^p - v_2^{p^2} c t_2^p) \\ &\equiv v_2^{pa_{2k+1}} x_{2k+1}^{p-1} t_1^p + v_2^{e(3)p^{2k}-e(2)p} v_3^{(p^2-e(2))p^{2k}} (2z^p - v_2^{p^2-1} z - v_2^{p^2+p-1} t_2) \end{aligned}$$

and

$$d_0(-2v_2^{e(3)p^{2k}-e(3)} v_3^{(p^2-e(2))p^{2k}+p+1}) \equiv -2v_2^{e(3)p^{2k}-e(3)} v_3^{(p^2-e(2))p^{2k}} (v_2 z^p - v_2^{p^2} z).$$

Therefore

$$d_0(x_{2k+2}) \equiv v_2^{pa_{2k+1}} x_{2k+1}^{p-1} t_1^p + v_2^{e(3)p^{2k}-e(2)} v_3^{(p^2-e(2))p^{2k}} (z - v_2^p t_2).$$

These complete the induction.

Put $d_0(x_i) \equiv v_2^{a_i} (x_{i-1}^{p-1} t_1^{p^{e_i}} + B_i + v_2^{a_{i-1}+1} C) \pmod{J_{e(3)}p^{i-1}+1}$ for a cochain C . It is easy to see

$$d_1(v_2^{a_i} (x_{i-1}^{p-1} t_1^{p^{e_i}} + B_i)) \equiv 0 \pmod{J_{e(3)}p^{i-1}+1}.$$

It follows that C is a cocycle of $\Omega^1 M_3^0$, and so C represents a cohomology class $av_3^{c(i)} \zeta_3 \in H^1 M_3^0$ for some $a \in \mathbb{Z}/p$ by (2.3). \square

Put

$$d_0(x_i) \equiv v_2^{a_i} A_i + v_2^{a_i} B_i \quad \text{for } A_i = x_{i-1}^{p-1} t_1^{p^{\varepsilon_i}},$$

where $\varepsilon_i = \frac{1}{2}(1 + (-1)^i)$. We introduce elements $y_{s,i}$ and $y'_{s,i} \in \Omega^1 E(3)_*$ by

$$y_{s,i} = x_i^s t_1^{p^{\varepsilon_i+1}} - s x_i^{s-p+1} B_{i+1} \quad \text{and} \quad y'_{s,i} = x_i^s t_1^{p^{\varepsilon_i}} + \frac{1}{2} s v_2^{a_i} x_i^{s-1} A_i t_1^{p^{\varepsilon_i}}.$$

Lemma 5.2 For the elements $y_{s,i}$ and $y'_{s,i}$,

$$\begin{aligned} d_1(y_{s,0}) &\equiv s(s+1)v_2^2 v_3^{s-p-1} G_2, & d_1(y_{s,1}) &\equiv s(s+1)v_2^{2p} v_3^{s-p-2} G_0, \\ d_1(y_{s,2}) &\equiv -s(s+1)v_2^{2p^2-p} v_3^{s-p^2-2p} (t_1^p \otimes z - v_2^p x), \\ d_1(y_{s,i}) &\equiv \begin{cases} -s(s+1)v_2^{2a_{2k+1}-p} x_{2k}^{s-p-2} (t_1 \otimes z - v_2^p G_0) & i = 2k+1, \\ -s(s+1)v_2^{2a_{2k+2}-p} x_{2k+1}^{s-p-2} (2t_1^p \otimes z - v_2^p K'_0) & i = 2k+2, \end{cases} \\ d_1(y'_{s,1}) &\equiv -s v_2^{p+1} v_3^{s-p-2p} K_2, & d_1(y'_{s,2}) &\equiv -s v_2^{p^2+p} v_3^{s-p^2-p-1} K_0, \\ d_1(y'_{s,3}) &\equiv s v_2^{a_3+p^2-p} v_3^{s-p^3-p^2-p} (z \otimes t_1 - v_2^p x'), \\ d_1(y'_{s,i}) &\equiv \begin{cases} s v_2^{e(3)p^{i-2}-p-1} v_3^{(sp^2-p-1)p^{2k-2}} (z \otimes t_1^p - v_2^p K_0) & i = 2k, \\ s v_2^{e(3)p^{i-2}-p-1} v_3^{(sp^2-p-1)p^{2k-1}} (2z \otimes t_1 - v_2^p G'_0) & i = 2k+1. \end{cases} \end{aligned}$$

Here $x = (t_2 + t_1^{p+1}) \otimes t_1^p + t_1^p \otimes t_1^{p+1} + \frac{1}{2} t_1^{2p} \otimes t_1$ and $x' = t_1^{p+1} \otimes t_1 + \frac{1}{2} t_1^p \otimes t_1^2$, and these congruences are considered modulo J_{a+1} , where a is the largest power of v_2 in each congruence. Furthermore, replace K'_0 and K_0 in the congruences on $d_1(y_{s,2k+2})$ and $d_1(y'_{s,2k})$ by $K'_0 + v_2 t_1^p \otimes Z'$ and $K_0 + v_2 Z' \otimes t_1^p$, respectively, if we consider the congruences modulo J_{a+2} .

Proof We note that

$$\begin{aligned} d_1(B_{i+1}) &\equiv -d_1(A_{i+1}) \equiv -d_0(x_i^{p-1}) \otimes t_1^{p^{\varepsilon_i+1}} \pmod{I_2}, \\ d_0(x_i^s) + s x_i^{s+1-p} d_0(x_i^{p-1}) &\equiv \binom{s+1}{2} x_i^{s-2} d_0(x_i)^2 \pmod{J_{3a_i}}. \end{aligned}$$

Indeed, $d_0(x_i^s) \equiv s x_i^{s-1} d_0(x_i) + \binom{s}{2} x_i^{s-2} d_0(x_i)^2 \pmod{J_{3a_i}}$. Also,

$$d_1(A_i t_1^{p^{\varepsilon_i}}) \equiv d_0(x_{i-1}^{p-1}) \otimes t_1^{2p^{\varepsilon_i}} - 2 x_{i-1}^{p-1} t_1^{p^{\varepsilon_i}} \otimes t_1^{p^{\varepsilon_i}} \equiv d_0(x_{i-1}^{p-1}) \otimes t_1^{2p^{\varepsilon_i}} - 2 A_i \otimes t_1^{p^{\varepsilon_i}} \pmod{J_{a_{i-1}+2}}.$$

Then we calculate

$$d_1(y_{s,i}) \equiv d_0(x_i^s) \otimes t_1^{p^{\varepsilon_i+1}} - s d_0(x_i^{s+1-p}) \otimes B_{i+1} + s x_i^{s+1-p} d_0(x_i^{p-1}) \otimes t_1^{p^{\varepsilon_i+1}} \quad (\text{by (4.2)})$$

$$\equiv \binom{s+1}{2} x_i^{s-2} d_0(x_i)^2 \otimes t_1^{p^{\varepsilon_i+1}} - s(s+1) x_i^{s-p} d_0(x_i) \otimes B_{i+1} \pmod{J_{2a_i+p}},$$

$$d_1(y'_{s,i}) \equiv s x_i^{s-1} d_0(x_i) \otimes t_1^{p^{\varepsilon_i}} + \frac{1}{2} s v_2^{a_i} x_i^{s-1} d_0(x_{i-1}^{p-1}) \otimes t_1^{2p^{\varepsilon_i}} - s v_2^{a_i} x_i^{s-1} A_i \otimes t_1^{p^{\varepsilon_i}} \quad (\text{by (4.2)})$$

$$\equiv s v_2^{a_i} x_i^{s-1} (B_i \otimes t_1^{p^{\varepsilon_i}} + \frac{1}{2} d_0(x_{i-1}^{p-1}) \otimes t_1^{2p^{\varepsilon_i}}) \pmod{J_{e(3)p^{i-2}+1}}.$$

Now we obtain the lemma from Lemma 5.1. \square

6 Proof of Lemma 3.4

In this section, we define the cochains $(t_1^{p^i})_s$ and verify their d_1 -differential.

6.1 The cochains $(t_1)_{sp^{2k}}$ and $(t_1^p)_{sp^{2k+1}}$ for $s \in \mathbb{Z}_0$

We define the cochains by

$$\begin{aligned}(t_1)_s &= y_{s,0}, (t_1^p)_{sp} = y_{s,1}, \quad (t_1)_{sp^2} = y_{s,2} - s(s+1)v_2^{2p^2-p}v_3^{sp^2-2p}\omega_2, \\ (t_1^p)_{sp^{2k+1}} &= y_{s,2k+1} - s(s+1)v_2^{2a_{2k+1}-p}x_{2k}^{sp-2}\omega_1, \\ (t_1)_{sp^{2k+2}} &= y_{s,2k+2} - s(s+1)v_2^{2a_{2k+2}-p^2-p}x_{2k+1}^{sp-2}(2\tilde{\omega}_2 + v_2^{p^2}(2zt_1^p + v_2^p\epsilon_0)),\end{aligned}$$

for $k \geq 1$. Then the lemma for this case follows immediately from Lemmas 5.2, 5.1 and 4.24 together with (4.21). Note also $2a_{2k+1} - p + 2 = 2pa_{2k} + p$. For example, for the case $p = 3$ and $k \geq 2$, we compute modulo $J_{2a_{2k}+2}$

$$\begin{aligned}d_1((t_1)_{3^{2k}s}) &\equiv d_1(y_{s,2k}) - s(s+1)v_2^{2a_{2k}-12}x_{2k-1}^{3s-2}d_1(2\tilde{\omega}_2 + v_2^9(2zt_1^3 + v_2^3\epsilon_0)) \\ &\equiv -s(s+1)v_2^{2a_{2k}-3}x_{2k-1}^{3s-2}\frac{(2t_1^3 \otimes z - v_2^3(K'_0 + v_2t_1^3 \otimes Z'))}{a} - s(s+1)v_2^{2a_{2k}-12}x_{2k-1}^{3s-2} \\ &\quad \cdot \left(2\left(\frac{v_2^9z \otimes t_1^3}{c} + \frac{2v_2^{12}K_0}{d} + v_2^{13}b_{1,0}\right) + v_2^9\left(-2\left(\frac{z \otimes t_1^3}{c} + \frac{t_1^3 \otimes z}{a}\right) + v_2^3\left(\frac{K'_0}{b} - \frac{K_0}{d}\right)\right)\right),\end{aligned}$$

where the second equivalence follows by Lemmas 5.2 and 4.24, and Equation (4.21)

6.2 The cochains $(t_1)_{sp^{2k}}$ and $(t_1^p)_{sp^{2k+1}}$ for $s \in \mathbb{Z}_1$

We put $s = tp^2 - 1$, and define the cochains $(t_1)_{(tp^2-1)p^{2k}}$ and $(t_1^p)_{(tp^2-1)p^{2k+1}}$ by

$$\begin{aligned}v_2^{a_{2k+1}}(t_1)_{(tp^2-1)p^{2k}} &= -v_3^{(t-1)p^{2k+2}}w^{p^{2k+1}} - d_0(v_2^{p^{2k+1}-p^{2k-2}}v_3^{(tp^2-1)p^{2k}}\sigma_k) \\ &\quad + v_2^{p^{2k+2}-p^{2k-1}}v_3^{(tp-1)p^{2k+1}}W_k, \\ (t_1^p)_{(tp^2-1)p^{2k+1}} &= (t_1)_{(tp^2-1)p^{2k}}^p,\end{aligned}$$

for the elements σ_k in Lemma 4.5, w in (4.7) and W_k in Corollary 4.11. Then this case follows from Lemmas 4.5 and 4.10, Corollary 4.11 and (2.8). We also use relations $w^{p^{2k+1}} \equiv -v_2^{p^{2k+1}}v_3^{p^{2k+2}-p^{2k}}t_1^{p^{2k}} \pmod{J_{a_{2k+1}+1}}$ by Lemma 4.10 and (4.6), and $b_{1,2}^{p^{2k+1}} \equiv b_{1,2k+3} \equiv v_3^{(p-1)p^{2k+1}}b_{1,2k} \pmod{I_3}$ by (4.6). For example,

$$\begin{aligned}v_2^{a_{2k+1}}(t_1)_{(tp^2-1)p^{2k}} &\equiv v_3^{(t-1)p^{2k+2}}(v_2^{p^{2k+1}}v_3^{p^{2k+2}-p^{2k}}t_1^{p^{2k}}) \quad (\text{by Lemma 4.5}) \\ &\quad - v_2^{p^{2k+1}-p^{2k-2}}v_3^{(tp^2-1)p^{2k}}(v_2^{p^{2k-2}}t_1^{p^{2k}} - v_2^{a_{2k}-1}t_1) \\ &\equiv v_2^{a_{2k+1}}v_3^{(tp^2-1)p^{2k}}t_1 \pmod{J_{a_{2k+1}+1}},\end{aligned}$$

since $p^{2k+1} - p^{2k-2} + a_{2k-1} = a_{2k+1}$ in (2.8), and

$$\begin{aligned} & v_2^{a_{2k+1}} d_1((t_1)_{(tp^2-1)p^{2k}}) \\ & \equiv \frac{v_2^{p^{2k+2}} v_3^{(t-1)p^{2k+2}} b_{1,2}^{p^{2k+1}}}{a} + v_2^{p^{2k+2}-p^{2k-1}} v_3^{(tp-1)p^{2k+1}} \left(-\frac{v_2^{p^{2k-1}} b_{1,2k}}{a} + v_2^{a_{2k}} b_{1,0} \right) \pmod{J_{a_{2k+2}+1}} \end{aligned}$$

by Lemma 4.10 and Corollary 4.11. Since $p^{2k+2} - p^{2k-1} + a_{2k} = a_{2k+2}$ in (2.8), we obtain the case for $(t_1)_{sp^{2k}}$.

6.3 The cochains $(t_1)_{sp^{2k+1}}$ and $(t_1^p)_{sp^{2k}}$ for $s \in \mathbb{Z}^{(p)}$

We begin by defining

$$(t_1^p)_s = v_3^s t_1^p + s v_2 v_3^{s-1} c t_2^p - s(s-1) v_2^2 v_3^{s-2} \mathfrak{t}_1.$$

Then we calculate by (4.2), (4.4), (4.8) and (4.22), and obtain

$$d_1((t_1^p)_s) \equiv s(s-1) v_2^2 v_3^{s-2} K_1 \pmod{J_3}.$$

Now we consider the cases for $p \mid s(s-1)$.

6.3.1 The cochains $(t_1^p)_{tp^k+1}$ for $k \geq 1$

We define the cochains by

$$\begin{aligned} (t_1^p)_{tp+1} &= v_3^{tp} z + t v_2^p v_3^{tp} t_2 - t v_2^{p+1} v_3^{tp-p} c t_3^p, \\ (t_1^p)_{tp^2+1} &= x_2^t z + t v_2^{a_2} v_3^{(tp-1)p} \omega_2, \\ (t_1^p)_{tp^{2k+1}+1} &= x_{2k+1}^t z + t v_2^{a_{2k+1}} v_3^{(tp-1)p^{2k}} \omega_1 + t v_2^{a_{2k}+p+1} (t_1^p)_{(tp^2-1)p^{2k-1}}, \\ (t_1^p)_{tp^{2k+2}+1} &= x_{2k+2}^t z + t v_2^{a_{2k+2}-p^2} v_3^{(tp-1)p^{2k+1}} (\tilde{\omega}_2 + v_2^2 z t_1^p), \end{aligned}$$

in $\Omega^1 E(3)_*$ for $k \geq 1$, $t \in \mathbb{Z}^{(p)}$, x_n in 5.1, z in (4.18) and ω_i in Lemma 4.24. We verify this case by a routine calculation using (4.2), (4.4), (4.18), (4.8) and (4.9). We see that

$$t_1^{p^3} \otimes z \equiv \eta_R(v_3) t_1^{p^3} \otimes t_1^p + v_2 t_1^{p^3} \otimes c t_2^p - v_2^p v_3^{p-1} t_1 \otimes t_2 \quad \text{and} \quad \eta_R(v_3) t_1^{p^3} \equiv v_3^p t_1 + v_2 c t_2^{p^2} \pmod{J_{p+1}}$$

by (4.18), (4.4) and (4.6). It follows that $t_1^{p^3} \otimes z \equiv -d_1(v_3^p t_2) + v_2 d_1(c t_3^p) \pmod{J_{p+1}}$, and then $d_1(t_1^{tp} z) \equiv t v_2^p v_3^{tp-p} (-d_1(v_3^p t_2) + v_2 d_1(c t_3^p)) + \binom{t}{2} v_2^{2p} v_3^{tp-1} t_1^2 \otimes t_1^p \pmod{J_{2p+1}}$. Thus we obtain $d_1((t_1^p)_{tp+1})$.

The congruences on $d_1((t_1^p)_{tp^k+1})$ for $k \geq 2$ follow directly from Lemmas 5.1 and 4.24 and the results on $d_1((t_1^p)_{(tp^2-1)p^{2k-1}})$ shown in the previous subsection. For example,

$$\begin{aligned} d_1((t_1^p)_{tp^{2k+1}+1}) &\equiv d_1(x_{2k+1}^t) \otimes z + t v_2^{a_{2k+1}} v_3^{(tp-1)p^{2k}} d_1(\omega_1) + t v_2^{a_{2k}+p+1} d_1((t_1^p)_{(tp^2-1)p^{2k-1}}) \\ &\equiv \frac{t v_2^{a_{2k+1}} v_3^{(tp-1)p^{2k}} t_1 \otimes z + t v_2^{a_{2k+1}} v_3^{(tp-1)p^{2k}} (-\frac{t_1 \otimes z}{a} - \frac{v_2^2 b_{1,1}}{b} - 2 v_2^p G_0)}{b} \\ &\quad + \frac{t v_2^{a_{2k}+p+1} + p a_{2k} - p a_{2k-1}}{b} v_3^{(tp-1)p^{2k}} b_{1,1} \pmod{J_{a_{2k+1}+p+1}}, \end{aligned}$$

where the second equivalence follows by Lemmas 5.1, 4.24 and 3.4(2).

6.3.2 The cochains $(t_1^p)_{tp^k+e(k)}$ for $k \geq 2$ We put $r = 2n - 1 + \varepsilon$ ($\varepsilon \in \{0, 1\}$), and

$$(t_1^p)'_{tp^r+e(r)} = x_r^t(w_{r+1} + v_2^{p^r-p^{r-3}}w_r\eta_R(\sigma_{n-1}^{p^\varepsilon}) + v_2^{a_r}w_rt_1^{p^\varepsilon})$$

for w_r in (4.15). Note that $w_r \equiv v_3^{pe(r-2)}w_2 \equiv -v_3^{pe(r-2)}z \pmod{J_p}$ by (4.15) and (4.18). Then $(t_1^p)'_{tp^r+e(r)} \equiv x_r^t w_{r+1} \equiv -v_3^{tp^r+e(r)}t_1^p \pmod{I_3}$. Furthermore, we calculate

$$\begin{aligned} d_1((t_1^p)'_{tp^r+e(r)}) &\equiv \underbrace{tv_2^{a_r}v_3^{(tp-1)p^{r-1}}t_1^{p^\varepsilon} \otimes w_{r+1}}_{\text{(by Lemmas 5.1 and 4.5 and Corollary 4.17)}} \\ &\quad + x_r^t \left(\frac{v_2^{p^r}w_r \otimes t_1^{p^{r-1}}}{a} - v_2^{p^r-p^{r-3}}w_r \otimes \left(\frac{v_2^{p^{2n-4+\varepsilon}}t_1^{p^{2n-2+\varepsilon}}}{a} - \frac{v_2^{a_{2n-3+\varepsilon}}t_1^{p^\varepsilon}}{b} \right) \right. \\ &\quad \left. - v_2^{a_r} \frac{(w_r \otimes t_1^{p^\varepsilon})}{b} + \underbrace{t_1^{p^\varepsilon} \otimes w_r \right) \\ &\equiv -(t-1)v_2^{a_r}v_3^{tp^r+pe(r-2)}t_1^{p^\varepsilon} \otimes z \pmod{J_{a_r+p}} \end{aligned}$$

together with (4.2) and (2.8). This case now follows from Lemma 4.24 by setting $(t_1^p)_{tp^r+e(r)} = -(t_1^p)'_{tp^r+e(r)} + (t-1)v_2^{a_r}v_3^{tp^r+pe(r-2)}\omega_{1+\varepsilon}$.

6.3.3 The cochains $(t_1^p)_{sp^{2k}}$ for $k \geq 1$ and $(t_1)_{sp^{2k+1}}$ for $k \geq 0$ We define $(t_1^{\varepsilon i})_{spi}$ by

$$\begin{aligned} (t_1)_{sp} &= y'_{s,1}, \quad (t_1^p)_{sp^2} = y'_{s,2}, \quad (t_1)_{sp^3} = y'_{s,3} + sv_2^{e(3)p-p-1}v_3^{(sp^2-p-1)p}(zt_1 - \omega_1), \\ (t_1^p)_{sp^4} &= y'_{s,4} - \frac{1}{2}sv_2^{e(3)p^2-p^2-p-1}v_3^{(sp^2-p-1)p^2}(\tilde{\omega}'_2 - v_2^{p^2}zt_1^p), \\ (t_1)_{sp^{2k+1}} &= y'_{s,2k+1} + 2sv_2^{e(3)p^{2k-1}-p-1}v_3^{(sp^2-p-1)p^{2k-1}}(zt_1 - \omega_1), \\ (t_1^p)_{sp^{2k+2}} &= y'_{s,2k+2} - sv_2^{e(3)p^{2k}-p^2-p-1}v_3^{(sp^2-p-1)p^{2k}}\tilde{\omega}_2, \end{aligned}$$

where $\tilde{\omega}'_2 = \tilde{\omega}_2 - v_2^{p^2+p}t_1^p t_2 - v_2^{e(3)}v_3^{-p^2}ct_4^p$. Except for $d_1((t_1^p)_{sp^4})$, the lemma for this case follows from Lemmas 5.2 and 4.24 with (4.2).

For $d_1((t_1^p)_{sp^4})$, we make a calculation:

$$\begin{aligned} \tilde{\omega}_2 &\equiv -w_3 && \text{(by Lemma 4.24)} \\ &\equiv t_1^p\eta_R(v_3^{p+1}) - v_2t_2^p\eta_R(v_3^p) + v_2^{p+1}t_3^p && \text{(by (4.15) and Proposition 4.12)} \\ &\equiv (z + \frac{v_2t_2^p}{a} - v_2^pct_2)\eta_R(v_3^p) - \frac{v_2t_2^p\eta_R(v_3^p)}{a} + v_2^{p+1}t_3^p && \text{(by (4.19))} \\ &\equiv v_3^p(z + v_2^pt_2) - v_2^{p+1}ct_3^p \pmod{J_{p+2}}. && \text{(by (4.4))} \end{aligned}$$

Applying the Hopf conjugation c to the congruences of (4.6) shows the relations

$$(6.1) \quad t_1^{p^3}\eta_R(v_3) \equiv v_3^pt_1 + v_2ct_2^{p^2} \quad \text{and} \quad ct_2^{p^3}\eta_R(v_3) \equiv v_3^{p^2}ct_2 - v_2ct_3^{p^2} \pmod{J_{p+1}}.$$

Then, modulo J_{p+2} ,

$$\begin{aligned}
 t_1^{p^4} \otimes v_3^p z &\equiv t_1^{p^4} \eta_R(v_3)^p \otimes z \\
 &\equiv (v_3^{p^2} t_1^p + v_2^p c t_2^{p^3}) \otimes z && \text{(by (6.1))} \\
 &\equiv v_3^{p^2} t_1^p \otimes z + v_2^p c t_2^{p^3} \eta_R(v_3) \otimes t_1^p + v_2^{p+1} c t_2^{p^3} \otimes c t_2^p && \text{(by (4.18))} \\
 &\equiv v_3^{p^2} t_1^p \otimes z + v_2^p (v_3^{p^2} c t_2 - v_2 c t_3^{p^2}) \otimes t_1^p + v_2^{p+1} c t_2^{p^3} \otimes c t_2^p, && \text{(by (6.1))} \\
 t_1^{p^4} \otimes v_2^p v_3^p t_2 &\equiv v_2^p t_1^{p^4} \eta_R(v_3)^p \otimes t_2 && \text{(by (6.1))} \\
 &\equiv v_2^p v_3^{p^2} t_1^p \otimes t_2.
 \end{aligned}$$

Therefore

$$\begin{aligned}
 d_1(v_2^p v_3^{p^2} t_1^p t_2 + v_2^{p+1} c t_4^p) &\equiv -v_2^p v_3^{p^2} \underbrace{(t_1^p \otimes t_2 + t_2 \otimes t_1^p + t_1^{p+1} \otimes t_1^p + t_1 \otimes t_1^{2p})}_a \\
 &\quad + v_2^{p+1} \underbrace{(t_1^{p^4} \otimes c t_3^p - c t_2^{p^3} \otimes c t_2^p + c t_3^{p^2} \otimes t_1^p - v_3^p b_{1,2}^p)}_{b \quad c \quad d}, \\
 t_1^{p^4} \otimes \tilde{\omega}_2 &\equiv v_3^{p^2} t_1^p \otimes z + v_2^p \underbrace{(v_3^{p^2} c t_2 - v_2 c t_3^{p^2})}_d \otimes t_1^p + \underbrace{v_2^{p+1} c t_2^{p^3} \otimes c t_2^p}_c \\
 &\quad + \underbrace{v_2^p v_3^{p^2} t_1^p \otimes t_2}_a - \underbrace{v_2^{p+1} t_1^{p^4} \otimes c t_3^p}_b,
 \end{aligned}$$

where the first equation follows from (4.2), (4.8) and (4.9). The sum of the wavy underlined terms is $-v_2^p v_3^{p^2} (2t_2 \otimes t_1^p + t_1 \otimes t_1^{2p}) = -v_2^p v_3^{p^2} K_0$, and $b_{1,2}^p \equiv v_3^{p^2-p} b_{1,0} \pmod{I_3}$ by (4.6). Then, modulo J_{p+2} ,

$$(6.2) \quad t_1^{p^4} \otimes \tilde{\omega}_2 + d_1(v_2^p v_3^{p^2} t_1^p t_2 + v_2^{p+1} c t_4^p) \equiv v_3^{p^2} t_1^p \otimes z - 2v_2^p v_3^{p^2} K_0 - v_2^{p+1} v_3^{p^2} b_{1,0}.$$

Now we calculate $d_1((t_1^p)_{s p^4}) \pmod{J_{e(3)p^2+1}}$ for odd prime p as

$$d_1(y'_{s,4}) \equiv s v_2^{e(3)p^2-p-1} v_3^{(s p^2-p-1)p^2} \underbrace{(z \otimes t_1^p - v_2^p (K_0 + v_2 Z' \otimes t_1^p))}_a \quad \text{(by 5.2)}$$

and

$$\begin{aligned}
 &d_1\left(-\frac{1}{2} s v_2^{e(3)p^2-p^2-p-1} v_3^{(s p^2-p-1)p^2} (\tilde{\omega}'_2 - v_2^{p^2} z t_1^p)\right) \\
 &\equiv \frac{1}{2} s v_2^{e(3)p^2-p-1} v_3^{(s p^2-p-2)p^2} \underbrace{t_1^{p^4} \otimes \tilde{\omega}_2}_{\sim} - \frac{1}{2} s v_2^{e(3)p^2-p^2-p-1} v_3^{(s p^2-p-1)p^2} \\
 &\quad \cdot (v_2^{p^2} z \otimes t_1^p + 2v_2^{p^2+p} K_0 + v_2^{p^2+p+1} b_{1,0} - \underbrace{d_1(v_2^{p^2+p} t_1^p t_2 + v_2^{e(3)} v_3^{-p^2} c t_4^p)}_{\sim} + v_2^{p^2} (z \otimes t_1^p + t_1^p \otimes z)) \\
 &\equiv \frac{1}{2} s v_2^{e(3)p^2-p-1} v_3^{(s p^2-p-2)p^2} \underbrace{(v_3^{p^2} t_1^p \otimes z - 2v_2^p v_3^{p^2} K_0 - v_2^{p+1} v_3^{p^2} b_{1,0})}_b \quad \text{(by (6.2))} \\
 &\quad - \frac{1}{2} s v_2^{e(3)p^2-p^2-p-1} v_3^{(s p^2-p-1)p^2} \underbrace{(v_2^{p^2} z \otimes t_1^p + 2v_2^{p^2+p} K_0 + v_2^{p^2+p+1} b_{1,0} + v_2^{p^2} \underbrace{(z \otimes t_1^p + t_1^p \otimes z)}_{\sim})}_a \underbrace{\quad}_b.
 \end{aligned}$$

6.4 The cochains $(t_1^{p^2})_{tp-1}$ for $t \in \mathbb{Z}$

Put

$$(t_1^{p^2})_{tp-1} = -v_2^{-1}v_3^{(t-1)p}w.$$

Then the lemma for this case follows from [Lemma 4.10](#).

References

- [1] **H R Miller, D C Ravenel**, *Morava stabilizer algebras and the localization of Novikov's E_2 -term*, Duke Math. J. 44 (1977) 433–447 [MR](#) [Zbl](#)
- [2] **H R Miller, D C Ravenel, W S Wilson**, *Periodic phenomena in the Adams–Novikov spectral sequence*, Ann. of Math. 106 (1977) 469–516 [MR](#) [Zbl](#)
- [3] **H Nakai**, *The chromatic E_1 -term $H^0M_1^2$ for $p > 3$* , New York J. Math. 6 (2000) 21–54 [MR](#) [Zbl](#)
- [4] **S Oka**, *Multiplicative structure of finite ring spectra and stable homotopy of spheres*, from “Algebraic topology”, Lecture Notes in Math. 1051, Springer (1984) 418–441 [MR](#) [Zbl](#)
- [5] **S Oka, K Shimomura**, *On products of the β -elements in the stable homotopy of spheres*, Hiroshima Math. J. 12 (1982) 611–626 [MR](#) [Zbl](#)
- [6] **D C Ravenel**, *The structure of BP_*BP modulo an invariant prime ideal*, Topology 15 (1976) 149–153 [MR](#) [Zbl](#)
- [7] **D C Ravenel**, *The cohomology of the Morava stabilizer algebras*, Math. Z. 152 (1977) 287–297 [MR](#) [Zbl](#)
- [8] **D C Ravenel**, *Complex cobordism and stable homotopy groups of spheres*, 2nd edition, Amer. Math. Soc., Providence, RI (2004) [Zbl](#)
- [9] **K Shimomura**, *Note on the right unit map and some elements of the Brown–Peterson homology*, J. Fac. Educ. Tottori Univ. (Nat. Sci.) 38 (1989) 77–89
- [10] **K Shimomura**, *The chromatic E_1 -term $H^1M_2^1$ and its application to the homology of the Toda–Smith spectrum $V(1)$* , J. Fac. Educ. Tottori Univ. (Nat. Sci.) 39 (1990) 63–83 Correction in 41 (1992) 7–11
- [11] **K Shimomura, H Tamura**, *Nontriviality of some compositions of β -elements in the stable homotopy of the Moore spaces*, Hiroshima Math. J. 16 (1986) 121–133 [MR](#) [Zbl](#)
- [12] **H Toda**, *On spectra realizing exterior parts of the Steenrod algebra*, Topology 10 (1971) 53–65 [MR](#) [Zbl](#)

Department of Mathematics, Faculty of Science, Kochi University
Kochi, Japan

Department of Mathematics, Faculty of Science and Technology, Kochi University
Kochi, Japan

katsumi@kochi-u.ac.jp, mao.shimomura@outlook.jp

Received: 7 November 2022 Revised: 23 July 2023

ALGEBRAIC & GEOMETRIC TOPOLOGY

msp.org/agt

EDITORS

PRINCIPAL ACADEMIC EDITORS

John Etnyre
etnyre@math.gatech.edu
Georgia Institute of Technology

Kathryn Hess
kathryn.hess@epfl.ch
École Polytechnique Fédérale de Lausanne

BOARD OF EDITORS

Julie Bergner	University of Virginia jeb2md@eservices.virginia.edu	Christine Lescop	Université Joseph Fourier lescop@ujf-grenoble.fr
Steven Boyer	Université du Québec à Montréal cohf@math.rochester.edu	Robert Lipshitz	University of Oregon lipshitz@uoregon.edu
Tara E. Brendle	University of Glasgow tara.brendle@glasgow.ac.uk	Norihiko Minami	Yamato University minami.norihiko@yamato-u.ac.jp
Indira Chatterji	CNRS & Univ. Côte d'Azur (Nice) indira.chatterji@math.cnrs.fr	Andrés Navas	Universidad de Santiago de Chile andres.navas@usach.cl
Alexander Dranishnikov	University of Florida dranish@math.ufl.edu	Robert Oliver	Université Paris 13 bobol@math.univ-paris13.fr
Tobias Ekholm	Uppsala University, Sweden tobias.ekholm@math.uu.se	Jessica S Purcell	Monash University jessica.purcell@monash.edu
Mario Eudave-Muñoz	Univ. Nacional Autónoma de México mario@matem.unam.mx	Birgit Richter	Universität Hamburg birgit.richter@uni-hamburg.de
David Futer	Temple University dfuter@temple.edu	Jérôme Scherer	École Polytech. Féd. de Lausanne jerome.scherer@epfl.ch
John Greenlees	University of Warwick john.greenlees@warwick.ac.uk	Vesna Stojanoska	Univ. of Illinois at Urbana-Champaign vesna@illinois.edu
Ian Hambleton	McMaster University ian@math.mcmaster.ca	Zoltán Szabó	Princeton University szabo@math.princeton.edu
Matthew Hedden	Michigan State University mhedden@math.msu.edu	Maggy Tomova	University of Iowa maggy-tomova@uiowa.edu
Hans-Werner Henn	Université Louis Pasteur henn@math.u-strasbg.fr	Chris Wendl	Humboldt-Universität zu Berlin wendl@math.hu-berlin.de
Daniel Isaksen	Wayne State University isaksen@math.wayne.edu	Daniel T Wise	McGill University, Canada daniel.wise@mcgill.ca
Thomas Koberda	University of Virginia thomas.koberda@virginia.edu	Lior Yanovski	Hebrew University of Jerusalem lior.yanovski@gmail.com
Markus Land	LMU München markus.land@math.lmu.de		

See inside back cover or msp.org/agt for submission instructions.

The subscription price for 2024 is US \$705/year for the electronic version, and \$1040/year (+\$70, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Algebraic & Geometric Topology is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [Current Mathematical Publications](#) and the [Science Citation Index](#).

Algebraic & Geometric Topology (ISSN 1472-2747 printed, 1472-2739 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840.

AGT peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing
<https://msp.org/>

© 2024 Mathematical Sciences Publishers

ALGEBRAIC & GEOMETRIC TOPOLOGY

Volume 24

Issue 7 (pages 3571–4137)

2024

Geography of bilinearized Legendrian contact homology	3571
FRÉDÉRIC BOURGEOIS and DAMIEN GALANT	
The deformation spaces of geodesic triangulations of flat tori	3605
YANWEN LUO, TIANQI WU and XIAOPING ZHU	
Finite presentations of the mapping class groups of once-stabilized Heegaard splittings	3621
DAIKI IGUCHI	
On the structure of the top homology group of the Johnson kernel	3641
IGOR A SPIRIDONOV	
The Heisenberg double of involutory Hopf algebras and invariants of closed 3–manifolds	3669
SERBAN MATEI MIHALACHE, SAKIE SUZUKI and YUJI TERASHIMA	
A closed ball compactification of a maximal component via cores of trees	3693
GIUSEPPE MARTONE, CHARLES OUYANG and ANDREA TAMBURELLI	
An algorithmic discrete gradient field and the cohomology algebra of configuration spaces of two points on complete graphs	3719
EMILIO J GONZÁLEZ and JESÚS GONZÁLEZ	
Spectral diameter of Liouville domains	3759
PIERRE-ALEXANDRE MAILHOT	
Classifying rational G –spectra for profinite G	3801
DAVID BARNES and DANNY SUGRUE	
An explicit comparison between 2–complicial sets and Θ_2 –spaces	3827
JULIA E BERGNER, VIKTORIYA OZORNOVA and MARTINA ROVELLI	
On products of beta and gamma elements in the homotopy of the first Smith–Toda spectrum	3875
KATSUMI SHIMOMURA and MAO-NO-SUKE SHIMOMURA	
Phase transition for the existence of van Kampen 2–complexes in random groups	3897
TSUNG-HSUAN TSAI	
A qualitative description of the horoboundary of the Teichmüller metric	3919
AITOR AZEMAR	
Vector fields on noncompact manifolds	3985
TSUYOSHI KATO, DAISUKE KISHIMOTO and MITSUNOBU TSUTAYA	
Smallest nonabelian quotients of surface braid groups	3997
CINDY TAN	
Lattices, injective metrics and the $K(\pi, 1)$ conjecture	4007
THOMAS HAETTEL	
The real-oriented cohomology of infinite stunted projective spaces	4061
WILLIAM BALDERRAMA	
Fourier transforms and integer homology cobordism	4085
MIKE MILLER EISMEIER	
Profinite isomorphisms and fixed-point properties	4103
MARTIN R BRIDSON	
Slice genus bound in DTS^2 from s –invariant	4115
QIUYU REN	
Relatively geometric actions of Kähler groups on $\text{CAT}(0)$ cube complexes	4127
COREY BREGMAN, DANIEL GROVES and KEJIA ZHU	