

AG
T

*Algebraic & Geometric
Topology*

Volume 24 (2024)

Fourier transforms and integer homology cobordism

MIKE MILLER EISMEIER



Fourier transforms and integer homology cobordism

MIKE MILLER EISMEIER

We explore the *Fourier transform* of the d -invariants, which is particularly well behaved with respect to connected sum. As corollaries, we show that lens spaces are cancellable in the monoid of 3-manifolds up to integer homology cobordism, and we recover a theorem of González-Acuña and Short on Alexander polynomials of knots with reducible surgeries.

57K31, 57R90

1. Introduction	4085
2. Weighted torsors and Fourier transforms	4088
3. A theorem of González-Acuña and Short	4094
4. d -invariants of 3-manifolds	4095
5. Questions	4099
References	4100

1 Introduction

The relation of homology cobordism between 3-manifolds has a long and interesting history. Fix a ring R . Let Y and Y' be closed oriented 3-manifolds, and suppose W is a compact oriented 4-manifold whose boundary ∂W is oriented diffeomorphic to $Y' - Y$. If both maps

$$i_*: H_*(Y; R) \rightarrow H_*(W; R), \quad i'_*: H_*(Y'; R) \rightarrow H_*(W; R)$$

are isomorphisms, we say that W is an R -homology cobordism and that Y and Y' are R -homology cobordant.

This relation is most well studied when $H_*(Y; R) \cong H_*(S^3; R)$, in which case Y is called an R -homology sphere. The set of R -homology spheres modulo R -homology cobordism form a group Θ_R^3 called the “ R -homology cobordism group”. The group operation is connected sum, the neutral element is $[S^3]$, and the inverse of $[Y]$ is $[-Y]$.

If one instead considers the set of all 3–manifolds modulo R –homology cobordism, the resulting object is a *monoid*, which we denote by $\widehat{\Theta}_R^3$; the R –homology cobordism group is the submonoid of invertible elements.¹ Some things are known about this monoid; for instance, every equivalence class contains an irreducible 3–manifold [Livingston 1981] or better yet a hyperbolic 3–manifold [Myers 1983]. In another direction, there are obstructions to finding a Seifert-fibered manifold in a given equivalence class [Cochran and Tanner 2014] or more generally to finding a graph manifold whose graph is a tree in a given equivalence class [Doig and Horn 2017].²

In this note, we will investigate integer homology cobordism between a class of 3–manifolds which are not integer homology spheres. In what follows, we suppress the ring $R = \mathbb{Z}$ from notation, and write integer homology and cohomology groups as $H_*(Y)$ and $H^*(Y)$.

Theorem 1 *Suppose L and L' are connected sums of lens spaces. If L and L' are integer homology cobordant by a cobordism W , then L is oriented diffeomorphic to L' , and the induced map $W_*: H_1(L) \rightarrow H_1(L')$ respects the natural direct sum decompositions.³*

Further, if Y is any closed, oriented 3–manifold and $L \# Y$ is integer homology cobordant to $L' \# Y$, then L is oriented diffeomorphic to L' .

The first part of the result, that the oriented diffeomorphism type of L and L' is determined by their integer homology cobordism type, is not new. It follows from the more general results of [Greene 2013] on alternating links, and indeed Greene’s results imply that double-branched covers of alternating links are determined by their homology cobordism type. Independent proofs of the more restrictive claim that the oriented diffeomorphism type of a lens space is determined by its d –invariants have also appeared [Doig and Wehrli 2015; Némethi 2005]. Even before then, the integer homology cobordism classification of lens spaces of odd order goes back to [Fintushel and Stern 1987].

Our argument is independent of [Greene 2013], depending only on the computation of Reidemeister torsion for lens spaces and its relationship to their d –invariants established in [Némethi 2005]. Theorem 1 is also stronger in two ways: first, it constraints the structure of the span $H_1(L) \leftarrow H_1(W) \rightarrow H_1(L')$ of any homology cobordism relating L and L' ; second, it establishes that connected sums of lens spaces are *cancellable* in $\widehat{\Theta}_{\mathbb{Z}}^3$.

The proof of this theorem is presented in Section 4. The key point is that — provided a certain nonvanishing property holds — one can recover (a reduced version of) the d –invariants of a connected summand from those of a connected sum.

¹If $H_1(Y; R)$ is nonzero, then Y is not invertible in $\widehat{\Theta}_R^3$; the purported inverse Y' should support an R –homology cobordism between $Y \# Y'$ and S^3 , but $|H_1(Y \# Y'; R)| \geq |H_1(Y; R)| > 1 = |H_1(S^3; R)|$.

²Though see the MR review of [Doig and Horn 2017] for some errata and [Suciu 2022, Proposition 9.2] for a simplified argument.

³This is meant in an unordered sense. Precisely, suppose $H_1(L) \cong \bigoplus_i H_1(L_i)$ and similarly for L' . Then W_* sends each $H_1(L_i)$ isomorphically onto some $H_1(L'_j)$.

To recover the d -invariants of the summands, we find it useful to pass to the *Fourier transform*. Given a function $f: A \rightarrow \mathbb{C}$ on a finite abelian group, its Fourier transform is instead a function on the *dual group* $A^\vee = \text{Hom}(A, S^1)$, defined by $\hat{f}(\phi) = \frac{1}{|A|} \sum_{a \in A} f(a) \overline{\phi(a)}$. Here A will be $H^2(Y) \cong H_1(Y)$, and we will pick a base spin^c structure to consider the d -invariants as a function on $H_1(Y)$.

The use of Fourier transforms is well established in the theory of Reidemeister torsion: analytic interpretations of the Reidemeister torsion (eg [Fried 1987; Ray and Singer 1971]) interpret the torsion as a function of oriented flat bundles, and hence take as input a representation $\phi: H_1(Y; \mathbb{Z}) \rightarrow S^1$. It is also profitable to rephrase the surgery relation in terms of Fourier transforms; Nicolaescu [2004] used this to compare Reidemeister torsion to a Seiberg–Witten invariant. See also the discussion peppered throughout [Nicolaescu 2003].

Here we use a simple property of Fourier transforms. Given two groups A and A' and functions $f: A \rightarrow \mathbb{C}$ and $f': A' \rightarrow \mathbb{C}$, the *direct sum* $(f \oplus f')(a, a') = f(a) + f'(a')$ on $A \times A'$ has an especially simple Fourier transform; one can effectively read off the values of $\hat{f}(\phi)$ and $\hat{f}'(\psi)$ from the knowledge of $\widehat{f \oplus f'}$ whenever ϕ or ψ are nontrivial homomorphisms. See Proposition 12 for a precise statement.

Applying this purely algebraic observation to d -invariants, one can recover (a reduced version of) the d -invariants of summands from those of a connected sum. Provided they satisfy a certain nonvanishing property, this recovery process is well defined up to automorphism of $H^2(Y) \times H^2(Y')$, and thus preserved by integer homology cobordisms. Because the d -invariants of lens spaces satisfy this nonvanishing property, and these reduced d -invariants — equivalent to Reidemeister torsion for L -spaces — classify lens spaces up to oriented diffeomorphism, the main theorem follows.

In fact, the proof of Theorem 1 yields a stronger claim: there exist monoid homomorphisms $c_{p,q}: \hat{\Theta}_{\mathbb{Z}}^3 \rightarrow \mathbb{N}$ with $c_{p,q}(L(p, q)) = 1$ and $c_{p,q}(L(r, s)) = 0$ unless $L(r, s)$ is oriented diffeomorphic to $L(p, q)$. The purely algebraic part of this claim is the content of Corollary 19 in Section 2, while the relevant computation for lens spaces is given in Proposition 26 in Section 4.

If one considers the *Grothendieck group* $\text{Gr}(\hat{\Theta}_{\mathbb{Z}}^3)$, the group whose elements are pairs $([Y], [Z])$ with $([Y], [Z]) = ([Y'], [Z'])$ if there is an integer homology cobordism $Y \# Z' \sim Y' \# Z$, the existence of these homomorphisms $c_{p,q}$ shows that lens spaces (up to oriented diffeomorphism) span a \mathbb{Z}^∞ summand of the Grothendieck group.

As an aside, the perspective of Fourier transforms appears useful whenever one has an invariant which is additive in the sense above, including both d -invariants and Reidemeister torsion. To demonstrate this, in Section 3 we reprove [González-Acuña and Short 1986, Theorem 2.2]: if K is a knot with reducible surgery $S_n(K) \cong Y \# Y'$ with $H_1(Y) = \mathbb{Z}/p$ and $H_1(Y') = \mathbb{Z}/q$, the Alexander polynomial of K is divisible by that of the (p, q) torus knot; $\Delta(T_{p,q}) \mid \Delta(K)$. We hope that Fourier transforms can be a useful organizing tool in other contexts as well.

Acknowledgements The author would like to thank Danny Ruberman for a useful discussion during the preparation of this note, as well as Tye Lidman for comments on an early draft and suggesting that one might reprove [González-Acuña and Short 1986, Theorem 2.2] using the Fourier transform technique. The basic algebraic observation used here came about during conversations with Aiden Sagerman on a related question about products of punctured lens spaces.

2 Weighted torsors and Fourier transforms

In this section we cover the purely algebraic aspects of the main result: the context of weighted torsors, the definition of the Fourier transform, and the process of recovering summands of the direct sum of weighted torsors using Fourier transforms.

2.1 Weighted torsors

Here we rapidly define the relevant algebraic objects. Though the initial discussion about torsors is valid for any group, the relevant groups for us will be abelian, so we use additive notation.

Definition 2 A *torsor* is a pair (A, S) , where A is a group and S carries a free and transitive (right) action by A .

A choice of element $s \in S$ gives rise to a bijection $m_s : A \cong S$, given by sending $a \mapsto s + a$; for a different choice of element $s' \in S$ with $s' = s + a_0$, the bijections differ by $(m_s^{-1}m_{s'})(a) = a_0 + a$.

This observation shows that one may think of a torsor as a group where one has forgotten which element is the identity, or as A “modulo translation”.

Definition 3 An *isomorphism of torsors* is a pair $(f, g) : (A, S) \rightarrow (A', S')$, where $f : A \rightarrow A'$ is a group isomorphism and $g : S \rightarrow S'$ is a function satisfying

$$g(s + a) = g(s) + f(a).$$

By transitivity of the group actions, g is necessarily a bijection. If one chooses basepoints $s \in S$ and $s' \in S'$, and we have $g(s) = s' + a'$, then the map $m_{s'}^{-1}g m_s : A \rightarrow A'$ is given by

$$(m_{s'}^{-1}g m_s)(a) = m_{s'}^{-1}g(s + a) = m_{s'}^{-1}(g(s) + f(a)) = m_{s'}^{-1}(s' + a' + f(a)) = a' + f(a).$$

Thinking of S as a group where we’ve forgotten the identity element (or as a sort of affine space), one should imagine g to be an affine function whose “linear part” is the homomorphism f ; indeed, one can recover f from g .

Remark 4 The group of automorphisms of (A, S) is a group sometimes called the *holomorph* of A , and can be understood as the group of affine automorphisms of A .

Definition 5 A *weighted torsor* is a torsor (A, S) equipped with a function $d : S \rightarrow \mathbb{C}$. If $S = A$, we call (A, d) a *weighted group*.

If one chooses a basepoint $s \in S$, we write $d_s: A \rightarrow \mathbb{C}$ for the function $d_s(a) = (dm_s)(a) = d(s + a)$. For a different choice of basepoint $s' = s + a_0$, we have $d_{s'}(a) = d_s(a + a_0)$.

Definition 6 An isomorphism of weighted torsors $(A, S, d) \rightarrow (A', S', d')$ is a torsor isomorphism $(f, g): (A, S) \rightarrow (A', S')$ which has $d'(g(s)) = d(s)$.

If one prefers to think entirely in terms of the group A (having chosen an arbitrary basepoint), a weighted torsor is a function $d: A \rightarrow \mathbb{C}$, with d considered equivalent to $d_b(a) = d(a + b)$ for any $b \in A$. In this perspective, an isomorphism of weighted torsors $(A, d) \cong (A', d')$ amounts to an affine isomorphism $f + a': A \rightarrow A'$ which has

$$d'(f(a) + a') = d(a).$$

2.2 Fourier transforms and weighted duals

Given an abelian group A , its Pontryagin dual is the group $A^\vee = \text{Hom}(A, S^1)$.

Convention For the rest of this note, abelian groups A and torsors (A, S) are assumed to be finite. This is true in all cases of interest to us, and simplifies discussions of Fourier transforms.

Given a function $d: A \rightarrow \mathbb{C}$, we can take its Fourier transform $\hat{d}: A^\vee \rightarrow \mathbb{C}$, defined as

$$\hat{d}(\phi) = \frac{1}{|A|} \sum_{a \in A} d(a) \overline{\phi(a)}.$$

Remark 7 This differs from Nicolaescu's definition [2003, Section 1.6] by a scalar factor of $1/|A|$. Our definition makes some important formulas later slightly simpler.

If $d': A \rightarrow \mathbb{C}$ is defined by $d'(a) = d(a + a')$, then

$$\hat{d}'(\phi) = \frac{1}{|A|} \sum_{a \in A} d'(a) \overline{\phi(a)} = \frac{1}{|A|} \sum_{a \in A} d'(a - a') \overline{\phi(a - a')} = \frac{1}{|A|} \sum_{a \in A} d(a) \overline{\phi(a)} \phi(a') = \hat{d}(\phi) \phi(a').$$

This computation inspires the following definition, which we phrase intrinsically on the dual $B = A^\vee$; the statement below implicitly uses the isomorphism $(A^\vee)^\vee \cong A$ for finite A . The terminology follows [Nicolaescu 2003, Definition 3.22] (though notice that Nicolaescu allows for a sign ambiguity, and we do not).

Definition 8 If B is an abelian group equipped with weights \hat{d} and \hat{d}' , and there exists some $\psi \in B^\vee$ such that $\hat{d}'(b) = \hat{d}(b)\psi(b)$ for all $b \in B$, we say that d and d' are t -equivalent. We say that weighted groups (B, \hat{d}) and (B', \hat{d}') are t -isomorphic if there exists an isomorphism $f: B \rightarrow B'$ and element $\phi \in B^\vee$ such that

$$\hat{d}'(f(b)) = \hat{d}(b)\phi(b)$$

for all $b \in B$.

The discussion above shows that given a weighted torsor (A, S, d) , choosing a basepoint $s \in S$ and taking the Fourier transform of d_s gives us a weighted group (A^\vee, \hat{d}_s) , well defined up to t -equivalence. Furthermore, it is clear that isomorphic weighted torsors give rise to t -isomorphic weighted groups.

Notice that $\hat{d}(1) = \sum_{a \in A} d(a)$, and if \hat{d} is t -isomorphic to \hat{d}' , then $\hat{d}(1) = \hat{d}'(1)$. Later, this term will cause us some minor irritation, so we do what we can to remove it.

Definition 9 A weighted torsor (A, S, d) is *reduced* if $\sum_{s \in S} d(s) = 0$; equivalently, $\hat{d}(1) = 0$. Given any weighted torsor (A, S, d) , its *reduced part* is given by (A, S, d^r) , where

$$d^r(s) = d(s) - \frac{1}{|A|} \sum_{s' \in S} d(s');$$

that is, we subtract off the average value.

It is clear that d^r is reduced, and that reduction doesn't change the value of the Fourier transform at any $\phi \neq 1$. To see this, we need a small useful lemma.

Lemma 10 If A is a finite abelian group and $\phi : A \rightarrow S^1$ is a homomorphism, then

$$\sum_{a \in A} \phi(a) = \begin{cases} 0 & \text{if } \phi \neq 1, \\ |A| & \text{if } \phi = 1. \end{cases}$$

Proof This is a special case of the orthogonality relations for irreducible characters [Serre 1977, Theorem 2.3.3]. The proof is included for completeness.

If ϕ is trivial this is obvious. For ϕ nontrivial, write ζ for a generator of $\phi(A)$ so that ζ is a primitive m^{th} root of unity for some $m > 1$. We have

$$\sum_{a \in A} \phi(a) = \frac{|A|}{m} \sum_{k=0}^{m-1} \zeta^k.$$

But $\sum_{k=0}^{m-1} \zeta^k = (1 - \zeta^m)/(1 - \zeta) = 0$ for $\zeta \neq 1$ a nontrivial m^{th} root of unity. □

It follows that if two weighted torsors have $d'(a) = d(a) + c$ for all $a \in A$ and some constant c , then

$$\hat{d}'(\phi) = \frac{1}{|A|} \sum_{a \in A} d(a)\overline{\phi(a)} + \frac{1}{|A|} \sum_{a \in A} c\overline{\phi(a)} = \begin{cases} \hat{d}(1) + c & \text{if } \phi = 1, \\ \hat{d}(\phi) & \text{if } \phi \neq 1. \end{cases}$$

Corollary 11 If (A, S, d) is a weighted torsor, its reduced part (A, S, d^r) satisfies

$$\hat{d}^r(\phi) = \begin{cases} 0 & \text{if } \phi = 1, \\ \hat{d}(\phi) & \text{if } \phi \neq 1. \end{cases}$$

2.3 Direct sums of weighted torsors

Given two weighted torsors (A, S, d) and (A', S', d') we say their *direct sum* is the weighted torsor $(A \times A', S \times S', d \oplus d')$, where

$$(d \oplus d')(s, s') = d(s) + d'(s').$$

Notice that the $(A \times A')^\vee$ is naturally isomorphic to $A^\vee \times (A')^\vee$; if $\phi: A \rightarrow S^1$ and $\psi: A' \rightarrow S^1$ are homomorphisms, these give rise to the homomorphism $\phi\psi: A \times A' \rightarrow S^1$ by pointwise multiplication, $(\phi\psi)(a, a') = \phi(a)\psi(a')$.

The observation which motivated the present note is the following calculation of the Fourier transform of a direct sum of weighted torsors.

Proposition 12 *If $d \oplus d': A \times A' \rightarrow \mathbb{Q}$ is the direct sum of two weighted groups, the Fourier transform satisfies*

$$\widehat{(d \oplus d')}(\phi\psi) = \begin{cases} \hat{d}(\phi) & \text{if } \phi \neq 1 \text{ and } \psi = 1, \\ \hat{d}'(\psi) & \text{if } \phi = 1 \text{ and } \psi \neq 1, \\ \hat{d}(1) + \hat{d}'(1) & \text{if } \phi = \psi = 1, \\ 0 & \text{otherwise.} \end{cases}$$

Proof We have

$$\begin{aligned} \widehat{(d \oplus d')}(\phi\psi) &= \frac{1}{|A||A'|} \sum_{a,a'} (d \oplus d')(a, a') \overline{\phi\psi(a, a')} \\ &= \frac{1}{|A||A'|} \sum_{a,a'} (d(a) + d'(a')) \overline{\phi(a)\psi(a')} \\ &= \frac{1}{|A||A'|} \left(\sum_{a,a'} d(a) \overline{\phi(a)\psi(a')} \right) + \frac{1}{|A||A'|} \left(\sum_{a,a'} d'(a') \overline{\psi(a')\phi(a)} \right) \\ &= \left(\frac{1}{|A|} \sum_a d(a) \overline{\phi(a)} \right) \left(\frac{1}{|A'|} \sum_{a'} \overline{\psi(a')} \right) + \left(\frac{1}{|A'|} \sum_{a'} d'(a') \overline{\psi(a')} \right) \left(\frac{1}{|A|} \sum_a \overline{\phi(a)} \right). \end{aligned}$$

By Lemma 10, the first term vanishes when $\psi \neq 1$ and is $|A'| \hat{d}(\phi)$ when $\psi = 1$, while the second term vanishes when $\phi \neq 1$ and is $\hat{d}'(\psi)$ when $\phi = 1$. This gives the stated claim in all cases except $\phi, \psi = 1$; in that case, it gives $\hat{d}(1) + \hat{d}'(1)$. □

In particular, for *nontrivial* ϕ and ψ , one can read off the values of $\hat{d}(\phi)$ and $\hat{d}'(\psi)$ from the Fourier transform of the direct sum $\widehat{d \oplus d'}$. In the nonreduced case, the fact that $\hat{d}(1)$ and $\hat{d}'(1)$ are combined in $\widehat{(d \oplus d')}(1)$ means that we cannot recover this information from the Fourier transform of the direct sum. This is why we restrict attention to reduced weighted torsors.

2.4 Nonvanishing properties and recovering invariants of summands

We will now be more precise about the process of recovering the summands of a direct sum of weighted torsors in a way which is well defined up to isomorphism. To do so, we must make some further assumptions.

Definition 13 A weighted group (B, \hat{d}) has the *nonvanishing property* if $\hat{d}(b) \neq 0$ for all *nontrivial* elements $b \in B$.

Notice that this property is well defined up to t -isomorphism, because if (B, \hat{d}) and (B', \hat{d}') are t -isomorphic, we have an isomorphism $f: B \rightarrow B'$, an element $\phi \in B^\vee$, and an equality $\hat{d}'(f(b)) = \hat{d}(b)\phi(b)$. Because b is nontrivial if and only if $f(b)$ is, and $\phi(b) \in S^1$ is nonzero, \hat{d}' has the nonvanishing property if and only if \hat{d} does.

Definition 14 If (B, \hat{d}) is a weighted group, a *special subgroup* is a *nontrivial* subgroup $C \subset B$ such that $\hat{d}(c) \neq 0$ for all nontrivial $c \in C$. A *maximal special subgroup* is a special subgroup which is maximal among special subgroups.

Notice that special subgroups are well defined up to t -equivalence of weights, and that t -isomorphism preserves maximal special subgroups: if $f: (B, \hat{d}) \rightarrow (B', \hat{d}')$ is a t -isomorphism and $C \subset B$ is a special subgroup, then $f(C)$ is too, and vice versa. Further, a t -isomorphism maps $(C, \hat{d}|_C)$ t -isomorphically onto $(f(C), \hat{d}'|_{f(C)})$. In particular, the maximal special subgroups (considered as weighted groups up to t -isomorphism) are t -isomorphism invariants of (B, \hat{d}) .

It immediately follows from this that isomorphisms between direct sums of weighted torsors with the nonvanishing property are rather constrained.

Corollary 15 Suppose $\{(A_i, d_i)\}_{i=1}^n$ is a collection of weighted groups whose Fourier transforms satisfy the nonvanishing property, and similarly with $\{(A'_j, d'_j)\}_{j=1}^m$. Write $(A, d) = \bigoplus_{i=1}^n (A_i, d_i)$ and similarly for (A', d') . If there is an affine isomorphism of weighted groups $\varphi + a': (A, d) \cong (A', d')$, then $n = m$ and the map φ preserves the direct sum decompositions, in the sense that for all i we have $\varphi(A_i) = A'_j$ for some j .

Proof By Proposition 12, the maximal special subgroups of $A^\vee = A_1^\vee \oplus \cdots \oplus A_n^\vee$ are precisely the coordinate axes A_i^\vee (where all coordinates but the i^{th} are nonzero). Comparing the number of maximal special subgroups, we see that $m = n$. Because φ^\vee maps maximal special subgroups bijectively to maximal special subgroups, for some permutation σ we have $\varphi^\vee((A'_{\sigma(i)})^\vee) = A_i^\vee$ for all i . That is, if $\psi': A' \rightarrow S^1$ is any homomorphism, then ψ' factors through $\pi'_{\sigma(i)}: A' \rightarrow A'_{\sigma(i)}$ if and only if there exists some $\psi: A_i \rightarrow S^1$ with

$$\psi \pi_i = \psi' \pi'_{\sigma(i)} \varphi.$$

It will follow that $\varphi(A_i) \subset A'_{\sigma(i)}$, and then equality follows because these have the same cardinality (their duals do) and φ is injective. To see this first claim, pick $x_i \in A_i$, and consider $y_j = \pi_j \varphi(x_i)$. If y_j is nonzero, there is some homomorphism $\psi'_j: A'_j \rightarrow S^1$ with $\psi'_j(y_j) \neq 0$. By the discussion above,

$$\psi'_j(y_j) = \psi'_j \pi'_j \varphi(x_i) = \psi \pi_{\sigma^{-1}(j)}(x_i).$$

This can only be nonzero if $\sigma^{-1}(j) = i$ by assumption, so $\varphi(x_i)$ indeed lies in $A'_{\sigma(i)}$. \square

We will prove the main theorem similarly, by counting maximal special subgroups (considered as weighted groups up to t -isomorphism); we introduce notation for this special concept.

Definition 16 Given a weighted group (B, \hat{d}) , we associate the multiset

$$MS(B, \hat{d}) = \{[C, \hat{d}|_C] \mid C \subset B \text{ is a maximal special subgroup}\}$$

of special subgroups of B equipped with the restriction of \hat{d} , considered up to t -isomorphism.

Recall here that a multiset M is a set (by an abuse of notation written with the same name M) where each element $x \in M$ is equipped with a weight $w_M(x) \geq 1$ labeling how many times it occurs in the multiset.

Notice that if (B, \hat{d}) is t -isomorphic to (B', \hat{d}') , then the multisets $MS(B, \hat{d})$ and $MS(B', \hat{d}')$ are isomorphic (there is a weight-preserving bijection between them). If (A, S, d) is a weighted torsor, the multiset $MS(A^\vee, \hat{d}_s)$ is an invariant of (A, S, d) ; isomorphic weighted torsors give rise to isomorphic multisets. We write $MS(A, S, d)$ for $MS(A^\vee, \hat{d}_s)$ for some choice of $s \in S$.

If M and N are multisets, we write $M \cup N$ for the multiset whose underlying set is the union of the underlying sets of M and N , and whose weight is $w_{M \cup N}(x) = w_M(x) + w_N(x)$. (Here we write $w_N(x) = 0$ if x does not lie in N , and similarly with M .)

The crucial observation, almost immediate from Proposition 12, is that this multiset is additive.

Proposition 17 If (A, S, d) and (A', S', d') are **reduced** weighted torsors, then

$$MS(A \times A', S \times S', d \oplus d') = MS(A, S, d) \cup MS(A', S', d').$$

Proof For convenience, we write $d^\oplus = (d \oplus d')$, and \hat{d}^\oplus for its Fourier transform.

As mentioned above, Proposition 12 implies that the maximal special subgroups of $(A \times A')^\vee \cong A^\vee \times (A')^\vee$ are precisely the maximal special subgroups of $A^\vee \times \{1\}$ and $\{1\} \times (A')^\vee$.

Given a maximal special subgroup $C \subset A^\vee$ (or similarly $C' \subset (A')^\vee$), what remains is to compare the restriction of \hat{d} to C with the restriction of \hat{d}^\oplus to $C \times \{1\}$, but

$$\hat{d}^\oplus|_{C \times \{1\}} = \hat{d}|_C$$

by the formula from Proposition 12 and the assumption that d and d' are *reduced* weighted torsors. \square

We can now define monoid homomorphisms from the appropriate monoid to \mathbb{N} .

Definition 18 We write $\hat{\Theta}_{\text{WT}}$ for the monoid whose elements are weighted torsors up to isomorphism, and whose product operation is direct sum.

There is a corresponding monoid $\hat{\Theta}_{\text{RWT}}$ of reduced weighted torsors, and the map $d \mapsto (d^r, \text{avg } d)$ defines a monoid isomorphism $\hat{\Theta}_{\text{WT}} \cong \hat{\Theta}_{\text{RWT}} \times \mathbb{C}$.

Write RWTN for the set of reduced weighted torsors whose Fourier transforms satisfy the nonvanishing property, considered up to isomorphism; because these are considered up to isomorphism, we may think of these as weighted groups up to affine isomorphism and drop the torsor S from notation.

For each $[A, d] \in \text{RWTN}$, we define a map $c_{A,d}: \widehat{\Theta}_{\text{WT}} \rightarrow \mathbb{N}$ by

$$c_{[A,d]}(B, T, f) = \# \text{ of occurrences of } [A^\vee, \hat{d}] \text{ in } MS(B, T, f^r).$$

That is, $c_{[A,d]}(B, T, f)$ is the weight $w_{MS(B,T,f^r)}([A^\vee, \hat{d}])$.

Corollary 19 *The functions $c_{[A,d]}$ are monoid homomorphisms. If (A', d') is another reduced weighted torsor whose Fourier transform has the nonvanishing property, then*

$$c_{[A,d]}(A', d') = \begin{cases} 1 & \text{if } (A, d) \text{ is isomorphic to } (A', d'), \\ 0 & \text{otherwise.} \end{cases}$$

Proof A maximal special subgroup is defined to be nontrivial, so for the trivial weighted torsor with underlying group 1 and zero weighting, $MS(1, 0) = \emptyset$; so $c_{[A,d]}(1, 0) = 0$ and thus c sends neutral element to neutral element. Additivity follows immediately from Proposition 17 and the fact that taking the reduced part $d \mapsto d'$ commutes with direct sums. So $c_{[A,d]}$ is a monoid homomorphism.

Because the Fourier transform of (A', d') has the nonvanishing property, $MS(A, d) = \{[(A')^\vee, \hat{d}']\}$. If $[A^\vee, \hat{d}]$ appears in this singleton set, then in fact $((A')^\vee, \hat{d}')$ is t -isomorphic to (A^\vee, \hat{d}) , and hence $(A, d) \cong (A', d')$. \square

It follows that the functions c assemble into a surjective monoid homomorphism $c: \widehat{\Theta}_{\text{WT}} \rightarrow \mathbb{N}^{\text{RWTN}}$, which behaves particularly well on reduced weighted torsors whose Fourier transforms have the nonvanishing property: there is a map $\mathbb{N}^{\text{RWTN}} \rightarrow \widehat{\Theta}_{\text{WTN}}$ whose composition with c is the identity.

3 A theorem of González-Acuña and Short

Before moving on to the main theorem, we use this opportunity to give an alternative proof of [González-Acuña and Short 1986, Theorem 2.2], suggested to the author by Tye Lidman.

For context, if Σ is a homology sphere and $K = C_{p,q}(K')$ is the cable of another knot $K' \subset \Sigma$, then the pq -surgery satisfies $\Sigma_{pq}(K) \cong \Sigma_{p/q}(K') \# L(q, p)$. When $\Sigma = S^3$, the cabling conjecture [González-Acuña and Short 1986, Conjecture A] predicts that this construction gives the *only examples* of knots with reducible surgery. Among their evidence was the following theorem.

Theorem 20 *Let $K \subset \Sigma$ be a knot in an integer homology sphere with reducible surgery*

$$\Sigma_{n/m}(K) \cong Y_1 \# Y_2,$$

where $|H_1(Y_1)| = p > 1$ and $|H_1(Y_2)| = q > 1$. Then the polynomial

$$\Delta_{p,q} = \frac{(t^{pq} - 1)(t - 1)}{(t^p - 1)(t^q - 1)}$$

divides the Alexander polynomial Δ_K .

Note that the Alexander polynomial of a cable knot $K = C_{p,q}(K')$ satisfies $\Delta_K = \Delta_{p,q} \Delta_{K'}$.

Proof The proof makes use of the Reidemeister torsion of a 3–manifold, and we quickly recall some properties from [Nicolaescu 2003, Section 3.7]. When Y is a rational homology sphere, its Reidemeister torsion $T_Y: H_1(Y) \rightarrow \mathbb{Q}$ makes $H_1(Y)$ into a weighted torsor, well defined up to isomorphism and multiplication by ± 1 . When $H_1(Y) \cong \mathbb{Z}$, by contrast, $T_Y(t)$ should be understood as a rational function $T_Y \in \mathbb{Q}(t)$, well defined up to multiplication by $\pm t^k$. The Fourier transform $\widehat{T}_Y: \mathbb{C} \rightarrow \mathbb{C}$ is a meromorphic function defined by evaluating T_Y on a given complex number, and is well defined up to a variation on t –equivalence, $\widehat{T}'(z) \sim \pm z^k \widehat{T}(z)$.

In the case of knot complements, the Reidemeister torsion is related to the Alexander polynomial by the formula

$$T_{\Sigma \setminus K}(t) = \pm \frac{t^k \Delta_K(t)}{1-t};$$

this first appeared as [Milnor 1962, Theorem 4].

We will use the surgery formula for the Fourier-transformed Reidemeister torsion as stated in [Nicolaescu 2003, Theorem 3.23]: if ζ is a primitive n^{th} root of unity, then

$$\widehat{T}_{\Sigma_{n/m}(K)}(\zeta) = \frac{\widehat{T}_{\Sigma \setminus K}(\zeta)}{(1-\zeta)^{-1}} = \pm \frac{\zeta^k \Delta_K(\zeta)}{(1-\zeta^{-1})(1-\zeta)}.$$

In particular, the zeroes of $\widehat{T}_{\Sigma_{n/m}(K)}$ are identified with the n^{th} roots of unity ζ for which $\Delta_K(\zeta) = 0$.

Here we use the canonical isomorphism $H_1(\Sigma_{n/m}(K)) \cong \mathbb{Z}/n$, sending a meridian of the knot in $\Sigma \setminus K$ to 1 to identify H_1^\vee with the group of n^{th} roots of unity.

Suppose K is a knot as in the statement of the theorem. The isomorphism $\mathbb{Z}/n \cong \mathbb{Z}/p \times \mathbb{Z}/q$ induced by the connected sum decomposition sends the elements (i, j) with i and j nontrivial to n^{th} roots of unity which are neither p^{th} nor q^{th} roots of unity. For rational homology spheres Y_1 and Y_2 we have $T_{Y_1 \# Y_2}(i, j) = T_{Y_1}(i) + T_{Y_2}(j)$ [Turaev 2002, Theorem XII.1.2]. It follows from Proposition 12 that $\widehat{T}_{Y_1 \# Y_2}(\zeta) = 0$ for any n^{th} root of unity ζ which is neither a p^{th} nor q^{th} root of unity. Thus $\Delta_K(\zeta) = 0$ for all such roots of unity, so Δ_K is divisible by

$$\prod_{\substack{\zeta = e^{2\pi i k/n} \\ 0 \leq k < n \\ p \nmid k \text{ and } q \nmid k}} (t - \zeta) = \frac{(t^{pq} - 1)(t - 1)}{(t^p - 1)(t^q - 1)} = \Delta_{p,q}$$

as claimed. □

4 d –invariants of 3–manifolds

If Y is a 3–manifold, there is a naturally associated torsor $(H^2(Y)_{\text{tors}}; \text{Spin}^c(Y)_{\text{tors}})$, where the latter is the set of spin^c structures with torsion first Chern class. When Y is a rational homology sphere, every spin^c structure is torsion.

When we refer to a homology cobordism, we mean a pair (W, φ) of a compact oriented connected 4–manifold and a *chosen* orientation-preserving diffeomorphism $\varphi: \partial W \cong Y - Y'$ such that the corresponding maps $Y \rightarrow W$ and $Y' \rightarrow W$ induce isomorphisms on all integer cohomology groups.

Given a homology cobordism $W: Y \rightarrow Y'$ there is an induced isomorphism of torsors

$$(W_*, W_*^c): (H^2(Y)_{\text{tors}}, \text{Spin}_{\text{tors}}^c(Y)) \rightarrow (H^2(Y')_{\text{tors}}, \text{Spin}_{\text{tors}}^c(Y')).$$

We will make use of three weighted torsors; the first and third are associated to rational homology spheres, while the second is associated to an arbitrary 3–manifold:

- the *d*–invariant $d_Y: \text{Spin}^c(Y) \rightarrow \mathbb{Q}$ [Ozsváth and Szabó 2003, Definition 4.1];
- the *twisted d*–invariant $\underline{d}_Y: \text{Spin}_{\text{tors}}^c(Y) \rightarrow \mathbb{Q}$ [Behrens and Golla 2018, Definition 3.1];
- the *Turaev–Reidemeister torsion* $T_Y: \text{Spin}^c(Y) \rightarrow \mathbb{Q}$ [Turaev 2002, Chapter X].

Remark 21 The Turaev–Reidemeister torsion is often written as an $H^2(Y)$ –equivariant map

$$\text{Spin}^c(Y) \rightarrow \mathbb{Q}[H^2(Y)],$$

eg [Turaev 2002, Chapter I.4.1]. This gives rise to the function T_Y above by extracting the coefficient of $0 \in H^2(Y)$. This can be extended to an arbitrary 3–manifold, but the discussion is somewhat more intricate when $H^2(Y)$ is infinite: instead, the torsion defines an $H^2(Y)$ –equivariant map to the fraction field $\mathbb{Q}(H^2(Y))$.

The twisted *d*–invariant is only used for a technical reason, to allow connected sums with arbitrary 3–manifolds instead of merely rational homology spheres. When Y is a rational homology sphere, we have the tautological equality $\underline{d}_Y(\mathfrak{s}) = d_Y(\mathfrak{s})$. The Turaev–Reidemeister torsion — and the relation to *d*–invariants — will be used exclusively for calculation.

First, we establish the relationship to the work from Section 2.

Lemma 22 *The assignment $Y \mapsto (H^2(Y)_{\text{tors}}, \text{Spin}^c(Y)_{\text{tors}}, \underline{d}_Y)$ defines a monoid homomorphism $\hat{\Theta}_{\mathbb{Z}} \rightarrow \hat{\Theta}_{\text{WT}}$.*

Proof This amounts to three claims: that $\underline{d}_Y(S^3) = 0$ (tautological), that the assignment $Y \mapsto \underline{d}_Y$ sends integer homology cobordisms to isomorphisms of weighted torsors (an immediate corollary of [Behrens and Golla 2018, Corollary 4.2]), and that \underline{d}_Y is additive, in the sense that

$$\underline{d}_{Y\#Y'}(\mathfrak{s} \# \mathfrak{s}') = \underline{d}_Y(\mathfrak{s}) + \underline{d}_{Y'}(\mathfrak{s}'),$$

which is [Behrens and Golla 2018, Proposition 3.7]. □

From this, we can immediately show that *provided the \hat{d} –invariants of the summands satisfy the nonvanishing property*, integer homology cobordisms between connected sums preserve the natural direct sum decomposition of their homology groups.

Corollary 23 Suppose $Y = \#_{i=1}^n Y_i$ and $Y' = \#_{j=1}^m Y'_j$ are connected sums of 3-manifolds such that the \hat{d} -invariants of each Y_i and Y'_j satisfy the nonvanishing property. If $W : Y \rightarrow Y'$ is a homology cobordism, then $n = m$ and the induced map $W_* : H_1(Y)_{\text{tors}} \rightarrow H_1(Y')_{\text{tors}}$ preserves the natural (unordered) direct sum decompositions.

Proof As mentioned above, if $W : Y \rightarrow Y'$ is a homology cobordism, it induces an isomorphism of weighted torsors $(\text{Spin}_{\text{tors}}^c(Y), \underline{d}) \cong (\text{Spin}_{\text{tors}}^c(Y'), \underline{d}')$. The statement follows immediately from Corollary 15. □

The following corollary is simply an application of Corollary 19, applied to these particular weighted torsors.

Corollary 24 Suppose Y_i is a collection of 3-manifolds indexed by some set S such that

- the groups $H^2(Y_i)$ are nontrivial;
- the Fourier transforms \hat{d}_{Y_i} satisfy the nonvanishing property;
- the weighted torsors $(H^2(Y_i), \text{Spin}_{\text{tors}}^c(Y_i), \underline{d}')$ are pairwise nonisomorphic.

Then there is a homomorphism $c : \hat{\Theta}_{\mathbb{Z}} \rightarrow \mathbb{N}^S$ with $c_i(Y_i) = 1$ and $c_i(Y_j) = 0$ for $i \neq j$. In particular, the Y_i are linearly independent in $\hat{\Theta}_{\mathbb{Z}}$ and span a \mathbb{Z}^S -summand of the Grothendieck group $\text{Gr}(\hat{\Theta}_{\mathbb{Z}})$.

Here \underline{d}' is the reduced part of \underline{d} as in Definition 9.

To prove Theorem 1, we need to show that the lens spaces $L(p, q)$ —considered up to oriented diffeomorphism—satisfy the assumptions of the corollary. This is classical; the crucial observation is that the reduced d -invariant recovers the Turaev–Reidemeister torsion.

Lemma 25 If Y is an L -space, then

$$T(\mathfrak{s}) = \frac{1}{2}(\underline{d}'(\mathfrak{s})).$$

This follows immediately from [Rustamov 2005, Theorems 5.3.3–4]. For lens spaces (and thus their connected sums, as both sides of this equality are additive) this was proven earlier in [Némethi 2005, Section 10.7], and indeed Némethi’s result is used in Rustamov’s argument.

The Turaev–Reidemeister torsion of lens spaces is classical, and the sign-refined version only slightly less so. For an appropriate choice of base spin^c structure and an appropriate isomorphism $H^2(L(p, q)) \cong \mathbb{Z}/p$, we have [Némethi and Nicolaescu 2002, Section 7.1] for any nontrivial p^{th} root of unity ζ

$$\hat{T}(\zeta) = \frac{1}{p(1 - \zeta^{-1})(1 - \zeta^{-q})}.$$

Notice that this formula has an extra factor of $1/p$ compared to Nicolaescu’s, owing to the change of convention discussed in Remark 7.

We write $d_{p,q}^r : \text{Spin}^c(L(p, q)) \rightarrow \mathbb{Q}$ for the reduced d -invariants of the lens space $L(p, q)$.

The lemma above establishes that (after choosing an appropriate base spin^c structure and isomorphism $L(p, q) \cong \mathbb{Z}/p$) we have for all nontrivial p^{th} roots of unity

$$(\hat{d}_{p,q}^r)(\zeta) = \frac{1}{2p(1-\zeta^{-1})(1-\zeta^{-q})}.$$

Proposition 26 *The Fourier transforms of the weighted torsors $d_{p,q}^r$ have the nonvanishing property. Furthermore, if $d_{p,q}^r \cong d_{r,s}^r$, then $p = r$ and $s \equiv q^{\pm 1} \pmod p$, so $L(p, q)$ and $L(r, s)$ are oriented diffeomorphic.*

Proof That these have the nonvanishing property is obvious.

What remains is the essentially classical claim that signed Turaev–Reidemeister torsion classifies lens spaces up to oriented diffeomorphism; we include a short proof for completeness. We may as well assume $p > 2$.

If $d_{p,q}^r \cong d_{r,s}^r$ then $\mathbb{Z}/p \cong \mathbb{Z}/r$, so $p = r$. If $\hat{d}_{p,q}^r$ is t -isomorphic to $\hat{d}_{p,s}^r$, such an isomorphism induces a t -isomorphism $\hat{f}_{p,q} \cong \hat{f}_{p,s}$ between the simpler functions

$$\hat{f}_{p,q}(\zeta) = (1-\zeta)(1-\zeta^q) = 1 - \zeta - \zeta^q + \zeta^{q+1}.$$

If $q = 1$ this is the Fourier transform of the function $f_{p,1} = (1, -2, 1, 0, \dots, 0)$, where we list off the values $f_{p,q}(i)$ in order starting at 0. If $q = p-1$ we have instead $f_{p,p-1} = (2, -1, 0, \dots, 0, -1)$. For $1 < q < p-1$, we have $f_{p,q}(0) = f_{p,q}(q+1) = 1$, while $f_{p,q}(1) = f_{p,q}(q) = -1$, and all other values are zero.

It is transparent that there is no affine isomorphism of \mathbb{Z}/p taking $f_{p,1}$ or $f_{p,p-1}$ to any of the other functions above, as the values are different; this reduces us to the case $1 < q < p-1$.

Now suppose there exists some integer k prime to p and some integer ℓ such that

$$(1) \quad f_{p,s}(ki + \ell) = f_{p,q}(i)$$

for all i . Since $f_{p,q}(0) = 1$, we have $f_{p,s}(\ell) = 0$ and thus either $\ell \equiv 0$ or $\ell \equiv s+1$. Because $i \mapsto ki + \ell$ is a bijection, in the former case we must have $k(q+1) \equiv s+1$ and in the latter case $k(q+1) + s+1 \equiv 0$. We handle these two cases separately.

- (i) If $\ell \equiv 0$, then applying (1) to $i \equiv 1$, we have either $k \equiv 1$ (which gives $q+1 \equiv s+1$ and hence $q \equiv s$) or $k \equiv s$ (which gives $qs + s \equiv s+1$ so $qs \equiv 1$).
- (ii) If $\ell \equiv s+1$, then applying (1) to $i \equiv 1$, we have either $k+s+1 \equiv 1$ (in which case $k \equiv -s$ so $-sq - s + s + 1 \equiv 0$ and thus $qs \equiv 1$) or $k+s+1 \equiv s$ (in which case $k \equiv -1$ so $-q-1 + s + 1 \equiv 0$ and $q \equiv s$).

In any of the four possibilities for the values of k and ℓ modulo p , we see that the desired claim holds. \square

The main theorem follows immediately from this proposition, as well as Corollaries 23 and 24.

5 Questions

We close with a handful of questions inspired by the results above.

Question 1 Which collections of 3–manifolds satisfy the hypotheses of Corollary 24? Does this class include spherical 3–manifolds, or double-branched covers of alternating links?

It would follow that manifolds in this class are integer homology cobordant if and only if they are diffeomorphic. For spherical 3–manifolds, an argument might proceed by an explicit computation of their d –invariants (or of their Reidemeister torsions); for alternating links an argument might follow through the lattice-theoretic techniques of [Greene 2013].

In another direction, recall that [Lisca 2007] provides a complete classification of connected sums of lens spaces up to rational homology cobordism. It would be interesting to determine the intermediate case of R –homology cobordism for, say, $R = \mathbb{Z}/p$.

Question 2 Determine the R –homology cobordism classification of connected sums of lens spaces for various rings R .

The author has made no attempt to investigate this. It would similarly be interesting to follow up on the classification of rational homology ribbon cobordisms between connected sum of lens spaces given in [Huber 2021, Theorem 1.3] and classify the R –homology ribbon cobordisms for various rings R .

It would be interesting to understand better the interaction between the integer homology *group* and the larger integer homology *monoid*. In the integer homology *group*, because all elements are invertible, all elements are also cancellative: if $[Y] + [Z] = [Z]$ we have $[Y] = 0$. This is not true in an arbitrary monoid, so one might ask if integer homology spheres *remain* cancellative when we pass to the integer homology monoid.

This question can be phrased in terms of the Grothendieck group $\text{Gr}(\widehat{\Theta}_{\mathbb{Z}})$ as follows.

Question 3 Does the map $\Theta_{\mathbb{Z}} \rightarrow \text{Gr}(\widehat{\Theta}_{\mathbb{Z}})$ have nontrivial kernel? That is, can one find an integer homology 3–sphere Y and a closed oriented 3–manifold Z so that Y is not integer homology cobordant to S^3 , but $Y \# Z$ is integer homology cobordant to Z ?

One might imagine that the behavior of integer homology spheres under integer homology cobordism is somehow orthogonal to the behavior of rational homology spheres; very optimistically, one might believe that $j : \Theta_{\mathbb{Z}} \rightarrow \widehat{\Theta}_{\mathbb{Z}}$ *splits*, meaning that there is a homomorphism $\mu : \widehat{\Theta}_{\mathbb{Z}} \rightarrow \Theta_{\mathbb{Z}}$ with $\mu j = 1$, and one might then try to understand the structure of the Grothendieck group in terms of $\ker(\mu)$ and $\Theta_{\mathbb{Z}}$. If such a splitting exists at the level of monoids, we also have such a splitting at the level of groups when we pass to the Grothendieck group.

One way to guarantee this is *impossible*, while also giving an element in the kernel discussed above, is to find indivisible elements in $\Theta_{\mathbb{Z}}$ which become divisible in $\text{Gr}(\widehat{\Theta}_{\mathbb{Z}})$. We pose the existence of such homology spheres as a question.

Question 4 Is there an integer homology sphere Y with the following properties?

- (i) Y is indivisible in the integer homology cobordism group: if $n > 1$ and Y' is another integer homology sphere, there is no integer homology cobordism between Y and $\#^n Y'$.
- (ii) There *does* exist an $n > 1$, an integer homology sphere Y' , and a 3-manifold Z such that $Y \# Z$ is integer homology cobordant to $(\#^n Y') \# Z$.

Note that if this held, then $-Y \# nY'$ would give an infinite-order element in the kernel of $\Theta_{\mathbb{Z}} \rightarrow \text{Gr}(\widehat{\Theta}_{\mathbb{Z}})$, answering Question 3 in the positive.

References

- [Behrens and Golla 2018] **S Behrens, M Golla**, *Heegaard Floer correction terms, with a twist*, *Quantum Topol.* 9 (2018) 1–37 MR Zbl
- [Cochran and Tanner 2014] **T D Cochran, D Tanner**, *Homology cobordism and Seifert fibered 3-manifolds*, *Proc. Amer. Math. Soc.* 142 (2014) 4015–4024 MR Zbl
- [Doig and Horn 2017] **M I Doig, P D Horn**, *On the intersection ring of graph manifolds*, *Trans. Amer. Math. Soc.* 369 (2017) 1185–1203 MR Zbl
- [Doig and Wehrli 2015] **M Doig, S Wehrli**, *A combinatorial proof of the homology cobordism classification of lens spaces*, preprint (2015) arXiv 1505.06970
- [Fintushel and Stern 1987] **R Fintushel, R Stern**, *Rational homology cobordisms of spherical space forms*, *Topology* 26 (1987) 385–393 MR Zbl
- [Fried 1987] **D Fried**, *Lefschetz formulas for flows*, from “The Lefschetz centennial conference, III” (A Verjovsky, editor), *Contemp. Math.* 58, Amer. Math. Soc., Providence, RI (1987) 19–69 MR Zbl
- [González-Acuña and Short 1986] **F González-Acuña, H Short**, *Knot surgery and primeness*, *Math. Proc. Cambridge Philos. Soc.* 99 (1986) 89–102 MR Zbl
- [Greene 2013] **J E Greene**, *Lattices, graphs, and Conway mutation*, *Invent. Math.* 192 (2013) 717–750 MR Zbl
- [Huber 2021] **M Huber**, *Ribbon cobordisms between lens spaces*, *Pacific J. Math.* 315 (2021) 111–128 MR Zbl
- [Lisca 2007] **P Lisca**, *Sums of lens spaces bounding rational balls*, *Algebr. Geom. Topol.* 7 (2007) 2141–2164 MR Zbl
- [Livingston 1981] **C Livingston**, *Homology cobordisms of 3-manifolds, knot concordances, and prime knots*, *Pacific J. Math.* 94 (1981) 193–206 MR Zbl
- [Milnor 1962] **J Milnor**, *A duality theorem for Reidemeister torsion*, *Ann. of Math.* 76 (1962) 137–147 MR Zbl
- [Myers 1983] **R Myers**, *Homology cobordisms, link concordances, and hyperbolic 3-manifolds*, *Trans. Amer. Math. Soc.* 278 (1983) 271–288 MR Zbl

- [Némethi 2005] **A Némethi**, *On the Ozsváth–Szabó invariant of negative definite plumbed 3–manifolds*, *Geom. Topol.* 9 (2005) 991–1042 MR Zbl
- [Némethi and Nicolaescu 2002] **A Némethi, L I Nicolaescu**, *Seiberg–Witten invariants and surface singularities*, *Geom. Topol.* 6 (2002) 269–328 MR Zbl
- [Nicolaescu 2003] **L I Nicolaescu**, *The Reidemeister torsion of 3–manifolds*, *De Gruyter Studies in Mathematics* 30, de Gruyter, Berlin (2003) MR Zbl
- [Nicolaescu 2004] **L I Nicolaescu**, *Seiberg–Witten invariants of rational homology 3–spheres*, *Commun. Contemp. Math.* 6 (2004) 833–866 MR Zbl
- [Ozsváth and Szabó 2003] **P Ozsváth, Z Szabó**, *Absolutely graded Floer homologies and intersection forms for four-manifolds with boundary*, *Adv. Math.* 173 (2003) 179–261 MR Zbl
- [Ray and Singer 1971] **D B Ray, I M Singer**, *R–torsion and the Laplacian on Riemannian manifolds*, *Advances in Math.* 7 (1971) 145–210 MR Zbl
- [Rustamov 2005] **R Rustamov**, *On Heegaard Floer homology of three-manifolds*, PhD thesis, Princeton University (2005) MR Available at <https://www.proquest.com/docview/305385735>
- [Serre 1977] **J-P Serre**, *Linear representations of finite groups*, *Graduate Texts in Math.* 42, Springer (1977) MR Zbl
- [Suciu 2022] **A I Suciu**, *Cohomology jump loci of 3–manifolds*, *Manuscripta Math.* 167 (2022) 89–123 MR Zbl
- [Turaev 2002] **V Turaev**, *Torsions of 3–dimensional manifolds*, *Progr. Math.* 208, Birkhäuser, Basel (2002) MR Zbl

*Department of Mathematics and Statistics, University of Vermont
Burlington, VT, United States*

`mike.miller-eismeier@uvm.edu`

Received: 5 June 2023 Revised: 14 August 2023

ALGEBRAIC & GEOMETRIC TOPOLOGY

msp.org/agt

EDITORS

PRINCIPAL ACADEMIC EDITORS

John Etnyre
etnyre@math.gatech.edu
Georgia Institute of Technology

Kathryn Hess
kathryn.hess@epfl.ch
École Polytechnique Fédérale de Lausanne

BOARD OF EDITORS

Julie Bergner	University of Virginia jeb2md@eservices.virginia.edu	Christine Lescop	Université Joseph Fourier lescop@ujf-grenoble.fr
Steven Boyer	Université du Québec à Montréal cohf@math.rochester.edu	Robert Lipshitz	University of Oregon lipshitz@uoregon.edu
Tara E Brendle	University of Glasgow tara.brendle@glasgow.ac.uk	Norihiko Minami	Yamato University minami.norihiko@yamato-u.ac.jp
Indira Chatterji	CNRS & Univ. Côte d'Azur (Nice) indira.chatterji@math.cnrs.fr	Andrés Navas	Universidad de Santiago de Chile andres.navas@usach.cl
Alexander Dranishnikov	University of Florida dranish@math.ufl.edu	Robert Oliver	Université Paris 13 bobol@math.univ-paris13.fr
Tobias Ekholm	Uppsala University, Sweden tobias.ekholm@math.uu.se	Jessica S Purcell	Monash University jessica.purcell@monash.edu
Mario Eudave-Muñoz	Univ. Nacional Autónoma de México mario@matem.unam.mx	Birgit Richter	Universität Hamburg birgit.richter@uni-hamburg.de
David Futер	Temple University dfuter@temple.edu	Jérôme Scherer	École Polytech. Féd. de Lausanne jerome.scherer@epfl.ch
John Greenlees	University of Warwick john.greenlees@warwick.ac.uk	Vesna Stojanoska	Univ. of Illinois at Urbana-Champaign vesna@illinois.edu
Ian Hambleton	McMaster University ian@math.mcmaster.ca	Zoltán Szabó	Princeton University szabo@math.princeton.edu
Matthew Hedden	Michigan State University mhedden@math.msu.edu	Maggy Tomova	University of Iowa maggy-tomova@uiowa.edu
Hans-Werner Henn	Université Louis Pasteur henn@math.u-strasbg.fr	Chris Wendl	Humboldt-Universität zu Berlin wendl@math.hu-berlin.de
Daniel Isaksen	Wayne State University isaksen@math.wayne.edu	Daniel T Wise	McGill University, Canada daniel.wise@mcgill.ca
Thomas Koberda	University of Virginia thomas.koberda@virginia.edu	Lior Yanovski	Hebrew University of Jerusalem lior.yanovski@gmail.com
Markus Land	LMU München markus.land@math.lmu.de		

See inside back cover or msp.org/agt for submission instructions.

The subscription price for 2024 is US \$705/year for the electronic version, and \$1040/year (+\$70, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Algebraic & Geometric Topology is indexed by Mathematical Reviews, Zentralblatt MATH, Current Mathematical Publications and the Science Citation Index.

Algebraic & Geometric Topology (ISSN 1472-2747 printed, 1472-2739 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840.

AGT peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<https://msp.org/>

© 2024 Mathematical Sciences Publishers

ALGEBRAIC & GEOMETRIC TOPOLOGY

Volume 24 Issue 7 (pages 3571–4137) 2024

Geography of bilinearized Legendrian contact homology	3571
FRÉDÉRIC BOURGEOIS and DAMIEN GALANT	
The deformation spaces of geodesic triangulations of flat tori	3605
YANWEN LUO, TIANQI WU and XIAOPING ZHU	
Finite presentations of the mapping class groups of once-stabilized Heegaard splittings	3621
DAIKI IGUCHI	
On the structure of the top homology group of the Johnson kernel	3641
IGOR A SPIRIDONOV	
The Heisenberg double of involutory Hopf algebras and invariants of closed 3-manifolds	3669
SERBAN MATEI MIHALACHE, SAKIE SUZUKI and YUJI TERASHIMA	
A closed ball compactification of a maximal component via cores of trees	3693
GIUSEPPE MARTONE, CHARLES OUYANG and ANDREA TAMBURELLI	
An algorithmic discrete gradient field and the cohomology algebra of configuration spaces of two points on complete graphs	3719
EMILIO J GONZÁLEZ and JESÚS GONZÁLEZ	
Spectral diameter of Liouville domains	3759
PIERRE-ALEXANDRE MAILHOT	
Classifying rational G -spectra for profinite G	3801
DAVID BARNES and DANNY SUGRUE	
An explicit comparison between 2-complicial sets and Θ_2 -spaces	3827
JULIA E BERGNER, VIKTORIYA OZORNOVA and MARTINA ROVELLI	
On products of beta and gamma elements in the homotopy of the first Smith–Toda spectrum	3875
KATSUMI SHIMOMURA and MAO-NO-SUKE SHIMOMURA	
Phase transition for the existence of van Kampen 2-complexes in random groups	3897
TSUNG-HSUAN TSAI	
A qualitative description of the horoboundary of the Teichmüller metric	3919
AITOR AZEMAR	
Vector fields on noncompact manifolds	3985
TSUYOSHI KATO, DAISUKE KISHIMOTO and MITSUNOBU TSUTAYA	
Smallest nonabelian quotients of surface braid groups	3997
CINDY TAN	
Lattices, injective metrics and the $K(\pi, 1)$ conjecture	4007
THOMAS HAETTEL	
The real-oriented cohomology of infinite stunted projective spaces	4061
WILLIAM BALDERRAMA	
Fourier transforms and integer homology cobordism	4085
MIKE MILLER EISMEIER	
Profinite isomorphisms and fixed-point properties	4103
MARTIN R BRIDSON	
Slice genus bound in $DT S^2$ from s -invariant	4115
QIUYU REN	
Relatively geometric actions of Kähler groups on CAT(0) cube complexes	4127
COREY BREGMAN, DANIEL GROVES and KEJIA ZHU	