

AG
T

*Algebraic & Geometric
Topology*

Volume 24 (2024)

Cartesian fibrations of $(\infty, 2)$ -categories

ANDREA GAGNA
YONATAN HARPAZ
EDOARDO LANARI



Cartesian fibrations of $(\infty, 2)$ -categories

ANDREA GAGNA

YONATAN HARPAZ

EDOARDO LANARI

We introduce four variance flavors of (co)cartesian fibrations of ∞ -bicategories with ∞ -bicategorical fibers, in the framework of scaled simplicial sets. Given a map $p: \mathcal{E} \rightarrow \mathcal{B}$ of ∞ -bicategories, we define p -(co)cartesian arrows and inner/outer triangles by means of lifting properties against p , leading to a notion of 2-inner/outer (co)cartesian fibrations as those maps with enough (co)cartesian lifts for arrows and enough inner/outer lifts for triangles, together with a compatibility property with respect to whiskerings in the outer case. By doing so, we also recover in particular the case of ∞ -bicategories fibered in ∞ -categories studied in previous work. We also prove that equivalences of such fibrations can be tested fiberwise. As a motivating example, we show that the domain projection $d: \text{Fun}^{\text{gr}}(\Delta^1, \mathcal{C}) \rightarrow \mathcal{C}$ is a prototypical example of a 2-outer cartesian fibration, where $\text{Fun}^{\text{gr}}(X, Y)$ denotes the ∞ -bicategory of functors, lax natural transformations and modifications. We then define 2-inner and 2-outer flavors of (co)cartesian fibrations of categories enriched in ∞ -categories, and we show that a fibration $p: \mathcal{E} \rightarrow \mathcal{B}$ of such enriched categories is a (co)cartesian 2-inner/outer fibration if and only if the corresponding map $N^{\text{sc}}(p): N^{\text{sc}}\mathcal{E} \rightarrow N^{\text{sc}}\mathcal{B}$ is a fibration of this type between ∞ -bicategories.

18N65, 18N50, 55U35

Introduction	4731
1. Preliminaries	4734
2. 2-Inner/outer cartesian fibrations	4744
3. The domain projection	4758
4. Enriched cartesian fibrations	4766
References	4777

Introduction

This paper is part of an ongoing series of works on the theory of $(\infty, 2)$ -categories. We will generally refer to these as ∞ -bicategories, and identify them with scaled simplicial sets satisfying suitable extension properties. Our goal here is to set up the fundamentals of a theory of (co)cartesian fibrations $\mathcal{E} \rightarrow \mathcal{B}$ of ∞ -bicategories, whose fibers encode a family of ∞ -bicategories \mathcal{E}_b depending functorially on $b \in \mathcal{B}$.

When \mathcal{E} and \mathcal{B} are ∞ -categories the corresponding notion of (co)cartesian fibration was set up by Lurie in [12]. Generalizing classical work of Grothendieck, Lurie showed that such fibrations over a base ∞ -category \mathcal{B} are in complete correspondence with functors $\mathcal{B} \rightarrow \mathcal{Cat}_\infty$ in the cocartesian case, and contravariant such functors in the cartesian case. This Grothendieck–Lurie correspondence plays a key role in higher category theory, as it permits the handling of highly coherent pieces of structure, such as functors, in a relatively accessible manner.

When coming to consider the ∞ -bicategorical counterpart, an immediate difference presents itself: here we have not just two different variances, but four, depending on whether or not the functorial dependence on 2-morphisms is covariant or contravariant. This additional axis of symmetry is already visible when the base is an ∞ -bicategory and the fibers \mathcal{E}_b are ∞ -categories, since \mathcal{Cat}_∞ itself has a nontrivial ∞ -bicategorical structure. A thorough treatment of this case was taken up in our previous work [6], where we have used the term inner (co)cartesian fibrations to indicate those cases where the dependence on 2-morphisms matches the one on 1-morphisms, and outer (co)cartesian fibration for those where these two dependencies have opposite variance. The division into these two types is also visible on the level of mapping ∞ -categories: when both variances match (the inner case) the induced functor on mapping ∞ -bicategories is a right fibration, and when they don't (the outer case) the induced functor on mapping ∞ -categories is a left fibration. One of the main results of [6] is an ∞ -bicategorical Grothendieck–Lurie correspondence for all the four variances. The proof crucially relies on the work of Lurie [13] in the inner cocartesian case, and the straightening–unstraightening Quillen equivalence he proved in that setting.

Our goal in the present paper is to extend these notions of fibrations to the setting where both the base \mathcal{B} and the fibers \mathcal{E}_b are ∞ -bicategories. This requires defining, in addition to a notion of (co)cartesian 1-morphisms, a suitable notion of (co)cartesian 2-morphism. Working in the setting of scaled simplicial sets, such 2-morphisms are determined by triangles, though this correspondence is not perfect (the same 2-morphism can be encoded by different triangles corresponding to different factorizations of the target 1-morphism), and one needs to be a bit more careful about how to define this. To match with the notation for fibrations, we use the terms *inner* and *outer* to describe triangles which roughly correspond to cartesian and cocartesian 2-morphisms. For somewhat technical reasons one needs to separate those further into two types, which we call left and right inner/outer triangles. This distinction turns out to behave slightly differently in the inner and outer cases, an issue due to the built-in asymmetry of triangles, which encode a 2-morphism together with a factorization of its target 1-morphism, but not of its domain.

In the future continuation of the present work we plan to establish a complete Grothendieck–Lurie correspondence for such fibrations over a fixed base \mathcal{B} , showing that they encode the four possible variances of functors $\mathcal{B} \rightarrow \mathbf{BiCat}_\infty$. Though we will not arrive to this goal yet here, we do lay the foundations and prove the basic properties we expect to need, based on our current work in progress in that direction. In particular, we prove that equivalences between such fibrations are detected on fibers, and construct a certain universal example in the form of the domain projection $\mathbf{Fun}^{\mathrm{gr}}(\Delta^1, \mathcal{B}) \rightarrow \mathcal{B}$, where $\mathbf{Fun}^{\mathrm{gr}}(\Delta^1, \mathcal{B})$ is the ∞ -bicategory of arrows in \mathcal{B} and lax squares between them. We then establish

a comparison between the notions of inner/outer (co)cartesian fibrations and an analogous one in the setting of categories enriched in marked simplicial sets. Such a comparison played an important role in [6] by implying that the $(\mathbb{Z}/2 \times \mathbb{Z}/2)$ -symmetry of the theory of ∞ -bicategories switched between the four types of fibrations. This allowed one to reduce the Grothendieck–Lurie correspondence in [6] to the inner cocartesian case, and we expect a similar role to be played in a future generalization of that correspondence to fibrations.

After finishing the work on the present paper, we became aware of independent work of Abellán García and Stern [1], which investigates the outer variant of these fibrations using a model category structure on marked biscaled simplicial sets. By contrast, our approach here consistently covers all four variances, and also interacts well with the framework we set up in [6], a property we expect would be useful in a future proof of a complete ∞ -bicategorical Grothendieck–Lurie correspondence for inner/outer (co)cartesian fibrations. Finally, let us also note that a Grothendieck–Lurie type correspondence for outer cartesian fibrations is sketched by Gaitsgory and Rozenblyum in [9, Appendix], though the argument relies in various parts on unproven statements. The extensive treatment of derived algebraic geometry developed in the body of their volumes [9; 10] based on that appendix makes for a powerful motivation for obtaining the ∞ -bicategorical Grothendieck–Lurie correspondence rigorously. We also plan to pursue the study of inner/outer *locally* (co)cartesian fibrations, which encode lax/oplax 2-functors from an ∞ -bicategory \mathcal{B} to BiCat_∞ , as we defined in [7, Section 3]. This will enable one to compare that last definition with that of [9; 10], which should also allow for the comparison of the two notions of Gray products, thus establishing many key unproven statements involving Gray products made in the appendix of loc. cit. The results of the present work are fundamental preliminaries for all these applications.

Organization of the paper

We begin with a preliminary section where we give pointers to the necessary definitions and results concerning marked and scaled simplicial sets, and recall the framework of inner/outer (co)cartesian fibrations set up in [6], which we rename 1-inner/outer (co)cartesian fibration in order to better distinguish between them and the type of fibrations introduced in the present paper. Next, we introduce the main concepts of this paper, namely those of left/right inner/outer triangles and the corresponding notions of fibrations. We establish fundamental results concerning closure properties of inner/outer triangles, as well as stability properties and a fiberwise criterion to test equivalences of fibrations. Finally, we prove Theorem 3.0.7, which concerns the domain projection on an ∞ -bicategory and provides a prototypical example of a 2-outer cartesian fibration.

In Section 4 we briefly recall the theory of (co)cartesian fibrations for ordinary 2-categories, as developed by Buckley in [5], as a motivation for its enhancement to the setting of categories enriched in marked simplicial sets. We then provide an extension of the results of [6, Section 3], by showing that enriched 2-inner/outer (co)cartesian fibrations identify via the coherent nerve functor with the respective notion of fibrations between ∞ -bicategories; see Theorem 4.2.4 for the precise statement.

Vistas and applications

A theory of fibrations provides the backbone to define symmetric monoidal $(\infty, 2)$ -categories, as a suitable class of inner cocartesian fibrations over Fin_* . We expect symmetric monoidal ∞ -bicategories to play a fundamental role as their 1-dimensional counterpart, providing extra expressive power thanks to the 2-dimensional structure. For instance, they can be used to encode the relevant dualities of Ind-coherent sheaves of (derived) schemes of finite type, as in [9, Chapter 9].

Furthermore, inner/outer (locally) (co)cartesian fibrations are used as a tool in [9, Chapters 11 and 12] to establish an adjoint theorem that plays a crucial role in extending the quasicohherent sheaves functor from derived affine schemes to derived prestacks; see [9, Chapter 11, Sections 3.1.1 and 3.2.1].

Finally, the theory of fibrations can be used to define a simplicial version of the notion of (relative) $(\infty, 2)$ -operad; see Batanin [2; 3]. Possible applications in this direction come from symplectic geometry, and in particular from the functoriality of the Fukaya category; see for instance Bottman and Carmeli [4].

Acknowledgements Gagna and Lanari gratefully acknowledge the support of Praemium Academiae of M Markl and RVO:67985840. Lanari is also grateful to Rune Haugseng and Nick Rozenblyum for fruitful conversations during his stay at MSRI.

1 Preliminaries

In this section we establish notation and recall some preliminary definitions and results that will be used.

1.1 Marked simplicial sets and enriched categories

Recall that a marked simplicial set is a pair (X, E_X) where X is a simplicial set and E_X is a collection of edges in X , called the *marked* edges, containing all degenerate edges. A map of marked simplicial sets $f: (X, E_X) \rightarrow (Y, E_Y)$ is a map of simplicial sets $f: X \rightarrow Y$ satisfying $f(E_X) \subseteq E_Y$. When denoting an explicit marked simplicial set we will often omit the reference to the degenerate edges. For example, we will write $(\Delta^n, \Delta^{\{0,1\}})$ for the marked simplicial set whose underlying simplicial set is the n -simplex and whose marked edges are all the degenerate edges together with the edge $\Delta^{\{0,1\}}$. The category of marked simplicial sets will be denoted by Set_Δ^+ . It is locally presentable and cartesian closed. For more background on marked simplicial sets we refer the reader to the comprehensive treatment in [12].

We will denote by $\text{Set}_\Delta^+ \text{-Cat}$ the category of categories enriched in Set_Δ^+ with respect to the cartesian product on Set_Δ^+ , to which we will refer as *marked simplicial categories*. For a marked simplicial category \mathcal{C} and two objects $x, y \in \mathcal{C}$, we will denote by $\mathcal{C}(x, y) \in \text{Set}_\Delta^+$ the associated mapping marked simplicial set. By an arrow $e: x \rightarrow y$ in a marked simplicial category \mathcal{C} we will simply mean a vertex $e \in \mathcal{C}(x, y)_0$ in the corresponding mapping object.

We will generally consider $\text{Set}_\Delta^+ \text{-Cat}$ together with its associated *Dwyer–Kan model structure*; see [12, Section A.3.2]. In this model structure the weak equivalences are the Dwyer–Kan equivalences, that is, the maps which are essentially surjective on homotopy categories and induce marked categorical equivalences on mapping objects. The fibrant objects are the enriched categories \mathcal{C} whose mapping objects $\mathcal{C}(x, y)$ are all fibrant, that is, ∞ -categories marked by their equivalences. The model category $\text{Set}_\Delta^+ \text{-Cat}$ is then a presentation of the theory of $(\infty, 2)$ -categories, and is Quillen equivalent to other known models; see Section 1.2 below.

We say that $\mathcal{E} \in \text{Set}_\Delta^+ \text{-Cat}$ is a *Cat $_\infty$ -category* if it is fibrant in the Dwyer–Kan model structure, ie if it is enriched over ∞ -categories, with marking given by equivalences. A *fibration* of *Cat $_\infty$ -categories* is a fibration between fibrant objects in the Dwyer–Kan model structure on $\text{Set}_\Delta^+ \text{-Cat}$.

1.2 Scaled simplicial sets and ∞ -bicategories

Definition 1.2.1 [13] A *scaled simplicial set* is a pair (X, T_X) where X is a simplicial set and T_X is a subset of the set of triangles of X , called *thin triangles*, containing the degenerate ones. A map of scaled simplicial sets $f : (X, T_X) \rightarrow (Y, T_Y)$ is a map of simplicial sets $f : X \rightarrow Y$ satisfying $f(T_X) \subseteq T_Y$.

We will denote by $\text{Set}_\Delta^{\text{sc}}$ the category of scaled simplicial sets. It is locally presentable and cartesian closed. When denoting an explicit scaled simplicial set we will often omit the reference to the degenerate edges. For example, we will write $(\Delta^n, \Delta^{\{0,1,n\}})$ for the scaled simplicial set whose underlying simplicial set is the n -simplex and whose thin triangles are all the degenerate triangles together with the triangle $\Delta^{\{0,1,n\}}$.

Definition 1.2.2 The set of *generating scaled anodyne maps* \mathcal{S} is the set of maps of scaled simplicial sets consisting of

- (i) the inner horns inclusions

$$(\Lambda_i^n, \{\Delta^{\{i-1,i,i+1\}}\}_{\Lambda_i^n}) \rightarrow (\Delta^n, \{\Delta^{\{i-1,i,i+1\}}\}) \quad \text{for } n \geq 2 \text{ and } 0 < i < n;$$

- (ii) the map

$$(\Delta^4, T) \rightarrow (\Delta^4, T \cup \{\Delta^{\{0,3,4\}}, \Delta^{\{0,1,4\}}\}),$$

where

$$T \stackrel{\text{def}}{=} \{\Delta^{\{0,2,4\}}, \Delta^{\{1,2,3\}}, \Delta^{\{0,1,3\}}, \Delta^{\{1,3,4\}}, \Delta^{\{0,1,2\}}\};$$

- (iii) the set of maps

$$\left(\Lambda_0^n \coprod_{\Delta^{\{0,1\}}} \Delta^0, \{\Delta^{\{0,1,n\}}\} \right) \rightarrow \left(\Delta^n \coprod_{\Delta^{\{0,1\}}} \Delta^0, \{\Delta^{\{0,1,n\}}\} \right) \quad \text{for } n \geq 3.$$

A general map of scaled simplicial set is said to be *scaled anodyne* if it belongs to the weakly saturated closure of \mathcal{S} .

Definition 1.2.3 An *∞ -bicategory* is a scaled simplicial set \mathcal{C} which admits extensions along all maps in \mathcal{S} .


To avoid confusion we point out that scaled simplicial sets as in Definition 1.2.3 are referred to in [13] as *weak ∞ -bicatogories*, while the term ∞ -bicatogory was used to designate the stronger property of being fibrant in the *bicategorical model structure* on $\text{Set}_{\Delta}^{\text{sc}}$ constructed in loc. cit., whose cofibrations are the monomorphisms, and which serves as a model for the theory of $(\infty, 2)$ -categories. It is related to the model of marked simplicial categories mentioned above via a Quillen equivalence

$$\text{Set}_{\Delta}^{\text{sc}} \begin{array}{c} \xrightarrow{\mathcal{C}^{\text{sc}}} \\ \perp \\ \xleftarrow{N^{\text{sc}}} \end{array} \text{Set}_{\Delta}^{+}\text{-Cat},$$


in which the right functor N^{sc} is called as the *scaled coherent nerve*. Nonetheless, as we have shown in [8], the weak and strong notions of ∞ -bicatogory in fact coincide. In particular, the fibrant objects in the bicategorical model structure can be characterized by the extension property of Definition 1.2.3, and the notion of weak ∞ -bicatogory will not be further mentioned in the present paper.

Notation 1.2.4 We will refer to the weak equivalences in the bicategorical model structure as *bicategorical weak equivalences*. Since all the objects in the bicategorical model structure are cofibrant, the left Quillen equivalence \mathcal{C}^{sc} preserves and detects weak equivalences.

Notation 1.2.5 In a drawing, every 2-simplex filled by a 2-cell with the equivalence symbol, or simply filled by an equivalence symbol such as in the triangles




is a thin 2-simplex. As for 3-simplices, we will often draw them as planarized tetrahedra




where an additional equivalence symbol can appear in some of the triangles to indicate their thinness.

Definition 1.2.6 Given a 3-simplex $\rho: \Delta^3 \rightarrow X$ of the form



we will say that ρ exhibits σ' as the *left whiskering* of σ by f .

Similarly, a 3-simplex $\rho: \Delta^3 \rightarrow X$ of the form



will be said to exhibit σ' as the *right whiskering* of σ by f .

Notation 1.2.7 Let X be a simplicial set. We will denote by $X_b = (X, \text{deg}_2(X))$ the scaled simplicial set where the thin triangles of X are the degenerate 2-simplices, and by $X_\# = (X, X_2)$ the scaled simplicial set where all the triangles of X are thin. The assignments

$$X \mapsto X_b \quad \text{and} \quad X \mapsto X_\#$$

are left and right adjoint, respectively, to the forgetful functor $\text{Set}_\Delta^{\text{sc}} \rightarrow \text{Set}_\Delta$.

Definition 1.2.8 Given a scaled simplicial set X , we define its *core* to be the simplicial set X^{th} spanned by those n -simplices of X whose 2-dimensional faces are all thin triangles. The assignment $X \mapsto X^{\text{th}}$ is then right adjoint to the functor $(-)_\#: \text{Set}_\Delta \rightarrow \text{Set}_\Delta^{\text{sc}}$.

Warning 1.2.9 In [14, Tag 01XA], Lurie uses the term *pith* in place of core, and denotes it by $\text{Pith}(\mathcal{C})$.

Remark 1.2.10 If \mathcal{C} is an ∞ -bicategory then its core \mathcal{C}^{th} is an ∞ -category.

Definition 1.2.11 Let \mathcal{C} be an ∞ -bicategory. We will say that an edge in \mathcal{C} is *invertible* if it is invertible when considered in the ∞ -category \mathcal{C}^{th} , that is, if its image in the homotopy category of \mathcal{C}^{th} is an isomorphism. We will sometimes refer to invertible edges in \mathcal{C} as *equivalences*. We will denote by $\mathcal{C}^\simeq \subseteq \mathcal{C}^{\text{th}}$ the subsimplicial set spanned by the invertible edges. Then \mathcal{C}^\simeq is an ∞ -groupoid (that is, a Kan complex), which we call the *core groupoid* of \mathcal{C} . It can be considered as the ∞ -groupoid obtained from \mathcal{C} by discarding all noninvertible 1-cells and 2-cells. If X is an arbitrary scaled simplicial set then we will say that an edge in X is *invertible* if its image in \mathcal{C} is invertible for any bicategorical equivalence $X \rightarrow \mathcal{C}$ such that \mathcal{C} is an ∞ -bicategory. This does not depend on the choice of the ∞ -bicategory replacement \mathcal{C} .

Notation 1.2.12 Let \mathcal{C} be an ∞ -bicategory and let $x, y \in \mathcal{C}$ be two vertices. In [13, Section 4.2], Lurie gives an explicit model for the mapping ∞ -category from x to y in \mathcal{C} that we now recall. Let $\text{Hom}_{\mathcal{C}}(x, y)$ be the marked simplicial set whose n -simplices are given by maps $f: \Delta^n \times \Delta^1 \rightarrow \mathcal{C}$ such that $f|_{\Delta^n \times \{0\}}$ is constant on x , $f|_{\Delta^n \times \{1\}}$ is constant on y , and the triangle $f|_{\Delta^{\{(i,0),(i,1),(j,1)\}}}$ is thin for every $0 \leq i \leq j \leq n$. An edge $f: \Delta^1 \times \Delta^1 \rightarrow \mathcal{C}$ of $\text{Hom}_{\mathcal{C}}(x, y)$ is marked exactly when the triangle $f|_{\Delta^{\{(0,0),(1,0),(1,1)\}}}$ is thin. The assumption that \mathcal{C} is an ∞ -bicategory implies that the marked simplicial set $\text{Hom}_{\mathcal{C}}(x, y)$ is *fibrant* in the marked categorical model structure, that is, it is an ∞ -category whose marked edges are exactly the equivalences.

Definition 1.2.13 We will denote by $\mathcal{C}at_\infty$ the scaled coherent nerve of the (fibrant) marked simplicial subcategory $(\text{Set}_\Delta^+)^{\circ} \subseteq \text{Set}_\Delta^+$ spanned by the fibrant marked simplicial sets. We will refer to $\mathcal{C}at_\infty$ as the ∞ -bicategory of ∞ -categories.

Definition 1.2.14 We define BiCat_∞ to be the scaled coherent nerve of the (large) marked simplicial category BiCat_Δ whose objects are the ∞ -bicategories and whose mapping marked simplicial set, for $\mathcal{C}, \mathcal{D} \in \text{BiCat}_\Delta$, is given by $\text{BiCat}_\Delta(\mathcal{C}, \mathcal{D}) := \text{Fun}^{\text{th}}(\mathcal{C}, \mathcal{D})^{\natural}$. Here $\text{Fun}^{\text{th}}(\mathcal{C}, \mathcal{D})$ is the core ∞ -category of the internal hom scaled simplicial set $\text{Fun}(\mathcal{C}, \mathcal{D})$, which happens to be an ∞ -bicategory whenever \mathcal{D} is (see [13, Proposition 3.1.8 and Lemma 4.2.6]), and by $(-)^{\natural}$ we mean the associated marked simplicial set in which the marked arrows are the equivalences. We will refer to BiCat_∞ as the ∞ -bicategory of ∞ -bicategories.

Since the scaled coherent nerve functor \mathbb{N}^{sc} is a right Quillen equivalence, it determines an equivalence

$$(1) \quad (\text{Set}_\Delta^+ \text{-Cat})_\infty \xrightarrow{\simeq} \text{BiCat}_\infty^{\text{th}}$$

between the ∞ -category associated to the model category $\text{Set}_\Delta^+ \text{-Cat}$ and the core ∞ -category of BiCat_∞ . In the model $\text{Set}_\Delta^+ \text{-Cat}$ the $(\mathbb{Z}/2)^2$ -action on the theory of $(\infty, 2)$ -categories can be realized by an action of $(\mathbb{Z}/2)^2$ on $\text{Set}_\Delta^+ \text{-Cat}$ via model category isomorphisms: the operation $\mathcal{C} \mapsto \mathcal{C}^{\text{op}}$, which inverts only the direction of 1-morphisms, is realized by setting $\mathcal{C}^{\text{op}}(x, y) = \mathcal{C}(y, x)$, while the operation $\mathcal{C} \mapsto \mathcal{C}^{\text{co}}$ of inverting only the direction of 2-morphisms is realized by setting $\mathcal{C}^{\text{co}}(x, y) = \mathcal{C}(x, y)^{\text{op}}$, where the right-hand side denotes the operation of taking opposites in marked simplicial sets. Through the equivalence (1) these two involutions induce a $(\mathbb{Z}/2)^2$ -action on the core ∞ -category $\text{BiCat}_\infty^{\text{th}}$, which we then denote by the same notation. In particular, we have involutions

$$(-)^{\text{op}}: \text{BiCat}_\infty^{\text{th}} \rightarrow \text{BiCat}_\infty^{\text{th}} \quad \text{and} \quad (-)^{\text{co}}: \text{BiCat}_\infty^{\text{th}} \rightarrow \text{BiCat}_\infty^{\text{th}},$$

the first inverting the direction of 1-morphisms and the second the direction of 2-morphisms.

Remark 1.2.15 The $(\mathbb{Z}/2)^2$ -action on $\text{BiCat}_\infty^{\text{th}}$ does not extend to an action of $(\mathbb{Z}/2)^2$ on the ∞ -bicategory BiCat_∞ . Instead, it extends to a *twisted* action, that is, $(-)^{\text{co}}$ and $(-)^{\text{op}}$ extend to equivalences of the form

$$(-)^{\text{co}}: \text{BiCat}_\infty \xrightarrow{\simeq} \text{BiCat}_\infty \quad \text{and} \quad (-)^{\text{op}}: \text{BiCat}_\infty \xrightarrow{\simeq} \text{BiCat}_\infty^{\text{co}}.$$

1.3 Join and slice

In [8, Section 2.2] and [6, Section 2.1] we used join and slice constructions in the setting of marked scaled simplicial sets, that is, simplicial sets X endowed both with a distinguished collection $E_X \subseteq X_1$ of marked edges and a distinguished collection $T_X \subseteq X_2$ of thin triangles. The category of marked scaled simplicial sets will be denoted by $\text{Set}_\Delta^{+, \text{sc}}$. Though we will need only a limited amount of the generality used in [6], let us recall the construction and terminology used there for the sake of consistency. Given two marked

scaled simplicial sets $(X, E_X, T_X), (Y, E_Y, T_Y)$, their join is the *scaled* simplicial set $(X * Y, T_{X*Y})$ whose underlying simplicial set $X * Y$ is the usual join of simplicial sets and the collection of thin triangles is

$$T_{X*Y} := [T_X \times Y_0] \cup [E_X \times E_Y] \cup [X_0 \times T_Y] \subseteq [X_2 \times Y_0] \cup [X_1 \times Y_1] \cup [X_0 \times Y_2] = (X * Y)_2.$$

For a fixed marked scaled simplicial set (Y, E_Y, T_Y) , the functor $(X, E_X, T_X) \mapsto (X * Y, E_{X*Y})$ is a colimit-preserving functor from marked scaled simplicial sets to scaled simplicial sets under (Y, T_Y) , and admits a right adjoint, that is, an associated slice construction. Given a marked scaled simplicial set (K, E_K, T_K) and a map of scaled simplicial set $p: (K, T_K) \rightarrow (Z, T_Z)$, we will denote by $(Z, T_Z)_{/p}$, or simply $Z_{/p}$ for brevity, the valued of this right adjoint at p . In particular, $Z_{/p}$ is the marked scaled simplicial set characterized by the property that maps of marked scaled simplicial sets $(X, E_X, T_X) \rightarrow Z_{/p}$ correspond to maps of scaled simplicial sets $(X * K, T_{X*K}) \rightarrow (Z, T_Z)$ under (K, E_K, T_K) . We then write $\bar{Z}_{/p}$ for the *underlying scaled simplicial set* of $Z_{/p}$, obtained by forgetting the marking. To avoid confusion, let us emphasize that the thin triangles in $\bar{Z}_{/p}$ do depend on the marking E_K of K , and not only on the map of scaled simplicial sets p . Similarly, we denote by $Z_{p/}$ the marked scaled simplicial set representing the functor

$$(X, E_X, T_X) \mapsto \text{Map}_{(\text{Set}_\Delta^{\text{sc}})_{(K, T_K)}}((K * X, T_{K*X}), T_{X*K}, (Z, T_Z)),$$

and by $\bar{Z}_{p/}$ its underlying scaled simplicial set.

We will mostly be interested in the case where the target Z is an ∞ -bicategory \mathcal{C} , and K is either Δ^0 or $\# \Delta^1$, the latter being the 1-simplex endowed with the maximal marking and the (unique) trivial scaling. In the latter case we will sometimes make use of [6, Notation 2.3.4], which we now recall for the convenience of the reader.

Notation 1.3.1 Given a scaled simplicial set (Z, T_Z) and an edge $e: x \rightarrow y$ in Z , we will denote by $Z_{/e^\#} \in \text{Set}_\Delta^{+, \text{sc}}$ the result of the slice construction above applied to the marked scaled simplicial set $\# \Delta^1$ and the map of scaled simplicial sets $\Delta_b^1 \rightarrow (Z, T_Z)$ determined by e . Explicitly, the set of n -simplices in $Z_{/e^\#}$ is given by

$$(Z_{/e^\#})_n \stackrel{\text{def}}{=} \{\alpha: {}^b \Delta^n * \# \Delta^1 \rightarrow Z \mid \alpha|_{\Delta^{\{n+1, n+2\}}} = e\},$$

where ${}^b \Delta^n$ denotes the n -simplex with minimal marking and minimal scaling. The marked edges of $Z_{/e^\#}$ are those which factor through $\# \Delta^1 * \# \Delta^1$, and the thin triangles are those which factor through $(\Delta^2)_\#^b * \# \Delta^1$, where $(\Delta^2)_\#^b$ is the 2-simplex with minimal marking and maximal scaling, that is, its unique nondegenerate triangle is thin. As above, we will write $\bar{Z}_{/e^\#}$ for the underlying scaled simplicial set of $Z_{/e^\#}$.

When $K = \Delta^0$ the map p corresponds to a vertex $y \in \mathcal{C}$, and we will denote the associated marked scaled slice by $\mathcal{C}_{/y}$. The fiber $(\mathcal{C}_{/y})_x$ of the projection $\mathcal{C}_{/y} \rightarrow \mathcal{C}$ over a vertex x of \mathcal{C} is then a marked scaled

simplicial set all of whose triangles are thin. Its underlying marked simplicial set, denoted by $\text{Hom}_{\mathcal{C}}^{\triangleright}(x, y)$ in [7, Section 2.3], is fibrant in the marked categorical model structure, so that one can treat it as an ∞ -category (marked by its equivalences). As such, it is a model for the mapping ∞ -category of \mathcal{C} ; see [8, Proposition 2.23]. We then write $\overline{\text{Hom}}_{\mathcal{C}}^{\triangleright}(x, y)$ for the underlying simplicial set of $\text{Hom}_{\mathcal{C}}^{\triangleright}(x, y)$.

1.4 1-Inner/outer (co)cartesian fibrations

The theory of inner and outer (co)cartesian fibrations of ∞ -bicategories was developed in [6] as an analogue of the usual notion of (co)cartesian fibrations of ∞ -categories. As in the latter case, such a fibration encodes the data of a family of ∞ -categories functorially parametrized by the base \mathcal{B} , only that in the ∞ -bicategorical setting there are four different variance flavors for this functorial dependence one can consider. Specifically, inner (resp. outer) cocartesian fibrations encode a covariant dependence on the 1-morphisms of \mathcal{B} and a covariant (resp. contravariant) dependence on the level of 2-morphisms. Similarly, inner (resp. outer) cartesian fibrations encode a contravariant dependence on the 1-morphisms of \mathcal{B} and a contravariant (resp. covariant) dependence on the level of 2-morphisms.

Below we recall the main definitions. We refer the reader to loc. cit. for a comprehensive treatment.

Definition 1.4.1 We will say that a map of scaled simplicial sets $X \rightarrow Y$ is a *weak fibration* if it has the right lifting property with respect to the following types of maps:

- (i) All scaled inner horn inclusions of the form

$$(\Lambda_i^n, \{\Delta^{\{i-1, i, i+1\}}\}_{|\Lambda_i^n}) \subseteq (\Delta^n, \{\Delta^{\{i-1, i, i+1\}}\}) \quad \text{for } n \geq 2 \text{ and } 0 < i < n.$$

- (ii) The scaled horn inclusions of the form

$$\left(\Lambda_0^n \coprod_{\Delta^{\{0,1\}}} \Delta^0, \{\Delta^{\{0,1,n\}}\}_{|\Lambda_0^n} \right) \subseteq \left(\Delta^n \coprod_{\Delta^{\{0,1\}}} \Delta^0, \{\Delta^{\{0,1,n\}}\} \right) \quad \text{for } n \geq 2.$$

- (iii) The scaled horn inclusions of the form

$$\left(\Lambda_n^n \coprod_{\Delta^{\{n-1,n\}}} \Delta^0, \{\Delta^{\{0,n-1,n\}}\}_{|\Lambda_n^n} \right) \subseteq \left(\Delta^n \coprod_{\Delta^{\{n-1,n\}}} \Delta^0, \{\Delta^{\{0,n-1,n\}}\} \right) \quad \text{for } n \geq 2.$$

Remark 1.4.2 The maps appearing in Definition 1.4.1 are all trivial cofibrations with respect to the bicategorical model structure. This means that any bicategorical fibration $\mathcal{E} \rightarrow \mathcal{B}$ is in particular a weak fibration. For example, if \mathcal{E} is an ∞ -bicategory then the terminal map $\mathcal{E} \rightarrow \Delta^0$ is a weak fibration.

Definition 1.4.3 Given a weak fibration $f: X \rightarrow Y$, we will say that f is

- a *1-inner fibration* if it detects thin triangles and the underlying map of simplicial sets is an inner fibration, that is, satisfies the right lifting property with respect to inner horn inclusions;

- a 1-outer fibration if it detects thin triangles and the underlying map of simplicial sets satisfies the right lifting property with respect to the inclusions

$$\Lambda_0^n \coprod_{\Delta^{\{0,1\}}} \Delta^0 \subseteq \Delta^n \coprod_{\Delta^{\{0,1\}}} \Delta^0 \quad \text{and} \quad \Lambda_n^n \coprod_{\Delta^{\{n-1,n\}}} \Delta^0 \subseteq \Delta^n \coprod_{\Delta^{\{n-1,n\}}} \Delta^0 \quad \text{for } n \geq 2.$$

Note that the collection of 1-inner fibrations is closed under the $(-)^{\text{op}}$ duality, and the same holds for the collection of 1-outer fibrations.

Remark 1.4.4 In [8] and [6] we used the terms inner and outer fibration for what we called above 1-inner and 1-outer fibrations, respectively. The reason for the terminology update is the desire to more clearly distinguish between these notions and those of 2-inner and 2-outer fibrations introduced in the present paper. In principle, 1-inner and 1-outer fibrations between ∞ -bicategories are the functors which induce right and left fibrations, respectively, on the level of mapping ∞ -categories. The notions of 2-inner and 2-outer fibration between ∞ -bicategories correspond in turn to functors which induce cartesian and cocartesian fibrations on the level of mapping ∞ -categories, together with the condition that composition of arrows preserves cocartesian edges; see Section 4.2.

Remark 1.4.5 In [14, Tag 01WF], Lurie uses the term *interior fibration* to encode what we just defined as 1-outer fibrations. Our choice in [6] (which already appeared in [8]) is motivated by the intent of highlighting that *special outer horns* admit fillers against such maps.

Definition 1.4.6 Let $p: X \rightarrow Y$ be a weak fibration. We will say that an edge $e: \Delta^1 \rightarrow X$ is *p-cartesian* if the dotted lift exists in any diagram of the form

$$\begin{array}{ccc} (\Lambda_n^n, \{\Delta^{\{0,n-1,n\}}\}_{|\Lambda_n^n}) & \xrightarrow{\sigma} & (X, T_X) \\ \downarrow & \nearrow \text{dotted} & \downarrow p \\ (\Delta^n, \{\Delta^{\{0,n-1,n\}}\}) & \longrightarrow & (Y, T_Y) \end{array}$$

with $n \geq 2$ and $\sigma_{|\Delta^{\{n-1,n\}}} = e$. We will say that e is *p-cocartesian* if $e^{\text{op}}: \Delta^1 \rightarrow X^{\text{op}}$ is *p^{op}-cartesian*.

As in [6, Definition 2.3.1], we will also say that the edge $e: \Delta^1 \rightarrow X$ is *strongly f-(co)cartesian* if it is a (co)cartesian edge with respect to the underlying map of simplicial sets.

Remark 1.4.7 If $p: \mathcal{E} \rightarrow \mathcal{B}$ is weak fibration between ∞ -bicategories, then any equivalence in \mathcal{E} is both *p-cartesian* and *p-cocartesian*, see [6, Corollary 2.3.10]. On the other hand, if $e: x \rightarrow y$ is either a *p-cartesian* or *p-cocartesian* edge in \mathcal{E} such that pe is an equivalence in \mathcal{B} then e is an equivalence. To see this, let $g: y \rightarrow x$ be an inverse to pe in \mathcal{B} , equipped with thin triangles of the following forms:

$$\begin{array}{ccc} px & \xlongequal{\quad} & px \\ pe \searrow & \simeq & \nearrow g \\ & y & \end{array} \qquad \begin{array}{ccc} py & \xlongequal{\quad} & py \\ g \searrow & \simeq & \nearrow pe \\ & px & \end{array}$$

If e is p -cartesian then we can lift the right-hand side triangle to a triangle in \mathcal{E} of the form

$$\begin{array}{ccc}
 y & \xrightarrow{\quad \cong \quad} & y \\
 g' \searrow & & \nearrow e \\
 & x &
 \end{array}$$

producing in particular a right inverse g' to e . But then g' is also p -cartesian by [6, Lemma 2.3.8], and since $pg' = g$ is invertible in \mathcal{B} we deduce from the same argument that g' also has a right inverse in \mathcal{E} . It then follows from standard arguments (which can be applied on the level of the core ∞ -category \mathcal{C}^{th}) that e and g' are homotopy inverses, and in particular e is an equivalence. If e is assumed instead to be p -cocartesian then one proceeds in the same manner by lifting the left-hand side triangle to \mathcal{E} .

Remark 1.4.8 If $p: \mathcal{E} \rightarrow \mathcal{B}$ is a weak fibration and $e, e': \Delta^1 \rightarrow \mathcal{E}$ are two arrows which are equivalent in $\text{Fun}(\Delta^1, \mathcal{E})$, then e is p -cartesian if and only if e' is. To see this, note that in this case one has both an equivalence going from e to e' and an equivalence going from e' to e , and so it will suffice to show that if we have an equivalence $e \Rightarrow e'$ and e' is p -cartesian, then e is p -cartesian. Indeed, such an equivalence is given by a map $H: \Delta_b^1 \times \Delta_b^1 \rightarrow \mathcal{E}$ such that $H|_{\Delta_b^1 \times \Delta^{\{0\}}} = e$ and $H|_{\Delta_b^1 \times \Delta^{\{1\}}} = e'$. Since p -cartesian edges are closed under composition [6, Lemma 2.3.9] and every equivalence is p -cartesian (Remark 1.4.7), we get that H sends the diagonal edge $\Delta^{\{(0,0),(1,1)\}} \subseteq \Delta^1 \times \Delta^1$ to a p -cartesian edge. Then, from the partial two-out-of-three property for p -cartesian edges of [6, Lemma 2.3.8], we get that e is p -cartesian.

Passing to opposites, we also obtain from this argument that e is p -cocartesian if and only if e' is. In addition, if p is a 1-outer fibration then by [6, Proposition 2.3.7] the collection of strongly p -(co)cartesian arrows coincides with that of p -(co)cartesian arrows, and hence e is strongly p -(co)cartesian if and only if e' is so — alternatively, when p is a 1-outer fibration the statements of [6, Lemmas 2.3.8 and 2.3.9] also apply to strongly p -(co)cartesian arrows, so that the above argument can simply be carried out verbatim.

Definition 1.4.9 Let $f: X \rightarrow Y$ be a weak fibration of scaled simplicial sets. We will say that f is a *cartesian fibration* if for every $x \in X$ and an edge $e: y \rightarrow f(x)$ in Y there exists a f -cartesian edge $\tilde{e}: \tilde{y} \rightarrow x$ such that $f(\tilde{e}) = e$. Dually, we will say that $f: X \rightarrow Y$ is a *cocartesian fibration* if $f^{\text{op}}: X^{\text{op}} \rightarrow Y^{\text{op}}$ is a cartesian fibration.

Definition 1.4.10 Let $f: X \rightarrow Y$ be a weak fibration of scaled simplicial sets. We will say that f is a 1-inner (resp. 1-outer) cartesian fibration if it is both a 1-inner (resp. 1-outer) fibration and a cartesian fibration. Dually, we will say that f is a 1-inner (resp. 1-outer) cocartesian fibration if p^{op} is a 1-inner (resp. 1-outer) cartesian fibration.

Remark 1.4.11 The classes of weak fibrations, 1-inner/outer fibrations and 1-inner/outer (co)cartesian fibrations are all closed under base change.

Example 1.4.12 If \mathcal{C} is an ∞ -bicategory, then the projection $\bar{\mathcal{C}}_{/x} \rightarrow \mathcal{C}$ is an example of a 1–outer cartesian fibration, where the cartesian edges are exactly those whose corresponding triangle in \mathcal{C} is thin (equivalently, those corresponding to marked edges in $\mathcal{C}_{/x}$); see [6, Corollary 2.4.7]. Similarly, $\bar{\mathcal{C}}_{x/} \rightarrow \mathcal{C}$ is a 1–outer cocartesian fibration, with the cocartesian edges again corresponding to thin triangles. More generally, for any map of scaled simplicial sets $p: (K, T_K) \rightarrow \mathcal{C}$, the associated slice projections $\bar{\mathcal{C}}_{/p} \rightarrow \bar{\mathcal{C}}$ and $\bar{\mathcal{C}}_{p/} \rightarrow \mathcal{C}$ are 1–outer cartesian and cocartesian fibrations, respectively.

Proposition 1.4.13 Any (co)cartesian fibration between ∞ -bicategories is a fibration in the bicategorical model structure on $\text{Set}_\Delta^{\text{sc}}$.

Proof We prove the cartesian case, from which the cocartesian case can be deduced by passing to opposites. Let $p: \mathcal{E} \rightarrow \mathcal{B}$ be a cartesian fibration between ∞ -bicategories. To show that p is a bicategorical fibration we need to produce the dotted lift in any square of the form

$$(2) \quad \begin{array}{ccc} K & \xrightarrow{f} & \mathcal{E} \\ \downarrow & \nearrow & \downarrow \\ L & \xrightarrow{g} & \mathcal{B} \end{array}$$

such that $K \rightarrow L$ is a bicategorical trivial cofibration of scaled simplicial sets. Since \mathcal{E} is an ∞ -bicategory it is in particular fibrant in the bicategorical model structure (see discussion in Section 1.2), and hence we can extend f to a map $h: L \rightarrow \mathcal{E}$. This is not yet a solution to the above lifting problem since the composite ph might be different from g . The two maps ph and g agree however on K by construction. Since the bicategorical model structure is cartesian closed and \mathcal{B} is fibrant, we may solve the lifting problem

$$\begin{array}{ccc} [K \times \Delta_b^1] \coprod_{K \times \partial \Delta^1} [L \times \partial \Delta^1] & \longrightarrow & \mathcal{B} \\ \downarrow & \nearrow & \\ L \times \Delta_b^1 & & \end{array}$$

yielding a natural transformation $H: L \times \Delta_b^1 \rightarrow \mathcal{B}$ from g and ph which is constant on K . Since $K \rightarrow L$ is a trivial cofibration and \mathcal{B} is fibrant the induced functor $\text{Fun}(L, \mathcal{B}) \rightarrow \text{Fun}(K, \mathcal{B})$ is an equivalence of ∞ -bicategories, and since the arrow in $\text{Fun}(L, \mathcal{B})$ associated to H maps to an identity arrow in $\text{Fun}(K, \mathcal{B})$ we deduce that it must be invertible in $\text{Fun}(L, \mathcal{B})$. In particular, the restriction of H to $\{l\} \times \Delta_b^1$ is an invertible arrow of \mathcal{B} for every vertex $l \in L$. Now since p is a cartesian fibration we can choose for each $l \in L_0 \setminus K_0$ a p -cartesian lift $e_l: \{l\} \times \Delta_b^1 \rightarrow \mathcal{E}$ of $H|_{\{l\} \times \Delta_b^1}$. Then each e_l is a p -cartesian lift of an equivalence, and is hence itself an equivalence, see Remark 1.4.7. Let $L' \subseteq L$ be the scaled simplicial subset whose underlying simplicial set is that of L and whose thin triangles are only those thin triangles which are contained in K . Applying [8, Proposition 2.38] we may solve the lifting problem

$$\begin{array}{ccc} [K \times \Delta_b^1] \coprod_{K \times \Delta^{\{1\}}} [L' \times \Delta^{\{1\}}] & \longrightarrow & \mathcal{E} \\ \downarrow & \nearrow G & \downarrow \\ L' \times \Delta_b^1 & \xrightarrow{H} & \mathcal{B} \end{array}$$

to yield a natural transformation $G: L' \times \Delta_b^1 \rightarrow \mathcal{E}$ such that $G|_{\{l\} \times \Delta_b^1} = e_l$ for every $l \in L_0 \setminus K_0$, where we point out that the assumption made in [8, Proposition 2.38] that p detects thin triangles is not needed since L' does not contain any thin triangles that are not in K , see [8, Remark 2.40]. In particular, G is a levelwise invertible natural transformation. We wish to show that $G|_{L' \times \Delta^{\{0\}}}$ extends to $L \times \Delta^{\{0\}}$, thus providing a solution to the original lifting problem (2). Indeed, if $\sigma: \Delta_b^2 \rightarrow L'$ is a triangle that is thin in L then the composed map

$$H_\sigma: \Delta_b^2 \times \Delta_b^1 \xrightarrow{\sigma \times \text{id}} L' \times \Delta_b^1 \xrightarrow{G} \mathcal{E}$$

is a levelwise invertible natural transformation between two triangles, one of which is thin (since $G|_{L' \times \Delta^{\{0\}}}$ extends to $L \times \Delta_b^1$ by construction), and hence the other one is thin as well by [8, Corollary 3.5]. \square

Over a fixed base \mathcal{B} , the collection of 1–inner cocartesian fibrations, and cocartesian edges preserving functors between them, can be organized into an ∞ –bicategory $\text{coCar}^{\text{inn}}(\mathcal{B})$. This ∞ –bicategory can be presented by a suitable model structure on the category of marked simplicial sets over the underlying simplicial set of \mathcal{B} , developed in [13, Section 3.2] using the machinery of categorical patterns. Lurie then constructs in loc. cit. a straightening–unstraightening Quillen equivalence between this model structure and the projective model structure on $\text{Fun}(\mathcal{C}^{\text{sc}}(\mathcal{B}), \text{Set}_\Delta^+)$. In [6] we used this to establish the following ∞ –bicategorical form of the Grothendieck–Lurie correspondence, for both cartesian and cocartesian, inner and outer flavors of fibrations:

Theorem 1.4.14 [6, Corollary 3.3.3] *For an ∞ –bicategory $\mathcal{B} \in \text{BiCat}_\infty$ there are natural equivalences of ∞ –bicategories*

$$\begin{aligned} \text{coCar}^{\text{inn}}(\mathcal{B}) &\simeq \text{Fun}(\mathcal{B}, \text{Cat}_\infty), & \text{coCar}^{\text{out}}(\mathcal{B}) &\simeq \text{Fun}(\mathcal{B}^{\text{co}}, \text{Cat}_\infty), \\ \text{Car}^{\text{inn}}(\mathcal{B}) &\simeq \text{Fun}(\mathcal{B}^{\text{coop}}, \text{Cat}_\infty), & \text{Car}^{\text{out}}(\mathcal{B}) &\simeq \text{Fun}(\mathcal{B}^{\text{op}}, \text{Cat}_\infty). \end{aligned}$$

2 2–Inner/outer cartesian fibrations

In this section we will define the principal notion of this paper, namely that of 2–inner/outer (co)cartesian fibrations, and study their basic properties.

2.1 Inner and outer triangles

Definition 2.1.1 Let $p: \mathcal{E} \rightarrow \mathcal{B}$ be a weak fibration of ∞ –bicategories and $\sigma: \Delta^2 \rightarrow \mathcal{E}$ a triangle.

- We will say that σ is *left p –inner* if the corresponding arrow in $\bar{\mathcal{E}}_{/\sigma(2)}$ is strongly cartesian with respect to the projection $\bar{\mathcal{E}}_{/\sigma(2)} \rightarrow \mathcal{E} \times_{\mathcal{B}} \bar{\mathcal{B}}_{/p\sigma(2)}$.
- We will say that σ is *right p –inner* if the corresponding arrow in $\bar{\mathcal{E}}_{\sigma(0)/}$ is strongly cocartesian with respect to the projection $\bar{\mathcal{E}}_{\sigma(0)/} \rightarrow \mathcal{E} \times_{\mathcal{B}} \bar{\mathcal{B}}_{p\sigma(0)/}$.
- We will say that σ is *left p –outer* if the corresponding arrow in $\bar{\mathcal{E}}_{/\sigma(2)}$ is strongly cocartesian with respect to the projection $\bar{\mathcal{E}}_{/\sigma(2)} \rightarrow \mathcal{E} \times_{\mathcal{B}} \bar{\mathcal{B}}_{/p\sigma(2)}$.

- We will say that σ is *right p -outer* if the corresponding arrow in $\bar{\mathcal{E}}_{\sigma(0)}$ is strongly cartesian with respect to the projection $\bar{\mathcal{E}}_{\sigma(0)} \rightarrow \mathcal{E} \times_{\mathcal{B}} \bar{\mathcal{B}}_{p\sigma(0)}$.

Remark 2.1.2 Unwinding the definitions, we see that σ is left p -inner if and only if, for $n \geq 3$, every commutative square of the form

$$(3) \quad \begin{array}{ccc} & \Delta^{\{n-2, n-1, n\}} & \\ & \swarrow \quad \searrow \sigma & \\ \Lambda_{n-1}^n & \xrightarrow{\quad} & \mathcal{E} \\ \downarrow & \nearrow \text{dotted} & \downarrow p \\ \Delta^n & \xrightarrow{\quad} & \mathcal{B} \end{array}$$

admits a diagonal filler as displayed by the dotted arrow, and right p -inner if the same holds for diagrams as above where Λ_{n-1}^n is replaced by Λ_1^n and $\Delta^{\{n-2, n-1, n\}}$ by $\Delta^{\{0, 1, 2\}}$. On the other hand, σ is left p -outer if and only if every commutative square of the form

$$(4) \quad \begin{array}{ccc} & \Delta^{\{0, 1, n\}} & \\ & \swarrow \quad \searrow \sigma & \\ \Lambda_0^n & \xrightarrow{\quad} & \mathcal{E} \\ \downarrow & \nearrow \text{dotted} & \downarrow p \\ \Delta^n & \xrightarrow{\quad} & \mathcal{B} \end{array}$$

admits a diagonal filler as displayed by the dotted arrow, and right p -outer if and only if the same holds for diagrams as above where Λ_0^n is replaced by Λ_n^n and $\Delta^{\{0, 1, n\}}$ by $\Delta^{\{0, n-1, n\}}$.

Remark 2.1.3 It follows from Remark 2.1.2 that any thin triangle in \mathcal{E} is both left and right p -inner. On the other hand, any thin triangle whose left leg is p -cocartesian is left p -outer and any thin triangle whose right leg p -cartesian is right p -outer.

Remark 2.1.3 admits a type of a converse statement:

Proposition 2.1.4 *Let $p: \mathcal{E} \rightarrow \mathcal{B}$ be a weak fibration of ∞ -bicategories. Suppose $\sigma: \Delta^2 \rightarrow \mathcal{E}$ is a triangle such that $p(\sigma)$ is thin in \mathcal{B} . If σ is either left or right p -inner then σ is thin in \mathcal{E} . The same holds if we assume that σ is left p -outer and left-degenerate or that σ is right p -outer and right-degenerate.*

Proof Write $x = \sigma(0), z = \sigma(2)$. Suppose first that σ is left p -inner. The condition that $p(\sigma)$ is thin means that the arrow determined by $p(\sigma)$ in $\bar{\mathcal{B}}_{/pz}$ is cartesian with respect to the projection $\bar{\mathcal{B}}_{/pz} \rightarrow \mathcal{B}$; see Example 1.4.12. By base change it then follows that the arrow determined by σ in $\mathcal{E} \times_{\mathcal{B}} \bar{\mathcal{B}}_{/pz}$ is cartesian with respect to the projection $\mathcal{E} \times_{\mathcal{B}} \bar{\mathcal{B}}_{/pz} \rightarrow \mathcal{E}$. At the same time, the arrow determined by σ in $\bar{\mathcal{E}}_{/z}$ is cartesian with respect to the projection $\bar{\mathcal{E}}_{/z} \rightarrow \mathcal{E} \times_{\mathcal{B}} \bar{\mathcal{B}}_{/pz}$ (by the definition of being left p -inner)

and so we conclude that the arrow determined by σ in $\bar{\mathcal{E}}_{/z}$ is cartesian with respect to the composed projection $\bar{\mathcal{E}}_{/z} \rightarrow \mathcal{E}$. By Example 1.4.12 we then get that σ is thin. The dual argument using $\bar{\mathcal{E}}_{x/}$ and $\bar{\mathcal{B}}_{px/}$ applies to the case where σ is right p -inner.

Now suppose that σ is left p -outer and left-degenerate. Then $p\sigma$ is left-degenerate and since $p\sigma$ is assumed thin it follows that the arrow in $\bar{\mathcal{B}}_{/pz}$ determined by $p(\sigma)$ is invertible (indeed, by Example 1.4.12 it is a cocartesian arrow with respect to the 1-outer fibration $\bar{\mathcal{B}}_{/pz} \rightarrow \bar{\mathcal{B}}$ lying over an equivalence). Since σ is left-degenerate we then have that the arrow determined by σ in $\mathcal{E} \times_{\mathcal{B}} \bar{\mathcal{B}}_{/pz}$ is invertible as well. Since σ is left p -outer it now follows that the arrow in $\bar{\mathcal{E}}_{/z}$ determined by σ is a cocartesian lift of an equivalence along the 1-outer fibration $\bar{\mathcal{E}}_{/z} \rightarrow \mathcal{E} \times_{\mathcal{B}} \bar{\mathcal{B}}_{/pz}$, and is hence itself an invertible arrow $\bar{\mathcal{E}}_{/z}$. As such, this arrow is in particular cocartesian with respect to the projection $\bar{\mathcal{E}}_{/z} \rightarrow \mathcal{E}$ (Remark 1.4.7), and so we conclude that σ is thin by Example 1.4.12. The dual argument using $\bar{\mathcal{E}}_{x/}$ and $\bar{\mathcal{B}}_{px/}$ applies to the case where σ is right p -outer and right-degenerate. □

Remark 2.1.5 By [6, Lemma 2.3.9 and Lemma 2.3.8] the collection of strongly (co)cartesian arrows in a given 1-outer fibration is closed under composition and has a partial two-out-of-three property. More precisely, if $\mathcal{C} \rightarrow \mathcal{D}$ is a 1-outer fibration of ∞ -bicategories and

$$\begin{array}{ccc} & y & \\ f \nearrow & & \searrow g \\ x & \xrightarrow{h} & z \end{array}$$

is a thin triangle in \mathcal{C} such that g is strongly cartesian then f is strongly cartesian if and only if h is strongly cartesian. Dually, if f is strongly cocartesian then g is strongly cocartesian if and only if h is strongly cocartesian. Now for any weak fibration $p: \mathcal{E} \rightarrow \mathcal{B}$ of ∞ -bicategories, both $\bar{\mathcal{E}}_{x/} \rightarrow \mathcal{E} \times_{\mathcal{B}} \bar{\mathcal{B}}_{x/}$ and $\bar{\mathcal{E}}_{/x} \rightarrow \mathcal{E} \times_{\mathcal{B}} \bar{\mathcal{B}}_{/x}$ are 1-outer fibrations for every $x \in \mathcal{E}$ by Remark 1.4.11 and Example 1.4.12. The above partial two-out-of-three property for strongly (co)cartesian edges then translates to a certain partial two-out-of-three property for inner/outer triangles. More precisely, suppose given a 3-simplex $\rho: \Delta^3 \rightarrow \mathcal{E}$ of the form

$$\begin{array}{ccc} x_0 & \xrightarrow{\quad} & x_3 \\ \downarrow & \searrow \sigma & \uparrow \\ x_1 & \xrightarrow{f} & x_2 \end{array} \quad \Longrightarrow \quad \begin{array}{ccc} x_0 & \xrightarrow{\quad} & x_3 \\ \downarrow & \searrow \tau & \uparrow \\ x_1 & \xrightarrow{f} & x_2 \end{array}$$

If θ is thin then we may consider ρ as encoding a thin triangle in $\bar{\mathcal{E}}_{/x_3}$ exhibiting the edge associated to σ as the composite of those associated to τ and η , whereas if η is thin then we may consider ρ as encoding thin a triangle in $\bar{\mathcal{E}}_{x_0/}$ exhibiting the edge associated to τ as the composite of those associated to θ and σ . We hence conclude the following:

- (i) If θ is thin and η is left p -inner then τ is left p -inner if and only if σ is.
- (ii) If η is thin and θ is right p -inner then σ is right p -inner if and only if τ is.

- (iii) If θ is thin and τ is left p -outer then η is left p -outer if and only if σ is.
- (iv) If η is thin and σ is right p -outer then θ is right p -outer if and only if τ is.

Combining this with Remark 2.1.3, we conclude that the collection of left p -inner triangles in \mathcal{E} is closed under left whiskering with 1-morphisms and the collection of right p -inner triangles is closed under right whiskering with 1-morphisms. On the other hand, the collection of left p -outer triangles is only closed under left whiskering with p -cocartesian 1-morphisms and the collection of right p -outer triangles is only closed under right whiskering with p -cartesian arrows.

Combining Remark 2.1.5(4) with Remark 2.1.3 we get that if

$$\begin{array}{ccc}
 x_0 & \longrightarrow & x_3 \\
 \downarrow & \searrow \theta & \uparrow g \\
 x_1 & \longrightarrow & x_2
 \end{array}
 \xrightarrow{\cong}
 \begin{array}{ccc}
 x_0 & \longrightarrow & x_3 \\
 \downarrow & \searrow \tau & \uparrow g \\
 x_1 & \longrightarrow & x_2
 \end{array}$$

is a 3-simplex such that g is p -cartesian then θ is right p -outer if and only if τ is. The “only if” direction of this implication also holds for right p -inner triangles by Remark 2.1.5(2) and Remark 2.1.3. The following lemma shows that on the other hand, the “if” direction of this implication actually holds for left p -inner and left p -outer triangles:


Lemma 2.1.6 *Let $p: \mathcal{E} \rightarrow \mathcal{B}$ be a weak fibration. Given a 3-simplex $\rho: \Delta^3 \rightarrow \mathcal{E}$ as above with g being p -cartesian, if τ is left p -inner or left p -outer then so is θ .*

Proof Consider the commutative diagram

$$\begin{array}{ccccc}
 \bar{\mathcal{E}}_{/x_2} & \xleftarrow{\cong} & \bar{\mathcal{E}}_{/g^\#} & \xrightarrow{\quad} & \bar{\mathcal{E}}_{/x_3} \\
 \downarrow q_2 & & \downarrow q_{2,3} & & \downarrow q_3 \\
 \bar{\mathcal{E}} \times_{\mathcal{B}} \bar{\mathcal{B}}_{/px_2} & \xleftarrow{\cong} & \mathcal{E} \times_{\mathcal{B}} \bar{\mathcal{B}}_{/pg^\#} & \xrightarrow{\quad} & \mathcal{E} \times_{\mathcal{B}} \bar{\mathcal{B}}_{/px_3} \\
 & & \downarrow & & \downarrow \\
 & & \bar{\mathcal{B}}_{/pg^\#} & \xrightarrow{\quad} & \bar{\mathcal{B}}_{/px_3}
 \end{array}$$

in which the left-pointing horizontal arrows are trivial fibrations by [6, Lemma 2.4.6]. Now since the two faces of ρ leaning on g are thin we have that the 3-simplex ρ determines an arrow e in $\bar{\mathcal{E}}_{/g^\#}$, whose image in $\bar{\mathcal{E}}_{/x_2}$ is the arrow associated to θ . We conclude that θ is left p -inner (resp. left p -outer) if and only if e is $q_{2,3}$ -cartesian (resp. $q_{2,3}$ -cocartesian). Now the image of e in $\bar{\mathcal{E}}_{/x_3}$ is the arrow associated to τ , and since τ is assumed to be left p -inner (resp. left p -outer) its associated arrow is q_3 -cartesian (resp. q_3 -cocartesian) in $\bar{\mathcal{E}}_{/x_3}$. At the same time, the assumption that g is p -cartesian implies that the vertical external square on the right is homotopy cartesian by [6, Lemma 2.3.6], and since the bottom right square is also cartesian we get from the pasting lemma that the top right square is homotopy cartesian. We hence conclude that e is $q_{2,3}$ -cartesian (resp. $q_{2,3}$ -cocartesian), as desired. \square

Remark 2.1.7 Inner and outer lifts in a given weak fibration are unique up to equivalence (once they exist). For example, suppose we have two left p -inner triangles α, α' in \mathcal{E} whose restrictions to Λ_1^2 and whose images under p coincide. By Remark 2.1.2 (applied for σ' and $n = 3$) we can find a 3-simplex H of the form



where β lives over a degenerate triangle in \mathcal{B} . By Remark 2.1.5 we then have that β is left p -inner as well, and hence thin by Proposition 2.1.4. We may consider H as exhibiting an equivalence between α and α' : its leftmost leg is invertible, the two faces leaning on this leg are thin, and the remaining two faces are α and α' . In a very similar manner, if α and α' are assumed instead to be right p -outer and such that their restrictions to Λ_2^2 and images under p coincide, then we construct the same type of 3-simplex H , only that this time we will take its $\Delta^{\{0,1,3\}}$ -face to be degenerate and extend H from its right outer horn using the right p -outerness of α . Replacing p with $p^{\text{op}}: \mathcal{E}^{\text{op}} \rightarrow \mathcal{B}^{\text{op}}$ we get the analogous statements for the uniqueness of *right* inner and *left* outer lifts.

Remark 2.1.8 Let $p: \mathcal{E} \rightarrow \mathcal{B}$ be a weak fibration of ∞ -bicategories. For given $x, z \in \mathcal{E}$, if we base change the weak fibration $\bar{\mathcal{E}}_{/z} \rightarrow \mathcal{E} \times_{\mathcal{B}} \bar{\mathcal{B}}_{/pz}$ along the map $\{x\} \times_{\mathcal{B}} \bar{\mathcal{B}}_{/pz} \rightarrow \mathcal{E} \times_{\mathcal{B}} \bar{\mathcal{B}}_{/pz}$ then we get the map of maximally scaled simplicial sets, whose underlying map of simplicial sets is

$$p_{x,z}^{\triangleright}: \overline{\text{Hom}}_{\mathcal{E}}^{\triangleright}(x, z) \rightarrow \overline{\text{Hom}}_{\mathcal{B}}^{\triangleright}(px, pz),$$

which is a model for the induced map on mapping ∞ -categories by [8, Section 2.3]; see in particular [8, Proposition 2.23]. Since base change maps detect (co)cartesian edges, it follows that for a triangle $\sigma: \Delta^2 \rightarrow \mathcal{E}$ such that $\sigma|_{\Delta^{\{0,1\}}}$ is degenerate, we have:

- If σ is left p -inner then the corresponding edge of $\overline{\text{Hom}}_{\mathcal{E}}^{\triangleright}(x, z)$ is $p_{x,z}^{\triangleright}$ -cartesian.
- If σ is left p -outer then the corresponding edge of $\overline{\text{Hom}}_{\mathcal{E}}^{\triangleright}(x, z)$ is $p_{x,z}^{\triangleright}$ -cocartesian.

2.2 2-Inner and 2-outer fibrations

Definition 2.2.1 Let $p: \mathcal{E} \rightarrow \mathcal{B}$ be a weak fibration and $\sigma: \Delta^2 \rightarrow \mathcal{B}$ a triangle.

- We say that σ has a *sufficient supply of left (resp. right) p -inner lifts* if for every $\rho: \Lambda_1^2 \rightarrow \mathcal{E}$ lifting $\sigma|_{\Lambda_1^2}$ there exists a left (resp. right) p -inner triangle $\tau: \Delta^2 \rightarrow \mathcal{E}$ such that $p\tau = \sigma$ and $\tau|_{\Lambda_1^2} = \rho$.
- We say that σ has a *sufficiently supply of left p -outer lifts* if for every $\rho: \Lambda_0^2 \rightarrow \mathcal{E}$ lifting $\sigma|_{\Lambda_0^2}$ and such that $\rho|_{\Delta^{\{0,1\}}}$ is p -cocartesian, there exists a left p -outer triangle $\tau: \Delta^2 \rightarrow \mathcal{E}$ such that $p\tau = \sigma$ and $\tau|_{\Lambda_0^2} = \rho$.

- We say that σ has a *sufficient supply of right p -outer lifts* if for every $\rho: \Lambda_2^2 \rightarrow \mathcal{E}$ lifting $\sigma|_{\Lambda_0^2}$ and such that $\rho|_{\Delta_{\{1,2\}}}$ is p -cartesian, there exists a right p -outer triangle $\tau: \Delta^2 \rightarrow \mathcal{E}$ such that $p\tau = \sigma$ and $\tau|_{\Lambda_2^2} = \rho$.

Definition 2.2.2 A weak fibration of scaled simplicial sets $p: \mathcal{E} \rightarrow \mathcal{B}$ is said to be a *2-inner fibration* if every triangle in \mathcal{B} admits both a sufficient supply of left p -inner lifts and a sufficient supply of right p -inner lifts.

We say that p is a *2-inner cartesian fibration* if it is both a 2-inner fibration and a cartesian fibration (in the sense of Definition 1.4.9). We say that p is a *2-inner cocartesian fibration* if $p^{\text{op}}: \mathcal{E}^{\text{op}} \rightarrow \mathcal{B}^{\text{op}}$ is a 2-inner cartesian fibration.

Definition 2.2.3 A weak fibration of scaled simplicial sets $p: \mathcal{E} \rightarrow \mathcal{B}$ is said to be a *2-outer fibration* if the following conditions are satisfied:

- Every triangle in \mathcal{B} admits both a sufficient supply of left p -outer lifts and a sufficient supply of right p -outer lifts.
- The collection of left p -outer triangles in \mathcal{E} is closed under right whiskering and the collection of right p -outer triangles is closed under left whiskering.

We say that p is a *2-outer cartesian fibration* if it is both a 2-outer fibration and a cartesian fibration (in the sense of Definition 1.4.9). Dually, we say that p is a *2-outer cocartesian fibration* if $p^{\text{op}}: \mathcal{E}^{\text{op}} \rightarrow \mathcal{B}^{\text{op}}$ is a 2-outer cartesian fibration.

Remark 2.2.4 For the previously introduced classes of fibrations, the class of 2-inner/outer (co)cartesian fibrations is readily seen to be closed under base change.

Definition 2.2.5 Given 2-inner/outer (co)cartesian fibrations $q: \mathcal{D} \rightarrow \mathcal{A}$ and $p: \mathcal{E} \rightarrow \mathcal{B}$, a *morphism of 2-inner/outer (co)cartesian fibrations* from q to p is a commutative square

$$\begin{array}{ccc} \mathcal{D} & \xrightarrow{g} & \mathcal{E} \\ q \downarrow & & \downarrow p \\ \mathcal{A} & \xrightarrow{f} & \mathcal{B} \end{array}$$

such that g sends q -(co)cartesian arrows to p -(co)cartesian arrows and left/right p -inner/outer triangles to left/right q -inner/outer triangles.

Proposition 2.2.6 Let $p: \mathcal{E} \rightarrow \mathcal{B}$ be a weak fibration of ∞ -bicategories. Then:

- If p is a 2-inner fibration then the induced map $p_{x,y}: \overline{\text{Hom}}_{\mathcal{E}}^{\triangleright}(x, y) \rightarrow \overline{\text{Hom}}_{\mathcal{B}}^{\triangleright}(px, py)$ is a cartesian fibration of ∞ -categories for every $x, y \in \mathcal{E}$.
- If p is a 2-outer fibration then the induced map $p_{x,y}: \overline{\text{Hom}}_{\mathcal{E}}^{\triangleleft}(x, y) \rightarrow \overline{\text{Hom}}_{\mathcal{B}}^{\triangleleft}(px, py)$ is a cocartesian fibration of ∞ -categories for every $x, y \in \mathcal{E}$.

Proof The condition that p is a weak fibration implies that the induced map $\bar{\mathcal{E}}_{/y} \rightarrow \mathcal{E} \times_{\mathcal{B}} \bar{\mathcal{B}}_{/py}$ is a 1–outer fibration (see Remark 1.4.11 and Example 1.4.12), and hence its base change $(\bar{\mathcal{E}}_{/y})_x \rightarrow (\bar{\mathcal{B}}_{/py})_{px}$ is a 1–outer fibration as well, and in particular a weak fibration. This last map is between ∞ –bicategories in which every triangle is thin, and hence its underlying map of simplicial sets

$$p_{x,y}: \overline{\text{Hom}}_{\mathcal{E}}^{\triangleright}(x, y) \rightarrow \overline{\text{Hom}}_{\mathcal{B}}^{\triangleright}(px, py)$$

is an inner fibration between ∞ –categories. We also note that arrows in $\overline{\text{Hom}}_{\mathcal{B}}^{\triangleright}(px, py)$ correspond to triangles $\sigma: \Delta^2 \rightarrow \mathcal{B}$ such that $\sigma|_{\Delta\{0,1\}}$ is degenerate on px and $\sigma|_{\Delta\{2\}} = py$. More precisely, these are arrows from $\sigma|_{\Delta\{0,2\}}$ to $\sigma|_{\Delta\{1,2\}}$, considered as vertices in $\overline{\text{Hom}}_{\mathcal{B}}^{\triangleright}(px, py)$.

Now if p is a 2–inner fibration, then for any choice of an edge $g: x \rightarrow y$ in \mathcal{E} lifting $\sigma|_{\Delta\{1,2\}}$, there exists a left p –inner triangle τ such that $\tau|_{\Delta\{0,1\}}$ is degenerate, $\tau|_{\Delta\{1,2\}} = g$ and $p\tau = \sigma$. By Remark 2.1.8 the triangle τ determines a $p_{x,y}$ –cartesian lift with target g of the arrow in $\overline{\text{Hom}}_{\mathcal{B}}^{\triangleright}(px, py)$ determined by σ . Since g was arbitrary we conclude that $p_{x,y}$ is a cartesian fibration. Similarly, if p is a 2–outer fibration we have that for any choice of an edge $g: x \rightarrow y$ in \mathcal{E} lifting $\sigma|_{\Delta\{0,2\}}$, there exists a left p –outer triangle τ such that $\tau|_{\Delta\{0,1\}}$ is degenerate, $\tau|_{\Delta\{0,2\}} = g$ and $p\tau = \sigma$. By Remark 2.1.8, the triangle τ determined a $p_{x,y}$ –cocartesian lift with domain g of the arrow in $\overline{\text{Hom}}_{\mathcal{B}}^{\triangleright}(px, py)$ determined by σ , and so $p_{x,y}$ is now a cocartesian fibration. □

2.3 Congruent triangles


Our goal in the present subsection is to establish some preliminary results showing that for most questions about 2–outer (co)cartesian fibrations between ∞ –bicategories, one may restrict attention to left/right outer triangles which are left/right-degenerate in the following sense:

Definition 2.3.1 We say that a triangle $\sigma: \Delta^2 \rightarrow X$ is *left degenerate* if the edge $\sigma|_{\Delta\{0,1\}}$ of X is degenerate. The triangle σ is said to be *right degenerate* if the edge $\sigma|_{\Delta\{1,2\}}$ is degenerate.

Warning 2.3.2 To avoid confusion, let us emphasize that left (or right) degenerate triangles in the sense of Definition 2.3.1 are not necessarily themselves degenerate. This terminology is also used in [1].

We will make use of the following construction.

Definition 2.3.3 Let X be a scaled simplicial set. Given a 3–simplex $\rho: \Delta^3 \rightarrow X$ of the form



we say that ρ exhibits σ as *left congruent* to σ' , and σ' as *right congruent* to σ .

Lemma 2.3.4 *Let $p: \mathcal{E} \rightarrow \mathcal{B}$ be a weak fibration and let σ and σ' be two triangles in \mathcal{E} such that σ is left congruent to σ' via a 3-simplex ρ as in Definition 2.3.3. Then the following holds:*

- (i) *The triangle σ is left/right p -inner if and only if σ' is.*
- (ii) *If f is p -cocartesian and σ' is left p -outer, then σ is left p -outer.*
- (iii) *If f is p -cartesian and σ is right p -outer, then σ' is right p -outer.*
- (iv) *If f is an equivalence, then σ is left/right p -outer if and only if σ' is.*

Proof The first three statements follows directly from the partial two-out-of-three properties elaborated in Remark 2.1.5 together with the fact that thin triangles are always left and right p -inner and that thin triangles with p -cocartesian left leg are left p -outer, while thin triangles with p -cartesian right leg are right p -outer; see Remark 2.1.3. To prove the last claim, we note that if f is an equivalence then the 3-simplex ρ determines in particular an equivalence between the arrows associated to σ and σ' in $\bar{\mathcal{E}}_{x_0/}$, where $x_0 = \sigma(0) = \sigma'(0)$, and hence each of these arrows is strongly cartesian with respect to the projection $\bar{\mathcal{E}}_{x_0/} \rightarrow \mathcal{E} \times_{\mathcal{B}} \mathcal{B}_{p x_0/}$ if and only if the other is so; see Remark 1.4.8. Similarly, ρ determines an equivalence between the arrows associated to σ and σ' in $\bar{\mathcal{E}}_{/x_3}$, where $x_3 = \sigma(2) = \sigma'(2)$, and hence each of these arrows is strongly cocartesian with respect to the projection $\bar{\mathcal{E}}_{/x_3} \rightarrow \mathcal{E} \times_{\mathcal{B}} \mathcal{B}_{/p x_3}$ if and only if the other is so. □

Lemma 2.3.5 *Let $p: \mathcal{E} \rightarrow \mathcal{B}$ be a weak fibration between ∞ -bicategories and let σ and σ' be two triangles in \mathcal{B} such that σ is left congruent to σ' . Then the following holds:*

- (i) *If the left leg of σ' admits a sufficient supply of p -cocartesian lifts and σ' admits a sufficient supply of left p -outer lifts, then σ admits a sufficient supply of left p -outer lifts.*
- (ii) *If the right leg of σ admits a sufficient supply of p -cartesian lifts and σ admits a sufficient supply of right p -outer lifts, then σ' admits a sufficient supply of right p -outer lifts.*

Proof We prove the first claim, the second claim then follows by applying the first claim to \mathcal{E}^{op} and switching the roles of σ and σ' . Let $\rho: \Delta^3 \rightarrow \mathcal{E}$ be a 3-simplex as above exhibiting σ as left congruent to σ' . In particular, $\sigma = \rho|_{\Delta^{\{0,2,3\}}}$. We need to show that for every pair of arrows

$$\begin{array}{ccc}
 y_0 & \xrightarrow{e_{0,3}} & y_3 \\
 & \searrow & \\
 & & y_2
 \end{array}$$

of \mathcal{E} lifting $\sigma|_{\Delta^{\{0,2\}}}$ and $\sigma|_{\Delta^{\{0,3\}}}$, respectively, with $e_{0,2}$ being p -cocartesian, there exists a left p -outer triangle lifting σ :

$$\begin{array}{ccc}
 y_0 & \xrightarrow{e_{0,3}} & y_3 \\
 & \searrow & \Downarrow \\
 & & x_2 \\
 & \searrow & \uparrow e_{2,3} \\
 & & x_2
 \end{array}$$

Now by assumption, the left leg of σ' admits a p -cocartesian lift $e_{0,1}: y_0 \rightarrow y_1$. Since σ' is assumed to have a sufficient supply of left p -outer lifts, we may now find a left p -outer lift τ' of σ' , depicted as

$$\begin{array}{ccc} y_0 & \xrightarrow{e_{0,3}} & y_3 \\ e_{0,1} \downarrow & \Downarrow & \nearrow e_{1,3} \\ y_1 & & \end{array}$$

At the same time, since $\rho|_{\Delta^{\{0,1,2\}}}$ is thin and $e_{0,1}$ is p -cocartesian the pair $e_{0,1}, e_{0,2}$ extends to a thin triangle $\theta: \Delta^{\{0,1,2\}} \rightarrow \mathcal{E}$ such that $p\theta = \rho|_{\Delta^{\{0,1,2\}}}$. Set $e_{1,2} = \theta|_{\Delta^{\{1,2\}}}$. By the two-out-of-three property for p -cocartesian edges — eg the dual of [6, Lemma 2.3.8] — we get that $e_{1,2}$ is p -cocartesian. Since $\rho|_{\Delta^{\{1,2,3\}}}$ is thin we may now lift it to a triangle $\eta: \Delta^{\{1,2,3\}} \rightarrow \mathcal{E}$ extending $e_{1,2}$ and $e_{1,3}$. The triangles τ', θ and η now glue to give a map $\alpha: \Lambda_1^3 \rightarrow \mathcal{E}$ lifting $\rho|_{\Lambda_1^3}$, where α sends $\Delta^{\{0,1,2\}}$ and $\Delta^{\{1,2,3\}}$ to thin triangles by construction. Since p is a weak fibration we may extend α to a map $\beta: \Delta^3 \rightarrow \mathcal{E}$ lifting ρ , so that $\tau := \beta|_{\Delta^{\{0,2,3\}}}$ gives in particular a triangle lifting σ and extending $e_{0,2}, e_{0,3}$. The 3-simplex β now exhibits τ as left congruent to τ' and hence τ is left p -outer by Lemma 2.3.4(2). \square

Remark 2.3.6 In the proof of Lemma 2.3.5(1) we have complete freedom in choosing the p -cocartesian lift $e_{0,1}$ of $\sigma'|_{\Delta^{\{0,1\}}}$. We may consequently slightly weaken the assumption on σ' : it suffices to assume that for every choice of lift of $e_{1,3}$ of $\sigma'|_{\Delta^{\{1,2\}}}$ there exists some left p -outer lift τ' of σ' such that $\tau|_{\Delta^{\{1,2\}}} = e_{1,3}$ and $\tau|_{\Delta^{\{0,1\}}}$ is p -cartesian (as opposed to assuming this for any choice of p -cocartesian lift of $\tau|_{\Delta^{\{0,1\}}}$). For example, if σ' is left degenerate then it suffices to assume that it has a sufficient supply of *left degenerate* left p -outer lifts (that is, for each choice of a lift of $\sigma'|_{\Delta^{\{1,2\}}}$). A similar statement holds for the right p -outer case of Lemma 2.3.5(2).

Lemma 2.3.7 *Let \mathcal{B} be an ∞ -bicategory. Then the following holds:*

- (i) *Any triangle σ in \mathcal{B} is left congruent to a left degenerate triangle σ' .*
- (ii) *Any triangle σ' in \mathcal{B} is right congruent to a right degenerate triangle σ .*


Proof We prove the first claim. The second claim then follows by applying the first statement to \mathcal{B}^{op} and switching the roles of σ and σ' . Let us depict σ as

$$\begin{array}{ccc} x_0 & \xrightarrow{h} & x_3 \\ & \searrow f & \Downarrow \\ & & x_2 \end{array} \quad \begin{array}{c} \uparrow g \\ \uparrow g \\ \uparrow g \end{array}$$

Let $K \subseteq \Delta^3$ be the simplicial subset spanned by the faces $\Delta^{\{0,2,3\}}$ and $\Delta^{\{0,1,2\}}$ and let $\rho: K \rightarrow \mathcal{B}$ be the map which sends $\Delta^{\{0,2,3\}}$ to σ and $\Delta^{\{0,1,2\}}$ to the degenerate triangle whose left leg is degenerate and whose other two legs are f . We may then visualize ρ as

$$\begin{array}{ccc} x_0 & \xrightarrow{h} & x_3 \\ \parallel & \searrow f & \Downarrow \\ \parallel & = f & \uparrow g \\ x_1 & \xrightarrow{f} & x_2 \end{array}$$

Now since \mathcal{B} is an ∞ -bicategory we may extend ρ to a map $\rho' : \Lambda_2^3 \rightarrow \mathcal{B}$ which sends the triangle $\Delta^{\{1,2,3\}}$ to a thin triangle, and then proceed to extend ρ' to a full 3-simplex $\Delta^3 \rightarrow \mathcal{B}$, which we can depict as



This 3-simplex then exhibits σ as left congruent to σ' , and σ' is left degenerate, as desired. □

Corollary 2.3.8 *Let $p : \mathcal{E} \rightarrow \mathcal{B}$ be a weak fibration. If every left degenerate triangle in \mathcal{B} has a sufficient supply of left p -outer lifts, then every triangle in \mathcal{B} has a sufficient supply of left p -outer lifts. Similarly, if every right degenerate triangle in \mathcal{B} has a sufficient supply of right p -outer lifts then every triangle in \mathcal{B} has a sufficient supply of right p -outer lifts.*

Proof Combine Lemma 2.3.7 with Lemma 2.3.5 using the fact that any degenerate arrow in \mathcal{B} has a sufficient supply of (co)cartesian lifts. □

Using Remark 2.3.6 we may also obtain the following strengthening of Corollary 2.3.8:

Corollary 2.3.9 *Let $p : \mathcal{E} \rightarrow \mathcal{B}$ be a weak fibration. If every left degenerate triangle in \mathcal{B} has a sufficient supply of left degenerate left p -outer lifts, then every triangle in \mathcal{B} has a sufficient supply of left p -outer lifts. Similarly, if every right degenerate triangle in \mathcal{B} has a sufficient supply of right-degenerate right p -outer lifts then every triangle in \mathcal{B} has a sufficient supply of right p -outer lifts.*

2.4 Homotopy invariance of fibrations

Our goal in this section is to prove the following homotopy invariance property for (co)cartesian fibrations.

Proposition 2.4.1 *Let*

$$\begin{array}{ccc}
 \mathcal{D} & \xrightarrow{\cong} & \mathcal{E} \\
 q \downarrow & & \downarrow p \\
 \mathcal{A} & \xrightarrow{\cong} & \mathcal{B}
 \end{array}$$

be a commutative diagram of ∞ -bicategories whose vertical maps are both bicategorical fibrations and whose horizontal maps are bicategorical equivalences. Then p is a 2-inner/outer (co)cartesian fibration if and only if q is. In addition, an edge in \mathcal{D} is q -(co)cartesian if and only if its image in \mathcal{E} is p -(co)cartesian, and similarly a triangle in \mathcal{D} is left/right q -inner/outer if and only if its image in \mathcal{E} is so with respect to p .

The proof of Proposition 2.4.1 will require a couple of lemmas, and will be given at the end of the section.

Lemma 2.4.2 *Let*

$$\begin{array}{ccc} \mathcal{D} & \xrightarrow{\cong} & \mathcal{E} \\ q \downarrow & & \downarrow p \\ \mathcal{A} & \xrightarrow{\cong} & \mathcal{B} \end{array}$$

be a commutative diagram of ∞ -bicategories whose vertical maps are both bicategorical fibrations and whose horizontal maps are bicategorical equivalences. Then an arrow in \mathcal{D} is q -(co)cartesian if and only if its image in \mathcal{E} is p -(co)cartesian.

Proof By [6, Proposition 2.3.7] we may replace the property of being cartesian with that of being weakly (co)cartesian. The desired claim now follows from the characterization [6, Proposition 2.3.3] of weakly (co)cartesian arrows in terms of mapping spaces. \square

In any weak fibration $p: \mathcal{E} \rightarrow \mathcal{B}$ between ∞ -bicategories, the collection of p -(co)cartesian arrows is closed under equivalences in the arrow category; see [6, Remark 2.3.12]. We now show that a similar property holds for inner/outer triangles:


Lemma 2.4.3 *Let $p: \mathcal{E} \rightarrow \mathcal{B}$ be a weak fibration between ∞ -bicategories, and let $H: \Delta_b^1 \times \Delta_b^2 \rightarrow \mathcal{E}$ be levelwise invertible natural transformation between triangles. Then $\sigma_0 := H|_{\Delta_{\{0\}} \times \Delta^2}$ is left/right p -inner/outer if and only if $\sigma_1 := H|_{\Delta_{\{1\}} \times \Delta^2}$ is so.*

Proof To fix ideas we prove the left inner and left outer cases, the right inner and right outer cases then follow by replacing p with opposite. In addition, it will suffice to prove that σ_0 is left p -inner/outer as soon as σ_1 is such, since we can get the other direction by replacing H by the inverse equivalence. We hence assume that σ_1 is left p -inner/outer.

For $i = 0, 1, 2$, let $\rho_i: \Delta^3 \rightarrow \Delta^1 \times \Delta^2$ be the 3-simplex given by


$$\rho_i(j) = \begin{cases} (0, j) & \text{if } j \leq i, \\ (1, j - 1) & \text{if } j > i. \end{cases}$$

We may then write $H\rho_0$ as




for some triangle τ . Since σ_1 is left p -inner/outer we get from the two-out-of-three properties of Remark 2.1.5 that τ is left p -inner/outer (in the outer case, we also point out that the triangle $H\rho_0|_{\Delta_{\{0,1,3\}}}$

is left p -outer by Remarks 2.1.3 and 1.4.7, since it is thin and its left leg is an equivalence). We now write $H\rho_1$ as



for some triangle τ' . Since τ is left p -inner/outer we now get from Points (1) and (4) of Lemma 2.3.4 that τ' is left p -inner/outer. Finally, we may write $H\rho_2$ as




and so by Lemma 2.1.6 we conclude that σ_0 is left p -inner/outer as well, as desired. □


Proof of Proposition 2.4.1 We first note that since both vertical arrows are fibrant and cofibrant in the arrow category (with respect to the projective model structure), the existence of a levelwise equivalence from q to p implies the existence of a levelwise equivalence from p to q . It will hence suffice to show that if p is a 2-inner/outer (co)cartesian fibration then so is q . Now since weak equivalences between fibrant objects are preserved under base change along fibrations it follows from the two-out-of-three property that the map $\mathcal{D} \rightarrow \mathcal{E} \times_{\mathcal{B}} \mathcal{A}$ is an equivalence of ∞ -bicategories. The map $\mathcal{E}' := \mathcal{E} \times_{\mathcal{B}} \mathcal{A} \rightarrow \mathcal{A}$ is then again a 2-inner/outer (co)cartesian fibration (Remark 2.2.4). We now factor the weak equivalence $\mathcal{D} \xrightarrow{\simeq} \mathcal{E}'$ as a composite

$$\mathcal{D} \xrightarrow{\simeq} \mathcal{D}' \xrightarrow{\simeq} \mathcal{E}'$$

where the first map is a trivial cofibration and the second a trivial fibration. Then $\mathcal{D}' \rightarrow \mathcal{E}' \rightarrow \mathcal{A}$ is a composite of a 2-inner/outer (co)cartesian fibration and a trivial fibration of scaled simplicial sets, and is hence itself again a 2-inner/outer (co)cartesian fibration. On the other hand, since $\mathcal{D} \rightarrow \mathcal{D}'$ is a trivial cofibration and $\mathcal{D} \rightarrow \mathcal{A}$ is in particular a bicategorical fibration we may solve the lifting problem



so that we obtain a retract diagram of arrows



Now by Lemma 2.4.2 the map $\mathcal{D}' \rightarrow \mathcal{D}$ preserves (co)cartesian arrows over \mathcal{A} , and hence the fact that \mathcal{D}' has a sufficient supply of q' –(co)cartesian edges implies that \mathcal{D} has a sufficient supply of q –(co)cartesian edges. It is thus left to show that \mathcal{D} also has a sufficient supply of left/right q –inner/outer triangles. Solving a lifting problem of the form

$$\begin{array}{ccc}
 [\Delta^1 \times \mathcal{D}] \amalg_{\partial \Delta^1 \times \mathcal{D}'} [\partial \Delta^1 \times \mathcal{D}] & \xrightarrow{\quad} & \mathcal{D}' \\
 \downarrow & \dashrightarrow^H & \downarrow q' \\
 \mathcal{D}' \times \Delta^1 & \xrightarrow{\quad} & \mathcal{A}
 \end{array}$$

we obtain a natural transformation H over \mathcal{A} from $i \circ r : \mathcal{D}' \rightarrow \mathcal{D}'$ to the identity $\text{id}_{\mathcal{D}'}$, whose restriction to \mathcal{D} is constant on i . Since i is a trivial cofibration, it is in particular essentially surjective, and so the natural transformation H is levelwise invertible. We now prove the inner case. Suppose we are given a triangle $\sigma : \Delta^2 \rightarrow \mathcal{A}$ and a lift $\rho : \Lambda_1^2 \rightarrow \mathcal{D}$ of $\sigma|_{\Lambda_1^2}$. Since q' is a left/right inner fibration we may extend $i\rho$ to a left/right q' –inner triangle $\tau : \Delta^2 \rightarrow \mathcal{D}'$ lifting σ . Evaluating the levelwise invertible natural transformation H at τ we obtain an equivalence $i r \tau \Rightarrow \tau$ covering the identity transformation on τ and restricting to the identity transformation from $i\rho$ to itself on Λ_1^2 . By Lemma 2.4.3 we deduce that $i r \tau$ is also q' –inner. Since i admits a retraction and by the explicit description of left/right inner triangles in terms of lifting properties as in Remark 2.1.2, we then deduce that $r \tau$ itself is a left/right q –inner extension of ρ lifting σ . In the outer case, the argument for the existence of a sufficient supply of left/right q –outer triangles is completely analogous, but one also needs to show the closure of q –outer triangles under whiskering as required in Definition 2.2.3(2). But this again follows from the corresponding property for q' and the fact that $i : \mathcal{D} \rightarrow \mathcal{D}'$ detects left/right outer triangles, as it admits a retraction over \mathcal{A} . \square

2.5 Equivalences of cartesian fibrations

This section is devoted to the proof of a fiberwise criterion to test equivalences of fibrations.

Theorem 2.5.1 *Consider a morphism of 2–inner/outer (co)cartesian fibrations (in the sense of Definition 2.2.5) given by a commutative diagram of the form*

$$(5) \quad \begin{array}{ccc}
 \mathcal{E} & \xrightarrow{r} & \mathcal{E}' \\
 p \downarrow & & \downarrow q \\
 \mathcal{B} & \xrightarrow{f} & \mathcal{B}'
 \end{array}$$

where p and q are 2–inner/outer cartesian fibrations between ∞ –bicategories. Suppose that f is an equivalence of ∞ –bicategories. Then r is an equivalence if and only if the induced map $r_b : \mathcal{E}_b \rightarrow \mathcal{E}'_{f(b)}$ on the level of fibers is an equivalence of ∞ –bicategories for all $b \in \mathcal{B}$.

Before proving this, we need a preliminary result, generalizing [12, Proposition 2.4.4.2]. Here we make use of the results from [6] involving the slice ∞ –bicategories associated with marked scaled simplicial sets; see Section 1.3 and Notation 1.3.1.

Lemma 2.5.2 *Let $p: \mathcal{C} \rightarrow \mathcal{D}$ be a 2-inner/outer cartesian fibration of ∞ -bicategories. Let $x, y \in \mathcal{C}$ be two objects, $\bar{e}: px \rightarrow py$ an arrow between their images in \mathcal{D} , and $e: x' \rightarrow y$ a p -cartesian lift of \bar{e} . Then the (homotopy) fiber of the map*

$$\phi: \overline{\text{Hom}}_{\mathcal{C}}^{\triangleright}(x, y) \rightarrow \overline{\text{Hom}}_{\mathcal{D}}^{\triangleright}(px, py)$$

at \bar{e} is naturally equivalent to the mapping ∞ -category $\overline{\text{Hom}}_{\mathcal{C}_{px}}^{\triangleright}(x, x')$, where \mathcal{C}_{px} denotes the (homotopy) fiber of p over px .

Proof To begin with we note that the map ϕ is a cartesian (or cocartesian in the outer case) fibration of ∞ -categories by Proposition 2.2.6 and hence the homotopy fiber in question is equivalent to the corresponding strict fiber. We now consider the following diagram

$$\begin{array}{ccc} & \overline{\mathcal{C}}_{/e\#} & \\ \cong \swarrow & & \searrow \cong \\ \overline{\mathcal{C}}_{/x'} \times_{\overline{\mathcal{D}}/px'} \overline{\mathcal{D}}_{/\bar{e}\#} & & \overline{\mathcal{C}}_{/y} \times_{\overline{\mathcal{D}}/py} \overline{\mathcal{D}}_{/\bar{e}\#} \end{array}$$

where the left diagonal map is a trivial fibration by [6, Lemma 2.4.6] and the right diagonal map is a trivial fibration by [6, Lemma 2.3.6] and the assumption that e is p -cartesian (and is hence in particular weakly p -cartesian, in the sense of loc. cit.). Taking fibers over $(x, s_1(\bar{e})) \in \mathcal{C} \times_{\mathcal{D}} \mathcal{D}_{/\bar{e}\#}$ we thus get a zig-zag of equivalences

$$\begin{array}{ccc} & \overline{\mathcal{C}}_{/e\#} \times_{\mathcal{C} \times_{\mathcal{D}} \mathcal{D}_{/\bar{e}\#}} \{(x, s_1(\bar{e}))\} & \\ \cong \swarrow & & \searrow \cong \\ \overline{\text{Hom}}_{\mathcal{C}_{p(x)}}^{\triangleright}(x, x')_{\#} & & \phi^{-1}(\bar{e})_{\#} \end{array}$$

relating the fiber of ϕ over \bar{e} to $\overline{\text{Hom}}_{\mathcal{C}_{p(x)}}^{\triangleright}(x, x')$, as desired. □

Proof of Theorem 2.5.1 We prove the 2-inner cartesian case. The proof for the remaining three variance flavors proceeds in exactly the same manner. Let us begin by proving the “only if” direction of the statement. Every object in the square (5) is fibrant, and the vertical maps are fibrations by Proposition 1.4.13, therefore every pullback along these maps is automatically a homotopy pullback. Consider the following commutative cube:

$$\begin{array}{ccccc} & \mathcal{E}_b & \xrightarrow{r_b} & \mathcal{E}'_{f(b)} & \\ & \swarrow & \downarrow r & \swarrow & \downarrow \\ \mathcal{E} & \xrightarrow{\quad} & \mathcal{E}' & & \\ p \downarrow & & \downarrow & & \downarrow \\ \mathcal{B} & \xrightarrow{f} & \mathcal{B}' & & \\ \{b\} \swarrow & & \downarrow q & & \swarrow \{f(b)\} \\ & \Delta^0 & \xrightarrow{\quad} & \Delta^0 & \end{array}$$

The front face and the side faces are homotopy pullbacks by assumption, so that the back one must also be such. Since the bottom horizontal map is an equivalence, the map r_b must be an equivalence as well.

Assume now that r is a fiberwise equivalence, and let us prove it is essentially surjective on objects and fully faithful. To see that r is essentially surjective, factor it as a composite $\mathcal{E} \rightarrow \mathcal{E}' \times_{\mathcal{B}'} \mathcal{B} \rightarrow \mathcal{E}'$, where the first map is essentially surjective since $r_b: \mathcal{E}_b \rightarrow \mathcal{E}'_{fb}$ is essentially surjective for every $b \in \mathcal{B}$, and the second is essentially surjective because $f: \mathcal{B} \rightarrow \mathcal{B}'$ is so. Concerning full-faithfulness, we consider the following commutative square of ∞ -categories:

$$(6) \quad \begin{array}{ccc} \overline{\text{Hom}}_{\mathcal{E}}^{\triangleright}(x, y) & \xrightarrow{r} & \overline{\text{Hom}}_{\mathcal{E}'}^{\triangleright}(rx, ry) \\ p \downarrow & & \downarrow q \\ \overline{\text{Hom}}_{\mathcal{B}}^{\triangleright}(px, py) & \xrightarrow{f} & \overline{\text{Hom}}_{\mathcal{B}'}^{\triangleright}(qrx, qry) \end{array}$$

Here, we have used the fact that $qr = fp$, and we denoted the induced action between the ∞ -categories of morphisms with the same letter as that of the map between the relevant ∞ -bicategories. Now, the vertical maps are cartesian fibrations of ∞ -categories by Proposition 2.2.6, and the bottom horizontal map is an equivalence since f is fully faithful. Therefore, the top horizontal map is an equivalence if and only if the square (6) is a homotopy pullback, which happens precisely if the maps induced between the (strict) fibers of the vertical maps are equivalences. Thanks to Lemma 2.5.2, the fiber of the left-hand side map over $\bar{e}: p(x) \rightarrow p(y)$ coincides with $\overline{\text{Hom}}_{\mathcal{E}_{px}}^{\triangleright}(x, x')$ for some p -cartesian 1-simplex $e: x' \rightarrow y$ that lifts \bar{e} . Since r is assumed to preserve cartesian edges we have that $re: rx' \rightarrow ry$ is a q -cartesian lift of $f\bar{e}$, and so the fiber of q over $f(\bar{e})$ is given by $\overline{\text{Hom}}_{\mathcal{E}'_{qrx}}^{\triangleright}(rx, rx')$. We can now finish the proof by observing that the induced map

$$\overline{\text{Hom}}_{\mathcal{E}_{px}}^{\triangleright}(x, x') \rightarrow \overline{\text{Hom}}_{\mathcal{E}'_{qrx}}^{\triangleright}(rx, rx')$$

is an equivalence of ∞ -categories, since $\mathcal{E}_{px} \rightarrow \mathcal{E}'_{fpx} \simeq \mathcal{E}'_{qrx}$ is assumed to be an equivalence of ∞ -bicategories. □

3 The domain projection

In this section we analyze a key example of a 2-outer cartesian fibration: the domain projection $d: \text{Fun}^{\text{gr}}(\Delta^1, \mathcal{C}) \rightarrow \mathcal{C}$ induced by the inclusion $\{0\} \hookrightarrow \Delta^1$, where $\text{Fun}^{\text{gr}}(\Delta^1, \mathcal{C})$ is the ∞ -bicategory whose objects are the arrows in \mathcal{C} and whose morphisms are the lax-commutative squares. It is characterized by the property that for every scaled simplicial set (K, T_K) , maps $(K, T_K) \rightarrow \text{Fun}^{\text{gr}}(\Delta^1, \mathcal{C})$ correspond to maps of scaled simplicial sets

$$\Delta_b^1 \otimes (K, T_K) \rightarrow \mathcal{C},$$

where \otimes denotes the *Gray product* of scaled simplicial sets [7, Definition 2.1]. Explicitly, $\Delta_b^1 \otimes (K, T_K)$ is the scaled simplicial set whose underlying simplicial set is the cartesian product $\Delta^1 \times K$, and where a

triangle $\sigma: \Delta^2 \rightarrow \Delta^1 \times K$ is thin if and only if its image in K belongs to T_K and either $\sigma|_{\Delta^{\{0,1\}}}$ maps to a degenerate edge in K or $\sigma|_{\Delta^{\{1,2\}}}$ maps to a degenerate edge in Δ^1 . Here, we have used the fact that all the triangles in Δ_b^1 are thin, otherwise the additional condition of projecting to a thin triangle in the first factor would have been necessary.

Remark 3.0.1 By the main result of [7] the functor $- \otimes -$ is a left Quillen bifunctor. It then follows that for any ∞ -bicategory \mathcal{C} , the domain projection

$$d: \text{Fun}^{\text{gr}}(\Delta^1, \mathcal{C}) \rightarrow \mathcal{C}$$

is a bicategorical fibration of ∞ -bicategories, and in particular a weak fibration; see Remark 1.4.2.

Our argument to show that the domain fibration is a 2–outer cartesian fibrations is based on the following extension lemma.

Lemma 3.0.2 *Let \mathcal{C} be an ∞ -bicategory, and for $n \geq 2$ suppose we are given an extension problem of the form*

$$\begin{array}{ccc} \left([\Delta^1 \times \Lambda_n^n] \coprod_{\partial \Delta^1 \times \Lambda_n^n} [\partial \Delta^1 \times \Delta^n], T' \right) & \xrightarrow{\rho} & \mathcal{C} \\ \downarrow & \nearrow \text{dotted} & \\ (\Delta^1 \times \Delta^n, T) & & \end{array}$$

where

$$T = \{ \Delta^{\{(0,n-1),(0,n),(1,n)\}}, \Delta^{\{(0,0),(1,n-1),(1,n)\}} \}$$

and T' is its restriction to the top left corner. If ρ sends $\Delta^{\{1\}} \times \Delta^{\{n-1,n\}}$ to an invertible edge in \mathcal{C} , then the dotted extension exists.

Proof We define a sequence of scaled maps

$$\tau_1, \dots, \tau_{n-1}: (\Delta^n, \Delta^{\{0,n-1,n\}}) \rightarrow (\Delta^1 \times \Delta^n, T)$$

in the following manner:

$$\tau_i(j) = \begin{cases} (0, j) & \text{if } j < i, \\ (1, j) & \text{if } j \geq i. \end{cases}$$

Set $K_0 := ([\Delta^1 \times \Lambda_n^n] \coprod_{\partial \Delta^1 \times \Lambda_n^n} [\partial \Delta^1 \times \Delta^n], T')$ and $K_i := K_{i-1} \cup \tau_i$ for $1 \leq i \leq n-1$, so that for every $1 \leq i \leq n-1$ we have pushout diagrams of the form

$$\begin{array}{ccc} (\Lambda_n^n, \{ \Delta^{\{0,n-1,n\}} \}) & \longrightarrow & K_{i-1} \\ \downarrow & & \downarrow \\ (\Delta^n, \{ \Delta^{\{0,n-1,n\}} \}) & \xrightarrow{\tau_i} & K_i \end{array}$$

These all have the property that the edge $\Delta^{\{n-1,n\}}$ is mapped to an equivalence in \mathcal{C} . All these maps admit lifts against \mathcal{C} by [6, Corollary 2.3.10], and so we can extend the map ρ to a map $\rho': K_{n-1} \rightarrow \mathcal{C}$. Consider now the $(n + 1)$ -simplices $\sigma_0, \dots, \sigma_n: \Delta^{n+1} \rightarrow \Delta^1 \times \Delta^n$ defined as

$$\sigma_i(j) = \begin{cases} (0, j) & \text{if } j \leq i, \\ (1, j - 1) & \text{if } j > i. \end{cases}$$

Observe that $\sigma_0, \dots, \sigma_{n-1}$ can be promoted to maps

$$(\Delta^{n+1}, \{\Delta^{\{0,n,n+1\}}\}) \rightarrow (\Delta^1 \times \Delta^n, T)$$

whereas σ_n can be promoted to a map

$$(\Delta^{n+1}, \{\Delta^{\{n-1,n,n+1\}}\}) \rightarrow (\Delta^1 \times \Delta^n, T)$$

We now set $K_{n+k} = K_{n-1+k} \cup \sigma_{n-k}$ for $0 \leq k \leq n$, and observe that $K_{2n} = (\Delta^1 \times \Delta^n, T)$. We then have a pushout diagram of the form

$$\begin{array}{ccc} (\Delta_n^{n+1}, \{\Delta^{\{n-1,n,n+1\}}\}|_{\Delta_n^{n+1}}) & \longrightarrow & K_{n-1} \\ \downarrow & & \downarrow \\ (\Delta^{n+1}, \{\Delta^{\{n-1,n,n+1\}}\}) & \xrightarrow{\sigma_n} & K_n \end{array}$$

and, for every $k > 0$, there are pushout diagrams of the form

$$\begin{array}{ccc} (\Delta_{n+1}^{n+1}, \{\Delta^{\{0,n,n+1\}}\}) & \longrightarrow & K_{n-1+k} \\ \downarrow & & \downarrow \\ (\Delta^{n+1}, \{\Delta^{\{0,n,n+1\}}\}) & \xrightarrow{\sigma_{n-k}} & K_{n+k} \end{array}$$

where the corresponding image of the edge $\Delta^{\{n,n+1\}}$ in \mathcal{C} is an equivalence. As before, this is enough to prove the existence of an extension to \mathcal{C} . □

Remark 3.0.3 Applying Lemma 3.0.2 to \mathcal{C}^{op} one obtains the following dual form of its statement, which we spell out for the convenience of the reader: suppose given an extension problem of the form

$$\begin{array}{ccc} \left([\Delta^1 \times \Lambda_0^n] \coprod_{\partial \Delta^1 \times \Lambda_0^n} [\partial \Delta^1 \times \Delta^n], T' \right) & \xrightarrow{\rho} & \mathcal{C} \\ \downarrow & \dashrightarrow & \\ (\Delta^1 \times \Delta^n, T) & & \end{array}$$

where

$$T = \{ \Delta^{\{(0,0),(1,0),(1,1)\}}, \Delta^{\{(0,0),(0,1),(1,n)\}} \}$$

and T' is its restriction to the top left corner. If ρ sends $\Delta^{\{1\}} \times \Delta^{\{0,1\}}$ to an invertible edge in \mathcal{C} then the dotted lift exists.

We now consider the question of cartesian 1-simplices for the domain projection.

Lemma 3.0.4 *Let \mathcal{C} be an ∞ -bicategory and consider the domain projection $d: \text{Fun}^{\text{gr}}(\Delta^1, \mathcal{C}) \rightarrow \mathcal{C}$ as above. Then a 1-simplex $\alpha: \Delta^1 \rightarrow \text{Fun}^{\text{gr}}(\Delta^1, \mathcal{C})$ is d -cartesian if its transpose $\hat{\alpha}: \Delta^1 \otimes \Delta^1 \rightarrow \mathcal{C}$ corresponds to a commutative square (ie both its nondegenerate triangles are thin) with the side $\hat{\alpha}(\{1\} \times \Delta^1)$ being an equivalence in \mathcal{C} . Pictorially, $\hat{\alpha}$ looks like*

$$\begin{array}{ccc} a & \xrightarrow{\quad} & x \\ g \downarrow & \searrow \simeq \nearrow & \downarrow \simeq \\ & k & \\ b & \xrightarrow{\quad} & y \\ & \swarrow \simeq \nwarrow & \end{array}$$

Proof Consider a 1-simplex $\alpha: \Delta^1 \rightarrow \text{Fun}^{\text{gr}}(\Delta^1, \mathcal{C})$, as described above. Given $n \geq 2$ and a solid square of the form

$$\begin{array}{ccc} & \Delta^{\{n-1, n\}} & \\ & \swarrow \quad \searrow \alpha & \\ (\Lambda_n^n, \Lambda_n^n \cap \Delta^{\{0, n-1, n\}}) & \xrightarrow{\quad} & \text{Fun}^{\text{gr}}(\Delta^1, \mathcal{C}) \\ \downarrow & \searrow \gamma & \downarrow d \\ (\Delta^n, \Delta^{\{0, n-1, n\}}) & \xrightarrow{\quad} & \mathcal{C} \end{array}$$

we have to exhibit a filler as indicated by the dotted arrow. Note that this lifting problem corresponds to one of the form

$$(7) \quad \begin{array}{ccc} \left(\Delta^1 \times \Lambda_n^n \coprod_{\Delta^{\{0\}} \times \Lambda_n^n} \Delta^{\{0\}} \times \Delta^n, T' \right) & \xrightarrow{f} & \mathcal{C} \\ \downarrow & \searrow & \\ (\Delta^1 \times \Delta^n, T) & & \end{array}$$

in which $\Delta^1 \times \Delta^{\{n-1, n\}}$ is mapped to $\hat{\alpha}$ under the adjunction $\Delta^1 \otimes - \dashv \text{Fun}^{\text{gr}}(\Delta^1, -)$. Here T is the union of the triangles which are thin in $\Delta_b^1 \otimes (\Delta^n, \{\Delta^{\{0, n-1, n\}}\})$, together with $\Delta^{\{(0, n-1), (0, n), (1, n)\}}$, and $T' \subseteq T$ is the subset of those triangles which are contained in the domain of the vertical arrow in (7). Since the edge $\Delta^{\{1\}} \times \Delta^{\{n-1, n\}}$ maps to an invertible edge of \mathcal{C} by our assumption on α and T contains $\Delta^{\{1\}} \times \Delta^{\{0, n-1, n\}}$ we can extend the map f in (7) to a map

$$f': \left([\Delta^1 \times \Lambda_n^n] \coprod_{\partial \Delta^1 \times \Lambda_n^n} [\partial \Delta^1 \times \Delta^n], T'' \right) \rightarrow \mathcal{C},$$

where T'' is the intersection of T with the triangles in $[\Delta^1 \times \Lambda_n^n] \coprod_{\partial \Delta^1 \times \Lambda_n^n} [\partial \Delta^1 \times \Delta^n]$. Let $S = T'' \cup \{\Delta^{\{(0, n-1), (0, n), (1, n)\}}, \Delta^{\{(0, 0), (1, n-1), (1, n)\}}\}$. Then S is contained in T and by Lemma 3.0.2 we may extend f' to $(\Delta^1 \times \Delta^n, S)$. When $n \geq 3$ we have that $T = T'' = S$ and so the proof is complete. In the case $n = 2$, one needs to additionally verify that the resulting extension sends $\Delta^{\{(0, 0), (1, 0), (1, 1)\}}$ to a thin triangle. Indeed, this follows from [8, Proposition 3.4] since S contains $\Delta^{\{(0, 0), (1, 1), (1, 2)\}}, \Delta^{\{(0, 0), (1, 0), (1, 2)\}}$ and $\Delta^{\{(1, 0), (1, 1), (1, 2)\}}$, and $\Delta^{\{(1, 1), (1, 2)\}}$ maps to an invertible edge in \mathcal{C} by assumption. \square

Lemma 3.0.5 *The domain projection*

$$d: \text{Fun}^{\text{gr}}(\Delta^1, \mathcal{C}) \rightarrow \mathcal{C}$$

of an ∞ -bicategory \mathcal{C} has enough cartesian edges.

Proof A lifting problem of the form

$$\begin{array}{ccc} \Delta^{\{1\}} & \xrightarrow{h} & \text{Fun}^{\text{gr}}(\Delta^1, \mathcal{C}) \\ \downarrow & & \downarrow d \\ \Delta^1 & \xrightarrow{f} & \mathcal{C} \end{array}$$

corresponds to the following data in \mathcal{C} :

$$\begin{array}{ccc} x & & \\ f \downarrow & & \\ y & \xrightarrow{h} & z \end{array}$$

This can be extended to a 1-simplex in $\text{Fun}^{\text{gr}}(\Delta^1, \mathcal{C})$, depicted as

$$\begin{array}{ccc} x & \xrightarrow{hf} & z \\ f \downarrow & \searrow hf & \parallel \\ y & \xrightarrow{h} & z \end{array}$$

where the bottom triangle is given by an extension along $\Lambda_1^2 \rightarrow \Delta_{\#}^2$ and the upper triangle is degenerate. Such a 1-simplex is then d-cartesian by Lemma 3.0.4. □

Lemma 3.0.6 *Let $\alpha: \Delta^2 \rightarrow \text{Fun}^{\text{gr}}(\Delta^1, \mathcal{C})$ be a triangle with transpose $\hat{\alpha}: \Delta^1 \otimes \Delta^2 \rightarrow \mathcal{C}$.*

- (i) *If $\hat{\alpha}|_{\Delta^{\{1\}} \times \Delta^2}$ is thin and $\alpha|_{\Delta^{\{1,2\}}}$ satisfies the assumption of Lemma 3.0.4 (so that it is d-cartesian by that lemma), then α is right d-outer.*
- (ii) *If $\hat{\alpha}|_{\Delta^{\{1\}} \times \Delta^2}$ is thin and $\alpha|_{\Delta^{\{0,1\}}}$ is invertible, then α is left d-outer.*

Proof We first prove (i). By Remark 2.1.2 we need to consider a lifting problem of the form

$$\begin{array}{ccc} & \Delta_b^{\{0,n-1,n\}} & \\ & \swarrow & \searrow \alpha \\ (\Lambda_n^n)_b & \xrightarrow{\quad} & \text{Fun}^{\text{gr}}(\Delta^1, \mathcal{C}) \\ \downarrow & \nearrow \text{dotted} & \downarrow d \\ \Delta_b^n & \xrightarrow{\quad} & \mathcal{C} \end{array}$$

for $n \geq 3$, where we have to exhibit a diagonal filler as indicated by the dotted arrow.

We obtain an equivalent lifting problem of the form

$$\begin{array}{ccc} \left(\Delta^1 \times \Lambda_n^n \coprod_{\Delta^{\{0\}} \times \Lambda_n^n} \Delta^{\{0\}} \times \Delta^n, T \right) & \xrightarrow{f} & \mathcal{C} \\ \downarrow & \nearrow & \\ (\Delta^1 \times \Delta^n, T) & & \end{array}$$

where T is the set of thin triangles in $\Delta_b^1 \otimes \Delta_b^n$ together with $\Delta^{\{(0,n-1),(0,n),(1,n)\}}$ and $\Delta^{\{1\}} \times \Delta^{\{0,n-1,n\}}$ (all of whom are contained in the top left corner, since $n \geq 3$). Our assumption on α implies that the edge $\Delta^{\{1\}} \times \Delta^{\{n-1,n\}}$ is invertible in \mathcal{C} and so by [6, Corollary 2.3.10] we can extend the map f to a map

$$f': \left(\Delta^1 \times \Lambda_n^n \coprod_{\partial \Delta^1 \times \Lambda_n^n} \partial \Delta^1 \times \Delta^n, T \right) \rightarrow \mathcal{C}.$$

Now since T contains $\Delta^{\{(0,0),(1,0),(1,n-1)\}}$, $\Delta^{\{(0,0),(1,0),(1,n)\}}$ and $\Delta^{\{1\}} \times \Delta^{\{0,n-1,n\}}$ and \mathcal{C} is an ∞ -bicategory, the map f' must also send $\Delta^{\{(0,0),(1,n-1),(1,n)\}}$ to a thin triangle; see [13, Remark 3.1.4]. We may consequently apply Lemma 3.0.2 in order to extend f' to all of $(\Delta^1 \times \Delta^n, T)$, as desired.

We now prove (ii). By Remark 2.1.2 we now need to consider a lifting problem of the form

$$\begin{array}{ccc} & \Delta_b^{\{0,1,n\}} & \\ & \swarrow \alpha & \searrow \\ (\Lambda_0^n)_b & \xrightarrow{\quad} & \text{Fun}^{\text{gr}}(\Delta^1, \mathcal{C}) \\ \downarrow & \nearrow & \downarrow d \\ \Delta_b^n & \xrightarrow{\quad} & \mathcal{C} \end{array}$$

with $n \geq 3$. We obtain an equivalent lifting problem of the form

$$\begin{array}{ccc} \left(\Delta^1 \times \Lambda_0^n \coprod_{\Delta^{\{0\}} \times \Lambda_0^n} \Delta^{\{0\}} \times \Delta^n, T \right) & \xrightarrow{f} & \mathcal{C} \\ \downarrow & \nearrow & \\ (\Delta^1 \times \Delta^n, T) & & \end{array}$$

where T is the set of thin triangles in $\Delta_b^1 \otimes \Delta_b^n$ together with $\Delta^{\{(0,0),(0,1),(1,1)\}}$ and $\Delta^{\{1\}} \times \Delta^{\{0,1,n\}}$. Since the image of the edge $\Delta^{\{1\}} \times \Delta^{\{0,1\}}$ is invertible in \mathcal{C} we can extend the map f to a map

$$f': \left([\Delta^1 \times \Lambda_0^n] \coprod_{\partial \Delta^1 \times \Lambda_0^n} [\partial \Delta^1 \times \Delta^n], T \right) \rightarrow \mathcal{C}.$$

Now since T contains $\Delta^{\{(0,0),(1,0),(1,1)\}}$, $\Delta^{\{(0,0),(1,0),(1,n)\}}$ and $\Delta^{\{1\}} \times \Delta^{\{0,1,n\}}$ and \mathcal{C} is an ∞ -bicategory, the map f' must also send $\Delta^{\{(0,0),(1,1),(1,n)\}}$ to a thin triangle. Since f' also sends $\Delta^{\{(0,0),(0,1),(1,1)\}}$ and $\Delta^{\{(0,1),(1,1),(1,n)\}}$ to thin triangles, the same holds for $\Delta^{\{(0,0),(0,1),(1,n)\}}$. We may consequently apply the dual form of Lemma 3.0.2 (see Remark 3.0.3), in order to extend f' to all of $(\Delta^1 \times \Delta^n, T)$, as desired. \square

Finally, we are ready to prove the main result of this section.

Theorem 3.0.7 *Given an ∞ -bicategory \mathcal{C} , the domain projection*

$$d: \text{Fun}^{\text{gr}}(\Delta^1, \mathcal{C}) \rightarrow \mathcal{C}$$

is a 2-outer cartesian fibration, whose p -cartesian 1-simplices are those described in Lemma 3.0.4. In addition, the right p -outer triangles whose right legs are cartesian are those described in Lemma 3.0.6(1), and the left p -outer triangles whose left leg is invertible are those described in Lemma 3.0.6(2).

Proof By Remark 3.0.1, the map d is a weak fibration and by Lemma 3.0.4 it has a sufficient supply of d -cartesian lifts for 1-morphisms. By the essential uniqueness of d -cartesian lifts we deduce that all d -cartesian arrows are of the form described in Lemma 3.0.4.

We now show that the triangles in \mathcal{C} have a sufficient supply of right p -outer lifts. This translates into a lifting problem of the form

$$\begin{array}{ccc} \Lambda_2^2 & \xrightarrow{\rho} & \text{Fun}^{\text{gr}}(\Delta^1, \mathcal{C}) \\ \downarrow & \nearrow \gamma & \downarrow d \\ \Delta^2 & \xrightarrow{\gamma} & \mathcal{C} \end{array}$$

with $\rho|_{\Delta^{\{1,2\}}}$ a d -cartesian edge, and so (as argued just above) of the form described in Lemma 3.0.4. This means in particular that $\rho|_{\Delta^{\{1,2\}}}$ corresponds to a map $\Delta_b^1 \otimes \Delta_b^{\{1,2\}} \rightarrow \mathcal{C}$ which sends both triangles to thin triangles and the edge $\Delta^{\{1\}} \times \Delta^{\{1,2\}}$ to an invertible edge in \mathcal{C} . Solving this lifting problem in a way that produces a triangle of $\text{Fun}^{\text{gr}}(\Delta^1, \mathcal{C})$ of the form described in Lemma 3.0.6(1) then corresponds to solving a lifting problem of the form

$$(8) \quad \begin{array}{ccc} (\Delta^1 \times \Lambda_2^2 \coprod_{\Delta^{\{0\}} \times \Lambda_2^2} \Delta^{\{0\}} \times \Delta^2, T') & \xrightarrow{\widehat{\rho} \cup \gamma} & \mathcal{C} \\ \downarrow & \nearrow & \\ (\Delta^1 \times \Delta^2, T) & & \end{array}$$

where T' is obtained by intersection from T , which in turn contains all triangles which are thin in $\Delta_b^1 \otimes \Delta_b^2$ as well as $\Delta^{\{1\}} \times \Delta^2$ and $\Delta^{\{(0,1),(0,2),(1,2)\}}$, and $\widehat{\rho}$ sends $\Delta^{\{1\}} \times \Delta^{\{1,2\}}$ to an invertible edge in \mathcal{C} . Since \mathcal{C} is an ∞ -bicategory we may then extend $\widehat{\rho} \cup \gamma$ to a map

$$g: \left(\Delta^1 \times \Lambda_2^2 \coprod_{\partial \Delta^1 \times \Lambda_2^2} \partial \Delta^1 \times \Delta^2, T'' \right) \rightarrow \mathcal{C},$$

where T'' is obtained from T by intersection. Applying Lemma 3.0.2 we may extend g to a map $g': (\Delta^1 \times \Delta^2, T'' \cup \{\Delta^{\{(0,0),(1,1),(1,2)\}}\}) \rightarrow \mathcal{C}$. We then observe that there is exactly one triangle in T which is not in $T'' \cup \{\Delta^{\{(0,0),(1,1),(1,2)\}}\}$, and that is the triangle $\Delta^{\{(0,0),(1,0),(1,1)\}}$. But this triangle is sent by g to a thin triangle in \mathcal{C} by [8, Proposition 3.4], since T contains $\Delta^{\{(0,0),(1,1),(1,2)\}}$, $\Delta^{\{(0,0),(1,0),(1,2)\}}$ and $\Delta^{\{(1,0),(1,1),(1,2)\}}$, and g sends $\Delta^{\{1\}} \times \Delta^{\{1,2\}}$ to an invertible edge.

We now show that the triangles in \mathcal{C} have a sufficient supply of left p -outer lifts. Invoking Corollary 2.3.8, it will suffice to test this only for triangles in \mathcal{B} whose left leg is degenerate. This then translates into a lifting problem of the form

$$\begin{array}{ccc} \Lambda_0^2 & \xrightarrow{\rho} & \text{Fun}^{\text{gr}}(\Delta^1, \mathcal{C}) \\ \downarrow & \nearrow \gamma & \downarrow d \\ \Delta^2 & \xrightarrow{\gamma} & \mathcal{C} \end{array}$$

with $\rho|_{\Delta^{\{0,1\}}}$ a d -cocartesian lift of an identity, and hence an invertible edge of $\text{Fun}^{\text{gr}}(\Delta^1, \mathcal{C})$. This means in particular that $\rho|_{\Delta^{\{0,1\}}}$ corresponds to a map $\Delta_b^1 \otimes \Delta_b^{\{0,1\}} \rightarrow \mathcal{C}$ which sends both triangles to thin triangles and the edges $\Delta^{\{0\}} \times \Delta^{\{0,1\}}$ and $\Delta^{\{1\}} \times \Delta^{\{0,1\}}$ to invertible edges in \mathcal{C} . Solving this lifting problem in a way that produces a triangle of $\text{Fun}^{\text{gr}}(\Delta^1, \mathcal{C})$ of the form described in Lemma 3.0.6(2) then corresponds to solving a lifting problem of the form

$$(9) \quad \begin{array}{ccc} \left(\Delta^1 \times \Lambda_0^2 \coprod_{\Delta^{\{0\}} \times \Lambda_0^2} \Delta^{\{0\}} \times \Delta^2, T' \right) & \xrightarrow{\widehat{\rho} \cup \gamma} & \mathcal{C} \\ \downarrow & \nearrow & \\ (\Delta^1 \times \Delta^2, T) & & \end{array}$$

where T' is obtained by intersection from T , which in turn contains all triangles which are thin in $\Delta_b^1 \otimes \Delta_b^2$ as well as the triangles $\Delta^{\{1\}} \times \Delta^2$ and $\Delta^{\{(0,0),(0,1),(1,1)\}}$, and $\widehat{\rho}$ sends $\Delta^{\{\varepsilon\}} \times \Delta^{\{0,1\}}$ to an invertible edge in \mathcal{C} for $\varepsilon = 0, 1$. Since \mathcal{C} is an ∞ -bicategory we may then extend $\widehat{\rho} \cup \gamma$ to a map

$$g: \left(\Delta^1 \times \Lambda_0^2 \coprod_{\partial \Delta^1 \times \Lambda_0^2} \partial \Delta^1 \times \Delta^2, T'' \right) \rightarrow \mathcal{C},$$

where T'' is obtained from T by intersection. Applying the dual of Lemma 3.0.2 (Remark 3.0.3) we may extend g to a map $g': (\Delta^1 \times \Delta^2, T'' \cup \{\Delta^{\{(0,0),(0,1),(1,2)\}}\}) \rightarrow \mathcal{C}$. We then observe that there are exactly two triangles in T which are not in T'' , namely, $\Delta^{\{(0,0),(1,1),(1,2)\}}$ and $\Delta^{\{(0,1),(1,1),(1,2)\}}$. To finish the proof we will show that these two triangles are sent by g to thin triangles in \mathcal{C} . For the first one, we note that since g sends $\Delta^{\{(0,0),(1,0),(1,1)\}}$, $\Delta^{\{(0,0),(1,0),(1,2)\}}$ and $\Delta^{\{1\}} \times \Delta^2$ to thin triangles, then it must also send $\Delta^{\{(0,0),(1,1),(1,2)\}}$ to a thin triangle. Then, since g also sends $\Delta^{\{(0,0),(0,1),(1,1)\}}$ and $\Delta^{\{(0,0),(0,1),(1,2)\}}$ to thin triangles, and the edge $\Delta^{\{(0,0),(0,1)\}}$ to an equivalence, then it must send $\Delta^{\{(0,1),(1,1),(1,2)\}}$ to a thin triangle as well; see [8, Proposition 3.4].

Having provided sufficiently many left and right d -outer lifts of the form appearing in Lemma 3.0.6, the uniqueness of d -outer lifts, as expressed for example in Remark 2.1.7, shows that all right d -outer triangles whose right leg is cartesian are of the form described in 3.0.6(1), and all left d -outer triangles whose left leg is invertible are of the form described in 3.0.6(2). To show that the collection of right (resp. left) d -outer triangles is closed under right (resp. left) whiskering, it will hence suffice to show that the collection of triangles $\alpha: \Delta^2 \rightarrow \text{Fun}^{\text{gr}}(\Delta^1, \mathcal{C})$ whose adjoint $\widehat{\alpha}: \Delta^1 \otimes \Delta^2 \rightarrow \mathcal{C}$ sends $\Delta^{\{1\}} \times \Delta^2$

to a thin triangle, is closed under both left and right whiskering. Indeed, this is exactly the preimage in $\text{Fun}^{\text{gr}}(\Delta^1, \mathcal{C})$ under the *codomain projection* of the collection of thin triangles in \mathcal{C} , and the collection of thin triangles is closed under whiskering from both sides. \square

Remark 3.0.8 Passing to opposites, Theorem 3.0.7 implies that the codomain projection

$$\text{cod}: \text{Fun}^{\text{opgr}}(\Delta^1, \mathcal{C}) \rightarrow \mathcal{C}$$

is 2–outer cocartesian fibration, whose cocartesian arrows and outer triangles admit similar descriptions. Note that we have switched not only from domain to codomain but also from Fun^{gr} to Fun^{opgr} . If we only switch from domain to codomain then the resulting projection

$$\text{cod}: \text{Fun}^{\text{gr}}(\Delta^1, \mathcal{C}) \rightarrow \mathcal{C}$$

is a 2–inner cocartesian fibration. This claim does not formally follow from the outer statement, but its proof can be obtained using a completely analogous argument, replacing the key extension result of Lemma 3.0.2 with one involving the pushout product of $\partial\Delta^1 \rightarrow \Delta^1$ and a suitable *inner* horn inclusion. Similarly, the projection

$$\text{d}: \text{Fun}^{\text{opgr}}(\Delta^1, \mathcal{C}) \rightarrow \mathcal{C}$$

is a 2–inner cartesian fibration.

4 Enriched cartesian fibrations

In this section we define the notion of cartesian fibration in the context of marked simplicial categories. The definition is motivated by the one given for 2–categories by Buckley in [5], which we recall in Section 4.1.

4.1 Recollection: fibrations of 2–categories

Fibrations of 2–categories were initially introduced by Hermida [11]. A suitably modified definition was later given by Buckley [5], who also proved an (un)straightening-type result. In what follows we give a concise summary of the main results of loc. cit., to be considered as a motivation for the discussion of simplicial categories in Section 4.2.

Definition 4.1.1 Let $p: \mathcal{E} \rightarrow \mathcal{B}$ be a 2–functor between 2–categories.

- A 1–cell $f: x \rightarrow y$ in \mathcal{E} is *p–cartesian* if the following square is a pullback of categories for every $a \in \mathcal{E}$:

$$\begin{array}{ccc} \mathcal{E}(a, x) & \xrightarrow{f \circ -} & \mathcal{E}(a, y) \\ p_{a,x} \downarrow & & \downarrow p_{a,y} \\ \mathcal{B}(pa, px) & \xrightarrow{p(f) \circ -} & \mathcal{B}(pa, py) \end{array}$$

- A 2–cell $\alpha: f \Rightarrow g: x \rightarrow y$ in \mathcal{E} is *p–cartesian* if it is cartesian with respect to the induced functor $p_{x,y}: \mathcal{E}(x, y) \rightarrow \mathcal{B}(px, py)$.

The notion of cartesian fibration for 2-categories amounts to the existence of enough cartesian lifts, as in the 1-dimensional case, but it also requires an additional property: cartesian 2-cells must be closed under horizontal composition.

Definition 4.1.2 A 2-functor between 2-categories $p: \mathcal{E} \rightarrow \mathcal{B}$ is called a 2-fibration if it satisfies the following properties:

- (i) For every object $e \in \mathcal{E}$ and every 1-cell $f: b \rightarrow p(e)$ there exists a p -cartesian 1-cell $h: a \rightarrow e$ in \mathcal{E} such that $p(h) = f$.
- (ii) For every pair of objects x, y in \mathcal{E} , the map $p_{x,y}: \mathcal{E}(x, y) \rightarrow \mathcal{B}(px, py)$ is a cartesian fibration of categories.
- (iii) Cartesian 2-cells are closed under horizontal composition, ie for every triple of objects (x, y, z) in \mathcal{E} , the functor $\circ_{x,y,z}: \mathcal{E}(y, z) \times \mathcal{E}(x, y) \rightarrow \mathcal{E}(x, z)$ sends $p_{y,z} \times p_{x,y}$ -cartesian 1-cells to $p_{x,z}$ -cartesian ones.

Replacing cartesian lifts for 1-cells by cocartesian lifts and similarly for 2-cells one may obtain four different variants of fibration, corresponding to the four possible types of variance for pseudofunctors $\mathcal{B} \rightarrow 2\text{-Cat}$.

Remark 4.1.3 Condition (iii) of Definition 4.1.2 is equivalent to requiring that given 1-cells in \mathcal{E} of the form $f: w \rightarrow x$ and $g: y \rightarrow z$, the whiskering functors

$$- \circ f: \mathcal{E}(x, y) \rightarrow \mathcal{E}(w, y) \quad \text{and} \quad g \circ -: \mathcal{E}(x, y) \rightarrow \mathcal{E}(x, z)$$

preserve cartesian 2-cells. This follows from the fact that horizontal composition can be obtained from vertical composition and whiskering composition.

The following result appears as Theorem 2.2.11 in [5].

Theorem 4.1.4 *There exists an equivalence of 3-categories between $2\text{Fib}_s(\mathcal{B})$ and $[\mathcal{B}_{\text{co}}^{\text{op}}, 2\text{-Cat}]$, the former being the 3-category of fibrations equipped with a choice of cartesian lifts compatible with composition, while the latter is the 3-category of (strict) 2-functors into 2-Cat , natural transformations and modifications.*

In the same paper, the author proves several weakenings of this statement, by looking at fibrations without a choice of lifts (which correspond to *pseudofunctors*) and fibrations of bicategories.

4.2 Cartesian fibrations of enriched categories

We now consider cartesian fibrations in the setting of Cat_∞ -categories. All definitions and statements can be dualized to the case of cocartesian fibrations by replacing all Cat_∞ -categories by their opposites.

Definition 4.2.1 Let $p: \mathcal{E} \rightarrow \mathcal{B}$ be a map of marked simplicial categories. A morphism $h \in \mathcal{E}(e', e)_0$ is said to be p -cartesian if the induced square

$$\begin{array}{ccc} \mathcal{E}(a, e') & \xrightarrow{h \circ -} & \mathcal{E}(a, e) \\ p_{a, e'} \downarrow & & \downarrow p_{a, e} \\ \mathcal{B}(pa, pe') & \xrightarrow{p(h) \circ -} & \mathcal{B}(pa, pe) \end{array}$$

is a homotopy pullback for the model structure on marked simplicial sets.

Definition 4.2.2 Let $p: \mathcal{E} \rightarrow \mathcal{B}$ be a fibration of $\mathcal{C}at_\infty$ -categories. We say that p is an *enriched cartesian fibration* if for any $e \in \mathcal{E}$ and any morphism $f \in \mathcal{B}(a, p(e))_0$ there exists a p -cartesian morphism $h \in \mathcal{E}(e', e)_0$, for some $e' \in \mathcal{E}$, such that $p(h) = f$.

Definition 4.2.3 Let $p: \mathcal{E} \rightarrow \mathcal{B}$ be a fibration of $\mathcal{C}at_\infty$ -categories. We say that p is an *enriched 2-inner (resp. 2-outer) fibration* if it satisfies the following properties:

- (i) For every pair of objects (x, y) in \mathcal{E} , the map $p_{x, y}: \mathcal{E}(x, y) \rightarrow \mathcal{B}(px, py)$ is a cartesian (resp. co-cartesian) fibration on the level of underlying simplicial sets.
- (ii) For every pair of 0-simplices $u: y \rightarrow z$ and $v: w \rightarrow x$ in \mathcal{E} , the commutative squares

$$\begin{array}{ccc} \mathcal{E}(x, y) & \xrightarrow{u \circ -} & \mathcal{E}(x, z) & & \mathcal{E}(x, y) & \xrightarrow{- \circ v} & \mathcal{E}(w, y) \\ p_{x, y} \downarrow & & \downarrow p_{x, z} & & \downarrow p_{x, y} & & \downarrow p_{w, y} \\ \mathcal{B}(px, py) & \xrightarrow{p(u) \circ -} & \mathcal{B}(px, pz) & & \mathcal{B}(px, py) & \xrightarrow{- \circ p(v)} & \mathcal{B}(pw, py) \end{array}$$

are morphisms of cartesian (resp. cocartesian) fibrations, ie the top horizontal maps in both squares preserve cartesian (resp. cocartesian) edges.

We will say that p is an *enriched 2-inner (resp. 2-outer) cartesian fibration* if it is an enriched 2-inner (resp. 2-outer) fibration and an enriched cartesian fibration.

Our goal in the present section is to prove the following:

Theorem 4.2.4 Let $p: \mathcal{E} \rightarrow \mathcal{B}$ be a fibration of $\mathcal{C}at_\infty$ -categories. Then $p: \mathcal{E} \rightarrow \mathcal{B}$ is an enriched 2-inner (resp. 2-outer) cartesian fibration in the sense of Definition 4.2.2 if and only if

$$N^{sc}(p): N^{sc}\mathcal{E} \rightarrow N^{sc}\mathcal{B}$$

is a 2-inner (resp. 2-outer) cartesian fibration of ∞ -bicategories.

The remainder of this section is devoted to the proof of Theorem 4.2.4.

Definition 4.2.5 We will denote by $\square^n = (\Delta^1)^n$ the *simplicial n -cube* and by $\partial\square^n$ its *boundary*, so that the inclusion $\partial\square^n \subseteq \square^n$ can be identified with the pushout-product of $\partial\Delta^1 \hookrightarrow \Delta^1$ with itself n times. For $i = 1, \dots, n$ we denote by $\prod_{\varepsilon}^{n,i} \hookrightarrow \square^n$ the iterated pushout-product

$$[\partial\Delta^1 \hookrightarrow \Delta^1] \square \dots \square [\Delta^{\{\varepsilon\}} \hookrightarrow \Delta^1] \square \dots \square [\partial\Delta^1 \hookrightarrow \Delta^1],$$

where $\varepsilon \in \{0, 1\}$ and $[\Delta^{\{\varepsilon\}} \hookrightarrow \Delta^1]$ appears in the i^{th} factor.

Lurie introduces in [12, Definition 3.1.1.1] the class of *cartesian anodyne* morphisms (called *marked anodyne* in loc. cit.), which is the smallest weakly saturated class generated by a certain set of monomorphisms of marked simplicial sets listed in [12, Definition 3.1.1.1], which contain, in particular, the collection of all (minimally marked) inner horn inclusions, as well as the marked outer horn inclusion $(\Lambda_n^n, \{\Delta^{\{n-1, n\}}\}) \hookrightarrow (\Delta^n, \{\Delta^{\{n-1, n\}}\})$. Dually, we shall call *cocartesian anodyne* maps the smallest weakly saturated class generated by the opposites of those maps (or simply those maps whose opposites are cartesian anodyne). By [12, Proposition 3.1.2.3], (co)cartesian anodyne maps are closed under pushout-product with monomorphisms.

Lemma 4.2.6 For $\varepsilon \in \{0, 1\}$, $n \geq 1$ and $1 \leq i \leq n$ let $E_{\varepsilon}^i \subseteq (\square^n)_1$ be the set of all degenerate edges together with the edge $(\varepsilon, \dots, \varepsilon) \times \Delta^1 \times (\varepsilon, \dots, \varepsilon)$, where Δ^1 sits in the i^{th} place. Then the inclusion of marked simplicial sets $(\prod_1^{n,i}, E_1^i) \hookrightarrow (\square^n, E_1^i)$ is cartesian anodyne, and the inclusion of marked simplicial sets $(\prod_0^{n,i}, E_0^i) \hookrightarrow (\square^n, E_0^i)$ is cocartesian anodyne.

Proof We note that the $\varepsilon = 0$ and $\varepsilon = 1$ statements imply each other by passing to opposites. We hence just prove the cartesian case. Ignoring the order of factors, we may identify the map $\prod_1^{n,i} \hookrightarrow \square^n$ with the box product of $\partial\square^{n-1} \hookrightarrow \square^{n-1}$ and $\Delta^{\{1\}} \hookrightarrow \Delta^1$. The nondegenerate m -simplices of \square^{n-1} which are not in $\partial\square^{n-1}$ correspond to maps of posets $[m] \rightarrow [1]^{n-1}$ whose projection to each factor $[1]$ is surjective. It then follows that the initial vertex of such an m -simplex must be $(0, \dots, 0)$, and the final vertex must be $(1, \dots, 1)$. Adding these nondegenerate simplices one by one in an order that respects dimensions (that is, first all the 1-dimensional ones, then all the 2-dimensional ones, etc, up to dimension $n - 1$) results in a factorization of the map $(\prod_1^{n,i}, E_1^i) \hookrightarrow (\square^n, E_1^i)$ into a finite composite of pushouts of maps of the form

$$\left(\Delta^{\{1\}} \times \Delta^m \prod_{\Delta^{\{1\}} \times \partial\Delta^m} \Delta^1 \times \partial\Delta^m, \{\Delta^1 \times \Delta^{\{m\}}\} \right) \rightarrow (\Delta^1 \times \Delta^m, \{\Delta^1 \times \Delta^{\{m\}}\})$$

for $m \geq 1$. It will hence suffice to show that each of these maps is cartesian anodyne. For $\ell = 0, \dots, m$, let

$$\tau_{\ell}: \Delta^{m+1} \rightarrow \Delta^1 \times \Delta^m$$

be the map given on vertices by the formula

$$\tau_{\ell}(j) = \begin{cases} (0, j) & \text{if } j \leq \ell, \\ (1, j - 1) & \text{if } j > \ell, \end{cases}$$

and for $k = 0, \dots, m + 1$ let $Z_k \subseteq (\Delta^1 \times \Delta^m, \{\Delta^1 \times \Delta^{\{m\}}\})$ be the marked simplicial subset obtained as the union of $[\Delta^1 \times \partial\Delta^m] \coprod_{\Delta^{\{1\}} \times \partial\Delta^m} [\Delta^{\{1\}} \times \Delta^m]$ and the simplices τ_ℓ for $0 \leq \ell < k$. Set

$$Z_0 \stackrel{\text{def}}{=} \left([\Delta^1 \times \partial\Delta^m] \coprod_{\Delta^{\{1\}} \times \partial\Delta^m} [\Delta^{\{1\}} \times \Delta^m], \{\Delta^1 \times \Delta^{\{m\}}\} \right).$$

We then have an ascending filtration of marked simplicial sets

$$Z_0 \subseteq Z_1 \subseteq \dots \subseteq Z_{m+1} = (\Delta^1 \times \Delta^m, \{\Delta^1 \times \Delta^{\{m\}}\}).$$

For each $k = 0, \dots, m - 1$ we then find a pushout square of marked simplicial sets

$$\begin{array}{ccc} (\Lambda_{k+1}^{m+1})^b & \longrightarrow & Z_k \\ \downarrow & & \downarrow \\ (\Delta^{m+1})^b & \longrightarrow & Z_{k+1} \end{array}$$

so that $Z_k \rightarrow Z_{k+1}$ is inner anodyne, and in particular cartesian anodyne. Finally, in the last step $k = m$ we find a pushout square of the form

$$\begin{array}{ccc} (\Lambda_{m+1}^{m+1}, \{\Delta^{\{m,m+1\}}\}) & \longrightarrow & Z_m \\ \downarrow & & \downarrow \\ (\Delta^{m+1}, \{\Delta^{\{m,m+1\}}\}) & \longrightarrow & Z_{m+1} \end{array}$$

so that $Z_m \rightarrow Z_{m+1}$ is cartesian anodyne, as desired. □

We recall the comparison of the notion of (co)cartesian 1-cells between the enriched and scaled models:

Proposition 4.2.7 [6, Proposition 3.1.3] *Given a fibration $p: \mathcal{E} \rightarrow \mathcal{B}$ of Cat_∞ -categories, an arrow in \mathcal{E} is p -cartesian if and only if the corresponding 1-simplex of $\text{N}^{\text{sc}}\mathcal{E}$ is $\text{N}^{\text{sc}}(p)$ -cartesian.*

Since the morphisms in a given Cat_∞ -category are in bijection with the edges of its scaled coherent nerve we readily obtain:

Corollary 4.2.8 *A fibration $p: \mathcal{E} \rightarrow \mathcal{B}$ of Cat_∞ -categories is cartesian if and only if $\text{N}^{\text{sc}}(p)$ is a cartesian fibration of ∞ -bicategories (in the sense of Definition 1.4.9).*

We now consider the analogous question on the level of triangles.

Remark 4.2.9 Let \mathcal{E} be a Cat_∞ -category and consider two 2-simplices α and α' of $\text{N}^{\text{sc}}\mathcal{E}$. If a 3-simplex ρ of $\text{N}^{\text{sc}}\mathcal{E}$ exhibits α as left-congruent to α' (cf Definition 2.3.3) in the form

$$\begin{array}{ccc} \begin{array}{ccc} x & \xrightarrow{f} & z \\ \parallel & \searrow g & \uparrow h \\ & & y \end{array} & \xrightarrow{\quad} & \begin{array}{ccc} x & \xrightarrow{f} & z \\ \parallel & \swarrow \alpha' & \uparrow h \\ & i & \searrow \approx \\ x & \xrightarrow{g} & y \end{array} \end{array}$$

then the 2-simplex ρ^* of $\mathcal{E}(x, z)$ corresponding to ρ goes from the 1-simplex α^* of $\mathcal{E}(x, z)$ corresponding to α to the composition of the 1-simplex $(\alpha')^*$ corresponding to α' followed by an equivalence. Hence, in the arrow ∞ -category of $\mathcal{E}(x, z)$ the 3-simplex ρ induces an equivalence between α^* and the composition of $(\alpha')^*$ with an arrow of $\mathcal{E}(x, z)$ which is an equivalence.

Lemma 4.2.10 *Let $p: \mathcal{E} \rightarrow \mathcal{B}$ be a fibration of Cat_∞ -categories. Then a triangle $\alpha: \Delta^2 \rightarrow \text{N}^{\text{sc}}\mathcal{E}$ of the form*

$$\begin{array}{ccc} x & \xrightarrow{h} & z \\ & \searrow f & \downarrow \alpha \\ & & y \\ & & \nearrow g \end{array}$$

is left $\text{N}^{\text{sc}}(p)$ -inner if and only if the corresponding 1-simplex $\alpha^*: h \rightarrow gf$ in $\mathcal{E}(x, z)$ is $p_{x,z}$ -cartesian and maps to a $p_{x',z}$ -cartesian arrow in $\mathcal{E}(x', z)$ after precomposing with any arrow $x' \rightarrow x$. Similarly, α is right $\text{N}^{\text{sc}}(p)$ -inner if and only if the corresponding 1-simplex $\alpha^*: h \rightarrow gf$ in $\mathcal{E}(x, z)$ is $p_{x,z}$ -cartesian and maps to a $p_{x,z'}$ -cartesian arrow in $\mathcal{E}(x, z')$ after postcomposing with any arrow $z \rightarrow z'$.

Proof We prove the left inner case. The proof for right inner triangles is completely analogous. Suppose first that α^* is $p_{x,z}$ -cartesian in $\mathcal{E}(x, z)$ and maps to a $p_{x',z}$ -cartesian arrow in $\mathcal{E}(x', z)$ after precomposing with any arrow $x' \rightarrow x$. By Remark 2.1.2 we have to provide a solution for every lifting problem of the form

$$\begin{array}{ccc} & \Delta^{\{n-2, n-1, n\}} & \\ & \swarrow & \searrow \alpha \\ \Lambda_{n-1}^n & \xrightarrow{\quad} & \text{N}^{\text{sc}}\mathcal{E} \\ \downarrow & & \downarrow \text{N}^{\text{sc}}(p) \\ \Delta^n & \xrightarrow{\quad} & \text{N}^{\text{sc}}\mathcal{B} \end{array}$$

for $n \geq 3$. Transposing along the adjunction $\mathcal{E}^{\text{sc}} \dashv \text{N}^{\text{sc}}$, this corresponds to a lifting problem of the form

$$(10) \quad \begin{array}{ccc} & \mathcal{E}^{\text{sc}} \Delta^{\{n-2, n-1, n\}} & \\ & \swarrow & \searrow \alpha^* \\ \mathcal{E}^{\text{sc}} \Lambda_{n-1}^n & \xrightarrow{f} & \mathcal{E} \\ \downarrow & & \downarrow p \\ \mathcal{E}^{\text{sc}} \Delta^n & \xrightarrow{g} & \mathcal{B} \end{array}$$

where we have committed a small abuse of language denoting by α^* also the transpose map $\mathcal{E}^{\text{sc}} \Delta^2 \rightarrow \mathcal{E}$ induced by α . As a straightforward calculation shows, the lifting problem in (10) corresponds, at the level of marked simplicial sets, to the lifting problem

$$\begin{array}{ccc} (\prod_1^{n-1, n-1})^b & \longrightarrow & \mathcal{E}(x', z) \\ \downarrow & & \downarrow p_{x', z} \\ (\square^{n-1})^b & \longrightarrow & \mathcal{B}(px', pz) \end{array}$$

where $x' := f(0)$. Moreover, the edge $(1, \dots, 1) \times \Delta^1$ is mapped by the top horizontal map to the whiskering of the 1-simplex α^* by a sequence of 0-simplices connecting x' and x , and it is therefore $p_{x',z}$ -cartesian in $\mathcal{E}(x', z)$. The desired lift hence exists by Lemma 4.2.6.

We now prove the “only if” direction, and so we assume that α is left p -inner. By Lemma 2.3.7 we may find a left degenerate triangle α' such that α is left congruent to α' , so that α' is also left p -inner by Lemma 2.3.4. Then α' has the same first and last vertex as α and hence determines an edge $(\alpha')^*$ in $\mathcal{E}(x, z)$. Furthermore, as pointed out in the previous remark the 3-simplex exhibiting α' as left congruent to α also determines an equivalence in the arrow ∞ -category of $\mathcal{E}(x, z)$ between the edge α^* and the edge given by $(\alpha')^*$ followed by an equivalence. Since every equivalence is $p_{x,z}$ -cartesian and moreover the property of being $p_{x,z}$ -cartesian is invariant under equivalences we may replace α by α' , so that we may simply assume that α is left degenerate. We now consider the commutative diagram of marked simplicial sets

$$(11) \quad \begin{array}{ccccc} \mathrm{Hom}_{\mathrm{N}^{\mathrm{sc}}\mathcal{E}}^{\triangleright}(x, z) & \xrightarrow{\cong} & \mathrm{Hom}_{\mathrm{N}^{\mathrm{sc}}\mathcal{E}}(x, z) & \xleftarrow{\cong} & \mathcal{E}(x, z) \\ \downarrow p_{x,y}^{\triangleright} & & \downarrow & & \downarrow p_{x,y} \\ \mathrm{Hom}_{\mathrm{N}^{\mathrm{sc}}\mathcal{B}}^{\triangleright}(x, z) & \xrightarrow{\cong} & \mathrm{Hom}_{\mathrm{N}^{\mathrm{sc}}\mathcal{B}}(x, z) & \xleftarrow{\cong} & \mathcal{B}(x, z) \end{array}$$

in which the horizontal maps are marked categorical equivalences and the vertical maps are fibrations between fibrant objects in the marked categorical model structure. Here, the horizontal equivalences in the right column are given by the canonical isomorphism $\mathrm{Hom}_{\mathrm{N}^{\mathrm{sc}}\mathcal{E}}(x, z) \cong \mathrm{Un}^{\mathrm{sc}} \mathcal{E}(x, z)$, see [13, Remark 4.2.1], and the horizontal equivalences in the left column are established in [8, Proposition 2.33]. The triangle α determines arrows in both the top left and top right marked simplicial sets, and the images of these two arrows coincides in $\mathrm{Hom}_{\mathrm{N}^{\mathrm{sc}}\mathcal{E}}(x, z)$ by direct inspection. We hence obtain that the arrow α^* is $p_{x,z}$ -cartesian in $\mathcal{E}(x, z)$ if and only if the arrow determined by α in $\mathrm{Hom}_{\mathrm{N}^{\mathrm{sc}}\mathcal{E}}^{\triangleright}(x, z)$ is $p_{x,z}^{\triangleright}$ -cartesian. The latter, and hence the former, indeed holds when α is left p -inner by Remark 2.1.8. The desired implication is now a consequence of the closure of left p -inner triangles under left whiskering; see Remark 2.1.5. □

We now come to the outer counterpart of Lemma 4.2.10.

Lemma 4.2.11 *Let $p: \mathcal{E} \rightarrow \mathcal{B}$ be a fibration of Cat_{∞} -categories. Let α^* be a triangle in $\mathrm{N}^{\mathrm{sc}}\mathcal{E}$ of the form*

$$\begin{array}{ccc} x & \xrightarrow{h} & z \\ & \searrow f & \nearrow g \\ & & y \end{array} \quad \Downarrow \alpha^*$$

Then the following holds:

- (i) *If the associated 1-simplex $\alpha^*: h \rightarrow gf$ is $p_{x,z}$ -cocartesian in $\mathcal{E}(x, z)$ and $f: x \rightarrow y$ is $\mathrm{N}^{\mathrm{sc}}(p)$ -cocartesian, then α is left $\mathrm{N}^{\mathrm{sc}}(p)$ -outer.*
- (ii) *If the associated 1-simplex $\alpha^*: h \rightarrow gf$ is $p_{x,z}$ -cocartesian in $\mathcal{E}(x, z)$ and $g: y \rightarrow z$ is $\mathrm{N}^{\mathrm{sc}}(p)$ -cartesian, then α is right $\mathrm{N}^{\mathrm{sc}}(p)$ -outer.*

- (iii) If α is left $N^{\text{sc}}(p)$ -outer and f is a degenerate edge, then α^* is $p_{x,z}$ -cocartesian.
- (iv) If α is right $N^{\text{sc}}(p)$ -outer and g is a degenerate edge, then α^* is $p_{x,z}$ -cocartesian.

Proof Statements (iii) and (iv) follow from Remark 2.1.8 by using the commutative diagram (11) as in the proof of Lemma 4.2.10, and statements (i) and (ii) imply each other by passing to opposites. We now prove (i). Suppose that f is $N^{\text{sc}}(p)$ -cocartesian and that α^* is $p_{x,z}$ -cocartesian in $\mathcal{E}(x, z)$, and let us prove that α is left $N^{\text{sc}}(p)$ -outer. By Remark 2.1.2 we have to provide a solution for every lifting problem of the form

$$(12) \quad \begin{array}{ccc} & \Delta^{\{0,1,n\}} & \\ & \swarrow \alpha^* & \searrow \\ \Lambda_0^n & \xrightarrow{\quad} & N^{\text{sc}} \mathcal{E} \\ \downarrow & & \downarrow N^{\text{sc}}(p) \\ \Delta^n & \xrightarrow{\quad} & N^{\text{sc}} \mathcal{B} \end{array}$$

Transposing along the adjunction $\mathcal{C}^{\text{sc}} \dashv N^{\text{sc}}$ we obtain an equivalent lifting problem of the form

$$\begin{array}{ccc} & \mathcal{C}^{\text{sc}} \Delta^{\{0,1,n\}} & \\ & \swarrow \alpha^* & \searrow \\ \mathcal{C}^{\text{sc}} \Lambda_0^n & \xrightarrow{f} & \mathcal{E} \\ \downarrow & & \downarrow p \\ \mathcal{C}^{\text{sc}} \Delta^n & \xrightarrow{g} & \mathcal{B} \end{array}$$

A simple combinatorial analysis shows that this amounts to compatibly solving the lifting problems determined by the back and front faces of the cube

$$(13) \quad \begin{array}{ccccc} & & \square_0^{n-1,1} & \xrightarrow{f_{0,n}} & \mathcal{E}(x, z) \\ & \nearrow -\circ\{0,1\} & \downarrow f_{1,n} & \nearrow -\circ f & \downarrow p_{x,z} \\ \partial \square^{n-2} & \xrightarrow{\quad} & \square^{n-1} & \xrightarrow{g_{0,n}} & \mathcal{E}(y, z) \\ \downarrow & & \downarrow & & \downarrow p_{y,z} \\ & \nearrow -\circ\{0,1\} & \square^{n-2} & \xrightarrow{g_{1,n}} & \mathcal{B}(px, pz) \\ \square^{n-2} & \xrightarrow{\quad} & \square^{n-2} & \xrightarrow{g_{1,n}} & \mathcal{B}(py, pz) \\ & & & & \downarrow -\circ p(f) \end{array}$$

where we have identified

$$\mathcal{C}^{\text{sc}} \Delta^n(0, n) = \square^{n-1}, \quad \mathcal{C}^{\text{sc}} \Lambda_0^n(0, n) = \square_0^{n-1,1}, \quad \mathcal{C}^{\text{sc}} \Delta^n(1, n) = \square^{n-2}, \quad \mathcal{C}^{\text{sc}} \Lambda_0^n(1, n) = \partial \square^{n-2}.$$

Equivalently, we may consider this as a lifting problem in the arrow category of marked simplicial sets, involving the morphism between arrows encoded by the left square against the morphism between arrows

encoded by the right square. We may endow this arrow category with the projective model structure, so that cofibrations are Reedy cofibrations and fibrations are levelwise. We then find that in the lifting problem encoded by the above cube, the left arrow constitutes a cofibration between cofibrant objects, while the right arrow is a fibration between fibrant objects (by our assumption that p is a fibration of $\mathcal{C}at_\infty$ -categories).

Now, it is a general fact concerning model categories that the lifting property in a given square involving a cofibration between cofibrant objects against a fibration between fibrant objects is a *homotopy invariant property*, that is, it does not change if one replaces the given square by a levelwise weakly equivalent one of the same nature. In particular, in proving the existence of a lift we may as well replace the cube (13) with a levelwise weakly equivalent one, as long as we make sure it also has the property that its left square is Reedy cofibrant and its right square is levelwise fibrant with vertical legs fibrations. We now choose to make such a modification by simply replacing the corner $\mathcal{E}(y, z)$ with the fiber product

$$\mathcal{X} := \mathcal{E}(x, z) \times_{\mathcal{B}(px, pz)} \mathcal{B}(py, pz),$$

which is also a homotopy fiber product since the vertical legs are fibrations between fibrant objects. The map $\mathcal{E}(y, z) \rightarrow \mathcal{X}$ is an equivalence: indeed, by Proposition 4.2.7, $f \in \mathcal{E}(x, y)_0$ is a cocartesian arrow, and hence the right square in (13) is homotopy cartesian. We conclude that the new cube is levelwise equivalent to the old one, while clearly still keeping the same property of having its left square Reedy cofibrant and its right square levelwise fibrant with vertical maps fibrations. At the same time, by its construction, the data of a lift in the modified cube is the same as a lift in its back square

$$\begin{array}{ccc} \square_0^{n-1,1} & \longrightarrow & \mathcal{E}(x, z) \\ \downarrow & & \downarrow p_{x,z} \\ \square^{n-1} & \longrightarrow & \mathcal{B}(px, pz) \end{array}$$

where the edge corresponding to $\Delta^1 \times (0, \dots, 0)$ in $\square_0^{n-1,1}$ is sent to a $p_{x,z}$ -cocartesian edge in $\mathcal{E}(x, z)$ by assumption. A solution therefore exists by Lemma 4.2.6, thus concluding the proof of the proposition. \square


Proposition 4.2.12 *Let $p: \mathcal{E} \rightarrow \mathcal{B}$ be a fibration of $\mathcal{C}at_\infty$ -categories. Then p is an enriched 2-inner (resp. 2-outer) fibration if and only if*

$$N^{sc}(p): N^{sc}\mathcal{E} \rightarrow N^{sc}\mathcal{B}$$

is a 2-inner (resp. 2-outer) fibration of ∞ -bicategories.

Proof We first note that since p is a fibration between fibrant objects the same holds for $N^{sc}(p)$, and so the latter is always a weak fibration. We now recall that the vertices in $N^{sc}(\mathcal{E})$ correspond exactly to the objects of \mathcal{E} , and the edges $f: x \rightarrow y$ in $N^{sc}(\mathcal{E})$ correspond exactly to a pair of objects $x, y \in \mathcal{E}$ and a

vertex $f \in \mathcal{E}(x, y)$. In addition, triangles



correspond bijectively to triples of objects $x, y, z \in \mathcal{E}$, triples of vertices $f \in \mathcal{E}(x, y)$, $g \in \mathcal{E}(y, z)$, $h \in \mathcal{E}(x, z)$ and an edge $\tau: h \Rightarrow g \circ f$ in $\mathcal{E}(x, z)$.

Considering inner lifts of triangles, it follows directly from Lemma 4.2.10 that the triangles in $\mathcal{N}^{\text{sc}}(\mathcal{B})$ admits a sufficient supply of left (resp. right) $\mathcal{N}^{\text{sc}}(p)$ -inner lifts if and only if each $p_{x,y}: \mathcal{E}(x, y) \rightarrow \mathcal{B}(x, y)$ has a sufficient supply of cartesian lifts, and these cartesian lifts are closed under precomposition (resp. postcomposition) in \mathcal{E} . We may then conclude that $\mathcal{N}^{\text{sc}}(p)$ is an 2-inner fibration if and only if \mathcal{E} is a 2-inner fibration of \mathcal{Cat}_{∞} -categories.

We now consider the outer case. By Corollary 2.3.9 we may restrict attention to triangles in \mathcal{B} with one leg degenerate and lifts whose same leg is degenerate in \mathcal{E} . We then deduce from Lemma 4.2.11 that the triangles in $\mathcal{N}^{\text{sc}}(\mathcal{B})$ admits a sufficient supply of left (resp. right) p -outer lifts if and only if each $p_{x,y}: \mathcal{E}(x, y) \rightarrow \mathcal{B}(x, y)$ has a sufficient supply of cocartesian lifts. Since the definition of 2-outer fibration contains explicitly the closure under left/right whiskering (which corresponds, up to equivalence, to pre/post composition), we can conclude that $\mathcal{N}^{\text{sc}}(p)$ is a 2-outer fibration if and only if \mathcal{E} is an enriched 2-outer fibration of \mathcal{Cat}_{∞} -categories, as desired. \square

We can now deduce our main result of interest:

Proof of Theorem 4.2.4 Our goal is to compare enriched 2-inner cartesian fibrations of \mathcal{Cat}_{∞} -categories with 2-inner cartesian fibrations of ∞ -bicategories. The cartesian fibrational part is given by Corollary 4.2.8 while the 2-inner fibrational part is dealt with by Proposition 4.2.12. Combining the two results the theorem is thereby proven. \square

We finish this section by collecting a few corollaries which can be easily deduced from the comparison of Theorem 4.2.4.

Corollary 4.2.13 *Let $p: \mathcal{E} \rightarrow \mathcal{B}$ be a 2-inner (co)cartesian fibration. Then a triangle $\sigma: \Delta^2 \rightarrow \mathcal{E}$ is left p -inner if and only if it is right p -inner.*

Proof Since \mathcal{N}^{sc} is a right Quillen equivalence there exists a fibration $q: \mathcal{C} \rightarrow \mathcal{D}$ of \mathcal{Cat}_{∞} -categories which fits in a commutative diagram of the form

$$(14) \quad \begin{array}{ccc}
 \mathcal{E} & \xrightarrow{\cong} & \mathcal{N}^{\text{sc}}\mathcal{C} \\
 p \downarrow & & \downarrow \mathcal{N}^{\text{sc}}q \\
 \mathcal{B} & \xrightarrow{\cong} & \mathcal{N}^{\text{sc}}\mathcal{D}
 \end{array}$$

where the horizontal maps are equivalences of ∞ -bicategories. Now the right vertical arrow is a bicategorical fibration, being the image under a right Quillen functor of a Dwyer–Kan fibration, while the

left vertical arrow is a bicategorical fibration by Proposition 1.4.13. Applying Proposition 2.4.1 we now obtain that the right vertical map is a 2–inner/outer (co)cartesian fibration and that the top horizontal map preserves and detects left/right inner triangles. It will hence suffice to prove the left and right $N^{sc}(p)$ –inner triangles coincide in $N^{sc}(\mathcal{C})$. By Lemma 4.2.10 this amounts to showing that for every $x, y \in \mathcal{C}$ and morphism $e \in \mathcal{C}(x, y)_1$ in the mapping ∞ –category, the condition that e is cartesian with respect to $q_{x,y}: \mathcal{C}(x, y) \rightarrow \mathcal{D}(qx, qy)$ and remains cartesian after postcomposing with any morphism $y \rightarrow z$ is equivalent to the condition that e is $q_{x,y}$ –cartesian and remains cartesian after precomposing with any morphism $w \rightarrow x$. Indeed, by Theorem 4.2.4 we have that $q: \mathcal{C} \rightarrow \mathcal{D}$ is an enriched 2–inner cartesian fibration of Cat_∞ –categories, and so both conditions are equivalent to e simply being $q_{x,y}$ –cartesian. \square

Corollary 4.2.14 *Let \mathcal{E} and \mathcal{B} two ∞ –bicategories.*

- (i) *A given 2–inner (co)cartesian fibration $p: \mathcal{E} \rightarrow \mathcal{B}$ is equivalent to a 1–inner (co)cartesian fibration if and only if every triangle is left and right inner.*
- (ii) *A given 2–outer (co)cartesian fibration $p: \mathcal{E} \rightarrow \mathcal{B}$ is equivalent to a 1–outer (co)cartesian fibration if and only if every triangle whose left leg is p –cocartesian is left outer and every triangle whose right leg is p –cartesian is right outer.*

Proof From the explicit description of Remark 2.1.2 one immediately finds that if p is a 1–inner (co)cartesian fibration then every triangle is both left and right p –inner. Similarly, if p is a 1–outer cartesian fibration then any (co)cartesian arrow is automatically strongly (co)cartesian by [6, Proposition 2.3.7] (and its dual), and so every triangle whose left leg is p –cocartesian is left p –outer and any triangle whose right leg is p –cartesian is right p –outer.

To prove the “if” direction, we now invoke the fact that N^{sc} is a right Quillen equivalence to deduce the existence of fibration $q: \mathcal{C} \rightarrow \mathcal{D}$ of Cat_∞ –categories which fits in a commutative diagram of the form

$$(15) \quad \begin{array}{ccc} \mathcal{E} & \xrightarrow{\cong} & N^{sc}\mathcal{C} \\ p \downarrow & & \downarrow N^{sc}q \\ \mathcal{B} & \xrightarrow{\cong} & N^{sc}\mathcal{D} \end{array}$$

whose horizontal maps are equivalences of ∞ –bicategories. Now the right vertical arrow is a bicategorical fibration, being the image under a right Quillen functor of a Dwyer–Kan fibration, while the left vertical arrow is a bicategorical fibration by Proposition 1.4.13. Applying Proposition 2.4.1 we now obtain that the right vertical map is a 2–inner/outer (co)cartesian fibration, and hence by Theorem 4.2.4 the functor $q: \mathcal{C} \rightarrow \mathcal{D}$ is an enriched 2–inner/outer (co)cartesian fibration of Cat_∞ –categories. For every $x, y \in \mathcal{E}$ with images $x', y' \in N^{sc}\mathcal{C}$ (which we can identify with objects of \mathcal{C}) we may then consider the commutative diagram

$$(16) \quad \begin{array}{ccccccc} \text{Hom}_{\mathcal{E}}^{\triangleright}(x, y) & \xrightarrow{\cong} & \text{Hom}_{N^{sc}\mathcal{C}}^{\triangleright}(x', y') & \xrightarrow{\cong} & \text{Hom}_{N^{sc}\mathcal{C}}(x', y') & \xleftarrow{\cong} & \mathcal{C}(x', y') \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ \text{Hom}_{\mathcal{B}}^{\triangleright}(px, py) & \xrightarrow{\cong} & \text{Hom}_{N^{sc}\mathcal{D}}^{\triangleright}(qx', qy') & \xrightarrow{\cong} & \text{Hom}_{N^{sc}\mathcal{D}}(qx', qy') & \xleftarrow{\cong} & \mathcal{D}(qx', qy') \end{array}$$

in which all vertical arrows are cartesian fibrations in the inner case and cocartesian fibrations in the outer case. Our assumption implies in particular that every left degenerate triangle in \mathcal{E} is left p -inner/outer, which by Remark 2.1.8 implies that the leftmost vertical map in (16) is a left fibration in the outer case and a right fibration in the inner case. The same consequently holds for all vertical maps in (16), which implies that $q: \mathcal{C} \rightarrow \mathcal{D}$ is an enriched 1-inner/outer (co)cartesian fibration of $\mathcal{C}at_\infty$ -categories. By [6, Proposition 3.1.3] this means that $N^{\text{sc}}q: N^{\text{sc}}\mathcal{C} \rightarrow N^{\text{sc}}\mathcal{D}$ is a 1-inner/outer (co)cartesian fibration. Since the square (15) is an equivalence between two bicategorical fibrations between fibrant objects, these two fibrations satisfy the same right lifting properties. Applying this to the right lifting properties of Definition 1.4.3, we conclude that if the right one is a 1-inner fibration then so is the left, and if the right one is a 1-outer fibration then so is the left. We may consequently deduce that $p: \mathcal{E} \rightarrow \mathcal{B}$ is a 1-inner/outer (co)cartesian fibration, as desired. \square

References

- [1] **F Abellán García, WH Stern**, *2-Cartesian fibrations, I: A model for ∞ -bicatogories fibred in ∞ -bicatogories*, Appl. Categ. Structures 30 (2022) 1341–1392 MR Zbl
- [2] **MA Batanin**, *Monoidal globular categories as a natural environment for the theory of weak n -categories*, Adv. Math. 136 (1998) 39–103 MR Zbl
- [3] **MA Batanin**, *The Eckmann–Hilton argument and higher operads*, Adv. Math. 217 (2008) 334–385 MR Zbl
- [4] **N Bottman, S Carmeli**, *$(A_\infty, 2)$ -categories and relative 2-operads*, High. Struct. 5 (2021) 401–421 MR Zbl
- [5] **M Buckley**, *Fibred 2-categories and bicategories*, J. Pure Appl. Algebra 218 (2014) 1034–1074 MR Zbl
- [6] **A Gagna, Y Harpaz, E Lanari**, *Fibrations and lax limits of $(\infty, 2)$ -categories*, preprint (2020) arXiv 2012.04537
- [7] **A Gagna, Y Harpaz, E Lanari**, *Gray tensor products and lax functors of $(\infty, 2)$ -categories*, Adv. Math. 391 (2021) art. id. 107986 MR Zbl
- [8] **A Gagna, Y Harpaz, E Lanari**, *On the equivalence of all models for $(\infty, 2)$ -categories*, J. Lond. Math. Soc. 106 (2022) 1920–1982 MR Zbl
- [9] **D Gaitsgory, N Rozenblyum**, *A study in derived algebraic geometry, I: Correspondences and duality*, Math. Surv. Monogr. 221, Amer. Math. Soc., Providence, RI (2017) MR Zbl
- [10] **D Gaitsgory, N Rozenblyum**, *A study in derived algebraic geometry, II: Deformations, Lie theory and formal geometry*, Math. Surv. Monogr. 221, Amer. Math. Soc., Providence, RI (2017) MR Zbl
- [11] **C Hermida**, *Some properties of Fib as a fibred 2-category*, J. Pure Appl. Algebra 134 (1999) 83–109 MR Zbl
- [12] **J Lurie**, *Higher topos theory*, Ann. of Math. Stud. 170, Princeton Univ. Press (2009) MR Zbl
- [13] **J Lurie**, *$(\infty, 2)$ -categories and the Goodwillie calculus, I*, preprint (2009) Available at <https://www.math.ias.edu/~lurie/papers/GoodwillieI.pdf>

- [14] **J Lurie**, *Kerodon: an online resource for homotopy-coherent mathematics* (2018) Available at <https://kerodon.net/>

*Institute of Mathematics, Czech Academy of Sciences
Prague, Czech Republic*

*Institut Galilée, Université Paris 13
Villetaneuse, France*

*Institute of Mathematics, Czech Academy of Sciences
Prague, Czech Republic*

`gagna@math.cas.cz`, `harpaz@math.univ-paris13.fr`, `edoardo.lanari.e1@gmail.com`

<https://sites.google.com/view/andreagagna/home>,

<https://www.math.univ-paris13.fr/~harpaz>, <https://edolana.github.io/>

Received: 27 July 2021 Revised: 5 February 2023

ALGEBRAIC & GEOMETRIC TOPOLOGY

msp.org/agt

EDITORS

PRINCIPAL ACADEMIC EDITORS

John Etnyre
etnyre@math.gatech.edu
Georgia Institute of Technology

Kathryn Hess
kathryn.hess@epfl.ch
École Polytechnique Fédérale de Lausanne

BOARD OF EDITORS

Julie Bergner	University of Virginia jeb2md@eservices.virginia.edu	Christine Lescop	Université Joseph Fourier lescop@ujf-grenoble.fr
Steven Boyer	Université du Québec à Montréal cohf@math.rochester.edu	Robert Lipshitz	University of Oregon lipshitz@uoregon.edu
Tara E Brendle	University of Glasgow tara.brendle@glasgow.ac.uk	Norihiko Minami	Yamato University minami.norihiko@yamato-u.ac.jp
Indira Chatterji	CNRS & Univ. Côte d'Azur (Nice) indira.chatterji@math.cnrs.fr	Andrés Navas	Universidad de Santiago de Chile andres.navas@usach.cl
Alexander Dranishnikov	University of Florida dranish@math.ufl.edu	Robert Oliver	Université Paris 13 bobol@math.univ-paris13.fr
Tobias Ekholm	Uppsala University, Sweden tobias.ekholm@math.uu.se	Jessica S Purcell	Monash University jessica.purcell@monash.edu
Mario Eudave-Muñoz	Univ. Nacional Autónoma de México mario@matem.unam.mx	Birgit Richter	Universität Hamburg birgit.richter@uni-hamburg.de
David Futer	Temple University dfuter@temple.edu	Jérôme Scherer	École Polytech. Féd. de Lausanne jerome.scherer@epfl.ch
John Greenlees	University of Warwick john.greenlees@warwick.ac.uk	Vesna Stojanoska	Univ. of Illinois at Urbana-Champaign vesna@illinois.edu
Ian Hambleton	McMaster University ian@math.mcmaster.ca	Zoltán Szabó	Princeton University szabo@math.princeton.edu
Matthew Hedden	Michigan State University mhedden@math.msu.edu	Maggy Tomova	University of Iowa maggy-tomova@uiowa.edu
Hans-Werner Henn	Université Louis Pasteur henn@math.u-strasbg.fr	Chris Wendl	Humboldt-Universität zu Berlin wendl@math.hu-berlin.de
Daniel Isaksen	Wayne State University isaksen@math.wayne.edu	Daniel T Wise	McGill University, Canada daniel.wise@mcgill.ca
Thomas Koberda	University of Virginia thomas.koberda@virginia.edu	Lior Yanovski	Hebrew University of Jerusalem lior.yanovski@gmail.com
Markus Land	LMU München markus.land@math.lmu.de		


See inside back cover or msp.org/agt for submission instructions.

The subscription price for 2024 is US \$705/year for the electronic version, and \$1040/year (+\$70, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Algebraic & Geometric Topology is indexed by Mathematical Reviews, Zentralblatt MATH, Current Mathematical Publications and the Science Citation Index.

Algebraic & Geometric Topology (ISSN 1472-2747 printed, 1472-2739 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840.

AGT peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<https://msp.org/>

© 2024 Mathematical Sciences Publishers

ALGEBRAIC & GEOMETRIC TOPOLOGY

Volume 24 Issue 9 (pages 4731–5219) 2024

Cartesian fibrations of $(\infty, 2)$ -categories	4731
ANDREA GAGNA, YONATAN HARPAZ and EDOARDO LANARI	
On the profinite distinguishability of hyperbolic Dehn fillings of finite-volume 3-manifolds	4779
PAUL RAPOPORT	
Index-bounded relative symplectic cohomology	4799
YUHAN SUN	
Heegaard Floer homology, knotifications of links, and plane curves with noncuspidal singularities	4837
MACIEJ BORODZIK, BEIBEI LIU and IAN ZEMKE	
Classifying spaces of infinity-sheaves	4891
DANIEL BERWICK-EVANS, PEDRO BOAVIDA DE BRITO and DMITRI PAVLOV	
Chern character for infinity vector bundles	4939
CHEYNE GLASS, MICAH MILLER, THOMAS TRADLER and MAHMOUD ZEINALIAN	
Derived character maps of group representations	4991
YURI BEREST and AJAY C RAMADOSS	
Instanton knot invariants with rational holonomy parameters and an application for torus knot groups	5045
HAYATO IMORI	
On the invariance of the Dowlin spectral sequence	5123
SAMUEL TRIPP and ZACHARY WINKELER	
Monoidal properties of Franke's exotic equivalence	5161
NIKITAS NIKANDROS and CONSTANZE ROITZHEIM	
Characterising quasi-isometries of the free group	5211
ANTOINE GOLDSBOROUGH and STEFANIE ZBINDEN	