

AG  
T

*Algebraic & Geometric  
Topology*

Volume 25 (2025)

**Bridge trisections and Seifert solids**

JASON JOSEPH

JEFFREY MEIER

MAGGIE MILLER

ALEXANDER ZUPAN



# Bridge trisections and Seifert solids

JASON JOSEPH

JEFFREY MEIER

MAGGIE MILLER

ALEXANDER ZUPAN

We adapt Seifert’s algorithm for classical knots and links to the setting of triplane diagrams for bridge trisected surfaces in the 4-sphere. Our approach allows for the construction of a Seifert solid that is described by a Heegaard diagram. The Seifert solids produced can be assumed to have exteriors that can be built without 3-handles; in contrast, we give examples of Seifert solids (not coming from our construction) whose exteriors require arbitrarily many 3-handles. We conclude with two classification results. The first shows that surfaces admitting doubly standard shadow diagrams are unknotted. The second says that a  $b$ -bridge trisection in which some sector contains at least  $b - 1$  patches is completely decomposable, thus the corresponding surface is unknotted. This settles affirmatively a conjecture of the second and fourth authors.

[57K45](#); [57K10](#)

## 1 Introduction

One of the most important avenues available for study in knotted surfaces in 4-space is the analysis of the 3-dimensional Seifert solids bounded by such surfaces. There are many situations in which information about such a Seifert solid gives rise to useful information about the corresponding knotted surface. Examples, ranging from classical to modern, include Gordon’s proof that 2-knots are not determined by their complements [7], Cochran’s characterization of fibered homotopy-ribbon 2-knots [3], and recent work of Dai and Miller analyzing the relevance of homology cobordism invariants of Seifert solids [4].

Here we show how topological information about a knotted surface can be recovered from a bridge trisection of the surface, which allows for the diagrammatic study of knotted surfaces and their Seifert solids. A *bridge trisection* of a surface  $\mathcal{S}$  in  $S^4$  is a certain decomposition of  $(S^4, \mathcal{S})$  into three trivial disk systems  $(B_1^4, \mathcal{D}_1)$ ,  $(B_2^4, \mathcal{D}_2)$ , and  $(B_3^4, \mathcal{D}_3)$  that can be encoded diagrammatically either as a triple of tangles called a *triplane diagram* or as a corresponding *shadow diagram*.

In [Section 3](#), we give a version of Seifert’s algorithm for bridge-trisected surfaces, showing how a triplane diagram can be used to produce a 3-manifold bounded by a connected surface  $\mathcal{S}$  with normal Euler number zero.

**Theorem 3.4** *If  $S$  is connected and  $e(S) = 0$ , then there is a procedure to produce a Seifert solid for  $S$  that takes as input a triplane diagram for  $S$ .*

In [Section 3.2](#), we give an explicit procedure for constructing a Heegaard diagram for such a 3-manifold when  $S \cong S^2$ . As a corollary of the work in building Seifert solids, we recover a combinatorial proof of the existence of Seifert solids. Although the literature already contains a method for producing a Heegaard diagram for a Seifert solid — namely, the work of Carter and Saito [\[2\]](#) — the procedure described here is a bit more practical. In [\[2, Section 3\]](#), the authors employ their methods to take a broken surface diagram and produce a genus 21 Heegaard diagram for a punctured  $L(3, 1) \# (\#^3(S^1 \times S^2))$  bounded by the 2-twist spun trefoil, noting that this solid is nonminimal, since the same 2-knot also bounds a punctured  $L(3, 1)$ . In contrast, in [Section 3.3](#) we use our procedure to find genus three Heegaard diagrams for Seifert solids bounded by the spun trefoil and 1-twist spun trefoil, where these solids are minimal. For the 2-twist spun trefoil, the procedure yields a genus four Heegaard diagram for a Seifert solid (calculations omitted here). The 2-dimensional data contained in a triplane diagram can often be easier to manipulate and simplify than the data in a broken surface diagram; as such, both the solids and their Heegaard diagrams produced by [Theorem 3.4](#) are likely to be less complicated.

We also show that certain bridge trisected surfaces are unknotted.

**Theorem 3.3** *If a surface  $S$  has a doubly standard shadow diagram, then  $S$  is unknotted.*

In practice, [Theorem 3.3](#) offers a new and effective method to show unknottedness for bridge trisected surfaces. The doubly standard criterion has considerable potential to aid in the tabulation of low-complexity knotted surfaces, since verifying that a shadow diagram is doubly standard can be much easier than proving unknottedness via other methods.

One of the key features of trisection theory is that it provides a vehicle to adapt 3-dimensional ideas to dimension four, and in [Section 4](#), we prove another result that fits into this line of research. It is well-known that the complement of every *canonical* Seifert surface (ie one obtained from Seifert’s algorithm) is a handlebody. Thus, it is natural to attempt to extend this notion to dimension four. In this vein, we call a Seifert solid *canonical* if it is obtained from the procedure presented in [Section 3](#), and we call a Seifert solid *spinal* if its exterior in  $S^4$  can be built without 3-handles. We prove the following two results relating (and distinguishing) these concepts:

**Theorem 4.1** *If a surface-knot  $S$  admits a Seifert solid, then it admits a canonical Seifert solid that is spinal.*

In fact, modulo some additional, easily satisfied connectivity conditions, every canonical Seifert solid is spinal. The next result shows that some Seifert solids (in contrast to canonical Seifert solids and many standard examples) are “far” from being spinal.

**Theorem 4.2** *Given any  $n \in \mathbb{N}$ , there exists a 2-knot  $\mathcal{K}$  that bounds a Seifert solid  $Y$  homeomorphic to  $(S^1 \times S^2)^\circ$  such that  $S^4 \setminus \nu(Y)$  requires at least  $n$  4-dimensional 3-handles.*

Finally, in [Section 5](#) we prove the following standardness result, affirmatively settling Conjecture 4.3 of Meier and Zupan [\[15\]](#).

**Theorem 5.2** *Let  $\mathfrak{T}$  be a  $(b; c)$ -bridge trisection with  $c_i = b - 1$  for some  $i \in \mathbb{Z}_3$ . Then  $\mathfrak{T}$  is completely decomposable, and the underlying surface-link is either the unlink of  $\min\{c_i\}$  2-spheres or the unlink of  $\min\{c_i\}$  2-spheres and one projective plane, depending on whether  $|c_{i-1} - c_{i+1}| = 1$  or 0.*

The proof relies on theorems of Scharlemann [\[19\]](#) and Bleiler and Scharlemann [\[1\]](#) regarding planar surfaces in 3-manifolds. The methods of the proof are somewhat unrelated to the methods used in the preceding sections and may be of independent interest. The second and fourth authors previously handled this case when  $c_i = b$  for some  $i \in \mathbb{Z}_3$  [\[15, Proposition 4.1\]](#). [Theorem 5.2](#) can be seen as the analog for bridge trisections of Theorem 1.2 of Meier, Schirmer and Zupan [\[13\]](#), which establishes a similar standardness result for trisections of closed manifolds; as such, our theorem fills an important gap in the trisections literature and provides yet another avenue to verify that a surface in  $S^4$  is unknotted.

## Acknowledgements

This paper began following discussions at the workshop *Unifying 4-dimensional knot theory*, which was hosted by the Banff International Research Station in November 2019, and the authors would like to thank BIRS for providing an ideal space for sparking collaboration. We are grateful to Masahico Saito for sharing his interest in adapting Seifert’s algorithm to bridge trisections and motivating this paper. The authors would like to thank Román Aranda, Scott Carter, and Peter Lambert-Cole for helpful conversations. The authors are grateful to the referee for their careful reading and useful comments. Joseph was supported by MPIM during part of this project, as well as NSF grants DMS-1664567 and DMS-1745670. Meier was supported by NSF grants DMS-1933019 and DMS-2006029. Miller was supported by MPIM during part of this project, as well as NSF grants DGE-1656466 (at Princeton) and DMS-2001675 (at MIT) and a research fellowship from the Clay Mathematics Institute (at Stanford). Zupan was supported by MPIM during part of this project, by NSF grants DMS-1664578 and DMS-2005518, and by a Simons Fellowship.

## 2 Preliminaries

We work in the smooth category. This section includes an abbreviated introduction to the concepts relevant to this paper, but the interested reader is encouraged to consult [\[5\]](#) for further information about 4-manifold trisections and [\[9, Section 2; 15\]](#) for more detailed discussions of bridge trisections. We limit our work here to surfaces in  $S^4$ , but there is also a theory of bridge trisections in arbitrary 4-manifolds; see [\[16\]](#).

## 2.1 Bridge trisections

Let  $S$  be an embedded surface in  $S^4$ , let  $b$  be a positive integer, and let  $\mathbf{c} = (c_1, c_2, c_3)$  be a triple of positive integers. A  $(b; \mathbf{c})$ -*bridge trisection* of  $(S^4, S)$  is a decomposition

$$(S^4, S) = (X_1, \mathcal{D}_1) \cup (X_2, \mathcal{D}_2) \cup (X_3, \mathcal{D}_3)$$

such that

- (1) each  $\mathcal{D}_i$  is a collection of  $c_i$  boundary-parallel disks in the 4-ball  $X_i$ ,
- (2) each intersection  $\mathcal{T}_i = \mathcal{D}_{i-1} \cap \mathcal{D}_i$  is a boundary-parallel tangle in the 3-ball  $H_i = X_{i-1} \cap X_i$  (with indices considered mod 3),
- (3) the triple intersection  $\mathcal{D}_1 \cap \mathcal{D}_2 \cap \mathcal{D}_3$  is a collection of  $b$  points in the 2-sphere  $\Sigma = X_1 \cap X_2 \cap X_3$ .

In [15], it was proved that every surface  $S$  admits a  $(b; \mathbf{c})$ -bridge trisection for some  $(b; \mathbf{c})$ . We choose orientations so that  $\partial(X_i, \mathcal{D}_i) = (H_i, \mathcal{T}_i) \cup (\bar{H}_{i+1}, \bar{\mathcal{T}}_{i+1})$ . When we wish to be succinct, we use  $\mathfrak{T}$  to represent a bridge trisection, with components labeled as above.

## 2.2 Diagrams for bridge trisections

The existence of bridge trisections gives rise to a new diagrammatic theory for surfaces in  $S^4$ , using an object called a *triplane diagram*, a triple  $\mathbb{D} = (\mathbb{D}_1, \mathbb{D}_2, \mathbb{D}_3)$  of trivial planar diagrams with the additional condition that each  $\mathbb{D}_i \cup \bar{\mathbb{D}}_{i+1}$  is a classical diagram for an unlink. In [15], it was shown that every triplane diagram determines a bridge trisection  $\mathfrak{T}$ . Conversely, given a bridge trisection  $\mathfrak{T}$  of  $(S^4, S)$ , we can choose a triple of disks  $E_i \subset H_i$  with common boundary and project the tangles  $\mathcal{T}_i$  onto  $E_i$  to obtain a triplane diagram. Of course, the choices of disks and projections are not unique, but any two triplane diagrams corresponding the same bridge trisection  $\mathfrak{T}$  are related by a finite collection of *interior Reidemeister moves* and *mutual braid transpositions*, while any two bridge trisections  $\mathfrak{T}$  and  $\mathfrak{T}'$  for the same surface  $S$  are related by *perturbation* and *deperturbation* moves.

In addition, bridge trisections yield another type of diagram: each trivial tangle  $\mathcal{T}_i$  can be isotoped rel-boundary into the surface  $\Sigma$ , yielding a triple  $(A, B, C)$  of pairwise disjoint collections of arcs called a *shadow diagram*, which has the property that  $\partial A = \partial B = \partial C$ , and the pairwise unions of any two of the tangles  $\mathcal{T}_A, \mathcal{T}_B$ , and  $\mathcal{T}_C$  determined by the arcs are unlinks. As with triplane diagrams, any shadow diagram determines a bridge trisection. Further details about shadow diagrams can be found in [14].

Here we consider special types of shadow diagrams. We say that a pair of collections of arcs in a shadow diagram is *standard* if their union is embedded. Any bridge trisection admits a shadow diagram  $(A, B, C)$  in which one of the pairs is standard. If two or three pairs of shadows in a shadow diagram  $(A, B, C)$  are standard, then we say that  $(A, B, C)$  is *doubly standard* or *triply standard*, respectively. [Theorem 3.3](#) says that doubly standard (and thus triply standard) diagrams always describe unknotted surfaces.




Figure 1: Triplane diagrams for  $P_+$  and  $P_-$ .

## 2.3 Unknotted surfaces

In this subsection, we review standard notions of unknottedness for surfaces in  $S^4$ . A closed connected orientable surface  $S$  in  $S^4$  is *unknotted* if it bounds an embedded 3-dimensional handlebody  $H \subset S^4$ . For nonorientable surfaces, the definition is slightly more involved. We define the two unknotted projective planes,  $P_{\pm}$ , to be the two standard projective planes in  $S^4$ , pictured via their triplane diagrams in Figure 1, where  $e(P_{\pm}) = \pm 2$ .

In general, for a nonorientable surface  $S$ , we say that  $S$  is *unknotted* if  $S$  is isotopic to a connected sum of some number of copies of  $P_+$  and  $P_-$ . See [9, Remark 2.6] for a detailed discussion of the orientation conventions used here.

## 3 Seifert solids

Classical results of Gluck [6] (resp. Gordon–Litherland [8]) assert that every orientable surface  $S$  (resp. surface  $S$  with  $e(S) = 0$ ) in  $S^4$  bounds an embedded 3-manifold, called a *Seifert solid* in the orientable case. In the setting of broken surface diagrams, Carter and Saito provided a procedure that in many respects mimics Seifert’s algorithm for classical knots [2]. In this section, we describe an extension of Seifert’s algorithm that takes an oriented triplane diagram  $\mathbb{D}$  and produces a Seifert solid whose intersection with  $\partial X_i$  agrees with the classical Seifert algorithm performed on the oriented unlink diagram  $\mathbb{D}_i \cup \overline{\mathbb{D}}_{i+1}$ . We also obtain alternative proofs of the theorems of Gluck and Gordon–Litherland mentioned above.

### 3.1 Existence of Seifert solids

Given a spanning surface  $F$  for an unlink  $U$ , we define the *cap-off*  $\mathcal{F}$  of  $F$  to be the closed surface  $\mathcal{F} \subset S^4$  obtained by gluing a collection of trivial disks in  $B_-^4$  to  $F$  along  $U$ . (There is a unique such choice of disks up to isotopy rel-boundary in  $B_-^4$  by eg [10] or [12].) Let  $F_+ \subset S^3$  denote the Möbius band bounded by the unknot so that  $F_+$  contains a positive half-twist and has boundary slope  $+2$ , and let  $F_- \subset S^3$  denote the Möbius band bounded by the unknot with a negative half-twist and boundary slope  $-2$ . For  $n > 0$ , let  $F_n$  be the connected surface obtained by attaching  $n - 1$  trivial bands to the split union of  $n$  copies of  $F_+$ ; that is,  $F_n$  is obtained by taking the boundary connected sum of  $n$  copies of  $F_+$ . For  $n < 0$ , let  $F_n$  be obtain by taking the boundary connected sum of  $(-n)$  copies of  $F_-$ . Finally, let  $F_0$  be the disk bounded by the unknot in  $S^3$ . Additionally, let  $\mathcal{F}_n$  be the cap-off of  $F_n$ . In Figure 1, the negative Möbius band is shown to cap off into  $B_+^4$  to obtain  $P_+$ ; see also [9, Figure 2]. Here, we are capping off into  $B_-^4$ , so that by definition the cap-off  $\mathcal{F}_{-1}$  of the negative Möbius band  $F_-$  is  $P_-$ . In

contrast, the cap-off  $\mathcal{F}_1$  of the positive Möbius band  $F_+$  is  $P_+$ . (Recall that  $P_+$  and  $P_-$  denote the two unknotted projective planes in  $S^4$ ; see [Section 2.3](#).) It follows that

$$\mathcal{F}_n = \begin{cases} \text{a connected sum of } n \text{ copies of } P_+ & \text{if } n > 0, \\ \text{a connected sum of } -n \text{ copies of } P_- & \text{if } n < 0, \\ \text{an unknotted 2-sphere} & \text{if } n = 0. \end{cases}$$

The intent of the cap-off notation is to emphasize the way in which  $\mathcal{F}_n$  can be obtained from a specific surface in  $S^3$ , which will be useful in the rest of this section—especially given the following lemma:

**Lemma 3.1** *Every incompressible spanning surface  $F$  for the unknot is isotopic to  $\mathcal{F}_n$  for some  $n \in \mathbb{Z}$ .*

**Proof** First, we argue that  $F_n$  is incompressible for all  $n$ . This follows from [20], but we include a proof here. Certainly,  $F_0$  and  $F_{\pm 1}$  are incompressible, since a compression increases Euler characteristic by two. Suppose now that  $F_n$  is compressible for some  $n > 1$ , and let  $F'_n$  be the component of the surface obtained by compressing  $F_n$  so that  $\partial F'_n = \partial F_n$ . In addition, let  $\mathcal{F}'_n \subset S^4$  be the cap-off of  $F'_n$ . Then the embedded surface  $\mathcal{F}_n$  can be obtained from  $\mathcal{F}'_n$  by a 1-handle attachment, and thus  $e(\mathcal{F}'_n) = e(\mathcal{F}_n) = 2n$ . However, since the nonorientable genus of  $\mathcal{F}'_n$  is strictly less than  $n$ , this contradicts the Whitney–Massey theorem (see discussion in [9]). We conclude that  $F_n$  is incompressible.

On the other hand, suppose that  $F$  is an arbitrary incompressible spanning surface for the unknot  $U$ . The exterior of  $U$  is a solid torus  $V$ , and every simple closed curve  $c \subset \partial V$  is homotopic to a  $(p, q)$ -curve, where a  $(0, 1)$ -curve is the boundary of a meridian disk of  $V$  and a  $(1, 0)$ -curve is the boundary of a meridian disk of  $N(U)$ . The boundary of  $F$  is a  $(2k, 1)$ -curve for some integer  $k$ . (The spanning surface  $F$  intersects the disk bounded by  $U$  in some number of arcs, the endpoints of which correspond to the intersections of the  $(p, q)$ -curve with the  $(0, 1)$ -curve.) If  $F$  is orientable, then it is well-known that  $F$  is isotopic to the meridian disk  $F_0$ .

Suppose that  $F$  is nonorientable. By [20, Corollary 12], the nonorientable genus of  $F$  is equal to  $|k|$ . Assuming that  $\partial F$  and  $\partial F_0$  meet efficiently, isotope  $F$  so that it intersects  $F_0$  minimally. By standard cut-and-paste arguments, an arc of  $F \cap F_0$  which is outermost in  $F_0$  gives rise to a boundary-compressing disk  $\Delta$  for  $F$ . Since  $\partial F$  and  $\partial F_0$  meet efficiently, the result  $F'$  of boundary-compressing  $F$  along  $\Delta$  has a single boundary component and nonorientable genus  $k - 1$ . Reversing the process, we see that  $F$  can be obtained from  $F'$  by attaching a boundary-parallel band to  $F'$  along opposite sides of  $\partial F'$ . Note that  $\partial V \setminus \partial F'$  is an annulus and the band is determined by a spanning arc. Working rel-boundary, all choices of spanning arcs are related by Dehn twists about  $\partial F'$ , and so it follows that up to isotopy, there is a unique band taking  $F'$  to  $F$ .

Finally, we claim that  $F$  is isotopic to  $F_k$ , and we prove this fact by inducting on  $k$ . If  $k = \pm 1$ , then  $F$  has genus one and is obtained from the disk  $F' = F_0$  by a single boundary tubing. By the above argument, there is precisely one way to do this, and thus  $F = F_{\pm 1}$ . Now suppose that  $k > 1$  and the claim holds for  $j = k - 1$ . As above, isotope  $F$  to meet  $F_0$  minimally, and since  $k > 1$ , there are at least two arcs  $a_0$  and  $a_1$  of  $F \cap F_0$  that are outermost in  $F_0$ . Let  $\ell$  be a  $(0, 1)$ -curve that meets  $\partial F$  in a single point contained

in  $a_0$ . Then  $a_1$  gives rise to a boundary-compressing disk  $\Delta_1$  and the result  $F'$  of boundary-compressing  $F$  along  $\Delta_1$  also satisfies  $|\partial F' \cap \ell| = 1$ , since the modification was carried out away from the arc  $a_0$ . We conclude that  $F'$  has genus  $k - 1$  and boundary slope  $(2(k - 1), 1)$ . By induction  $F' = F_{k-1}$ , and since there is a unique way to obtain  $F$  from  $F'$  by boundary-tubing, it follows that  $F = F_k$ . The case  $k < -1$  follows symmetrically.  $\square$

In the next proposition, we use [Lemma 3.1](#) to understand the cap-off of any spanning surface  $F$  for an unlink in  $S^3$ :

**Proposition 3.2** *Let  $F$  be a spanning surface for an unlink  $U$  in  $S^3$ .*

- (1) *If every component of  $\partial F$  has slope 0, then the cap-off  $\mathcal{F}$  bounds a (possibly nonorientable, possibly disconnected) handlebody  $V \subset B^4$  such that  $V \cap \partial B^4 = F$ .*
- (2) *The normal Euler number  $e(\mathcal{F})$  is equal to the sum of the slopes of the boundary components of  $F$ .*
- (3) *The cap-off  $\mathcal{F}$  is a split union of unknotted surfaces in  $S^4$ .*

**Proof** Suppose  $F$  and  $F'$  are two spanning surfaces for an unlink  $U$  in  $S^3$  such that  $F'$  is isotopic relative to  $U$  to the surface obtained by surgering  $F$  along a compressing disk  $D$  for  $F$ . Then there is a compression body  $C \subset S^3 \times [0, 1]$  such that

- $C \cap (S^3 \times \{1\}) = F \times \{1\}$ ,
- $C \cap (S^3 \times \{0\}) = F' \times \{0\}$ , and
- $\partial C = (F \times \{1\}) \cup (\overline{F' \times \{0\}}) \cup (U \times [0, 1])$ ,

and  $C$  has a single critical point (of index 1) with respect to the Morse function  $S^3 \times [0, 1] \rightarrow [0, 1]$ , which we assume lies in  $S^3 \times \{\frac{1}{2}\}$ . Note that  $C$  is a product cobordism above and below  $S^3 \times \{\frac{1}{2}\}$ .

Any spanning surface  $F$  for  $U$  can be reduced to  $F'$ , a union of 2-spheres and incompressible spanning surfaces for components of  $U$  via a sequence of compressions and isotopies. If each component of  $\partial F$  has slope 0, then  $F'$  is a collection of disks and spheres. Applying the compression body construction described above for each compression taking  $F$  to  $F'$  and stacking the results, we get a compression body  $C$  cobounded by  $F$  and  $F'$ . Since  $F'$  is a collection of disks and spheres, there is a handlebody with boundary  $\mathcal{F} = F \cup \mathcal{D}$ , where  $\mathcal{D} = \overline{F'} \cup (U \times [0, 1])$  is a collection of properly embedded disks in  $B^4$ : simply cap-off the sphere components of  $C$  with 3-balls whose interiors are pushed sufficiently deep into  $B^4$ . This handlebody is nonorientable (resp. disconnected) if and only if  $F$  is. This establishes (1).

Let  $F$  be any spanning surface for an unlink  $U = \bigsqcup_{i=1}^n U_i$ . Let  $B = \bigsqcup_{i=1}^n B_i$  be a collection of disjoint 3-balls with  $U_i \subset \text{Int}(B_i)$ . Let  $F' = \bigsqcup_{i=1}^n F_i$  be a split union of incompressible spanning surfaces for the components of  $U$ , with  $F_i \subset \text{Int}(B_i)$ , so that the slopes of  $F$  and  $F'$  agree at each component of  $U$ . Let  $F''$  be the result of surgering  $F'$  along a collection of arcs so that  $F''$  and  $F$  have the same homeomorphism type relative to  $U$ ; moreover, assume that every arc of the collection intersects each component of  $\partial B$  in at most one point. It follows that  $F''$  decomposes as a split union of connected sums of surfaces, each summand of which is either a torus or an incompressible spanning surface for an unknot. Therefore, the



cap-off  $\mathcal{F}''$  is the split union of connected sums of surfaces, each summand of which is an unknotted surface in  $S^4$ . Livingston showed that  $F$  and  $F''$  are isotopic rel-boundary in  $B^4$  [12]. It follows that the cap-off  $\mathcal{F}$  will be isotopic to the cap-off  $\mathcal{F}''$ , which completes the proof of (3). Since (2) holds for  $\mathcal{F}_1$  and  $\mathcal{F}_{-1}$ , and since the normal Euler number is additive under connected sum, (2) follows, as well.  $\square$

Recall that a shadow diagram is doubly standard if two of the pairings of arcs yield embedded curves. We can use Proposition 3.2 to obtain the following classification result for doubly standard diagrams:

**Theorem 3.3** *If  $S$  has a doubly standard shadow diagram, then  $S$  is unknotted.*

Note that Theorem 3.3 also applies to surfaces with triply standard shadow diagrams, as a special class of doubly standard shadow diagrams.

**Proof** Suppose  $S$  has a shadow diagram  $(A, B, C)$  such that the pairings  $(A, B)$  and  $(B, C)$  are standard. Consider the standard Heegaard splitting  $\partial X_3 = S^3 = H_+ \cup_\Sigma H_-$ , and let  $\Sigma_\pm$  be a parallel copy of  $\Sigma$  pushed slightly into  $H_\pm$ . Note that  $A \cup B$  may have nested components (so that components of  $A \cup B$  don't necessarily bound a collection of disjoint disks). After a sequence of arc slides, however, performed only on the arcs in  $A$ , we obtain arcs  $A'$  such that the embedded curves  $A' \cup B$  bound a pairwise disjoint collection of disks. We perform a similar procedure with  $B \cup C$  to obtain  $B \cup C'$ . Now embed parallel copies  $A'_+ \cup B_+$  of the curves  $A' \cup B$  in  $\Sigma_+$  so that they bound a pairwise disjoint collection  $D_+$  of disks in  $\Sigma_+$ , and embed parallel copies  $B_- \cup C'_-$  of the curves  $B \cup C'$  in  $\Sigma_-$  so that they bound a pairwise disjoint collection  $D_-$  of disks in  $\Sigma_-$ . In  $H_+$ , there is an isotopy of  $B_+$  to  $B \subset \Sigma$  taking the disks  $D_+$  to disks  $D_1 \subset H_+$  such that  $D_1 \cap \Sigma = B$ . The tangle  $\mathcal{T}_1 = S \cap (H_+)$  is the image of  $A'_+$  under this isotopy. Similarly, in  $H_-$  there is an isotopy of  $B_-$  to  $B$  taking the disks  $D_-$  to disks  $D_2 \subset H_-$  such that  $D_2 \cap \Sigma = B$ . The tangle  $\mathcal{T}_3 = S \cap H_-$  is the image of  $C'_-$  under this isotopy. See Figure 2.

By construction  $D_1 \cap D_2 = B$ , so that  $F = D_1 \cup D_2$  is a spanning surface for the unlink  $\mathcal{T}_1 \cup \mathcal{T}_3$ . Note further that  $D_1$  is a trivial disk system for  $\mathcal{T}_1 \cup B$ , and  $D_2$  is a trivial disk system for  $B \cup \mathcal{T}_3$ ; hence,  $S$  is




Figure 2: Left: a doubly standard shadow diagram  $(A, B, C)$ ; the pairings  $(A, B)$  and  $(B, C)$  are standard. Middle: disks in  $\Sigma_+$  and  $\Sigma_-$  bounded by parallel copies of  $A \cup B$  and  $B \cup C$ , respectively. Right: a spanning surface  $F$  for  $\mathcal{T}_1 \cup \mathcal{T}_2$  in  $\partial X_3 = S^3$ .

the union of  $D_1$ ,  $D_2$ , and  $D_3$ , where  $D_3$  is a trivial disk system for  $\mathcal{T}_1 \cup \mathcal{T}_3$  pushed into  $B^4$ . However, since  $F = D_1 \cup D_2 \subset S^3$ , it follows that  $S$  is also isotopic to the cap-off  $\mathcal{F}$  of  $F$ , which is unknotted by [Proposition 3.2](#).  $\square$

We are now ready to prove our main result:

**Theorem 3.4** *If  $S$  is connected and  $e(S) = 0$ , then there is a procedure to produce a Seifert solid for  $S$  that takes as input a triplane diagram for  $S$ .*

**Proof** The proof follows from the proofs of [Propositions 3.5](#) and [3.6](#) below.  $\square$

In [Section 3.2](#), we show that there is a procedure to produce a Heegaard splitting for the Seifert solid when  $S$  is a 2-knot.

In addition to providing the proof of the above theorem, the next two propositions provide alternative proofs of the results in [\[6; 8\]](#) mentioned above.

**Proposition 3.5** *Every orientable surface-link  $S$  bounds a Seifert solid in  $S^4$ .*

**Proof** Let  $\mathbb{D}$  be a triplane diagram for  $S$ , with induced orientation on the bridge points  $\mathbf{x}$ . Perform mutual braid transpositions so that the bridge points alternate sign (orientation). Then there are  $b$  pairwise disjoint arcs  $\varepsilon$  contained in the equator  $e$  connecting bridge points of opposite signs, so that  $\mathbb{D}_i \cup \varepsilon$  is an oriented link diagram. Let  $F_i$  be the Seifert surface obtained by performing Seifert's procedure on the diagram  $\mathbb{D}_i \cup \varepsilon$ , and let  $\hat{F}_i = F_i \cup \bar{F}_{i+1}$  be the spanning surface obtained by gluing  $F_i$  to  $\bar{F}_{i+1}$  along  $\varepsilon$ . By [Proposition 3.2](#), there exists a handlebody  $V_i \subset X_i$  such that  $\partial V_i = \hat{F}_i \cup \mathcal{D}_i$  and  $V_i \cap \partial X_i = \hat{F}_i$ . Finally,  $Y = V_1 \cup V_2 \cup V_3$  is an embedded 3-manifold whose boundary is  $\mathcal{D}_1 \cup \mathcal{D}_2 \cup \mathcal{D}_3 = S$ , and so  $Y$  is a Seifert solid for  $S$ .  $\square$

**Proposition 3.6** *If  $S$  is connected and  $e(S) = 0$ , then  $S$  bounds a spanning solid in  $S^4$ .*

**Proof** Consider a bridge trisection  $\mathfrak{T}$  of  $S$ , with  $U_i = \partial \mathcal{D}_i$  and  $\tau = \mathcal{T}_1 \cup \mathcal{T}_2 \cup \mathcal{T}_3$ . By taking, for example, a triplane diagram  $\mathbb{D}$  and compatible checkerboard surfaces in  $\mathbb{D}_i$ , we can produce spanning surfaces  $\hat{F}_i$  for  $U_i$  such that  $\hat{F}_i \cap H_i = \hat{F}_{i-1} \cap H_i$ . Let  $F_i$  denote  $\hat{F}_i \cap H_i$ . For each component  $J$  of  $U_i = \partial \hat{F}_i$ , let  $\iota_{\hat{F}_i}(J)$  denote the induced boundary slope on the curve  $J$  by the surface  $\hat{F}_i$ . Then by [Proposition 3.2](#), we have

$$\sum_{J \subset U_1 \cup U_2 \cup U_3} \iota_{\hat{F}_i}(J) = 0.$$

Choose a triple of spanning surfaces  $\hat{F}_i$  such that  $\sum |\iota_{\hat{F}_i}(J)|$  is minimal over all possible choices. We claim that  $\sum |\iota_{\hat{F}_i}(J)| = 0$ . If not, then there exist boundary curves  $J_+$  and  $J_-$  such that  $\iota_{\hat{F}_i}(J_+) > 0$  and  $\iota_{\hat{F}_i}(J_-) < 0$ . Noting that the surface  $S$  contains all curves  $J \subset U_i \subset \tau$ , push each curve  $J \subset U_i$  slightly off of  $\tau$  into the corresponding disk component of  $\mathcal{D}_i$ , so that the collection of curves  $J$  is embedded in  $S$  and disjoint from  $\tau$ . Choose a path  $\gamma \subset S$  from  $J_+$  to  $J_-$ , avoiding the bridge points, noting that  $|\gamma \cap \tau| > 0$ .

At each point of  $\gamma \cap \tau$ , modify the corresponding component of  $F_i$  by taking the boundary connected sum of  $F_i$  with a trivial Möbius band to obtain new surfaces  $\hat{F}'_i$  and  $F'_i$ , so that the corresponding boundary curves satisfy  $\iota_{\hat{F}'}(J'_+) = \iota_{\hat{F}}(J_+) - 2$ ,  $\iota_{\hat{F}'}(J'_-) = \iota_{\hat{F}}(J_-) + 2$ , and  $\iota_{\hat{F}'}(J') = \iota_{\hat{F}}(J)$  for all other curves  $J'$ . It follows that  $\sum |\iota_{\hat{F}'}(J')| < \sum |\iota_{\hat{F}}(J)|$ , contradicting our assumption of minimality. (Note that  $\iota_{\hat{F}}(J)$  is always even, since it represents the number of intersection points between the boundary curves of spanning surfaces; see the proof of [Lemma 3.1](#).)

We conclude that  $\iota_{\hat{F}}(J) = 0$  for all curves  $J$ , and thus by [Proposition 3.2](#), each spanning surface  $\hat{F}_i$  cobounds a (possibly) nonorientable handlebody  $V_i \subset X_i$  with the disks  $\mathcal{D}_i$ . It follows that  $V_1 \cup V_2 \cup V_3$  is a spanning solid for  $\mathcal{S}$  in  $S^4$ .  $\square$

### 3.2 Procedure to find a Heegaard diagram for a Seifert solid

In this subsection, we describe a procedure for finding a Heegaard diagram for the Seifert solid coming from a bridge trisection  $\mathfrak{T}$  of a 2-knot  $\mathcal{S}$ . We use labels consistent with those appearing above in the proof of [Proposition 3.5](#). The process is illustrated in [Figures 3–6](#).

**Step 1** Given a triplane diagram  $\mathbb{D}$  for  $\mathcal{S}$ , perform interior Reidemeister moves and mutual braid transpositions so that the induced Seifert surfaces satisfy the following conditions:

- (a) Each of  $F_1$ ,  $F_2$ , and  $\hat{F}_1$  is a collection of disks.
- (b) Surfaces  $\hat{F}_2$  and  $\hat{F}_3$  are connected.
- (c)  $g(\hat{F}_2) = g(F_3)$ .

See [Figure 3](#). Note that attaining condition (a) is possible since any triplane diagram can be converted to one in which two of the tangles have no crossings. Condition (b) can be attained by performing interior Reidemeister moves on the diagram  $\mathbb{D}_3$ . Attaining condition (c) is possible since we can arrange so that  $F_2$  is a collection of  $b$  bridge disks, in which case  $\hat{F}_2$  deformation retracts onto  $F_3$  (although in general, we need not assume that  $F_2$  has  $b$  components, as shown below).

**Step 2** Following the proof of [Proposition 3.2](#), the surfaces  $\hat{F}_2$  and  $\hat{F}_3$  compress completely to disks in  $S^3$ . Let  $\alpha$  be a complete collection of pairwise disjoint compressing curves in  $\hat{F}_3$ , and let  $\beta$  be a complete collection of pairwise disjoint compressing curves in  $\hat{F}_2$ . See [Figure 4](#) (top row).

**Step 3** If necessary, slide the curves  $\beta$  over the components of  $\partial\mathcal{D}_2$  to obtain a collection of curves  $\beta' \subset F_3$ . Note that since  $g(F_3) = g(\hat{F}_2)$ , as curves in  $\mathcal{F}_2 = \hat{F}_2 \cup \mathcal{D}_2$ , the collection  $\beta$  can be isotoped to be contained in  $F_3$ , and any isotopy of a curve over a disk component of  $\mathcal{D}_2$  can be realized as a slide over  $\partial\mathcal{D}_2$ . Thus, such a sequence of slides exists. See [Figure 4](#) (middle row).

**Step 4** Let  $P = \mathcal{D}_1 \cup \mathcal{D}_2$ , so that  $P$  is a planar surface with  $c_3$  boundary components, let  $Q$  be the surface obtained by gluing  $P$  to  $\hat{F}_3$  along their boundaries, and let  $\alpha^*$  be a choice of  $c_3 - 1$  boundary components of  $P$  and some minimal number of curves in  $\alpha$  such that  $\alpha^*$  forms a cut system for  $Q$ .



Figure 3: To perform the Seifert solid procedure on a triplane diagram, we first perform mutual braid transposition until the tangle diagrams in  $V_1$  and  $V_2$  have no crossings. Then we perform the usual Seifert's procedure for knot diagrams to obtain surfaces  $F_1$ ,  $F_2$ , and  $F_3$  that agree in the bridge sphere  $\Sigma$ , with  $F_1$ ,  $F_2$ , and  $\hat{F}_1$  all collections of disks and  $g(\hat{F}_2) = g(F_3)$ .

**Step 5** Let  $\beta^*$  be the union of  $\beta'$  and a collection of curves in  $Q$  obtained by the following instructions: For each component of  $J$  of  $\partial\mathcal{D}_1$ , suppose that  $J$  meets  $d$  disk components of  $F_2$ . Choose  $d - 1$  of these components, isotope them off of  $F_2$  in  $\mathcal{F}_2 = F_2 \cup F_3 \cup \mathcal{D}_2$ , and add these  $d - 1$  curves to  $\beta^*$ . Discard any superfluous curves of  $\beta'$  so that  $\beta^*$  is a cut system for  $Q$ .

**Proposition 3.7** Using the procedure described above,  $\mathcal{S}$  bounds a punctured copy of the 3-manifold determined by the Heegaard diagram  $(Q; \alpha^*, \beta^*)$ .

**Proof** Suppose that  $\mathbb{D}$  is a triplane diagram satisfying conditions (a), (b), and (c) given in Step 1 above. Following the proofs of Propositions 3.2 and 3.5, for each  $i$ , the surface  $\hat{F}_i \cup \mathcal{D}_i$  bounds a handlebody  $V_i$ , where  $V_1$  is a collection of 3-balls, say  $B_1, \dots, B_n$ , and  $V_2$  and  $V_3$  are connected. Moreover,  $\alpha$  contains a




Figure 4: Top: we find complete sets of compressing curves  $\alpha$  and  $\beta$  for  $\hat{F}_3$  and  $\hat{F}_2$ , respectively. Middle: we slide  $\alpha$  and  $\beta$  (with slides indicated in top row) over  $\partial\hat{F}_3$  and  $\partial\hat{F}_2$  to obtain curve systems  $\alpha'$  and  $\beta'$  that are each completely within  $F_3$ . Bottom: We obtain  $\alpha^*$  (red and purple curves) by adding boundary curves as in Step (4) of Section 3.2. We obtain  $\beta^*$  by adding arcs as in Step (5). Then  $(Q; \alpha^*, \beta^*)$  is a Heegaard diagram for a (closure of a) Seifert solid for the 2-knot described by the initial triplane diagram.

cut system for  $V_3$  and  $\beta$  contains a cut system for  $V_2$ . Since  $\beta'$  is homotopic to  $\beta$  in  $\partial V_2$ , it follows that  $\beta'$  also contains a cut system for  $V_2$ . Thus, the Seifert solid bounded by  $\mathcal{S}$  is equal to  $V_2 \cup V_3 \cup B_1 \cup \dots \cup B_n$ . Let  $Y$  be the closed 3-manifold obtained by capping off the boundary  $\mathcal{S}$  of this Seifert solid with an abstract 3-ball  $B_0$ . We will show that  $(Q; \alpha^*, \beta^*)$  is a Heegaard diagram for  $Y$ .




Figure 5: We start performing the Seifert solid procedure (Section 3.2) on the triplane diagram in the top row.

To this end, consider  $W = V_3 \cup B_0$  and  $W' = V_2 \cup B_1 \cup \cdots \cup B_n$ . Considering that  $\partial V_2 = F_2 \cup F_3 \cup \mathcal{D}_2$  and  $\partial(B_1 \cup \cdots \cup B_n) = F_1 \cup F_2 \cup \mathcal{D}_1$ , we have that

$$\partial W' = F_3 \cup F_1 \cup \mathcal{D}_2 \cup \mathcal{D}_1 = \hat{F}_3 \cup P = Q.$$

Additionally, the 3-balls  $B_i$  are attached to  $V_2$  along  $F_2$ , which is a collection of disks by condition (a). It follows that the curves  $\beta' \cup \partial F_2$  bound compressing disks in  $W'$  cutting  $W'$  into a collection of 3-balls, so  $W'$  is a handlebody. In addition, choosing all but one curve of  $\partial F_2$  for each component  $B_i$  and a subset of  $\beta'$  as in Step 5 above yields a cut system  $\beta^*$  for  $W'$ .

Turning our attention to  $W$ , we have  $\partial V_3 = \hat{F}_3 \cup \mathcal{D}_3$  and  $\partial B_0 = \mathcal{D}_1 \cup \mathcal{D}_2 \cup \mathcal{D}_3$ , so  $\partial W = \hat{F}_3 \cup \mathcal{D}_1 \cup \mathcal{D}_2 = Q$ , and in addition, the curves  $\alpha$  and  $\partial \mathcal{D}_3$  bound disks cutting  $W$  into 3-balls. Choosing  $\alpha^*$  to contain all but one curve of  $\partial \mathcal{D}_3$  and a subset of  $\alpha$  as in Step 4, the curves in  $\alpha^*$  bound disks cutting  $W$  into a single 3-ball, so  $\alpha^*$  is a cut system for  $W$ . We conclude that  $(Q; \alpha^*, \beta^*)$  is a Heegaard diagram for  $Y$ , as desired.  $\square$

**Remark 3.8** It may be the case that the surface  $F_3$  compresses in  $H_3$ , in which case  $\alpha$  and  $\beta$  could have one or more curves in  $F_3$  in common. Following the procedure with such  $\alpha$  and  $\beta$  produces one or more extra  $S^1 \times S^2$  summands for the 3-manifold  $Y$ , and a simpler Seifert solid can be obtained by first compressing  $F_3$  maximally in  $H_3$ .




Figure 6: Left: the curves  $\alpha'$  in  $F_3$ . Center left: the curves  $\beta'$  in  $F_2$ . Center right: we add some boundary curves of  $F_2$  to  $\alpha$  to obtain  $\alpha^*$  and some arc to  $\beta'$  to obtain  $\beta^*$ . Right: we simplify the resulting Heegaard diagram  $(\Sigma; \alpha^*, \beta^*)$  to see that it is a diagram of  $S^3$ . Thus, the initial 2-knot bounds a copy of  $B^3$  in  $S^4$ , so is unknotted.

**Remark 3.9** The procedure above can be generalized: We can relax conditions (a), (b), and (c) from Step 1; the only assumption necessary to ensure that  $V_1 \cup V_2$  is a handlebody is that their intersection  $F_2$  is a collection of disks. However, the weaker conditions make it somewhat more difficult to draw the diagram, since we are no longer guaranteed the existence of the slides of Step 3 — it may be the case that  $\beta$  curves necessarily intersect the disks  $\mathcal{D}_1$  and  $\mathcal{D}_2$ .

**Remark 3.10** The observant reader might notice that we call our process the *Seifert solid procedure*, rather than *algorithm*. An algorithm gives an output completely determined from the input, independent of further choices. A procedure may require additional choices for the output to be determined. In the procedure we give in this section to find a description of a Seifert solid for a 2-knot, we are forced to choose compressing circles for surfaces in  $S^3$ . These circles are generally not unique (and in fact, different choices can determine different Seifert solids), so we do not refer to this procedure as an algorithm.

### 3.3 Some examples

In this subsection, we carry out the procedure described above for a couple of specific examples. The first is the spun trefoil. In Figure 3, we see a triplane diagram for the spun trefoil coming from [15], followed by the result of performing triplane moves so that the induced Seifert surfaces  $F_i$  satisfy conditions (a), (b), and (c) from Step 1 above.

In the top row of Figure 4, we find the compressing curves  $\alpha$  on  $\hat{F}_3$  and  $\beta$  on  $\hat{F}_2$ . Note that in this case  $\mathcal{D}_3$  contains two disks, so that  $P = \mathcal{D}_1 \cup \mathcal{D}_2$  is an annulus, and  $Q = \hat{F}_3 \cup P$  can be obtained by identifying the two boundary components of  $\hat{F}_3$ . Under this identification, the identified boundary components constitute the third curve in the cut system  $\alpha^*$ . In the second row at left, we slide the two curves of  $\alpha$  over the third curve of  $\alpha^*$  in  $Q$ . In the second row at right, we slide the two curves of  $\beta$  over a boundary component as shown to get the curves  $\beta' \subset F_3$  (which are identical to the image of  $\alpha$  under the slides described above). Finally, the third curve of  $\beta^*$  consists of the teal arc depicted in  $F_3$  and a spanning arc

in the annulus  $A$ , or equivalently, we can identify the endpoints of the teal arc. In the lower row, we see the diagram for the Seifert solid, the standard (once-stabilized) Heegaard diagram for  $\#^2(S^1 \times S^2)$ .

**Remark 3.11** These diagrams and arguments easily generalize to produce the Seifert solid  $\#^{p-1}(S^1 \times S^2)$  for the spun  $(p, 2)$ -torus knot. Miyazaki proved that the degree of the Alexander polynomial (over  $\mathbb{Q}[t, t^{-1}]$ ) is a lower bound for the second Betti number of any Seifert solid [17]. Since the degree of the Alexander polynomial of  $T(2, p)$  is  $p - 1$ , these solids are minimal in the sense that the corresponding 2-knots cannot bound any 3-manifold with a smaller second Betti number, eg fewer  $S^1 \times S^2$  summands.

For the second example, we find a Seifert solid for the 1-twist spun trefoil (which is unknotted by [23]). In Figure 5, we include a simplified triplane diagram for the 1-twist spun trefoil along with the surfaces  $\hat{F}_2$  and  $\hat{F}_3$  this diagram generates.

Next, we find the compressing curves  $\alpha$  for  $\hat{F}_3$  and  $\beta$  for  $\hat{F}_2$ . As in the spun trefoil example above,  $P = \mathcal{D}_1 \cup \mathcal{D}_2$  is an annulus, so we view  $Q$  as being obtained by identifying the two boundary components of  $\hat{F}_3$ , with this identified boundary the third curve in  $\alpha^*$ . Figure 6 shows the curves  $\alpha$ ,  $\beta$ , and the union of the sets in  $Q$ , yielding the standard diagram for  $S^3$ , in which the third curve of  $\beta^*$  appears as a teal arc with boundary points identified (as above). Note that the existence of the curves  $\alpha$  and  $\beta$  is guaranteed by Proposition 3.2; in practice, however, these curves are found using ad hoc methods.

## 4 Spinal Seifert solids

A natural aspect of the study of Seifert surfaces for links in the 3-sphere is the consideration of their exteriors. We call a Seifert surface  $F$  for  $L$  *canonical* if it is isotopic to a surface obtained by applying Seifert's procedure to a diagram for  $L$ . We call a Seifert surface  $F$  *free* if its exterior  $S^3 \setminus \nu(F)$  is a 3-dimensional handlebody — equivalently, has free fundamental group. It is an easy exercise to see that a canonical Seifert surface is free, provided that it is connected; so every link admits a free Seifert surface, by the application of Seifert's algorithm to a nonsplit diagram. However, such a surface can be far from minimal genus. M Kobayashi and T Kobayashi showed that the difference between the genus of a knot and the minimal genus of a free Seifert surface for the knot can be arbitrarily large, and that moreover the difference between the minimal genus of a free Seifert surface for a knot and the minimal genus of a canonical Seifert surface can also be arbitrarily large [11]. (In fact, they show that both of these differences can be made arbitrarily large at the same time.)

In this section, we introduce 4-dimensional analogs of the notions of canonical and free Seifert surfaces. Going forward, let  $S \subset S^4$  be a surface-link admitting a Seifert solid. (This is equivalent to the condition that  $S$  be orientable or have normal Euler number zero.) We call a Seifert solid  $Y$  *canonical* if it is isotopic to a Seifert solid obtained by the procedure given in Section 3.1 (see Propositions 3.5 and 3.6). We call a Seifert solid  $Y$  *spinal* if  $S^4 \setminus \nu(Y)$  deformation retracts onto a finite 2-complex. Equivalently,  $S^4 \setminus \nu(Y)$  can be built with handles of index at most two.



**Theorem 4.1** *If a surface-knot  $\mathcal{S}$  admits a Seifert solid, then it admits a canonical Seifert solid that is spinal.*

**Proof** First, note that in the proof of Propositions 3.5 and 3.6, it is possible to arrange that each Seifert surface  $F_i$  is connected: for example, this is assured if each  $\mathbb{D}_i \cup \varepsilon$  is nonsplit. Let  $Y$  be a canonical Seifert solid for  $\mathcal{S}$  given by Proposition 3.5 or Proposition 3.6 such that the canonical surface  $F_i = Y \cap H_i$  is connected for each  $i \in \mathbb{Z}_3$ . We make use of the notation of the proof of Proposition 3.5 in what follows.

Recall that  $V_i = X_i \cap Y$  is a handlebody with  $\partial V_i = \hat{F}_i \cup \mathcal{D}_i$ . Moreover,  $V_i$  is built relative to  $\hat{F}_i$  by attaching 3-dimensional 2- and 3-handles. It follows that  $X_i \setminus \nu(V_i)$  can be built with 4-dimensional 0-, 1-, and 2-handles.

Next, recall that  $F_i$  is a canonical Seifert surface for the link  $\mathbb{D}_i \cup \varepsilon$ , considered in  $S^3 = H_i \cup_\Sigma B^3$ . Since we have assumed  $F_i$  is connected,  $F_i$  is free in  $H_i \cup_\Sigma B^3$ . Since  $\varepsilon \subset \partial H_i$ , it follows that  $H_i \setminus F_i$  is also a 3-dimensional handlebody.

Finally, we can build  $S^4 \setminus \nu(Y)$  by taking the  $X_i \setminus \nu(V_i)$  and gluing them along the  $H_i \setminus \nu(F_i)$ . Since the three gluings occur along 3-dimensional handlebodies,  $S^4 \setminus \nu(Y)$  is obtained from the disjoint union of the  $X_i \setminus \nu(V_i)$  by attaching 4-dimensional 1- and 2-handles. Because each of the  $X_i \setminus \nu(V_i)$  were built with 4-dimensional handles of index at most two, the same is true for  $S^4 \setminus \nu(Y)$ . This shows that  $Y$  is spinal, as desired.  $\square$

When studying Seifert surfaces, the genus of the surface is the obvious measure of complexity that one might try to minimize. In contrast, there are many ways one might try to quantify the complexity of a Seifert solid  $Y$  for a surface-knot; indeed, any complexity one might associate to a 3-manifold could be interesting to consider. Here we content ourselves to give some examples showing that there is at least one sense in which a simple Seifert solid for a surface-knot can be arbitrarily far from being spinal.

**Theorem 4.2** *Given any  $n \in \mathbb{N}$ , there exists a 2-knot  $\mathcal{K}$  that bounds a Seifert solid  $Y$  homeomorphic to  $(S^1 \times S^2)^\circ$  such that  $S^4 \setminus \nu(Y)$  requires at least  $n$  4-dimensional 3-handles.*

**Proof** Let  $J$  be an arbitrary knot, and let  $K = \text{Wh}_0(J \# \bar{J})$  be the untwisted Whitehead double of the connected sum of  $J$  with its mirror. Let  $F$  be the standard genus one Seifert surface for  $K$ , and let  $\gamma$  be the curve on  $F$  that is isotopic to  $J \# \bar{J}$ . (Alternatively,  $F$  is obtained by taking a 0-framed annular thickening of a curve  $\gamma$  isotopic to  $J \# \bar{J}$  and plumbing on a Hopf band.)

Let  $E$  be the standard ribbon disk for  $\gamma$ , so that  $(B^4, E) = (S^3, J)^\circ \times I$ . The surface  $F$  can be surgered along  $E$  in the 4-ball to get a slice disk  $D$  for  $K$ , and the trace of this surgery yields a solid torus  $V$  with  $\partial V = F \cup D$ .

Let  $\mathcal{K} = D \cup_K \bar{D}$  be the 2-knot obtained by doubling  $D$ , and let  $Y = V \cup_F \bar{V}$  be the double of  $V$  along  $F$ . Then  $Y$  is a Seifert solid for  $\mathcal{K}$  and  $Y \cong (S^1 \times S^2)^\circ$ .

We claim that  $\pi_1(S^4 \setminus \nu(Y)) \cong \pi_1(S^3 \setminus \nu(J))$ . First,  $\pi_1(S^4 \setminus \nu(Y)) \cong \pi_1(B^4 \setminus V)$ , since the former exterior is the double of the latter exterior along the exterior of  $F$  in  $S^3$  and  $\pi_1(S^3 \setminus \nu(F))$  surjects onto  $\pi_1(B^4 \setminus V)$  under inclusion. Next, by construction,  $V$  is obtained by thickening the slice disk  $E$  and attaching a trivial 3-dimensional 1-handle. It follows that

$$\pi_1(B^4 \setminus \nu(V)) \cong \pi_1(B^4 \setminus \nu(E)) \cong \pi_1(S^3 \setminus \nu(J)),$$

as desired.

To complete the proof, let  $n \in \mathbb{N}$  be given, and choose  $J$  to be any knot with  $\text{rank}(\pi_1(S^3 \setminus \nu(J))) \geq n + 2$  (eg take  $J$  to be a connected sum of  $n + 1$  trefoils [21]). The exterior  $S^4 \setminus \nu(Y)$  can be built relative to  $\partial(S^4 \setminus \nu(Y)) \cong (S^1 \times S^2) \# (S^1 \times S^2)$  with some number of 4-dimensional 1-, 2-, 3-, and 4-handles. Since the 1-handles correspond to generators of the fundamental group, at least  $n$  are required; the boundary  $\partial(S^4 \setminus \nu(Y))$  contributes only two to the rank of the fundamental group. Similarly, since we can obtain another presentation of  $\pi_1(S^4 \setminus \nu(Y))$  with generators corresponding to 3-handles, the number of 3-handles in this decomposition is at least  $n + 2$ .  $\square$

We note that the construction of  $\mathcal{K}$  given in the above proof is closely related to an interesting construction of 2-knots given by Cochran [3].

Next, we observe that many important examples of Seifert solids are, in fact, spinal:

- (1) Every ribbon 2-knot bounds a Seifert solid  $Y$  that is homeomorphic to  $(\#^m(S^1 \times S^2))^\circ$  for some  $m$  [22]. The manifold  $Y$  is obtained by taking a Seifert surface  $F$  for some ribbon knot in an equatorial  $S^3$ , thickening it, and attaching trivial 2-handles above and below the equator. By attaching tubes to  $F$  (at the cost of increasing  $m$ ), we can arrange for  $F$  to be free. Then  $Y$  is spinal.
- (2) If  $\mathcal{K}$  is fibered with fiber  $Y$ , then  $S^4 \setminus \nu(Y) \cong Y \times I$  is spinal, since  $Y$  is a punctured 3-manifold.
- (3) Connected Seifert solids arising from broken surface diagrams via the construction given by Carter and Saito [2] are spinal. Recall that a connected canonical Seifert surface is free because it deformation retracts to a graph so that on each edge, there is one local maximum and no local minima with respect to the radial height function on  $S^3$ . (Here the vertices of the graph correspond to the disks produced in Seifert's procedure while the edges correspond to the half-twisted bands.) This ensures that the exterior of a canonical surface can be built with 0- and 1-handles. Similarly, a Seifert solid constructed à la [2] deformation retracts to a 2-complex with one local maximum and no other critical points in the interior of each 1- and 2-cell. Thus, the exterior of such a Seifert solid can be built with 0-, 1-, and 2-handles.

Finally, we can formulate a question analogous to the 3-dimensional results in [11] in the setting of surface-knots.

**Question 4.3** Define the **genus** of an orientable surface-knot  $S$  in  $S^4$  to be the minimal first Betti number of any Seifert solid bounded by  $S$ , and define the **spinal genus** and **canonical genus** similarly, using spinal Seifert solids and canonical Seifert solids, respectively. Do there exist surface-knots for which these three measures of complexity differ?

We remark that using techniques as in the proof of [Theorem 4.2](#), one can show that for some of the known classical knots  $K$  whose genus and free genus are sufficiently different (see [\[18\]](#), for example), the spun knots  $\mathcal{S}(K)$  admit low-complexity nonspinal Seifert solids, whereas the obvious spinal and canonical Seifert solids have greater complexity. However, it is likely to be considerably more difficult to obstruct the existence of low-complexity spinal or canonical Seifert solids, even for these examples.

## 5 On standardness of bridge trisections

The goal of this section is to prove [Theorem 5.2](#), which states that a  $(b; c_1, c_2, c_3)$ -bridge trisection that satisfies  $c_i \geq b - 1$  for some  $i \in \mathbb{Z}_3$  can be completely decomposed into standard pieces. This proves Conjecture 4.3 of [\[15\]](#), and the theorem can be viewed as the bridge trisection analog of the main result in [\[13\]](#), which states that every  $(g; k_1, k_2, k_3)$ -trisection with  $k_i \geq g - 1$  for some  $i$  is standard in that it decomposes into genus one summands.

We encourage the reader to recall the notions of *perturbation* and *connected summation* for bridge trisections. The former was first introduced in [\[15, Section 6\]](#), where it was referred to as stabilization, and the latter can be reviewed in [\[15, Subsection 2.2\]](#). See also [\[14, Section 3\]](#) for a succinct description of these concepts.

We call a surface-link an *unlink* if it is the split union of unknotted surface-knots, though we allow the topology of each component to vary. For example, one might have a 2-component unlink that is the split union of an unknotted 2-sphere and an unknotted projective plane. (See [\[14, Subsection 2.2\]](#) and [Section 2.3](#) above for a brief discussion of unknotted surface-knots.)

Before proving [Theorem 5.2](#) in generality, we recall the case in which  $c_i = b$  for some  $i \in \mathbb{Z}_3$ . This was addressed as [\[15, Proposition 4.1\]](#). A bridge trisection is called *completely decomposable* if it is a disjoint union of perturbations of one-bridge and two-bridge trisections.

**Proposition 5.1** [\[15, Proposition 4.1\]](#) *Let  $\mathfrak{T}$  be a  $(b; c_1, c_2, c_3)$ -bridge trisection with  $c_i = b$  for some  $i \in \mathbb{Z}_3$ . Then  $\mathfrak{T}$  is completely decomposable, and the underlying surface-link is the unlink of  $\min_i \{c_i\}$  2-spheres.*

Note that if  $c_i = b$  for some  $i \in \mathbb{Z}_3$ , then  $c_{i-1} = c_{i+1}$ . Similarly, in what follows we will see that if  $c_i = b - 1$  for some  $i \in \mathbb{Z}_3$ , then  $|c_{i-1} - c_{i+1}| \leq 1$ . We now present and prove the main result of this section:

**Theorem 5.2** *Let  $\mathfrak{T}$  be a  $(b; c_1, c_2, c_3)$ -bridge trisection with  $c_i = b - 1$  for some  $i \in \mathbb{Z}_3$ . Then  $\mathfrak{T}$  is completely decomposable, and the underlying surface-link is either the unlink of  $\min\{c_i\}$  2-spheres or the unlink of  $\min\{c_i\}$  2-spheres and one projective plane, depending on whether  $|c_{i-1} - c_{i+1}| = 1$  or  $c_{i-1} = c_{i+1}$ .*

The key ingredient in the proof of the theorem is a pair of results of Bleiler and Scharlemann about planar surfaces in 3-manifolds [1; 19]. We refer the reader to Section 1 of each of these papers, as we will adopt the notation of [1, Theorem 1.3; 19, Theorem 1.1] in the proof below.

**Proof of Theorem 5.2** We induct on the bridge number  $b$  of the bridge trisection. When  $b = 1$  or  $b = 2$ , there is an easy classification of  $b$ -bridge trisections [15, Subsection 4.3], which we take as the base case. Assume the theorem holds when the bridge number is less than  $b$ , and let  $\mathfrak{T}$  be a  $(b; c_1, c_2, c_3)$ -bridge trisection. Assume without loss of generality that  $c_3 = b - 1$ .

Suppose that  $\mathcal{T}_1$ ,  $\mathcal{T}_2$ , and  $\mathcal{T}_3$  are the three tangles comprising the spine of the bridge trisection. Every  $b$ -bridge splitting of a  $c$ -component unlink with  $b > c$  is a perturbation of the standard  $c$ -bridge splitting of the  $c$ -component unlink, which is itself unique up to isotopy [15, Proposition 2.3]. It follows that there exist collections  $\Delta_1$  and  $\Delta_3$  of bridge disks for  $\mathcal{T}_1$  and  $\mathcal{T}_3$ , respectively, such that the shadows  $\Delta_1^* = \Delta_1 \cap \Sigma$  and  $\Delta_3^* = \Delta_3 \cap \Sigma$  have the property that  $\Delta_1^* \cup \Delta_3^*$  is an embedded collection of  $b - 2$  bigons and a single quadrilateral. Let  $\alpha_0^*$  denote one of the arcs of  $\Delta_1^*$  in the quadrilateral.

Let  $L = \mathcal{T}_2 \cup \overline{\mathcal{T}_3}$ , and let  $\mathfrak{b}$  be the band for  $L$  that is framed by  $\Sigma$  and whose core is  $\alpha_0^*$ . Then the data  $(\Sigma, L, \mathfrak{b})$  encodes a banded  $b$ -bridge splitting, since the resolution  $L_{\mathfrak{b}}$  is the unlink  $L' = \mathcal{T}_2 \cup \overline{\mathcal{T}_1}$ . (Here, we think of  $\mathfrak{b}$  as being slightly perturbed to lie in the 3-ball containing  $\mathcal{T}_3$ .) We refer the reader to Section 3 of [15], especially Lemma 3.3, for more details about banded bridge splittings and how they arise from bridge trisections.

Assume without loss of generality that  $c_2 = |L|$  is greater than or equal to  $c_1 = |L'|$ . We break the remainder of the proof into two cases: Either  $c_2 > c_1$  or  $c_2 = c_1$ . Note that since there is only one band present, we must have  $c_2 - c_1 \leq 1$ . The proofs of the two cases are very similar, except that we apply [19, Theorem 1.1] in the first case and [1, Theorem 1.3] in the second.

**Case 1** If  $c_2 = c_1 + 1$ , then  $\mathfrak{b}$  connects distinct components  $K_1$  and  $K_2$  of  $L$ . Let  $K'$  denote the component of  $L'$  obtained as the resolution  $(K_1 \cup K_2)_{\mathfrak{b}}$ . We now translate this setup into the notation of [19, Section 1]. Let  $N = \overline{\nu(K_1 \cup \mathfrak{b} \cup K_2)}$ , a genus two handlebody, and let  $M = S^3 \setminus \nu(L \setminus (K_1 \sqcup K_2))$ . Let  $E_1$  denote the spanning disk bounded by  $K_1$ . Let  $P' = \partial \nu(E_1)$ , a 2-sphere disjoint from  $K_1 \sqcup K_2$  in  $M$ . Let  $Q'$  denote a spanning disk bounded by  $K'$  in  $M$ . Let  $P = \overline{P' \setminus N}$ , and let  $Q = \overline{Q' \setminus N}$ .

It is clear from this setup that  $P \cap \partial N$  is a collection of  $m$  parallel separating curves  $A_m$  for some odd  $m$ , since  $P'$  was disjoint from  $K_1$  and  $K_2$ , but intersects  $\mathfrak{b}$  transversely; see [19, Figure 1]. Similarly,  $Q \cap \partial N$  agrees with the curves  $B_n$ , since  $\partial Q' = K'$  and  $Q'$  may crash through  $\mathfrak{b}$  in arcs parallel to its core. Thus,  $M$ ,  $N$ ,  $P$ , and  $Q$  satisfy the hypotheses of [19, Theorem 1.1]. The relevant conclusion is that  $A_1$  and  $B_0$  bound embedded disks  $E$  and  $F$  in  $\overline{M \setminus N}$  that intersect in a single arc; compare with the proof of [19, Main Theorem].

Translating this conclusion back into the setting of interest, the disk  $E$  is properly embedded in  $S^3 \setminus \nu(\mathfrak{b})$  and  $F$  is a spanning disk for  $K'$ . This implies that the pair  $(B^3, T) = (S^3, L) \setminus (\nu(\mathfrak{b}), \nu(L \cap \mathfrak{b}))$  is the

split union of a trivial tangle and an unlink: the strands of the trivial tangle are parallel into pushoffs of  $E$  via the components of  $F \setminus \nu(E)$ , at which point they are parallel into  $\partial\nu(\mathfrak{b})$  via the pushoffs of  $E$ .

The bridge sphere  $\Sigma$  induces a bridge splitting  $(B^3, T)$ . By [24, Theorem 2.2],  $\Sigma$  is either minimal for  $(B^3, T)$  or perturbed.<sup>1</sup> If the splitting were minimal, we would have  $b = c_2$ , so  $\mathfrak{T}$  would be completely decomposable by Proposition 5.1. If the splitting is perturbed, then  $\mathfrak{T}$  is perturbed, since each bridge arc of  $\mathcal{T}_3$  that is disjoint from  $\nu(\mathfrak{b})$  is a strand of a 1-bridge splitting of a component of  $L_3 = \mathcal{T}_3 \cup \overline{\mathcal{T}}_1$ . After deperturbing  $\mathfrak{T}$ , we find that  $\mathfrak{T}$  is completely decomposable, by the inductive hypothesis.

**Case 2** If  $c_2 = c_1$ , then  $\mathfrak{b}$  connects a component  $K$  of  $L$  to itself. Let  $K' = K_{\mathfrak{b}}$ . We now translate this setup into the notation of [1, Section 1], abbreviating the discourse where it is overly repetitive of the previous case. Let  $M = S^3 \setminus \nu(L \setminus K)$ , and let  $N = \nu(K \cup \mathfrak{b})$ . Let  $P'$  be a spanning disk bounded by  $K$  in  $M$ , and let  $Q'$  be a spanning disk bounded by  $K'$  in  $M$ . Let  $P = \overline{P'} \setminus \overline{N}$ , and let  $Q = \overline{Q'} \setminus \overline{N}$ .

It is clear from the setup that the hypotheses of [1, Theorem 1.3] are satisfied, so we can conclude that some  $A_0$  and  $B_0$  bound embedded disks  $E_P$  and  $E_Q$ , respectively, in  $\overline{M \setminus N}$ . Moreover, there is a properly embedded disk  $D$  in  $\overline{M \setminus N}$ , disjoint from  $E_P$  and  $E_Q$ , that runs once over one of the handles of  $N$  and is disjoint from the other handle. We can extend  $E_P$  to a spanning disk  $F$  for  $K$ ; compare with the proof of [1, Theorem 1.8].

The strands of  $K \setminus \nu(\mathfrak{b})$  are parallel into pushoffs of  $D$  via the components of  $E_P \setminus \nu(D)$ , at which point they are parallel into  $\partial\nu(\mathfrak{b})$  via the pushoffs of  $D$ . It follows that the tangle  $(B^3, T) = (S^3, L) \setminus (\nu(\mathfrak{b}), \nu(\mathfrak{b} \cap K))$  is the split union of a trivial tangle and an unlink, and  $\Sigma$  gives rise to a bridge splitting of  $(B^3, T)$ . As before, this splitting is either minimal or perturbed. The case that the splitting is perturbed has the same consequence as in Case 1 above.

If the splitting is minimal, then it is a split union of a 2-bridge splitting of the trivial tangle and a  $(b-2)$ -bridge splitting of an unlink. It follows that the bridge trisection is a split union:  $\mathfrak{T} = \mathfrak{T}' \sqcup \mathfrak{T}''$ , where  $\mathfrak{T}'$  is a  $(2, 1)$ -bridge trisection (of a projective plane, necessarily), and  $\mathfrak{T}''$  is a  $(b-2; c_1-1, c_2-1, b-2)$ -bridge trisection (of an unlink of 2-spheres, necessarily). The latter is completely decomposable by Proposition 5.1.  $\square$

We can also use Theorem 5.2 to understand surface-links with particular banded link presentations, where a *banded link presentation*  $(L, \nu)$  consists of an unlink  $L \subset S^3$  and a collection of bands  $\nu$  such that the resolution  $L_\nu$  of  $L$  along  $\nu$  is also an unlink. Every banded link presentation gives rise to a surface  $\mathcal{S}$  in  $S^4$ , and conversely, every surface-link  $\mathcal{S}$  in  $S^4$  can be presented by a banded link [10].

In [15, Section 3], the authors introduced the notion of *banded bridge splitting* of  $(L, \nu)$ , a bridge splitting of  $L$  such that the bands  $\nu$  are isotopic into the bridge sphere with the surface framing and are dual to a collection of bridge disks on one side. They showed that  $(S^4, \mathcal{S})$  admits a  $(b; \mathbf{c})$ -bridge trisection if

<sup>1</sup>Although [24, Theorem 2.2], as stated, applies to a closed 3-manifold  $M$  and a link  $K$  in  $M$ , a verbatim proof establishes the more general case where the 3-manifold  $M$  is replaced by a punctured 3-manifold and the link  $K$  is a tangle.

and only if a banded link presentation  $(L, v)$  of  $\mathcal{S}$  admits a banded  $b$ -bridge splitting such that  $|L| = c_1$ ,  $|v| = b - c_2$ , and  $|L_v| = c_3$ . As a corollary to [Theorem 5.2](#), we obtain the following, which states, in essence, that a surface is unknotted if the bands are attached in a relatively simple way to the maxima or minima disks.

**Corollary 5.3** *Suppose a surface-link  $\mathcal{S}$  in  $S^4$  is presented by a banded link  $(L, v)$  with a banded  $b$ -bridge splitting such that  $b = |L| + 1$  or  $b = |L_v| + 1$ . Then  $\mathcal{S}$  is an unlink of 2-spheres or an unlink of 2-spheres and an unknotted projective plane.*

The corollary exploits a feature of trisection theory called *handle triality*: If  $(L, v)$  admits a banded bridge splitting as in the corollary, then it admits a  $(b, c)$ -bridge trisection such that  $c_1 = b - 1$  or  $c_3 = b - 1$ . By the three-fold symmetry of the trisection setup, we can extract a different banded link presentation with a single band, as in the proof of [Theorem 5.2](#), and now we rely on known results about surface-links built with a single band to classify  $\mathcal{S}$ . The result can be interpreted as an analog for knotted surfaces of [\[13, Theorem 1.2\]](#).

## References

- [1] **S Bleiler, M Scharlemann**, *A projective plane in  $\mathbb{R}^4$  with three critical points is standard: strongly invertible knots have property P*, *Topology* 27 (1988) 519–540 [MR](#) [Zbl](#)
- [2] **J S Carter, M Saito**, *A Seifert algorithm for knotted surfaces*, *Topology* 36 (1997) 179–201 [MR](#) [Zbl](#)
- [3] **T Cochran**, *Ribbon knots in  $S^4$* , *J. Lond. Math. Soc.* 28 (1983) 563–576 [MR](#) [Zbl](#)
- [4] **I Dai, M Miller**, *The 0-concordance monoid admits an infinite linearly independent set*, *Proc. Amer. Math. Soc.* 151 (2023) 3601–3609 [MR](#) [Zbl](#)
- [5] **D Gay, R Kirby**, *Trisecting 4-manifolds*, *Geom. Topol.* 20 (2016) 3097–3132 [MR](#) [Zbl](#)
- [6] **H Gluck**, *The embedding of two-spheres in the four-sphere*, *Trans. Amer. Math. Soc.* 104 (1962) 308–333 [MR](#) [Zbl](#)
- [7] **C M Gordon**, *Knots in the 4-sphere*, *Comment. Math. Helv.* 51 (1976) 585–596 [MR](#) [Zbl](#)
- [8] **C M Gordon, R A Litherland**, *On the signature of a link*, *Invent. Math.* 47 (1978) 53–69 [MR](#) [Zbl](#)
- [9] **J Joseph, J Meier, M Miller, A Zupan**, *Bridge trisections and classical knotted surface theory*, *Pacific J. Math.* 319 (2022) 343–369 [MR](#) [Zbl](#)
- [10] **A Kawauchi, T Shibuya, S Suzuki**, *Descriptions on surfaces in four-space, I: Normal forms*, *Math. Sem. Notes Kobe Univ.* 10 (1982) 75–125 [MR](#) [Zbl](#)
- [11] **M Kobayashi, T Kobayashi**, *On canonical genus and free genus of knot*, *J. Knot Theory Ramifications* 5 (1996) 77–85 [MR](#) [Zbl](#)
- [12] **C Livingston**, *Surfaces bounding the unlink*, *Michigan Math. J.* 29 (1982) 289–298 [MR](#) [Zbl](#)
- [13] **J Meier, T Schirmer, A Zupan**, *Classification of trisections and the generalized property R conjecture*, *Proc. Amer. Math. Soc.* 144 (2016) 4983–4997 [MR](#) [Zbl](#)

- [14] **J Meier, A Thompson, A Zupan**, *Cubic graphs induced by bridge trisections*, Math. Res. Lett. 30 (2023) 1207–1231 [MR](#) [Zbl](#)
- [15] **J Meier, A Zupan**, *Bridge trisections of knotted surfaces in  $S^4$* , Trans. Amer. Math. Soc. 369 (2017) 7343–7386 [MR](#) [Zbl](#)
- [16] **J Meier, A Zupan**, *Bridge trisections of knotted surfaces in 4-manifolds*, Proc. Natl. Acad. Sci. USA 115 (2018) 10880–10886 [MR](#) [Zbl](#)
- [17] **K Miyazaki**, *On the relationship among unknotting number, knotting genus and Alexander invariant for 2-knots*, Kobe J. Math. 3 (1986) 77–85 [MR](#) [Zbl](#)
- [18] **Y Moriah**, *On the free genus of knots*, Proc. Amer. Math. Soc. 99 (1987) 373–379 [MR](#) [Zbl](#)
- [19] **M Scharlemann**, *Smooth spheres in  $\mathbb{R}^4$  with four critical points are standard*, Invent. Math. 79 (1985) 125–141 [MR](#) [Zbl](#)
- [20] **C M Tsau**, *A note on incompressible surfaces in solid tori and in lens spaces*, from “Knots 90”, de Gruyter, Berlin (1992) 213–229 [MR](#) [Zbl](#)
- [21] **R Weidmann**, *On the rank of amalgamated products and product knot groups*, Math. Ann. 312 (1998) 761–771 [MR](#) [Zbl](#)
- [22] **T Yanagawa**, *On ribbon 2-knots: the 3-manifold bounded by the 2-knots*, Osaka Math. J. 6 (1969) 447–464 [MR](#) [Zbl](#)
- [23] **E C Zeeman**, *Twisting spun knots*, Trans. Amer. Math. Soc. 115 (1965) 471–495 [MR](#) [Zbl](#)
- [24] **A Zupan**, *Bridge and pants complexities of knots*, J. Lond. Math. Soc. 87 (2013) 43–68 [MR](#) [Zbl](#)

Department of Mathematics, North Carolina School of Science and Mathematics  
Morganton, NC, United States

Department of Mathematics, Western Washington University  
Bellingham, WA, United States

Department of Mathematics, University of Texas at Austin  
Austin, TX, United States

Department of Mathematics, University of Nebraska–Lincoln  
Lincoln, NE, United States

[jason.joseph@ncssm.edu](mailto:jason.joseph@ncssm.edu), [jeffrey.meier@wwu.edu](mailto:jeffrey.meier@wwu.edu), [maggie.miller.math@gmail.com](mailto:maggie.miller.math@gmail.com),  
[zupan@unl.edu](mailto:zupan@unl.edu)

Received: 4 November 2022      Revised: 28 August 2023



# ALGEBRAIC & GEOMETRIC TOPOLOGY

[msp.org/agt](https://msp.org/agt)

## EDITORS

### PRINCIPAL ACADEMIC EDITORS

John Etnyre  
[etnyre@math.gatech.edu](mailto:etnyre@math.gatech.edu)  
Georgia Institute of Technology

Kathryn Hess  
[kathryn.hess@epfl.ch](mailto:kathryn.hess@epfl.ch)  
École Polytechnique Fédérale de Lausanne

### BOARD OF EDITORS

Julie Bergner	University of Virginia <a href="mailto:jeb2md@eservices.virginia.edu">jeb2md@eservices.virginia.edu</a>	Thomas Koberda	University of Virginia <a href="mailto:thomas.koberda@virginia.edu">thomas.koberda@virginia.edu</a>
Steven Boyer	Université du Québec à Montréal <a href="mailto:cohf@math.rochester.edu">cohf@math.rochester.edu</a>	Markus Land	LMU München <a href="mailto:markus.land@math.lmu.de">markus.land@math.lmu.de</a>
Tara E. Brendle	University of Glasgow <a href="mailto:tara.brendle@glasgow.ac.uk">tara.brendle@glasgow.ac.uk</a>	Christine Lescop	Université Joseph Fourier <a href="mailto:lescop@ujf-grenoble.fr">lescop@ujf-grenoble.fr</a>
Indira Chatterji	CNRS & Univ. Côte d'Azur (Nice) <a href="mailto:indira.chatterji@math.cnrs.fr">indira.chatterji@math.cnrs.fr</a>	Norihiko Minami	Yamato University <a href="mailto:minami.norihiko@yamato-u.ac.jp">minami.norihiko@yamato-u.ac.jp</a>
Octav Cornea	Université de Montréal <a href="mailto:cornea@dms.umontreal.ca">cornea@dms.umontreal.ca</a>	Andrés Navas	Universidad de Santiago de Chile <a href="mailto:andres.navas@usach.cl">andres.navas@usach.cl</a>
Alexander Dranishnikov	University of Florida <a href="mailto:dranish@math.ufl.edu">dranish@math.ufl.edu</a>	Robert Oliver	Université Paris 13 <a href="mailto:bobol@math.univ-paris13.fr">bobol@math.univ-paris13.fr</a>
Tobias Ekholm	Uppsala University, Sweden <a href="mailto:tobias.ekholm@math.uu.se">tobias.ekholm@math.uu.se</a>	Jessica S Purcell	Monash University <a href="mailto:jessica.purcell@monash.edu">jessica.purcell@monash.edu</a>
Mario Eudave-Muñoz	Univ. Nacional Autónoma de México <a href="mailto:mario@matem.unam.mx">mario@matem.unam.mx</a>	Birgit Richter	Universität Hamburg <a href="mailto:birgit.richter@uni-hamburg.de">birgit.richter@uni-hamburg.de</a>
David Futер	Temple University <a href="mailto:dfuter@temple.edu">dfuter@temple.edu</a>	Jérôme Scherer	École Polytech. Féd. de Lausanne <a href="mailto:jerome.scherer@epfl.ch">jerome.scherer@epfl.ch</a>
John Greenlees	University of Warwick <a href="mailto:john.greenlees@warwick.ac.uk">john.greenlees@warwick.ac.uk</a>	Vesna Stojanoska	Univ. of Illinois at Urbana-Champaign <a href="mailto:vesna@illinois.edu">vesna@illinois.edu</a>
Ian Hambleton	McMaster University <a href="mailto:ian@math.mcmaster.ca">ian@math.mcmaster.ca</a>	Zoltán Szabó	Princeton University <a href="mailto:szabo@math.princeton.edu">szabo@math.princeton.edu</a>
Matthew Hedden	Michigan State University <a href="mailto:mhedden@math.msu.edu">mhedden@math.msu.edu</a>	Maggy Tomova	University of Iowa <a href="mailto:maggy-tomova@uiowa.edu">maggy-tomova@uiowa.edu</a>
Kristen Hendricks	Rutgers University <a href="mailto:kristen.hendricks@rutgers.edu">kristen.hendricks@rutgers.edu</a>	Daniel T Wise	McGill University, Canada <a href="mailto:daniel.wise@mcgill.ca">daniel.wise@mcgill.ca</a>
Hans-Werner Henn	Université Louis Pasteur <a href="mailto:henn@math.u-strasbg.fr">henn@math.u-strasbg.fr</a>	Lior Yanovski	Hebrew University of Jerusalem <a href="mailto:lior.yanovski@gmail.com">lior.yanovski@gmail.com</a>
Daniel Isaksen	Wayne State University <a href="mailto:isaksen@math.wayne.edu">isaksen@math.wayne.edu</a>		

---

See inside back cover or [msp.org/agt](https://msp.org/agt) for submission instructions.

The subscription price for 2025 is US \$760/year for the electronic version, and \$1110/year (+\$75, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Algebraic & Geometric Topology is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [Current Mathematical Publications](#) and the [Science Citation Index](#).

Algebraic & Geometric Topology (ISSN 1472-2747 printed, 1472-2739 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840.

---

AGT peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**

nonprofit scientific publishing

<https://msp.org/>

© 2025 Mathematical Sciences Publishers



# ALGEBRAIC & GEOMETRIC TOPOLOGY

Volume 25      Issue 3 (pages 1265–1915)      2025

---

A-polynomials, Ptolemy equations and Dehn filling	1265
JOSHUA A HOWIE, DANIEL V MATHEWS and JESSICA S PURCELL	
The Alexandrov theorem for $2 + 1$ flat radiant spacetimes	1321
LÉO MAXIME BRUNSWIC	
Real algebraic overtwisted contact structures on 3-spheres	1377
ŞEYMA KARADERELI and FERİT ÖZTÜRK	
Fully augmented links in the thickened torus	1411
ALICE KWON	
Unbounded $\mathfrak{sl}_3$ -laminations and their shear coordinates	1433
TSUKASA ISHIBASHI and SHUNSUKE KANO	
Bridge trisections and Seifert solids	1501
JASON JOSEPH, JEFFREY MEIER, MAGGIE MILLER and ALEXANDER ZUPAN	
Random Artin groups	1523
ANTOINE GOLDSBOROUGH and NICOLAS VASKOU	
A deformation of Asaeda–Przytycki–Sikora homology	1545
ZHENKUN LI, YI XIE and BOYU ZHANG	
Cubulating a free-product-by-cyclic group	1561
FRANÇOIS DAHMANI and SURAJ KRISHNA MEDA SATISH	
Virtual domination of 3-manifolds, III	1599
HONGBIN SUN	
The Kakimizu complex for genus one hyperbolic knots in the 3-sphere	1667
LUIS G VALDEZ-SÁNCHEZ	
Band diagrams of immersed surfaces in 4-manifolds	1731
MARK HUGHES, SEUNGWON KIM and MAGGIE MILLER	
Anosov flows and Liouville pairs in dimension three	1793
THOMAS MASSONI	
Hamiltonian classification of toric fibres and symmetric probes	1839
JOÉ BRENDEL	
An example of higher-dimensional Heegaard Floer homology	1877
YIN TIAN and TIANYU YUAN	
Fibered 3-manifolds and Veech groups	1897
CHRISTOPHER J LEININGER, KASRA RAFI, NICHOLAS ROUSE, EMILY SHINKLE and YVON VERBERNE	