

AG
T

*Algebraic & Geometric
Topology*

Volume 25 (2025)

Random Artin groups

ANTOINE GOLDSBOROUGH

NICOLAS VASKOU

Random Artin groups

ANTOINE GOLDSBOROUGH

NICOLAS VASKOU

We introduce a new model of random Artin groups. The two variables we consider are the rank of the Artin groups and the set of permitted coefficients of their defining graphs.

The heart of our model is to control the speed at which we make that set of permitted coefficients grow relatively to the growth of the rank of the groups, as it turns out different speeds yield very different results. We describe these speeds by means of (often polynomial) functions. In this model, we show that for a large range of such functions, a random Artin group satisfies most conjectures about Artin groups asymptotically almost surely.

Our work also serves as a study of how restrictive the commonly studied families of Artin groups are, as we compute explicitly the probability that a random Artin group belongs to various families of Artin groups, such as the classes of 2-dimensional Artin groups, FC-type Artin groups, large-type Artin groups, and others.

[20F36](#), [20F65](#), [20F69](#), [20P05](#); [20F67](#)

1 Introduction

Artin groups are a family of groups that have drawn an increasing interest in the past few decades. They are defined as follows. Let Γ be a *defining graph*, that is a simplicial graph with vertex set $V(\Gamma)$ and edge set $E(\Gamma)$, such that every edge e_{ab} of Γ connecting two vertices a and b is given a coefficient $m_{ab} \in \{2, 3, \dots\}$. Then Γ defines an *Artin group*:

$$A_\Gamma := \langle V(\Gamma) \mid \underbrace{aba \cdots}_{m_{ab} \text{ terms}} = \underbrace{bab \cdots}_{m_{ab} \text{ terms}}, \forall e_{ab} \in E(\Gamma) \rangle.$$

The cardinality of $V(\Gamma)$, that is the number of *standard generators* of A_Γ , is called the *rank* of A_Γ . When a and b are not connected by an edge we set $m_{ab} := \infty$.

One of the main reasons why Artin groups have become of such great interest is because of the amount of (often easily stated) conjectures and problems about them that are still to be solved. While some of these conjectures are algebraic (torsion, centres), some others are more geometric (acylindrical hyperbolicity, CAT(0)-ness), algorithmic (word and conjugacy problems, biautomaticity), or even topological. Although close to none of these conjectures or problems has been answered in the most general case, there has

been progress on each of them. A common theme towards proving these conjectures has been to prove them for smaller families of Artin groups.

The goal of this paper is to consider Artin groups with a probabilistic approach. One might wonder what a typical Artin group looks like, and hence want to define a notion of randomness for Artin groups. By computing the different “sizes” of the most commonly studied classes of Artin groups, we give a way to quantify how restrictive these different classes really are. In light of that, our model provides a novel and explicit way of quantifying the state of the common knowledge about the aforementioned conjectures and problems about Artin groups.

Although Artin groups are defined using defining graphs, it is not known in general when two defining graphs give rise to isomorphic Artin groups. This problem, known as the *isomorphism problem*, is actually quite hard to solve even for restrictive classes of Artin groups. With our current knowledge, any (reachable) theory of randomness for Artin groups must then be based on the randomness of defining graphs, and not of the Artin groups themselves.

Random right-angled Coxeter (and Artin) groups have been studied by several authors in the literature (see Behrstock, Hagen and Sisto [1] and Charney and Farber [4]), using the Erdős–Rényi model. While in [4] the authors fix the probability of apparition of an edge as some constant $0 \leq p \leq 1$, in [1] this model is refined: $p = p(N)$ depends on the rank N of the group. That said, these models restrict to right-angled groups, where the associated defining graphs are not labelled. In [7], Deibel introduces a model of randomness for Coxeter groups in general. There are similarities between this model and ours, although the former revolves more about making the probabilities of apparition of specific coefficients vary. In particular, this model is not very well suited to provide insights on the “sizes” of the most commonly studied classes of Coxeter and Artin groups. On the contrary, this is a central goal of our model.

The two variables that come to mind when thinking about Artin groups are their rank, that is the number of vertices of the defining graph, as well as the choice of the associated coefficients. A first step in the theory is to consider what happens if we restrict ourselves to the family $\mathcal{G}^{N,M}$ of all the defining graphs with N vertices and with coefficients in $\{\infty, 2, 3, \dots, M\}$, for some $N \geq 1$ and $M \geq 2$. As we want any possible rank and any possible coefficient to eventually appear in a random Artin group, a convenient way to think about randomness is to pick a defining graph at random in the family $\mathcal{G}^{N,M}$, and then to make N and M grow to infinity. Note that isomorphic labelled graphs may be counted multiples times in $\mathcal{G}^{N,M}$.

As it turns out, randomness of defining graphs highly depends on the speed at which N and M grow. A prime example of this is that the probability for a defining graph of $\mathcal{G}^{N,M}$ to give an Artin group of large-type (meaning that none of the coefficients is 2) tends to 1 when M grows much faster than N , and tends to 0 when N grows much faster than M . To solve this problem, we decide to relate N and M through a function f such that $M := f(N)$. This way, we only have to look at the family $\mathcal{G}^{N,f(N)}$ when N goes to infinity.

If $A_{\mathcal{F}}$ is a family of Artin groups coming from a family of defining graphs \mathcal{F} , a way of measuring the “size” of $A_{\mathcal{F}}$ is to compute the limit

$$\lim_{N \rightarrow \infty} \frac{\#(\mathcal{F} \cap \mathcal{G}^{N,f(N)})}{\#(\mathcal{G}^{N,f(N)})}.$$

Of course, this ratio depends on the choice we make for the function f . When the above limit is 1, that is when the probability that a graph picked at random in $\mathcal{G}^{N,f(N)}$ will give an Artin group that belongs to the said family $A_{\mathcal{F}}$ tends to 1, we say that a random Artin group (with respect to f) is *asymptotically almost surely* in $A_{\mathcal{F}}$.

One may wonder why our model only considers graphs of rank N , and not all graphs with rank at most N . As it turns out, the size of the set of all graphs with at most N vertices (and coefficients in $\{\infty, 2, \dots, f(N)\}$) is asymptotically the same as the size of $\mathcal{G}^{N,f(N)}$, in the sense that the quotient of the two values tends to 1 when N approaches ∞ . Thus asymptotically it is not an actual restriction to only consider graphs with precisely N vertices.

Now, there are families $A_{\mathcal{F}}$ of Artin groups for which the above limit tends to 1 no matter what (sensible) choice we make for the function f . We say that such a family is *uniformly large* (resp. *uniformly small* if that limit is always 0). Our first result concern such families of Artin groups:

Theorem 1.1 *The family of irreducible Artin groups and the family of Artin groups with connected defining graphs are uniformly large. On the other hand, the family of Artin groups of type FC is uniformly small. In particular, the same applies to the families of RAAGs and triangle-free Artin groups.*

As mentioned earlier, there are numerous families of Artin groups whose “size” depends on the choice of function f . When f is large enough, which means that the choice of possible coefficients for the defining graphs grows fast enough compared to the rank of the Artin group, we obtain much stronger results. This is made explicit in the next two theorems.

For two nondecreasing divergent functions $f, g: \mathbb{N} \rightarrow \mathbb{N}$ we say that $f \preceq g$ if the limit

$$\lim_{N \rightarrow \infty} \frac{f(N)}{g(N)}$$

exists and is finite. If $f \preceq g$ and $f \succeq g$ then we will write $f \simeq g$. Finally if $f \preceq g$ but $f \not\succeq g$ then we will write $f < g$, and similarly for $f > g$.

Theorem 1.2 *Let $A_{\mathcal{F}}$ be any family of Artin groups defined by forbidding a finite number of coefficients from their defining graphs, and consider a function $f: \mathbb{N} \rightarrow \mathbb{N}$. Let Γ be a graph picked at random in $\mathcal{G}^{N,f(N)}$.*

- (1) *If $f(N) > N^2$, then A_{Γ} asymptotically almost surely belongs to $A_{\mathcal{F}}$.*
- (2) *If $f(N) < N^2$, then A_{Γ} asymptotically almost surely does not belong to $A_{\mathcal{F}}$.*
- (3) *If $f(N) \simeq N^2$, then the probability that A_{Γ} belongs to $A_{\mathcal{F}}$ is strictly between 0 and 1.*

Note that the previous theorem applies to the families of large-type, extra-large-type, or large-type and free-of-infinity Artin groups. There are strong results in the literature about these families of Artin groups, as most of the famous conjectures and problems about Artin groups have been solved for at least one of them (see [Section 2](#)).

While these different families of Artin groups have the same threshold at $f(N) \simeq N^2$ no matter how many coefficients we forbid, the class of 2-dimensional Artin groups turns out to be substantially bigger. Studying this class, we obtain the following result:

Theorem 1.3 *Consider a nondecreasing divergent function $f: \mathbb{N} \rightarrow \mathbb{N}$. Let Γ be a graph picked at random in $\mathcal{G}^{N, f(N)}$.*

- (1) *If $f(N) \succ N^{3/2}$, then A_Γ asymptotically almost surely is 2-dimensional.*
- (2) *If $f(N) < N^{3/2}$, then A_Γ asymptotically almost surely is not 2-dimensional.*

A consequence of the two previous theorems is that we are able, when f grows fast enough, to show that a random Artin group asymptotically almost surely satisfies most of the main conjectures about Artin groups:

Theorem 1.4 *Let $f: \mathbb{N} \rightarrow \mathbb{N}$ be such that $f(N) \succ N^{3/2}$, and let Γ be a graph picked at random in $\mathcal{G}^{N, f(N)}$. Then asymptotically almost surely, the following properties hold:*

- (1) *A_Γ is torsion-free;*
- (2) *A_Γ has trivial centre;*
- (3) *A_Γ has solvable word and conjugacy problem;*
- (4) *A_Γ satisfies the $K(\pi, 1)$ -conjecture;*
- (5) *the set of parabolic subgroups of A_Γ is closed under (arbitrary) intersections;*
- (6) *A_Γ is acylindrically hyperbolic;*
- (7) *A_Γ satisfies the Tits alternative;*
- (8) *A_Γ is not virtually cocompactly cubulated.*

Moreover, if $f(N) \succ N^2$ then asymptotically almost surely the following properties also hold:

- (1) *A_Γ is CAT(0);*
- (2) *A_Γ is hierarchically hyperbolic;*
- (3) *A_Γ is systolic and thus biautomatic;*
- (4) *A_Γ is rigid;*
- (5) *$\text{Aut}(A_\Gamma) \cong A_\Gamma \rtimes \text{Out}(A_\Gamma)$, where $\text{Out}(A_\Gamma) \cong \text{Aut}(\Gamma) \times (\mathbb{Z}/2\mathbb{Z})$ is finite.*




Figure 1: The axis represents various (polynomial) functions f . Above the main axis are described the classes of Artin groups that we obtain asymptotically almost surely with respect to f , while under this axis we list the properties that we know these groups will satisfy asymptotically almost surely.

At last, we also prove interesting results for families of Artin groups in which the number M of permitted coefficients grows “slowly enough” compared to the rank N . We focus on the class of Artin groups A_Γ whose associated graphs Γ are not cones, and we prove that for most (nondecreasing divergent) functions, the probability that a random Artin group is acylindrically hyperbolic and has trivial centre tends to 1.

Theorem 1.5 *Let $\alpha \in (0, 1)$ and let $f : \mathbb{N} \rightarrow \mathbb{N}$ be a nondecreasing divergent function satisfying $f(N) \prec N^{1-\alpha}$. Let now Γ be a graph picked at random in $\mathcal{G}^{N, f(N)}$. Then the associated Artin group A_Γ is acylindrically hyperbolic and has trivial centre asymptotically almost surely.*

The results of the above theorems for polynomial functions is encapsulated in Figure 1.

The previous results shows that we are very close to being able to state that “almost all Artin groups are acylindrically hyperbolic and have trivial centres”. It is conjectured that all irreducible nonspherical Artin groups are acylindrically hyperbolic; see Charney and Morris-Wright [5]. Although proving this conjecture for all Artin groups seems to be a difficult problem, some progress has been made in recent years; see Kato and Oguni [13] and Vaskou [17]. It would seem to be an interesting line of research to try to expand the spectrum of families of Artin groups for which one can prove acylindrical hyperbolicity, in order to “fill in” the gap of functions at which a random Artin group is acylindrically hyperbolic. This leads to the following question.

Question 1.6 Construct a family $A_{\mathcal{F}}$ of acylindrically hyperbolic Artin groups or of Artin groups with trivial centres for which the following holds:

There exists an $\alpha \in (0, 1)$ such that for all functions $f: \mathbb{N} \rightarrow \mathbb{N}$ satisfying $N^{1-\alpha} \preceq f(N) \preceq N^{3/2}$, a graph Γ picked at random in $\mathcal{G}^{N, f(N)}$ is such that A_{Γ} asymptotically almost surely belongs to $A_{\mathcal{F}}$.

Acknowledgements

The authors would like to thank Alessandro Sisto for helpful conversations and for pointing out the strategy of the second moment method, and Alexandre Martin for useful comments. We would also like to thank the Institut Henri Poincaré (UAR 839 CNRS-Sorbonne Université), and LabEx CARMIN (ANR-10-LABX-59-01) for providing a warm environment for this project to begin. We thank the referee for carefully reading the article and for their detailed comments. The work of Goldsborough was supported by the EPSRC-UKRI studentship EP/V520044/1.

2 Preliminaries and first results

In this section we bring more details about some of the notions discussed in the introduction. This includes discussions about most of the commonly studied classes of Artin groups, as well as discussions regarding open conjectures related to Artin groups.

Throughout this paper, we will often call a *triangle* in a graph Γ any subgraph of Γ that is generated by 3 vertices. This notion will be convenient, although one must note that with this definition, triangles may have strictly fewer than 3 edges, as subgraphs of Γ .

Most of the main conjectures about Artin groups are still open in general. That said, many of them have been proved for smaller families of Artin groups. Two important of these families are the families of 2-dimensional Artin groups and the family of Artin groups of type FC. These two families have been extensively studied following the work of Charney and Davis [3]. The other well-studied families are usually subfamilies of these.

Before coming to these definitions, we first recall what a parabolic subgroup of an Artin group is. Let A_{Γ} be any Artin group, and let Γ' be a full subgraph of Γ . A standard result about Artin groups states that the subgroup of A_{Γ} generated by the vertices of Γ' is also an Artin group, that is isomorphic to $A_{\Gamma'}$ [14]. Such a subgroup is called a *standard parabolic subgroup* of A_{Γ} . The conjugates of these subgroups are called the *parabolic subgroups* of A_{Γ} .

Definition 2.1 (0) An Artin group A_{Γ} is said to be *spherical* if the associated Coxeter group W_{Γ} is finite.

(1) An Artin group A_Γ is said to be *2-dimensional* if for every triplet of distinct standard generators $a, b, c \in V(\Gamma)$, the subgraph Γ' spanned by a, b and c corresponds to an Artin group $A_{\Gamma'}$ that is *not* spherical. By a result of [3], this is equivalent to requiring that

$$\frac{1}{m_{ab}} + \frac{1}{m_{ac}} + \frac{1}{m_{bc}} \leq 1.$$

We let \mathcal{D} be the set of graphs Γ such that the above condition is satisfied. We let $A_{\mathcal{D}}$ be the set of 2-dimensional Artin groups. The family of 2-dimensional Artin groups contains the well-studied families of *large-type* Artin groups (every coefficient is at least 3), *extra-large-type* Artin groups (every coefficient is at least 4), or *XXL* Artin groups (every coefficient is at least 5).

(2) An Artin group A_Γ is said to be of *type FC* if every complete subgraph $\Gamma' \subseteq \Gamma$ generates an Artin group $A_{\Gamma'}$ that is spherical. Let \mathcal{FC} be the set of graphs Γ that give rise to an Artin group of type FC and let $A_{\mathcal{FC}}$ be the set of Artin groups of type FC.

The family of Artin groups of type FC contains the family of right-angled Artin groups, also called *RAAGs* (the only permitted coefficients are 2 and ∞), the family of spherical Artin groups, and the family of *triangle-free* Artin groups (the Artin groups whose associated graphs don't contain any 3-cycles). Being triangle-free is actually equivalent to being both of type FC and 2-dimensional.

We now move towards the main conjectures related to Artin groups. For each conjecture, we will briefly describe the state of the common research towards proving it, by mentioning the one or two result(s) that will turn out to be the more “probabilistically relevant” in our model — in other words, the results that cover the largest classes.

Conjecture 2.2 Let A_Γ be any Artin group. Then:

- (1) A_Γ is torsion-free.
 \hookrightarrow This was proved for 2-dimensional Artin groups [3].
- (2) If A_Γ is irreducible and nonspherical, then A_Γ has trivial centre.
 \hookrightarrow This was proved for 2-dimensional Artin groups [17], and for Artin groups whose graph is not the cone of a single vertex [5].
- (3) A_Γ has solvable word and conjugacy problems.
 \hookrightarrow This was proved for 2-dimensional Artin groups [11].
- (4) A_Γ satisfies the $K(\pi, 1)$ -conjecture.
 \hookrightarrow This was proved for 2-dimensional Artin groups [3].
- (5) Intersections of parabolic subgroups of A_Γ give parabolic subgroups of A_Γ .
 \hookrightarrow This was proved for large-type Artin groups [6] and more generally for $(2, 2)$ -free 2-dimensional Artin groups [2].
- (6) A_Γ is CAT(0).
 \hookrightarrow This was proved for XXL Artin groups [9].

- (7) If A_Γ is irreducible and nonspherical, then A_Γ is acylindrically hyperbolic.
 - \hookrightarrow This was proved for 2-dimensional Artin groups [17], and for Artin groups whose graph is not the cone of a single vertex [13].
- (8) A_Γ is hierarchically hyperbolic.
 - \hookrightarrow This was proved for extra-large-type Artin groups [10].
- (9) A_Γ is systolic and biautomatic.
 - \hookrightarrow This was proved for large-type Artin groups [12].
- (10) A_Γ satisfies the Tits alternative.
 - \hookrightarrow This was proved for 2-dimensional Artin groups [15].

In addition to these conjectures, the following questions have been raised:

Question 2.3 Let A_Γ be any Artin group.

- (1) When is A_Γ not virtually cocompactly cubulated?
 - \hookrightarrow This was proved to be the case when A_Γ is 2-dimensional and satisfies the condition of [8, Conjecture B].
- (2) When is $\text{Out}(A_\Gamma)$ finite?
 - \hookrightarrow This was proved to be the case for large-type free-of-infinity Artin groups [18].
- (3) When is A_Γ rigid, in the sense of [16]?
 - \hookrightarrow This was proved to be the case for large-type Artin groups that have no separating edges [16, Theorem B]. This includes the class of large-type free-of-infinity Artin groups.

Definition 2.4 Let \mathcal{F} be a family of defining graphs and let $A_{\mathcal{F}}$ be the corresponding class of Artin groups. Let $f : \mathbb{N} \rightarrow \mathbb{N}$ be a nondecreasing divergent function. We say that a random Artin group (with respect to f) A_Γ belongs to $A_{\mathcal{F}}$ with probability

$$\mathbb{P}_f[A_\Gamma \in A_{\mathcal{F}}] := \lim_{N \rightarrow \infty} \mathbb{P}[\Gamma \in \mathcal{F} \mid \Gamma \in \mathcal{G}^{N, f(N)}] = \lim_{N \rightarrow \infty} \frac{\#(\mathcal{F} \cap \mathcal{G}^{N, f(N)})}{\#(\mathcal{G}^{N, f(N)})},$$

when the limit exists. Furthermore, we say that a random Artin group A_Γ (with respect to f) is *asymptotically almost surely* in $A_{\mathcal{F}}$ if $\mathbb{P}_f[A_\Gamma \in A_{\mathcal{F}}] = 1$. Similarly, we say that A_Γ is *asymptotically almost surely not* in $A_{\mathcal{F}}$ if $\mathbb{P}_f[A_\Gamma \in A_{\mathcal{F}}] = 0$.

Definition 2.5 Let $A_{\mathcal{F}}$ be a family of Artin groups. Then we say that $A_{\mathcal{F}}$ is *uniformly large* if for every nondecreasing divergent function $f : \mathbb{N} \rightarrow \mathbb{N}$, a random Artin group A_Γ (with respect to f) is asymptotically almost surely in $A_{\mathcal{F}}$. We say that \mathcal{F} is *uniformly small* if A_Γ is asymptotically almost surely not in $A_{\mathcal{F}}$.

We now move towards our first results. The first thing we will prove is that the family of irreducible Artin groups and the family of Artin groups with connected defining graphs are uniformly large. This is

important as many results regarding Artin groups assume that the corresponding groups are irreducible and/or have a connected defining graph. Our work shows that these two hypotheses are very much not restrictive.

Definition 2.6 Let Γ_1 and Γ_2 be two defining graphs. The graph $\Gamma_1 *_k \Gamma_2$ is the graph obtained by attaching every vertex of Γ_1 to every vertex of Γ_2 by an edge with label k (with $k \in \{\infty, 2, 3, \dots\}$).

Let now Γ be any defining graph. Then Γ is called a k -join relative to Γ_1 and Γ_2 if there are two subgraphs $\Gamma_1, \Gamma_2 \subseteq \Gamma$ such that $V(\Gamma_1) \sqcup V(\Gamma_2) = V(\Gamma)$ and such that $\Gamma = \Gamma_1 *_k \Gamma_2$.

We will denote by $\mathcal{A}_{\mathcal{F}_k}$ the class of Artin groups whose defining graphs decompose as k -joins.

Remark 2.7 (1) If $\Gamma \in \mathcal{F}_2$ then A_Γ decomposes as a direct product $A_{\Gamma_1} \times A_{\Gamma_2}$ in an obvious way. In that case, Γ is called *reducible*. The class \mathcal{F}_2^C of *irreducible* defining graphs will be denoted by Irr .
 (2) If $\Gamma \in \mathcal{F}_\infty$ then it is disconnected. The class \mathcal{F}_∞^C of connected defining graphs will be denoted by Con .

Lemma 2.8 For all $k \in \{\infty, 2, 3, \dots\}$, the family $\mathcal{A}_{\mathcal{F}_k}$ is uniformly small. In particular, the classes A_{Irr} and A_{Con} of Artin groups are both uniformly large.

Proof We will count the number of decompositions of the graph Γ as $\Gamma = \Gamma_1 *_k \Gamma_2$. Without loss of generality, we will let Γ_1 denote the subgraph with the lower rank, so that $|V(\Gamma_1)| \leq \lfloor N/2 \rfloor$. Let $f: \mathbb{N} \rightarrow \mathbb{N}$ be a nondecreasing divergent function and consider the family \mathcal{F}_k . For a given $N \geq 1$,

$$\begin{aligned} \mathbb{P}[\Gamma \in \mathcal{F}_k \mid \Gamma \in \mathcal{G}^{N, f(N)}] &= \mathbb{P}[\exists \Gamma_1, \Gamma_2 \text{ with } |V(\Gamma_1)| \leq N/2 \text{ such that } \Gamma = \Gamma_1 *_k \Gamma_2 \mid \Gamma \in \mathcal{G}^{N, f(N)}] \\ &\leq \sum_{j=1}^{\lfloor N/2 \rfloor} \mathbb{P}[\exists \Gamma_1, \Gamma_2 \text{ with } |V(\Gamma_1)| = j \text{ such that } \Gamma = \Gamma_1 *_k \Gamma_2 \mid \Gamma \in \mathcal{G}^{N, f(N)}] \\ &= \sum_{j=1}^{\lfloor N/2 \rfloor} \binom{N}{j} \left(\frac{1}{f(N)}\right)^{j(N-j)} \\ &\leq \sum_{j=1}^{\lfloor N/2 \rfloor} \left(\frac{Ne}{j f(N)^{N/2}}\right)^j \leq \frac{Ne}{f(N)^{N/2}} \cdot \left(\frac{1 - (Ne/f(N)^{N/2})^{N/2+1}}{1 - Ne/f(N)^{N/2}}\right) \end{aligned}$$

where we used the bound

$$\binom{N}{j} \leq \left(\frac{Ne}{j}\right)^j.$$

Now $\lim_{N \rightarrow \infty} Ne/f(N)^{N/2} = 0$ for any nondecreasing divergent function f , so we obtain

$$\mathbb{P}_f[A_\Gamma \in \mathcal{A}_{\mathcal{F}_k}] = \lim_{N \rightarrow \infty} \mathbb{P}[\Gamma \in \mathcal{F}_k \mid \Gamma \in \mathcal{G}^{N, f(N)}] = 0 \cdot \left(\frac{1-0}{1-0}\right) = 0.$$

This proves the main statement of the lemma. The second then directly follows from [Remark 2.7](#). □

Our next result concerns the class of Artin groups of type FC.

Lemma 2.9 *The family $A_{\mathcal{F}\mathcal{C}}$ of Artin groups of type FC is uniformly small. In particular, the family of triangle-free Artin groups, the family of spherical Artin groups and the family of RAAGs are also uniformly small.*

Proof Let f be any nondecreasing divergent function, and let $\Gamma \in \mathcal{G}^{N,f(N)}$. We want to bound the probability that Γ belongs to $\mathcal{F}\mathcal{C} \cap \mathcal{G}^{N,f(N)}$. Let a, b and c be three vertices of Γ . The probability that any of the three corresponding coefficients m_{ab}, m_{ac} and m_{bc} is not 2 or ∞ is precisely $(f(N) - 2)/f(N)$, and hence the probability that the three coefficients are not 2 nor ∞ is $((f(N) - 2)/f(N))^3$. Note that when this happens, the subgraph $\Gamma' \subseteq \Gamma$ spanned by a, b and c is complete but generates an Artin group $A_{\Gamma'}$ which is nonspherical (the sum of the inverses of the three corresponding coefficients is ≤ 1). In particular, Γ is not of type FC. We obtain

$$\mathbb{P}_f[A_\Gamma \notin A_{\mathcal{F}\mathcal{C}}] = \lim_{N \rightarrow \infty} \frac{\#(\mathcal{G}^{N,f(N)} \setminus \mathcal{F}\mathcal{C})}{\#(\mathcal{G}^{N,f(N)})} \geq \lim_{N \rightarrow \infty} \left(\frac{f(N) - 2}{f(N)} \right)^3 = \lim_{N \rightarrow \infty} \left(1 - \frac{2}{f(N)} \right)^3 = 1. \quad \square$$

As mentioned in the introduction, there are interesting classes of Artin groups for which the probability that a graph taken at random will belong to the class highly depends on the choice of function f . Some examples are given through the following theorem.

Theorem 2.10 *Let \mathcal{F} be any family of graphs defined by forbidding a finite number k of coefficients and let $A_{\mathcal{F}}$ be the family of corresponding Artin groups. Consider a function $f : \mathbb{N} \rightarrow \mathbb{N}$. Let A_Γ be a random Artin group (with respect to f).*

- (1) *If $f(N) \succ N^2$, then A_Γ asymptotically almost surely belongs to $A_{\mathcal{F}}$.*
- (2) *If $f(N) \prec N^2$, then A_Γ asymptotically almost surely does not belong to $A_{\mathcal{F}}$.*
- (3) *If $f(N) \simeq N^2$ then asymptotically we have $\mathbb{P}_f[A_\Gamma \in A_{\mathcal{F}}] \in (0, 1)$. Moreover, if $f(N) = \lambda N^2$ for some $\lambda > 0$, then $\mathbb{P}_f[A_\Gamma \in A_{\mathcal{F}}] = e^{-k/2\lambda}$.*

Proof A graph with N vertices has $\frac{1}{2}N(N - 1)$ pairs of vertices, each of which is given one of $f(N)$ possible coefficients. Hence, direct computations on the possible number of graphs give

$$\#\mathcal{G}^{N,f(N)} = (f(N))^{\frac{N(N-1)}{2}}.$$

Similarly, we have

$$\#(\mathcal{F} \cap \mathcal{G}^{N,f(N)}) = (f(N) - k)^{\frac{N(N-1)}{2}}.$$

And thus we obtain

$$\begin{aligned} \mathbb{P}_f[A_\Gamma \in A_{\mathcal{F}}] &= \lim_{N \rightarrow \infty} \frac{\#(\mathcal{F} \cap \mathcal{G}^{N,f(N)})}{\#(\mathcal{G}^{N,f(N)})} = \lim_{N \rightarrow \infty} \left(\frac{f(N) - k}{f(N)} \right)^{\frac{N(N-1)}{2}} \\ &= \lim_{N \rightarrow \infty} \left(\frac{f(N) - k}{f(N)} \right)^{f(N) \left(\frac{N(N-1)}{2f(N)} \right)}. \end{aligned}$$

Observe that

$$\lim_{N \rightarrow \infty} \left(\frac{f(N) - k}{f(N)} \right)^{f(N)} = e^{-k}.$$

In particular, for any $\epsilon > 0$ there is a big enough N_ϵ such that for all $N \geq N_\epsilon$ we have

$$e^{-k} - \epsilon \leq \left(\frac{f(N) - k}{f(N)} \right)^{f(N)} \leq e^{-k} + \epsilon.$$

Hence for $N \geq N_\epsilon$,

$$(e^{-k} - \epsilon)^{r(N)} \leq \left(\frac{f(N) - k}{f(N)} \right)^{f(N) \left(\frac{N(N-1)}{2f(N)} \right)} \leq (e^{-k} + \epsilon)^{r(N)},$$

where $r(N) = N(N - 1)/(2f(N))$.

Therefore, if $f(N) \succ N^2$, there is a function h with $\lim_{N \rightarrow \infty} h(N) = \infty$ such that $f(N) = h(N)N^2$, and hence $r(N) = (N - 1)/(2Nh(N))$ which tends to 0 as $N \rightarrow +\infty$. Thus, in this case

$$\mathbb{P}_f[A_\Gamma \in A_{\mathcal{F}}] = \lim_{N \rightarrow \infty} \left(\frac{f(N) - k}{f(N)} \right)^{f(N)r(N)} = 1.$$

If $f(N) \prec N^2$, there exists a function h with $\lim_{N \rightarrow \infty} h(N) = \infty$ such that $f(N)h(N) = N^2$, and here $r(N) = (N - 1)h(N)/(2N)$ which tends to ∞ as $N \rightarrow \infty$, so in this case

$$\mathbb{P}_f[A_\Gamma \in A_{\mathcal{F}}] = \lim_{N \rightarrow \infty} \left(\frac{f(N) - k}{f(N)} \right)^{f(N)r(N)} = 0.$$

If $f(N) \simeq N^2$, then $\lim_{N \rightarrow \infty} f(N)/N^2$ is a nonzero constant and hence $\lim_{N \rightarrow \infty} r(N) = M$ for some constant $M > 0$. Thus in this case,

$$\mathbb{P}_f[A_\Gamma \in A_{\mathcal{F}}] = e^{-kM}.$$

Finally, if $f(N) = \lambda N^2$, we obtain $r(N) \rightarrow 1/(2\lambda) =: M$ and the result follows. □

The previous theorem has many consequences, as it can be applied to the families of large-type, extra-large-type, XXL or free-of-infinity Artin groups, for which much is known. Before stating an explicit result in [Corollary 2.12](#), we prove the following small lemma:

Lemma 2.11 *Let $A_{\mathcal{F}}$ and $A_{\mathcal{H}}$ be two families of Artin groups, let $f : \mathbb{N} \rightarrow \mathbb{N}$ be a nondecreasing divergent function, and suppose that $\mathbb{P}_f[A_\Gamma \in A_{\mathcal{H}}] = 1$. Then*

$$\mathbb{P}_f[A_\Gamma \in A_{\mathcal{F}}] = \mathbb{P}_f[A_\Gamma \in A_{\mathcal{F}} \cap A_{\mathcal{H}}].$$

Proof This is straightforward:

$$\begin{aligned} \mathbb{P}_f[A_\Gamma \in A_{\mathcal{F}}] &= \mathbb{P}_f[A_\Gamma \in A_{\mathcal{F}} \cap A_{\mathcal{H}}] + \underbrace{\mathbb{P}_f[A_\Gamma \in A_{\mathcal{F}} \cup A_{\mathcal{H}}]}_{=1} - \underbrace{\mathbb{P}_f[A_\Gamma \in A_{\mathcal{H}}]}_{=1} \\ &= \mathbb{P}_f[A_\Gamma \in A_{\mathcal{F}} \cap A_{\mathcal{H}}]. \end{aligned}$$
□

Corollary 2.12 *Let $f : \mathbb{N} \rightarrow \mathbb{N}$ be a function satisfying $f(N) \succ N^2$. Then a random Artin group A_Γ (with respect to f) satisfies any of the following properties asymptotically almost surely:*

- (1) A_Γ is CAT(0);
- (2) A_Γ is hierarchically hyperbolic;
- (3) A_Γ is systolic and biautomatic;
- (4) A_Γ is rigid;
- (5) $\text{Aut}(A_\Gamma) \cong A_\Gamma \rtimes \text{Out}(A_\Gamma)$, where $\text{Out}(A_\Gamma) \cong \text{Aut}(\Gamma) \times (\mathbb{Z}/2\mathbb{Z})$ is finite.

Proof Let $A_{\mathfrak{K}}$ be the class of XXL free-of-infinity Artin groups, and let $A_{\mathfrak{L}} := A_{\text{Irr}} \cap A_{\text{Con}} \cap A_{\mathfrak{K}}$. Using Lemmas 2.8 and 2.11 we can see that $\mathbb{P}_f[A_\Gamma \in A_{\mathfrak{L}}] = \mathbb{P}_f[A_\Gamma \in A_{\mathfrak{K}}]$. The class $A_{\mathfrak{K}}$ has been defined as forbidding 4 coefficients from the defining graph; hence by Theorem 2.10 a random Artin group A_Γ (with respect to f) is asymptotically almost surely in $A_{\mathfrak{K}}$ and therefore asymptotically almost surely in $A_{\mathfrak{L}}$. The various results given in Conjecture 2.2 concern families of Artin groups that all contain the family $A_{\mathfrak{L}}$. In particular, every Artin group in $A_{\mathfrak{L}}$ satisfies the ten points of Conjecture 2.2. The results given in items (6), (8) and (9) of Conjecture 2.2 are precisely those needed for items (1), (2) and (3) of Corollary 2.12. Similarly, every Artin group in $A_{\mathfrak{K}}$ is rigid, as per item (3) of Question 2.3. This proves point (4) of Corollary 2.12. For item (5), this follows from [18, Theorem A] which shows that this result holds for large-type free-of-infinity Artin groups, and in particular for Artin groups in the family $A_{\mathfrak{K}}$. \square

Remark 2.13 The previous corollary proves the five points in the second half of Theorem 1.4. Note that at this point, we could already prove the eight points in the first half of Theorem 1.4 for $f(N) \succ N^2$. We did not include this proof as it will be extended to all functions $f(N) \succ N^{3/2}$ in the following section.

3 Two-dimensional Artin groups

This section aims at studying from our probabilistic point of view the family of 2-dimensional Artin groups. This family is particularly important in the study of Artin groups, and many authors in the literature have obtained strong results for this class (see Conjecture 2.2).

Our goal will be to show that if $f(N) \succ N^{3/2}$ then asymptotically almost surely a random Artin group (with respect to f) will be 2-dimensional and if $f(N) \prec N^{3/2}$ then asymptotically almost surely a random Artin group (with respect to f) will not be 2-dimensional. In particular, we will be able to improve the result of Corollary 2.12, thus proving Theorem 1.4.

The condition of being 2-dimensional (see Definition 2.1(1)) is quite specific, which makes it hard to compute the “size” of the family. As it turns out, the size of this family is comparable to the size of another family of Artin groups, which is slightly easier to compute (see Lemma 3.2 and Theorem 3.3). This other family resembles the family introduced in [2]. We introduce it here:

Definition 3.1 We say an Artin group A_Γ is $(2, 2)$ -free if Γ does not have any two adjacent edges labelled by 2. We denote by \mathcal{B} the set of graphs that do not have two adjacent edges labelled by 2. We define $A_{\mathcal{B}}$ to be the family of $(2, 2)$ -free Artin groups.

Recall that in Definition 2.1(1), we have defined the set of graphs \mathcal{D} and the set of Artin groups $A_{\mathcal{D}}$. The following lemma is a key result. It will allow us to restrict to the study of $(2, 2)$ -free Artin groups, as asymptotically this family has the same size as the family $A_{\mathcal{D}}$.

Lemma 3.2 For all nondecreasing divergent functions $f : \mathbb{N} \rightarrow \mathbb{N}$,

- $\mathbb{P}_f[A_\Gamma \in A_{\mathcal{D}}] \leq \mathbb{P}_f[A_\Gamma \in A_{\mathcal{B}}]$;
- further, if $f(N) \succ N$, then $\mathbb{P}_f[A_\Gamma \in A_{\mathcal{D}}] = \mathbb{P}_f[A_\Gamma \in A_{\mathcal{B}}]$.

Proof The probability that a random Artin group A_Γ gives rise to a 2-dimensional Artin group can be found by conditioning on the event “ $\Gamma \in \mathcal{B}$ ”:

$$(*) \quad \mathbb{P}[\Gamma \in \mathcal{D} \mid \Gamma \in \mathcal{G}^{N, f(N)}] = \mathbb{P}[\Gamma \in \mathcal{D} \mid (\Gamma \in \mathcal{B}) \cap (\Gamma \in \mathcal{G}^{N, f(N)})] \mathbb{P}[\Gamma \in \mathcal{B} \mid \Gamma \in \mathcal{G}^{N, f(N)}] + \mathbb{P}[\Gamma \in \mathcal{D} \mid (\Gamma \notin \mathcal{B}) \cap (\Gamma \in \mathcal{G}^{N, f(N)})] \mathbb{P}[\Gamma \notin \mathcal{B} \mid \Gamma \in \mathcal{G}^{N, f(N)}].$$

Note that once we have two adjacent edges e_1 and e_2 labelled by 2, then the probability that the triangle spanned by $\{e_1, e_2\}$ generates an Artin groups of spherical type is exactly the probability that the last edge is not labelled by ∞ . This probability is $(f(N) - 1)/f(N)$; hence we have

$$\mathbb{P}[\Gamma \in \mathcal{D} \mid (\Gamma \notin \mathcal{B}) \cap (\Gamma \in \mathcal{G}^{N, f(N)})] \leq 1 - \frac{f(N) - 1}{f(N)} = \frac{1}{f(N)}.$$

Whence we get the following upper bound for $(*)$, for any nondecreasing function f :

$$\mathbb{P}[\Gamma \in \mathcal{D} \mid \Gamma \in \mathcal{G}^{N, f(N)}] \leq \mathbb{P}[\Gamma \in \mathcal{B} \mid \Gamma \in \mathcal{G}^{N, f(N)}] + \mathbb{P}[\Gamma \notin \mathcal{B} \mid \Gamma \in \mathcal{G}^{N, f(N)}] \cdot \frac{1}{f(N)}.$$

By noting that for any nondecreasing divergent function f we have that $1/f(N) \rightarrow 0$, we get

$$\mathbb{P}_f[A_\Gamma \in A_{\mathcal{D}}] = \lim_{N \rightarrow \infty} \mathbb{P}[\Gamma \in \mathcal{D} \mid \Gamma \in \mathcal{G}^{N, f(N)}] \leq \lim_{N \rightarrow \infty} \mathbb{P}[\Gamma \in \mathcal{B} \mid \Gamma \in \mathcal{G}^{N, f(N)}] = \mathbb{P}_f[A_\Gamma \in A_{\mathcal{B}}].$$

We now deal with the lower bound. The probability that a given triangle Δ is of spherical type is the quotient

$$(**) \quad \frac{\# \text{ ways that } \Delta \text{ can be spherical}}{\# \text{ possible coefficients on } \Delta}.$$

In general, the only triangles that give spherical Artin groups are of the form $(2, 3, 3)$, $(2, 3, 4)$, $(2, 3, 5)$, and $(2, 2, k)$ for $k \geq 2$. In our case, it is given that A_Γ is $(2, 2)$ -free, so the only triangles which are of spherical type are of the form $(2, 3, 3)$, $(2, 3, 4)$ or $(2, 3, 5)$. When considering the possible permutations of the order of the coefficients, this gives 15 possibilities. This yields the numerator of $(**)$.

In order to find an upper bound for $(**)$, it remains to find a lower bound for the denominator. In a graph Γ that we know is $(2, 2)$ -free, a triangle whose edges are all labelled by coefficients other than 2 will always be a possible combination of coefficients for a triangle Δ of Γ . Hence the number of possible

coefficients for a triangle Δ of a $(2, 2)$ -free graph is at least $(f(N) - 1)^3$. This yields

$$(***) \quad \frac{\# \text{ ways that } \Delta \text{ can be spherical}}{\# \text{ possible coefficients on } \Delta} \leq \frac{15}{(f(N) - 1)^3}.$$

Hence, by an union bound we get

$$\begin{aligned} \mathbb{P}[\Gamma \notin \mathcal{D} \mid (\Gamma \in \mathcal{B}) \cap (\Gamma \in \mathcal{G}^{N, f(N)})] &\leq \sum_{\Delta \text{ triangle in } \Gamma} \mathbb{P}[\Delta \text{ is of spherical type} \mid (\Gamma \in \mathcal{B}) \cap (\Gamma \in \mathcal{G}^{N, f(N)})] \\ &\leq \binom{N}{3} \frac{15}{(f(N) - 1)^3}. \end{aligned}$$

Therefore, by $(*)$ we get

$$(****) \quad \mathbb{P}[\Gamma \in \mathcal{D} \mid \Gamma \in \mathcal{G}^{N, f(N)}] \geq \left(1 - \binom{N}{3} \frac{15}{(f(N) - 1)^3}\right) \mathbb{P}[\Gamma \in \mathcal{B} \mid \Gamma \in \mathcal{G}^{N, f(N)}].$$

Hence, if $f(N) \succ N$, we have

$$\lim_{N \rightarrow \infty} \left(\binom{N}{3} \frac{15}{(f(N) - 1)^3} \right) = 0.$$

This means that

$$\begin{aligned} \mathbb{P}_f[A_\Gamma \in A_{\mathcal{D}}] &= \lim_{N \rightarrow \infty} \mathbb{P}[\Gamma \in \mathcal{D} \mid \Gamma \in \mathcal{G}^{N, f(N)}] \\ &\geq \lim_{N \rightarrow \infty} \mathbb{P}[\Gamma \in \mathcal{B} \mid \Gamma \in \mathcal{G}^{N, f(N)}] \quad (\text{by } (***)) \\ &= \mathbb{P}_f[A_\Gamma \in A_{\mathcal{B}}]. \end{aligned} \quad \square$$

We now move towards determining for which (nondecreasing divergent) functions a random Artin group is asymptotically almost surely 2-dimensional, or not 2-dimensional. In view of [Lemma 3.2](#), looking at $(2, 2)$ -free Artin groups will be enough to give a conclusion for 2-dimensional Artin groups. The result we want to prove is the following:

Theorem 3.3 *Let $f : \mathbb{N} \rightarrow \mathbb{N}$; then, for a random Artin group A_Γ (with respect to f):*

- (1) *If $f(N) \succ N^{3/2}$, then asymptotically almost surely A_Γ is 2-dimensional.*
- (2) *If $f(N) \prec N^{3/2}$, then asymptotically almost surely A_Γ is not 2-dimensional.*
- (3) *If $f(N) \simeq N^{3/2}$ then $\mathbb{P}_f[A_\Gamma \in A_{\mathcal{D}}] < 1$. Moreover, if $f(N) = N^{3/2}$ then $\mathbb{P}_f[A_\Gamma \in A_{\mathcal{D}}] \leq 2/3$.*

Proof Let f be any nondecreasing, divergent function. We need to compute $\mathbb{P}_f[A_\Gamma \in A_{\mathcal{D}}]$. In view of [Lemma 3.2](#), it is enough to compute $\mathbb{P}_f[A_\Gamma \in A_{\mathcal{B}}]$, ie the probability that an Artin group A_Γ picked at random is $(2, 2)$ -free. To do this, we will use the second moment method.

Let us consider a graph $\Gamma \in \mathcal{G}^{N, f(N)}$. For any ordered triplet (v_1, v_2, v_3) of distinct vertices of Γ , we let $I_{(v_1, v_2, v_3)} : \mathcal{G}^{N, f(N)} \rightarrow \{0, 1\}$ be the random variable which takes 1 on $\Gamma \in \mathcal{G}^{N, f(N)}$ precisely when (v_1, v_2, v_3) spans a triangle with $m_{v_1, v_2} = m_{v_1, v_3} = 2$. We let

$$X = \left(\sum_{(v_1, v_2, v_3) \in V(\Gamma)^3} I_{(v_1, v_2, v_3)} \right) : \mathcal{G}^{N, f(N)} \rightarrow \mathbb{N},$$

where the sum is taken over all triplets of distinct vertices. The variable X counts the number of pairs of adjacent edges labelled by a 2, twice (because of the permutation of these edges).

We can compute the expectation $\mathbb{E}[I_{(v_1, v_2, v_3)}] = f(N)^{-2}$ and hence

$$\mathbb{E}[X] = \sum_{(v_1, v_2, v_3)} \mathbb{E}[I_{(v_1, v_2, v_3)}] = N(N-1)(N-2)f(N)^{-2} \sim N^3 f(N)^{-2}.$$

Now, we use the second moment method, as in [4, Theorem 6]:

$$\mathbb{P}[X \neq 0] \geq \frac{\mathbb{E}[X]^2}{\mathbb{E}[X^2]}.$$

We have already computed $\mathbb{E}[X]$, so we now compute $\mathbb{E}[X^2]$ by dividing into several cases the sum

$$X^2 = \sum I_{(v_1, v_2, v_3)} I_{(w_1, w_2, w_3)}.$$

Note that the sum is taken over all ordered triplets (v_1, v_2, v_3) and (w_1, w_2, w_3) of vertices, where the v_i are distinct, and the w_i are distinct. Also note that if one of the two triangles does not have two edges labelled by 2, then the corresponding term in the sum is trivial. In other words, it is enough to only sum over pairs of triangles that both have at least two edges labelled by 2. In a triangle (v_1, v_2, v_3) such that $m_{v_1 v_2} = m_{v_1 v_3} = 2$, we shall call v_1 the *central* vertex of the triangle. The different cases are treated below. They can be seen in Figure 2.

Case 1 Let X_1 denote the sum of products $I_{(v_1, v_2, v_3)} I_{(w_1, w_2, w_3)}$ such that no vertex appears in both triples. Then

$$\mathbb{E}[X_1] = \frac{N!}{(N-6)!} f(N)^{-4} \sim N^6 f(N)^{-4}.$$

Case 2 Let X_2 denote the sum of products $I_{(v_1, v_2, v_3)} I_{(w_1, w_2, w_3)}$ such that these two triangles share exactly one vertex and the vertex they share is central in both triangles (ie $v_1 = w_1$). Then we have

$$\mathbb{E}[X_2] = \frac{N!}{(N-5)!} f(N)^{-4} \sim N^5 f(N)^{-4}.$$

Case 3 Let X_3 denote the sum of products $I_{(v_1, v_2, v_3)} I_{(w_1, w_2, w_3)}$ such that these two triangles share exactly one vertex, where this vertex is the central vertex for one triangle and not a central vertex for the other triangle (for example $v_2 = w_1$). In this case

$$\mathbb{E}[X_3] = 4 \frac{N!}{(N-5)!} f(N)^{-4} \sim 4N^5 f(N)^{-4}.$$

Case 4 Let X_4 denote the sum of products $I_{(v_1, v_2, v_3)} I_{(w_1, w_2, w_3)}$ such that these two triangles share exactly one vertex, where this vertex is not central for either triangle (for example $v_2 = w_2$). Then

$$\mathbb{E}[X_4] = 4 \frac{N!}{(N-5)!} f(N)^{-4} \sim 4N^5 f(N)^{-4}.$$




Figure 2: From top-left to bottom-right: the ten cases described in the proof of [Theorem 3.3](#). The edges that are not explicitly labelled by 2 can be labelled by any coefficient, including ∞ .

Case 5 Let X_5 denote the sum of products $I_{(v_1, v_2, v_3)} I_{(w_1, w_2, w_3)}$ such that these two triangles share exactly two vertices and these two vertices are not central for either triangle (for example $v_2 = w_2$ and $v_3 = w_3$). In this case

$$\mathbb{E}[X_5] = 2 \frac{N!}{(N-4)!} f(N)^{-4} \sim 2N^4 f(N)^{-4}.$$

Case 6 Let X_6 denote the sum of products $I_{(v_1, v_2, v_3)} I_{(w_1, w_2, w_3)}$ such that these two triangles share exactly two vertices and one of these is central in both triangles and the other is not (for example $v_1 = w_1$ and $v_3 = w_2$). In this case

$$\mathbb{E}[X_6] = 4 \frac{N!}{(N-4)!} f(N)^{-3} \sim 4N^4 f(N)^{-3}.$$

Case 7 Let X_7 denote the sum of products $I_{(v_1, v_2, v_3)} I_{(w_1, w_2, w_3)}$ such that these two triangles share exactly two vertices where one of these is central for the triangle (v_1, v_2, v_3) but not for (w_1, w_2, w_3) , and the other vertex is central for the triangle (w_1, w_2, w_3) but not for (v_1, v_2, v_3) (for example $v_1 = w_3$ and $w_1 = v_3$). In this case

$$\mathbb{E}[X_7] = 4 \frac{N!}{(N-4)!} f(N)^{-3} \sim 4N^4 f(N)^{-3}.$$

Case 8 Let X_8 denote the sum of products $I_{(v_1, v_2, v_3)} I_{(w_1, w_2, w_3)}$ such that these two triangles share exactly two vertices where one of these is central for the triangle (v_1, v_2, v_3) but none of the two vertices is central for (w_1, w_2, w_3) (for example $v_1 = w_2$ and $v_3 = w_3$). In this case

$$\mathbb{E}[X_8] = 4 \frac{N!}{(N-4)!} f(N)^{-4} \sim 4N^4 f(N)^{-4}.$$

Case 9 Let X_9 denote the sum of products $I_{(v_1, v_2, v_3)} I_{(w_1, w_2, w_3)}$ such that these two triangles share all three vertices, and such the central vertices of both triangles are the same (ie $v_1 = w_1$). In this case

$$\mathbb{E}[X_9] = 2 \frac{N!}{(N-3)!} f(N)^{-2} \sim 2N^3 f(N)^{-2}.$$

Case 10 Let X_{10} denote the sum of products $I_{(v_1, v_2, v_3)} I_{(w_1, w_2, w_3)}$ such that these two triangles share all three vertices, and such that the central vertex of the first triangle is not the central vertex of the second triangle (for example $v_1 = w_2$). We get

$$\mathbb{E}[X_{10}] = 4 \frac{N!}{(N-3)!} f(N)^{-3} \sim 2N^3 f(N)^{-3}.$$

Therefore, we have

$$\begin{aligned} \frac{\mathbb{E}[X^2]}{\mathbb{E}[X]^2} &= \sum_{i=1}^8 \frac{\mathbb{E}[X_i]}{\mathbb{E}[X]^2} \\ &\sim \frac{N^6 f(N)^{-4} + 9N^5 f(N)^{-4} + 6N^4 f(N)^{-4} + 8N^4 f(N)^{-3} + 2N^3 f(N)^{-3} + 2N^3 f(N)^{-2}}{N^6 f(N)^{-4}} \\ &\sim 1 + \frac{9}{N} + \frac{6}{N^2} + \frac{8f(N)}{N^2} + \frac{2f(N)}{N^3} + \frac{2f(N)^2}{N^3}. \end{aligned}$$

Hence, if $f(N) < N^{3/2}$ then by definition there exists a nondecreasing divergent function h such that $f(N)h(N) = N^{3/2}$. In this case we get

$$\mathbb{P}[X \neq 0] \geq \left(\frac{\mathbb{E}[X^2]}{\mathbb{E}[X]^2} \right)^{-1} \sim \left(1 + \frac{9}{N} + \frac{6}{N^2} + \frac{8}{h(N)N^{1/2}} + \frac{4}{h(N)N^{3/2}} + \frac{2}{h(N)^2} \right)^{-1}.$$

When $f(N) < N^{3/2}$, we obtain

$$\mathbb{P}_f[A_\Gamma \in A_{\mathfrak{B}}] = \lim_{N \rightarrow \infty} \mathbb{P}[\Gamma \in \mathfrak{B} \mid \Gamma \in \mathcal{G}^{N, f(N)}] = \lim_{N \rightarrow \infty} \mathbb{P}[X = 0] = 1 - \lim_{N \rightarrow \infty} \mathbb{P}[X \neq 0] = 0.$$

Thus asymptotically almost surely A_Γ is not (2, 2)-free. In view of [Lemma 3.2](#), this also means that asymptotically almost surely A_Γ is not of dimension 2, this proves item (2) in [Theorem 3.3](#).

If $f(N) \simeq N^{3/2}$ then the quotient $f(N)/N^{3/2}$ tends to M for some constant $M > 0$. Hence in this case,

$$\mathbb{P}[X \neq 0] \gtrsim \left(1 + \frac{9}{N} + \frac{6}{N^2} + \frac{8f(N)}{N^2} + \frac{2f(N)}{N^3} + \frac{2f(N)^2}{N^3} \right)^{-1} \sim (1 + 2M^2)^{-1} > 0.$$

Therefore $\mathbb{P}_f[A_\Gamma \in A_{\mathfrak{B}}] < 1$ at $f(N) \simeq N^{3/2}$ and hence by [Lemma 3.2](#) we have that $\mathbb{P}_f[A_\Gamma \in A_{\mathfrak{D}}] < 1$.

We note that the above calculation allows us to find a better upper bound for $\mathbb{P}_f[A_\Gamma \in A_{\mathfrak{B}}]$ at $f(N) = N^{3/2}$. Indeed, this implies that $M = 1$ and hence we get $\mathbb{P}[X \neq 0] \gtrsim \frac{1}{3}$, and so at $f(N) = N^{3/2}$ we have $\mathbb{P}_f[A_\Gamma \in A_{\mathfrak{B}}] \leq \frac{2}{3}$. Hence by [Lemma 3.2](#), this proves item (3) in the theorem.

We note that $\mathbb{P}[\Gamma \in \mathfrak{B} \mid \Gamma \in \mathcal{G}^{N, f(N)}] = 1 - \mathbb{P}[X \geq 1]$ and by the Markov inequality,

$$\mathbb{P}[X \geq 1] \leq \mathbb{E}[X] \leq N^3 f(N)^{-2}.$$

Hence if $f(N) \succ N^{3/2}$ then we can write $f(N) = N^{3/2}g(N)$ for some nondecreasing divergent function $g: \mathbb{N} \rightarrow \mathbb{N}$ and in this case

$$\mathbb{P}[X \geq 1] \leq \frac{1}{g(N)^2}.$$

Therefore, for $f(N) \succ N^{3/2}$ we have

$$\mathbb{P}_f[A_\Gamma \in A_{\mathfrak{B}}] = \lim_{N \rightarrow \infty} \mathbb{P}[\Gamma \in \mathfrak{B} \mid \mathcal{G}^{N, f(N)}] = 1 - \lim_{N \rightarrow \infty} \mathbb{P}[X \geq 1] \geq 1 - \lim_{N \rightarrow \infty} \frac{1}{g(N)^2} = 1.$$

In particular, asymptotically almost surely A_Γ is $(2, 2)$ -free. By applying Lemma 3.2 (as $f(N) \succ N$), we get that asymptotically almost surely A_Γ is 2-dimensional. This proves item (1) and hence Theorem 3.3. \square

Before stating a corollary which will be a refinement of Corollary 2.12, we prove a small lemma which will allow us to study the problem of (virtual) cocompact cubulation of random Artin groups. We note that the class \mathcal{P} defined below is point 3 in [8, Conjecture B].

Lemma 3.4 *Let \mathcal{P} be the class of defining graphs Γ for which there exist 4 distinct $a, b, c, d \in V(\Gamma)$ such that $m_{ab} \notin \{2, \infty\}, m_{ac}, m_{bd} \neq \infty$ and $m_{ad}, m_{bc} \neq 2$. Then $A_{\mathcal{P}}$ is uniformly large.*

Proof Let $f: \mathbb{N} \rightarrow \mathbb{N}$ be any nondecreasing, divergent function. Fix a, b, c and d to be any distinct vertices. The probability that these vertices and their corresponding coefficients satisfy the defining condition of \mathcal{P} is at exactly

$$\left(\frac{f(N)-1}{f(N)}\right)^4 \left(\frac{f(N)-2}{f(N)}\right).$$

This tends to 1 for all nondecreasing divergent functions f . \square

Corollary 3.5 *Let $f: \mathbb{N} \rightarrow \mathbb{N}$ be a function satisfying $f(N) \succ N^{3/2}$. Then a random Artin group A_Γ (with respect to f) satisfies any of the following properties asymptotically almost surely:*

- (1) A_Γ is torsion-free;
- (2) A_Γ has trivial centre;
- (3) A_Γ has solvable word and conjugacy problems;
- (4) A_Γ satisfies the $K(\pi, 1)$ -conjecture;
- (5) the set of parabolic subgroups of A_Γ is closed under arbitrary intersections;
- (6) A_Γ is acylindrically hyperbolic;
- (7) A_Γ satisfies the Tits alternative;
- (8) A_Γ is not virtually cocompactly cubulated.

Proof By Theorem 3.3, A_Γ is asymptotically almost surely 2-dimensional. Using Lemma 3.2, A_Γ is also asymptotically almost surely $(2, 2)$ -free. Using Lemma 2.8, we also know that A_Γ is asymptotically almost surely irreducible. By Lemma 3.4 we know that A_Γ is asymptotically almost surely in $A_{\mathcal{P}}$. Using Lemma 2.11 three times, this ensures that A_Γ is asymptotically almost surely in the class

$$A_{\mathcal{H}} := A_{\text{Irr}} \cap A_{\mathfrak{D}} \cap A_{\mathfrak{B}} \cap A_{\mathcal{P}}.$$

Note that the results given for points (1), (2), (3), (4), (5), (7) and (10) of [Conjecture 2.2](#) concern families of Artin groups that all contain $A_{\mathfrak{A}}$. In particular, every Artin group of $A_{\mathfrak{A}}$ satisfies the first seven points of the [Corollary 3.5](#). For point (8) of [Corollary 3.5](#), we note that by [[8](#), Theorem E], if $A_\Gamma \in A_{\mathfrak{A}} \cap A_{\mathfrak{B}}$ then A_Γ is not virtually cocompactly cubulated. \square

Finding out the exact probability for an Artin group to be 2-dimensional (or equivalently, (2, 2)-free) at $f(N) = N^{3/2}$ requires more work. In [Theorem 3.3](#), we gave an upper bound for this probability. The goal of the following lemma is to give an explicit formula for the value of $\mathbb{P}_f[A_\Gamma \in A_{\mathfrak{B}}]$ at $f(N) = N^{3/2}$. Later, we give a conjecture on the exact value.

Lemma 3.6 *For all nondecreasing, divergent functions $f : \mathbb{N} \rightarrow \mathbb{N}$ we have that*

$$\mathbb{P}_f[A_\Gamma \in A_{\mathfrak{B}}] = \lim_{N \rightarrow \infty} \left(\frac{f(N) - 1}{f(N)} \right)^{\binom{N}{2}} \left(\sum_{k=1}^{\lfloor N/2 \rfloor} \frac{N!(f(N) - 1)^{-k}}{(N - 2k)! k! 2^k} + 1 \right).$$

Proof Let E_k be the family of defining graphs that have exactly k edges labelled by a 2, and consider the associated family A_{E_k} of Artin groups. Note that each edge is attached to two vertices, so by the pigeonhole principle, if $k > N/2$ then $\mathbb{P}_f[\Gamma \in \mathfrak{B} \cap E_k] = 0$. Hence

$$\mathbb{P}[\Gamma \in \mathfrak{B} \mid \Gamma \in \mathcal{G}^{N, f(N)}] = \sum_{k=0}^{\lfloor N/2 \rfloor} \mathbb{P}[\Gamma \in \mathfrak{B} \cap E_k \mid \Gamma \in \mathcal{G}^{N, f(N)}].$$

As usual, the total number of graphs in $\mathcal{G}^{N, f(N)}$ is $f(N) \binom{N}{2}$. On the other hand, we must compute how many of these graphs have exactly k edges labelled by a 2, while these edges are never adjacent.

First of all, when $k = 0$, we have $\mathbb{P}[\Gamma \in \mathfrak{B} \cap E_k \mid \Gamma \in \mathcal{G}^{N, f(N)}] = ((f(N) - 1)/f(N)) \binom{N}{2}$.

For the case when $0 < k \leq \lfloor N/2 \rfloor$, we look at how many ways we have of placing the k edges labelled by a 2. For the first such edge, we have $\binom{N}{2}$ choices. The two vertices of the first edge must not appear in any other edge labelled by a 2, so for the second edge we only have $\binom{N-2}{2}$ choices left. This goes on until the k^{th} edge labelled by a 2, for which we have $\binom{N-2(k-1)}{2}$ choices. As the order in which we have chosen these edges do not matter, we must divide this product by $k!$. Now for the remaining $\binom{N}{2} - k$ edges, we can use any label other than a 2. Hence we multiply the previous product by $(f(N) - 1) \binom{N}{2}^{-k}$. Hence, for $0 < k \leq \lfloor N/2 \rfloor$, we have

$$\mathbb{P}[\Gamma \in \mathfrak{B} \cap E_k \mid \Gamma \in \mathcal{G}^{N, f(N)}] = \frac{(f(N) - 1) \binom{N}{2}^{-k} \cdot \prod_{i=0}^{k-1} \binom{N-2i}{2}}{f(N) \binom{N}{2} \cdot k!}.$$

Therefore,

$$\begin{aligned} \mathbb{P}_f[A_\Gamma \in A_{\mathfrak{B}}] &= \lim_{N \rightarrow \infty} \sum_{k=1}^{\lfloor N/2 \rfloor} \mathbb{P}[\Gamma \in \mathfrak{B} \cap E_k \mid \Gamma \in \mathcal{G}^{N, f(N)}] + \mathbb{P}[\Gamma \in \mathfrak{B} \cap E_0 \mid \Gamma \in \mathcal{G}^{N, f(N)}] \\ &= \lim_{N \rightarrow \infty} \sum_{k=1}^{\lfloor N/2 \rfloor} \frac{(f(N) - 1) \binom{N}{2}^{-k} \cdot \prod_{i=0}^{k-1} \binom{N-2i}{2}}{f(N) \binom{N}{2} \cdot k!} + \left(\frac{f(N) - 1}{f(N)} \right) \binom{N}{2} \end{aligned}$$

$$= \lim_{N \rightarrow \infty} \left(\frac{f(N) - 1}{f(N)} \right)^{\binom{N}{2}} \left(\sum_{k=1}^{\lfloor N/2 \rfloor} \frac{N!(f(N) - 1)^{-k}}{(N - 2k)! k! 2^k} + 1 \right),$$

where we go from the second to the third line by noting that

$$\prod_{i=0}^{k-1} \binom{N - 2i}{2} = \frac{1}{2^k} N(N - 1)(N - 2) \cdots (N - 2(k - 1))(N - 2(k - 1) - 1) = \frac{N!}{(N - 2k)! 2^k}. \quad \square$$

Now, by Lemma 3.2 at $f(N) = N^{3/2}$ we have $\mathbb{P}_f[A_\Gamma \in A_{\mathfrak{A}}] = \mathbb{P}_f[A_\Gamma \in A_{\mathfrak{B}}]$; hence Lemma 3.6 also holds for $\mathbb{P}_f[A_\Gamma \in A_{\mathfrak{A}}]$. We have computed this expression in *Python* for N up to 190, which leads us to the following conjecture.

Conjecture 3.7 For $f(N) = N^{3/2}$ we have

$$\mathbb{P}_f[A_\Gamma \in A_B] = 1 - e^{-1}.$$

In particular, we also have

$$\mathbb{P}_f[A_\Gamma \in A_{\mathfrak{A}}] = 1 - e^{-1}.$$

4 Acylindrical hyperbolicity and centres

Two open questions in the study of Artin groups is whether all irreducible nonspherical Artin groups are acylindrically hyperbolic and have trivial centres (see Conjecture 2.2). In this section, we study these two aspects of Artin groups for another family of Artin groups, that we will denote $A_{\mathcal{C}}$. The families of Artin groups studied in Sections 2 and 3 are very large when $f(N)$ grows fast enough compared to N . While the spirit of this section resembles that of Sections 2 and 3, $A_{\mathcal{C}}$ will turn out to be very large when $f(N)$ grows slowly enough compared to N .

Definition 4.1 A graph Γ is said to be a cone if it has a join decomposition as a subgraph consisting of a single vertex v_0 and a subgraph Γ' such that $\Gamma = v_0 * \Gamma'$. Let \mathcal{C} be the class of defining graphs that are cones and \mathcal{C}^C the class of defining graphs which are not cones.

Recall that *Irr* is the class of irreducible graphs. By [13, Theorem 1.4], we have that if Γ has at least 3 vertices, is irreducible and is not a cone then A_Γ is acylindrically hyperbolic. Hence it suffices to find the probability that a random Artin group is irreducible and is not a cone.

Proposition 4.2 For all $\alpha \in (0, 1)$ and all nondecreasing functions $f(N) \prec N^{1-\alpha}$ we have that

$$\mathbb{P}_f[A_\Gamma \in A_{\mathcal{C}^C}] = 1.$$

Proof Fix $\alpha \in (0, 1)$ and $f(N) \prec N^{1-\alpha}$ a nondecreasing divergent function. Then, by definition, there exists a nondecreasing divergent function h such that $f(N)h(N) = N^{1-\alpha}$.

By the definition of a cone and by a union bound, we get

$$\begin{aligned} \mathbb{P}[\Gamma \in \mathcal{C} \mid \Gamma \in \mathcal{G}^{N,f(N)}] &\leq \sum_{v_0 \in V(\Gamma)} \mathbb{P}[\forall u \in V(\Gamma) - v_0 : m_{u,v_0} \neq \infty \mid \Gamma \in \mathcal{G}^{N,f(N)}] \\ &= \sum_{v_0 \in V(\Gamma)} \left(\frac{f(N) - 1}{f(N)} \right)^{N-1} \\ &= N \left(\frac{f(N) - 1}{f(N)} \right)^{N-1} \\ &= N \left(\left(\frac{f(N) - 1}{f(N)} \right)^{f(N)} \right)^{h(N)N^\alpha} \left(\frac{f(N)}{f(N) - 1} \right). \end{aligned}$$

Thus,

$$\mathbb{P}_f[A_\Gamma \in A_{\mathcal{C}}] = \lim_{N \rightarrow \infty} \mathbb{P}[\Gamma \in \mathcal{C} \mid \Gamma \in \mathcal{G}^{N,f(N)}] = \lim_{N \rightarrow \infty} N e^{-N^\alpha h(N)} = 0.$$

Hence for $f(N) \prec N^{1-\alpha}$ we have $\mathbb{P}_f[A_\Gamma \in A_{\mathcal{C}}] = 1$, proving the proposition. □

Corollary 4.3 *Let $\alpha \in (0, 1)$ and let $f(N) \prec N^{1-\alpha}$ be a nondecreasing divergent function. Then a random Artin group (with respect to f) asymptotically almost surely is acylindrically hyperbolic and has a trivial centre.*

Proof We note that by Lemmas 2.8 and 2.11 we have $\mathbb{P}_f[A_\Gamma \in A_{\text{irr}} \cap A_{\mathcal{C}}] = \mathbb{P}_f[A_\Gamma \in A_{\mathcal{C}}]$. As we noted above, by [13, Theorem 1.4], if Γ is irreducible and not a cone then A_Γ is acylindrically hyperbolic. Hence, by Proposition 4.2, for a function f as in the statement of the corollary, we get that a random Artin group (relatively to f) is asymptotically almost surely irreducible and a cone, hence asymptotically almost surely acylindrically hyperbolic.

Further, by [5, Theorem 3.3], we have that if Γ is irreducible and not a cone then A_Γ has trivial centre. Hence a random Artin group (relatively to f) asymptotically almost surely has a trivial centre. □

Let $\alpha \in (0, 1)$, by Corollary 4.3 and Corollary 3.5-(6), we have shown that for all nondecreasing divergent functions f such that either

- $f(N) \prec N^{1-\alpha}$, or
- $f(N) \succ N^{3/2}$,

a random Artin group A_Γ (relatively to f) is asymptotically almost surely acylindrically hyperbolic and has trivial centre. This motivates the following:

Question 4.4 For which nondecreasing divergent functions f do we have that a random Artin group (relatively to f) is asymptotically almost surely acylindrically hyperbolic and has trivial centre?

References

[1] **J Behrstock, MF Hagen, A Sisto**, *Thickness, relative hyperbolicity, and randomness in Coxeter groups*, *Algebr. Geom. Topol.* 17 (2017) 705–740 [MR](#) [Zbl](#)

- [2] **MA Blufstein**, *Parabolic subgroups of two-dimensional Artin groups and systolic-by-function complexes*, Bull. Lond. Math. Soc. 54 (2022) 2338–2350 [MR](#) [Zbl](#)
- [3] **R Charney, M W Davis**, *The $K(\pi, 1)$ -problem for hyperplane complements associated to infinite reflection groups*, J. Amer. Math. Soc. 8 (1995) 597–627 [MR](#) [Zbl](#)
- [4] **R Charney, M Farber**, *Random groups arising as graph products*, Algebr. Geom. Topol. 12 (2012) 979–995 [MR](#) [Zbl](#)
- [5] **R Charney, R Morris-Wright**, *Artin groups of infinite type: trivial centers and acylindrical hyperbolicity*, Proc. Amer. Math. Soc. 147 (2019) 3675–3689 [MR](#) [Zbl](#)
- [6] **M Cumplido, A Martin, N Vaskou**, *Parabolic subgroups of large-type Artin groups*, Math. Proc. Cambridge Philos. Soc. 174 (2023) 393–414 [MR](#) [Zbl](#)
- [7] **A Deibel**, *Random Coxeter groups*, Int. J. Algebra Comput. 30 (2020) 1305–1321 [MR](#) [Zbl](#)
- [8] **T Haettel**, *Virtually cocompactly cubulated Artin–Tits groups*, Int. Math. Res. Not. 2021 (2021) 2919–2961 [MR](#) [Zbl](#)
- [9] **T Haettel**, *XXL type Artin groups are CAT(0) and acylindrically hyperbolic*, Ann. Inst. Fourier (Grenoble) 72 (2022) 2541–2555 [MR](#) [Zbl](#)
- [10] **M Hagen, A Martin, A Sisto**, *Extra-large type Artin groups are hierarchically hyperbolic*, Math. Ann. 388 (2024) 867–938 [MR](#) [Zbl](#)
- [11] **J Huang, D Osajda**, *Metric systolicity and two-dimensional Artin groups*, Math. Ann. 374 (2019) 1311–1352 [MR](#) [Zbl](#)
- [12] **J Huang, D Osajda**, *Large-type Artin groups are systolic*, Proc. Lond. Math. Soc. 120 (2020) 95–123 [MR](#) [Zbl](#)
- [13] **M Kato, S-i Oguni**, *Acylindrical hyperbolicity of Artin groups associated with graphs that are not cones*, Groups Geom. Dyn. 18 (2024) 1291–1316 [MR](#) [Zbl](#)
- [14] **H van der Lek**, *The homotopy type of complex hyperplane complements*, PhD thesis, Katholieke Universiteit te Nijmegen (1983) Available at <https://repository.ubn.ru.nl/handle/2066/148301>
- [15] **A Martin**, *The Tits alternative for two-dimensional Artin groups and Wise’s power alternative*, J. Algebra 656 (2024) 294–323 [MR](#) [Zbl](#)
- [16] **A Martin, N Vaskou**, *Characterising large-type Artin groups*, Bull. Lond. Math. Soc. (online publication August 2024)
- [17] **N Vaskou**, *Acylindrical hyperbolicity for Artin groups of dimension 2*, Geom. Dedicata 216 (2022) art. id. 7 [MR](#) [Zbl](#)
- [18] **N Vaskou**, *Automorphisms of large-type free-of-infinity Artin groups*, Geom. Dedicata 219 (2025) art. no. 16 [MR](#) [Zbl](#)

School of Mathematics & Computer Sciences, Heriot-Watt University
Edinburgh, United Kingdom

Department of Mathematics, University of Bristol
Bristol, United Kingdom

ag2017@hw.ac.uk, nicolas.vaskou@bristol.ac.uk

Received: 25 January 2023 Revised: 30 October 2023

ALGEBRAIC & GEOMETRIC TOPOLOGY

msp.org/agt

EDITORS

PRINCIPAL ACADEMIC EDITORS

John Etnyre
etnyre@math.gatech.edu
Georgia Institute of Technology

Kathryn Hess
kathryn.hess@epfl.ch
École Polytechnique Fédérale de Lausanne

BOARD OF EDITORS

Julie Bergner	University of Virginia jeb2md@eservices.virginia.edu	Thomas Koberda	University of Virginia thomas.koberda@virginia.edu
Steven Boyer	Université du Québec à Montréal cohf@math.rochester.edu	Markus Land	LMU München markus.land@math.lmu.de
Tara E Brendle	University of Glasgow tara.brendle@glasgow.ac.uk	Christine Lescop	Université Joseph Fourier lescop@ujf-grenoble.fr
Indira Chatterji	CNRS & Univ. Côte d'Azur (Nice) indira.chatterji@math.cnrs.fr	Norihiko Minami	Yamato University minami.norihiko@yamato-u.ac.jp
Octav Cornea	Université de Montreal cornea@dms.umontreal.ca	Andrés Navas	Universidad de Santiago de Chile andres.navas@usach.cl
Alexander Dranishnikov	University of Florida dranish@math.ufl.edu	Robert Oliver	Université Paris 13 bobol@math.univ-paris13.fr
Tobias Ekholm	Uppsala University, Sweden tobias.ekholm@math.uu.se	Jessica S Purcell	Monash University jessica.purcell@monash.edu
Mario Eudave-Muñoz	Univ. Nacional Autónoma de México mario@matem.unam.mx	Birgit Richter	Universität Hamburg birgit.richter@uni-hamburg.de
David Futer	Temple University dfuter@temple.edu	Jérôme Scherer	École Polytech. Féd. de Lausanne jerome.scherer@epfl.ch
John Greenlees	University of Warwick john.greenlees@warwick.ac.uk	Vesna Stojanoska	Univ. of Illinois at Urbana-Champaign vesna@illinois.edu
Ian Hambleton	McMaster University ian@math.mcmaster.ca	Zoltán Szabó	Princeton University szabo@math.princeton.edu
Matthew Hedden	Michigan State University mhedden@math.msu.edu	Maggy Tomova	University of Iowa maggy-tomova@uiowa.edu
Kristen Hendricks	Rutgers University kristen.hendricks@rutgers.edu	Daniel T Wise	McGill University, Canada daniel.wise@mcgill.ca
Hans-Werner Henn	Université Louis Pasteur henn@math.u-strasbg.fr	Lior Yanovski	Hebrew University of Jerusalem lior.yanovski@gmail.com
Daniel Isaksen	Wayne State University isaksen@math.wayne.edu		

See inside back cover or msp.org/agt for submission instructions.

The subscription price for 2025 is US \$760/year for the electronic version, and \$1110/year (+\$75, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Algebraic & Geometric Topology is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [Current Mathematical Publications](#) and the [Science Citation Index](#).

Algebraic & Geometric Topology (ISSN 1472-2747 printed, 1472-2739 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840.

AGT peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**

nonprofit scientific publishing

<https://msp.org/>

© 2025 Mathematical Sciences Publishers

ALGEBRAIC & GEOMETRIC TOPOLOGY

Volume 25 Issue 3 (pages 1265–1915) 2025

A-polynomials, Ptolemy equations and Dehn filling	1265
JOSHUA A HOWIE, DANIEL V MATHEWS and JESSICA S PURCELL	
The Alexandrov theorem for $2 + 1$ flat radiant spacetimes	1321
LÉO MAXIME BRUNSWIC	
Real algebraic overtwisted contact structures on 3-spheres	1377
ŞEYMA KARADERELI and FERİT ÖZTÜRK	
Fully augmented links in the thickened torus	1411
ALICE KWON	
Unbounded \mathfrak{sl}_3 -laminations and their shear coordinates	1433
TSUKASA ISHIBASHI and SHUNSUKE KANO	
Bridge trisections and Seifert solids	1501
JASON JOSEPH, JEFFREY MEIER, MAGGIE MILLER and ALEXANDER ZUPAN	
Random Artin groups	1523
ANTOINE GOLDSBOROUGH and NICOLAS VASKOU	
A deformation of Asaeda–Przytycki–Sikora homology	1545
ZHENKUN LI, YI XIE and BOYU ZHANG	
Cubulating a free-product-by-cyclic group	1561
FRANÇOIS DAHMANI and SURAJ KRISHNA MEDA SATISH	
Virtual domination of 3-manifolds, III	1599
HONGBIN SUN	
The Kakimizu complex for genus one hyperbolic knots in the 3-sphere	1667
LUIS G VALDEZ-SÁNCHEZ	
Band diagrams of immersed surfaces in 4-manifolds	1731
MARK HUGHES, SEUNGWON KIM and MAGGIE MILLER	
Anosov flows and Liouville pairs in dimension three	1793
THOMAS MASSONI	
Hamiltonian classification of toric fibres and symmetric probes	1839
JOÉ BRENDEL	
An example of higher-dimensional Heegaard Floer homology	1877
YIN TIAN and TIANYU YUAN	
Fibered 3-manifolds and Veech groups	1897
CHRISTOPHER J LEININGER, KASRA RAFI, NICHOLAS ROUSE, EMILY SHINKLE and YVON VERBERNE	