

AG
T

*Algebraic & Geometric
Topology*

Volume 25 (2025)

A minimality property for knots without Khovanov 2-torsion

ONKAR SINGH GUJRAL

JOSHUA WANG



A minimality property for knots without Khovanov 2-torsion

ONKAR SINGH GUJRAL

JOSHUA WANG

A conjecture of Shumakovitch states that every nontrivial knot has 2-torsion in its Khovanov homology. We show that if a knot K has no 2-torsion in its Khovanov homology, then the rank of its reduced Khovanov homology is minimal among all knots obtainable from K by a proper rational tangle replacement. It follows, for example, that unknotting number 1 knots have 2-torsion in their Khovanov homology.

57K18

Shumakovitch conjectured that every nontrivial knot has an element of order 2 in its Khovanov homology [10, Conjecture 1]. The conjecture has been verified for some infinite families of knots (see, for example, [2; 9; 10]) and has withstood large computational searches. In this note, we provide topological evidence for the conjecture, and we verify the conjecture for a large class of knots that include all unknotting number 1 knots.

Two links differ by a *rational tangle replacement* if they agree outside of a ball, and if within the ball, each is a rational tangle. A rational tangle replacement is *proper* if the arcs of the two rational tangles connect the same end points [4; 8]. Changing a crossing is an example of a proper rational tangle replacement, while resolving a crossing is an example of a nonproper rational tangle replacement. In the following statement, $\text{Kh}(K)$ and $\overline{\text{Kh}}(K)$ denote the unreduced and reduced Khovanov homology groups of K , respectively, thought of as abelian groups with bigradings suppressed.

Theorem 1 *Suppose K is a knot such that there is no 2-torsion in $\text{Kh}(K)$. If J is a knot that differs from K by a proper rational tangle replacement, then*

$$\text{rank } \overline{\text{Kh}}(K) \leq \text{rank } \overline{\text{Kh}}(J).$$

Corollary 2 *Any knot whose unknotting number is 1 has 2-torsion in its Khovanov homology. More generally, if K is a nontrivial knot that can be obtained from the unknot or a trefoil by a proper rational tangle replacement, then $\text{Kh}(K)$ contains 2-torsion.*

Proof of Corollary 2 Let J be the unknot or a trefoil, and let K be obtained from J by a proper rational tangle replacement. If there is no 2-torsion in $\text{Kh}(K)$, then $\text{rank } \overline{\text{Kh}}(K) \leq \text{rank } \overline{\text{Kh}}(J) \leq 3$ by Theorem 1. The rank of $\overline{\text{Kh}}(K)$ cannot be 3 since then K would be a trefoil [3, Theorem 1.4], which has 2-torsion in its Khovanov homology. Since the rank of $\overline{\text{Kh}}(K)$ is odd, it must be 1, and so K is the unknot [6]. \square

Our proof of Theorem 1 combines the main result of Iltgen, Lewark, and Marino [4] with an observation of Kotelskiy, Watson, and Zibrowius [5, Proposition 9.3] using the following lemma.

Lemma 3 *Let \mathbb{F} be a field, and suppose M and N are finitely generated modules over the polynomial ring $\mathbb{F}[X]$ of the form*

$$M = (\mathbb{F}[X])^r \oplus \bigoplus_{i=1}^m \frac{\mathbb{F}[X]}{X^{a_i}}, \quad N = (\mathbb{F}[X])^s \oplus \bigoplus_{i=1}^n \frac{\mathbb{F}[X]}{X^{b_i}},$$

where $r, m, s, n \geq 0$ and $a_1, \dots, a_m, b_1, \dots, b_n \geq 1$. Furthermore, suppose $f: M \rightarrow N$ and $g: N \rightarrow M$ are $\mathbb{F}[X]$ -module maps for which $f \circ g = X$ and $g \circ f = X$. If the numbers a_1, \dots, a_m are all at least 2, then $m \leq n$.

Proof Let X_M and X_N denote the structural maps $X: M \rightarrow M$ and $X: N \rightarrow N$, respectively. Our aim is to establish $m = \dim_{\mathbb{F}} \ker X_M \leq \dim_{\mathbb{F}} \ker X_N = n$.

Setting $C := g^{-1}(\ker X_M)$, we first claim that $g|_C: C \rightarrow \ker X_M$ is surjective. Since the numbers a_1, \dots, a_m are all at least two, any element y in the kernel of X_M lies in the image of X_M , and therefore may be written as $y = X_M z = g(f(z))$, which proves the claim. Next, note that g sends $X_N C$ to zero, and so $g|_C$ induces a surjection $C/X_N C \rightarrow \ker X_M$. Thus

$$\dim_{\mathbb{F}} \ker X_M \leq \dim_{\mathbb{F}} C - \dim_{\mathbb{F}} X_N C = \dim_{\mathbb{F}} \ker(X_N|_C) \leq \dim_{\mathbb{F}} \ker X_N. \quad \square$$

Proof of Theorem 1 Let $\overline{\text{BN}}(K)$ denote the reduced Bar-Natan homology of K with rational coefficients. It is a rank-1 finitely generated graded module over $\mathbb{Q}[H]$ where H has nonzero degree, so we may write

$$\overline{\text{BN}}(K) \cong \mathbb{Q}[H] \oplus \bigoplus_{i=1}^m \frac{\mathbb{Q}[H]}{H^{a_i}}, \quad \overline{\text{BN}}(J) \cong \mathbb{Q}[H] \oplus \bigoplus_{i=1}^n \frac{\mathbb{Q}[H]}{H^{b_i}},$$

where $a_1, \dots, a_m, b_1, \dots, b_n$ are positive. By hypothesis, there is no 2-torsion in $\text{Kh}(K)$, so Proposition 9.3 of [5] implies that the numbers a_1, \dots, a_m are all at least 2. Furthermore, [5, Proof of Proposition 9.3] also gives $\text{rk } \overline{\text{Kh}}(K) = 1 + 2m$ and $\text{rk } \overline{\text{Kh}}(J) = 1 + 2n$.

By [4, Proof of Theorem 1.1], there are $\mathbb{Q}[H]$ -module maps $f: \overline{\text{BN}}(K) \rightarrow \overline{\text{BN}}(J)$ and $g: \overline{\text{BN}}(J) \rightarrow \overline{\text{BN}}(K)$ satisfying $f \circ g = H$ and $g \circ f = H$. We note that the complex $[[D]]$ over $\mathbb{Z}[G]$ associated to a diagram D considered in [4] recovers the reduced Bar-Natan complex as $[[D]] \otimes_{\mathbb{Z}[G]} \mathbb{Q}[H]$ where $\mathbb{Z}[G] \rightarrow \mathbb{Q}[H]$ sends G to $-H$. By Lemma 3, we obtain

$$\text{rk } \overline{\text{Kh}}(K) = 1 + 2m \leq 1 + 2n = \text{rk } \overline{\text{Kh}}(J). \quad \square$$

Remark 4 Suppose K is a knot such that $\overline{\text{BN}}(K)$ does not contain $\mathbb{Q}[H]/H$ as a direct summand. Our proof of Theorem 1 implies that the conclusion of Theorem 1 holds for K . This observation gives evidence in favor of the affirmative for [5, Question 9.4], which suggests that the reduced Bar-Natan homology

of any nontrivial knot contains $\mathbb{Q}[H]/H$ as a direct summand. We note that the existence of such a summand in reduced Bar-Natan homology implies the existence of 2-torsion in Khovanov homology [5, Proposition 9.3]. An analogous question is raised in [1; 7] in the context of the Floer homology of rational homology spheres.

Acknowledgments We thank Akram Alishahi, John Baldwin, Artem Kotelskiy, Lukas Lewark, Tom Mrowka, and Raphael Zentner for interesting discussions. Gujral thanks his advisor Lisa Piccirillo for useful conversations, support, and advice, and Wang thanks Peter Kronheimer for his continued guidance, support, and encouragement. We also thank the referee for helpful suggestions. Gujral and Wang were partially supported by the Simons Collaboration on New Structures in Low-Dimensional Topology, and Wang was also partially supported by the NSF MSPRF grant DMS-2303401.

References

- [1] **A Alfieri, F Binns**, *Is the geography of Heegaard Floer homology restricted or the L-space conjecture false?*, preprint (2024) arXiv 2404.00490
- [2] **MM Asaeda, JH Przytycki**, *Khovanov homology: torsion and thickness*, from “Advances in topological quantum field theory”, NATO Sci. Ser. II Math. Phys. Chem. 179, Kluwer, Dordrecht (2004) 135–166 MR
- [3] **JA Baldwin, S Sivek**, *Khovanov homology detects the trefoils*, *Duke Math. J.* 171 (2022) 885–956 MR
- [4] **D Ittgen, L Lewark, L Marino**, *Khovanov homology and rational unknotting*, preprint (2021) arXiv 2110.15107
- [5] **A Kotelskiy, L Watson, C Zibrowius**, *Immersed curves in Khovanov homology*, preprint (2019) arXiv 1910.14584
- [6] **PB Kronheimer, TS Mrowka**, *Khovanov homology is an unknot-detector*, *Publ. Math. Inst. Hautes Études Sci.* 113 (2011) 97–208 MR
- [7] **F Lin**, *A remark on taut foliations and Floer homology*, preprint (2023) arXiv 2309.01222
- [8] **D McCoy, R Zentner**, *The Montesinos trick for proper rational tangle replacement*, *Proc. Amer. Math. Soc.* 151 (2023) 1811–1822 MR
- [9] **JH Przytycki, R Sazdanović**, *Torsion in Khovanov homology of semi-adequate links*, *Fund. Math.* 225 (2014) 277–304 MR
- [10] **AN Shumakovitch**, *Torsion of Khovanov homology*, *Fund. Math.* 225 (2014) 343–364 MR

Department of Mathematics, Massachusetts Institute of Technology
Cambridge, MA, United States

Department of Mathematics, Massachusetts Institute of Technology
Cambridge, MA, United States

onkar@mit.edu, joshuaxw@princeton.edu

Received: 24 November 2023 Revised: 1 September 2024

ALGEBRAIC & GEOMETRIC TOPOLOGY

msp.org/agt

EDITORS

PRINCIPAL ACADEMIC EDITORS

John Etnyre
etnyre@math.gatech.edu
Georgia Institute of Technology

Kathryn Hess
kathryn.hess@epfl.ch
École Polytechnique Fédérale de Lausanne

BOARD OF EDITORS

Julie Bergner	University of Virginia jeb2md@eservices.virginia.edu	Thomas Koberda	University of Virginia thomas.koberda@virginia.edu
Steven Boyer	Université du Québec à Montréal cohf@math.rochester.edu	Markus Land	LMU München markus.land@math.lmu.de
Tara E Brendle	University of Glasgow tara.brendle@glasgow.ac.uk	Christine Lescop	Université Joseph Fourier lescop@ujf-grenoble.fr
Indira Chatterji	CNRS & Univ. Côte d'Azur (Nice) indira.chatterji@math.cnrs.fr	Norihiko Minami	Yamato University minami.norihiko@yamato-u.ac.jp
Octav Cornea	Université de Montreal cornea@dms.umontreal.ca	Andrés Navas	Universidad de Santiago de Chile andres.navas@usach.cl
Alexander Dranishnikov	University of Florida dranish@math.ufl.edu	Jessica S Purcell	Monash University jessica.purcell@monash.edu
Tobias Ekholm	Uppsala University, Sweden tobias.ekholm@math.uu.se	Birgit Richter	Universität Hamburg birgit.richter@uni-hamburg.de
Mario Eudave-Muñoz	Univ. Nacional Autónoma de México mario@matem.unam.mx	Jérôme Scherer	École Polytech. Féd. de Lausanne jerome.scherer@epfl.ch
David Futер	Temple University dfuter@temple.edu	Vesna Stojanoska	Univ. of Illinois at Urbana-Champaign vesna@illinois.edu
John Greenlees	University of Warwick john.greenlees@warwick.ac.uk	Zoltán Szabó	Princeton University szabo@math.princeton.edu
Matthew Hedden	Michigan State University mhedden@math.msu.edu	Maggy Tomova	University of Iowa maggy-tomova@uiowa.edu
Kristen Hendricks	Rutgers University kristen.hendricks@rutgers.edu	Daniel T Wise	McGill University, Canada daniel.wise@mcgill.ca
Hans-Werner Henn	Université Louis Pasteur henn@math.u-strasbg.fr	Lior Yanovski	Hebrew University of Jerusalem lior.yanovski@gmail.com
Daniel Isaksen	Wayne State University isaksen@math.wayne.edu		


See inside back cover or msp.org/agt for submission instructions.

The subscription price for 2025 is US \$760/year for the electronic version, and \$1110/year (+\$75, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Algebraic & Geometric Topology is indexed by Mathematical Reviews, Zentralblatt MATH, Current Mathematical Publications and the Science Citation Index.

Algebraic & Geometric Topology (ISSN 1472-2747 printed, 1472-2739 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004.

AGT peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<https://msp.org/>

© 2025 Mathematical Sciences Publishers

ALGEBRAIC & GEOMETRIC TOPOLOGY

Volume 25 Issue 7 (pages 3789–4436) 2025

Algebras for enriched ∞ -operads	3789
RUNE HAUGSENG	
An overtwisted convex hypersurface in higher dimensions	3813
RIVER CHIANG and KLAUS NIEDERKRÜGER-EID	
Presheaves of groupoids as models for homotopy types	3833
LÉONARD GUETTA	
Product and coproduct on fixed point Floer homology of positive Dehn twists	3875
YUAN YAO and ZIWEN ZHAO	
Multitwists in big mapping class groups	3921
GEORGE DOMAT, FEDERICA FANONI and SEBASTIAN HENSEL	
Cofibrantly generated model structures for functor calculus	3931
LAUREN BANDKLAYDER, JULIA E BERGNER, RHIANNON GRIFFITHS, BRENDA JOHNSON and REKHA SANTHANAM	
Endomorphisms of Artin groups of type D	3975
FABRICE CASTEL and LUIS PARIS	
Linear bounds of the crosscap number of knots	4009
ROB MCCONKEY	
Atiyah–Segal completion for the Hermitian K-theory of symplectic groups	4037
JENS HORNBOSTEL, HERMAN ROHRBACH and MARCUS ZIBROWIUS	
A minimality property for knots without Khovanov 2-torsion	4073
ONKAR SINGH GUJRAL and JOSHUA WANG	
Local indicability of groups with homology circle presentations	4077
AGUSTÍN NICOLÁS BARRETO and ELÍAS GABRIEL MINIAN	
Cubulating drilled bundles over graphs	4095
MAHAN MJ and BISWAJIT NAG	
Infinitely many homeomorphic hyperbolic plugs with the same basic sets	4147
FANGFANG CHEN	
One-point compactifications of configuration spaces and the self duality of the little disks operad	4163
CONNOR MALIN	
Braided multitwists	4185
RODRIGO DE POOL	
Positive intermediate Ricci curvature on connected sums	4209
PHILIPP REISER and DAVID J WRAITH	
The \mathcal{S}_n -equivariant Euler characteristic of the moduli space of graphs	4229
MICHAEL BORINSKY and JOS VERMASEREN	
Classification of metric fibrations	4257
YASUHIKO ASAO	
Spaces over BO are thickened manifolds	4287
HIRO LEE TANAKA	
Representation-graded Bredon homology of elementary abelian 2-groups	4321
MARKUS HAUSMANN and STEFAN SCHWEDE	
The \mathbb{Z}/p -equivariant cohomology of the genus-zero Deligne–Mumford space with $1 + p$ marked points	4341
DAIN KIM and NICHOLAS WILKINS	
Geometry of free extensions of free groups via automorphisms with fixed points on the complex of free factors	4357
PRITAM GHOSH and FUNDA GÜLTEPE	
Coarse cohomology of configuration space and coarse embedding	4391
ARKA BANERJEE	
Correction to the article An algebraic model for finite loop spaces	4427
CARLES BROTO, RAN LEVI and BOB OLIVER	