

AG
T

*Algebraic & Geometric
Topology*

Volume 25 (2025)

Local indicability of groups with homology circle presentations

AGUSTÍN NICOLÁS BARRETO

ELÍAS GABRIEL MINIAN

Local indicability of groups with homology circle presentations

AGUSTÍN NICOLÁS BARRETO

ELÍAS GABRIEL MINIAN

We investigate conditions that guarantee local indicability of groups that admit presentations with the homology of a circle, generalizing a result of J Howie for two-relator presentations. We apply our results to investigate local indicability of LOT groups and some classes of non-cycle-free Adian presentations, extending previous results in that direction by J Howie and D Wise.

20F05, 20F65, 57M05, 57M07, 57M10

1 Introduction

Recall that a group G is locally indicable if each of its finitely generated nontrivial subgroups admits a homomorphism onto the infinite cyclic group \mathbb{Z} . This class of groups appeared in the work of Higman on the units of group rings [8] and has been intensively studied in the last four decades since the works of Brodskii [3] and Howie [9; 10]. The theory of locally indicable groups has connections with equations over groups [3; 9; 14], with complexes of nonpositive curvature and coherence [13; 16; 17; 18], and with dynamics and left orderable groups (see, for example, [4; 5; 15]). They are also related to the study of asphericity of 2-complexes and Whitehead’s conjecture, since any connected 2-complex X with $\pi_1(X)$ locally indicable and $H_2(X) = 0$ is aspherical (see [10; 11]). In fact, locally indicable groups are \mathbb{Z} -conservative [6; 12].

It is known that torsion-free one-relator groups are locally indicable [3]. Moreover one-relator products of locally indicable groups are locally indicable if the relator is not a proper power (see [10]). In [11, Theorem 6.2] Howie proved local indicability of groups admitting presentations

$$\mathcal{P} = \langle a, b, c \mid a^{-1}w_1^{-1}bw_1, b^{-1}w_2^{-1}cw_2 \rangle$$

with three generators and two relators, where w_1 is a specific type of word called a sloping word. These presentations arise naturally when studying (weak) labeled oriented trees (LOTs), and the result is used to prove local indicability of LOT groups for LOTs of diameter 3 and some families of LOTs of diameter 4 (see [11]).

One of the main results of this paper is a generalization of Howie’s result [11, Theorem 6.2] to homology circle presentations of any length where the relators satisfy some extra hypotheses (weaker than Howie’s

original result for two-relator presentations). It is clear that some extra condition is needed to derive local indicability, since one can construct homology circle presentations of nonlocally indicable groups just by adding a new generator to any balanced presentation of a perfect group. The condition that we require depends on the minima or maxima of the total exponents of the initial subwords of all but one of the relators (see Definitions 2.2 and 2.4).

Theorem 2.5 *Let $\mathcal{P} = \langle a_1, \dots, a_{k+2} \mid r_1, \dots, r_k, s \rangle$ be a presentation (of deficiency 1) of a group G with $H_1(G) = \mathbb{Z}$ (for some $k \geq 0$), where all the relators are cyclically reduced and have total exponent 0. If the multisets of minima $m(r_1), \dots, m(r_k)$ are concatenable, then G is locally indicable.*

The proof essentially follows Howie's original proof of [11, Theorem 6.2], adapted to a more general context. One can recover Howie's result as a particular case.

The condition on the relators having total exponent 0 can be dropped by considering any surjective homomorphism $\varphi: G \rightarrow \mathbb{Z}$ and the sequences of minima or maxima with respect to φ .

Theorem 2.9 *Let $\mathcal{P} = \langle a_1, \dots, a_{k+2} \mid r_1, \dots, r_k, s \rangle$ be a presentation (of deficiency 1) of a group G with $H_1(G) = \mathbb{Z}$ (for some $k \geq 0$), where all the relators are cyclically reduced. If there is a surjective homomorphism $\varphi: G \rightarrow \mathbb{Z}$ such that $m_\varphi(r_1), \dots, m_\varphi(r_k)$ are concatenable, then G is locally indicable.*

For two-relator presentations of deficiency 1 one can widely extend the result by using relative maxima or minima (see Definition 2.11).

Corollary 2.12 *Let $\mathcal{P} = \langle a, b, c \mid r, s \rangle$ be a presentation of a group G with $H_1(G) = \mathbb{Z}$ where the relators are cyclically reduced. Let $\varphi: G \rightarrow \mathbb{Z}$ be a surjective homomorphism with $\varphi(c) \neq 0$. If r attains a unique relative minimum at a , then G is locally indicable.*

In Theorem 2.18 we extend our methods to investigate presentations with the homology of a wedge of circles.

Our main applications are related to LOT groups and Adian presentations. Labeled oriented trees give rise to homology circle presentations that generalize Wirtinger presentations for knots. It is well known that knot groups are locally indicable (see [10]). Local indicability of LOT groups is an open problem (that would imply asphericity of the associated presentations). In fact, it is not even known whether all LOT groups are torsion-free. In [11] Howie associated to any LOT Γ two graphs. The right graph $I(\Gamma)$ is obtained from Γ by interchanging the initial vertex and the label of every edge in Γ . Analogously, the left graph $T(\Gamma)$ is constructed by interchanging the final vertex with the label. A similar construction was investigated by Gersten in [7] in the more general context of Adian presentations $\mathcal{P} = \langle A \mid u_i = v_i \ (i \in J) \rangle$ (where u_i and v_i are nontrivial positive words). Howie proved that, if $I(\Gamma)$ or $T(\Gamma)$ has no cycles, then the group $G(\Gamma)$ is locally indicable (and, in particular, Γ is aspherical) [11, Theorem 10.1]. For

Adian presentations, Gersten showed that if both $T(\mathcal{P})$ and $I(\mathcal{P})$ have no cycles then the associated 2-complex $\mathcal{K}_{\mathcal{P}}$ is diagrammatically reducible and, in particular, aspherical [7, Proposition 4.12]. Moreover, when $\ell(u_i) = \ell(v_i)$ for every i (as in the case of LOT presentations), if either $T(\mathcal{P})$ or $I(\mathcal{P})$ has no cycles then $\mathcal{K}_{\mathcal{P}}$ is diagrammatically reducible (and hence aspherical) [7, Proposition 4.15]. Here $\ell(w)$ denotes the length of the word w . More recently, Wise proved that the 2-complexes associated to cycle-free Adian presentations (ie those for which both graphs are forests) have nonpositive sectional curvature, which implies that the corresponding groups are locally indicable [17, Theorem 11.4]. For LOT presentations we obtain the following extension of Howie's result.

Corollary 3.1 *Let Γ be a LOT. If either $I(\Gamma)$ or $T(\Gamma)$ has at most one cycle, then $G(\Gamma)$ is locally indicable.*

In [Corollary 3.3](#) we extend this result to a wider class of Adian presentations, generalizing, for a certain subfamily, Wise's result [17, Theorem 11.4].

Finally, [Theorem 3.6](#) describes a strategy to study local indicability of LOT groups when both graphs $I(\Gamma)$ and $T(\Gamma)$ have more than one cycle. This result is a consequence of [Theorem 2.18](#).

Acknowledgements

We would like to thank Jim Howie and Jonathan Barmak for useful suggestions and comments during the preparation of this article.

The authors are researchers of CONICET. This research was partially supported by grants PICT 2019-2338 and UBACYT 20020190100099BA.

2 Main results

The results of this section are based on some ideas and techniques developed by Jim Howie in the proof of [11, Theorem 6.2] and use well-known properties of locally indicable groups. Given a presentation \mathcal{P} of a group $G = G(\mathcal{P})$, we denote by $\mathcal{K}_{\mathcal{P}}$ its standard 2-complex which has a single 0-cell, one 1-cell for each generator and one 2-cell for each relator. Recall from [10] that an elementary reduction (X, Y) is a pair of 2-complexes such that $Y \subset X$ and $X - Y$ consists of exactly one 1-cell e^1 and at most one 2-cell e^2 , where e^2 properly involves e^1 , which means that its attaching map is not homotopic in $Y \cup e^1$ to a map into Y . A 2-complex K is reducible if for every finite subcomplex X there exists an elementary reduction (X, Y) . Also, a group presentation \mathcal{P} is reducible if its associated 2-complex $\mathcal{K}_{\mathcal{P}}$ is. We will use the following known fact.

Theorem 2.1 (Howie [10]) *If G has a reducible presentation in which no relator is a proper power, then G is locally indicable.*

Let F be a free group on a set A . Recall that the total exponent of a word $S = a_1^{\varepsilon_1} \dots a_n^{\varepsilon_n}$ (with $\varepsilon_i \in \{\pm 1\}$) is $\exp(S) = \varepsilon_1 + \dots + \varepsilon_n$. The length of a word S will be denoted by $\ell(S)$. An initial subword of a word $S = a_1^{\varepsilon_1} \dots a_n^{\varepsilon_n}$ is a word of the form $a_1^{\varepsilon_1} \dots a_i^{\varepsilon_i}$ for some $1 \leq i \leq n$. Suppose \mathcal{P} is a finite presentation of a group G such that the total exponents of all its relators are zero. Let $\varphi: G \rightarrow \mathbb{Z}$ be the surjective homomorphism which sends every generator of \mathcal{P} to $1 \in \mathbb{Z}$. Let $\tilde{\mathcal{K}}$ be the infinite cyclic cover of $\mathcal{K}_{\mathcal{P}}$ determined by φ . Similarly as in [10], we index its 0-cells with the integers. Via $\varphi: G \rightarrow \mathbb{Z}$ every generator a in \mathcal{P} determines one 1-cell a_j oriented from j to $j + 1$ for each $j \in \mathbb{Z}$. Given a relator r , we denote by R the corresponding 2-cell in $\mathcal{K}_{\mathcal{P}}$, and by R_j , the 2-cell in $\tilde{\mathcal{K}}$ that covers R and whose minimum traversed 0-cell is j .

For any relator r , the sequence of total exponents of its initial subwords represents the indices of the 0-cells traversed by the attaching path of the corresponding 2-cell in $\tilde{\mathcal{K}}$ starting at vertex 0. Suppose that the minimum value attained in the sequence is $m \in \mathbb{Z}$. Note that this minimum can be attained more than once and that the rightmost letter of an initial subword where m is attained must have exponent -1 . This letter represents a 1-cell joining the vertices m and $m + 1$ (and it is traversed backwards). The letters with exponent 1 where the value $m + 1$ is attained represent 1-cells oriented from m to $m + 1$ and they are traversed forwards. Note that if we invert the relator these roles are interchanged. This motivates the following definition.

Definition 2.2 Let r be a relator of a presentation \mathcal{P} and let m be the minimum value of the sequence of total exponents of its initial subwords. We define the *multiset of minima* $m(r)$ as the multiset that contains the letters of r with exponent -1 that attain the value m and the letters that appear with exponent 1 in r that attain the value $m + 1$.

For example, if $r = a^{-1}b^{-1}cb^{-1}ab$, the total exponents are $-1, -2, -1, -2, -1, 0$ and $m(r) = \{b, c, b, a\}$.

Remark 2.3 We can formulate an analogous definition for maxima instead of minima. In fact, every result in this paper that holds for minima has also an analogous version for maxima.

Definition 2.4 Given multisets A_1, \dots, A_n we say that they are *concatenable* if there is an ordering A_{i_1}, \dots, A_{i_n} such that, for every $1 \leq j \leq n$, there exists an element in A_{i_j} with multiplicity 1 that does not belong to the union of $A_{i_1}, \dots, A_{i_{j-1}}$.

The following result is our first generalization of Howie's [11, Theorem 6.2].

Theorem 2.5 Let $\mathcal{P} = \langle a_1, \dots, a_{k+2} \mid r_1, \dots, r_k, s \rangle$ be a presentation (of deficiency 1) of a group G with $H_1(G) = \mathbb{Z}$ (for some $k \geq 0$), where all the relators are cyclically reduced and have total exponent 0. If the multisets of minima $m(r_1), \dots, m(r_k)$ are concatenable, then G is locally indicable.

We recover Howie's result [11, Theorem 6.2] as a particular case.

Corollary 2.6 (Howie [11]) Let Γ be a weakly labeled oriented tree with 3 vertices, in which at least one label is a reduced sloping word. Then $G(\Gamma)$ is locally indicable.

The corollary follows from the fact that the standard presentation of the group of a weakly labeled oriented tree with three vertices $\mathcal{P} = \langle a, b, c \mid r, s \rangle$ is a homology circle and the sloping word condition assures that $m(r)$ attains a unique minimum or a unique maximum in its sequence. In particular, it is concatenable. In fact, for a two-relator presentation $\mathcal{P} = \langle a, b, c \mid r, s \rangle$ it is not required that $m(r)$ has a unique minimum in order to be concatenable. We just need that there exists a generator that appears only once in $m(r)$. In [Corollary 2.12](#) we will show a generalization of this result for two-relator presentations.

The proof of [Theorem 2.5](#) uses the following technical lemma.

Lemma 2.7 *Let F be a free group on a set A and T a cyclically reduced word in F with $\exp(T) = 0$. Then T cannot be factored as a product*

$$T = w_0 S_0 w_0^{-1} w_r S_r w_r^{-1} w_{r-1} S_{r-1} w_{r-1}^{-1} \dots w_1 S_1 w_1^{-1}$$

with $w_0, w_1, \dots, w_r, S_0, S_1, \dots, S_r \in F$ ($r \geq 0$), satisfying

- $\exp(T) = \exp(S_i) = 0$ for $0 \leq i \leq r$,
- $\exp(w_i) \geq 1$ for $1 \leq i \leq r$ and $\exp(w_0) = 1$,
- the total exponents of all the initial subwords of T, S_i and w_i are nonnegative ($0 \leq i \leq r$).

Proof Suppose there exists such a factorization for some cyclically reduced word T . Over all possible factorizations of T and T^{-1} satisfying these conditions, we consider the cyclically reduced words $w_0, w_1, \dots, w_r, S_0, S_1, \dots, S_r$ that minimize the sum

$$L = \sum_{i=0}^r (\ell(w_i) + \ell(S_i)).$$

We can assume that the factorization that minimizes L is one of T . Otherwise, we change T by T^{-1} . We rewrite the equality as

$$w_0^{-1} T w_1 S_1^{-1} w_1^{-1} w_2 S_2^{-1} w_2^{-1} \dots w_r S_r^{-1} w_r^{-1} w_0 = S_0.$$

Note that S_i^{-1} satisfies the same properties as S_i for every $0 \leq i \leq r$.

Let c be the rightmost letter in T . Note that it must appear as c^{-1} (with negative exponent) since the initial subwords of T have nonnegative total exponent and $\exp(T) = 0$. The key observation is that this appearance of c^{-1} in T must cancel with a c , since, in any other case, $w_0^{-1} T$ would be an initial subword of S_0 with total exponent -1 , which contradicts the hypotheses. There are five types of possible cancellations for c^{-1} .

Case 1 (the rightmost letter c^{-1} in T cancels with a c in some S_i^{-1} with $i \geq 1$) In this case we can write $S_i^{-1} = U c V$ for some (possibly empty) words U and V . The word located between the letter c^{-1} in T and the c in S_i^{-1} must be trivial, so $w_1 S_1^{-1} w_1^{-1} \dots w_{i-1} S_{i-1}^{-1} w_{i-1}^{-1} w_i U$ is trivial, and in particular this word has total exponent 0. Since the total exponent of $w_1 S_1^{-1} w_1^{-1} \dots w_{i-1} S_{i-1}^{-1} w_{i-1}^{-1}$ is 0 and $\exp(w_i) \geq 1$, we have that $\exp(U) \leq -1$, but this is a contradiction since the subwords of S_i^{-1} have nonnegative total exponent.

Case 2 (the rightmost letter c^{-1} in T cancels with a c in some $w_i = UcV$ with $i \geq 1$) In this case, the hypothesis implies that $w_1 S_1^{-1} w_1^{-1} \dots w_{i-1} S_{i-1}^{-1} w_{i-1}^{-1} U$ is trivial, so we can erase this subword from the original equality, replace $w_i^{-1} = V^{-1} c^{-1} U^{-1}$, and obtain

$$w_0^{-1} T c V S_i^{-1} V^{-1} c^{-1} U^{-1} w_{i+1} \dots w_r S_r^{-1} w_r^{-1} w_0 = S_0.$$

Note that $\exp(cV) = 1$, since $\exp(w_i) = \exp(UcV) \geq 1$ and $\exp(U) = 0$. Also, the total exponents of the initial subwords of cV are nonnegative because this property holds for w_i and $\exp(U) = 0$. If U is empty, we obtain a strictly shorter writing of T where cV plays the role of w_i , which is a contradiction. If U is not empty, we can replace in the last equality $U^{-1} = w_1 S_1^{-1} w_1^{-1} \dots w_{i-1} S_{i-1}^{-1} w_{i-1}^{-1}$ obtaining a new factorization for T where cV plays the role of w_i . Since cV has a strictly shorter length than w_i , this would be a contradiction.

Case 3 (the rightmost letter c^{-1} in T cancels with a c in some $w_i^{-1} = V^{-1} c U^{-1}$ with $i \geq 1$) This is similar to case 2.

Case 4 (the rightmost letter c^{-1} in T cancels with a c in $w_0^{-1} = V^{-1} c U^{-1}$) This condition implies $cU^{-1} = T^{-1}$. After replacing, we get

$$V^{-1} w_1 S_1^{-1} w_1^{-1} \dots w_r S_r^{-1} w_r^{-1} T V = S_0.$$

Since $\exp(w_0) = \exp(TV) = 1$ and $\exp(T) = 0$ we get that $\exp(V) = 1$. Also, the total exponents of initial subwords of V are nonnegative because w_0 satisfies that condition and $\exp(T) = 0$. Taking inverses at both sides of the equality we get

$$V^{-1} T^{-1} w_r S_r w_r^{-1} \dots w_1 S_1 w_1^{-1} V = S_0^{-1},$$

and since T and the S_i have total exponent 0, this provides a factorization of T^{-1} satisfying the conditions, with a strictly smaller value of L , which is again a contradiction.

Case 5 (the rightmost letter c^{-1} in T cancels with a c in $w_0 = UcV$) This is similar to case 4. □

Proof of Theorem 2.5 We follow the proof of [11, Theorem 6.2]. If $k = 0$, G is a torsion-free one-relator group and the result follows from [3] (see also [10, Corollary 4.3]). Now suppose that $k \geq 1$. Let R_1, \dots, R_k, S be the 2-cells of $\mathcal{K}_{\mathcal{P}}$ corresponding to r_1, \dots, r_k and s , respectively. Let $\varphi: G \rightarrow \mathbb{Z}$ be the (surjective) homomorphism that sends every generator to 1 and let $\tilde{\mathcal{K}}$ be the corresponding infinite cyclic cover of $\mathcal{K}_{\mathcal{P}}$. It suffices to prove that $\pi_1(\tilde{\mathcal{K}})$ is locally indicable.

We index the 0-cells of $\tilde{\mathcal{K}}$ with the integers and denote $a_{i,j}$ the 1-cell that covers a_i and joins the 0-cells j and $j + 1$. It is oriented from j to $j + 1$. We denote by $R_{i,j}$ and S_j the 2-cells that cover R_i and S , respectively, for which the minimum 0-cell is j . Since $\pi_1(\tilde{\mathcal{K}})$ is the direct limit of the fundamental groups of its finite connected subcomplexes, it suffices to prove that for every finite connected subcomplex $K \subset \tilde{\mathcal{K}}$, there is connected subcomplex $L \subset \tilde{\mathcal{K}}$ with $K \subset L$ and $\pi_1(L)$ locally indicable. The strategy is to include K in a finite subcomplex L that is homotopy equivalent to a 2-complex L' , which in turn collapses to a reducible 2-complex and then use Theorem 2.1.

We define first the iterated rewrites of the cells S_j . Since $m(r_1), \dots, m(r_k)$ are concatenable, we can assume without loss of generality that $a_i \in m(r_i)$ with multiplicity 1 and $a_i \notin \bigcup_{j=1}^{i-1} m(r_j)$ ($1 \leq i \leq k$). Equivalently, for every $1 \leq i \leq k$, $a_{i,0}$ appears exactly once in the attaching path of $R_{i,0}$ and does not appear in the attaching path of $R_{j,0}$ for any $j < i$. Also $a_{i,0}$ passes through the 0-cell 0 once. We can then write the attaching path of $R_{i,0}$ as $a_{i,0} p_{i,0}^{-1}$ where $p_{i,0}$ is a path which does not involve $a_{i,0}$. Let $S_0^{(1)}$ be the path obtained after replacing, for each $1 \leq i \leq k$ in decreasing order, every occurrence of $a_{i,0}$ in the attaching path of S_0 by $p_{i,0}$ and cyclically reducing. Note that $S_0^{(1)}$ does not pass through $a_{i,0}$ for any $1 \leq i \leq k$. Analogously, $a_{i,1}$ appears just once in the attaching path of $R_{i,1}$ and so it can be written as $a_{i,1} p_{i,1}^{-1}$ where $p_{i,1}$ is a path that does not involve $a_{i,1}$. Let $S_0^{(2)}$ be the path obtained after replacing in $S_0^{(1)}$ every occurrence of $a_{i,1}$ by $p_{i,1}$, for each $1 \leq i \leq k$ in decreasing order, and cyclically reducing. Denote $S_1^{(1)}$ the path obtained by doing the same in S_1 . By repeating this procedure, we define the iterated rewrite $S_l^{(j)}$ for each $l \geq 0$ and $j \geq 1$.

Let α_1 be the minimum 0-cell traversed by $S_0^{(1)}$. Note that $\alpha_1 \geq 0$. In general, for every j , we denote by α_j the minimum 0-cell of $S_0^{(j)}$. Note that $\alpha_j \geq \alpha_{j-1}$ and that the minimum 0-cell of $S_l^{(j)}$ is $\alpha_j + l$. We will show first that for $j \in \mathbb{N}$ big enough, the sequence $\{\alpha_j\}_{j \geq 1}$ stabilizes. Since the sequence is increasing, we only need to prove that it is bounded above. We show that $\alpha_n < \rho + \sigma + 1$ for every n , where $\rho = \sum_{i=1}^k \rho_i$ and $\rho_1, \dots, \rho_k, \sigma$ are the maximum 0-cells traversed by the attaching paths of $R_{1,0}, \dots, R_{k,0}, S_0$, respectively.

Suppose that there exists $j \in \mathbb{N}$ such that $\alpha_j \geq \rho + \sigma + 1$. Take $n \in \mathbb{N}$ big enough such that $\alpha_{n-\rho-\sigma} \geq \rho + \sigma + 1$. Let Y be the full subcomplex of $\widetilde{\mathcal{K}}$ containing all 0-cells j for $0 \leq j \leq n + \rho + \sigma$. This complex contains the 1-cells $a_{i,j}$ for each $1 \leq i \leq k + 2$ and $0 \leq j \leq n + \rho + \sigma - 1$, the 2-cells $R_{i,j}$ for $1 \leq i \leq k$ and $0 \leq j \leq n + \sigma + \rho - \rho_i$ and the 2-cells S_j with $0 \leq j \leq n + \rho$. Note that the number of 0-cells in Y is $n + \rho + \sigma + 1$, the number of 1-cells is $(k + 2)(n + \rho + \sigma)$, for each $1 \leq i \leq k$ it has $n + \sigma + \rho - \rho_i + 1$ 2-cells corresponding to the $R_{i,j}$ with $0 \leq j \leq n + \sigma + \rho - \rho_i$, and $n + \rho + 1$ 2-cells corresponding to the S_j with $0 \leq j \leq n + \rho$. Therefore its Euler characteristic is $\chi(Y) = -\rho - \sigma + k + 2$.

We construct a 2-complex Y' homotopy equivalent to Y by replacing each S_j by a 2-cell attached via the path $S_j^{(n-j)}$ ($0 \leq j \leq \rho + \sigma$). Note that this 2-complex collapses to a subcomplex Y'' by removing $R_{i,j}$ and $a_{i,j}$ for every $1 \leq i \leq k$ and $0 \leq j \leq \rho + \sigma$. Let Z be the 1-subcomplex of Y'' that contains the 0-cells from 0 to $\rho + \sigma + 1$ and the 1-cells $a_{i,j}$ with $i = k + 1, k + 2$ and $0 \leq j \leq \rho + \sigma$. Since, for every j , the minimum 0-cell of $S_j^{(n-j)}$ is $j + \alpha_{n-j} \geq j + \alpha_{n-\rho-\sigma} \geq j + \rho + \sigma + 1$, the identity $Z \mapsto Z$ can be continuously extended to a retraction $Y'' \mapsto Z$ which sends all of $Y'' - Z$ to the 0-cell $\rho + \sigma + 1$. It follows that $\beta_1(Y) = \beta_1(Y'') \geq \beta_1(Z) = \rho + \sigma + 1$. Here β_1 denotes the first Betti number. Note that $\beta_0(Y) = 1$ since Y is connected. Note also that $\mathcal{K}_{\mathcal{P}}$ is a homology circle because $H_1(\mathcal{K}_{\mathcal{P}}) = H_1(G) = \mathbb{Z}$ and the deficiency of the presentation is 1. This implies that $\beta_2(Y) = 0$ since $\widetilde{\mathcal{K}}$ is an infinite cyclic cover (see [1, Proposition 1]). It follows that $\chi(Y) \leq -\rho - \sigma$, which is a contradiction. Therefore, the sequence $\{\alpha_j\}_{j \geq 1}$ is bounded above by $\rho + \sigma + 1$.

Now let $K \subset \widetilde{\mathcal{K}}$ be a finite connected subcomplex. We can assume that the minimum 0-cell in K is 0. Let q be the maximum 0-cell in K and take n large enough such that $\alpha_i = \alpha$ for every $i \geq n$. Take L to be the subcomplex of $\widetilde{\mathcal{K}}$ containing every 0-cell between 0 and $q + n + \rho - \sigma$, every 1-cell $a_{i,j}$ with $1 \leq i \leq k + 2$ and $0 \leq j \leq q + n + \rho - \sigma - 1$, the 2-cells $R_{i,j}$ with $1 \leq i \leq k$ and $0 \leq j \leq q + n + \rho - \rho_i - \sigma$ and the 2-cells S_j with $0 \leq j \leq q - \sigma$. Note that $K \subset L$.

In order to prove that $\pi_1(L)$ is locally indicable, we replace each S_j by a 2-cell attached via $S_j^{(q+n-\sigma-j)}$ and obtain a new 2-complex L' homotopy equivalent to L which collapses to a 2-subcomplex L'' by removing every 2-cell $R_{i,j}$ and every 1-cell $a_{i,j}$ for each $1 \leq i \leq k$ and $0 \leq j \leq q + n + \rho - \rho_i - \sigma$. We will show that the fundamental group of the 2-complex L'' admits a reducible presentation. For $0 \leq j \leq q - \sigma$, the minimum 0-cell of $S_j^{(q+n-\sigma-j)}$ is $j + \alpha$. The 2-cell $S_j^{(q+n-\sigma-j)}$ involves any of the 1-cells in its attaching path incident to the 0-cell $j + \alpha$, and these 1-cells are not in the attaching paths of the remaining 2-cells $S_l^{(q+n-\sigma-l)}$ (for $l > j$). We take the quotient of L'' by a spanning tree in its 1-skeleton. The quotient can be seen as the 2-complex associated to a presentation \mathcal{P} of $\pi_1(L'')$. In order to prove that this presentation is reducible it suffices to show that any subpresentation $\mathcal{P}' \subseteq \mathcal{P}$ with at least one relator has an elementary reduction. We take a subpresentation \mathcal{P}' and consider the relator s_l associated to the 2-cell $S_l^{(q+n-\sigma-l)}$ with minimum l . At least one of the 1-cells of $S_l^{(q+n-\sigma-l)}$ incident to the 0-cell $l + \alpha$ is not in the spanning tree and therefore it represents a generator g of the presentation. Lemma 2.7 applied to $T = s_l$ (where the words S_1, \dots, S_r in Lemma 2.7 play the role of the remaining relators of \mathcal{P}') shows that s_l is not a conjugate in $H * \langle g \rangle$ to an element in the group H whose presentation is obtained from \mathcal{P}' by removing the generator g and the relator s_l . This implies that s_l properly involves g (see [10, page 449]). It follows that $\pi_1(L) = \pi_1(L'')$ is locally indicable by Theorem 2.1. Note that the no-proper-power condition follows from [1, Proposition 1] and the fact that $\widetilde{\mathcal{K}}$ is an infinite cyclic cover of a homology circle 2-complex (see also [11, page 290]). \square

Example 2.8 Consider

$$\mathcal{P} = \langle a, b, c, d \mid a^{-1}c^{-1}b^{-1}a^{-1}badcb^{-1}d, c^{-1}a^{-1}b^{-1}d^{-1}badcb^{-1}d, a^{-1}b^{-1}c^{-1}acdb^{-1}c^{-1}ad \rangle.$$

Name the relators r_1, r_2 and r_3 , respectively. Note that $m(r_1) = \{a, b\}$ and $m(r_2) = \{d, b\}$. Theorem 2.5 (with $a_1 = a$ and $a_2 = d$) proves that $G(\mathcal{P})$ is locally indicable.

Theorem 2.5 can be generalized to presentations \mathcal{P} where the total exponents of the relators are not necessarily 0. Let \mathcal{P} be a presentation and $\varphi: G(\mathcal{P}) \rightarrow \mathbb{Z}$ a surjective homomorphism. By changing, if necessary, a generator by its inverse, we can assume that $\varphi(a)$ is nonnegative for every generator a of \mathcal{P} . Given a word w on the generators, the weight of w (with respect to φ) is $\varphi(w)$ (where w is viewed as an element of $G(\mathcal{P})$). Similarly as before, we can consider the sequence of weights (with respect to φ) of initial subwords of a relator r and define the multiset $m_\varphi(r)$ of minima of r . If the minimum of the sequence is m , $m_\varphi(r)$ contains a copy of the letter a for every initial subword wa^{-1} with weight m and every initial subword wa with weight $m + \varphi(a)$. In particular, if φ sends every generator to 1,

$m_\varphi(r) = m(r)$. For example, if $\mathcal{P} = \langle a, b, c \mid abc^{-1}b^2, ab^{-3}a \rangle$ we can define $\varphi: G(\mathcal{P}) \rightarrow \mathbb{Z}$ by $\varphi(a) = 3$, $\varphi(b) = 2$ and $\varphi(c) = 9$. In this case, the sequence for the relator $r = abc^{-1}b^2$ is 3, 5, -4, -2, 0.

Theorem 2.9 *Let $\mathcal{P} = \langle a_1, \dots, a_{k+2} \mid r_1, \dots, r_k, s \rangle$ be a presentation (of deficiency 1) of a group G with $H_1(G) = \mathbb{Z}$ (for some $k \geq 0$), where all the relators are cyclically reduced. If there is a surjective homomorphism $\varphi: G \rightarrow \mathbb{Z}$ such that $m_\varphi(r_1), \dots, m_\varphi(r_k)$ are concatenable, then G is locally indicable.*

Proof We adapt the proof of Theorem 2.5 to the more general setting. Let $\tilde{\mathcal{K}}$ be the infinite cyclic cover of $\mathcal{K}_\mathcal{P}$ determined by $\varphi: G \rightarrow \mathbb{Z}$. The 1-cells $a_{i,j}$ are oriented from j to $j + \varphi(a_i)$ (recall that we assume that the weights $\varphi(a_i)$ are nonnegative). As before, ρ_i and σ are the maximum 0-cells of $R_{i,0}$ and S_0 , respectively, where $R_{i,j}$ and S_j are the 2-cells of $\tilde{\mathcal{K}}$ covering R_i and S with minimum 0-cell j .

We will assume again that $a_i \in m_\varphi(r_i)$ with multiplicity 1 and $a_i \notin \bigcup_{j=1}^{i-1} m_\varphi(r_j)$. This allows us to write the attaching path of $R_{i,j}$ as $a_{i,j} p_{i,j}^{-1}$ where $p_{i,j}$ is a path that does not involve $a_{i,j}$. Let $S_j^{(t)}$ be the iterated rewrite defined as in the proof of Theorem 2.5 and again let α_j be the minimum 0-cell of $S_0^{(j)}$.

In order to prove that the sequence $\{\alpha_j\}_{j \geq 1}$ stabilizes one has to consider a slightly different complex Y . We show that $\alpha_j < \rho + \sigma + \varphi(a_s)$ where $\varphi(a_s) = \max\{\varphi(a_i)\}$. Suppose $\alpha_j \geq \rho + \sigma + \varphi(a_s)$ for some j . Take $n \in \mathbb{N}$ big enough such that $\alpha_{n-\rho-\sigma} \geq \rho + \sigma + \varphi(a_s)$. Let \tilde{Y} be the full subcomplex of $\tilde{\mathcal{K}}$ containing the 0-cells between 0 and $n + \rho + \sigma$. It contains the 1-cells $a_{i,j}$ for $1 \leq i \leq k + 2, 0 \leq j \leq n + \rho + \sigma - \varphi(a_i)$, the 2-cells $R_{i,j}$ for $1 \leq i \leq k$ and $0 \leq j \leq n + \rho + \sigma - \rho_i$ and the 2-cells S_j for $0 \leq j \leq n + \rho$. The complex Y is obtained from \tilde{Y} by attaching an extra 1-cell between j and $j + 1$ for every $0 \leq j \leq \rho + \sigma + \varphi(a_s) - 1$.

Now we construct a homotopy equivalent 2-complex Y' by replacing each S_j by a 2-cell attached via the path $S_j^{(n-j)}$ for each $0 \leq j \leq \rho + \sigma$. Note that Y' collapses to a subcomplex Y'' after removing $a_{i,j}$ and $R_{i,j}$ for every $1 \leq i \leq k$ and $0 \leq j \leq \rho + \sigma + \varphi(a_s) - \varphi(a_i)$. The attaching of the extra 1-cells guarantees that the full subcomplex Z of Y'' that contains the 0-cells between 0 and $\rho + \sigma + \varphi(a_s)$ is connected, and so is Y . Note that Z is a 1-subcomplex since $\alpha_{n-\rho-\sigma} \geq \rho + \sigma + \varphi(a_s)$. By a similar argument to that of Theorem 2.5, $\beta_1(Y) \geq \beta_1(Z) = 2\rho + 2\sigma + 2\varphi(a_s) - \varphi(a_{k+1}) - \varphi(a_{k+2}) + 2$. As before, this implies that $\chi(Y) \leq -1 - 2\rho - 2\sigma - 2\varphi(a_s) + \varphi(a_{k+1}) + \varphi(a_{k+2})$. On the other side, a direct computation shows that

$$\chi(Y) = -2\rho - 2\sigma - \varphi(a_s) + \sum_{i=1}^{k+2} \varphi(a_i),$$

which implies $\sum_{i=1}^k \varphi(a_i) + \varphi(a_s) \leq -1$, a contradiction.

Now the proof proceeds similarly as in Theorem 2.5. □

Example 2.10 Let $\mathcal{P} = \langle a, b, c \mid c^{-1}b^{-1}c^{-1}abca, b^{-1}c^{-1}aca^{-1}bab^{-1} \rangle$. Using the homomorphism $\varphi: G(\mathcal{P}) \rightarrow \mathbb{Z}$ defined by $\varphi(a) = \varphi(b) = 1$ and $\varphi(c) = 2$, the multiset of minima of the first relator is $\{c, a\}$. By Theorem 2.9, $G_\mathcal{P}$ is locally indicable.

A generalization for two-relator presentations

For presentations $\mathcal{P} = \langle a, b, c \mid r, s \rangle$, [Theorem 2.9](#) requires the existence of a surjective homomorphism $\varphi: G \rightarrow \mathbb{Z}$ such that $m_\varphi(r)$ (or $m_\varphi(s)$) is concatenable, which essentially means that there exists a generator that attains the minimum of the whole word (and only once). We show now that, by applying some Andrews–Curtis moves (or extended Nielsen transformations) to \mathcal{P} , this hypothesis can be relaxed: we only require the existence of a generator with a unique relative minimum.

Definition 2.11 Let \mathcal{P} be a presentation and $\varphi: G(\mathcal{P}) \rightarrow \mathbb{Z}$ a surjective homomorphism. By changing, if necessary, a generator by its inverse, we assume that $\varphi(a)$ is nonnegative for every generator a of \mathcal{P} . We say that a relator r attains a *unique relative minimum at a* if it has an initial subword of the form $w = \tilde{w}a^{-1}$ or $w = \tilde{w}a$, with $\varphi(w) = m_a$, that satisfies the following condition. If $w = \tilde{w}a^{-1}$, then $\varphi(\widehat{w}a^{-1}) > m_a$ and $\varphi(\widehat{w}a) > m_a + \varphi(a)$ for every other initial subwords $\widehat{w}a^{-1}$ and $\widehat{w}a$. If $w = \tilde{w}a$, then $\varphi(\widehat{w}a) > m_a$ and $\varphi(\widehat{w}a^{-1}) > m_a - \varphi(a)$ for every other initial subwords $\widehat{w}a^{-1}$ and $\widehat{w}a$.

For example, if $r = a^{-1}bac^{-1}bbc^{-1}b$ and $\varphi(a) = 1, \varphi(b) = 2, \varphi(c) = 4$, the sequence of weights is $-1, 1, 2, -2, 0, 2, -2, 0$. Note that $m_\varphi(r) = \{c, c, b, b\}$ is not concatenable but there is a unique relative minimum at a (with value -1).

Corollary 2.12 Let $\mathcal{P} = \langle a, b, c \mid r, s \rangle$ be a presentation of a group G with $H_1(G) = \mathbb{Z}$ where the relators are cyclically reduced. Let $\varphi: G \rightarrow \mathbb{Z}$ be a surjective homomorphism with $\varphi(c) \neq 0$. If r attains a unique relative minimum at a , then G is locally indicable.

Proof Let $m \in \mathbb{N}$ be arbitrary. Suppose that r attains the relative minimum in an initial subword of the form $\tilde{w}a^{-1}$. We apply the following Nielsen transformations to \mathcal{P} : we add a new generator d together with a new relator $ac^{-m}d^{-1}c^m$, then we replace all the occurrences of the generator a in r and s by $c^{-m}dc^m$ (and cyclically reduce, if necessary) and obtain new relators \tilde{r} and \tilde{s} . Finally we remove the relator $ac^{-m}d^{-1}c^m$ together with the generator a . We end up with an equivalent presentation $\mathcal{P}' = \langle b, c, d \mid \tilde{r}, \tilde{s} \rangle$. Since the presentations are equivalent, we have $H_*(\mathcal{P}') = H_*(\mathcal{P})$ and $G(\mathcal{P}') = G(\mathcal{P})$. Note that $\varphi(d) = \varphi(a)$. We will show that, for $m \in \mathbb{N}$ big enough, \mathcal{P}' satisfies the hypotheses of [Theorem 2.9](#).

For every subword wa^{-1} with weight k in r there are $2m + 1$ subwords

$$wc^{-1}, \dots, wc^{-m}, wc^{-m}d^{-1}, wc^{-m}d^{-1}c, \dots, wc^{-m}d^{-1}c^m$$

in \tilde{r} with weights $k + \varphi(a) - \varphi(c), \dots, k + \varphi(a) - m\varphi(c), k - m\varphi(c), k - (m - 1)\varphi(c), \dots, k$. Similarly, for every subword wa with weight k in r , we have $2m + 1$ subwords

$$wc^{-1}, \dots, wc^{-m}, wc^{-m}d, wc^{-m}dc, \dots, wc^{-m}dc^m$$

with weights $k - \varphi(a) - \varphi(c), \dots, k - \varphi(a) - m\varphi(c), k - m\varphi(c), k - (m - 1)\varphi(c), \dots, k$. Since there is no initial subword of the form $\tilde{w}a$ with weight $m_a + \varphi(a)$ or less, we can take m big enough such that the minimum in the sequence of weights of \tilde{r} is unique and is attained at the d that appears after replacing the rightmost a^{-1} of the unique subword $\tilde{w}a$ of weight m_a . The corresponding subword has weight $m_a - m\varphi(c)$.

If r attains the relative minimum in an initial subword of the form $\tilde{w}a$, replace r by r^{-1} . □

Example 2.13 Consider the presentation

$$\mathcal{P} = \langle a, b, c \mid a^{-1}b^{-1}abaab^{-1}cb^{-1}a^{-1}a^{-1}a^{-1}baac^{-1}, c^{-1}b^{-1}cbccb^{-1}cb^{-1}c^{-1}c^{-1}c^{-1}baca^{-1} \rangle.$$

If we take $\varphi: G(\mathcal{P}) \rightarrow \mathbb{Z}$ defined by $\varphi(a) = \varphi(b) = \varphi(c) = 1$, the first relator attains a unique relative minimum at c . By [Corollary 2.12](#), $G(\mathcal{P})$ is locally indicable. One can check that the previous methods do not work for this presentation.

Weakly concatenable relators and I -test

[Theorems 2.5](#) and [2.9](#) apply to presentations with the homology of S^1 . Both results impose conditions on all the relators of the presentation but one. This fact will be used in the next section to extend a well-known criterion of Howie on LOT groups (see [Corollary 3.1](#)). We now investigate local indicability of groups with presentations with the homology of a wedge of circles, ie groups G with nontrivial free abelian $H_1(G)$ that admit presentations $\mathcal{P} = \langle a_1, \dots, a_n \mid r_1, \dots, r_k \rangle$ where $n - k = \text{rank}(H_1(G))$. The next result imposes weaker conditions to all the relators of the presentation.

It will be convenient to use the notion of I -values. This notion is inspired in the I -test developed in [\[2\]](#). We will work with sequences of minima. As before, one can formulate and prove analogous results for maxima.

Definition 2.14 Let \mathcal{P} be a presentation of a group G and $\varphi: G \rightarrow \mathbb{Z}$ a surjective homomorphism. By changing, if necessary, a generator by its inverse, we can assume that $\varphi(a)$ is nonnegative for every generator a of \mathcal{P} . Given a word r on the generators, the sequence of I -values of r with respect to φ is obtained by assigning to each initial subword w with the form $w = \tilde{w}x$ (where x is a generator) the value $\varphi(\tilde{w})$ and to each initial subword of the form $w = \tilde{w}x^{-1}$ the value $\varphi(\tilde{w}x^{-1})$.

This sequence is useful when considering the covering of $\mathcal{K}_{\mathcal{P}}$ corresponding to φ . The I -values of a relator r are precisely the initial vertices of the 1-cells traversed by the attaching path of R_m (where m is the minimum of its sequence of weights).

Remark 2.15 The unique relative minimum condition of [Definition 2.11](#) is equivalent to the relator having an initial subword wa^{-1} or wa with minimum I -value m_a over all initial subwords of the form $\tilde{w}a$ or $\tilde{w}a^{-1}$.

Remark 2.16 Given a relator r , the multiset of letters in which the minima of the sequence of I -values of r are attained is exactly $m_{\varphi}(r)$.

Definition 2.17 A family of relators $\{r_1, \dots, r_k\}$ is *weakly concatenable* (with respect to $\varphi: G \rightarrow \mathbb{Z}$), if there is an ordering $m_\varphi(r_{i_1}), \dots, m_\varphi(r_{i_k})$ of their multisets of minima such that, for every $1 \leq j \leq k$, there exists an element $x \in m_\varphi(r_{i_j})$ such that $x \notin \bigcup_{s=1}^{j-1} m_\varphi(r_{i_s})$ and, if the minimum I -value of r_i is m , the number of subwords wx^{-1} of r_i with I -value m is different from the number of subwords wx with I -value m .

The condition on the number of subwords with rightmost letter x in the previous definition will be used in the next result to prove that a certain 2-cell $R_{i,j}$ corresponding to the relator r_i properly involves a 1-cell corresponding to the generator x .

Theorem 2.18 Let $\mathcal{P} = \langle a_1, \dots, a_n \mid r_1, \dots, r_k \rangle$ be a group presentation with cyclically reduced relators of a group G with $H_1(G)$ nontrivial free abelian of rank $n - k$. Let $\varphi: G \rightarrow \mathbb{Z}$ be a surjective homomorphism. If $\{r_1, \dots, r_k\}$ is weakly concatenable with respect to φ , then G is locally indicable.

Proof We consider as before the covering $\tilde{\mathcal{K}}$ of $\mathcal{K}_{\mathcal{P}}$ corresponding to φ . Again, the 1-cells $a_{i,j}$ are oriented from j to $j + \varphi(a_i)$ (recall that we assume that the weights $\varphi(a_i)$ are nonnegative) and $R_{i,j}$ are the 2-cells covering R_i with minimum 0-cell j . By Theorem 2.1 it suffices to prove that every finite connected 2-subcomplex $K \subset \tilde{\mathcal{K}}$ is reducible. Note again that the no-proper-power condition follows from [1, Proposition 1] and the fact that $\tilde{\mathcal{K}}$ is an infinite cyclic cover of a 2-complex $\mathcal{K}_{\mathcal{P}}$ with the homology of a wedge of circles.

Given a finite and connected 2-subcomplex K , let $j \in \mathbb{Z}$ be the minimum 0-cell in K that is incident to some 2-cell of K . Let $R_{i_1,j}, \dots, R_{i_s,j}$ with $i_1 < \dots < i_s$ be the 2-cells incident to the 0-cell j . By hypothesis, $a_{i_s,j}$ is a face of the 2-cell $R_{i_s,j}$ and is not face of any other 2-cell $R_{i,l}$ of K . This 1-cell is properly involved in $R_{i_s,j}$ by definition of weakly concatenable, and therefore it is an elementary reduction. \square

Example 2.19 Consider the presentation $\mathcal{P} = \langle a, b, c \mid r_1, r_2 \rangle$ with relators

$$r_1 = a^{-1}c^{-1}aaab^{-1}b^{-1}c^{-1}ab, \quad r_2 = c^{-1}b^{-1}cb^{-1}caba^{-1}ba^{-1}.$$

Let $\varphi: G \rightarrow \mathbb{Z}$ be the homomorphism defined by $\varphi(a) = \varphi(b) = \varphi(c) = 1$. It is easy to verify that the multisets of minima are $m_\varphi(r_1) = \{c, a, c, a\}$ and $m_\varphi(r_2) = \{b, c, b, c\}$, and that $\{r_1, r_2\}$ is weakly concatenable with the ordering $m_\varphi(r_2), m_\varphi(r_1)$. By Theorem 2.18, $G(\mathcal{P})$ is locally indicable.

3 Applications to labeled oriented trees

We now apply the results of the previous section to derive new results on LOT groups. The first one is a straightforward application of Theorem 2.5 and is one of the main results of this paper. It is an extension of a well known result of Howie [11, Theorem 10.1] on the left and right graphs associated to a labeled oriented tree, and is also related to similar results by Gersten [7, Proposition 4.15] and Wise [17] on Adian presentations.

Recall that a labeled oriented graph (LOG) Γ consists of two sets $V(\Gamma)$ and $E(\Gamma)$ of vertices and edges, and three functions $i, t, \lambda: E \rightarrow V$ that map each edge to its initial vertex, terminal vertex and label, respectively. It is called a labeled oriented tree (LOT) when the underlying graph is a tree. The standard presentation $P(\Gamma)$ (of a group $G(\Gamma)$) associated to each LOG Γ is

$$P(\Gamma) = \langle V(\Gamma) \mid \{t(e)^{-1}\lambda(e)^{-1}i(e)\lambda(e) : e \in E(\Gamma)\} \rangle$$

Note that all relators have total exponent 0 and, if Γ is a LOT, $P(\Gamma)$ has deficiency 1 and it is a homology circle. Every LOT can be changed, by a finite sequence of transformations, to a reduced one with the same homotopy type (see [11, Section 3]). In particular one can assume that $\lambda(e) \neq i(e)$ and $\lambda(e) \neq t(e)$ for every edge e , which implies that all the relators of $P(\Gamma)$ are cyclically reduced.

In this section we will use the homomorphism $\varphi: G(\Gamma) \rightarrow \mathbb{Z}$ that sends every generator of $P(\Gamma)$ to 1. The sequence of I -values with respect to φ will be called I -sequence. Note that, given any relator r (corresponding to an edge e of Γ), its I -sequence attains exactly two minima, one at $\lambda(e)$ and the other one at $i(e)$, then $m(r) = \{\lambda(e), i(e)\}$. Analogously, it attains two maxima, one at $\lambda(e)$ and the other one at $t(e)$.

In [11] Howie introduced two graphs $I(\Gamma)$ and $T(\Gamma)$ associated to any LOG Γ . The vertex set of the left graph $T(\Gamma)$ is $V(\Gamma)$ and, for every edge e in Γ , we put an (unoriented) edge $\{\lambda(e), i(e)\}$ (these are precisely the letters in which the minima of the I -sequence are attained). Similarly, the right graph $I(\Gamma)$ has an edge $\{\lambda(e), t(e)\}$ for every edge e in Γ . In [7] Gersten investigated these graphs in the more general context of Adian presentations. An Adian presentation is one of the form


$$\mathcal{P} = \langle A \mid u_i = v_i \ (i \in J) \rangle,$$

where u_i and v_i are nontrivial positive words on A . In this context, the edges of the graph $T(\mathcal{P})$ (resp $I(\mathcal{P})$) connect the first (resp last) letter of u_i to that of v_i . Howie proved that, if $I(\Gamma)$ or $T(\Gamma)$ is a tree then $G(\Gamma)$ is locally indicable (and, in particular, Γ is aspherical) [11, Theorem 10.1]. In the case of Adian presentations, Gersten proved that if the presentation is cycle-free (ie if both graphs are forests) then $\mathcal{K}_{\mathcal{P}}$ is DR (diagrammatically reducible) and, in particular, aspherical [7, Proposition 4.12]. Moreover, when $\ell(u_i) = \ell(v_i)$ for every i (as in the case of LOTs), if either $T(\mathcal{P})$ or $I(\mathcal{P})$ has no cycles then $\mathcal{K}_{\mathcal{P}}$ is DR (and hence aspherical) [7, Proposition 4.15]. More recently, Wise showed that the 2-complexes associated to cycle-free Adian presentations have nonpositive sectional curvature, which implies that the corresponding groups are locally indicable [17, Theorem 11.4].

Note that [Theorem 2.5](#) imposes conditions to all the relators of the presentation but one. If either $I(\Gamma)$ or $T(\Gamma)$ has only one cycle, we can remove an edge of the cycle to get a forest. Then, as an immediate application of [Theorem 2.5](#) (or the analogous result for maxima), we obtain the following generalization of [11, Theorem 10.1].

Corollary 3.1 *Let Γ be a LOT. If either $I(\Gamma)$ or $T(\Gamma)$ has at most one cycle, then $G(\Gamma)$ is locally indicable.*

Example 3.2 Consider the following LOT Γ :



Neither $I(\Gamma)$ nor $T(\Gamma)$ is a tree, so Howie’s (or Gersten’s) result do not apply. Note that $T(\Gamma)$ has exactly one cycle and then, by [Corollary 3.1](#), the group $G(\Gamma)$ is locally indicable (and Γ is aspherical).

In [\[11\]](#) Howie studied some subfamilies of LOTs of diameter 4 with 3 nonextremal vertices. Using [Corollary 3.1](#) one can show local indicability for most examples of the remaining subfamilies. [Example 3.2](#) belongs to one of them, in which one of the extremal vertices appears three times as a label.

[Corollary 3.1](#) can be generalized to a more general class of Adian presentations. Suppose

$$\mathcal{P} = \langle a_1, \dots, a_n \mid u_1 = v_1, \dots, u_{n-1} = v_{n-1} \rangle$$

is an Adian presentation of deficiency 1 with $H_1(G(\mathcal{P})) = \mathbb{Z}$ (where the relators are cyclically reduced). Suppose further that there exists a surjective homomorphism $\varphi: G(\mathcal{P}) \rightarrow \mathbb{Z}$ with $\varphi(a_i) > 0$ for every a_i . For any such homomorphism φ , the multiset of minima $m_\varphi(r_i)$ of each relator r_i is the set consisting of the first letter of u_i and the first letter of v_i . Then, if either $I(\mathcal{P})$ or $T(\mathcal{P})$ has at most one cycle, as an immediate application of [Theorem 2.9](#), we obtain the following result, which extends, for a subfamily of Adian presentations, Wise’s result [\[17, Theorem 11.4\]](#).

Corollary 3.3 *Let $\mathcal{P} = \langle a_1, \dots, a_n \mid u_1 = v_1, \dots, u_{n-1} = v_{n-1} \rangle$ be an Adian presentation of deficiency 1 with $H_1(G(\mathcal{P})) = \mathbb{Z}$. Suppose further that there exists a surjective homomorphism $\varphi: G(\mathcal{P}) \rightarrow \mathbb{Z}$ with $\varphi(a_i) > 0$ for every a_i . If either $I(\mathcal{P})$ or $T(\mathcal{P})$ has at most one cycle, then $G(\mathcal{P})$ is locally indicable.*

The condition on the existence of the homomorphism φ in the previous corollary is automatically satisfied when $\ell(u_i) = \ell(v_i)$ for every i .

We finish the paper by analyzing a strategy that can be useful to study local indicability of LOT groups for which $I(\Gamma)$ and $T(\Gamma)$ have more than one cycle. The main idea is to apply convenient extended Nielsen transformations to the LOT presentation $P(\Gamma)$ to obtain an equivalent presentation \mathcal{P} which satisfies the hypotheses of [Theorem 2.18](#).

For now on we consider the labeled oriented (original) versions of $I(\Gamma)$ and $T(\Gamma)$: for every edge $a \xrightarrow{c} b$ of a LOT Γ , we put an edge $a \xrightarrow{b} c$ in $T(\Gamma)$ and an edge $c \xrightarrow{a} b$ in $I(\Gamma)$. We state the result for $T(\Gamma)$, the corresponding result for $I(\Gamma)$ is analogous.

Definition 3.4 Let Γ be a LOT. Let C be a cycle in $T(\Gamma)$ and e an edge in C . We say that e is *properly labeled* in C if the number of edges in C with the same label as e oriented clockwise is different from the number of edges with the same label as e oriented counterclockwise.

Lemma 3.5 *Let Γ be a LOT. Let e be an edge in a simple cycle C of $T(\Gamma)$ with label x . Suppose further that x does not appear as a vertex in C . Then, by applying extended Nielsen transformations, one can replace the relator r corresponding to e in $P(\Gamma)$ by another relator \tilde{r} which contains the generator x and other generators that appear as labels or vertices in C , the new relator \tilde{r} has a constant I -sequence (it attains a minimum at every letter), and in particular it attains a minimum at x . Moreover, if e is properly labeled in C , the number of initial subwords of \tilde{r} of the form wx and the number of initial subwords of the form wx^{-1} are different.*

Proof We say that a word is a *zigzag* if its length is even and the exponent of the i^{th} letter is $(-1)^{i+1}$ (ie it has the form $x_1x_2^{-1} \dots x_{2s}^{-1}$). Suppose r_1 and r_2 are relators and a is a letter in which both relators attain a minimum in the I -sequence. Let b and c be the other letters in which r_1 and r_2 attain the other minimum. Up to cyclic permutation and inversions, we can assume $r_1 = bw_1a^{-1}$ and $r_2 = aw_2c^{-1}$ with w_1, w_2 zigzags. Now, we can change r_1 by $r_1r_2 = bw_1w_2c^{-1}$ which attains exactly two minima in its I -sequence, one at b and the other one at c .

Let $C = a_1, a_2, a_3, \dots, a_l, a_1$ be the cycle of $T(\Gamma)$ and $e = (a_1, a_2)$. Let r_i be the relator corresponding to (a_i, a_{i+1}) for every $1 \leq i \leq l$ (where the indices are taken modulo l). Up to cyclic permutations and inversions, we can assume that $r_i = a_iw_i a_{i+1}^{-1}$ where w_i is a zigzag word. Now we can replace r_1 by $r_1r_2 \dots r_l = a_1w_1w_2 \dots w_l a_1^{-1}$, which, after a cyclic permutation, is $w_1w_2 \dots w_l$. Note that $r = w_1w_2 \dots w_l$ is a zigzag. Then every letter attains the same I -value (the minimum). Finally, since (a_1, a_2) is properly labeled, the number of occurrences of x in this new relator is different from the number of occurrences of x^{-1} . □

Theorem 3.6 *Let Γ be a LOT. Denote by X_0, X_1, \dots, X_k the connected components of $T(\Gamma)$. Suppose that C_1, \dots, C_k are generating simple cycles in X_0 and the other connected components X_1, \dots, X_k are trees. If for every $1 \leq i \leq k$ there is at least one properly labeled edge of C_i that has a vertex of X_i as a label and for $j \geq i$ there is no other vertex of X_j that appears as a label in the edges of C_i , then G_Γ is locally indicable.*

Proof To illustrate the idea of the proof, we show the case $k = 1$. Let e_1 be a properly labeled edge in the cycle C_1 , which is labeled with a vertex x_1 of X_1 . By Lemma 3.5 we can replace the corresponding relator r_1 by a relator \tilde{r}_1 , which attains a minimum at the vertex x_1 and in which the number of initial subwords wx_1 with minimum I -value and the number of initial subwords wx_1^{-1} with minimum I -value are different. Note that the new presentation \mathcal{P} is Andrews–Curtis equivalent to $P(\Gamma)$.

Now, the relators of the new presentation \mathcal{P} are weakly concatenable and the result follows from Theorem 2.18. The ordering for a weak concatenation from back to front is the following: first follow the order of collapses of all the edges of the tree X_1 to the vertex x_1 , then continue with the new relator \tilde{r}_1 , and finally use any ordering of weak concatenation for the remaining relators of X_0 (note that $X_0 - e_1$ is a tree).





Figure 1: $I(\Gamma)$.

The general case follows by induction in a similar way. We require that for $j \geq i$ there is no other vertex of X_j that appears as a label in the edges of C_i . This condition is necessary to guarantee weak concatenability, since by Lemma 3.5 the new relators \tilde{r}_i attain a minimum at every letter that appears in the cycle C_i . \square

Example 3.7 Consider the following LOT Γ :



Both $I(\Gamma)$ and $T(\Gamma)$ have two cycles, so we cannot apply Corollary 3.1. The graph $I(\Gamma)$ is displayed in Figure 1.

Note that $I(\Gamma)$ satisfies the hypotheses of Theorem 3.6 with X_0 the subgraph with vertices $\{b, d, 1, e, a, f\}$, $X_1 = \{0\}$ and X_2 the tree with vertices $\{2, c\}$. In the cycle $C_1 = 1, b, d, 1$ we consider the edges properly labeled with $x_1 = 0$ and in the cycle $C_2 = 1, a, e, 1$ we take the edge properly labeled with $x_2 = 2$. This shows that $G(\Gamma)$ is locally indicable.

References

- [1] **J F Adams**, *A new proof of a theorem of WH Cockcroft*, J. Lond. Math. Soc. 30 (1955) 482–488 [MR](#)
- [2] **J A Barmak, E G Minian**, *A new test for asphericity and diagrammatic reducibility of group presentations*, Proc. Roy. Soc. Edinburgh Sect. A 150 (2020) 871–895 [MR](#)
- [3] **SD Brodskii**, *Equations over groups and groups with one defining relation*, Uspekhi Mat. Nauk 35 (1980) 183 [MR](#) In Russian; translated in [Russian Math. Surv.](#) 35 (1980) 165
- [4] **A Clay, D Rolfsen**, *Ordered groups and topology*, Graduate Studies in Math. 176, Amer. Math. Soc., Providence, RI (2016) [MR](#)

- [5] **B Derooin, A Navas, C Rivas**, *Groups, orders, and dynamics*, preprint (2014) [arXiv 1408.5805](#)
- [6] **S M Gersten**, *Conservative groups, indicability, and a conjecture of Howie*, *J. Pure Appl. Algebra* 29 (1983) 59–74 [MR](#)
- [7] **S M Gersten**, *Reducible diagrams and equations over groups*, from “Essays in group theory”, *Math. Sci. Res. Inst. Publ.* 8, Springer (1987) 15–73 [MR](#)
- [8] **G Higman**, *The units of group-rings*, *Proc. Lond. Math. Soc.* 46 (1940) 231–248 [MR](#)
- [9] **J Howie**, *On pairs of 2-complexes and systems of equations over groups*, *J. Reine Angew. Math.* 324 (1981) 165–174 [MR](#)
- [10] **J Howie**, *On locally indicable groups*, *Math. Z.* 180 (1982) 445–461 [MR](#)
- [11] **J Howie**, *On the asphericity of ribbon disc complements*, *Trans. Amer. Math. Soc.* 289 (1985) 281–302 [MR](#)
- [12] **J Howie, H R Schreebeli**, *Homological and topological properties of locally indicable groups*, *Manuscripta Math.* 44 (1983) 71–93 [MR](#)
- [13] **J Howie, H Short**, *Coherence and one-relator products of locally indicable groups*, *Groups Geom. Dyn.* 17 (2023) 1085–1113 [MR](#)
- [14] **A A Klyachko, M A Mikheenko**, *Yet another Freiheitssatz: mating finite groups with locally indicable ones*, *Glasg. Math. J.* 65 (2023) 337–344 [MR](#)
- [15] **A Rhemtulla, D Rolfsen**, *Local indicability in ordered groups: braids and elementary amenable groups*, *Proc. Amer. Math. Soc.* 130 (2002) 2569–2577 [MR](#)
- [16] **H Wilton**, *Rational curvature invariants for 2-complexes*, *Proc. A.* 480 (2024) art. id. 20240025 [MR](#)
- [17] **D T Wise**, *Sectional curvature, compact cores, and local quasiconvexity*, *Geom. Funct. Anal.* 14 (2004) 433–468 [MR](#)
- [18] **D T Wise**, *Coherence, local indicability and nonpositive immersions*, *J. Inst. Math. Jussieu* 21 (2022) 659–674 [MR](#)

*Departamento de Matematica, Universidad de Buenos Aires
Buenos Aires, Argentina*

*Departamento de Matematica, Universidad de Buenos Aires
Buenos Aires, Argentina*

abarreto@dm.uba.ar, gminian@dm.uba.ar

Received: 11 December 2023 Revised: 24 May 2024

ALGEBRAIC & GEOMETRIC TOPOLOGY

msp.org/agt

EDITORS

PRINCIPAL ACADEMIC EDITORS

John Etnyre
etnyre@math.gatech.edu
Georgia Institute of Technology

Kathryn Hess
kathryn.hess@epfl.ch
École Polytechnique Fédérale de Lausanne

BOARD OF EDITORS

Julie Bergner	University of Virginia jeb2md@eservices.virginia.edu	Thomas Koberda	University of Virginia thomas.koberda@virginia.edu
Steven Boyer	Université du Québec à Montréal cohf@math.rochester.edu	Markus Land	LMU München markus.land@math.lmu.de
Tara E Brendle	University of Glasgow tara.brendle@glasgow.ac.uk	Christine Lescop	Université Joseph Fourier lescop@ujf-grenoble.fr
Indira Chatterji	CNRS & Univ. Côte d'Azur (Nice) indira.chatterji@math.cnrs.fr	Norihiko Minami	Yamato University minami.norihiko@yamato-u.ac.jp
Octav Cornea	Université de Montréal cornea@dms.umontreal.ca	Andrés Navas	Universidad de Santiago de Chile andres.navas@usach.cl
Alexander Dranishnikov	University of Florida dranish@math.ufl.edu	Jessica S Purcell	Monash University jessica.purcell@monash.edu
Tobias Ekholm	Uppsala University, Sweden tobias.ekholm@math.uu.se	Birgit Richter	Universität Hamburg birgit.richter@uni-hamburg.de
Mario Eudave-Muñoz	Univ. Nacional Autónoma de México mario@matem.unam.mx	Jérôme Scherer	École Polytech. Féd. de Lausanne jerome.scherer@epfl.ch
David Futер	Temple University dfuter@temple.edu	Vesna Stojanoska	Univ. of Illinois at Urbana-Champaign vesna@illinois.edu
John Greenlees	University of Warwick john.greenlees@warwick.ac.uk	Zoltán Szabó	Princeton University szabo@math.princeton.edu
Matthew Hedden	Michigan State University mhedden@math.msu.edu	Maggy Tomova	University of Iowa maggy-tomova@uiowa.edu
Kristen Hendricks	Rutgers University kristen.hendricks@rutgers.edu	Daniel T Wise	McGill University, Canada daniel.wise@mcgill.ca
Hans-Werner Henn	Université Louis Pasteur henn@math.u-strasbg.fr	Lior Yanovski	Hebrew University of Jerusalem lior.yanovski@gmail.com
Daniel Isaksen	Wayne State University isaksen@math.wayne.edu		

See inside back cover or msp.org/agt for submission instructions.

The subscription price for 2025 is US \$760/year for the electronic version, and \$1110/year (+\$75, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Algebraic & Geometric Topology is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [Current Mathematical Publications](#) and the [Science Citation Index](#).

Algebraic & Geometric Topology (ISSN 1472-2747 printed, 1472-2739 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004.

AGT peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**

nonprofit scientific publishing

<https://msp.org/>

© 2025 Mathematical Sciences Publishers

ALGEBRAIC & GEOMETRIC TOPOLOGY

Volume 25 Issue 7 (pages 3789–4436) 2025

Algebras for enriched ∞ -operads	3789
RUNE HAUGSENG	
An overtwisted convex hypersurface in higher dimensions	3813
RIVER CHIANG and KLAUS NIEDERKRÜGER-EID	
Presheaves of groupoids as models for homotopy types	3833
LÉONARD GUETTA	
Product and coproduct on fixed point Floer homology of positive Dehn twists	3875
YUAN YAO and ZIWEN ZHAO	
Multitwists in big mapping class groups	3921
GEORGE DOMAT, FEDERICA FANONI and SEBASTIAN HENSEL	
Cofibrantly generated model structures for functor calculus	3931
LAUREN BANDKLAYDER, JULIA E BERGNER, RHIANNON GRIFFITHS, BRENDA JOHNSON and REKHA SANTHANAM	
Endomorphisms of Artin groups of type D	3975
FABRICE CASTEL and LUIS PARIS	
Linear bounds of the crosscap number of knots	4009
ROB MCCONKEY	
Atiyah–Segal completion for the Hermitian K -theory of symplectic groups	4037
JENS HORNOSTEL, HERMAN ROHRBACH and MARCUS ZIBROWIUS	
A minimality property for knots without Khovanov 2-torsion	4073
ONKAR SINGH GUJRAL and JOSHUA WANG	
Local indicability of groups with homology circle presentations	4077
AGUSTÍN NICOLÁS BARRETO and ELÍAS GABRIEL MINIAN	
Cubulating drilled bundles over graphs	4095
MAHAN MJ and BISWAJIT NAG	
Infinitely many homeomorphic hyperbolic plugs with the same basic sets	4147
FANGFANG CHEN	
One-point compactifications of configuration spaces and the self duality of the little disks operad	4163
CONNOR MALIN	
Braided multitwists	4185
RODRIGO DE POOL	
Positive intermediate Ricci curvature on connected sums	4209
PHILIPP REISER and DAVID J WRAITH	
The \mathbb{S}_n -equivariant Euler characteristic of the moduli space of graphs	4229
MICHAEL BORINSKY and JOS VERMASEREN	
Classification of metric fibrations	4257
YASUHIKO ASAO	
Spaces over BO are thickened manifolds	4287
HIRO LEE TANAKA	
Representation-graded Bredon homology of elementary abelian 2-groups	4321
MARKUS HAUSMANN and STEFAN SCHWEDE	
The \mathbb{Z}/p -equivariant cohomology of the genus-zero Deligne–Mumford space with $1+p$ marked points	4341
DAIN KIM and NICHOLAS WILKINS	
Geometry of free extensions of free groups via automorphisms with fixed points on the complex of free factors	4357
PRITAM GHOSH and FUNDA GÜLTEPE	
Coarse cohomology of configuration space and coarse embedding	4391
ANKA BANERJEE	