

AG
T

*Algebraic & Geometric
Topology*

Volume 25 (2025)

**Geometry of free extensions of free groups via automorphisms
with fixed points on the complex of free factors**

PRITAM GHOSH

FUNDA GÜLTEPE



Geometry of free extensions of free groups via automorphisms with fixed points on the complex of free factors

PRITAM GHOSH
FUNDA GÜLTEPE

We give conditions of an extension of a free group to be hyperbolic and relatively hyperbolic using the dynamics of the action of $\text{Out}(\mathbb{F})$ on the complex of free factors combined with weak attraction theory. We work with subgroups of exponentially growing outer automorphisms and instead of using a standard ping-pong argument with loxodromics, we allow fixed points for the action and investigate the geometry of the extension group when the fixed points of the automorphisms on the complex of free factors are sufficiently far apart.

20E06, 20F28, 20F67, 57M07

1 Introduction

Let \mathbb{F} be a free group of finite rank ≥ 3 . Similar to that of mapping classes for a surface, there is a dichotomy for elements of the group of outer automorphisms $\text{Out}(\mathbb{F})$ of the free group \mathbb{F} , in terms of their *growth*. An element $\phi \in \text{Out}(\mathbb{F})$ is called *exponentially growing* if for some conjugacy class $[w]$ of an element $w \in \mathbb{F}$, the word length of $\phi^i([w])$ grows exponentially with i for any fixed generating set of \mathbb{F} . We will call a subgroup of $\text{Out}(\mathbb{F})$ exponentially growing if all of its elements are. For an exponentially growing subgroup $\mathcal{Q} < \text{Out}(\mathbb{F})$, we are interested in understanding the geometry of the extension $E_{\mathcal{Q}}$ given by the short exact sequence

$$1 \rightarrow \mathbb{F} \rightarrow E_{\mathcal{Q}} \rightarrow \mathcal{Q} \rightarrow 1$$

that is induced from the sequence

$$1 \rightarrow \mathbb{F} \rightarrow \text{Aut}(\mathbb{F}) \rightarrow \text{Out}(\mathbb{F}) \rightarrow 1.$$

The extension group $E_{\mathcal{Q}}$ is the pullback of \mathcal{Q} to $\text{Aut}(\mathbb{F})$, and hence a subgroup of $\text{Aut}(\mathbb{F})$.

When \mathcal{Q} is free group of rank ≥ 2 , which is our main focus for this paper, we say that $E_{\mathcal{Q}}$ is a free-by-free group and it is known that $E_{\mathcal{Q}} \cong \mathbb{F} \rtimes \widehat{\mathcal{Q}}$, where $\widehat{\mathcal{Q}}$ is some (any) lift of \mathcal{Q} to $\text{Aut}(\mathbb{F})$. In this paper we give necessary and sufficient conditions for $E_{\mathcal{Q}}$ to be hyperbolic using the dynamical information of the generators of $\mathcal{Q} < \text{Out}(\mathbb{F})$ obtained from train-track maps of free group automorphisms, as well as via their actions on the complex of free factors \mathcal{FF} .

An outer automorphism ϕ is *atoroidal* (or hyperbolic) if no power of ϕ fixes a nontrivial conjugacy class in \mathbb{F} . Atoroidal automorphisms are a special class of exponentially growing elements of $\text{Out}(\mathbb{F})$. Using train-track theory, Bestvina–Feighn–Handel [6] developed dynamical invariants for exponentially growing outer automorphisms of \mathbb{F} , called *attracting and repelling laminations*. Namely, associated to each exponentially growing outer automorphism ϕ (respectively, ϕ^{-1}), we have finitely many invariant sets of biinfinite *lines* in the (compactified Cayley graph of) Gromov hyperbolic space $\mathbb{F} \cup \partial\mathbb{F}$. These sets of lines are called *attracting laminations* (respectively, *repelling laminations*) and denoted by $\mathcal{L}^+(\phi)$ (respectively, $\mathcal{L}^-(\phi)$). Atoroidal outer automorphisms are characterized by the property that for every conjugacy class $[w]$, the sequence $\phi^i([w])$ converges to some element of $\mathcal{L}^+(\phi)$ as $i \rightarrow \infty$ [18, Lemma 3.1]. Ghosh showed that these sets were central to describing the Cannon–Thurston laminations for the extension group $\mathbb{F} \rtimes \langle \phi \rangle$ [18, Lemma 4.4]. We use $\mathcal{L}^\pm(\phi)$ to denote the union of the two lamination sets $\mathcal{L}^+(\phi)$ and $\mathcal{L}^-(\phi)$.

We show that $\mathcal{L}^\pm(\phi)$ is instrumental in characterizing hyperbolicity of the extension group E_Q , when Q is a free group of rank ≥ 2 . Our Theorem 1.1 shows that in this case the extension group is hyperbolic, provided we enforce the condition that the attracting and repelling laminations associated to the generators of Q to be pairwise disjoint. We also require throughout the paper that our outer automorphisms are *rotationless* (see Section 2.6), a condition that ensures that we will be able to use special relative train-track maps, which can be achieved by taking a uniform power of automorphisms [16].

Theorem 1.1 *Let ϕ_1, \dots, ϕ_k be a collection of exponentially growing outer automorphisms such that no pair of automorphisms have a common power. Then the following are equivalent:*

- (1) *Each ϕ_i is an atoroidal outer automorphism and $\mathcal{L}^\pm(\phi_i) \cap \mathcal{L}^\pm(\phi_j) = \emptyset$ for all $i \neq j$.*
- (2) *There exists $M > 0$ such that for all $m_i \geq M$, $Q = \langle \phi_1^{m_1}, \dots, \phi_k^{m_k} \rangle$ is a free group and the extension group $\mathbb{F} \rtimes \widehat{Q}$ is hyperbolic (where \widehat{Q} is any lift of Q).*

Theorem 1.1 is comparable to Theorem 1.3 of Farb and Mosher [15] and the main Theorem of [22] where hyperbolicity of the extension group $E_Q = \pi_1(S) \rtimes Q$ is characterized by the *convex cocompactness* of a free subgroup Q of the mapping class group $\text{MCG}(S)$ of a closed surface S .

We say that $\phi_i, \phi_j, i \neq j$ are *independent* if generic leaves of elements of $\mathcal{L}^\pm(\phi_i)$ and $\mathcal{L}^\pm(\phi_j)$ are not asymptotic to each other. Hence, Theorem 1.1 can be written as;

Theorem *Let ϕ_1, \dots, ϕ_k be a collection of exponentially growing outer automorphisms such that no pair of automorphisms have a common power. Then, ϕ_1, \dots, ϕ_k is a purely atoroidal (all infinite-order elements are atoroidal), pairwise independent collection if and only if the extension group $\mathbb{F} \rtimes \widehat{Q}$ for a free group $Q = \langle \phi_1^{m_1}, \dots, \phi_k^{m_k} \rangle$ generated by high enough m_i 's is hyperbolic.*

The interest in the geometry of extensions of groups started with Thurston [30] where the \mathbb{Z} -extension of $\pi_1(S)$ of a closed surface S is hyperbolic if and only if the cyclic group is generated by a pseudo-Anosov element of the mapping class group. Bestvina–Feighn [2] and Brinkmann [8] have shown that when Q is an infinite cyclic subgroup of $\text{Out}(\mathbb{F})$, E_Q is hyperbolic if and only if ϕ is atoroidal.

We do not need Q to be free to prove that the collection is independent and purely atoroidal.

1.1 Fixed points on \mathcal{FF} and hyperbolicity of the extension

Free factor complex \mathcal{FF} of the free group \mathbb{F} is a simplicial complex whose vertices are conjugacy classes of nontrivial free factors of \mathbb{F} . Two vertices corresponding to free factors are connected by an edge if one includes the other as subgroup, up to conjugation. \mathcal{FF} is Gromov-hyperbolic [3; 24] and in many other ways shows resemblance to the curve complex. Hence the action of $\text{Out}(\mathbb{F})$ on \mathcal{FF} is reminiscent of the action of the mapping class group on the curve complex.

An element of $\text{Out}(\mathbb{F})$ is *fully irreducible* if and only if it has no periodic orbits on \mathcal{FF} . It is well known that fully irreducible elements of $\text{Out}(\mathbb{F})$ act loxodromically on \mathcal{FF} [3] and this was exploited by Dowdall–Taylor [13] to produce hyperbolic extensions of free groups. Their theorem can be compared to those of Kent–Leininger [25], and Hamenstädt [22] characterizing convex cocompactness, hence the hyperbolicity of the extension, with quasi-isometric embedding of the group in the curve complex. As such the automorphisms of [13] are fully irreducible.

We take a different approach in this paper and produce hyperbolic extensions of groups whose elements are not necessarily fully irreducible, hence we allow automorphisms to have fixed points on \mathcal{FF} . The following result shows that the only condition one needs for hyperbolicity is that they do not have a common fixed point on \mathcal{FF} .

Theorem 1.2 *Let ϕ_1, \dots, ϕ_k be a collection of atoroidal elements which do not have a common power. If no pair $\phi_i, \phi_j, i \neq j$, fixes a common vertex in the free factor complex \mathcal{FF} , then*

- (1) *there exists $M > 0$ such that for all $m_i \geq M$, $Q = \langle \phi_1^{m_1}, \dots, \phi_k^{m_k} \rangle$ is a free group,*
- (2) *the extension group $\mathbb{F} \rtimes \widehat{Q}$ is a hyperbolic group (where \widehat{Q} is any lift of Q).*

Our Theorem 1.2 hence responds to the question of what happens when a pair ϕ, ψ in a collection of automorphisms do have common fixed points for their action on \mathcal{FF} .

1.2 Sufficiently different automorphisms, nonattracting sink and the geometry of the extension group

One of the motivations for this work is to establish a connection between the dynamics of the action of $\text{Out}(\mathbb{F})$ on the free factor complex and the dynamical data that we have from the train-track theory. We believe that this is a theme that has barely been explored and a lot can be learned if we are able exploit this connection.

We call a collection of automorphisms *sufficiently different* if no pair of automorphisms have a common power and for each pair of outer automorphisms in the collection, the distance between the fixed vertices in \mathcal{FF} is at least 2 (if such vertices exist) (see Section 3.1). Lemma 3.2 gives a simple criterion for constructing free subgroups of $\text{Out}(\mathbb{F})$ using sufficiently different elements.

Given a short exact sequence $1 \rightarrow \mathbb{F} \rightarrow E \rightarrow Q \rightarrow 1$ of finitely generated groups, we say that E has a *cuspid-preserving* relatively hyperbolic structure if there exists a collection of finitely generated subgroups $\{H_i\}$ of \mathbb{F} such that Q preserves conjugacy class of each H_i and E is (strongly) hyperbolic relative to the collection $\{N_E(H_i)\}$ of normalizers of H_i in E . Our main aim in Section 3.2 is to address the question of when a free subgroup Q of $\text{Out}(\mathbb{F})$ can yield relatively hyperbolic extension E with the cuspid-preserving property.

In the proposition below we conclude that being sufficiently different is an obstruction to cuspid-preserving relative hyperbolicity of the extension group, except in one particular case. Equivalently, it is an obstruction to having an *admissible subgroup system* [20, Section 2.8] for the collection of ϕ_i 's. An admissible subgroup system is a malnormal collection of subgroups with some special properties, and the *nonattracting sink* \mathcal{K}_ϕ^* of an automorphism $\phi \in \text{Out}(\mathbb{F})$, which is developed in [20] as an example of an admissible subgroup system, carries all conjugacy classes which do not grow exponentially under iteration by ϕ (see Lemma 3.6). A nonattracting sink can be computed explicitly using relative train-track maps.

For a collection ϕ_1, \dots, ϕ_k of pairwise sufficiently different exponentially growing outer automorphisms, let \mathcal{K}_i^* be the nonattracting sink of ϕ_i .

Proposition 3.10 *Let ϕ_1, \dots, ϕ_k be a collection of exponentially growing, sufficiently different outer automorphisms. If all the nonattracting sinks are nonempty then exactly one of following are true:*

- (1) ϕ_1, \dots, ϕ_k is a collection of fully irreducible geometric outer automorphisms induced by some pseudo-Anosov homomorphisms of the same compact surface with one boundary component. Moreover, all sufficiently high powers of ϕ_i 's generate a free group Q such that E_Q has a cuspid-preserving relatively hyperbolic structure.
- (2) $\mathcal{K}_i^* \neq \mathcal{K}_j^*$ for some $i \neq j$ and the extension E_Q of a free group Q , which is generated by sufficiently high powers of ϕ_i 's, cannot have a cuspid preserving relatively hyperbolic structure.

We have the following theorem characterizing the hyperbolicity of E_Q using sinks:

Theorem 1.3 *Let ϕ_1, \dots, ϕ_k be a collection of exponentially growing and sufficiently different outer automorphisms. Let Q be a free group generated by sufficiently high powers of ϕ_i 's. Then, $\mathbb{F} \rtimes \widehat{Q}$ is a hyperbolic group if and only if $\mathcal{K}_i^* = \emptyset$ for all $i \in \{1, \dots, k\}$.*

In this theorem Q is not necessarily a *convex cocompact* subgroup of $\text{Out}(\mathbb{F})$ in the sense of [11; 12; 21], since our groups do not quasi-isometrically embed in \mathcal{FF} . Hence our theorem cannot be obtained via their work. In fact, as far as our knowledge goes, this theorem cannot be obtained by studying the action of Q on any known hyperbolic simplicial complex with a nice $\text{Out}(\mathbb{F})$ action.

In a similar vein, we obtain the following necessary and sufficient condition in terms of nonattracting sinks, regarding cusp-preserving relative hyperbolicity of the extension group when Q is generated by sufficiently different collections of outer automorphisms.

Theorem 5.1 *Let ϕ_1, \dots, ϕ_k be a collection of pairwise sufficiently different and exponentially growing outer automorphisms. Let \mathcal{K}_i^* be the nonattracting sink of $\phi_i, i \in \{1, \dots, k\}$, and assume that $\mathcal{K}_j^* \neq \emptyset$ for some fixed j . Then the following are equivalent:*

- (1) $\mathbb{F} \rtimes \widehat{Q}$ has a cusp preserving relatively hyperbolic structure where Q is a free group generated by sufficiently large exponents of ϕ_i 's.
- (2) $\mathcal{K}_i^* = \mathcal{K}_j^*$ for all i .

To summarize, given a pair of sufficiently different, exponentially growing elements of $\text{Out}(\mathbb{F})$, all sinks empty gives hyperbolic extensions (Theorem 1.3). No sink empty gives relatively hyperbolic extensions with cusp-preserving structure only for geometric fully irreducibles (Proposition 3.10). Some sink nonempty gives relatively hyperbolic extensions with cusp-preserving structure only if all sinks are nonempty and are equal (Theorem 5.1).

1.3 Plan of the paper

Section 2 collects all the definitions and the tools we use.

In Section 3 we investigate obstructions to (relative) hyperbolicity using the dynamics of sufficiently different automorphisms on \mathcal{FF} . We also discuss the notion of a sink and include proof of Proposition 3.10.

In Section 4 we continue working with a specific type of exponentially growing automorphism; a reducible automorphism that is fully irreducible on a free factor (a *partial* fully irreducible). We investigate the conditions on the small displacement sets on \mathcal{FF} of partial fully irreducible automorphisms that determine the geometry of the extension groups they generate.

Section 5 includes proofs of Theorems 1.1, 1.2 and 5.1. In this section we discuss further the notion of sink and how the sinks of automorphisms determine the geometry of an extension group generated by those automorphisms and prove Theorem 1.3. We work with partial and relatively fully irreducible elements directly to exhibit examples of relatively hyperbolic free group extensions.

Section 6 discusses further problems such as characterization of nonrelative hyperbolicity; in this section we give an example which could help characterize nonrelative hyperbolicity of a free group extension.

Acknowledgements We thank Spencer Dowdall, Chris Leininger, Lee Mosher, Pranab Sardar, and Sam Taylor for their remarks and comments on the earlier version of the paper. We also would like to thank the anonymous referee for carefully reading the paper and providing numerous suggestions that have improved the exposition.

Ghosh was supported by the Ashoka University faculty research grant. Gültepe was partially supported by NSF grant DMS-2137611.

2 Preliminaries

2.1 Marked graphs, circuits and path

A *marked graph* is a graph G which is a core graph (a graph with no valence-1 vertices) that is equipped with a homotopy equivalence to the rose $m: G \rightarrow R_n$ (where $n = \text{rank}(\mathbb{F})$). Thus the fundamental group of G can be identified with \mathbb{F} (up to inner automorphisms). A *circuit* in a marked graph is a locally injective and continuous map of S^1 into G . The set of circuits in G can be identified with the set of conjugacy classes in \mathbb{F} .

A *path* is an immersion of the interval $[0, 1]$ into G with endpoints at vertices of G . Every path can be written as a finite concatenation of edges of G , so that there is no backtracking.

A *line* ℓ is a bi-infinite concatenation $\ell = \dots E_{i-1} E_i E_{i+1} \dots$, $i \in \mathbb{Z}$, of edges of G without backtracking, ie so that $E_{j-1}^{-1} \neq E_j \neq E_{j+1}^{-1}$ (where E_s^{-1} is E_s traveled in opposite direction). A *ray* γ is “one-sided” infinite concatenation of edges $\gamma = E_i E_{i+1} \dots$, $i \in \mathbb{Z}_{\geq 0}$, in G without backtracking. Two lines are said to be asymptotic if they have a common subray. Any continuous map f from S^1 or $[0, 1]$ to G can be *tightened* to a circuit or path, in other words is freely homotopic to a locally injective and continuous map. Tightened image of a path α under f will be denoted by $f_{\#}(\alpha)$ and we will not distinguish between circuits or paths that differ by a homeomorphism of their respective domains.

2.2 Topological representative, EG strata, NEG strata

A *filtration* of a marked graph G is a strictly increasing sequence $G_0 \subset G_1 \subset \dots \subset G_k = G$ of subgraphs G_r with no isolated vertices. The filtration is *f-invariant* if $f(G_r) \subset G_r$ for all r .

The *stratum of height r* is a subgraph $H_r = G_r \setminus G_{r-1}$ together with the endpoints of edges. The minimum r such that a subset of G is contained in G_r is called the *height* of the subset.

For an *f*-invariant filtration, the square matrix M_r whose j^{th} column records the number of times the image of an edge e_j under f intersects the other edges in H_r is called the *transition matrix* of the stratum H_r . M_r is said to be irreducible if for each i, j , the (i, j) entry of some power of M_r is nonzero. In this case we say that the associated stratum H_r is also *irreducible*. When H_r is irreducible, the Perron–Frobenius theorem states that the matrix M_r has a unique eigenvalue $\lambda \geq 1$, called the *Perron–Frobenius eigenvalue*, for which some associated eigenvector has positive entries. If $\lambda = 1$ then H_r is a *nonexponentially growing (NEG) stratum* whereas if $\lambda > 1$ we say that H_r is an *exponentially growing (EG) stratum*.

An automorphism $\phi \in \text{Out}(\mathbb{F})$ can be represented by a homotopy equivalence $f: G \rightarrow G$ that takes vertices to vertices and edges to edge-paths of a marked graph G with marking $\rho: R_n \rightarrow G$, called a *topological representative*. Topological representative f preserves the marking of G , in other words $\bar{\rho} \circ f \circ \rho: R_n \rightarrow R_n$ represents R_n . A nontrivial path α in G is a *periodic Nielsen path* if $f_{\#}^k(\alpha) = \alpha$ for some k , where the smallest such k is called the *period*. α is a Nielsen path if $k = 1$. A periodic Nielsen path is *indivisible (iNP)* if it cannot be written as a concatenation of nontrivial periodic Nielsen paths.

Given a topological representative $f : G \rightarrow G$ let Tf be such that $Tf(E)$ is the first edge in the edge path associated to $f(E)$. A *turn* is a pair of edges $\{E_i, E_j\}$ of edges and we let $Tf(E_i, E_j) = (Tf(E_i), Tf(E_j))$ making Tf a map that takes turns to turns. We say that a nondegenerate (i.e. $i \neq j$) turn is *illegal* if for some iterate of Tf the turn becomes degenerate; otherwise the turn is legal. A path is said to be *legal path* if it contains only legal turns and it is r -legal if it is of height r and all its illegal turns are in G_{r-1} .

Relative train-track map Given $\phi \in \text{Out}(\mathbb{F})$ and a topological representative $f : G \rightarrow G$ with a filtration $G_0 \subset G_1 \subset \dots \subset G_k$ which is preserved by f , we say that f is a relative train-track map if the following conditions are satisfied:

- (1) f maps r -legal paths to r -legal paths.
- (2) If γ is a path in G_{r-1} with endpoints in $G_{r-1} \cap H_r$ then $f_{\#}(\gamma)$ is nontrivial.
- (3) If E is an edge in H_r then $Tf(E)$ is an edge in H_r . In particular, every turn consisting of a direction of height r and one of height $< r$ is legal.

By [6, Theorem 5.1.5] every $\phi \in \text{Out}(\mathbb{F})$ has a relative train-track map representative which satisfies some useful conditions in addition to the ones we listed above.

2.3 Weak topology

Given a graph G , we define an equivalence relation on the set of all paths, rays, lines, and circuits in G by declaring two of paths/rays/lines/circuits to be equivalent if they differ by a homeomorphism of their domains.

Let $\widehat{\mathcal{B}}(G)$ denote the compact space of equivalence classes of circuits and finite paths, rays and lines in a graph G , whose endpoints (if any) are vertices of G . For each finite path γ in G , we denote by $\widehat{N}(G, \gamma)$ the set of all paths and circuits in $\widehat{\mathcal{B}}(G)$ which have γ as its subpath. The collection of all such sets gives a basis for a topology on $\widehat{\mathcal{B}}(G)$ called *weak topology*. Let $\mathcal{B}(G) \subset \widehat{\mathcal{B}}(G)$ be the compact subspace of all lines in G with the induced topology.

Two distinct points in $\partial\mathbb{F}$ determine a *line*, up to reversing the direction. Let $\widetilde{\mathcal{B}} = \{\partial\mathbb{F} \times \partial\mathbb{F} - \Delta\}/(\mathbb{Z}/2\mathbb{Z})$ be the set of pairs of boundary points of \mathbb{F} where Δ is the diagonal and \mathbb{Z}_2 acts by interchanging factors. We can give the weak topology to $\widetilde{\mathcal{B}}$, induced by the Cantor topology on $\partial\mathbb{F}$.

\mathbb{F} acts on $\widetilde{\mathcal{B}}$ and the quotient space $\mathcal{B} = \widetilde{\mathcal{B}}/\mathbb{F}$ is compact but non-Hausdorff as such the topology is called *weak*. The quotient topology is also called the *weak topology*. For any marked graph G , there is a natural identification $\mathcal{B} \approx \mathcal{B}(G)$.

We call the elements of \mathcal{B} *lines* as well. A lift of a line $\gamma \in \mathcal{B}$ is an element $\tilde{\gamma} \in \widetilde{\mathcal{B}}$ that projects to γ under the quotient map and the two elements of $\tilde{\gamma}$ are called its endpoints.

A line (or a path) γ is said to be *weakly attracted* to a line (path) β under the action of $\phi \in \text{Out}(\mathbb{F})$, if for some k $\phi^k(\gamma)$ converges to β in the weak topology, in other words, if any given finite subpath of β is

contained in $\phi^k(\gamma)$ for some k . Similarly if we have a homotopy equivalence $f: G \rightarrow G$, a line(path) γ is said to be *weakly attracted* to a line(path) β under the action of $f_\#$ if $f_\#^k(\gamma)$ weakly converges to β . the *accumulation set* of γ is the set of lines $l \in \mathcal{B}(G)$ that are elements of the weak closure of a ray γ in G . This is the set of lines l such that every finite subpath of l occurs infinitely many times as a subpath γ . Similarly, the weak accumulation set of some point $\xi \in \partial\mathbb{F}$ is the set of lines in the weak closure of any of the asymptotic rays in its equivalence class.

2.4 Free factor systems and malnormal subgroup systems

A finite collection $\mathcal{K} = \{[K_1], [K_2], \dots, [K_s]\}$ of conjugacy classes of nontrivial, finite-rank subgroups $K_s < \mathbb{F}$ is called a *subgroup system*. A subgroup $K < \mathbb{F}$ is *malnormal* if $xK_sx^{-1} \cap K$ is trivial for all $x \in \mathbb{F} - K$. A subgroup system is called *malnormal* if each of its subgroups $[K_s] \in \mathcal{K}$ is malnormal and for all $[K_s], [K_t] \in \mathcal{K}$ if $[K_s] \cap [K_t]$ is nontrivial then $s = t$. Given two malnormal subgroup systems $\mathcal{K}, \mathcal{K}'$ we define a partial ordering \sqsubset on the set of subgroup systems by $\mathcal{K} \sqsubset \mathcal{K}'$ if for each conjugacy class of subgroup $[K] \in \mathcal{K}$ there exists some conjugacy class of subgroup $[K'] \in \mathcal{K}'$ such that $K < K'$.

Given a finite collection $\{K_1, K_2, \dots, K_s\}$ of subgroups of \mathbb{F} , we say that this collection determines a *free factorization* of \mathbb{F} if $\mathbb{F} = K_1 * K_2 * \dots * K_s$. A *free factor system* $\mathcal{F} := \{[F_1], [F_2], \dots, [F_p]\}$ of \mathbb{F} is a finite collection of conjugacy classes of subgroups such that there is a free factorization of \mathbb{F} of the form $\mathbb{F} = F_1 * F_2 * \dots * F_p * B$, where B is some (possibly trivial) finite-rank subgroup of \mathbb{F} . Every free factor system is a malnormal subgroup system.

A malnormal subgroup system \mathcal{K} *carries* a conjugacy class $[c] \in \mathbb{F}$ if there exists some $[K] \in \mathcal{K}$ such that $c \in K$. We say that \mathcal{K} carries a line γ if one of the following equivalent conditions hold:

- (1) γ is the weak limit of a sequence of conjugacy classes carried by \mathcal{K} .
- (2) There exists some $[K] \in \mathcal{K}$ and a lift $\tilde{\gamma}$ of γ so that the endpoints of $\tilde{\gamma}$ are in ∂K .

For any marked graph G and any subgraph $H \subset G$, the fundamental groups of the noncontractible components of H form a free factor system, denoted by $[\pi_1(H)]$. Every free factor system \mathcal{F} can be realized as $[\pi_1(H)]$ for some nontrivial core subgraph H of some marked graph G . An equivalent way of saying that a line or circuit γ is carried by \mathcal{F} is that for any marked graph G and a subgraph $H \subset G$ with $[\pi_1(H)] = \mathcal{F}$, the realization of γ in G is contained in H .

We have the following fact:

Lemma 2.1 [23, Fact 1.8] *Given a subgroup system \mathcal{K} the set of lines/circuits carried by \mathcal{K} is a closed set in the weak topology.*

As a consequence, given a malnormal subgroup system \mathcal{K} and a sequence of lines/circuits $\{\gamma_n\}$, if \mathcal{K} carries every weak limit of every subsequence of $\{\gamma_n\}$, then γ_n is carried by \mathcal{K} for all sufficiently large n [23, Lemma 1.11].

Given two free factor systems, an extension of the notion of “intersection ” of subgroups, extended to free factor systems is called the “meet” of free factors:

Lemma 2.2 [6, Section 2.6] *Every collection $\{\mathcal{F}_i\}$ of free factor systems has a well-defined meet $\bigwedge \{\mathcal{F}_i\}$, which is the unique maximal free factor system \mathcal{F} such that $\mathcal{F} \sqsubset \mathcal{F}_i$ for all i . Moreover, for any free factor $F < \mathbb{F}$ we have $[F] \in \bigwedge \{\mathcal{F}_i\}$ if and only if there exists an indexed collection of subgroups $\{A_i\}_{i \in I}$ such that $[A_i] \in \mathcal{F}_i$ for each i and $F = \bigcap_{i \in I} A_i$.*

The free factor support $\mathcal{F}_{\text{supp}}(B)$ of a set of lines B in \mathcal{B} is defined as the meet of all free factor systems that carry B [6]. If B is a single line then $\mathcal{F}_{\text{supp}}(B)$ is single free factor. We say that a set of lines B is filling if $\mathcal{F}_{\text{supp}}(B) = \mathbb{F}$.

2.5 Attracting laminations and nonattracting subgroup systems

For any marked graph G , there is a natural identification $\mathcal{B} \approx \mathcal{B}(G)$ which induces a bijection between the closed subsets of $\mathcal{B}(\mathcal{G})$ of \mathcal{B} . A lamination Λ is a closed subset of any of these two sets. Given a lamination $\Lambda \subset \mathcal{B}$ we look at the corresponding lamination in $\mathcal{B}(G)$ as the realization of Λ in G . An element $\lambda \in \Lambda$ is called a leaf of the lamination.

A lamination Λ is called an attracting lamination for ϕ if it is the weak closure of a line ℓ (called the generic leaf of Λ) satisfying:

- ℓ is birecurrent leaf of Λ : every finite subpath of ℓ occurs infinitely many times as subpath in both directions.
- There is a neighborhood V^+ such that every line in V^+ is weakly attracted to ℓ in the weak topology. V^+ is called an attracting neighborhood of ℓ (see [6, Definition 3.1.1]).
- no lift $\tilde{\ell} \in \mathcal{B}$ of ℓ is the axis of a generator of a rank-1 free factor of \mathbb{F} .

Attracting neighborhoods of Λ are defined by choosing sufficiently long segments of generic leaves of Λ . If V^+ is an attracting neighborhood, then $\phi(V^+) \subset V^+$.

Associated to each $\phi \in \text{Out}(\mathbb{F})$ is a finite set $\mathcal{L}^+(\phi)$ of laminations, called the set of attracting laminations of ϕ [6]. Similarly we define the set of attracting laminations (or the set of repelling laminations) $\mathcal{L}^-(\phi)$ of ϕ^{-1} (or of ϕ). Both $\mathcal{L}^+(\phi)$ and $\mathcal{L}^-(\phi)$ are ϕ -invariant [6, Lemmas 3.1.13 and 3.1.6]. Given a relative train-track map $f : G \rightarrow G$ representing ϕ , then there is a bijection between the set $\mathcal{L}^+(\phi)$ and the set of exponentially growing strata in G which is determined by the height of a generic leaf of $\Lambda^+ \in \mathcal{L}^+(\phi)$ [6, Section 3].

Free factor support of an element Λ of $\mathcal{L}^+(\phi)$ or $\mathcal{L}^-(\phi)$ is the smallest free factor system that carries each leaf of Λ , denoted by $\mathcal{F}_{\text{supp}}(\Lambda)$ [6, Section 3.2]. Free factor support of $\Lambda^+ \in \mathcal{L}^+(\phi)$ is equal to free factor support of a generic leaf of Λ^+ [6, Corollaries 2.6.5 and 3.1.11]. If $\Lambda_1, \Lambda_2 \in \mathcal{L}^+(\phi)$ then $\Lambda_1 = \Lambda_2 \Leftrightarrow \mathcal{F}_{\text{supp}}(\Lambda_1) = \mathcal{F}_{\text{supp}}(\Lambda_2)$ [23, Fact 1.14].

An element of $\mathcal{L}^+(\phi)$ is dual to an element of $\mathcal{L}^-(\phi)$ if these two elements have the same free factor support. This imposes a bijection between these two sets [6, Lemma 3.2.4].

A line/circuit γ is said to be *weakly attracted* to $\Lambda_1 \in \mathcal{L}^+(\phi)$ if γ is weakly attracted to some (hence every) generic leaf of Λ under action of ϕ . No leaf of any element of $\mathcal{L}^-(\phi)$ is ever attracted to any leaf of any element of $\mathcal{L}^+(\phi)$. No Nielsen path is weakly attracted to either an element of $\mathcal{L}^+(\phi)$ or an element of $\mathcal{L}^-(\phi)$.

An attracting lamination Λ^+ and a repelling lamination Λ^- of ϕ cannot have leaves which are asymptotic, for otherwise it would violate the fact that no leaf of Λ^- is weakly attracted to Λ^+ . This allows us to choose sufficiently long subpaths of generic leaves of Λ^+ , Λ^- and construct attracting and repelling neighborhoods V^+ , V^- so that $V^+ \cap V^- = \emptyset$.

An element of $\Lambda \in \mathcal{L}^+(\phi)$ is said to be *topmost* if there does not exist any $\Lambda_j \in \mathcal{L}^+(\phi)$ such that $\Lambda \subset \Lambda_j$. Elements of $\mathcal{L}^+(\phi)$ can be divided into two distinct classes. If there exists a finite, ϕ -invariant collection of distinct, nontrivial conjugacy classes $\mathcal{C} = \{[c_1], [c_2], \dots, [c_s]\}$ such that $\mathcal{F}_{\text{supp}}(\mathcal{C}) = \mathcal{F}_{\text{supp}}(\Lambda^+)$, then $\Lambda^+ \in \mathcal{L}^+(\phi)$ is said to be *geometric* [23, Definition 2.19]. It is said to be *nongeometric* otherwise.

The *nonattracting subgroup system* of an attracting lamination was first introduced by Bestvina–Feighn–Handel in [6] and later explored in great details by Handel–Mosher in [23, Part III, pages 192–202]. Nonattracting subgroup system records the information about the lines and circuits which are not attracted to the lamination. We will list some of the properties which are central to our proofs.

Lemma 2.3 [23, Lemmas 1.5 and 1.6] *Let $\phi \in \text{Out}(\mathbb{F})$ and $\Lambda^+ \in \mathcal{L}^+(\phi)$ be an attracting lamination such that $\phi(\Lambda^+) = \Lambda^+$. Then there exists a subgroup system $\mathcal{A}_{\text{na}}(\Lambda^+)$ such that:*

- (1) $\mathcal{A}_{\text{na}}(\Lambda^+)$ is a malnormal subgroup system and the set of lines carried by $\mathcal{A}_{\text{na}}(\Lambda^+)$ is closed in the weak topology.
- (2) A conjugacy class $[c]$ is not attracted to Λ^+ if and only if it is carried by $\mathcal{A}_{\text{na}}(\Lambda^+)$.
- (3) If each conjugacy class of a finite-rank subgroup $B < \mathbb{F}$ is not weakly attracted to Λ^+ , then there exists some $A < \mathbb{F}$ such that $B < A$ and $[A] \in \mathcal{A}_{\text{na}}(\Lambda^+)$.
- (4) If Λ^- and Λ^+ are dual to each other, we have $\mathcal{A}_{\text{na}}(\Lambda^+) = \mathcal{A}_{\text{na}}(\Lambda^-)$.
- (5) $\mathcal{A}_{\text{na}}(\Lambda^+)$ is a free factor system if and only if Λ^+ is not geometric.
- (6) If $\{\gamma_n\}_{n \in \mathbb{N}}$ is a sequence of lines such that every weak limit of every subsequence of $\{\gamma_n\}$ is carried by $\mathcal{A}_{\text{na}}(\Lambda^+)$ then for all sufficiently large n , $\{\gamma_n\}$ is carried by $\mathcal{A}_{\text{na}}(\Lambda^+)$.

A pseudo-Anosov on a surface with nonempty boundary induces a fully irreducible automorphism called *geometric fully irreducible*. Geometricity was relativized by Bestvina–Feighn–Handel [6] by extending it so that it is a property of an EG stratum of a relative train-track map. The definition of geometricity is expressed in terms of the existence of what we call a “geometric model” (see [23] for details of constructions of geometric models.)

When Λ^+ is geometric, the geometric model gives us the structure of the nonattracting subgroup system [23, Definition 2.1, Lemma 2.5]. In particular, there is a unique closed indivisible Nielsen path ρ_r of height r [23, Fact 1.42].

The facts listed above show that when Λ^+ is geometric, $\mathcal{A}_{\text{na}}(\Lambda^+) = \mathcal{F} \cup \{[F_m]\}$, where \mathcal{F} is some free-factor system such that $[\pi_1 G_{r-1}] \sqsubset \mathcal{F}$. Moreover, ϕ restricted to F_m has polynomial growth and if $[c]$ is the conjugacy class determined by ρ_r , then $[c]$ is carried by $[F_m]$. We shall refer to the component $[F_m]$ as a *geometric component* of $\mathcal{A}_{\text{na}}(\Lambda^+)$. In the case H_r is a top stratum, $\mathcal{A}_{\text{na}}(\Lambda^+) = \mathcal{F} \cup \{[c]\}$.

2.6 Completely split improved relative train-track (CT) maps

A *splitting* of a line, a path or a circuit α is a concatenation $\dots \alpha_0 \alpha_1 \dots \alpha_k \dots$ of subpaths of α in G such that for all $i \geq 1$, $f_{\#}^i(\alpha) = \dots f_{\#}^i(\alpha_0) f_{\#}^i(\alpha_1) \dots f_{\#}^i(\alpha_k) \dots$, for a relative train-track map $f: G \rightarrow G$. The subpath α_i 's in a splitting are called *terms* or *components* of the splitting of α . The notation $\alpha \cdot \beta$ will denote a splitting and $\alpha\beta$ will denote a concatenation of nontrivial paths α, β .

A splitting $\alpha_1 \cdot \alpha_2 \cdots \alpha_k$ of a nontrivial path or circuit α is a *complete splitting* if each component α_i is either a single edge in an irreducible stratum, an indivisible Nielsen path, an exceptional path (see [16, Definition 4.1]) or a *taken* connecting path in a zero stratum (see [16, Definition 4.4]).

Completely split improved relative train-track (CT) maps are topological representatives with particularly nice properties. In particular, for each edge E in each irreducible stratum, the path $f_{\#}(E)$ is completely split. Moreover, for each taken connecting path α in each zero stratum, the path $f_{\#}(\alpha)$ is completely split. CT maps are guaranteed to exist for *rotationless* (see [16, Definition 3.13]) outer automorphisms, as has been shown in the following.

Lemma 2.4 [16, Theorem 4.28] *For each rotationless $\phi \in \text{Out}(\mathbb{F})$ and each increasing sequence \mathcal{F} of ϕ -invariant free factor systems, there exists a CT map $f: G \rightarrow G$ that is a topological representative for ϕ and f realizes \mathcal{F} .*

Feighn–Handel [16] showed that there exists some $k > 0$ such that, given any $\phi \in \text{Out}(\mathbb{F})$, ϕ^k is rotationless. So given any outer automorphism ϕ , some finite power of ϕ has a completely split improved relative train-track representative. In the body of our work, we will use CT's defined in [16]. In particular, we will use the properties given in [23, Definition 1.29] assume all exponentially growing outer automorphisms are rotationless.

2.7 Critical constant

Let $\phi \in \text{Out}(\mathbb{F})$ be exponentially growing and $f: G \rightarrow G$ be a CT map representing ϕ . Below we will adapt a *bounded cancellation* notion coming from Bestvina–Feighn–Handel's bounded cancellation lemma for train-track representatives of automorphisms of \mathbb{F} [5]; which is inspired by Cooper's same

named lemma [9]. To summarize, if we have a path in G which has some r -legal “central” subsegment of length greater than the critical constant given below, then this segment is protected by the bounded cancellation lemma and the length of this segment grows exponentially under iteration.

Definition 2.5 Let H_r be an exponentially growing stratum with associated Perron–Frobenius eigenvalue λ_r and let $\text{BCC}(f)$ denote the bounded cancellation constant for f . Then,

$$\frac{2 \text{BCC}(f)}{\lambda_r - 1}$$

is called the *critical constant* for H_r .

It is easy to see that for every number $C > 0$ that exceeds the critical constant, there is some $1 \geq \mu > 0$ such that if $\alpha\beta\gamma$ is a concatenation of r -legal paths where β is some r -legal segment of length $\geq C$, then the r -legal leaf segment of $f_{\#}^k(\alpha\beta\gamma)$ corresponding to β has length at least $\mu\lambda^k|\beta|_{H_r}$ (see [5, page 219]). Here $|\beta|_{H_r}$ is the length of an edge-path $\beta \subset G$ is the number of the edges of β that remain only in H_r . For the rest of this paper, let C be a number larger than the maximum of all critical constants corresponding to EG strata of $f: G \rightarrow G$.

2.8 Admissible subgroup systems and relative hyperbolicity

Given a group G and a collection $\{K_\alpha\}$ of subgroups $K_\alpha < G$, we obtain the *coned-off Cayley graph* of G or the *electrocuted G* relative to the collection $\{K_\alpha\}$ by assigning a vertex v_α for each left coset of K_α on the Cayley graph of G such that each point of a left coset of K_α is joined to (or coned-off at) v_α by an edge of length $\frac{1}{2}$. The resulting metric space is denoted by $(\widehat{G}, |\cdot|_{\text{el}})$.

A group G is said to be (weakly) relatively hyperbolic relative to the collection of subgroups $\{K_\alpha\}$ if \widehat{G} is a δ -hyperbolic metric space. G is said to be strongly hyperbolic relative to the collection $\{K_\alpha\}$ if the coned-off space \widehat{G} is weakly hyperbolic relative to $\{K_\alpha\}$ and it satisfies a certain *bounded coset penetration* property (see [14]). In this paper, when we say “relative hyperbolicity” we always mean “strong relative hyperbolicity”.

Definition 2.6 Given $\phi \in \text{Out}(\mathbb{F})$, let $\mathcal{K} = \{[K_1], [K_2], \dots, [K_p]\}$ be a subgroup system and $\mathcal{L}_{\mathcal{K}}^+(\phi)$ (respectively, $\mathcal{L}_{\mathcal{K}}^-(\phi)$) denote the collection of attracting (respectively, repelling) laminations of ϕ whose generic leaves are not carried by \mathcal{K} . Assume:

- (1) $\mathcal{L}_{\mathcal{K}}^+(\phi), \mathcal{L}_{\mathcal{K}}^-(\phi)$ are both nonempty.
- (2) \mathcal{K} is a malnormal subgroup system.
- (3) $\phi(K_s) = K_s$ for each $s \in \{1, \dots, p\}$.
- (4) Let V^+ denote the union of attracting neighborhoods of elements of $\mathcal{L}_{\mathcal{K}}^+(\phi)$ defined by generic leaf segments of length $\geq 2C$. Define V^- similarly for $\mathcal{L}_{\mathcal{K}}^-(\phi)$. By increasing C if necessary,

$$V^+ \cap V^- = \emptyset.$$

- (5) Every conjugacy class which is not carried by \mathcal{K} is weakly attracted to some element of $\mathcal{L}_{\mathcal{K}}^+(\phi)$.

We will call subgroup systems satisfying the properties above an *admissible subgroup system* for ϕ (where $\phi \in \text{Out}(\mathbb{F})$ is exponentially growing). If we are given some finitely generated group $Q = \langle \phi_1, \phi_2, \dots, \phi_k \rangle$ we say that \mathcal{K} is an *admissible subgroup system* for Q if \mathcal{K} is an admissible subgroup system for each ϕ_i . We will simply write “admissible subgroup system” when the context is clear.

In our previous work, we proved:

Theorem 2.7 [20] *Let ϕ_1, \dots, ϕ_k be a collection of exponentially growing outer automorphisms of \mathbb{F} such that ϕ_i, ϕ_j do not have a common power whenever $i \neq j$. Then the following are equivalent:*

- (1) *There exists a subgroup system $\mathcal{K} = \{[K_1], \dots, [K_p]\}$ which is admissible for each ϕ_i and for which $\mathcal{L}_{\mathcal{K}}^{\pm}(\phi_i) \cap \mathcal{L}_{\mathcal{K}}^{\pm}(\phi_j) = \emptyset$ whenever $i \neq j$.*
- (2) *For every free group Q generated by sufficiently high powers of ϕ_i 's, $\mathbb{F} \rtimes \widehat{Q}$ is hyperbolic relative to a collection $\{K_s \rtimes \widehat{Q}_s\}_{s=1}^p$ where \widehat{Q}_s is a lift of Q that fixes K_s .*

3 Detecting obstructions to (relative) hyperbolicity on the complex of free factors

Why high powers? Throughout this paper we have worked with a subgroup Q which becomes a free group after taking high powers of outer automorphisms, which satisfy certain conditions. We wish to clarify the reasons for resorting to high powers. Firstly, there does not exist a good (for our purposes) hyperbolic metric space where hyperbolic outer automorphisms act loxodromically. The second necessity comes from a technicality, which we describe via the following example:

Let ϕ be the outer automorphism class of the automorphism corresponding to

$$\Phi: a \mapsto ad, b \mapsto a, c \mapsto b, d \mapsto c$$

and let ψ be the outer automorphism class of the automorphism corresponding to

$$\Psi: a \mapsto ac, b \mapsto a, c \mapsto b, d \mapsto db.$$

Observe that both ϕ and ψ are positive automorphisms and one can get a relative train-track map (in fact a CT map) on the standard rose with four petals. ϕ has only one strata (the full graph) and ψ has an NEG strata (namely d) sitting on top of an EG strata (namely $\{a, b, c\}$), but the NEG edge d grows exponentially under iteration. In both cases, we have exactly one EG strata and the corresponding attracting lamination has a trivial nonattracting subgroup system by design.

Thus ϕ, ψ are both atoroidal outer automorphisms (since nonattracting subgroup system for the unique attracting lamination is trivial in both cases). Also, they have disjoint laminations since by design the free factor support of the attracting lamination of ψ is $[\langle a, b, c \rangle]$ whereas ϕ is fully irreducible, hence the associated lamination fills. These automorphisms are sufficiently different and have empty sinks.

If one looks at the extension group E_Q associated to the group $Q = \langle \phi, \psi \rangle$, then it follows that $\Phi^{-1}\Psi$ fixes both b, c and hence the group E_Q has multiple copies of $\mathbb{Z} \oplus \mathbb{Z}$, so E_Q cannot be hyperbolic. Now consider $H = \langle \Phi^{-1}\Psi, b, c \rangle$. If E_Q were relatively hyperbolic, then H would be conjugate into some peripheral subgroup (call it P) in this relatively hyperbolic structure. But $b \in \Phi H \Phi^{-1}$ together with malnormality implies that H and $\Phi H \Phi^{-1}$ are both contained in P . Hence $\Phi b \Phi^{-1} \in P \implies a \in P$. Applying the same argument again, $\Phi a \Phi^{-1} \in P \implies ad \in P \implies d \in P$. As a result, $\mathbb{F} < P$. Hence, E_Q cannot be relatively hyperbolic.

However, as we shall see later if the group Q , whose generators are sufficiently different with empty corresponding sinks is generated instead by *sufficiently high powers of ϕ and ψ* , its extension E_Q is hyperbolic –which is something one expects. This example highlights a tractable but key issue due to which we need to pass to high powers throughout this paper.

3.1 Sufficiently different outer automorphisms and their dynamics on \mathcal{FF}

In this section we will let our exponentially growing automorphisms have fixed points, but require those fixed points to be sufficiently apart from each other in \mathcal{FF} . We begin our analysis on \mathcal{FF} and combine dynamics of automorphisms on \mathcal{FF} with the dynamics on the set of laminations via weak attraction theory.

Definition 3.1 A collection of outer automorphisms will be called *sufficiently different* if

- no pair of the automorphisms have a common power (hence they do not generate a virtually cyclic (elementary) subgroup of $\text{Out}(\mathbb{F})$), and
- fixed vertices (if any) of these outer automorphisms on \mathcal{FF} are distance at least 2 from each other in \mathcal{FF} .

Given that \mathcal{FF} has infinite diameter, it is natural to expect that two randomly chosen exponentially growing outer automorphisms will be sufficiently different. Hence, we study their dynamics in relation to one another.

Lemma 3.2 Let ϕ_1, \dots, ϕ_k be a collection of exponentially growing outer automorphisms of \mathbb{F} that are sufficiently different. Then,

- (1) No generic leaf of any attracting or repelling lamination of a ϕ_i is carried by the nonattracting subgroup system of any attracting or repelling lamination of ϕ_j , for $i \neq j$.
- (2) There exists some M such that whenever $m_i \geq M$, for all i , $\langle \phi_1^{m_1}, \dots, \phi_k^{m_k} \rangle$ is a free group.

Proof If no ϕ_i fixes any vertex in the free factor complex, then they are all fully irreducible and the attracting and repelling laminations associated to them fill \mathbb{F} (see Lemma 2.2 and the paragraph afterwards). The result then follows from an induction argument applied to [5, Proposition 3.7].

Suppose that some ϕ_i fixes some vertex in \mathcal{FF} . Let Λ_j^+ be an attracting lamination of ϕ_j , $j \neq i$. If $\mathcal{F}_{\text{supp}}(\Lambda_j^+)$ fills then we are done as in this case no generic leaf of Λ_j^+ can be carried by the nonattracting subgroup system of any attracting or repelling lamination of ϕ_i . Suppose then that $\mathcal{F}_{\text{supp}}(\Lambda_j^+)$ is proper and Λ_j^+ is carried by $\mathcal{A}_{\text{na}}(\Lambda_i^+)$ for some attracting lamination Λ_i^+ of ϕ_i . This would imply that $\mathcal{F}_{\text{supp}}(\Lambda_j^+)$ is carried by a free factor component $[B] \in \mathcal{A}_{\text{na}}(\Lambda_i^+)$. This violates the distance requirement of being sufficiently different. So the result follows.

To prove (2), we claim that there exists an attracting (respectively, repelling) lamination Λ_i^+ (respectively, Λ_i^-) of ϕ_i and an attracting (respectively, repelling) lamination Λ_j^+ (respectively, Λ_j^-) of ϕ_j such that no generic leaf of Λ_i^+ or Λ_i^- is asymptotic to any generic leaf of Λ_j^+ or Λ_j^- . Assuming the claim to be true we are done by [23, Corollary 2.17] and applying induction to [6, Lemma 3.4.2] (detecting \mathbb{F}_2 via laminations).

To prove the claim, suppose first that there exists some attracting lamination Λ of ϕ_i such that $\mathcal{F}_{\text{supp}}(\Lambda)$ is proper. If some generic leaf of Λ is asymptotic to some generic leaf of an attracting or repelling lamination of ϕ_j then birecurrency of generic leaves implies that both ends of a leaf would have the same height, and as a result these lines would have the same free factor support. Hence, $\mathcal{F}_{\text{supp}}(\Lambda)$ will be invariant under both ϕ_i and ϕ_j , thus violating the distance requirement in \mathcal{FF} between their fixed vertices. So the claim is proved in the case of existence of any attracting or repelling lamination for either ϕ_i or ϕ_j with a proper free factor support. Suppose then neither ϕ_i nor ϕ_j have an attracting lamination with a proper free factor support. This implies that there exist unique attracting laminations Λ_i^+ , Λ_j^+ for ϕ_i, ϕ_j , respectively, that fill \mathbb{F} . The filling condition implies that Λ_i^+, Λ_j^+ are necessarily topmost. If Λ_i^+, Λ_j^+ have asymptotic generic leaves, say ℓ_i, ℓ_j , respectively, then ℓ_i cannot be carried by the nonattracting subgroup system of Λ_j^+ . Also, asymptoticity will imply that ℓ_i cannot be weakly attracted to Λ_j^- . Hence by the weak attraction theorem [6, Theorem 6.0.1] ℓ_i must be a generic leaf of Λ_j^+ . But by being a generic leaf, closure of ℓ_i is all of Λ_j^+ (and it is also all of Λ_i^+ by the nature of ℓ_i). Hence $\Lambda_i^+ = \Lambda_j^+$. As a result, in this case the following proposition completes the proof of our claim by giving us a contradiction with the hypothesis that no pair ϕ_i, ϕ_j have common power. \square

The following is a technical result which identifies dynamical conditions for detecting when a subgroup of $\text{Out}(\mathbb{F})$ is virtually cyclic.

Proposition 3.3 *Let ϕ_1, \dots, ϕ_k be a collection of exponentially growing automorphisms of $\text{Out}(\mathbb{F})$ for which the following conditions are satisfied:*

- (1) *No pair ϕ_i, ϕ_j with $i \neq j$ have a common invariant free factor.*
- (2) *There exists a common attracting lamination Λ^+ such that Λ^+ fills.*

Then all ϕ_i are fully irreducibles with common powers.

Proof We will prove the theorem for $k = 2$ to avoid extra notation as the general case is the same. Consider the group $\mathcal{H} = \langle \phi_1, \phi_2 \rangle < \text{Out}(\mathbb{F})$. Then $\mathcal{H} < \text{Stab}(\Lambda^+)$. If either ϕ_1 or ϕ_2 is fully irreducible, then $\text{Stab}(\Lambda^+)$ is virtually cyclic by [5, Theorem 2.14]. So \mathcal{H} must be virtually cyclic and hence ϕ_1, ϕ_2 are both fully irreducible and must have common powers.

Suppose then that both ϕ_1, ϕ_2 are reducible. Then the group \mathcal{H} is fully irreducible, since ϕ_1, ϕ_2 do not have any common invariant free factor. By applying [23, Theorem C', page 2] with $\mathcal{F} = \emptyset$, we see that there exists a fully irreducible $\xi \in \mathcal{H} < \text{Stab}(\Lambda^+)$. Let $\Lambda_\xi^+, \Lambda_\xi^-$ be the unique attracting and repelling laminations of ξ and ℓ^+ be a generic leaf of Λ_ξ^+ . Then Λ_ξ^+ and Λ_ξ^- both fill \mathbb{F} . If $\Lambda^+ = \Lambda_\xi^+$ or Λ_ξ^- , then we are reduced to the earlier case. So suppose Λ^+ is distinct from Λ_ξ^+ and Λ_ξ^- . Choose a train-track map $f: G \rightarrow G$ for ξ . Let V_ξ^+ be some attracting neighborhood of Λ_ξ^+ , which is described by some sufficiently long generic leaf segment of ℓ^+ . Similarly choose a repelling neighborhood V_ξ^- . Now, by our assumption, if γ^+ is a generic leaf of Λ^+ , then γ^+ fills, and it is not a generic leaf of Λ_ξ^- . Then by the weak attraction theorem [23, Corollary 2.17] together with [23, Theorem G], there exists some $k > 0$ such that $f_\#^k(\gamma^+) \in V_\xi^+$. Since $\xi \in \text{Stab}(\Lambda^+)$ and $f_\#^k(\gamma^+)$ is always birecurrent, $f_\#^k(\gamma^+) \in \Lambda^+$ is a birecurrent leaf. Hence we conclude that some birecurrent leaf of Λ^+ contains the subpath of ℓ^+ which described V_ξ^+ . As γ^+ was chosen to be generic leaf, we have $\overline{\gamma^+} = \Lambda^+$. Hence the chosen subpath of ℓ^+ is also a subpath of γ^+ . Since this subpath was chosen arbitrarily, we deduce that ℓ^+ is in the closure of generic leaves of Λ^+ , implying that $\Lambda_\xi^+ \subset \Lambda^+$.

Now consider the dual lamination Λ^- of Λ^+ . Since ℓ^+ is a filling line, by the weak attraction theorem of [6, Theorem 6.0.1] either ℓ^+ is weakly attracted to Λ^- or ℓ^+ is a generic leaf of Λ^+ . Since being weakly attracted to a lamination is an open condition and every subpath of ℓ^+ is a subpath of some generic leaf of Λ^+ , the first option is not possible. Hence ℓ^+ must be a generic leaf of Λ^+ , implying that $\Lambda^+ = \Lambda_\xi^+$, which is a contradiction to our assumption that they are distinct. Therefore $\Lambda^+ = \Lambda_\xi^+$ or $\Lambda^+ = \Lambda_\xi^-$ and we are done by the first case. \square

Remark 3.4 When Λ_1^\pm and Λ_2^\pm are both geometric and ϕ_1, ϕ_2 satisfy the hypothesis of Lemma 3.2, it may seem that no additional restriction is placed on the conjugacy class representing the unique indivisible Nielsen path associated to the laminations. However, we shall see in Lemma 3.9 that the hypothesis that the fixed vertex sets of ϕ_1 and ϕ_2 are of distance at least 2 in \mathcal{FF} puts some very strong restrictions on them.

3.2 Cusp preserving relative hyperbolicity

Given a short exact sequence of finitely generated groups $1 \rightarrow \mathbb{F} \rightarrow E \rightarrow Q \rightarrow 1$, we say that E has a *cusp-preserving* relatively hyperbolic structure if there exists a collection of finitely generated subgroups $\{H_i\}$ of \mathbb{F} such that Q preserves conjugacy class of each H_i and E is (strongly) hyperbolic relative to the collection $\{N_E(H_i)\}$ (where $N_E(H_i)$ is the normalizer of H_i in E) (see [20, Theorem 5.1] for motivation behind this property).

Recall that an outer automorphism ϕ is said to be *polynomially growing* if for each automorphism $\Phi \in \text{Aut}(\mathbb{F})$ representing ϕ and each $c \in \mathbb{F}$ the cyclically reduced word length of $\phi^i(c)$ is bounded above by a polynomial of i .

A subgroup $Q < \text{Out}(\mathbb{F})$ is called *upper polynomially growing (UPG)* if each $\phi \in Q$ is polynomially growing and Q has unipotent image in $\text{GL}_n(\mathbb{Z})$. If $Q < \text{Out}(\mathbb{F})$ is UPG we will call every $\phi \in Q$ also UPG.

Recall that if Λ_1, Λ_2 are attracting laminations for ϕ with nonattracting subgroup systems $\mathcal{K}_1, \mathcal{K}_2$, respectively, then we define the *meet* (Lemma 2.2) $\mathcal{K}_1 \wedge \mathcal{K}_2$ as follows:

$$\mathcal{K}_1 \wedge \mathcal{K}_2 = \{[A \cap B^w] : [A] \in \mathcal{K}_1, [B] \in \mathcal{K}_2, w \in \mathbb{F}, A \cap B^w \neq \{\text{id}\}\},$$

where B^w denotes the group $w^{-1}Bw$, with $w \in \mathbb{F}$.

Definition 3.5 (sink of an automorphism [20, Section 4.3]) Given any exponentially growing outer automorphism ϕ of \mathbb{F} , consider the full list of attracting laminations $\{\Lambda_q\}_{q=1}^r$ and let

$$\mathcal{K}_\phi^* = \mathcal{K}_1 \wedge \mathcal{K}_2 \wedge \dots \wedge \mathcal{K}_r, \quad \text{where } \mathcal{K}_q = \mathcal{A}_{\text{na}}(\Lambda_q) \text{ for } q = 1, \dots, r.$$

If ϕ is not exponentially growing define $\mathcal{K}_\phi^* = \{[\mathbb{F}]\}$. We shall call \mathcal{K}_ϕ^* the *nonattracting sink* of ϕ .

The nonattracting sink \mathcal{K}_ϕ^* of ϕ is thus a ϕ -invariant malnormal subgroup system which carries all conjugacy classes which are either fixed or grow polynomially under iteration by ϕ . Thus, by construction, the nonattracting sink is designed to trap obstructions to hyperbolicity properties of the extension groups.

Lemma 3.6 Given an exponentially growing $\phi \in \text{Out}(\mathbb{F})$ and its nonattracting sink \mathcal{K}_ϕ^* , we have:

- (1) The nonattracting sink of ϕ is an admissible subgroup system for ϕ .
- (2) A conjugacy class is carried by \mathcal{K}_ϕ^* if and only if it is not weakly attracted to any attracting lamination of ϕ .
- (3) A conjugacy class is carried by \mathcal{K}_ϕ^* if and only if it is either fixed by ϕ or grows polynomially under iteration by ϕ .
- (4) $\mathcal{K}_\phi^* = \mathcal{F} \cup \{[F_1], \dots, [F_p]\}$, for some free factor system \mathcal{F} and finite-rank ϕ -invariant subgroups $F_i \in \mathbb{F}$, each of which is a geometric component of the nonattracting subgroup system of some geometric lamination of ϕ .
- (5) If \mathcal{K}_ϕ^* has only infinite cyclic components, then no conjugacy class has nontrivial polynomial growth under iteration by ϕ .

Proof (1) and (2) follows from [20, Corollary 4.13]. (3) follows from the fact that any conjugacy class grows exponentially under iteration by ϕ if and only if it is weakly attracted to some attracting lamination of ϕ under iteration.

(4) follows from the proof of part (1) of [20, Proposition 4.12].

To prove (5) note that if $[w]$ is a conjugacy class that grows polynomially under iteration by ϕ , then there must be at least a rank-2 subgroup which carries both $[w]$ and a twistor to generate at least linear-order growth. This proves that if \mathcal{K}_ϕ^* has only infinite cyclic components, then no conjugacy class can have nontrivial polynomial growth. \square

Remark If $\mathcal{K}_\phi^* = \mathcal{F} \cup \{[F_1], \dots, [F_p]\}$ then each nontrivial $[F_i]$ arises due to presence of a geometric strata and restriction of ϕ to each nontrivial F_i is polynomially growing. Moreover, each nontrivial $[F_i]$ carries the conjugacy of the unique closed indivisible Nielsen path associated to a geometric strata. This is an essential ingredient in the proof of [20, Proposition 4.12]

We now proceed with the following proposition which identifies a sufficient condition for a collection of outer automorphisms to not have a common admissible subgroup system.

Proposition 3.7 *Let ϕ_1, \dots, ϕ_k be a collection of exponentially growing automorphisms so that no pair of elements have a common power. Let Q be any free group generated by some powers of ϕ_i 's. Let \mathcal{K}_i^* be the nonattracting sink of ϕ_i and assume that the following conditions hold:*

- (1) $\mathcal{K}_i^* \neq \emptyset$ for some $i \in \{1, \dots, k\}$.
- (2) If $\mathcal{K}_i^* \neq \emptyset$, then \mathcal{K}_i^* is not carried by $\mathcal{A}_{\text{na}}(\Lambda_j^+)$ for any attracting lamination Λ_j^+ of ϕ_j for some $j \neq i$.

Then we have the following:

- (a) There is no admissible subgroup system for Q .
- (b) The extension group $\mathbb{F} \rtimes \widehat{Q}$ is not hyperbolic relative to any collection of subgroups so that the cusps are preserved.

Proof Suppose that (1) and (2) hold. Without loss of generality assume that $\mathcal{K}_1^* \neq \emptyset$. Using item (3) of Lemma 2.3, condition (2) implies that \mathcal{K}_1^* cannot be carried by any admissible subgroup system for ϕ_j , for some $j \neq 1$. If \mathcal{K} were an admissible subgroup system for Q , then every conjugacy class not carried by \mathcal{K} would get weakly attracted to some element of $\mathcal{L}_{\mathcal{K}}^+(\phi_s)$ (hence would grow exponentially under iteration by ϕ_s), for each $1 \leq s \leq k$. Hence (a) holds.

Now suppose that $\mathbb{F} \rtimes \widehat{Q}$ is hyperbolic relative to some collection of peripheral subgroups. Let $[g]$ be carried by \mathcal{K}_i^* . Then $[g]$ is not weakly attracted to any attracting lamination of ϕ_i and hence by definition either $[g]$ is fixed by ϕ_i or $|\phi_i^n([g])|$ grows polynomially. So g must be conjugate to some element in a peripheral subgroup. As a consequence, the peripheral subgroup system in Theorem 2.7 is nonempty. Moreover, whenever $\mathbb{F} \rtimes \widehat{Q}$ preserves cusps, the same theorem ensures the existence of an admissible subgroup system, contradicting (a) above. This completes proof of (b). \square

Corollary 3.8 *Let ϕ_1, \dots, ϕ_k be a collection of exponentially growing and sufficiently different outer automorphisms of \mathbb{F} . Then if some ϕ_i fixes the conjugacy class of some rank-1 free factor $[\langle g \rangle]$, then $[g]$ can not be carried by a free factor component of the nonattracting subgroup system of any attracting lamination of ϕ_j , for any $j \neq i$.*

Moreover, if the distance between the fixed points sets in \mathcal{FF} of any distinct pair of ϕ_i, ϕ_j is at least 5, then $[g]$ is weakly attracted to every attracting lamination of ϕ_j for every $j \neq i$.

Proof For concreteness, suppose ϕ_1 fixes the conjugacy class of a rank-1 free factor $[\langle g \rangle]$. Let Λ_j^+ be any attracting lamination of ϕ_j , $1 \neq j$. If $[g]$ is carried by $[B] \in \mathcal{A}_{na}(\Lambda_j^+) = \mathcal{F}_j \cup [F_{m_j}]$, then either $[B]$ is a free factor component of \mathcal{F}_j or $[B] = [F_{m_j}]$. The former case directly violates the sufficiently different hypothesis.

To complete the proof, it remains to show that if the distance between the fixed points sets in \mathcal{FF} of any distinct pair of ϕ_i, ϕ_j is at least 5, then $[g]$ is weakly attracted to every attracting lamination of ϕ_j for every $j \neq i$. To see this, observe that if $[g]$ is carried by $[F_{m_j}]$, then the free factor conjugacy class $[\langle g \rangle]$ is disjoint from $\mathcal{F}_{supp}(\Lambda_j^+)$ and hence their distance in \mathcal{FF} is at most 4 — violating our hypothesis. \square

In Proposition 3.7 we identified sufficient conditions for a collection of exponentially growing outer automorphisms to not have a common admissible subgroup system. In the following lemma, we have a hypothesis which negates that sufficient condition and examine the impact of our hypothesis on the nonattracting sinks for a collection of sufficiently different outer automorphisms. As it turns out, the nonattracting sinks in such cases will have a very rigid structure.

For an automorphism ϕ , closed Nielsen paths are circuits which are in bijective correspondence with conjugacy classes which are fixed by ϕ . There are three primary cases in our analysis where Nielsen paths appear. First one (and the most important for us) as an indivisible Nielsen path associated to a geometric EG strata. The second one is from the “twistor” in Dehn twist parts. The third case is when there is an isolated invariant rank-1 free factor, not covered by the previous two cases. All Nielsen paths are carried by the nonattracting sink. In case of absence of a free factor component in the nonattracting sink, which is essentially what the hypothesis of Lemma 3.9 ensures, the second and third cases of Nielsen paths are ruled out.

Lemma 3.9 *Let ϕ_1, \dots, ϕ_k be a collection of sufficiently different exponentially growing outer automorphisms of $\text{Out}(\mathbb{F})$. Let \mathcal{K}_i^* be the nonattracting sink of ϕ_i for each i and suppose that the following condition holds:*

- *If $\mathcal{K}_i^* \neq \emptyset$ then for each $j \neq i$, there exists an attracting lamination Λ_j^+ of ϕ_j such that \mathcal{K}_i^* is carried by $\mathcal{A}_{na}(\Lambda_j^+)$.*

Then the following are true:

- (a) *If $\mathcal{K}_i^* \neq \emptyset$ for some i , then $\mathcal{K}_i^* \neq \emptyset$ for every i . Moreover, each \mathcal{K}_i^* has exactly one component.*
- (b) *If $\mathcal{K}_i^* \neq \emptyset$ for some i , then there exists a finite-rank subgroup F such that $\mathcal{K}_i^* = [F]$ for each i .*
- (c) *There exists an admissible subgroup system for the collection ϕ_1, \dots, ϕ_k .*

Proof For concreteness assume that $\mathcal{K}_i^* = \mathcal{F}_i \cup [F_{i_1}] \cup \cdots \cup [F_{i_n}]$ is nonempty and let $j \neq i$. Let Λ_j^+ be an attracting lamination of ϕ_j such that $\mathcal{A}_{\text{na}}(\Lambda_j^+) = \mathcal{F}_j \cup [F_j]$ carries \mathcal{K}_i^* . Sufficiently different condition implies that each component of the free factor system \mathcal{F}_i must be carried by $[F_j]$. Moreover, if the conjugacy class $[g_{i_k}]$ carried by the geometric component $[F_{i_k}]$, which represents the indivisible Nielsen path associated to the corresponding geometric strata, is a free factor conjugacy class, then $[F_{i_k}]$ must be carried by $[F_j]$ due to sufficiently different condition. On the other hand if $[g_{i_k}]$ is not a free factor conjugacy class and is carried by some component of \mathcal{F}_j , then the same component will also carry the free factor support of $[g_{i_k}]$, hence the free factor support of the corresponding geometric lamination (which is invariant under ϕ_i)—violating sufficiently different condition. Thus each $[F_{i_k}]$ (and hence entire \mathcal{K}_i^*) must be carried by $[F_j]$. This also proves that \mathcal{K}_j^* is nonempty as $[F_j]$ is one of the components of the sink of ϕ_j .

Now, switching the roles of i, j in the above argument, we see that there exists an attracting lamination Λ_i^+ with a nonattracting subgroup system $\mathcal{F}'_i \cup [F_{i_l}]$ for some $1 \leq l \leq n$ and entire \mathcal{K}_j^* is carried by $[F_{i_l}]$. But we know that $[F_j]$ carries each $[F_{i_k}]$, thus proving that $[F_j] = [F_{i_l}]$ and $[F_{i_k}]$ is trivial if $k \neq l$. For the final step, observe that we have reduced to the case $\mathcal{K}_i^* = \mathcal{F}_i \cup [F_j]$ is carried by $\mathcal{A}_{\text{na}}(\Lambda_j^+) = \mathcal{F}_j \cup [F_j]$. If $[B] \in \mathcal{F}_i$ is nontrivial free factor component, as seen earlier—it must be carried by $[F_j]$, which violates malnormality of \mathcal{K}_i^* . Hence $\mathcal{K}_i^* = [F_j]$ has a single component. Since i was chosen arbitrarily, this proves (a).

To prove (b), observe that in the paragraph above, we have actually shown that $\mathcal{K}_i^* = [F_j] = [F_{i_l}] = \mathcal{K}_j^*$. Since i, j are arbitrary here, we may let $[F] = [F_j]$ and complete the proof of (b).

For the final part, we note that if all of the nonattracting sinks are empty, then we take $\mathcal{K} = \emptyset$ to be our admissible subgroup system, otherwise we may take $\mathcal{K} = \{[F]\}$ as our admissible subgroup system. \square

Having identified the structure of the nonattracting sinks for a sufficiently different collection of exponentially growing outer automorphisms, we proceed to answering the question about when we can expect extension groups to have a cusp-preserving relative hyperbolic structure. The answer breaks down into three cases. The first case that we deal with is the following proposition, when all the nonattracting sinks are nonempty. The remaining two cases when either all nonattracting are empty (Theorem 1.3) or the mixed case with some being empty (Theorem 5.1) are dealt with later.

Proposition 3.10 *Let ϕ_1, \dots, ϕ_k be a collection of exponentially growing, sufficiently different outer automorphisms. If all the nonattracting sinks are nonempty then exactly one of following are true:*

- (1) ϕ_1, \dots, ϕ_k is a collection of fully irreducible geometric outer automorphisms induced by some pseudo-Anosov homomorphisms of the same compact surface with one boundary component. Moreover, all sufficiently high powers of ϕ_i 's generate a free group Q such that E_Q has a cusp-preserving relatively hyperbolic structure.
- (2) $\mathcal{K}_i^* \neq \mathcal{K}_j^*$ for some $i \neq j$ and the extension E_Q of a free group Q , which is generated by sufficiently high powers of ϕ_i 's, cannot have a cusp preserving relatively hyperbolic structure.

Proof When (1) holds, the work of Bowditch [7] and Mj–Reeves [27] for mapping classes of once punctured surfaces shows that sufficiently high powers of ϕ_i 's generate a free group Q of rank k and the corresponding extension group $\mathbb{F} \rtimes \widehat{Q}$ is hyperbolic relative to the collection $\{\langle \sigma \rangle \oplus \langle \phi_i \rangle\}$, where σ is the conjugacy class corresponding to the boundary of the surface. This gives us a cusp-preserving relatively hyperbolic structure and an admissible subgroup system $\{\langle \sigma \rangle\}$ for Q , implying that (2) fails.

Suppose that (1) is false. We argue by contradiction to show that (2) must be true. Our argument involves showing that we cannot satisfy the hypothesis of Lemma 3.9, hence we must satisfy the hypothesis of Proposition 3.7. This will allow us to conclude that admissible subgroup system does not exist. Suppose on the contrary that we do indeed satisfy the hypothesis of Lemma 3.9. Conclusion (2) of Lemma 3.9 proves the existence of a finite-rank subgroup $[F]$ such that $\mathcal{K}_i^* = [F]$ for each i . Moreover, the proof shows that this $[F]$ is actually a geometric component of \mathcal{K}_i^* , for each i , ie, for each ϕ_i there exists some (geometric) attracting lamination Λ_i^+ such that $\mathcal{A}_{\text{na}}(\Lambda_i^+) = \mathcal{F}_i \cup [F]$ for some free factor system \mathcal{F}_i .

If F is not infinite cyclic, then this means there are NEG edges attached to the boundary circle of the surface which supports the lamination Λ_i^+ for each i . Using the fact that $\mathcal{A}_{\text{na}}(\Lambda_i^+)$ cannot carry the free factor support of Λ_i^+ , we may conclude that if $[F]$ is not cyclic, then there is a common free factor conjugacy class (namely, the free factor support of the surface noted earlier) which is left invariant by each ϕ_i — thus violating sufficiently different hypothesis. Thus F must be infinite cyclic, say $\langle c \rangle$, where $[c]$ represents the conjugacy class of the word representing the boundary of the surface which supports all the laminations Λ_i^+ 's. If the free factor support of this surface was proper, then we would violate sufficiently different hypothesis. Hence the surface and therefore $[c]$ must fill, ie $\mathcal{F}_{\text{supp}}([c]) = \mathbb{F}$. This implies that the geometric stratum corresponding to the Λ_i^+ is a top stratum for each i . Moreover, a geometric lamination being bottommost [23, Proposition 2.15] implies that any of the lower strata cannot be EG, since the lamination fills. But since the sink of each ϕ_i is just $\langle [c] \rangle$, the lower filtration is just trivial (contractible) in each case. Hence $\mathcal{A}_{\text{na}}(\Lambda_i^+) = [F] = [c]$ for each i .

Using [17, Lemma 5.3], this implies that each ϕ_i is a geometric fully irreducible map which is induced by a pseudo-Anosov homeomorphism on the same compact surface (with one boundary component), bringing us back to (1) — which we had assumed to fail — hence giving us a contradiction. In conclusion, if (1) fails, then we must satisfy the hypothesis of Proposition 3.7, and hence (2) follows using Theorem 2.7. \square

4 Small displacement sets, weak attraction theory and further obstructions to (relative) hyperbolicity

To further our analysis and suggested by the previous section, we will consider a *small displacement sets* in \mathcal{FF} for exponentially growing outer automorphisms. We will then investigate interactions between attracting/repelling laminations, their corresponding nonattracting subgroup systems and the small displacements sets.

4.1 Small displacement sets of partial fully irreducibles

Definition 4.1 Let F be some free-factor of \mathbb{F} which is invariant under $\psi \in \text{Out}(\mathbb{F})$. If the restriction of ψ to F is fully irreducible element of $\text{Out}(F)$, then we say that ψ is partial fully irreducible on F .

Consider the isometric action of $\text{Out}(\mathbb{F})$ on the free factor complex, \mathcal{FF} . For an automorphism ϕ and constant $C > 0$, we define the *small displacement set* of $\langle \phi \rangle$ on \mathcal{FF} to be

$$\mathcal{S}_C = \{x \in \mathcal{FF} : \exists k \neq 0 \text{ such that } d(x, \phi^k(x)) \leq C\}.$$

Below we will prove that a small displacement set of a partial fully irreducible outer automorphism has a diameter bounded from above. For that we will need a few definitions regarding \mathcal{FF} .

A *marking* of a graph G is a homotopy equivalence $\mathcal{R}_n \rightarrow G$ from a rose \mathcal{R}_n . A metric on G is a function that assigns a positive number (length) to each edge of G . The (unprojectivized) *outer space* is a space of marked metric graphs which is introduced by Culler and Vogtmann in [10] as an analog of Teichmüller space. We will denote by CV the *projectivized* outer space, in which the graphs will all have total volume 1. For the details we refer the reader to [10; 31].

We will use the coarse projection $\pi : \text{CV} \rightarrow \mathcal{FF}$ defined as follows. For each proper subgraph Γ_0 of a marked graph G that contains a circle, its image in \mathcal{FF} is the conjugacy class of the smallest free factor containing Γ_0 . Now by [3], for two such proper subgraphs Γ_1 and Γ_2 , $d_{\mathcal{FF}}(\pi(\Gamma_1), \pi(\Gamma_2)) \leq 4$ [3, Lemma 3.1]. Then for $G \in \text{CV}$ we define

$$\pi(G) := \{\pi(\Gamma) \mid \Gamma \text{ is a proper, connected, noncontractible subgraph of } G\}$$

We will call the induced map $\text{CV} \rightarrow \mathcal{FF}$ also π which is clearly a *coarse* projection in that the diameter of each $\pi(G)$ is bounded by 4.

For a point $G \in \text{CV}$ and $A < \mathbb{F}$ we consider the core graph $A|G$ corresponding to the conjugacy class of A . We will denote the projection of this core subgraph to $\mathcal{FF}(A)$ by $\pi_A(G)$.

In other words, $\pi_A(G) = \pi(A|G)$ where $\pi : \text{CV} \rightarrow \mathcal{FF}$ is the coarse Lipschitz map that takes a graph to the collection of all the proper subgraphs it contains. The image of this projection is of diameter at most 4, by Bestvina–Feighn [3]. The pulled back metric via the immersion $p : A|G \rightarrow G$ gives that $A|G \in \text{CV}(A)$, the outer space of A .

Let A and B be conjugacy classes of two free factors. Assume that A and B are not disjoint, in other words they are not free factors of a common free splitting of \mathbb{F} and one of them is not included in the other. In this case A and B said to *overlap*. For two overlapping conjugacy classes of free factors, we have the following definition.

Definition 4.2 [4; 28; 29] Let A, B be two overlapping free factors with rank of A at least 2. Then the *subfactor projection* $\pi_A(B)$ of B to A is defined to be

$$\bigcup \{\pi_A(G) : G \in \text{CV} \text{ and } B|G \subset G\},$$

where $B|G \subset G$ will mean embedding of the core graph $B|G$ in G .

In other words, given any tree T with a vertex stabilizer B and \mathbb{F} action, A fixes a tree T^A . If T^A is not degenerate, which is guaranteed by the overlapping condition, the induced action of A on T^A gives a set of vertex stabilizers by the Bass–Serre theory. We define $\pi_A(B)$ to be this set of vertex stabilizers.

Remark Any two rank-1 free factors of \mathbb{F} are either disjoint or equal. There is no possibility of overlap. Hence the definition of subfactor projections uses $\text{rank} \geq 2$.

Lemma 4.3 (bounded geodesic image theorem [4; 28]) *For a free group of rank $n \geq 3$ there is a number $M_0 \geq 0$ such that if A is a free factor and γ is a geodesic in \mathcal{FF} such that every vertex of γ meets A , then*

$$\text{diam}\{\pi_A(\gamma)\} \leq M_0.$$

Proposition 4.4 *Let ϕ be partially fully irreducible automorphism with respect to a free factor F^1 with minimum translation length λ on F^1 . Then, whenever $\lambda \geq M_0$ and for every $C > 0$, there exists a sufficiently large constant $D = D(C, \lambda)$ such that the diameter of the small displacement set*

$$\mathcal{S}_{\phi,C} = \{x \in \mathcal{FF} : \exists k \neq 0 \text{ such that } d_{\mathcal{FF}}(x, \phi^k(x)) \leq C\}$$

corresponding to $\langle \phi \rangle$ is bounded above by D .

Proof Let $x \in \mathcal{S}_C$. Then, by definition of this set, for some $k > 0$ we have $d_{\mathcal{FF}}(x, \phi^k(x)) \leq C$. Let also $G \in \text{CV}$ such that $\pi(G) = x$. Now, let $\{G_t\}$ be a folding path in CV between G and $\phi^k(G)$. The projection path $\pi(\{G_t\})$ in \mathcal{FF} is an unparametrized quasigeodesic between x and $\phi^k(x)$ and it is Hausdorff close to a geodesic [3].

Now, since F^1 is of rank at least 2, the subfactor projection to the free factor complex $\mathcal{FF}(F^1)$ is coarsely defined. Moreover, by the hypothesis (or for a suitable power $> k$ of ϕ), $d_{F^1}(\tilde{x}, \phi^k_{|F^1}(\tilde{x})) > M_0$ where \tilde{x} is the subfactor projection of x to F^1 and M_0 is the constant from the Lemma 4.3. By Lemma 4.3 again, there is a vertex F^2 along $\pi(\{G_t\})$ which does not project to $\mathcal{FF}(F^1)$. By [28] this means that either F^1 and F^2 are disjoint or one is included in the other as subgroups. Hence we have $d_{\mathcal{FF}}(F^1, F^2) \leq 4$.

Use triangle inequality to deduce that $d_{\mathcal{FF}}(x, F^1) \leq 4 + C + C_1$ where C_1 is the Hausdorff distance between $\pi(\{G_t\})$ and the geodesic between x and $\phi^k(x)$. Hence, $D = 2(4 + C + C_1)$. □

The following proposition captures the impact of the separation of small displacement sets of automorphisms $\phi_i, \phi_j, i \neq j$, on the weak attraction property of their corresponding lamination sets. We continue using the same notation we set up at the beginning of this subsection (see the paragraph before Proposition 5.3).

Thus, let \mathcal{S}_ϕ denote the small displacement set of $\langle \phi \rangle$ in \mathcal{FF} . We define the distance between the small displacement sets of ϕ_i and ϕ_j in \mathcal{FF} to be

$$d_{\mathcal{FF}}(\mathcal{S}_{\phi_i}, \mathcal{S}_{\phi_j}) := \min\{d_{\mathcal{FF}}([A], [B]) : A \in \mathcal{S}_{\phi_i}, B \in \mathcal{S}_{\phi_j}\}$$

Proposition 4.5 Let ϕ_i be partially fully irreducible on F^i for all $i \in \{1, \dots, k\}$. Assume that $d_{\mathcal{FF}}(\mathcal{S}_{\phi_i}, \mathcal{S}_{\phi_j}) \geq 5$ for all $i \neq j$.

- (1) There exists some M such that whenever $m_i \geq M$, for all i , $\langle \phi_1^{m_1}, \dots, \phi_k^{m_k} \rangle$ is a free group.
- (2) For $i \neq j$, ϕ_i and ϕ_j do not have any common invariant free factors.
- (3) For all $i \in \{1, \dots, k\}$, let Λ_i^\pm be any attracting lamination of ϕ_i and $\mathcal{A}_{\text{na}}(\Lambda_i^\pm)$ be nontrivial. For a pair $\phi_i, \phi_j, i \neq j$, let $\{[A_k^i]\}_{k=1}^p = \mathcal{A}_{\text{na}}(\Lambda_i^\pm)$ and $\{[B_s^j]\}_{s=1}^q = \mathcal{A}_{\text{na}}(\Lambda_j^\pm)$. Then, $\{[A_k^i \cap B_s^j]\}_{k,s}$ is a malnormal subgroup system.
- (4) No generic leaf of any attracting or repelling lamination of a ϕ_i is carried by a nonattracting subgroup system for **any** nongeometric dual lamination pair of ϕ_j , for $j \neq i$.
- (5) If $[g]$ is a nontrivial conjugacy class of a free factor which is fixed by ϕ_i , then $[g]$ is cannot be carried by the nonattracting subgroup system of any attracting lamination of any ϕ_j for $j \neq i$.

Proof (1) is straightforward using Lemma 3.2.

Any invariant free factor of ϕ_i is contained in the respective small displacement set. When the respective small displacement sets are disjoint, the distance between any invariant free factor of ϕ_i and any invariant free factor of ϕ_j is greater than 0. Hence we cannot have a common invariant free factor. This proves (2).

Every free factor component of $\mathcal{A}_{\text{na}}(\Lambda_i^\pm)$ is a vertex in the small displacement set of ϕ_i . When the small displacement sets are of distance at least 5, every free factor component of $\mathcal{A}_{\text{na}}(\Lambda_i^\pm)$ of rank ≥ 2 meets every free factor component of $\mathcal{A}_{\text{na}}(\Lambda_j^\pm)$ of rank ≥ 2 . Malnormality of the collection of subgroups $\{[A_k^i \cap B_s^j]\}_{k,s}$ follows directly from the malnormality of $\mathcal{A}_{\text{na}}(\Lambda_i^\pm)$, for all i . This proves (3).

Let Λ_j^+ be any attracting lamination for ϕ_j with nontrivial nonattracting subgroup system $\mathcal{A}_{\text{na}}(\Lambda_j^\pm)$. By definition every free factor component of $\mathcal{A}_{\text{na}}(\Lambda_j^\pm)$ is invariant under ϕ_j . Hence every free factor component of $\mathcal{A}_{\text{na}}(\Lambda_j^\pm)$ is an element of the small displacement set of ϕ_j . Let ℓ be a generic leaf of some attracting lamination $\Lambda_i^+, i \neq j$, and let $[B] \in \mathcal{A}_{\text{na}}(\Lambda_j^\pm)$ be some free factor component of rank ≥ 2 . Assume ℓ is carried by $[B]$. Then Λ_i^+ is carried by $[B]$ which implies that $\mathcal{F}_{\text{supp}}(\Lambda_i^+) < [B]$, since $\mathcal{F}_{\text{supp}}(\Lambda_i^+)$ is the smallest free factor system that contains Λ_i^+ . But $\mathcal{F}_{\text{supp}}(\Lambda_i^+)$ is invariant under ϕ_i and hence $\mathcal{F}_{\text{supp}}(\Lambda_i^+) \in \mathcal{S}_{\phi_i}$ which violates $d_{\mathcal{FF}}(\mathcal{S}_{\phi_i}, \mathcal{S}_{\phi_j}) \geq 5$ when $i \neq j$. This completes the proof of (4).

Let $[g]$ be the conjugacy class of a rank-1 free factor fixed by ϕ_1 . If $j \neq 1$ and Λ_j^+ is an attracting lamination of ϕ_j , with $\mathcal{A}_{\text{na}}(\Lambda_j^\pm) = \mathcal{F}_j \cup [F_{m_j}]$. If $[g]$ is carried by \mathcal{F}_j , we would violate the distance between small displacement sets ≥ 5 condition. So suppose that $[g]$ is carried by $[F_{m_j}]$. In this case, note that the free factor support of Λ_j^+ is not carried by $[F_{m_j}]$ and hence the free factor conjugacy classes $[g]$ and $\mathcal{F}_{\text{supp}}(\Lambda_j^+)$ are disjoint and therefore violate that the small displacement sets have distance ≥ 5 . Therefore, $[g]$ cannot be carried by $\mathcal{A}_{\text{na}}(\Lambda_j^\pm)$. \square

The following proposition produces examples similar to Example 6.2 when the small displacement sets are sufficiently separated.

Proposition 4.6 *Let ϕ_i be partially fully irreducible on F^i and assume $d_{\mathcal{FF}}(\mathcal{S}_{\phi_i}, \mathcal{S}_{\phi_j}) \geq 5$ for all $j \neq i$. Suppose that some ϕ_i fixes a rank-1 free factor $[g]$. Then $\mathbb{F} \rtimes \widehat{Q}$ is not hyperbolic relative to any collection of subgroups so that cusps are preserved.*

Proof By Proposition 4.5(5), $[g]$ is not carried by the nonattracting subgroup system of any attracting lamination of any ϕ_j for $j \neq i$. This tells us that $[g]$ is weakly attracted to every attracting lamination of ϕ_j , $j \neq i$. Using the fact that $[g]$ is carried by \mathcal{K}_i^* , Proposition 3.7 now completes the proof. \square

5 Geometry of free by free extensions

5.1 Hyperbolicity of the extension from independent atoroidal automorphisms

Now we are ready to characterize hyperbolicity of the extension group when automorphisms are atoroidal.

Theorem 1.1 *Let ϕ_1, \dots, ϕ_k be a collection of exponentially growing outer automorphisms such that no pair of automorphisms have a common power. Then the following are equivalent:*

- (1) *Each ϕ_i is an atoroidal outer automorphism and $\mathcal{L}^\pm(\phi_i) \cap \mathcal{L}^\pm(\phi_j) = \emptyset$ for all $i \neq j$.*
- (2) *There exists $M > 0$ such that for all $m_i \geq M$, $Q = \langle \phi_1^{m_1}, \dots, \phi_k^{m_k} \rangle$ is a free group and the extension group $\mathbb{F} \rtimes \widehat{Q}$ is hyperbolic (where \widehat{Q} is any lift of Q).*

Proof We prove the result for $k = 2$ for the sake of simplicity of the notation. The general case is an exact replica of the argument.

(1) \implies (2) Since ϕ_1, ϕ_2 are both hyperbolic, by [18, Lemma 3.1] every conjugacy class grows exponentially under iteration by both of them. So $\mathcal{K}_1^* = \mathcal{K}_2^* = \emptyset$. We will use Theorem 2.7 for $\mathcal{K} = \mathcal{K}_1^* = \mathcal{K}_2^* = \emptyset$ (there is no restriction on \mathcal{K} being nonempty in that theorem). To this end, all that remains is to show that the set of laminations are pairwise disjoint. Let $f: G \rightarrow G$ be a CT-map representing ϕ_1 . If a generic leaf $\ell_1 \in \Lambda_1^+ \in \mathcal{L}^\pm(\phi_1)$ has a common end with a generic leaf $\ell_2 \in \Lambda_2^+ \in \mathcal{L}^\pm(\phi_2)$, then birecurrence implies that both ends of ℓ_i have height s , where H_s is the exponentially growing strata corresponding to Λ_1^+ and $\ell_1, \ell_2 \subset G_s$. Since ℓ_2 is asymptotic to ℓ_1 and being weakly attracted to the dual lamination Λ_1^- is an open condition, ℓ_2 cannot be weakly attracted to Λ_1^- under iteration by ϕ_1^{-1} . Using [6, Proposition 6.0.8], we get that ℓ_2 is a generic leaf of Λ_1^+ . Hence $\Lambda_1^+ = \bar{\ell}_2 = \Lambda_2^+$, violating our hypothesis. Hence (2) follows from Theorem 2.7 as $\mathbb{F} \rtimes \widehat{Q}$ is hyperbolic relative to empty sets by this theorem.

(2) \implies (1) Hyperbolicity of $\mathbb{F} \rtimes \widehat{Q}$ implies that the Cannon–Thurston map for the inclusion $\iota: \mathbb{F} \rightarrow \mathbb{F} \rtimes \widehat{Q}$ exists. Since ϕ_1, ϕ_2 do not have a common power, $\phi_1^\infty, \phi_2^\infty$ represent two distinct points in the boundary of $\mathbb{F} \rtimes \widehat{Q}$ corresponding the forward end of the axis generated by the elements ϕ_1, ϕ_2 , respectively. Using [26, Proposition 5.1], we see that the ending lamination sets of corresponding to ϕ_1^∞ and ϕ_2^∞

must be disjoint. Since [18, Theorem 3.1, Lemma 4.4] guarantees that generic leaves of all attracting laminations for ϕ_i is contained in the ending lamination set corresponding to ϕ_i^∞ and we know that the ending lamination set corresponding to any boundary point is a closed set, we get that ϕ_1 and ϕ_2 have no common attracting lamination. Similar arguments, while working with pairs of distinct boundary points $\{\phi_1^\infty, \phi_2^{-\infty}\}, \{\phi_1^{-\infty}, \phi_2^{-\infty}\}, \{\phi_1^{-\infty}, \phi_2^\infty\}$, gives us $\mathcal{L}^\pm(\phi_1) \cap \mathcal{L}^\pm(\phi_2) = \emptyset$. \square

A simple application of Theorem 2.7 given below demonstrates a way to construct free-by-free hyperbolic groups. We neither need the automorphisms to be fully irreducible, nor do we need quasi-isometrically embedded orbits for the quotient group for the hypothesis. The converse to the corollary is obviously false.

Theorem 1.2 *Let ϕ_1, \dots, ϕ_k be a collection of atoroidal elements which do not have a common power. If no pair $\phi_i, \phi_j, i \neq j$, fixes a common vertex in the free factor complex \mathcal{FF} , then*

- (1) *there exists $M > 0$ such that for all $m_i \geq M$, $Q = \langle \phi_1^{m_1}, \dots, \phi_k^{m_k} \rangle$ is a free group,*
- (2) *the extension group $\mathbb{F} \rtimes \widehat{Q}$ is a hyperbolic group (where \widehat{Q} is any lift of Q).*

Proof Let $\Lambda^\pm \in \mathcal{L}^\pm(\phi_i) \cap \mathcal{L}^\pm(\phi_j)$ be an attracting lamination for some $i \neq j$. If $\mathcal{F}_{\text{supp}}(\Lambda)$ is proper, then ϕ_i, ϕ_j both fix the vertex corresponding to $\mathcal{F}_{\text{supp}}(\Lambda)$, giving us a contradiction. If Λ fills \mathbb{F} , then Proposition 3.3 gives us a contradiction. Hence $\mathcal{L}^\pm(\phi_i) \cap \mathcal{L}^\pm(\phi_j) = \emptyset$ and the conclusions follow from Theorem 1.1. \square

5.2 Hyperbolicity and relative hyperbolicity from sufficiently different automorphisms

In this section we will focus more on the action on \mathcal{FF} of our outer automorphisms; fixing some free factors.

Theorem 1.3 *Let ϕ_1, \dots, ϕ_k be a collection of exponentially growing and sufficiently different outer automorphisms. Let Q be a free group generated by sufficiently high powers of ϕ_i 's. Then, $\mathbb{F} \rtimes \widehat{Q}$ is a hyperbolic group if and only if $\mathcal{K}_i^* = \emptyset$ for all $i \in \{1, \dots, k\}$.*

Proof Suppose that $\mathcal{K}_i^* = \emptyset$ for all $i \in \{1, \dots, k\}$. If none of the ϕ_i fixes a vertex in \mathcal{FF} , then they are all fully irreducibles which do not have a common power, and we are done by [5, Theorem 5.2]. Suppose some ϕ_i fixes a vertex in \mathcal{FF} and $\mathcal{K}_i^* = \emptyset$ for all $i \in \{1, \dots, k\}$. To apply Theorem 1.1 it is enough to show that $\mathcal{L}^\pm(\phi_i) \cap \mathcal{L}^\pm(\phi_j) = \emptyset$ for all $j \neq i$.

Since \mathcal{K}_i^* is trivial every conjugacy class in \mathbb{F} is weakly attracted to some attracting lamination of ϕ_i . Hence ϕ_i is hyperbolic (atoroidal) outer automorphism for all $i \in \{1, \dots, k\}$.

Case 1 Let Λ_i^+ be an attracting lamination of ϕ_i such that $\mathcal{F}_{\text{supp}}(\Lambda_i^+)$ is a proper free factor. Let $j \neq i$. If ϕ_j is fully irreducible, then Λ_i^+ cannot be an attracting or repelling lamination of ϕ_j since ϕ_j has unique attracting and repelling laminations which fill. If ϕ_j is reducible, then Λ_i^+ cannot be an attracting or repelling lamination of ϕ_j for otherwise $\mathcal{F}_{\text{supp}}(\Lambda_i^+)$ would be a common invariant free factor and this contradicts the fact that any vertex fixed by ϕ_i is distance at least 2 from any vertex fixed by ϕ_j .

Case 2 Suppose that Λ_i^+ fills. Pick some $j \neq i$. Then Λ_i^+ is not an attracting lamination of ϕ_j for otherwise hypotheses of Proposition 3.3 are satisfied, which contradicts with the fact that ϕ_i, ϕ_j do not have a common power. Same contradiction occurs if Λ_i^+ is a repelling lamination of ϕ_j by replacing ϕ_j with ϕ_j^{-1} in Proposition 3.3. Therefore ϕ_i and ϕ_j do not have any common attracting or repelling laminations. Now apply Theorem 1.1 to get hyperbolicity of $\mathbb{F} \rtimes \widehat{Q}$.

The converse part easily follows from the observation that hyperbolicity of $\mathbb{F} \rtimes \widehat{Q}$ implies ϕ_i, ϕ_j are both hyperbolic outer automorphisms for all $i \neq j, i, j \in \{1, \dots, k\}$, and using [18, Lemma 3.1]. \square

The following theorem gives a necessary and sufficient condition for $\mathbb{F} \rtimes \widehat{Q}$ to have a cusp preserving relative hyperbolic structure when Q is generated by a collection of sufficiently different outer automorphisms. One should read the following result as a relative hyperbolic analog of Theorem 1.3.

Theorem 5.1 *Let ϕ_1, \dots, ϕ_k be a collection of pairwise sufficiently different and exponentially growing outer automorphisms. Let \mathcal{K}_i^* be the nonattracting sink of $\phi_i, i \in \{1, \dots, k\}$, and assume that $\mathcal{K}_j^* \neq \emptyset$ for some fixed j . Then the following are equivalent:*

- (1) $\mathbb{F} \rtimes \widehat{Q}$ has a cusp preserving relatively hyperbolic structure where Q is a free group generated by sufficiently large exponents of ϕ_i 's.
- (2) $\mathcal{K}_i^* = \mathcal{K}_j^*$ for all i .

Proof Lemma 3.2 ensures high enough m_i 's so that $Q = \langle \phi_1^{m_1}, \dots, \phi_k^{m_k} \rangle$ is a free group. If (2) holds, then (1) immediately follows from Theorem 2.7.

Suppose next that for a free group Q generated by sufficiently large exponents of ϕ_i 's, $\mathbb{F} \rtimes \widehat{Q}$ has a relatively hyperbolic structure where cusps are preserved. By Proposition 3.7 \mathcal{K}_j^* must be carried by nonattracting subgroup system of some $\Lambda_i \in \mathcal{L}^+(\phi_i)$ for each $i \in \{1, \dots, k\}, i \neq j$. Lemma 3.9(b) completes the proof. \square

5.3 Relatively hyperbolic extensions using partially fully irreducibles

In this section we will give examples of relatively hyperbolic extensions using specific types of exponentially growing outer automorphisms. Our first example is for an exponentially growing outer automorphism which is *partially* fully irreducible.

Let ϕ_1, \dots, ϕ_k be a collection of partially fully irreducibles, each with respect to a free factor F^i with $[F^i] \neq [F^j]$ whenever $i \neq j$. We denote the corresponding dual attracting and repelling lamination pair of each ϕ_i by Λ_i^+ and Λ_i^- . By partial fully irreducibility we have $\mathcal{F}_{\text{supp}}(\Lambda_i^\pm) = [F^i]$. We use these conventions below.

Lemma 5.2 *Let ϕ_i be partially fully irreducible on $F^i, i \in \{1, \dots, k\}$. Then Λ_i^\pm and Λ_j^\pm have asymptotic leaves if and only if $[F^i] = [F^j]$.*

Proof Since the restriction of ϕ_i to $[F^i]$ is fully irreducible, every leaf of Λ_i^+ and of Λ_i^- is generic. If $\ell_i \in \Lambda_i^+$ and $\ell_j \in \Lambda_j^+$ have a common end, birecurrence of generic leaves implies that ℓ_i is carried by $[F^j] = \mathcal{F}_{\text{supp}}(\Lambda_j^+)$. But since $\mathcal{F}_{\text{supp}}(\Lambda_i^+) = \mathcal{F}_{\text{supp}}(\ell_i) = [F^i]$, we must have $[F^i] = [F^j]$. The converse direction is obvious. \square

Proposition 5.3 *Let ϕ_i be partially fully irreducible on F^i , where $[F^i] \neq [F^j]$ whenever $i \neq j$, $i, j \in \{1, \dots, k\}$.*

- (1) *For all $i \in \{1, \dots, k\}$, $[F^i]$ and free factor components of $\mathcal{A}_{\text{na}}(\Lambda_i^\pm)$ are close in the free factor complex \mathcal{FF} .*
- (2) *For $i \neq j$, $\mathcal{A}_{\text{na}}(\Lambda_i^\pm)$ carries a leaf of Λ_j^+ if and only if $[F^j]$ is carried by $\mathcal{A}_{\text{na}}(\Lambda_i^\pm)$. Equivalently, $\mathcal{A}_{\text{na}}(\Lambda_i^\pm)$ carries a leaf of Λ_j^+ if and only if $[F^j]$ is distance ≤ 1 from some free factor component of $\mathcal{A}_{\text{na}}(\Lambda_i^\pm)$.*
- (3) *If $[F^i]$ is at distance at least 2 from every free factor component of $\mathcal{A}_{\text{na}}(\Lambda_j^\pm)$, then any generic leaf of Λ_i^+ is weakly attracted to Λ_j^+ under the action of ϕ_i (ϕ_i^{-1}), for all $i, j \in \{1, \dots, k\}$, $i \neq j$.*

Proof Fix i . To prove (1), we show that any representative of $[F^i]$ and any representative of any free factor component of $\mathcal{A}_{\text{na}}(\Lambda_i^\pm)$ intersect at identity only. If they are not, there is some nontrivial conjugacy class $[g]$ of F^i which is carried also by $\mathcal{A}_{\text{na}}(\Lambda_i^\pm)$. Since ϕ_i restricted to F^i is fully irreducible, this implies that Λ_i^+ must be geometric and $[g]$ must be a representative of the unique closed indivisible Nielsen path corresponding to Λ_i^+ . But this implies that $[g]$ cannot be a free factor of \mathbb{F} and hence cannot be carried by any free factor component of $\mathcal{A}_{\text{na}}(\Lambda_i^\pm)$. The contradiction proves our claim.

Since the restriction of each ϕ_i to F^i is fully irreducible, every leaf of Λ_i^+ is generic and hence not a circuit. If $\ell \in \Lambda_i^+$ is any leaf which is carried by $\mathcal{A}_{\text{na}}(\Lambda_j^\pm)$, $i \neq j$, malnormality of $\mathcal{A}_{\text{na}}(\Lambda_j^\pm)$ implies that there must be some free factor conjugacy class $[H] \in \mathcal{A}_{\text{na}}(\Lambda_j^\pm)$ which carries Λ_i^+ (since the set of lines carried by $[H]$ is closed and $\bar{\ell} = \Lambda_i^+$). By definition of free factor support we then get $F^i \leq H$ up to conjugation, which is equivalent to saying that the distance between $[F^i]$ and $[H]$ in the free factor complex of \mathbb{F} is ≤ 1 . The converse follows directly from definitions.

The proof of (3) is exactly the same as the proof of [19, Lemma 5.2] (the cited proof does not anywhere use the assumptions of being atoroidal on the automorphisms) or one can see [23, Remark 2.8, page 208]. \square

Lemma 5.4 *Let $\psi \in \text{Out}(\mathbb{F})$ be partially fully irreducible with respect to F . Then there exists a ψ -invariant, malnormal subgroup system $\mathcal{K} = \{[K_1], \dots, [K_p]\}$ such that the extension group $\mathbb{F} \rtimes \langle \psi \rangle$ is strongly hyperbolic relative to the collection of subgroups $\{K_s \rtimes_{\Psi_s} \mathbb{Z}\}_{s=1}^p$, where Ψ_s is a chosen lift of ϕ such that $\Psi_s(K_s) = K_s$.*

Proof If ψ is partially fully irreducible with respect to F , then restriction of ψ to F is fully irreducible and hence there exists an attracting lamination Λ^+ supported by $[F]$.

Let \mathcal{K} be the nonattracting subgroup system associated to Λ^+ . Now apply the main result of [19] with \mathcal{K} and ψ . \square

Theorem 5.5 Let ϕ_i be partially fully irreducible on F^i , and $[F^i] \neq [F^j]$, $i, j \in \{1, \dots, k\}$. Suppose that $[F^i]$ is distance ≥ 2 from every free-factor component of $\mathcal{A}_{\text{na}}(\Lambda_j^\pm)$ for all $i = 1, \dots, k$ and $i \neq j$; if such free factor components exist. Then there exists $M > 0$ such that for every $m_i \geq M$, $Q = \langle \phi_1^{m_1}, \dots, \phi_k^{m_k} \rangle$ is a free group of rank k .

Moreover, if $\mathcal{A}_{\text{na}}(\Lambda_1^\pm) = \mathcal{A}_{\text{na}}(\Lambda_2^\pm) = \dots = \mathcal{A}_{\text{na}}(\Lambda_k^\pm)$, then for any lift \widehat{Q} of Q , $\mathbb{F} \rtimes \widehat{Q}$ is relatively hyperbolic group.

Proof Lemma 5.2 tells us that Λ_i^\pm and Λ_j^\pm do not have any asymptotic leaves. This implies that we can choose long generic-leaf segments of Λ_i^+ and Λ_i^- and construct attracting and repelling neighborhoods V_i^+ and V_i^- for each $i = 1, \dots, k$ so that $V_i^+, V_i^-, V_j^+, V_j^-$ are pairwise disjoint for $i \neq j$.

Since generic leaves of Λ_i^+ are not carried by $\mathcal{A}_{\text{na}}(\Lambda_j^\pm)$ for $i \neq j$, by Proposition 5.3 generic leaves of Λ_i^+ are weakly attracted to Λ_j^+ (respectively, to Λ_j^-) under the action of ϕ_j (respectively, of ϕ_j^{-1}).

The conclusion about Q being free group now follows directly using [6, Lemma 3.4.2, page 551]. For the relative hyperbolicity of $\mathbb{F} \rtimes \widehat{Q}$, let $\mathcal{K} = \mathcal{A}_{\text{na}}(\Lambda_1^\pm)$ and apply of Theorem 2.7. □

The above theorem remains true if we relax the requirement of $\mathcal{A}_{\text{na}}(\Lambda_1^\pm) = \mathcal{A}_{\text{na}}(\Lambda_2^\pm) = \dots = \mathcal{A}_{\text{na}}(\Lambda_k^\pm)$ and replace it by the requirement of Q having an admissible subgroup system.

5.4 Relatively hyperbolic extensions using relative fully irreducibles

Given a collection of free factors F^1, F^2, \dots, F^p of \mathbb{F} such that $\mathbb{F} = F^1 * F^2 * \dots * F^p * B$ with B possibly trivial, we say that the collection forms a *free factor system*, written as $\mathcal{F} := \{[F^1], [F^2], \dots, [F^p]\}$ and we say that a conjugacy class $[c]$ of a word $c \in \mathbb{F}$ is carried by \mathcal{F} if there exists some $1 \leq s \leq p$ and a representative H^s of F^s such that $c \in H^s$.

Definition 5.6 We say that an automorphism $\phi \in \text{Out}(\mathbb{F})$ is *fully irreducible relative to a free factor system* \mathcal{F} if \mathcal{F} is ϕ -invariant and there is no proper ϕ -invariant free factor system \mathcal{F}' such that $\mathcal{F} \sqsubset \mathcal{F}'$.

Relative fully irreducibles were developed by Handel and Mosher [23] to find a better analog of Ivanov’s theorem on subgroups of mapping class groups of a surface so that the analogous theorem for subgroups of $\text{Out}(\mathbb{F})$ works “inductively” on free factors that are fixed and that the behavior of the automorphism in between free factors is better understood. Below we will describe how subgroups made of relative fully irreducibles give us easy ways to construct relatively hyperbolic extensions, under some mild restrictions on the generators.

Let $\mathcal{F} \sqsubset \mathcal{F}'$ be free factor systems, where \mathcal{F} is a proper free factor system. If there exists a marked graph G and realizations $H \subset H' \subset G$ of these free factor systems such that $H' \setminus H$ is a single edge, then \mathcal{F}' is said to be a one-edge extension of \mathcal{F} . If no such realization exist, then \mathcal{F}' is said to be a *multi-edge extension* of \mathcal{F} .

We record the following lemma which follows from various nontrivial results in [23] and use the conclusion as our working definition of relative fully irreducible outer automorphism.

Lemma 5.7 *Suppose $\mathcal{F} \sqsubset \{\mathbb{F}\}$ is a multiedge extension invariant under ϕ and every component of \mathcal{F} is ϕ -invariant. If ϕ is fully irreducible rel \mathcal{F} then there exists ϕ -invariant dual lamination pair Λ_ϕ^\pm such that the following hold:*

- (1) $\mathcal{F}_{\text{supp}}(\mathcal{F}, \Lambda^\pm) = [\mathbb{F}]$.
- (2) If Λ_ϕ^\pm is nongeometric then $\mathcal{A}_{\text{na}}(\Lambda_\phi^\pm) = \mathcal{F}$.
- (3) If Λ_ϕ^\pm is geometric then there exists a root free $\sigma \in \mathbb{F}$ such that $\mathcal{A}_{\text{na}}(\Lambda_\phi^\pm) = \mathcal{F} \cup \{[\sigma]\}$.

Conversely, there exists a ϕ -invariant dual lamination pair such that if (1) and (2) hold or if (1) and (3) hold then ϕ is fully irreducible rel \mathcal{F} .

Proof Let $f: G \rightarrow G$ be an improved relative train-track map representing ϕ and G_{r-1} be the filtration element realizing \mathcal{F} . Apply [23, Proposition 2.2] to get all the conclusions for some iterate ϕ^k of ϕ .

We claim that Λ_ϕ^\pm obtained by applying [23, Proposition 2.2] must be ϕ -invariant. Otherwise, by the definition of being fully irreducible relative to \mathcal{F} $\phi(\Lambda_\phi^+)$ will be an attracting lamination which is properly contained in G_{r-1} and hence is carried by \mathcal{F} which in turn is carried by the nonattracting subgroup system for Λ_ϕ^+ . This is a contradiction, hence Λ_ϕ^+ is ϕ -invariant. Similar arguments work for Λ_ϕ^- .

The converse part follows from the case analysis in the proof of [23, Theorem I, pages 18–19]. \square

Theorem 5.8 *Let ϕ, ψ be fully irreducible relative to the multiedge extension \mathcal{F} , with corresponding invariant lamination pairs $\Lambda_\phi^\pm, \Lambda_\psi^\pm$ (as in the equivalence Lemma 5.7). If no leaf of $\Lambda_\phi^+ \cup \Lambda_\phi^-$ is asymptotic to any leaf of $\Lambda_\psi^+ \cup \Lambda_\psi^-$, then there exists an integer $M \geq 1$ such that:*

- (1) $Q = \langle \phi^m, \psi^n \rangle$ is free group of rank 2 for all $m, n \geq M$.
- (2) If $\Lambda_\phi^\pm, \Lambda_\psi^\pm$ are both nongeometric then the extension group $\mathbb{F} \rtimes \widehat{Q}$ is hyperbolic relative to the finite collection of subgroups $\{F_s \rtimes \widehat{Q}_s\}_{s=1}^p$, where \widehat{Q}_s is a lift that preserves F_s .
- (3) If both Λ_ϕ^\pm and Λ_ψ^\pm are geometric laminations which come from the same surface, both fixing the conjugacy class $[\sigma]$ representing the surface boundary, then $\mathbb{F} \rtimes \widehat{Q}$ is hyperbolic relative to the finite collection of subgroups $\{[\sigma] \rtimes \widehat{Q}_\sigma\} \cup \{F_s \rtimes \widehat{Q}_s\}_{s=1}^p$, where \widehat{Q}_s is a lift that preserves F_s and \widehat{Q}_σ is a lift that fixes σ .

Proof To prove (1) we apply Theorem 2.7 with $\mathcal{K} = \mathcal{F}$ and to prove (2) we apply Theorem 2.7 with $\mathcal{K} = \mathcal{F} \cup \{[\sigma]\}$. \square

As a corollary of the theorem for $\mathcal{F} = \emptyset$, we recover the case for surface group with punctures, which was proved in [27, Theorem 4.9].

6 Further discussion: nonrelative hyperbolicity

From the results we have so far we gather a necessary condition of nonrelative hyperbolicity:

Proposition 6.1 *Let $\mathbb{F} \rtimes \widehat{Q}$ be NRH for every free group Q which is generated by nonzero powers of ϕ_i 's which are sufficiently different. Then, there exists $i \in \{1, \dots, k\}$ such that $\mathcal{K}_i^* \neq \emptyset$ and $\mathcal{K}_i^* \neq \mathcal{K}_j^*$ for some $j \neq i$.*

A sufficient condition for NRH remains to be unknown. However, we investigate the following example.

Example 6.2 Let $\mathbb{F} = \langle a, b, c, d, e \rangle$, $F^1 = \langle a, b, c \rangle$, $F^2 = \langle a, b, c, d \rangle$. Define ϕ_1 as the outer automorphism class of the map $\Phi_1: a \mapsto ac, b \mapsto a, c \mapsto b, d \mapsto dc, e \mapsto ec$ and ϕ_2 to be the outer automorphism class of the map $\Phi_2: a \mapsto ad, b \mapsto a, c \mapsto b, d \mapsto c, e \mapsto e$. Then $\mathcal{A}_{\text{na}}(\Lambda_1^\pm) = \emptyset$ and $\mathcal{A}_{\text{na}}(\Lambda_2^\pm) = \{\langle e \rangle\}$. In any relatively hyperbolic structure, the subgroup $\mathbb{Z} \oplus \mathbb{Z} \cong \langle e, \Phi_2 \rangle$ must be contained in some peripheral subgroup. A simple computation shows that $\mathbb{F} \leq \langle \Phi_1^{-1} \Phi_2, b, c \rangle$ and hence the extension group $1 \rightarrow \mathbb{F} \rightarrow E \rightarrow \langle \phi_1, \phi_2 \rangle \rightarrow 1$ is not relatively hyperbolic (NRH as in [1]).

The hypothesis for freeness in the above theorem is easily checked, and so there exists M such that for every $m, n \geq M$, the group $Q = \langle \phi_1^m, \phi_2^n \rangle$ is a free group of rank 2. Once we raise the automorphisms to sufficiently high powers, the abnormality with peripheral subgroups goes away. We ask the question: is $\mathbb{F} \rtimes \widehat{Q}$ hyperbolic relative to $\langle e, \Phi_2 \rangle$ when Q is generated by high enough powers of ϕ_1, ϕ_2 ?

Remark 6.3 Note that $[F^i]$ is a fixed vertex for the action of ϕ_i on the free-factor complex \mathcal{FF} , for $i = 1, 2$. Also the free-factors in the set $\mathcal{A}_{\text{na}}(\Lambda_i^\pm)$ are fixed vertices for the same action. If we translate the hypothesis of Theorem 5.5 in the framework of the free factor complex \mathcal{FF} of \mathbb{F} , then what we require is that $[F^1]$ and $[F^2]$ are distinct points in \mathcal{FF} and $\mathcal{A}_{\text{na}}(\Lambda_1^\pm) = \mathcal{A}_{\text{na}}(\Lambda_2^\pm)$. Using Proposition 5.3, the second hypothesis in turn implies that $[F^1]$ and the free-factor components of $\mathcal{A}_{\text{na}}(\Lambda_2^\pm)$ are close in \mathcal{FF} . Similarly free-factor components of $\mathcal{A}_{\text{na}}(\Lambda_1^\pm)$ and $[F^2]$ are also close in \mathcal{FF} . By partial fully irreducibility $[F^i]$ and free-factor components of $\mathcal{A}_{\text{na}}(\Lambda_i^\pm)$ are also close. Therefore the entire setup of Theorem 5.5 is in a small (bounded) region of \mathcal{FF} . The obvious question that comes to mind is: if we assume that $[F^1]$ and $[F^2]$ are far apart in Theorem 5.5 can we still expect relative hyperbolicity of $\mathbb{F} \rtimes \widehat{Q}$?

References

- [1] **J Behrstock, C Druţu, L Mosher**, *Thick metric spaces, relative hyperbolicity, and quasi-isometric rigidity*, Math. Ann. 344 (2009) 543–595 MR
- [2] **M Bestvina, M Feighn**, *A combination theorem for negatively curved groups*, J. Differential Geom. 35 (1992) 85–101 MR
- [3] **M Bestvina, M Feighn**, *Hyperbolicity of the complex of free factors*, Adv. Math. 256 (2014) 104–155 MR

- [4] **M Bestvina, M Feighn**, *Subfactor projections*, *J. Topol.* 7 (2014) 771–804 MR
- [5] **M Bestvina, M Feighn, M Handel**, *Laminations, trees, and irreducible automorphisms of free groups*, *Geom. Funct. Anal.* 7 (1997) 215–244 MR
- [6] **M Bestvina, M Feighn, M Handel**, *The Tits alternative for $\text{Out}(F_n)$, I: Dynamics of exponentially-growing automorphisms*, *Ann. of Math.* 151 (2000) 517–623 MR
- [7] **B H Bowditch**, *The Cannon–Thurston map for punctured-surface groups*, *Math. Z.* 255 (2007) 35–76 MR
- [8] **P Brinkmann**, *Hyperbolic automorphisms of free groups*, *Geom. Funct. Anal.* 10 (2000) 1071–1089 MR
- [9] **D Cooper**, *Automorphisms of free groups have finitely generated fixed point sets*, *J. Algebra* 111 (1987) 453–456 MR
- [10] **M Culler, K Vogtmann**, *Moduli of graphs and automorphisms of free groups*, *Invent. Math.* 84 (1986) 91–119 MR
- [11] **S Dowdall, I Kapovich, S J Taylor**, *Cannon–Thurston maps for hyperbolic free group extensions*, *Israel J. Math.* 216 (2016) 753–797 MR
- [12] **S Dowdall, S J Taylor**, *The co-surface graph and the geometry of hyperbolic free group extensions*, *J. Topol.* 10 (2017) 447–482 MR
- [13] **S Dowdall, S J Taylor**, *Hyperbolic extensions of free groups*, *Geom. Topol.* 22 (2018) 517–570 MR
- [14] **B Farb**, *Relatively hyperbolic groups*, *Geom. Funct. Anal.* 8 (1998) 810–840 MR
- [15] **B Farb, L Mosher**, *Convex cocompact subgroups of mapping class groups*, *Geom. Topol.* 6 (2002) 91–152 MR
- [16] **M Feighn, M Handel**, *The recognition theorem for $\text{Out}(F_n)$* , *Groups Geom. Dyn.* 5 (2011) 39–106 MR
- [17] **P Ghosh**, *Applications of weak attraction theory in $\text{Out}(\mathbb{F}_N)$* , *Geom. Dedicata* 181 (2016) 1–22 MR
- [18] **P Ghosh**, *Limits of conjugacy classes under iterates of hyperbolic elements of $\text{Out}(\mathbb{F})$* , *Groups Geom. Dyn.* 14 (2020) 177–211 MR
- [19] **P Ghosh**, *Relative hyperbolicity of free-by-cyclic extensions*, *Compos. Math.* 159 (2023) 153–183 MR
- [20] **P Ghosh, F Gültepe**, *Relative hyperbolicity of free extensions of free groups*, preprint (2023) arXiv 2307.09674 To appear in *Trans. Amer. Math. Soc.*
- [21] **U Hamenstädt, S Hensel**, *Stability in outer space*, *Groups Geom. Dyn.* 12 (2018) 359–398 MR
- [22] **U Hamenstaedt**, *Word hyperbolic extensions of surface groups*, preprint (2005) arXiv math/0505244
- [23] **M Handel, L Mosher**, *Subgroup decomposition in $\text{Out}(F_n)$* , *Mem. Amer. Math. Soc.* 1280, Amer. Math. Soc., Providence, RI (2020) MR
- [24] **A Hatcher, K Vogtmann**, *The complex of free factors of a free group*, *Quart. J. Math. Oxford Ser.* 49 (1998) 459–468 MR
- [25] **R P Kent, IV, C J Leininger**, *Shadows of mapping class groups: capturing convex cocompactness*, *Geom. Funct. Anal.* 18 (2008) 1270–1325 MR
- [26] **M Mitra**, *Ending laminations for hyperbolic group extensions*, *Geom. Funct. Anal.* 7 (1997) 379–402 MR
- [27] **M Mj, L Reeves**, *A combination theorem for strong relative hyperbolicity*, *Geom. Topol.* 12 (2008) 1777–1798 MR
- [28] **S J Taylor**, *A note on subfactor projections*, *Algebr. Geom. Topol.* 14 (2014) 805–821 MR

- [29] **S J Taylor**, *Right-angled Artin groups and $\text{Out}(\mathbb{F}_n)$, I: Quasi-isometric embeddings*, *Groups Geom. Dyn.* 9 (2015) 275–316 MR
- [30] **W P Thurston**, *Hyperbolic structures on 3-manifolds, II: Surface groups and 3-manifolds which fiber over the circle*, preprint (1986) arXiv math/9801045 Reprinted as pages 79–110 in his *Collected works with commentary, II*, Amer. Math. Soc., Providence, RI (2022)
- [31] **K Vogtmann**, *Automorphisms of free groups and outer space*, from “Proceedings of the Conference on Geometric and Combinatorial Group Theory, I” (L Mosher, M Sageev, editors), volume 94 (2002) 1–31 MR

*Department of Mathematics, Ashoka University
Sonipat, India*

*Department of Mathematics and Statistics, University of Toledo
Toledo, OH, United States*

pritam.ghosh@ashoka.edu.in, funda.gultepe@utoledo.edu

Received: 13 May 2024 Revised: 30 September 2024

ALGEBRAIC & GEOMETRIC TOPOLOGY

msp.org/agt

EDITORS

PRINCIPAL ACADEMIC EDITORS

John Etnyre
etnyre@math.gatech.edu
Georgia Institute of Technology

Kathryn Hess
kathryn.hess@epfl.ch
École Polytechnique Fédérale de Lausanne

BOARD OF EDITORS

Julie Bergner	University of Virginia jeb2md@eservices.virginia.edu	Thomas Koberda	University of Virginia thomas.koberda@virginia.edu
Steven Boyer	Université du Québec à Montréal cohf@math.rochester.edu	Markus Land	LMU München markus.land@math.lmu.de
Tara E Brendle	University of Glasgow tara.brendle@glasgow.ac.uk	Christine Lescop	Université Joseph Fourier lescop@ujf-grenoble.fr
Indira Chatterji	CNRS & Univ. Côte d'Azur (Nice) indira.chatterji@math.cnrs.fr	Norihiko Minami	Yamato University minami.norihiko@yamato-u.ac.jp
Octav Cornea	Université de Montreal cornea@dms.umontreal.ca	Andrés Navas	Universidad de Santiago de Chile andres.navas@usach.cl
Alexander Dranishnikov	University of Florida dranish@math.ufl.edu	Jessica S Purcell	Monash University jessica.purcell@monash.edu
Tobias Ekholm	Uppsala University, Sweden tobias.ekholm@math.uu.se	Birgit Richter	Universität Hamburg birgit.richter@uni-hamburg.de
Mario Eudave-Muñoz	Univ. Nacional Autónoma de México mario@matem.unam.mx	Jérôme Scherer	École Polytech. Féd. de Lausanne jerome.scherer@epfl.ch
David Futер	Temple University dfuter@temple.edu	Vesna Stojanoska	Univ. of Illinois at Urbana-Champaign vesna@illinois.edu
John Greenlees	University of Warwick john.greenlees@warwick.ac.uk	Zoltán Szabó	Princeton University szabo@math.princeton.edu
Matthew Hedden	Michigan State University mhedden@math.msu.edu	Maggy Tomova	University of Iowa maggy-tomova@uiowa.edu
Kristen Hendricks	Rutgers University kristen.hendricks@rutgers.edu	Daniel T Wise	McGill University, Canada daniel.wise@mcgill.ca
Hans-Werner Henn	Université Louis Pasteur henn@math.u-strasbg.fr	Lior Yanovski	Hebrew University of Jerusalem lior.yanovski@gmail.com
Daniel Isaksen	Wayne State University isaksen@math.wayne.edu		


See inside back cover or msp.org/agt for submission instructions.

The subscription price for 2025 is US \$760/year for the electronic version, and \$1110/year (+\$75, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Algebraic & Geometric Topology is indexed by Mathematical Reviews, Zentralblatt MATH, Current Mathematical Publications and the Science Citation Index.

Algebraic & Geometric Topology (ISSN 1472-2747 printed, 1472-2739 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004.

AGT peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<https://msp.org/>

© 2025 Mathematical Sciences Publishers

ALGEBRAIC & GEOMETRIC TOPOLOGY

Volume 25 Issue 7 (pages 3789–4436) 2025

Algebras for enriched ∞ -operads	3789
RUNE HAUGSENG	
An overtwisted convex hypersurface in higher dimensions	3813
RIVER CHIANG and KLAUS NIEDERKRÜGER-EID	
Presheaves of groupoids as models for homotopy types	3833
LÉONARD GUETTA	
Product and coproduct on fixed point Floer homology of positive Dehn twists	3875
YUAN YAO and ZIWEN ZHAO	
Multitwists in big mapping class groups	3921
GEORGE DOMAT, FEDERICA FANONI and SEBASTIAN HENSEL	
Cofibrantly generated model structures for functor calculus	3931
LAUREN BANDKLAYDER, JULIA E BERGNER, RHIANNON GRIFFITHS, BRENDA JOHNSON and REKHA SANTHANAM	
Endomorphisms of Artin groups of type D	3975
FABRICE CASTEL and LUIS PARIS	
Linear bounds of the crosscap number of knots	4009
ROB MCCONKEY	
Atiyah–Segal completion for the Hermitian K-theory of symplectic groups	4037
JENS HORNBOSTEL, HERMAN ROHRBACH and MARCUS ZIBROWIUS	
A minimality property for knots without Khovanov 2-torsion	4073
ONKAR SINGH GUJRAL and JOSHUA WANG	
Local indicability of groups with homology circle presentations	4077
AGUSTÍN NICOLÁS BARRETO and ELÍAS GABRIEL MINIAN	
Cubulating drilled bundles over graphs	4095
MAHAN MJ and BISWAJIT NAG	
Infinitely many homeomorphic hyperbolic plugs with the same basic sets	4147
FANGFANG CHEN	
One-point compactifications of configuration spaces and the self duality of the little disks operad	4163
CONNOR MALIN	
Braided multitwists	4185
RODRIGO DE POOL	
Positive intermediate Ricci curvature on connected sums	4209
PHILIPP REISER and DAVID J WRAITH	
The \mathcal{S}_n -equivariant Euler characteristic of the moduli space of graphs	4229
MICHAEL BORINSKY and JOS VERMASEREN	
Classification of metric fibrations	4257
YASUHIKO ASAO	
Spaces over BO are thickened manifolds	4287
HIRO LEE TANAKA	
Representation-graded Bredon homology of elementary abelian 2-groups	4321
MARKUS HAUSMANN and STEFAN SCHWEDE	
The \mathbb{Z}/p -equivariant cohomology of the genus-zero Deligne–Mumford space with $1 + p$ marked points	4341
DAIN KIM and NICHOLAS WILKINS	
Geometry of free extensions of free groups via automorphisms with fixed points on the complex of free factors	4357
PRITAM GHOSH and FUNDA GÜLTEPE	
Coarse cohomology of configuration space and coarse embedding	4391
ARKA BANERJEE	
Correction to the article An algebraic model for finite loop spaces	4427
CARLES BROTO, RAN LEVI and BOB OLIVER	