

AG
T

*Algebraic & Geometric
Topology*

Volume 25 (2025)

**The structure of relatively hyperbolic groups
in convex real projective geometry**

MITUL ISLAM
ANDREW ZIMMER



The structure of relatively hyperbolic groups in convex real projective geometry

MITUL ISLAM
ANDREW ZIMMER

We prove a general structure theorem for relatively hyperbolic groups (with arbitrary peripheral subgroups) acting naive convex cocompactly on properly convex domains in real projective space. We also establish a characterization of such groups in terms of the existence of an invariant collection of closed unbounded convex subsets with good isolation properties. This is a real projective analogue of results of Hindawi–Hruska–Kleiner for CAT(0) spaces. We also obtain an equivariant homeomorphism between the Bowditch boundary of the group and a quotient of the ideal boundary.

20F65, 20F67, 22E40, 57N16, 57S20

1 Introduction

Let \mathbb{H}^d denote real hyperbolic d -space and recall that a discrete subgroup $\Gamma \leq \text{Isom}(\mathbb{H}^d)$ is called *convex cocompact* if there exists a nonempty Γ -invariant geodesically convex closed subset $\mathcal{C} \subset \mathbb{H}^d$ where the quotient $\Gamma \backslash \mathcal{C}$ is compact. The Beltrami–Klein model realizes \mathbb{H}^d as a properly convex domain $\mathbb{B}^d \subset \mathbb{P}(\mathbb{R}^{d+1})$ (namely the Euclidean unit ball in a standard affine chart) in such a way that the isometry group $\text{Isom}(\mathbb{H}^d)$ coincides with the subgroup of $\text{PGL}_{d+1}(\mathbb{R})$ which preserves \mathbb{B}^d (namely $\text{PO}(d, 1)$ up to conjugation). Further, in this model, a subset being geodesically convex is equivalent to being convex in some (hence any) affine chart that contains \mathbb{B}^d .

The Beltrami–Klein model perspective allows one to naturally generalize the classical notion of convex cocompact groups. In particular, one can consider a general properly convex domain $\Omega \subset \mathbb{P}(\mathbb{R}^d)$ and the group $\text{Aut}(\Omega) \leq \text{PGL}_d(\mathbb{R})$ of projective automorphisms which preserve Ω . Then a discrete subgroup $\Gamma \leq \text{Aut}(\Omega)$ is called *naive convex cocompact* if there exists a nonempty Γ -invariant closed convex subset $\mathcal{C} \subset \Omega$ where the quotient $\Gamma \backslash \mathcal{C}$ is compact. In this case, we say that $(\Omega, \mathcal{C}, \Gamma)$ is a *naive convex cocompact triple*.

When the group Γ is word hyperbolic (eg in the classical real hyperbolic setting), there is a close connection between this notion of naive convex cocompact groups and Anosov representations/higher Teichmüller theory; see [7; 27]. As one moves beyond the word hyperbolic case, the structure of these discrete groups becomes more mysterious.

In this paper, we study naive convex cocompact groups which are (intrinsically) relatively hyperbolic groups. This is a rich class with many examples, for instance where:

- (a) Γ is a projective reflection group generated by reflection along faces of a projective Coxeter polytope à la Vinberg and Γ is irreducible as a Coxeter group; see [8; 19].
- (b) Γ is isomorphic to $\pi_1(M)$ where M is a closed three-manifold such that each geometric component in its JSJ decomposition supports a hyperbolic structure; see [1; 17].

In both of these cases, the groups are relatively hyperbolic with respect to peripheral subgroups which are virtually abelian of rank at least two. However, there also exist examples where Γ is relatively hyperbolic with respect to non-virtually abelian subgroups; see the discussion in [23, Section 2.6.3]. The primary goal of this paper is to describe the structure of such examples. One of our main results is proving that the relative hyperbolicity of Γ is equivalent to the existence of a so-called *peripheral family*, a Γ -invariant collection of closed convex subsets with good isolation properties.

This investigation extends some recent work. Previously in [15], we considered the special case where the peripheral subgroups were virtually abelian of rank at least two. For such groups, we proved that relative hyperbolicity is equivalent to the existence of a collection of properly embedded simplices with good isolation properties. This geometric description is analogous to the case of CAT(0) spaces with isolated flats [13]. We also showed that the boundary of these simplices are, in a technical sense, the only places where the boundary is irregular; see [15, Theorem 1.19(6), Theorem 1.8(7) and (8)]. Shortly after, Weisman [23] considered convex cocompact groups (a more restrictive class than naive convex cocompact) who were relatively hyperbolic and whose peripheral subgroups were also convex cocompact, but not necessarily virtually abelian. For such groups he established a similar result about the irregular boundary points and also showed that the Bowditch boundary could be realized as a quotient of the boundary.

There are a number of other results in the literature concerning relatively hyperbolic groups acting on properly convex domains; see for instance [3; 4; 5; 6]. These results consider the case when $\Gamma \backslash \mathcal{C}$ is noncompact and characterize when Γ is relatively hyperbolic with respect to the fundamental groups of the ends (under some geometric assumptions on the ends and \mathcal{C}). There is some similarity between the statements in this paper and the statements in [3; 4; 5], but to the best of our knowledge, there is no actual overlap between the results.

We extend the results in [15; 23] to the case of general peripheral subgroups and the case of general naive convex cocompact groups. The proofs build upon ideas from both papers.

We will now introduce the notation required to precisely state our main results. Given a properly convex domain $\Omega \subset \mathbb{P}(\mathbb{R}^d)$, we will let d_Ω denote the Hilbert metric, which is a natural and classical $\text{Aut}(\Omega)$ -invariant, proper, and complete metric on Ω ; see Section 2.2 below. This allows us to speak of the diameter $\text{diam}_\Omega(A)$ and r -open neighborhood $\mathcal{N}_\Omega(A, r) \subset \Omega$ of a subset $A \subset \Omega$ relative to the Hilbert metric.

Definition 1.1 Suppose that $\Omega \subset \mathbb{P}(\mathbb{R}^d)$ is a properly convex domain, $\mathcal{C} \subset \Omega$ is a closed¹ convex subset, $\Gamma \leq \text{Aut}(\Omega)$, and \mathcal{X} is a collection of closed unbounded convex subsets of \mathcal{C} . Then we say that:

- (1) \mathcal{X} is Γ -invariant if $\Gamma \cdot \mathcal{X} = \mathcal{X}$.
- (2) \mathcal{X} is *strongly isolated* if for every $r > 0$ there exists $D_1(r) > 0$ such that if $X_1, X_2 \in \mathcal{X}$ are distinct, then

$$\text{diam}_\Omega(\mathcal{N}_\Omega(X_1, r) \cap \mathcal{N}_\Omega(X_2, r)) \leq D_1(r).$$

- (3) \mathcal{X} *coarsely contains the properly embedded simplices of \mathcal{C}* if there exists $D_2 > 0$ such that if $S \subset \mathcal{C}$ is a properly embedded simplex of dimension at least two, then there exists $X \in \mathcal{X}$ with $S \subset \mathcal{N}_\Omega(X, D_2)$.

When $(\Omega, \mathcal{C}, \Gamma)$ is a naive convex cocompact triple and \mathcal{X} satisfies all three of the above conditions, we say that \mathcal{X} is a *peripheral family of $(\Omega, \mathcal{C}, \Gamma)$* .

Given a closed convex subset $\mathcal{C} \subset \Omega$ of a properly convex domain, the *ideal boundary* of \mathcal{C} is $\partial_i \mathcal{C} := \bar{\mathcal{C}} \cap \partial \Omega$ (see Section 2.3). Also, given $x \in \bar{\Omega}$ we will let $F_\Omega(x)$ denote the open face of x in $\bar{\Omega}$ and given $A \subset \bar{\Omega}$ we will let $F_\Omega(A) = \bigcup_{x \in A} F_\Omega(x)$. Using these boundary faces and a peripheral family, one can define a natural quotient of the ideal boundary:

Definition 1.2 Suppose that \mathcal{X} is a peripheral family of $(\Omega, \mathcal{C}, \Gamma)$. Let

$$[\partial_i \mathcal{C}]_{\mathcal{X}} := \partial_i \mathcal{C} / \sim$$

be the topological quotient where $x \sim y$ if and only if

- (1) $x, y \in F_\Omega(\partial_i X)$ for some $X \in \mathcal{X}$, or
- (2) $F_\Omega(y) = F_\Omega(x)$.

Remark 1.3 We remark that similar boundary quotients are considered by Choi [5] and Weisman [23].

We need one more piece of notation to state our main result: the *limit set* of a subgroup $G \leq \text{Aut}(\Omega)$ is

$$\mathcal{L}_\Omega(G) := \partial \Omega \cap \bigcup_{p \in \Omega} \overline{G \cdot p}.$$

This definition ensures that $\mathcal{L}_\Omega(G)$ is base point independent—unlike real hyperbolic geometry, the accumulation points of a single orbit in Ω may depend on the base point.

Our main result—Theorem 1.4 below—is essentially an equivalence statement for a naive convex cocompact triple $(\Omega, \mathcal{C}, \Gamma)$: the relative hyperbolicity of the group Γ is equivalent to the existence of a peripheral family \mathcal{X} . Theorem 1.4(1) shows that the group-theoretic property of relative hyperbolicity implies the geometric property of admitting a peripheral family. Theorem 1.4(2) shows the converse.

¹We make it our convention that, “ $Y \subset \Omega$ is a closed subset” always means that Y is closed in the subspace topology inherited from Ω .

Theorem 1.4 Suppose that $(\Omega, \mathcal{C}, \Gamma)$ is a naive convex cocompact triple. Then the following hold:

- (1) If Γ is relatively hyperbolic with respect to $\mathcal{P} = \{P_1, \dots, P_N\}$ and X_j is the closed convex hull of $\mathcal{L}_\Omega(P_j) \cap \partial_i \mathcal{C}$ in Ω , then

$$\mathcal{X} := \Gamma \cdot \{X_1, \dots, X_N\}$$

is a peripheral family of $(\Omega, \mathcal{C}, \Gamma)$.

- (2) If \mathcal{X} is a peripheral family of $(\Omega, \mathcal{C}, \Gamma)$ and $\mathcal{P} := \{P_1, \dots, P_N\}$ is a set of representatives of the Γ -conjugacy classes in $\{\text{Stab}_\Gamma(X) : X \in \mathcal{X}\}$, then Γ is relatively hyperbolic with respect to \mathcal{P} .

Moreover, when either of the above conditions are satisfied, then:

- (a) (\mathcal{C}, d_Ω) is relatively hyperbolic with respect to $\mathcal{X} = \Gamma \cdot \{X_1, \dots, X_N\}$.
 (b) There is a Γ -equivariant homeomorphism between the Bowditch boundary $\partial(\Gamma, \mathcal{P})$ and $[\partial_i \mathcal{C}]_{\mathcal{X}}$.
 (c) Each (Ω, X_j, P_j) is a naive convex cocompact triple.
 (d) There exists $L > 0$ such that if $x \in \partial_i \mathcal{C}$ and $\text{diam}_{F_\Omega(x)}(F_\Omega(x) \cap \partial_i \mathcal{C}) \geq L$, then $x \in F_\Omega(\partial_i X)$ for some $X \in \mathcal{X}$.
 (e) There exists $R > 0$ such that if $X \in \mathcal{X}$ and $x \in \partial_i X$, then

$$d_{F_\Omega(x)}^{\text{Haus}}(F_\Omega(x) \cap \partial_i X, F_\Omega(x) \cap \partial_i \mathcal{C}) \leq R.$$

Remark 1.5 Properties (d) and (e) are somewhat technical. Informally, property (d) states that any boundary face of Ω that $\partial_i \mathcal{C}$ intersects in a “large set” must intersect the ideal boundary of an element in \mathcal{X} . Property (e) informally states that if $X \in \mathcal{X}$, then $\partial_i X$ coarsely contains any boundary face of $\partial_i \mathcal{C}$ that it intersects.

Theorem 1.4(1) and (2) can be viewed as a real projective analogue of the results of Hruska–Kleiner and Hindawi–Hruska–Kleiner [13] in the setting of CAT(0)-geometry. This earlier work motivated the results in this paper, but the methods of proof are very different. In particular, we should also note that an old result of Kelly–Straus [18] says that a Hilbert geometry (Ω, d_Ω) is CAT(0) if and only if it is isometric to real hyperbolic $(d-1)$ -space (in which case Ω coincides, up to a change of coordinates, with the Beltrami–Klein model of real hyperbolic $(d-1)$ -space).

As mentioned above, in previous work [15] we proved a version of Theorem 1.4 in the special case when \mathcal{X} consisted of properly embedded simplices of dimension at least two and the subgroups in \mathcal{P} were virtually abelian of rank at least two. In this case, using the simple structure of simplices and abelian subgroups in naive convex cocompact groups (see [14]), one can weaken the strongly isolated assumption to only assuming that \mathcal{X} is closed and discrete. In fact, a substantial portion of this earlier work was building a strongly isolated collection of properly embedded simplices from a closed and discrete collection. It seems unlikely to us that such a weakening is possible in the general case.

1.1 Convex cocompact groups

As mentioned above, the class of naive convex cocompact groups includes the more restrictive, but still interesting, class of convex cocompact groups. For this class of groups, Theorem 1.4 can be restated in a much simpler way.

Given a properly convex domain $\Omega \subset \mathbb{P}(\mathbb{R}^d)$ and a discrete subgroup $\Gamma \leq \text{Aut}(\Omega)$, let $\mathcal{C}_\Omega(\Gamma) \subset \Omega$ denote the closed convex hull of $\mathcal{L}_\Omega(\Gamma)$ in Ω .

Definition 1.6 A discrete subgroup $\Gamma \leq \text{Aut}(\Omega)$ is called *convex cocompact* if $\mathcal{C}_\Omega(\Gamma)$ is nonempty and the quotient $\Gamma \backslash \mathcal{C}_\Omega(\Gamma)$ is compact.

Every convex cocompact subgroup is clearly naive convex cocompact, but the converse is not true; see [15, Section 2.3] for some examples. One key difference between the two definitions is that if $\Gamma \leq \text{Aut}(\Omega)$ is convex cocompact, then any open boundary face of Ω which intersects $\partial_i \mathcal{C}_\Omega(\Gamma)$ is actually contained in $\partial_i \mathcal{C}_\Omega(\Gamma)$. On the other hand, if $(\Omega, \mathcal{C}, \Gamma)$ is a naive convex cocompact triple, then it is possible for $\partial_i \mathcal{C}$ to intersect a boundary face of Ω in a small set. This seemingly small difference makes convex cocompact groups much easier to study.

For convex cocompact groups, Theorem 1.4(a)–(e) can be restated and expanded as follows.

Theorem 1.7 (see Section 10) *Suppose that $\Omega \subset \mathbb{P}(\mathbb{R}^d)$ is a properly convex domain and $\Gamma \leq \text{Aut}(\Omega)$ is a convex cocompact subgroup. If Γ is relatively hyperbolic with respect to $\mathcal{P} = \{P_1, \dots, P_N\}$, then:*

(a) $(\mathcal{C}_\Omega(\Gamma), d_\Omega)$ is relatively hyperbolic with respect to

$$\mathcal{X} := \Gamma \cdot \{\mathcal{C}_\Omega(P_1), \dots, \mathcal{C}_\Omega(P_N)\}.$$

(b) Let $[\partial_i \mathcal{C}_\Omega(\Gamma)]_{\mathcal{P}}$ denote the quotient of $\partial_i \mathcal{C}_\Omega(\Gamma)$ obtained by collapsing each limit set $\mathcal{L}_\Omega(\gamma P_j \gamma^{-1})$ to a point (where $\gamma \in \Gamma$ and $P_j \in \mathcal{P}$). Then there is a Γ -equivariant homeomorphism between the Bowditch boundary $\partial(\Gamma, \mathcal{P})$ and $[\partial_i \mathcal{C}_\Omega(\Gamma)]_{\mathcal{P}}$.

(c) Each P_j is a convex cocompact subgroup of $\text{Aut}(\Omega)$.

(d) If $x \in \partial_i \mathcal{C}_\Omega(\Gamma)$ is not a \mathcal{C}^1 -smooth point of Ω (ie Ω does not have a unique supporting hyperplane at x), then $x \in \mathcal{L}_\Omega(\gamma P_j \gamma^{-1})$ for some $\gamma \in \Gamma$ and $P_j \in \mathcal{P}$.

(e) If $\ell \subset \partial_i \mathcal{C}_\Omega(\Gamma)$ is a nontrivial line segment, then $\ell \subset \mathcal{L}_\Omega(\gamma P_j \gamma^{-1})$ for some $\gamma \in \Gamma$ and $P_j \in \mathcal{P}$.

We should note that the main new content of Theorem 1.7 is part (c). In particular:

- (1) Part (a) is an immediate consequence of part (c).
- (2) In [15] we previously proved parts (d) and (e) in the case where each P_j is virtually abelian with rank at least two, and once part (c) is known, the same argument works in the more general setting.
- (3) Weisman [23] established parts (b) and (e) with part (c) as an assumption.

In the context of convex cocompact groups, peripheral families can also be defined in terms of their boundary behavior.

Proposition 1.8 (see Section 10) *Suppose that $\Omega \subset \mathbb{P}(\mathbb{R}^d)$ is a properly convex domain and $\Gamma \leq \text{Aut}(\Omega)$ is a convex cocompact subgroup. If \mathcal{X} is a Γ -invariant collection of closed unbounded convex subsets of Ω , then the following are equivalent:*

- (1) \mathcal{X} is a peripheral family of $(\Omega, \mathcal{C}_\Omega(\Gamma), \Gamma)$.
- (2) \mathcal{X} has the following properties:
 - (a) \mathcal{X} is closed in the local Hausdorff topology induced by the Hilbert metric d_Ω .
 - (b) If $X_1, X_2 \in \mathcal{X}$ are distinct, then $\partial_i X_1 \cap \partial_i X_2 = \emptyset$.
 - (c) If $\ell \subset \partial_i \mathcal{C}_\Omega(\Gamma)$ is a nontrivial line segment, then $\ell \subset \partial_i X$ for some $X \in \mathcal{X}$.

Moreover, when the above conditions are satisfied, then

$$(1-1) \quad [\partial_i \mathcal{C}_\Omega(\Gamma)]_{\mathcal{P}} = [\partial_i \mathcal{C}_\Omega(\Gamma)]_{\mathcal{X}},$$

where \mathcal{P} is a set of representatives of the Γ -conjugacy classes in $\{\text{Stab}_\Gamma(X) : X \in \mathcal{X}\}$.

Previously, Weisman [23, Theorem 1.16] characterized when a convex cocompact subgroup is hyperbolic relative to a collection of convex cocompact subgroups in terms of the behavior of the limit sets of the subgroups. We note that combining Proposition 1.8 and Theorem 1.4 provides a “subset” version of this characterization.

1.2 The case of Gromov hyperbolic groups

A word hyperbolic group is relatively hyperbolic with respect to the empty set. Thus if $(\Omega, \mathcal{C}, \Gamma)$ is a naive convex cocompact triple and Γ is a word hyperbolic group, then Theorem 1.4 holds with $\mathcal{X} = \emptyset$.

However, the choice of this peripheral family \mathcal{X} is not canonical even for a convex cocompact group Γ (unlike our work in [15] where \mathcal{X} was the set of all maximal properly embedded simplices in $\mathcal{C}_\Omega(\Gamma)$ of dimension at least two). For instance, suppose that $\Gamma := \langle a, b \rangle$ is a convex cocompact subgroup of $\text{PSO}(2, 1)$ that is isomorphic to a free group on two generators. Then Γ is relatively hyperbolic with respect to $\mathcal{P} := \{g\}$ where $g := aba^{-1}b^{-1}$. Then $\mathcal{X} := \Gamma \cdot (g^+, g^-)$ where (g^+, g^-) is the unique g -invariant projective line in the Beltrami–Klein model of \mathbb{H}^2 . On the other hand, we can also choose $\mathcal{X} = \emptyset$ (when $\mathcal{P} = \emptyset$).

1.3 Outline of the paper and proofs

Sections 2 and 3 are expository. In Section 2 we recall the basic definitions and results about properly convex domains that we will require and in Section 3 we recall some properties of relatively hyperbolic spaces and groups.

Sections 4 through 9 are devoted to the proof of Theorem 1.4. The difficult part of Theorem 1.4 is part (2). Our general strategy is based on combining ideas from [15] and [23]. In particular, as in [23], we will use Yaman’s characterization of relative hyperbolicity to show that Γ is relatively hyperbolic. We will verify the conditions in Yaman’s theorem by further developing the ideas used in [15] to study closed and discrete collections of properly embedded simplices.

We should also note that Choi [5] (which predates the work of Weisman) used Yaman’s characterization to verify that the fundamental groups of certain noncompact convex real projective manifolds were relatively hyperbolic with respect to the fundamental groups of their ends.

The arguments used to verify the conditions of Yaman’s theorem in this paper are somewhat similar to the analogous ones in [5; 23]. But given the different setups, there does not seem to be an easy way to reduce the proofs in this paper to any lemmas in these two works.

The content of these sections is as follows:

- (1) In Section 4 we prove a quantitative result which informally states that long line segments in an open boundary face (relative to the Hilbert metric of the face) imply the existence of nearby properly embedded simplices.
- (2) In Section 5 we establish a number of useful properties of peripheral families.
- (3) In Section 6 we prove that peripheral subgroups of a relatively hyperbolic naive convex cocompact subgroup are themselves naive convex cocompact.
- (4) In Section 7 we prove a technical result which will allow us to show that nonconical limit points are bounded parabolic points. A key tool here is a “closest subset” projection map.
- (5) In Section 8 we prove some basic properties of the quotient space $[\partial_i \mathcal{C}]_{\mathcal{X}}$, including a sufficient condition for a point to be a conical limit point.
- (6) In Section 9 we put everything together and prove Theorem 1.4.

Finally, in Section 10, we use Theorem 1.4 to prove Theorem 1.7.

Acknowledgements

MIslam thanks Louisiana State University for hospitality during a visit in March 2020 where work on this project started.

MIslam was partially supported by Emmy Noether Project 427903332 (funded by the DFG), DMS-1607260 (National Science Foundation), and Rackham Predoctoral Fellowship (University of Michigan). A Zimmer was partially supported by grants DMS-2105580 and DMS-2104381 from the National Science Foundation.

The authors thank the reviewers for their thorough reading, comments, and suggestions.

2 Preliminaries

2.1 Notation

If $V \subset \mathbb{R}^d$ is a nonzero linear subspace, we will let $\mathbb{P}(V) \subset \mathbb{P}(\mathbb{R}^d)$ denote its projectivization. In most other cases, we will use $[o]$ to denote the projective equivalence class of an object o , for instance:

- (1) If $v \in \mathbb{R}^d \setminus \{0\}$, then $[v]$ denotes the image of v in $\mathbb{P}(\mathbb{R}^d)$.
- (2) If $\phi \in \text{GL}_d(\mathbb{R})$, then $[\phi]$ denotes the image of ϕ in $\text{PGL}_d(\mathbb{R})$.
- (3) If $T \in \text{End}(\mathbb{R}^d) \setminus \{0\}$, then $[T]$ denotes the image of T in $\mathbb{P}(\text{End}(\mathbb{R}^d))$.

We also standardize some metric notations. If (X, d) is a metric space, $p \in X$, $A \subset X$, and $r > 0$, then:

- (1) $\mathcal{N}_X(A, r) := \{x \in X : d(x, A) < r\}$.
- (2) $\mathcal{B}_X(p, r) := \{x \in X : d(x, p) < r\}$.
- (3) $\text{diam}_X(A) := \sup\{d(x, x') : x, x' \in A\}$.

2.2 Convexity and the Hilbert metric

A subset $C \subset \mathbb{P}(\mathbb{R}^d)$ is

- (1) *properly convex* if there exists an affine chart \mathbb{A} of $\mathbb{P}(\mathbb{R}^d)$ where $C \subset \mathbb{A}$ is a bounded convex subset;
- (2) a *properly convex domain* if C is properly convex and open in $\mathbb{P}(\mathbb{R}^d)$.

Given a properly convex set $C \subset \mathbb{P}(\mathbb{R}^d)$ and a subset $X \subset \bar{C}$, we define its *convex hull* as

$$\text{ConvHull}_C(X) := \bigcap \{Y : Y \text{ is a closed convex subset such that } X \subset Y \subset \bar{C}\}.$$

A *line segment* in $\mathbb{P}(\mathbb{R}^d)$ is a connected subset of a projective line. Given two points $x, y \in \mathbb{P}(\mathbb{R}^d)$ there is no canonical line segment with endpoints x and y , but we will use the following convention: if $C \subset \mathbb{P}(\mathbb{R}^d)$ is a properly convex set and $x, y \in \bar{C}$, then (when the context is clear) we will let $[x, y]$ denote the closed line segment joining x to y which is contained in \bar{C} . In this case, we will also let $(x, y) = [x, y] \setminus \{x, y\}$, $[x, y) = [x, y] \setminus \{y\}$, and $(x, y] = [x, y] \setminus \{x\}$.

Let $\Omega \subset \mathbb{P}(\mathbb{R}^d)$ be a properly convex domain. If $x, y \in \Omega$ are distinct, let $[a, b] := \mathbb{P}(\text{Span}\{x, y\}) \cap \bar{\Omega}$ where a and b are labeled such that $x \in [a, y]$ (ie the points are ordered a, x, y, b along $[a, b]$). Then the *Hilbert distance* between x and y is defined to be

$$d_\Omega(x, y) := \frac{1}{2} \log[a, x, y, b],$$

where

$$[a, x, y, b] := \frac{|x - b| |y - a|}{|x - a| |y - b|}$$

is the cross-ratio (here $|\cdot|$ is some (any) norm in some (any) affine chart which contains a, x, y, b). Then (Ω, d_Ω) is a complete geodesic metric space and $\text{Aut}(\Omega)$ acts properly and by isometries on Ω ; see for instance [2, Section 28]. Further, the projective line segment $[x, y]$ is a geodesic for the Hilbert distance.

2.3 Boundaries and faces

Suppose that $C \subset \mathbb{P}(\mathbb{R}^d)$ is a properly convex set. The *relative interior* of C , denoted by $\text{rel-int}(C)$, is the interior of C in $\mathbb{P}(\text{Span } C)$. We will say that C is *open in its span* if $C = \text{rel-int}(C)$.

The boundary of C is $\partial C := \bar{C} \setminus \text{rel-int}(C)$, the ideal boundary of C is $\partial_i C := \partial C \setminus C$, and the nonideal boundary of C is $\partial_n C := \partial C \cap C$.

Definition 2.1 If $C \subset \mathbb{P}(\mathbb{R}^d)$ is a properly convex set which is open in its span and $x \in \bar{C}$, let $F_C(x)$ denote the open face of x :

$$F_C(x) = \{x\} \cup \{y \in \bar{C} : \text{there exists an open line segment in } \bar{C} \text{ containing } x \text{ and } y\}.$$

We have the following observation about faces; see [16, Appendix] for a proof.

Observation 2.2 Suppose that $\Omega \subset \mathbb{P}(\mathbb{R}^d)$ is a properly convex domain.

- (1) $F_\Omega(x) = \Omega$ when $x \in \Omega$.
- (2) $F_\Omega(x)$ is open in its span.
- (3) $y \in F_\Omega(x)$ if and only if $x \in F_\Omega(y)$ if and only if $F_\Omega(x) = F_\Omega(y)$.
- (4) If $x, y \in \bar{\Omega}$, $z \in (x, y)$, $p \in F_\Omega(x)$, and $q \in F_\Omega(y)$, then

$$(p, q) \subset F_\Omega(z).$$

In particular, $(p, q) \subset \Omega$ if and only if $(x, y) \subset \Omega$.

If $B \subset C \subset \mathbb{P}(\mathbb{R}^d)$ are properly convex sets, then we say that B is properly embedded in C if $B \hookrightarrow C$ is a proper map with respect to the subspace topology. Note that B is properly embedded in C if and only if $\partial_i B \subset \partial_i C$.

2.4 Limits of automorphisms

Every $T \in \mathbb{P}(\text{End}(\mathbb{R}^d))$ induces a map

$$\mathbb{P}(\mathbb{R}^d) \setminus \mathbb{P}(\ker T) \rightarrow \mathbb{P}(\mathbb{R}^d)$$

defined by $x \mapsto T(x)$. We will frequently use the following observation.

Observation 2.3 If $\{T_n\}$ is a sequence in $\mathbb{P}(\text{End}(\mathbb{R}^d))$ converging to $T \in \mathbb{P}(\text{End}(\mathbb{R}^d))$, then

$$T(x) = \lim_{n \rightarrow \infty} T_n(x)$$

for all $x \in \mathbb{P}(\mathbb{R}^d) \setminus \mathbb{P}(\ker T)$. The convergence is uniform on compact subsets of $\mathbb{P}(\mathbb{R}^d) \setminus \mathbb{P}(\ker T)$.

We can view $\mathbb{P}(\text{End}(\mathbb{R}^d))$ as a compactification of $\text{PGL}_d(\mathbb{R})$ and then consider limits of automorphisms in this compactification.

Proposition 2.4 [14, Proposition 5.6] Suppose that $\Omega \subset \mathbb{P}(\mathbb{R}^d)$ is a properly convex domain, $p_0 \in \Omega$, and $\{g_n\}$ is a sequence in $\text{Aut}(\Omega)$ such that:

- (1) $g_n(p_0) \rightarrow x \in \partial\Omega$.
- (2) $g_n^{-1}(p_0) \rightarrow y \in \partial\Omega$.
- (3) $g_n \rightarrow T$ in $\mathbb{P}(\text{End}(\mathbb{R}^d))$.

Then image $T \subset \text{Span } F_\Omega(x)$, $\mathbb{P}(\ker T) \cap \Omega = \emptyset$, and $y \in \mathbb{P}(\ker T)$.

In the case of “nontangential” convergence we can say more.

Proposition 2.5 [14, Proposition 5.7] *Suppose that $\Omega \subset \mathbb{P}(\mathbb{R}^d)$ is a properly convex domain, $p_0 \in \Omega$, $x \in \partial\Omega$, $\{p_n\}$ is a sequence in $[p_0, x)$ converging to x , and $\{g_n\}$ is a sequence in $\text{Aut}(\Omega)$ such that*

$$\sup_{n \geq 1} d_\Omega(g_n(p_0), p_n) < +\infty.$$

If $g_n \rightarrow T \in \mathbb{P}(\text{End}(\mathbb{R}^d))$, then $T(\Omega) = F_\Omega(x)$.

Proposition 5.7 in [14] is stated differently and a proof of the statement above can be found in [28, Proposition 2.13].

2.5 Projective simplices

A subset $S \subset \mathbb{P}(\mathbb{R}^d)$ is called a k -dimensional simplex in $\mathbb{P}(\mathbb{R}^d)$ if there exists $g \in \text{PGL}_d(\mathbb{R})$ such that

$$gS = \{[x_1 : \cdots : x_{k+1} : 0 : \cdots : 0] \in \mathbb{P}(\mathbb{R}^d) : x_1, \dots, x_{k+1} > 0\}.$$

In this case, we call the $k + 1$ points

$$g^{-1}[1 : 0 : \cdots : 0], g^{-1}[0 : 1 : 0 : \cdots : 0], \dots, g^{-1}[0 : \cdots : 0 : 1 : 0 : \cdots : 0] \in \partial S$$

the vertices of S .

The Hilbert metric on a simplex can be explicitly computed (see [11; 20, Proposition 1.7; 22]) and from this explicit form one obtains the following.

Proposition 2.6 *If $\Omega \subset \mathbb{P}(\mathbb{R}^d)$ is a properly convex domain and $S \subset \Omega$ is a properly embedded simplex, then (S, d_Ω) is quasi-isometric to $\mathbb{R}^{\dim S}$.*

Remark 2.7 By the definition of the Hilbert metric, $d_S = d_\Omega|_{S \times S}$. So the quasi-isometry constants only depend on $\dim S$.

2.6 The Hausdorff distance and local Hausdorff topology

When (X, d) is a metric space, the Hausdorff distance between two subsets $A, B \subset X$ is defined by

$$d_X^{\text{Haus}}(A, B) = \max\left\{\sup_{a \in A} \inf_{b \in B} d(a, b), \sup_{b \in B} \inf_{a \in A} d(a, b)\right\}.$$

When (X, d) is a complete metric space, d_X^{Haus} is a complete metric on the set of nonempty compact subsets of X .

The local Hausdorff topology is a natural topology on the set of nonempty closed sets in X . For a nonempty closed set C_0 , a base point $x_0 \in X$, and $r_0, \epsilon_0 > 0$, define the set $U(C_0, x_0, r_0, \epsilon_0)$ to consist of all closed subsets $C \subset X$ where

$$d_X^{\text{Haus}}(C_0 \cap \mathcal{B}_X(x_0, r_0), C \cap \mathcal{B}_X(x_0, r_0)) < \epsilon_0.$$

The *local Hausdorff topology* on the set of nonempty closed subsets of X is the topology generated by the sets $U(\cdot, \cdot, \cdot, \cdot)$. This is the same topology as the one called the *local Hausdorff convergence topology* in [16, Section 2.5].

2.7 Distance estimates for the Hilbert metric

The asymptotic behavior of the Hilbert distance connects naturally with the structure of open faces in the boundary.

Proposition 2.8 *Suppose that $\Omega \subset \mathbb{P}(\mathbb{R}^d)$ is a properly convex domain and $\{p_n\}, \{q_n\}$ are sequences in Ω where $p_n \rightarrow p \in \bar{\Omega}$ and $q_n \rightarrow q \in \bar{\Omega}$. If*

$$\liminf_{n \rightarrow \infty} d_{\Omega}(p_n, q_n) < +\infty,$$

then $F_{\Omega}(p) = F_{\Omega}(q)$ and

$$d_{F_{\Omega}(p)}(p, q) \leq \liminf_{n \rightarrow \infty} d_{\Omega}(p_n, q_n).$$

Proof This is a well-known property of the Hilbert metric, but we provide the proof for the readers' convenience.

Passing to a subsequence we can suppose that

$$\liminf_{n \rightarrow \infty} d_{\Omega}(p_n, q_n) = \lim_{n \rightarrow \infty} d_{\Omega}(p_n, q_n).$$

Since d_{Ω} is proper, if at least one of p, q is in Ω , then both must be in Ω . Further, if $p, q \in \Omega$, then $F_{\Omega}(p) = F_{\Omega}(q) = \Omega$ and the continuity of the Hilbert metric implies that

$$d_{F_{\Omega}(p)}(p, q) = d_{\Omega}(p, q) = \lim_{n \rightarrow \infty} d_{\Omega}(p_n, q_n).$$

Hence we can assume that $p, q \in \partial\Omega$. It suffices to consider $p \neq q$ since otherwise $d_{F_{\Omega}(p)}(p, q) = 0$.

For each $n \geq 1$, let $a_n, b_n \in \partial\Omega$ be the points where $p_n, q_n \in (a_n, b_n)$ and labeled so that the points are in the order a_n, p_n, q_n, b_n along (a_n, b_n) . Passing to a subsequence, we can assume that $a_n \rightarrow a$ and $b_n \rightarrow b$ in $\partial\Omega$. Then $p, q \in [a, b]$. Since $p \neq q$ and

$$\lim_{n \rightarrow \infty} \frac{1}{2} \log[a_n, p_n, q_n, b_n] = \lim_{n \rightarrow \infty} d_{\Omega}(p_n, q_n) < +\infty,$$

the definition of the cross-ratio implies that a, p, q, b are all distinct. Hence $p, q \in (a, b)$ and so $F_{\Omega}(p) = F_{\Omega}(q)$. Finally, since $(a, b) \subset F_{\Omega}(p)$, the monotonicity of the Hilbert metric implies that

$$d_{F_{\Omega}(p)}(p, q) \leq d_{(a,b)}(p, q) = \liminf_{n \rightarrow \infty} d_{\Omega}(p_n, q_n). \quad \square$$

We will frequently use the following fact about the Hausdorff distance between line segments.

Proposition 2.9 [14, Proposition 5.3] *Suppose that $\Omega \subset \mathbb{P}(\mathbb{R}^d)$ is a properly convex domain. If $p_1, p_2, q_1, q_2 \in \bar{\Omega}$ are such that $(p_1, p_2) \subset \Omega$, $F_\Omega(p_1) = F_\Omega(q_1)$, and $F_\Omega(p_2) = F_\Omega(q_2)$, then*

$$d_\Omega^{\text{Haus}}([p_1, p_2] \cap \Omega, [q_1, q_2] \cap \Omega) \leq \max\{d_{F_\Omega(p_1)}(p_1, q_1), d_{F_\Omega(p_2)}(p_2, q_2)\}.$$

Using induction, Proposition 2.9 can be upgraded as follows.

Proposition 2.10 *Suppose that $\Omega \subset \mathbb{P}(\mathbb{R}^d)$ is a properly convex domain, $q_1, \dots, q_m \in \bar{\Omega}$, and*

$$z \in \text{rel-int}(\text{ConvHull}_\Omega(q_1, \dots, q_m)).$$

If $p_1, \dots, p_m \in \bar{\Omega}$ and $F_\Omega(p_j) = F_\Omega(q_j)$ for all $1 \leq j \leq m$, then

$$d_F^{\text{Haus}}(\text{ConvHull}_\Omega(q_1, \dots, q_m) \cap F, \text{ConvHull}_\Omega(p_1, \dots, p_m) \cap F) \leq \max_{1 \leq j \leq m} d_{F_\Omega(q_j)}(q_j, p_j),$$

where $F := F_\Omega(z)$.

Proof For $1 \leq k \leq m$, let

$$S_k := \text{rel-int}(\text{ConvHull}_\Omega(q_1, \dots, q_k)) \quad \text{and} \quad S'_k := \text{rel-int}(\text{ConvHull}_\Omega(p_1, \dots, p_k)).$$

Since S_k, S'_k are open in their span, there exist faces F_k, F'_k of Ω such that $S_k \subset F_k$ and $S'_k \subset F'_k$.

For each $1 \leq k \leq m$, we claim that $F_k = F'_k$ and

$$d_{F_k}^{\text{Haus}}(S_k, S'_k) \leq \max_{1 \leq j \leq k} d_{F_\Omega(q_j)}(q_j, p_j).$$

We induct on k . The base case $k = 1$ is by definition.

Fix $k > 1$. Then fix $w \in S_k$. We will show that $F_k = F'_k$ and

$$(2-1) \quad d_{F_k}(w, S'_k) \leq \max_{1 \leq j \leq k} d_{F_\Omega(q_j)}(q_j, p_j).$$

If $w = q_k$, then we must have $q_k \in S_{k-1}$. Then $q_k \in F_{k-1}$, which implies that $p_k \in F_{k-1}$. So $S_k \cup S'_k \subset F_{k-1}$ and $F_k = F_{k-1} = F'_k$ by the induction hypothesis. Further,

$$d_{F_k}(w, S'_k) \leq d_{F_k}(q_k, p_k) = d_{F_\Omega(q_k)}(q_k, p_k).$$

If $w \neq q_k$, then there exists $q' \in S_{k-1}$ such that $w \in (q', q_k)$. Indeed, if $q_k \in \bar{S}_{k-1}$, then $S_k = S_{k-1}$ and then this is obvious as we can extend $(q_k, w]$ beyond $w \in S_k$. Otherwise, if $q_k \notin \bar{S}_{k-1}$, then every point in \bar{S}_k is a nontrivial convex combination of a point $q'' \in \bar{S}_{k-1}$ and q_k . Among all such points in \bar{S}_k , $S_k = \text{rel-int}(\bar{S}_k)$ is precisely the set of points for which $q'' \in S_{k-1}$. In particular, as $w \in S_k$, there exists $q' \in S_{k-1}$ such that $w \in (q_k, q')$.

Since $q' \in S_{k-1}$, by induction there exists $p' \in S'_{k-1}$ with

$$d_{F_{k-1}}(q', p') \leq \max_{1 \leq j \leq k-1} d_{F_\Omega(q_j)}(q_j, p_j).$$

There are two cases to consider depending on whether p' equals p_k or not.

Case 1 Assume $p' \neq p_k$. Then Observation 2.2(4) implies that $(q', q_k) \cup (p', p_k) \subset F_k$. Since $(p', p_k) \subset S_k$, we then have $F_k = F'_k$. Further, by Proposition 2.9,

$$d_{F_k}(w, S'_k) \leq d_{F_k}^{\text{Haus}}((q', q_k), (p', p_k)) \leq \max_{1 \leq j \leq k} d_{F_{\Omega}(q_j)}(q_j, p_j).$$

Case 2 Assume $p' = p_k$. Then $p_k = p' \in S'_{k-1} \subset F_{k-1}$, which implies that $q_k \in F_{k-1}$. Therefore $S_k \cup S'_k \subset F_{k-1}$ and $F_k = F_{k-1} = F'_k$ by the induction hypothesis. Further, by Proposition 2.9,

$$d_{F_k}(w, S'_k) \leq d_{F_k}([q', q_k], [p_k, p_k]) \leq \max(d_{F_k}(q', p_k), d_{F_k}(q_k, p_k)) \leq \max_{1 \leq j \leq k} d_{F_{\Omega}(q_j)}(q_j, p_j).$$

Thus in all cases $F_k = F'_k$ and (2-1) holds. Since $w \in S_k$ was arbitrary, we see that

$$\sup_{w \in S_k} d_{F_k}(w, S'_k) \leq \max_{1 \leq j \leq k} d_{F_{\Omega}(q_j)}(q_j, p_j).$$

Repeating the same argument with $w' \in S'_k$ shows that

$$\sup_{w' \in S'_k} d_{F_k}(w', S_k) \leq \max_{1 \leq j \leq k} d_{F_{\Omega}(q_j)}(q_j, p_j).$$

Hence,

$$d_{F_k}^{\text{Haus}}(S_k, S'_k) \leq \max_{1 \leq j \leq k} d_{F_{\Omega}(q_j)}(q_j, p_j).$$

This completes the proof of the claim.

Now, to prove the proposition, we apply the claim when $k = m$. Since $z \in S_m$, we have $F = F_{\Omega}(z) = F_m$. By the claim, $S_m \cup S'_m \subset F_m$ and

$$d_{F_m}^{\text{Haus}}(S_m, S'_m) \leq \max_{1 \leq j \leq m} d_{F_{\Omega}(q_j)}(q_j, p_j).$$

Finally, S_m (resp. S'_m) is dense in $\text{ConvHull}_{\Omega}(q_1, \dots, q_m) \cap F$ (resp. $\text{ConvHull}_{\Omega}(p_1, \dots, p_m) \cap F$). So the result follows. \square

3 Relatively hyperbolic spaces and groups

In this expository section we recall some basic properties of relatively hyperbolic groups and spaces. We define relative hyperbolic spaces and groups in terms of Druţu and Sapir’s tree-graded spaces; see [10, Definition 2.1].

Definition 3.1 (1) A complete geodesic metric space (X, d) is said to be *relatively hyperbolic with respect to a collection of subsets* \mathcal{S} if all its asymptotic cones, with respect to a fixed nonprincipal ultrafilter, are tree-graded with respect to the collection of ultralimits of the elements of \mathcal{S} .

(2) A finitely generated group G is said to be *relatively hyperbolic with respect to a family of subgroups* $\{H_1, \dots, H_N\}$ if the Cayley graph of G with respect to some (hence any) finite set of generators is relatively hyperbolic with respect to the collection of left cosets $\{gH_i : g \in G, i = 1, \dots, N\}$.

Remark 3.2 These are one among several equivalent definitions of relatively hyperbolic spaces/groups; see [10] for more details.

Recall that if (X, d) is a metric space, $A \subset X$, and $r > 0$, then

$$\mathcal{N}_X(A, r) := \{x \in X : d(x, A) < r\}.$$

We will frequently use the following properties of relatively hyperbolic spaces. These results are taken from Druţu–Sapir [10].

Theorem 3.3 *Suppose that (X, d_X) is relatively hyperbolic with respect to \mathcal{S} .*

- (1) [10, Corollary 5.8] *For any $A \geq 1$ and $B \geq 0$ there exists $M = M(A, B)$ such that if $k \geq 2$ and $f: \mathbb{R}^k \rightarrow X$ is an (A, B) -quasi-isometric embedding, then there exists some $S \in \mathcal{S}$ such that*

$$f(\mathbb{R}^k) \subset \mathcal{N}_X(S, M).$$

- (2) [10, Lemma 4.15] *For any $A \geq 1$ and $B \geq 0$ there exists $t = t(A, B)$ such that if $S \in \mathcal{S}$, $r \geq 1$, and $\gamma: [0, T] \rightarrow X$ is an (A, B) -quasigeodesic with $\gamma(0), \gamma(T) \in \mathcal{N}_X(S, r)$, then*

$$\gamma([0, T]) \subset \mathcal{N}_X(S, tr).$$

- (3) [10, Theorem 4.1] *For any $r > 0$ there exists $D = D(r) > 0$ such that if $S_1, S_2 \in \mathcal{S}$ are distinct, then*

$$\text{diam}_X(\mathcal{N}_X(S_1, r) \cap \mathcal{N}_X(S_2, r)) \leq D.$$

3.1 Immediate consequences in convex projective geometry

As an immediate consequence of the general theory of relatively hyperbolic spaces, we have the following.

Proposition 3.4 *Suppose that $(\Omega, \mathcal{C}, \Gamma)$ is a naive convex cocompact triple and \mathcal{X} is a Γ -invariant collection of closed unbounded convex subsets of \mathcal{C} . If (\mathcal{C}, d_Ω) is relatively hyperbolic with respect to \mathcal{X} , then \mathcal{X} is a peripheral family of $(\Omega, \mathcal{C}, \Gamma)$ (ie Γ -invariant, strongly isolated, and coarsely contains the properly embedded simplices of \mathcal{C}).*

Proof Theorem 3.3(3) implies that \mathcal{X} is strongly isolated. Proposition 2.6 and Theorem 3.3(1) imply that \mathcal{X} coarsely contains the properly embedded simplices of \mathcal{C} . \square

3.2 Yaman’s characterization

We recall Yaman’s characterization of relatively hyperbolic groups in terms of convergence group actions [21; 26].

Recall that when M is a nonempty compact Hausdorff metrizable space, a subgroup $G \leq \text{Homeo}(M)$ is called a *convergence group* if for every sequence $\{g_n\}$ of distinct elements in G there exists a subsequence $\{g_{n_j}\}$ and points $x, y \in M$ such that $g_{n_j}|_{M - \{y\}}$ converges locally uniformly to the constant map x . For such a subgroup:

- (1) The *limit set* $\mathcal{L}(G) \subset M$ is the set of points $x \in M$ where there exist $y \in M$ and a sequence $\{g_n\}$ in G where $g_n|_{M-\{y\}}$ converges locally uniformly to the constant map x .
- (2) A point $x \in \mathcal{L}(G)$ is a *conical limit point* if there exist distinct points $a, b \in M$ and a sequence of elements $\{g_n\}$ in G where $\lim_{n \rightarrow \infty} g_n(x) = a$ and $\lim_{n \rightarrow \infty} g_n(y) = b$ for all $y \in M - \{x\}$.
- (3) A point $x \in \mathcal{L}(G)$ is a *parabolic point* if $\text{Stab}_G(x)$ is infinite and nonloxodromic (ie x is the unique fixed point in $\mathcal{L}(G)$ for every $g \in \text{Stab}_G(x)$ of infinite order).
- (4) A parabolic point $x \in \mathcal{L}(G)$ is a *bounded parabolic point* if $\text{Stab}_G(x)$ acts cocompactly on $\mathcal{L}(G) - \{x\}$.

Finally, we say that a convergence group $G \leq \text{Homeo}(M)$ is *geometrically finite* if $\mathcal{L}(G)$ is a nonempty perfect set (ie $\#\mathcal{L}(G) \geq 3$) and every point in $\mathcal{L}(G)$ is either a conical limit point or a bounded parabolic point.

Theorem 3.5 (Yaman [26, Theorem 0.1]) *Suppose that $G \leq \text{Homeo}(M)$ is a geometrically finite convergence group and $B \subset \mathcal{L}(G)$ is the set of bounded parabolic fixed points. If B has finitely many G -orbits and $\text{Stab}_G(b) := \{h \in G : h(b) = b\}$ is finitely generated for every $b \in B$, then:*

- (1) G is relatively hyperbolic with respect to $\mathcal{P} = \{\text{Stab}_G(p_1), \dots, \text{Stab}_G(p_N)\}$ where $B = \bigsqcup_{i=1}^N G(p_i)$.
- (2) The Bowditch boundary $\partial(G, \mathcal{P})$ is equivariantly homeomorphic to $\mathcal{L}(G)$.

If Γ is relatively hyperbolic with respect to \mathcal{P} , then Γ acts as a geometrically finite convergence group on the Bowditch boundary.

4 Finding properly embedded simplices

For convex cocompact groups, if there is a line segment in the ideal boundary of the convex hull, then there is a properly embedded simplex in the convex hull [7, Theorem 1.15]. This fails for naive convex cocompact subgroups; see [15, Section 2.3]. Instead we will show that if there exists a line segment in the ideal boundary which is sufficiently long (relative to the Hilbert metric of the face containing it), then there exists a properly embedded simplex.

In the following proposition, $d_{\mathbb{P}}$ denotes some distance on $\mathbb{P}(\mathbb{R}^d)$ induced by a Riemannian metric.

Proposition 4.1 *Suppose that $(\Omega, \mathcal{C}, \Gamma)$ is a naive convex cocompact triple and $q \in \mathcal{C}$. For any $r, \epsilon > 0$ there exists $L = L(q, r, \epsilon) \geq 0$ with the following property: if*

- (1) $a, b \in \partial_i \mathcal{C}$ are contained in a boundary face F of Ω with $d_F(a, b) \geq L$,
- (2) $m \in (a, b)$ is the midpoint of $[a, b]$ relative to d_F , and
- (3) $p \in \mathbb{P}(\text{Span}\{a, b, q\}) \cap \mathcal{C}$ is sufficiently close to m (in the metric $d_{\mathbb{P}}$),

then there exists a properly embedded simplex $S \subset \mathcal{C}$ of dimension at least two with

$$\mathbb{P}(\text{Span}\{a, b, q\}) \cap \mathcal{B}_{\Omega}(p, r) \subset \mathcal{N}_{\Omega}(S, \epsilon).$$

Proof Suppose for a contradiction that the proposition is false for some choice of $r, \epsilon > 0$. Then for every $n \in \mathbb{N}$ there exist $a_n, b_n \in \partial_i \mathcal{C}$ and a sequence $\{p_{n,j}\}_{j \geq 1}$ where:

- (1) a_n, b_n are contained in a boundary face F_n and $d_{F_n}(a_n, b_n) > n$.
- (2) If $m_n \in (a_n, b_n)$ is the midpoint of $[a_n, b_n]$ relative to d_{F_n} , then $m_n = \lim_{j \rightarrow \infty} p_{n,j}$.
- (3) $p_{n,j} \in \mathbb{P}(\text{Span}\{a_n, b_n, q\}) \cap \mathcal{C}$ for all $j \geq 1$.
- (4) $\mathbb{P}(\text{Span}\{a_n, b_n, q\}) \cap \mathcal{B}_\Omega(p_{n,j}, r)$ is not contained in the ϵ -neighborhood of any properly embedded simplex in \mathcal{C} of dimension at least two.

We first claim that for every n , there exists j_n such that

$$(4-1) \quad d_\Omega(p_{n,j_n}, (a_n, q] \cup [q, b_n)) \geq \frac{1}{2}n.$$

Fix n and suppose not. Then for every $j \in \mathbb{N}$ we can find $a_{n,j} \in (a_n, q]$ and $b_{n,j} \in (b_n, q]$ with

$$d_\Omega(p_{n,j}, \{a_{n,j}, b_{n,j}\}) \leq \frac{1}{2}n.$$

Since $\lim_{j \rightarrow \infty} p_{n,j} = m_n$ and d_Ω is proper, we must have $\lim_{j \rightarrow \infty} a_{n,j} = a_n$ and $\lim_{j \rightarrow \infty} b_{n,j} = b_n$. Further, by the definition of the Hilbert metric,

$$\frac{1}{2}n \geq \limsup_{j \rightarrow \infty} d_\Omega(p_{n,j}, \{a_{n,j}, b_{n,j}\}) \geq d_{F_n}(m_n, \{a_n, b_n\}) > \frac{1}{2}n.$$

So we have a contradiction and hence for each n such a j_n exists.

After possibly passing to a subsequence we can find a sequence $\{\gamma_n\}$ in Γ such that as $n \rightarrow \infty$,

$$\gamma_n(p_{n,j_n}) \rightarrow p_\infty \in \mathcal{C}, \quad \gamma_n(a_n) \rightarrow a_\infty \in \bar{\mathcal{C}}, \quad \gamma_n(b_n) \rightarrow b_\infty \in \bar{\mathcal{C}}, \quad \gamma_n(q) \rightarrow q_\infty \in \bar{\mathcal{C}}.$$

By construction, $[a_\infty, b_\infty] \subset \partial_i \mathcal{C}$, and by (4-1), we see that

$$[a_\infty, q_\infty] \cup [q_\infty, b_\infty] \subset \partial_i \mathcal{C}.$$

Hence $a_\infty, b_\infty, q_\infty$ are the vertices of a properly embedded two-dimensional simplex $S \subset \mathcal{C}$ which contains p_∞ . So for n large, we have

$$\mathbb{P}(\text{Span}\{\gamma_n(a_n), \gamma_n(b_n), \gamma_n(q)\}) \cap \mathcal{B}_\Omega(\gamma_n(p_{n,j_n}), r) \subset \mathcal{N}_\Omega(S, \epsilon),$$

which implies that

$$\mathbb{P}(\text{Span}\{a_n, b_n, q\}) \cap \mathcal{B}_\Omega(p_{n,j_n}, r) \subset \mathcal{N}_\Omega(\gamma_n^{-1}S, \epsilon).$$

Thus we have a contradiction. □

5 Properties of peripheral families

Theorem 5.1 *Suppose that $(\Omega, \mathcal{C}, \Gamma)$ is a naive convex cocompact triple and \mathcal{X} is a peripheral family of $(\Omega, \mathcal{C}, \Gamma)$ (ie \mathcal{X} is Γ -invariant, strongly isolated, and coarsely contains all properly embedded simplices in \mathcal{C}). Then:*

- (1) Γ has finitely many orbits in \mathcal{X} .
- (2) If $X \in \mathcal{X}$, then $\text{Stab}_\Gamma(X)$ acts cocompactly on X . In particular, $\text{Stab}_\Gamma(X)$ is finitely generated.

- (3) If $X_1, X_2 \in \mathcal{X}$ are distinct, then $F_\Omega(\partial_i X_1) \cap F_\Omega(\partial_i X_2) = \emptyset$.
- (4) There exists $L > 0$ such that if $x \in \partial_i \mathcal{C}$ and $\text{diam}_{F_\Omega(x)}(F_\Omega(x) \cap \partial_i \mathcal{C}) \geq L$, then $x \in F_\Omega(\partial_i X)$ for some $X \in \mathcal{X}$.
- (5) There exists $R > 0$ such that if $X \in \mathcal{X}$ and $x \in \partial_i X$, then

$$d_{F_\Omega(x)}^{\text{Haus}}(F_\Omega(x) \cap \partial_i \mathcal{C}, F_\Omega(x) \cap \partial_i X) \leq R.$$
- (6) If we have $X \in \mathcal{X}$, $x \in \partial_i X$, and $[x, y] \subset \partial_i \mathcal{C}$, then $[x, y] \subset F_\Omega(\partial_i X)$.
- (7) If $[x, y] \subset \partial_i \mathcal{C}$, then either $F_\Omega(x) = F_\Omega(y)$, or there exists $X \in \mathcal{X}$ such that $x, y \in F_\Omega(\partial_i X)$.
- (8) If $X \in \mathcal{X}$, then $\text{Stab}_\Gamma(X)$ acts cocompactly on

$$\text{ConvHull}_\Omega(F_\Omega(\partial_i X) \cap \partial_i \mathcal{C}) \cap \Omega$$

and this set is nonempty.

For the rest of the section, fix $\Omega, \mathcal{C}, \Gamma$, and \mathcal{X} satisfying the hypotheses of the theorem.

Since \mathcal{X} coarsely contains all properly embedded simplices in \mathcal{C} , there exists $D_2 > 0$ such that if $S \subset \mathcal{C}$ is a properly embedded simplex of dimension at least two, then there exists $X \in \mathcal{X}$ with

$$(5-1) \quad S \subset \mathcal{N}_\Omega(X, D_2).$$

Lemma 5.2 *If $K \subset \mathcal{C}$ is compact, then the set $\{X \in \mathcal{X} : X \cap K \neq \emptyset\}$ is finite.*

Proof Suppose not. Then there exist an infinite sequence of pairwise distinct elements $\{X_n\}$ in \mathcal{X} where $X_n \cap K \neq \emptyset$ for all n . As each X_n is unbounded, we can fix $k_n \in X_n \cap K$ and $x_n \in \partial_i X_n$. Passing to a subsequence we can suppose that $k_n \rightarrow k \in K$ and $x_n \rightarrow x \in \partial_i \mathcal{C}$. Then

$$\liminf_{n, \ell \rightarrow \infty} \text{diam}_\Omega(\mathcal{N}_\Omega(X_n, 1) \cap \mathcal{N}_\Omega(X_\ell, 1)) \geq \liminf_{n, \ell \rightarrow \infty} \text{diam}_\Omega(\mathcal{N}_\Omega([k_n, x_n], 1) \cap \mathcal{N}_\Omega([k_\ell, x_\ell], 1)) = \infty,$$

and so by the strong isolation property $X_n = X_\ell$ for sufficiently large n, ℓ . So we have a contradiction. \square

Lemma 5.3 (part (1)) *Γ has finitely many orbits in \mathcal{X} .*

Proof Since Γ acts cocompactly on \mathcal{C} , there exists a compact set $K \subset \mathcal{C}$ such that $\Gamma \cdot K = \mathcal{C}$. Then for each $X \in \mathcal{X}$ there exists $g \in \Gamma$ such that $K \cap gX \neq \emptyset$. So by Lemma 5.2, there exist X_1, \dots, X_m such that $\mathcal{X} = \bigsqcup_{i=1}^m \Gamma \cdot X_i$. \square

Lemma 5.4 (part (2)) *If $X \in \mathcal{X}$, then $\text{Stab}_\Gamma(X)$ acts cocompactly on X .*

Proof This argument is standard; see [12, Theorem 3.7; 13, Section 3.1; 25, Proposition 4.0.4].

Fix a compact set $K \subset \mathcal{C}$ such that $\Gamma \cdot K = \mathcal{C}$. Let

$$\mathcal{G} := \{g \in \Gamma : X \cap gK \neq \emptyset\}.$$

Then $X \subset \mathcal{G} \cdot K$. Also, by the previous lemma

$$\{g^{-1}X : g \in \mathcal{G}\} = \{h_1^{-1}X, \dots, h_k^{-1}X\}$$

for some $h_1, \dots, h_k \in \mathcal{G}$. Notice that $g^{-1}X = h_j^{-1}X$ if and only if $gh_j^{-1} \in \text{Stab}_\Gamma(X)$; hence

$$\mathcal{G} = \bigcup_{j=1}^k \text{Stab}_\Gamma(X)h_j.$$

Finally, if $\hat{K} := \bigcup_{j=1}^k h_j K$, then \hat{K} is compact and $X \subset \text{Stab}_\Gamma(X) \cdot \hat{K}$. So $X = \text{Stab}_\Gamma(X) \cdot (X \cap \hat{K})$. Since X is closed, $X \cap \hat{K}$ is compact and thus $\text{Stab}_\Gamma(X)$ acts cocompactly on X .

Now, since X is convex, the metric space (X, d_Ω) is geodesic. So by the fundamental lemma of geometric group theory, $\text{Stab}_\Gamma(X)$ is finitely generated. \square

Lemma 5.5 (part (3)) *If $X_1, X_2 \in \mathcal{X}$ are distinct, then $F_\Omega(\partial_i X_1) \cap F_\Omega(\partial_i X_2) = \emptyset$.*

Proof Suppose $X_1, X_2 \in \mathcal{X}$ and $x \in F_\Omega(\partial_i X_1) \cap F_\Omega(\partial_i X_2)$. There exists $x_1 \in \partial_i X_1$ and $x_2 \in \partial_i X_2$ with

$$x_1, x_2 \in F_\Omega(x).$$

Fix $q_1 \in X_1$ and $q_2 \in X_2$. Then Proposition 2.9 implies that

$$d_\Omega^{\text{Haus}}([q_1, x_1], [q_2, x_2]) \leq \max\{d_\Omega(q_1, q_2), d_{F_\Omega(x)}(x_1, x_2)\}.$$

So for $r > \max\{d_\Omega(q_1, q_2), d_{F_\Omega(x)}(x_1, x_2)\}$, we have

$$\text{diam}_\Omega(\mathcal{N}_\Omega(X_1, r) \cap \mathcal{N}_\Omega(X_2, r)) = \infty.$$

Since \mathcal{X} is strongly isolated, then $X_1 = X_2$. \square

Recall that D_2 is the constant in (5-1).

Lemma 5.6 *There exists $L > 0$ such that if*

- (1) $a, b \in \partial_i \mathcal{C}$ are in the same boundary face F of Ω ,
- (2) $d_F(a, b) \geq L$, and
- (3) m is the midpoint of $[a, b]$ with respect to d_F ,

then there exists $X \in \mathcal{X}$ and $x \in \partial_i X \cap F$ such that

$$d_F(m, x) \leq D_2 + 1.$$

Proof Since \mathcal{X} is strongly isolated, there exists $r > 0$ such that

$$(5-2) \quad \text{diam}_\Omega(\mathcal{N}_\Omega(X_1, D_2 + 1) \cap \mathcal{N}_\Omega(X_2, D_2 + 1)) < r$$

for all distinct pairs $X_1, X_2 \in \mathcal{X}$. Fix $q \in \mathcal{C}$ and let $L > 0$ satisfy Proposition 4.1 for $q \in \mathcal{C}$ and constants $\epsilon = 1$ and r as above. Now suppose

- (1) $a, b \in \partial_i \mathcal{C}$ are in the same boundary face F of Ω ,
- (2) $d_F(a, b) \geq L$, and
- (3) m is the midpoint of $[a, b]$ with respect to d_F .

By our choice of L , there exists $q' \in (m, q]$ such that if $p \in (m, q']$ then there exists some properly embedded simplex $S_p \subset \mathcal{C}$ of dimension at least two such that

$$\mathbb{P}(\text{Span}\{a, b, q\}) \cap \mathcal{B}_\Omega(p, r) \subset \mathcal{N}_\Omega(S_p, 1).$$

Then, by our choice of D_2 in (5-1), there exists $X_p \in \mathcal{X}$ such that

$$\mathbb{P}(\text{Span}\{a, b, q\}) \cap \mathcal{B}_\Omega(p, r) \subset \mathcal{N}_\Omega(X_p, D_2 + 1).$$

We claim that X_p does not depend on $p \in (m, q']$. To verify this it is enough to fix $p_1, p_2 \in (m, q']$ with $d_\Omega(p_1, p_2) < r$ and show that X_{p_1} and X_{p_2} coincide. For such $p_1, p_2 \in (m, q']$ we have

$$\mathbb{P}(\text{Span}\{a, b, q\}) \cap \mathcal{B}_\Omega(p_1, r) \cap \mathcal{B}_\Omega(p_2, r) \subset \mathcal{N}_\Omega(X_{p_1}, D_2 + 1) \cap \mathcal{N}_\Omega(X_{p_2}, D_2 + 1),$$

and

$$\text{diam}_\Omega(\mathbb{P}(\text{Span}\{a, b, q\}) \cap \mathcal{B}_\Omega(p_1, r) \cap \mathcal{B}_\Omega(p_2, r)) \geq r.$$

So $X_{p_1} = X_{p_2}$ by our choice of r ; see (5-2).

Now let $X := X_p$ for $p \in (m, q']$. Then

$$(m, q'] \subset \mathcal{N}_\Omega(X, D_2 + 1)$$

which implies, by Proposition 2.8, that there exists $x \in \partial_i X \cap F$ such that

$$d_F(m, x) \leq D_2 + 1. \quad \square$$

Lemma 5.7 (part (4)) *If $x \in \partial_i \mathcal{C}$ and*

$$\text{diam}_{F_\Omega(x)}(F_\Omega(x) \cap \partial_i \mathcal{C}) \geq L,$$

then $x \in F_\Omega(\partial_i X)$ for some $X \in \mathcal{X}$.

Proof This follows immediately from Lemma 5.6. □

Lemma 5.8 (part (5)) *If $X \in \mathcal{X}$ and $x \in \partial_i X$, then*

$$d_{F_\Omega(x)}^{\text{Haus}}(F_\Omega(x) \cap \partial_i \mathcal{C}, F_\Omega(x) \cap \partial_i X) \leq \max\{L, 2D_2 + 2\}.$$

Proof Fix $X \in \mathcal{X}$ and $x \in \partial_i X$. Since $F_\Omega(x) \cap \partial_i X \subset F_\Omega(x) \cap \partial_i \mathcal{C}$, it suffices to show that

$$\sup_{y \in F_\Omega(x) \cap \partial_i \mathcal{C}} d_{F_\Omega(x)}(y, F_\Omega(x) \cap \partial_i X) \leq \max\{L, 2D_2 + 2\}.$$

Fix $y \in F_\Omega(x) \cap \partial_i \mathcal{C}$ and suppose for a contradiction that

$$d_{F_\Omega(x)}(y, F_\Omega(x) \cap \partial_i X) > \max\{L, 2D_2 + 2\}.$$

By changing x we may suppose that

$$d_{F_\Omega(x)}(y, F_\Omega(x) \cap \partial_i X) = d_{F_\Omega(x)}(y, x).$$

Let m be the midpoint of $[x, y]$ with respect to $d_{F_\Omega(x)}$. Then by Lemma 5.6 there exist $X' \in \mathcal{X}$ and $x' \in \partial_i X'$ such that

$$d_{F_\Omega(x)}(m, x') \leq D_2 + 1.$$

However, then $x \in F_\Omega(\partial_i X) \cap F_\Omega(\partial_i X') \neq \emptyset$ and so by Lemma 5.5 we have $X = X'$. So

$$D_2 + 1 < \frac{1}{2}d_{F_\Omega(x)}(y, F_\Omega(x) \cap \partial_i X) = d_{F_\Omega(x)}(m, F_\Omega(x) \cap \partial_i X) \leq d_{F_\Omega(x)}(m, x') \leq D_2 + 1,$$

and we have a contradiction. □

Lemma 5.9 (part (7)) *If $[x, y] \subset \partial_i C$, then either $F_\Omega(x) = F_\Omega(y)$, or there exists $X \in \mathcal{X}$ such that $x, y \in F_\Omega(\partial_i X)$.*

Proof We suppose that $F_\Omega(x) \neq F_\Omega(y)$ and show that $x, y \in F_\Omega(\partial_i X)$ for some $X \in \mathcal{X}$. Fix $z \in (x, y)$. Since $F_\Omega(x) \neq F_\Omega(y)$,

$$\text{diam}_{F_\Omega(z)}((x, y)) = \infty.$$

So by Lemma 5.7 there exists $X \in \mathcal{X}$ with $z \in F_\Omega(\partial_i X)$. Then Lemma 5.8 implies that

$$(x, y) \subset \mathcal{N}_{F_\Omega(z)}(F_\Omega(z) \cap \partial_i X, \max\{L, 2D_2 + 2\}).$$

Finally Proposition 2.8 implies that $x, y \in F_\Omega(\partial_i X)$. □

Lemma 5.10 (part (6)) *If we have $X \in \mathcal{X}$, $x \in \partial_i X$, and $[x, y] \subset \partial_i C$, then $[x, y] \subset F_\Omega(\partial_i X)$.*

Proof This follows immediately from Lemmas 5.9 and 5.5. □

Observation 5.11 *If $X \in \mathcal{X}$, then $\text{ConvHull}_\Omega(\partial_i X) \cap \Omega$ is nonempty.*

Proof By assumption X is unbounded and so $\partial_i X \neq \emptyset$. Fix $w_2 \in \partial_i X$ and $w_1 \in X$. Also fix a sequence $\{w'_n\} \subset [w_1, w_2)$ such that $w'_n \rightarrow w_2$. As $\text{Stab}_\Gamma(X)$ acts cocompactly on X , for each $n \geq 1$ there exists $\gamma_n \in \text{Stab}_\Gamma(X)$ such that $\{\gamma_n w'_n\}$ lies in a compact subset of X . Up to passing to a subsequence, we can assume that $\gamma_n w'_n \rightarrow z' \in X$ and $\gamma_n w_i \rightarrow z_i \in \bar{X}$ for $i = 1, 2$. Then $z_2 \in \partial_i X$ as $\partial_i X$ is compact and $z_1 \in \partial_i X$ as $d_\Omega(z', z_1) = \lim_{n \rightarrow \infty} d_\Omega(\gamma_n w'_n, \gamma_n w_1) = \infty$. Then $(z_1, z_2) \subset \Omega$, as $z' \in (z_1, z_2) \cap \Omega$. □

Lemma 5.12 (part (8)) *If $X \in \mathcal{X}$, then $F_\Omega(\partial_i X) \cap \partial_i C$ is closed in $\mathbb{P}(\mathbb{R}^d)$ and $\text{Stab}_\Gamma(X)$ acts cocompactly on the nonempty set $\text{ConvHull}_\Omega(F_\Omega(\partial_i X) \cap \partial_i C) \cap \Omega$.*

Proof We first verify that $F_\Omega(\partial_i X) \cap \partial_i C$ is closed. Suppose $\{x_n\}$ is a sequence in $F_\Omega(\partial_i X) \cap \partial_i C$ converging to some $x \in \partial_i C$ (note $\partial_i C$ is closed). Then for each n there exists $x'_n \in \partial_i X$ with $x_n \in F_\Omega(x'_n)$. In particular, $[x_n, x'_n] \subset \partial_i C$. Passing to a subsequence we can suppose that $x'_n \rightarrow x'$. Then $x' \in \partial_i X$ and $[x, x'] \subset \partial_i C$. Then Lemma 5.10 implies that $x \in F_\Omega(\partial_i X)$. Hence $F_\Omega(\partial_i X) \cap \partial_i C$ is closed.

Then the set

$$\mathcal{C}_X := \text{ConvHull}_\Omega(F_\Omega(\partial_i X) \cap \partial_i C) \cap \Omega$$

is closed in \mathcal{C} . As $\text{ConvHull}_\Omega(\partial_i X) \cap \Omega \subset \mathcal{C}_X$, Observation 5.11 implies that \mathcal{C}_X is nonempty. As $\text{Stab}_\Gamma(X)$ acts cocompactly on X , it suffices to show that \mathcal{C}_X is contained in a bounded neighborhood of X . Fix $p \in \mathcal{C}_X$. Then there exist $p_1, \dots, p_\ell \in F_\Omega(\partial_i X) \cap \partial_i C$ such that $p \in \text{ConvHull}_\Omega(p_1, \dots, p_\ell)$. Then by Lemma 5.8 there exist $p'_1, \dots, p'_\ell \in \partial_i X$ such that

$$d_{F_\Omega(p_j)}(p_j, p'_j) \leq \max\{L, 2D_2 + 2\} \quad \text{for } j = 1, \dots, \ell.$$

Then by Proposition 2.10,

$$\begin{aligned} d_{\Omega}(p, X) &\leq d_{\Omega}^{\text{Haus}}(\text{ConvHull}_{\Omega}(p_1, \dots, p_{\ell}) \cap \Omega, \text{ConvHull}_{\Omega}(p'_1, \dots, p'_{\ell}) \cap \Omega) \\ &\leq \max_{1 \leq j \leq \ell} d_{F_{\Omega}(p_j)}(p_j, p'_j) \leq \max\{L, 2D_2 + 2\}. \end{aligned}$$

Since $p \in \mathcal{C}_X$ was arbitrary, $\mathcal{C}_X \subset \overline{\mathcal{N}_{\Omega}(X, \max\{L, 2D_2 + 2\})}$. □

6 Naive convex cocompactness of peripheral subgroups

Proposition 6.1 *Suppose that $(\Omega, \mathcal{C}, \Gamma)$ is a naive convex cocompact triple and Γ is relatively hyperbolic with respect to $\{P_1, \dots, P_N\}$. For $1 \leq j \leq N$, let*

$$X_j := \text{ConvHull}_{\Omega}(\mathcal{L}_{\Omega}(P_j) \cap \partial_i \mathcal{C}) \cap \Omega.$$

Then:

- (1) *Each (Ω, X_j, P_j) is a naive convex cocompact triple.*
- (2) *$(\mathcal{C}, d_{\Omega})$ is relatively hyperbolic with respect to $\mathcal{X} := \Gamma \cdot \{X_1, \dots, X_N\}$.*

The rest of the section is devoted to the proof of this proposition. Suppose $(\Omega, \mathcal{C}, \Gamma)$ is a naive convex cocompact triple and Γ is relatively hyperbolic with respect to $\{P_1, \dots, P_N\}$.

Fix a point $p_0 \in \mathcal{C}$. The fundamental lemma of geometric group theory implies that $(\mathcal{C}, d_{\Omega})$ is relatively hyperbolic with respect to $\Gamma \cdot \{P_1(p_0), \dots, P_N(p_0)\}$.

For each $1 \leq j \leq N$, let $\mathcal{L}_j \subset \partial_i \mathcal{C}$ denote the accumulation points of the orbit $P_j(p_0)$ and let

$$\hat{X}_j = \text{ConvHull}_{\Omega}(\mathcal{L}_j) \cap \Omega.$$

Lemma 6.2 *\hat{X}_j is nonempty and P_j acts cocompactly on \hat{X}_j .*

Proof For $n \geq 1$ let $Y_j^{(n)}$ be the points which are contained in the convex hull of at most n points in $P_j(p_0)$. By Carathéodory's convex hull theorem, $Y_j^{(n)} = Y_j^{(d)}$ for all $n \geq d$. Further, the closure of $Y_j^{(d)}$ in Ω contains \hat{X}_j .

By Theorem 3.3(2) there exists $t > 0$ such that if $r \geq 1$ and $p, q \in \mathcal{C} \cap \mathcal{N}_{\Omega}(P_j(p_0), r)$, then we have $[p, q] \subset \mathcal{N}_{\Omega}(P_j(p_0), tr)$.

We claim that

$$(6-1) \quad Y_j^{(n)} \subset \mathcal{N}_{\Omega}(P_j(p_0), (n-1)t)$$

for all $n \geq 2$. The base case follows by our choice of t . Suppose (6-1) holds for some n . If $p \in Y_j^{(n+1)}$, then there exist $p_1 \in P_j(p_0)$ and $p_2 \in Y_j^{(n)}$ such that $p \in [p_1, p_2]$. Then, by induction there exists $q \in P_j(p_0)$ with $d_{\Omega}(p_2, q) < (n-1)t$. Then

$$[p_1, q] \subset Y_j^{(2)} \subset \mathcal{N}_{\Omega}(P_j(p_0), t),$$

and by Proposition 2.9,

$$d_{\Omega}(p, [p_1, q]) \leq d_{\Omega}^{\text{Haus}}([p_1, p_2], [p_1, q]) \leq d_{\Omega}(p_2, q) < (n - 1)t.$$

So $p \in \mathcal{N}_{\Omega}(P_j(p_0), nt)$. Since $p \in Y_j^{(n+1)}$ was arbitrary, this completes the induction step. Thus (6-1) holds for all $n \geq 2$.

Then

$$\hat{X}_j \subset \overline{Y_j^{(d)}} \subset \overline{\mathcal{N}_{\Omega}(P_j(p_0), (d - 1)t)},$$

and so

$$\hat{X}_j = P_j \cdot \overline{(\mathcal{B}_{\Omega}(p_0, (d - 1)t) \cap \hat{X}_j)}.$$

Thus P_j acts cocompactly on \hat{X}_j .

We now show that $\hat{X}_j \neq \emptyset$. Since P_j is infinite and acts properly on Ω , there exists $\{g_k\}$ in P_j such that

$$\lim_{k \rightarrow \infty} d_{\Omega}(p_0, g_k(p_0)) = \infty.$$

Let m_k be the midpoint of $[p_0, g_k(p_0)]$ with respect to d_{Ω} . By (6-1),

$$m_k \in Y_j^{(2)} \subset \mathcal{N}_{\Omega}(P_j(p_0), t).$$

Thus there exists $\{h_k\}$ in P_j such that $h_k(m_k) \in \mathcal{B}_{\Omega}(p_0, t)$. Then, up to passing to a subsequence, we can assume that $h_k(m_k) \rightarrow m \in \Omega$ and $h_k[p_0, g_k(p_0)] \rightarrow [x, y]$. Then $x, y \in \mathcal{L}_j$ and so

$$m \in (x, y) \subset \hat{X}_j. \quad \square$$

The fundamental lemma of geometric group theory implies that $(\mathcal{C}, d_{\Omega})$ is relatively hyperbolic with respect to $\Gamma \cdot \{\hat{X}_1, \dots, \hat{X}_N\}$. In particular, by Proposition 3.4 and Theorem 5.1(5) there exists $R > 0$ such that if $1 \leq j \leq N$ and $x \in \partial_i \hat{X}_j$, then

$$(6-2) \quad d_{F_{\Omega}(x)}^{\text{Haus}}(F_{\Omega}(x) \cap \partial_i \mathcal{C}, F_{\Omega}(x) \cap \partial_i \hat{X}_j) \leq R.$$

Lemma 6.3 *If $1 \leq j \leq N$, then*

$$d_{F_{\Omega}(x)}^{\text{Haus}}(\hat{X}_j, X_j) \leq R.$$

Hence P_j acts cocompactly on X_j and $(\mathcal{C}, d_{\Omega})$ is relatively hyperbolic with respect to $\Gamma \cdot \{X_1, \dots, X_N\}$.

Proof First note that $\hat{X}_j \subset X_j$, as $\mathcal{L}_j \subset \mathcal{L}_{\Omega}(P_j)$. By the previous lemma, P_j acts cocompactly on \hat{X}_j . Thus it suffices to show that

$$X_j \subset \overline{\mathcal{N}_{\Omega}(\hat{X}_j, R)}.$$

Fix $q \in X_j$. Then there exist $q_1, \dots, q_{\ell} \in \mathcal{L}_{\Omega}(P_j) \cap \partial_i \mathcal{C}$ where

$$q \in \text{ConvHull}_{\Omega}(q_1, \dots, q_{\ell}).$$

For each $1 \leq i \leq \ell$, there exist $p_i \in \Omega$ and a sequence $\{\gamma_{i,n}\}_{n \geq 1}$ in P_j with $\gamma_{i,n}(p_i) \rightarrow q_i$. Passing to subsequences we can suppose that

$$\gamma_{i,n}(p_0) \rightarrow q'_i \in \mathcal{L}_j \subset \partial_i \hat{X}_j,$$

for all $1 \leq i \leq \ell$. By Proposition 2.8, $F_\Omega(q_i) = F_\Omega(q'_i)$. So by (6-2) there exist $q''_i \in \partial_i \widehat{X}_j$ such that $d_{F_\Omega(q_i)}(q_i, q''_i) \leq R$. Then by Proposition 2.10,

$$d_\Omega(q, \widehat{X}_j) \leq d_\Omega^{\text{Haus}}(\text{ConvHull}_\Omega(q_1, \dots, q_\ell) \cap \Omega, \text{ConvHull}_\Omega(q''_1, \dots, q''_\ell) \cap \Omega) \leq \max_{1 \leq i \leq \ell} d_{F_\Omega(q_i)}(q_i, q''_i) \leq R. \quad \square$$

This finishes the proof of Proposition 6.1.

7 Cocompact boundary actions

Given a naive convex cocompact triple (Ω, X, G) , it seems natural to ask if G acts cocompactly on $\partial\Omega - F_\Omega(\partial_i X)$ or more generally on $\partial_i \mathcal{C} - F_\Omega(\partial_i X)$ when $\mathcal{C} \subset \Omega$ is a closed G -invariant convex subset containing X . The next theorem gives a sufficient condition for this to occur.

Theorem 7.1 *Suppose that (Ω, X, G) is a naive convex cocompact triple and $\mathcal{C} \subset \Omega$ is a closed G -invariant convex subset containing X with the property that if $x \in \partial_i X$ and $[x, y] \subset \partial_i \mathcal{C}$, then $[x, y] \subset F_\Omega(\partial_i X)$. Then G acts cocompactly on $\partial_i \mathcal{C} - F_\Omega(\partial_i X)$.*

Letting $\mathcal{C} = \Omega$ and changing notation, we obtain:

Corollary 7.2 *Suppose that $(\Omega, \mathcal{C}, \Gamma)$ is a naive convex cocompact triple with the property that if $x \in \partial_i \mathcal{C}$ and $[x, y] \subset \partial\Omega$, then $[x, y] \subset F_\Omega(\partial_i \mathcal{C})$. Then Γ acts cocompactly on $\partial\Omega - F_\Omega(\partial_i \mathcal{C})$.*

Remark 7.3 The main application of Theorem 7.1 will be in Section 9 as we now explain. Suppose that $(\Omega, \mathcal{C}, \Gamma)$ is a naive convex cocompact triple, \mathcal{X} is a peripheral family of $(\Omega, \mathcal{C}, \Gamma)$, and $X \in \mathcal{X}$. Then $(\Omega, X, \text{Stab}_\Gamma(X))$ and \mathcal{C} satisfy the conditions in Theorem 7.1; see Theorem 5.1(2) and (6). We will use this in the proof of Theorem 1.4(2). In particular, we use this to show that nonconical points are bounded parabolic points.

For the rest of the section, fix $\Omega, \mathcal{C}, X, G$ satisfying the hypotheses of Theorem 7.1. The key idea will be to study the following “closest point projection”.

Definition 7.4 Define $\pi_X : \bar{\mathcal{C}} \rightarrow \{\text{subsets of } X\}$ as follows:

- For $p \in \mathcal{C}$, let

$$\pi_X(p) := \{x \in X : d_\Omega(p, x) = d_\Omega(p, X)\}$$
 denote the points in X closest to p .
- For $y \in \partial_i \mathcal{C}$ let $\pi_X(y) \subset X$ denote the set of all points $x \in X$ where there exist sequences $\{p_n\}$ and $\{x_n\}$ such that $p_n \in \mathcal{C}$, $x_n \in \pi_X(p_n)$, $p_n \rightarrow y$, and $x_n \rightarrow x$.

We start by observing the following.

Observation 7.5 (1) *If $p \in \mathcal{C}$, then $\pi_X(p)$ is a compact convex subset of X .*

(2) *If $p \in \mathcal{C}$ and $x \in \pi_X(p)$, then $x \in \pi_X(q)$ for all $q \in [x, p]$.*

(3) *If $g \in G$, then $\pi_X \circ g = g \circ \pi_X$.*

Proof Notice that if $p \in \mathcal{C}$, then

$$\pi_X(p) = X \cap \{x \in \Omega : d_\Omega(p, x) \leq d_\Omega(p, X)\}.$$

Part (1) is then a consequence of the fact that closed metric balls in the Hilbert metric are convex and compact; see Proposition 2.9 above. Part (2) follows from the fact that $[x, p]$ can be parametrized to be a geodesic in the Hilbert metric and part (3) is by definition. \square

Lemma 7.6 *If $y \in \partial_i \mathcal{C} - F_\Omega(\partial_i X)$, then $\pi_X(y) \subset X$ is a nonempty bounded set.*

Proof It suffices to fix sequences $\{p_n\}$ and $\{x_n\}$ such that $p_n \in \mathcal{C}$, $x_n \in \pi_X(p_n)$, $p_n \rightarrow y$, and $x_n \rightarrow x \in \bar{X}$, then prove that $x \in X$.

Suppose for a contradiction that $x \in \partial_i X$. Since $y \in \partial_i \mathcal{C} - F_\Omega(\partial_i X)$ our hypothesis on X says that $(x, y) \subset \Omega$. Fix $v \in (x, y)$. Then pick a sequence $v_n \in (x_n, p_n)$ with $v_n \rightarrow v$. Then $x_n \in \pi_X(v_n)$ by Observation 7.5(2). However, we then have

$$d_\Omega(v, X) = \lim_{n \rightarrow \infty} d_\Omega(v_n, X) = \lim_{n \rightarrow \infty} d_\Omega(v_n, x_n) = \lim_{n \rightarrow \infty} d_\Omega(v, x_n) = \infty,$$

and we have a contradiction. \square

Lemma 7.7 *If $y \in \partial_i \mathcal{C} \cap F_\Omega(\partial_i X)$, then $\pi_X(y) = \emptyset$.*

Proof Suppose for a contradiction that there exist sequences $\{p_n\}$ and $\{x_n\}$ where $p_n \in \mathcal{C}$, $x_n \in \pi_X(p_n)$, $p_n \rightarrow y$, and $x_n \rightarrow x \in X$.

By hypothesis, there exists $w \in \partial_i X$ with $y \in F_\Omega(w)$. Fix $q \in (x, y)$ such that $d_\Omega(q, x) > d_{F_\Omega(w)}(y, w)$. Then fix $q_n \in (x_n, p_n)$ such that $q_n \rightarrow q$. By Observation 7.5(2),

$$d_\Omega(q_n, X) = d_\Omega(q_n, x_n),$$

and so

$$d_\Omega(q, X) = \lim_{n \rightarrow \infty} d_\Omega(q_n, X) = d_\Omega(q, x) > d_{F_\Omega(w)}(y, w).$$

However, this is impossible since Proposition 2.9 implies that

$$d_\Omega(q, X) \leq d_\Omega(q, (x, w)) \leq d_\Omega^{\text{Haus}}((x, y), (x, w)) \leq d_{F_\Omega(w)}(y, w).$$

So we have a contradiction. \square

Lemma 7.8 *If $K \subset X$ is compact, then the set*

$$\hat{K} := \{y \in \partial_i \mathcal{C} : \pi_X(y) \cap K \neq \emptyset\}$$

is compact.

Proof Consider a sequence $\{y_n\}$ in \hat{K} where $y_n \rightarrow y \in \partial_i \mathcal{C}$. Fix $x_n \in \pi_X(y_n) \cap K$. Passing to a subsequence, we can suppose that $x_n \rightarrow x \in K$. By the definition of π_X , we can also find sequences $\{p_{n,m}\}$ and $\{x_{n,m}\}$ such that $p_{n,m} \in \mathcal{C}$, $x_{n,m} \in \pi_X(p_{n,m})$, and

$$y_n = \lim_{m \rightarrow \infty} p_{n,m} \quad \text{and} \quad x_n = \lim_{m \rightarrow \infty} x_{n,m}.$$

Then we can pick subsequences $\{p_{n_j, m_j}\}$ and $\{x_{n_j, m_j}\}$ such that

$$y = \lim_{j \rightarrow \infty} p_{n_j, m_j} \quad \text{and} \quad x = \lim_{j \rightarrow \infty} x_{n_j, m_j}.$$

So $x \in \pi_X(y) \cap K$ and hence $y \in \widehat{K}$. □

Proof of Theorem 7.1 Fix a compact set $K \subset X$ such that $G \cdot K = X$. Then Lemmas 7.8 and 7.7 imply that

$$\widehat{K} := \{y \in \partial_i \mathcal{C} : \pi_X(y) \cap K \neq \emptyset\}$$

is compact and contained in $\partial_i \mathcal{C} - F_\Omega(\partial_i X)$.

We claim that

$$G \cdot \widehat{K} = \partial_i \mathcal{C} - F_\Omega(\partial_i X).$$

Fix $y \in \partial_i \mathcal{C} - F_\Omega(\partial_i X)$. By Lemma 7.6 there exists $x \in \pi_X(y)$. There exists $g \in G$ such that $g(x) \in K$. Then $g(x) \in \pi_X(g(y))$ and so $g(y) \in \widehat{K}$. So $y \in G \cdot \widehat{K}$. Since y was arbitrary, this proves the claim and the theorem. □

8 Basic properties of boundary quotients

For the rest of the section fix a properly convex domain $\Omega \subset \mathbb{P}(\mathbb{R}^d)$, a closed convex subset $\mathcal{C} \subset \Omega$, and a discrete subgroup $\Gamma \leq \text{Aut}(\Omega)$ which preserves \mathcal{C} . We do not assume that Γ acts cocompactly on \mathcal{C} .

Also fix a Γ -invariant equivalence relation \sim on $\partial_i \mathcal{C}$ such that the set

$$R := \{(x, y) \in \partial_i \mathcal{C} \times \partial_i \mathcal{C} : x \sim y\}$$

is closed and $\#(\partial_i \mathcal{C} / \sim) \geq 3$. For each $x \in \partial_i \mathcal{C}$, let $[x]$ denote the equivalence class of x and let

$$(8-1) \quad \mathcal{C}_x := \text{ConvHull}_\Omega([x]) \cap \Omega.$$

Notice that it is possible for \mathcal{C}_x to be empty.

We consider two conditions:

Condition (1) Whenever $x, y \in \partial_i \mathcal{C}$ and $[x, y] \subset \partial \Omega$, we have $x \sim y$.

Condition (2) There exist $r, D > 0$ such that if $x \not\sim y$, then

$$\text{diam}_\Omega(\mathcal{N}_\Omega(\mathcal{C}_x, r) \cap \mathcal{N}_\Omega(\mathcal{C}_y, r)) < D.$$

We will prove the following results about the quotient space $\partial_i \mathcal{C} / \sim$. These arguments are similar to analogous arguments of Choi [5] and Weisman [23].

Proposition 8.1 *The quotient $\partial_i \mathcal{C} / \sim$ is a compact Hausdorff metrizable space.*

Proposition 8.2 *If Condition (1) holds, then Γ acts as a convergence group on $\partial_i \mathcal{C} / \sim$.*

To state the final result, we need a definition.

Definition 8.3 A point $x \in \partial \Omega$ is a *uniformly conical limit point* of Γ acting on Ω if for any $p_0 \in \Omega$ the image of $[p_0, x]$ in $\Gamma \backslash \Omega$ is relatively compact.

Proposition 8.4 *If Conditions (1) and (2) hold, $x \in \partial_i \mathcal{C}$ is a uniformly conical limit point of Γ acting on Ω , and $[x] \subset F_\Omega(x)$, then $[x] \in \partial_i \mathcal{C}/\sim$ is a conical limit point of Γ (in the convergence group sense; see Section 3.2).*

8.1 Proof of Proposition 8.1

The proof is an exercise in point set topology. We include it for the convenience of the reader. We also note that similar arguments appear in [5, Proposition 10.3.8; 23, Proposition 8.10]. For the rest of this proof, let $\pi: \partial_i \mathcal{C} \rightarrow \partial_i \mathcal{C}/\sim$ denote the quotient map.

We begin the proof with an useful observation:

Observation 8.5 *Let $U \subset \partial_i \mathcal{C}$ be an open set, $U^* := \{x \in \partial_i \mathcal{C} : [x] \subset U\}$, and $V := \pi(U^*)$. Then U^* and V are open subsets of $\partial_i \mathcal{C}$ and $\partial_i \mathcal{C}/\sim$ respectively.*

Proof of Observation 8.5 If U^* is empty, then V is empty and the statement is true. So assume that U^* is nonempty. To show that U^* is open, we show that $\partial_i \mathcal{C} - U^*$ is closed. Consider any sequence $\{z_n\}$ in $\partial_i \mathcal{C} - U^*$ such that $z_n \rightarrow z \in \partial_i \mathcal{C}$. Since $z_n \notin U^*$, there exists $z'_n \notin U$ such that $(z_n, z'_n) \in R$. As $\partial_i \mathcal{C} - U$ is compact, we can pass to a subsequence and assume that $z'_n \rightarrow z' \in \partial_i \mathcal{C} - U$. Since R is closed and $(z_n, z'_n) \rightarrow (z, z')$, we have $(z, z') \in R$. Thus, $z \in \partial_i \mathcal{C} - U^*$ and hence $\partial_i \mathcal{C} - U^*$ is closed.

Since U^* is a union of equivalence classes, we have

$$\pi^{-1}(V) = \pi^{-1}(\pi(U^*)) = U^*,$$

and so, by the definition of the quotient topology, V is open. □

$\partial_i \mathcal{C}/\sim$ is Hausdorff Since $\partial_i \mathcal{C}$ is a compact Hausdorff space and R is closed, it is well known that $\partial_i \mathcal{C}/\sim$ is Hausdorff; eg [9, Proposition 1.4.4]. To see this, fix $x, y \in \partial_i \mathcal{C}$ such that $\pi(x) \neq \pi(y)$. Since R is closed, $[w] = \{z \in \partial_i \mathcal{C} : (w, z) \in R\}$ is a closed, and hence compact, subset of $\partial_i \mathcal{C}$ for any $w \in \partial_i \mathcal{C}$. Then $[x]$ and $[y]$ are disjoint compact subsets of a Hausdorff space $\partial_i \mathcal{C}$. So we can separate $[x]$ and $[y]$ using disjoint open sets $U_x, U_y \subset \partial_i \mathcal{C}$.

Let $U_w^* := \{z \in \partial_i \mathcal{C} : [z] \subset U_w\}$ and $V_w := \pi(U_w^*)$ for $w = x, y$. By Observation 8.5, U_w^* are disjoint open sets with $[w] \subset U_w$ for $w = x, y$. Then, by Observation 8.5, V_x and V_y are disjoint nonempty open subsets of $\partial_i \mathcal{C}/\sim$ that separate $\pi(x)$ and $\pi(y)$. So $\partial_i \mathcal{C}/\sim$ is Hausdorff.

$\partial_i \mathcal{C}/\sim$ is compact and metrizable Since $\partial_i \mathcal{C}/\sim$ is Hausdorff and $\partial_i \mathcal{C}/\sim = \pi(\partial_i \mathcal{C})$ is the continuous image of a compact metric space, it is compact and metrizable; eg [24, Corollary 23.2]. Once again, we include the argument.

Since $\partial_i \mathcal{C}$ is compact and π is continuous, then $\partial_i \mathcal{C}/\sim$ is compact. Then, by the Uryshon metrization theorem, it suffices to show that $\partial_i \mathcal{C}/\sim$ has a countable basis to prove that it is metrizable. Since $\partial_i \mathcal{C}$ is

a compact metrizable space, we can choose a countable basis $\mathcal{U} := \{U_n : n \in \mathbb{N}\}$ for $\partial_i \mathcal{C}$. Without loss of generality, we can assume that \mathcal{U} is closed under finite unions. For $n \geq 1$, let

$$U_n^* := \{x \in \partial_i \mathcal{C} : [x] \subset U_n\} \quad \text{and} \quad V_n := \pi(U_n^*).$$

By Observation 8.5, these are open sets. Further, since any equivalence class is compact and \mathcal{U} is closed under finite unions, for any $x \in \partial_i \mathcal{C}$ there exists U_n with $[x] \subset U_n$. Then $x \in [x] \subset U_n^*$ and $[x] \in V_n$. Thus $\mathcal{U}^* := \{U_n^* : U_n^* \neq \emptyset, n \in \mathbb{N}\}$ is an open cover of $\partial_i \mathcal{C}$ and $\mathcal{V} := \{V_n : U_n^* \neq \emptyset, n \in \mathbb{N}\}$ is an open cover of $\partial_i \mathcal{C}/\sim$.

We will show that \mathcal{V} a basis for the topology on $\partial_i \mathcal{C}/\sim$. Let $w \in W \subset \partial_i \mathcal{C}/\sim$ where W is an open set. Then $\pi^{-1}(w) = [w]$ is an equivalence class and hence compact. Then, since \mathcal{U} is closed under finite unions, there exists U_n with $[w] \subset U_n \subset \pi^{-1}(W)$. Then $w \in V_n^* \subset W$. Thus \mathcal{V} is a basis for the topology.

8.2 Proof of Proposition 8.2

We also note that similar arguments appear in [5, Theorem 10.3.1; 23, Proposition 8.8].

Notice that Condition (1) implies that

$$\partial_i \mathcal{C} \cap \overline{F_\Omega(x')} \subset [x']$$

for all $x' \in \partial_i \mathcal{C}$.

Suppose that $\{\gamma_n\}$ is a sequence of distinct elements in Γ . Fix $p_0 \in \mathcal{C}$. Passing to a subsequence we can suppose that $\gamma_n(p_0) \rightarrow x \in \partial_i \mathcal{C}$, $\gamma_n^{-1}(p_0) \rightarrow y \in \partial_i \mathcal{C}$, and $\gamma_n \rightarrow S$ in $\mathbb{P}(\text{End}(\mathbb{R}^d))$. Then, by Observation 2.3,

$$\lim_{n \rightarrow \infty} \gamma_n(z) = S(z)$$

for all $z \in \mathbb{P}(\mathbb{R}^d) - \mathbb{P}(\ker S)$ and the convergence is locally uniform. By Proposition 2.4,

$$S(\bar{\Omega} - \mathbb{P}(\ker S)) \subset \bar{\Omega} \cap \text{Span } F_\Omega(x) = \overline{F_\Omega(x)}.$$

Then

$$S(\bar{\mathcal{C}} - \mathbb{P}(\ker S)) \subset \partial_i \mathcal{C} \cap \overline{F_\Omega(x)} \subset [x].$$

Proposition 2.4 also implies that $\mathbb{P}(\ker S) \cap \Omega = \emptyset$ and $y \in \mathbb{P}(\ker S)$. So if $z \in \partial_i \mathcal{C} \cap \mathbb{P}(\ker S)$, then $[y, z] \subset \partial \Omega$ and then Condition (1) implies that $z \in [y]$. Hence

$$\gamma_n|_{(\partial_i \mathcal{C}/\sim) - [y]}$$

converges locally uniformly to the constant map $[x]$.

8.3 Proof of Proposition 8.4

We also note that similar arguments appear in [5, Theorem 10.3.1; 23, Proposition 8.17].

Recall the notation \mathcal{C}_x from (8-1). We start with a lemma.

Lemma 8.6 *If $y \in \partial_i \mathcal{C}$ and $z \in F_\Omega(\partial_i \mathcal{C}_y) \cap \partial_i \mathcal{C}$, then $z \sim y$.*

Proof By hypothesis, there exists some $z' \in \partial_i \mathcal{C}_y$ with $z \in F_\Omega(z')$.

Since $[y]$ is closed, the extreme points of $\text{ConvHull}_\Omega([y])$ are contained in $[y]$. Then there exist $y_1, \dots, y_m \in [y]$ such that

$$z' \in \text{rel-int}(\text{ConvHull}_\Omega(\{y_1, \dots, y_m\})).$$

Thus $y_1, \dots, y_m \in \overline{F_\Omega(z')}$. Then $[z, y_1] \subset \partial\Omega$ where $z, y_1 \in \partial_i \mathcal{C}$. Condition (1) implies that $z \sim y_1$. Thus $z \sim y$. \square

Now suppose $x \in \partial_i \mathcal{C}$ is a uniform conical limit point of Γ acting on Ω . Fix $p_0 \in \mathcal{C}$ and a sequence $\{p_n\}$ in $[p_0, x]$ with $p_n \rightarrow x$.

By Condition (2), there exist $r, D > 0$ such that if $y_1 \not\sim y_2$, then

$$\text{diam}_\Omega(\mathcal{N}_\Omega(\mathcal{C}_{y_1}, r) \cap \mathcal{N}_\Omega(\mathcal{C}_{y_2}, r)) < D.$$

Lemma 8.7 For every $n \in \mathbb{N}$, there exists $q_n \in [p_n, x]$ such that

$$\mathcal{B}_\Omega(q_n, 2D) \cap (p_0, x) \not\subset \mathcal{N}_\Omega(\mathcal{C}_y, r)$$

for all $y \in \partial_i \mathcal{C}$.

Proof Fix $n \in \mathbb{N}$ and suppose not. Then for every $q \in [p_n, x]$ there exists $y(q) \in \partial_i \mathcal{C}$ such that

$$\mathcal{B}_\Omega(q, 2D) \cap (p_0, x) \subset \mathcal{N}_\Omega(\mathcal{C}_{y(q)}, r).$$

Note that this implies that $\mathcal{C}_{y(q)}$ is a nonempty set. We claim that $\mathcal{C}_{y(q)}$ does not depend on q . To show this it is enough to fix $q', q'' \in [p_n, x]$ with $d_\Omega(q', q'') \leq D$ and show that $\mathcal{C}_{y(q')} = \mathcal{C}_{y(q'')}$. In this case, $\mathcal{B}_\Omega(q', D) \subset \mathcal{B}_\Omega(q'', 2D)$; hence

$$\mathcal{B}_\Omega(q', D) \cap (p_0, x) \subset \mathcal{N}_\Omega(\mathcal{C}_{y(q')}, r) \cap \mathcal{N}_\Omega(\mathcal{C}_{y(q'')}, r).$$

Then by our choice of D , we have $\mathcal{C}_{y(q')} = \mathcal{C}_{y(q'')}$. Thus $\mathcal{C}_{y(q)}$ does not depend on $q \in [p_n, x]$. Thus

$$(p_n, x) \subset \mathcal{N}_\Omega(\mathcal{C}_{y(p_n)}, r)$$

Then Proposition 2.8 implies that $x \in F_\Omega(\partial_i \mathcal{C}_{y(p_n)})$. So by the previous lemma $x \sim y(p_n)$. Then $\mathcal{C}_x = \mathcal{C}_{y(p_n)}$ is nonempty, which contradicts the assumption that $[x] \subset F_\Omega(x)$. \square

We finish the proof of Proposition 8.4. Fix a sequence $\{q_n\}$ as in the above lemma. Since x is a uniformly conical limit point, there exists a sequence $\{\gamma_n\}$ in Γ such that $\{\gamma_n(q_n)\}$ is relatively compact in Ω . Passing to a subsequence we can suppose that $\gamma_n(p_0) \rightarrow b$, $\gamma_n^{-1}(p_0) \rightarrow c$, and $\gamma_n(x) \rightarrow a$. By the proof of Proposition 8.2,

$$\gamma_n|_{(\partial_i \mathcal{C}/\sim)-[c]}$$

converges locally uniformly to the constant map $[b]$. Also $\gamma_n[x] \rightarrow [a]$. So, to show that $[x]$ is a conical limit point in the convergence group sense, we need to prove that $[x] = [c]$ and $[a] \neq [b]$.

Suppose for a contradiction that $[a] = [b]$. Since $\{\gamma_n(q_n)\}$ is relatively compact in Ω and $\gamma_n(q_n) \in (\gamma_n(x), \gamma_n(p_0))$, we have $(a, b) \subset \Omega$. Hence $(a, b) \subset \mathcal{C}_a$ and so for n large,

$$\mathcal{B}_\Omega(\gamma_n(q_n), 2D) \cap (\gamma_n(p_0), \gamma_n(x)) \subset \mathcal{N}_\Omega(\mathcal{C}_a, r).$$

This implies that

$$\mathcal{B}_\Omega(q_n, 2D) \cap (p_0, x) \subset \mathcal{N}_\Omega(\mathcal{C}_{\gamma_n^{-1}(a)}, r),$$

which is a contradiction. So $[a] \neq [b]$.

Since $\gamma_n([x]) \rightarrow [a]$ and $[a] \neq [b]$, we must have $[x] = [c]$.

9 Proof of Theorem 1.4

In this section we prove Theorem 1.4. Suppose that $(\Omega, \mathcal{C}, \Gamma)$ is a naive convex cocompact triple.

Theorem 1.4(1) Suppose that Γ is relatively hyperbolic with respect to $\mathcal{P} := \{P_1, \dots, P_N\}$ and

$$X_j := \text{ConvHull}_\Omega(\mathcal{L}_\Omega(P_j) \cap \partial_i \mathcal{C}) \cap \Omega, \quad \text{where } j = 1, \dots, N.$$

Then Proposition 6.1 implies that (\mathcal{C}, d_Ω) is relatively hyperbolic with respect to

$$\mathcal{X} = \Gamma \cdot \{X_1, \dots, X_N\}.$$

Proposition 3.4 implies that \mathcal{X} is a peripheral family of $(\Omega, \mathcal{C}, \Gamma)$.

Theorem 1.4(2) Suppose that \mathcal{X} is a peripheral family of $(\Omega, \mathcal{C}, \Gamma)$ and \mathcal{P} is a set of representatives of the Γ -conjugacy classes in $\{\text{Stab}_\Gamma(X) : X \in \mathcal{X}\}$. Let

$$[\partial_i \mathcal{C}]_{\mathcal{X}} := \partial_i \mathcal{C} / \sim$$

be as in Definition 1.2. We claim that the action of Γ on $[\partial_i \mathcal{C}]_{\mathcal{X}}$ satisfies Theorem 3.5.

Lemma 9.1 *The set $R := \{(x, y) \in \partial_i \mathcal{C} \times \partial_i \mathcal{C} : x \sim y\}$ is closed.*

Proof Suppose that $\{(x_n, y_n)\}$ is a sequence in R converging to (x, y) in $\partial_i \mathcal{C} \times \partial_i \mathcal{C}$.

Case 1 Assume $(x, y) \subset \Omega$. Then $(x_n, y_n) \subset \Omega$ for n sufficiently large, and for such n there exists $X_n \in \mathcal{X}$ with $x_n, y_n \in F_\Omega(\partial_i X_n)$. By Theorem 5.1(5) there is some $R > 0$ such that for every n there exist $x'_n, y'_n \in \partial_i X_n$ where

$$d_{F_\Omega(x_n)}(x_n, x'_n) \leq R \quad \text{and} \quad d_{F_\Omega(y_n)}(y_n, y'_n) \leq R.$$

Then by Proposition 2.9,

$$d_\Omega^{\text{Haus}}((x_n, y_n), (x'_n, y'_n)) \leq R.$$

Since $(x_n, y_n) \rightarrow (x, y)$, this implies that

$$\lim_{n, k \rightarrow \infty} \text{diam}_\Omega(\mathcal{N}_\Omega(X_n, R+1) \cap \mathcal{N}_\Omega(X_k, R+1)) = \infty.$$

Since \mathcal{X} is strongly isolated, $X := X_n = X_k$ for n, k sufficiently large. Then

$$(x, y) \subset \mathcal{N}_\Omega(X, R + 1),$$

and so $x, y \in F_\Omega(\partial_i X)$ by Proposition 2.8. Thus $x \sim y$.

Case 2 Assume $[x, y] \subset \partial\Omega$. Then $x \sim y$ by Theorem 5.1(7). \square

Let

$$B := \{[F_\Omega(\partial_i X) \cap \partial_i \mathcal{C}] : X \in \mathcal{X}\} \subset [\partial_i \mathcal{C}]_\mathcal{X}.$$

Lemma 9.2 (1) $[\partial_i \mathcal{C}]_\mathcal{X}$ is a compact Hausdorff metrizable space.

(2) Γ acts as a convergence group on $[\partial_i \mathcal{C}]_\mathcal{X}$.

(3) If $z \in [\partial_i \mathcal{C}]_\mathcal{X} - B$, then z is a conical limit point of Γ .

Proof We use the results of Section 8. Part (1) follows from the previous lemma and Proposition 8.1.

Notice that Theorem 5.1(7) implies that the quotient $[\partial_i \mathcal{C}]_\mathcal{X}$ satisfies Condition (1) from Section 8. We claim that the quotient $[\partial_i \mathcal{C}]_\mathcal{X}$ satisfies Condition (2) from Section 8. First notice that if $z \in [\partial_i \mathcal{C}]_\mathcal{X} - B$, then $\mathcal{C}_z = \emptyset$. So it suffices to consider elements in B . Further, if $b = [F_\Omega(\partial_i X) \cap \partial_i \mathcal{C}] \in B$, then by Theorem 5.1(2) and (8), $\text{Stab}_\Gamma(X)$ acts cocompactly on both \mathcal{C}_b and X . So \mathcal{C}_b is contained in a bounded neighborhood of X . By Theorem 5.1(1), we can choose this bound to be independent of b . Then since \mathcal{X} is strongly isolated, $[\partial_i \mathcal{C}]_\mathcal{X}$ satisfies Condition (2).

Then Γ acts as a convergence group on $[\partial_i \mathcal{C}]_\mathcal{X}$ by Proposition 8.2. If $z \in [\partial_i \mathcal{C}]_\mathcal{X} - B$, then we have $z = [F_\Omega(x) \cap \partial_i \mathcal{C}]$ for some $x \in \partial_i \mathcal{C}$ and so Proposition 8.4 implies that z is a conical limit point of Γ . This proves parts (2) and (3). \square

Lemma 9.3 (1) Γ has finitely many orbits in B .

(2) If $b = [F_\Omega(\partial_i X) \cap \partial_i \mathcal{C}] \in B$, then $\text{Stab}_\Gamma(b) = \text{Stab}_\Gamma(X)$. In particular, $\text{Stab}_\Gamma(b)$ is finitely generated.

(3) If $b \in B$, then b is a bounded parabolic fixed point.

Proof (1) This follows immediately from Theorem 5.1(1).

(2) It is clear that $\text{Stab}_\Gamma(b) \supset \text{Stab}_\Gamma(X)$. For the other inclusion, if $\gamma \in \text{Stab}_\Gamma(b)$, then

$$F_\Omega(\partial_i(\gamma X)) = \gamma F_\Omega(\partial_i X) = F_\Omega(\partial_i X).$$

So $\gamma X = X$ by Theorem 5.1(3). Hence $\gamma \in \text{Stab}_\Gamma(X)$ and so $\text{Stab}_\Gamma(b) = \text{Stab}_\Gamma(X)$. The “in particular” part then follows from Theorem 5.1(2).

(3) The proof is an application of Theorem 7.1, as outlined in Remark 7.3. We now provide the details.

Fix $b = [F_\Omega(\partial_i X) \cap \partial_i \mathcal{C}] \in B$. By Theorem 5.1(2), $\text{Stab}_\Gamma(X)$ acts cocompactly on X . Furthermore, by Theorem 5.1(6), if $x \in \partial_i X$ and $[x, y] \subset \partial_i \mathcal{C}$, then $[x, y] \in F_\Omega(\partial_i X)$.

Then Theorem 7.1 implies that $\text{Stab}_\Gamma(X)$ acts cocompactly on $\partial_i \mathcal{C} - F_\Omega(\partial_i X)$. Thus $\text{Stab}_\Gamma(b)$ acts cocompactly on $[\partial_i \mathcal{C}]_\mathcal{X} - \{b\}$. \square

Then Theorem 3.5 implies that Γ is relatively hyperbolic with respect to \mathcal{P} and there exists a Γ -equivariant homeomorphism $\partial(\Gamma, \mathcal{P}) \rightarrow [\partial_i \mathcal{C}]_{\mathcal{X}}$.

Theorem 1.4(a)–(e) Now suppose that at least one of the conditions is satisfied. Proposition 6.1 implies parts (a) and (c). Part (b) was established in the proof of Theorem 1.4(2); see the last paragraph. Parts (d) and (e) follow from Theorem 5.1.

10 The convex cocompact case

We prove Theorem 1.7 and Proposition 1.8. As mentioned in the introduction, the key result which allows our results to simplify in the convex cocompact case is the following observation; see [7, Corollaries 4.10 and 4.13].

Observation 10.1 *Suppose that $\Omega \subset \mathbb{P}(\mathbb{R}^d)$ is a properly convex domain and $\Gamma \leq \text{Aut}(\Omega)$ is a convex cocompact subgroup.*

- (1) *If $x \in \partial\Omega$ and $F_{\Omega}(x) \cap \partial_i \mathcal{C}_{\Omega}(\Gamma) \neq \emptyset$, then $F_{\Omega}(x) \subset \partial_i \mathcal{C}_{\Omega}(\Gamma)$.*
- (2) *$\mathcal{L}_{\Omega}(\Gamma) = \partial_i \mathcal{C}_{\Omega}(\Gamma)$. In particular, $\mathcal{L}_{\Omega}(\Gamma)$ is closed.*

Proof (1) Fix $x \in \partial\Omega$ with $F_{\Omega}(x) \cap \partial_i \mathcal{C}_{\Omega}(\Gamma) \neq \emptyset$. Then Proposition 2.5 implies that $F_{\Omega}(x) \subset \mathcal{L}_{\Omega}(\Gamma)$ and we have $\mathcal{L}_{\Omega}(\Gamma) \subset \partial_i \mathcal{C}_{\Omega}(\Gamma)$ by the definition of $\mathcal{C}_{\Omega}(\Gamma)$.

(2) Since Γ acts cocompactly on \mathcal{C} , we have $\partial_i \mathcal{C}_{\Omega}(\Gamma) \subset \mathcal{L}_{\Omega}(\Gamma)$ and the reverse inclusion is by definition of $\mathcal{C}_{\Omega}(\Gamma)$. □

10.1 Proof of Theorem 1.7

Suppose that $\Gamma \leq \text{Aut}(\Omega)$ is a convex cocompact group such that Γ is relatively hyperbolic with respect to $\mathcal{P} := \{P_1, \dots, P_N\}$. Then by definition $(\Omega, \mathcal{C}_{\Omega}(\Gamma), \Gamma)$ is a naive convex cocompact triple. Let $\mathcal{C} = \mathcal{C}_{\Omega}(\Gamma)$ and

$$\mathcal{X} := \Gamma \cdot \{\mathcal{C}_{\Omega}(P_1), \dots, \mathcal{C}_{\Omega}(P_N)\}.$$

Lemma 10.2 (parts (a) and (c)) *Each P_j is a convex cocompact subgroup of $\text{Aut}(\Omega)$ and $(\mathcal{C}, d_{\Omega})$ is relatively hyperbolic with respect to \mathcal{X} .*

Proof Since $\mathcal{L}_{\Omega}(P_j) \subset \mathcal{L}_{\Omega}(\Gamma) \subset \partial_i \mathcal{C}$, then

$$\mathcal{C}_{\Omega}(P_j) = \text{ConvHull}_{\Omega}(\mathcal{L}_{\Omega}(P_j)) \cap \Omega = \text{ConvHull}_{\Omega}(\mathcal{L}_{\Omega}(P_j) \cap \partial_i \mathcal{C}) \cap \Omega.$$

Proposition 6.1 implies that P_j acts cocompactly on $\mathcal{C}_{\Omega}(P_j)$. Then $(\mathcal{C}, d_{\Omega})$ is relatively hyperbolic with respect to \mathcal{X} . □

If $X \in \mathcal{X}$, then $X = \gamma \mathcal{C}_{\Omega}(P_j) = \mathcal{C}_{\Omega}(\gamma P_j \gamma^{-1})$ for some $\gamma \in \Gamma$ and $1 \leq j \leq N$. Then, since P_j is a convex cocompact subgroup of $\text{Aut}(\Omega)$, Observation 10.1 implies that

$$(10-1) \quad F_{\Omega}(\partial_i X) = \partial_i X = \mathcal{L}_{\Omega}(\gamma P_j \gamma^{-1}).$$

Lemma 10.3 (part (e)) *If $\ell \subset \partial_i \mathcal{C}_\Omega(\Gamma)$ is a nontrivial line segment, then $\ell \subset \mathcal{L}_\Omega(\gamma P_j \gamma^{-1})$ for some $\gamma \in \Gamma$ and $P_j \in \mathcal{P}$.*

Proof Fix $x \in \text{rel-int}(\ell)$. Then $\dim F_\Omega(x) \geq 1$ and so, by Observation 10.1,

$$\text{diam}_{F_\Omega(x)}(F_\Omega(x) \cap \partial_i \mathcal{C}) = \text{diam}_{F_\Omega(x)} F_\Omega(x) = \infty.$$

Thus by Theorem 1.4(d), there exist $\gamma \in \Gamma$ and $P_j \in \mathcal{P}$ such that

$$x \in F_\Omega(\partial_i X),$$

where $X = \mathcal{C}_\Omega(\gamma P_j \gamma^{-1}) = \gamma \mathcal{C}_\Omega(P_j)$. Then (10-1) implies that

$$\text{rel-int}(\ell) \subset F_\Omega(x) \subset F_\Omega(\partial_i X) = \mathcal{L}_\Omega(\gamma P_j \gamma^{-1}).$$

Since $\mathcal{L}_\Omega(\gamma P_j \gamma^{-1})$ is closed (see Observation 10.1), this completes the proof. \square

Recall that, by definition, $[\partial_i \mathcal{C}_\Omega(\Gamma)]_{\mathcal{P}}$ is obtained from $\partial_i \mathcal{C}_\Omega(\Gamma)$ by collapsing each $\mathcal{L}_\Omega(\gamma P_j \gamma^{-1})$ to a point. Hence (10-1) and Lemma 10.3 imply that

$$(10-2) \quad [\partial_i \mathcal{C}]_{\mathcal{X}} = [\partial_i \mathcal{C}]_{\mathcal{P}}.$$

Lemma 10.4 (part (b)) *There is a Γ -equivariant homeomorphism*

$$\partial(\Gamma, \mathcal{P}) \rightarrow [\partial_i \mathcal{C}_\Omega(\Gamma)]_{\mathcal{P}}.$$

Proof This follows immediately from Theorem 1.4(b) and (10-2). \square

To prove Theorem 1.7(d) we will use the following lemma.

Lemma 10.5 [15, Lemma 15.5] *Assume that $x \in \partial_i \mathcal{C}$ is not a C^1 -smooth point of $\partial\Omega$ and $q \in \mathcal{C}$. For any $r > 0$ and $\epsilon > 0$ there exists $q_{r,\epsilon} \in (x, q]$ with the following property: if $p \in (x, q_{r,\epsilon}]$, then there exists a properly embedded simplex $S = S(p) \subset \mathcal{C}$ of dimension at least two such that*

$$(10-3) \quad \mathcal{B}_\Omega(p, r) \cap (x, q] \subset \mathcal{N}_\Omega(S, \epsilon).$$

Let us now prove part (d).

Lemma 10.6 (part (d)) *If $x \in \partial_i \mathcal{C}_\Omega(\Gamma)$ is not a C^1 -smooth point of Ω , then $x \in \mathcal{L}_\Omega(\gamma P_j \gamma^{-1})$ for some $\gamma \in \Gamma$ and $P_j \in \mathcal{P}$.*

Proof Fix $q \in \mathcal{C}$. By Theorem 1.4, \mathcal{X} is a peripheral family of $(\Omega, \mathcal{C}, \Gamma)$. So there exist constants $D_1, D_2 > 0$ such that:

- (a) If $S \subset \mathcal{C}$ is a properly embedded simplex of dimension at least two, then there exists $X \in \mathcal{X}$ with $S \subset \mathcal{N}_\Omega(X, D_2)$.
- (b) If $X_1, X_2 \in \mathcal{X}$ are distinct, then

$$\text{diam}_\Omega(\mathcal{N}_\Omega(X_1, D_2 + 1) \cap \mathcal{N}_\Omega(X_2, D_2 + 1)) < D_1.$$

Let $q' \in (x, q]$ satisfy Lemma 10.5 for the constants $r = D_1 + 1$ and $\epsilon = 1$. Fix a sequence $\{p_n\}$ in $(x, q']$ such that $p_n \rightarrow x$ and $d_\Omega(p_n, p_{n+1}) \leq 1$. For each $n \geq 1$, fix $S_n \subset \mathcal{C}$ a properly embedded simplex of dimension at least two such that

$$\mathcal{B}_\Omega(p_n, D_1 + 1) \cap (x, q] \subset \mathcal{N}_\Omega(S_n, 1).$$

Then fix $X_n \in \mathcal{X}$ such that

$$S_n \subset \mathcal{N}_\Omega(X_n, D_2).$$

Then

$$\begin{aligned} \mathcal{B}_\Omega(p_n, D_1) \cap (x, q] &\subset (\mathcal{B}_\Omega(p_n, D_1 + 1) \cap \mathcal{B}_\Omega(p_{n+1}, D_1 + 1)) \cap (x, q] \\ &\subset \mathcal{N}_\Omega(X_n, D_2 + 1) \cap \mathcal{N}_\Omega(X_{n+1}, D_2 + 1). \end{aligned}$$

Since $\text{diam}_\Omega(\mathcal{B}_\Omega(p_n, D_1) \cap (x, q]) \geq D_1$, this implies that there exists $X \in \mathcal{X}$ such that $X = X_n$ for all n large enough. Then

$$(x, q'] \subset \mathcal{N}_\Omega(X, D_2 + 1).$$

Now $X = \mathcal{C}_\Omega(\gamma P_j \gamma^{-1})$ for some $\gamma \in \Gamma$ and $P_j \in \mathcal{P}$. Finally, Proposition 2.8 and (10-1) imply that $x \in \partial_i \mathcal{C}_\Omega(\gamma P_j \gamma^{-1}) = \mathcal{L}_\Omega(\gamma P_j \gamma^{-1})$. \square

10.2 Proof of Proposition 1.8

Suppose that $\Gamma \leq \text{Aut}(\Omega)$ is a convex cocompact subgroup and \mathcal{X} is a Γ -invariant collection of closed unbounded convex subsets of Ω .

(1) \implies (2) Suppose that \mathcal{X} is a peripheral family of $(\Omega, \mathcal{C}_\Omega(\Gamma), \Gamma)$.

Lemma 10.7 \mathcal{X} is closed in the local Hausdorff topology induced by the Hilbert metric d_Ω .

Proof This follows immediately from Lemma 5.2. \square

Lemma 10.8 If $X_1, X_2 \in \mathcal{X}$ are distinct, then $\partial_i X_1 \cap \partial_i X_2 = \emptyset$.

Proof We prove the contrapositive. Suppose $x \in \partial_i X_1 \cap \partial_i X_2$. Fix $q_1 \in X_1$ and $q_2 \in X_2$. Then

$$d_\Omega^{\text{Haus}}((x, q_1], (x, q_2]) \leq d_\Omega(q_1, q_2)$$

by Proposition 2.9. So

$$(x, q_1] \subset \mathcal{N}_\Omega(X_1, d_\Omega(q_1, q_2) + 1) \cap \mathcal{N}_\Omega(X_2, d_\Omega(q_1, q_2) + 1),$$

and hence $X_1 = X_2$. \square

Lemma 10.9 If $\ell \subset \partial_i \mathcal{C}_\Omega(\Gamma)$ is a nontrivial line segment, then $\ell \subset \partial_i X$ for some $X \in \mathcal{X}$.

Proof Fix $x \in \text{rel-int}(\ell)$. Then $\dim F_\Omega(x) \geq 1$ and so by Observation 10.1,

$$\text{diam}_{F_\Omega(x)}(F_\Omega(x) \cap \partial_i \mathcal{C}) = \text{diam}_{F_\Omega(x)} F_\Omega(x) = \infty.$$

Thus by Theorem 5.1(4) there exists $X \in \mathcal{X}$ with $x \in F_\Omega(\partial_i X)$. We claim that $F_\Omega(x) \subset \partial_i X$ which will imply the lemma.

To show that $F_\Omega(x) \subset \partial_i X$, it suffices to fix an extreme point $e \in \partial F_\Omega(x)$ of $F_\Omega(x)$ and show that $\partial_i X$ contains e . By Observation 10.1 and Theorem 5.1(5) there exists $R > 0$ such that

$$d_{F_\Omega(x)}^{\text{Haus}}(F_\Omega(x), \partial_i X \cap F_\Omega(x)) \leq R.$$

Since $F_{F_\Omega(x)}(e) = \{e\}$, then Proposition 2.8 implies that $e \in \partial_i X$. \square

(2) \implies (1) Suppose that \mathcal{X} has the following properties:

- (a) \mathcal{X} is closed in the local Hausdorff topology induced by the Hilbert metric d_Ω .
- (b) If $X_1, X_2 \in \mathcal{X}$ are distinct, then $\partial_i X_1 \cap \partial_i X_2 = \emptyset$.
- (c) If $\ell \subset \partial_i \mathcal{C}_\Omega(\Gamma)$ is a nontrivial line segment, then $\ell \subset \partial_i X$ for some $X \in \mathcal{X}$.

Lemma 10.10 *If $X_1, X_2 \in \mathcal{X}$ and $F_\Omega(\partial_i X_1) \cap F_\Omega(\partial_i X_2) \neq \emptyset$, then $X_1 = X_2$.*

Proof By hypothesis there exists $x_1 \in \partial_i X_1$ and $x_2 \in \partial_i X_2$ with $F_\Omega(x_1) = F_\Omega(x_2)$. If $x_1 = x_2$, then property (b) implies that $X_1 = X_2$. Otherwise, $[x_1, x_2] \subset F_\Omega(x_1) \subset \partial\Omega$ and so by property (c) there exists X_3 with $[x_1, x_2] \subset \partial_i X_3$. But then by property (b), $X_1 = X_3 = X_2$. \square

Lemma 10.11 *\mathcal{X} is discrete in the local Hausdorff topology induced by the Hilbert metric d_Ω .*

Proof Fix a sequence $\{X_n\}$ in \mathcal{X} converging to some closed subset X . Since \mathcal{X} is closed, $X \in \mathcal{X}$. Fix $p \in X$ and $x \in \partial_i X$.

We claim that $X_n = X$ when n is sufficiently large. Suppose not. Then after passing to a subsequence we can suppose that $X_n \neq X$ for all n . For each n fix $p_n \in X_n$ such that $p_n \rightarrow p$. Passing to a tail of our sequence we can suppose that $d_\Omega(p_n, p) < 1$ for all n . The previous lemma implies that $x \notin F_\Omega(\partial_i X_n)$ and so by Proposition 2.8,

$$\lim_{q \in [p_n, x], q \rightarrow x} d_\Omega(q, X_n) = +\infty.$$

So for each n there exists $q_n \in [p_n, x)$ with $d_\Omega(q_n, X_n) = 1$. Notice that

$$\lim_{n \rightarrow \infty} d_\Omega(q_n, X) \leq \lim_{n \rightarrow \infty} d_\Omega^{\text{Haus}}((x, p_n], (x, p]) \leq \lim_{n \rightarrow \infty} d_\Omega(p_n, p) = 0$$

by Proposition 2.9. Then, since $X_n \rightarrow X$ in the local Hausdorff topology and $d_\Omega(q_n, X_n) = 1$, the sequence $\{q_n\}$ must leave every compact subset of Ω . So $q_n \rightarrow x$.

Since $q_n \in \mathcal{C}$, there exists $\{\gamma_n\}$ in Γ such that $\{\gamma_n(q_n)\}$ is relatively compact in Ω . Passing to a subsequence, we can suppose that $\gamma_n(q_n) \rightarrow q_\infty \in \mathcal{C}$, $\gamma_n(p) \rightarrow p_\infty \in \bar{\mathcal{C}}$, $\gamma_n(X) \rightarrow X_\infty$, and $\gamma_n(X_n) \rightarrow Y_\infty$. Since $q_n \rightarrow x$, we must have $p_\infty \in \partial_i \mathcal{C}$. Since \mathcal{X} is Γ -invariant and closed in the local Hausdorff topology, $X_\infty, Y_\infty \in \mathcal{X}$.

By construction, $q_\infty \in X_\infty$ and $d_\Omega(q_\infty, Y_\infty) = 1$. So $X_\infty \neq Y_\infty$. Also

$$\lim_{n \rightarrow \infty} d_\Omega(\gamma_n(p), \gamma_n(p_n)) = \lim_{n \rightarrow \infty} d_\Omega(p_n, p) = 0,$$

so Proposition 2.8 implies that $\gamma_n(p_n) \rightarrow p_\infty$. Then, by Proposition 2.9,

$$[q_\infty, p_\infty) \subset \mathcal{N}_\Omega(Y_\infty, 2),$$

since $[q_n, p_n] \subset \mathcal{N}_\Omega(X_n, 2)$ for all n . So Proposition 2.8 implies that $p_\infty \in F_\Omega(\partial_i Y_\infty)$. However, by construction, $p_\infty \in \partial_i X_\infty$ and so Lemma 10.10 implies that $X_\infty = Y_\infty$. So we have a contradiction. \square

Lemma 10.12 \mathcal{X} is strongly isolated.

Proof Fix $r > 0$ and suppose for a contradiction that for every $n \in \mathbb{N}$ there exist $X_n, Y_n \in \mathcal{X}$ distinct such that

$$\text{diam}_\Omega(\mathcal{N}_\Omega(X_n, r) \cap \mathcal{N}_\Omega(Y_n, r)) \geq n.$$

Proposition 2.9 implies that $C_n := \mathcal{N}_\Omega(X_n, r) \cap \mathcal{N}_\Omega(Y_n, r)$ is convex.

We claim that each C_n is bounded in (\mathcal{C}, d_Ω) . If not, there exists $x \in \bar{C}_n \cap \partial\Omega$. Then Proposition 2.8 implies that $x \in F_\Omega(\partial_i X_n) \cap F_\Omega(\partial_i Y_n)$. So $X_n = Y_n$ by Lemma 10.10. Thus we have a contradiction and so each C_n is bounded in (\mathcal{C}, d_Ω) .

For each n let $(p_n, q_n) \subset C_n$ denote an open line segment with maximal length (with respect to the Hilbert metric). Then $d_\Omega(p_n, q_n) \geq n$.

Now fix a sequence $\{\gamma_n\}$ in Γ such that $\{\gamma_n(p_n)\}$ is relatively compact in Ω . Passing to a subsequence, we can assume that $\gamma_n(p_n) \rightarrow p \in \Omega$, $\gamma_n(q_n) \rightarrow q \in \bar{\Omega}$, $\gamma_n(X_n) \rightarrow X$, and $\gamma_n(Y_n) \rightarrow Y$. Then $q \in \partial_i \mathcal{C}$, as $d_\Omega(p, q_n) \rightarrow \infty$ and $q_n \in \mathcal{C}$. Furthermore,

$$(q, p) \subset \mathcal{N}_\Omega(X, r + 1) \cap \mathcal{N}_\Omega(Y, r + 1),$$

and $X, Y \in \mathcal{X}$. Proposition 2.8 implies that

$$q \in F_\Omega(\partial_i X) \cap F_\Omega(\partial_i Y).$$

So Lemma 10.10 implies that $X = Y$. Then Lemma 10.11 implies that

$$\gamma_n(X_n) = X = Y = \gamma_n(Y_n)$$

for n large. So $X_n = Y_n$ for n large and we have a contradiction. \square

The following result implies that \mathcal{X} coarsely contains the properly embedded simplices of \mathcal{C} .

Lemma 10.13 If $S \subset \mathcal{C}$ is a properly embedded simplex with dimension at least two, then there exists $X \in \mathcal{X}$ with $S \subset X$.

Proof Fix a properly embedded simplex $S \subset \mathcal{C}$ with dimension at least two. Let v_1, \dots, v_k denote the vertices of S . By property (c), for each $2 \leq j \leq k$ there exists some $X_j \in \mathcal{X}$ with $[v_1, v_j] \subset \partial_i X_j$. Then by property (b), $X_2 = X_3 = \dots = X_k$. So by convexity $S \subset X_2$. \square

Proof of (1-1) in Proposition 1.8 Suppose that (1) and (2) hold, and let \mathcal{P} be a set of representatives of the Γ -conjugacy classes in $\{\text{Stab}_\Gamma(X) : X \in \mathcal{X}\}$. To show that the quotients $[\partial_i \mathcal{C}]_\mathcal{X}$ and $[\partial_i \mathcal{C}]_\mathcal{P}$ coincide it suffices to prove the following.

Lemma 10.14 *If $X \in \mathcal{X}$, then $\partial_i X = F_\Omega(\partial_i X) = \mathcal{L}_\Omega(\text{Stab}_\Gamma(X))$.*

Proof Let $P = \text{Stab}_\Gamma(X)$.

We first observe that $\partial_i X = F_\Omega(\partial_i X)$. By definition $\partial_i X \subset F_\Omega(\partial_i X)$. For the other inclusion, fix $x \in F_\Omega(\partial_i X)$. Then fix $x' \in \partial_i X$ with $x \in F_\Omega(x')$. Observation 10.1 implies that $x \in \partial_i \mathcal{C}$. So, if $x' \neq x$, then there exists $Y \in \mathcal{X}$ with $[x', x] \subset \partial_i Y$. But then $\partial_i X \cap \partial_i Y \neq \emptyset$ which implies that $X = Y$. So $x \in \partial_i X$. Thus $\partial_i X = F_\Omega(\partial_i X)$.

Next we show that $\mathcal{L}_\Omega(P)$ is a subset of $\partial_i X$. Fix $x \in \mathcal{L}_\Omega(P)$, then there exist $p \in \Omega$ and a sequence $\{g_n\}$ in P such that $g_n(p) \rightarrow x$. Fix $q \in X$. By passing to a subsequence we may suppose that $g_n(q) \rightarrow x' \in \partial_i X$. Then Proposition 2.8 implies that $x \in F_\Omega(x') \subset \partial_i X$.

Finally, Theorem 5.1(2) implies that P acts cocompactly on X and so $\partial_i X \subset \mathcal{L}_\Omega(P)$ by Proposition 2.5. \square

References

- [1] **Y Benoist**, *Convexes divisibles IV: Structure du bord en dimension 3*, Invent. Math. 164 (2006) 249–278 MR
- [2] **H Busemann, P J Kelly**, *Projective geometry and projective metrics*, Academic Press, New York (1953) MR
- [3] **S Choi**, *A classification of radial or totally geodesic ends of real projective orbifolds I: a survey of results*, from “Hyperbolic geometry and geometric group theory” (K Fujiwara, S Kojima, K Ohshika, editors), Adv. Stud. Pure Math. 73, Math. Soc. Japan, Tokyo (2017) 69–134 MR
- [4] **S Choi**, *The convex real projective orbifolds with radial or totally geodesic ends: a survey of some partial results*, from “In the tradition of Ahlfors–Bers, VII” (A S Basmajian, Y N Minsky, A W Reid, editors), Contemp. Math. 696, Amer. Math. Soc., Providence, RI (2017) 51–85 MR
- [5] **S Choi**, *Real projective orbifolds with ends and their deformation theory: the deformation theory for nicest ones*, preprint (2025) arXiv 2503.00438v2
- [6] **D Cooper, D D Long, S Tillmann**, *On convex projective manifolds and cusps*, Adv. Math. 277 (2015) 181–251 MR
- [7] **J Danciger, F Guéritaud, F Kassel**, *Convex cocompact actions in real projective geometry*, Ann. Sci. Éc. Norm. Supér. 57 (2024) 1753–1843 MR
- [8] **J Danciger, F Guéritaud, F Kassel, G-S Lee, L Marquis**, *Convex cocompactness for Coxeter groups*, J. Eur. Math. Soc. 27 (2025) 119–181 MR
- [9] **T tom Dieck**, *Algebraic topology*, Eur. Math. Soc., Zürich (2008) MR
- [10] **C Druţu, M Sapir**, *Tree-graded spaces and asymptotic cones of groups*, Topology 44 (2005) 959–1058 MR
- [11] **P de la Harpe**, *On Hilbert’s metric for simplices*, from “Geometric group theory, I” (G A Niblo, M A Roller, editors), London Math. Soc. Lecture Note Ser. 181, Cambridge Univ. Press (1993) 97–119 MR
- [12] **G C Hruska**, *Geometric invariants of spaces with isolated flats*, Topology 44 (2005) 441–458 MR

- [13] **G C Hruska, B Kleiner**, *Hadamard spaces with isolated flats*, *Geom. Topol.* 9 (2005) 1501–1538 MR
With an appendix joint with M Hindawi. Correction in 13 (2009) 699–707
- [14] **M Islam, A Zimmer**, *A flat torus theorem for convex co-compact actions of projective linear groups*, *J. Lond. Math. Soc.* 103 (2021) 470–489 MR
- [15] **M Islam, A Zimmer**, *Convex cocompact actions of relatively hyperbolic groups*, *Geom. Topol.* 27 (2023) 417–511 MR
- [16] **M Islam, A Zimmer**, *Convex co-compact groups with one-dimensional boundary faces*, *Groups Geom. Dyn.* 18 (2024) 1145–1183 MR
- [17] **M Islam, A Zimmer**, *Convex co-compact representations of 3-manifold groups*, *J. Topol.* 17 (2024) art. id. e12332 MR
- [18] **P Kelly, E Straus**, *Curvature in Hilbert geometries*, *Pacific J. Math.* 8 (1958) 119–125 MR
- [19] **G-S Lee, L Marquis**, *Discrete Coxeter groups*, from “Surveys in geometry II” (A Papadopoulos, editor), Springer (2024) 337–359 MR
- [20] **R D Nussbaum**, *Hilbert’s projective metric and iterated nonlinear maps*, *Mem. Amer. Math. Soc.* 391, Amer. Math. Soc., Providence, RI (1988) MR
- [21] **P Tukia**, *Conical limit points and uniform convergence groups*, *J. Reine Angew. Math.* 501 (1998) 71–98 MR
- [22] **C Vernicos**, *On the Hilbert geometry of convex polytopes*, from “Handbook of Hilbert geometry” (A Papadopoulos, M Troyanov, editors), IRMA Lect. Math. Theor. Phys. 22, Eur. Math. Soc., Zürich (2014) 111–125 MR
- [23] **T Weisman**, *Dynamical properties of convex cocompact actions in projective space*, *J. Topol.* 16 (2023) 990–1047 MR
- [24] **S Willard**, *General topology*, Dover, Mineola, NY (2004) MR
- [25] **D T Wise**, *Non-positively curved squared complexes: aperiodic tilings and non-residually finite groups*, PhD thesis, Princeton University (1996) MR Available at <https://www.proquest.com/docview/304259249>
- [26] **A Yaman**, *A topological characterisation of relatively hyperbolic groups*, *J. Reine Angew. Math.* 566 (2004) 41–89 MR
- [27] **A Zimmer**, *Projective Anosov representations, convex cocompact actions, and rigidity*, *J. Differential Geom.* 119 (2021) 513–586 MR
- [28] **A Zimmer**, *A higher-rank rigidity theorem for convex real projective manifolds*, *Geom. Topol.* 27 (2023) 2899–2936 MR

MI: *Mathematisches Institut
Heidelberg, Germany*

Current address: *Max Planck Institute for Mathematics in the Sciences
Leipzig, Germany*

AZ: *Department of Mathematics, University of Wisconsin–Madison
Madison, WI, United States*

mitul.islam@gmail.com, amzimmer2@wisc.edu

Received: 13 July 2023 Revised: 23 August 2024

ALGEBRAIC & GEOMETRIC TOPOLOGY

msp.org/agt

EDITORS

PRINCIPAL ACADEMIC EDITORS

John Etnyre
etnyre@math.gatech.edu
Georgia Institute of Technology

Kathryn Hess
kathryn.hess@epfl.ch
École Polytechnique Fédérale de Lausanne

BOARD OF EDITORS

Julie Bergner	University of Virginia jeb2md@eservices.virginia.edu	Thomas Koberda	University of Virginia thomas.koberda@virginia.edu
Steven Boyer	Université du Québec à Montréal cohf@math.rochester.edu	Markus Land	LMU München markus.land@math.lmu.de
Tara E Brendle	University of Glasgow tara.brendle@glasgow.ac.uk	Christine Lescop	Université Joseph Fourier lescop@ujf-grenoble.fr
Indira Chatterji	CNRS & Univ. Côte d'Azur (Nice) indira.chatterji@math.cnrs.fr	Norihiko Minami	Yamato University minami.norihiko@yamato-u.ac.jp
Octav Cornea	Université de Montreal cornea@dms.umontreal.ca	Andrés Navas	Universidad de Santiago de Chile andres.navas@usach.cl
Alexander Dranishnikov	University of Florida dranish@math.ufl.edu	Jessica S Purcell	Monash University jessica.purcell@monash.edu
Tobias Ekholm	Uppsala University, Sweden tobias.ekholm@math.uu.se	Birgit Richter	Universität Hamburg birgit.richter@uni-hamburg.de
Mario Eudave-Muñoz	Univ. Nacional Autónoma de México mario@matem.unam.mx	Jérôme Scherer	École Polytech. Féd. de Lausanne jerome.scherer@epfl.ch
David Futер	Temple University dfuter@temple.edu	Vesna Stojanoska	Univ. of Illinois at Urbana-Champaign vesna@illinois.edu
John Greenlees	University of Warwick john.greenlees@warwick.ac.uk	Zoltán Szabó	Princeton University szabo@math.princeton.edu
Matthew Hedden	Michigan State University mhedden@math.msu.edu	Maggy Tomova	University of Iowa maggy-tomova@uiowa.edu
Kristen Hendricks	Rutgers University kristen.hendricks@rutgers.edu	Daniel T Wise	McGill University, Canada daniel.wise@mcgill.ca
Hans-Werner Henn	Université Louis Pasteur henn@math.u-strasbg.fr	Lior Yanovski	Hebrew University of Jerusalem lior.yanovski@gmail.com
Daniel Isaksen	Wayne State University isaksen@math.wayne.edu		


See inside back cover or msp.org/agt for submission instructions.

The subscription price for 2025 is US \$760/year for the electronic version, and \$1110/year (+\$75, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Algebraic & Geometric Topology is indexed by Mathematical Reviews, Zentralblatt MATH, Current Mathematical Publications and the Science Citation Index.

Algebraic & Geometric Topology (ISSN 1472-2747 printed, 1472-2739 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004.

AGT peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<https://msp.org/>

© 2025 Mathematical Sciences Publishers

ALGEBRAIC & GEOMETRIC TOPOLOGY

Volume 25 Issue 9 (pages 5175–5754) 2025

Orbifolds, orbispaces and global homotopy theory	5175
BRANKO JURAN	
Higher symplectic capacities	5205
KYLER SIEGEL	
The cohomology of biquotients via a product on the two-sided bar construction	5279
JEFFREY D CARLSON	
Envelopes for algebraic patterns	5319
SHAUL BARKAN, RUNE HAUGSENG and JAN STEINEBRUNNER	
On the topology of character varieties of once-punctured torus bundles	5389
STEPHAN TILLMANN and YOUHENG YAO	
Segalification and the Boardman–Vogt tensor product	5439
SHAUL BARKAN and JAN STEINEBRUNNER	
Small cancellation and outer automorphisms of Kazhdan groups acting on hyperbolic spaces	5463
IONUȚ CHIFAN, ADRIAN IOANA, DENIS OSIN and BIN SUN	
The structure of relatively hyperbolic groups in convex real projective geometry	5503
MITUL ISLAM and ANDREW ZIMMER	
Characterizations of stability via Morse limit sets	5541
JACOB GARCIA	
Pre-Lie algebras with divided powers and the Deligne groupoid in positive characteristic	5567
MARVIN VERSTRAETE	
A note on rational maps with three branching points on the Riemann sphere	5607
ZHIQIANG WEI, YINGYI WU and BIN XU	
Loop homology of moment-angle complexes in the flag case	5619
FEDOR VYLEGZHANIN	
Algebraicity in monochromatic homotopy theory	5665
TORGEIR AAMBØ	
Twisted spectra revisited	5693
ALICE HEDENLUND and TASOS MOULINOS	