AGT

Algebraic & Geometric Topology

Volume 25 (2025)

A note on rational maps with three branching points on the Riemann sphere

ZHIQIANG WEI
YINGYI WU
BIN XU

msp

msp

Algebraic & Geometric Topology **25**:9 (2025) 5607–5617
DOI: 10.2140/agt.2025.25.5607
Published: 18 December 2025

A note on rational maps with three branching points on the Riemann sphere

ZHIQIANG WEI
YINGYI WU
BIN XU

Studying the existence of rational maps with given branching datum is a classical problem in the field of complex analysis and algebraic geometry. This problem dates back to Hurwitz and remains open to this day. We utilize complex analysis to establish a property of rational maps with 3 branching points on the Riemann sphere.

Given two compact Riemann surfaces M and N, a pair $(d, \mathcal{D})$ of an integer $d \geq 2$ and a collection $\mathcal{D}$ of nontrivial partitions of d is called a candidate branching datum if it satisfies the Riemann–Hurwitz formula. A candidate branching datum is considered exceptional when there is no holomorphic map realization for it. As applications of our main theorem, we present some new types of exceptional branching datum which cover some previously known results. We also deduce the realizability of a certain type of candidate branching datum on the Riemann sphere.

57M12

1 Introduction

Studying the existence of branched covers between two compact Riemann surfaces with given branching datum is an important problem in the field of complex analysis and algebraic geometry. This problem is commonly referred to as the Hurwitz existence problem. It asks whether, given two compact Riemann surfaces M and N, along with a collection $\mathcal{D}$ of partitions of a positive integer d, there exists a degree-d branched cover $f: M \to N$ with $\mathcal{D}$ as the branching datum.

Let M and N be compact Riemann surfaces, and let their respective Euler characteristics be denoted by $\chi(M)$ and $\chi(N)$. A smooth map $f: M \to N$ is a degree-d branched cover if for each $x \in N$ there is a partition $\mu(x) = [\alpha_1, \ldots, \alpha_r]$ (here the square brackets are used to denote an unordered set with repetitions) of d such that, over a neighborhood of x in N, f is equivalent to the map $\widetilde{f}: \{1, \ldots, r\} \times \mathbb{D} \to \mathbb{D}$ where $\widetilde{f}(j, z) = z^{\alpha_j}$ and x corresponds to 0 in the disc $\mathbb{D} = \{z \mid |z| < 1\} \subseteq \mathbb{C}$. The points $x \in N$ for which $\mu(x)$ is not the trivial partition $[1, 1, \ldots, 1]$ of d constitute the branching set B_f of f, and it is a finite set. The collection $\mathcal{D} = \{\mu(x) \mid x \in B_f\}$ (with repetitions allowed) is called the branching datum of f. As is well known, the degree-d, Euler characteristics $\chi(M)$, $\chi(N)$ and the branching datum $\mathcal{D}$ of f should satisfy the Riemann–Hurwitz formula

$$\nu(\mathcal{D}) = d \cdot \chi(N) - \chi(M), \tag{1}$$

© 2025 MSP (Mathematical Sciences Publishers). Distributed under the Creative Commons Attribution License 4.0 (CC BY). Open Access made possible by subscribing institutions via Subscribe to Open.

where $\nu(\mathcal{D})$ denotes the total branching of f, which is defined as follows. Let $B_f = \{x_1, \dots, x_n\} \subseteq N$ and $\mathcal{D} = \{[\alpha_1^1, \dots, \alpha_1^{r_1}], \dots, [\alpha_n^1, \dots, \alpha_n^{r_n}]\}$ represent the branching set and the branching datum of f, respectively. For each x_k, its preimage under f consists of a finite number of points $y_k^1, \dots, y_k^{r_k} \in M$, and near each y_k^j the map f is equivalent to $\widetilde{f}(z) = z^{\alpha_k^j}$. The integer α_k^j is usually referred to as the local degree or multiplicity of f at the point y_k^j. The total branching of f is then defined as $\nu(\mathcal{D}) = \sum_{k=1}^{n} \sum_{j=1}^{r_k} (\alpha_k^j - 1)$. Since

$$\sum_{j=1}^{r_k} \alpha_k^j = d \quad \text{for all } k = 1, \dots, n, \tag{2}$$

the Riemann–Hurwitz formula (1) can be expressed as

$$\sum_{k=1}^{n} (d - r_k) = d \cdot \chi(N) - \chi(M). \tag{3}$$

Given two compact Riemann surfaces M and N, a pair $(d, \mathcal{D})$ of an integer $d \geq 2$ and a collection $\mathcal{D}$ of nontrivial partitions of d is called a candidate branching datum if it satisfies the Riemann–Hurwitz formula. In his classical work [7], Hurwitz reduced the Hurwitz problem to a problem involving partitions realized by suitable permutations in symmetric groups. In [5], Edmonds, Kulkarni, and Stong proved that all candidate branching datum is realizable when $\chi(N) \leq 0$. However, when $N = S^2$ the problem becomes much more complex. It is well known that there exist candidate branching datum $(d, \mathcal{D})$ that cannot be realized by a branched cover, hence called exceptional. For example, this is the case for $d = 4$ and $\mathcal{D} = \{[3, 1], [2, 2], [2, 2]\}$. Characterizing all of such exceptional candidate branching datum remains an open problem to this day. In [25], Zheng determined all exceptional candidate branching datum with 3 branching points and degree $d \leq 22$ by computer for the cases where $M = N = S^2$ and $M = T^2$, $N = S^2$.

Finding new types of exceptional candidate branching datum is of interest as it may provide insights towards establishing a universal criterion. However, the general pattern of realizable data remains unclear. Various approaches such as dessins d'enfant, Speiser graphs and the monodromy approach have been explored to attack the problem. For more details, see [1; 2; 4; 5; 6; 8; 9; 11; 12; 13; 14; 15; 16; 17; 18; 19; 20; 21; 22; 24; 25]. Particularly, we refer the reader to [4; 19; 21] for a review of available results and techniques. In [5], a conjecture proposing connections with number-theoretic facts has been put forward, which is supported by strong evidence in [15; 16].

Conjecture 1.1 (prime-degree conjecture) Suppose $(d, \mathcal{D})$ is a candidate branching datum. If d is a prime, then $(d, \mathcal{D})$ is realizable.

In [5], Edmonds, Kulkarni, and Stong reduced the prime-degree conjecture to the case of candidate branching datum with exactly 3 partitions. Up to now, this conjecture is still open. In this paper we investigate the properties of branched covers for the cases where $M = N = S^2$. Regarding $S^2 \cong \overline{\mathbb{C}}$ as the Riemann sphere, by Stoïlow's theorem (see [23] or [10]), the Hurwitz problem reduces to the

existence of rational maps on $\overline{\mathbb{C}}$ with the given branching datum. Thus we can use complex analysis techniques to attack the problem. Motivation by the recent work of Song and Xu [22], we use their idea of expressing a rational map as a composition of a rational map and a power map. Our main result is as follows and can be viewed as a generalization of a result in [18, Theorems 1.4 and 1.5].

Theorem 1.2 *Suppose $f:\overline{\mathbb{C}}\to\overline{\mathbb{C}}$ is a rational map of degree $d=rk$, $r\geq 2$, $k\geq 2$, with 3 branching points and associated branching datum*

$$\{\mu_1=[\alpha_1,\dots,\alpha_A],\mu_2=[rx_1,\dots,rx_B],\mu_3=[ry_1,\dots,ry_C]\}.$$

Then, up to two Möbius transformations on $\overline{\mathbb{C}}$, there is a rational map $g:\overline{\mathbb{C}}\to\overline{\mathbb{C}}$ of degree k such that $f=g^r$, with the property that its branching datum consists of $[x_1,\dots,x_B]$ and $[y_1,\dots,y_C]$, together with data arising by splitting the partition of $[\alpha_1,\dots,\alpha_A]$ into r partitions of k.

Remark In Theorem 1.2, the sentence "up to two Möbius transformations" means that $f=\varphi\circ g^k\circ\phi$, where $\varphi,\phi:\overline{\mathbb{C}}\to\overline{\mathbb{C}}$ are two Möbius transformations.

Remark In Theorem 1.2, if $x_1=\cdots=x_B=1$ or $y_1=\cdots=y_C=1$ then the corresponding partition of k should be expunged from the branching datum of g.

Then, under the assumption of Theorem 1.2, one would have immediate corollaries for realizable data.

Corollary 1.3 *Each $\alpha_j\leq k$.*

Corollary 1.4 *If $A=2$, then $r=2$, $\mu_1=[k,k]$.*

Corollary 1.5 *If $A=r$, then $\mu_1=[k,\dots,k]$.*

As applications of these results, we offer new proofs of some of the results presented in [18; 26] and give some new type of exceptional candidate branching datum (Propositions 1.9 and 1.10).

Proposition 1.6 [26] *Suppose $d=2k$ with $k\geq 2$. Then a candidate branching datum of the form*

$$(d,\{[k_1,k_2],[\underbrace{2,\dots,2}_{k}],[\underbrace{2,\dots,2}_{k}]\})$$

is exceptional if $k_1\neq k_2$. Suppose in addition $k\geq 3$. Then a candidate branching datum of the form

$$(d,\{[\underbrace{2,\dots,2}_{k}],[\underbrace{2,\dots,2}_{j_1},2k-2j_1],[\underbrace{2,\dots,2}_{j_2},2k-2j_2]\})$$

is exceptional if $j_1\neq j_2$.

Proof Since $k_1\neq k_2$ and $k_1+k_2=2k$, $k_1,k_2\neq k$. By Corollary 1.4, $(d,\mathcal{D})$ is exceptional.

Suppose $k\geq 3$, and note that $j_1+j_2=k$ and $j_1\neq j_2$. Then, $2j_1<k$ or $2j_2<k$. If $2j_1<k$, then $2k-2j_1>k$; If $2j_2<k$, then $2k-2j_2>k$. Thus, by Corollary 1.3, $(d,\mathcal{D})$ is exceptional. □

Proposition 1.7 [26] *Suppose $d = 3k$ with $k \geq 3$. Then a candidate branching datum of the form*

$$(d, \{[k-2, \underbrace{2, \dots, 2}_{k+1}], [\underbrace{3, \dots, 3}_{k}], [\underbrace{3, \dots, 3}_{k}]\})$$

is exceptional if k is odd.

Proof Since k is odd and 2 is even, there is just one odd in $[k-2, 2, \dots, 2]$. Thus $[k-2, 2, \dots, 2]$ cannot be split into 3 partitions of k. Therefore, by Theorem 1.2, $(d, \mathcal{D})$ is exceptional. □

Proposition 1.8 [26] *Suppose $d = rk$ with $r \geq 2$, $k \geq 2$. Then a candidate branching datum of the form*

$$(d, \{[j_1, j_2, \underbrace{1, \dots, 1}_{(r-2)k}], [\underbrace{r, \dots, r}_{k}], [\underbrace{r, \dots, r}_{k}]\})$$

is exceptional if $j_1 \neq j_2$.

Proof Since $j_1 + j_2 = 2k$ and $j_1 \neq j_2$, it follows that either $j_1 > k$ or $j_2 > k$. Thus, by Corollary 1.3, $(d, \mathcal{D})$ is exceptional. □

Proposition 1.9 *Suppose $d = 3k$ with $k \geq 3$ and $k \equiv 1 \bmod 2$. Then a candidate branching datum of the form*

$$(d, \{[j_1, j_2, \underbrace{2, \dots, 2}_{k}], [\underbrace{3, \dots, 3}_{k}], [\underbrace{3, \dots, 3}_{k}]\})$$

is exceptional if j_1, j_2 have different parity.

Proof Since j_1, j_2 have different parity, there is just one odd in the partition

$$[j_1, j_2, \underbrace{2, \dots, 2}_{k}].$$

Since k is odd, $[k-2, 2, \dots, 2]$ cannot be split into 3 partitions of k. Thus by Theorem 1.2, $(d, \mathcal{D})$ is exceptional. □

Proposition 1.10 *Suppose $d = 3k$ with $k = 2 + 3l$, $l \geq 1$. Then a candidate branching datum of the form*

$$\big(d, \{[\underbrace{3, \dots, 3}_{k-1}, \underbrace{1, 1, 1}_{3}], [\underbrace{3, \dots, 3}_{k}], [\underbrace{3, \dots, 3}_{k}]\}\big)$$

is exceptional.

Proof Since $k = 2 + 3l \geq 5$ and the number of 1's in

$$[\underbrace{3, \dots, 3}_{k-1}, \underbrace{1, 1, 1}_{3}]$$

equals 3,

$$[\underbrace{3, \dots, 3}_{k-1}, \underbrace{1, 1, 1}_{3}]$$

cannot be split into 3 partitions of k. Thus by Theorem 1.2, $(d, \mathcal{D})$ is exceptional. □

From the constructed exceptional datum mentioned above, we obtain an additional result regarding nonprime degrees on the case $M = N = S^2$, which has been proven in [5; 26] using different methods.

Corollary 1.11 [5; 26] *For every nonprime d, there exists at least one candidate branching datum $(d, \mathcal{D})$ that is exceptional.*

2 Proof of Theorem 1.2

The key to proving Theorem 1.2 is to use suitable Möbius transformations to simplify the representation of f. This simplification then facilitates the construction of the mapping g, derived from the transformed expression. The final step involves calculating the branching datum of g. To begin, we establish the following lemma.

Lemma 2.1 *Suppose $g : \mathbb{C} \to \mathbb{C}$ is a nonconstant holomorphic function. For a point $z_0 \in \mathbb{C}$, if $g(z_0) \neq 0$, then g and g^r, for every integer r, have the same local degree at z_0.*

Proof Call n the local degree of g at z_0. Then, knowing that $g(z_0) \neq 0$, the order-n Taylor expansion of g at z_0 has the form

$$g(z) = a + b(z - z_0)^n + o((z - z_0)^n),$$

with $a, b \neq 0$. It follows that

$$g^r(z) = a^r + rab(z - z_0)^n + o((z - z_0)^n),$$

and $rab \neq 0$, so g^r also has local degree n at z_0. □

Now, the proof of Theorem 1.2 is as follows.

Since Möbius transformations act in a triply transitive fashion on $\overline{\mathbb{C}}$, one can assume that:

(i) 0 in the target $\overline{\mathbb{C}}$ is the branching point corresponding to the partition $[rx_1, \dots, rx_B]$.

(ii) ∞ in the target $\overline{\mathbb{C}}$ is the branching point corresponding to the partition $[ry_1, \dots, ry_C]$.

(iii) 1 in the target $\overline{\mathbb{C}}$ is not the branching point, and $f(\infty) = 1$.

Thus, the expression for f can be written as

$$f(z) = \frac{(z - z_1)^{rx_1}(z - z_2)^{rx_2} \cdots (z - z_B)^{rx_B}}{(z - w_1)^{ry_1}(z - w_2)^{ry_2} \cdots (z - w_C)^{ry_C}},$$

where $z_1, \dots, z_B, w_1, \dots, w_C \in \mathbb{C}$ are distinct complex numbers.

Define

$$g(z) = \frac{(z - z_1)^{x_1}(z - z_2)^{x_2} \cdots (z - z_B)^{x_B}}{(z - w_1)^{y_1}(z - w_2)^{y_2} \cdots (z - w_C)^{y_C}}.$$

Then g is a rational map with degree k, and $f = g^r$ on $\overline{\mathbb{C}}$. It is evident that 0 is a branching point of g if and only if $x_j \geq 2$ for some index j, and ∞ is a branching point of g if and only if $y_l \geq 2$ for some index l.

For any $z_0 \in \mathbb{C} \setminus \{z_1, \dots, z_B, w_1, \dots, w_C\}$, by Lemma 2.1, g and $f = g^r$ have the same local degree at z_0. Consequently, the branching datum of g is

$$\{[\alpha_1^1, \dots, \alpha_1^{l_1}], \dots, [\alpha_s^1, \dots, \alpha_s^{l_s}], [x_1, \dots, x_B], [y_1, \dots, y_C]\},$$

for some $1 \leq s \leq r$, where $\alpha_1^1, \dots, \alpha_1^{l_1}, \dots, \alpha_s^1, \dots, \alpha_s^{l_s}$ belong to $\alpha_1, \dots, \alpha_A$ which means that up to a permutation $(\alpha_1^1, \dots, \alpha_1^{l_1}, \dots, \alpha_s^1, \dots, \alpha_s^{l_s}, 1, \dots, 1) = (\alpha_1, \dots, \alpha_A)$.

In summary, we have completed the proof of the Theorem 1.2.

By Lemma 2.1, the proof of the following theorem is easy.

Theorem 2.2 *Suppose $g : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ is a rational map with degree $d \geq 3$ and branching datum*

$$\{[\alpha_1, \dots, \alpha_A], [\beta_1, \dots, \beta_B], [\gamma_1, \dots, \gamma_C]\}.$$

Then, for any integer $k \geq 2$, there is a Möbius transformation φ such that $f = (\varphi \circ g)^k$ is a rational map with degree kd and branching datum

$$\{[\alpha_1, \dots, \alpha_A, \underbrace{1, \dots, 1}_{(k-1)d}], [k\beta_1, \dots, k\beta_B], [k\gamma_1, \dots, k\gamma_C]\}.$$

3 Some existence results

We conclude the paper by presenting some existence results for a certain type of rational maps on $\overline{\mathbb{C}}$ with 3 branching points. First, we give a new proof of the following theorem which was proved in [5] and which is a special case in [4, Theorem 3.5].

Theorem 3.1 [5] *There exists a rational map $f : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ with branching datum*

$$\{[\alpha_1, \alpha_2], [\underbrace{2, \dots, 2}_{k}], [\underbrace{2, \dots, 2}_{k}]\},$$

where $k \geq 2$ and $\alpha_1 + \alpha_2 = 2k$, if and only if $\alpha_1 = \alpha_2 = k$.

Proof Necessity follows from Corollary 1.4. The proof of sufficiency is as follows.

Define $g(z) = (z^k - 1)/(z^k + 1)$. By direct calculation, the branching points of g are -1 and 1. Additionally, $g^{-1}(-1) = \{0\}$, $g^{-1}(1) = \{\infty\}$ and the branching datum of g is $\{[k], [k]\}$. Set $f = g^2$. By direct calculation, we find that the branching points of f are 1, 0, ∞, and the branching datum of f is

$$\{[k, k], [\underbrace{2, \dots, 2}_{k}], [\underbrace{2, \dots, 2}_{k}]\}. \qquad \square$$

Second, to give more existence results, we recall a result by Boccara [3].

Theorem 3.2 [3] *Let p and q be two positive integers. Consider a candidate branching datum*

$$\mathcal{D} = \{[\alpha_1, \dots, \alpha_p], [\beta_1, \dots, \beta_q], [p+q-1, 1, \dots, 1]\},$$

which contains 3 partitions of a positive integer d. Then there exists a rational map $f : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ with $\mathcal{D}$ being its branching datum if and only if

$$p + q - 1 \leq \frac{d}{\mathrm{GCD}(\alpha_1, \dots, \alpha_p, \beta_1, \dots, \beta_q)}.$$

Theorem 3.3 *Suppose that $k \geq 3$. There exists a degree-$2k$ rational map $f : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ with branching datum*

$$\{[\alpha_1, \alpha_2, \alpha_3], [\underbrace{2, \dots, 2}_{k}], [\underbrace{2, \dots, 2}_{k-2}, 4]\}$$

if and only if one of the α_j's is k.

Proof **Sufficiency** Without loss of generality, we can suppose that $\alpha_1 = k$. By Theorem 3.2, there exists a rational map $g : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ with branching datum

$$\{[k], [\alpha_2, \alpha_3], [\underbrace{1, \dots, 1}_{k-2}, 2]\}.$$

Using a suitable Möbius transformation, we may suppose that:

(i) -1 is the branching point of g corresponding to the partition $[k]$.

(ii) 1 is the branching point of g corresponding to the partition $[\alpha_2, \alpha_3]$.

(iii) 0 is the branching point of g corresponding to the partition $[\underbrace{1, \dots, 1}_{k-2}, 2]$.

Then $f = g^2$ is the desired function.

Necessity Suppose $f : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ is a rational map with branching datum

$$\{[\alpha_1, \alpha_2, \alpha_3], [\underbrace{2, \dots, 2}_{k}], [\underbrace{2, \dots, 2}_{k-2}, 4]\}.$$

Then, by Theorem 1.2, $[\alpha_1, \alpha_2, \alpha_3]$ can be split into 2 partitions of k. Thus one of the α_j's is k. □

In general, we obtain the following theorem.

Theorem 3.4 *Suppose that $k \geq 3$ and $x \geq 1$. There exists a degree-$2k$ rational map $f : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ with branching datum*

$$\{[\alpha_1, \alpha_2, \dots, \alpha_{x+1}], [\underbrace{2, \dots, 2}_{k}], [\underbrace{2, \dots, 2}_{k-x}, 2x]\}$$

if and only if $[\alpha_1, \alpha_2, \dots, \alpha_{x+1}]$ can be split into two partitions of k and $k/\mathrm{GCD}(\alpha_1, \alpha_2, \dots, \alpha_{x+1}) \geq x$.

Proof **Necessity** Suppose $f:\overline{\mathbb{C}}\to\overline{\mathbb{C}}$ is a rational map with branching datum

$$\{[\alpha_1,\alpha_2,\dots,\alpha_{x+1}],[\underbrace{2,\dots,2}_{k}],[\underbrace{2,\dots,2}_{k-x},2x]\}.$$

Then, by Theorem 1.2, there exists a rational map g of degree k with branching datum

$$\{[\alpha_1^1,\dots,\alpha_1^{r_1}],[\alpha_2^1,\dots,\alpha_2^{r_2}],[1,\dots,1,x]\}$$

such that $f=g^2$, where $[\alpha_1^1,\dots,\alpha_1^{r_1},\alpha_2^1,\dots,\alpha_2^{r_2}]=[\alpha_1,\alpha_2,\dots,\alpha_{x+1}]$. By Theorem 3.2,

$$\frac{k}{\mathrm{GCD}(\alpha_1,\alpha_2,\dots,\alpha_{x+1})}\geq x.$$

Sufficiency Suppose $[\alpha_1,\alpha_2,\dots,\alpha_{x+1}]$ can be split into two partitions of k. Denote the two partitions by $[\alpha_1^1,\dots,\alpha_1^{r_1}]$ and $[\alpha_2^1,\dots,\alpha_2^{r_2}]$. Since

$$\mathrm{GCD}(\alpha_1^1,\dots,\alpha_1^{r_1},\alpha_2^1,\dots,\alpha_2^{r_2})=\mathrm{GCD}(\alpha_1,\alpha_2,\dots,\alpha_{x+1}),$$

we have

$$\frac{k}{\mathrm{GCD}(\alpha_1^1,\dots,\alpha_1^{r_1},\alpha_2^1,\dots,\alpha_2^{r_2})}\geq x.$$

Thus by a result of Boccara [3], there exists a rational map $g:\overline{\mathbb{C}}\to\overline{\mathbb{C}}$ with branching datum

$$\{[\alpha_1^1,\dots,\alpha_1^{r_1}],[\alpha_2^1,\dots,\alpha_2^{r_2}],[1,\dots,1,x]\}.$$

Using a suitable Möbius transformation, we may suppose that:

(i) -1 is the branching point of g corresponding to the partition $[\alpha_1^1,\dots,\alpha_1^{r_1}]$.

(ii) 1 is the branching point of g corresponding to the partition $[\alpha_2^1,\dots,\alpha_2^{r_2}]$.

(iii) 0 is the branching point of g corresponding to the partition $[1,\dots,1,x]$.

Then $f=g^2$ is the desired function. □

Finally, we provide more explicit examples to explain our results.

Example 3.5 There exists a rational map $f:\overline{\mathbb{C}}\to\overline{\mathbb{C}}$ with branched data

$$\{[\alpha_1,\alpha_2,\alpha_3,2,2],[3,3,3],[3,3,3]\},$$

where $\alpha_1+\alpha_2+\alpha_3=5$ and $\alpha_1\geq\alpha_2\geq\alpha_3$, if and only if $\alpha_1=3$.

Proof **Necessity** Since $\alpha_1+\alpha_2+\alpha_3=5$ and $\alpha_1\geq\alpha_2\geq\alpha_3$, it follows that $\alpha_1=2$ or 3. If $\alpha_1=2$, then $\alpha_2=2$ and $\alpha_3=1$. But $[\alpha_1,\alpha_2,\alpha_3,2,2]=[2,2,1,2,2]$ cannot be split into 3 partitions of 3. Thus $\alpha_1=3$.

Sufficiency If $\alpha_1=3$, then $\alpha_2=\alpha_3=1$. By Theorem 3.2, there exists a rational map g with degree 3 and branching datum

$$\{[3],[2,1],[2,1]\}.$$

Using a suitable Möbius transformation, we may suppose that:

(i) 1 is the branching point of g corresponding to the partition $[3]$.

(ii) $e^{\frac{2\pi}{3}i}$ is the branching point of g corresponding to the partition $[2,1]$.

(iii) $e^{\frac{4\pi}{3}i}$ is the branching point of g corresponding to the partition $[2,1]$.

Then $f = g^3$ is a rational map with branching datum

$$\{[3,1,1,2,2],[3,3,3],[3,3,3]\}. \qquad \square$$

Example 3.6 There exists a rational map $f:\overline{\mathbb{C}}\to\overline{\mathbb{C}}$ with branching datum

$$\{[\alpha_1,\alpha_2,\alpha_3,2,2,2,2],[3,3,3,3,3],[3,3,3,3,3]\},$$

where $\alpha_1+\alpha_2+\alpha_3=7$ and $\alpha_1\geq\alpha_2\geq\alpha_3$, if and only if $\alpha_1=5$ or $\alpha_1=\alpha_2=3$.

Proof **Necessity** Since $\alpha_1+\alpha_2+\alpha_3=7$ and $\alpha_1\geq\alpha_2\geq\alpha_3$, it follows that $\alpha_1=3,4$ or 5.

When $\alpha_1=3$, it follows that $(\alpha_2,\alpha_3)=(2,2)$ or $(3,1)$. When $\alpha_1=3,\alpha_2=2$ and $\alpha_3=2$, since $[3,2,2,2,2,2,2]$ cannot be split into 3 partitions of 5, we exclude this case. When $\alpha_1=3,\alpha_2=3$ and $\alpha_3=1$, it is evident that $[3,3,1,2,2,2,2]$ can be split into 3 partitions of 5.

When $\alpha_1=4$, since $\alpha_2+\alpha_3=3$, there is just one odd in $[4,\alpha_2,\alpha_3,2,2,2,2]$. Thus $[4,\alpha_2,\alpha_3,2,2,2,2]$ cannot be split into 3 partitions of 5. Thus we exclude this case.

When $\alpha_1=5$, it follows that $\alpha_2=\alpha_3=1$. It is evident that $[5,1,1,2,2,2,2]$ can be split into 3 partitions of 5.

In summary, $\alpha_1=5$ or $\alpha_1=\alpha_2=3$.

Sufficiency When $\alpha_1=5$, it follows that $\alpha_2=\alpha_3=1$. By Theorem 3.2, there exists a rational map g with branching datum

$$\{[5],[2,2,1],[2,2,1]\}.$$

Using a suitable Möbius transformation, we may suppose that:

(i) 1 is the branching point of g corresponding to the partition $[5]$.

(ii) $e^{\frac{2\pi}{3}i}$ is the branching point of g corresponding to the partition $[2,2,1]$.

(iii) $e^{\frac{4\pi}{3}i}$ is the branching point of g corresponding to the partition $[2,2,1]$.

Then $f=g^3$ is a rational map with branching datum

$$\{[5,1,1,2,2,2,2],[3,3,3,3,3],[3,3,3,3,3]\}.$$

When $\alpha_1=\alpha_2=3$, it follows that $\alpha_3=1$. By a result of Zheng [25] there exists a rational map g with branching datum

$$\{[3,2],[3,2],[2,2,1]\}.$$

Using a suitable Möbius transformation, we may suppose that:

(i) 1 is the branching point of g corresponding to the partition $[3, 2]$.

(ii) $e^{\frac{2\pi}{3}i}$ is the branching point of g corresponding to the partition $[3, 2]$.

(iii) $e^{\frac{4\pi}{3}i}$ is the branching point of g corresponding to the partition $[2, 2, 1]$.

Then $f = g^3$ is a rational map with branching datum

$$\{[3, 3, 1, 2, 2, 2, 2], [3, 3, 3, 3, 3, 3], [3, 3, 3, 3, 3, 3]\}. \qquad \square$$

Acknowledgments

We would like to express our gratitude to the reviewers for constructive comments and suggestions, which have helped to improve the quality and clarity of this article. This work was initiated during Wei's visit to the Chern Institute of Mathematics in the fall of 2023. Wei would like to express his gratitude to Professor Zizhou Tang for the warm hospitality he received. Wei is partially supported by the National Natural Science Foundation of China (Grant No. 12171140). Wu is partially supported by the Fundamental Research Funds for the Central Universities and the CAS Project for Young Scientists in Basic Research (Grant No. YSBR-001). Xu is supported in part by the National Natural Science Foundation of China (Grant Nos. 12271495, 11971450, and 12071449) and the CAS Project for Young Scientists in Basic Research (Grant No. YSBR-001).

References

[1] **K Barański**, *On realizability of branched coverings of the sphere*, Topology Appl. 116 (2001) 279–291 MR

[2] **F Baroni**, **C Petronio**, *Solution of the Hurwitz problem with a length-2 partition*, Illinois J. Math. 68 (2024) 479–511 MR

[3] **G Boccara**, *Cycles comme produit de deux permutations de classes données*, Discrete Math. 38 (1982) 129–142 MR

[4] **W Cui**, **J Hu**, *Totally ramified rational maps*, Conform. Geom. Dyn. 26 (2022) 208–234 MR

[5] **A L Edmonds**, **R S Kulkarni**, **R E Stong**, *Realizability of branched coverings of surfaces*, Trans. Amer. Math. Soc. 282 (1984) 773–790 MR

[6] **S M Gersten**, *On branched covers of the 2-sphere by the 2-sphere*, Proc. Amer. Math. Soc. 101 (1987) 761–766 MR

[7] **A Hurwitz**, *Ueber Riemann'sche Flächen mit gegebenen Verzweigungspunkten*, Math. Ann. 39 (1891) 1–60 MR

[8] **D H Husemoller**, *Ramified coverings of Riemann surfaces*, Duke Math. J. 29 (1962) 167–174 MR

[9] **A G Khovanskii**, **S Zdravkovska**, *Branched covers of S^2 and braid groups*, J. Knot Theory Ramifications 5 (1996) 55–75 MR

[10] **R Luisto**, **P Pankka**, *Stoïlow's theorem revisited*, Expo. Math. 38 (2020) 303–318 MR

[11] **A D Mednykh**, *Nonequivalent coverings of Riemann surfaces with a prescribed ramification type*, Sibirsk. Mat. Zh. 25 (1984) 120–142 MR In Russian; translated in Sib. Math. J. 25 (1984) 606–625

[12] **A D Mednykh**, *Branched coverings of Riemann surfaces whose branch orders coincide with the multiplicity*, Comm. Algebra 18 (1990) 1517–1533 MR

[13] **S Monni**, **J S Song**, **Y S Song**, *The Hurwitz enumeration problem of branched covers and Hodge integrals*, J. Geom. Phys. 50 (2004) 223–256 MR

[14] **A Okounkov**, **R Pandharipande**, *Gromov–Witten theory, Hurwitz theory, and completed cycles*, Ann. of Math. 163 (2006) 517–560 MR

[15] **F Pakovich**, *Solution of the Hurwitz problem for Laurent polynomials*, J. Knot Theory Ramifications 18 (2009) 271–302 MR

[16] **M A Pascali**, **C Petronio**, *Surface branched covers and geometric 2-orbifolds*, Trans. Amer. Math. Soc. 361 (2009) 5885–5920 MR

[17] **M A Pascali**, **C Petronio**, *Branched covers of the sphere and the prime-degree conjecture*, Ann. Mat. Pura Appl. 191 (2012) 563–594 MR

[18] **E Pervova**, **C Petronio**, *On the existence of branched coverings between surfaces with prescribed branch data, I*, Algebr. Geom. Topol. 6 (2006) 1957–1985 MR

[19] **E Pervova**, **C Petronio**, *On the existence of branched coverings between surfaces with prescribed branch data, II*, J. Knot Theory Ramifications 17 (2008) 787–816 MR

[20] **E Pervova**, **C Petronio**, *Realizability and exceptionality of candidate surface branched covers: methods and results*, from "Geometry Seminars" (S Coen, editor), Univ. Stud. Bologna, Bologna (2010) 105–120 MR

[21] **C Petronio**, *The Hurwitz existence problem for surface branched covers*, Winter Braids Lect. Notes 7 (2020) art. id. 2 MR

[22] **J Song**, **B Xu**, *On rational functions with more than three branch points*, Algebra Colloq. 27 (2020) 231–246 MR

[23] **S Stoïlow**, *Sur les transformations continues et la topologie des fonctions analytiques*, Ann. Sci. École Norm. Sup. 45 (1928) 347–382 MR

[24] **R Thom**, *L'équivalence d'une fonction différentiable et d'un polynome*, Topology 3 (1965) 297–307 MR

[25] **H Zheng**, *Realizability of branched coverings of S^2*, Topology Appl. 153 (2006) 2124–2134 MR

[26] **X Zhu**, *Spherical conic metrics and realizability of branched covers*, Proc. Amer. Math. Soc. 147 (2019) 1805–1815 MR

ZW: *School of Mathematics and Statistics, Henan University*
Kaifeng, China
Center for Applied Mathematics of Henan Province, Henan University
Zhengzhou, China
YW: *School of Mathematical Sciences, University of Chinese Academy of Sciences*
Beijing, China
BX: *CAS Wu Wen-Tsun Key Laboratory of Mathematics and School of Mathematical Sciences*
University of Science and Technology of China
Hefei, China
weizhiqiang15@mails.ucas.edu.cn, wuyy@ucas.ac.cn, bxu@ustc.edu.cn

Received: 23 January 2024 Revised: 3 September 2024

ALGEBRAIC & GEOMETRIC TOPOLOGY

msp.org/agt

EDITORS

PRINCIPAL ACADEMIC EDITORS

John Etnyre
etnyre@math.gatech.edu
Georgia Institute of Technology

Kathryn Hess
kathryn.hess@epfl.ch
École Polytechnique Fédérale de Lausanne

BOARD OF EDITORS

Julie Bergner	University of Virginia jeb2md@eservices.virginia.edu	Thomas Koberda	University of Virginia thomas.koberda@virginia.edu
Steven Boyer	Université du Québec à Montréal cohf@math.rochester.edu	Markus Land	LMU München markus.land@math.lmu.de
Tara E Brendle	University of Glasgow tara.brendle@glasgow.ac.uk	Christine Lescop	Université Joseph Fourier lescop@ujf-grenoble.fr
Indira Chatterji	CNRS & Univ. Côte d'Azur (Nice) indira.chatterji@math.cnrs.fr	Norihiko Minami	Yamato University minami.norihiko@yamato-u.ac.jp
Octav Cornea	Université' de Montreal cornea@dms.umontreal.ca	Andrés Navas	Universidad de Santiago de Chile andres.navas@usach.cl
Alexander Dranishnikov	University of Florida dranish@math.ufl.edu	Jessica S Purcell	Monash University jessica.purcell@monash.edu
Tobias Ekholm	Uppsala University, Sweden tobias.ekholm@math.uu.se	Birgit Richter	Universität Hamburg birgit.richter@uni-hamburg.de
Mario Eudave-Muñoz	Univ. Nacional Autónoma de México mario@matem.unam.mx	Jérôme Scherer	École Polytech. Féd. de Lausanne jerome.scherer@epfl.ch
David Futer	Temple University dfuter@temple.edu	Vesna Stojanoska	Univ. of Illinois at Urbana-Champaign vesna@illinois.edu
John Greenlees	University of Warwick john.greenlees@warwick.ac.uk	Zoltán Szabó	Princeton University szabo@math.princeton.edu
Matthew Hedden	Michigan State University mhedden@math.msu.edu	Maggy Tomova	University of Iowa maggy-tomova@uiowa.edu
Kristen Hendricks	Rutgers University kristen.hendricks@rutgers.edu	Daniel T Wise	McGill University, Canada daniel.wise@mcgill.ca
Hans-Werner Henn	Université Louis Pasteur henn@math.u-strasbg.fr	Lior Yanovski	Hebrew University of Jerusalem lior.yanovski@gmail.com
Daniel Isaksen	Wayne State University isaksen@math.wayne.edu		

See inside back cover or msp.org/agt for submission instructions.

The subscription price for 2025 is US $760/year for the electronic version, and $1110/year (+$75, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Algebraic & Geometric Topology is indexed by Mathematical Reviews, Zentralblatt MATH, Current Mathematical Publications and the Science Citation Index.

Algebraic & Geometric Topology (ISSN 1472-2747 printed, 1472-2739 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004.

AGT peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

mathematical sciences publishers

nonprofit scientific publishing

https://msp.org/

© 2025 Mathematical Sciences Publishers

ALGEBRAIC & GEOMETRIC TOPOLOGY

Volume 25 Issue 9 (pages 5175–5754) 2025

Orbifolds, orbispaces and global homotopy theory 5175

BRANKO JURAN

Higher symplectic capacities 5205

KYLER SIEGEL

The cohomology of biquotients via a product on the two-sided bar construction 5279

JEFFREY D CARLSON

Envelopes for algebraic patterns 5319

SHAUL BARKAN, RUNE HAUGSENG and JAN STEINEBRUNNER

On the topology of character varieties of once-punctured torus bundles 5389

STEPHAN TILLMANN and YOUHENG YAO

Segalification and the Boardman–Vogt tensor product 5439

SHAUL BARKAN and JAN STEINEBRUNNER

Small cancellation and outer automorphisms of Kazhdan groups acting on hyperbolic spaces 5463

IONUȚ CHIFAN, ADRIAN IOANA, DENIS OSIN and BIN SUN

The structure of relatively hyperbolic groups in convex real projective geometry 5503

MITUL ISLAM and ANDREW ZIMMER

Characterizations of stability via Morse limit sets 5541

JACOB GARCIA

Pre-Lie algebras with divided powers and the Deligne groupoid in positive characteristic 5567

MARVIN VERSTRAETE

A note on rational maps with three branching points on the Riemann sphere 5607

ZHIQIANG WEI, YINGYI WU and BIN XU

Loop homology of moment-angle complexes in the flag case 5619

FEDOR VYLEGZHANIN

Algebraicity in monochromatic homotopy theory 5665

TORGEIR AAMBØ

Twisted spectra revisited 5693

ALICE HEDENLUND and TASOS MOULINOS