

AG  
T

*Algebraic & Geometric  
Topology*

Volume 26 (2026)

**Kontsevich's characteristic classes as topological invariants  
of configuration space bundles**

XUJIA CHEN





# Kontsevich’s characteristic classes as topological invariants of configuration space bundles

XUJIA CHEN

Kontsevich’s characteristic classes are invariants of framed smooth fiber bundles with homology sphere fibers. It was shown by Watanabe that they can be used to distinguish smooth  $S^4$ -bundles that are all trivial as topological fiber bundles. In this article we show that this ability of Kontsevich’s classes is a manifestation of the following principle: the “real blow-up” construction on a smooth manifold essentially depends on its smooth structure and thus, given a smooth manifold (or smooth fiber bundle)  $M$ , the topological invariants of spaces constructed from  $M$  by real blow-ups could potentially differentiate smooth structures on  $M$ . The main theorem says that Kontsevich’s characteristic classes of a smooth framed bundle  $\pi$  are determined by the topology of the 2-point configuration space bundle of  $\pi$  and framing data.

1. Introduction	349
2. Information from the graph	357
3. Defining various spaces, all having a $\mathcal{G}$ -action	360
4. Reconstructing Kontsevich’s characteristic classes	373
5. Equivalence with the original definition	382
6. Some remarks about the condition in Theorem 1.2	389
Acknowledgements	394
References	395

## 1 Introduction

Given the following data:

- a smooth fiber bundle  $E \xrightarrow{\pi} B$  whose fibers are homology spheres,
- a smooth section  $s_\infty : B \rightarrow E$  and a trivialization  $t$  of  $\pi$  in a neighborhood  $U$  of  $s_\infty(B)$ ,
- a vertical framing  $F$  on  $E - s_\infty(B)$  which is standard (i.e., looks like the standard framing on  $\mathbb{R}^n$  near  $\infty$ ) with respect to  $t$  in  $U$ ,

Kontsevich’s characteristic classes are a collection of cohomology classes in  $H^*(B; \mathbb{R})$ , parameterized by some combinatorial data (“graph homology”). They were introduced by Kontsevich [7] and have been exploited by various authors thereafter; see [11] for a good introduction. In [19], Watanabe constructed smooth (trivialized near a section and framed, in the above sense)  $S^4$ -bundles with nontrivial Kontsevich’s characteristic classes, implying that as smooth fiber bundles (with fixed trivialization near a section) they are nontrivial, while as topological fiber bundles they are trivial.

MSC2020: 57N16, 57R20.

© 2026 The Author, under license to MSP (Mathematical Sciences Publishers). Distributed under the Creative Commons Attribution License 4.0 (CC BY). Open Access made possible by subscribing institutions via [Subscribe to Open](#).

We would like to understand why Kontsevich’s characteristic classes are able to differentiate smooth fiber bundles that are topologically the same. These classes are constructed by considering the configuration space bundles associated to  $E \xrightarrow{\pi} B$ , which are obtained by doing a sequence of real (oriented) blow-up operations fiberwise, and then doing some sort of intersection in the total space of the configuration space bundle to get an intersection number. Since intersection theoretical invariants usually do not depend on the smooth structure, while the real blow-up operations do, it is plausible that different smooth structures on the original bundle  $E \xrightarrow{\pi} B$  yield different topological structures on the induced configuration space bundles, and Kontsevich’s characteristic classes only depend on the topological bundle structure of the configuration space bundles (together with some information from the framing). The purpose of the present article is to make this statement precise and to give a detailed proof. The main theorem, Theorem 1.2, says that the topological information from the 2-point configuration space bundle of  $E \xrightarrow{\pi} B$ , together with a framing, determines Kontsevich’s characteristic classes of  $E \xrightarrow{\pi} B$ .

**Remark 1.1** The homotopy type of the real oriented blow-up of a manifold  $X$  along a submanifold  $Y$ ,  $\text{Bl}_Y X$ , does not depend on the smooth structure on  $X$ , since it is just  $X - Y$ ; but the topological structure of  $\partial \text{Bl}_Y X$  as a sphere bundle over  $Y$  and how  $\partial \text{Bl}_Y X$  is attached to  $X - Y$  do depend on the smooth structure in an essential way; and a framing on the normal bundle of  $Y$  can be used to capture this structure.

For example, the simplest Kontsevich’s characteristic classes, when it is a number ( $\Theta$ -graph invariants), can be viewed as the triple intersection number of a cohomology class (called “propagator class” in Section 4) in a space  $\overline{C}_2(\pi)/\sim_F$ , where  $\overline{C}_2(\pi)$  is the total space of the 2-point configuration space bundle associated to  $E \xrightarrow{\pi} B$ , and  $\sim_F$  is an equivalence relation on  $\partial^v \overline{C}_2(\pi)$  (the vertical boundary of  $\overline{C}_2(\pi)$ )—it uses the framing data  $F$  to “pinch”  $\partial^v \overline{C}_2(\pi)$ , making it lower-dimensional.<sup>1</sup>

## 1.1 Statement of main result

Throughout Sections 1–5, let  $M$  be a closed smooth  $d$ -dimensional manifold whose  $R$ -homology groups are the same as that of the  $d$ -sphere, where  $R = \mathbb{Z}$  or  $\mathbb{R}$ , and let  $\infty$  be a fixed point in  $M$ . Let  $\overline{C}_2(M, \infty)$  be the configuration space of two ordered, distinct points in  $M$ , neither of which is  $\infty$ , compactified by the Fulton–MacPherson compactification. More precisely,  $\overline{C}_2(M, \infty)$  is obtained from  $M \times M$  by first blowing up  $\infty \times \infty$  and then blowing up the strict transforms<sup>2</sup> of  $\infty \times M$ ,  $M \times \infty$  and the diagonal  $\Delta$ . (All the blow-ups we use are oriented.) Denote by  $f_+, f_- : \overline{C}_2(M, \infty) \rightarrow M$  the two forgetful maps lifting

$$M^2 \longrightarrow M, \quad f_+(x_1, x_2) = x_2, \quad f_-(x_1, x_2) = x_1,$$

respectively. So,  $(f_-, f_+) : \overline{C}_2(M, \infty) \rightarrow M \times M$  is the blow-down map.

<sup>1</sup>This idea of constructing  $\overline{C}_2(\pi)/\sim_F$  and viewing the propagator as a cohomology class in it has already been explored in the early work of Kuperberg and Thurston [10].

<sup>2</sup>Suppose  $N$  and  $N'$  are submanifolds of  $M$ . The *strict transform* of  $N'$  in  $\text{Bl}_N M$  is the closure of  $N' \setminus N$ , which is a subset of  $M \setminus N \subset \text{Bl}_N M$ .

Denote by  $\text{Diff}_+(M)$  (resp.  $\text{Homeo}_+(M)$ ) the group of orientation-preserving diffeomorphisms (resp. homeomorphisms) of  $M$ , with the Whitney (resp. compact-open) topology, and define

$$\begin{aligned} \text{Diff}_+(M, N_\infty) &:= \{g \in \text{Diff}_+(M) \mid \exists \text{ neighborhood } U \ni \infty \text{ such that } g|_U = \text{id}\}, \\ \text{Homeo}_+(M, N_\infty) &:= \{g \in \text{Homeo}_+(M) \mid \exists \text{ neighborhood } U \ni \infty \text{ such that } g|_U = \text{id}\}, \\ \mathcal{G} &:= \{(\tilde{g}, g) \in \text{Homeo}(\bar{C}_2(M, \infty)) \times \text{Homeo}_+(M, N_\infty) \mid g \circ f_\pm = f_\pm \circ \tilde{g}\}. \end{aligned}$$

By a smooth  $(M, \infty)$ -bundle  $E \xrightarrow{\pi} B$  we mean a fiber bundle with typical fiber  $(M, \infty)$  and structure group  $\text{Diff}_+(M, N_\infty)$ . It has a canonical section  $s_\infty$  and a canonical germ of trivializations

$$\begin{array}{ccc} t : B \times U & \xrightarrow{\sim} & \tilde{U} \\ & \searrow \text{projection to } B & \swarrow \pi \\ & & B \end{array}$$

of some neighborhood  $\tilde{U} \supset s_\infty(B)$  ( $U$  is some neighborhood of  $\infty$  in  $M$ ). Abusing notation we also denote the image  $s_\infty(B)$  by  $s_\infty$ .

A framing  $F$  on a smooth  $(M, \infty)$ -bundle  $E \xrightarrow{\pi} B$  is a continuous choice of basis for the vertical tangent space at every point of  $E - s_\infty$ , such that there are neighborhoods  $E \supset \tilde{U} \supset s_\infty$  and  $M \supset U \supset \infty$ , a diffeomorphism  $(U, \infty) \approx ((\mathbb{R}^d - 0) \sqcup \infty, \infty)$ , satisfying that  $t^*(F|_{\tilde{U}})$  is the standard framing on  $\mathbb{R}^d$  under this diffeomorphism.

Let  $\bar{C}_2(\pi) \rightarrow B$  be the associated  $\bar{C}_2(M, \infty)$ -bundle, let  $f_+, f_- : \bar{C}_2(\pi) \rightarrow E$  be the two forgetful maps and let  $\partial^v \bar{C}_2(\pi)$  consist of boundaries of every fiber. So

$$\partial^v \bar{C}_2(\pi) = (f_-, f_+)^{-1}(\Delta(\pi) \cup s_\infty \times_B E \cup E \times_B s_\infty),$$

where  $\times_B$  denotes fiber product over  $B$  and  $\Delta(\pi) \subset E \times_B E$  is the fiberwise diagonal. The framing  $F$  induces a map  $(f_-, f_+)^{-1}(\Delta(\pi)) \rightarrow S^{d-1}$  which at each point  $x \in E - s_\infty$  maps

$$(f_-, f_+)^{-1}(x, x) = S\mathcal{N}_{\Delta(\pi)}^v E|_x \approx ST_x^v E$$

to  $S^{d-1}$  using  $F : T_x^v E \approx \mathbb{R}^d$ ; here  $ST^v E$  denotes the sphere bundle of the vertical tangent bundle of  $E$  and  $S\mathcal{N}_{\Delta(\pi)}^v E$  denotes the sphere bundle of the vertical normal bundle of  $\Delta(\pi)$  in  $E$ . Using the trivialization  $t$ , this map can be extended to a map  $\partial^v \bar{C}_2(\pi) \rightarrow S^{d-1}$  that we still denote by  $F$ , abusing notation; see for example [20, Section 2.3] for the detailed definition.

Kontsevich's invariants, as cohomology classes in  $B$  parameterized by graph homology, are defined for smooth framed  $(M, \infty)$ -bundles with smooth base  $B$  and  $R = \mathbb{R}$ . In this article we need to restrict to the case of having only trivalent graphs,<sup>3</sup> so, throughout this article, Kontsevich's invariant for a bundle

<sup>3</sup>The general definition of Kontsevich's invariants doesn't need to assume this, but for us, because of a nuanced technicality (more precisely, we need the very last sentence of Section 2 to hold, which is needed in the proof of Lemma 3.20), the proof of Theorem 1.2 only works if all the vertices of the graph have the same valency. Since it is customary to consider trivalent graphs, we will just assume that.

$\pi : E \rightarrow B$  is a map

$$\{\text{formal sum of trivalent graphs, closed in graph homology}\} \longrightarrow H^*(B; R).$$

One of the by-products of this article is to extend Kontsevich’s invariants to the case where  $B$  is any paracompact Hausdorff space and  $R = \mathbb{Z}$ . The main theorem is

**Theorem 1.2** *Let  $E' \xrightarrow{\pi'} B'$ ,  $E'' \xrightarrow{\pi''} B''$  be smooth  $(M, \infty)$ -fiber bundles and  $s'_\infty, s''_\infty$  their canonical sections. Let  $F'$  and  $F''$  be framings on  $\pi'$  and  $\pi''$ , respectively. If there exist **continuous** maps*

$$\tilde{h} : \bar{C}_2(\pi') \longrightarrow \bar{C}_2(\pi''), \quad h : E' \longrightarrow E'', \quad h_B : B' \longrightarrow B'', \quad h_S : S^{d-1} \xrightarrow{\text{homeomorphism}} S^{d-1}$$

such that the following diagrams commute

$$\begin{array}{ccc} \bar{C}_2(\pi') & \xrightarrow{\tilde{h}} & \bar{C}_2(\pi'') \\ f'_- \downarrow \downarrow f'_+ & & f''_- \downarrow \downarrow f''_+ \\ E' & \xrightarrow{h} & E'' \\ s'_\infty \uparrow \downarrow \pi' & & s''_\infty \uparrow \downarrow \pi'' \\ B' & \xrightarrow{h_B} & B'' \end{array} \quad \begin{array}{ccc} \partial^v \bar{C}_2(\pi') & \xrightarrow{\tilde{h}|_{\partial^v \bar{C}_2(\pi')}} & \partial^v \bar{C}_2(\pi'') \\ \downarrow F' & & \downarrow F'' \\ S^{d-1} & \xrightarrow{h_S} & S^{d-1} \end{array}$$

and for every point  $b \in B$ ,  $\tilde{h}$  and  $h$  restrict to orientation-preserving homeomorphisms on the fibers over  $b$ . Then the Kontsevich’s characteristic classes of  $\pi''$  pull back to those of  $\pi'$  by  $h_B$ .

**Remark 1.3** Without the commutativity condition of the diagram on the right, the assumption of this theorem is the same as saying  $(\tilde{h}, h)$  is a  $\mathcal{G}$ -bundle map. See Section 6 for some discussions about the condition in this theorem. In particular, Theorem 6.2 is a restatement and extension of Theorem 1.2.

**Remark 1.4** Theorem 1.4 in [12] (this preprint appeared a few months after the preprint of the present paper) implies Theorem 1.2, at least in the case  $M$  is a sphere. See Theorem 6.2 — a restatement of Theorem 1.2 (and the second to last paragraph above it, as well as Proposition 6.5) — for why this is the case.

### 1.2 Outline of the proof

We reconstruct Kontsevich’s characteristic classes in a way that all the definitions are made using only the topological bundle structure on  $\bar{C}_2(\pi)$ ,  $\pi$  and the maps  $f_\pm$ , avoiding using the smooth structure in definitions. This will make Theorem 1.2 automatic. Sections 2–4 are devoted to this reconstruction, which is really just a translation of the original construction. In Section 5 we show that the new definition is equivalent to the original one.

To make such a reconstruction, the natural strategy is to translate the original construction using differential forms into the language of some topological cohomology theory (here we use Čech cochains) and thus avoid using the smooth structure. However, the cochains in all topological cohomology theories are rather cumbersome to work with, so we need to find the appropriate spaces so that we actually work

with cohomology classes. The seemingly unmotivated definitions in Section 3 are for this purpose. This approach is very similar to the work of Kuperberg and Thurston [10] (and some later works, e.g., [8]), but with some major differences; see Remark 1.5 below.

We describe in a bit more detail how the reconstruction is done below. First, in this paragraph, we briefly recall how the original construction roughly goes. Let  $E \xrightarrow{\pi} B$  be a smooth  $(M, \infty)$ -bundle with smooth, compact base  $B$ , and  $F$  a framing on  $\pi$ . Let  $\Gamma$  be a trivalent graph that is closed in graph homology;<sup>4</sup> denote its vertex set and edge set by  $V(\Gamma)$  and  $E(\Gamma)$ , respectively. Denote by  $\overline{C}_{V(\Gamma)}(\pi)$  the Fulton–MacPherson compactification of

$$C_{V(\Gamma)}(\pi) := \{(x_v \in E)_{v \in V(\Gamma)} \mid x_v \notin s_\infty, \pi(x_v) = \pi(x_w), x_v \neq x_w, \forall v \neq w \in V(\Gamma)\}.$$

It is a manifold with boundaries and corners. For every edge  $e$  of  $\Gamma$ , there is a forgetful map

$$f_e : \overline{C}_{V(\Gamma)}(\pi) \rightarrow \overline{C}_2(\pi)$$

forgetting everything but the two points labeled by the vertices adjacent to  $e$ . Take a closed  $(d-1)$ -form  $\omega$  (called *propagator*) on  $\overline{C}_2(\pi)$  satisfying  $\omega|_{\partial^v \overline{C}_2(\pi)} = F^* \text{vol}$  for some form  $\text{vol}$  on  $S^{d-1}$  such that  $\int_{S^{d-1}} \text{vol} = 1$ . Then the desired characteristic class (with parameter  $\Gamma$ ) is defined to be the class represented by the push-forward of  $\bigwedge_e f_e^* \omega_e$  to  $B$ . This pushed-forward form is not automatically closed or independent of the choice of  $\omega$ ; all the trouble here is that  $\overline{C}_{V(\Gamma)}(\pi)$  has boundary. The codimension-1 boundary strata of  $\overline{C}_{V(\Gamma)}(\pi)$  are in correspondence with subsets  $A \subset \{\infty\} \sqcup V(\Gamma)$  having at least 2 elements. Denote by  $\overline{\mathcal{P}}_A$  the closed boundary stratum corresponding to  $A$ ; it represents the configurations where the points with labels in  $A$  all coincide and “bubble off to a screen”. These boundary strata are divided into 4 types, and treated separately.

- (1) The subgraph  $\Gamma_A$  of  $\Gamma$  spanned by vertices in  $A$  has a zero- or univalent vertex, and is not of the form as in the 4-th type below. Then  $\overline{\mathcal{P}}_A$  is contracted by the map to  $B$  and thus does not contribute.
- (2)  $\Gamma_A$  has a bivalent vertex; then there is an involution on  $\overline{\mathcal{P}}_A$ , making the contribution from  $\overline{\mathcal{P}}_A$  cancel with itself.
- (3)  $\infty \in A$  or  $A = V(\Gamma)$ . This case relies on the framing  $F$ , the trivialization of  $\pi$  near  $s_\infty$ , and that  $\omega$  is defined to be compatible with  $F$ . A dimension count shows that  $\overline{\mathcal{P}}_A$  does not contribute either.
- (4)  $\Gamma_A$  has two vertices connected by an edge, then these  $\overline{\mathcal{P}}_A$  correspond to boundary terms of  $\Gamma$  in graph cohomology. Since  $\Gamma$  is closed, their total contribution is 0.

In our reconstruction, we no longer use differential form propagators; instead, define the space  $\overline{C}_2(\pi)/\sim_F$  obtained from  $\overline{C}_2(\pi)$  by contracting  $\partial^v \overline{C}_2(\pi)$  to  $S^{d-1}$  using the framing  $F$ , and the propagator can be naturally replaced by a *propagator class*  $\Omega \in H^{d-1}(\overline{C}_2(\pi)/\sim_F)$ . Denote by  $q : \overline{C}_2(\pi) \rightarrow \overline{C}_2(\pi)/\sim_F$  the quotient map.

<sup>4</sup>Throughout this paper we work with a single graph  $\Gamma$  instead of a formal sum of graphs, but this is just for simplicity. All the arguments can be modified to work if we assume  $\Gamma$  is a formal sum of graphs instead; see Remark 3.26.

Our construction also treats the boundary strata of  $\bar{C}_{V(\Gamma)}(\pi)$  type by type; each type is treated in the same spirit as the original arguments.

(1) Instead of using  $\bar{C}_{V(\Gamma)}(\pi)$ , we use a “weaker” compactification of  $C_{V(\Gamma)}(\pi)$ , denoted by  $\bar{C}_\Gamma(\pi)$ . When marked points coincide,  $\bar{C}_{V(\Gamma)}(\pi)$  records the colliding directions of every pair of points, as well as the relative colliding speed of each triple of points.<sup>5</sup> But  $\bar{C}_\Gamma(\pi)$  only records the colliding directions of pairs of points whose labels in  $V(\Gamma)$  are connected by an edge of  $\Gamma$ ;  $\bar{C}_\Gamma(\pi)$  also does not record the relative colliding speed of triples of points. Indeed,  $\bar{C}_\Gamma(\pi)$  is defined to be the closure of the image of

$$C_{V(\Gamma)}(\pi) \xrightarrow{(f_e)_{e \in E(\Gamma)}} \bar{C}_2(\pi)^{E(\Gamma)}.$$

We denote the image of each  $\bar{\mathcal{P}}_A$  in  $\bar{C}_\Gamma(\pi)$  by  $\bar{\mathcal{P}}_A^\Gamma$ . Those  $\bar{\mathcal{P}}_A \subset \bar{C}_{V(\Gamma)}(M, \infty)$  for  $A$  of type 1 are “contracted”, i.e.,  $\bar{\mathcal{P}}_A^\Gamma \subset \bar{C}_\Gamma(\pi)$  has codimension-2 or higher.

(2) and (4) The original arguments in these two cases tell us that each  $\bar{\mathcal{P}}_A$  of type 2 has an involution and each  $\bar{\mathcal{P}}_A$  of type 4 can be paired-up and cancel each other. Morally, we want to define a new space obtained from  $\bar{C}_\Gamma(\pi)$  by gluing each  $\bar{\mathcal{P}}_A^\Gamma$  of type 2 to itself by the involution, and glue each type 4 pair  $\bar{\mathcal{P}}_{A_1}^\Gamma, \bar{\mathcal{P}}_{A_2}^\Gamma$  to each other; thus they are no longer boundaries. However, the gluing procedure involves much technicality, so we instead just use the space

$$X_\Gamma(\pi) := \bigcup_{\sigma \in \tilde{S}_{E(\Gamma)}} \sigma \cdot \bar{C}_\Gamma(\pi) \subset \bar{C}_2(\pi)^{E(\Gamma)},$$

where  $\tilde{S}_{E(\Gamma)}$  is the signed permutation group of the set  $E(\Gamma)$  (a slight generalization of the usual permutation group), which acts on  $\bar{C}_2(\pi)^{E(\Gamma)}$  by permuting the factors. Taking the union of all the translates of  $\bar{C}_\Gamma(\pi)$  by  $\tilde{S}_{E(\Gamma)}$  makes the type 2 and type 4 boundary strata of them coincide and cancel with each other. We will explain in Section 3.4 that after removing a codimension-2 subset  $T_2(\pi)$  from  $X_\Gamma(\pi)$ , it is (fiberwise) a “manifold with boundary and bindings” (see Figure 1 for what binding means), where the pages adjacent to a binding, when counted with sign, sum up to 0. Since  $X_\Gamma(\pi)$  is a subspace of  $\bar{C}_2(\pi)^{E(\Gamma)}$ , for each edge  $e$  of  $\Gamma$ , we still have the forgetful map  $f_e : X_\Gamma(\pi) \rightarrow \bar{C}_2(\pi)$ .

(3) Denote by  $S(\pi) \subset X_\Gamma$  the union of all  $\bar{\mathcal{P}}_A^\Gamma$  of type 3. Then the boundary of  $X_\Gamma(\pi) - T_2(\pi)$  is contained in  $S(\pi)$ . We show that the cup product  $\Omega_\Gamma := \bigcup_{e \in E(\Gamma)} f_e^* q^* \Omega$  actually lands in the relative group  $H^*(X_\Gamma(\pi), S(\pi))$ .

<sup>5</sup>This perspective is most manifest in the construction of the Fulton–MacPherson compactification as taking closure in a big ambient space, as in [15, Definition 1.3]. There, for a manifold  $M$ , the compactified  $n$ -point configuration space  $\bar{C}_n(M)$  is defined as follows. Arbitrarily choose an embedding of  $M$  into a big Euclidean space  $\mathbb{R}^m$ . Then  $\bar{C}_n(M)$  is the closure of the uncompactified space  $C_n(M)$  in  $M^n \times (S^{m-1})^{\binom{n}{2}} \times [0, \infty]^{\binom{n}{3}}$ , where the embedding of  $C_n(M)$  into this product space is given by: the  $M^n$  factor records the position of the  $n$  marked points in  $M$ ; the  $(S^{m-1})^{\binom{n}{2}}$  factor records the direction between each pair of marked points; the  $[0, \infty]^{\binom{n}{3}}$  factor records the relative distance  $|x_1 - x_2|/|x_1 - x_3|$  among each triple of points. For a manifold  $M$  and a graph  $\Gamma$ , our weaker compactification here can be defined in a similar way with the big product space replaced by  $M^{V(\Gamma)} \times (S^{m-1})^{E(\Gamma)}$ ; namely, only the directions between pairs of points connected by edges of  $\Gamma$  are recorded. The  $\bar{C}_{V(\Gamma)}(\pi)$  is the bundle version (family version) of this.




Figure 1: Left: a binding with 4 adjacent pages. Right: an illustration of  $X_\Gamma$ .

Since  $X_\Gamma(\pi)$  can be roughly thought of as a manifold with boundary  $S(\pi)$  and bindings with pages summing up to 0,  $H^*(X_\Gamma(\pi), S(\pi))$  is not trivial and  $\Omega_\Gamma$  contains all the information we need. We push it forward to  $B$  (cohomology push-forward is defined in Section 4.2.1 using the Leray–Serre spectral sequence) to obtain a class in  $H^*(B)$ , which is the desired Kontsevich’s characteristic class with parameter  $\Gamma$ .

**Remark 1.5** Our approach above to modify  $\bar{C}_{V(\Gamma)}(\pi)$  is very similar to that of [10] and [8]. The similarities include: considering the space  $\bar{C}_2(\pi)/\sim_F$  and view the propagator as a cohomology class of it; using a smaller configuration space to deal with boundary strata of type (1); and using a gluing construction to deal with boundary strata of type (2) and (4). The main difference concerns the construction of the “smaller configuration space”. In [8; 10], it is constructed by blowing up fewer diagonals in  $M^{V(\Gamma)}$  (denote by  $M$  the 3-manifold whose configuration space is to be constructed); thus the “smaller configuration space” obtained is a smooth manifold, and when doing the gluing, how the corners glue together nicely is analyzed. Here, we construct the “smaller configuration space” in a much simpler way (the closure of the image of the forgetful maps), and define the glued space  $X_\Gamma$  in a very simple way as well. As a trade-off, the downside of this approach is that these spaces are not smooth; therefore we have to make an effort to analyze that the part that is not necessarily smooth is of at least codimension-2. Yet it is because the definition of  $X_\Gamma(\pi)$  is so simple in our approach—essentially, only the topology of  $\bar{C}_2(\pi)$  is used—that we are able to prove Theorem 1.2 which only involves the 2-point configuration space instead of the  $n$ -point configuration spaces for bigger  $n$ .

**Remark 1.6** The reason we use Čech instead of singular or simplicial cochains is completely technical. As mentioned above, the spaces we work with are only topological spaces instead of manifolds, but we need a careful control of dimension, so we use the “covering dimension” which is defined for topological spaces. Čech cochains are the most convenient to work with in this situation.

### 1.3 Some auxiliary notation

For a set  $S$  let  $|S|$  denote the number of elements in  $S$ . For a set  $I$  and a space  $X$ , denote  $X^I = \prod_{i \in I} X$ ; for  $n \in \mathbb{Z}^{>0}$ , denote

$$X^n = \underbrace{X \times X \times \cdots \times X}_{n \text{ times}}.$$

Denote by  $\Delta_{\text{big}} \subset X^I$  the big diagonal. For a map  $f : X \rightarrow Y$ , denote

$$f^I : X^I \rightarrow Y^I, \quad f^I((x_i)_{i \in I}) = (f(y_i))_{i \in I}.$$

By the real blow-up of a smooth manifold  $X$  along a submanifold  $Y$  we mean the oriented blow-up: replacing  $Y$  by the sphere normal bundle of  $Y$  in  $X$ .

We will be using Čech cochains and Čech cohomology. For a space  $X$  with an open cover  $\mathcal{U}$ , and a coefficient ring  $R$ , we write  $\check{C}_{\mathcal{U}}^*(X; R)$  for the  $R$ -module of Čech cochains on  $X$  with respect to  $\mathcal{U}$ , valued in the constant sheaf  $R \times X$ . By a skew-symmetric Čech cochain  $\alpha \in \check{C}_{\mathcal{U}}^n(X; R)$  we mean  $\alpha$  is such that

$$\alpha(U_0, \dots, U_i, U_{i+1}, \dots, U_n) = -\alpha(U_0, \dots, U_{i+1}, U_i, \dots, U_n)$$

for all  $U_0, \dots, U_n \in \mathcal{U}$  and  $0 \leq i < n$ . If  $A \subset X$  is closed, we denote  $\check{C}_{\mathcal{U}}^*(X, A; R) = \check{C}_{\mathcal{U}}^*(X; R|_{X-A})$ , where  $R|_{X-A}$  is the sheaf obtained by restricting  $R \times X$  to  $X - A$  and then extending it by 0 to  $X$ . So,  $\check{C}_{\mathcal{U}}^*(X, A; R)$  can be identified with a submodule of  $\check{C}_{\mathcal{U}}^*(X; R)$ : it consists of Čech cochains that vanish on  $A$ . Denote by  $H_{\mathcal{U}}^*(X; R)$  and  $H_{\mathcal{U}}^*(X, A; R)$  the cohomology of  $\check{C}_{\mathcal{U}}^*(X; R)$  and  $\check{C}_{\mathcal{U}}^*(X, A; R)$ , respectively.

If  $\mathcal{U}'$  is a refinement of  $\mathcal{U}$ , then there is a well-defined map

$$H_{\mathcal{U}}^*(X; R) \rightarrow H_{\mathcal{U}'}^*(X; R)$$

(resp.  $H_{\mathcal{U}}^*(X, Y; R) \rightarrow H_{\mathcal{U}'}^*(X, Y; R)$ ), independent of the choice of refinements. Write  $H^*(X; R)$  (resp.  $H^*(X, Y; R)$ ) for the cohomology of  $X$  (resp. the pair  $(X, Y)$ ) with  $R$  coefficients; it is the direct limit of  $H_{\mathcal{U}}^*(X; R)$  (resp.  $H_{\mathcal{U}}^*(X, Y; R)$ ) as  $\mathcal{U}$  gets more and more refined. In Sections 3 and 4, the coefficient ring  $R$  is assumed to be  $\mathbb{Z}$  or  $\mathbb{R}$ , and is often suppressed from notation; in Section 5,  $R = \mathbb{R}$ .

We will also be using compactly supported Čech cohomology (and very occasionally Čech cohomology with some other support  $\Phi$ ; usually  $\Phi$  is the collection of compact subsets inside a subspace). The last two paragraphs hold in these cases as well. The notation in the compactly supported case will be an added subscript “ $c$ ”, e.g.,  $\check{C}_{c, \mathcal{U}}^*(X; R)$ ,  $H_{c, \mathcal{U}}^*(X; R)$ ,  $H_c^*(X; R)$ . (And the notation in the case with support  $\Phi$  will be an added subscript “ $\Phi$ ”.)

We follow [17, Section 2.4] for the definition of fiber bundles. When we say  $p : B \rightarrow X$  is a fiber bundle with fiber  $Y$  and structure group  $G$ , we mean a fiber bundle as in [17, Definition 2.3], with given (maximal) coordinate functions that we do not explicitly mention. Given a  $G$ -action on some other space  $Y'$  (resp. and a  $G$ -equivariant map  $f : Y' \rightarrow Y$ ), by *the associated bundle of  $p$  with fiber  $Y'$*  we mean a fiber bundle  $p' : B' \rightarrow X$  with fiber  $Y'$  and structure group  $G$  (resp. and a  $G$ -bundle map, see [17, Definition 2.5] for the definition,  $\tilde{f} : B' \rightarrow B$ ,  $p \circ \tilde{f} = p'$ ) built up by gluing coordinate charts in the same way as  $p$ .<sup>6</sup> Then  $p'$  (resp.  $(p', \tilde{f})$ ) is unique up to equivalence.

In Section 1.1 we defined a smooth  $(M, \infty)$ -bundle to be a fiber bundle with typical fiber  $(M, \infty)$  and structure group  $\text{Diff}_+(M, N_\infty)$ . If  $B$  is a smooth manifold, then it is the same as saying (by smooth approximation theorems we can make the transition maps smooth)  $E \xrightarrow{\pi} B$  is a smooth submersion between smooth manifolds, with a smooth section  $s_\infty : B \rightarrow E$ , such that for all  $b \in B$ ,  $(\pi^{-1}(b), s_\infty(b))$  is diffeomorphic to  $(M, \infty)$ , together with a neighborhood  $\tilde{U} \subset E$  of  $s_\infty(B)$  and a trivialization

$$t : B \times (U, \infty) \xrightarrow{\sim} (\tilde{U}, s_\infty)$$

where  $U \subset M$  is some neighborhood of  $\infty$ .

Throughout this article we assume the reader is familiar with the Fulton–MacPherson compactification — having the picture in their mind; see [5] or [11] for reference. Some familiarity with the original definition of Kontsevich's characteristic classes would help (see any one of [7; 11; 20, Section 2] for reference), but it will not be needed until Section 5.

## 2 Information from the graph

In this section we summarize notation and statements regarding graphs that will be used later.

Say a graph is *directed* if its edges have directions, *ordered* if both of its vertex set and edge set are ordered. If a vertex or edge is the  $i$ -th one in the ordering, we call  $i$  its *label*. Given such a graph  $\Gamma$ , we denote by  $V(\Gamma)$  its vertex set and  $E(\Gamma)$  its edge set. Denote by  $e_i^\Gamma$  the  $i$ -th edge of  $\Gamma$  and  $v_i^\Gamma$  the  $i$ -th vertex of  $\Gamma$ . Conversely, given an edge  $e$  or vertex  $v$  of  $\Gamma$ , we denote by  $o_\Gamma(e)$  or  $o_\Gamma(v)$  its label. (So  $o_\Gamma(e_i^\Gamma) = o_\Gamma(v_i^\Gamma) = i$ .) For an edge  $e$  of  $\Gamma$ , denote by  $v_+(e)$  and  $v_-(e)$  the output and input vertex connected to  $e$ , respectively.

Suppose  $\Gamma_1$  and  $\Gamma_2$  are directed, ordered graphs,  $\alpha : \Gamma_1 \rightarrow \Gamma_2$  an undirected, unordered graph isomorphism, then we denote by  $\text{sgn}(\alpha, \text{vertex})$ ,  $\text{sgn}(\alpha, \text{edge}) \in \{+1, -1\}$  the permutation signs of  $\alpha$  on the set of vertices and edges, respectively, and denote

$$\begin{aligned} \text{sgn}(\alpha, \rightarrow) &= (-1)^{\text{number of edges whose direction is reversed by } \alpha}, \\ \text{sgn}_d(\alpha) &:= \begin{cases} \text{sgn}(\alpha, \text{edge}) & \text{for } d \in \mathbb{Z} \text{ even,} \\ \text{sgn}(\alpha, \text{vertex}) + \text{sgn}(\alpha, \rightarrow) & \text{for } d \in \mathbb{Z} \text{ odd.} \end{cases} \end{aligned}$$

<sup>6</sup>If  $p$  is a principal  $G$ -bundle (i.e., the  $G$  action on  $Y$  is free and transitive), then  $p'$  can be written as  $(Y' \times B)/G \rightarrow X$ . Here  $(Y' \times B)/G$  means the quotient of  $Y' \times B$  by the  $G$ -action  $g \cdot (y, b) = (yg, g^{-1}b)$ .

For a graph  $\Gamma$ , define  $\text{Aut}^u(\Gamma)$  to be the group of automorphisms of  $\Gamma$  as an undirected, unlabeled graph. Define

$$|\text{Aut}^u(\Gamma)|_d^\pm = \sum_{\alpha \in \text{Aut}^u(\Gamma)} \text{sgn}_d(\alpha) \in \mathbb{Z}.$$

For a set  $I$ , denote by  $S_I$  the permutation group of  $I$  and  $\tilde{S}_I$  the signed permutation group of  $I$ . Namely,  $\tilde{S}_I$  is the group of bijections from  $\bigsqcup_{i \in I} \{i^+, i^-\}$  to itself satisfying that, if  $i^+$  is mapped to  $j^+$  or  $j^-$ , then  $i^-$  is also mapped to  $j^+$  or  $j^-$ . There is an obvious map from  $\tilde{S}_I$  to  $S_I$ . For  $\sigma \in \tilde{S}_I$ , we denote by  $\text{sgn}(\sigma)$  the sign of its image in  $S_I$ ; denote  $\text{sgn}'(\sigma) := (-1)^{|\{i \in I \mid \sigma(i^+) = i^-\}|}$ .

Given an element  $\alpha \in \text{Aut}^u(\Gamma)$ , we denote by  $\alpha_V \in S_{V(\Gamma)}$  and  $\alpha_E \in S_{E(\Gamma)}$  the permutations of the vertices and edges induced by  $\alpha$ , respectively. Define

$$\psi_\Gamma : \text{Aut}^u(\Gamma) \longrightarrow \tilde{S}_{E(\Gamma)}, \quad \psi_\Gamma(\alpha)(e^\pm) = \alpha_E(e)^\pm \quad \text{if } \alpha_V(v_+(e)) = \alpha_V(v_\pm(e)).$$

Intuitively, one can think of  $e^\pm$  as the two half-edges of  $e$ .


If  $\Gamma$  has no isolated vertex, this is an injective group homomorphism.

### 2.1 Quick review of graph homology

Let  $\mathfrak{G}$  be the free abelian group generated by directed, ordered graphs that are also nonempty, connected and such that every vertex is at least trivalent. Define two equivalence relations on  $\mathfrak{G}$ ,  $\sim_{\text{odd}}$  and  $\sim_{\text{even}}$ , generated by the following: for directed ordered graphs  $\Gamma_1$  and  $\Gamma_2$ , if there exists an isomorphism  $\phi : \Gamma_1 \rightarrow \Gamma_2$  as unoriented unordered graphs, then

$$\Gamma_1 \sim_{\text{odd}} (\text{sgn}(\phi, \text{vertex}) \cdot \text{sgn}(\phi, \rightarrow))\Gamma_2, \quad \Gamma_1 \sim_{\text{even}} \text{sgn}(\phi, \text{edge})\Gamma_2.$$

For a directed, ordered graph  $\Gamma$  and an edge  $e$  of  $\Gamma$ , we define  $\Gamma/e$  to be the graph obtained from  $\Gamma$  by contracting  $e$ , with edge directions unchanged, vertices ordered in the same way as  $\Gamma$ , except for the new vertex, now put in the very front; edges ordered in the same way as  $\Gamma$ ; as below:



Define  $\mathbb{Z}$ -linear maps

$$\delta_{\text{odd}} : \mathfrak{G}/\sim_{\text{odd}} \longrightarrow \mathfrak{G}/\sim_{\text{odd}}, \quad \delta_{\text{even}} : \mathfrak{G}/\sim_{\text{even}} \longrightarrow \mathfrak{G}/\sim_{\text{even}}$$

induced by, for a directed ordered graph  $\Gamma$ ,

$$\delta_{\text{odd}}(\Gamma) := \sum_{e \in E(\Gamma)} (-1)^{o_\Gamma(v_+(e)) - o_\Gamma(v_-(e)) +} \begin{cases} 0 & \text{if } o_\Gamma(v_+(e)) > o_\Gamma(v_-(e)) \\ 1 & \text{if } o_\Gamma(v_+(e)) < o_\Gamma(v_-(e)) \end{cases} \Gamma/e,$$

$$\delta_{\text{even}}(\Gamma) := \sum_{e \in E(\Gamma)} (-1)^{o_\Gamma(e)} \Gamma/e.$$

It can easily be seen that  $\delta_{\text{odd}}^2, \delta_{\text{even}}^2 = 0$ . Graph homology is defined to be the homology of  $(\mathfrak{G}/\sim_{\text{odd}}, \delta_{\text{odd}})$  and  $(\mathfrak{G}/\sim_{\text{even}}, \delta_{\text{even}})$ . For shorter notation, for an edge  $e$  in a graph  $\Gamma$ , we define

$$\text{sgn}_{\text{odd}, \Gamma}(e) = (-1)^{o_{\Gamma}(v_+(e)) - o_{\Gamma}(v_-(e)) + \begin{cases} 0 & \text{if } o_{\Gamma}(v_+(e)) > o_{\Gamma}(v_-(e)) \\ 1 & \text{if } o_{\Gamma}(v_+(e)) < o_{\Gamma}(v_-(e)) \end{cases}}$$

The main takeaway we need is the following statement, which follows directly from definition: if  $\Gamma_1 + \dots + \Gamma_n \in \mathfrak{G}$ ,  $[\delta_{\text{odd}}(\sum_{i=1}^n \Gamma_i)]_{\sim_{\text{odd}}} = 0$  or  $[\delta_{\text{even}}(\sum_{i=1}^n \Gamma_i)]_{\sim_{\text{even}}} = 0$  (“ $[\cdot]$ ” denotes taking equivalence class with respect to  $\sim_{\text{odd}}, \sim_{\text{even}}$ ), then there exists an (often not unique) pairing between the edges of  $\Gamma_1, \dots, \Gamma_n$ , such that, if  $e_i^{\Gamma_a}, e_j^{\Gamma_b}$  is a pair, then there is an undirected, unordered graph isomorphism

$$\alpha : \Gamma_a / e_i^{\Gamma_a} \longrightarrow \Gamma_b / e_j^{\Gamma_b}$$

such that

$$\text{sgn}(\alpha, \text{vertex}) \cdot \text{sgn}(\alpha, \rightarrow) \cdot \text{sgn}_{\text{odd}, \Gamma}(e_i^{\Gamma_a}) \cdot \text{sgn}_{\text{odd}, \Gamma}(e_j^{\Gamma_b}) = -1$$

or

$$\text{sgn}(\alpha, \text{edge}) \cdot (-1)^{o_{\Gamma_a}(e_i^{\Gamma_a}) + o_{\Gamma_b}(e_j^{\Gamma_b})} = -1,$$

respectively.

## 2.2 Information from the graph

Let  $d \geq 3$  be an integer. Denote by  $\bar{d} \in \{\text{odd}, \text{even}\}$  the parity of  $d$ . Let  $\Gamma$  be a connected, directed, ordered graph such that all vertices are at least trivalent, and  $[\delta_{\bar{d}}(\Gamma)]_{\sim_{\bar{d}}} = 0$ . For example, the tetrahedron graph—the graph with 4 vertices and 1 edge between each pair of vertices, arbitrarily directed and ordered—satisfies these conditions when  $d$  is even.

It is easy to see that all the arguments in this section work if we assume  $\Gamma$  is a formal sum of graphs instead.

For  $V' \subset V(\Gamma)$ , we denote by  $\Gamma_{V'}$  the subgraph of  $\Gamma$  spanned by the vertices in  $V'$ , and by  $\Gamma/V'$  the graph obtained from  $\Gamma$  by contracting  $\Gamma_{V'}$  to one single vertex  $[V']_v$ . The order of edges in  $\Gamma_{V'}$ ,  $\Gamma/V'$  and the order of vertices in  $\Gamma_{V'}$  are the same as in  $\Gamma$ . The order of vertices in  $\Gamma/V'$  is defined as:  $[V']_v$  is the first one, and the rest are ordered as in  $\Gamma$ .

Suppose  $A \subset \{\infty\} \sqcup V(\Gamma)$ ,  $|A| \geq 2$ . Define

$$\Gamma_A = \begin{cases} \Gamma_A \text{ as in the previous paragraph} & \text{if } \infty \notin A, \\ \Gamma/(A - \infty) & \text{if } \infty \in A, \end{cases} \quad \Gamma/\Gamma_A = \begin{cases} \Gamma/A & \text{if } \infty \notin A, \\ \Gamma_{A-\infty} & \text{if } \infty \in A. \end{cases}$$

We say  $A$  is of

- type 1, if  $\Gamma_A$  has a zero-valent or univalent vertex, and either  $|A| \geq 3$  or  $\Gamma_A$  has no edge;
- type 2, if  $\Gamma_A$  has a bivalent vertex but no uni- or zero-valent vertex;
- type 3, if all vertices of  $\Gamma_A$  are at least trivalent;
- type 4, if  $\Gamma_A$  has exactly 2 vertices with 1 edge connecting them.

Notice that since all vertices of  $\Gamma$  are at least trivalent,  $\infty \in A$  implies  $A$  is of type 3.

Suppose  $A$  as above is of type 2. We fix  $v_A$  a bivalent vertex in  $A$ . Denote by  $e_A^1, e_A^2 \in E(\Gamma)$  the two edges connected to  $v_A$ . Define  $\sigma_A \in \tilde{S}_{E(\Gamma)}$  as follows:

$$\begin{aligned} \sigma_A(e^\pm) &= e^\pm && \text{if } e \neq e_A^1, e_A^2, \\ \sigma_A(e_A^{1\pm}) &= e_A^{2\mp} && \text{if } e_A^1 \text{ and } e_A^2 \text{ both start or both end at } v_A, \\ \sigma_A(e_A^{1\pm}) &= e_A^{2\pm} && \text{if one of } e_A^1 \text{ or } e_A^2 \text{ starts at } v_A \text{ and the other ends at } v_A. \end{aligned}$$

Now we look at  $A$  of type 4 above. Since  $[\delta_{\bar{d}}(\Gamma)]_{\sim_{\bar{d}}} = 0$ , there exists a pairing (given  $\Gamma$ , let us choose it once and for all, and call it a  $\Gamma$ -pairing) between type 4  $A$ 's such that, if  $A_1, A_2$  are paired (call them a  $\Gamma$ -pair), then there exists an undirected, unordered graph isomorphism  $\alpha_{A_1 A_2} : \Gamma/A_1 \rightarrow \Gamma/A_2$ , as in the last paragraph of Section 2.1. Denote by  $e_1, e_2$  the edge in  $\Gamma$  between the two vertices in  $A_1, A_2$ , respectively. We call  $e_1, e_2$  a  $\Gamma$ -pair as well, and denote  $\alpha_{A_1 A_2}$  also by  $\alpha_{e_1 e_2}$ . We define  $\sigma_{e_1 e_2} \in \tilde{S}_{E(\Gamma)}$  to be

$$\begin{aligned} \sigma_{e_1 e_2}(e_1^\pm) &= e_2^\pm, \\ \sigma_{e_1 e_2}(e_i^{\Gamma\pm}) &= e_j^{\Gamma\pm} && \text{if } \alpha_{A_1 A_2}(e_i^\Gamma) = e_j^\Gamma, \text{ preserving direction,} \\ \sigma_{e_1 e_2}(e_i^{\Gamma\pm}) &= e_j^{\Gamma\mp} && \text{if } \alpha_{A_1 A_2}(e_i^\Gamma) = e_j^\Gamma, \text{ reversing direction.} \end{aligned}$$

Note that we implicitly identified edges in  $E_\Gamma - \{e_1\}$  (resp.  $E_\Gamma - \{e_2\}$ ) with edges in  $\Gamma/A_1$  (resp.  $\Gamma/A_2$ ). Evidently  $\sigma_{e_2 e_1} = \sigma_{e_1 e_2}^{-1}$ .

If  $\Gamma$  is trivalent, then  $[A_1]_v$  and  $[A_2]_v$  are the only vertices of valency 4 in  $\Gamma/A_1$  and  $\Gamma/A_2$ , respectively. So  $\alpha_{A_1 A_2}$  must map  $[A_1]_v$  to  $[A_2]_v$ .

### 3 Defining various spaces, all having a $\mathcal{G}$ -action

As mentioned in Section 1.2, we would like to (at least partially) upgrade the construction of Kontsevich's classes from the chain level to the cohomology level, and this is done by replacing the configuration space  $C_V(\Gamma)(\pi)$  with another space  $X_\Gamma(\pi)$ . In this section we construct the "single fiber version" of it,  $X_\Gamma$ , and prove various statements regarding its structure. In Section 4 we will consider the fiber bundle version. That the fiber bundles we consider later are  $\mathcal{G}$ -bundles (see Section 1.1) means that the spaces we construct here need to have  $\mathcal{G}$ -actions, which is clear from the definitions given in this section.

Let  $(M, \infty)$  be as in Section 1.1. Denote by  $\Delta \subset M \times M$  the diagonal. Denote by  $C_I(M) = M^I - \Delta_{\text{big}}$  the configuration space of distinct marked points on  $M$  labeled by  $I$ . If  $I$  is ordered, let  $C_I(M)$  be oriented by the product orientation on  $M^I$ . Denote by  $\bar{C}_I(M)$  its Fulton–MacPherson compactification [5]. It can be constructed from  $M^I$  by a sequence of real blow-ups along various diagonals. For example,  $\bar{C}_{\{1,2\}}(M)$  is the real blow-up of  $M \times M$  along  $\Delta$ . The space  $\bar{C}_I(M)$  has the structure of a smooth manifold with boundaries and corners. For  $I' \subset I$  we have a smooth forgetful map  $f_{I'} : \bar{C}_I(M) \rightarrow \bar{C}_{I'}(M)$  lifting the map

$$M^I \rightarrow M^{I'}, \quad (x_i)_{i \in I} \rightarrow (x_i)_{i \in I'}.$$

If  $I'$  has only one element  $i$ , we also denote  $f_i = f_{I'}$ .

Denote  $C_I(M, \infty) = \{(x_i)_{i \in I} \in C_I(M) \mid x_i \neq \infty, \forall i\}$ , oriented in the same way as  $C_I(M)$ . Denote by  $\bar{C}_I(M, \infty)$  its Fulton–MacPherson compactification, which is defined to be the preimage of  $\infty$  under the forgetful map  $f_{\{*\}} : \bar{C}_{I \sqcup \{*\}}(M) \rightarrow M$ . For simplicity we write  $\bar{C}_n(M, \infty) := \bar{C}_{\{1, \dots, n\}}(M, \infty)$ . For example,  $\bar{C}_2(M, \infty)$  is described in Section 1.1. Define  $\tau : \bar{C}_2(M, \infty) \rightarrow \bar{C}_2(M, \infty)$  to be the map swapping the two marked points.

For  $A \subset \{\infty\} \sqcup I$ ,  $|A| \geq 2$ , we denote by  $\mathring{\mathcal{G}}_A \subset \bar{C}_I(M, \infty)$  the (open) boundary stratum corresponding to that the marked points with labels in  $A$  coincide. Denote by  $\bar{\mathcal{G}}_A$  its closure in  $\bar{C}_I(M, \infty)$ .

### 3.1 Defining various spaces

Recall in Section 1.1 we defined

$$\mathcal{G} := \{(\tilde{g}, g) \in \text{Homeo}(\bar{C}_2(M, \infty)) \times \text{Homeo}_+(M, N_\infty) \mid g \circ f_\pm = f_\pm \circ \tilde{g}\}.$$

It is easy to see that the action of  $\mathcal{G}$  commutes with the point-swapping map  $\tau$ . We call the action of  $\mathcal{G}$  on  $\bar{C}_2(M, \infty)^I$  by acting simultaneously on every factor the *diagonal action*. All the actions of  $\mathcal{G}$  we talk about below are action by homeomorphisms.

Let  $\Gamma$  be a graph as in Section 2.2. Assume  $\Gamma$  is trivalent. For an edge  $e$  of  $\Gamma$ , we denote by  $f_e : \bar{C}_{V(\Gamma)}(M, \infty) \rightarrow \bar{C}_2(M, \infty)$  the forgetful map lifting

$$M^{V(\Gamma)} \longrightarrow M^2, \quad (x_v)_{v \in V(\Gamma)} \longrightarrow (x_{v_\pm(e)}, x_{v_\mp(e)}).$$

**Definition 3.1** Denote  $f_\Gamma = (f_e)_{e \in E(\Gamma)} : \bar{C}_{V(\Gamma)}(M, \infty) \rightarrow \bar{C}_2(M, \infty)^{E(\Gamma)}$ . Define

$$\bar{C}_\Gamma(M, \infty) = \text{image}(f_\Gamma), \quad C_\Gamma(M, \infty) = f_\Gamma(C_{V(\Gamma)}(M, \infty)).$$

So,  $C_\Gamma(M, \infty) \subset \bar{C}_\Gamma(M, \infty) \subset \bar{C}_2(M, \infty)^{E(\Gamma)}$ .

Since  $\Gamma$  is connected,  $f_\Gamma|_{C_{V(\Gamma)}(M, \infty)}$  is an embedding. Thus it gives a diffeomorphism

$$C_{V(\Gamma)}(M, \infty) \rightarrow C_\Gamma(M, \infty).$$

Since  $V(\Gamma)$  is ordered, this also gives  $C_\Gamma(M, \infty)$  an orientation. Since  $C_\Gamma(M, \infty)$  can be written as

$$\{(z_e)_{e \in E(\Gamma)} \in \bar{C}_2(M, \infty)^{E(\Gamma)} \mid \forall e, e' \in E(\Gamma), \forall s, s' \in \{+, -\}, f_s(z_e) = f_{s'}(z_{e'}) \iff v_s(e) = v_{s'}(e')\},$$

it is invariant under the diagonal  $\mathcal{G}$ -action.

**Lemma 3.2**  $\bar{C}_\Gamma(M, \infty)$  is the closure of  $C_\Gamma(M, \infty)$  in  $\bar{C}_2(M, \infty)^{E(\Gamma)}$ .

**Proof** Since  $\bar{C}_{V(\Gamma)}(M, \infty)$  is compact,  $f_\Gamma(\bar{C}_{V(\Gamma)}(M, \infty))$  is closed and thus contains the closure of  $C_\Gamma(M, \infty)$ . On the other hand,  $C_{V(\Gamma)}(M, \infty)$  is dense in  $\bar{C}_{V(\Gamma)}(M, \infty)$ , so

$$C_\Gamma(M, \infty) = f_\Gamma(C_{V(\Gamma)}(M, \infty))$$

is dense in  $f_\Gamma(\bar{C}_{V(\Gamma)}(M, \infty))$ . □

Abusing notation, we denote by the projection  $\bar{C}_\Gamma(M, \infty) \rightarrow \bar{C}_2(M, \infty)$  to the  $e$ -th coordinate still by  $f_e$  and denote the inclusion map  $\bar{C}_\Gamma(M, \infty) \rightarrow \bar{C}_2(M, \infty)^{E(\Gamma)}$  still by  $f_\Gamma$ . It follows directly from Lemma 3.2 that  $\bar{C}_\Gamma(M, \infty)$  is invariant under the diagonal action of  $\mathcal{G}$  on  $\bar{C}_2(M, \infty)^{E(\Gamma)}$ .

**Definition 3.3** For  $I$  an ordered set, define an action  $\phi$  of  $\tilde{S}_I$  on  $\bar{C}_2(M, \infty)^I$  by diffeomorphisms: for  $\sigma \in \tilde{S}_I$ ,

$$\phi(\sigma) : \bar{C}_2(M, \infty)^I \longrightarrow \bar{C}_2(M, \infty)^I, \quad \phi(\sigma)(z_i)_{i \in I} = \left( \begin{cases} z_j & \text{if } \sigma(i^\pm) = j^\pm \\ \tau(z_j) & \text{if } \sigma(i^\pm) = j^\mp \end{cases} \right)_{i \in I}.$$

It is clear from definition that every  $\phi(\sigma)$  is equivariant with respect to the diagonal action of  $\mathcal{G}$  on  $\bar{C}_2(M, \infty)^I$ .

**Definition 3.4**  $X_\Gamma := \bigcup_{\sigma \in \tilde{S}_{E(\Gamma)}} \phi(\sigma)(\bar{C}_\Gamma(M, \infty)) \subset \bar{C}_2(M, \infty)^{E(\Gamma)}$ .

We still denote the inclusion map  $X_\Gamma \rightarrow \bar{C}_2(M, \infty)^{E(\Gamma)}$  by  $f_\Gamma$  and its  $e$ -th factors by  $f_e$ .

It follows from the  $\mathcal{G}$ -invariance of  $\bar{C}_\Gamma(M, \infty)$  and the  $\mathcal{G}$ -equivariance of each  $\phi(\sigma)$  that  $X_\Gamma$  is invariant under the diagonal action of  $\mathcal{G}$  on  $\bar{C}_2(M, \infty)^{E(\Gamma)}$ . We therefore define the action of  $\mathcal{G}$  on  $X_\Gamma$  to be the restriction of the diagonal action. The maps  $f_e$  are clearly  $\mathcal{G}$ -equivariant.

It seems that  $X_\Gamma$  (and its bundle version) is the appropriate space that accommodates Kontsevich’s characteristic classes. It is defined this way—taking the  $|\tilde{S}_{E(\Gamma)}|$  copies of  $\bar{C}_\Gamma(M, \infty)$  inside of  $\bar{C}_2(M, \infty)^{E(\Gamma)}$  and taking their union—because this makes the type 2 and 4 boundary strata of the many copies of  $\bar{C}_\Gamma(M, \infty)$  cancel with each other; see Figure 1. On the other hand, being a subspace of  $\bar{C}_2(M, \infty)^{E(\Gamma)}$  automatically makes it (and all its subspaces) metrizable, as a topological space, which is needed for the various arguments regarding covering dimension and Čech cohomology below.

We first make a remark that in the definition of  $X_\Gamma$  as a union, the “main stratum” parts either coincide or do not intersect. This is the content of Lemma 3.5 below.

Recall  $\text{Aut}^u(\Gamma)$  is the group of automorphisms of  $\Gamma$  as an undirected, unlabeled graph. An element  $\alpha \in \text{Aut}^u(\Gamma)$  consists of permutations  $\alpha_V \in S_{V(\Gamma)}$  and  $\alpha_E \in S_{E(\Gamma)}$ . Denote by

$$\gamma : \text{Aut}^u(\Gamma) \longrightarrow \text{Diff}(\bar{C}_{V(\Gamma)}(M, \infty))$$

the action of  $\text{Aut}^u(\Gamma)$  on  $\bar{C}_{V(\Gamma)}(M, \infty)$  by permuting marked points according to  $\alpha_V$ , namely,  $\gamma(\alpha)$  lifts the map

$$M^{V(\Gamma)} \longrightarrow M^{V(\Gamma)}, \quad (x_v)_{v \in V(\Gamma)} \longrightarrow (x_{\alpha_V(v)})_{v \in V(\Gamma)}.$$

Then (recall  $\psi_\Gamma : \text{Aut}^u(\Gamma) \rightarrow \tilde{S}_{E(\Gamma)}$  defined right before Section 2.1)

$$(1) \quad f_\Gamma \circ \gamma(\alpha) = \phi(\psi_\Gamma(\alpha)) \circ f_\Gamma.$$

**Lemma 3.5** For  $\sigma \in \tilde{S}_{E(\Gamma)}$ , if  $\sigma \notin \text{image}(\psi_\Gamma)$ , then  $\phi(\sigma)(C_\Gamma(M, \infty)) \cap \bar{C}_\Gamma(M, \infty) = \emptyset$ ; if  $\sigma = \psi_\Gamma(\alpha)$  for some  $\alpha \in \text{Aut}^u(\Gamma)$ , then  $\phi(\sigma)(C_\Gamma(M, \infty)) = C_\Gamma(M, \infty)$  and  $\phi(\sigma)$  changes its orientation by  $\text{sgn}(\alpha, \text{vertex})^d$ .

**Proof** This lemma is intuitively quite simple: an edge  $e$  of  $\Gamma$  contains the information of the two vertices  $v_{\pm}(e)$ ; so, since every vertex of  $\Gamma$  is connected to some edge, all the vertices of  $\Gamma$  (and hence  $\Gamma$  itself) can be recovered from its edges. We spell out the details below:

For simplicity we prove the lemma in the case  $\Gamma$  has no repeated edges; it can easily be generalized to the other cases as well. Given an element  $z = (z_e)_{e \in E(\Gamma)} \in \bar{C}_2(M, \infty)^{E(\Gamma)}$ , we define its *set of vertex positions*  $\bigcup_e \{f_+(z_e), f_-(z_e)\} \subset M$ . It does not change under the  $\phi(\sigma)$  action which only permutes factors. For  $x \in \bar{C}_{V(\Gamma)}(M, \infty)$ , we denote by  $\{f_v(x)\}_{v \in V(\Gamma)}$  the set of vertex positions of  $f_{\Gamma}(x)$ , which has exactly  $|V(\Gamma)|$  (distinct) elements if  $x \in C_{V(\Gamma)}(M, \infty)$  and less otherwise. Thus, if we suppose  $\phi(\sigma)(C_{\Gamma}(M, \infty)) \cap \bar{C}_{\Gamma}(M, \infty) \neq \emptyset$ , and if  $x, y \in C_{V(\Gamma)}(M, \infty)$  are such that  $f_{\Gamma}(x) = \phi(\sigma)f_{\Gamma}(y)$ , then there is a permutation  $\alpha_V \in S_{V(\Gamma)}$  such that  $f_{\alpha_V(v)}(y) = f_v(x)$ . Since  $\Gamma$  has no repeated edge, there is a unique  $\alpha \in \text{Aut}^u(\Gamma)$  with  $\alpha_V$  as given. So  $\gamma(\alpha)(x) = y$ . So,

$$f_{\Gamma}(x) = \phi(\sigma)f_{\Gamma}(y) = \phi(\sigma)(f_{\Gamma}(\gamma(\alpha)(x))) = \phi(\sigma) \circ \phi(\psi_{\Gamma}(\alpha))(f_{\Gamma}(x)) = \phi(\sigma\psi_{\Gamma}(\alpha))(f_{\Gamma}(x)).$$

Since  $\phi(\sigma\psi_{\Gamma}(\alpha))$  acts by permuting factors of  $\bar{C}_2(M, \infty)^{E(\Gamma)}$  composed with  $\tau$ 's, and no two factors of  $f_{\Gamma}(x)$  are the same even modulo  $\tau$ , because  $\Gamma$  has no repeated edges, we must have  $\sigma\psi_{\Gamma}(\alpha) = \text{id}$ . So  $\sigma = \psi_{\Gamma}(\alpha^{-1})$ . This proves the lemma except for the statement about orientation. The orientation statement is straightforward.  $\square$

**Definition 3.6** For  $A \subset \{\infty\} \sqcup V(\Gamma)$ ,  $|A| \geq 2$ , define

$$C_{V(\Gamma):A}(M, \infty) = \begin{cases} \{(x_v)_{v \in V(\Gamma)} \in (M - \{\infty\})^{V(\Gamma)} \mid x_v = x_w \iff (v, w \in A \text{ or } v = w)\} & \text{if } \infty \notin A, \\ p_{V(\Gamma)-A}^{-1}(C_{V(\Gamma)-A}(M, \infty)) \cap p_A^{-1}(\infty, \dots, \infty) & \text{if } \infty \in A, \end{cases}$$

where  $p_{V(\Gamma)-A} : M^{V(\Gamma)} \rightarrow M^{V(\Gamma)-A}$  and  $p_A : M^{V(\Gamma)} \rightarrow M^{A-\{\infty\}}$  are the projections. Then we have the following commutative diagram:

$$\begin{array}{ccc} \mathring{\mathcal{P}}_A & \xrightarrow{f_{\Gamma}} & \bar{C}_2(M, \infty)^{E(\Gamma)} \\ \downarrow & & \downarrow (f_-, f_+)^{E(\Gamma)} \\ C_{V(\Gamma):A}(M, \infty) & \xrightarrow{f'_{\Gamma}} & (M^2)^{E(\Gamma)} \end{array}$$

Define

$$\mathcal{P}_A^{\Gamma} = (((f_-, f_+)^{E(\Gamma)})^{-1} f'_{\Gamma}(C_{V(\Gamma):A}(M, \infty))) \cap \bar{C}_{\Gamma}(M, \infty).$$

Denote by  $\bar{\mathcal{P}}_A^{\Gamma}$  the closure of  $\mathcal{P}_A^{\Gamma}$ . For an edge  $e$  of  $\Gamma$ , denote  $\mathcal{P}_e^{\Gamma} := \mathcal{P}_{\{v_+^{\Gamma}(e), v_-^{\Gamma}(e)\}}^{\Gamma}$ .

Intuitively, elements of  $\mathcal{P}_A^{\Gamma}$  are configurations of points such that those in  $A$  all coincide in  $M$  and no other two points coincide. Notice that  $\bar{\mathcal{P}}_A = f_{\Gamma}^{-1}(\bar{\mathcal{P}}_A^{\Gamma})$ ,  $\mathring{\mathcal{P}}_A \subset f_{\Gamma}^{-1}(\mathcal{P}_A^{\Gamma}) \subset \bar{\mathcal{P}}_A$ , and the inclusions are often strict.

Recall that (see Figure 1) in  $X_{\Gamma}$  as a union of many copies of  $\bar{C}_{\Gamma}(M, \infty)$  inside of  $\bar{C}_2(M, \infty)^{E(\Gamma)}$ , we want those codimension-1 strata,  $\mathcal{P}_A^{\Gamma}$ , of type 2 or 4 in the various copies of  $\bar{C}_{\Gamma}(M, \infty)$  to glue together to form bindings. For example, if the binding has only two pages, it should locally be two topological

manifolds with boundary gluing together along their boundaries, forming a topological manifold. This is only true if we remove the nonnice, “singular” parts of those  $\mathcal{P}_A^\Gamma$  and only leave the nice part, which is what we call  $\overset{\circ}{\mathcal{P}}_A^\Gamma$  in Definition 3.7 below.

**Definition 3.7** For  $A \subset \{\infty\} \sqcup V(\Gamma)$ ,  $|A| \geq 2$ , define  $\overset{\circ}{\mathcal{P}}_A^\Gamma \subset \mathcal{P}_A^\Gamma$  to be the set of points such that locally  $\bar{C}_\Gamma(M, \infty)$  is a topological manifold with boundary, i.e.,

$$\overset{\circ}{\mathcal{P}}_A^\Gamma := \{x \in \mathcal{P}_A^\Gamma \mid \exists U \subset \bar{C}_\Gamma(M, \infty) \text{ open neighborhood of } x$$

and homeomorphism  $\nu : U \rightarrow \mathbb{R}^{d|V(\Gamma)|-1} \times \mathbb{R}^{\geq 0}$  such that  $\nu(\mathcal{P}_A^\Gamma \cap U) = \mathbb{R}^{d|V(\Gamma)|-1} \times \{0\}\}$ .

Moreover, define

$$\overset{\circ}{\mathcal{P}}_A^\Gamma = \overset{\circ}{\mathcal{P}}_A^\Gamma - \bigcup_{\sigma \in \tilde{\mathcal{S}}_{E(\Gamma)}} \bigcup_{\substack{A' \subset \{\infty\} \sqcup V(\Gamma) \\ |A'| \geq 2}} \phi(\sigma)(\bar{\mathcal{P}}_{A'}^\Gamma - \overset{\circ}{\mathcal{P}}_{A'}^\Gamma).$$

In Definition 3.8 below we give names to the various parts of  $X_\Gamma$ . The “good” part, consisting of the main strata and the “nonsingular” parts of the type 2 and 4 codimension-1 strata, will function like a topological manifold with bindings as in Figure 1. The remaining of  $X_\Gamma$  can again be divided into two parts:  $S$ , consisting of all  $\bar{\mathcal{P}}_A^\Gamma$  of type 3, and  $T_2$ , consisting of everything else. We will show that neither  $S$  or  $T_2$  causes an issue in our argument, but for different reasons: we will show that  $T_2$  is of codimension at least 2 (hence the “2” in the name “ $T_2$ ”), therefore not contributing to an intersection-theoretical argument; although  $S$  has a larger dimension, it can be shown (in Section 4.1) that the cohomology class we care about is actually in the relative cohomology  $H^*(X_\Gamma, S)$ .

**Definition 3.8** Define the following subsets of  $X_\Gamma$ :

$$X_\Gamma^{\text{good}} = \bigcup_{\sigma \in \tilde{\mathcal{S}}_{E(\Gamma)}} \phi(\sigma) \left( C_\Gamma(M, \infty) \cup \bigcup_{A \text{ of type 2 or 4}} \overset{\circ}{\mathcal{P}}_A^\Gamma \right), \quad T_1 = X_\Gamma - X_\Gamma^{\text{good}},$$

$$S = \bigcup_{\sigma \in \tilde{\mathcal{S}}_{E(\Gamma)}} \bigcup_{A \text{ of type 3}} \phi(\sigma)(\bar{\mathcal{P}}_A^\Gamma), \quad T_2 = \bigcup_{\sigma \in \tilde{\mathcal{S}}_{E(\Gamma)}} \bigcup_A \phi(\sigma)(\bar{\mathcal{P}}_A^\Gamma - \overset{\circ}{\mathcal{P}}_A^\Gamma).$$

$A$  in the above formulas are subsets of  $\{\infty\} \sqcup V(\Gamma)$  such that  $|A| \geq 2$ .

Notice that if  $A$  is of type 1, then  $\overset{\circ}{\mathcal{P}}_A^\Gamma = \emptyset$  since  $\bar{\mathcal{P}}_A \subset \bar{C}_{V(\Gamma)}(M, \infty)$  is contracted by  $f_\Gamma$ . So  $T_1 \supset T_2 \supset T_1 - S$ . Also  $T_2, T_1$  and  $S$  are invariant under the  $\tilde{\mathcal{S}}_{E(\Gamma)}$ -action  $\phi$ . And  $T_1$  and  $T_2$  are closed, since every point in  $\overset{\circ}{\mathcal{P}}_A^\Gamma$  (resp.  $\overset{\circ}{\mathcal{P}}_A^\Gamma$ ) has a neighborhood lying in  $C_\Gamma(M, \infty) \cup \overset{\circ}{\mathcal{P}}_A^\Gamma$  (resp.  $C_\Gamma(M, \infty) \cup \overset{\circ}{\mathcal{P}}_A^\Gamma$ ). It is evident from the above three definitions that for every  $A, \mathcal{P}_A^\Gamma, \bar{\mathcal{P}}_A^\Gamma, \overset{\circ}{\mathcal{P}}_A^\Gamma, \overset{\circ}{\mathcal{P}}_A^\Gamma \subset \bar{C}_\Gamma(M, \infty)$  are invariant under the  $\mathcal{G}$ -action, and therefore  $S, T_1$  and  $T_2$  are invariant under the  $\mathcal{G}$ -action too.

Recall the *covering dimension* of a topological space  $X$  is the biggest integer  $N$  satisfying that any open cover of  $X$  has a refinement  $\mathcal{U}$  such that if  $U_0, \dots, U_{N+1} \in \mathcal{U}$  and  $U_i \neq U_j$  for all  $i, j$ , then  $U_0 \cap \dots \cap U_{N+1} = \emptyset$ . Below we write  $\text{dim}_t(X)$  for the covering dimension of  $X$  (“ $t$ ” stands for “topological”). Notice that everything we have defined so far are subspaces of  $\bar{C}_2(M, \infty)^{E(\Gamma)}$ , thus are all metrizable topological spaces.

In Sections 3.2 and 3.3 below, we show that

- $\dim_t(T_2) \leq d|V(\Gamma)| - 2$  and  $\dim_t(T_1) \leq d|V(\Gamma)| - 1$ ;
- $H^{d|V(\Gamma)|}(X_\Gamma, S; R)$  admits a nontrivial  $\mathcal{G}$ -equivariant map to  $R$ . (Recall  $R = \mathbb{Z}$  or  $\mathbb{R}$  is the coefficient ring; we take the trivial  $\mathcal{G}$ -action on  $R$ .) This statement is the goal of Section 3 and is all that we need to reconstruct Kontsevich's classes. In all that follows we will call this map  $\rho$ .

### 3.2 Dimension control

We list some basic properties of the covering dimension that will be used later in this subsection. Let  $X$  and  $X'$  be nonempty metrizable spaces.

- If  $Y \subset X$  is closed, then  $\dim_t(Y) \leq \dim_t(X)$ . This follows from the definition.
- If  $Y_1, \dots, Y_n \subset X$  are closed and  $\dim_t(Y_i) \leq m$  for all  $i$ , then  $\dim_t(Y_1 \cup \dots \cup Y_n) \leq m$ ; see [13, Theorem 9-10].
- $\dim_t(X \times X') \leq \dim_t(X) + \dim_t(X')$ ; see [13, Theorem 12-14].

**Lemma 3.9** *Let  $f : X \rightarrow Y$  be a smooth map between smooth manifolds (possibly with boundary and corners). Assume  $X$  is compact. Denote*

$$A_r = \{x \in X \mid \text{rank}(d_x f) \leq r\}, \quad fA_r = f(A_r).$$

*Then  $\dim_t(fA_r) \leq r$ . Specifically,  $\dim_t(f(X)) \leq \dim(X)$ .*

**Proof** This follows from two celebrated theorems. By [14, Corollary on page 169], there exist countably many charts  $(U_i \subset Y, \phi_i : U_i \rightarrow \mathbb{R}^n)_{i=1}^\infty$  of  $Y$  such that  $fA_r \subset \bigcup_i U_i$  and the Hausdorff dimension of  $\phi_i(U_i \cap fA_r)$  is at most  $r$  for all  $i$ . By [18], the covering dimension of a metrizable, separable space (which  $\phi_i(U_i \cap fA_r)$  is) is no bigger than its Hausdorff dimension. So,  $\dim_t(U_i \cap fA_r) \leq r$  for all  $i$ . Since  $X$  is compact and  $A_r$  is closed in  $X$ ,  $fA_r$  is compact, and so it is covered by the charts  $U_1, \dots, U_m$  for some  $m$ . By shrinking each  $U_i$  a little bit, we have closed subsets  $V_i$  with  $V_i \subset U_i \subset Y$  for  $i = 1, \dots, m$ , which still cover  $fA_r$ . And  $\dim_t(V_i \cap fA_r) \leq r$  for all  $i \leq m$ . Since  $V_i \cap fA_r$  are closed subsets of  $fA_r$  and  $fA_r$  is their union,  $\dim_t(fA_r) \leq r$ . □

**Corollary 3.10**  $\dim_t(X_\Gamma) \leq d|V(\Gamma)|$ ,  $\dim_t(S) \leq d|V(\Gamma)| - 1$ ,  $\dim_t(T_1) \leq d|V(\Gamma)| - 1$ .

**Definition 3.11** Given a graph  $\Gamma'$ , define  $\bar{V}_{\Gamma'} \subset (S^{d-1})^{E(\Gamma')}$  to be the image of the map

$$f_{\Gamma'}^{\mathbb{R}^d} = (f'_e)_{e \in E(\Gamma')} : \bar{C}_{V(\Gamma')}^{\text{quo}}(\mathbb{R}^d) \rightarrow (S^{d-1})^{E(\Gamma')},$$

where  $\bar{C}_{V(\Gamma')}^{\text{quo}}(\mathbb{R}^d)$  is the Fulton–MacPherson space of configurations of  $V(\Gamma')$ -marked points in  $\mathbb{R}^d$ , modulo translation and scaling (so  $\dim(\bar{C}_{V(\Gamma')}^{\text{quo}}(\mathbb{R}^d)) = d|V(\Gamma')| - d - 1$ ; “quo” stands for “quotient”), and  $f'_e$  is the unique map induced from

$$(\mathbb{R}^d)^{V(\Gamma')} - \Delta_{\text{big}} \rightarrow S^{d-1}, \quad (x_v)_{v \in V(\Gamma')} \rightarrow \frac{x_{v_+(e)} - x_{v_-(e)}}{|x_{v_+(e)} - x_{v_-(e)}|},$$

i.e.,  $f'_e$  is the direction between the points marked by the vertices adjacent to  $e$ . Notice the  $GL(d)$  action on  $\mathbb{R}^d$  induces  $GL(d)$ -actions on  $S^{d-1}$ ,  $\bar{C}_{V(\Gamma')}^{\text{quo}}(\mathbb{R}^d)$  and  $\bar{V}_{\Gamma'}$ .

Let  $C_{V(\Gamma')}^{\text{quo}}(\mathbb{R}^d)$  be the quotient of  $(\mathbb{R}^d)^{V(\Gamma')} - \Delta_{\text{big}}$  by translation and scaling.

**Lemma 3.12** *If  $x, y \in C_{V(\Gamma')}^{\text{quo}}(\mathbb{R}^d)$ ,  $x \neq y$  and  $f_{\Gamma'}^{\mathbb{R}^d}(x) = f_{\Gamma'}^{\mathbb{R}^d}(y)$ , then  $d_x f_{\Gamma'}^{\mathbb{R}^d}$  and  $d_y f_{\Gamma'}^{\mathbb{R}^d}$  are not injective.*

**Proof** Since  $C_{V(\Gamma')}^{\text{quo}}(\mathbb{R}^d)$  is a quotient, let us take representatives  $x' \in (\mathbb{R}^d)^{V(\Gamma')}$  of  $x$  and  $y' \in (\mathbb{R}^d)^{V(\Gamma')}$  of  $y$  such that the marked point labeled by  $v_1^{\Gamma'}$  is the origin and the marked point labeled by  $v_2^{\Gamma'}$  has norm 1. It is easy to see that for all  $0 < \lambda < 1$ , if  $\lambda x' + (1 - \lambda)y' \in (\mathbb{R}^d)^{V(\Gamma')} - \Delta_{\text{big}}$ , then  $f_{\Gamma'}^{\mathbb{R}^d}([\lambda x' + (1 - \lambda)y']) = f_{\Gamma'}^{\mathbb{R}^d}(x) = f_{\Gamma'}^{\mathbb{R}^d}(y)$ . So the differential of  $f_{\Gamma'}^{\mathbb{R}^d}$  at  $x$  or  $y$  vanishes in at least one direction. □

For  $A \subset \{\infty\} \sqcup V(\Gamma)$ ,  $|A| \geq 2$ , recall the definitions of  $\Gamma_A$  and  $\Gamma/\Gamma_A$  from Section 2.2. From the construction of Fulton–MacPherson compactification, we have:

**Fact 3.13**  $\bar{\mathcal{P}}_A$  is a fiber bundle over  $\bar{C}_{V(\Gamma/A)}(M, \infty)$  with fiber  $\bar{C}_{V(\Gamma_A)}^{\text{quo}}(\mathbb{R}^d)$  and structure group  $SL(d)$ .

Notice  $\bar{\mathcal{P}}_A^\Gamma = f_\Gamma(\bar{\mathcal{P}}_A)$ .

**Corollary 3.14**  $\bar{\mathcal{P}}_A^\Gamma$  is a fiber bundle over  $\bar{C}_{\Gamma/A}(M, \infty)$  with fiber  $\bar{V}_{\Gamma_A}$  and structure group  $SL(d)$ . The map  $f_\Gamma|_{\bar{\mathcal{P}}_A}$  is an  $SL(d)$ -bundle map covering  $f_{\Gamma/A} : \bar{C}_{V(\Gamma/A)}(M, \infty) \rightarrow \bar{C}_{\Gamma/A}(M, \infty)$ , which is  $f_{\Gamma_A}^{\mathbb{R}^d}$  on each fiber.

Notice that the covering dimension of the total space of a fiber bundle with compact base is no more than the sum of the covering dimensions of its fiber and base: over each chart, the bundle is a product, so this inequality holds; and since covering dimension does not increase when taking finite unions of closed subsets, it holds for the whole total space.

**Corollary 3.15** *Suppose  $A \subset \{\infty\} \sqcup V(\Gamma)$  is of type 1. Then  $\dim_t(\bar{\mathcal{P}}_A^\Gamma) \leq d|V(\Gamma)| - 2$ .*

**Proof** Since  $\Gamma_A$  has a zero-valent vertex  $v_0$  or a univalent vertex  $v_1$  adjacent to  $v'_1$ ,

$$f_{\Gamma_A}^{\mathbb{R}^d} : \bar{C}_{V(\Gamma_A)}^{\text{quo}}(\mathbb{R}^d) \rightarrow \bar{V}_{\Gamma_A}$$

factors through  $f_{v_0} : \bar{C}_{V(\Gamma_A)}^{\text{quo}}(\mathbb{R}^d) \rightarrow \bar{C}_{V(\Gamma_A) - \{v_0\}}^{\text{quo}}(\mathbb{R}^d)$  which forgets the point labeled by  $v_0$ , or through  $f_{v_1} : \bar{C}_{V(\Gamma_A)}^{\text{quo}}(\mathbb{R}^d) \rightarrow S^{d-1} \times \bar{C}_{V(\Gamma_A) - \{v_1\}}^{\text{quo}}(\mathbb{R}^d)$  which forgets the distance between the points  $v_1$  and  $v'_1$ . So,  $\dim_t(\bar{V}_{\Gamma_A}) < \dim(\bar{C}_{V(\Gamma_A)}^{\text{quo}}(\mathbb{R}^d))$ . □

**Lemma 3.16**  $\dim_t(\bar{\mathcal{P}}_A^\Gamma - \mathring{\mathcal{P}}_A^\Gamma) \leq d|V(\Gamma)| - 2$  for all  $A \subset \{\infty\} \sqcup V(\Gamma)$ ,  $|A| \geq 2$ .

**Proof** Denote by  $p_A : \bar{\mathcal{P}}_A \rightarrow \bar{C}_{V(\Gamma/A)}(M, \infty)$  and  $p_A^\Gamma : \bar{\mathcal{P}}_A^\Gamma \rightarrow \bar{C}_{\Gamma/A}(M, \infty)$  the fiber bundles in Fact 3.13 and Corollary 3.14, respectively. Denote

$$Z_A = \{x \in \bar{\mathcal{P}}_A \mid \text{rank } d_x(f_\Gamma|_{p_A^{-1}(p_A(x))}) < d|V(\Gamma_A)| - d - 1\}, \quad R_A = f_\Gamma(\bar{\mathcal{P}}_A - Z_A).$$

(Notice that  $p_A^{-1}(p_A(x))$  is just the fiber of  $p_A$  containing  $x$ .) In other words,  $Z_A$  consists of points  $x \in \bar{\mathcal{P}}_A$  at which  $f_\Gamma|_{p_A^{-1}(p_A(x))}$  is not an immersion.  $\bar{\mathcal{P}}_A^\Gamma - R_A$  is also a fiber bundle over  $\bar{C}_{\Gamma/A}(M, \infty)$  with fiber

$$f_{\Gamma_A}^{\mathbb{R}^d}(\{x \in \bar{C}_{V(\Gamma_A)}^{\text{quo}}(\mathbb{R}^d) \mid \text{rank } d_x f_{\Gamma_A}^{\mathbb{R}^d} < d|V(\Gamma_A)| - d - 1\}),$$

so  $\dim_t(\bar{\mathcal{P}}_A^\Gamma - R_A) \leq d|V(\Gamma)| - 2$  by Lemma 3.9, and  $\bar{\mathcal{P}}_A^\Gamma - R_A$  is closed in  $\bar{\mathcal{P}}_A^\Gamma$ . Since  $\bar{\mathcal{P}}_A - \mathring{\mathcal{P}}_A$  is covered by codimension-2 or higher strata of  $\bar{C}_{V(\Gamma)}(M, \infty)$ ,  $\dim_t(f_\Gamma(\bar{\mathcal{P}}_A - \mathring{\mathcal{P}}_A)) \leq d|V(\Gamma)| - 2$ . We claim that

$$(2) \quad \bar{\mathcal{P}}_A^\Gamma - (f_\Gamma(\bar{\mathcal{P}}_A - \mathring{\mathcal{P}}_A) \cup (\bar{\mathcal{P}}_A^\Gamma - R_A)) \subset \mathring{\mathcal{P}}_A^\Gamma.$$

If this is true, then  $\bar{\mathcal{P}}_A^\Gamma - \mathring{\mathcal{P}}_A^\Gamma$  is contained in the union of two closed subsets of  $\dim_t \leq d|V(\Gamma)| - 2$ ; and since  $\bar{\mathcal{P}}_A^\Gamma - \mathring{\mathcal{P}}_A^\Gamma$  is itself closed in  $\bar{\mathcal{P}}_A^\Gamma$  (that  $\mathring{\mathcal{P}}_A^\Gamma$  is open in  $\bar{\mathcal{P}}_A^\Gamma$  follows easily from the definition of  $\mathring{\mathcal{P}}_A^\Gamma$ ), we are done.

Let  $x$  be in the left-hand side of (2). Since  $x \in R_A$  and  $x \notin f_\Gamma(\bar{\mathcal{P}}_A - \mathring{\mathcal{P}}_A)$ , by Lemma 3.12,  $f_\Gamma^{-1}(x)$  consists of a single element  $y \in \mathring{\mathcal{P}}_A$ . We next show that  $f_\Gamma$  is a homeomorphism onto its image in a neighborhood  $U \subset \bar{C}_{V(\Gamma)}(M, \infty)$  of  $y$ . Since  $p_A(\mathring{\mathcal{P}}_A) = C_{V(\Gamma/A)}(M, \infty) = C_{\Gamma/A}(M, \infty)$ ,  $f_\Gamma|_{\mathring{\mathcal{P}}_A}$  is locally the product of  $f_{\Gamma_A}^{\mathbb{R}^d}$  with a diffeomorphism. So  $f_\Gamma|_{\mathring{\mathcal{P}}_A}$  is an immersion at  $y$ . Since  $f_\Gamma|_{C_{V(\Gamma)}(M, \infty)}$  is injective,  $f_\Gamma$  is injective in an open neighborhood  $U \subset \bar{C}_{V(\Gamma)}(M, \infty)$  of  $y$ . Since  $f_\Gamma$  is a closed map ( $\bar{C}_{V(\Gamma)}(M, \infty)$  is compact),  $f_\Gamma|_U$  is a homeomorphism onto  $f_\Gamma(U)$ . Since  $f_\Gamma(\bar{C}_{V(\Gamma)}(M, \infty) - U)$  is a closed subset of  $\bar{C}_\Gamma(M, \infty)$  not containing  $x$ , there is a neighborhood  $V \subset \bar{C}_\Gamma(M, \infty)$  of  $x$  such that  $V \cap f_\Gamma(\bar{C}_{V(\Gamma)}(M, \infty) - U) = \emptyset$ , so  $V \subset f_\Gamma(U)$ . This shows that  $\bar{C}_\Gamma(M, \infty)$  has the structure of a topological manifold with boundary in the neighborhood  $V$  of  $x$ , completing the proof.  $\square$

**Corollary 3.17**  $\dim_t(T_2) \leq d|V(\Gamma)| - 2.$

**Lemma 3.18** *Let  $Y$  be a compact metrizable space with  $\dim_t(Y) = n$ , and  $Y_2 \subset Y_1 \subset Y$  be closed subspaces such that  $\dim_t(Y_1) \leq n - 1$  and  $\dim_t(Y_2) \leq n - 2$ . Then, for every open cover  $\mathfrak{U}$  of  $Y$ , there exists a refinement  $\mathfrak{U}'$  of  $\mathfrak{U}$  such that*

(\*) *there are open neighborhoods  $N_{Y_1}$  of  $Y_1$  and  $N_{Y_2}$  of  $Y_2$  such that for all  $U_0, \dots, U_n \in \mathfrak{U}'$ , pairwise distinct,*

$$(U_0 \cap \dots \cap U_n) \cap N_{Y_1} = \emptyset, \quad (U_0 \cap \dots \cap U_{n-1}) \cap N_{Y_2} = \emptyset.$$

Hence, if  $S \subset Y_1$  is a closed subset such that  $Y_1 - Y_2 \subset S$ , then there are canonical isomorphisms

$$H^n(Y, S) \approx H^n(Y, Y_1) \approx H_c^n(Y - Y_1).$$

**Proof** The proof of [9, Lemma 21.2.1] goes through almost verbatim here and gives us the first statement. (We first use [13, Proposition 12-9(1)(3)] where  $Y_1, Y_2$  are plugged in as  $C_1, C_2$ , and then use [13, Proposition 9-3].) The second statement easily follows using standard arguments in Čech cohomology. For the first isomorphism: the restriction map  $\check{C}_{\mathfrak{U}}^i(Y, Y_1) \rightarrow \check{C}_{\mathfrak{U}}^i(Y, S)$  is an equality for all  $i \geq n - 1$  and all open covers  $\mathfrak{U}$  of  $Y$  satisfying (\*), so  $\varinjlim_{\mathfrak{U} \text{ satisfying } (*)} H_{\mathfrak{U}}^n(Y, Y_1) = \varinjlim_{\mathfrak{U} \text{ satisfying } (*)} H_{\mathfrak{U}}^n(Y, S)$ ; since every open cover of  $Y$  has a refinement satisfying (\*), these two limits are equal to  $H^n(Y, Y_1), H^n(Y, S)$ ,

respectively. For the second isomorphism: this follows from [2, Proposition 12-3]. Alternatively, let  $\Phi$  be the collection of compact subsets of  $Y - Y_1$ , viewed as subsets of  $Y$ , and it is not hard to check directly that the natural restriction maps  $H_{\Phi}^n(Y) \rightarrow H_c^n(Y - Y_1)$  and  $H_{\Phi}^n(Y) \rightarrow H^n(Y, Y_1)$  are isomorphisms.  $\square$

**Corollary 3.19**  $H^{d|V(\Gamma)|}(X_{\Gamma}, S) \approx H^{d|V(\Gamma)|}(X_{\Gamma}, T_1) \approx H_c^{d|V(\Gamma)|}(X_{\Gamma} - T_1)$  via  $\mathcal{G}$ -equivariant isomorphisms.

### 3.3 Gluing codimension-1 faces

The goal of this Section is to construct a nontrivial map  $\rho : H_c^{d|V(\Gamma)|}(X_{\Gamma} - T_1) \rightarrow R$ , and the method is to realize  $X_{\Gamma} - T_1$  as the image of a proper map from an oriented topological manifold of dimension  $d|V(\Gamma)|$ . An alternative approach for constructing  $\rho$  is given in Section 3.4 (which we will only sketch), and it is a better and more canonical approach. But the method here is less technical and easier to write, so we use it instead.

Recall at the end of Section 2.2 we defined, for every  $A \subset \{\infty\} \sqcup V(\Gamma)$  of type 2,  $\sigma_A \in \tilde{S}_{E(\Gamma)}$ , and for every  $\Gamma$ -pair  $A_1, A_2 \in E(\Gamma)$ ,  $\sigma_{A_1 A_2} \in \tilde{S}_{E(\Gamma)}$ . Thus, we have

$$\phi(\sigma_A), \phi(\sigma_{A_1 A_2}) : \bar{C}_2(M, \infty)^{E(\Gamma)} \longrightarrow \bar{C}_2(M, \infty)^{E(\Gamma)}$$

as in Definition 3.3.

**Lemma 3.20**  $\phi(\sigma_A)(\bar{\mathcal{P}}_A^{\Gamma}) = \bar{\mathcal{P}}_A^{\Gamma}$  and  $\phi(\sigma_{A_1 A_2})(\bar{\mathcal{P}}_{A_1}^{\Gamma}) = \bar{\mathcal{P}}_{A_2}^{\Gamma}$ .

**Proof** Let  $A$  be of type 2 with chosen bivalent vertex  $v_A$ , and vertices  $v_A^1$  and  $v_A^2$  adjacent to it. There is a dense open subset  $\mathring{\mathcal{P}}'_A \subset \mathring{\mathcal{P}}_A$  on which we can define an involution  $\phi'_A : \mathring{\mathcal{P}}'_A \rightarrow \mathring{\mathcal{P}}'_A$  which fixes all other marked points and reflects the point labeled by  $v_A$  along the midpoint of the line segment between the points labeled by  $v_A^1$  and  $v_A^2$ , on the screen that these marked points lie on. (This argument is in Kontsevich’s original paper [7, Lemma 2.1]; see e.g., [20, Figure 16] for a nice picture. Notice here we use  $\mathring{\mathcal{P}}'_A$  because  $\phi'_A$  is not well defined on  $\mathring{\mathcal{P}}_A$ , due to cases when the new position of the point labeled by  $v_A$  coincides with other points.) Clearly  $\phi(\sigma_A) \circ f_{\Gamma} = f_{\Gamma} \circ \phi'_A$ . So

$$\phi(\sigma_A)(\bar{\mathcal{P}}_A^{\Gamma}) = \phi(\sigma_A)(f_{\Gamma}(\bar{\mathcal{P}}_A)) = \overline{\phi(\sigma_A)(f_{\Gamma}(\mathring{\mathcal{P}}'_A))} = \overline{f_{\Gamma}(\phi'_A(\mathring{\mathcal{P}}'_A))} = \overline{f_{\Gamma}(\mathring{\mathcal{P}}'_A)} = \bar{\mathcal{P}}_A^{\Gamma}.$$

Let  $A_1, A_2$  be a  $\Gamma$ -pair. Since for  $A$  of type 4,  $\mathring{\mathcal{P}}_A$  is an  $S^{d-1}$ -bundle over  $C_{V(\Gamma/A)}(M, \infty)$  with fiber over  $x$  canonically identified with  $ST_{f_{[A]_v}(x)}M$ , we can define  $\phi'_{A_1 A_2} : \mathring{\mathcal{P}}_{A_1} \rightarrow \mathring{\mathcal{P}}_{A_2}$  by lifting the map  $C_{V(\Gamma/A_1)}(M) \rightarrow C_{V(\Gamma/A_2)}(M)$  switching marked points in the same way as  $\alpha_{A_1 A_2}$  (defined by the end of Section 2.2) maps vertices of  $\Gamma/A_1$  to vertices of  $\Gamma/A_2$ . Since  $[A_1]_v$  is mapped to  $[A_2]_v$ , the fibers are canonically identified. By the definition of  $\sigma_{A_1 A_2}$ ,  $\phi(\sigma_{A_1 A_2}) \circ f_{\Gamma} = f_{\Gamma} \circ \phi'_{A_1 A_2}$ . So

$$\begin{aligned} \phi(\sigma_{A_1 A_2})(\bar{\mathcal{P}}_{A_1}^{\Gamma}) &= \phi(\sigma_{A_1 A_2})(f_{\Gamma}(\bar{\mathcal{P}}_{A_1})) \\ &= \overline{\phi(\sigma_{A_1 A_2})(f_{\Gamma}(\mathring{\mathcal{P}}_{A_1}))} = \overline{f_{\Gamma}(\phi'_{A_1 A_2}(\mathring{\mathcal{P}}_{A_1}))} = \overline{f_{\Gamma}(\mathring{\mathcal{P}}_{A_2})} = \bar{\mathcal{P}}_{A_2}^{\Gamma}. \end{aligned} \quad \square$$

**Corollary 3.21**  $\phi(\sigma_A)(\mathring{\mathring{\mathcal{P}}}_A^{\Gamma}) = \mathring{\mathring{\mathcal{P}}}_A^{\Gamma}$  and  $\phi(\sigma_{A_1 A_2})(\mathring{\mathring{\mathcal{P}}}_{A_1}^{\Gamma}) = \mathring{\mathring{\mathcal{P}}}_{A_2}^{\Gamma}$ .

Denote

$$C'_\Gamma(M, \infty) := C_\Gamma(M, \infty) \cup \bigcup_{A \text{ of type 2 or 4}} \overset{\circ\circ}{\mathcal{P}}_A^\Gamma \subset \bar{C}_\Gamma(M, \infty).$$

Then by the definition of  $\overset{\circ\circ}{\mathcal{P}}_A^\Gamma$ ,  $C'_\Gamma(M, \infty)$  is a topological manifold with boundary.

**Definition 3.22** Take  $2^{|E(\Gamma)|} |E(\Gamma)|!$  copies of  $C'_\Gamma(M, \infty)$ , labeled by elements in  $\tilde{S}_{E(\Gamma)}$ . We write  $C'_\Gamma(M, \infty)^\sigma$  for the copy labeled by  $\sigma \in \tilde{S}_{E(\Gamma)}$ , and similarly write  $(\overset{\circ\circ}{\mathcal{P}}_A^\Gamma)^\sigma$ ,  $x^\sigma$ , etc., for its subspaces and elements. We orient  $C'_\Gamma(M, \infty)^\sigma$  by twisting the orientation on  $C_\Gamma(M, \infty)$  by  $(-1)^{(d-1)\text{sgn}(\sigma)+d\text{sgn}'(\sigma)}$ .

Define

$$\tilde{X}_\Gamma := \left( \bigsqcup_{\sigma \in \tilde{S}_{E(\Gamma)}} C'_\Gamma(M, \infty)^\sigma \right) / \sim_\Gamma,$$

where  $\sim_\Gamma$  is the following equivalence relation (gluing boundary components pairwise):

- for all  $A \subset \{\infty\} \sqcup V(\Gamma)$  of type 2,  $x \in \overset{\circ\circ}{\mathcal{P}}_A^\Gamma$  and  $\sigma \in \tilde{S}_{E(\Gamma)}$ ,  $x^\sigma \sim_\Gamma (\phi(\sigma_A)(x))^{\sigma\sigma_A^{-1}}$ ;
- for all  $A_1, A_2$  a  $\Gamma$ -pair,  $x \in \overset{\circ\circ}{\mathcal{P}}_{A_1}^\Gamma$  and  $\sigma \in \tilde{S}_{E(\Gamma)}$ ,  $x^\sigma \sim_\Gamma (\phi(\sigma_{A_1 A_2})(x))^{\sigma\sigma_{A_1 A_2}^{-1}}$ .

Moreover, define

$$\tilde{f} = (\tilde{f}_e)_{e \in E(\Gamma)} : \tilde{X}_\Gamma \longrightarrow \bar{C}_2(M, \infty)^{E(\Gamma)}, \quad \tilde{f}|_{C'_\Gamma(M, \infty)^\sigma} = \phi(\sigma) \circ f_\Gamma.$$

It is well defined since  $\phi(\sigma\sigma_A^{-1}) \circ \phi(\sigma_A) = \phi(\sigma)$  and  $\phi(\sigma\sigma_{A_1 A_2}^{-1}) \circ \phi(\sigma_{A_1 A_2}) = \phi(\sigma)$ .

It can be easily seen that  $\text{image}(\tilde{f}) = X_\Gamma - T_1$ . It follows from Lemma 3.5 and the definition above that  $\tilde{f}|_{\bigsqcup_\sigma C'_\Gamma(M, \infty)^\sigma}$  is a covering map (onto its image) of degree  $|\text{Aut}^u(\Gamma)|_d^\pm$ . Since  $\phi(\sigma)$  is  $\mathcal{G}$ -equivariant for all  $\sigma \in \tilde{S}_{E(\Gamma)}$ , the diagonal  $\mathcal{G}$ -action on  $\bar{C}_2(M, \infty)^{E(\Gamma)}$  lifts to an action of  $\mathcal{G}$  on  $\tilde{X}_\Gamma$ , so that  $\tilde{f}$  is equivariant.

**Lemma 3.23**  $\tilde{X}_\Gamma$  is an oriented topological manifold of dimension  $d|V(\Gamma)|$ .

**Proof** That it is a topological manifold of dimension  $d|V(\Gamma)|$  follows from that  $\phi(\sigma_A) : \overset{\circ\circ}{\mathcal{P}}_A^\Gamma \rightarrow \overset{\circ\circ}{\mathcal{P}}_A^\Gamma$  and  $\phi(\sigma_{A_1 A_2}) : \overset{\circ\circ}{\mathcal{P}}_{e_1}^\Gamma \rightarrow \overset{\circ\circ}{\mathcal{P}}_{e_2}^\Gamma$  are homeomorphisms and  $\sim_\Gamma$  glues together these boundary components of  $\bigsqcup_{\sigma \in \tilde{S}_{E(\Gamma)}} C'_\Gamma(M, \infty)^\sigma$  pairwise. It is not difficult to verify that  $\phi(\sigma_A)$  and  $\phi(\sigma_{A_1 A_2})$  are orientation-reversing, if the  $(\overset{\circ\circ}{\mathcal{P}}_A^\Gamma)^\sigma$ 's are oriented as boundaries of  $C'_\Gamma(M, \infty)$ .  $\square$

The purpose of the following lemma is to show that  $\tilde{f}$  induces a map

$$\tilde{f}^* : H_c^{d|V(\Gamma)|}(X_\Gamma - T_1) \longrightarrow H_c^{d|V(\Gamma)|}(\tilde{X}_\Gamma).$$

**Lemma 3.24**  $\tilde{f} : \tilde{X}_\Gamma \longrightarrow X_\Gamma - T_1$  is a proper map.

**Proof** Let  $K \subset X_\Gamma - T_1$  be compact. To show  $\tilde{f}^{-1}(K)$  is compact, let  $\{x_n\}_{n=1}^\infty$  be a sequence of points in  $\tilde{f}^{-1}(K)$ . There is a subsequence (still call it  $\{x_n\}$ ) and some  $\sigma \in \tilde{S}_{E(\Gamma)}$  such that  $x_n \in C'_\Gamma(M, \infty)^\sigma$  for all  $n$ . After possibly passing to a subsequence,  $\tilde{f}(x_n)$  converges to some  $y \in \phi(\sigma)(\bar{C}_\Gamma(M, \infty)) \cap K$ , since  $\tilde{f}(x_n) \in \phi(\sigma)(C'_\Gamma(M, \infty))$  and  $\bar{C}_\Gamma(M, \infty)$  is closed. But  $y \notin \phi(\sigma)(\overset{\circ\circ}{\mathcal{P}}_A^\Gamma - \overset{\circ\circ}{\mathcal{P}}_A^\Gamma)$  for any  $A$ , since  $\phi(\sigma)(\overset{\circ\circ}{\mathcal{P}}_A^\Gamma - \overset{\circ\circ}{\mathcal{P}}_A^\Gamma) \subset T_1$ . Thus,  $y \in \phi(\sigma)(C'_\Gamma(M, \infty))$ . Since  $\tilde{f}$  maps  $C'_\Gamma(M, \infty)^\sigma$  homeomorphically

onto  $\phi(\sigma)(C'_\Gamma(M, \infty))$ , by the definition of  $C'_\Gamma(M, \infty)$ ,  $\{x_n\}$  converges to the unique element  $x \in \tilde{f}^{-1}(y) \cap C'_\Gamma(M, \infty)^\sigma$ .  $\square$

**Definition 3.25** Define  $\rho : H^{d|V(\Gamma)|}(X_\Gamma, S; R) \longrightarrow R$  to be the composition

$$H^{d|V(\Gamma)|}(X_\Gamma, S) \xrightarrow{\text{Corollary 3.19}} H_c^{d|V(\Gamma)|}(X_\Gamma - T_1) \xrightarrow{\tilde{f}^*} H_c^{d|V(\Gamma)|}(\tilde{X}_\Gamma) \longrightarrow R,$$

where the last arrow is by taking the cap product with the fundamental class of  $\tilde{X}_\Gamma$  (in the sense of Borel–Moore homology).

Since the  $\mathcal{G}$ -action on  $\tilde{X}_\Gamma$  is orientation-preserving, the last map is  $\mathcal{G}$ -equivariant (where  $R$  is equipped with the trivial action). So  $\mathcal{G}$  acts on all the objects involved in this definition and all maps involved are equivariant, implying that  $\rho$  is  $\mathcal{G}$ -equivariant.

**Remark 3.26** If  $\Gamma$  is not a single graph but a formal sum of graphs,  $\sum_{i=1}^m \Gamma_i$ , the arguments in Section 3 need to be modified as follows. First, note that if two graphs share a common boundary term, then they must have the same number of edges; therefore without loss of generality we can assume all the  $\Gamma_i$  have the same number of edges, say  $n$ . Then, let us fix, for each  $\Gamma_i$ , a bijection between  $E(\Gamma_i)$  and  $\{1, \dots, n\}$ . Since we will soon be summing over all permutations of edges, this choice does not matter. Our “ambient space” in this case will be  $\bar{C}_2(M, \infty)^n$ , in place of  $\bar{C}_2(M, \infty)^{E(\Gamma)}$  in the single-graph case. Now, for each  $\Gamma_i$ , we can define subsets  $C_{\Gamma_i}(M, \infty), \bar{C}_{\Gamma_i}(M, \infty) \subset \bar{C}_2(M, \infty)^n$ , just like in the single-graph case, using the chosen bijection. Similar to Lemma 3.5, it is easy to see that the elements in  $\{\phi(\sigma)(C_{\Gamma_i}(M, \infty))\}_{i \in \{1, \dots, m\}, \sigma \in \tilde{S}_{\{1, \dots, n\}}}$  are disjoint from each other. We define

$$X_{\sum_{i=1}^m \Gamma_i} = \bigcup_{\sigma \in \tilde{S}_{\{1, \dots, n\}}, i \in \{1, \dots, m\}} \phi(\sigma)(\bar{C}_{\Gamma_i}(M, \infty)) \subset \bar{C}_2(M, \infty)^n.$$

The rest of the arguments in the paper all generalize to this case in a straightforward way.

### 3.4 Digression: what $X_\Gamma$ looks like

This subsection can be skipped. It is here to justify Figure 1: after removing a codimension-2 subset from  $X_\Gamma$ , it looks like a manifold with boundary and bindings, where each binding component is connected to an even number of pages, summing up to 0 when counted with sign. This statement would also allow us to define  $\rho$  in a different (more canonical) way than in Section 3.3. But working out everything precisely is quite technically involved, so we will only sketch such an approach in this subsection.

By Lemma 3.5,

$$X_\Gamma = \bigsqcup_{[\sigma] \in \tilde{S}_{E(\Gamma)}/\text{image}(\psi_\Gamma)} \phi(\sigma)(C_\Gamma(M, \infty)) \sqcup \bigcup_{\sigma \in \tilde{S}_{E(\Gamma)}} \phi(\sigma)(\bar{C}_\Gamma(M, \infty) - C_\Gamma(M, \infty)).$$

Denote the first term above by  $\overset{\circ}{X}_\Gamma$ .

First we define “binding points” (those  $p$  in the definition below).

**Definition 3.27** For  $A \subset \{\infty\} \sqcup V(\Gamma)$ ,  $|A| \geq 2$ , define  $\overset{\infty}{\mathcal{P}}_A^\Gamma \subset \mathcal{P}_A^\Gamma$  to be the set of points  $p$  satisfying that there exists a neighborhood  $U \subset \bar{C}_2(M, \infty)^{E(\Gamma)}$  of  $p$ , such that for every  $\sigma \in \tilde{S}_{E(\Gamma)}$ , either

(1)  $\phi(\sigma)(\bar{C}_\Gamma(M, \infty)) \cap U = \emptyset$ , or

(2)  $p = \sigma(q)$  for some  $A' \subset \{\infty\} \sqcup V(\Gamma)$ ,  $|A'| \geq 2$ ,  $q \in \mathcal{P}_{A'}^\Gamma$ , and

- there is a homeomorphism

$$\nu : U \cap \phi(\sigma)(\bar{C}_\Gamma(M, \infty)) \longrightarrow \mathbb{R}^{N-1} \times \mathbb{R}^{\geq 0} \quad \text{such that } \nu(U \cap \phi(\sigma)(\mathcal{P}_{A'}^\Gamma)) = \mathbb{R}^{N-1} \times \{0\}$$

(i.e.,  $\phi(\sigma)(\bar{C}_\Gamma(M, \infty))$  is a topological manifold with boundary near  $p$ );

- $U \cap \phi(\sigma)(\mathcal{P}_{A'}^\Gamma) = U \cap \mathcal{P}_A^\Gamma$ .

For  $p \in \overset{\infty}{\mathcal{P}}_A^\Gamma$ , define the *signed count of pages at  $p$*  to be the signed count of elements in  $\tilde{S}_{E(\Gamma)}$ : those  $\sigma$  of case (1) above are counted with 0; those  $\sigma$  of case (2) above are counted with  $\pm 1$ ,  $+1$  if the boundary orientations of  $\mathcal{P}_A^\Gamma$  near  $p$ , as boundary of  $\phi(\sigma)(\bar{C}_\Gamma(M, \infty))$  and as boundary of  $\bar{C}_\Gamma(M, \infty)$ , agree, and  $-1$  if they disagree.

**Lemma 3.28** *If  $p \in \overset{\infty}{\mathcal{P}}_A^\Gamma$  where  $A$  is of type 2 or 4, then the signed count of pages at  $p$  is always 0.*

**Sketch of proof** For type 2  $A$ 's, the pages come in pairs of opposite signs; see Lemma 3.20. For type 4  $A$ 's, the pages sum up to 0 because  $\Gamma$  is closed in graph homology.  $\square$

In this subsection,  $\overset{\infty}{\mathcal{P}}_A^\Gamma$  replaces the role played by  $\mathcal{P}_A^\Gamma$  in the previous section. We define  $S$ ,  $T_1$  and  $T_2$  verbatim as in Definition 3.8, just with  $\overset{\infty}{\mathcal{P}}_A^\Gamma$  replaced by  $\mathcal{P}_A^\Gamma$ . The statements in the paragraph below Definition 3.8 still hold. We next show that analogues of Corollaries 3.17 and 3.19 still hold.

**Lemma 3.29** *There is a  $\mathcal{G}$ -equivariant surjective map  $H_c^{d|V(\Gamma)|}(X_\Gamma - T_1; R) \rightarrow R$ , where  $\mathcal{G}$  acts on  $R$  trivially.*

**Sketch of proof**  $X_\Gamma - T_1$  consists of two parts:  $\overset{\circ}{X}_\Gamma$  is an open subset of  $X_\Gamma - T_1$  which is also an  $d|V(\Gamma)|$ -dimensional (oriented) topological manifold, and  $Y := X_\Gamma - T_1 - \overset{\circ}{X}_\Gamma$  is a closed subset of  $X_\Gamma - T_1$  which is also a  $(d|V(\Gamma)|-1)$ -dimensional topological manifold (this follows from the definition of book binding points). It also follows from the definition of book binding points that  $X_\Gamma - T_1$  is locally contractible. So we have the long exact sequence of compactly supported cohomology,

$$\dots \longrightarrow H_c^{d|V(\Gamma)|-1}(Y) \xrightarrow{\delta} H_c^{d|V(\Gamma)|}(\overset{\circ}{X}_\Gamma) \longrightarrow H_c^{d|V(\Gamma)|}(X_\Gamma - T_1) \longrightarrow H_c^{d|V(\Gamma)|}(Y) \longrightarrow \dots,$$

where the last term is 0. Denote by  $J_1$  and  $J_2$  the set of connected components of  $Y$  and  $\overset{\circ}{X}_\Gamma$ , respectively. Then

$$H_c^{d|V(\Gamma)|-1}(Y) \approx R^{\oplus J_1}, \quad H_c^{d|V(\Gamma)|}(\overset{\circ}{X}_\Gamma) \approx R^{\oplus J_2},$$

and  $\delta$  is the coboundary map. So, by Lemma 3.28, the image of  $\delta$  is contained in  $\{(r_i)_{i \in J_2} \mid \sum_i r_i = 0\}$ . Therefore, the map

$$H_c^{d|V(\Gamma)|}(\overset{\circ}{X}_\Gamma) \longrightarrow R, \quad (r_i)_{i \in J_2} \longrightarrow \sum_{i \in J_2} r_i$$

induces a surjective map from the quotient  $H_c^{d|V(\Gamma)|}(\overset{\circ}{X}_\Gamma)/\text{image}(\delta) \approx H_c^{d|V(\Gamma)|}(X_\Gamma - T_1)$  to  $R$ , as desired.  $\square$

**Lemma 3.30**  $\dim_t(T_2) \leq d|V(\Gamma)| - 2.$

Together with Lemma 3.18, Lemma 3.30 implies that  $H^{d|V(\Gamma)|}(X_\Gamma, S) \approx H_c^{d|V(\Gamma)|}(X_\Gamma - T_1)$ , and we can thus define  $\rho : H^{d|V(\Gamma)|}(X_\Gamma, S) \rightarrow R$  using the map in Lemma 3.29. The rest of this subsection is devoted to the following

**Sketch of proof of Lemma 3.30** Recall  $T_2$  consists of points in  $X_\Gamma - \overset{\circ}{X}_\Gamma$  that are not binding points. So, we need to analyze, for  $A, A' \in V(\Gamma) \sqcup \{\infty\}$  and  $\sigma \in \tilde{S}_{E(\Gamma)}$ , how  $\mathcal{G}_A^\Gamma$  and  $\phi(\sigma)(\mathcal{G}_{A'}^\Gamma)$  intersect. Suppose  $A, A'$  and  $\sigma$  are such that they do intersect. Then, by the same reasoning as in Lemma 3.5, there exists an unordered, unoriented graph isomorphism  $\Gamma/\Gamma_{A'} \rightarrow \Gamma/\Gamma_A$  whose edge permutation is given by the restriction of  $\sigma$  to  $E(\Gamma/\Gamma_{A'})$ . So  $\sigma$  also restricts to a bijection  $E(\Gamma_{A'}) \rightarrow E(\Gamma_A)$ . Abusing notation we still denote by  $\phi(\sigma)$  the map  $(S^{d-1})^{E(\Gamma_{A'})} \rightarrow (S^{d-1})^{E(\Gamma_A)}$ , permuting factors according to  $\sigma$  and composing with the antipodal map when there is a negative sign. Recall the notation “ $f_{\Gamma'}^{\mathbb{R}^d}$ ” in Definition 3.11. Denote

$$V_{A'}^\sigma = \phi(\sigma)(f_{\Gamma_{A'}}^{\mathbb{R}^d}(C_{V(\Gamma_{A'})}^{\text{quo}}(\mathbb{R}^d))) \subset (S^{d-1})^{E(\Gamma_A)}, \quad V_A = f_{\Gamma_A}^{\mathbb{R}^d}(C_{V(\Gamma_A)}^{\text{quo}}(\mathbb{R}^d)) \subset (S^{d-1})^{E(\Gamma_A)}.$$

Then  $\mathcal{G}_A^\Gamma \cap \phi(\sigma)(\mathcal{G}_{A'}^\Gamma)$  is a fiber bundle over  $C_{\Gamma/A}(M, \infty)$  with fiber  $V_A \cap V_{A'}^\sigma \subset (S^{d-1})^{E(\Gamma_A)}$ .

**Lemma 3.31** *Let  $\Gamma'$  be a graph. Then  $f_{\Gamma'}^{\mathbb{R}^d}(C_{V(\Gamma')}^{\text{quo}}(\mathbb{R}^d)) \subset (S^{d-1})^{E(\Gamma')} \subset (\mathbb{R}^d)^{E(\Gamma')}$ , where  $S^{d-1}$  is viewed as the unit sphere in  $\mathbb{R}^d$ , is semialgebraic.*

**Proof** It is the image of the composition of a linear map  $(\mathbb{R}^d)^{V(\Gamma')} - \Delta_{\text{big}} \rightarrow (\mathbb{R}^d - \{0\})^{E(\Gamma')}$  with the projection map  $(\mathbb{R}^d - \{0\})^{E(\Gamma')} \rightarrow (S^{d-1})^{E(\Gamma')}$ . Both of these maps’ graphs are semialgebraic. So the image is also semialgebraic by Tarski–Seidenberg theorem.  $\square$

By the above lemma,  $V_{A'}^\sigma$  and  $V_A$  are semialgebraic. They are also open subsets of  $(S^{d-1})^{E(\Gamma_A)}$ . Let  $Y_{A'}^\sigma, Y_A \subset (S^{d-1})^{E(\Gamma_A)}$  be minimal algebraic sets containing  $V_{A'}^\sigma, V_A$ , respectively. Then  $V_{A'}^\sigma \subset Y_{A'}^\sigma$  and  $V_A \subset Y_A$  are open (in Euclidean topology), and the Krull dimensions of both  $Y_A$  and  $Y_{A'}^\sigma$  are  $d|V(\Gamma_A)| - d - 1$ . Denote by  $Z(A, A', \sigma)$  the union of irreducible components of  $Y_{A'}^\sigma \cap Y_A$  whose Krull dimension is less than  $d|V(\Gamma_A)| - d - 1$ . Suppose  $p \in V_A \cap V_{A'}^\sigma - Z(A, A', \sigma)$ , then  $p$  is in some irreducible component  $Y$  of  $Y_A \cap Y_{A'}^\sigma$ , whose Krull dimension is  $d|V(\Gamma_A)| - d - 1$ , so  $Y$  must also be an irreducible component of both  $Y_A$  and  $Y_{A'}^\sigma$ . Since  $p \notin Z(A, A', \sigma)$ , there exists a neighborhood  $U_p \subset (S^{d-1})^{E(\Gamma_A)}$  of  $p$  such that  $V_A \cap U_p = Y \cap U_p = V_{A'}^\sigma \cap U_p$ .

Now, define  $\tilde{Z}(A, A', \sigma)$  to be the subfiber bundle of  $\mathcal{G}_A^\Gamma$  over  $C_{\Gamma/A}(M, \infty)$  whose fibers are

$$Z(A, A', \sigma) \cap V_A.$$

For a given  $A \subset V(\Gamma) \sqcup \{\infty\}$ , define  $\mathcal{G}_A^{\Gamma, \text{rmv}} = \bigcup_{\sigma, A'} \tilde{Z}(A, A', \sigma)$ , where “rmv” stands for “remove”. It can be shown that the Krull dimension of an algebraic subset of  $\mathbb{R}^n$  is equal to its covering dimension in Euclidean topology, so  $\dim_t(\mathcal{G}_A^{\Gamma, \text{rmv}}) \leq d|V(\Gamma)| - 2$ . By the conclusion of the previous paragraph,

every point in  $\bar{\mathcal{G}}_A^\Gamma$  which is not in (1)  $\mathcal{G}_A^{\Gamma, \text{rmv}}$  or (2)  $\mathcal{G}_A^\Gamma - \overset{\circ}{\mathcal{G}}_A^\Gamma$  (as in Definition 3.27) or (3) the image of some codimension at least 2 stratum of  $\bar{C}_{V(\Gamma)}(M, \infty)$  under  $f_\Gamma$ , is a binding point, by the definition of binding points. By Lemma 3.16,  $\dim_t(\mathcal{G}_A^\Gamma - \overset{\circ}{\mathcal{G}}_A^\Gamma) \leq d|V(\Gamma)| - 2$ . So, the union of the above three sets has covering dimension at most  $d|V(\Gamma)| - 2$ .  $\square$

### 4 Reconstructing Kontsevich's characteristic classes

In this section we consider the fiber bundle version of the spaces in Section 3 and give a definition of Kontsevich's classes using only the data allowed in Theorem 1.2.

Recall  $R = \mathbb{Z}$  or  $\mathbb{R}$  and all cohomology in this section are with  $R$ -coefficients, which we omit. Let  $\Gamma$  be as in Section 2.2; assume also that  $\Gamma$  is trivalent. Although we consider a single graph for simplicity, the content of this section generalizes verbatim to when  $\Gamma$  is a linear combination of graphs, as long as the spaces are modified according to Remark 3.26. Let  $M$  be as in Section 1.1. Assume  $d \geq 3$ . Let  $\pi : E \rightarrow B$  be a framed smooth  $(M, \infty)$ -fiber bundle as in Section 1.1. We assume  $B$  can be given a CW-structure. Using CW-approximation, Definition 4.10 and thus Corollary 4.11 generalize to cases where  $B$  is just a paracompact Hausdorff space.

Let

$$\begin{array}{cccccccccccc}
 C_2(\pi) & \bar{C}_2(\pi) & \bar{C}_2^{E(\Gamma)}(\pi) & \bar{C}_\Gamma(\pi) & \mathcal{G}_A(\pi) & \mathcal{G}_A^\Gamma(\pi) & \bar{\mathcal{G}}_A^\Gamma(\pi) & X_\Gamma(\pi) & T_1(\pi) & T_2(\pi) & S(\pi) & \tilde{X}_\Gamma(\pi) \\
 \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow^{\pi_X} & \downarrow & \downarrow & \downarrow & \downarrow^{\pi_{\tilde{X}}} \\
 B & B & B & B & B & B & B & B & B & B & B & B
 \end{array}$$

be the associated bundles of  $\pi$  with fibers  $C_2(M, \infty)$ ,  $\bar{C}_2(M, \infty)$ ,  $\bar{C}_2(M, \infty)^{E(\Gamma)}$ ,  $\bar{C}_\Gamma(M, \infty)$ ,  $\mathcal{G}_A$ ,  $\mathcal{G}_A^\Gamma$ ,  $\bar{\mathcal{G}}_A^\Gamma$ ,  $X_\Gamma$ ,  $T_1$ ,  $T_2$ ,  $S$ ,  $\tilde{X}_\Gamma$ , respectively. (All of these spaces, except for  $\tilde{X}_\Gamma$ , are defined in Section 3.1;  $\tilde{X}_\Gamma$  is defined in Definition 3.22.) Correspondingly, the maps  $f_\pm$ ,  $f_\Gamma$ ,  $\tilde{f}$  in Section 3.1 induce bundle maps. Abusing notation, we still denote them by  $f_\pm$ ,  $f_\Gamma$ ,  $\tilde{f}$ . Notice that  $\bar{C}_2^{E(\Gamma)}(\pi)$  denotes the fiber product

$$\underbrace{\bar{C}_2(\pi) \times_B \bar{C}_2(\pi) \times_B \cdots \times_B \bar{C}_2(\pi)}_{E(\Gamma) \text{ times}}$$

while  $\bar{C}_2(\pi)^{E(\Gamma)}$  denotes the direct product of the total space

$$\underbrace{\bar{C}_2(\pi) \times \bar{C}_2(\pi) \times \cdots \times \bar{C}_2(\pi)}_{E(\Gamma) \text{ times}},$$

ignoring the fiber bundle structure.

**Lemma 4.1** *Under the condition of Theorem 1.2,  $(\tilde{h}, h)$  induces  $\mathcal{G}$ -bundle maps between the  $\pi'$  and  $\pi''$  version of all the bundles above.*

**Proof** By the existence of  $\tilde{h}$ ,  $h$  is a  $\mathcal{G}$ -bundle map. Since every space defined in Section 3.1 has induced  $\mathcal{G}$ -action and the maps between them defined in Section 3.1 are all  $\mathcal{G}$ -equivariant,

$$\tilde{h}^{E(\Gamma)} : \bar{C}_2(\pi')^{E(\Gamma)} \longrightarrow \bar{C}_2(\pi'')^{E(\Gamma)}$$

restricts to  $\mathcal{G}$ -bundle maps

$$\begin{aligned} \tilde{h}^{E(\Gamma)} : \bar{C}_2^{E(\Gamma)}(\pi') &\rightarrow \bar{C}_2^{E(\Gamma)}(\pi''), & \tilde{h}_\Gamma : \bar{C}_\Gamma(\pi') &\rightarrow \bar{C}_\Gamma(\pi''), \\ \tilde{h}_X : X_\Gamma(\pi') &\rightarrow X_\Gamma(\pi''), & \tilde{h}_S : S(\pi') &\rightarrow S(\pi''), \quad \text{etc.} \end{aligned} \quad \square$$

The framing on  $\pi$  induces a map  $F : \partial^v \bar{C}_2(\pi) \rightarrow S^{d-1}$ , as in [20, Section 2.4.3] (it is called  $p(\tau_E)$  there). Define  $\sim_F$  to be the following equivalence relation on  $\bar{C}_2(\pi)$ :

$$x \sim_F y \quad \text{if } x = y \in C_2(\pi) = \bar{C}_2(\pi) - \partial^v \bar{C}_2(\pi) \quad \text{or} \quad F(x) = F(y), \quad x, y \in \partial^v \bar{C}_2(\pi).$$

Denote by  $q : \bar{C}_2(\pi) \rightarrow \bar{C}_2(\pi)/\sim_F$  the quotient map by  $\sim_F$ , where the target is equipped with the quotient topology. Let us denote  $S_\pi^{d-1} = q(\partial^v \bar{C}_2(\pi))$  and denote  $(\bar{C}_2(\pi)/\sim_F) - S_\pi^{d-1}$  still by  $C_2(\pi)$ ; then  $\bar{C}_2(\pi)/\sim_F = C_2(\pi) \sqcup S_\pi^{d-1}$  as a set. It is not hard to see that  $\bar{C}_2(\pi)/\sim_F$  is Hausdorff and  $S_\pi^{d-1}$  is a deformation retract of some neighborhood of it (hint: since  $\bar{C}_2(\pi)$  restricted to each cell of  $B$  is a manifold with compact boundary, using a cell-by-cell construction we can find a collar neighborhood of  $\partial^v \bar{C}_2(\pi)$  in  $\bar{C}_2(\pi)$ ). Notice that the orientation on  $M$  specifies an orientation on  $S_\pi^{d-1}$ .

**Definition 4.2** The propagator class  $\Omega(\pi) \in H^{d-1}(\bar{C}_2(\pi)/\sim_F; R)$  is the unique class satisfying that  $\Omega(\pi)|_{S_\pi^{d-1}} \in H^{d-1}(S_\pi^{d-1}; R)$  is the Poincaré dual of the point class.

The existence and uniqueness of such a class follows from the exact sequence

$$H^{d-1}(\bar{C}_2(\pi)/\sim_F, S_\pi^{d-1}) \rightarrow H^{d-1}(\bar{C}_2(\pi)/\sim_F) \rightarrow H^{d-1}(S_\pi^{d-1}) \rightarrow H^d(\bar{C}_2(\pi)/\sim_F, S_\pi^{d-1})$$

and the vanishing of its first and last terms: for  $n = d, d - 1$ ,

$$H^n(\bar{C}_2(\pi)/\sim_F, S_\pi^{d-1}) \approx H^n((\bar{C}_2(\pi)/\sim_F)/S_\pi^{d-1}) = H^n(\bar{C}_2(\pi)/\partial^v \bar{C}_2(\pi)) \approx H^n(\bar{C}_2(\pi), \partial^v \bar{C}_2(\pi))$$

and the vanishing of the last term follows from the proof of [20, Lemma 2.10]. The definition of propagator class here is completely analogous to that in [10].

We next show that  $\Omega(\pi)$  gives us a class  $\Omega_\Gamma(\pi) \in H^{|E(\Gamma)|(d-1)}(X_\Gamma(\pi), S(\pi))$ .

### 4.1 Defining $\Omega_\Gamma$

Define the composition

$$\iota : X_\Gamma(\pi) \subset \bar{C}_2^{E(\Gamma)}(\pi) \subset \bar{C}_2(\pi)^{E(\Gamma)} \xrightarrow{q, \dots, q} (\bar{C}_2(\pi)/\sim_F)^{E(\Gamma)}.$$

We remark that  $\bar{C}_2(\pi)^{E(\Gamma)}$  is the direct product and  $\bar{C}_2^{E(\Gamma)}(\pi)$  is the fiber product. For  $I \subset E(\Gamma)$ , denote by  $p_I : (\bar{C}_2(\pi)/\sim_F)^{E(\Gamma)} \rightarrow (\bar{C}_2(\pi)/\sim_F)^I$  the projection to the  $I$ -factors. If  $I = \{e\}$  we also denote  $p_e = p_I$ . For  $\sigma \in \tilde{S}_{E(\Gamma)}$  and  $A \subset \{\infty\} \sqcup V(\Gamma)$ , define  $I_{\sigma, A} = \sigma(E(\Gamma_A)) \subset E(\Gamma)$ , where we implicitly identify edges of  $\Gamma_A$  also as edges of  $\Gamma$  and abuse notation to still write  $\sigma$  for its image in  $S_{E(\Gamma)}$ .

**Lemma 4.3** Suppose  $A \subset \{\infty\} \sqcup V(\Gamma)$  and  $\sigma \in \tilde{S}_{E(\Gamma)}$ . Denote

$$\bar{V}_A^\sigma = p_{I_{\sigma, A}}(\iota(\phi(\sigma)(\bar{\mathcal{F}}_A^\Gamma(\pi)))) \subset (\bar{C}_2(\pi)/\sim_F)^{I_{\sigma, A}}.$$

Then  $\bar{V}_A^\sigma \subset (S_\pi^{d-1})^{I_{\sigma, A}}$  and it is the image of  $\bar{C}_{V(\Gamma_A)}^{\text{quo}}(\mathbb{R}^d)$  under a smooth map.

**Proof** This is a consequence of the framing  $F$  on  $\pi$  (in the case  $\infty \notin A$ ) and the trivialization of  $\pi$  near  $s_\infty(B)$  (in the case  $\infty \in A$ ). For simplicity we only consider the case  $\sigma = \text{id}$ ; the other cases follow easily. First assume  $\infty \notin A$ . Recall  $\bar{V}_{\Gamma_A} \subset (S^{d-1})^{E(\Gamma_A)}$  as in Definition 3.11. Since the framing  $F$  identifies the vertical tangent space of  $E$  at each point not in  $s_\infty(B)$  with  $\mathbb{R}^d$ ,

$$\mathcal{P}_A^\Gamma(\pi) \approx C_{\Gamma/A}(\pi) \times \bar{V}_{\Gamma_A} \subset C_2(\pi)^{E(\Gamma/A)} \times (S_\pi^{d-1})^{E(\Gamma_A)} \subset (\bar{C}_2(\pi)/\sim_F)^{E(\Gamma/A)} \times (\bar{C}_2(\pi)/\sim_F)^{E(\Gamma_A)},$$

and  $(p_{E(\Gamma_A)} \circ \iota)(\mathcal{P}_A^\Gamma(\pi)) = \bar{V}_{\Gamma_A}$ . Since  $\bar{V}_{\Gamma_A}$  is closed,  $(p_{E(\Gamma_A)} \circ \iota)(\bar{\mathcal{P}}_A^\Gamma(\pi)) = \bar{V}_{\Gamma_A}$  as well.

Now assume  $\infty \in A$ . Since  $\pi$  is trivialized near  $s_\infty(B)$ , the vertical tangent spaces of  $E$  at points in  $s_\infty(B)$  are all identified with  $T_\infty M$ . So

$$\bar{\mathcal{P}}_A(\pi) \approx \bar{C}_{V(\Gamma/A)}(\pi) \times \bar{C}_{V(\Gamma_A)}^{\text{quo}}(T_\infty M),$$

and  $(p_{E(\Gamma_A)} \circ \iota)(\bar{\mathcal{P}}_A^\Gamma(\pi)) = (p_{E(\Gamma_A)} \circ \iota \circ f_\Gamma)(\bar{\mathcal{P}}_A(\pi))$ ; the map  $p_{E(\Gamma_A)} \circ \iota \circ f_\Gamma$  factors through the projection to the second factor above. □

By Lemmas 4.6 and 4.7 below (these are two somewhat technical lemmas whose statements and proofs are postponed until after Proposition 4.5, due to their lengths), the cup product

$$\cup : H^{d-1}(\bar{C}_2(\pi)/\sim_F)^{\otimes E(\Gamma)} \longrightarrow H^{|E(\Gamma)|(d-1)}((\bar{C}_2(\pi)/\sim_F)^{E(\Gamma)}),$$

$$\bigotimes_{i=1}^{|E(\Gamma)|} \sigma_{e_i^\Gamma} \longrightarrow p_{e_1^\Gamma}^* \sigma_{e_1^\Gamma} \cup \dots \cup p_{e_{|E(\Gamma)|}^\Gamma}^* \sigma_{e_{|E(\Gamma)|}^\Gamma}$$

factors through  $H^{|E(\Gamma)|(d-1)}((\bar{C}_2(\pi)/\sim_F)^{E(\Gamma)}, \bigcup_{\sigma, A} p_{I_{\sigma, A}}^{-1}(\bar{V}_A^\sigma))$ , where the union ranges over all  $\sigma \in \tilde{S}_{E(\Gamma)}$  and  $A \subset \{\infty\} \sqcup V(\Gamma)$  of type 3 (same below when we write “ $\bigcup_{\sigma, A}$ ”).

**Definition 4.4**

$$\Omega'_\Gamma(\pi) := p_{e_1^\Gamma}^* \Omega(\pi) \cup \dots \cup p_{e_{|E(\Gamma)|}^\Gamma}^* \Omega(\pi) \in H^{|E(\Gamma)|(d-1)}\left((\bar{C}_2(\pi)/\sim_F)^{E(\Gamma)}, \bigcup_{\sigma, A} p_{I_{\sigma, A}}^{-1}(\bar{V}_A^\sigma)\right),$$

$$\Omega_\Gamma(\pi) := \iota^* \Omega'_\Gamma(\pi) \in H^{|E(\Gamma)|(d-1)}(X_\Gamma(\pi), S(\pi)).$$

**Proposition 4.5** Under the assumptions of Theorem 1.2, let  $\tilde{h}_X : X_\Gamma(\pi') \rightarrow X_\Gamma(\pi'')$  be the  $\mathcal{G}$ -bundle map as in Lemma 4.1. Then  $\tilde{h}_X^*(\Omega_\Gamma(\pi'')) = \Omega_\Gamma(\pi')$ .

**Proof** By the second commutative diagram in Theorem 1.2,  $\tilde{h}$  induces

$$\tilde{h}_q : \bar{C}_2(\pi')/\sim_{F'} \longrightarrow \bar{C}_2(\pi'')/\sim_{F''}$$

which commutes with  $q$  and restricts to the homeomorphism  $h_S : S_{\pi'}^{d-1} \rightarrow S_{\pi''}^{d-1}$ . It follows that  $\tilde{h}_q^* \Omega(\pi'') = \Omega(\pi')$ . Since  $\iota$  commutes with  $\tilde{h}_X$  and  $(\tilde{h}_q)^{E(\Gamma)}$ , by the naturality statement in Lemma 4.6 below,  $\tilde{h}_X^*(\Omega_\Gamma(\pi'')) = \Omega_\Gamma(\pi')$ . □

**Lemma 4.6** Let  $Y = Y_1 \times \cdots \times Y_n$  be a product of paracompact Hausdorff spaces. Let  $s, r \in \mathbb{Z}^{>0}$ . Suppose for all  $i = 1, \dots, r$ , we have  $I_i = \{a_1^i, \dots, a_{m_i}^i\} \subset \{1, \dots, n\}$  and a closed subset  $V_i \subset Y_{a_1^i} \times \cdots \times Y_{a_{m_i}^i}$  satisfying the following condition: every open cover of  $Y$  has a refinement of the form

$$\mathcal{U}_1 \times \cdots \times \mathcal{U}_n := \{U_1 \times \cdots \times U_n \mid U_j \in \mathcal{U}_j\}, \quad \text{where } \mathcal{U}_j \text{ is an open cover of } Y_j$$

such that

(†) for all  $i = 1, \dots, r$ ,  $s'_1, \dots, s'_{m_i} = s$  or  $s - 1$ , with at most one of them being  $s - 1$ ,

$$((U_{a_1^i}^0 \cap \cdots \cap U_{a_1^i}^{s'_1}) \times \cdots \times (U_{a_{m_i}^i}^0 \cap \cdots \cap U_{a_{m_i}^i}^{s'_{m_i}})) \cap V_i = \emptyset,$$

where  $U_{a_j^i}^0, \dots, U_{a_j^i}^{s'_j}$  are any pairwise distinct elements of  $\mathcal{U}_{a_j^i}$ , for every  $j = 1, \dots, m_i$ .

Denote  $p_{I_i} : Y \rightarrow Y_{a_1^i} \times \cdots \times Y_{a_{m_i}^i}$  the projection. Then there is a map

$$\Xi : H^s(Y_1) \otimes \cdots \otimes H^s(Y_n) \longrightarrow H^{sn} \left( Y, \bigcup_{i=1}^r p_{I_i}^{-1}(V_i) \right)$$

such that  $\Xi$  composed with the restriction to  $H^{sn}(Y)$  is the cup product. And  $\Xi$  is natural in the following sense: if  $Y' = Y'_1 \times \cdots \times Y'_n$ ,  $\{V'_i \subset Y'_{a_1^i} \times \cdots \times Y'_{a_{m_i}^i}\}_{i=1}^r$  satisfy the same condition as  $Y, \{Y_i\}, \{V_i\}$  above, with  $\Xi'$  the corresponding map, and there are continuous maps  $\{f_i : Y'_i \rightarrow Y_i\}_{i=1}^r$  such that  $(f_{a_1^i} \times \cdots \times f_{a_{m_i}^i})(V'_i) \subset V_i$  for all  $i$ , then  $\Xi' \circ (f_1^* \otimes \cdots \otimes f_n^*) = (f_1 \times \cdots \times f_n)^* \circ \Xi$ .

**Proof** First we define  $\Xi$ . Given  $\sigma_1 \in H^s(Y_1), \dots, \sigma_n \in H^s(Y_n)$ , take an open cover  $\mathcal{U} = \mathcal{U}_1 \times \cdots \times \mathcal{U}_n$  of  $Y$  satisfying (†) and such that for all  $i$ ,  $\sigma_i$  is represented by some skew-symmetric Čech cochain  $\alpha_i \in \check{C}_{\mathcal{U}_i}^s(Y_i)$  (see e.g., [3] for the equivalence between usual and skew-symmetric Čech cohomology; in particular, every (usual) Čech cohomology class has a skew-symmetric cochain representative). Denote by  $p_i : Y \rightarrow Y_i$  projection to the  $i$ -th factor. Define

$$(3) \quad \alpha := p_1^* \alpha_1 \cup \cdots \cup p_n^* \alpha_n \in \check{C}_{\mathcal{U}}^{sn}(Y).$$

We claim  $\alpha|_{\bigcup_{i=1}^r p_{I_i}^{-1}(V_i)} = 0$ . Suppose  $x \in p_{I_i}^{-1}(V_i)$  for some  $i$  and

$$x \in (U_1^0 \times \cdots \times U_n^0) \cap (U_1^1 \times \cdots \times U_n^1) \cap \cdots \cap (U_1^{sn} \times \cdots \times U_n^{sn}), \quad U_i^j \in \mathcal{U}_i.$$

Then

$$\begin{aligned} \alpha((U_1^0 \times \cdots \times U_n^0), (U_1^1 \times \cdots \times U_n^1), \dots, (U_1^{sn} \times \cdots \times U_n^{sn}))(x) \\ = \alpha_1(U_1^0, \dots, U_1^s)(p_1(x)) \cdots \alpha_n(U_n^{s(n-1)}, \dots, U_n^{sn})(p_n(x)) \end{aligned}$$

contains the factor

$$(4) \quad \alpha_{a_1^i}(U_{a_1^i}^{s(a_1^i-1)}, \dots, U_{a_1^i}^{sa_1^i})(p_{a_1^i}(x)) \cdots \alpha_{a_{m_i}^i}(U_{a_{m_i}^i}^{s(a_{m_i}^i-1)}, \dots, U_{a_{m_i}^i}^{sa_{m_i}^i})(p_{a_{m_i}^i}(x)).$$

Since  $\alpha_1, \dots, \alpha_n$  are skew-symmetric, for (4) to be nonzero,  $U_{a_j^{i_j-1}}^{s(a_j^{i_j-1})}, \dots, U_{a_j^{i_j}}^{s(a_j^{i_j})}$  must be pairwise different for all  $j = 1, \dots, m_i$ . But  $p_{I_i}(x) \in V_i$ . So by  $(\dagger)$ , (4)  $\neq 0$  is impossible. This proves  $\alpha|_{\bigcup_{i=1}^r p_{I_i}^{-1}(V_i)} = 0$ , so  $\alpha \in \check{C}_{\mathcal{U}}^{sn}(Y, \bigcup_{i=1}^r p_{I_i}^{-1}(V_i))$ . We define

$$\Xi(\sigma_1, \dots, \sigma_n) = \text{the direct image of } [\alpha] \text{ in } H^{sn}\left(Y, \bigcup_{i=1}^r p_{I_i}^{-1}(V_i)\right).$$

Then  $\Xi$  is clearly linear.

Next we show that  $[\alpha]$  above does not depend on the choices of  $\alpha_i$ . Suppose  $\alpha_i, \alpha'_i \in \check{C}_{\mathcal{U}_i}^s(Y_i)$  are both skew-symmetric cocycles,  $[\alpha_i] = [\alpha'_i] = \sigma_i$ . Then there exists a skew-symmetric  $\tilde{\alpha}_i \in \check{C}_{\mathcal{U}_i}^{s-1}(Y_i)$ ,  $d\tilde{\alpha}_i = \alpha_i - \alpha'_i$ . Define

$$\tilde{\alpha} := p_1^* \alpha_1 \cup \dots \cup p_{i-1}^* \alpha_{i-1} \cup p_i^* \tilde{\alpha}_i \cup p_{i+1}^* \alpha_{i+1} \cup \dots \cup p_n^* \alpha_n \in \check{C}_{\mathcal{U}}^{sn-1}(Y).$$

By the same argument as above,  $\tilde{\alpha}$  vanishes on  $\bigcup_{i=1}^r p_{I_i}^{-1}(V_i)$ , so  $\tilde{\alpha} \in \check{C}_{\mathcal{U}}^{sn-1}(Y, \bigcup_{i=1}^r p_{I_i}^{-1}(V_i))$ . Since all  $\alpha_j$  are cocycles,

$$d\tilde{\alpha} = (-1)^i (p_1^* \alpha_1 \cup \dots \cup d(p_i^* \tilde{\alpha}_i) \cup \dots \cup p_n^* \alpha_n) = (-1)^i (p_1^* \alpha_1 \cup \dots \cup p_i^* (\alpha_i - \alpha'_i) \cup \dots \cup p_n^* \alpha_n).$$

Therefore  $[\alpha]$  does not depend on the choice of  $\alpha_i$ .

We then show that  $\Xi(\sigma_1, \dots, \sigma_n)$  does not depend on the choices of  $\mathcal{U}_i$ . Suppose  $\mathcal{U}' = \mathcal{U}'_1 \times \dots \times \mathcal{U}'_n$  also satisfy  $(\dagger)$  and for all  $i$ ,  $\mathcal{U}'_i$  is a refinement of  $\mathcal{U}_i$ . Let  $\{\mu_i : \mathcal{U}'_i \rightarrow \mathcal{U}_i\}_{i=1}^n$  be some refinement maps. Denote  $\mu = (\mu_1, \dots, \mu_n) : \mathcal{U}' \rightarrow \mathcal{U}$ . Then for  $U_j^i \in \mathcal{U}'_j$ ,

$$\begin{aligned} \mu^* \alpha & \left( (U_1^0 \times \dots \times U_n^0), (U_1^1 \times \dots \times U_n^1), \dots, (U_1^{sn} \times \dots \times U_n^{sn}) \right) (x) \\ & = \alpha \left( (\mu_1(U_1^0) \times \dots \times \mu_n(U_n^0)), (\mu_1(U_1^1) \times \dots \times \mu_n(U_n^1)), \dots, (\mu_1(U_1^{sn}) \times \dots \times \mu_n(U_n^{sn})) \right) (x) \\ & = \alpha_1(\mu_1(U_1^0), \dots, \mu_1(U_1^{sn})) (p_1(x)) \cdots \alpha_n(\mu_n(U_n^{s(n-1)}), \dots, \mu_n(U_n^{sn})) (p_n(x)) \\ & = \mu_1^* \alpha_1(U_1^0, \dots, U_1^{sn}) (p_1(x)) \cdots \mu_n^* \alpha_n(U_n^{s(n-1)}, \dots, U_n^{sn}) (p_n(x)). \end{aligned}$$

This says  $\mu^* \alpha = p_1^*(\mu_1^* \alpha_1) \cup \dots \cup p_n^*(\mu_n^* \alpha_n) \in \check{C}_{\mathcal{U}' }^{sn}(Y, \bigcup_{i=1}^r p_{I_i}^{-1}(V_i))$ . So, if we define  $\Xi(\sigma_1, \dots, \sigma_n)$  using  $\mathcal{U}'$ , then it is the direct image of  $\mu^*[\alpha] = [\mu^* \alpha] \in H_{\mathcal{U}'}^{sn}(Y, \bigcup_{i=1}^r p_{I_i}^{-1}(V_i))$  which is the same as the direct image of  $[\alpha]$ . Now, if we assume  $\mathcal{U}'$  satisfies  $(\dagger)$  but not necessarily a refinement of  $\mathcal{U}$ , then by the assumption of the lemma we can find a common refinement of  $\mathcal{U}, \mathcal{U}'$  that also satisfies  $(\dagger)$ . This implies that  $\Xi(\sigma_1, \dots, \sigma_n)$  is independent of the choice of  $\mathcal{U}$ , and thus  $\Xi$  is well defined.

Naturality follows immediately from the definition of  $\Xi$ : let  $Y', \{Y'_i\}, \{f_i\}, \{V'_i\}$  be as in the lemma and  $\mathcal{U}, \alpha, \alpha_i$  as in the first paragraph of the proof; then  $\mathcal{U}' := f_1^{-1} \mathcal{U}_1 \times \dots \times f_n^{-1} \mathcal{U}_n$  satisfies  $(\dagger)$ . Take  $\alpha'_i = f_i^*(\alpha_i) \in \check{C}_{\mathcal{U}'_i}^s(Y')$ . Then  $\alpha'$  defined using (3) is the same as  $(f_1 \times \dots \times f_n)^*(\alpha)$  and the conclusion follows.  $\square$

**Lemma 4.7** Every open cover of  $(\bar{C}_2(\pi)/\sim_F)^{E(\Gamma)}$  has a refinement of the form

$$\mathcal{U}_{e_\Gamma} \times \dots \times \mathcal{U}_{e_{\Gamma(E(\Gamma))}}, \quad \text{where each } \mathcal{U}_{e_\Gamma} \text{ is an open cover of } \bar{C}_2(\pi)/\sim_F$$

such that

(\*) for all  $A$  of type 3,  $\sigma \in \tilde{S}_{E(\Gamma)}$ ,  $s'_e = d - 1$  or  $d - 2$  with at most one being  $d - 2$ ,

$$\bar{V}_A^\sigma \cap \prod_{e \in I_{\sigma,A}} (U_e^0 \cap \dots \cap U_e^{s'_e}) = \emptyset,$$

where  $U_e^0, \dots, U_e^{s'_e}$  are any pairwise distinct elements of  $\mathcal{U}_e$ , for every  $e \in I_{\sigma,A}$ .

**Proof** Since  $A$  is of type 3,  $|V(\Gamma_A)| \leq (2/3)|E(\Gamma_A)|$ . Since  $d \geq 3$ ,  $2d/3 \leq d - 1$ , and

$$\dim \bar{C}_{V(\Gamma_A)}^{\text{quo}}(\mathbb{R}^d) = d|V(\Gamma_A)| - d - 1 \leq (2d/3)|E(\Gamma_A)| - (d + 1) \leq \dim((S^{d-1})^{E(\Gamma_A)}) - 4.$$

So by Lemma 4.3  $\bar{V}_A^\sigma$  is the image of a smooth map  $f_{\sigma,A} : \bar{C}_{V(\Gamma_A)}^{\text{quo}}(\mathbb{R}^d) \rightarrow (S^{d-1})^{I_{\sigma,A}}$ , where the domain is of codimension at least 4 with respect to the target. (Indeed, codimension-2 would suffice for this lemma.)

Fix a metric  $D$  on  $\bar{C}_2(\pi)/\sim_F$ . Let  $K_0 \subset K_1 \subset \dots$  be a sequence of compact subsets of  $\bar{C}_2(\pi)/\sim_F$ , such that  $K_0$  contains a neighborhood of  $S_\pi^{d-1}$  and  $\bigcup_{i=1}^\infty K_i = \bar{C}_2(\pi)/\sim_F$ .

By Lebesgue's number lemma, it suffices to show that for any  $\epsilon_0, \epsilon_1, \dots > 0$ , there exists  $(\mathcal{U}_e)_{e \in E(\Gamma)}$  such that for all  $e$  and  $U \in \mathcal{U}_e$ , there exists  $i$  such that  $U \subset K_i$  and  $\text{diameter}_D(U) < \epsilon_i$ , satisfying (\*).

We first reduce the lemma to the following statement: for all  $\epsilon > 0$ , there exist  $(\mathcal{U}_e)_{e \in E(\Gamma)}$ , where each  $\mathcal{U}_e$  is an open covers of  $S_\pi^{d-1}$ , such that for all  $U \in \bigcup_e \mathcal{U}_e$ ,  $\text{diameter}_D(U) < \epsilon/4$ , satisfying (\*). Suppose this is true. Plug in  $\epsilon_0$  for  $\epsilon$ . We can enlarge each  $U \in \mathcal{U}_e$  a little bit to get an open subset  $l(U)$  of  $\bar{C}_2(\pi)/\sim_F$ , still contained in  $K_0$  and of diameter less than  $\epsilon_0$ . (For example, take  $l(U) = \{x \in C_2(\pi) \mid D(x, U) < \epsilon_0/2\} \cap \overset{\circ}{K}_0 \cup U$ .) Denote  $l(\mathcal{U}_e) = \{l(U)\}_{U \in \mathcal{U}_e}$ . We can cover

$$\bar{C}_2(\pi)/\sim_F = \bigcup_{U \in \mathcal{U}_e} l(U)$$

by locally finitely many open subsets of  $C_2(\pi)$ , such that each of these open subsets is contained in some  $K_i$  and is of diameter less than  $\epsilon_i$ ; call this collection of open sets  $\mathcal{U}'_e$ . Then for every  $e$ ,  $l(\mathcal{U}_e) \cup \mathcal{U}'_e$  is an open cover of  $\bar{C}_2(\pi)/\sim_F$ , in which each open set is contained in some  $K_i$  and of diameter less than  $\epsilon_i$ . Since for every  $A$ ,  $\bar{V}_A^\sigma$  is contained in  $(S_\pi^{d-1})^{I_{\sigma,A}}$  whereas elements in  $\mathcal{U}'_e$  do not intersect  $S_\pi^{d-1}$ , that (\*) is satisfied by  $(l(\mathcal{U}_e) \cup \mathcal{U}'_e)_{e \in E(\Gamma)}$  follows from that it is satisfied by  $(\mathcal{U}_e)_{e \in E(\Gamma)}$ .

We then reduce the statement at the beginning of last paragraph to the following: for any  $\epsilon > 0$  there exist triangulations  $(T_e)_{e \in E(\Gamma)}$  of  $S^{d-1}$ , compatible with the smooth structure on  $S^{d-1}$ , such that the diameter of each simplex is less than  $\epsilon/4$ , satisfying

(\*\*) for all  $\sigma, A$  of type 3, and all collections of simplices  $(S_e : \text{asimplexin } T_e)_{e \in I_{\sigma,A}}$  such that  $S_e$  is of dimension 0 or 1, at most one of dimension 1,  $\bar{V}_A^\sigma \cap \prod_{e \in I_{\sigma,A}} \bar{S}_e = \emptyset$ .

If this is true, we can take  $\mathcal{U}_e$  to be obtained from  $T_e$  by slightly enlarging each top-dimensional simplex  $S$  to an open neighborhood  $U(S)$  of its closure, so that:

- (1) (for a nontop dimensional simplex  $S$ , still denote by  $U(S) = \bigcap_{S'} U(S')$  where  $S'$  runs through the top-dimensional simplices that  $S$  is a face of)

$$\bar{V}_A^\sigma \cap \prod_{e \in I_{\sigma,A}} U(S_e) = \emptyset \quad \text{whenever} \quad \bar{V}_A^\sigma \cap \prod_{e \in I_{\sigma,A}} \bar{S}_e = \emptyset;$$

- (2) for any finite collection,  $S_1, \dots, S_k$ , of simplices in  $T_e$ ,

$$S_1 \cap \dots \cap S_k = \emptyset \implies U(S_1) \cap \dots \cap U(S_k) = \emptyset.$$

(These are clearly easy to satisfy. For (2), for every  $S$  in  $T_e$ , take  $U(S)$  contained in the union of the stars of the barycenters of its faces in the barycentric subdivision of  $T$ . For (1), take  $U(S)$  contained in the  $\epsilon'$ -neighborhood of  $S$ , where  $\epsilon'$  is the minimal distance between some  $\bar{V}_A^\sigma$  and the union of products of closed simplices  $\bar{V}_A^\sigma$  does not intersect.) We also require  $U(S)$  to be contained in the  $\epsilon/8$ -neighborhood of  $S$ . Then  $\{u_e\}$  satisfies (\*).

It remains to prove the statement at the beginning of the last paragraph. Take arbitrary triangulations  $(T_e^0)_{e \in E(\Gamma)}$  of  $S^{d-1}$  with diameter smaller than  $\epsilon/8$ , we perturb them simplex by simplex to satisfy (\*\*). The point is that every time we perturb a simplex away from  $\bar{V}_A^\sigma$ , we do it so slightly that no new unwanted intersection appears. The rest of the proof consists of technical details of this.

Below by “distance” we mean the restriction of  $D$  to  $S_\pi^{d-1}$  and its products. By a triangulation  $T$  of  $S^{d-1}$  we mean a homeomorphism  $T : |K_T| \rightarrow S^{d-1}$ , where  $K_T$  is a finite simplicial complex and  $|K_T|$  its realization. Given  $\mathcal{T} = (T_e)_{e \in E(\Gamma)}$  a tuple of triangulations of  $S^{d-1}$ , denote

$$\epsilon(\mathcal{T}) = \text{minimal distance between} \quad \prod_{e \in I_{\sigma,A}} \bar{S}_e \quad \text{and} \quad \bar{V}_A^\sigma,$$

where the minimum is taken over all  $\sigma, A, (\bar{S}_e)_{e \in I_{\sigma,A}}$  such that  $\bar{V}_A^\sigma \cap \prod_{e \in I_{\sigma,A}} \bar{S}_e = \emptyset$ . There are only finitely many of them. And  $\epsilon(\mathcal{T}) > 0$ . Denote by  $\mathcal{J}_1(\mathcal{T})$  (resp.  $\mathcal{J}_0(\mathcal{T})$ ) the set of tuples  $(\sigma, A, (x_e)_{e \in I_{\sigma,A}})$  such that  $x_e$  is the image of a 0- or 1-simplex of  $T_e$ , exactly one of them (resp. none of them) being a 1-simplex, and  $\bar{V}_A^\sigma \cap \prod_{e \in I_{\sigma,A}} x_e \neq \emptyset$ . Then  $\mathcal{J}_0(\mathcal{T})$  and  $\mathcal{J}_1(\mathcal{T})$  are finite.

Now take arbitrary triangulations  $\mathcal{T}^0 = (T_e^0)_{e \in E(\Gamma)}$  such that the diameter of any simplex is smaller than  $\epsilon/8$ . Denote by  $\epsilon'$  the smallest diameter of a simplex of  $\mathcal{T}^0$ . For the rest of the paragraph, for all  $\mathcal{T}$ , define  $\epsilon'(\mathcal{T}) = \min\{\epsilon'/4, \epsilon(\mathcal{T})\}$ . Take an element  $(\sigma, A, (x_e)_{e \in I_{\sigma,A}}) \in \mathcal{J}_0(\mathcal{T})$ . Since  $\bar{V}_A^\sigma$  cannot cover a neighborhood of  $\prod_{e \in I_{\sigma,A}} x_e$ , we can find  $\prod_{e \in I_{\sigma,A}} x'_e$  in the  $\epsilon'(\mathcal{T}^0)/2$ -neighborhood of it that does not meet  $\bar{V}_A^\sigma$ . For each  $e$  we can perturb the triangulation map  $T_e^0$  in a neighborhood of  $(T_e^0)^{-1}(x_e)$  to get  $T_e^1$ , in such a way that  $T_e^1((T_e^0)^{-1}(x_e)) = x'_e$ , and the distance between old and new images of any point is less than  $\epsilon'(\mathcal{T}^0)/|E(\Gamma)|$ . For  $e \notin I_{\sigma,A}$ , define  $T_e^1 = T_e^0$ . Define  $\mathcal{T}^1 = (T_e^1)_{e \in E(\Gamma)}$ . Then  $|\mathcal{J}_0(\mathcal{T}^1)| < |\mathcal{J}_0(\mathcal{T}^0)|$  and  $|\mathcal{J}_1(\mathcal{T}^1)| \leq |\mathcal{J}_1(\mathcal{T}^0)|$ . We do this one by one for all elements of  $\mathcal{J}_0(\mathcal{T}^0)$ , getting new triangulations  $\mathcal{T}^k = (T_e^k)_{e \in E(\Gamma)}$  by the end. Now take  $(\sigma, A, (x_e)_{e \in I_{\sigma,A}}) \in \mathcal{J}_1(\mathcal{T}^k)$  where  $e_* \in I_{\sigma,A}$  is such that  $x_{e_*}$  is the (image of a) 1-simplex. Let  $p : \bar{V}_A^\sigma \subset (S^{d-1})^{I_{\sigma,A}} \rightarrow (S^{d-1})^{I_{\sigma,A}-e_*}$  be the projection forgetting the  $e_*$ -factor. Recall  $f_{\sigma,A} : \bar{C}_{V(\Gamma_A)}^{\text{quo}}(\mathbb{R}^d) \rightarrow (S^{d-1})^{I_{\sigma,A}}$  defined at the beginning

of this proof. By possibly perturbing  $(x_e)_{e \in I_{\sigma,A}-e_*}$  in the same way as above, we can assume that  $(x_e)_{e \in I_{\sigma,A}-e_*}$  is a regular value of  $p \circ f_{\sigma,A}$ . So  $\bar{V}_A^\sigma \cap p^{-1}((x_e)_{e \in I_{\sigma,A}-e_*}) \subset S^{d-1}$  is the smooth image of a manifold of dimension less than  $d - 2$ . So we can perturb  $T_{e_*}^k$  in a neighborhood of  $(T_{e_*}^k)^{-1}(x_{e_*})$ , fixing some neighborhoods of the 0-simplices (since  $\mathcal{F}_0 = \emptyset$  now), to get a new triangulation  $T_{e_*}^{k+1}$ , so that  $T_{e_*}^{k+1}((T_{e_*}^k)^{-1}(x_{e_*}))$  does not intersect  $\bar{V}_A^\sigma \cap p^{-1}((x_e)_{e \in I_{\sigma,A}-e_*})$ , and the new and old image of any point has distance less than  $\epsilon'(\mathcal{T}^k)/|E(\Gamma)|$ . For  $e \in I_{\sigma,A}$  where  $x_e$  is not perturbed, or  $e \notin I_{\sigma,A}$ , define  $T_e^{k+1} = T_e^k$ . Define  $\mathcal{T}^{k+1} = (T_e^{k+1})_{e \in E(\Gamma)}$ . Then  $|\mathcal{F}_1(\mathcal{T}^{k+1})| < |\mathcal{F}_1(\mathcal{T}^k)|$ . Keep doing this one by one for all elements in  $\mathcal{F}_1(\mathcal{T}^k)$ . By the end, for some  $l$ , we obtain a tuple of triangulations  $\mathcal{T}^{k+l}$  satisfying (\*\*). □

### 4.2 Defining Kontsevich’s characteristic classes

To “push-forward”  $\Omega_\Gamma(\pi)$  to a cohomology class on the base  $B$ , the Leray–Serre spectral sequence is a convenient tool to formulate it. We follow [6] for the definition of Leray–Serre spectral sequence. First we make a general definition.

**4.2.1 Cohomology push-forward** Suppose  $B$  is a CW complex and  $X \xrightarrow{\pi} B$  a fiber bundle with fiber  $F$ . Denote by  $B_p$  the  $p$ -skeleton of  $B$  and  $X_p = \pi^{-1}(B_p)$ . Suppose there is  $k_0 > 0$  such that  $H^k(F) = 0$  for all  $k > k_0$ . Then, for any integers  $n$  and  $p < n - k_0$ , in the Leray–Serre spectral sequence for  $X_p \xrightarrow{\pi} B_p$ ,  $E_2^{a,b} = 0$  for all  $a + b = n$ , so  $H^n(X_p) = 0$ .

The Leray–Serre spectral sequence for  $X \xrightarrow{\pi} B$  tells us the following (cf. [6, Theorem 5.15]; note here we use the local coefficient version; cf. [16]): suppose  $n \geq k_0 \in \mathbb{Z}$ ; then

- $H^n(X)$  has a filtration by subgroups  $F_p^n = \ker(H^n(X) \rightarrow H^n(X_{p-1}))$  and  $E_\infty^{p,n-p} \approx F_p^n / F_{p+1}^n$ ;
- $E_2^{p,q} \approx H^p(B; H^q(F))$ , where the latter is understood as cohomology with local coefficients;
- $d_r : E_r^{p,q} \rightarrow E_r^{p+r,q+1-r}$ .

Suppose  $r \geq 2$ . Since  $H^k(F) = 0$  for all  $k > k_0$ ,

$$E_2^{n-k_0-r,k_0+r-1} \approx H^{n-k_0-r}(B, H^{k_0+r-1}(F)) = \{0\}.$$

Since  $E_r^{p,q}$  is obtained from  $E_2^{p,q}$  by taking subgroups and quotients,  $E_r^{n-k_0-r,k_0+r-1} = \{0\}$  for all  $r \geq 2$ . Therefore, all the  $d_r$ ’s mapping into  $E_r^{n-k_0,k_0}$  vanish and  $E_\infty^{n-k_0,k_0}$  is a subgroup of  $E_2^{n-k_0,k_0} \approx H^{n-k_0}(B; H^{k_0}(F))$ . Since  $H^n(X_{n-k_0-1}) = 0$ ,  $H^n(X) = F_{n-k_0}^n$ . This identifies a map

$$(5) \quad \pi_* : H^n(X) \longrightarrow H^{n-k_0}(B; H^{k_0}(F)).$$

**Definition 4.8** We call  $\pi_*$  the *cohomology push-forward* of the fiber bundle  $X \xrightarrow{\pi} B$ .

By the naturality of the Leray–Serre spectral sequence [6, pages 537–538],  $\pi_*$  does not depend on the choice of the CW structure on  $B$ , and is natural: suppose  $X' \xrightarrow{\pi'} B'$  is another fiber bundle with fiber  $F'$  such that  $H^k(F') = 0$  for all  $k > k_0$  and  $(\tilde{f} : X' \rightarrow X, f : B' \rightarrow B)$  is a bundle map (so

$H^{k_0}(F') \approx f^* H^{k_0}(F)$  as local systems over  $B'$ ); then  $f^* \circ \pi_* = \pi'_* \circ \tilde{f}^*$ . Using CW approximation,  $\pi_*$  can be generalized to the case where  $B$  is an arbitrary space.

The above procedure can be generalized to the relative version:  $X \xrightarrow{\pi} B$  has a subbundle  $Y \xrightarrow{\pi} B$  with fiber  $A \subset F$ . Replacing  $F$  with the pair  $(F, A)$  and  $X$  with  $(X, Y)$  everywhere, everything goes through without change.

**Remark 4.9** (This remark will be used later in Section 5.2.) An explicit description of  $\pi_*$  can be obtained by carefully unwinding the definition; we follow [6] for the construction of Leray–Serre spectral sequences and specifically, what we do below comes from the diagram on page 526 and  $\Phi$  in the proof of Theorem 5.3 in [6]. Denote  $p_0 = n - k_0$  for simplicity. Given an element  $\sigma \in H^n(X)$ , first restrict it to  $H^n(X_{p_0})$ ; the image will lie in the image of  $H^n(X_{p_0}, X_{p_0-1})$ . Take such a preimage, say  $\sigma'$ . For each  $p_0$ -cell  $e : (D^{p_0}, \partial D^{p_0}) \rightarrow (X_{p_0}, X_{p_0-1})$  ( $D^{p_0}$  being the standard  $p_0$ -dimensional ball),  $e^* \sigma' \in H^n(e^*(X), (e|_{\partial D^{p_0}})^*(X))$ . Since  $e^* \pi$  is trivializable, the Künneth formula gives

$$H^n(e^*(X), (e|_{\partial D^{p_0}})^*(X)) \approx H^{p_0}(D^{p_0}, \partial D^{p_0}) \otimes H^{k_0}(F_b),$$

where  $F_b = \pi^{-1}(e(b))$  is the fiber over an arbitrarily fixed point  $b \in \mathring{D}^{p_0}$ . Other fibers over  $D^{p_0}$  are identified with  $F_b$  via the trivialization. Notice that the Künneth isomorphism does not depend on the choice of the trivialization of  $e^* \pi$ : the Künneth map is determined by the projection map  $e^*(X) \rightarrow F_b$  (note that  $\partial D^{p_0}$  is not involved here); any two trivializations give homotopic projection maps since  $D^{p_0}$  is contractible. Let us denote by  $\sigma(e) \in H^{k_0}(F_b)$  the image of  $e^* \sigma'$  under the Künneth map, where  $H^{p_0}(D^{p_0}, \partial D^{p_0}) \approx R$  identified using the canonical orientation on  $D^{p_0}$ . Then  $\{e \rightarrow \sigma(e)\}_e$ , where  $e$  ranges over all  $p_0$ -cells of  $B$ , gives a cellular cochain on  $B$  with coefficients in the local system  $H^{k_0}(F)$ . It is a cocycle and represents a cohomology class in  $H^{p_0}(B; H^{k_0}(F))$ , which is  $\pi_*(\sigma)$ . The relative version is similar.

**4.2.2 Defining Kontsevich's characteristic classes** Applying the construction above to the fiber bundle  $(X_\Gamma(\pi), S(\pi)) \xrightarrow{\pi_X} B$  with  $k_0 = d|V(\Gamma)|$ , we get

$$(6) \quad \pi_{X*} : H^{|E(\Gamma)|(d-1)}(X_\Gamma(\pi), S(\pi)) \longrightarrow H^{|E(\Gamma)|(d-1)-d|V(\Gamma)|}(B; H^{d|V(\Gamma)|}(X_\Gamma, S)).$$

The map  $\rho : H^{d|V(\Gamma)|}(X_\Gamma, S) \rightarrow R$  in Definition 3.25 induces the corresponding map of local systems on  $B$ . So we get an induced map

$$(7) \quad H^*(B; H^{d|V(\Gamma)|}(X_\Gamma, S)) \longrightarrow H^*(B; R).$$

**Definition 4.10** Define  $K_{\Gamma, \pi, F} \in H^{|E(\Gamma)|(d-1)-d|V(\Gamma)|}(B; R)$ , to be the image of  $\Omega_\Gamma(\pi)$  under (6) and (7).

The corollary below is a direct consequence of Proposition 4.5 and the naturality of cohomology push-forward.

**Corollary 4.11** Under the assumptions of Theorem 1.2,  $K_{\Gamma, \pi', F'} = h_B^* K_{\Gamma, \pi'', F''}$ .

### 5 Equivalence with the original definition

In this section the coefficient ring  $R = \mathbb{R}$ . All open covers are assumed to be locally finite. The goal of this section is to prove the following

**Proposition 5.1** *Suppose  $(E \xrightarrow{\pi} B, F)$  is a framed smooth  $(M, \infty)$  bundle over a smooth manifold  $B$ . Then  $K_{\Gamma, \pi, F}$  defined above agrees, up to scaling by a constant depending only on  $\Gamma$ ,<sup>7</sup> with the usual definition of Kontsevich’s characteristic classes for  $(E \xrightarrow{\pi} B, F)$ ; see, for example, [20] for definition.*

#### 5.1 Čech to de Rham preliminary

First we state some general facts translating Čech to de Rham cohomology. Let  $Y$  be a smooth manifold. Denote by  $\mathcal{A}_Y^q$  the sheaf of differential  $q$ -form germs on  $Y$ ,  $\mathcal{Z}_Y^q$  the subsheaf of closed  $q$ -form germs and  $A^q(Y)$  the space of global  $q$ -forms on  $Y$ . Let  $\mathcal{U}$  be an open cover of  $Y$ . Let  $\underline{l} = \{l_U : Y \rightarrow \mathbb{R}^{\geq 0}\}_{U \in \mathcal{U}}$  be a partition of unity subordinate to  $\mathcal{U}$ . For any  $p, q \in \mathbb{Z}^{\geq 0}$  define

$$h_{p,q}^l : \check{C}_{\mathcal{U}}^p(Y; \mathcal{A}_Y^q) \longrightarrow \check{C}_{\mathcal{U}}^{p-1}(Y; \mathcal{A}_Y^{q+1}),$$

$$h_{p,q}^l(\sigma)(U_0, \dots, U_{p-1}) = (-1)^p \sum_{U \in \mathcal{U}} d(l_U \cdot \sigma(U, U_0, \dots, U_{p-1})).$$

By  $l_U \cdot \sigma(U, \dots, U_{p-1})$ , we mean a form on  $U_0 \cap \dots \cap U_{p-1}$  which is given by this formula on  $U \cap U_0 \cap \dots \cap U_{p-1}$  and 0 elsewhere; it is smooth since  $l_U$  vanishes in a neighborhood of  $\partial U$ . Clearly  $\text{image}(h_{p,q}^l) \subset \check{C}_{\mathcal{U}}^{p-1}(Y; \mathcal{Z}_Y^{q+1})$ . For each  $p$ , define

$$h^l : \check{C}_{\mathcal{U}}^p(Y; \mathbb{R}) \longrightarrow \check{C}_{\mathcal{U}}^0(Y; \mathcal{Z}_Y^p), \quad h^l(\sigma) = (-1)^p (h_{1,p-1}^l \circ h_{2,p-2}^l \circ \dots \circ h_{p,0}^l)(\sigma).$$

By [1, Proposition 9.8], if  $\sigma \in \check{C}_{\mathcal{U}}^p(Y; \mathbb{R})$  is a Čech cocycle, then  $h^l(\sigma)$  is a global closed form; if  $\mathcal{U}$  is such that any finite intersection of elements has trivial cohomology, then  $\sigma \rightarrow h^l(\sigma)$  induces the canonical isomorphism between  $\check{H}^p(Y; \mathbb{R})$  and  $H_{\text{deRham}}^p(Y)$ . If  $\Phi$  is a family of supports on  $Y$ , the arguments in [1, Section 8] still go through if all differential forms are assumed to have supports in  $\Phi$  (i.e.,  $\check{C}_{\mathcal{U}}^p$  is replaced with its subspace of  $\Phi$ -supported cochains  $\check{C}_{\mathcal{U}, \Phi}^p(Y; \mathcal{A}_Y^q)$ ); so  $h^l$  still induces the canonical isomorphism between  $\check{H}_{\Phi}^p(Y; \mathbb{R})$  and  $H_{\text{deRham}, \Phi}^p(Y)$ , where  $H_{\text{deRham}, \Phi}^p(Y)$  is the cohomology of the cochain complex of  $\Phi$ -supported differential forms on  $Y$ .

Suppose  $\mathcal{U}$  and  $\mathcal{U}'$  are two open covers of  $Y$ ,  $\mathcal{U}$  is finer than  $\mathcal{U}'$  and  $\mu : \mathcal{U} \rightarrow \mathcal{U}'$  is a refinement map. Let  $\underline{l} = \{l_U\}_{U \in \mathcal{U}}$  be a partition of unity subordinate to  $\mathcal{U}$ , then

$$\mu_* \underline{l} := \left\{ l_{U'} := \sum_{\substack{U \in \mathcal{U} \\ \mu(U) = U'}} l_U : Y \longrightarrow \mathbb{R} \right\}_{U' \in \mathcal{U}'}$$

<sup>7</sup>Again,  $\Gamma$  here is a linear combination of graphs. In the proof we only consider the case when  $\Gamma$  is a single graph but the argument generalizes verbatim.

is a partition of unity subordinate to  $\mathcal{U}'$ . It is easy to check that

$$(8) \quad h_{p,q}^{\mu * \underline{l}}(\mu^* \sigma) = \mu^*(h_{p,q}^{\underline{l}}(\sigma)) \quad \text{for all } p, q \text{ and } \sigma \in \check{C}_{\mathcal{U}'}^p(Y; \mathcal{A}_Y^q).$$

Define cup product

$$\cup : \check{C}_{\mathcal{U}}^{p_1}(Y; \mathcal{A}_Y^{q_1}) \otimes \check{C}_{\mathcal{U}}^{p_2}(Y; \mathcal{A}_Y^{q_2}) \longrightarrow \check{C}_{\mathcal{U}}^{p_1+p_2}(Y; \mathcal{A}_Y^{q_1+q_2}),$$

$$(\sigma_1 \cup \sigma_2)(U_0, \dots, U_{p_1+p_2}) = (-1)^{q_1 p_2} \sigma_1(U_0, \dots, U_{p_1}) \wedge \sigma_2(U_{p_1}, \dots, U_{p_1+p_2}),$$

where the two forms on the right-hand side are restricted to  $U_0 \cap \dots \cap U_{p_1+p_2}$ . For simplicity we omit the notation for restriction; same below. When restricted to  $\check{C}_{\mathcal{U}}^{p_1}(Y; \mathcal{E}_Y^0) \otimes \check{C}_{\mathcal{U}}^{p_2}(Y; \mathcal{E}_Y^0)$ , this is the usual cup product for Čech cochains.

**Lemma 5.2** *Let  $Y = Y_1 \times \dots \times Y_m$  be a product of smooth manifolds. Denote by  $\pi_i : Y \rightarrow Y_i$  the projection to the  $i$ -th factor. For every  $i = 1, \dots, m$ , let  $\mathcal{U}_i$  be an open cover of  $Y_i$  and  $\underline{l}_i = \{l_U\}_{U \in \mathcal{U}_i}$  be a partition of unity. Denote  $\mathcal{U} = \mathcal{U}_1 \times \dots \times \mathcal{U}_m$  the product open cover of  $Y$ . Then*

$$\underline{l} = \{l_{U_1 \times \dots \times U_m} := (l_{U_1} \circ \pi_1) \cdots (l_{U_m} \circ \pi_m) : Y \longrightarrow \mathbb{R}\}_{U_1 \in \mathcal{U}_1, \dots, U_m \in \mathcal{U}_m}$$

is a partition of unity of  $Y$  subordinate to  $\mathcal{U}$ . Let  $p = \sum_{i=1}^m p_i$  and  $q = \sum_{i=1}^m q_i$  be nonnegative integers. Let  $\sigma_i \in \check{C}_{\mathcal{U}_i}^{p_i}(Y_i; \mathcal{E}_{Y_i}^{q_i})$  be Čech cocycles. Define

$$\sigma = \pi_1^* \sigma_1 \cup \dots \cup \pi_m^* \sigma_m \in \check{C}_{\mathcal{U}}^p(Y; \mathcal{E}_Y^q).$$

Then,

(1) if  $m = 2$ ,

$$h_{p,q}^{\underline{l}}(\sigma) = \begin{cases} \pi_1^* h_{p_1,q_1}^{l_1}(\sigma_1) \cup \pi_2^*(\sigma_2) & \text{if } p_1 > 0, \\ \pi_1^* \sigma_1 \cup \pi_2^* h_{p_2,q_2}^{l_2}(\sigma_2) & \text{if } p_1 = 0, \end{cases}$$

(2) if  $q_i = 0$  for all  $i$ ,  $h^{\underline{l}}(\sigma) = \pi_1^* h^{l_1}(\sigma_1) \wedge \dots \wedge \pi_m^* h^{l_m}(\sigma_m)$ .

**Proof** This is direct computation. For (1), when  $p_1 > 0$ ,

$$\begin{aligned} & h_{p,q}^{\underline{l}}(\sigma)(U_1^0 \times U_2^0, \dots, U_1^{p-1} \times U_2^{p-1}) \\ &= (-1)^p \sum_{U_1 \in \mathcal{U}_1, U_2 \in \mathcal{U}_2} ((\pi_1^* dl_{U_1})(l_{U_2} \circ \pi_2) + (l_{U_1} \circ \pi_1)(\pi_2^* dl_{U_2})) \\ & \quad \wedge (-1)^{q_1 p_2} \pi_1^* \sigma_1(U_1, U_1^0, \dots, U_1^{p_1-1}) \wedge \pi_2^* \sigma_2(U_2^{p_1-1}, \dots, U_2^{p-1}) \\ &= (-1)^{p+q_1 p_2} \left( \left( \sum_{U_1} (\pi_1^* dl_{U_1}) \wedge \pi_1^* \sigma_1(U_1, \dots, U_1^{p_1-1}) \right) \wedge \left( \sum_{U_2} (l_{U_2} \circ \pi_2) \pi_2^* \sigma_2(U_2^{p_1-1}, \dots, U_2^{p-1}) \right) \right) \\ & \quad + \left( \sum_{U_1} (-1)^{q_1} (l_{U_1} \circ \pi_1) \pi_1^* \sigma_1(U_1, \dots, U_1^{p_1-1}) \right) \wedge \underbrace{\left( \sum_{U_2} (\pi_2^* dl_{U_2}) \wedge \pi_2^* \sigma_2(U_2^{p_1-1}, \dots, U_2^{p-1}) \right)}_{=0} \\ &= (-1)^{p+p_1+q_1 p_2} \pi_1^* h_{p_1,q_1}^{l_1}(\sigma_1)(U_1^0, \dots, U_1^{p_1-1}) \wedge \pi_2^* \sigma_2(U_2^{p_1-1}, \dots, U_2^{p-1}) \\ &= (\pi_1^* h_{p_1,q_1}^{l_1}(\sigma_1) \cup \pi_2^* \sigma_2)(U_1^0 \times U_2^0, \dots, U_1^{p-1} \times U_2^{p-1}). \end{aligned}$$



### 5.2 Proof of Proposition 5.1

We continue with the notation in Sections 3 and 4. The comparison between  $K_{\Gamma, \pi, F}$  and the standard definition is done by using the simplicial cohomology of  $B$ .

**5.2.1 Careful choices of open cover refinements in  $\bar{C}_2^{E(\Gamma)}(\pi)$**  Let  $\mathcal{U}'$  be an open cover of  $\bar{C}_2(\pi)/\sim_F$  such that there exists  $\check{\omega} \in \check{C}_{\mathcal{U}'}^{d-1}(\bar{C}_2(\pi)/\sim_F)$ , a cocycle representative of  $\Omega(\pi)$  ( $\Omega(\pi)$  is defined in Definition 4.2). Note the following commutative square, where  $\hat{\iota}$  and  $\iota$  are inclusion maps:

$$\begin{CD} \partial^v \bar{C}_2(\pi) @>\hat{\iota}>> \bar{C}_2(\pi) \\ @V F VV @VV q V \\ S^{d-1} @>\iota>> \bar{C}_2(\pi)/\sim_F \end{CD}$$

Since the Čech-to-de Rham procedure is done on a smooth manifold, we need to pull  $\check{\omega}$  back to  $\bar{C}_2(\pi)$  to produce a propagator. For this purpose, We need to make a careful choice of an open cover and a partition of unity on  $\bar{C}_2(\pi)$  in the next lemma.

**Lemma 5.3** *There exist an open cover  $\mathcal{U}$  of  $\bar{C}_2(\pi)$  such that all nonempty intersections of its elements are contractible, a refining map  $\mu : \mathcal{U} \rightarrow q^*\mathcal{U}'$ , and partitions of unity  $\underline{l}$  on  $\bar{C}_2(\pi)$  subordinate to  $\mathcal{U}$ ,  $\underline{l}_S$  on  $S^{d-1}$  subordinate to  $\iota^*\mathcal{U}'$ , such that  $\hat{\iota}^*(\mu_*\underline{l}) = F^*\underline{l}_S$  (they are both subordinate to  $\hat{\iota}^*q^*\mathcal{U}' = F^*\iota^*\mathcal{U}'$  on  $\partial^v \bar{C}_2(\pi)$ ).*

**Proof** By the way  $F$  is defined, it can be extended to a smooth map  $F : N_\partial \rightarrow S^{d-1}$ , where  $N_\partial \subset \bar{C}_2(\pi)$  is a neighborhood of  $\partial^v \bar{C}_2(\pi)$ . Still denote by  $\hat{\iota} : N_\partial \rightarrow \bar{C}_2(\pi)$  the inclusion. Let  $\underline{l}_S$  be a partition of unity on  $S^{d-1}$  subordinate to  $\iota^*\mathcal{U}'$ . Then  $F^*\underline{l}_S =: \{F^*\underline{l}_S(U)\}_{U \in \mathcal{U}'}$ , where  $F^*\underline{l}_S(U)$  is supported on  $q^{-1}(U) \cap N_\partial$ , is a partition of unity on  $N_\partial$  subordinate to  $\hat{\iota}^*q^*\mathcal{U}'$ . Our next goal is to find a partition of unity  $\underline{l}'$  on  $\bar{C}_2(\pi)$  subordinate to  $q^*\mathcal{U}'$ , such that  $\hat{\iota}^*\underline{l}' = F^*\underline{l}_S$ .

We can find a smooth function  $g : \bar{C}_2(\pi) \rightarrow \mathbb{R}^{\geq 0}$  supported in  $N_\partial$  such that  $g|_{\partial^v \bar{C}_2(\pi)} \equiv 1$ . Let  $K \subset \bar{C}_2(\pi)$  be a compact subset containing  $\bar{C}_2(\pi) - N_\partial$ , and let  $N'_\partial \subset \bar{C}_2(\pi) - K$  be a neighborhood of  $\partial^v \bar{C}_2(\pi)$ ; let  $\{V_i \subset \bar{C}_2(\pi) - N'_\partial\}_{i \in I}$  be an open cover of  $K$ ; let  $\{g_i\}_{i \in I} \sqcup \{g'\}$ , where  $g_i$  is supported in  $V_i$  and  $g'$  is supported in  $\bar{C}_2(\pi) - K$ , be a partition of unity on  $\bar{C}_2(\pi)$  subordinate to  $\{V_i\}_i \sqcup \{\bar{C}_2(\pi) - K\}$ . Then  $(\sum_i g_i)|_K \equiv 1$  and  $(\sum_i g_i)|_{N'_\partial \cap \bar{C}_2(\pi)} \equiv 0$ ; so  $g := 1 - \sum_i g_i$ , extended by 1 to  $\partial^v \bar{C}_2(\pi)$ , satisfies the requirement. For each  $U \in \mathcal{U}'$ , define  $h_U = g \cdot F^*\underline{l}_S(U) : N_\partial \rightarrow \mathbb{R}$ . Then it can be smoothly extended to the entire  $\bar{C}_2(\pi)$ , taking value 0 out of  $N_\partial$ ; so  $h_U|_{\partial^v \bar{C}_2(\pi)} = F^*\underline{l}_S(U)|_{\partial^v \bar{C}_2(\pi)}$ ,  $h_U$  is supported in  $q^{-1}(U)$  and  $(\sum_U h_U)|_{N'_\partial} \equiv 1$ .

Let  $K' \subset \bar{C}_2(\pi)$  be a compact subset containing  $\bar{C}_2(\pi) - N'_\partial$ . Let  $\{G_U \subset q^{-1}(U) \cap \bar{C}_2(\pi)\}_{U \in \mathcal{U}'}$  be compact subsets that still cover  $K'$ . For each  $U \in \mathcal{U}'$ , take  $\phi_U : \bar{C}_2(\pi) \rightarrow \mathbb{R}^{\geq 0}$  that is supported in  $q^{-1}(U) \cap \bar{C}_2(\pi)$  and  $\phi_U|_{G_U} \equiv 1$ . Then  $\sum_{U \in \mathcal{U}'} (\phi_U + h_U)$ , as a function on  $\bar{C}_2(\pi)$ , is positive everywhere and equals to 1 on  $\partial^v \bar{C}_2(\pi)$ . Define  $l'_U = (\phi_U + h_U) / (\sum_U (\phi_U + h_U))$ . Then  $l' := \{l'_U\}_{U \in \mathcal{U}'}$  is a partition of unity as required.

Next we define  $\mathcal{U}$ . Fix a Riemannian metric on  $\bar{C}_2(\pi)$  (which is a smooth manifold with boundary and corners). For every  $U \in \mathcal{U}'$ , let  $\{V_i^U \subset q^{-1}(U) \subset \bar{C}_2(\pi)\}_{i \in I_U}$  be a finite collection of geodesically convex open subsets, such that  $\text{supp}(l'_U) \subset \bigcup_{i \in I_U} V_i^U$ . Then  $\mathcal{U} := \{V_i^U\}_{U \in \mathcal{U}', i \in I_U}$  is an open cover of  $\bar{C}_2(\pi)$  whose intersections are all contractible. Define  $\mu : \mathcal{U} \rightarrow q^*\mathcal{U}'$ ,  $\mu(V_i^U) = q^{-1}(U)$ ; it is a refinement map.

Now, we construct a partition of unity  $\underline{l}$  on  $\bar{C}_2(\pi)$  subordinate to  $\mathcal{U}$  such that  $\mu_*\underline{l} = \underline{l}'$ . Using the same argument used to find the  $g_i$ s above, for every  $U \in \mathcal{U}'$  we can find smooth functions  $\psi_i^U : \bar{C}_2(\pi) \rightarrow \mathbb{R}^{\geq 0}$  supported in  $V_i^U$ , for every  $i \in I_U$ , such that  $(\sum_{i \in I_U} \psi_i^U)|_{\text{supp}(l'_U)} \equiv 1$ . Define  $l_i^U = \psi_i^U \cdot l'_U$ . Then  $\sum_{i \in I_U} l_i^U = l'_U$ . So  $\underline{l} := \{l_i^U\}_{U \in \mathcal{U}', i \in I_U}$  is a partition of unity as required.  $\square$

**Corollary 5.4**  $h^L(\mu^*q^*\check{\omega})$  is a closed  $(d-1)$ -form on  $\bar{C}_2(\pi)$  such that  $\hat{t}^*h^L(\mu^*q^*\check{\omega}) = F^*\alpha$  for some closed form  $\alpha \in A^{d-1}(S^{d-1})$  with  $\int_{S^{d-1}} \alpha = 1$ . In other words,  $h^L(\mu^*q^*\check{\omega})$  is a propagator.

**Proof** That it is a closed  $(d-1)$ -form is clear. And

$$\hat{t}^*h^L(\mu^*q^*\check{\omega}) \stackrel{(8)}{=} \hat{t}^*h^{\mu_*L}(q^*\check{\omega}) \stackrel{(9)}{=} h^{\hat{t}^*\mu_*L}(\hat{t}^*q^*\check{\omega}) \stackrel{\text{Lemma 5.3}}{=} h^{F^*L_S}(F^*l^*\check{\omega}) \stackrel{(9)}{=} F^*h^{L_S}(l^*\check{\omega}).$$

Since  $[\check{\omega}] = \Omega(\pi) \in H^{d-1}(\bar{C}_2(\pi)/\sim_F)$  and  $h^{L_S}$  induces the canonical isomorphism between Čech and de Rham cohomology,  $[h^{L_S}(l^*\check{\omega})] \in H^{d-1}(S^{d-1}; \mathbb{R})$  is the Poincaré dual of the point class by the definition of  $\Omega(\pi)$ . So  $\int_{S^{d-1}} h^{L_S}(l^*\check{\omega}) = 1$ .  $\square$

Now, let  $\{\mathcal{U}'_e\}_{e \in E(\Gamma)}$  be a collection of open covers of  $\bar{C}_2(\pi)/\sim_F$  given by Lemma 4.7. For every  $e$ , applying the above argument with  $\mathcal{U}'$  replaced by  $\mathcal{U}'_e$ , we get an open cover  $\mathcal{U}_e$  with refinement map  $\mu_e : \mathcal{U}_e \rightarrow q^*\mathcal{U}'_e$  and partition of unity  $\underline{l}_e$  on  $\bar{C}_2(\pi)$  subordinate to  $\mathcal{U}_e$ , such that  $h^{L_e}(\mu_e^*q^*\check{\omega})$  is a propagator. Denote  $\omega_e = h^{L_e}(\mu_e^*q^*\check{\omega})$ . Denote by  $\text{pr}_e : \bar{C}_2(\pi)^{E(\Gamma)} \rightarrow \bar{C}_2(\pi)$  the projection to the  $e$ -th factor. (Recall  $\bar{C}_2^{E(\Gamma)}(\pi)$  is the fiber product and  $\bar{C}_2(\pi)^{E(\Gamma)}$  is the direct product of the total space.) Denote

$$\tilde{\mathcal{U}} := \text{pr}_{e_1}^*\mathcal{U}_{e_1} \times \cdots \times \text{pr}_{e_{|E(\Gamma)|}}^*\mathcal{U}_{e_{|E(\Gamma)|}}$$

the product open cover on  $\bar{C}_2(\pi)^{E(\Gamma)}$ . Define

$$\check{\omega}'_\Gamma := \text{pr}_{e_1}^*\mu_{e_1}^*q^*\check{\omega} \cup \text{pr}_{e_2}^*\mu_{e_2}^*q^*\check{\omega} \cup \cdots \cup \text{pr}_{e_{|E(\Gamma)|}}^*\mu_{e_{|E(\Gamma)|}}^*q^*\check{\omega} \in \check{C}_{\tilde{\mathcal{U}}}^{|E(\Gamma)|(d-1)}(\bar{C}_2(\pi)^{E(\Gamma)}, S(\pi))$$

(that it is supported away from  $S(\pi)$  follows from the choice of  $\{\mathcal{U}'_e\}$  given by Lemma 4.7); its restriction to  $X_\Gamma$  represents the class  $\Omega_\Gamma(\pi) \in H^{|E(\Gamma)|(d-1)}(X_\Gamma(\pi), S(\pi))$ . Define

$$\omega_\Gamma = \text{pr}_{e_1}^*\omega_{e_1} \wedge \cdots \wedge \text{pr}_{e_{|E(\Gamma)|}}^*\omega_{e_{|E(\Gamma)|}} \in A^{|E(\Gamma)|(d-1)}(\bar{C}_2(\pi)^{E(\Gamma)});$$

then the push-forward of  $\omega_\Gamma|_{C_\Gamma(\pi)}$  to  $B$  represents Kontsevich’s class in the usual definition. By Lemma 5.2,  $\omega_\Gamma = h^{\tilde{l}}\check{\omega}'_\Gamma$ , where  $\tilde{l}$  is the partition of unity on  $\bar{C}_2(\pi)^{E(\Gamma)}$  subordinate to  $\tilde{\mathcal{U}}$  given by taking the “product” of the  $\underline{l}_e$ ’s as in Lemma 5.2. Below we denote the restriction of  $\omega_\Gamma$  to  $\bar{C}_2^{E(\Gamma)}(\pi)$  still by  $\omega_\Gamma$ .

Fix a triangulation on  $B$  and denote by  $B_p$  the  $p$ -skeleton of  $B$  with respect to this triangulation. Denote

$$p_0 = |E(\Gamma)|(d - 1) - d|V(\Gamma)| = \deg \Omega_\Gamma(\pi) - \dim X_\Gamma.$$

Recall that  $f = (f_e)_{e \in E(\Gamma)} : \bar{C}_{V(\Gamma)}(\pi) \rightarrow \bar{C}_2^{E(\Gamma)}(\pi)$  is the forgetful map, and  $\phi$  is the factor-permuting action of  $\tilde{S}_{E(\Gamma)}$  on  $\bar{C}_2^{E(\Gamma)}(\pi)$ . Denote by

$$\pi_V : \bar{C}_{V(\Gamma)}(\pi) \longrightarrow B, \quad \hat{\pi} : \bar{C}_2^{E(\Gamma)}(\pi) \longrightarrow B, \quad \pi_X : X_\Gamma(\pi) \longrightarrow B$$

the bundle projection maps (the first two were both denoted by  $\pi$ ; here we want to distinguish them to avoid confusion). Define

$$\mathring{X}_\Gamma = \bigcup_{\sigma \in \tilde{S}_{E(\Gamma)}} \phi(\sigma)(C_\Gamma(M, \infty)), \quad T = X_\Gamma - \mathring{X}_\Gamma \subset \bar{C}_2(M, \infty)^{E(\Gamma)},$$

and denote  $\mathring{X}_\Gamma(\pi), T(\pi) \subset \bar{C}_2^{E(\Gamma)}(\pi)$  the bundle version of them. For all  $p$ , set  $X_p = \hat{\pi}^{-1}(B_p) \cap X_\Gamma(\pi)$ .<sup>8</sup>

By Lemma 3.18, we can find an open cover  $\hat{\mathcal{U}}$  of  $\bar{C}_2(\pi)^{E(\Gamma)}$  refining  $\tilde{\mathcal{U}}$ , such that there exists a neighborhood  $N$  of  $(T(\pi) \cap X_{p_0}) \cup X_{p_0-1}$  in  $\bar{C}_2(\pi)^{E(\Gamma)}$ , satisfying

$$U_0 \cap \cdots \cap U_{|E(\Gamma)|(d-1)} \cap N = \emptyset \quad \text{for all } U_0 \neq \cdots \neq U_{|E(\Gamma)|(d-1)} \in \hat{\mathcal{U}}.$$

Let  $\hat{\mu} : \hat{\mathcal{U}} \rightarrow \tilde{\mathcal{U}}$  be a refinement map. Then  $\hat{\mu}^* \check{\omega}'_\Gamma$  is supported away from  $N$ . Define  $\check{\omega}_\Gamma = (\hat{\mu}^* \check{\omega}'_\Gamma)|_{X_\Gamma(\pi)}$ .

Let  $\hat{l}$  be a partition of unity on  $\bar{C}_2(\pi)^{E(\Gamma)}$  subordinate to  $\hat{\mathcal{U}}$ ; then  $\hat{\mu}_* \hat{l}$  is a partition of unity subordinate to  $\tilde{\mathcal{U}}$ . Define  $\bar{\omega}_\Gamma := h^{\hat{\mu}_* \hat{l}}(\check{\omega}'_\Gamma) = h^{\hat{l}}(\hat{\mu}^* \check{\omega}'_\Gamma)$ . Then  $\bar{\omega}_\Gamma|_{X_\Gamma(\pi)} = h^{\hat{l}|_{X_\Gamma(\pi)}}(\check{\omega}_\Gamma)$ . Since all the intersections of elements of  $\tilde{\mathcal{U}}$  are contractible (since the same is true for each  $\mathcal{U}_e$  and  $\tilde{\mathcal{U}}$  is their product), both  $h^{\hat{\mu}_* \hat{l}}$  and  $h^{\hat{l}}$  induce the isomorphism between Čech and de Rham cohomology (here we let the family of supports be the collection of compact subsets in  $\bar{C}_2(\pi)^{E(\pi)}$  that do not intersect  $S(\pi)$ ), so

$$(10) \quad [\bar{\omega}_\Gamma] = [\check{\omega}'_\Gamma] = [\omega_\Gamma] \in H^*(\bar{C}_2(\pi)^{E(\Gamma)}, S(\pi)).$$

Denote the restriction of  $\bar{\omega}_\Gamma$  to  $\bar{C}_2^{E(\Gamma)}(\pi)$  still by  $\bar{\omega}_\Gamma$ ; then, pulling back the above equation by restriction,

$$(11) \quad [\bar{\omega}_\Gamma] = [\omega_\Gamma] \in H^*(\bar{C}_2^{E(\Gamma)}(\pi), S(\pi)).$$

**5.2.2 Passing to the simplicial cohomology on  $B$**  Define  $s(\sigma) = (d - 1) \operatorname{sgn}(\sigma) + d \operatorname{sgn}'(\sigma)$ . For a differential form  $\alpha \in A^m(\bar{C}_2^{E(\Gamma)}(\pi))$ , define  $\pi_*^s(\alpha)$  to be the degree  $(m - d|V(\Gamma)|)$  simplicial cochain on  $B$  that sends a dimension  $m - d|V(\Gamma)|$  simplex  $\Delta$  to

$$\sum_{\sigma \in \tilde{S}_{E(\Gamma)}} (-1)^{s(\sigma)} \int_{\pi_V^{-1}(\Delta)} (\phi(\sigma) \circ f)^* \alpha = \sum_{\sigma \in \tilde{S}_{E(\Gamma)}} (-1)^{s(\sigma)} \int_{\pi_V^{-1}(\Delta)} f^* \phi(\sigma)^* \alpha.$$

<sup>8</sup>The reason for defining  $T$  instead of using the previously defined  $T_1$  is the following: we did not establish anything regarding smoothness in Section 3.3 when gluing the various copies of  $\bar{C}_\Gamma(M, \infty)$  together to form  $\tilde{X}_\Gamma$ ; now we are working with differential forms, we want our Čech cochains to be supported away from this “joint” part, as shown in the next paragraph.

Then, since  $\phi(\sigma)^*\omega_\Gamma = (-1)^s\omega_\Gamma$ ,  $[\pi_*^s\omega_\Gamma] \in H^{p_0}(B)$  is  $2^{|E(\Gamma)|}|E(\Gamma)|!$  times Kontsevich's class in the usual definition.

Recall that  $\pi_{\tilde{X}} : \tilde{X}_\Gamma(\pi) \rightarrow B$  and  $\tilde{f} : \tilde{X}_\Gamma(\pi) \rightarrow \bar{C}_2^{E(\Gamma)}(\pi)$  are the bundle versions of  $\tilde{X}_\Gamma$  and  $\tilde{f}$  in Definition 3.22. In Definition 3.22, by choosing a collar neighborhood when gluing the copies of  $C'_\Gamma(M, \infty)$ ,  $\tilde{X}_\Gamma$  can be given a smooth structure so that  $\tilde{f} : \tilde{X}_\Gamma(\pi) \rightarrow \bar{C}_2^{E(\Gamma)}(\pi)$  is piecewise smooth (smooth away from  $\tilde{f}^{-1}(T(\pi))$ ); so pulling back a differential form by  $\tilde{f}$  is still well defined (the result would be a piecewise-smooth form), and the usual differential form push-forward  $(\pi_{\tilde{X}})_*$  is also well defined for these forms (by integrating along each piece of the fiber and summing up). It follows from the definition that  $\pi_*^s(\alpha) = (\pi_{\tilde{X}})_*\tilde{f}^*\alpha$ .

**Lemma 5.5** *Suppose  $\alpha_1, \alpha_2 \in A^*(\bar{C}_2^{E(\Gamma)}(\pi))$  are such that there exists  $\tilde{\alpha} \in A^{*-1}(\bar{C}_2^{E(\Gamma)}(\pi))$  supported away from  $S(\pi)$  and  $d\tilde{\alpha} = \alpha_1 - \alpha_2$ . Then  $\pi_*^s\alpha_1 - \pi_*^s\alpha_2$  is a coboundary.*

**Proof** For a simplex  $\Delta$  in  $B$ ,

$$\begin{aligned} & \sum_{\sigma \in \tilde{S}_{E(\Gamma)}} (-1)^{s(\sigma)} \int_{\pi_V^{-1}(\Delta)} (\phi(\sigma) \circ f)^*(\alpha_1 - \alpha_2) \\ &= \sum_{\sigma \in \tilde{S}_{E(\Gamma)}} (-1)^{s(\sigma)} \int_{\partial\pi_V^{-1}(\Delta)} (\phi(\sigma) \circ f)^*\tilde{\alpha} \\ &= \sum_{\sigma \in \tilde{S}_{E(\Gamma)}} (-1)^{s(\sigma)} \int_{\pi_V^{-1}(\partial\Delta)} (\phi(\sigma) \circ f)^*\tilde{\alpha} + \sum_{\sigma \in \tilde{S}_{E(\Gamma)}} (-1)^{s(\sigma)} \sum_A \int_{\pi_V^{-1}(\Delta) \cap \mathcal{G}_A} (\phi(\sigma) \circ f)^*\tilde{\alpha}, \end{aligned}$$

where  $A \subset V(\Gamma) \cup \{\infty\}$ ,  $|A| \geq 2$ . The second term vanishes: for  $A$  of type 1,  $f(\mathcal{G}_A)$  has positive codimension; for  $A$  of type 2 or 4, contributions to this term cancel with each other when summed over  $\sigma$ ; for  $A$  of type 3,  $\tilde{\alpha}$  vanishes on  $S(\pi)$  by assumption. So,

$$(\pi_*^s\alpha_1 - \pi_*^s\alpha_2)(\Delta) = (\pi_*^s\tilde{\alpha})(\partial\Delta) = (\delta\pi_*^s\tilde{\alpha})(\Delta). \quad \square$$

Since  $[\bar{\omega}_\Gamma] = [\omega_\Gamma] \in H^*(\bar{C}_2^{E(\Gamma)}(\pi), S(\pi))$  by (11) and  $[\pi_*^s\omega_\Gamma] \in H^{p_0}(B)$  is  $2^{|E(\Gamma)|}|E(\Gamma)|!$  times Kontsevich's class in the usual definition (last sentence in the first paragraph of Section 5.2.2), by Lemma 5.5,  $\pi_*^s(\bar{\omega}_\Gamma)$  also represents  $2^{|E(\Gamma)|}|E(\Gamma)|!$  times Kontsevich's class in the usual definition. We are now ready to finish the proof of Proposition 5.1.

We apply Remark 4.9 to the situation here. For each  $p_0$ -simplex  $\Delta^{p_0}$  in  $B$ ,  $\check{\omega}_\Gamma|_{\pi_X^{-1}(\Delta^{p_0})}$  is supported away from  $(T(\pi) \cap \pi_X^{-1}(\Delta^{p_0})) \cup \pi_X^{-1}(\partial\Delta^{p_0})$ . So  $\check{\omega}_\Gamma|_{\pi_X^{-1}(\Delta^{p_0})}$  is a cocycle in

$$\check{C}_{\check{q}|_{\pi_X^{-1}(\Delta^{p_0})}}^{|E(\Gamma)|(d-1)}(X_\Gamma(\pi)|_{\Delta^{p_0}}, X_\Gamma(\pi)|_{\partial\Delta^{p_0}} \cup T(\pi)|_{\Delta^{p_0}}).$$

By the Künneth formula,

$$H^{|E(\Gamma)|(d-1)}(X_\Gamma(\pi)|_{\Delta^{p_0}}, X_\Gamma(\pi)|_{\partial\Delta^{p_0}} \cup T(\pi)|_{\Delta^{p_0}}) \approx H^{p_0}(\Delta^{p_0}, \partial\Delta^{p_0}) \otimes H^{d|V(\Gamma)}(X_{\Gamma,b}, T_b),$$

where  $b \in \overset{\circ}{\Delta}^{p_0}$  is an arbitrary point and  $(X_{\Gamma,b}, T_b)$  is the fiber of  $(X_{\Gamma}(\pi), T(\pi))$  over  $b$ . Let  $\chi'(\Delta^{p_0}) \in H^{d|V(\Gamma)}(X_{\Gamma,b}, T_b)$  be such that  $[\check{\omega}_{\Gamma}|_{\pi_{\bar{X}}^{-1}(\Delta^{p_0})}] \approx 1 \otimes \chi'(\Delta^{p_0})$  under the Künneth isomorphism, where  $1 \in \mathbb{R} \approx H^{p_0}(\Delta^{p_0}, \partial\Delta^{p_0})$  (notice the identification uses the orientation of  $B$ ). Then  $\{\Delta^{p_0} \rightarrow \chi'(\Delta^{p_0})\}_{\Delta^{p_0}}$  is a simplicial cocycle on  $B$  with coefficients in the local system  $H^{d|V(\Gamma)}(X_{\Gamma}, T)$ . Its restriction to  $H^{d|V(\Gamma)}(X_{\Gamma}, T_1)$  (note  $T_1 \subset T$ ) represents  $(\pi_X)_*\Omega_{\Gamma}(\pi)$ . Let

$$\chi(\Delta^{p_0}) = \tilde{f}^* \chi'(\Delta^{p_0}) \in H_c^{d|V(\Gamma)}(\tilde{X}_{\Gamma,b}) \approx \mathbb{R}$$

(the last identification uses the orientation on  $\tilde{X}_{\Gamma}$ ); then  $\{\Delta^{p_0} \rightarrow \chi(\Delta^{p_0})\}_{\Delta^{p_0}}$  is a representative of  $K_{\Gamma,\pi,F}$  by the definition of  $K_{\Gamma,\pi,F}$  in Section 4.2.2.

Note the bundle map

$$\begin{array}{ccccc} \tilde{X}_{\Gamma}(\pi) & \xrightarrow{\tilde{f}} & X_{\Gamma}(\pi) - T_1(\pi) & \hookrightarrow & X_{\Gamma}(\pi) \\ & \searrow \pi_{\tilde{X}} & \downarrow \pi_X & \swarrow \pi_X & \\ & & B & & \end{array}$$

Since  $\check{\omega}_{\Gamma} = 0$  in a neighborhood of  $T(\pi) \cap X_{p_0}$ ,  $\tilde{f}^* \check{\omega}_{\Gamma}|_{\pi_{\bar{X}}^{-1}(B_{p_0})}$  is compactly supported. For each  $p_0$ -simplex  $\Delta^{p_0}$  of  $B$ , by the naturality of the Künneth formula,

$$[\tilde{f}^* \check{\omega}_{\Gamma}|_{\pi_{\bar{X}}^{-1}(\Delta^{p_0})}] \approx 1 \otimes \tilde{f}^* \chi'(\Delta^{p_0}) = 1 \otimes \chi(\Delta^{p_0})$$

under the Künneth isomorphism

$$H_c^{E(\Gamma)|(d-1)}(\tilde{X}_{\Gamma}(\pi)|_{\Delta^{p_0}}, \tilde{X}_{\Gamma}(\pi)|_{\partial\Delta^{p_0}}) \approx H^{p_0}(\Delta^{p_0}, \partial\Delta^{p_0}) \otimes H_c^{d|V(\Gamma)}(\tilde{X}_{\Gamma,b}).$$

Since  $\bar{\omega}_{\Gamma} = h^{\hat{l}}(\hat{\mu}^* \check{\omega}'_{\Gamma})$  and  $\hat{\mu}^* \check{\omega}'_{\Gamma}$  is supported away from  $N$ ,  $\bar{\omega}_{\Gamma}$  is also supported away from  $N$ . So  $[\bar{\omega}_{\Gamma}] = [\check{\omega}'_{\Gamma}] \in H^*(\bar{C}_2(\pi)^{E(\Gamma)}, N)$ , and  $\tilde{f}^* \bar{\omega}_{\Gamma}$  is a smooth form. For every simplex  $\Delta^{p_0}$  in  $B$ , pulling back the cohomology equality to  $\tilde{X}_{\Gamma}(\pi)$  and restricting to  $\Delta^{p_0}$  in  $B$ ,

$$\begin{aligned} [\tilde{f}^* \bar{\omega}_{\Gamma}|_{\pi_{\bar{X}}^{-1}(\Delta^{p_0})}] &= [\tilde{f}^* \check{\omega}_{\Gamma}|_{\pi_{\bar{X}}^{-1}(\Delta^{p_0})}] \\ &= 1 \otimes \chi(\Delta^{p_0}) \in H_c^{E(\Gamma)|(d-1)}(\tilde{X}_{\Gamma}(\pi)|_{\Delta^{p_0}}, \tilde{X}_{\Gamma}(\pi)|_{\partial\Delta^{p_0}}) \\ &\approx H^{p_0}(\Delta^{p_0}, \partial\Delta^{p_0}) \otimes H_c^{d|V(\Gamma)}(\tilde{X}_{\Gamma,b}) \\ &\approx \mathbb{R} \otimes \mathbb{R}. \end{aligned}$$

Therefore,  $\int_{\pi_{\bar{X}}^{-1}(\Delta^{p_0})} \tilde{f}^* \bar{\omega}_{\Gamma} = \chi(\Delta^{p_0})$ . This completes the proof of Proposition 5.1.

## 6 Some remarks about the condition in Theorem 1.2

Theorem 1.2 can be formulated in a different way.

For open subsets  $U, V \subset \mathbb{R}^d$  and a continuous map  $f : U \rightarrow V$ , say  $f$  is *almost differentiable* if the map  $(f, f) : U \times U \rightarrow V \times V$  lifts to a continuous map<sup>9</sup>  $\tilde{f} : \text{Bl}_{\Delta}(U \times U) \rightarrow \text{Bl}_{\Delta}(V \times V)$ , where  $\Delta$

denotes the diagonal in  $U \times U$  and  $V \times V$ , respectively, and  $\text{Bl}_\Delta$  denotes real oriented blow-up along  $\Delta$ . We can define an *almost differentiable manifold* to be a topological manifold together with a maximal collection of charts where the transition maps are almost differentiable homeomorphisms whose inverses are also almost differentiable. So, if  $M$  is an almost differentiable manifold, then  $\text{Bl}_\Delta(M \times M)$  is well defined. The corresponding automorphism group  $\text{Aut}^{ad}(M)$  in this category consists of homeomorphisms  $f : M \rightarrow M$  such that  $(f, f) : M \times M \rightarrow M \times M$  lifts to a homeomorphism  $\text{Bl}_\Delta(M \times M) \rightarrow \text{Bl}_\Delta(M \times M)$ . Denote by  $\pi : \text{Bl}_\Delta(M \times M) \rightarrow M \times M$  the blow-down map.

**Remark 6.1** (This was pointed out by an anonymous referee.) The notion of almost differentiability can be rephrased as follows: for open subsets  $U, V \subset \mathbb{R}^d$ , a continuous map  $f : U \rightarrow V$  is *almost differentiable* if

- for each  $a \in U$  and  $v \in S^{d-1}$ , the limit

$$SDf_a(v) = \lim_{t \rightarrow 0, t > 0} \frac{f(a + tv) - f(a)}{|f(a + tv) - f(a)|} \in ST_{f(a)}V$$

exists, and

- $SDf : STU \rightarrow STV$ ,  $SDf(a, v) := SDf_a(v)$  is continuous.

Given two real vector spaces  $T_1$  and  $T_2$ , and a linear isomorphism  $f : T_1 \rightarrow T_2$ , since  $f(\lambda v) = \lambda f(v)$  for all  $v \in T_1 - 0$  and  $\lambda \in \mathbb{R} - 0$ ,  $f$  induces a homeomorphism  $ST_1 \rightarrow ST_2$ , where  $ST_i = (T_i - 0)/\text{scaling}$  denotes the unit sphere in  $T_i$ .

Suppose  $M$  is a  $d$ -dimensional almost differentiable manifold. Define a *framing*  $F$  on  $M$  to be a continuous map  $F : \partial\text{Bl}_\Delta(M \times M) \rightarrow S^{d-1}$  such that for every  $x \in M$ ,  $F|_{\pi^{-1}(x,x)}$  satisfies that if  $\phi : \mathbb{R}^d \supset U \xrightarrow{\sim} N \subset M$ ,  $\phi(0) = x$  is a chart of  $M$  near  $x$ , then  $F|_{\pi^{-1}(x,x)} := ST_x U \rightarrow S^{d-1}$  is a homeomorphism induced from a linear map  $T_x U \rightarrow \mathbb{R}^d$ . By Proposition 6.5 below, this condition doesn't depend on the choice of such a chart  $\phi$ .

Suppose  $M$  is an almost differentiable manifold and  $\infty \in M$  a fixed point. Then the group  $\mathcal{G}$  defined in Section 1.1 can be similarly defined here:

$$\mathcal{G} := \text{Aut}_\infty^{ad}(M) := \{g \in \text{Aut}^{ad}(M) \mid \exists \text{ neighborhood } N \ni \infty \text{ such that } g|_N = \text{id}\}.$$

Define an *almost differentiable*  $(M, \infty)$ -*bundle* to be a fiber bundle with fiber  $M$  and structure group  $\text{Aut}_\infty^{ad}(M)$ . Such a bundle  $\pi : E \rightarrow B$  has an associated vertical sphere tangent bundle  $\pi' : ST^v E \rightarrow E$ . Define a *framing* on  $\pi : E \rightarrow B$  to be a topological trivialization

$$\begin{array}{ccc} F : ST^v E|_{E-s_\infty} & \xrightarrow{\text{homeomorphism}} & S^{d-1} \times (E - s_\infty) \\ & \searrow \pi' & \swarrow \text{projection} \\ & & E - s_\infty \end{array}$$

<sup>9</sup>Although the notation  $\tilde{f}$  has a completely different meaning in Sections 3–5, Section 6 is relatively independent from the others, so this abuse of notation should not cause confusion.

of  $\pi'|_{E-s_\infty}$ , standard near  $s_\infty$ , and such that for each point  $p \in E - s_\infty$ ,  $F|_p : ST_p^v E \rightarrow S^{d-1}$  is induced from a linear map (with respect to any chart of  $E|_p$  near  $p$ ). When  $M$  is smooth, this definition of framing is equivalent to a topological trivialization

$$T^v(E - \infty) \xrightarrow{\approx} \mathbb{R}^d \times (E - s_\infty)$$

that is a linear isomorphism over each point in  $E - s_\infty$  but varies only continuously instead of smoothly as the point in  $E - s_\infty$  varies.

The arguments in Sections 3 and 4 did not actually use the assumption that  $\pi : E \rightarrow B$  is a smooth fiber bundle, and everything goes through if  $\pi$  is only assumed to be a framed almost differentiable  $(M, \infty)$ -bundle, where  $M$  is a smooth homology sphere.<sup>10</sup> Therefore, we have:<sup>11</sup>

**Theorem 6.2** (restatement of Theorem 1.2) *Suppose  $M$  is a smooth homology sphere and  $\infty \in M$  is a point. Then Kontsevich's characteristic classes can be defined for framed almost differentiable  $(M, \infty)$ -bundles; when the bundle is smooth, the definition agrees with the original one.*

### 6.1 Auxiliary observations on almost differentiability

The almost differentiability condition is actually quite strong. I do not know at the time of writing if an almost differentiable manifold (respectively, an almost differentiable bundle) necessarily has a unique  $C^1$  structure, if one exists. We close this section with three auxiliary observations. Example 6.3 below shows that almost differentiability does not imply  $C^1$ . Proposition 6.4 below shows that almost differentiability implies quasiconformal. Proposition 6.5 shows that an almost differentiable map induces a linear map between tangent bundles, modulo scaling by a positive smooth function.

**Example 6.3** Let  $B_\epsilon^n$  be the standard ball of radius  $\epsilon < 1/(2e)$  in  $\mathbb{R}^n$ . Define

$$f : B_\epsilon^n \rightarrow \mathbb{R}^n, \quad f(x) = -2 \log(|x|) \cdot x.$$

Then  $f$  maps  $B_\epsilon^n$  homeomorphically onto its image, and  $f$  is almost differentiable, but not continuously differentiable at 0. See [21] for a detailed proof—the point being that the function  $-2 \log |x|$  approaches  $\infty$  slow enough as  $x \rightarrow 0$ .

The following definition of quasiconformal of a map is copied from [4]. A homeomorphism  $f : U \rightarrow \mathbb{R}^d$  from an open subset  $U$  to its image is  $k$ -quasiconformal if for all  $x \in U$

$$H_f(x) = \limsup_{r \rightarrow 0} \frac{\max\{|f(y) - f(x)| \mid |y - x| = r\}}{\min\{|f(y) - f(x)| \mid |y - x| = r\}} \leq k.$$

$f$  is quasiconformal if it is  $k$ -quasiconformal for some  $k \geq 1$ .

<sup>10</sup>That  $M$  is smooth instead of just almost differentiable is needed in the proof, to have the structure of the Fulton–MacPherson compactification of configuration spaces. However, this is mainly used in Section 3.2; specifically, to get Corollary 3.17. The argument there can probably go through by considering everything locally—trying to show that for every point  $p \in T_2$  (resp.  $T_1$ ), there is a neighborhood  $U$  of  $p$  such that  $\dim_t(T_2 \cap U)$  (resp.  $\dim_t(T_1 \cap U)$ ) is bounded as desired. If this works, the statement of Theorem 6.2 can potentially be generalized to the case when  $M$  is only almost differentiable.

<sup>11</sup>I would like to thank the anonymous referee for pointing this out.

**Proposition 6.4** *Let  $U \subset \mathbb{R}^d$  be open and  $f : U \rightarrow \mathbb{R}^d$  be an almost differentiable homeomorphism to its image. Then for every compact subset  $K \subset U$ ,  $f$  is quasiconformal on  $K$ .*


**Proof** First notice that  $f$  being almost differentiable implies: for every point  $x \in U$  there is a map  $f'_x : S^{d-1} \rightarrow S^{d-1}$  such that for any sequence of pairs of points  $\{(x_n, y_n) \in U \times U - \Delta\}_{n=1}^\infty$ ,

$$\lim_{n \rightarrow \infty} (x_n, y_n) = (x, x), \quad \lim_{n \rightarrow \infty} \frac{y_n - x_n}{|y_n - x_n|} = v \implies \lim_{n \rightarrow \infty} \frac{f(y_n) - f(x_n)}{|f(y_n) - f(x_n)|} = f'_x(v).$$

Suppose  $f$  is not quasiconformal on some  $K$ . Then for every  $k > 0$  there exists  $x_k \in K$  such that, for all  $\epsilon > 0$ , there exist  $y_k^b, y_k^s \in U$  (the superscripts stand for “big” and “small”) satisfying

$$|y_k^b - x_k| = |y_k^s - x_k| < \epsilon, \quad \frac{|f(y_k^b) - f(x_k)|}{|f(y_k^s) - f(x_k)|} > k.$$

Let  $k$  range over  $\mathbb{Z}^{>0}$ . Since  $K$  is compact, by possibly passing to a subsequence, we can assume  $x_k \rightarrow x$  as  $k \rightarrow \infty$  for some  $x \in K$ . Plugging in  $\epsilon = 1/k$  above, for every  $k$  we get a tuple of points  $x_k, y_k^b, y_k^s$ , all limit to  $x$  as  $k \rightarrow \infty$ . For each  $k$ , denote by  $S_k$  the sphere centered at  $x_k$  on which  $y_k^s, y_k^b$  lie. Define  $z_k \in S_k$  to be the midpoint of the shortest geodesic (if not unique, take an arbitrary one) between  $y_k^b, y_k^s$  on  $S_k$ . This implies that the angle between the vectors  $y_k^b - x_k$  and  $y_k^b - z_k$  is at least  $\pi/4$ ; same for the vectors  $z_k - y_k^s$  and  $z_k - x_k$ :



Since

$$\frac{|f(y_k^b) - f(x_k)|}{|f(z_k) - f(x_k)|} \cdot \frac{|f(z_k) - f(x_k)|}{|f(y_k^s) - f(x_k)|} > k,$$

one of the factors must be bigger than  $\sqrt{k}$ . In the case it is the first factor, define  $z_k^b = y_k^b, z_k^s = z_k$ ; in the case it is the second factor, define  $z_k^b = z_k, z_k^s = y_k^s$ . Now we have a sequence of tuples  $(x_k, z_k^b, z_k^s)$ , all converging to  $x$  as  $k \rightarrow \infty$ , and

$$\lim_{k \rightarrow \infty} \frac{f(z_k^b) - f(x_k)}{|f(z_k^b) - f(x_k)|} = \lim_{k \rightarrow \infty} \frac{f(z_k^b) - f(z_k^s)}{|f(z_k^b) - f(z_k^s)|},$$

because in the triangle  $(f(z_k^b), f(z_k^s), f(x_k))$ , the ratio between the lengths of the edges  $f(z_k^s)f(x_k)$  and  $f(z_k^b)f(x_k)$  goes to 0, implying that the angle between the edges  $f(z_k^b)f(z_k^s)$  and  $f(z_k^b)f(x_k)$  goes

to 0. Now, since  $f$  is almost differentiable and thus so is  $f^{-1}$ ,

$$\begin{aligned} \lim_{k \rightarrow \infty} \frac{z_k^b - x_k}{|z_k^b - x_k|} &= (f^{-1})'_{f(x)} \left( \lim_{k \rightarrow \infty} \frac{f(z_k^b) - f(x_k)}{|f(z_k^b) - f(x_k)|} \right) \\ &= (f^{-1})'_{f(x)} \left( \lim_{k \rightarrow \infty} \frac{f(z_k^b) - f(z_k^s)}{|f(z_k^b) - f(z_k^s)|} \right) = \lim_{k \rightarrow \infty} \frac{z_k^b - z_k^s}{|z_k^b - z_k^s|}, \end{aligned}$$

which contradicts that the angle between the two vectors is at least  $\pi/4$ . □

The converse to Proposition 6.4 is not true. For example,  $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ ,  $f(x) = |x|^{-1/2} \cdot x$ , is quasiconformal but not almost differentiable.

**Proposition 6.5** *Let  $U_1$  and  $U_2$  be open subsets of  $\mathbb{R}^d$  and  $f : U_1 \rightarrow U_2$  be an almost differentiable homeomorphism. For every point  $x \in U$ , denote  $\tilde{f}_x : ST_x U_1 \rightarrow ST_{f(x)} U_2$  the homeomorphism given by restricting  $\tilde{f}$  to  $\pi^{-1}(x, x)$ , where  $ST$  denotes the unit sphere in the tangent space. Then  $\tilde{f}_x$  is induced from a linear isomorphism  $T_x U_1 \rightarrow T_x U_2$ .*

**Proof** Given a directed, ordered graph  $\Gamma$  and  $U \subset \mathbb{R}^d$  an open subset, define

$$g_\Gamma^U : U^{V(\Gamma)} - \Delta_{\text{big}} \rightarrow (U \times U - \Delta)^{E(\Gamma)}, \quad (x_v)_{v \in V(\Gamma)} \longrightarrow ((x_{v_\Gamma^-(e)}, x_{v_\Gamma^+(e)}))_{e \in E(\Gamma)}.$$

Then, by the construction of Fulton–MacPherson compactification,  $g_\Gamma^U$  extends to a map

$$\bar{g}_\Gamma^U : \bar{C}_{V(\Gamma)}(U) \longrightarrow \text{Bl}_\Delta(U \times U)^{E(\Gamma)},$$

where  $\bar{C}_{V(\Gamma)}(U)$  is the Fulton–MacPherson configuration space of  $V(\Gamma)$ -labeled marked points in  $U$ . Similar to the proof of Lemma 3.2,  $\text{image}(\bar{g}_\Gamma^U)$  is the closure of  $\text{image}(g_\Gamma^U)$  in  $\text{Bl}_\Delta(U \times U)^{E(\Gamma)}$ . Let  $\pi : \text{Bl}_\Delta(U \times U)^{E(\Gamma)} \rightarrow (U \times U)^{E(\Gamma)}$  be the blow-down map. For a point  $x \in U$ ,

$$\pi^{-1}(((x, x))_{e \in E(\Gamma)}) = (ST_x U)^{E(\Gamma)} = (S^{d-1})^{E(\Gamma)}$$

and, by the construction of Fulton–MacPherson compactification,

$$\pi^{-1}(((x, x))_{e \in E(\Gamma)}) \cap \text{image}(\bar{g}_\Gamma^U) = \bar{V}_\Gamma \subset (S^{d-1})^{E(\Gamma)};$$

recall  $\bar{V}_\Gamma$  is defined in Definition 3.11. Now plug in  $U_1, U_2$  for  $U$ . From the definition of  $g_\Gamma^U$ ,

$$(f, f)^{E(\Gamma)} \circ g_\Gamma^{U_1} = g_\Gamma^{U_2} \circ f^{V(\Gamma)},$$

so  $\text{image}(g_\Gamma^{U_2}) = (f, f)^{E(\Gamma)}(\text{image}(g_\Gamma^{U_1}))$ . Passing to their closures in  $\text{Bl}_\Delta(U_i \times U_i)^{E(\Gamma)}$ , we have  $\text{image}(\bar{g}_\Gamma^{U_2}) = \tilde{f}^{E(\Gamma)}(\text{image}(\bar{g}_\Gamma^{U_1}))$ . Therefore, for each  $x \in U_1$ ,

$$(\tilde{f}_x)^{E(\Gamma)} : (S^{d-1})^{E(\Gamma)} = (ST_x U_1)^{E(\Gamma)} \longrightarrow (ST_{f(x)} U_2)^{E(\Gamma)} = (S^{d-1})^{E(\Gamma)}$$

maps  $\bar{V}_\Gamma$  to  $\bar{V}_\Gamma$ . The conclusion of the proposition follows from Lemma 6.6 below. □

**Lemma 6.6** Suppose  $f : S^{d-1} \rightarrow S^{d-1}$  is a homeomorphism such that for any directed, ordered graph  $\Gamma$ ,  $\bar{V}_\Gamma$  is invariant under  $f^{E(\Gamma)} := (f, \dots, f) : (S^{d-1})^{E(\Gamma)} \rightarrow (S^{d-1})^{E(\Gamma)}$ . Then  $f$  is induced by a map  $F \in \text{GL}(d)$ .

**Proof** (This proof is given by Fabian Gundlach.) The strategy is to define an increasing sequence of subsets  $\{A_n \subset S^{d-1}\}_{n=1}^\infty$ ,  $A_n \subset A_{n+1}$ , such that  $\bigcup_n A_n$  is dense in  $S^{d-1}$ , and show that for each  $n$ ,  $f|_{A_n}$  is induced by a map  $F_n \in \text{GL}(d)$ . For each  $n \in \mathbb{Z}^{>0}$ , let  $\Gamma_n$  be the complete graph with  $(n+1)^d$  vertices labeled by elements in  $L_n := \{0, \dots, n\}^d$ . Then, putting the vertex of  $\Gamma_n$  labeled by  $(m_1, \dots, m_d) \in L_n$  at  $(m_1, \dots, m_d) \in \mathbb{R}^d$  gives an element in  $C_{V(\Gamma_n)}^{\text{quo}}(\mathbb{R}^d)$ . Since  $\bar{V}_{\Gamma_n}$  is invariant under  $f^{E(\Gamma_n)}$ , there is an element  $x = (x_{\underline{m}} \in \mathbb{R}^d)_{\underline{m} \in L_n} \in (\mathbb{R}^d)^{V(\Gamma_n)}$  such that for any  $\underline{m}_1 \neq \underline{m}_2 \in L_n$ ,  $x_{\underline{m}_1} \neq x_{\underline{m}_2}$  and

$$(12) \quad \frac{x_{\underline{m}_1} - x_{\underline{m}_2}}{|x_{\underline{m}_1} - x_{\underline{m}_2}|} = f\left(\frac{\underline{m}_1 - \underline{m}_2}{|\underline{m}_1 - \underline{m}_2|}\right).$$

Denote  $e_j = (0, \dots, 0, 1, 0, \dots, 0) \in L_n$  where 1 is at the  $j$ -th place. We also view  $e_j$  as an element in  $S^{d-1}$ . For all  $\underline{m} \in L_n$  and  $j \in \{1, \dots, d\}$ ,  $x_{\underline{m}+e_j} - x_{\underline{m}}$  has direction  $f(e_j)$  by (12). We next show that  $|x_{\underline{m}+e_j} - x_{\underline{m}}|$  does not depend on  $\underline{m}$  either. For  $k \neq j$ , since  $x_{\underline{m} \pm e_k} - x_{\underline{m}}$  is parallel to  $x_{\underline{m}+e_j \pm e_k} - x_{\underline{m}+e_j}$ , the points  $x_{\underline{m}}, x_{\underline{m}+e_j}, x_{\underline{m} \pm e_k}, x_{\underline{m}+e_j \pm e_k}$  form a parallelogram, so we must have  $x_{\underline{m} \pm e_k + e_j} - x_{\underline{m} \pm e_k} = x_{\underline{m}+e_j} - x_{\underline{m}}$ . Therefore, plugging in  $k$  for  $j$ , we have  $x_{\underline{m}+e_k} - x_{\underline{m}} = x_{\underline{m}-e_j+e_k} - x_{\underline{m}-e_j}$  as well. So, in the two triangles  $(x_{\underline{m}}, x_{\underline{m}+e_j}, x_{\underline{m}+e_k})$  and  $(x_{\underline{m}-e_j}, x_{\underline{m}}, x_{\underline{m}-e_j+e_k})$ , one of the pairs of corresponding edges are equal as vectors. The other two pairs of corresponding edges are both parallel by (12), so they must both be equal. This shows  $x_{\underline{m}+e_j} - x_{\underline{m}} = x_{\underline{m}} - x_{\underline{m}-e_j}$ . Therefore,  $x_{\underline{m}+e_j} - x_{\underline{m}}$  does not depend on the choice of  $\underline{m}$ . Without loss of generality we can assume  $x_{(0, \dots, 0)} = (0, \dots, 0) \in \mathbb{R}^d$ . Then  $x_{\underline{m}+e_j} - x_{\underline{m}} = x_{e_j}$  for all  $\underline{m}, j$ . So,  $x_{\underline{m}_1 + \underline{m}_2} = x_{\underline{m}_1} + x_{\underline{m}_2}$  for all  $\underline{m}_1, \underline{m}_2$ . This shows that the map  $F_n \in \text{GL}(d)$  defined by “for all  $i$ ,  $F_n(e_i) = x_{e_i}$ ” maps  $\underline{m}$  to  $x_{\underline{m}}$  for all  $\underline{m} \in L_n$ . For  $F \in \text{GL}(d)$ , denote by  $\hat{F} : S^{d-1} \rightarrow S^{d-1}$  the homeomorphism induced by  $F$ . Now define

$$A_n = \left\{ \frac{\underline{m}_1 - \underline{m}_2}{|\underline{m}_1 - \underline{m}_2|} \right\}_{\underline{m}_1 \neq \underline{m}_2 \in L_n} \subset S^{d-1}.$$

Then  $f|_{A_n} = \hat{F}_n|_{A_n}$ . On the other hand, the condition  $\hat{F}_n|_{A_n} = f|_{A_n}$  uniquely determines  $F$  up to scaling: since  $\{e_i\}_{i=1}^d \subset A_n$ , there exists  $(0 \neq \lambda_i \in \mathbb{R})_{i=1}^d$  such that  $F_n(e_i) = \lambda_i f(e_i)$ ; since  $e_i + e_j \in A_n$ , the direction of  $F(e_i + e_j) = \lambda_i f(e_i) + \lambda_j f(e_j)$  is determined by  $f$ , so  $\lambda_i/\lambda_j$  is determined. Therefore, for different  $n$ ,  $F_n$  differ only by scaling. This determines a map  $F \in \text{GL}(d)$  up to scaling, satisfying  $\hat{F}|_{\bigcup_n A_n} = f|_{\bigcup_n A_n}$ . Since  $\bigcup_n A_n$  is dense in  $S^{d-1}$ ,  $\hat{F} = f$ . □

**Acknowledgements**

I thank Peter Kronheimer for many very helpful discussions throughout the course of the work. I thank Fabian Gundlach for helpful discussions regarding Example 6.3 and for providing the proof of Lemma 6.6. I would also like to thank the referees who carefully read this paper and made many valuable comments and suggestions.

## References

- [1] **R Bott, L W Tu**, *Differential forms in algebraic topology*, Graduate Texts in Math. 82, Springer (1982) MR
- [2] **G E Bredon**, *Sheaf theory*, 2nd edition, Graduate Texts in Math. 170, Springer (1997) MR
- [3] **B Conrad**, *Cech cohomology and alternating cochains*, Stanford University Available at <https://math.stanford.edu/~conrad/papers/cech.pdf>
- [4] **S K Donaldson, D P Sullivan**, *Quasiconformal 4-manifolds*, Acta Math. 163:3-4 (1989) 181–252 MR
- [5] **W Fulton, R MacPherson**, *A compactification of configuration spaces*, Ann. of Math. 139:1 (1994) 183–225 MR
- [6] **A Hatcher**, *Spectral sequences*, book project (2004) Available at <https://pi.math.cornell.edu/~hatcher/AT/SSpage.html>
- [7] **M Kontsevich**, *Feynman diagrams and low-dimensional topology*, from “First European Congress of Mathematics, II” (Paris, 1992) (A Joseph, F Mignot, F Murat, B Prum, R Rentschler, editors), Progr. Math. 120, Birkhäuser, Basel (1994) 97–121 MR
- [8] **R Koytcheff**, *Bott–Taubes–Vassiliev cohomology classes by cut-and-paste topology*, Internat. J. Math. 30:10 (2019) art. id. 1950047 MR
- [9] **P Kronheimer, T Mrowka**, *Monopoles and three-manifolds*, New Mathematical Monographs 10, Cambridge Univ. Press (2007) MR
- [10] **G Kuperberg, D P Thurston**, *Perturbative 3-manifold invariants by cut-and-paste topology*, preprint (1999) arXiv math/9912167
- [11] **C Lescop**, *Invariants of links and 3-manifolds from graph configurations*, EMS Press, Berlin (2024) MR
- [12] **J Lin, Y Xie**, *Configuration space integrals and formal smooth structures*, preprint (2023) arXiv 2310.14156
- [13] **K Nagami**, *Dimension theory*, Pure and Applied Mathematics 37, Academic Press, New York (1970) MR
- [14] **A Sard**, *Hausdorff measure of critical images on Banach manifolds*, Amer. J. Math. 87 (1965) 158–174 MR
- [15] **D P Sinha**, *Manifold-theoretic compactifications of configuration spaces*, Selecta Math. 10:3 (2004) 391–428 MR
- [16] **N E Steenrod**, *Homology with local coefficients*, Ann. of Math. 44 (1943) 610–627 MR
- [17] **N Steenrod**, *The topology of fibre bundles*, Princeton Mathematical Series 14, Princeton Univ. Press (1951) MR
- [18] **E Szpilrajn**, *La dimension et la mesure*, Fundam. Math. 28 (1937) 81–89
- [19] **T Watanabe**, *Some exotic nontrivial elements of the rational homotopy groups of  $\text{Diff}(S^4)$* , preprint (2018) arXiv 1812.02448
- [20] **T Watanabe**, *Addendum to: Some exotic nontrivial elements of the rational homotopy groups of  $\text{Diff}(S^4)$  (homological interpretation)*, preprint (2021) arXiv 2109.01609
- [21] **Xujia**, *Example of homeomorphism that lifts to real blow up but not  $C^1$ ?*, MathOverflow answer (2022) Available at <https://mathoverflow.net/q/436642>

XUJIA CHEN xchen@mpim-bonn.mpg.de  
 Max Planck Institute for Mathematics, Bonn, Germany

Received: April 30, 2024      Revised: January 28, 2025



# ALGEBRAIC & GEOMETRIC TOPOLOGY

[msp.org/agt](https://msp.org/agt)

## EDITORS

### PRINCIPAL ACADEMIC EDITORS

John Etnyre  
etnyre@math.gatech.edu  
Georgia Institute of Technology

Kathryn Hess  
kathryn.hess@epfl.ch  
École Polytechnique Fédérale de Lausanne

### BOARD OF EDITORS

Julie Bergner	University of Virginia jeb2md@eservices.virginia.edu	Thomas Koberda	University of Virginia thomas.koberda@virginia.edu
Steven Boyer	Université du Québec à Montréal cohf@math.rochester.edu	Markus Land	LMU München markus.land@math.lmu.de
Tara E Brendle	University of Glasgow tara.brendle@glasgow.ac.uk	Christine Lescop	Université Joseph Fourier lescop@ujf-grenoble.fr
Indira Chatterji	CNRS & Univ. Côte d'Azur (Nice) indira.chatterji@math.cnrs.fr	Norihiko Minami	Yamato University minami.norihiko@yamato-u.ac.jp
Octav Cornea	Université de Montreal cornea@dms.umontreal.ca	Andrés Navas	Universidad de Santiago de Chile andres.navas@usach.cl
Alexander Dranishnikov	University of Florida dranish@math.ufl.edu	Jessica S Purcell	Monash University jessica.purcell@monash.edu
Tobias Ekholm	Uppsala University, Sweden tobias.ekholm@math.uu.se	Birgit Richter	Universität Hamburg birgit.richter@uni-hamburg.de
Mario Eudave-Muñoz	Univ. Nacional Autónoma de México mario@matem.unam.mx	Jérôme Scherer	École Polytech. Féd. de Lausanne jerome.scherer@epfl.ch
David Futer	Temple University dfuter@temple.edu	Vesna Stojanoska	Univ. of Illinois at Urbana-Champaign vesna@illinois.edu
John Greenlees	University of Warwick john.greenlees@warwick.ac.uk	Zoltán Szabó	Princeton University szabo@math.princeton.edu
Matthew Hedden	Michigan State University mhedden@math.msu.edu	Maggy Tomova	University of Iowa maggy-tomova@uiowa.edu
Kristen Hendricks	Rutgers University kristen.hendricks@rutgers.edu	Daniel T Wise	McGill University, Canada daniel.wise@mcgill.ca
Hans-Werner Henn	Université Louis Pasteur henn@math.u-strasbg.fr	Lior Yanovski	Hebrew University of Jerusalem lior.yanovski@gmail.com
Daniel Isaksen	Wayne State University isaksen@math.wayne.edu		

---

See inside back cover or [msp.org/agt](https://msp.org/agt) for submission instructions.


The subscription price for 2026 is US \$795/year for the electronic version, and \$1170/year (+\$80, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Algebraic & Geometric Topology is indexed by Mathematical Reviews, Zentralblatt MATH, Current Mathematical Publications and the Science Citation Index.

Algebraic & Geometric Topology (ISSN 1472-2747 printed, 1472-2739 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004.

---

AGT peer review and production are managed by EditFlow<sup>®</sup> from MSP.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<https://msp.org/>

© 2026 Mathematical Sciences Publishers

# ALGEBRAIC & GEOMETRIC TOPOLOGY

Volume 26 Issue 1 (pages 1–410) 2026

---

The Borel cohomology of free iterated loop spaces	1
ISHAN LEVY and JUSTIN WU	
The K-theory cochains of H-spaces and height-1 chromatic homotopy theory	29
SVEN VAN NIGTEVECHT	
Cup-one algebras and 1-minimal models	65
RICHARD D. PORTER and ALEXANDER I. SUCIU	
Cubical approximation for directed topology, II	135
SANJEEVI KRISHNAN	
Flat semigroups and weighted homogeneous surface singularities	201
ZSOLT BAJA and TAMÁS LÁSZLÓ	
Interleaving distances, monoidal actions and 2-categories	227
PATRICK K. MCFADDIN and TOM NEEDHAM	
Suspension splittings of 5-dimensional Poincaré duality complexes and their applications	283
STEVEN AMELOTTE, TYRONE CUTLER and TSELEUNG SO	
Bernoulli shifts on additive categories and algebraic $K$ -theory of wreath products	321
JULIAN KRANZ and SHINTARO NISHIKAWA	
Kontsevich's characteristic classes as topological invariants of configuration space bundles	349
XUJIA CHEN	
Profinite completions of products	397
PETER J. HAINE	