

AG
T

*Algebraic & Geometric
Topology*

Volume 26 (2026)

Profinite completions of products

PETER J. HAINE



Profinite completions of products

PETER J. HAINE

A source of difficulty in profinite homotopy theory is that the profinite completion functor does not preserve finite products. We provide a new, checkable criterion on prospaces X and Y that guarantees that the profinite completion of $X \times Y$ agrees with the product of the profinite completions of X and Y . Using this criterion, we show that profinite completion preserves products of étale homotopy types of qcqs schemes. This fills a gap in Chough’s proof of the Künneth formula for the étale homotopy type of a product of proper schemes over a separably closed field.

0. Introduction	397
1. Universality of colimits	399
2. Completions of products	403
Appendix. Classifying prospaces via geometric realizations	407
Acknowledgements	409
References	409

0 Introduction

Write \mathbf{Spc}_π for the ∞ -category of π -finite spaces. Given a set Σ of primes, write $\mathbf{Spc}_\Sigma \subset \mathbf{Spc}_\pi$ for the full subcategory spanned by those π -finite spaces whose homotopy groups have orders divisible only by primes in Σ . Write

$$(-)_{\Sigma}^{\wedge} : \mathrm{Pro}(\mathbf{Spc}) \rightarrow \mathrm{Pro}(\mathbf{Spc}_{\Sigma})$$

for the Σ -completion functor, i.e., the left adjoint to the inclusion $\mathrm{Pro}(\mathbf{Spc}_{\Sigma}) \subset \mathrm{Pro}(\mathbf{Spc})$. One source of difficulty in profinite homotopy theory is that the Σ -completion functor does not preserve finite limits, or even finite products (see [5, Remark 3.10; 15, Remark E.5.2.6]).

As far as we are aware, given connected spaces X and Y , the only general condition to check that the profinite completion of $X \times Y$ is the product of the profinite completions of X and Y is to check that the homotopy groups of X and Y are *good* in the sense of Serre [5, Proposition 3.9]. However, it is generally quite difficult to check if a group is good, and there are hard conjectures about whether or not naturally occurring groups are good. For example, Deligne and Morava’s conjecture that the mapping class group $\Gamma_{g,n}$ of a genus- g curve with n marked points is good [19, problem on p. 94] is still open.

The purpose of this note is to provide the following new, checkable criterion on prospaces X and Y that guarantees that the natural map $(X \times Y)_{\Sigma}^{\wedge} \rightarrow X_{\Sigma}^{\wedge} \times Y_{\Sigma}^{\wedge}$ is an equivalence.

0.1 Theorem (Theorem 2.13) *Let X and Y be prospaces and let Σ be a set of primes. If X and Y are both in the smallest full subcategory of $\text{Pro}(\mathbf{Spc})$ containing $\text{Pro}(\mathbf{Spc}_\pi)$ that is closed under geometric realizations of simplicial objects, retracts, and cofiltered limits, then the natural map*

$$(X \times Y)_\Sigma^\wedge \rightarrow X_\Sigma^\wedge \times Y_\Sigma^\wedge$$

is an equivalence.

In fact, [Theorem 2.13](#) is a bit more general. However, the more general statement requires introducing a bit of terminology, so we refer the reader to later in the text for the statement.

Now we state the two key applications. The first is that if one is already in the setting of profinite homotopy theory, then Σ -completion preserves products:

0.2 Corollary (Corollary 2.9) *Let Σ be a set of primes. Then the Σ -completion functor restricted to profinite spaces*

$$(-)_\Sigma^\wedge : \text{Pro}(\mathbf{Spc}_\pi) \rightarrow \text{Pro}(\mathbf{Spc}_\Sigma)$$

preserves products.

The second is that in the setting of étale homotopy theory, Σ -completion preserves finite products. Given a scheme X , write $\Pi_\infty^{\text{ét}}(X) \in \text{Pro}(\mathbf{Spc})$ for the étale homotopy type of X . Our work with Barwick and Glasman [[1](#), Theorems 10.2.3 and 12.5.1] provides a description of the (protruncated) étale homotopy type of a qcqs scheme as the geometric realization of an explicit simplicial profinite space. The existence of this presentation implies:

0.3 Corollary (Example 2.17) *Let Σ be a set of primes and let X and Y be qcqs schemes. Then the natural map of profinite spaces*

$$(\Pi_\infty^{\text{ét}}(X) \times \Pi_\infty^{\text{ét}}(Y))_\Sigma^\wedge \rightarrow \Pi_\infty^{\text{ét}}(X)_\Sigma^\wedge \times \Pi_\infty^{\text{ét}}(Y)_\Sigma^\wedge$$

is an equivalence.

0.4 Remark [Corollary 0.3](#) fills a gap in Chough's proof of the Künneth formula for the étale homotopy type of a product of proper schemes over a separably closed field [[7](#), Theorem 5.3]. Chough's proof cites the false claim that profinite completion preserves finite limits. However, what Chough actually uses is [Corollary 0.3](#) (with Σ the set of all primes). In particular, the conclusion of [[7](#), Theorem 5.3] remains valid.

In our work with Holzschuh and Wolf [[9](#), §4], [Corollary 0.3](#) is a key ingredient used to prove other Künneth formulas in étale homotopy theory.

0.5 Proof Strategy The first observation is that since protruncation preserves limits [[8](#), Proposition 3.9], it suffices to prove a variant of [Theorem 0.1](#) where X and Y are already protruncated. Since equivalences of protruncated spaces are detected on all truncations (which is not true for general prospaces), this simplifies the situation. Second, since Σ -completion preserves cofiltered limits, [Theorem 0.1](#) would follow if we knew that geometric realizations preserved finite products in the ∞ -categories of protruncated and Σ -profinite spaces. See [Lemma 2.7](#) and [Corollary 2.8](#) for the key categorical result explaining this.

With these reductions, the main technical step in our argument is to show the stronger claim that geometric realizations are *universal* in the ∞ -categories of protruncated and Σ -profinite spaces. In fact, we show that the colimit of any diagram which can be evaluated as a finite colimit in every n -category for $n < \infty$ (see [Definition 1.9](#)) is universal in these ∞ -categories. See [Proposition 1.17](#) and [Corollary 1.18](#).

0.6 Linear Overview [Section 1](#) proves that geometric realizations are universal in the ∞ -categories of protruncated and profinite spaces. In particular, geometric realizations preserve finite products in these ∞ -categories. [Section 2](#) proves [Corollaries 0.2](#) and [0.3](#). It is immediate from [[1](#), Theorem 10.2.3] that the protruncated étale homotopy type can be written as the geometric realization of a simplicial profinite space. However, for ease of reference we have provided a detailed explanation of this fact in the [Appendix](#).

0.7 Conventions Throughout, we use the notational conventions of [[8](#), §§1 and 3]. In an effort to keep this note short, we do not recapitulate them here.

1 Universality of colimits

In this section, we prove that geometric realizations are universal in the ∞ -categories of protruncated and Σ -profinite spaces. We accomplish this by proving a more general fact: colimits over diagrams that can be computed as finite colimits when valued in an n -category (see [Definition 1.9](#)) are universal in the ∞ -categories of protruncated and Σ -profinite spaces ([Proposition 1.17](#) and [Corollary 1.18](#)).

The first observation is that finite colimits are universal in protruncated spaces.

1.1 Lemma *Let \mathcal{C} be an ∞ -category with pullbacks and finite colimits. If finite colimits are universal in \mathcal{C} , then finite colimits are universal in $\text{Pro}(\mathcal{C})$.*

Proof By (the dual of) [[12](#), [Proposition 5.3.5.15](#)], pullbacks, pushouts, and finite coproducts are computed “levelwise” in $\text{Pro}(\mathcal{C})$. Thus the assumption that finite colimits are universal in \mathcal{C} implies that pushouts and finite coproducts are universal in $\text{Pro}(\mathcal{C})$. Since a functor that preserves pushouts and finite coproducts preserves finite colimits, we deduce that all finite colimits are universal in $\text{Pro}(\mathcal{C})$. \square

1.2 Example Finite colimits are universal in $\text{Pro}(\mathbf{Spc})$. For each integer $n \geq 0$, finite colimits are universal in $\text{Pro}(\mathbf{Spc}_{\leq n})$.

1.3 Recollection A localization $L : \mathcal{C} \rightarrow \mathcal{D}$ is *locally cartesian* if for any cospan $X \rightarrow Z \leftarrow Y$ such that $X, Y \in \mathcal{D}$, the natural map $L(X \times_Z Y) \rightarrow X \times_Z L(Y)$ is an equivalence.

1.4 Example [[8](#), [Proposition 3.18](#)] For any set Σ of primes, the localization

$$(-)_{\Sigma}^{\wedge} : \text{Pro}(\mathbf{Spc}_{< \infty}) \rightarrow \text{Pro}(\mathbf{Spc}_{\Sigma})$$

is locally cartesian. However, $(-)_{\Sigma}^{\wedge}$ does not generally preserve finite products.

The following is immediate from the definitions:

1.5 Lemma Let \mathcal{J} be an ∞ -category, \mathcal{C} an ∞ -category with pullbacks and \mathcal{J} -shaped colimits, and let $L : \mathcal{C} \rightarrow \mathcal{D}$ be a locally cartesian localization. If \mathcal{J} -shaped colimits are universal in \mathcal{C} , then \mathcal{J} -shaped colimits are universal in \mathcal{D} .

1.6 Example Since the pro-truncation functor $\tau_{<\infty} : \text{Pro}(\mathbf{Spc}) \rightarrow \text{Pro}(\mathbf{Spc}_{<\infty})$ preserves limits [8, Proposition 3.9], Example 1.2 and Lemma 1.5 show that finite colimits are universal in $\text{Pro}(\mathbf{Spc}_{<\infty})$.

Now we formulate the key property of the category $\mathbf{\Delta}^{\text{op}}$ that we need.

1.7 Definition Let $n \geq 0$ be an integer. A functor between ∞ -categories $c : \mathcal{J} \rightarrow \mathcal{I}$ is n -colimit-cofinal if for every n -category \mathcal{C} and functor $f : \mathcal{I} \rightarrow \mathcal{C}$, the following conditions are satisfied:

- (1) The colimit $\text{colim}_{\mathcal{J}} f$ exists if and only if the colimit $\text{colim}_{\mathcal{I}} fc$ exists.
- (2) If the colimit $\text{colim}_{\mathcal{I}} fc$ exists, then the natural map $\text{colim}_{\mathcal{I}} fc \rightarrow \text{colim}_{\mathcal{J}} f$ is an equivalence.

1.8 Example For an integer $n \geq 0$, write $\mathbf{\Delta}_{\leq n} \subset \mathbf{\Delta}$ for the full subcategory spanned by those nonempty linearly ordered finite sets of cardinality $\leq n + 1$. By [10, Proposition A.1], the inclusion $\mathbf{\Delta}_{\leq n}^{\text{op}} \subset \mathbf{\Delta}^{\text{op}}$ is n -colimit-cofinal.

1.9 Definition Let \mathcal{J} be an ∞ -category. We say that \mathcal{J} is *almost finite* if for each integer $n \geq 0$, there exists a *finite* ∞ -category \mathcal{J}_n and an n -colimit-cofinal functor $c_n : \mathcal{J}_n \rightarrow \mathcal{J}$.

Here are a number of important examples of almost finite ∞ -categories.

1.10 Example If \mathcal{J} is an ∞ -category that admits a colimit-cofinal functor from a finite ∞ -category, then \mathcal{J} is almost finite.

1.11 Example For each $n \geq 0$, the category $\mathbf{\Delta}_{\leq n}^{\text{op}}$ is a finite ∞ -category [6, Example 6.5.3]. Hence the category $\mathbf{\Delta}^{\text{op}}$ is almost finite: the inclusion $\mathbf{\Delta}_{\leq n}^{\text{op}} \hookrightarrow \mathbf{\Delta}^{\text{op}}$ is an n -colimit-cofinal functor from a finite ∞ -category.

1.12 Definition Let K be a simplicial set. The ∞ -category *presented by* K is the image of K under the natural functor $\mathbf{sSet} \rightarrow \mathbf{Cat}_{\infty}$ obtained by inverting the weak equivalences in the Joyal model structure. The *space presented by* K is the image of K under the natural functor $\mathbf{sSet} \rightarrow \mathbf{Spc}$ obtained by inverting the weak equivalences in the Kan–Quillen model structure.

1.13 Example Let K be a simplicial set with finitely many simplices in each dimension and let \mathcal{J} be the ∞ -category presented by K . Then \mathcal{J} is almost finite: we take \mathcal{J}_n to be the ∞ -category presented by the $(n+1)$ -skeleton of $\text{sk}_{n+1} K$ and $c_n : \mathcal{J}_n \rightarrow \mathcal{J}$ the functor induced by the inclusion $\text{sk}_{n+1} K \subset K$.

1.14 Recollection A space X is *almost π -finite* if $\pi_0(X)$ is finite and all homotopy groups of X are finite. An almost π -finite space admits a presentation by a Kan complex with finitely many simplices in each dimension [15, Lemma E.1.6.5]. Hence:

1.15 Example As a special case of Example 1.13, every almost π -finite space is an almost finite ∞ -category.

1.16 Definition Let \mathcal{C} be an ∞ -category with pullbacks. We say that *almost finite colimits are universal in \mathcal{C}* if for each almost finite ∞ -category \mathcal{J} , the ∞ -category \mathcal{C} admits \mathcal{J} -shaped colimits and \mathcal{J} -shaped colimits are universal in \mathcal{C} .

The argument that Lurie gives in the proof of [15, Theorem E.6.3.1] essentially shows that almost finite colimits are universal in $\text{Pro}(\mathbf{Spc}_\Sigma)$. However, the statement about universality of colimits is less general and the result is only stated when Σ is the set of all primes. We also need to know that almost finite colimits are also universal in $\text{Pro}(\mathbf{Spc}_{<\infty})$. The strategy is the same as Lurie’s proof: we use that equivalences are checked on truncations to reduce to the case of finite colimits.

1.17 Proposition *Almost finite colimits are universal in $\text{Pro}(\mathbf{Spc}_{<\infty})$.*

Moreover, Proposition 1.17 strengthens [15, Theorem E.6.3.1]:

1.18 Corollary *Let Σ be a set of primes. Then almost finite colimits are universal in $\text{Pro}(\mathbf{Spc}_\Sigma)$.*

Proof of Corollary 1.18 Since the localization $(-)_\Sigma^\wedge : \text{Pro}(\mathbf{Spc}_{<\infty}) \rightarrow \text{Pro}(\mathbf{Spc}_\Sigma)$ is locally cartesian [8, Proposition 3.18], this follows from Lemma 1.5 and Proposition 1.17. \square

Proof of Proposition 1.17 Let \mathcal{J} be an almost finite ∞ -category, let $f : X \rightarrow Z$ be a morphism in $\text{Pro}(\mathbf{Spc}_{<\infty})$, and let

$$g : \mathcal{J} \rightarrow \text{Pro}(\mathbf{Spc}_{<\infty})/Z$$

be a diagram of protruncated spaces over Z . To prove the claim, it suffices to show that for each integer $n \geq 0$, the induced map

$$\tau_{\leq n}(\text{colim}_{i \in \mathcal{J}} X \times_Z g(i)) \rightarrow \tau_{\leq n}(X \times_Z \text{colim}_{i \in \mathcal{J}} g(i))$$

is an equivalence in $\text{Pro}(\mathbf{Spc}_{\leq n})$. Since \mathcal{J} is almost finite, there exists a finite ∞ -category \mathcal{J}_{n+2} and $(n+2)$ -colimit-cofinal functor $c_{n+2} : \mathcal{J}_{n+2} \rightarrow \mathcal{J}$. Consider the commutative diagram

$$\begin{array}{ccccc} \text{colim}_{j \in \mathcal{J}_{n+2}} \tau_{\leq n}(X \times_Z g c_{n+2}(j)) & \xrightarrow{\sim} & \tau_{\leq n}(\text{colim}_{j \in \mathcal{J}_{n+2}} X \times_Z g c_{n+2}(j)) & \longrightarrow & \tau_{\leq n}(X \times_Z \text{colim}_{j \in \mathcal{J}_{n+2}} g c_{n+2}(j)) \\ \downarrow & & \downarrow & & \downarrow \\ \text{colim}_{i \in \mathcal{J}} \tau_{\leq n}(X \times_Z g(i)) & \xrightarrow{\sim} & \tau_{\leq n}(\text{colim}_{i \in \mathcal{J}} X \times_Z g(i)) & \longrightarrow & \tau_{\leq n}(X \times_Z \text{colim}_{i \in \mathcal{J}} g(i)) \end{array}$$

(Here, the colimits in the leftmost column are computed in $\text{Pro}(\mathbf{Spc}_{\leq n})$.) Since $\text{Pro}(\mathbf{Spc}_{\leq n})$ is an $(n+1)$ -category and $c_{n+2} : \mathcal{J}_{n+2} \rightarrow \mathcal{J}$ is $(n+2)$ -colimit-cofinal, the leftmost vertical map is an equivalence. Thus the central vertical map is also an equivalence. Since \mathcal{J}_{n+2} is finite and finite colimits are universal in $\text{Pro}(\mathbf{Spc}_{<\infty})$ (Example 1.6), the top right-hand horizontal map is an equivalence.

To complete the proof, it suffices to show that the rightmost vertical map is an equivalence. For this, consider the commutative square

$$\begin{array}{ccc}
 \operatorname{colim}_{j \in \mathcal{J}_{n+2}} \tau_{\leq n+1} g c_{n+2}(j) & \xrightarrow{\sim} & \tau_{\leq n+1}(\operatorname{colim}_{j \in \mathcal{J}_{n+2}} g c_{n+2}(j)) \\
 \downarrow & & \downarrow \\
 \operatorname{colim}_{i \in \mathcal{J}} \tau_{\leq n+1} g(i) & \xrightarrow{\sim} & \tau_{\leq n+1}(\operatorname{colim}_{i \in \mathcal{J}} g(i))
 \end{array}$$

where the colimits in the left-hand column are computed in $\operatorname{Pro}(\mathbf{Spc}_{\leq n+1})$. Since $\operatorname{Pro}(\mathbf{Spc}_{\leq n+1})$ is an $(n+2)$ -category and c_{n+2} is $(n+2)$ -colimit-cofinal, the left-hand vertical map is an equivalence. Hence the right-hand vertical map is also an equivalence. As a consequence, the map

$$\operatorname{colim}_{j \in \mathcal{J}_{n+2}} g c_{n+2}(j) \rightarrow \operatorname{colim}_{i \in \mathcal{J}} g(i)$$

is n -connected. Thus the base change

$$X \times_Z \operatorname{colim}_{j \in \mathcal{J}_{n+2}} g c_{n+2}(j) \rightarrow X \times_Z \operatorname{colim}_{i \in \mathcal{J}} g(i)$$

is also n -connected. Hence the map

$$\tau_{\leq n}(X \times_Z \operatorname{colim}_{j \in \mathcal{J}_{n+2}} g c_{n+2}(j)) \rightarrow \tau_{\leq n}(X \times_Z \operatorname{colim}_{i \in \mathcal{J}} g(i))$$

is an equivalence, as desired. □

The universality of geometric realizations implies that geometric realizations preserve finite products:

1.19 Lemma *Let \mathcal{J} be a sifted ∞ -category and let \mathcal{C} be an ∞ -category with finite limits and \mathcal{J} -shaped colimits. If \mathcal{J} -shaped colimits are universal in \mathcal{C} , then the functor*

$$\operatorname{colim}_{\mathcal{J}} : \operatorname{Fun}(\mathcal{J}, \mathcal{C}) \rightarrow \mathcal{C}$$

preserves finite products.

Proof Let $X_{\bullet}, Y_{\bullet} : \mathcal{J} \rightarrow \mathcal{C}$ be functors. We have natural equivalences

$$\begin{aligned}
 \operatorname{colim}_{i \in \mathcal{J}} X_i \times Y_i &\xrightarrow{\sim} \operatorname{colim}_{(i,j) \in \mathcal{J} \times \mathcal{J}} X_i \times Y_j && (\mathcal{J} \text{ is sifted}) \\
 &\simeq \operatorname{colim}_{i \in \mathcal{J}} \operatorname{colim}_{j \in \mathcal{J}} (X_i \times Y_j) \\
 &\xrightarrow{\sim} \operatorname{colim}_{i \in \mathcal{J}} (X_i \times \operatorname{colim}_{j \in \mathcal{J}} Y_j) && (\mathcal{J}\text{-shaped colimits are universal}) \\
 &\xrightarrow{\sim} (\operatorname{colim}_{i \in \mathcal{J}} X_i) \times (\operatorname{colim}_{j \in \mathcal{J}} Y_j) && (\mathcal{J}\text{-shaped colimits are universal}). \quad \square
 \end{aligned}$$

1.20 Corollary *Let Σ be a set of primes. Then geometric realizations preserve finite products in the ∞ -categories $\operatorname{Pro}(\mathbf{Spc}_{\Sigma})$ and $\operatorname{Pro}(\mathbf{Spc}_{< \infty})$.*

Proof Combine [Proposition 1.17](#), [Corollary 1.18](#), and [Lemma 1.19](#). □

2 Completions of products

The goal of this section is to prove Corollaries 0.2 and 0.3. We do this by noting that in the setting of both of these results, the prospaces of interest can be written as geometric realizations of simplicial profinite spaces. We begin by axiomatizing the situation:

2.1 Definition Let \mathcal{C} be an ∞ -category, let $\mathcal{D} \subset \mathcal{C}$ be a full subcategory, and let $X \in \mathcal{C}$. A \mathcal{D} -resolution of X is a simplicial object $X_\bullet : \mathbf{\Delta}^{\text{op}} \rightarrow \mathcal{D}$ together with an equivalence

$$X \simeq \text{colim}(\mathbf{\Delta}^{\text{op}} \xrightarrow{X_\bullet} \mathcal{D} \hookrightarrow \mathcal{C}).$$

We say that an object X admits a \mathcal{D} -resolution if there exists \mathcal{D} -resolution of X .

2.2 Notation Write $\mathbf{Set}_{\text{fin}}$ for the category of finite sets and all maps.

Finite and almost π -finite spaces admit $\mathbf{Set}_{\text{fin}}$ -resolutions:

2.3 Lemma Let X be a space which is finite¹ or almost π -finite. Then:

- (1) As an object of \mathbf{Spc} , the space X admits a $\mathbf{Set}_{\text{fin}}$ -resolution.
- (2) When regarded as a constant object of $\text{Pro}(\mathbf{Spc})$, the space X admits a $\mathbf{Set}_{\text{fin}}$ -resolution.
- (3) The protruncated space $\tau_{<\infty}(X)$ admits a $\mathbf{Set}_{\text{fin}}$ -resolution.
- (4) The profinite space X_π^\wedge admits a $\mathbf{Set}_{\text{fin}}$ -resolution.

Proof For (1), if X is finite, then X can be written as the geometric realization of a simplicial set with finitely many nondegenerate simplices [21, Proposition 2.5].² If X is almost π -finite, then X can be written as the geometric realization of a Kan complex with finitely many simplices in each dimension [15, Lemma E.1.6.5].

Now note that (2)–(4) follow from (1) and that each functor in the diagram

$$\mathbf{Spc} \hookrightarrow \text{Pro}(\mathbf{Spc}) \xrightarrow{\tau_{<\infty}} \text{Pro}(\mathbf{Spc}_{<\infty}) \xrightarrow{(-)^\wedge_\pi} \text{Pro}(\mathbf{Spc}_\pi)$$

preserves colimits. □

Algebraic geometry also gives rise to many examples of protruncated spaces admitting profinite resolutions.

2.4 Notation (shapes) Given an ∞ -topos X , we write $\Pi_\infty(X) \in \text{Pro}(\mathbf{Spc})$ for the *shape* of X . We write $\Pi_{<\infty}(X)$ for the protruncation of $\Pi_\infty(X)$.

¹I.e., in the smallest subcategory of \mathbf{Spc} containing the point and the empty set and closed under pushouts. Equivalently, a space X is finite if and only if X is represented by a finite CW complex.

²In fact, every finite space is equivalent to the classifying space of a finite poset; see [12, Proposition 4.1.1.3(3) and Variant 4.2.3.16; 16, Tag 02MU; 18, Theorem 1].

2.5 Recall that an ∞ -topos X is *spectral* in the sense of [1, Definition 9.2.1] if X is bounded coherent and the ∞ -category $\text{Pt}(X)$ of points of X has the property that every endomorphism of an object of $\text{Pt}(X)$ is an equivalence. The most important example of a spectral ∞ -topos is the étale ∞ -topos of a qcqs scheme [1, Example 9.2.4]. **Example A.9** explains why the protruncated shape $\Pi_{<\infty}(X)$ of a spectral ∞ -topos admits a natural $\text{Pro}(\mathbf{Spc}_\pi)$ -resolution.

The next few results are the key categorical input we need. To state them, we axiomatize the abstract properties of the subcategory $\text{Pro}(\mathbf{Spc}_\pi) \subset \text{Pro}(\mathbf{Spc}_{<\infty})$.

2.6 Notation Let \mathcal{C} be an ∞ -category with geometric realizations and finite products, and let $L : \mathcal{C} \rightarrow \mathcal{D}$ be a localization. Assume that geometric realizations preserve finite products in both \mathcal{C} and \mathcal{D} . Write $\mathcal{C}_{|\mathcal{D}|} \subset \mathcal{C}$ for the smallest full subcategory containing $\mathcal{D} \subset \mathcal{C}$ and closed under geometric realizations and retracts.

2.7 Lemma *In the setting of Notation 2.6, let $X_\bullet, Y_\bullet : \Delta^{\text{op}} \rightarrow \mathcal{C}$ be simplicial objects with colimits X and Y . Assume that for each $n \geq 0$, the natural map*

$$L(X_n \times Y_n) \rightarrow L(X_n) \times L(Y_n)$$

is an equivalence. Then the natural map $L(X \times Y) \rightarrow L(X) \times L(Y)$ is an equivalence.

Proof We compute

$$\begin{aligned} L(X \times Y) &\simeq L(\text{colim}_{m \in \Delta^{\text{op}}} X_m \times \text{colim}_{n \in \Delta^{\text{op}}} Y_n) \\ &\simeq L(\text{colim}_{n \in \Delta^{\text{op}}} X_n \times Y_n) && \text{(geometric realizations preserve finite products in } \mathcal{C} \text{)} \\ &\simeq \text{colim}_{n \in \Delta^{\text{op}}} L(X_n \times Y_n) \\ &\xrightarrow{\simeq} \text{colim}_{n \in \Delta^{\text{op}}} L(X_n) \times L(Y_n) && \text{(assumption)} \\ &\simeq \text{colim}_{m \in \Delta^{\text{op}}} L(X_m) \times \text{colim}_{n \in \Delta^{\text{op}}} L(Y_n) && \text{(geometric realizations preserve finite products in } \mathcal{D} \text{)} \\ &\simeq L(X) \times L(Y). \end{aligned}$$

□

2.8 Corollary *In the setting of Notation 2.6:*

(1) *The full subcategory of $\mathcal{C} \times \mathcal{C}$ spanned by those objects (X, Y) such that the natural map*

$$L(X \times Y) \rightarrow L(X) \times L(Y)$$

is an equivalence is closed under geometric realizations and retracts.

(2) *If $X, Y \in \mathcal{C}_{|\mathcal{D}|}$, then the natural map $L(X \times Y) \rightarrow L(X) \times L(Y)$ is an equivalence.*

(3) *If $X, Y \in \mathcal{C}$ admit \mathcal{D} -resolutions, then the natural map $L(X \times Y) \rightarrow L(X) \times L(Y)$ is an equivalence.*

Proof Item (1) follows from **Lemma 2.7** and the fact that equivalences are closed under retracts. Now (2) is an immediate consequence of (1) and the definition of $\mathcal{C}_{|\mathcal{D}|}$ as the closure of $\mathcal{D} \subset \mathcal{C}$ under geometric realizations and retracts. Finally, (3) is a special case of (2). □

We now record some consequences of [Corollary 2.8](#).

2.9 Corollary *Let Σ be a set of prime numbers. Then the Σ -completion functor*

$$(-)_{\Sigma}^{\wedge} : \text{Pro}(\mathbf{Spc}_{\pi}) \rightarrow \text{Pro}(\mathbf{Spc}_{\Sigma})$$

preserves products.

Proof Since Σ -completion preserves cofiltered limits and the terminal object, it suffices to show that Σ -completion preserves binary products of profinite spaces. Again because Σ -completion preserves cofiltered limits, we are reduced to showing that if X and Y are π -finite spaces, then the natural map

$$(X \times Y)_{\Sigma}^{\wedge} \rightarrow X_{\Sigma}^{\wedge} \times Y_{\Sigma}^{\wedge}$$

is an equivalence. Since π -finite spaces admit $\mathbf{Set}_{\text{fin}}$ -resolutions ([Lemma 2.3](#)) and geometric realizations preserve finite products in $\text{Pro}(\mathbf{Spc}_{\pi})$ and $\text{Pro}(\mathbf{Spc}_{\Sigma})$ ([Corollary 1.20](#)), the claim follows from [Corollary 2.8](#). \square

2.10 Warning The functor $(-)_{\Sigma}^{\wedge} : \text{Pro}(\mathbf{Spc}_{\pi}) \rightarrow \text{Pro}(\mathbf{Spc}_{\Sigma})$ does not generally preserve pullbacks, or even loop objects.

We now introduce a slight enlargement of the subcategory of protruncated spaces admitting a $\text{Pro}(\mathbf{Spc}_{\pi})$ -resolution on which profinite completion preserves finite products.

2.11 Notation Let

$$\text{Pro}(\mathbf{Spc}_{<\infty})' \subset \text{Pro}(\mathbf{Spc}_{<\infty})$$

denote the smallest full subcategory containing $\text{Pro}(\mathbf{Spc}_{\pi})$ and closed under geometric realizations, retracts, and cofiltered limits.

2.12 Observation (procompact spaces) Write $\mathbf{Spc}^{\omega} \subset \mathbf{Spc}$ for the full subcategory spanned by the compact objects.³ Then by [Lemma 2.3](#), the image of

$$\tau_{<\infty} : \text{Pro}(\mathbf{Spc}^{\omega}) \rightarrow \text{Pro}(\mathbf{Spc}_{<\infty})$$

is contained in $\text{Pro}(\mathbf{Spc}_{<\infty})'$.

The following is the main result of this note.

2.13 Theorem *Let Σ be a set of primes and let $X, Y \in \text{Pro}(\mathbf{Spc}_{<\infty})'$. Then the natural map*

$$(X \times Y)_{\Sigma}^{\wedge} \rightarrow X_{\Sigma}^{\wedge} \times Y_{\Sigma}^{\wedge}$$

is an equivalence.

³A space X is compact if and only if X is a retract of a finite space. In more classical terminology, a space X is compact if and only if X is represented by a *finitely dominated* CW complex.

Proof By [Corollary 2.9](#), the Σ -completion functor

$$(-)_{\Sigma}^{\wedge} : \text{Pro}(\mathbf{Spc}_{\pi}) \rightarrow \text{Pro}(\mathbf{Spc}_{\Sigma})$$

preserves products. Hence it suffices to prove the claim in the special case where Σ is the set of all primes. For this, note that cofiltered limits preserve finite products in $\text{Pro}(\mathbf{Spc}_{<\infty})$ and $\text{Pro}(\mathbf{Spc}_{\pi})$; moreover, the profinite completion functor preserves cofiltered limits. Thus it suffices to treat the case where X and Y are in the smallest full subcategory of $\text{Pro}(\mathbf{Spc}_{<\infty})$ containing $\text{Pro}(\mathbf{Spc}_{\pi})$ and closed under geometric realizations and retracts. In this case, since geometric realizations preserve finite products in $\text{Pro}(\mathbf{Spc}_{<\infty})$ and $\text{Pro}(\mathbf{Spc}_{\pi})$ ([Corollary 1.20](#)), [Corollary 2.8](#) completes the proof. \square

2.14 Example If X and Y are protruncated spaces that admit $\text{Pro}(\mathbf{Spc}_{\pi})$ -resolutions, then the natural map $(X \times Y)_{\Sigma}^{\wedge} \rightarrow X_{\Sigma}^{\wedge} \times Y_{\Sigma}^{\wedge}$ is an equivalence.

2.15 Example If $X, Y \in \mathbf{Spc}$ are compact, then the natural map $(X \times Y)_{\Sigma}^{\wedge} \rightarrow X_{\Sigma}^{\wedge} \times Y_{\Sigma}^{\wedge}$ is an equivalence.

We conclude by recording two applications of [Theorem 2.13](#).

2.16 Corollary Let Σ be a set of primes and let X and Y be spectral ∞ -topoi. Then the natural map

$$(\Pi_{\infty}(X) \times \Pi_{\infty}(Y))_{\Sigma}^{\wedge} \rightarrow \Pi_{\infty}(X)_{\Sigma}^{\wedge} \times \Pi_{\infty}(Y)_{\Sigma}^{\wedge}$$

is an equivalence.

Proof [Example A.9](#) shows that $\Pi_{<\infty}(X)$ and $\Pi_{<\infty}(Y)$ admit $\text{Pro}(\mathbf{Spc}_{\pi})$ -resolutions. Since protruncation preserves products, the claim now follows from [Theorem 2.13](#). \square

2.17 Example Let Σ be a set of primes and let X and Y be qcqs schemes. Then the natural map

$$(\Pi_{\infty}^{\acute{e}t}(X) \times \Pi_{\infty}^{\acute{e}t}(Y))_{\Sigma}^{\wedge} \rightarrow \Pi_{\infty}^{\acute{e}t}(X)_{\Sigma}^{\wedge} \times \Pi_{\infty}^{\acute{e}t}(Y)_{\Sigma}^{\wedge}$$

is an equivalence.

The second application is to ∞ -operads. Recently, it has become clear that it is important to consider profinite completions of ∞ -operads. For example, Boavida de Brito, Horel, and Robertson explained a beautiful relationship between the Grothendieck–Teichmüller group and the profinite completion of the genus-0 surface operad [\[5\]](#). However, since profinite completion does not preserve products, profinite completions of ∞ -operads generally cannot be computed “levelwise”. In general, more care is required to work with profinite completions of ∞ -operads; this is the topic of recent work of Blom and Moerdijk [\[3; 4\]](#).

An immediate consequence of [Theorem 2.13](#) is that profinite completions *can* be computed levelwise more generally than was previously known:

2.18 Corollary Let Σ be a set of primes and let $\{\mathcal{O}(n)\}_{n \geq 0}$ be an ∞ -operad in spaces. Assume that for each $n \geq 0$, we have

$$\tau_{<\infty} \mathcal{O}(n) \in \text{Pro}(\mathbf{Spc}_{<\infty})'.$$

Then the levelwise Σ -completion $\{\mathcal{O}(n)_{\Sigma}^{\wedge}\}_{n \geq 0}$ defines an ∞ -operad in $\text{Pro}(\mathbf{Spc}_{\Sigma})$.

Appendix Classifying prospaces via geometric realizations

The purpose of this appendix is to explain why the protruncated shape of a spectral ∞ -topos (e.g., the étale ∞ -topos of a qcqs scheme) admits a presentation as a geometric realization of a simplicial profinite space (Example A.9). Using the description of the protruncated shape of a spectral ∞ -topos as a protruncated classifying prospace given by Barwick–Glasman–Haine [1, Theorem 10.2.3], this is an exercise in the definitions. For the ease of the reader, we spell out the details here.

We make use of the description of ∞ -categories as simplicial spaces.

A.1 Recollection (∞ -categories as simplicial spaces) The nerve functor

$$N : \mathbf{Cat}_\infty \rightarrow \mathbf{Fun}(\mathbf{\Delta}^{\text{op}}, \mathbf{Spc}), \quad \mathcal{C} \mapsto [I \mapsto \mathbf{Fun}(I, \mathcal{C})^{\sim}],$$

is fully faithful [11; 13, §1; 14, Proposition A.7.10; 15, §A.8.2; 20]. One can explicitly identify its image; objects in the image of this embedding are often called *complete Segal spaces* or *categories internal to spaces*. Under this embedding, the subcategory $\mathbf{Spc} \subset \mathbf{Cat}_\infty$ corresponds to the constant functors $\mathbf{\Delta}^{\text{op}} \rightarrow \mathbf{Spc}$. Moreover, the localization $B : \mathbf{Cat}_\infty \rightarrow \mathbf{Spc}$ is given by geometric realization.

A.2 Notation Let \mathcal{C} be an ∞ -category with finite limits. We write

$$\mathbf{Cat}(\mathcal{C}) \subset \mathbf{Fun}(\mathbf{\Delta}^{\text{op}}, \mathcal{C})$$

for the full subcategory spanned by the *categories internal to \mathcal{C}* . See [1, Definition 13.1.1; 17, Proposition 3.2.7] for the definition.

A.3 Notation Write $\mathbf{Cat}_{<\infty} \subset \mathbf{Cat}_\infty$ for the full subcategory spanned by those ∞ -categories \mathcal{C} for which there exists an integer $n \geq 0$ such that \mathcal{C} is an n -category. Write $\mathbf{Cat}_{\infty, \pi} \subset \mathbf{Cat}_{<\infty}$ for the full subcategory spanned by those ∞ -categories with the property that there are finitely many objects up to equivalence and all mapping spaces are π -finite.

A.4 Observation The nerve $N : \mathbf{Cat}_\infty \xrightarrow{\sim} \mathbf{Cat}(\mathbf{Spc})$ restricts to equivalences

$$\mathbf{Cat}_{<\infty} \xrightarrow{\sim} \mathbf{Cat}(\mathbf{Spc}_{<\infty}) \quad \text{and} \quad \mathbf{Cat}_{\infty, \pi} \xrightarrow{\sim} \mathbf{Cat}(\mathbf{Spc}_\pi).$$

In order to describe protruncated classifying spaces via geometric realizations, it is useful to describe pro- ∞ -categories as category objects in prospaces.

A.5 Observation By [1, Proposition 13.1.12; 12, Proposition 5.3.5.11], the composite

$$\mathbf{Cat}_{<\infty} \xrightarrow[\mathbf{N}]{\sim} \mathbf{Cat}(\mathbf{Spc}_{<\infty}) \hookrightarrow \mathbf{Cat}(\mathbf{Pro}(\mathbf{Spc}_{<\infty}))$$

extends along cofiltered limits to a fully faithful right adjoint

$$N : \mathbf{Pro}(\mathbf{Cat}_{<\infty}) \hookrightarrow \mathbf{Cat}(\mathbf{Pro}(\mathbf{Spc}_{<\infty})).$$

This functor restricts to a fully faithful right adjoint

$$N : \mathbf{Pro}(\mathbf{Cat}_{\infty, \pi}) \hookrightarrow \mathbf{Cat}(\mathbf{Pro}(\mathbf{Spc}_\pi)).$$

A.6 Remark We do not know if the embedding $N : \mathbf{Cat}_\infty \hookrightarrow \mathbf{Cat}(\mathbf{Pro}(\mathbf{Spc}))$ extends along cofiltered limits to a fully faithful functor $\mathbf{Pro}(\mathbf{Cat}_\infty) \rightarrow \mathbf{Cat}(\mathbf{Pro}(\mathbf{Spc}))$.

A.7 Observation It is immediate from the definitions that the following diagram of fully faithful right adjoints commutes:

$$\begin{array}{ccccccc}
 \mathbf{Pro}(\mathbf{Spc}_\pi) & \xleftarrow[\perp]{B_\pi^\wedge} & \mathbf{Pro}(\mathbf{Cat}_{\infty,\pi}) & \xleftarrow[\perp]{N} & \mathbf{Cat}(\mathbf{Pro}(\mathbf{Spc}_\pi)) & \xleftarrow[\perp]{} & \mathbf{Fun}(\Delta^{\text{op}}, \mathbf{Pro}(\mathbf{Spc}_\pi)) \\
 \uparrow \downarrow & & \uparrow \downarrow & & \uparrow \downarrow & & \uparrow \downarrow \\
 \mathbf{Pro}(\mathbf{Spc}_{<\infty}) & \xleftarrow[\perp]{B_{<\infty}} & \mathbf{Pro}(\mathbf{Cat}_{<\infty}) & \xleftarrow[\perp]{N} & \mathbf{Cat}(\mathbf{Pro}(\mathbf{Spc}_{<\infty})) & \xleftarrow[\perp]{} & \mathbf{Fun}(\Delta^{\text{op}}, \mathbf{Pro}(\mathbf{Spc}_{<\infty}))
 \end{array}$$

$(-)^{\wedge}_\pi$ $(-)^{\wedge}_\pi \circ -$

The long composite left adjoints

$$\mathbf{Fun}(\Delta^{\text{op}}, \mathbf{Pro}(\mathbf{Spc}_{<\infty})) \rightarrow \mathbf{Pro}(\mathbf{Spc}_{<\infty}) \quad \text{and} \quad \mathbf{Fun}(\Delta^{\text{op}}, \mathbf{Pro}(\mathbf{Spc}_\pi)) \rightarrow \mathbf{Pro}(\mathbf{Spc}_\pi)$$

are simply the colimit functors. Since the diagram of left adjoints also commutes we deduce:

A.8 Corollary Let $\mathcal{C} \in \mathbf{Pro}(\mathbf{Cat}_{\infty,\pi})$ be a profinite ∞ -category. Then there are natural equivalences

$$\begin{aligned}
 B_{<\infty}(\mathcal{C}) &\simeq \text{colim}(\Delta^{\text{op}} \xrightarrow{N(\mathcal{C})} \mathbf{Pro}(\mathbf{Spc}_\pi) \hookrightarrow \mathbf{Pro}(\mathbf{Spc}_{<\infty})), \\
 B_\pi^\wedge(\mathcal{C}) &\simeq \text{colim}(\Delta^{\text{op}} \xrightarrow{N(\mathcal{C})} \mathbf{Pro}(\mathbf{Spc}_\pi)).
 \end{aligned}$$

A.9 Example Let X be a spectral ∞ -topos. Through [1, Theorem 9.3.1], Barwick–Glasman–Haine refined the ∞ -category $\mathbf{Pt}(X)$ of points of X to a profinite ∞ -category

$$\widehat{\Pi}_{(\infty,1)}(X) \in \mathbf{Pro}(\mathbf{Cat}_{\infty,\pi})$$

called the *stratified shape* of X . In [1, Theorem 10.2.3] they show that there is a natural equivalence

$$\Pi_{<\infty}(X) \xrightarrow{\sim} B_{<\infty}(\widehat{\Pi}_{(\infty,1)}(X)).$$

That is, the protruncated shape of X can be recovered as the protruncated classifying space of the stratified shape $\widehat{\Pi}_{(\infty,1)}(X)$. Hence Corollary A.8 shows that the protruncated shape $\Pi_{<\infty}(X)$ admits a natural $\mathbf{Pro}(\mathbf{Spc}_\pi)$ -resolution in the sense of Definition 2.1.

A.10 Remark Using the proétale topology [2], one can show that the protruncated étale homotopy type of a qcqs scheme admits a $\mathbf{Pro}(\mathbf{Set}_{\text{fin}})$ -resolution.

To see this, first note that given a qcqs scheme X , the pullback functor $\nu^* : X_{\text{ét}} \rightarrow X_{\text{proét}}$ from the étale ∞ -topos of X to the proétale ∞ -topos of X is fully faithful when restricted to truncated objects. (This observation generalizes [2, Lemma 5.1.2 and Corollary 5.1.6].) As a result, the protruncated shapes of $X_{\text{ét}}$ and $X_{\text{proét}}$ are equivalent. Hence the protruncated étale homotopy type is a hypercomplete proétale cosheaf.

Recall that the w-contractible affine schemes form a basis for the proétale topology; in particular, every qcqs scheme admits a proétale hypercover by w-contractible affine schemes. Moreover, for each

w-contractible affine scheme U , the prospace $\Pi_{<\infty}^{\text{ét}}(U)$ is naturally identified with the profinite set $\pi_0(U)$ of connected components of U . Hence the protruncated étale homotopy type

$$\Pi_{<\infty}^{\text{ét}} : \mathbf{Sch}^{\text{qcqs}} \rightarrow \text{Pro}(\mathbf{Spc}_{<\infty})$$

is the unique hypercomplete proétale cosheaf whose restriction to w-contractible affines is given by $U \mapsto \pi_0(U)$. Given a qcqs scheme X , choose a proétale hypercover U_{\bullet} of X by w-contractible affines. Then the simplicial object $\pi_0(U_{\bullet})$ is a $\text{Pro}(\mathbf{Set}_{\text{fin}})$ -resolution of $\Pi_{<\infty}^{\text{ét}}(X)$.

Acknowledgements

We thank Luciana Basualdo Bonatto, Chang-Yeon Chough, Tim Holzschuh, Marcy Robertson, and Sebastian Wolf for many enlightening discussions around the contents of this note. We gratefully acknowledge support from the NSF Mathematical Sciences Postdoctoral Research Fellowship under Grant #DMS-2102957 and a grant from the Simons Foundation (816048, LC).

References

- [1] **C Barwick, S Glasman, P Haine**, *Exodromy*, preprint (2020) [arXiv 1807.03281](#)
- [2] **B Bhatt, P Scholze**, *The pro-étale topology for schemes*, from “De la géométrie algébrique aux formes automorphes, I” (J-B Bost, P Boyer, A Genestier, L Lafforgue, S Lysenko, S Morel, B C Ngo, editors), Astérisque 369, Soc. Math. France, Paris (2015) 99–201 [MR](#)
- [3] **T Blom, I Moerdijk**, *Profinite ∞ -operads*, Adv. Math. 408 (2022) art. id. 108601 [MR](#)
- [4] **T Blom, I Moerdijk**, *Profinite completions of topological operads*, preprint (2024) [arXiv 2312.12567](#)
- [5] **P Boavida de Brito, G Horel, M Robertson**, *Operads of genus zero curves and the Grothendieck–Teichmüller group*, Geom. Topol. 23:1 (2019) 299–346 [MR](#)
- [6] **B Calmès, E Dotto, Y Harpaz, F Hebestreit, M Land, K Moi, D Nardin, T Nikolaus, W Steimle**, *Hermitian K-theory for stable ∞ -categories, I: Foundations*, Selecta Math. (N.S.) 29:1 (2023) art. id. 10 [MR](#)
- [7] **C-Y Chough**, *Proper base change for étale sheaves of spaces*, J. Eur. Math. Soc. 24:11 (2022) 3679–3702 [MR](#)
- [8] **P J Haine, T Holzschuh, S Wolf**, *The fundamental fiber sequence in étale homotopy theory*, Int. Math. Res. Not. 2024:1 (2024) 175–196 [MR](#)
- [9] **P J Haine, T Holzschuh, S Wolf**, *Nonabelian basechange theorems and étale homotopy theory*, J. Topol. 17:4 (2024) art. id. e70009 [MR](#)
- [10] **L Hesselholt, P Pstragowski**, *Dirac geometry, I: Commutative algebra* (2023) [arXiv 2207.09256](#)
- [11] **A Joyal, M Tierney**, *Quasi-categories vs Segal spaces*, from “Categories in algebra, geometry and mathematical physics” (A Davydov, M Batanin, M Johnson, S Lack, A Neeman, editors), Contemp. Math. 431, Amer. Math. Soc., Providence, RI (2007) 277–326 [MR](#)
- [12] **J Lurie**, *Higher topos theory*, Annals of Mathematics Studies 170, Princeton Univ. Press (2009) [MR](#)
- [13] **J Lurie**, *($\infty, 2$)-categories and the Goodwillie calculus, I*, preprint (2009) Available at <http://www.math.ias.edu/~lurie/papers/GoodwillieI.pdf>
- [14] **J Lurie**, *Higher algebra*, preprint (2017) Available at <http://www.math.ias.edu/~lurie/papers/HA.pdf>
- [15] **J Lurie**, *Spectral algebraic geometry*, preprint (2018) Available at <http://www.math.ias.edu/~lurie/papers/SAG-rootfile.pdf>
- [16] **J Lurie**, *Kerodon*, online resource (2024) Available at <https://kerodon.net>
- [17] **L Martini**, *Yoneda’s lemma for internal higher categories* (2022) [arXiv 2103.17141](#)
- [18] **M C McCord**, *Singular homology groups and homotopy groups of finite topological spaces*, Duke Math. J. 33 (1966) 465–474 [MR](#)

- [19] **T Oda**, *Etale homotopy type of the moduli spaces of algebraic curves*, from “Geometric Galois actions, 1” (L Schneps, P Lochak, editors), London Math. Soc. Lecture Note Ser. 242, Cambridge Univ. Press (1997) 85–95 [MR](#)
- [20] **C Rezk**, *A model for the homotopy theory of homotopy theory*, Trans. Amer. Math. Soc. 353:3 (2001) 973–1007 [MR](#)
- [21] **M Volpe**, *Verdier duality on conically smooth stratified spaces*, Algebr. Geom. Topol. 25:2 (2025) 919–950 [MR](#)

PETER J. HAINE phaine@usc.edu

Department of Mathematics, University of Southern California, Los Angeles, CA, United States

Received: July 3, 2024 Revised: January 27, 2025

ALGEBRAIC & GEOMETRIC TOPOLOGY

msp.org/agt

EDITORS

PRINCIPAL ACADEMIC EDITORS

John Etnyre
etnyre@math.gatech.edu
Georgia Institute of Technology

Kathryn Hess
kathryn.hess@epfl.ch
École Polytechnique Fédérale de Lausanne

BOARD OF EDITORS

Julie Bergner	University of Virginia jeb2md@eservices.virginia.edu	Thomas Koberda	University of Virginia thomas.koberda@virginia.edu
Steven Boyer	Université du Québec à Montréal cohf@math.rochester.edu	Markus Land	LMU München markus.land@math.lmu.de
Tara E Brendle	University of Glasgow tara.brendle@glasgow.ac.uk	Christine Lescop	Université Joseph Fourier lescop@ujf-grenoble.fr
Indira Chatterji	CNRS & Univ. Côte d'Azur (Nice) indira.chatterji@math.cnrs.fr	Norihiko Minami	Yamato University minami.norihiko@yamato-u.ac.jp
Octav Cornea	Université de Montreal cornea@dms.umontreal.ca	Andrés Navas	Universidad de Santiago de Chile andres.navas@usach.cl
Alexander Dranishnikov	University of Florida dranish@math.ufl.edu	Jessica S Purcell	Monash University jessica.purcell@monash.edu
Tobias Ekholm	Uppsala University, Sweden tobias.ekholm@math.uu.se	Birgit Richter	Universität Hamburg birgit.richter@uni-hamburg.de
Mario Eudave-Muñoz	Univ. Nacional Autónoma de México mario@matem.unam.mx	Jérôme Scherer	École Polytech. Féd. de Lausanne jerome.scherer@epfl.ch
David Futер	Temple University dfuter@temple.edu	Vesna Stojanoska	Univ. of Illinois at Urbana-Champaign vesna@illinois.edu
John Greenlees	University of Warwick john.greenlees@warwick.ac.uk	Zoltán Szabó	Princeton University szabo@math.princeton.edu
Matthew Hedden	Michigan State University mhedden@math.msu.edu	Maggy Tomova	University of Iowa maggy-tomova@uiowa.edu
Kristen Hendricks	Rutgers University kristen.hendricks@rutgers.edu	Daniel T Wise	McGill University, Canada daniel.wise@mcgill.ca
Hans-Werner Henn	Université Louis Pasteur henn@math.u-strasbg.fr	Lior Yanovski	Hebrew University of Jerusalem lior.yanovski@gmail.com
Daniel Isaksen	Wayne State University isaksen@math.wayne.edu		

See inside back cover or msp.org/agt for submission instructions.

The subscription price for 2026 is US \$795/year for the electronic version, and \$1170/year (+\$80, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Algebraic & Geometric Topology is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [Current Mathematical Publications](#) and the [Science Citation Index](#).

Algebraic & Geometric Topology (ISSN 1472-2747 printed, 1472-2739 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004.

AGT peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing
<https://msp.org/>

© 2026 Mathematical Sciences Publishers

ALGEBRAIC & GEOMETRIC TOPOLOGY

Volume 26 Issue 1 (pages 1–410) 2026

The Borel cohomology of free iterated loop spaces	1
ISHAN LEVY and JUSTIN WU	
The K-theory cochains of H-spaces and height-1 chromatic homotopy theory	29
SVEN VAN NIGTEVECHT	
Cup-one algebras and 1-minimal models	65
RICHARD D. PORTER and ALEXANDER I. SUCIU	
Cubical approximation for directed topology, II	135
SANJEEVI KRISHNAN	
Flat semigroups and weighted homogeneous surface singularities	201
ZSOLT BAJA and TAMÁS LÁSZLÓ	
Interleaving distances, monoidal actions and 2-categories	227
PATRICK K. MCFADDIN and TOM NEEDHAM	
Suspension splittings of 5-dimensional Poincaré duality complexes and their applications	283
STEVEN AMELOTTE, TYRONE CUTLER and TSELEUNG SO	
Bernoulli shifts on additive categories and algebraic K -theory of wreath products	321
JULIAN KRANZ and SHINTARO NISHIKAWA	
Kontsevich's characteristic classes as topological invariants of configuration space bundles	349
XUJIA CHEN	
Profinite completions of products	397
PETER J. HAINE	