

AG
T

*Algebraic & Geometric
Topology*

Volume 26 (2026)

Stellar subdivisions, wedges and Buchstaber numbers

SUYOUNG CHOI AND HYEONTAE JANG

Stellar subdivisions, wedges and Buchstaber numbers

SUYOUNG CHOI AND HYEONTAE JANG

The Buchstaber number of a simplicial complex K is a significant invariant in toric topology. In particular, when K is a (polytopal) PL sphere, the maximal Buchstaber number is closely connected to several important objects, such as toric manifolds, quasitoric manifolds, and topological toric manifolds. A PL sphere is called a *seed* if it cannot be obtained from another PL sphere through a wedge operation. The *toric colorable seed inequality*, established by Choi and Park in 2017, bounds the maximal number of vertices of a seed with a maximal Buchstaber number. This inequality plays a key role in characterizing PL spheres that achieve maximal Buchstaber numbers.

We prove that the inequality is tight. Specifically, we show how to construct larger seeds from existing ones using stellar subdivisions and wedges, while preserving both the maximality of Buchstaber numbers and polytopality.

1 Introduction

Let K be an $(n-1)$ -dimensional simplicial complex on $[m] = \{1, 2, \dots, m\}$ and (X, Y) a topological pair. The *polyhedral product* $(\underline{X}, \underline{Y})^K$ is a subspace of X^m with respect to K :

$$(\underline{X}, \underline{Y})^K := \bigcup_{\sigma \in K} \{(x_1, \dots, x_m) \in X^m \mid x_i \in Y \text{ when } i \notin \sigma\}.$$

One of our main considerations is the *moment-angle complex*

$$\mathcal{Z}_K := (\underline{D}^2, \underline{S}^1)^K$$

of K , where $D^2 = \{z \in \mathbb{C} \mid |z| \leq 1\}$ is the 2-dimensional disk, and $S^1 = \partial D^2$ is its boundary circle. The canonical action of S^1 on D^2 induces the action of the m -dimensional torus $T^m = S^1 \times \dots \times S^1$ on \mathcal{Z}_K . The *Buchstaber number* $s(K)$ of K is the maximum number r such that there is an r -dimensional subtorus H of T^m freely acting on \mathcal{Z}_K .

It is known that the following inequality holds by Buchstaber and Panov [2] and Erokhovets [8]:

$$1 \leq s(K) \leq m - n.$$

The case when K is a PL sphere with $s(K) = m - n$, particularly when it is a polytopal PL sphere, is of special interest because many significant toric spaces, including complete nonsingular toric varieties, simply called *toric manifolds*, and their topological generalizations such as *quasitoric manifolds*, defined by Davis and Januszkiewicz [6], and *topological toric manifolds*, defined by Ishida, Fukukawa and Masuda [14], are formed in \mathcal{Z}_K/H , where K is a (polytopal) PL sphere and H is a free torus action of

MSC2020: primary 57S12; secondary 14M25.

© 2026 MSP (Mathematical Sciences Publishers). Distributed under the Creative Commons Attribution License 4.0 (CC BY). Open Access made possible by subscribing institutions via [Subscribe to Open](#).

rank $m - n$ on \mathcal{Z}_K . Hence, such PL spheres are said to be *toric colorable* and the characterization of toric colorable PL spheres is of considerable importance.

The breakthrough in addressing this problem is the approach by Choi and Park. The *wedge* is an operation on simplicial complexes that preserves the PL sphereness and polytopality, and a PL sphere K is called a *seed* if it cannot be obtained by a wedge operation from any other PL sphere. Ewald [9] and Bahri, Bendersky, Cohen and Gitler [1] showed an advantageous property of the wedge operation in toric topology that it preserves the maximality of the Buchstaber number. On the other hand, the wedge operation does not change the difference between the number of vertices and the dimension. Therefore, since every toric colorable PL sphere can be obtained from toric colorable seeds by a sequence of wedges, it is enough to consider toric colorable seeds. In [5], Choi and Park proved that for a fixed $m - n$, the number of $(n-1)$ -dimensional toric colorable seeds K with m vertices is finite. More precisely, K must satisfy the inequality, which we refer to as the *toric colorable seed inequality*,

$$(1-1) \quad m \leq 2^{m-n} - 1.$$

This inequality transforms the problem of finding all PL spheres with a maximal Buchstaber number into a finite problem for each fixed $p := m - n$. An interesting question is whether this inequality is tight. If the upper bound of (1-1) can be reduced, it would further transform the problem into a smaller finite problem. It is known that the inequality is tight for $p = 3$ and 4. Denote by $C^4(7)$ the cyclic polytope of dimension 4 with seven vertices. The boundary complex of $C^4(7)$ is a seed, and, by Erokhovets [7], its Buchstaber number is maximal. Hence, the inequality is tight for $p = 3$. According to Choi, Jang and Vallée [3], there are four PL spheres that confirms the tightness for $p = 4$ as well. However, in general, it is not known whether the inequality is tight. Cyclic and stacked polytopes are often considered to provide an example due to their simplicity in construction, but they do not provide extremal examples for this inequality. When n is significantly larger than p , the former are not toric colorable by Hasui [13], and the latter are not seeds.

In this paper, we construct a new polytopal toric colorable seed from an old one using two operations, the stellar subdivision and the wedge on PL spheres, that preserve the maximality of Buchstaber numbers and polytopality. [Theorem 3.1](#) is our main theorem.

Theorem 3.1 *Let K be a seed on $[m]$ and $J = (j_1, j_2, \dots, j_m)$ a positive integer tuple such that $j_v \leq 2$ for all $v \in [m]$. For any assembled face σ of $K(J)$, a new seed is obtained in two ways depending on the cardinality of σ .*

- (1) *If $\sigma = \{v\}$ for a vertex v of K , then $Ss_{\{v,v\}}(K(J))$ is a suspended seed.*
- (2) *If $|\sigma| > 1$ and K is nonsuspended, then $Ss_{\sigma}(K(J))$ is a nonsuspended seed.*

By using this result, we can show that (1-1) is tight for all $p \geq 3$, even for the class of polytopal PL spheres. In other words, one can precisely determine the conditions on m and n for an $(n-1)$ -dimensional seed K with m vertices to support quasitoric manifolds or topological toric manifolds.

Corollary 3.2 Assume that $p \geq 3$. For any $m \leq 2^p - 1$ and $n \geq 2$ with $p = m - n$, there exists a polytopal $(n-1)$ -dimensional seed K with m vertices such that $s(K) = p$.

We end the introduction with a discussion on the characterization of PL spheres that support toric manifolds. To characterize them, it is sufficient to consider the class of nonsuspended seeds that support a toric manifold. Let t_p denote the sharp upper bound of m for $(n-1)$ -dimensional nonsuspended seeds supporting a toric manifold with $m - n = p$. However, in this class, (1-1) is not tight. For instance, when $p = 3$, $t_3 = 5$, which is less than $2^3 - 1 = 7$, as noted by Gretenkort, Kleinschmidt and Sturmfels [12].

Since a stellar subdivision and a wedge operation are stable within this class, our main theorem provides the following meaningful lower bound of t_p :

$$3 \cdot 2^{p-2} - 1 \leq t_p \leq 2^p - 1.$$

Problem 1.1 Let K be the underlying complex of a nonsingular complete fan with m rays in \mathbb{R}^n . Assume that K is a seed. For a fixed $p = m - n$, find the sharp upper bound t_p of m .

2 Basic operations on simplicial complexes

Let K be a simplicial complex on $[m] = \{1, 2, \dots, m\}$. A nonempty element of K is called a *face* of K . Any singleton in K is often called a *vertex*, and a maximal face of K with respect to inclusion is called a *facet*. We often identify a vertex $\{v\}$ of K with its unique element v as well as with the simplicial complex consisting of the unique vertex v . The *dimension* of a face σ is $|\sigma| - 1$, and K is said to be *pure* if the dimensions of all facets are the same. The *dimension* of pure simplicial complex K is the dimension of its facet. Throughout this paper, every simplicial complex is assumed to be pure.

For any face σ of K , the *boundary complex* of σ is the simplicial complex $\partial\sigma = \{\tau \in K \mid \tau \subsetneq \sigma\}$, the *star* of σ in K is the simplicial complex

$$\text{St}_K(\sigma) = \{\tau \in K \mid \tau \cup \sigma \in K\},$$

and the *link* of σ in K is the simplicial complex

$$\text{Lk}_K(\sigma) = \{\tau \in K \mid \tau \cup \sigma \in K, \tau \cap \sigma = \emptyset\}.$$

For another simplicial complex L whose vertex set is disjoint to that of K , the *join* of K and L is the simplicial complex

$$K * L = \{\sigma \cup \tau \mid \sigma \in K, \tau \in L\}.$$

The *stellar subdivision* of K at $\sigma \in K$ is the simplicial complex

$$(2-1) \quad \text{Ss}_\sigma(K) = K \setminus \text{St}_K(\sigma) \cup v_\sigma * \partial\sigma * \text{Lk}_K(\sigma),$$

where v_σ is a new vertex. See Figure 1(c).




Figure 1: Operations on simplicial complexes.

Let v be a vertex of K , and I the 1-dimensional simplicial complex with two vertices v and v' . The wedge of K at v is the simplicial complex

$$(2-2) \quad \text{Wed}_v(K) = I * \text{Lk}_K(v) \cup \partial I * \{\sigma \in K \mid v \notin \sigma\}.$$

The face $\{v, v'\}$ of $\text{Wed}_v(K)$ is called a *wedged edge* of $\text{Wed}_v(K)$. Note that $\text{Lk}_{\text{Wed}_v(K)}(v') = K$, and $\text{Lk}_{\text{Wed}_v(K)}(v)$ is isomorphic to K by identifying v' and v . In this sense, v' can be regarded as a copy of v . See Figure 1(b).

There is another construction of a wedge using minimal nonfaces, known as the *J-construction*, established by Bahri, Bendersky, Cohen and Gitler [1]. The set of minimal nonfaces of $\text{Wed}_v(K)$ is obtained by duplicating v as v and v' in each minimal nonface of K . In this construction, one can observe that wedge operations are associative and commutative up to suitable vertex identification. Then we can use the notation $K(J)$ for a positive integer tuple $J = (j_1, j_2, \dots, j_m)$ to refer the simplicial complex obtained by a sequence of wedge operations with j_v copies $v = v^{(0)}, v' = v^{(1)} \dots, v^{(j_v-1)}$ of each vertex v . Using this notation, $\text{Wed}_v(K)$ can be expressed as $K(1, \dots, 1, 2, 1, \dots, 1)$, where 2 appears in the v -th position. To clarify, we note that K is a subcomplex of $K(J)$, that is, any face of K is a face of $K(J)$. Moreover, choosing $0 \leq s_v \leq j_v - 1$ for each $v \in [m]$, then for a face σ of $K(J)$, the set $\sigma^{(s_1, s_2, \dots, s_m)}$ obtained by replacing $v \in \sigma$ by $v^{(s_v)}$ is also a face of $K(J)$. In such a way, we denote the set $\{1^{(s_1)}, 2^{(s_2)}, \dots, m^{(s_m)}\}$ by $[m]^{(s_1, s_2, \dots, s_m)}$. The *assembled face* of $K(J)$ by (s_1, s_2, \dots, s_m) denotes the set $[m]^{(s_1, s_2, \dots, s_m)} \setminus \{v \in [m] \mid j_v = 1\}$. The following lemma shows that any assembled face is literally a face of $K(J)$.

Lemma 2.1 *Let K be a simplicial complex on $[m]$, and $J = (j_1, j_2, \dots, j_m)$ a positive integer tuple. Choose $0 \leq s_v \leq j_v - 1$ for each $v \in [m]$. Then $[m]^{(s_1, s_2, \dots, s_m)} \setminus \{v \in [m] \mid j_v = 1\}$ is a face of $K(J)$.*

Proof If we prove the case $s_1 = s_2 = \dots = s_m = 0$, then the statement holds by the above argument. In this case, it is enough to observe that for a vertex v and a face σ of K , the set $\sigma \cup \{v\}$ is a face of $\text{Wed}_v(K)$ by (2-2). □

A simplicial complex K is called a *PL sphere* if there exists a subdivision of K such that it can be expressed as a subdivision of the boundary complex of a simplex. A PL sphere is called a *seed* if it is not a wedge of any other PL sphere. Then any PL sphere K is written as $K = K'(J)$ for some seed K' and some positive integer tuple J .

Proposition 2.2 [3; 4; 11] *If K is a PL sphere, then both $Ss_\sigma(K)$ and $K(J)$ are also PL spheres for any $\sigma \in K$ and any positive integer tuple J . In particular, if K is polytopal, then so are $Ss_\sigma(K)$ and $K(J)$.*

For an $(n-1)$ -dimensional PL sphere with m vertices, the *Picard number* of the complex K is defined as $\text{Pic}(K) := m - n$. Note that the dimension of $Ss_\sigma(K)$ is $n - 1$, and the dimension of $\text{Wed}_v(K)$ of K is n , while both complexes have $m + 1$ vertexes. Therefore, the stellar subdivision increases the Picard number by 1, whereas the wedge operation preserves the Picard number:

$$\text{Pic}(K) = \text{Pic}(K(J)) = \text{Pic}(Ss_\sigma(K)) - 1.$$

A PL sphere is said to be *toric colorable* if its Buchstaber number is equal to its Picard number.

Proposition 2.3 [9; 10] *Let K be a PL sphere. If K is toric colorable, then both $Ss_\sigma(K)$ and $K(J)$ are also toric colorable for any $\sigma \in K$ and any positive integer tuple J :*

$$\text{Pic}(K) = s(K) = s(K(J)) = s(Ss_\sigma(K)) - 1.$$

The rest of this section is devoted to the introduction to the suspension operation. If the vertex v was a ghost vertex, then (2-2) becomes $\partial I * K$. This is called the *suspension* of K . The pair $\{v, v'\}$ of vertices of $\partial I * K$ is called a *suspended pair* of $\partial I * K$, and the vertices v and v' are called *suspended vertexes*.

Proposition 2.4 [5] *Let K be a PL sphere. For vertices v and w of K , every facet of K contains v or w if and only if K is the wedge with a wedged edge $\{v, w\}$ or the suspension with a suspended pair $\{v, w\}$.*

Lemma 2.5 *Let K be a seed, J a positive integer tuple, and v, w two distinct vertices of K . If every facet of $K(J)$ contains $v^{(s_v)}$ or $w^{(s_w)}$ for some $0 \leq s_v \leq j_v - 1$ and $0 \leq s_w \leq j_w - 1$, then $\{v, w\}$ is a suspended pair of K .*

Proof Without loss of generality, we may assume that $s_v = s_w = 0$. For any vertex $\neq v, w$ of $K(J)$, every facet of the link of the vertex still contains $v = v^{(0)}$ or $w = w^{(0)}$. The repeated link operations with vertices $u^{(s_u)}$ for all $u \in [m]$ and $s_u \geq 1$ gives K , and then every facet of K contains v or w . By Proposition 2.4, $\{v, w\}$ is a suspended pair of K since K has no wedged edge. \square

The set of minimal nonfaces of the join of two simplicial complexes is the union of the sets of minimal nonfaces of those. Then one can easily observe that join and wedge operations are associative and commutative. Hence we can write a PL sphere K as

$$(2-3) \quad K = \partial I_1(J_1) * \partial I_2(J_2) * \cdots * \partial I_\ell(J_\ell) * L(J_{\ell+1}),$$

where each I_k is a 1-simplex for $1 \leq k \leq \ell$, and L is a seed without a suspended pair. A PL sphere is said to be *nonsuspended* if it is not the suspension of any other PL sphere. The interpretation (2-3) of J -constructions involving suspensions explains the following.

Proposition 2.6 *Let K be a PL sphere.*

- K is a suspension if and only if so is its wedge at any vertex v of K not contained in a suspended pair.
- K is a seed if and only if so is the suspension of K .

3 Main construction

Let K be a seed on $[m]$ and $J = (j_1, \dots, j_m)$ a positive integer tuple. Recall that K is a subcomplex of $K(J)$. Then the wedged edge $\{v, v'\}$ is a face of $\text{Wed}_v(K)$ for each vertex v of K , and each assembled face σ of $K(J)$ is a face of $K(J)$ by Lemma 2.1. Therefore, both $\text{Ss}_{\{v,v'\}}(\text{Wed}_v(K))$ and $\text{Ss}_\sigma(K(J))$ are well defined.

Theorem 3.1 *Let K be a seed on $[m]$ and $J = (j_1, j_2, \dots, j_m)$ a positive integer tuple such that $j_v \leq 2$ for all $v \in [m]$. For any assembled face σ of $K(J)$, a new seed is obtained in two ways depending on the cardinality of σ .*

- (1) *If $\sigma = \{v\}$ for a vertex v of K , then $\text{Ss}_{\{v,v'\}}(K(J))$ is a suspended seed.*
- (2) *If $|\sigma| > 1$ and K is nonsuspended, then $\text{Ss}_\sigma(K(J))$ is a nonsuspended seed.*

Proof Without loss of generality, we may assume $\sigma = \{v \mid j_v = 2\}$. First, let $\sigma = \{v\}$. Since $\text{Ss}_{\{v,v'\}}(\text{Wed}_v(K))$ is isomorphic to the suspension of K , it is a seed by Proposition 2.6.

Next, let $|\sigma| > 1$ and let K be nonsuspended. Suppose that $\text{Ss}_\sigma(K(J))$ has two vertices x and y such that every facet of $\text{Ss}_\sigma(K(J))$ contains x or y . By Proposition 2.4, $\{x, y\}$ is a wedged edge or a suspended pair of $\text{Ss}_\sigma(K(J))$.

From the construction (2-1) of stellar subdivisions, one can observe that the facet set \mathcal{F} of $\text{Ss}_\sigma(K(J))$ is partitioned into two subsets

$$\begin{aligned} \mathcal{F}_1 &= \mathcal{F} \cap K(J) \setminus \text{St}_{K(J)}(\sigma), \\ \mathcal{F}_2 &= \mathcal{F} \cap v_\sigma * \partial\sigma * \text{Lk}_{K(J)}(\sigma). \end{aligned}$$

The subset \mathcal{F}_1 consists of the facets of $K(J)$ not containing σ , and the subset \mathcal{F}_2 is involved in the facets of $K(J)$ that contain σ . Let \mathcal{F}_σ be the set consisting of facets of $K(J)$ containing σ . Then the facet set of $K(J)$ is $\mathcal{F}_1 \cup \mathcal{F}_\sigma$. Since every facet in \mathcal{F}_2 contains x or y , there are four possibilities:

- (1) $x = v_\sigma$,
- (2) $x, y \in \sigma$,
- (3) $x \in \sigma$ and $y \notin \sigma$, and
- (4) $x, y \notin \sigma$.

We want to prove that all of these four cases lead to contradictions, so there is no such pairs of vertices x and y .

Case 1 ($x = v_\sigma$): There is no facet in \mathcal{F}_1 containing x , so every facet in \mathcal{F}_1 contains y . For any $z \in \sigma$ with $z \neq y$, every facet of $K(J)$ contains y or z since the facets in \mathcal{F}_1 contain y , and the facets in \mathcal{F}_σ contain $\sigma \ni z$. If $y \in \sigma$, then every facet of $K(J)$ contains y , but it is impossible that every facet of a PL sphere shares one vertex. Thus, y is not contained in σ .

Suppose that $y \neq z'$ for any $z \in \sigma$. Then $\text{Lk}_{K(J)}(\{z' \mid z \in \sigma\}) = K$ contains y or z for any $z \in \sigma$ as we discussed in the proof of Lemma 2.5. By Proposition 2.6, K is a suspension or a wedge, but we assumed

that K is a nonsuspended seed. Thus, there exists $y_0 \in \sigma$ such that $y = y'_0$. By Lemma 2.5, $\{y_0, z\}$ is a suspended pair of K for any other vertices $z \in \sigma$. The assumption that $|\sigma| > 1$ ensures the existence of such z , and this contradicts the assumption that K is nonsuspended.

Case 2 ($x, y \in \sigma$): The face σ of $K(J)$ contains $\{x, y\}$. Then $\{x, y\}$ is a wedged edge of $K(J)$ since any facet in \mathcal{F}_1 contains x or y , and any facet in \mathcal{F}_σ contains $\{x, y\}$. Note that σ is also a face of K . Then x and y are two distinct vertices of K , and then $\{x, y\}$ is a suspended pair of K by Lemma 2.5, but K is assumed to have no suspended pair.

Case 3 ($x \in \sigma$ and $y \notin \sigma$): Note that $\sigma \setminus \{x\}$ is a facet of the boundary complex $\partial\sigma$. Then for any facet τ of $\text{Lk}_{K(J)}(\sigma)$, the set $\{v_\sigma\} \cup \sigma \setminus \{x\} \cup \tau$ is a facet in \mathcal{F}_2 not containing x . Hence it must contain y , but there is a facet τ of $\text{Lk}_{K(J)}(\sigma)$ not containing y since it is a PL sphere, which contradicts the assumption that every facet of $\text{Ss}_\sigma(K(J))$ contains x or y .

Case 4 ($x, y \notin \sigma$): Every facet in $\text{Lk}_{K(J)}(\sigma)$ contains x or y . Since it is a PL sphere, both x and y have to appear in $\text{Lk}_{K(J)}(\sigma)$ as its vertices. By the assumption that $j_v \leq 2$ for any vertex v of K and the definition of σ , $\text{Lk}_{K(J)}(\sigma)$ is isomorphic to K . By Proposition 2.4, $\text{Lk}_{K(J)}(\sigma)$ is a wedge or a suspension, which contradicts the assumption that $\text{Lk}_{K(J)}(\sigma) \cong K$ is a nonsuspended seed. \square

Corollary 3.2 Assume that $p \geq 3$. For any $m \leq 2^p - 1$ and $n \geq 2$ with $p = m - n$, there exists a polytopal $(n-1)$ -dimensional seed K with m vertices such that $s(K) = p$.

Proof We prove by induction on $p \geq 3$ that there exists a polytopal toric colorable seed with m vertices for any $p + 2 \leq m \leq 2^p - 1$, and the one with $m = 2^p - 1$ is nonsuspended. Note that $p + 2 \leq m \leq 2^p - 1$ is equivalent to $2 \leq n \leq 2^p - p - 1$.

For $p = 3$, the face structures of a pentagon for $n = 2$, a cross-polytope for $n = 3$, and $C^4(7)$ for $n = 4$ are toric colorable seeds. These are all polytopal by Mani [15], and the one with $n = 4$ is nonsuspended.

Assume that the statement holds for some $p \geq 3$. Let K_n be an $(n-1)$ -dimensional polytopal toric colorable seed of Picard number p for $2 \leq n \leq 2^p - p - 1$. Now, it is enough to show the existence of a polytopal toric colorable seed of Picard number $p + 1$ for $2 \leq n \leq 2^{p+1} - p - 2$.

For $n = 2$, note that the face complex of an $(n+p+1)$ -gon is a polytopal toric colorable seed of Picard number $p + 1$.

For $3 \leq n \leq 2^p - p$, by (1) of Theorem 3.1, $\text{Ss}_{\{v,v'\}}(\text{Wed}_v(K_{n-1}))$ for any vertex $v \in K_{n-1}$ is an $(n-1)$ -dimensional seed of Picard number $p + 1$ that is polytopal by Proposition 2.2 and toric colorable by Proposition 2.3.

For $2^p - p + 1 \leq n \leq 2^{p+1} - p - 2$, we consider K_{2^p-p-1} that is nonsuspended by the induction hypothesis. Note that K_{2^p-p-1} has $2^p - 1$ vertices. Let $\sigma = \{1, 2, 3, \dots, k\}$ for $2 \leq k \leq 2^p - 1$, and J the positive integer tuple whose first k components are 2 and the other components are 1. Then the Picard number and the dimension of $L_k := \text{Ss}_\sigma(K_{2^p-p-1}(J))$ are $p + 1$ and $2^p - p + k - 2$, respectively. For each $2 \leq k \leq 2^p - 1$, L_k is indeed an example what we want for $n = 2^p - p + k - 1$ since L_k is a nonsuspended seed by (2) of Theorem 3.1, polytopal by Proposition 2.2, and toric colorable by Proposition 2.3. \square

Remark 3.3 If we restrict the assumption of the previous corollary to $p \geq 4$, then all the resulting polytopal seeds can be constructed to be nonsuspended as follows. First, one can easily observe that there exists at least one $(n-1)$ -dimensional polytopal toric colorable nonsuspended seed for each $n = 2, 3$, and for the dimensions, all PL spheres are toric colorable. Then for a Picard number p , if there exists an $(n-1)$ -dimensional polytopal toric colorable nonsuspended seed with m vertices for each $n \geq 2$ and $m \leq 2^p - 1$ with $p = m - n$, then we can replace the suspension process in the proof of [Corollary 3.2](#) with (2) of [Theorem 3.1](#) by constructing an $(n+1)$ -dimensional nonsuspended seed of Picard number $p+1$. The obstruction is that for $p = 3$ and $n = 3$, the only toric colorable seed is the boundary of cross-polytope, which is suspended. However, we can construct an $(n+1)$ -dimensional nonsuspended seed of Picard number 4 from the boundary of a pentagon by applying (2) of [Theorem 3.1](#) with $J = (2, 2, 2, 1, 1)$.

Remark 3.4 To achieve the optimal case where $m = 2^{m-n} - 1$, the operation $K(J)$ is applied, where all components of J are equal to 2. This operation is commonly referred to as the *doubling operation*. For more details, see work by Ustinovskii [\[17\]](#) or Park [\[16\]](#).

Acknowledgement

This research was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) (RS-2021-NR060141 and RS-2025-00521982).

References

- [1] **A Bahri, M Bendersky, F R Cohen, S Gitler**, *Operations on polyhedral products and a new topological construction of infinite families of toric manifolds*, Homology Homotopy Appl. 17:2 (2015) 137–160 [MR](#)
- [2] **V M Buchstaber, T E Panov**, *Toric topology*, Mathematical Surveys and Monographs 204, Amer. Math. Soc., Providence, RI (2015) [MR](#)
- [3] **S Choi, H Jang, M Vallée**, *The characterization of $(n-1)$ -spheres with $n+4$ vertices having maximal Buchstaber number*, J. Reine Angew. Math. 811 (2024) 267–292 [MR](#)
- [4] **S Choi, H Park**, *Wedge operations and torus symmetries*, Tohoku Math. J. (2) 68:1 (2016) 91–138 [MR](#)
- [5] **S Choi, H Park**, *Wedge operations and torus symmetries, II*, Canad. J. Math. 69:4 (2017) 767–789 [MR](#)
- [6] **M W Davis, T Januszkiewicz**, *Convex polytopes, Coxeter orbifolds and torus actions*, Duke Math. J. 62:2 (1991) 417–451 [MR](#)
- [7] **N Y Erokhovets**, *Moment-angle manifolds of simple n -dimensional polytopes with $n+3$ facets*, Uspekhi Mat. Nauk 66:5(401) (2011) 187–188 [MR](#)
- [8] **N Y Erokhovets**, *Buchstaber invariant theory of simplicial complexes and convex polytopes*, Proc. Steklov Inst. Math. 286:1 (2014) 128–187 [MR](#)
- [9] **G Ewald**, *Spherical complexes and nonprojective toric varieties*, Discrete Comput. Geom. 1:2 (1986) 115–122 [MR](#)
- [10] **G Ewald**, *Combinatorial convexity and algebraic geometry*, Graduate Texts in Mathematics 168, Springer (1996) [MR](#)
- [11] **G Ewald, G C Shephard**, *Stellar subdivisions of boundary complexes of convex polytopes*, Math. Ann. 210 (1974) 7–16 [MR](#)
- [12] **J Gretenkort, P Kleinschmidt, B Sturmfels**, *On the existence of certain smooth toric varieties*, Discrete Comput. Geom. 5:3 (1990) 255–262 [MR](#)
- [13] **S Hasui**, *On the classification of quasitoric manifolds over dual cyclic polytopes*, Algebr. Geom. Topol. 15:3 (2015) 1387–1437 [MR](#)

- [14] **H Ishida, Y Fukukawa, M Masuda**, *Topological toric manifolds*, Mosc. Math. J. 13:1 (2013) 57–98, 189–190 [MR](#)
- [15] **P Mani**, *Spheres with few vertices*, J. Combinatorial Theory Ser. A 13 (1972) 346–352 [MR](#)
- [16] **H Park**, *Wedge operations and doubling operations of real toric manifolds*, Chinese Ann. Math. Ser. B 38:6 (2017) 1321–1334 [MR](#)
- [17] **Y M Ustinovskii**, *The toral rank conjecture for moment-angle complexes*, Mat. Zametki 90:2 (2011) 300–305 [MR](#)

SUYOUNG CHOI schoi@ajou.ac.kr

Department of Mathematics, Ajou University, Suwon, South Korea

HYEONTAE JANG Hjang1112@kias.re.kr

School of Mathematics, Korea Institute for Advanced Study, Seoul, South Korea

Received: October 16, 2024 Revised: January 17, 2025

ALGEBRAIC & GEOMETRIC TOPOLOGY

msp.org/agt

EDITORS

PRINCIPAL ACADEMIC EDITORS

John Etnyre
etnyre@math.gatech.edu
Georgia Institute of Technology

Vesna Stojanoska
vesna@illinois.edu
University of Illinois at Urbana-Champaign

BOARD OF EDITORS

| | | | |
|------------------------|---|-------------------|---|
| Julie Bergner | University of Virginia jeb2md@eservices.virginia.edu | Daniel Isaksen | Wayne State University isaksen@math.wayne.edu |
| Steven Boyer | Université du Québec à Montréal cohf@math.rochester.edu | Thomas Koberda | University of Virginia thomas.koberda@virginia.edu |
| Tara E Brendle | University of Glasgow tara.brendle@glasgow.ac.uk | Markus Land | LMU München markus.land@math.lmu.de |
| Indira Chatterji | CNRS & Univ. Côte d'Azur (Nice) indira.chatterji@math.cnrs.fr | Christine Lescop | Université Joseph Fourier lescop@ujf-grenoble.fr |
| Octav Cornea | Université de Montreal cornea@dms.umontreal.ca | Norihiko Minami | Yamato University minami.norihiko@yamato-u.ac.jp |
| Alexander Dranishnikov | University of Florida dranish@math.ufl.edu | Andrés Navas | Universidad de Santiago de Chile andres.navas@usach.cl |
| Tobias Ekholm | Uppsala University, Sweden tobias.ekholm@math.uu.se | Jessica S Purcell | Monash University jessica.purcell@monash.edu |
| Mario Eudave-Muñoz | Univ. Nacional Autónoma de México mario@matem.unam.mx | Birgit Richter | Universität Hamburg birgit.richter@uni-hamburg.de |
| David Futер | Temple University dfuter@temple.edu | Jérôme Scherer | École Polytech. Féd. de Lausanne jerome.scherer@epfl.ch |
| John Greenlees | University of Warwick john.greenlees@warwick.ac.uk | Zoltán Szabó | Princeton University szabo@math.princeton.edu |
| Matthew Hedden | Michigan State University mhedden@math.msu.edu | Maggy Tomova | University of Iowa maggy-tomova@uiowa.edu |
| Kristen Hendricks | Rutgers University kristen.hendricks@rutgers.edu | Daniel T Wise | McGill University, Canada daniel.wise@mcgill.ca |
| Hans-Werner Henn | Université Louis Pasteur henn@math.u-strasbg.fr | Lior Yanovski | Hebrew University of Jerusalem lior.yanovski@gmail.com |
| Kathryn Hess | École Polytechnique Féd. de Lausanne kathryn.hess@epfl.ch | | |

See inside back cover or msp.org/agt for submission instructions.

The subscription price for 2026 is US \$795/year for the electronic version, and \$1170/year (+\$80, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Algebraic & Geometric Topology is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [Current Mathematical Publications](#) and the [Science Citation Index](#).

Algebraic & Geometric Topology (ISSN 1472-2747 printed, 1472-2739 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004.

AGT peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**

nonprofit scientific publishing

<https://msp.org/>

© 2026 Mathematical Sciences Publishers

| | |
|--|-----|
| Isospectrality of Margulis–Smilga spacetimes for irreducible representations of real split semisimple Lie groups | 411 |
| SOURAV GHOSH | |
| RO(G)-graded Bredon cohomology of Euclidean configuration spaces | 437 |
| DANIEL DUGGER and CHRISTY HAZEL | |
| KSp-characteristic classes determine Spin^h cobordism | 485 |
| JONATHAN BUCHANAN and STEPHEN MCKEAN | |
| Linear upper bounds on ribbonlength of knots and links | 553 |
| HYOUNGJUN KIM, SUNGJONG NO and HYUNGKEE YOO | |
| Profinite rigidity properties of central extensions of 2-orbifold groups | 565 |
| PAWEŁ PIWEK | |
| Magnitude homology equivalence of Euclidean sets | 599 |
| ADRIÁN DOÑA MATEO and TOM LEINSTER | |
| Characterising slopes for hyperbolic knots and Whitehead doubles | 625 |
| LAURA WAKELIN | |
| The quasi-isometry invariance of the coset intersection complex | 659 |
| CAROLYN ABBOTT and EDUARDO MARTÍNEZ-PEDROZA | |
| Symmetry in the cubical Joyal model structure | 699 |
| BRANDON DOHERTY | |
| Explicit formulas for the Hattori–Stong theorem and applications | 735 |
| PING LI and WANGYANG LIN | |
| Stellar subdivisions, wedges and Buchstaber numbers | 751 |
| SUYOUNG CHOI and HYEONTAE JANG | |
| An obstruction theory for strictly commutative algebras in positive characteristic | 761 |
| OISÍN FLYNN-CONNOLLY | |
| Spherical p -group complexes arising from finite groups of Lie type | 791 |
| KEVIN IVAN PITERMAN | |