

AG
T

*Algebraic & Geometric
Topology*

Volume 26 (2026)

Tautological rings of fibrations

NILS PRIGGE



Tautological rings of fibrations

NILS PRIGGE

We study the analogue of tautological rings of fibre bundles in the context of fibrations with Poincaré fibre, i.e., the ring obtained by fibre integrating powers of the fibrewise Euler class. We discuss how to compute the Euler ring with tools from rational homotopy theory and completely determine the tautological ring for even spheres, complex projective spaces and some products of odd spheres.

1 Introduction

Let $\pi : E \rightarrow B$ be a fibration with fibre X an oriented Poincaré duality space of formal dimension d (see [35, Chapter 1]), which if X is a simply connected finite CW complex just means that one can choose a fundamental class $[X] \in H_d(X; \mathbb{Z})$ that induces the Poincaré duality isomorphism for all local coefficient systems. It is called *oriented* if the corresponding local coefficient system $\mathcal{H}^d(X; \mathbb{Z})$ is trivial and we choose an isomorphism $\mathcal{H}^d(X; \mathbb{Z}) \cong \mathbb{Z}$. In [19, Section 3] the authors construct for such fibrations a *fibrewise Euler class* $e^{\text{fw}}(\pi) \in H^d(E; \mathbb{Z})$ which extends the construction for smooth fibre bundles, i.e., if X is a closed, oriented manifold and $\pi : E \rightarrow B$ is an oriented fibre bundle, then the fibrewise Euler class agrees with the Euler class of the vertical tangent bundle $T_\pi E \rightarrow E$ which, if π is a smooth submersion, is defined as the kernel of the differential of the projection map $T_\pi E := \ker(D\pi : TE \rightarrow TB) \subset TE$.

Using the fibrewise Euler class we can extend the construction of *tautological classes* of smooth fibre bundles [14; 16; 23; 27] to fibrations. Recall that given a smooth, oriented fibre bundle $\pi : E \rightarrow B$ with fibre M a closed manifold and a class $c \in H^{|c|}(\text{B SO}(d); \mathbb{Q})$, the fibre integral

$$\kappa_c(\pi) := \int_\pi c(T_\pi E) \in H^{|c|-d}(B; \mathbb{Q})$$

is a characteristic class of the bundle and the subring $R^*(\pi) \subset H^*(B; \mathbb{Q})$ generated by all tautological classes is called the *tautological ring*.

As we can define fibre integration more generally for oriented fibrations with Poincaré fibre (see [6, Section 8] and Section 3), we obtain analogous characteristic classes of oriented fibrations $\pi : E \rightarrow B$ with Poincaré fibre by setting

$$\kappa_i(\pi) := \pi_!(e^{\text{fw}}(\pi)^{i+1}) \in H^{i-d}(B; \mathbb{Z}),$$

where $\pi_! : H^*(E; \mathbb{Z}) \rightarrow H^{*-d}(B; \mathbb{Z})$ is another notation for the fibre integration map. In particular, for the universal oriented X -fibration that classifies oriented X -fibrations [30]

$$(1-1) \quad X \hookrightarrow E \xrightarrow{\pi} \text{BhAut}^+(X),$$

MSC2020: 55R40.

© 2026 MSP (Mathematical Sciences Publishers). Distributed under the Creative Commons Attribution License 4.0 (CC BY). Open Access made possible by subscribing institutions via [Subscribe to Open](#).

where the base $\text{BhAut}^+(X)$ is the classifying space of the monoid of orientation-preserving homotopy self-equivalences $\text{hAut}^+(X)$, we can study the subring generated by the tautological classes.

Note that while tautological classes of fibrations can be defined integrally, we only study the tautological ring rationally due to the limitations of the techniques we use, and from now on we always use rational coefficients which we omit from the notation.

Definition 1.1 Let X be an oriented Poincaré duality space of formal dimension d . The Euler ring $E^*(X)$ is the subring of $H^*(\text{BhAut}^+(X))$ generated by all tautological classes

$$(1-2) \quad \kappa_i := \pi_!(e^{\text{fw}}(\pi)^{i+1}) \in H^{i \cdot d}(\text{BhAut}^+(X)),$$

where $e^{\text{fw}}(\pi) \in H^d(E)$ is the fibrewise Euler class of the universal oriented X -fibration in (1-1).

Unlike the smooth tautological ring, the computation of the Euler ring $E^*(X)$ is a purely homotopy-theoretic problem. The main content of this paper is to work out how one can use the models from rational homotopy theory for fibrations to compute Euler rings. Specifically, we determine representatives of the fibre integration maps and fibrewise Euler classes in terms of the algebraic models from rational homotopy theory. The computation of the Euler ring turns out to be particularly tractable for rationally elliptic spaces and we determine the Euler ring for two classes of examples.

Theorem A The Euler ring of complex projective space is $E^*(\mathbb{C}P^n) \cong \mathbb{Q}[\kappa_1, \dots, \kappa_{n-1}, \kappa_{n+1}]$.

Theorem B Let X be either rationally equivalent to $S^{2k+1} \times \dots \times S^{2k+1}$ for $k \geq 1$ or a finite CW complex rationally equivalent to a product of two odd-dimensional simply connected spheres of different dimension. Then $E^*(X) = \mathbb{Q}$.

Some further computations of Euler rings can be found in the author's thesis [25, Section 4.2]. One can also extend the definition of the Euler ring for fibrations with extra structure to obtain, for example, better homotopy-theoretic approximations to the smooth tautological ring [2]. Finally, these techniques can be used to infer properties about smooth tautological rings [26].

2 Rational homotopy theory of fibrations

The classifying space $\text{BhAut}^+(X)$ is rarely simply connected even if X is, and therefore we cannot immediately apply the results from rational homotopy theory. Instead, for a simply connected Poincaré duality space X we study the universal 1-connected fibration

$$(2-1) \quad X \hookrightarrow E \xrightarrow{\pi} \text{BhAut}_0(X),$$

over the classifying space of the path component of the identity $\text{hAut}_0(X) \subset \text{hAut}^+(X)$ (or equivalently the induced fibration over the universal covering of $\text{BhAut}^+(X)$). Many people have studied rational models of (2-1) (see [21; 32; 33]) and fibrations in general (see [11; 18]), and we will use these algebraic models to compute the image of the Euler ring

$$(2-2) \quad E_0^*(X) \subset H^*(\text{BhAut}_0(X))$$

induced by the natural map $\text{BhAut}_0(X) \rightarrow \text{BhAut}^+(X)$. We discuss in Section 5 how one can in some cases upgrade the computation of $E_0(X)$ to a computation of the full Euler ring $E^*(X)$.

2.1 Rational models of fibrations

In the following, we use standard terminology from rational homotopy theory as discussed in [11]. Due to different conventions in the literature, we use cohomological grading conventions for cdgas and differential graded modules and homological grading conventions for dg Lie algebras.

Throughout this paper, we denote a cdga model of a map of connected spaces $\pi : E \rightarrow B$, typically a fibration with simply connected fibre, by $\pi^* : \mathbb{R} \rightarrow \mathbb{S}$ and we also assume, unless stated otherwise, that cdga models are connected (i.e., $\mathbb{R}^0 = \mathbb{S}^0 = \mathbb{Q}$). The following cdga models for fibrations enjoy good homotopical properties.

Definition 2.1 [11, Section 14] A *relative Sullivan algebra* is a cdga $(\mathbb{R} \otimes \Lambda V, D)$ so that $\text{id}_{\mathbb{R}} \otimes 1 : \mathbb{R} \rightarrow \mathbb{R} \otimes \Lambda V$ is a map of cdgas, and $V = \bigoplus_{p \geq 1} V^p$ is a graded vector space with an exhaustive filtration $V(0) \subset V(1) \subset \dots$ of graded subspaces so that $D|_{1 \otimes V(0)} : V(0) \rightarrow \mathbb{R}$ and $D|_{1 \otimes V(k)} : V(k) \rightarrow \mathbb{R} \otimes \Lambda V(k-1)$. A *relative Sullivan model* of a map of cdgas $\pi^* : \mathbb{R} \rightarrow \mathbb{S}$ is a relative Sullivan algebra $\mathbb{S} = (\mathbb{R} \otimes \Lambda V, D)$ together with a quasi-isomorphism $\mathbb{S}' \xrightarrow{\cong} \mathbb{S}$ of \mathbb{R} -algebras.

By [11, Proposition 14.3] any map $\pi^* : \mathbb{R} \rightarrow \mathbb{S}$ of connected cdgas admits a relative Sullivan model if $H(\pi^*) : H^1(\mathbb{R}) \rightarrow H^1(\mathbb{S})$ is injective. Since this condition is always satisfied if π^* is the model of a fibration $\pi : E \rightarrow B$ with connected fibre, we can always find relative Sullivan models in this case. Moreover, if either the base or fibre has finite type and $\pi_1(B)$ acts trivially on the cohomology of the fibre, then $\mathbb{S} \otimes_{\mathbb{R}} \mathbb{Q}$ is a Sullivan model of the fibre by [18, Theorem 20.3].

There is a convenient way to obtain relative Sullivan algebras via differential graded Lie algebras. We follow [2] in our convention (which in turn is based on [33]), in defining Chevalley–Eilenberg complex of a dg Lie algebra L as the dg coalgebra

$$\mathcal{C}^{\text{CE}}(L) = (\Lambda^c sL, D = d_1 + d_2),$$

where $\Lambda^c sL$ denotes the cofree, conilpotent, cocommutative coalgebra on the suspension $(sL)_* = L_{*-1}$ and the differentials are determined by their corestrictions

$$d_1(sl) = -sd_L(l), \quad d_2(sl_1 \wedge sl_2) = (-1)^{|l_1|} s[l_1, l_2].$$

If a dg Lie algebra acts on a cdga A through derivations, then the Chevalley–Eilenberg cochain complex is

$$\mathcal{C}_{\text{CE}}(L; A) := (\text{Hom}(\mathcal{C}^{\text{CE}}(L), A), \partial + t),$$

where

$$\begin{aligned} \partial(f) &= d_A \circ f - (-1)^{|f|} f \circ D, \\ (2-3) \quad t(f)(sl_1 \wedge \dots \wedge sl_n) &= \sum_{i=1}^n (-1)^{|sl_i|(|f|+|sl_1|+\dots+|sl_{i-1}|)} l_i \cdot f(sl_1 \wedge \dots \wedge \widehat{sl}_i \wedge \dots \wedge sl_n), \end{aligned}$$

and $\mathcal{C}_{\text{CE}}(L; A)$ is a cdga via the convolution product. An element $f \in \mathcal{C}_{\text{CE}}^{\text{CE}}(L; A)$ is an n -cochain if $f(sl_1 \wedge \cdots \wedge sl_k) = 0$ unless $k = n$ and we identify 0-cochains with A . Moreover, if A is a Sullivan algebra then $\mathcal{C}_{\text{CE}}(L, A)$ is a relative Sullivan algebra over the Chevalley–Eilenberg cochain complex $\mathcal{C}_{\text{CE}}(L) := \mathcal{C}_{\text{CE}}(L; \mathbb{Q})$ via the map on Chevalley–Eilenberg cochain complexes induced by the unit $\eta : \mathbb{Q} \rightarrow A$.

With these definitions in place, we can describe a cdga model of (2-1). Let X be a simply connected space of finite type and with minimal Sullivan model $(\Lambda V, d)$. Consider the dg Lie algebra $(\text{Der}^+(\Lambda V), [d, -])$ of positive-degree derivations, where a derivation $\theta \in \text{Der}(\Lambda V)$ has degree n if it *lowers*¹ the degree by n (additionally, $\theta \in \text{Der}^+(\Lambda V)_1$ only if $[d, \theta] = 0$). It was first proved by Sullivan [32] that $\text{Der}^+(\Lambda V)$ is a dg Lie model of $\text{BhAut}_0(X)$, and one can further describe a model of (2-1) as follows.

Theorem 2.2 *Let X be a 1-connected space of finite type with minimal Sullivan model $(\Lambda V, d)$ and unit $\eta : \mathbb{Q} \rightarrow \Lambda V$. Then*

$$(2-4) \quad \mathcal{C}_{\text{CE}}^*(\text{Der}^+(\Lambda V); \mathbb{Q}) \xrightarrow{\eta_*} \mathcal{C}_{\text{CE}}^*(\text{Der}^+(\Lambda V); \Lambda V)$$

is a relative Sullivan model of the universal oriented 1-connected fibration (2-1).

Remark 2.3 (i) In a previous version of this article we proved this result via a comparison of (2-4) with Tanré’s model [33] of the universal 1-connected fibration (see also [25]). It has been pointed out to the author by Andrey Lazarev that instead one can derive this result more directly from [21]. Another proof is due to Alexander Berglund and is based on rational models for the bar construction in terms of Chevalley–Eilenberg complexes [1]. The statement of Theorem 2.2 can be found as a special case of [2, Proposition 3.6] and we refer to these papers for a proof.

(ii) Theorem 2.2 is particularly useful for rationally elliptic spaces because the dg Lie algebra $\text{Der}^+(\Lambda V)$ is finite dimensional in contrast to Tanré’s model which is only of finite type. However, if X is rationally hyperbolic then $\text{Der}^+(\Lambda V)$ is not even of finite type and it seems more feasible to study other models of (2-1) instead, for example, based on a dg Lie algebra model of the fibre (see, for example, [3; 31]).

(iii) The theory of minimal models in its most general form [18, Theorem 20.3] applies more generally to give a model for (2-1) for any connected space X of finite type and minimal Sullivan model $(\Lambda V, d)$. It seems plausible to the author that the algebraic tools we develop in subsequent sections generalise to study the Euler ring $E_0^*(X)$ of nonsimply connected Poincaré duality spaces X . However, minimal Sullivan models of nonsimply connected spaces are considerably more complicated and the dg Lie algebra of derivations is intractable for computations, so that we have not worked in this generality.

Example 2.4 Let $X = S^{2n}$ with Sullivan model

$$A_n = (\Lambda(x, y), |x| = 2n, |y| = 4n - 1, d = x^2 \cdot \partial/\partial y).$$

¹This slightly confusing convention is due to using homological grading conventions for dg Lie algebras and cohomological grading conventions for cdgas and can be avoided if one sticks to just one.

Then $\text{Der}^+(A_n)$ is 3-dimensional with basis $\eta_{2n-1} := x \cdot \partial/\partial y$, $\eta_{2n} := \partial/\partial x$ and $\eta_{4n-1} := \partial/\partial y$ and differential $[d, \eta_{2n}] = -2\eta_{2n-1}$. Hence, the inclusion of the abelian Lie algebra with trivial differential $\mathfrak{g} := \mathbb{Q}\{\eta_{4n-1}\} \hookrightarrow \text{Der}^+(A_n)$ is a quasi-isomorphism of dg Lie algebras and we get a cdga quasi-isomorphism $\mathcal{C}_{\text{CE}}^*(\text{Der}^+(A_n); \mathbb{Q}) \rightarrow \mathcal{C}_{\text{CE}}^*(\mathfrak{g}; \mathbb{Q}) = (\Lambda z_{4n}, d = 0)$ as well as with coefficients in A_n . Thus, the cdga model of the universal 1-connected S^{2n} -fibration in Theorem 2.2 is equivalent to

$$(\Lambda(z_{4n}), d = 0) \rightarrow (\Lambda(z_{4n}) \otimes \Lambda(x, y), D(x) = 0, D(y) = x^2 + z_{4n}),$$

where z_{4n} is the 1-cochain dual to $s\eta_{4n-1}$.

3 Fibre integration in rational homotopy theory

Before we discuss rational models for fibre integration, we recall the definition suitable for oriented fibrations as a special case of the following construction: Let $\pi : E \rightarrow B$ be a fibration with fibre X and $H^*(X) = 0$ for $* > d$ and $H^d(X)$ nontrivial. Given a $\pi_1(B)$ -module homomorphism $\phi : \mathcal{H}^d(X) \rightarrow \mathbb{Q}$, we can define ϕ -integration as the composition

$$(3-1) \quad \phi_! : H^*(E) \twoheadrightarrow E_\infty^{*-d,d} \subset E_2^{*-d,d} = H^{*-d}(B; \mathcal{H}^d(X)) \xrightarrow{H(\phi)} H^{*-d}(B),$$

where we project $H^*(E)$ onto the d -th row of the E_∞ -page of the Serre spectral sequence, which is possible since $H^*(X) = 0$ for $* > d$, and $E_\infty^{*-d,d} \subset E_2^{*-d,d}$ as there are no differentials into this row.

Note that because the cohomological Serre spectral sequence is compatible with cup product, there is a push-pull identity

$$(3-2) \quad \phi_!(\pi^*(x) \smile y) = x \smile \phi_!(y)$$

for $x \in H^*(B)$ and $y \in H^*(E)$, and thus $\phi_!$ is a $H^*(B)$ -module map.

Definition 3.1 Let $\pi : E \rightarrow B$ be an orientable fibration with Poincaré fibre X of formal dimension d and let $\varepsilon_X : H^d(X) \rightarrow \mathbb{Q}$ be an orientation of X . Then

$$\pi_! := (\varepsilon_X)_! : H^*(E) \rightarrow H^{*-d}(B)$$

is called *fibre integration* of $\pi : E \rightarrow B$.

3.1 Chain level fibre integration

In the main result of this section we show that there exists a chain level representative of fibre integration. More precisely, given a relative Sullivan model $S = (R \otimes \Lambda V, D)$ of an oriented fibration $\pi : E \rightarrow B$ with Poincaré fibre X , there exists a R -module homomorphism $\Pi : S \rightarrow R$ of degree $-d$ which is unique up to homotopy and induces fibre integration on cohomology. This is most conveniently expressed in terms of differential Ext groups.

Let R be a connected cdga and M, N a differential graded R -modules. Then R -module homomorphisms $\text{Hom}_R(M, N)$ is a R -module with differential $D(f) := d_N f - (-1)^{|f|} f d_M$. Recall from [11, Section 7]

that M is a *semifree* R -module if there is an exhaustive filtration $M(0) \subset M(1) \subset \dots \subset M$ so that $M(0)$ and $M(k)/M(k-1)$ are free R -modules for all $k \geq 1$. A *semifree resolution* of a R -module M is a semifree R -module M' with a quasi-isomorphism $M' \xrightarrow{\cong} M$ and we denote by $M \otimes_{\mathbb{R}}^{\mathbb{L}} N$ the derived tensor product of two R -modules given by $M' \otimes_R N$ for some semifree resolution M' . Following [12], the differential Ext groups for R -modules M, N are defined as

$$(3-3) \quad \text{Ext}_{\mathbb{R}}(M, N) := H(\text{Hom}_{\mathbb{R}}(M', N))$$

for a semifree resolution $M' \xrightarrow{\cong} M$.

Proposition 3.2 *Let $\pi : E \rightarrow B$ be a fibration with connected base and total space and 1-connected fibre X . Assume that $\pi_1(B)$ acts trivially on $H^*(X)$ and that $H^*(X)$ is of finite type and nontrivial in degree d and vanishes for $* > d$. Let $\pi^* : R \rightarrow S$ be a cdga model of π . Then the augmentation induces an isomorphism*

$$(3-4) \quad \text{Ext}_{\mathbb{R}}^{-d}(S, \mathbb{R}) \xrightarrow{\cong} \text{Ext}_{\mathbb{Q}}^{-d}(S \otimes_{\mathbb{R}}^{\mathbb{L}} \mathbb{Q}, \mathbb{Q}) \cong \text{Hom}(H^d(X), \mathbb{Q}),$$

and given $\phi \in \text{Hom}(H^d(X), \mathbb{Q})$ the chain level representative $[\Phi] \in \text{Ext}_{\mathbb{R}}^{-d}(S, \mathbb{R})$ induces ϕ -integration.

Remark 3.3 Proposition 3.2 generalises [13, Theorem A] where they identify fibre integration as elements in differential Ext groups for fibrations over Poincaré duality spaces and for pullbacks from such fibrations. Moreover, fibre integration can be identified rationally as a map of parametrised suspension spectra $\pi_! : \Sigma_B^{\infty} B_+ \rightarrow \Sigma_B^{\infty-d} E_+$, and by [12, Theorem 1.1] the set of homotopy classes of such maps is given by differential Ext groups which is consistent with our result.

Proof Let $S' \xrightarrow{\cong} S$ be a relative Sullivan model. Then $\text{Ext}^*(S, \mathbb{R}) = H^*(\text{Hom}_{\mathbb{R}}(S', \mathbb{R}))$ since S' is a semifree resolution by [11, Lemma 14.1]. We consider the exhaustive filtration of $\text{Hom}_{\mathbb{R}}(S', \mathbb{R})$ given by $F^p = \text{Hom}_{\mathbb{R}}(S', \mathbb{R}^{\geq p})$. According to [5, Theorem 9.3] the corresponding spectral sequence is conditionally convergent to the completion

$$\begin{aligned} \varprojlim_p \text{Hom}_{\mathbb{R}}(S', \mathbb{R}) / \text{Hom}_{\mathbb{R}}(S', \mathbb{R}^{\geq p}) &\cong \varprojlim_p \text{Hom}_{\mathbb{R}}(S', \mathbb{R} / \mathbb{R}^{\geq p}) \\ &\cong \text{Hom}_{\mathbb{R}}(S', \varprojlim_p \mathbb{R} / \mathbb{R}^{\geq p}) \\ &\cong \text{Hom}_{\mathbb{R}}(S', \mathbb{R}), \end{aligned}$$

where we have used that S' is a projective R -module for the first isomorphism. The E_1 -page is $H(\text{Hom}_{\mathbb{R}}(S', \mathbb{Q})) \otimes B^p$. By the assumption on the fibration it follows from [18] that the Sullivan fibre $S' \otimes_{\mathbb{R}} \mathbb{Q} = (\Lambda V, d)$ is a cdga model of X . Hence, we have that the E_1 -page can be simplified as $E_1^{p,q} = H^q((\Lambda V)^{\vee}) \otimes R^p \cong \text{Hom}^q(H^*(X), \mathbb{Q}) \otimes R^p$. Since $\pi_1(B)$ acts trivially, the differential on the E_1 -page is given by $\text{id} \otimes d_R : E_1^{p,q} \rightarrow E_1^{p+1,q}$ (if R was simply connected this follows for degree reasons but in general it follows from the fact the cdga model of the universal 1-connected X -fibration in Theorem 2.2 has a simply connected base and by assumption the relative Sullivan model is obtained by base change along a map $\mathcal{C}_{\text{CE}}(\text{Der}^+(\Lambda V)) \rightarrow R$). Hence the E_2 -page is $E_2^{p,q} = H^p(\mathbb{R}) \otimes H^q((\Lambda V)^{\vee})$.

In particular, the spectral sequence vanishes for $q < -d$. Since the gradings are such that the differentials are $d_r : E_r^{p,q} \rightarrow E_r^{p+r,q-r+1}$, there are only finitely many nontrivial differentials. This implies that the derived E_∞ -page is zero and so by [5, Theorem 7.1] the spectral sequence converges strongly,

$$E_2^{p,q} = H^p(\mathbb{R}) \otimes H^q((\Lambda V)^\vee) \cong H^p(B) \otimes \text{Hom}^q(H^*(X), \mathbb{Q}) \Rightarrow H(\text{Hom}_{\mathbb{R}}(S', \mathbb{R})),$$

and we can recover $H(\text{Hom}_{\mathbb{R}}(S', \mathbb{R}))$ from the entries of the E_∞ -page. The only contribution with total degree $-d$ comes from $E_\infty^{0,-d} \cong E_2^{0,-d} \cong \text{Hom}(H^d(X), \mathbb{Q})$ which proves the first part of the statement.

It remains to show that for $\phi = H^d(\Phi \otimes_{\mathbb{R}} \mathbb{Q}) : H^d(\Lambda V, d) \rightarrow \mathbb{Q}$ the induced ϕ -integration map coincides with $H(\Phi) : H^*(S') \rightarrow H^*(\mathbb{R})$. First, we note that $S' = (\mathbb{R} \otimes \Lambda V, D)$ has a filtration $F^p = \mathbb{R}^{\geq p} \otimes \Lambda V$ and that the corresponding spectral sequence converges as $E_2^{p,q} = H^p(\mathbb{R}) \otimes H^q(\Lambda V) \Rightarrow H^{p+q}(S')$. In fact, \mathbb{R} also has an analogous filtration $G^p = \mathbb{R}^{\geq p}$ with only nontrivial differential on the E_1 -page. Then Φ induces a map between these two filtrations and the map on E_2 -pages is precisely ϕ -integration defined using this spectral sequence. As we have defined ϕ -integration using the Serre spectral sequence, it remains to show that this spectral sequence is isomorphic to the Serre spectral sequence. Grivel has shown in [17] that the above filtration gives rise to the Serre spectral sequence if the base is simply connected, and this has been generalised by Halperin [18] if $\pi_1(B)$ acts nilpotently on the cohomology of the fibre. More precisely, it follows from the proof of [17, Theorem 6.4], respectively [18, Section 20], that the comparison map of $\mathbb{R} \rightarrow S'$ with $A_{PL}(B) \rightarrow A_{PL}(E)$ is compatible with Dress' construction of the Serre spectral sequence [10] and induces an isomorphism on the E_2 -pages. \square

We can use Proposition 3.2 to build a representative of fibre integration for an oriented fibration $\pi : E \rightarrow B$ and oriented Poincaré fibre (X, ε_X) : Consider a relative Sullivan model $\pi^* : \mathbb{R} \rightarrow (\mathbb{R} \otimes \Lambda V, D)$. Pick a chain level representative ε of the orientation $\varepsilon_X \in H^{-d}(\text{Hom}(\Lambda V, \mathbb{Q}))$ of X . By Proposition 3.2 there is a cycle $\Pi \in \text{Hom}^{-d}(S', \mathbb{R})$ unique up to chain homotopy that satisfies

$$(3-5) \quad \Pi(1 \otimes \chi) = \varepsilon_X(\chi) \in \mathbb{R}^0 = \mathbb{Q}$$

for all $\chi \in (\Lambda V)^d$ and that induces fibre integration on cohomology

$$\pi_! : H^*(E; \mathbb{Q}) \cong H^*(S') \xrightarrow{H(\Pi)} H^{*-d}(\mathbb{R}) \cong H^{*-d}(B; \mathbb{Q}).$$

We demonstrate this technique in the following example.

Example 3.4 Recall the relative Sullivan model of the universal 1-connected fibration for an even-dimensional sphere $X = S^{2n}$ as discussed in Example 2.4. We choose as orientation $\varepsilon_X : A_n \rightarrow \mathbb{Q}$ the homomorphism determined by $\varepsilon_X(x) = 1$. For degree reasons $\Pi(yx^k) = 0$ and since Π has to be a chain map we have $0 = \Pi(D(yx^k)) = \Pi(x^{k+2} + z_{4n}x^k)$. This determines a $\Lambda(z_{4n})$ -module map $\Pi : (\Lambda(z_{4n}, x, y), D) \rightarrow (\Lambda(z_{4n}), d = 0)$ by

$$\Pi(yx^k) = 0 \quad \text{and} \quad \Pi(x^n) = \begin{cases} 0, & n = 2k, \\ (-1)^k z_{4n}^k, & n = 2k + 1, \end{cases}$$

which is a chain map by construction and induces fibre integration on cohomology as it satisfies (3-5).

4 The fibrewise Euler class

The definition of the fibrewise Euler class in [19, Definition 3.1.1] uses constructions in the category of parametrised spectra. And while there has been a lot of progress to adapt the tools from rational homotopy theory to the context of parametrised stable homotopy theory [7; 8], there is a simpler way to define the fibrewise Euler class with rational coefficients so that we can avoid discussing the category of parametrised spectra and their rational models altogether.

We begin by describing a special case of the definition of the fibrewise Euler class. Consider an oriented fibration $\pi : E \rightarrow B$ with fibre X so that both X and B are oriented Poincaré duality spaces of dimensions d and b , respectively. Then the total space is also an oriented Poincaré duality space [15], and we can define the Umkehr map of the fibrewise diagonal $\Delta : E \rightarrow E \times_B E$ by

$$(4-1) \quad \Delta_! : H^*(E) \xrightarrow{D_E} H_{b+d-*}(E) \xrightarrow{\Delta^*} H_{b+d-*}(E \times_B E) \xrightarrow{D_{E \times_B E}^{-1}} H^{*+d}(E \times_B E),$$

where D_E and $D_{E \times_B E}$ denote the Poincaré duality isomorphisms. It is shown in [19, Section 3] that the fibrewise Euler class of $\pi : E \rightarrow B$ agrees with

$$(4-2) \quad e^{\text{fw}}(\pi) = \Delta^*(\Delta_!(1)) \in H^d(E).$$

This is sufficient to define the fibrewise Euler class with rational coefficients, because for any space X the rational homology groups are isomorphic to rationalised stable framed bordism $H_*(X; \mathbb{Q}) \cong \Omega_*^{\text{sfr}}(X) \otimes \mathbb{Q}$, and so we can determine a rational cohomology class by defining its evaluation on framed bordism classes. Hence, denoting by E the total space of the universal 1-connected fibration (2-1), given a stably framed bordism class $[f : M^d \rightarrow E, \xi] \in \Omega_d^{\text{sfr}}(E) \otimes \mathbb{Q}$, we can consider the pullback of the X -fibration $p_1 : E \times_{\text{BhAut}_0(X)} E \rightarrow E$ along f . The pullback $\pi : f^*(E \times_{\text{BhAut}_0(X)} E) \rightarrow M$ has a section s via the diagonal, and we can associate to it an Euler class e_f via (4-2). Then the fibrewise Euler class $e^{\text{fw}}(\pi) \in H^d(E; \mathbb{Q})$ agrees with the class defined by the pairing $\langle e^{\text{fw}}(\pi), [f : M \rightarrow E, \xi] \rangle = \langle s^*(e_f), [M] \rangle$ since the construction in [19] coincides with (4-2) if the base is a Poincaré duality space.

4.1 A cocycle representative via rational homotopy theory

The idea to obtain cocycle representatives of the fibrewise Euler class is simply to construct a chain level representative of the Umkehr map $\Delta_! : H^*(E) \rightarrow H^{*+d}(E \times_B E)$ in terms of the algebraic models. In the following, we denote by $\mathbb{Q}[n]$ the graded vector space with \mathbb{Q} in degree n and for an \mathbb{R} -module M define $M[n] := M \otimes \mathbb{Q}[n]$.

Proposition 4.1 *Let $\mathbb{R} \rightarrow \mathbb{S}$ be a relative Sullivan model of an oriented fibration $\pi : E \rightarrow B$ with connected base and total space and simply connected Poincaré fibre X of formal dimension d . Let ε_X be the orientation and $\bar{\Pi} \in \text{Hom}_{\mathbb{R}}^{-d}(\mathbb{S}, \mathbb{R})$ be a cocycle representative of fibre integration. Then the map*

$$\bar{\Pi} : \mathbb{S}[-d] \rightarrow \text{Hom}_{\mathbb{R}}(\mathbb{S}, \mathbb{R}), \quad e \mapsto (e' \mapsto (-1)^{d+d \cdot |e|} \bar{\Pi}(e \cdot e')),$$

is a quasi-isomorphism of \mathbb{R} -modules.

Proof It is a simple check that $\overline{\Pi}$ defines a \mathbb{R} -module homomorphism. By assumption, $S = (\mathbb{R} \otimes \Lambda V, D)$ is a relative Sullivan algebra and thus has a filtration which induces the Serre spectral sequence as discussed in the proof of Proposition 3.2. In the same proof we have described a filtration of $\text{Hom}_{\mathbb{R}}(S, \mathbb{R})$ which strongly converges because there is a horizontal vanishing line. The map $\overline{\Pi}$ is compatible with the two filtrations and induces a map of the associated spectral sequences. The induced map on the E_2 -page is given by

$$E_2^{p,q} = H^p(\mathbb{R}) \otimes H^q(\Lambda V, d) \xrightarrow{\text{Id} \otimes \bar{\varepsilon}_X} H^p(\mathbb{R}) \otimes \text{Hom}(H^{q-d}(\Lambda V), \mathbb{Q}),$$

where $\bar{\varepsilon}_X : H^q(\Lambda V) \rightarrow \text{Hom}(H^{d-q}(\Lambda V), \mathbb{Q})$ is the adjoint of $H^q(\Lambda V) \otimes H^{d-q}(\Lambda V) \xrightarrow{\cup} H^d(\Lambda V) \xrightarrow{\varepsilon_X} \mathbb{Q}$. Since $(H^*(X; \mathbb{Q}), \varepsilon_X)$ is an oriented Poincaré duality algebra, $\overline{\Pi}$ induces an isomorphism of E_2 -pages. \square

This enables us to define an algebraic Umkehr map as follows. Let $\mathbb{R} \rightarrow \mathbb{S}$ be a relative Sullivan model of $\pi : E \rightarrow B$ and $\Pi : \mathbb{S} \rightarrow \mathbb{R}$ a chain level representative of fibre integration. Then $\mathbb{S} \otimes_{\mathbb{R}} \mathbb{S}$ is a Sullivan model of $E \times_B E$, the multiplication $\mu : \mathbb{S} \otimes_{\mathbb{R}} \mathbb{S} \rightarrow \mathbb{S}$ is a model of the fibrewise diagonal $\Delta : E \rightarrow E \times_B E$ and $\Pi \otimes \Pi : \mathbb{S} \otimes_{\mathbb{R}} \mathbb{S} \rightarrow \mathbb{R} \otimes_{\mathbb{R}} \mathbb{R} = \mathbb{R}$ is a chain level representative of fibre integration for $E \times_B E$. Since \mathbb{S} is \mathbb{R} -semifree, we can find a lift of \mathbb{R} -modules

$$(4-3) \quad \begin{array}{ccc} S'[-d] & \overset{\Delta_!}{\dashrightarrow} & S' \otimes_{\mathbb{R}} S'[-2d] \\ \overline{\Pi} \downarrow \simeq & & \overline{\Pi \otimes \Pi} \downarrow \simeq \\ \text{Hom}_{\mathbb{R}}(S', \mathbb{R}) & \xrightarrow{\mu^*} & \text{Hom}_{\mathbb{R}'}(S' \otimes_{\mathbb{R}} S', \mathbb{R}) \end{array}$$

which is unique up to homotopy and therefore obtain a well-defined class

$$(4-4) \quad [\mu(\Delta_!(1)) \in H^d(\mathbb{S})].$$

Proposition 4.2 *Let $\pi : E \rightarrow B$ be an oriented fibration with connected base and total space and simply connected Poincaré fibre X of formal dimension d so that $\pi_1(B)$ acts trivially on $H^*(X)$. If $\mathbb{R} \rightarrow \mathbb{S}$ is a relative Sullivan model and $\Delta_! : \mathbb{S} \rightarrow \mathbb{S} \otimes_{\mathbb{R}} \mathbb{S}$ an Umkehr map as in (4-3), then $\mu(\Delta_!(1)) \in S^d$ is a representative of the fibrewise Euler class $e^{\text{fw}}(\pi) \in H^d(E)$.*

Proof We first observe that the definition of the class in (4-4) is natural with respect to pullbacks: For a map $f : B' \rightarrow B$ with cdga model $\phi : \mathbb{R} \rightarrow \mathbb{R}'$, a model of the pullback $\pi : f^*E \rightarrow B'$ is given by $\mathbb{R}' \otimes_{\mathbb{R}} \mathbb{S}$ and a model on the map of total spaces is given by sending $\Phi(s) = 1 \otimes s \in \mathbb{R}' \otimes_{\mathbb{R}} \mathbb{S}$ for $s \in \mathbb{S}$ by [18, Section 20.6]. It follows from Proposition 3.2 that a cocycle representative of fibre integration is given by $\mathbb{R}' \otimes_{\mathbb{R}} \Pi : \mathbb{R}' \otimes_{\mathbb{R}} \mathbb{S} \rightarrow \mathbb{R}' \otimes_{\mathbb{R}} \mathbb{R} \cong \mathbb{R}'$ and therefore $\mathbb{R}' \otimes_B \Delta_!$ is a model of the Umkehr map so that (4-4) in this case is given by $[1 \otimes \mu(\Delta_!(1))] = \Phi(\mu(\Delta_!(1))) \in \mathbb{R}' \otimes_{\mathbb{R}} \mathbb{S}$.

Hence, it suffices to prove that (4-4) coincides with the class defined in (4-2) for fibrations where the base space is a Poincaré duality space (or even just for closed, stably framed manifolds). So let us assume that the base space is a closed manifold B or more generally a Poincaré duality space. Then $B \rightarrow *$ is a fibration with Poincaré fibre and we can apply the results of the previous section to get a chain level representative of fibre integration map $\Pi_B \in \text{Hom}_{\mathbb{Q}}(\mathbb{R}, \mathbb{Q})$ corresponding to evaluating a fundamental class.

Suppose $R \rightarrow S$ is a Sullivan model of the fibration and let $\Pi \in \text{Hom}_R(S, R)$ be a model of fibre integration of π . Then $([B] \cap -) \circ \pi_! : H^{b+d}(E) \rightarrow H^b(B) \rightarrow \mathbb{Q}$ is an orientation of the Poincaré algebra $H^*(E; \mathbb{Q})$ itself and therefore $\Pi_B \circ \Pi \in \text{Hom}_{\mathbb{Q}}(S, \mathbb{Q})$ is a cocycle representative. Define $\Pi_E := \Pi_B \circ \Pi$ so that

$$\begin{array}{ccc} \text{Hom}_R(S, R) & \xleftarrow{\cong} & S[-d] & \xrightarrow{\cong} & \text{Hom}_{\mathbb{Q}}(S, \mathbb{Q}) \\ & & \searrow & \swarrow & \\ & & & & \end{array} \quad \begin{array}{c} \xrightarrow{\quad (\Pi_B)_* \quad} \\ \end{array}$$

is a commuting diagram, where $(\Pi_B)_* f = \Pi_B \circ f$ for $f \in \text{Hom}_R(S, R)$. We can choose chain level representatives of fibre integration of $E \times_B E \rightarrow B$ and $E \times_B E \rightarrow *$ as $\Pi \otimes \Pi : S \otimes_R S \rightarrow R \otimes_R R \cong R$ and $\Pi_{E \times_B E} := \Pi_B \circ (\Pi \otimes \Pi)$ by the same arguments as above. We therefore have a diagram

$$\begin{array}{ccc} S[-d] & \overset{\Delta_!}{\dashrightarrow} & S \otimes_R S[-2d] \\ \downarrow \cong & & \downarrow \cong \\ \Pi_E \left(\begin{array}{ccc} \text{Hom}_R(S, R) & \xrightarrow{\Delta^*} & \text{Hom}_R(S \otimes_R S, R) \\ \downarrow (\Pi_B)_* & & \downarrow (\Pi_B)_* \\ \text{Hom}_{\mathbb{Q}}(S, \mathbb{Q}) & \xrightarrow{\Delta^*} & \text{Hom}_{\mathbb{Q}}(S \otimes_R S, \mathbb{Q}) \end{array} \right) \cong \Pi_{E \times_B E} \end{array}$$

where the dashed maps denote the Umkehr map from (4-3). The upper square commutes up to homotopy by construction and therefore so does the outer square by commutativity of the lower square. This shows that $\Delta_!$ is a cochain level representative of the Gysin map and thus $[\mu(\Delta_!(1))] \in H^d(S)$ agrees with (4-2). \square

4.2 The fibrewise Euler class of Leray–Hirsch fibrations

In the case of fibrations $\pi : E \rightarrow B$ with oriented Poincaré fibre (X, ε_X) which are Leray–Hirsch, i.e., where the restriction map $H^*(E) \rightarrow H^*(X)$ is surjective, the definition of fibre integration and the fibrewise Euler class can be simplified significantly. Surjectivity of the restriction map implies that $H^*(E)$ is a free $H^*(B)$ -module, and we may denote by $1, e_1, \dots, e_k \in H^*(E)$ a $H^*(B)$ -basis of the cohomology of E that restricts to a basis $1, x_1 = i^*(e_1), \dots, x_k = i^*(e_k) \in H^*(X)$ of the fibre. If X is a Poincaré complex of formal dimension d , we can order the basis such that $|e_k| = d$ and all other $|e_i|$ have lower degree. Since fibre integration is a $H^*(B)$ -module map. It suffices to determine $\pi_!$ on a basis and for degree reasons $\pi_!(e_i) = 0$ for $i < k$. If we set $\pi_!(e_k) = \varepsilon_X(x_k)$ this restricts to the orientation on the fibre hence determines fibre integration as

$$(4-5) \quad \pi_! \left(\sum_{i=0}^k b_i \cdot e_i \right) = \varepsilon_X(x_k) \cdot b_k$$

for $b_i \in H^*(B)$. Since the fibre is Poincaré, the (fibrewise) intersection pairing

$$(4-6) \quad \langle -, - \rangle : H^*(E) \otimes_{H^*(B)} H^*(E) \xrightarrow{\pi_!(- \cup -)} H^*(B)$$

is nondegenerate. This enables us to mimic the construction of the Euler class as the dual of the fibrewise diagonal.

Proposition 4.3 [27] *Let $\pi : E \rightarrow B$ be an oriented fibration with Poincaré fibre which is Leray–Hirsch. Let $e_0, \dots, e_k \in H^*(E)$ be an $H^*(B)$ -module basis and denote by $e_0^\#, \dots, e_k^\# \in H^*(E)$ the dual basis under the nondegenerate pairing (4-6). Then the fibrewise Euler class is*

$$(4-7) \quad e^{\text{fw}}(\pi) = \sum_{i=0}^k (-1)^{|e_i|} e_i e_i^\# \in H^d(E; \mathbb{Q}).$$

Example 4.4 Let $X = S^{2n}$ be an even-dimensional sphere and recall the cdga model and fibre integration from the Examples 2.4 and 3.4. Then 1 and x are a $H^*(\text{BhAut}_0(S^{2n}); \mathbb{Q})$ -basis of the cohomology of the total space that restricts to a basis of $H^*(S^{2n})$ on the fibre, i.e., the fibration is Leray–Hirsch. Note that the formula for fibre integration in (4-5) gives the same result as our construction of Π in Example 3.4. We can apply the above proposition to find a representative of the fibrewise Euler class. The dual basis with respect to the pairing induced by $\pi_! = H(\Pi)$ is $x^\# = 1$ and $1^\# = x$ (since $\pi_!(x \cdot 1^\#) = \pi_!(x^2) = 0$), and we find that the fibrewise Euler class is represented by $e^{\text{fw}}(\pi) = 2x$.

4.3 Fibrations with positively rationally elliptic fibre

A simply connected space X is called *rationally elliptic* if both $\dim H(X; \mathbb{Q})$ and $\dim \pi(X) \otimes \mathbb{Q}$ are finite-dimensional vector spaces. If moreover $\chi(X) > 0$ then X is called *positively rationally elliptic*. Algebraic models for (positively) elliptic spaces are quite rigid. For example, one can show that any positively rationally elliptic space satisfies rational Poincaré duality. The main result in this section is a simple, closed formula for the fibrewise Euler class of a fibration with positively rationally elliptic fibre.

The minimal Sullivan model of a rationally elliptic space $\Lambda = (\Lambda V, d)$ is free on a finite-dimensional vector space V . Hence, the dg Lie algebra model $\text{Der}^+(\Lambda)$ for $\text{BhAut}_0(X)$ is finite dimensional which makes the study of fibrations with rationally elliptic fibres tractable via Theorem 2.2. Moreover, a famous conjecture due to Halperin states that any fibration with positively elliptic fibre (and trivial holonomy action) is Leray–Hirsch. This conjecture is known to be true for a large number of examples [22; 28; 34] and by [22] it is equivalent to $\pi_{2i-1}(\text{BhAut}_0(X)) \otimes \mathbb{Q} = 0$ for all $i \in \mathbb{N}$. Since $\pi_{2i-1}(\text{BhAut}_0(X)) \otimes \mathbb{Q}$ is isomorphic to $H_{2i}(\text{Der}^+(\Lambda), [d, -])$, this condition can easily be checked in examples.

We recall a few results about positively rationally elliptic space from [11, Section 32]. A pure Sullivan algebra is a cdga $\Lambda = (\Lambda V, d)$ with $d(V^{\text{even}}) = 0$ and $d(V^{\text{odd}}) \subset \Lambda V^{\text{even}}$, and we define $P = V^{\text{odd}}$ and $Q = V^{\text{even}}$. Observe that pure Sullivan algebras Λ are bigraded with additional lower grading given by $\Lambda_k = \Lambda Q \otimes \Lambda^k P$. By [11, Proposition 32.10] a minimal Sullivan model of a positively rationally elliptic space is isomorphic to a pure Sullivan algebra with $\dim P = \dim Q$ and so that $d|_P$ maps a basis of P to a regular sequence of the graded polynomial ring ΛQ . In this case, the cohomology is concentrated in lower grading 0, i.e., $H^*(\Lambda) = H_0^*(\Lambda)$.

This bigrading is inherited by the dg Lie algebra $\text{Der}_{*,*}^+(\Lambda)$ where a derivation θ has bidegree (m, n) if θ lowers the internal degree by $|m|$ and $\theta(\Lambda Q \otimes \Lambda^k P) \subset \Lambda Q \otimes \Lambda^{k+n} P$. The following statement has been explained to the author by Alexander Berglund.

Lemma 4.5 *Let Λ be a pure Sullivan model for a positively rationally elliptic space that satisfies the Halperin conjecture. There exists an abelian dg Lie algebra with trivial differential $\mathfrak{a} \subset \text{Der}_{*,-1}^+(\Lambda)$ that is quasi-isomorphic to $\text{Der}^+(\Lambda)$.*

Proof If Λ is a pure Sullivan model for a positively rationally elliptic space, the projection map $\psi : \Lambda \rightarrow H(\Lambda)$ is a quasi-isomorphism. It induces a quasi-isomorphism of chain complexes $\text{Der}^+(\Lambda) \rightarrow \text{Der}_{\psi}^+(\Lambda, H(\Lambda))$ (analogous to [3, Lemma 3.5]). Since the $H(\Lambda) = H_0(\Lambda)$, we see that $H_{*,>0}(\text{Der}^+(\Lambda))$ is contained in the kernel and thus is trivial. The Halperin conjecture implies that $H_*(\text{Der}^+(\Lambda))$ is concentrated in odd degrees and therefore $H_*(\text{Der}^+(\Lambda)) \cong H_{\text{odd},-1}(\text{Der}^+(\Lambda))$ as claimed. Hence, any choice of representatives in $\text{Der}_{*,-1}^+(\Lambda)$ of a basis of $H_*(\text{Der}^+(\Lambda))$ spans an abelian dg Lie subalgebra with trivial differential that is quasi-isomorphic to $\text{Der}^+(\Lambda)$. \square

Halperin’s conjecture implies that $\text{BhAut}_0(X) \simeq_{\mathbb{Q}} \prod_{i=1}^k K(\mathbb{Q}, 2n_i)$ and so the cohomology of the total space E of (2-1) is a free module on $H^*(X; \mathbb{Q})$ over a positively and evenly graded polynomial ring. Moreover, it follows from Lemma 4.5 and Theorem 2.2 that as a ring $H^*(E)$ is a complete intersection over $H^*(\text{BhAut}_0(X); \mathbb{Q})$ (see proof of Theorem 4.7). Here, by a *complete intersection* over a commutative ring R we mean a finite R -algebra S that is isomorphic to $R[x_1, \dots, x_n]/(f_1, \dots, f_n)$ for $f_1, \dots, f_n \in R[x_1, \dots, x_n]$ (see [29]).

For a finite R -algebra S , we can define the trace of an endomorphism $\text{Hom}_R(S, S) \cong S \otimes_R \text{Hom}_R(S, R)$ via the evaluation $S \otimes_R \text{Hom}_R(S, R) \rightarrow R$. In particular, we can associate to any $s \in S$ the trace of the endomorphism $s \cdot - : S \rightarrow S$ and we obtain an element $\text{Tr}_{S/R} \in \text{Hom}_R(S, R)$. We will use the following result about complete intersections.

Proposition 4.6 [29] *Let R be a commutative ring and $f_1, \dots, f_n \in R[x_1, \dots, x_n]$ for a nonnegative integer n . Assume that $S = R[x_1, \dots, x_n]/(f_1, \dots, f_n)$ is a finite R -algebra. Then*

- (i) S is a projective R -module;²
- (ii) $\text{Hom}_R(S, R)$ is a free of rank 1 as an S -module;
- (iii) there is a generator λ of $\text{Hom}_R(S, R)$ as an S -module such that $\text{Tr}_{S/R} = \det(\partial f_i / \partial x_j) \cdot \lambda$.

We recognise (iii) as an analogue for complete intersections of the relation between fibre integration, the fibrewise Euler class and the Becker–Gottlieb transfer $\tau_{\pi} : \Sigma^{\infty} B_+ \rightarrow \Sigma^{\infty} E_+$ for fibrations with Poincaré fibres. The transfer map induces a map on cohomology $\text{trf}_{\pi}^* : H^*(E) \rightarrow H^*(B)$ and if the fibre is a Poincaré complex one can show that

$$\text{trf}_{\pi}^*(x) = \pi_!(e^{\text{fw}}(\pi) \cdot x)$$

for all $x \in H^*(E)$. If we consider the universal 1-connected fibration (2-1) of a positively rationally elliptic space that satisfies the Halperin conjecture, its algebraic model $R \rightarrow S$ is equivalent to a complete intersection over a polynomial ring and we can identify the Becker–Gottlieb transfer and fibre integration with the trace $\text{Tr}_{S/R}$ and λ , respectively, which therefore leads to an identification of the fibrewise Euler class.

²In our applications, R is a positively graded polynomial ring so that S is in fact free.

Theorem 4.7 *Let X be a simply connected, oriented Poincaré duality space that is positively rationally elliptic and satisfies the Halperin conjecture. Then a cdga model of (2-1) is given by a complete intersection $S = \mathbb{R}[x_1, \dots, x_n]/(f_1, \dots, f_n)$ over a polynomial ring \mathbb{R} and the fibrewise Euler class is given by*

$$(4-8) \quad e^{\text{fw}}(\pi) = \det\left(\frac{\partial f_i}{\partial x_j}\right) \in S.$$

Remark 4.8 The first part of the theorem improves a result by Kuribayashi [20, Theorem 1.1] where he showed that the cohomology ring of the total space of the universal 1-connected fibration (2-1) for positively rationally elliptic spaces that satisfy the Halperin conjecture is a complete intersection over the cohomology of the base.

Proof Let $\Lambda = (\mathbb{Q}[x_1, \dots, x_n] \otimes \Lambda(y_1, \dots, y_n), d)$ be a pure Sullivan model of X with differential $d(y_i) = \tilde{f}_i \in \mathbb{Q}[x_1, \dots, x_n]$ where $\tilde{f}_1, \dots, \tilde{f}_n$ is a regular sequence. The cohomology ring $H(\Lambda)$ is a Poincaré duality algebra and $\det(\partial \tilde{f}_i / \partial x_j) \in H^*(X)$ is a generator in top degree [24; 28], so that the orientation is defined by $\varepsilon_X(\det(\partial \tilde{f}_i / \partial x_j)) := \chi(X)$. Now let $\mathfrak{a} \subset \text{Der}_{*,-1}^+(\Lambda)$ be an abelian dg Lie subalgebra quasi-isomorphic to $\text{Der}^+(\Lambda)$ from Lemma 4.5. Then a model of the universal 1-connected fibration is given by

$$\mathbb{R} := \mathcal{C}_{\text{CE}}(\mathfrak{a}) \rightarrow \mathcal{C}_{\text{CE}}(\mathfrak{a}; \Lambda) \cong (\mathbb{R} \otimes \Lambda(x_i, y_i), D)$$

by Theorem 2.2. And because $\mathfrak{a} \subset \text{Der}_{*,-1}^+(\Lambda)$, we see that $D(x_i) = 0$ and $D(y_i) = f_i \in \mathbb{R}[x_1, \dots, x_n]$ are polynomials that satisfy $\tilde{f}_i = f_i \otimes_{\mathbb{R}} 1 \in \mathbb{R}[x_1, \dots, x_n] \otimes_{\mathbb{R}} \mathbb{Q}$. This implies $S := \mathbb{R}[x_1, \dots, x_n]/(f_1, \dots, f_n)$ is a complete intersection over \mathbb{R} and the projection map $\mathcal{C}_{\text{CE}}(\mathfrak{a}; \Lambda) \rightarrow S$ is a quasi-isomorphism.

One can show that the trace $\text{Tr}_{S/\mathbb{R}}$ and the transfer tr_{π}^* agree using the general theory in [9] or by directly checking that

$$(4-9) \quad \text{Tr}_{S/\mathbb{R}}(x) = \pi_!(e^{\text{fw}}(\pi) \cdot x),$$

as in [27, Lemma 2.3] using that S is a finite free \mathbb{R} -algebra. Now consider λ from Proposition 4.6. Then $\lambda(\det(\partial f_i / \partial x_j)) = \text{Tr}_{S/\mathbb{R}}(1) = \chi(X)$ by (4-9) which agrees with $\pi_!(\det(\partial f_i / \partial x_j)) = \varepsilon_X(\det(\partial \tilde{f}_i / \partial x_j))$. Therefore, it follows from degree reasons that $\pi_! = \lambda$ and we obtain (4-8) from the identity

$$\det(\partial f_i / \partial x_j) \cdot \pi_! = \det(\partial f_i / \partial x_j) \cdot \lambda = \text{Tr}_{E/B} = \text{tr}_{\pi}^* = e^{\text{fw}}(\pi) \cdot \pi_!$$

since $\text{Hom}_{\mathbb{R}}(S, \mathbb{R})$ is a free S -module by (ii). □

5 Computations

Before we come to the computation of the Euler ring, we record a few general facts. First we observe that in some cases it is sufficient to compute the Euler ring of the universal 1-connected fibration $E_0(X) \subset H^*(\text{BhAut}_0(X))$.

Lemma 5.1 *If $\pi_0(\text{hAut}^+(X))$ is finite, then $i : \text{BhAut}_0(X) \rightarrow \text{BhAut}^+(X)$ induces an isomorphism $E^*(X) \cong E_0^*(X)$.*

Proof By naturality of the fibrewise Euler class, $i : \text{BhAut}_0(X) \rightarrow \text{BhAut}^+(X)$ induces a surjection of Euler rings $E^*(X) \twoheadrightarrow E_0^*(X)$. It follows from the spectral sequence associated to the fibre sequence

$$\text{BhAut}_0(X) \rightarrow \text{BhAut}^+(X) \rightarrow \text{B}\pi_0(\text{hAut}^+(X))$$

that $H^*(\text{BhAut}(X); \mathbb{Q}) \cong H^*(\text{BhAut}_0(X); \mathbb{Q})^{\pi_0(\text{hAut}^+(X))}$ if $\pi_0(\text{hAut}^+(X))$ is finite. In particular, i induces an injection on rational cohomology. \square

For a positively rationally elliptic space there is a simple criterion to check if it has finitely many homotopy automorphisms.

Proposition 5.2 *Let X be a simply connected Poincaré duality space that is positively rationally elliptic space. If the cohomology ring $H^*(X; \mathbb{Q})$ has finitely many orientation-preserving automorphisms (i.e., algebra automorphisms that preserve the orientation $\varepsilon_X \in H^d(X; \mathbb{Q})^\vee$), then $\pi_0(\text{hAut}^+(X))$ is finite.*

Proof The statement follows if the kernel of $\mathcal{E}(X) \rightarrow \text{Aut}(H^*(X; \mathbb{Q}), \varepsilon_X)$ is finite. Rationalisation induces a map $\mathcal{E}(X) \rightarrow \mathcal{E}(X_{\mathbb{Q}})$ which has finite kernel [32, Theorem 10.2], so that it suffices to prove that the subgroup of $\mathcal{E}(X_{\mathbb{Q}})$ of homotopy equivalences that induce the identity on cohomology is trivial. By [32, 10.3], homotopy automorphism of $X_{\mathbb{Q}}$ are the same as homotopy classes of automorphisms of a minimal Sullivan model.

For a positively rationally elliptic space we can choose $\Lambda = (\Lambda(x_i, y_i)_{i=1, \dots, n}, d = \sum_i f_i \partial y_i)$ as a model where $f_1, \dots, f_n \in \Lambda(x_i)_{i=1, \dots, n}$ is a regular sequence. Let $\phi : \Lambda \rightarrow \Lambda$ be an automorphism so that $H(\phi) = \text{Id}$. Then $[x_i - \phi(x_i)] = 0$ and we can pick a coboundary $\xi_i \in \Lambda_{>0}$ satisfying $d\xi_i = x_i - \phi(x_i) \in (f_1, \dots, f_k) + \Lambda_{>0}$. We want to define a homotopy $H : \Lambda \rightarrow \Lambda \otimes \Lambda(t, dt)$ by setting $H(x_i) = \phi(x_i) + t(x_i - \phi(x_i)) - \xi_i dt$. Then $dH(x_i) = 0 = Hd(x_i)$ and $\varepsilon_1 \circ H(x_i) = x_i$ and $\varepsilon_0 \circ H(x_i) = \phi(x_i)$. It remains to define $H(y_i)$, i.e., we have to find a coboundary for $f_i(H(x_1), \dots, H(x_n)) \in \Lambda \otimes \Lambda(t, dt)$. Observe that $\Lambda \otimes \Lambda(t, dt)$ is a positively elliptic Sullivan algebra as well since f_1, \dots, f_n, t is a regular sequence in $\mathbb{Q}[x_1, \dots, x_n, t]$, and $f_i(H(x_1), \dots, H(x_n)) - d\phi(y_i) \in (\Lambda \otimes \Lambda^+(t, dt)) \oplus (\Lambda \otimes \Lambda(t, dt))_{>0}$. Since both summands are acyclic, there is $\zeta_i \in \Lambda \otimes \Lambda(t, dt)$ so that $d\zeta_i = f_i(H(x_1), \dots, H(x_n)) - d\phi(y_i)$, and we can set $H(y_i) := \zeta_i + \phi(y_i)$. This shows ϕ is homotopic to a map which is the identity on $\Lambda(x_i)$.

Given an automorphism $\psi : \Lambda \rightarrow \Lambda$ with $\psi(x_i) = x_i$, then $d\psi(y_i) = \psi(f_i) = f_i = d(y_i)$ and hence $y_i - \psi(y_i)$ is a cocycle in $\Lambda_{>0}$. Let ξ_i be a coboundary $d\xi_i = y_i - \psi(y_i)$. Then $\psi \simeq_H \text{Id}_\Lambda$ via $H(x_i) := x_i$ and $H(y_i) := \psi(y_i) + t(y_i - \psi(y_i)) - \xi_i dt$. \square

Corollary 5.3 *Let X be a simply connected Poincaré duality space with $H^*(X; \mathbb{Q}) \cong \mathbb{Q}[x]/(x^{n+1})$ for some n and $|x|$ even. Then $\pi_0(\text{hAut}^+(X))$ is finite.*

Proof For any choice of orientation of X , the group $\text{Aut}(H^*(X; \mathbb{Q}), \varepsilon_X)$ is trivial if n is odd and $\mathbb{Z}/2$ if n is even. Hence, $\pi_0(\text{hAut}^+(X))$ is finite by Proposition 5.2 \square

Finally, in some cases the Euler ring is finitely generated so that the computation of the Euler ring amounts to computing the ideal of relations among a generating set, and which simplifies some computations below.

Proposition 5.4 *Let X be a Poincaré duality space so that $H^*(X)$ is concentrated in even degrees and let $n = \dim H^*(X; \mathbb{Q})$. Then $E^*(X)$ is generated by $\kappa_1, \dots, \kappa_{n-2}, \kappa_n$.*

Proof This follows directly from the tools developed in [27]. Applying [27, Corollary 2.7] for $x = e^{fw}(\pi)$, the corresponding monic polynomial $\rho_x(z) \in H^*(\text{BhAut}^+(X); \mathbb{Q})[z]$ of degree n is given by

$$\rho_x(z) = \frac{(-1)^n}{n!} \sum_{\sigma \in \Sigma_{n+1}} \text{sgn}(\sigma) \kappa_{l(\gamma_2)} \cdots \kappa_{l(\gamma_q(\sigma))} \cdot z^{l(\gamma_1)-1},$$

where $\sigma = \gamma_1 \cdots \gamma_q(\sigma)$ is the cycle decomposition of σ , $l(\gamma_i)$ denotes the length of γ_i and 1 is contained in the support of γ_1 . Then [27, Corollary 2.7] implies that $\rho_x(e^{fw}(\pi)) = 0 \in H^*(E)$ and by fibre integrating $0 = e^{fw}(\pi)^i \cdot \rho(e^{fw}(\pi))$, one can decompose κ_{n+i} in terms of κ -classes of lower degree for $i \geq 0$; except when $i = 1$ as $\rho_x(z)$ has a constant term $-\kappa_n/n$ which cancels the leading term $\pi_!(e^{fw}(\pi)^{n+1})$ in $\pi_!(e^{fw}(\pi)\rho_x(e^{fw}(\pi)))$. □

5.1 The Euler ring of even spheres

Proposition 5.5 *The Euler ring of an even-dimensional sphere is $E^*(S^{2n}) \cong \mathbb{Q}[\kappa_2]$ where $\kappa_2^k = 2^{k-1}\kappa_{2k}$ and all odd κ_{2i+1} vanish.*

Proof We have seen in Example 4.4 that $e^{fw}(\pi) = 2x \in \Lambda(x, y, z_{4n})$ and in Example 3.4 that fibre integration is given by $\Pi(x^{2k}) = 0$ and $\Pi(x^{2k+1}) = (-1)^k z_{4n}^k$. Thus $\kappa_{2k} = 2^{1-k}(-2^3 z_{4n})^k = 2^{1-k}\kappa_{2k}$. Since $\pi_0(\text{hAut}^+(S^{2n}))$ is trivial, the 1-connected universal fibration is the universal fibration and the result follows. □

5.2 The Euler ring of complex projective space

A minimal model of $\mathbb{C}P^n$ is

$$P_n := (\Lambda(x, y), |x| = 2, |y| = 2n + 1, d = x^{n+1}\partial/\partial y)$$

with orientation $\varepsilon_{\mathbb{C}P^n}(x^n) = 1$ induced by integral Poincaré duality, and we apply Theorem 2.2 to compute a model for the minimal 1-connected $\mathbb{C}P^n$ -fibration.

Proposition 5.6 *A cdga model of the universal 1-connected $\mathbb{C}P^n$ -fibration is given by*

$$(5-1) \quad B_n := (\mathbb{Q}[x_2, \dots, x_{n+1}], |x_i| = 2i, d = 0)$$

$$\xrightarrow{\pi} E_n := \left(B_n[x] / \left(x^{n+1} + \sum_{i=2}^{n+1} x_i \cdot x^{n+1-i} \right), |x| = 2, d = 0 \right),$$

and the fibrewise Euler class in E_n is represented by

$$(5-2) \quad e^{fw}(\pi) = (n + 1) \cdot x^n + \sum_{i=2}^n (n + 1 - i) \cdot x_i \cdot x^{n-i} \in E_n.$$

Proof Note that $\text{Der}^+(P_n)$ has a (vector space) basis given by $\theta_i := x^{n+1-i} \partial/\partial y$ for $i = 1, \dots, n + 1$ of degree $2i - 1$ and $\eta := \partial/\partial x$ of degree 2. The only nontrivial differential on the derivation Lie algebra is given by $[d, \eta] = -(n + 1)\theta_1$. Since $\mathbb{C}P^n$ satisfies the Halperin conjecture, it follows from Lemma 4.5 that there exists a quasi-isomorphic abelian dg Lie subalgebra $\mathfrak{a}_n \subset \text{Der}^+(P_n)$ with trivial differential, which in this case is easy to identify as $\mathfrak{a}_n := \mathbb{Q}\{\theta_2, \dots, \theta_{n+1}\}$. The statement follows directly from Theorem 4.7 for this choice of \mathfrak{a}_n . \square

Theorem 5.7 *The Euler ring of complex projective space is $E^*(\mathbb{C}P^n) \cong \mathbb{Q}[\kappa_1, \dots, \kappa_{n-1}, \kappa_{n+1}]$.*

Proof By Corollary 5.3, $\pi_0(\text{hAut}^+(\mathbb{C}P^n))$ is finite (in fact one can easily see $\pi_0(\text{hAut}(\mathbb{C}P^n)) \cong \mathbb{Z}/2$), and so $E^*(\mathbb{C}P^n) \cong E_0^*(\mathbb{C}P^n)$ by Lemma 5.1. The Euler ring is generated by $\kappa_1, \dots, \kappa_{n-1}, \kappa_{n+1}$ by Proposition 5.4, so it remains to show they are algebraically independent which follows if $\det(\partial\kappa_i/\partial x_j)$ is nonzero.

It turns out that the polynomials representing the κ_i are quite complicated so that it is difficult to give a closed formula for the determinant of the Jacobian. We will resolve this issue by focussing on the terms containing x_{n+1} because it is the variable of the highest degree and it is not contained in $e^{\text{fw}}(\pi)$ so that it only arises through fibre integrating x^k for $k > n$. It will be sufficient to consider elements modulo decomposables, i.e., for $x, y \in B_n$ then $x \sim y$ if $x - y \in (B_n^+)^2$. We will start with the following observation about fibre integration.

Observation 1 *If $k = 2, \dots, n + 1$ then $\pi_!(x^{n+k}) \sim -x_k \in B_n$.*

Proof Rewriting x^{n+2} in terms of the module basis $\{1, x, \dots, x^n\}$, one can see that $\pi_!(x^{n+2}) = x_2$. Then $\pi_!(x^{n+k}) = \pi_!(x^{n+1} \cdot x^{k-1}) = -\sum_{i=2}^{n+1} x_i \cdot \pi_!(x^{n+k-i})$ and by induction over k , the only indecomposable contribution is for $i = k$. \square

Observation 2 *For $i = 1, \dots, n - 1$ the highest power of x_{n+1} in $\kappa_i(x_2, \dots, x_{n+1})$ is $i - 1$ and the coefficient $c_i \in \mathbb{Q}[x_2, \dots, x_n]$ of x_{n+1}^{i-1} satisfies $c_i \sim (-1)^i i (n + 1)^i (n - i) \cdot x_{n+1-i}$.*

Proof It follows from degree considerations that the highest power of x_{n+1} is $i - 1$ and c_i equals $A \cdot x_{n+1-i} + \text{decomposables}$. It remains to determine the coefficient A . When expanding $e^{\text{fw}}(\pi)$ using (5-2), the only relevant contributions are

$$\begin{aligned} (n + 1)^{i+1} x^{n(i+1)} - (i + 1)(n + 1 - (n + 1 - i)) x_{n+1-i} x^{n-(n+1-i)} \cdot (n + 1)^i x^{ni} \\ = (n + 1)^{i+1} (x^{n+1})^{i-1} \cdot x^{2n-i+1} - i(i + 1)(n + 1)^i x_{n+1-i} \cdot (x^{n+1})^{i-1} \cdot x^n. \end{aligned}$$

Now rewrite $x^{n+1} = \sum_{i=2}^{n+1} x_i x^{n+1-i}$ and collect all terms containing x_{n+1}^{i-1} and $x_{n+1}^{i-2} x_{n+1-i}$ (we can ignore the rest because it cannot contribute to A) to get

$$(n + 1)^{i+1} (x_{n+1}^{i-1} + (i - 1)x_{n+1}^{i-2} x_{n+1-i} \cdot x^i) \cdot x^{2n-i+1} - i(i + 1)(n + 1)^i x_{n+1-i} \cdot x_{n+1}^{i-1} \cdot x^n.$$

The statement follows by fibre integrating and discarding decomposables as in Observation 1 above. \square

Observation 3 *The highest contribution of x_{n+1} in κ_{n+1} is $(-1)^n (n + 1)^{n+2} \cdot x_{n+1}^n$.*

Proof The expression for $e^{\text{fw}}(\pi)^{n+2}$ contains $(n + 1)^{n+2} \cdot x^{n(n+2)} = (n + 1)^{n+2} \cdot (x^{n+1})^n \cdot x^n$. This is the only summand that fibre integrates to a multiple of x_{n+1}^n , i.e., $\kappa_{n+1} = (-1)^n (n + 1)^{n+2} \cdot x_{n+1}^n + \dots$ where we can ignore all other terms. \square

We can now analyse $\det(\partial\kappa_i/\partial x_j)$ which contains the summand

$$\frac{\partial\kappa_1}{\partial x_n} \cdot \frac{\partial\kappa_2}{\partial x_{n-1}} \dots \frac{\partial\kappa_{n-1}}{\partial x_2} \cdot \frac{\partial\kappa_{n+1}}{\partial x_{n+1}}.$$

It follows from Observations 2 and 3 that the above expression contains $C \cdot x_{n+1}^N$, where C is a nonzero constant and $N = \frac{1}{2}n(n - 1)$. This is the only possible way to get a monomial in $\det(\frac{\partial\kappa_i}{\partial x_j})$ that contains only x_{n+1} . Hence, the determinant does not vanish and the generating set $\kappa_1, \dots, \kappa_{n-1}, \kappa_{n+1}$ is algebraically independent. \square

Remark 5.8 Theorem 5.7 has been studied in the smooth case for $n = 2$ in [27] by studying the natural smooth 2-torus action on $\mathbb{C}P^2$ and that implies our result in this case as well. This has been extended by Dexter Chua to $n \leq 4$, but for large n the algebra becomes intractable.

5.3 The Euler ring of products of odd spheres

The main result of this section is the computation of $E_0^*(X)$ for a simply connected Poincaré duality space X that is rationally equivalent to a product of odd spheres $\prod_{i=1}^{2n} S^{2k_i+1}$ for some $n, k_i > 0$.

Theorem 5.9 *Let X be a simply connected Poincaré duality space that is rationally equivalent to a product of odd spheres $\prod_{i=1}^{2n} S^{2k_i+1}$ for some $n, k_i > 0$. Then the fibrewise Euler class of the universal 1-connected X -fibration $E \rightarrow \text{BhAut}_0(X)$ is trivial and hence $E_0^*(X) \cong \mathbb{Q}$.*

Proof The minimal model of X is given by an exterior algebra $A_X = (\Lambda(x_i)_{1 \leq i \leq 2n}, d = 0)$ and by Theorem 2.2 the model of the universal 1-connected fibration is given by

$$(5-3) \quad B_X := \mathcal{C}_{\text{CE}}^*(\text{Der}^+(A_X); \mathbb{Q}) \rightarrow E_X := \mathcal{C}_{\text{CE}}^*(\text{Der}^+(A_X); A_X) \cong (B_X \otimes A_X, D).$$

Observe that $\mathcal{C}_{\text{CE}}^*(\text{Der}^+(A_X); A_X)$ is a finitely generated B_X -module as the minimal Sullivan model is finite dimensional. Let $\varepsilon_X : A_X \rightarrow \mathbb{Q}$ denote a rational orientation, then $B_X \otimes \varepsilon_X : (B_X \otimes A_X, D) \rightarrow B_X$ is the only module homomorphism that restricts to ε_X on the fibre and thus is a representative of fibre integration by Proposition 3.2.

Moreover, $\text{Hom}_{B_X}(E_X, B_X)$ is a minimal semifree module and therefore the quasi-isomorphism $\overline{\Pi} : E_X[-d] \rightarrow \text{Hom}_{B_X}(E_X, B_X)$ from Proposition 4.1 is in fact an isomorphism by uniqueness of minimal free resolutions [11, Example 8, Chapter 6] (similarly for $\overline{\Pi} \otimes \overline{\Pi}$). Hence, the algebraic Umkehr map is given by

$$\Delta_! : E_X[-|F|] \xrightarrow{\cong} \overline{\Pi} \rightarrow \text{Hom}_{B_X}(E_X, B_X) \xrightarrow{\Delta^*} \text{Hom}_{B_X}(E_X \otimes_{B_X} E_X, B_X) \xrightarrow{\cong} \overline{(\Pi \otimes \Pi)}^{-1} \rightarrow E_X \otimes_{B_X} E_X[-2|F|].$$

The composition of $\overline{\Pi}$ with the vector space isomorphism $\text{Hom}_{B_X}(E_X, B_X) \cong (A_X)^\vee \otimes B_X$ is given by $\bar{\varepsilon}_X \otimes \text{Id}_{B_X}$ where $\bar{\varepsilon}_X : A_X \rightarrow (A_X)^\vee$ is the adjoint of $\varepsilon_X : A_X \otimes A_X \rightarrow \mathbb{Q}$. The same statement holds for

$\overline{\Pi} \otimes \overline{\Pi}$ with the appropriate choice of orientation on $X \times X$ given by $\varepsilon_{X \times X} := \varepsilon_X \otimes \varepsilon_X : (A_X \otimes A_X)^{\otimes 2} \rightarrow \mathbb{Q}$. Note that $\Delta^* \overline{\Pi}(1)$ is contained in $(A_X \otimes A_X)^\vee \otimes 1$ so that $(\overline{\Pi} \otimes \overline{\Pi})^{-1} \Delta^* \overline{\Pi}(1)$ is in $A_X \otimes A_X \otimes 1 \subset E_X \otimes_{B_X} E_X$. A direct computation shows that

$$\Delta_!(1) = \sum_{S_1 \sqcup S_2 = F} \pm x_{S_1} \otimes x_{S_2} \in E_X \otimes_{B_X} E_X$$

for some signs that can be worked out. Hence, the fibrewise Euler class is

$$e^{\text{fw}}(\pi) = \Delta^* \circ \Delta_!(1) = \sum_{S_1 \sqcup S_2 = F} \pm x_F$$

and since $\Pi(e^{\text{fw}}(\pi)) = \chi(X) = 0$, the summands must cancel. □

One can easily see that the group of homotopy self-equivalences $\pi_0(\text{hAut}^+(X))$ is not finite in most cases, and so we cannot infer that the full Euler ring is trivial in general except in the two cases below.

Theorem 5.10 *Let X be either rationally equivalent to $(S^{2k+1})^{\times n}$ or a finite CW complex rationally equivalent to $S^{2k+1} \times S^{2l+1}$ for $1 < k < l$ and n even. Then $E^*(X) = \mathbb{Q}$.*

Proof We start with the proof of the second case. By [32, Theorem 10.3] the group $\pi_0(\text{hAut}^+(X))$ is commensurable with an arithmetic subgroup of the homotopy classes automorphisms of A_X . If $X \simeq_{\mathbb{Q}} S^{2k+1} \times S^{2l+1}$ then the group of automorphisms of A_X modulo homotopy is $\mathbb{Q}^\times \times \mathbb{Q}^\times$ and the arithmetic subgroups of this linear algebraic group are finite and hence by commensurability so is $\pi_0(\text{hAut}^+(X))$. Hence, $E^*(X) \cong E_0^*(X) = \mathbb{Q}$ by Lemma 5.1 and Theorem 5.9.

We need to introduce some notation in the first case. Let $\pi : E \rightarrow \text{BhAut}^+(X)$ denote the universal orientated X -fibration and $\pi_0 : E_0 \rightarrow \text{BhAut}_0(X)$ the universal 1-connected X -fibration. The cdga model of the total space E_0 is a free algebra on generators $x_1, \dots, x_n, y^1, \dots, y^n$ with differential $D(x_i) = y^i$ and therefore $H(E_0) = \mathbb{Q}$. It is well known that E is homotopy equivalent to the classifying space of pointed homotopy automorphisms $\text{BhAut}_*^+(X)$ and there is a fibration sequence $E_0 \rightarrow E \xrightarrow{H} \text{B}\pi_0(\text{hAut}_*^+(X))$ which induces an isomorphism $H^*(E) \cong H^*(\text{B}\pi_0(\text{hAut}_*^+(X)))$ since the rational cohomology of E_0 is trivial. In particular, $e^{\text{fw}}(\pi) = H^*e$ for some $e \in H^{n \cdot (2k+1)}(\text{B}\pi_0(\text{hAut}_*^+(X)))$. It follows from the commutative diagram

$$\begin{array}{ccc} E & \xrightarrow[\simeq_{\mathbb{Q}}]{H} & \text{B}\pi_0(\text{hAut}_*^+(X)) \\ \downarrow \pi & & \parallel \\ \text{BhAut}^+(X) & \xrightarrow{h} & \text{B}\pi_0(\text{hAut}^+(X)) \end{array}$$

that the fibrewise Euler class $e^{\text{fw}}(\pi) = H^*e = \pi^*h^*e$ is pulled back from the base, and therefore that the fibre integrals $\pi_!(\pi^*(h^*e)^k) = (h^*e)^k \cdot \pi_!(1) = 0$ all vanish. □

Remark 5.11 There are more cases when the group of homotopy self-equivalences $\mathcal{E}(X)$ is finite for a space X that is rationally equivalent to a product of odd-dimensional spheres, and the second case in Theorem 5.10 is merely one of the simplest to establish. But we expect that the Euler ring vanishes in

general even if $\mathcal{E}(X)$ is not finite using recent results in [4], where Berglund and Zeman have given a rational description of $\text{BhAut}(X)$ via a fibre sequence

$$\text{BhAut}_u(X) \rightarrow \text{BhAut}(X) \rightarrow \text{B}\Gamma(X),$$

where $\Gamma(X)$ is a certain arithmetic group and $\text{BhAut}_u(X)$ is the classifying space of normal unipotent X -fibrations. They provide $\Gamma(X)$ -equivariant models for $\text{BhAut}_u(X)$ so that one can obtain an algebraic model for $\text{BhAut}(X)$. We expect that one can extend the results in this paper using their results to study Euler rings in more generality, and in particular that the Euler ring $E^*(X)$ vanishes for X a product of odd spheres in general.

Acknowledgements

This paper is part of my PhD project and I would like to thank my supervisor Oscar Randal-Williams for many enlightening discussions and his help and patience. I would also like to thank Alexander Berglund for many useful discussions and for his help with the revision of this article, and Andrey Lazarev for pointing out modern references for models of the universal fibration. For the revision of this article I have been supported by the Knut and Alice Wallenberg foundation through grant 2019.0519.

References

- [1] **A Berglund**, *Rational models for automorphisms of fiber bundles*, Doc. Math. 25 (2020) 239–265 MR
- [2] **A Berglund**, *Characteristic classes for families of bundles*, Selecta Math. (N.S.) 28:3 (2022) art. id. 51 MR
- [3] **A Berglund, I Madsen**, *Rational homotopy theory of automorphisms of manifolds*, Acta Math. 224:1 (2020) 67–185 MR
- [4] **A Berglund, T Zeman**, *Algebraic models for classifying spaces of fibrations*, Geom. Topol. 29:7 (2025) 3567–3634 MR
- [5] **J M Boardman**, *Conditionally convergent spectral sequences*, from “Homotopy invariant algebraic structures” (Baltimore, MD, 1998) (J-P Meyer, J Morava, W S Wilson, editors), Contemp. Math. 239, Amer. Math. Soc., Providence, RI (1999) 49–84 MR
- [6] **A Borel, F Hirzebruch**, *Characteristic classes and homogeneous spaces, I*, Amer. J. Math. 80 (1958) 458–538 MR
- [7] **V Braunack-Mayer**, *Strict algebraic models for rational parametrised spectra, I*, Algebr. Geom. Topol. 21:2 (2021) 917–1019 MR
- [8] **V Braunack-Mayer**, *Strict algebraic models for rational parametrised spectra, I*, Algebr. Geom. Topol. 21:2 (2021) 917–1019 MR
- [9] **A Dold, D Puppe**, *Duality, trace, and transfer*, from “Proceedings of the International Conference on Geometric Topology” (Warsaw, 1978) (K Borsuk, A Kirkor, editors), PWN, Warsaw (1980) 81–102 MR
- [10] **A Dress**, *Zur Spectralsequenz von Faserungen*, Invent. Math. 3 (1967) 172–178 MR
- [11] **Y Félix, S Halperin, J-C Thomas**, *Rational homotopy theory*, Graduate Texts in Mathematics 205, Springer (2001) MR
- [12] **Y Félix, A Murillo, D Tanré**, *Fibrewise stable rational homotopy*, J. Topol. 3:4 (2010) 743–758 MR
- [13] **Y Félix, J-C Thomas**, *String topology on Gorenstein spaces*, Math. Ann. 345:2 (2009) 417–452 MR
- [14] **S Galatius, I Grigoriev, O Randal-Williams**, *Tautological rings for high-dimensional manifolds*, Compos. Math. 153:4 (2017) 851–866 MR
- [15] **DH Gottlieb**, *Poincaré duality and fibrations*, Proc. Amer. Math. Soc. 76:1 (1979) 148–150 MR
- [16] **I Grigoriev**, *Relations among characteristic classes of manifold bundles*, Geom. Topol. 21:4 (2017) 2015–2048 MR
- [17] **P-P Grivel**, *Formes différentielles et suites spectrales*, Ann. Inst. Fourier (Grenoble) 29:3 (1979) 17–37 MR

- [18] **S Halperin**, *Lectures on minimal models*, Mém. Soc. Math. France 9-10, Soc. Math. France, Paris (1983) MR
- [19] **F Hebestreit, M Land, W Lück, O Randal-Williams**, *A vanishing theorem for tautological classes of aspherical manifolds*, Geom. Topol. 25:1 (2021) 47–110 MR
- [20] **K Kuribayashi**, *On the rational cohomology of the total space of the universal fibration with an elliptic fibre*, from “Homotopy theory of function spaces and related topics” (Y Félix, G Lupton, S B Smith, editors), Contemp. Math. 519, Amer. Math. Soc., Providence, RI (2010) 165–179 MR
- [21] **A Lazarev**, *Models for classifying spaces and derived deformation theory*, Proc. Lond. Math. Soc. (3) 109:1 (2014) 40–64 MR
- [22] **W Meier**, *Rational universal fibrations and flag manifolds*, Math. Ann. 258:3 (1981/82) 329–340 MR
- [23] **D Mumford**, *Towards an enumerative geometry of the moduli space of curves*, from “Arithmetic and geometry, II” (M Artin, J Tate, editors), Progr. Math. 36, Birkhäuser, Boston, MA (1983) 271–328 MR
- [24] **A Murillo**, *The top cohomology class of certain spaces*, J. Pure Appl. Algebra 84:2 (1993) 209–214 MR
- [25] **N Prigge**, *On tautological classes of fibre bundles and self-embedding calculus*, PhD thesis, University of Cambridge (2020) Available at <https://doi.org/10.17863/CAM.65008>
- [26] **N Prigge**, *Tautological rings of fake quaternionic spaces*, Selecta Math. (N.S.) 31:1 (2025) art. id. 1 MR
- [27] **O Randal-Williams**, *Some phenomena in tautological rings of manifolds*, Selecta Math. (N.S.) 24:4 (2018) 3835–3873 MR
- [28] **H Shiga, M Tezuka**, *Rational fibrations, homogeneous spaces with positive Euler characteristics and Jacobians*, Ann. Inst. Fourier (Grenoble) 37:1 (1987) 81–106 MR
- [29] **B de Smit, H W Lenstra, Jr.**, *Finite complete intersection algebras and the completeness radical*, J. Algebra 196:2 (1997) 520–531 MR
- [30] **J Stasheff**, *A classification theorem for fibre spaces*, Topology 2 (1963) 239–246 MR
- [31] **R Stoll**, *The stable cohomology of self-equivalences of connected sums of products of spheres*, Forum Math. Sigma 12 (2024) art. id. e1 MR
- [32] **D Sullivan**, *Infinitesimal computations in topology*, Inst. Hautes Études Sci. Publ. Math. :47 (1977) 269–331 MR
- [33] **D Tanré**, *Homotopie rationnelle: modèles de Chen, Quillen, Sullivan*, Lecture Notes in Mathematics 1025, Springer (1983) MR
- [34] **J-C Thomas**, *Rational homotopy of Serre fibrations*, Ann. Inst. Fourier (Grenoble) 31:3 (1981) 71–90 MR
- [35] **C T C Wall**, *Poincaré complexes, I*, Ann. of Math. (2) 86 (1967) 213–245 MR

NILS PRIGGE nils.prigge@gmail.com

Matematiska Institutionen, Stockholm University, Stockholm, Sweden

Received: October 11, 2024 Revised: April 9, 2025

ALGEBRAIC & GEOMETRIC TOPOLOGY

msp.org/agt

EDITORS

PRINCIPAL ACADEMIC EDITORS

John Etnyre
etnyre@math.gatech.edu
Georgia Institute of Technology

Vesna Stojanoska
vesna@illinois.edu
University of Illinois at Urbana-Champaign

BOARD OF EDITORS

Julie Bergner	University of Virginia jeb2md@eservices.virginia.edu	Daniel Isaksen	Wayne State University isaksen@math.wayne.edu
Steven Boyer	Université du Québec à Montréal cohf@math.rochester.edu	Thomas Koberda	University of Virginia thomas.koberda@virginia.edu
Tara E Brendle	University of Glasgow tara.brendle@glasgow.ac.uk	Markus Land	JGU Mainz mland@uni-mainz.de
Indira Chatterji	CNRS & Univ. Côte d'Azur (Nice) indira.chatterji@math.cnrs.fr	Christine Lescop	Université Joseph Fourier lescop@ujf-grenoble.fr
Octav Cornea	Université de Montreal cornea@dms.umontreal.ca	Norihiko Minami	OCAMI (Osaka Central Adv. Math. Inst.) norihikominami@gmail.com
Alexander Dranishnikov	University of Florida dranish@math.ufl.edu	Andrés Navas	Universidad de Santiago de Chile andres.navas@usach.cl
Tobias Ekholm	Uppsala University, Sweden tobias.ekholm@math.uu.se	Jessica S Purcell	Monash University jessica.purcell@monash.edu
Mario Eudave-Muñoz	Univ. Nacional Autónoma de México mario@matem.unam.mx	Birgit Richter	Universität Hamburg birgit.richter@uni-hamburg.de
David Futer	Temple University dfuter@temple.edu	Jérôme Scherer	École Polytech. Féd. de Lausanne jerome.scherer@epfl.ch
John Greenlees	University of Warwick john.greenlees@warwick.ac.uk	Zoltán Szabó	Princeton University szabo@math.princeton.edu
Matthew Hedden	Michigan State University mhedden@math.msu.edu	Maggy Tomova	University of Iowa maggy-tomova@uiowa.edu
Kristen Hendricks	Rutgers University kristen.hendricks@rutgers.edu	Daniel T Wise	McGill University, Canada daniel.wise@mcgill.ca
Hans-Werner Henn	Université Louis Pasteur henn@math.u-strasbg.fr	Lior Yanovski	Hebrew University of Jerusalem lior.yanovski@gmail.com
Kathryn Hess	École Polytechnique Féd. de Lausanne kathryn.hess@epfl.ch		


See inside back cover or msp.org/agt for submission instructions.

The subscription price for 2026 is US \$795/year for the electronic version, and \$1170/year (+\$80, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Algebraic & Geometric Topology is indexed by Mathematical Reviews, Zentralblatt MATH, Current Mathematical Publications and the Science Citation Index.

Algebraic & Geometric Topology (ISSN 1472-2747 printed, 1472-2739 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004.

AGT peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<https://msp.org/>

© 2026 Mathematical Sciences Publishers

ALGEBRAIC & GEOMETRIC TOPOLOGY

Volume 26 Issue 4 (pages 1229–1596) 2026

The Deligne–Mumford operad as a trivialization of the circle action	1229
ALEXANDRU OANCEA and DMITRY VAINTROB	
On embeddings of 4-manifolds in codimension 2	1293
ABHIJEET GHANWAT and DISHANT M PANCHOLI	
Recollements and stratification	1321
JAY SHAH	
The guts of nearly fibered knots	1385
ZHENKUN LI and FAN YE	
Finiteness properties of some groups of piecewise projective homeomorphisms	1395
DANIEL S. FARLEY	
Skew-rack cocycle invariants of closed 3-manifolds	1451
TAKEFUMI NOSAKA	
The rational abelianization of the Chillingworth subgroup of the mapping class group of a surface	1465
RYOTARO KOSUGE	
Acyindrical hyperbolicity for Artin groups with a visual splitting	1507
RUTH CHARNEY, ALEXANDRE MARTIN and ROSE MORRIS-WRIGHT	
Equivariant preimage theory for G -maps	1529
THAÍS F M MONIS and PETER WONG	
Homotopy commutativity in quasitoric manifolds	1549
SHO HASUI, DAISUKE KISHIMOTO, YICHEN TONG and MITSUNOBU TSUTAYA	
Tautological rings of fibrations	1565
NILS PRIGGE	
On the analog category of finite groups	1585
BEN KNUDSEN and SHMUEL WEINBERGER	