

AG  
T

*Algebraic & Geometric  
Topology*

Volume 26 (2026)

**A note on the involutive invariants of splices**

KRISTEN HENDRICKS, MATTHEW STOFFREGEN AND IAN ZEMKE

# A note on the involutive invariants of splices

KRISTEN HENDRICKS, MATTHEW STOFFREGEN AND IAN ZEMKE

A natural family of potentially 2-torsion elements in the integer homology cobordism group consists of splices of knots with their mirrors. We show that such 3-manifolds have locally trivial involutive Floer homology. We show some related families of splices also have locally trivial involutive Floer homology. Our arguments show that many gauge-theoretic invariants also vanish on these 3-manifolds.

## 1 Introduction

The integer homology cobordism group  $\Theta_{\mathbb{Z}}^3$  is the group of oriented homology three-spheres up to the equivalence relation of homology cobordism. In 2013, C. Manolescu used a  $\text{Pin}(2)$ -equivariant version of Seiberg–Witten Floer homology to show that  $\Theta_{\mathbb{Z}}^3$  contains no element  $Y$  of order two whose Rokhlin invariant  $\mu(Y)$  is 1 [18], which due to previous work of Galewski and Stern [7] and Matumoto [19] was sufficient to disprove the remaining outstanding cases of the triangulation conjecture. It remains unknown whether  $\Theta_{\mathbb{Z}}^3$  has any torsion elements; in particular, whether it contains a torsion element of order two. In order to produce an element of order two, it suffices to exhibit an oriented integer homology sphere  $Y$  with an orientation-reversing diffeomorphism  $Y \cong -Y$  with the property that  $Y$  takes a nontrivial value under any invariant of homology cobordism.

Three-manifolds obtained by splicing knot complements have attracted attention as a potential source of examples of elements of order two in  $\Theta_{\mathbb{Z}}^3$ . If  $K_0 \subseteq Y_0$  and  $K_1 \subseteq Y_1$  are two knots, a *splice* of  $K_0$  and  $K_1$  is a 3-manifold obtained by gluing  $Y_0 \setminus \nu(K_0)$  and  $Y_1 \setminus \nu(K_1)$  using an orientation-reversing diffeomorphism  $\phi$  of their boundaries. (Some authors require  $\phi$  to swap the meridian with the Seifert longitude, but we consider more general diffeomorphisms  $\phi$ ).

In this note, we consider two natural constructions of splices which produce homology 3-spheres with orientation-reversing diffeomorphisms:

- (Type-1) splices of a knot  $K \subseteq Y$  with its mirror  $K \subseteq -Y$  such that there is a diffeomorphism of the splice which swaps  $Y \setminus \nu(K)$  and  $-Y \setminus \nu(K)$ ;
- (Type-2) splices of two knots  $K_0, K_1$  in  $S^3$  such that there is a diffeomorphism of the splice which fixes the subsets  $S^3 \setminus \nu(K_0)$  and  $S^3 \setminus \nu(K_1)$  setwise, but is orientation reversing on each.

Not all knots or gluing maps will yield a splice which admits such a symmetry. We enumerate all the requirements in Section 2. In this paper, we will refer to the above families as *Type-1* and *Type-2 symmetric splices*, respectively. We will see in Section 2.1 that Type-1 splices must have  $(Y, K)$  *reversible*,

MSC2020: 57K18, 57K31, 57R58.

© 2026 The Authors, under license to MSP (Mathematical Sciences Publishers). Distributed under the Creative Commons Attribution License 4.0 (CC BY). Open Access made possible by subscribing institutions via [Subscribe to Open](#).

i.e., there must be a diffeomorphism  $(Y, K) \cong (Y, -K)$ , where  $-K$  denotes the knot with the opposite string orientation. In [Section 2.2](#) we will see that in a Type-2 symmetric splice, one of  $K_0$  and  $K_1$  must be negatively amphichiral and the other positively amphichiral. We will also enumerate all possible gluing maps.

*Involutive Heegaard Floer homology* is a shadow theory of  $\text{Pin}(2)$ -equivariant Seiberg–Witten Floer homology, introduced by Manolescu and Hendricks in 2015 and elaborated by Hendricks and Zemke with Manolescu [[12](#); [13](#)], which is conjecturally equivalent to  $\mathbb{Z}/4\mathbb{Z}$ -equivariant Seiberg–Witten Floer homology. Although involutive Heegaard Floer homology does not possess the technical power of a  $\text{Pin}(2)$ -equivariant theory, it enjoys the greater computability of the Heegaard Floer invariants, including a conveniently computable surgery formula [[11](#)], and has been a key tool in recent developments regarding the structure of the homology cobordism group [[4](#); [9](#); [10](#); [11](#)].

In this note we show that the homology cobordism involutive invariants of Type-1 and Type-2 splices are typically trivial. Recall for a homology sphere  $Z$ , these invariants consist of a pair  $(\text{CF}^-(Z), \iota)$  a chain complex together with a homotopy involution, together called an *iota-complex*; for more details, see [Section 3](#).

**Theorem 1.1** (1) *Suppose that  $K$  is a knot in an integer homology 3-sphere  $Y$ . If  $Z$  is a Type-1 symmetric splice of  $(Y, K)$  with  $(-Y, -K)$  then the *iota-complex*  $(\text{CF}^-(Z), \iota)$  is locally trivial.*

(2) *If  $Z$  is a Type-2 symmetric splice of  $(S^3, K_0)$  and  $(S^3, K_1)$  such that  $\text{CFK}^-(K_0)$  and  $\text{CFK}^-(K_1)$  are (noninvolutively) locally trivial, then the *iota-complex*  $(\text{CF}^-(Z), \iota)$  is locally trivial.*

**Remark 1.2** (1) It is not clear to the authors whether [Theorem 1.1\(2\)](#) extends to all amphichiral  $K_0$  and  $K_1$ , nor whether it can be extended to knots in homology 3-spheres other than  $S^3$ .

(2) As we mentioned above, in a symmetric splice of Type-2, one of  $K_0$  and  $K_1$  must be negative amphichiral, and the other positive amphichiral. To the best of our knowledge, all known amphichiral knots have locally trivial (noninvolutive) knot Floer complex  $\text{CFK}^-(K)$ . Also note if  $K$  is *strongly* negative amphichiral, i.e., if the pair  $(S^3, K)$  admits an orientation-reversing diffeomorphism  $\phi$  which has exactly two fixed points, both of which lie along  $K$ , then Kawauchi’s result [[15](#)] implies that  $K$  is rationally slice, and hence has locally trivial  $\text{CFK}^-(K)$ .

The key topological input to [Theorem 1.1](#) is the following result:

**Proposition 1.3** (1) *If  $Z$  is a Type-1 symmetric splice of  $(Y, K)$  with  $(-Y, -K)$ , then there is a negative definite Spin cobordism from  $Z$  to  $\mathbb{RP}^3$  which has  $b_2^- = 1$  and  $b_1 = 0$ .*

(2) *If  $Z$  is a Type-1 symmetric splice of  $(Y, K)$  with  $(-Y, -K)$ , then there is a negative definite (non-Spin) filling of  $Z$  with  $b_2^- = 2$  and  $b_1 = 0$ .*

(3) *If  $Z$  is a Type-2 symmetric splice of  $(Y_0, K_0)$  with  $(Y_1, K_1)$ , then there is a negative definite Spin cobordism from  $Z$  to  $(Y_0 \# Y_1)_{-2}(K_0 \# K_1)$  with  $b_2^- = 1$  and  $b_1 = 0$ .*

**Remark 1.4** In unpublished work, Mike Miller Eismeier independently proved [Proposition 1.3\(2\)](#) and used it to show that certain instanton-theoretic gauge-theoretic invariants are trivial on such splices.

We now sketch some ideas in the proof of [Theorem 1.1](#), assuming [Proposition 1.3](#). For Type-1 splices, the negative definite Spin cobordism  $W$  from  $Z$  to  $\mathbb{R}P^3$  has a unique self-conjugate  $\text{Spin}^c$  structure  $\mathfrak{s}$ . Furthermore, the Heegaard Floer grading shift  $d(W, \mathfrak{s})$  is equal to the correction term  $d(\mathbb{R}P^3, \mathfrak{s}|_{\mathbb{R}P^3})$ . Since  $\mathbb{R}P^3$  is a Heegaard Floer L-space, the cobordism map  $F_{W, \mathfrak{s}}$  can be viewed as a local map from  $(\text{CF}^-(Z), \iota)$  to the trivial  $\iota$ -complex. Since  $Z \cong -Z$ , we can dualize the map  $F_{W, \mathfrak{s}}$  to get a local map in the opposite direction.

For Type-2 splices, the cobordism  $W$  from  $Z$  to  $S^3_{-2}(K_0 \# K_1)$  also has a unique self-conjugate Spin structure  $\mathfrak{s}$ . In this case,  $\mathfrak{s}$  restricts to the  $\text{Spin}^c$  structure identified with  $[1] \in \mathbb{Z}/2 \cong \text{Spin}^c(S^3_{-2}(K_0 \# K_1))$  under the standard identification. We use [\[11, Theorem 1.6\(2\)\]](#), which implies that since  $\text{CFK}^-(K_0)$  and  $\text{CFK}^-(K_1)$  are (noninvolutively) locally trivial, the  $\iota$ -complex  $(\text{CF}^-(S^3_{-2}(K_0 \# K_1), [1]), \iota)$  is locally trivial up to an overall grading shift.

### 1.1 Other gauge-theoretic invariants

We note that the topological argument yielding [Theorem 1.1](#) applies equally well to the  $\text{Pin}(2)$ -equivariant Seiberg–Witten Floer spectra:

**Proposition 1.5** *The  $\text{Pin}(2)$ -equivariant Seiberg–Witten Floer spectra of symmetric splices of Type-1 are locally trivial.*

The same argument also implies the vanishing of Lin’s invariants  $\alpha(Z), \beta(Z), \gamma(Z)$  [\[16\]](#) for Type-1 symmetric splices.

Note that [Proposition 1.3\(2\)](#) can be applied to other invariants of gauge theory. In particular if  $A$  is a partially ordered set and

$$\omega : \Theta_{\mathbb{Z}}^3 \rightarrow A$$

is a homology cobordism invariant which is monotonic under negative definite cobordisms with  $b_1 = 0$ , then [Proposition 1.3\(2\)](#) implies that if  $Z$  is a Type-1 symmetric splice, then  $\omega(Z) \leq \omega(S^3)$  and  $\omega(S^3) \leq \omega(Z)$ , so in particular  $\omega(Z) = \omega(S^3)$ . This holds for many gauge-theoretic invariants of homology cobordism. For example, this argument applies to the  $r_s$ -invariants of Nozaki, Sato, Taniguchi [\[21\]](#) and Daemi’s  $\Gamma$  invariant [\[1\]](#). Compare also [\[2\]](#).

### Organization

This note is organized as follows. In [Section 2](#) we discuss the geometry of splices; in particular, we classify which symmetric splices of Types 1 and 2 are integer homology spheres and have a natural orientation-reversing diffeomorphism. In [Section 3](#) we briefly recall relevant aspects of Heegaard Floer theory, focusing on its interaction with homology cobordism and concordance, for the reader’s convenience. In [Section 4](#) we conclude with the proof of [Theorem 1.1](#).

## 2 Symmetric splices

In this section we recall some background on splices. Let  $K_0 \subseteq Y_0$  and  $K_1 \subseteq Y_1$  be oriented knots in integer homology 3-spheres. Let  $\phi \in \mathbb{Z}^2 \rightarrow \mathbb{Z}^2$  be a  $2 \times 2$  matrix with determinant  $-1$ . The map  $\phi$  determines an orientation-reversing diffeomorphism

$$\phi : \partial(Y_0 \setminus \nu(K_0)) \rightarrow \partial(Y_1 \setminus \nu(K_1)),$$

where we view the first component of  $\mathbb{Z}^2$  being an oriented meridian of  $K_j$ , and the second component being the oriented Seifert longitude. We define

$$\text{Sp}_\phi(K_0, K_1) := (Y_0 \setminus \nu(K_0)) \cup_\phi (Y_1 \setminus \nu(K_1)).$$

In this section, we will be interested in knots that have various symmetries. We use the following standard terminology:

**Definition 2.1** Let  $K$  be a knot in an oriented 3-manifold  $Y$ .

- (1)  $(Y, K)$  is *reversible* if  $(Y, K) \cong (Y, -K)$ .
- (2)  $(Y, K)$  is *negative amphichiral* if  $(Y, K) \cong (-Y, -K)$ .
- (3)  $(Y, K)$  is *positive amphichiral* if  $(Y, K) \cong (-Y, K)$ .

In the above,  $\cong$  means orientation-preserving diffeomorphic.

### 2.1 Type-1 symmetric splices

We now focus on Type-1 splices, i.e., splices of  $(Y, K)$  and  $(-Y, -K)$  which admit orientation-reversing diffeomorphisms which switch  $Y \setminus \nu(K)$  with  $-Y \setminus \nu(K)$  but fix  $\mathbb{T}^2 := \partial(Y \setminus \nu(K))$  setwise. We will write  $\text{Sp}_\phi(K, mK)$  for such splices. In this section we prove the following:

**Proposition 2.2** *Suppose  $K$  is a knot in an integer homology 3-sphere  $Y$ , and  $\phi \in \text{GL}_2^-(\mathbb{Z})$ . Assume that  $K$  is not an unknot in  $Y$ . Then the 3-manifold  $\text{Sp}_\phi(K, mK)$  admits an orientation-reversing diffeomorphism  $g$  which fixes  $\partial(Y \setminus \nu(K))$  setwise and such that the image of each of  $Y \setminus \nu(K)$  and  $-Y \setminus \nu(K)$  is the other, if and only if the following are satisfied:*

- (1)  $K$  is reversible.
- (2)  $\phi = \begin{pmatrix} n & \pm 1 \\ \pm(1+n^2) & n \end{pmatrix}$  for some  $n \in \mathbb{Z}$ . In this case we define  $\phi = \phi_n^\pm$ .

**Proof** It will be evident from the proof that if  $(Y, K)$  and  $\phi$  are as in the statement, then  $\text{Sp}_\phi(K, mK)$  will have a symmetry  $g$  as above. Hence, we assume that  $\text{Sp}_\phi(K, mK)$  admits an orientation-reversing diffeomorphism  $g$ , as in the statement, and we will show that  $K$  and  $\phi$  have the stated properties.

For our proof, it is somewhat easier write the gluing map in terms of a different basis. Note that if  $(\mu, \lambda)$  is our oriented basis for  $\partial Y \setminus \nu(K)$ , then  $\phi$  is written in terms of the basis  $(\mu, -\lambda)$  for  $-Y \setminus \nu(K)$ . For our purposes, it is more helpful to write  $\phi$  in terms of the basis  $(\mu, \lambda)$  for both  $Y \setminus \nu(K)$  and  $-Y \setminus \nu(K)$ , using the same longitude and meridian for both. Let us write  $\psi$  for the map  $\phi$  in this basis. Note that  $\det(\psi) = 1$ .

Additionally, to simplify the notation, we will view  $\text{Sp}_\phi(K, mK)$  as the union of two copies of  $Y \setminus \nu(K)$ , which we denote by  $X_0$  and  $X_1$ . We write

$$\text{Sp}_\phi(K, mK) = \frac{X_0 \sqcup X_1}{\sim},$$

where  $x \in \partial X_0$  is identified with  $\psi(x) \in \partial X_1$ . By assumption  $g$  is induced by some diffeomorphisms  $g_{10} : X_0 \rightarrow X_1$  and  $g_{01} : X_1 \rightarrow X_0$  as

$$\begin{array}{ccc} & g_{10} & \\ X_0 & \xrightarrow{\quad} & X_1 \\ & g_{01} & \end{array}$$

The diffeomorphism  $g_{10} \sqcup g_{01}$  descends to the quotient if and only if

$$(2-1) \quad g_{10}(x) = (\psi \circ g_{01} \circ \psi)(x)$$

for all  $x \in \partial X_0$ .

We now claim that

$$(2-2) \quad g_{01}|_{\partial X_1}, g_{10}|_{\partial X_0} \in \{\text{id}, -\text{id}\}.$$

To see this, note that both must map  $\lambda$  to  $\pm\lambda$ , because they must preserve the kernel of the map  $H_1(\partial X_i) \rightarrow H_1(X_i)$ . Less obviously, they must also send  $\mu$  to  $\pm\mu$ . Homology considerations imply that  $g_{01}$  and  $g_{10}$  map  $\mu$  to  $\pm\mu + j\lambda$  for some  $j \in \mathbb{Z}$ . This would imply that, up to composition with the elliptic involution,  $g_{01}|_{\partial X_0}$  is an  $j$ -fold composition of a Dehn twist parallel to the Seifert longitude, and similarly for  $g_{10}$ . It follows from [20, Theorem 1] that this can only happen if  $\lambda$  bounds a disk in  $Y \setminus \nu(K)$ , i.e.,  $K$  is an unknot, which we exclude by hypothesis.

Since  $g_{01}|_{\partial X_1}, g_{10}|_{\partial X_0} \in \{\text{id}, -\text{id}\}$ , these maps are central in  $\text{GL}_2(\mathbb{Z})$ , and hence (2-1) implies that  $\psi^2 = \pm \text{id}$ .

We now consider the map  $\psi$  in more detail. The Mayer–Vietoris exact sequence reads

$$H_1(\mathbb{T}^2) \rightarrow H_1(Y \setminus \nu(K)) \oplus H_1(Y \setminus \nu(K)) \rightarrow H_1(Z) \rightarrow 0.$$

In particular, we see that for  $Y$  to be a homology sphere, we need  $\psi(\lambda) = \pm\mu + n\lambda$ , for some  $n \in \mathbb{Z}$ . That is, we can write  $\psi$  as a matrix as

$$\psi = \begin{pmatrix} n_1 & \pm 1 \\ * & n_2 \end{pmatrix}.$$

The condition that  $\det \psi = 1$  imposes the restriction that

$$(2-3) \quad \psi = \begin{pmatrix} n_1 & \pm 1 \\ \mp(1-n_1n_2) & n_2 \end{pmatrix}.$$

It is straightforward to see that there are no such matrices of the above form which square to the identity matrix. On the other hand, the matrix in (2-3) squares to  $-\text{id}$  if and only if  $n_1 = -n_2$ . Setting  $n = n_1$  and then changing to the basis gives the expression for  $\phi$  in the statement.

Next, we observe that (2-1) now implies that  $g_{10}|_{\partial X_0} = -g_{01}|_{\partial X_1}$ . Therefore one map is the identity, while the other is the elliptic involution. Therefore  $Y \setminus \nu(K)$  admits an orientation-preserving diffeomorphism which restricts to the elliptic involution on the boundary. Equivalently, there is a diffeomorphism of pairs  $(Y, K) \cong (Y, -K)$ . □

**Lemma 2.3** *Suppose that  $(Y, K)$  is reversible. Then  $\text{Sp}_{\phi_n^+}(K, mK) \cong \text{Sp}_{\phi_{-n}^-}(K, mK)$ .*

**Proof** Since  $K$  is reversible, the elliptic involution of the boundary extends to an orientation-preserving diffeomorphism of  $Y \setminus \nu(K)$ . Therefore

$$\text{Sp}_{\phi_n^+}(K, mK) \cong \text{Sp}_{-\phi_n^+}(K, mK) = \text{Sp}_{\phi_{-n}^-}(K, mK). \quad \square$$

### 2.2 Type-2 symmetric splices

We now consider Type-2 splices. We say that a pair  $(Y, K)$  is *negative amphichiral* if  $(-Y, -K) \cong (Y, K)$ , and we say that  $(Y, K)$  is *positive amphichiral* if  $(-Y, K) \cong (Y, K)$ .

**Proposition 2.4** *Let  $(Y_0, K_0)$  and  $(Y_1, K_1)$  be knots and  $\phi \in \text{GL}_2^-(\mathbb{Z})$ , and furthermore suppose that  $\text{Sp}_\phi(K_0, K_1)$  is an integer homology 3-sphere. Then  $\text{Sp}_\phi(K_0, K_1)$  admits an orientation-reversing diffeomorphism  $g$  which preserves the subspaces  $Y_i \setminus \nu(K_i)$  setwise if and only if the following hold:*

- (1) *One of  $(Y_0, K_0), (Y_1, K_1)$  is negative amphichiral and the other is positive amphichiral.*
- (2)  $\phi = \pm \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ .

**Proof** We write

$$X_0 = Y_0 \setminus \nu(K_0) \quad \text{and} \quad X_1 = Y_1 \setminus \nu(K_1).$$

It will follow from the course of our proof that if  $\phi$  and  $(Y_i, K_i)$  are as in the statement, then there is an orientation-reversing diffeomorphism  $g$  as in the statement. Hence we will assume that such a  $g$  exists, and prove that it has the stated form.

We assume that  $g$  is induced by a pair of maps,  $g_0$  and  $g_1$ , as

$$g_0 \circlearrowleft X_0, \quad X_1 \circlearrowright g_1.$$

The maps  $g_0$  and  $g_1$  induce a map on the quotient space if and only if

$$(2-4) \quad \phi \circ g_0 = g_1 \circ \phi.$$

The proof of (2-2) shows that for  $i = 0, 1$ , we have  $g_i|_{\partial X_i} \in \{e, -e\}$  where

$$e = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

Equation (2-4) implies that

$$\phi \circ e = \pm e \circ \phi.$$

It is easy to see that this restricts  $\phi$  to be one of four matrices

$$\phi = \pm e \quad \text{and} \quad \phi = \pm \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

Note that if  $\phi = \pm e$ , then the splice  $\text{Sp}_\phi(K_0, K_1)$  has  $b_1 = 1$ , so we exclude this case and restrict to the second case. We observe that in the latter case, we have

$$\phi \circ e = -e \circ \phi.$$

In particular, we conclude from (2-4) that  $g_0 = -g_1$ . Note that this corresponds exactly to one of  $K_0$  and  $K_1$  being negative amphichiral, and the other being positive amphichiral.  $\square$

### 2.3 Factorizations in $\text{SL}_2(\mathbb{Z})$

In this section, we describe some straightforward algebra which will be used in the subsequent section on Kirby calculus.

We consider the elements  $\psi_n^+ \in \text{SL}_2(\mathbb{Z})$ , given by

$$\psi_n^+ = \begin{pmatrix} n & 1 \\ -(1+n^2) & -n \end{pmatrix}.$$

We define the following elements of  $\text{SL}_2(\mathbb{Z})$ :

$$T_n := \begin{pmatrix} 1 & n \\ 0 & 1 \end{pmatrix} \quad \text{and} \quad H := \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}.$$

**Lemma 2.5** *The map  $\psi_n^+$  may be written as*

$$\psi_n^+ = HT_{-n}HT_nH.$$

The proof is a straightforward computation, which we leave to the reader.

### 2.4 Kirby calculus

We can now translate Lemma 2.5 into Kirby calculus. Our main result is the following:

**Proposition 2.6** *Let  $K \subseteq Y$  be a knot. The manifold  $\text{Sp}_{\phi_n^+}(K, mK)$  has a Kirby diagram as shown in Figure 1.*

We begin with the following elementary topological lemma.

**Lemma 2.7** *Let  $(Y_0, K_0)$  and  $(Y_1, K_1)$  be knots with Morse framings  $\lambda_0$  and  $\lambda_1$ , respectively. Let  $\phi$  be the gluing map which identifies the meridian  $\mu_0$  with  $\mu_1$ , and which maps  $\lambda_0$  to  $-\lambda_1$ . (Here,  $-\lambda_2$  denotes the Morse framing  $\lambda_2$  with the parametrization reversed). Then*

$$(Y_0 \setminus \nu(K_0)) \cup_\phi (Y_1 \setminus \nu(K_1))$$

is equal to  $(Y_0 \# Y_1)_{\lambda_0 + \lambda_1}(K_0 \# K_1)$ .




Figure 1: The manifold  $\text{Sp}_{\phi_n^+}(K, mK)$ . We view the box labeled by  $K$  as being inside  $Y$ .

See [6, Lemma 6.1; 8, Lemma 7.1] for a proof. See also [26, Section 1.1.7].

The above lemma extends in a straightforward manner to link complements when we take the connected sum along a single component. We note that the Hopf link has complement  $\mathbb{T}^2 \times [0, 1]$ . The meridian of the first component of the Hopf link is equal to the longitude of the second up to sign, and vice versa. Therefore from a factorization of the gluing diffeomorphism  $\psi_n^+$ , as in Lemma 2.5, we may read off a Kirby calculus description of  $\text{Sp}_{\phi_n^+}(K, mK)$ . Namely, we start with  $K$ , which we give framing 0. Reading the factorization

$$\psi_n^+ = HT_{-n}HT_nH$$

from right to left we form a Kirby calculus presentation inductively as follows:

- (1) Start with  $K$ , given framing 0.
- (2) For each  $H$ , we take the connected sum with a  $(0, 0)$ -framed Hopf link.
- (3) For each  $T_n$ , we add  $n$  to the framing of the most-recently added component in this process.
- (4) We finish by taking the connected sum of the final unknot, which has framing 0, with  $mK$ .

Note that since we are assuming that  $K$  is reversible, we do not need to worry about the sign of the clasps that we add when taking the connected sum with Hopf links.

We now describe some Kirby calculus moves in our description of  $\text{Sp}_{\phi_n^+}(K, mK)$  which will be helpful later on.

**Lemma 2.8** *Let  $K$  be a reversible knot in an integer homology sphere  $Y$ , and  $n \in \mathbb{Z}$ . Then there are reversible knots  $K' \subseteq Y'$  and  $K'' \subseteq Y''$ , where  $Y'$  and  $Y''$  are also integer homology 3-spheres, so that*

$$\text{Sp}_{\phi_n^+}(K, mK) = \text{Sp}_{\phi_{n+1}^+}(K', mK') = \text{Sp}_{\phi_{n-1}^+}(K'', mK'').$$

**Proof** The proof is to take the Kirby calculus description in Figure 1, and blow up the clasps between  $K$  and the unknotted component clasped with it and between  $mK$  and the unknotted component clasped with it. In this manner,  $n$  can be increased or decreased. Then  $Y' = Y_{+1}(K)$  and  $K'$  is the dual knot of  $K$ . We give  $K'$  blackboard framing 1 in Figure 2, but note that this corresponds to the Seifert framing for  $K'$  if we view it as living in  $Y_{+1}(K)$ . Hence, we obtain a description of the same form as Figure 1, except with  $n$  replaced by  $n + 1$ . □




Figure 2: Left: an alternate surgery description of  $\text{Sp}_{\phi_n^+}(K, mK)$ , obtained by blowing up two clasps in Figure 1. This identifies  $\text{Sp}_{\phi_n^+}(K, mK)$  with  $\text{Sp}_{\phi_{n+1}^+}(K', mK')$  where  $K' \subseteq Y' = Y_{+1}(K)$  is the dual knot of  $K$ . Right: the knot  $K' \subseteq Y_{+1}(K)$ .

### 3 Heegaard Floer invariants of concordance and homology cobordism

In this section, we review some background on Heegaard Floer invariants of homology cobordism and knot concordance. We focus on Hendricks and Manolescu’s *involutive Heegaard Floer homology* [12], which we review in Section 3.1, as well as the notion of a knot-like complex, which we review in Section 3.2. We presume the reader is familiar with ordinary Heegaard Floer homology for three-manifolds [23; 24] and knots [22; 25].

#### 3.1 Iota-complexes and involutive Heegaard Floer homology

In this section we briefly introduce the structure of the involutive Heegaard Floer invariants, with a focus on the properties of local equivalence.

We begin with certain algebraic definitions. Throughout,  $\mathbb{F}$  denotes the field with two elements,  $U$  is a variable of degree  $-2$ , and  $\mathbb{F}[U]_d$  is the graded module such that  $\text{gr}(1) = d$ .

**Definition 3.1** An *iota-complex* (or  $\iota$ -complex)  $(C, \iota)$  is a chain complex  $C$ , which is free and finitely generated over  $\mathbb{F}[U]$ , equipped with an endomorphism  $\iota$ . Here  $\mathbb{F}$  is the field of 2 elements, and  $U$  is a formal variable with grading  $-2$ . Furthermore, the following hold:

- (1)  $C$  is equipped with a  $\mathbb{Z}$ -grading, compatible with the action of  $U$ . We call this grading the *Maslov or homological grading*.
- (2) There is a grading-preserving isomorphism  $U^{-1} H_*(C) \cong \mathbb{F}[U, U^{-1}]$ .
- (3)  $\iota$  is a grading-preserving chain map and  $\iota^2 \simeq \text{id}$ .

Given two iota-complexes  $(C_1, \iota_1)$  and  $(C_2, \iota_2)$ , a homogeneously graded  $\mathbb{F}[U]$ -chain map  $f : C_1 \rightarrow C_2$  is said to be an  $\iota$ -homomorphism if  $\iota_2 \circ f + f \circ \iota_1 \simeq 0$ . Two iota-complexes  $(C_1, \iota_1)$  and  $(C_2, \iota_2)$  are called  $\iota$ -equivalent if there is a homotopy equivalence  $\Phi : C_1 \rightarrow C_2$  which is an  $\iota$ -homomorphism.

Heegaard Floer homology associates to any closed oriented 3-manifold  $Y$  equipped with a  $\text{Spin}^c$  structure  $\mathfrak{s}$  an  $\mathbb{F}[U]$ -chain complex  $\text{CF}^-(Y, \mathfrak{s})$ , well defined up to homotopy equivalence. If  $\mathfrak{s}$  is self-conjugate, involutive Heegaard Floer homology considers the additional data of a homotopy involution  $\iota$  on  $\text{CF}^-(Y, \mathfrak{s})$ . In the case that  $Y$  is a rational homology 3-sphere,  $(\text{CF}^-(Y, \mathfrak{s}), \iota)$  is an iota-complex. Hendricks and Manolescu [12] prove that pair  $(\text{CF}^-(Y, \mathfrak{s}), \iota)$  is well defined up to the notion of iota-equivalence described above.

Continuing with algebra, the tensor product of iota-complexes  $(C_1, \iota_1)$  and  $(C_2, \iota_2)$  is given by

$$(3-1) \quad (C_1, \iota_1) \otimes (C_2, \iota_2) := (C_1 \otimes_{\mathbb{F}[U]} C_2, \iota_1 \otimes \iota_2).$$

Moreover, Hendricks, Manolescu, and Zemke [13] establish that

$$(\text{CF}^-(Y_1 \# Y_2, \mathfrak{s}_1 \# \mathfrak{s}_2), \iota) \simeq (\text{CF}^-(Y_1, \mathfrak{s}_1), \iota_1) \otimes (\text{CF}^-(Y_2, \mathfrak{s}_2), \iota_2),$$

where  $\simeq$  denotes homotopy-equivalence of iota-complexes.

**Definition 3.2** Suppose  $(C, \iota)$  and  $(C', \iota')$  are two iota-complexes.

- (1) A *local map* from  $(C, \iota)$  to  $(C', \iota')$  is a grading-preserving  $\iota$ -homomorphism  $F : C \rightarrow C'$ , which induces an isomorphism from  $U^{-1}H_*(C)$  to  $U^{-1}H_*(C')$ .
- (2) We say that  $(C, \iota)$  and  $(C', \iota')$  are *locally equivalent* if there is a local map from  $(C, \iota)$  to  $(C', \iota')$ , as well as a local map from  $(C', \iota')$  to  $(C, \iota)$ . We say that  $(C, \iota)$  is *locally trivial* if it is locally equivalent to  $(\mathbb{F}[U]_0, \text{Id})$ .

The set of local equivalence classes forms an abelian group, denoted by  $\mathfrak{J}$ , with product given by the operation  $\otimes$  in (3-1) [13, Section 8]. Inverses are given by dualizing both the chain complex  $C$  and the map  $\iota$  with respect to  $\mathbb{F}[U]$ . The map

$$Y \mapsto [(\text{CF}^-(Y), \iota)]$$

determines a homomorphism from  $\Theta_{\mathbb{Z}}^3$  to  $\mathfrak{J}$  [13, Theorem 1.8].

The local equivalence classes of nonzero integer surgeries on knots are computed in [11, Theorem 1.6(2)]. For our purposes, the important case is the following.

**Lemma 3.3** [11, Theorem 1.6(2)] *For  $n > 0$ , the local equivalence class of  $(\text{CF}^-(S_{2n}^3(K), [n]), \iota)$  has the form*

$$\begin{array}{ccc} A_n^-(K) & & A_n^-(K) \\ & \searrow v & \swarrow v' \\ & B_n^-(K) & \end{array}$$

where  $A_n^-(K)$  and  $B_n^-(K)$  are subcomplexes of the knot Floer complex of  $K$ ,  $v$  is a particular map between them, and the involution swaps the two copies of  $A_n^-(K)$ , and fixes  $B_n^-(K)$ . The gradings on the above are induced by the Maslov grading on the knot Floer complex, shifted up by the Heegaard Floer correction term  $d(L(2n, 1), [n])$  of the lens space  $L(2n, 1)$  in the corresponding  $\text{Spin}^c$  structure.

One straightforward corollary is the following:

**Corollary 3.4** *For  $n > 0$ , if  $K \subseteq S^3$  is a knot such that  $d(A_n^-(K)) = 0$ , then  $(\text{CF}^-(S_{2n}^3(K), [n]), \iota)$  is locally equivalent to  $(\mathbb{F}[U]_d, \text{id})$ , where  $d = d(L(2n, 1), [n])$ .*

**Proof** We note that  $B_n^-(K) \simeq \mathbb{F}[U]$ , so using the same logic as in the proof of [11, Proposition 3.24], we may replace it with a copy of  $\mathbb{F}[U]$ . By the classification theorem for finitely generated chain complexes over  $\mathbb{F}[U]$ , we can write  $A_n^-(K)$  as a sum of one tower  $\mathbb{F}[U]$ , as well as some number of 2-step complexes

of the form  $\mathbb{F}[U] \xrightarrow{U^i} \mathbb{F}[U]$ . Write  $\mathbf{x}_l$  and  $\mathbf{x}_r$  for tower generators of the two copies of  $A_n^-(K)$ , which is to say, generators for the copy of  $\mathbb{F}[U]$  in the basis chosen. The map  $v$  sends  $\mathbf{x}_l$  and  $\mathbf{x}_r$  to a nonzero element of the tower  $\mathbb{F}[U]$ . Since  $d(A_n^-(K)) = 0$ , we conclude that  $v(\mathbf{x}_l) = v(\mathbf{x}_r) = 1$ . On a two step subcomplex of the left copy of  $A_n^-(K)$ , say with generators  $\mathbf{a}$  and  $\mathbf{b}$  such that  $\partial(\mathbf{a}) = U^i \mathbf{b}$ , we must have  $v(\mathbf{b}) = 0$  because  $v$  is a chain map. If  $v(\mathbf{a})$  is nonzero, then perform a change of basis, adding a multiple of  $\mathbf{x}_l$  to  $\mathbf{a}$ . After this change of basis,  $v(\mathbf{a}) = 0$ . We do the same change of basis to the right copy of  $A_n^-(K)$ . After this change of basis, it becomes apparent that the complex in the statement of Lemma 3.3 is locally equivalent to

$$\begin{array}{ccc} \mathbb{F}[U]_d & & \mathbb{F}[U]_d \\ & \searrow 1 & \swarrow 1 \\ & \mathbb{F}[U]_d & \end{array}$$

where the involution is reflection, and  $d$  denotes  $d(L(2n, 1), [n])$ . The above is homotopy equivalent to  $(\mathbb{F}[U]_d, \text{id})$ . □

### 3.2 Knot-like complexes

We now recall the standard notion of knot-like complexes in Heegaard Floer theory. There are many different variations on the definition in the literature due to many different authors. The earliest version is Hom’s notion of  $\varepsilon$ -equivalence [14]. See also [3; 5; 28] for other variations.

**Definition 3.5** A *knot-like* complex  $C$  is a finitely generated, free chain complex over a 2-variable polynomial ring  $\mathbb{F}[U, V]$  satisfying the following:

- (1)  $C$  is equipped with a  $\mathbb{Z} \times \mathbb{Z}$ -valued bigrading, denoted by  $(\text{gr}_w, \text{gr}_z)$ , which has the property that  $(\text{gr}_w - \text{gr}_z)/2$  is integrally valued. The variable  $U$  has bigrading  $(-2, 0)$  and the variable  $V$  has bigrading  $(0, -2)$ .
- (2) There is a grading-preserving isomorphism  $(U, V)^{-1} H_*(C) \cong \mathbb{F}[U, V, U^{-1}, V^{-1}]$ .
- (3)  $\partial$  has bigrading  $(-1, -1)$ .

A local map from  $C_0$  to  $C_1$  (where  $C_i$  are knot-like complexes) consists of a grading-preserving  $\mathbb{F}[U, V]$  linear chain map  $F : C_0 \rightarrow C_1$  such that  $F$  induces an isomorphism from  $(U, V)^{-1} H_*(C_0)$  to  $(U, V)^{-1} H_*(C_1)$ . We say that  $C_0$  and  $C_1$  are *locally equivalent* if there exist local maps from  $C_0$  to  $C_1$  and from  $C_1$  to  $C_0$ . A knot-like complex  $C$  is *locally trivial* if it is locally equivalent to a rank-one complex  $\mathbb{F}[U, V]$  wherein  $1 \in \mathbb{F}[U, V]$  is concentrated in grading  $(0, 0)$ . Note that  $C$  is locally trivial if and only if there is an isomorphism

$$C \cong \mathbb{F}[U, V] \oplus A,$$

where  $A$  is a summand of  $C$  such that  $(U, V)^{-1} H_*(A) \cong 0$ .

If  $K \subseteq S^3$ , the full version of the knot Floer complex  $\text{CFK}^-(K)$  is a knot-like complex. If  $K$  is a slice knot, then  $\text{CFK}^-(K)$  is locally trivial.

### 4 Proofs of the main results

We first observe that if  $Z$  is a symmetric splice, since  $Z$  has order at most two in  $\Theta_{\mathbb{Z}}^3$ , we must have  $d(Z) = 0$  since  $d$  is a homomorphism.

**Lemma 4.1** *Suppose that  $Z$  is the homology sphere obtained by splicing  $(Y, K)$  and  $(-Y, -K)$  using the gluing map  $\phi_0^+$  from Proposition 2.2, where  $Y$  is a homology 3-sphere. Then there is a negative definite Spin cobordism  $W$  from  $Z$  to  $\mathbb{R}P^3$ . The 4-manifold  $W$  has a unique self-conjugate  $\text{Spin}^c$  structure  $\mathfrak{s}$ . Further, letting  $d(W, \mathfrak{s})$  denote the grading shift of the map associated to the cobordism  $(W, \mathfrak{s})$ , we have*

$$d(W, \mathfrak{s}) = d(\mathbb{R}P^3, \mathfrak{s}|_{\mathbb{R}P^3}) - d(Z).$$

**Proof** We begin with the Kirby calculus presentation from Figure 1. We can blow-down one of the unknots to obtain a Kirby calculus description of  $Z$  as Dehn surgery on a knot  $K \# H \# -K \subseteq Y \# -Y$ , where  $H$  denotes a Hopf link. That is, we add a clasp between  $K$  and  $-K$ . (Note that the sign of the clasp is not important since  $(Y, K)$  is reversible). The two components are given framing 0. We now blow up the clasp to obtain the 3-component link  $K \cup U \cup mK$  with clasps between the components. Each component is given framing  $-1$ . There is a cobordism  $X$  from  $Z$  to a manifold  $Z'$  by performing  $-1$  surgery on a meridian of the unknot  $U$ , where this knot is given Seifert framing  $-2$  inside of  $Z$ . The result is  $-2$  surgery on  $K \# -K \subseteq Y \# -Y$ . The pair  $(Y \# -Y, K \# -K)$  is homology concordant to  $(S^3, U)$ . Therefore  $Z'$  admits a homology cobordism to  $\mathbb{R}P^3$ , viewed as  $-2$  surgery on the unknot. Let  $W$  denote the composition of these two cobordisms. The cobordism  $W$  is shown in Figure 3.

We observe that  $d(Z) = 0$  since  $Z \cong -Z$ . On the other hand, we compute that the shift in grading for the Spin structure on  $W$  is

$$\frac{1}{4}(-2\chi(W) - 3\sigma(W)) = \frac{1}{4}.$$

We note that the  $d$ -invariants of the two  $\text{Spin}^c$  structures ( $=$  Spin structures) on  $\mathbb{R}P^3$  are  $\frac{1}{4}$  and  $-\frac{1}{4}$ . Since the Maslov grading takes values in a single coset of  $\mathbb{Q}/\mathbb{Z}$  in each  $\text{Spin}^c$  structure, it follows that the Spin structure on  $W$  restricts to the  $\text{Spin}^c$  structure on  $\mathbb{R}P^3$  which has  $d$ -invariant  $\frac{1}{4}$ . □

Using Lemma 4.1, we now prove Theorem 1.1(1), which concerns symmetric splices of Type-1:

**Proof of Theorem 1.1(1)** Let  $Z$  be a symmetric splice of Type-1. By Proposition 2.2,  $Z$  can be written as a splice  $\text{Sp}_{\phi_n^\pm}(K, mK)$  for some pair  $(Y, K)$ , where  $Y$  is a homology sphere and  $(Y, K)$  is reversible. By Lemmas 2.3 and 2.8, we may assume that  $n = 0$  by changing  $(Y, K)$  appropriately. Applying Lemma 4.1 gives a local map from  $(\text{CF}^-(Z), \iota)$  to  $(\text{CF}^-(\mathbb{R}P^3, \mathfrak{s}), \iota)$  for the Spin structure  $\mathfrak{s}$  with  $d(\mathbb{R}P^3, \mathfrak{s}) = \frac{1}{4}$ . Since the grading shift of the cobordism map is also  $\frac{1}{4}$ , and  $(\text{CF}^-(\mathbb{R}P^3, \mathfrak{s}), \iota) \cong (\mathbb{F}[U]_{1/4}, \text{id})$ , we conclude that there is a local map from  $(\text{CF}^-(Z), \iota)$  to the trivial complex. Dualizing and using the fact that  $Z \cong -Z$  gives a local map in the opposite direction. □

**Proof of Theorem 1.1(2)** The proof is similar to the proof of Theorem 1.1(1). By Proposition 2.4, the 3-manifold  $Z$  can be written as  $\text{Sp}_{\phi_0^+}(K_0, K_1)$  where  $K_0, K_1 \subseteq S^3$  are positive and negative amphichiral knots, respectively. By adapting the argument from Theorem 1.1(1), we obtain a negative definite Spin




Figure 3: The cobordism  $W$  from  $Z$  to  $\mathbb{R}P^3$ .

cobordism from  $Z$  to  $S_{-2}^3(K_0 \# K_1)$  which shifts the Maslov grading by  $\frac{1}{4}$ . Assuming (up to relabeling) that  $K_0$  is positive amphichiral and  $K_1$  is negative amphichiral, we observe that

$$S_{-2}^3(K_0 \# K_1) \cong -S_{+2}^3(K_0 \# -K_1),$$

where  $-K_1$  denotes  $K_1$  with its string orientation reversed. Therefore, [Corollary 3.4](#) implies that

$$(\text{CF}^-(S_{+2}^3(K_0 \# -K_1), [1]), \iota) \sim_{\text{loc}} (\mathbb{F}[U]_{-1/4}, \text{id}).$$

Dualizing, we obtain that

$$(\text{CF}^-(S_{-2}^3(K_0 \# K_1), [1]), \iota) \sim_{\text{loc}} (\mathbb{F}[U]_{1/4}, \text{id}).$$

It follows that there is a local map from  $(\text{CF}^-(Z), \iota)$  to  $(\mathbb{F}[U], \text{id})$ . Since  $Z \cong -Z$ , we conclude that  $(\text{CF}^-(Z), \iota)$  is locally trivial. □

We now prove [Proposition 1.3](#), most of which we have already proven:

**Proof of Proposition 1.3** Part (1) follows from [Lemma 4.1](#), above. Part (3) is similar, and is described in our proof of [Theorem 1.1\(2\)](#). Finally Part (2) is obtained by composing the cobordism from Part (1) with the natural negative definite cobordism from  $\mathbb{R}P^3$  to  $\emptyset$ , namely the disk bundle over  $S^2$  with Euler number  $-2$ . □

**Proof of Proposition 1.5** The argument is essentially identical to the proof of Part (1) of [Theorem 1.1](#), but where the notation is adjusted to be for Seiberg–Witten Floer spectra in the setting of [\[18\]](#). In particular,

local equivalence of  $\text{Pin}(2)$ -equivariant spectra is defined just as in [Definition 3.2](#) above, except  $\text{Pin}(2)$ -equivariant spectra take the place of  $\text{iota}$ -complexes (see [\[27, Definition 2.7\]](#) and surrounding discussion).

Let  $Z$  be a symmetric splice of Type-1. [Lemma 4.1](#) gives a local map

$$\Sigma^{\frac{1}{16}\mathbb{H}} \text{SWF}(Z) \rightarrow \text{SWF}(\mathbb{R}\mathbb{P}^3, \mathfrak{s}),$$

with  $\mathfrak{s}$  as in the proof of [Theorem 1.1](#). We refer the reader to [\[17\]](#) for the definition of the (formal) fractional suspension. Meanwhile,  $\text{SWF}(\mathbb{R}\mathbb{P}^3, \mathfrak{s}) = S^{\frac{1}{16}\mathbb{H}}$ , and so we have a local map  $\text{SWF}(Z) \rightarrow S^0$ . Using that  $Z \cong -Z$ , we have a local map  $\text{SWF}(-Z) \rightarrow S^0$ ; furthermore, for general integer homology spheres  $X$ , we have  $\text{SWF}(X)$  and  $\text{SWF}(-X)$  are Spanier–Whitehead dual. As a consequence, if there is a local map  $\text{SWF}(-Z) \rightarrow S^0$  then there is a local map  $S^0 \rightarrow \text{SWF}(Z)$ . Thus  $S^0 \leq \text{SWF}(Z) \leq S^0$  in local equivalence, and so  $\text{SWF}(Z)$  is locally trivial as a  $\text{Pin}(2)$ -spectrum.  $\square$

## Acknowledgements

We are grateful to Jen Hom for helpful conversations, and to the referee for helpful comments and corrections. Hendricks was partially supported by NSF grant DMS-2019396. Stoffregen was partially supported by NSF grant DMS-2203828. Zemke was partially supported by NSF grant DMS-2204375 and a Sloan Research Fellowship.

## References

- [1] **A Daemi**, *Chern–Simons functional and the homology cobordism group*, *Duke Math. J.* 169:15 (2020) 2827–2886 [MR](#)
- [2] **A Daemi**, **C Scaduto**, *Equivariant aspects of singular instanton Floer homology*, *Geom. Topol.* 28:9 (2024) 4057–4190 [MR](#)
- [3] **I Dai**, **J Hom**, **M Stoffregen**, **L Truong**, *More concordance homomorphisms from knot Floer homology*, *Geom. Topol.* 25:1 (2021) 275–338 [MR](#)
- [4] **I Dai**, **J Hom**, **M Stoffregen**, **L Truong**, *An infinite-rank summand of the homology cobordism group*, *Duke Math. J.* 172:12 (2023) 2365–2432 [MR](#)
- [5] **I Dai**, **J Hom**, **M Stoffregen**, **L Truong**, *Homology concordance and knot Floer homology*, *Math. Ann.* 390:4 (2024) 6111–6186 [MR](#)
- [6] **S Fukuhara**, **N Maruyama**, *A sum formula for Casson’s  $\lambda$ -invariant*, *Tokyo J. Math.* 11:2 (1988) 281–287 [MR](#)
- [7] **D E Galewski**, **R J Stern**, *Classification of simplicial triangulations of topological manifolds*, *Ann. of Math. (2)* 111:1 (1980) 1–34 [MR](#)
- [8] **C M Gordon**, *Dehn surgery and satellite knots*, *Trans. Amer. Math. Soc.* 275:2 (1983) 687–708 [MR](#)
- [9] **K Hendricks**, **J Hom**, **T Lidman**, *Applications of involutive Heegaard Floer homology*, *J. Inst. Math. Jussieu* 20:1 (2021) 187–224 [MR](#)
- [10] **K Hendricks**, **J Hom**, **M Stoffregen**, **I Zemke**, *On the quotient of the homology cobordism group by Seifert spaces*, *Trans. Amer. Math. Soc. Ser. B* 9 (2022) 757–781 [MR](#)
- [11] **K Hendricks**, **J Hom**, **M Stoffregen**, **I Zemke**, *Surgery exact triangles in involutive Heegaard Floer homology* (2025) [arXiv 2011.00113](#)
- [12] **K Hendricks**, **C Manolescu**, *Involutive Heegaard Floer homology*, *Duke Math. J.* 166:7 (2017) 1211–1299 [MR](#)
- [13] **K Hendricks**, **C Manolescu**, **I Zemke**, *A connected sum formula for involutive Heegaard Floer homology*, *Selecta Math. (N.S.)* 24:2 (2018) 1183–1245 [MR](#)
- [14] **J Hom**, *The knot Floer complex and the smooth concordance group*, *Comment. Math. Helv.* 89:3 (2014) 537–570 [MR](#)

- [15] **A Kawauchi**, *Rational-slice knots via strongly negative-amphicheiral knots*, *Commun. Math. Res.* 25:2 (2009) 177–192 [MR](#)
- [16] **F Lin**, *A Morse–Bott approach to monopole Floer homology and the triangulation conjecture*, *Mem. Amer. Math. Soc.* 1221, Amer. Math. Soc., Providence, RI (2018) [MR](#)
- [17] **C Manolescu**, *Seiberg–Witten–Floer stable homotopy type of three-manifolds with  $b_1 = 0$* , *Geom. Topol.* 7 (2003) 889–932 [MR](#)
- [18] **C Manolescu**, *Pin(2)-equivariant Seiberg–Witten Floer homology and the triangulation conjecture*, *J. Amer. Math. Soc.* 29:1 (2016) 147–176 [MR](#)
- [19] **T Matumoto**, *Triangulation of manifolds*, from “Algebraic and geometric topology, II” (Stanford, CA, 1976) (R J Milgram, editor), *Proc. Sympos. Pure Math.* XXXII, Amer. Math. Soc., Providence, RI (1978) 3–6 [MR](#)
- [20] **D McCullough**, *Homeomorphisms which are Dehn twists on the boundary*, *Algebr. Geom. Topol.* 6 (2006) 1331–1340 [MR](#)
- [21] **Y Nozaki, K Sato, M Taniguchi**, *Filtered instanton Floer homology and the homology cobordism group*, *J. Eur. Math. Soc.* 26:12 (2024) 4699–4761 [MR](#)
- [22] **P Ozsváth, Z Szabó**, *Holomorphic disks and knot invariants*, *Adv. Math.* 186:1 (2004) 58–116 [MR](#)
- [23] **P S Ozsváth, Z Szabó**, *Holomorphic disks and three-manifold invariants: properties and applications*, *Ann. of Math. (2)* 159:3 (2004) 1159–1245
- [24] **P Ozsváth, Z Szabó**, *Holomorphic disks and topological invariants for closed three-manifolds*, *Ann. of Math. (2)* 159:3 (2004) 1027–1158 [MR](#)
- [25] **J A Rasmussen**, *Floer homology and knot complements*, PhD thesis, Harvard University (2003) [MR](#) [arXiv math/0306378](#)
- [26] **N Saveliev**, *Invariants for homology 3-spheres*, *Encyclopaedia of Mathematical Sciences* 140, Springer (2002) [MR](#)
- [27] **M Stoffregen**, *Pin(2)-equivariant Seiberg–Witten Floer homology of Seifert fibrations*, *Compos. Math.* 156:2 (2020) 199–250 [MR](#)
- [28] **I Zemke**, *Connected sums and involutive knot Floer homology*, *Proc. Lond. Math. Soc. (3)* 119:1 (2019) 214–265 [MR](#)

KRISTEN HENDRICKS [kristen.hendricks@rutgers.edu](mailto:kristen.hendricks@rutgers.edu)

*Department of Mathematics, Rutgers University, Piscataway, NJ, United States*

MATTHEW STOFFREGEN [stoffre1@msu.edu](mailto:stoffre1@msu.edu)

*Department of Mathematics, Michigan State University, East Lansing, MI, United States*

IAN ZEMKE [izemke@uoregon.edu](mailto:izemke@uoregon.edu)

*Department of Mathematics, University of Oregon, Eugene, OR, United States*

Received: July 15, 2024      Revised: March 21, 2025

# ALGEBRAIC & GEOMETRIC TOPOLOGY

[msp.org/agt](https://msp.org/agt)

## EDITORS

### PRINCIPAL ACADEMIC EDITORS

John Etnyre  
[etnyre@math.gatech.edu](mailto:etnyre@math.gatech.edu)  
Georgia Institute of Technology

Vesna Stojanoska  
[vesna@illinois.edu](mailto:vesna@illinois.edu)  
University of Illinois at Urbana-Champaign

### BOARD OF EDITORS

Julie Bergner	University of Virginia <a href="mailto:jeb2md@eservices.virginia.edu">jeb2md@eservices.virginia.edu</a>	Daniel Isaksen	Wayne State University <a href="mailto:isaksen@math.wayne.edu">isaksen@math.wayne.edu</a>
Steven Boyer	Université du Québec à Montréal <a href="mailto:cohf@math.rochester.edu">cohf@math.rochester.edu</a>	Thomas Koberda	University of Virginia <a href="mailto:thomas.koberda@virginia.edu">thomas.koberda@virginia.edu</a>
Tara E Brendle	University of Glasgow <a href="mailto:tara.brendle@glasgow.ac.uk">tara.brendle@glasgow.ac.uk</a>	Markus Land	JGU Mainz <a href="mailto:mland@uni-mainz.de">mland@uni-mainz.de</a>
Indira Chatterji	CNRS & Univ. Côte d'Azur (Nice) <a href="mailto:indira.chatterji@math.cnrs.fr">indira.chatterji@math.cnrs.fr</a>	Christine Lescop	Université Joseph Fourier <a href="mailto:lescop@ujf-grenoble.fr">lescop@ujf-grenoble.fr</a>
Octav Cornea	Université de Montreal <a href="mailto:cornea@dms.umontreal.ca">cornea@dms.umontreal.ca</a>	Norihiko Minami	OCAMI (Osaka Central Adv. Math. Inst.) <a href="mailto:norihikominami@gmail.com">norihikominami@gmail.com</a>
Alexander Dranishnikov	University of Florida <a href="mailto:dranish@math.ufl.edu">dranish@math.ufl.edu</a>	Andrés Navas	Universidad de Santiago de Chile <a href="mailto:andres.navas@usach.cl">andres.navas@usach.cl</a>
Tobias Ekholm	Uppsala University, Sweden <a href="mailto:tobias.ekholm@math.uu.se">tobias.ekholm@math.uu.se</a>	Jessica S Purcell	Monash University <a href="mailto:jessica.purcell@monash.edu">jessica.purcell@monash.edu</a>
Mario Eudave-Muñoz	Univ. Nacional Autónoma de México <a href="mailto:mario@matem.unam.mx">mario@matem.unam.mx</a>	Birgit Richter	Universität Hamburg <a href="mailto:birgit.richter@uni-hamburg.de">birgit.richter@uni-hamburg.de</a>
David Futер	Temple University <a href="mailto:dfuter@temple.edu">dfuter@temple.edu</a>	Jérôme Scherer	École Polytech. Féd. de Lausanne <a href="mailto:jerome.scherer@epfl.ch">jerome.scherer@epfl.ch</a>
John Greenlees	University of Warwick <a href="mailto:john.greenlees@warwick.ac.uk">john.greenlees@warwick.ac.uk</a>	Zoltán Szabó	Princeton University <a href="mailto:szabo@math.princeton.edu">szabo@math.princeton.edu</a>
Matthew Hedden	Michigan State University <a href="mailto:mhedden@math.msu.edu">mhedden@math.msu.edu</a>	Maggy Tomova	University of Iowa <a href="mailto:maggy-tomova@uiowa.edu">maggy-tomova@uiowa.edu</a>
Kristen Hendricks	Rutgers University <a href="mailto:kristen.hendricks@rutgers.edu">kristen.hendricks@rutgers.edu</a>	Daniel T Wise	McGill University, Canada <a href="mailto:daniel.wise@mcgill.ca">daniel.wise@mcgill.ca</a>
Hans-Werner Henn	Université Louis Pasteur <a href="mailto:henn@math.u-strasbg.fr">henn@math.u-strasbg.fr</a>	Lior Yanovski	Hebrew University of Jerusalem <a href="mailto:lior.yanovski@gmail.com">lior.yanovski@gmail.com</a>
Kathryn Hess	École Polytechnique Féd. de Lausanne <a href="mailto:kathryn.hess@epfl.ch">kathryn.hess@epfl.ch</a>		

---

See inside back cover or [msp.org/agt](https://msp.org/agt) for submission instructions.


The subscription price for 2026 is US \$795/year for the electronic version, and \$1170/year (+\$80, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Algebraic & Geometric Topology is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [Current Mathematical Publications](#) and the [Science Citation Index](#).

Algebraic & Geometric Topology (ISSN 1472-2747 printed, 1472-2739 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004.

---

AGT peer review and production are managed by EditFlow<sup>®</sup> from MSP.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<https://msp.org/>

© 2026 Mathematical Sciences Publishers

# ALGEBRAIC & GEOMETRIC TOPOLOGY

Volume 26

Issue 5 (pages 1597–1963)

2026

---

On homology concordance in contractible manifolds and two-bridge links	1597
HUGO ZHOU	
On the mapping class groups of simply connected smooth 4-manifolds	1635
DAVID BARAGLIA	
Negative-definite spin filling and branched double covers	1655
SOHEIL AZARPENDAR	
On local fibrations of $(\infty, 2)$ -categories	1681
FERNANDO ABELLÁN	
Brauer–Wall groups and truncated Picard spectra of $K$ -theory	1749
JONATHAN BEARDSLEY, KIRAN LUECKE and JACK MORAVA	
Polyhedral coproducts	1781
STEVEN AMELOTTE, WILLIAM HORNSLIEN and LEWIS STANTON	
Homoclinic leaves, Hausdorff limits and homeomorphisms	1801
IAN BIRINGER and CYRIL LECUIRE	
Motivic real topological Hochschild spectrum	1867
DOOSUNG PARK	
A note on the involutive invariants of splices	1907
KRISTEN HENDRICKS, MATTHEW STOFFREGEN and IAN ZEMKE	
Lagrangian metric geometry with Riemannian bounds	1923
JEAN-PHILIPPE CHASSÉ	