

ANNALS OF K-THEORY

no. 2 vol 1 2016

**The surgery exact sequence, K-theory
and the signature operator**

Paolo Piazza and Thomas Schick



A JOURNAL OF THE K-THEORY FOUNDATION

The surgery exact sequence, K-theory and the signature operator

Paolo Piazza and Thomas Schick

The main result of this paper is a new and direct proof of the natural transformation from the surgery exact sequence in topology to the analytic K-theory sequence of Higson and Roe.

Our approach makes crucial use of analytic properties and new index theorems for the signature operator on Galois coverings with boundary. These are of independent interest and form the second main theme of the paper. The main technical novelty is the use of large-scale index theory for Dirac-type operators that are perturbed by lower-order operators.

1. Introduction	109
2. Index and rho-classes defined by perturbations	113
3. Delocalized APS-index theorem for perturbed operators	127
4. Mapping the surgery sequence to K-Theory	129
5. Proof of the delocalized APS index theorem	137
Acknowledgments	152
References	153

1. Introduction

Let V be a smooth, closed, oriented n -dimensional manifold. We consider $\Gamma := \pi_1(V)$ and the universal cover $\tilde{V} \rightarrow V$. Finally, we let $\tilde{u} : \tilde{V} \rightarrow E\Gamma$ be a Γ -equivariant map covering a classifying map $u : V \rightarrow B\Gamma$ for $\tilde{V} \rightarrow V$. Associated to these data there are two important exact sequences.

The first one, due to Browder, Novikov, Sullivan and Wall, is the surgery exact sequence in topology [Wall 1999; Ranicki 2002; Lück 2002]:

$$\cdots \rightarrow L_{n+1}(\mathbb{Z}\Gamma) \dashrightarrow \mathcal{S}(V) \rightarrow \mathcal{N}(V) \rightarrow L_n(\mathbb{Z}\Gamma). \quad (1.1)$$

The central object of interest in this sequence is the *structure set* $\mathcal{S}(V)$; elements in the set $\mathcal{S}(V)$ are given by homotopy manifold structures on V , i.e., orientation-preserving homotopy equivalences $f : M \rightarrow V$, with M a smooth oriented closed

MSC2010: 46L80, 46L87, 58J22.

Keywords: signature operator, K-theory, exact surgery sequence, index classes, rho-classes.

manifold, considered up to h -cobordism. $\mathcal{N}(V)$ is the set of degree-one normal maps $f : M \rightarrow V$ considered up to normal bordism. Finally, the abelian groups $L_*(\mathbb{Z}\Gamma)$, the L -groups of the integral group ring $\mathbb{Z}\Gamma$, are defined algebraically but have a geometric realization as cobordism groups of manifolds with boundary with additional structure on the boundary.

The surgery exact sequence (1.1) plays a fundamental role in the classification of high-dimensional smooth compact manifolds.

The second exact sequence associated to $\tilde{V} \rightarrow V$ is purely analytic and is due to Higson and Roe. Consider the C^* -algebra $C^*(\tilde{V})^\Gamma$, called the Roe algebra and obtained as the closure of the Γ -equivariant locally compact finite propagation operators; this is an ideal in $D^*(\tilde{V})^\Gamma$, the C^* -algebra obtained as the closure of the Γ -equivariant pseudolocal finite propagation operators. There is a short exact sequence of C^* -algebras

$$0 \rightarrow C^*(\tilde{V})^\Gamma \rightarrow D^*(\tilde{V})^\Gamma \rightarrow D^*(\tilde{V})^\Gamma / C^*(\tilde{V})^\Gamma \rightarrow 0$$

and thus a six-term long exact sequence in K -theory

$$\begin{aligned} \cdots \rightarrow K_{n+1}(C^*(\tilde{V})^\Gamma) &\rightarrow K_{n+1}(D^*(\tilde{V})^\Gamma) \\ &\rightarrow K_{n+1}(D^*(\tilde{V})^\Gamma / C^*(\tilde{V})^\Gamma) \xrightarrow{\partial} K_n(C^*(\tilde{V})^\Gamma) \rightarrow \cdots . \end{aligned}$$

There are canonical isomorphisms

$$K_{*+1}(D^*(\tilde{V})^\Gamma / C^*(\tilde{V})^\Gamma) = K_*(V) \quad \text{and} \quad K_*(C^*(\tilde{V})^\Gamma) = K_*(C_r^*\Gamma),$$

with $C_r^*\Gamma$ denoting, as usual, the reduced C^* -algebra of the group Γ . Thus we can rewrite the long exact sequence in K -theory as

$$\cdots \rightarrow K_{n+1}(C_r^*\Gamma) \rightarrow K_{n+1}(D^*(\tilde{V})^\Gamma) \rightarrow K_n(V) \rightarrow K_n(C_r^*\Gamma) \rightarrow \cdots . \quad (1.2)$$

This is the *analytic surgery sequence* of Higson and Roe.

By [Roe 2002], the map $K_*(V) \rightarrow K_*(C_r^*\Gamma)$ is precisely the (Baum–Connes) assembly map. This connects the Higson–Roe surgery sequence to fundamental questions such as the strong Novikov conjecture or the Baum–Connes conjecture.

Higson and Roe [2005a; 2005b; 2005c] constructed the following remarkable commutative diagram:

$$\begin{array}{ccccccc} L_{n+1}(\mathbb{Z}\Gamma) & \dashrightarrow & \mathcal{G}(V) & \longrightarrow & \mathcal{N}(V) & \longrightarrow & L_n(\mathbb{Z}\Gamma) \\ \downarrow \gamma & & \downarrow \alpha & & \downarrow \beta & & \downarrow \gamma \\ K_{n+1}(C_r^*\Gamma)[\frac{1}{2}] & \longrightarrow & K_{n+1}(D^*(\tilde{V})^\Gamma)[\frac{1}{2}] & \longrightarrow & K_n(V)[\frac{1}{2}] & \longrightarrow & K_n(C_r^*\Gamma)[\frac{1}{2}] \end{array} \quad (1.3)$$

where $A[\frac{1}{2}]$ is a shorthand for $A \otimes_{\mathbb{Z}} \mathbb{Z}[\frac{1}{2}]$ if A is any abelian group.

It is important to mention that both α and γ in (1.2) are constructed using fine properties of Poincaré spaces that are *not* smooth manifolds. The main ingredient is the use of homotopy equivalences to glue manifolds with boundary along their boundaries. The resulting objects are *not* manifolds, but still Poincaré complexes. These have well-defined (higher) signatures which then feature in the construction.

One main goal of this article is to give an alternative and direct route to the transformation from the smooth surgery exact sequence in topology to the analytic surgery sequence of Higson and Roe. The main point of our construction is that we use index-theoretic constructions, applied to the signature operator, throughout.

Our approach follows the one presented in our paper [PS 2014], where we showed how to map the exact sequence of Stolz for positive scalar curvature metrics to the Higson–Roe exact sequence. Throughout, we will follow the notations of our paper [PS 2014]; the current paper should be considered as a companion to the latter one, with the signature operator replacing the Dirac operator in a fundamental way.

Our main result, Theorem 4.10, is that there are natural (index-theoretic) maps Ind_Γ , ρ , β making the following diagram commutative:

$$\begin{array}{ccccccc}
 L_{n+1}(\mathbb{Z}\Gamma) & \dashrightarrow & \mathcal{S}(V) & \longrightarrow & \mathcal{N}(V) & \longrightarrow & L_n(\mathbb{Z}\Gamma) \\
 \downarrow \text{Ind}_\Gamma & & \downarrow \rho & & \downarrow \beta & & \downarrow \text{Ind}_\Gamma \\
 K_{n+1}(C_r^*\Gamma)[\frac{1}{2}] & \longrightarrow & K_{n+1}(D^*(\tilde{V})^\Gamma)[\frac{1}{2}] & \longrightarrow & K_n(V)[\frac{1}{2}] & \longrightarrow & K_n(C_r^*\Gamma)[\frac{1}{2}]
 \end{array} \tag{1.4}$$

The main technical novelty, compared to the companion paper [PS 2014], is that we have to use an orientation-preserving homotopy $f : M \rightarrow V$ to perturb the signature operator (on the disjoint union of M and V) to an invertible operator. We follow an explicit recipe for such a perturbation C_f initiated in the fundamental work [Hilsum and Skandalis 1992], and then modified in [PS 2007] and further modified in [Wahl 2013b]. We will use the latter version of this perturbation and for this reason we name it Wahl’s perturbation. One advantage of choosing Wahl’s perturbation is that it makes it possible to extend most of the results of Hilsum and Skandalis from closed manifolds to manifolds with boundary. See [Wahl 2013b, Theorem 8.4].

Notice that the perturbed operator loses several of the appealing properties of a Dirac-type operator (like unit propagation of the associated wave operator). Therefore, the second main theme of this work is the proof of a new coarse index theorem for the signature operator on Galois coverings with boundary. This is of independent interest. The main technical novelty is the use of large-scale index theory for Dirac-type operators that are perturbed by lower-order operators.

More explicitly, in (1.4) the map ρ is defined applying the following idea: if $f : M \rightarrow V$ is a homotopy equivalence then $\rho[f : M \rightarrow V]$ is defined in terms of

the projection onto the positive part of the spectrum of the self-adjoint invertible operator $D + C_f$, where D is the signature operator on the Galois covering defined by $(f \circ u) \sqcup (-u) : M \sqcup (-V) \rightarrow B\Gamma$ and C_f is Wahl's perturbation defined by f .

A geometrically given cycle for $L_n(\mathbb{Z}\Gamma)$ consists in particular of a manifold with boundary (made up of two components), and the extra datum of a homotopy equivalence of the boundaries of the pieces. Wahl's perturbation for this homotopy equivalence can then be used to perturb the signature operator to be invertible at the boundary. This allows for the definition of a generalized Atiyah–Patodi–Singer index class $\text{Ind}(D, f)$ in $K_{n+1}(C_r^*\Gamma)$, which in the end defines the map Ind_Γ .

Finally, β is defined as in Higson and Roe: if $f : M \rightarrow V$ defines a class in $\mathcal{N}(V)$ then its image through β is obtained as $f_*([D_M]) - [D_V] \in K_n(V)$, with $[D_M]$ and $[D_V]$ the fundamental classes associated to the signature operators on the smooth compact manifolds M and V .

We shall first treat the case in which V is odd-dimensional and only at the end indicate how the results of this paper and of [PS 2014] can also be extended to the even-dimensional case.

It should be added that the map Ind_Γ of our diagram, the one out of $L_*(\mathbb{Z}\Gamma)$, has already been constructed by Charlotte Wahl [2013b]. We will make use of her important results for parts of our program. The main novelty in the approach presented in this paper and in [Wahl 2013b], compared to [Higson and Roe 2005a; 2005b; 2005c], is therefore the definition of the map Ind_Γ , the definition of ρ and, crucially, the proof of well-definedness and of commutativity of the squares.

All this we prove by establishing and then employing a delocalized Atiyah–Patodi–Singer index theorem for *perturbed* Dirac operators:

Theorem 1.5 (Theorem 3.1). *Let W be an oriented manifold with free cocompact orientation-preserving action and with boundary $M_1 \sqcup M_2$. Let $f : M_1 \rightarrow M_2$ be an orientation-preserving Γ -equivariant homotopy equivalence. Then*

$$\iota_*(\text{Ind}(D, f)) = j_*(\rho(D_\partial + C_f)) \quad \text{in } K_0(D^*(\tilde{W})^\Gamma).$$

Here D is the signature operator on W ; $\text{Ind}(D, f)$ is the generalized APS index class associated to D and to the homotopy equivalence f ;

$$j : D^*(\partial\tilde{W})^\Gamma \rightarrow D^*(\tilde{W})^\Gamma$$

is the homomorphism induced by the inclusion $\partial\tilde{W} \rightarrow \tilde{W}$; and

$$\iota : C^*(\tilde{W})^\Gamma \rightarrow D^*(\tilde{W})^\Gamma$$

is the inclusion.

It should be added that this result generalizes to general Dirac-type operators with more abstract boundary perturbations making the boundary operator invertible. See Theorem 3.1 for a precise statement.

The main novelty in Theorem 4.10, i.e., the construction and the commutativity of the diagram (1.4), compared to the analogous result in [PS 2014], is the treatment of the technicalities which arise when dealing with perturbed Dirac-type operators.

Remark 1.6. By using the signature operator of Hilsum and Teleman it is possible to extend our results to Lipschitz manifolds and thus, by Sullivan’s theorem, to the surgery exact sequence in the category TOP of topological manifolds. We refer the reader to [Zenobi 2015] where stability results for the topological structure set under products are also discussed.

Remark 1.7. Because of their fundamental result, Higson and Roe name the group $K_0(D_\Gamma^*)$ the *analytic structure group*; they denote it $\mathcal{S}_1(\Gamma)$. Similarly, the analytic structure group $\mathcal{S}_0(\Gamma)$ is, by definition, the group $K_1(D_\Gamma^*)$. Deeley and Goffeng [2014] have introduced a geometric structure group $\mathcal{S}_1^{\text{geo}}(\Gamma)$, with cycles defined à la Baum–Douglas. Using the results of the present paper, Deeley and Goffeng have proved that their geometric definition is isomorphic to the analytic one given by Higson and Roe.

2. Index and rho-classes defined by perturbations

2A. The index homomorphism in L-theory. The map

$$\text{Ind}_\Gamma : L_{n+1}(\mathbb{Z}\Gamma) \rightarrow K_{n+1}(C_r^*\Gamma)$$

has been defined by Wahl, building on results of Hilsum and Skandalis [1992] and the authors [PS 2007]. We briefly describe it. Assume that $n + 1$ is even. Recall (see for example [Higson and Roe 2005c, Chapter 4] and the references therein) that an element $x \in L_{n+1}(\mathbb{Z}\Gamma)$ is represented by a quadruple

$$(W, F, X \times [0, 1], u : X \rightarrow B\Gamma)$$

with W a cobordism between two smooth orientable manifolds $\partial_1 W$ and $\partial_2 W$, X a smooth orientable manifold, $F : (W, \partial W) \rightarrow (X \times [0, 1], \partial(X \times [0, 1]))$ a degree-one normal map of pairs, $f_1 := F|_{\partial_1 W}$ and $f_2 := F|_{\partial_2 W}$ oriented homotopy equivalences and $u : X \rightarrow B\Gamma$ a classifying map. Let $f = f_1 \sqcup f_2$ denote the restriction of F to ∂W . Consider $Z := W \sqcup X \times [0, 1]$, a manifold with boundary. Let \mathcal{D}_Z be the signature operator on Z with coefficients in the Mishchenko bundle defined by $u : X \rightarrow B\Gamma$ and $u \circ F : W \rightarrow B\Gamma$, i.e., the bundle obtained as pullback of the $C_r^*\Gamma$ -module bundle $E\Gamma \times_\Gamma C_r^*\Gamma$ over $B\Gamma$. Then, proceeding as in [Wahl 2013b], we can construct a smoothing perturbation \mathcal{C}_f of the boundary operator $\mathcal{D}_{\partial Z}$ with the property that $\mathcal{D}_{\partial Z} + \mathcal{C}_f$ is invertible. This perturbation, a smoothing version of the original

one defined in [Hilsum and Skandalis 1992], is, first of all, a self-adjoint bounded operator on the $C_r^*\Gamma$ -Hilbert module $\mathcal{E}_M := L^2(Z, \Lambda^*Z \otimes \mathcal{F}_M)$ with \mathcal{F}_M denoting the Mishchenko bundle. Moreover, it is an element in $\Psi_{C_r^*\Gamma}^{-\infty}(Z, \Lambda^*Z \otimes \mathcal{F}_M)$, the smoothing operators in the Mishchenko–Fomenko calculus. Wahl’s perturbation \mathcal{C}_f is an example of what is called, in the literature, a smoothing trivializing perturbation; more generally, a *trivializing perturbation* for \mathcal{D} is a self-adjoint $C_r^*\Gamma$ -bounded operator \mathcal{C} with the property that $\mathcal{D} + \mathcal{C}$ is invertible as an unbounded operator on the Hilbert module \mathcal{E}_M . The original perturbation defined by Hilsum and Skandalis is an example of such a perturbation; see [Zenobi 2015, Proposition 3.1] for a proof. Notice that in this paper only the smoothing trivializing perturbation \mathcal{C}_f will be used; this corresponds to the choice $\epsilon \in (0, +\infty)$ in Wahl’s treatment.

We extend Wahl’s perturbation \mathcal{C}_f in the obvious way to the cylinder $\mathbb{R} \times \partial Z$ (we extend it to be constant in the cylindrical direction) and then use a cutoff function in order to graft this operator to the manifold with cylindrical end Z_∞ associated to Z . We denote this global perturbation by $\mathcal{C}_{f,\infty}$; this is the global perturbation chosen by Wahl and it is the one we shall take. (In previous work on higher APS index theory the global perturbation $\mathcal{C}_{f,\infty}$ was chosen to be b -pseudodifferential; see [Melrose and Piazza 1997a; Leichtnam and Piazza 1998]. While this choice would simplify some of our arguments in Section 2C, it would eventually make the proof of our main theorem more involved; this is why we have chosen the perturbation just explained.) Proceeding as in [Leichtnam and Piazza 1998] and [Leichtnam et al. 2000, Theorem 10.1], one proves that there is a well-defined index class associated to $\mathcal{D}_\infty + \mathcal{C}_{f,\infty}$, with \mathcal{D} denoting the Mishchenko–Fomenko signature operator. The index class is an element in $K_{n+1}(C_r^*\Gamma)$. See Section 2C2 below for further details. Thus, to the quadruple $(W, F, X \times [0, 1], u : X \rightarrow B\Gamma)$ we associate $\text{Ind}(\mathcal{D}_\infty + \mathcal{C}_{f,\infty}) \in K_{n+1}(C_r^*\Gamma)$.

Theorem 2.1. *This construction induces a well-defined group homomorphism*

$$\text{Ind}_\Gamma : L_{n+1}(\mathbb{Z}\Gamma) \rightarrow K_{n+1}(C_r^*\Gamma).$$

Proof. This is proved in [Wahl 2013b, Theorem 9.1]. □

We give more information on this index class in Section 2C.

2B. rho-classes. In this subsection we first fix a Γ -manifold \tilde{V} with a free cocompact action of Γ with quotient V , a smooth compact manifold without boundary. We fix a Γ -invariant metric on \tilde{V} ; we also fix a Γ -equivariant hermitian vector bundle \tilde{E} on \tilde{V} with quotient E on V . We assume the existence of a Γ -equivariant Clifford structure on \tilde{E} and we denote by D the corresponding Dirac-type operator

on \tilde{V} ; this is a Γ -equivariant operator. Notice that we *do not* employ the tilde-notation for the operators on the covering. We denote by \mathcal{D} the induced operator in the Mishchenko–Fomenko calculus.

Recall the main players in the Higson–Roe surgery sequence. We have the C^* -algebra $C^*(\tilde{V})^\Gamma$, called Roe algebra, obtained as the closure of the locally compact Γ -invariant finite-propagation operators, and we have the C^* -algebra $D^*(\tilde{V})^\Gamma$, obtained as the closure of Γ -invariant pseudolocal finite-propagation operators. Recall also that $C^*(\tilde{V})^\Gamma$ is an ideal in $D^*(\tilde{V})^\Gamma$. We refer to the companion paper [PS 2014] for the precise definitions and for the notation we adopt. One of the extra subtleties is that one has to stabilize the bundles by tensoring with $l^2(\mathbb{N})$ for the definition in particular of $D^*(\tilde{V})^\Gamma$ and let operators act on one corner $V \otimes Ce_1 \subset V \otimes l^2(\mathbb{N})$. See for example [PS 2014, §1.2]. We will suppress this throughout in the notation.

2B1. Operators on the covering. Recall from [PS 2014] that given a cocompact Galois covering $\tilde{V} \rightarrow V$ there is an isomorphism

$$\mathbb{K}(\mathcal{E}_M) \cong C^*(\tilde{V})^\Gamma, \quad (2.2)$$

where we recall that \mathcal{E}_M is the Mishchenko $C_r^*\Gamma$ -Hilbert module $L^2(V, E \otimes \mathcal{F}_M)$, with $\mathcal{F}_M = \tilde{V} \times_\Gamma C_r^*\Gamma$, the Mishchenko bundle. Consider now V and \tilde{V} as above. Let $\mathcal{C} \in \Psi_{C_r^*\Gamma}^{-\infty}(V, E \otimes \mathcal{F}_M)$ be a smoothing trivializing perturbation for a Dirac-type operator $\mathcal{D} \in \text{Diff}_{C_r^*\Gamma}^1(V, E \otimes \mathcal{F}_M)$. Obviously, from the Mishchenko–Fomenko calculus, we have that $\Psi_{C_r^*\Gamma}^{-\infty}(V, E \otimes \mathcal{F}_M) \subset \mathbb{K}(\mathcal{E}_M)$. Using (2.2) we obtain immediately that \mathcal{C} defines an element C in $C^*(\tilde{V})^\Gamma$.

In what follows we have two goals in mind: on the one hand we wish to generalize the assignment of the operator C to any self-adjoint bounded operator \mathcal{C} (not necessarily a smoothing operator in the Mishchenko–Fomenko calculus); on the other hand we wish to give a precise definition for such a C . To this end, let $\pi : C_r^*\Gamma \rightarrow \mathcal{B}(l^2(\Gamma))$ be the left regular representation. Recall for example from [Schick 2005] that tensoring with π (a faithful representation) induces an isomorphism of right Hilbert Γ -modules

$$\mathcal{E}_M \otimes_\pi l^2(\Gamma) \rightarrow L^2(\tilde{V}, \Lambda^* \tilde{V}). \quad (2.3)$$

Associating to each $\mathcal{C} \in \mathbb{B}(\mathcal{E}_M)$ the operator $\mathcal{C} \otimes_\pi \text{Id}_{l^2(\Gamma)}$ in $\mathcal{B}(\mathcal{E}_M \otimes_\pi l^2(\Gamma))$, we define a homomorphism of C^* -algebras

$$\mathbb{B}(\mathcal{E}_M) \rightarrow \mathcal{B}(\mathcal{E}_M \otimes_\pi l^2(\Gamma)), \quad \mathcal{C} \mapsto \mathcal{C} \otimes_\pi \text{Id}_{l^2(\Gamma)}.$$

Notice that the right-hand side is Γ -equivariant, given that \mathcal{C} is $C_r^*\Gamma$ -linear. Thus, conjugating with (2.3), we obtain a C^* -homomorphism

$$L_\pi : \mathbb{B}(\mathcal{E}_M) \rightarrow \mathcal{B}(L^2(\tilde{V}, \Lambda^* \tilde{V}))^\Gamma, \quad \mathcal{C} \mapsto C_\pi.$$

Proposition 2.4. L_π sends $\mathbb{K}(\mathcal{E}_M)$ isomorphically onto $C^*(\tilde{V})^\Gamma$. Moreover, it sends $\mathbb{B}(\mathcal{E}_M)$ into $\mathfrak{M}(C^*(\tilde{V})^\Gamma)$, the multiplier algebra of $C^*(\tilde{V})^\Gamma$.

Proof. L_π is injective, given that π is faithful. The statement about $L_\pi(\mathbb{K}(\mathcal{E}_M))$ follows by looking at the image under L_π of a dense set in $\mathbb{K}(\mathcal{E}_M)$. We can choose, for example, $\Psi_{C_\Gamma}^{-\infty}$. If $A \in \Psi_{C_\Gamma}^{-\infty}$ then $L_\pi(A)$ is nothing but the associated Γ -compactly supported smoothing operator on the covering (see [Lott 1992, Proposition 6]) and these operators are dense in $C^*(\tilde{V})^\Gamma$. Thus L_π sends $\mathbb{K}(\mathcal{E}_M)$ isomorphically onto $C^*(\tilde{V})^\Gamma$. The second statement is classical: indeed, Kasparov has proved (see [Wegge-Olsen 1993]) that $\mathbb{B}(\mathcal{E}_M)$ is the multiplier algebra of $\mathbb{K}(\mathcal{E}_M)$. The statement now follows from general arguments; indeed, an injective representation ϕ of a C^* -algebra K into the bounded operators of a Hilbert space H extends uniquely to a representation $\tilde{\phi}$ of the multiplier algebra of K , and $\tilde{\phi}(\mathfrak{M}(K)) = \mathfrak{M}(\phi(K))$. \square

We are now ready to define precisely the bounded operator C , on the covering \tilde{V} , corresponding to a self-adjoint operator $\mathcal{C} \in \mathbb{B}(\mathcal{E}_M)$. We simply set

$$C := L_\pi(\mathcal{C}). \quad (2.5)$$

We know that this is an element in $\mathfrak{M}(C^*(\tilde{V})^\Gamma)$, the multiplier algebra of $C^*(\tilde{V})^\Gamma$. If \mathcal{C} is, in addition, smoothing, i.e., $\mathcal{C} \in \Psi_{C_\Gamma}^{-\infty}(V, E \otimes \mathcal{F}_M)$, then $C \in C^*(\tilde{V})^\Gamma$.

Remark 2.6. If we apply all this to the smoothing trivializing perturbation \mathcal{C}_f defined by a homotopy equivalence $f : M \rightarrow V$, we immediately see that \mathcal{C}_f defines a perturbation $C_f := L_\pi(\mathcal{C}_f)$ in $C^*(\tilde{Z})^\Gamma$, with $Z = M \cup (-V)$ and \tilde{Z} the Galois covering defined by $f \circ u \cup (-u) : M \cup (-V) \rightarrow B\Gamma$, u denoting a classifying map for the universal cover of V .

2B2. *The rho-class associated to a perturbation.* Let $\mathcal{D} \in \text{Diff}_{C_\Gamma}^1(V, E \otimes \mathcal{F}_M)$ be a Dirac-type operator as above. We first assume V and therefore \tilde{V} to be odd-dimensional. Assume that there exists a trivializing perturbation \mathcal{C} for \mathcal{D} . We recall that this means that \mathcal{C} is a self-adjoint bounded operator on \mathcal{E}_M , the Mishchenko C_Γ^* -Hilbert module, with the property that the self-adjoint regular operator $\mathcal{D} + \mathcal{C}$ is invertible. Following [Melrose and Piazza 1997a], it is proved in [Leichtnam and Piazza 2003] that this is true if and only if the index class $\text{Ind}(\mathcal{D}) \in K_1(C_\Gamma^*)$ vanishes.

For the proof of the following proposition see, e.g., the proof of [Leichtnam et al. 2000, Lemma 2.1].

Proposition 2.7. *Let D be the Dirac operator on \tilde{V} corresponding to \mathcal{D} . Let $C := L_\pi(\mathcal{C})$. Then $D + C$ is self-adjoint and L^2 -invertible.*

Next we recall a result implicitly proved in [Higson and Roe 2005a, Proposition 5.9].

Proposition 2.8. *Let D be a self-adjoint unbounded operator on $H := L^2(\tilde{V}, \tilde{E})$, as above. Let \mathfrak{A} be a C^* -algebra in $\mathfrak{B}(H)$ and let \mathfrak{J} be an ideal in \mathfrak{A} . Let \mathfrak{M} be the multiplier algebra of \mathfrak{J} . Assume that S is a self-adjoint operator in \mathfrak{M} , that the resolvent of D is in \mathfrak{J} and that $D(1 + D^2)^{-1/2}$ is in \mathfrak{A} . Then the resolvent of $D + S$ is in \mathfrak{J} and $(D + S)(1 + (D + S)^2)^{-1/2} \in \mathfrak{A}$. Consequently,*

- $f(D + S) \in \mathfrak{J}$ for all $f \in C_0(\mathbb{R})$, and
- $f(D + S) \in \mathfrak{A}$ for all $f \in C([-\infty, \infty])$.

Proposition 2.9. *Let \mathfrak{D} and D be as above; let \mathfrak{C} be a trivialisizing perturbation for \mathfrak{D} and let $C := L_\pi(\mathfrak{C}) \in \mathfrak{M}(C^*(\tilde{V})^\Gamma)$ be the corresponding perturbation of D on the covering. Then*

$$\frac{D + C}{|D + C|} \text{ is an element in } D^*(\tilde{V})^\Gamma.$$

Proof. Choose $\mathfrak{A} = D^*(\tilde{V})^\Gamma$ and $\mathfrak{J} = C^*(\tilde{V})^\Gamma$. Then we can apply Proposition 2.8, given that $C \in \mathfrak{M}(C^*(\tilde{V})^\Gamma)$ (see Proposition 2.4). We thus obtain that $\chi(D + C)$ is an element in $D^*(\tilde{V})^\Gamma$ for any chopping function χ ; choosing χ equal to ± 1 on the spectrum of $D + C$ we are done. \square

This brings us to the definition of rho-classes.

Definition 2.10. Let $\tilde{V} \rightarrow V$ be a Γ -covering of a smooth compact orientable odd-dimensional manifold V without boundary. Let D be a Γ -equivariant Dirac operator on \tilde{V} acting on the sections of a Γ -equivariant bundle \tilde{E} . Let \mathfrak{M} be the multiplier algebra of $C^*(\tilde{V})^\Gamma$. For any self-adjoint operator $A \in \mathfrak{M}$ with the property that $D + A$ is L^2 -invertible we can consider the operator $(D + A)/|D + A|$. Then, proceeding as above, we have that

$$\frac{D + A}{|D + A|} \text{ is an element in } D^*(\tilde{V})^\Gamma.$$

The rho-class associated to D and to the trivialisizing perturbation A is, by definition, the idempotent defined by the involution $(D + A)/|(D + A)|$:

$$\rho(D + A) := \left[\frac{1}{2} \left(\frac{D + A}{|D + A|} + 1 \right) \right] \in K_0(D^*(\tilde{V})^\Gamma). \quad (2.11)$$

In the even-dimensional case, with a $\mathbb{Z}/2$ -graded bundle $\tilde{E} = \tilde{E}^+ \oplus \tilde{E}^-$, we can proceed analogously, once we have a $\mathbb{Z}/2$ -graded trivialisizing self-adjoint perturbation $A \in \mathfrak{M}(C^*(\tilde{V})^\Gamma)$, namely

$$D + A = \begin{pmatrix} 0 & D^- + A^- \\ D^+ + A^+ & 0 \end{pmatrix}. \quad (2.12)$$

A necessary and sufficient condition for such a perturbation to exist is that $\text{Ind}(\mathcal{D}) = 0$ in $K_0(C_r^*\Gamma)$; see [Melrose and Piazza 1997b; Leichtnam and Piazza 2003]. In order to define the rho-class associated to (2.12) we consider the space J of Γ -equivariant bundle isometries from \tilde{E}^- to \tilde{E}^+ (more precisely, on their stabilizations, obtained by tensoring the two bundles with $l^2(\mathbb{N})$ (see the beginning of Section 2B); this makes sure that the bundles are trivial). Any element u in J induces in a natural way an isometry $U : L^2(\tilde{V}, \tilde{E}^-) \rightarrow L^2(\tilde{V}, \tilde{E}^+)$. It is obvious that U commutes with the action of Γ and that it covers the identity in the D^* -sense; see for example [PS 2014, §1]. Thus we can set

$$\rho(D + A) := [U\chi(D + A)_+] \in K_1(D^*(\tilde{V})^\Gamma), \quad (2.13)$$

with χ an odd chopping function equal to the sign function on the spectrum of $D + A$. Since J is isomorphic, once we stabilize, to the space of $U(H)$ -valued functions on the quotient $V = \tilde{V}/\Gamma$ (H a separable Hilbert space), and since $U(H)$ is contractible, we see immediately that also J is contractible. Thus, by homotopy invariance of K-theory, the right-hand side in (2.13) does not depend on the choice of u in J .¹

Finally, if we consider the canonical map $u : \tilde{V} \rightarrow E\Gamma$, then we define

$$\rho_\Gamma(D + A) := u_*\rho(D + A) \in K_*(D_\Gamma^*). \quad (2.14)$$

Remark 2.15. The rho-classes do depend, in general, on the choice of the trivializing perturbation A . This will be clear from our delocalized APS index Theorem 3.1 for perturbed operators.

Remark 2.16. Note that $K_*(D_\Gamma^*)$, by the Baum–Connes conjecture, is expected to vanish for torsion-free groups, but is often nonzero otherwise. Therefore, the universal ρ -class ρ_Γ is of interest essentially only if Γ is a group with nontrivial torsion.

2B3. Fundamental examples of rho-classes. We present two fundamental examples of rho-classes. As we shall see later, these examples enjoy strong stability properties with respect to the trivializing perturbation.

Definition 2.17. Let (V, g) be an oriented smooth Riemannian manifold without boundary with fundamental group Γ . Let $u : V \rightarrow B\Gamma$ be the classifying map for the universal cover of V . Let (M, h) be another oriented Riemannian manifold without boundary and assume that $f : M \rightarrow V$ is an oriented homotopy equivalence.

¹In our previous paper [PS 2014] we allowed for an arbitrary Γ -equivariant isometry $U : L^2(\tilde{V}, \tilde{E}^-) \rightarrow L^2(\tilde{V}, \tilde{E}^+)$ covering the identity in the D^* -sense; see the discussion leading to Definition 1.11. This is not correct, for we could then change at will the class on the right-hand side of (2.13) by composing on the left with an arbitrary Γ -equivariant unitary $V \in U(L^2(\tilde{V}, \tilde{E}^-))$, covering the identity in the D^* -sense.

We consider $Z = M \sqcup (-V)$ with the obvious classifying map $u_Z : Z \rightarrow B\Gamma$ induced by u and by $u \circ f$. We then obtain

$$\rho(D + C_f) \in K_{n+1}(D^*(\tilde{Z})^\Gamma) \quad \text{and} \quad \rho_\Gamma(D + C_f) \in K_{n+1}(D_\Gamma^*),$$

with D equal to the signature operator on the covering $\tilde{Z} := u_Z^* E\Gamma$ and C_f the smoothing trivializing perturbation defined by the homotopy equivalence f . We choose Wahl's version of this perturbation. Notice that there is an obvious Γ -equivariant map $\tilde{\phi} : \tilde{Z} \rightarrow \tilde{V}$, $\tilde{\phi} := \tilde{f} \cup \text{id}_{(-\tilde{v})}$. We set

$$\begin{aligned} \rho(f) &:= \tilde{\phi}_* \rho(D + C_f) \in K_{n+1}(D^*(\tilde{V})^\Gamma); \\ \rho_\Gamma(f) &:= \rho_\Gamma(D + C_f) \in K_{n+1}(D_\Gamma^*). \end{aligned} \tag{2.18}$$

Observe that, by functoriality, $\rho_\Gamma(f) = \tilde{u}_* \rho(f)$.

Proof. We have to argue that $\rho(f)$ and $\rho_\Gamma(f)$ do not depend on the choices involved. The signature operator on Z depends on the choice of the Riemannian metrics on M and V . The perturbation \mathcal{C}_f depends on several choices (see [Wahl 2013b] for details); it also depends on the choice of the metric on M and V . Wahl proves (compare [ibid., §4]) that two different choices can be joined by a path of invertible operators $D(t) + B(t)$; thus, according to (an easy extension of) Proposition 2.33 below, the rho-class $\rho(D + C_f) \in K_{n+1}(D^*(\tilde{Z})^\Gamma)$ is independent of the choice of the Riemannian metrics and of the choices we have made in defining the trivializing perturbation C_f . Consequently, also $\rho(f)$ and $\rho_\Gamma(f)$ are independent of these choices. \square

Example 2.19. Let (M, g) be a closed oriented Riemannian manifold and $\tilde{M} \rightarrow M$ a Γ -cover. Let $2m$ or $2m + 1$ be the dimension of M , depending whether M is even- or odd-dimensional. We assume that the Laplacian on differential forms on the covering is L^2 -invertible in degree m . We shall say briefly that the Laplacian on the covering is invertible in middle degree. By the homotopy-invariance of L^2 -Betti numbers and of the Novikov–Shubin invariants this is a homotopy-invariant condition. Let D be the signature operator on \tilde{M} and let \mathcal{D} be the associated Mishchenko–Fomenko operator. As explained in [Leichtnam and Piazza 2000] there is then a class of smoothing trivializing perturbations \mathcal{S} , that were named *symmetric* there, and that enjoy strong stability properties, as we shall explain in a moment. We define the rho-class of a manifold M satisfying the above condition as the $K_*(D^*(\tilde{M})^\Gamma)$ -class

$$\rho(D + \mathcal{S}) \quad \text{with } \mathcal{S} \text{ a symmetric trivializing perturbation,}$$

with $*$ = $\dim M + 1$. We also define

$$\rho_\Gamma(D + \mathcal{S}) = \tilde{u}_*(\rho(D + \mathcal{S})) \in K_*(D_\Gamma^*).$$

We show in Remark 3.8 that

$$\rho_\Gamma(D + \mathcal{S}) = \rho_\Gamma(D + \mathcal{S}') \in K_*(D_\Gamma^*)$$

if $\mathcal{S}, \mathcal{S}'$ are symmetric trivializing perturbations. (2.20)

This brings us to the following definition:

Let $\tilde{M} \rightarrow M$ be a Γ -cover and assume that the Laplacian on differential forms on \tilde{M} is invertible in middle degree. Let D be the signature operator on \tilde{M} . We then define $\rho_\Gamma(\tilde{M}) \in K_(D_\Gamma^*)$ as*

$$\rho_\Gamma(\tilde{M}) := \rho_\Gamma(D + \mathcal{S}) \quad \text{for any symmetric trivializing perturbation } \mathcal{S}.$$

The notation is justified because, as in the previous example, the right-hand side is also independent of the choice of metric. Working a bit harder, one can extend these results to odd-dimensional Galois coverings $\tilde{M} \rightarrow M$, $\dim M = 2m + 1$, with the property that in degree m the reduced and unreduced cohomology with values in the local system defined by the Mishchenko–Fomenko bundle are equal. See [Leichtnam et al. 2000; Wahl 2010].

2C. Index classes. The index map $\text{Ind}_\Gamma : L_{n+1}(\mathbb{Z}\Gamma) \rightarrow K_{n+1}(C_r^*\Gamma)$ is constructed using APS higher index theory in the Mishchenko–Fomenko framework. We wish to frame the construction of APS index classes on manifolds with boundary and with cylindrical ends in coarse index theory.

2C1. Coarse index classes on manifolds with cylindrical ends. We will now consider manifolds with boundary and with cylindrical ends. First we recall the basic notation.

Notation 2.21. Let (W, g) be a Riemannian manifold with boundary $M := \partial W$. We shall always assume g to have product structure near the boundary. We denote by W_∞ the manifold W with an infinite semi-cylinder $[0, \infty) \times \partial M$ attached to the boundary. If \tilde{W} is a Γ -covering of W , then we similarly denote by \tilde{W}_∞ the manifold obtained from \tilde{W} by attaching an infinite semi-cylinder.

We will denote by P_0 the multiplication operator on \tilde{W}_∞ defined by the characteristic function of the subset $[0, \infty) \times \partial\tilde{W}$ in \tilde{W}_∞ ; similarly, we denote by P_R the multiplication operator defined by the characteristic function of the subset $[R, \infty) \times \partial\tilde{W}$ in \tilde{W}_∞ . We have similar operators defined in $\mathbb{R} \times \partial\tilde{W}$ and with a small abuse of notation we employ the same symbols.

We assume W to be even-dimensional.

We consider a Γ -equivariant Dirac-type operator D on \tilde{W} , acting on the sections of a Γ -equivariant $\mathbb{Z}/2$ -graded Hermitian vector bundle \tilde{E} . We denote the boundary operator by D_∂ and the natural extension of D to \tilde{W}_∞ by D_∞ . Notice once again that we do not employ the tilde-notation for the operators on the covering.

We denote by \mathfrak{D} , \mathfrak{D}_∂ and \mathfrak{D}_∞ the corresponding operators in the Mishchenko–Fomenko calculus. We adopt the Clifford and grading conventions of [Wahl 2013b].

We consider C_∂ , a norm limit of Γ -equivariant self-adjoint bounded finite-propagation operators on $L^2(\partial\tilde{W})$ such that $D_\partial + C_\partial$ is L^2 -invertible. For example, C_∂ could be equal to $L_\pi(\mathcal{C}_\partial)$, with \mathcal{C}_∂ a smoothing trivializing perturbation for \mathfrak{D}_∂ . We now define a global perturbation on \tilde{W}_∞ .

Definition 2.22. We extend C_∂ on $\mathbb{R} \times \partial\tilde{W}$ to be constant in the \mathbb{R} direction. We fix an inward collar on \tilde{W} diffeomorphic to $[-1, 0] \times \tilde{W}$; we consider \tilde{W}_∞ ; then, using a cutoff function equal to 1 on $[0, \infty)$ and equal to 0 at -1 , we graft this operator to a bounded operator on \tilde{W}_∞ , denoted C_∞^+ , acting from the sections of \tilde{E}^+ to the sections of \tilde{E}^- . We set $C_\infty^- := (C_\infty^+)^*$, the formal adjoint of C_∞^+ , and we consider

$$C_\infty := \begin{pmatrix} 0 & C_\infty^- \\ C_\infty^+ & 0 \end{pmatrix}.$$

We consider the unbounded operator $D_\infty + C_\infty$, which is odd with respect to the $\mathbb{Z}/2$ -grading induced by \tilde{E} . The operator C_∂ is, by assumption, a norm limit of finite-propagation operators. Therefore C_∞ also has this property.

More generally, let B_∞ be a bounded self-adjoint odd operator on W_∞ with the following two properties:

$$\begin{aligned} B_\infty &\text{ is a norm limit of finite-propagation } \Gamma\text{-operators, and} \\ P_0 B_\infty P_0 - P_0 C_\infty P_0 &\in C^*(\tilde{W} \subset \tilde{W}_\infty)^\Gamma. \end{aligned} \tag{2.23}$$

Notice that C_∞ has these properties.

Lemma 2.24. *The operator B_∞ belongs to the multiplier algebra $\mathfrak{M}(C^*(\tilde{W}_\infty)^\Gamma)$.*

Proof. Indeed, as is well known, every operator which is a norm limit of Γ -equivariant finite-propagation operators is a multiplier of $C^*(\tilde{W}_\infty)^\Gamma$. We give for completeness the easy argument: if A is such an operator with propagation R and if ϕ is a compactly supported function, then $\phi A = \phi A \psi$ for every compactly supported ψ which is equal to 1 on the R -neighborhood of the support of ϕ . Consequently, if $E \in C^*(\tilde{W}_\infty)^\Gamma$, then $\phi A E = \phi A \psi E$ is compact given that ψE is compact. Passing to norm limits, the general statement follows. \square

We can now apply the Higson–Roe result, as stated in Proposition 2.8, with $\mathfrak{A} = D^*(\tilde{W}_\infty)^\Gamma$, $\mathfrak{J} = C^*(\tilde{W}_\infty)^\Gamma$ and $S = B_\infty \in \mathfrak{M}(C^*(\tilde{W}_\infty)^\Gamma)$, obtaining:

Lemma 2.25. *For every $f \in C_0(\mathbb{R})$ we get $f(D_\infty + B_\infty) \in C^*(\tilde{W}_\infty)^\Gamma$. If $f \in C([-\infty, \infty])$ then $f(D_\infty + B_\infty) \in D^*(\tilde{W}_\infty)^\Gamma$.*

From Lemma 2.25 we learn that if χ is a chopping function then $\chi(D_\infty + B_\infty)$ is an element in $D^*(\tilde{W}_\infty)^\Gamma$ and an involution modulo $C^*(\tilde{W}_\infty)^\Gamma$. However, more is true.

Proposition 2.26. *If C_∂ and C_∞ are as above, if B_∞ is as in (2.23) and χ is a chopping function equal to the sign function on the spectrum of $D_\partial + C_\partial$, then $\chi(D_\infty + B_\infty)$ is an involution modulo $C^*(\tilde{W} \subset \tilde{W}_\infty)^\Gamma$.*

Proof. Basically, we prove this by comparing $\chi(D_\infty + B_\infty)$ to the corresponding operator, $D_{\text{cyl}} + C_\partial \otimes \text{Id}_\mathbb{R}$, on the two-sided cylinder $\partial\tilde{W} \times \mathbb{R}$.

We start with some general comparison results. For simplification, we use the following notation:

- We use D instead of D_∞ , and \bar{D} for the corresponding (translation-invariant) operator on $\mathbb{R} \times \partial\tilde{W}$.
- We use C instead of C_∞ and \bar{C} for the translation-invariant perturbation of \bar{D} on $\mathbb{R} \times \partial\tilde{W}$; note that then $\bar{D} + \bar{C}$ is invertible.
- We use B instead of B_∞ .
- We choose a cutoff function χ which is equal to ± 1 on the spectrum of $\bar{D} + \bar{C}$, setting $e := 1 - \chi^2$; then $e(\bar{D} + \bar{C}) = 0$.
- The subspace $[0, \infty) \times \partial\tilde{W}$ is contained isometrically in \tilde{W}_∞ and in $\mathbb{R} \times \partial\tilde{W}$; we freely identify sections supported on this part of both manifolds. We employ the multiplication operator P_R for $R \geq 0$; we remark that this is nothing but the orthogonal projection onto the sections supported on $[R, \infty) \times \partial\tilde{W}$. Once again, with a small abuse of notation, we employ the same symbols for this operator in \tilde{W}_∞ , and in $\mathbb{R} \times \partial\tilde{W}$. This way, in particular, we consider an operator like $P_0 f(\bar{D}) P_R$ as acting on $L^2(\tilde{W}_\infty)$ (instead of $L^2(\partial\tilde{W} \times \mathbb{R})$).

Lemma 2.27. *For $f \in C_0(\mathbb{R})$, we have*

$$P_0 f(D) P_0 - P_0 f(\bar{D}) P_0 \in C^*(\tilde{W} \subset \tilde{W}_\infty)^\Gamma.$$

Proof. We know that $f(D) \in C^*(\tilde{W}_\infty)^\Gamma$ and $f(\bar{D}) \in C^*(\mathbb{R} \times \partial\tilde{W})^\Gamma$ and that P_0 is in D^* . So it only remains to check the support condition, i.e., to show that for each $\epsilon > 0$ there is an $R > 0$ such that $\|P_0 f(D) P_R - P_0 f(\bar{D}) P_R\| < \epsilon$ (then also its adjoint $P_R f(D) P_0 - P_R f(\bar{D}) P_0$ has norm $< \epsilon$). For the latter, we approximate f in supremum norm by a function g whose Fourier transform has compact support (say in $[-R, R]$), and therefore approximate $f(D)$ in operator norm by $g(D)$ (and $f(\bar{D})$ by $g(\bar{D})$). The usual Fourier inversion formula then implies, using unit propagation speed for the wave operators of D and \bar{D} , that $P_0 g(D) P_R = P_0 g(\bar{D}) P_R$, and the statement follows. \square

We now generalize Lemma 2.27 to $D + B$. Note that we don't have unit propagation speed available. Instead, we give a proof which uses a comparison between $f(D)$ and $f(D + B)$.

Lemma 2.28. *For $f \in C_0(\mathbb{R})$ and B as in (2.23) we have that*

$$P_0 f(D + B)P_0 - P_0 f(\bar{D} + \bar{C})P_0 \in C^*(\tilde{W} \subset \tilde{W}_\infty)^\Gamma.$$

Proof. For the following simple argument, using the Neumann series, we are grateful to the referee. We have to show that for $f \in C_0(\mathbb{R})$ it holds that

$$\lim_{R \rightarrow \infty} \|P_R f(D + B)P_0 - P_R f(\bar{D} + \bar{C})P_0\| = 0; \quad (2.29)$$

the corresponding statement then holds for the adjoint, involving $P_0 f(D + B)P_R$.

Recall that if X is a norm limit of bounded finite-propagation operators then $\lim_{|R-S| \rightarrow \infty} \|(1 - P_S)X P_R\| = 0$. Therefore, if A, \bar{A}, B, \bar{B} are norm limits of Γ -equivariant finite-propagation operators and if

$$P_0 A P_0 - P_0 \bar{A} P_0, P_R B P_0 - P_R \bar{B} P_0 \in C^*(\tilde{W} \subset \tilde{W}_\infty)^\Gamma$$

in norm, then also

$$\begin{aligned} & P_{2R} A B P_0 - P_{2R} \bar{A} \bar{B} P_0 \\ &= (P_{2R} A P_R - P_{2R} \bar{A} P_R) P_R B P_0 + P_{2R} \bar{A} P_R (P_R B P_0 - P_R \bar{B} P_0) \\ &\quad + P_{2R} A ((1 - P_R) B P_0) - P_{2R} \bar{A} ((1 - P_R) \bar{B} P_0) \\ &\xrightarrow{R \rightarrow \infty} 0 \end{aligned}$$

since all four expressions in parentheses go to zero as $R \rightarrow \infty$.

Consequently, if (2.29) is true for f_1 and f_2 then it is also true for their product $f_1 f_2$. As a C^* -algebra, $C_0(\mathbb{R})$ is for any $\lambda > 0$ generated by the two functions $x \mapsto (x + \lambda i)^{-1}$ and $x \mapsto (x - \lambda i)^{-1}$. This means that it suffices to treat these two functions for $\lambda > 0$ large, and we concentrate for notational convenience on the first.

As λ is sufficiently large, we can then use the Neumann series

$$\begin{aligned} (D + B + \lambda i)^{-1} &= (D + \lambda i)^{-1} \sum_{k=0}^{\infty} (-B(D + \lambda i)^{-1})^k, \\ (\bar{D} + \bar{B} + \lambda i)^{-1} &= (\bar{D} + \lambda i)^{-1} \sum_{k=0}^{\infty} (-\bar{B}(\bar{D} + \lambda i)^{-1})^k. \end{aligned}$$

By Lemma 2.27,

$$P_0 (D + \lambda I)^{-1} P_0 - P_0 (\bar{D} + \lambda i)^{-1} P_0 \in C^*(\tilde{W} \subset \tilde{W}_\infty)^\Gamma,$$

and, by (2.23),

$$P_0 B P_0 - P_0 \bar{B} P_0 \in C^*(\tilde{W} \subset \tilde{W}_\infty)^\Gamma.$$

Using (2.29) we can pass to products and finite sums, showing that for each $N \in \mathbb{N}$ also

$$P_0 \left((D + \lambda i)^{-1} \sum_{k=0}^N (-B(D + \lambda i)^{-1})^k \right) P_0 \\ - P_0 \left((\bar{D} + \lambda i)^{-1} \sum_{k=0}^N (-\bar{B}(\bar{D} + \lambda i)^{-1})^k \right) P_0 \in C^*(\tilde{W} \subset \tilde{W}_\infty)^\Gamma.$$

Because of the norm convergence of the Neumann series, we also have

$$P_0(D + B + \lambda i)^{-1} P_0 - P_0(\bar{D} + \bar{B} + \lambda i)^{-1} P_0 \in C^*(\tilde{W} \subset \tilde{W}_\infty)^\Gamma. \quad \square$$

To finish the proof of Proposition 2.26 we only have to prove the support condition for $e(D + B)$, with $e(x) = 1 - \chi(x)^2$ as above; in particular, $e \in C_0(\mathbb{R})$. It follows from Lemma 2.28 that

$$P_0 e(D + B) P_0 - P_0 e(\bar{D} + \bar{C}) P_0 \in C^*(\tilde{W} \subset \tilde{W}_\infty)^\Gamma,$$

and by our choice of χ we have $e(\bar{D} + \bar{C}) = 0$. Finally, by finite propagation and the definition of P_0 , also $P_0 e(D + B)(1 - P_0)$, $(1 - P_0)e(D + B)(1 - P_0)$, $(1 - P_0)e(D + B)P_0 \in C^*(\tilde{W} \subset \tilde{W}_\infty)^\Gamma$. \square

Definition 2.30. These considerations imply that there is a well-defined coarse relative index class

$$\text{Ind}^{\text{rel}}(D_\infty + C_\infty) \in K_0(C^*(\tilde{W} \subset \tilde{W}_\infty)^\Gamma). \quad (2.31)$$

It is obtained by the standard construction in coarse index theory: with an isometry $U : L^2(\tilde{W}_\infty, \tilde{E}^-) \rightarrow L^2(\tilde{W}_\infty, \tilde{E}^+)$ which is Γ -equivariant and which covers the identity in the D^* -sense (see [PS 2014, §1]) the operator $U\chi(D)_+$ is invertible in $D^*(\tilde{W}_\infty)^\Gamma / C^*(\tilde{W} \subset \tilde{W}_\infty)^\Gamma$ and therefore defines an element in K_1 of this C^* -algebra. $\text{Ind}^{\text{rel}}(D_\infty + C_\infty)$ is its image under the boundary map of the associated long exact sequence in K-theory. We are also interested in the associated *coarse index class*

$$\text{Ind}(D, C) := c_*^{-1} \text{Ind}^{\text{rel}}(D_\infty + C_\infty) \in K_0(C^*(\tilde{W})^\Gamma). \quad (2.32)$$

Here we use the canonical inclusion $c : C^*(W)^\Gamma \rightarrow C^*(W \subset W_\infty)^\Gamma$ which induces an isomorphism in K-theory; see [PS 2014, Lemma 1.9].

Note that the right-hand side of (2.32) is just a notation; we have not really defined an operator C .

Proposition 2.33. *If B_∞ is the norm limit of Γ -equivariant bounded finite-propagation operators, C_∂ is a perturbation as above and we assume that*

$$P_0 B_\infty P_0 - P_0 C_\infty P_0 \in C^*(\tilde{W}_\infty \subset \tilde{W})^\Gamma,$$

then

$$\mathrm{Ind}^{\mathrm{rel}}(D_\infty + B_\infty) = \mathrm{Ind}^{\mathrm{rel}}(D_\infty + C_\infty).$$

If $[0, 1] \ni t \mapsto C_\partial^t$ is a continuous family of perturbations such that $D_\partial + C_\partial^t$ is invertible for all t , then $\mathrm{Ind}^{\mathrm{rel}}(D_\infty + C_\infty^t)$ is independent of t .

Finally, if $\tilde{M} \rightarrow M$ is a Γ -Galois covering of a closed compact manifold M and if $[0, 1] \ni t \mapsto G_t$ is a continuous family of trivializing perturbations for D , as in Definition 2.10, then $\rho(D + G_t)$ and $\rho_\Gamma(D + G_t)$ are independent of t .

Proof. For the first two statements, write either $A_t := tB_\infty + (1-t)C_\infty$ or $A_t = C_\infty^t$. Then $[0, 1] \ni t \mapsto D_\infty + A_t$ is norm resolvent continuous, as

$$(D_\infty + A_t + i)^{-1} - (D_\infty + A_s + i)^{-1} = (D_\infty + A_t + i)^{-1}(A_s - A_t)(D_\infty + A_s + i)^{-1}$$

and $\|(D_\infty + A_t + i)^{-1}\| \leq 1$.

As the C^* -algebra $C[-\infty, \infty]$ of bounded continuous functions on \mathbb{R} with limits at $\pm\infty$ is generated by $(x+i)^{-1}$, $(x-i)^{-1}$, 1 and $\phi(x) = x/\sqrt{1+x^2}$, and the first two are taken care of by norm resolvent continuity, it remains to show that $t \mapsto \phi(D_\infty + A_t)$ is norm-continuous to conclude that $t \mapsto \chi(D_\infty + A_t)$ is norm-continuous for arbitrary $\chi \in C[-\infty, \infty]$.

To this end we write $\|\phi(D_\infty + A_t) - \phi(D_\infty + A_s)\|$ as $\|\phi(G) - \phi(G + S_{s,t})\|$, with $G = D_\infty + A_t$ and $S_{s,t} = A_s - A_t$. We employ the following identity; see [Higson and Roe 2005a, Proposition 5.9] (this identity plays a fundamental role in the proof of Proposition 2.8):

$$\begin{aligned} \phi(G) - \phi(G + S_{s,t}) &= \frac{1}{\pi} \int_1^{+\infty} \frac{u}{\sqrt{u^2-1}} R(u) S_{s,t} R_{s,t}(u) du \\ &\quad + \frac{1}{\pi} \int_1^{+\infty} \frac{u}{\sqrt{u^2-1}} R(-u) S_{s,t} R_{s,t}(-u) du, \end{aligned}$$

where $R(u) = (G + iu)^{-1}$ and $R_{s,t}(u) = (G + S_{s,t} + iu)^{-1}$. Now, the norm of either of the two resolvents is bounded by $1/|u|$; thus

$$\|\phi(G) - \phi(G + S_{s,t})\| \leq \frac{2}{\pi} \|A_s - A_t\| \left(\int_1^\infty \frac{u}{\sqrt{u^2-1}} \frac{1}{u^2} du \right).$$

This establishes the required continuity. In particular, for any chopping function χ , $t \mapsto \chi(D_\infty + A_t)$ is a *norm-continuous* family of involutions in the quotient $D^*(\tilde{W}_\infty)^\Gamma / C^*(\tilde{W}_\infty \subset \tilde{W})^\Gamma$. By homotopy-invariance of K-theory classes, the first two assertions follow.

A similar proof applies to the last statement once we observe that we can use the same chopping function taking values ± 1 on the spectrum of all (invertible!)

operators in the compact family $D_\infty + A_t$, $t \in [0, 1]$. This follows from the norm-continuity of $t \mapsto (D_\infty + A_t)^{-1}$, using

$$(D_\infty + A_s)^{-1} = (D_\infty + A_t)^{-1}(1 + (A_s - A_t)(D_\infty + A_t)^{-1})^{-1}$$

and a Neumann series argument for the last term. \square

2C2. Compatibility of index classes. We now prove the compatibility of this relative index class in coarse geometry with the index class in the Mishchenko–Fomenko framework.

Let D_∞ be a Dirac-type operator on \widetilde{W}_∞ and let \mathcal{D}_∞ denote the associated operator on W_∞ with coefficients in the Mishchenko bundle \mathcal{F}_M . Let \mathcal{C}_∂ be a trivializing perturbation for \mathcal{D}_∂ and let be $\mathcal{D}_\infty + \mathcal{C}_\infty$ the associated operator on W_∞ , obtained by applying the same construction explained above for the covering \widetilde{W} ; see Definition 2.22. We concentrate directly on the case in which \mathcal{C}_∂ is smoothing (as far as compatibility is concerned, this is the case we are interested in). By [Leichtnam and Piazza 1998, Theorem 6.2] and [Leichtnam et al. 2000, Theorem 10.1], $(\mathcal{D}_\infty + \mathcal{C}_\infty)^+$ is invertible modulo $\mathbb{K}(\mathcal{E}_M)$. Further extensions of these results were subsequently given in [Wahl 2013a]. Consequently, we get an index class

$$\text{Ind}_{\text{MF}}^b(\mathcal{D}_\infty + \mathcal{C}_\infty) \in K_{n+1}(\mathbb{K}(\mathcal{E}_M)) = K_{n+1}(C_r^*\Gamma)$$

Remark 2.34. We can apply this construction to the manifold with boundary appearing in the definition of a cycle of $L_{n+1}(\mathbb{Z}\Gamma)$, choosing as a Dirac operator \mathcal{D} the signature operator and as a trivializing perturbation for the boundary operator \mathcal{D}_∂ the operator \mathcal{C}_f , the one induced by a homotopy equivalence f . We obtain an index class $\text{Ind}_{\text{MF}}^b(\mathcal{D}_\infty + \mathcal{C}_{f,\infty}) \in K_{n+1}(C_r^*\Gamma)$; this is precisely the index class we have considered in Section 2A.

We have already observed in [PS 2014] that there is an isomorphism $\mathbb{K}(\mathcal{E}_M) \cong C^*(\widetilde{W} \subset \widetilde{W}_\infty)^\Gamma$. One can prove, as in [PS 2014, §2], the following proposition.

Proposition 2.35. *Under the canonical isomorphism*

$$K_{n+1}(\mathbb{K}(\mathcal{E}_M)) \cong K_{n+1}(C^*(\widetilde{W} \subset \widetilde{W}_\infty)^\Gamma),$$

one has the equality

$$\text{Ind}_{\text{MF}}^b(\mathcal{D}_\infty + \mathcal{C}_\infty) = \text{Ind}^{\text{rel}}(D_\infty + C_\infty).$$

Remark 2.36. The index class $\text{Ind}_{\text{MF}}^b(\mathcal{D}_\infty + \mathcal{C}_\infty)$ heavily depends on the choice of trivializing perturbation \mathcal{C}_∂ . (Consequently, the same is true for $\text{Ind}^{\text{rel}}(D_\infty + C_\infty)$.) Indeed, it is proved in [Leichtnam and Piazza 2003, Theorem 6], inspired by [Melrose and Piazza 1997a], that if \mathcal{C}'_∂ is a different perturbation then

$$\text{Ind}_{\text{MF}}^b(\mathcal{D}_\infty + \mathcal{C}_\infty) - \text{Ind}_{\text{MF}}^b(\mathcal{D}_\infty + \mathcal{C}'_\infty) = [\mathcal{P}' - \mathcal{P}] \quad \text{in } K_{\dim W}(C_r^*\Gamma), \quad (2.37)$$

with $[\mathcal{P}' - \mathcal{P}]$ the difference class of the two projections $\mathcal{P} = \chi_{[0,\infty)}(\mathcal{D}_\partial + \mathcal{C})$ and $\mathcal{P}' = \chi_{[0,\infty)}(\mathcal{D}_\partial + \mathcal{C}')$.

3. Delocalized APS-index theorem for perturbed operators

Let $(\tilde{W}, g_{\tilde{W}})$ be an even-dimensional Riemannian orientable manifold with boundary $\partial\tilde{W}$. Assume that Γ acts freely, cocompactly and isometrically on \tilde{W} . We consider the associated manifold with cylindrical ends \tilde{W}_∞ . Let C_∂ be a trivializing perturbation as before (i.e., the norm limit of Γ -equivariant bounded self-adjoint finite-propagation operators such that $D_\partial + C_\partial$ is L^2 -invertible) and let C_∞ be the associated perturbation on \tilde{W}_∞ , as in Section 2C1. Recall the coarse index class $\text{Ind}(D, C) \in K_0(C^*(\tilde{W})^\Gamma)$ defined in (2.32). Our main tool in this paper will be the following ‘‘delocalized APS-index theorem for perturbed operators’’.

Theorem 3.1. *The following equality holds:*

$$\iota_*(\text{Ind}(D, C)) = j_*(\rho(D_\partial + C_\partial)) \quad \text{in } K_0(D^*(\tilde{W})^\Gamma). \quad (3.2)$$

Here we use $j : D^*(\partial\tilde{W})^\Gamma \rightarrow D^*(\tilde{W})^\Gamma$, induced by the inclusion $\partial\tilde{W} \rightarrow \tilde{W}$, and the inclusion $\iota : C^*(\tilde{W})^\Gamma \rightarrow D^*(\tilde{W})^\Gamma$.

Corollary 3.3. *By functoriality, using the canonical Γ -map $u : \tilde{W} \rightarrow E\Gamma$ we have*

$$\iota_* u_*(\text{Ind}(D, C)) = \rho_\Gamma(D_\partial + C_\partial) \quad \text{in } K_0(D_\Gamma^*).$$

If we define $\text{Ind}_\Gamma(D, C) := u_*(\text{Ind}(D, C))$ in $K_0(C_\Gamma^*)$, then the last equation reads

$$\iota_*(\text{Ind}_\Gamma(D, C)) = \rho_\Gamma(D_\partial + C_\partial) \quad \text{in } K_0(D_\Gamma^*). \quad (3.4)$$

We prove these results in Section 5.

Remark 3.5. Consider B_∞ , a bounded self-adjoint Γ -equivariant odd operator on W_∞ satisfying (2.23), i.e., B_∞ is a norm limit of Γ -equivariant finite-propagation operators and $P_0 B_\infty P_0 - P_0 C_\infty P_0 \in C^*(\tilde{W} \subset \tilde{W}_\infty)^\Gamma$. By Proposition 2.26, we have the coarse index class $\text{Ind}(D, B) := c_*^{-1} \text{Ind}^{\text{rel}}(D_\infty + B_\infty) \in K_{n+1}(C^*(\tilde{W})^\Gamma)$. Since $\text{Ind}(D, B) = \text{Ind}(D, C)$ by Proposition 2.33, we also have

$$\iota_*(\text{Ind}_\Gamma(D, B)) = \rho_\Gamma(D_\partial + C_\partial). \quad (3.6)$$

This remark brings us immediately to a version of bordism invariance for ρ -classes:

Corollary 3.7. *Let \tilde{M}_1 and \tilde{M}_2 be two free cocompact Γ -manifolds of dimension n . Let D_j be a Γ -equivariant Dirac-type operator on \tilde{M}_j , $C_j \in C^*(\tilde{M}_j)^\Gamma$ a trivializing perturbation ($j = 1, 2$). Assume that D_1 and D_2 are bordant, i.e., there exists a manifold \tilde{W} with free cocompact Γ -action such that $\partial\tilde{W} = \tilde{M}_1 \sqcup \tilde{M}_2$ and D a Γ -equivariant Dirac-type operator on W with product structure near the boundary (in the sense of Dirac-type operators) and with boundary operator*

$D_1 \sqcup -D_2$. Moreover, let B_∞ be a global perturbation which is a norm limit of finite-propagation operators and such that $P_0 B_\infty P_0 - P_0 C_\infty^\sqcup P_0 \in C^*(\tilde{W} \subset \tilde{W}_\infty)^\Gamma$, where $C_\infty^\sqcup := C_{1,\infty} \sqcup (-C_{2,\infty})$.

Assume in addition that $\text{Ind}_\Gamma(D, B) = 0$ in $K_{n+1}(C_\Gamma^*)$. (Note that this is satisfied, e.g., if B_∞ is a trivializing perturbation of D , i.e., if $D + B_\infty$ is invertible.) Then

$$\rho_\Gamma(D_1 + C_1) = \rho_\Gamma(D_2 + C_2) \in K_{n+1}(D_\Gamma^*).$$

Proof. The rho-class is additive for disjoint union. By assumption, $\text{Ind}_\Gamma(D, B) = 0$. The assertion now follows directly from (3.6). \square

Remark 3.8. We can now go back to Example 2.19 and explain why

$$\rho_\Gamma(D + \mathcal{S}_0) = \rho_\Gamma(D + \mathcal{S}_1) \quad \text{if } \mathcal{S}_0, \mathcal{S}_1 \text{ are two symmetric trivializing perturbations.}$$

Recall that we are considering $\tilde{M} \rightarrow M$, a Γ -cover such that the differential-form Laplacian on the covering is L^2 -invertible in degree m . The above equality takes place in $K_{\dim M+1}(D_\Gamma^*)$.

We consider $[0, 1] \times \tilde{M}$ with associated manifold with cylindrical ends. By the delocalized APS-index Theorem 3.1 we have that

$$\rho_\Gamma(D + \mathcal{S}_0) - \rho_\Gamma(D + \mathcal{S}_1) = \iota_*(\text{Ind}_\Gamma(D_\infty + \mathcal{S}_\infty^{0,1})),$$

with $\mathcal{S}_\infty^{0,1}$ the grafted perturbation associated to \mathcal{S}_0 on $\{0\} \times \tilde{M}$ and \mathcal{S}_1 on $\{1\} \times \tilde{M}$. Using in a fundamental way the hypothesis that the \mathcal{S}^j are symmetric we have (see [Leichtnam and Piazza 2000, Proposition 4.4; Wahl 2010])

$$\text{Ind}_\Gamma(D_\infty + \mathcal{S}_\infty^{0,1}) = \text{Ind}_\Gamma(D_\infty + \mathcal{S}_\infty^{0,0}) \quad \text{in } K_*(C_r^*\Gamma) = K_*(C_\Gamma^*).$$

Thus

$$\begin{aligned} & \rho_\Gamma(D + \mathcal{S}_0) - \rho_\Gamma(D + \mathcal{S}_1) \\ &= (\rho_\Gamma(D + \mathcal{S}_0) - \rho_\Gamma(D + \mathcal{S}_1)) - (\rho_\Gamma(D + \mathcal{S}_0) - \rho_\Gamma(D + \mathcal{S}_0)) \\ &= \iota_*(\text{Ind}_\Gamma(D_\infty + \mathcal{S}_\infty^{0,1})) - \iota_*(\text{Ind}_\Gamma(D_\infty + \mathcal{S}_\infty^{0,0})) = 0. \end{aligned}$$

Remark 3.9. As in [PS 2014], it is plausible that the methods which prove the delocalized APS-index theorem for perturbed operators also yield a secondary partitioned manifold index theorem.² We leave the precise formulation and proof to future investigations. A potential application of such a theorem would be the following: assume that M_1, M_2 are two oriented complete Riemannian manifolds with free proper isometric Γ -action and with Γ -invariant and Γ -cocompact separating

²We take this opportunity to point out that the secondary partitioned manifold index theorem proved in [PS 2014] can in fact be sharpened, with essentially the same proof, from an equality of *universal* rho-classes to an equality of rho-classes. We thank Vito Felice Zenobi and Rudolf Zeidler for alerting us to this sharpening.

hypersurfaces $N_1 \subset M_1$, $N_2 \subset M_2$ with product neighborhoods. Let $f : M_1 \rightarrow M_2$ be a Γ -equivariant homotopy equivalence which restricts to a homotopy equivalence $f| : N_1 \rightarrow N_2$. Assume that f coarsely preserves the signed distance to the hypersurfaces, i.e., that the two functions $d_1, d_2 \circ f : M_1 \rightarrow \mathbb{R}$ with $d_1(x) = \epsilon_1(x)d(x, N_1)$ and $d_2(y) = \epsilon_2(y)d(y, N_2)$ are coarsely equivalent to each other. Here $\epsilon_1(x) = \pm 1$ depending on whether x lies in the positive or negative half of M .

We call such a situation a “partitioned manifold homotopy equivalence”.

In this situation, possibly with further hypotheses on f , one should obtain a partitioned manifold rho-class $\rho^{\text{pm}}(f) \in K_*(D_\Gamma^*)$. Similarly, we defined the class $\rho_\Gamma(f|_{N_1}) \in K_*(D_\Gamma^*)$ in Definition 2.17. The secondary partitioned manifold index theorem would claim that these two classes coincide.

As a potential application, assume that $f : N_1 \rightarrow N_2$ and $g : X_1 \rightarrow N_2$ are homotopy equivalences and that $\rho_\Gamma(f) \neq \rho_\Gamma(g)$ so that

$$(N_1 \xrightarrow{f} N_2) \quad \text{and} \quad (X_1 \xrightarrow{g} N_2)$$

are not h-cobordant to one another. Then, also between the stabilizations

$$(N_1 \times \mathbb{R} \xrightarrow{f \times \text{id}_\mathbb{R}} N_2 \times \mathbb{R}) \quad \text{and} \quad (X_1 \times \mathbb{R} \xrightarrow{g \times \text{id}_\mathbb{R}} N_2 \times \mathbb{R})$$

there is no Riemannian h-cobordism such that the inclusion maps of the boundaries are continuous coarse homotopy equivalences. Otherwise, we expect

$$\rho^{\text{pm}}(f \times \text{id}_\mathbb{R}) = \rho^{\text{pm}}(g \times \text{id}_\mathbb{R});$$

however, by the secondary index theorem, this would contradict the hypothesis $\rho_\Gamma(f) \neq \rho_\Gamma(g)$.

4. Mapping the surgery sequence to K-Theory

4A. The structure set and the map ρ_Γ . Let V be an n -dimensional smooth closed oriented manifold with fundamental group Γ , $n > 4$.

Definition 4.1. The *structure set* $\mathcal{S}^h(V)$ consists of equivalence classes $[f : M \rightarrow V]$, where M is a smooth closed oriented manifold, and f is a homotopy equivalence. Two homotopy equivalences $f_1 : M_1 \rightarrow V$ and $f_2 : M_2 \rightarrow V$ are equivalent if there is an h-cobordism X between M_1 and M_2 and a map $F : X \rightarrow V \times [0, 1]$ (necessarily a homotopy equivalence) such that $F|_{M_1} = f_1$ and $F|_{M_2} = f_2$.

Let V be as above. Let $u : V \rightarrow B\Gamma$ be the classifying map for the universal cover of V and let $\tilde{u} : \tilde{V} \rightarrow E\Gamma$ be a Γ -equivariant lift of u . Let $f : M \rightarrow V$ be a cycle representing an element $[f : M \rightarrow V]$ in $\mathcal{S}^h(V)$. As in Definition 2.17 we consider $Z = M \sqcup (-V)$ with the obvious classifying map $u_Z : Z \rightarrow B\Gamma$ induced by u and by $u \circ f$. We observe that Z comes with a map ϕ to V , $\phi := f \sqcup (-\text{Id}_V)$, which is covered by a Γ -equivariant map $\tilde{\phi} : \tilde{Z} \rightarrow \tilde{V}$.

Definition 4.2. We define a map

$$\rho : \mathcal{G}^h(V) \rightarrow K_{n+1}(D^*(\tilde{V})^\Gamma), \quad [M \xrightarrow{f} V] \mapsto \rho(f), \quad (4.3)$$

where we recall that D is the signature operator on the covering $\tilde{Z} := u_Z^*E\Gamma$, C_f is the smoothing trivializing perturbation constructed using the homotopy equivalence f and $\rho(f) := \tilde{\phi}_*(\rho(D + C_f)) \in K_{n+1}(D^*(\tilde{V})^\Gamma)$.

Definition 4.4. We define

$$\rho_\Gamma : \mathcal{G}^h(V) \rightarrow K_{n+1}(D_\Gamma^*); \quad [M \xrightarrow{f} V] \mapsto \rho_\Gamma(f), \quad (4.5)$$

where we recall that

$$\rho_\Gamma(f) := \rho_\Gamma(D + C_f) := (\tilde{u}_Z)_*(\rho(D + C_f)) = \tilde{u}_*(\rho(f)).$$

If necessary, we shall denote the right-hand sides of (4.3) and (4.5) as $\rho[f : M \rightarrow V]$ and $\rho_\Gamma[f : M \rightarrow V]$, respectively.

From now on we assume our manifold V to be odd-dimensional.

Remark 4.6. Notice that we follow here different conventions with respect to [Higson and Roe 2005c]; indeed we follow the conventions of [PS 2007], slightly modified as in [Wahl 2013b]. In particular, the operator we use on an odd-dimensional manifold is not the odd signature operator as in the work of Atiyah, Patodi and Singer (and the one used by topologists to define the signature K-homology class; compare [Rosenberg and Weinberger 2006]). It is rather the direct sum of two (unitarily equivalent) versions of this operator, and comes up as the boundary operator of the signature operator on an even-dimensional manifold. This results in a possible loss of 2-torsion information. On the other hand, with this modification our diagrams commute without inverting 2.

However, a factor of 2 would, with our conventions, show up in the same portion of the sequence for an even-dimensional manifold V .

It would be interesting to improve the constructions of our invariants, in particular the rho-invariant, so that they deal with the signature operator on odd-dimensional manifolds as in the work of Atiyah, Patodi and Singer (potentially containing additional 2-torsion information). This involves understanding how the Hilsum–Skandalis–Wahl perturbation can be made compatible with the splitting of the operator alluded to above. It also involves establishing bordism-invariance of the rho-invariant to be obtained that way. This is a nontrivial task, as is shown in [Rosenberg and Weinberger 2006], where the corresponding problem of bordism-invariance of the K-homology class is solved.

Finally, we believe that our diagram can be extended to the left, provided that one inverts 2 (and multiplies all maps by the appropriate powers of 2, similar to

[Higson and Roe 2005c]); in such a situation the powers of 2 discussed above will inevitably appear to make all squares commutative.

Proposition 4.7. *The maps*

$$\mathcal{S}^h(V) \rightarrow K_{n+1}(D_\Gamma^*), \quad [M \xrightarrow{f} V] \mapsto \rho_\Gamma(f)$$

and

$$\mathcal{S}^h(V) \rightarrow K_{n+1}(D^*(\tilde{V})^\Gamma), \quad [M \xrightarrow{f} V] \mapsto \rho(f)$$

are well-defined.

Proof. This follows from [Wahl 2013b] and our delocalized APS index theorem. Let us first prove that $\rho(f) \in K_0(D^*(\tilde{V})^\Gamma)$ is well-defined. We assume as before that $[f_1 : M_1 \rightarrow V] = [f_2 : M_2 \rightarrow V]$ in $\mathcal{S}^h(V)$ and let $F : X \rightarrow V \times [0, 1]$ be the h-cobordism realizing this equality; we set again $W = X \sqcup V \times [0, 1]$. Then there is a map $\Phi : W \rightarrow V$ given by

$$F \sqcup \text{id}_{V \times [0,1]} : W \rightarrow V \times [0, 1],$$

followed by the projection π_1 onto the first factor of $V \times [0, 1]$; that is, the map $\Phi = \pi_1 \circ (F \sqcup \text{id}_{V \times [0,1]})$. Let

$$\phi_1 = f_1 \sqcup (-\text{Id}_V) : M_1 \sqcup (-V) \rightarrow V$$

and similarly for ϕ_2 . Set, as before, $Z_1 := M_1 \sqcup (-V)$ and $Z_2 := M_2 \sqcup (-V)$. We have observed that these manifolds come with classifying maps $u \circ \phi_j$ into $B\Gamma$ and we have denoted by \tilde{Z}_j the corresponding Γ -covers. As before, we consider the Γ -equivariant lifts $\tilde{\phi}_j : \tilde{Z}_j \rightarrow \tilde{V}$. Recall now that $\rho(f_1) \in K_0(D^*(\tilde{V})^\Gamma)$ is obtained by pushing forward through $(\tilde{\phi}_1)_*$ the rho-class of \tilde{Z}_1 , denoted $\rho(D\tilde{z}_1 + C_{f_1}) \in K_0(D^*(\tilde{Z}_1)^\Gamma)$. Similarly,

$$\rho(f_2) = (\tilde{\phi}_2)_* \rho(D\tilde{z}_2 + C_{f_2}).$$

Clearly Φ restricted to $M_1 \sqcup (-V) =: Z_1$ is ϕ_1 , and similarly for Φ restricted to $M_2 \sqcup (-V) =: Z_2$. We can now apply Φ_* to the delocalized APS index formula for W , which we write with self-explanatory notation as

$$(\iota_{\tilde{W}})_* \text{Ind}(D_{\tilde{W}}, C_{1,2}) = (j_1)_* \rho(D\tilde{z}_1 + C_{f_1}) - (j_2)_* \rho(D\tilde{z}_2 + C_{f_2}), \quad (4.8)$$

with j_1 and j_2 the obvious inclusions. This equality takes place in $K_0(D^*(\tilde{W})^\Gamma)$. Extending the work of Hilsum and Skandalis to manifolds with cylindrical ends, Wahl [2013b, Theorem 8.4] proved that the existence of the map F entering into the h-cobordism, i.e., of a global homotopy equivalence, implies the vanishing of the index class obtained by grafting the two smoothing perturbations C_{f_j} on the cylindrical ends (Wahl works in the Mishchenko–Fomenko framework but since we know that this is equivalent to the coarse framework we can directly state her

results in the way we just have). Since the latter is precisely the index class that was denoted $\text{Ind}(D_{\tilde{W}}, C_{1,2})$ in the left-hand side of (4.8), we obtain at once that

$$0 = (j_1)_*\rho(D_{\tilde{Z}_1} + C_{f_1}) - (j_2)_*\rho(D_{\tilde{Z}_2} + C_{f_2}) \quad \text{in } K_0(D^*(\tilde{W})^\Gamma).$$

We now apply Φ_* to both sides and we obtain the following equality in $K_0(D^*(\tilde{V})^\Gamma)$:

$$0 = \Phi_*((j_1)_*\rho(Z_1)) - \Phi_*((j_2)_*\rho(Z_2)).$$

However, by functoriality, the right-hand side is exactly $\rho(f_1) - \rho(f_2)$. Thus $0 = \rho(f_1) - \rho(f_2)$, and we are done.

This argument also proves that ρ_Γ is well-defined; indeed, it suffices to apply $\tilde{u}_* : K_0(D^*(\tilde{V})^\Gamma) \rightarrow K_0(D_\Gamma^*)$ to the equality $0 = \rho(f_1) - \rho(f_2)$ and recall that $\tilde{u}_*(\rho(f_j)) = \rho_\Gamma(f_j)$. See the remark after (2.18). \square

4B. The set $\mathcal{N}(V)$ and the map $\beta_\Gamma : \mathcal{N}(V) \rightarrow K_n(B\Gamma)$. Let V be an n -dimensional smooth closed oriented manifold with fundamental group Γ . We assume that $n > 4$ and that it is odd. Let $u : V \rightarrow B\Gamma$ be the classifying map for the universal covering of V and let $\tilde{V} := u^*E\Gamma$.

Definition 4.9. $\mathcal{N}(V)$ is the set of normal maps. Its elements are pairs $[f : M \rightarrow V]$ where M is again a closed smooth oriented manifold and f is a degree-1 normal map. Two such pairs are equivalent if there is a normal cobordism between them (for all this compare for example [Higson and Roe 2005c, §5] and the references therein).

The map β_Γ is the composition of $\beta : \mathcal{N}(V) \rightarrow K_*(V)$ and the group homomorphism $u_* : K_*(V) \rightarrow K_*(B\Gamma)$ induced by the classifying map of V . Here β maps $[f : M \rightarrow V]$ in $\mathcal{N}(V)$ to $f_*[D_M] - [D_V] \in K_*(V)$, where $[D_M], [D_V]$ are the K-homology classes of the signature operators. The map $\beta : \mathcal{N}(V) \rightarrow K_*(V)$ already appears in the work of Higson and Roe, where it is proved to be well-defined.

4C. Mapping the surgery sequence to the Higson–Roe sequence.

Theorem 4.10. *Let V be an n -dimensional smooth closed oriented manifold with fundamental group Γ . We assume that $n > 4$ is odd. Then there is a commutative diagram with exact rows*

$$\begin{array}{ccccccc} L_{n+1}(\mathbb{Z}\Gamma) & \dashrightarrow & \mathcal{S}(V) & \longrightarrow & \mathcal{N}(V) & \longrightarrow & L_n(\mathbb{Z}\Gamma) \\ \downarrow \text{Ind}_\Gamma & & \downarrow \rho & & \downarrow \beta & & \downarrow \text{Ind}_\Gamma \\ K_{n+1}(C_r^*\Gamma) & \longrightarrow & K_{n+1}(D^*(\tilde{V})^\Gamma) & \longrightarrow & K_n(V) & \longrightarrow & K_n(C_r^*\Gamma) \end{array} \quad (4.11)$$

By employing the classifying map $u : V \rightarrow B\Gamma$ for the universal cover \tilde{V} of V we also get a commutative diagram mapping into the universal Higson–Roe surgery

sequence:

$$\begin{array}{ccccccc}
 L_{n+1}(\mathbb{Z}\Gamma) & \dashrightarrow & \mathcal{S}(V) & \longrightarrow & \mathcal{N}(V) & \longrightarrow & L_n(\mathbb{Z}\Gamma) \\
 \downarrow \text{Ind}_\Gamma & & \downarrow \rho_\Gamma & & \downarrow \beta_\Gamma & & \downarrow \text{Ind}_\Gamma \\
 K_{n+1}(C_r^*\Gamma) & \xrightarrow{i_*} & K_{n+1}(D_\Gamma^*) & \longrightarrow & K_n(B\Gamma) & \longrightarrow & K_n(C_r^*\Gamma)
 \end{array} \quad (4.12)$$

Proof. Thanks to the work of Wahl (see Theorem 2.1) and Proposition 2.33 we know that the index homomorphism $\text{Ind} : L_*(\mathbb{Z}\Gamma) \rightarrow K_*(C_r^*\Gamma)$ is well-defined.³ By Proposition 4.7, ρ is well-defined, and we know from [Higson and Roe 2005c, Definition 5.2] that β is also well-defined. We now proceed to prove the commutativity of the squares.

Recall first of all the meaning of exactness of the surgery sequence at $\mathcal{S}(V)$; see for example [Higson and Roe 2005c]. Let $[f : M \rightarrow V] \in \mathcal{S}(V)$ and let $a \in L_{n+1}(\mathbb{Z}\Gamma)$. Then a can be represented by $(W, F, M \times [0, 1], u_M : M \rightarrow B\Gamma)$ with $F : W \rightarrow M \times [0, 1]$ a normal map of degree one between manifolds with boundary, and $u_M = u \circ f$. Here W is a manifold with two boundary components $\partial_0 W$ and $\partial_1 W$,

$$\partial_0 F := F|_{\partial_0 W} : \partial_0 W \rightarrow \{0\} \times M$$

is a diffeomorphism, and

$$\partial_1 F := F|_{\partial_1 W} : \partial_1 W \rightarrow \{1\} \times M$$

is a homotopy equivalence. For later use we set $X := W \sqcup M \times [0, 1]$ and we remark that X comes with a natural map $u_f : X \rightarrow V$ with $u_f := f \circ (\pi_1(F \circ \text{id}_{M \times [0, 1]}))$ and π_1 the projection onto the first factor. The action of $L_{n+1}(\mathbb{Z}\Gamma)$ on $\mathcal{S}(V)$ is as follows: if $a \in L_{n+1}(\mathbb{Z}\Gamma)$ and $[f : M \rightarrow V] \in \mathcal{S}(V)$ then

$$a \cdot [M \xrightarrow{f} V] := [\partial_1 W \xrightarrow{f \circ \partial_1 F} V].$$

Exactness of the surgery sequence at $\mathcal{S}(V)$ means the following: two elements in the structure set belong to the same orbit under the action of the group $L_{n+1}(\mathbb{Z}\Gamma)$ if and only if their images in $\mathcal{N}(V)$ coincide. By definition, in this situation where we deal with group actions instead of homomorphisms, commutativity of the first square means

$$\rho(a \cdot [M \xrightarrow{f} V]) - \rho[M \xrightarrow{f} V] = i_* \text{Ind}(a) \quad \text{in } K_{n+1}(D^*(\tilde{V})^\Gamma),$$

i.e., that

$$\rho[\partial_1 W \xrightarrow{f \circ \partial_1 F} V] - \rho[M \xrightarrow{f} V] = i_* \text{Ind}(a).$$

³Wahl concentrates on the case when $*$ is even, but for this particular result her arguments apply, with a minimum amount of work, to the case when $*$ is odd.

The homomorphism i_* appearing on the right-hand side is, up to a canonical isomorphism, the map induced by the inclusion $C^*(\tilde{V})^\Gamma \rightarrow D^*(\tilde{V})^\Gamma$, whereas the index class $\text{Ind}(a)$ can be taken to be the pushforward, from $K_{n+1}(C^*(\tilde{X})^\Gamma)$ to $K_{n+1}(C^*(\tilde{V})^\Gamma)$, of the index class defined by the perturbed signature operator on \tilde{X} :

$$\text{Ind}(a) = (\tilde{u}_f)_*(\text{Ind}(D_{\tilde{X}} + C_{\partial_0 F, \partial_1 F})), \quad (4.13)$$

with \tilde{u}_f the Γ -map covering $u_f : X \rightarrow V$. Now, the proof of [Wahl 2013b, Proposition 7.1], together with Proposition 2.33, shows that we have the following identity in $K_{*+1}(D^*(\tilde{V})^\Gamma)$:

$$\rho[L \xrightarrow{f \circ g} V] + \tilde{f}_*(\rho[M \xrightarrow{\text{Id}_M} M]) = \tilde{f}_*(\rho[L \xrightarrow{g} M]) + \rho[M \xrightarrow{f} V]. \quad (4.14)$$

In order to prove (4.14) we consider the manifold $Y := L \sqcup (-M) \sqcup M \sqcup (-V)$ and the natural inclusions

$$\begin{aligned} Z_1 &:= L \sqcup (-M) \xrightarrow{j_1} Y, & Z_2 &:= M \sqcup (-V) \xrightarrow{j_2} Y, \\ T_1 &:= L \sqcup (-M) \xrightarrow{i_1} Y, & T_2 &:= M \sqcup (-V) \xrightarrow{i_2} Y. \end{aligned}$$

One gets the invertible operators $D_{\tilde{Z}_1} + C_g$ and $D_{\tilde{Z}_2} + C_f$ and the associated rho-classes

$$\rho(D_{\tilde{Z}_1} + C_g) \in K_{n+1}(D^*(\tilde{Z}_1)^\Gamma); \quad \rho(D_{\tilde{Z}_2} + C_f) \in K_{n+1}(D^*(\tilde{Z}_2)^\Gamma).$$

Similarly, we obtain the invertible operators $D_{\tilde{T}_1} + C_{f \circ g}$ and $D_{\tilde{T}_2} + C_{\text{Id}}$ and the rho-classes

$$\rho(D_{\tilde{T}_1} + C_{f \circ g}) \in K_{n+1}(D^*(\tilde{T}_1)^\Gamma); \quad \rho(D_{\tilde{T}_2} + C_{\text{Id}}) \in K_{n+1}(D^*(\tilde{T}_2)^\Gamma).$$

We claim that

$$\begin{aligned} (\tilde{j}_1)_*(\rho(D_{\tilde{Z}_1} + C_g)) + (\tilde{j}_2)_*(\rho(D_{\tilde{Z}_2} + C_f)) \\ = (\tilde{i}_1)_*(\rho(D_{\tilde{T}_1} + C_{f \circ g})) + (\tilde{i}_2)_*(\rho(D_{\tilde{T}_2} + C_{\text{Id}})) \end{aligned} \quad (4.15)$$

in the group $K_{n+1}(D^*(\tilde{Y})^\Gamma)$. Indeed, by a simple functoriality argument, both sides are defined by rho-classes of suitable perturbations of the signature operator on \tilde{Y} ; Wahl shows [2013b, proof of Proposition 7.1] that there is a continuous family of trivializing perturbations, $[0, 1] \ni t \mapsto G_t$, interpolating from the perturbation defining the left-hand side to the perturbation defining the right-hand side. Thanks to Proposition 2.33 we then obtain immediately (4.15). We now consider the map $\Psi : Y \rightarrow V$ defined by

$$\Psi|_L := f \circ g, \quad \Psi|_M := f, \quad \Psi|_{-V} := \text{Id}_{-V},$$

and we observe that

$$\Psi \circ i_1 = (f \circ g) \sqcup \text{Id}_{(-V)}, \quad \Psi \circ i_2 = f \sqcup (-f), \quad \Psi \circ j_1 = f \circ (g \sqcup \text{Id}_{(-M)}).$$

By applying $\tilde{\Psi}_*$ to both sides of (4.15) we then get, by functoriality, that

$$\tilde{f}_*(\rho(g)) + \rho(f) = \rho(f \circ g) + \tilde{f}_*(\rho(\text{Id}_M)) \quad \text{in } K_{n+1}(D^*(\tilde{V})^\Gamma),$$

which is precisely (4.14). Granted formula (4.14) we obtain, as in [Wahl 2013b, Theorem 9.1],

$$\rho(a \cdot [M \xrightarrow{f} V]) - \rho[M \xrightarrow{f} V] = \tilde{f}_*(\rho[\partial_1 W \xrightarrow{\partial_1 F} M]) - \tilde{f}_*(\rho[\partial_0 W \xrightarrow{\partial_0 F} M]). \quad (4.16)$$

By applying the delocalized APS index Theorem 3.1 for perturbed operators and functoriality we recognize in the difference on the right-hand side the element $i_* \text{Ind}(a)$, with $\text{Ind}(a)$ as in (4.13). The commutativity of the first square is therefore established.

Next we tackle the commutativity of the second square, where we recall that the horizontal map $\mathcal{S}(V) \rightarrow \mathcal{N}(V)$ is the forgetful map. Let $[f : M \rightarrow V] \in \mathcal{S}(V)$ and consider $Z = M \sqcup (-V)$ with the usual maps $\phi : Z \rightarrow V$, $\phi := f \sqcup \text{Id}_{(-V)}$, and $u_Z : Z \rightarrow B\Gamma$, $u_Z := u \circ \phi$. We let C_f be the trivializing perturbation defined by f , so that $D_{\tilde{Z}} + C_f$ is invertible, with spectrum disjoint from the interval $[-2\epsilon, 2\epsilon]$. Recall that for any closed compact manifold Z of odd dimension, endowed with a classifying map $u_Z : Z \rightarrow B\Gamma$,

$$K_1(Z) = K_0(D^*Z/C^*Z) = K_0(D^*(\tilde{Z})^\Gamma/C^*(\tilde{Z})^\Gamma),$$

with $\tilde{Z} = u_Z^* E\Gamma$; in this realization of the K-homology groups, the K-homology signature class $[D_Z] \in K_1(Z)$ is obtained by considering the projection $[\frac{1}{2}(\chi(D_{\tilde{Z}}) + 1)]$ in $K_0(D^*(\tilde{Z})^\Gamma/C^*(\tilde{Z})^\Gamma)$, with χ a chopping function. The continuous path of operators

$$[0, 1] \rightarrow K_0(D^*(\tilde{Z})^\Gamma/C^*(\tilde{Z})^\Gamma), \quad t \mapsto [\frac{1}{2}(\chi(D_{\tilde{Z}} + tC_f) + 1)],$$

shows that for a chopping function χ equal to 1 on $[\epsilon, +\infty)$ and -1 on $(-\infty, \epsilon]$ we have the equality

$$\left[\frac{1}{2} \left(\frac{D_{\tilde{Z}} + C_f}{|D_{\tilde{Z}} + C_f|} + 1 \right) \right] = [\frac{1}{2}(\chi(D_{\tilde{Z}}) + 1)] \quad \text{in } K_0(D^*(\tilde{Z})^\Gamma/C^*(\tilde{Z})^\Gamma). \quad (4.17)$$

With $\phi = f \sqcup \text{Id}_{(-V)}$ we now apply $(\tilde{\phi})_*$ to both sides of this equality. On the left-hand side we obtain the class

$$\rho[M \xrightarrow{f} V] \in K_0(D^*(\tilde{V})^\Gamma/C^*(\tilde{V})^\Gamma) = K_1(V),$$

whereas on the right we obtain $\phi_*[D_Z] \in K_1(V)$, which is precisely $f_*[D_M] - [D_V]$. The commutativity of the second square is established.

Finally, we tackle the third square. The image of an element $[f : M \rightarrow V] \in \mathcal{N}(V)$ in $L_n(\mathbb{Z}\Gamma)$ is simply the class associated to the cycle represented by the closed odd-dimensional manifold $Z = M \cup (-V)$ endowed with the classifying map u_Z . This element in $L_n(\mathbb{Z}\Gamma)$ is the *surgery obstruction* associated to $[f : M \rightarrow V] \in \mathcal{N}(V)$, and one of the fundamental results in surgery theory states that a normal map $f : M \rightarrow V$ can be surgered to a homotopy equivalence if and only if its surgery obstruction in $L_n(\mathbb{Z}\Gamma)$ vanishes. This is the meaning of exactness of the surgery sequence at $\mathcal{N}(V)$. Going back to the commutativity of the third square, we observe that the map Ind sends this class, the surgery obstruction associated to $[f : M \rightarrow V]$, to the associated index class in $K_n(C_r^*\Gamma)$. By [Kasparov 1995, Theorem 2], this class is equal to the Mishchenko symmetric signature of the covering $u_Z^*E\Gamma \rightarrow Z \equiv M \cup (-V)$. This remark reduces the proof of the commutativity of the third square to the proof given in [Higson and Roe 2005c, Theorem 5.4]. \square

Remark 4.18. In [Xie and Yu 2014], a slightly different approach to the delocalized APS index theorem for spin Dirac operators is worked out. It uses the localization algebras invented in [Yu 1997], which allow one to formulate and prove product formulas for products between fundamental classes (primary invariants) and rho-classes (secondary invariants). Presumably, this approach can be generalized to the perturbed operators we have to use here; however, it is clear that one will face technical difficulties related to the nonlocality of the perturbed operators and the very definition of localization algebras (based on locality of the operators involved).

Remark 4.19. So far we have treated only the case in which the manifold V appearing in the structure set $\mathcal{G}^h(V)$ is odd-dimensional. Similarly, in [PS 2014] we have only treated the group $\text{Pos}_n^{\text{spin}}(X)$ for n odd. The problem, both here and in [PS 2014], is centered around the delocalized APS index theorem, a crucial tool which we prove only for *even-dimensional* manifolds with boundary.

However, there is a clear strategy to deduce the remaining case from the one we have established, which is also suggested in [Xie and Yu 2014]. One should develop a “suspension homomorphism”, mapping

$$K_*(C^*(W)^\Gamma) \quad \text{to} \quad K_{*+1}(C^*(W \times \mathbb{R})^{\Gamma \times \mathbb{Z}})$$

and

$$K_*(D^*(W)^\Gamma) \quad \text{to} \quad K_{*+1}(D^*(W \times \mathbb{R})^{\Gamma \times \mathbb{Z}})$$

in functorial and compatible ways (for manifolds W with and without boundary).

These homomorphisms have to map the rho-classes (in the case of a manifold without boundary) and index classes (in general) of the Dirac operators on W to

those on $W \times \mathbb{R}$, and they have to be injective. It is then an easy exercise to deduce the delocalized APS index theorem for one parity from the one for the other parity.

This program has now been implemented by Vito Felice Zenobi [2015], building on previous work of Paul Siegel [2012a; 2012b]. Consequently, the delocalized APS index theorem presented in this article is now established in every dimension. Similarly, Zenobi shows that the delocalized APS index theorem proved in [PS 2014] is valid in all dimensions. The latter is also a consequence of the treatment given in [Xie and Yu 2014].

Notice that a general treatment of the suspension homomorphism and the relevant compatibilities and product formulas, using Yu’s localization algebras, is carried out by Rudolf Zeidler [2015], making more explicit some of the constructions in [Xie and Yu 2014]. The methods in [Zeidler 2015] also cover the case of the signature operator and its rho-invariants and are therefore another route to derive the delocalized APS index theorem for odd-dimensional manifolds from the one for even-dimensional manifolds.

5. Proof of the delocalized APS index theorem

In this section we prove Theorem 3.1, the delocalized APS index theorem for perturbed operators. Recall from [PS 2014, §4] that the corresponding result for the Dirac operator of a spin manifold with positive scalar curvature on the boundary is proved in two steps: first we show how to reduce the validity of the theorem on \tilde{W}_∞ to a version of the theorem on $\mathbb{R} \times \partial\tilde{W}$ (which we call the “cylinder delocalized index theorem for perturbed operators”); next a detailed analysis, with explicit computations, is carried out on the cylinder in order to establish the cylinder delocalized index theorem. We will follow this strategy also here.

In this subsection we make more precise the notation given in the proof of Proposition 2.26; thus we denote by p the characteristic function of $[0, \infty) \times \partial\tilde{W}$ inside W_∞ and we denote by P the corresponding multiplication operator. Finally, as in the proof of Proposition 2.26, we denote by p_0 the characteristic function of $[0, \infty) \times \partial\tilde{W}$ inside the full cylinder $\mathbb{R} \times \partial\tilde{W}$ and by P_0 the corresponding multiplication operator.

5A. Reduction to the cylinder. Once we have at our disposal Proposition 2.26, together with its proof, it is elementary to check that the reduction to the cylinder proceeds exactly as in [PS 2014]. Thus:

- (1) We establish that $P_0\chi(\bar{D} + \bar{C})_+P_0$ is an involution in

$$D^*([0, \infty) \times \partial\tilde{W})^\Gamma / D^*({0} \times \partial\tilde{W}) \subset [0, \infty) \times \partial\tilde{W})^\Gamma,$$

where we recall that on the cylinder the bundle on which the signature operator acts can be identified with the direct sum of two copies of the same bundle, with

the obvious grading. We obtain

$$\partial[P_0\chi(\bar{D} + \bar{C})_+ P_0] \in K_0(D^*(\partial\tilde{W} \subset [0, \infty) \times \partial\tilde{W})^\Gamma).$$

(2) Similarly, we have a class $\partial[P\chi(\bar{D} + \bar{C})P] \in K_0(D^*(\tilde{W} \subset \tilde{W}_\infty)^\Gamma)$, where it is obvious how to consider $P\chi(\bar{D} + \bar{C})P$ as an element of $D^*(\tilde{W}_\infty)^\Gamma$.

(3) Next we establish that

$$\chi(D + C) - P\chi(\bar{D} + \bar{C})P \in D^*(\tilde{W} \subset \tilde{W}_\infty)^\Gamma. \quad (5.1)$$

We shall prove (5.1) at the end of this subsection, just below.

(4) Using the latter information, we show that if

$$\iota_* : K_*(C^*(\tilde{W} \subset \tilde{W}_\infty)^\Gamma) \rightarrow K_*(D^*(\tilde{W} \subset \tilde{W}_\infty)^\Gamma)$$

is induced by the inclusion, then

$$\iota_*(\text{Ind}^{\text{rel}}(D + C)) = \partial[P\chi(\bar{D} + \bar{C})_+ P] \in K_0(D^*(\tilde{W} \subset \tilde{W}_\infty)^\Gamma).$$

(5) We then show that if j_+ is the homomorphism

$$K_*(D^*(\partial\tilde{W} \subset [0, \infty) \times \partial\tilde{W})^\Gamma) \rightarrow K_0(D^*(\tilde{W} \subset \tilde{W}_\infty)^\Gamma),$$

then

$$j_+(\partial[P_0\chi(\bar{D} + \bar{C})_+ P_0]) = \partial[P\chi(\bar{D} + \bar{C})_+ P] \in K_0(D^*(\tilde{W} \subset \tilde{W}_\infty)^\Gamma).$$

(6) Next, we assume the following result:

Theorem 5.2 (cylinder delocalized index theorem for perturbed operators). *If j_∂ is the isomorphism $K_*(D^*(\partial\tilde{W})^\Gamma) \rightarrow K_*(D^*(\partial\tilde{W} \subset [0, \infty) \times \partial\tilde{W})^\Gamma)$, then*

$$\partial[P_0\chi(\bar{D} + \bar{C})_+ P_0] = j_\partial \rho(D_\partial + C_\partial). \quad (5.3)$$

(7) Finally, we recall the following commutative diagram from [PS 2014, Proposition 3.1]:

$$\begin{array}{ccc} K_*(D^*(\partial\tilde{W})^\Gamma) & \xrightarrow{j_*} & K_*(D^*(\tilde{W})^\Gamma) \\ \cong \downarrow j_\partial & & \cong \downarrow c \\ K_*(D^*(\partial\tilde{W} \subset [0, \infty) \times \partial\tilde{W})^\Gamma) & \xrightarrow{j_+} & K_*(D^*(\tilde{W} \subset \tilde{W}_\infty)^\Gamma) \end{array} \quad (5.4)$$

Granted these properties, one gets

$$\begin{aligned} \iota_*(\text{Ind}^{\text{rel}}(D + C)) &= \partial[P\chi(\bar{D} + \bar{C})_+ P] \\ &= j_+(\partial[P_0\chi(\bar{D} + \bar{C})_+ P_0]) \\ &= j_+(j_\partial \rho(D_\partial + C)). \end{aligned}$$

Once we apply c^{-1} to both sides we obtain precisely the equality in the theorem, i.e., $\iota_*(\text{Ind}(D, C)) = j_*(\rho(D_\partial + C_\partial))$.

Proof of (5.1). We go back to the notation adopted in the proof of Proposition 2.26; thus, with a small abuse of notation, we don't distinguish between P_0 and P . Recall that we identify the image of P_0 in $L^2(\mathbb{R} \times \tilde{W})$ with the corresponding subspace of $L^2(\tilde{W}_\infty)$ (in order to lighten the notation we do not write the vector bundles in the L^2 -spaces). This way, we consider $P_0\chi(\bar{D} + \bar{C})P_0$ as an operator on $L^2(\tilde{W}_\infty)$.

First of all we remark that thanks to Proposition 2.26 we only need to establish (5.1) for one specific chopping function; we therefore choose $\chi(x) = x/\sqrt{1+x^2}$. We know (see [PS 2014, Lemma 4.12]) that $\chi(D) - P_0\chi(\bar{D})P_0 \in D^*(\tilde{W} \subset \tilde{W}_\infty)^\Gamma$. We also know from [Higson and Roe 2005a, Lemma 5.8] that

$$\chi(\bar{D}) - \chi(\bar{D} + \bar{C}) \in C^*(\mathbb{R} \times \partial\tilde{W})^\Gamma; \quad \chi(D) - \chi(D + C) \in C^*(\tilde{W}_\infty)^\Gamma.$$

Then we write

$$\begin{aligned} \chi(D + C) - P_0\chi(\bar{D} + \bar{C})P_0 &= (\chi(D + C) - \chi(D)) \\ &\quad + (\chi(D) - P_0\chi(\bar{D})P_0) + (P_0\chi(\bar{D})P_0 - P_0\chi(\bar{D} + \bar{C})P_0). \end{aligned}$$

As already remarked, the second summand on the right-hand side is an element in $D^*(\tilde{W} \subset \tilde{W}_\infty)^\Gamma$. We will prove that the sum of the first and third term on the right-hand side is in $C^*(\tilde{W} \subset \tilde{W}_\infty)^\Gamma$. To this end we write an explicit expression for this sum, using that $\chi(x) = x/\sqrt{1+x^2}$. Indeed, using once again [ibid., Lemma 5.8], we can write this sum as the operator norm convergent integral

$$\begin{aligned} &\frac{1}{\pi} \int_1^\infty \frac{t}{\sqrt{t^2-1}} ((D+it)^{-1}C(D+C+it)^{-1} - P_0(\bar{D}+it)^{-1}\bar{C}(\bar{D}+\bar{C}+it)^{-1}P_0) dt \\ &+ \frac{1}{\pi} \int_1^\infty \frac{t}{\sqrt{t^2-1}} ((D-it)^{-1}C(D+C-it)^{-1} - P_0(\bar{D}-it)^{-1}\bar{C}(\bar{D}+\bar{C}-it)^{-1}P_0) dt. \end{aligned}$$

Therefore it suffices to show that the integrands are in $C^*(\tilde{W} \subset \tilde{W}_\infty)^\Gamma$. Let us consider the first integral and the t -dependent operator there:

$$(D+it)^{-1}C(D+C+it)^{-1} - P_0(\bar{D}+it)^{-1}\bar{C}(\bar{D}+\bar{C}+it)^{-1}P_0. \quad (5.5)$$

Writing $1 = P_0 + (1 - P_0)$ and reasoning as we did at the end of the proof of Proposition 2.26 (after the proof of Lemma 2.28), we see that (5.5) is equal, up to a term in $C^*(\tilde{W} \subset \tilde{W}_\infty)^\Gamma$, to

$$\begin{aligned} &P_0(D+it)^{-1}P_0CP_0(D+C+it)^{-1}P_0 - P_0(\bar{D}+it)^{-1}P_0\bar{C}P_0(\bar{D}+\bar{C}+it)^{-1}P_0 \\ &= P_0(D+it)^{-1}P_0CP_0(P_0(D+C+it)^{-1}P_0 - P_0(\bar{D}+\bar{C}+it)^{-1}P_0) \\ &\quad + (P_0(D+it)^{-1}P_0 - P_0(\bar{D}+it)^{-1}P_0)P_0CP_0(\bar{D}+\bar{C}+it)^{-1}P_0, \end{aligned}$$

since, by definition, $P_0\bar{C}P_0 = P_0CP_0$. We already know that

$$(P_0(D + C + it)^{-1}P_0 - P_0(\bar{D} + \bar{C} + it)^{-1}P_0)$$

and

$$(P_0(D + it)^{-1}P_0 - P_0(\bar{D} + it)^{-1}P_0)$$

are elements in $C^*(\tilde{W} \subset \tilde{W}_\infty)^\Gamma$. Since the other factors are all norm limits of finite-propagation operators, thus multipliers of $C^*(\tilde{W} \subset \tilde{W}_\infty)^\Gamma$, we see that (5.5) is in $C^*(\tilde{W} \subset \tilde{W}_\infty)^\Gamma$. The proof of (5.1) is complete. \square

5B. Proof of the cylinder delocalized index theorem for perturbed operators. We now prove Theorem 5.2, namely that if j_∂ is the isomorphism

$$K_*(D^*(\partial\tilde{W})^\Gamma) \rightarrow K_*(D^*(\partial\tilde{W} \subset \mathbb{R}_\geq \times \partial\tilde{W})^\Gamma)$$

induced by the inclusion, then

$$\partial[P_0\chi(\bar{D} + \bar{C})_+P_0] = j_\partial\rho(D_\partial + C_\partial) \quad \text{in } K_*(D^*(\partial\tilde{W} \subset \mathbb{R}_\geq \times \partial\tilde{W})^\Gamma),$$

where we abbreviate $\mathbb{R}_\geq := [0, \infty)$.

We wish to point out that our arguments, although somewhat lengthy, are elementary. It is certainly possible to envisage a proof based on the Volterra expansion for the wave operator of a perturbed Dirac operator. However, doing this properly does require some nontrivial work and, in addition, it would not generalize easily to Lipschitz manifolds. This is why we have followed the route presented below.

Notation 5.6. We set $\tilde{M} := \partial\tilde{W}$. We denote the Dirac operator on \tilde{M} , the cross-section of the cylinder, by D , and the Dirac operator on $\mathbb{R} \times \tilde{M}$ by \bar{D} . We denote a trivializing perturbation for D by C and we denote the resulting perturbation on $\mathbb{R} \times \tilde{M}$, the one obtained by extending C to be constant in the \mathbb{R} -direction, by \bar{C} . We use the symbol $\mathcal{L}^2(\tilde{M})$ for the L^2 -sections of the relevant Clifford bundle. Similarly we employ $\mathcal{L}^2(\mathbb{R} \times \tilde{M})$ and $\mathcal{L}^2(\mathbb{R}_\geq \times \tilde{M})$.

Proposition 5.7. *The bounded linear operator*

$$V : \mathcal{L}^2(\tilde{M}) \rightarrow \mathcal{L}^2([0, \infty) \times \tilde{M}), \quad s \mapsto Vs, \\ (Vs)(t) := \sqrt{2|D + C|}e^{-t|D + C|}(s), \quad (5.8)$$

covers in the D^ -sense the inclusion $i : M \hookrightarrow \mathbb{R}_\geq \times M, m \mapsto (0, m)$.*

Proof. We prove this in Section 5B1. \square

Proposition 5.9. *The operator $\frac{|D + C| + \partial_t}{(D + C) - \partial_t}$ belongs to $D^*(\mathbb{R} \times \tilde{M})^\Gamma$.*

Proof. We prove this in Section 5B2. \square

Granted these two propositions, we can proceed as in [PS 2014, §4.4]. We have the following proposition.

Proposition 5.10. *Set*

$$Q := P_0 \frac{|D + C| - \partial_t}{(D + C) + \partial_t} P_0 : \mathcal{L}^2(\mathbb{R}_{\geq} \times \tilde{M}) \rightarrow \mathcal{L}^2(\mathbb{R}_{\geq} \times \tilde{M}).$$

Then

$$\begin{aligned} \left(P_0 \frac{|D + C| + \partial_t}{(D + C) - \partial_t} P_0 \right) \circ Q &= \text{Id}_{\mathcal{L}^2(\mathbb{R}_{\geq} \times \tilde{M})}; \\ Q \circ \left(P_0 \frac{|D + C| + \partial_t}{(D + C) - \partial_t} P_0 \right) &= \text{Id}_{\mathcal{L}^2(\mathbb{R}_{\geq} \times \tilde{M})} - \Pi, \end{aligned} \quad (5.11)$$

with $\Pi := V \chi_{[0, \infty)}(D + C)V^*$.

This implies that

$$\begin{aligned} \partial \left[P_0 \frac{|D + C| + \partial_t}{(D + C) - \partial_t} P_0 \right] &= [V \chi_{[0, \infty)}(D + C)V^*] \\ &\text{in } K_0(D^*(\tilde{M} \subset \mathbb{R}_{\geq} \times \tilde{M})^\Gamma). \end{aligned} \quad (5.12)$$

The proof of Proposition 5.10 proceeds as in [PS 2014, §4.4], given that the arguments there are purely functional-analytic and rest ultimately on the Browder–Gårding decomposition for the self-adjoint operator $D + C$. Now, on the right-hand side of (5.12) we have, by definition, $j_\partial \rho(D_\partial + C_\partial)$, whereas the left-hand side is equal to $\partial[P_0 \chi(\bar{D} + \bar{C})_+ P_0]$. This assertion, namely that

$$\partial \left[P_0 \frac{|D + C| + \partial_t}{(D + C) - \partial_t} P_0 \right] = \partial[P_0 \chi(\bar{D} + \bar{C})_+ P_0], \quad (5.13)$$

is proved as in [PS 2014], using exactly the deformation argument explained there, after the proof of Proposition 4.33. Thus, assuming Propositions 5.7 and 5.9, we have proved the cylinder delocalized index Theorem 5.2.

5B1. *Proof of Proposition 5.7: the map V covers the inclusion in the D^* -sense.* We recall first of all that, by definition, $V : \mathcal{L}^2(\tilde{M}) \rightarrow \mathcal{L}^2([0, \infty) \times \tilde{M})$ covers in the D^* -sense the inclusion $i : \tilde{M} \hookrightarrow \mathbb{R}_{\geq} \times \tilde{M}$ if V is the norm limit of bounded Γ -equivariant maps U satisfying the following two conditions:

- There is an $R > 0$ such that $\phi U \psi = 0$ if $d(\text{Supp}(\phi), i(\text{Supp}(\psi))) > R$, for $\phi \in C_c(\mathbb{R}_{\geq} \times \tilde{M})$ and $\psi \in C_c(\tilde{M})$.
- $\phi U \psi$ is compact for each $\phi \in C_0(\mathbb{R}_{\geq} \times \tilde{M})$ and $\psi \in C_c(\tilde{M})$ such that $\text{Supp}(\phi) \cap i(\text{Supp}(\psi)) = \emptyset$.

Notation 5.14. In all this subsection we shall denote by M the total space of a Riemannian Γ -Galois covering with compact base M/Γ . As before, we write D for a Dirac-type operator on M , acting on sections of a Clifford module bundle. We will in the notation ignore this bundle.

Let $E : L^2(M) \rightarrow L^2(M)$ be a bounded equivariant self-adjoint operator that is a norm limit of equivariant finite-propagation operators and with the property that $D + E$ is invertible.

Recall that we set

$$\begin{aligned} V : L^2(M) &\rightarrow L^2([0, \infty) \times M) = L^2([0, \infty), L^2(M)); \\ (Vu)(t) &= \sqrt{2|D + E|} e^{-t|D + E|} u. \end{aligned}$$

Lemma 5.15. *Let H be a Hilbert space and $W : H \rightarrow L^2([0, \infty), H)$ be an operator of the form $W(u)(t) = f_t(A)u$ for a self-adjoint operator A on H and a measurable function $[0, \infty) \rightarrow C_b(\mathbb{R}); t \mapsto f_t$.*

Then $\|W\|^2 \leq \sup_{\lambda \in \text{Spec}(A)} \int_0^\infty |f_t(\lambda)|^2 dt$.

Proof. By definition, for $u \in H$ we have $|Wu|^2 = \int_0^\infty |f_t(A)u|_H^2 dt$. Use the Browder–Gårding decomposition as direct integral $H = \int H_\lambda d\mu(\lambda)$ according to the spectral decomposition of the self-adjoint operator A . Then $u = \int_{\text{Spec}(A)} u_\lambda d\mu(\lambda)$ with $u_\lambda \in H_\lambda$ and $|u|_H^2 = \int |u_\lambda|_{H_\lambda}^2 d\mu(\lambda)$. Moreover, for a function $f(\lambda)$ we have $f(A)u = \int_{\text{Spec}(A)} f(\lambda)u_\lambda d\mu(\lambda)$. Then

$$\begin{aligned} |Wu|^2 &= \int_0^\infty |f_t(A)u|^2 dt = \int_0^\infty \int_{\text{Spec}(A)} |f_t(\lambda)u_\lambda|_{H_\lambda}^2 d\mu(\lambda) dt \\ &\stackrel{\text{Fubini}}{=} \int_{\text{Spec}(A)} \int_0^\infty |f_t(\lambda)|^2 dt |u_\lambda|_{H_\lambda}^2 d\mu(\lambda) \\ &\leq \left(\sup_{\lambda \in \text{Spec}(A)} \int_0^\infty |f_t(\lambda)|^2 dt \right) \cdot \underbrace{\int_{\text{Spec}(A)} |u_\lambda|_{H_\lambda}^2 d\mu(\lambda)}_{=|u|_H^2}. \quad \square \end{aligned}$$

Lemma 5.16. *The operator $P_R V : L^2(M) \rightarrow L^2([0, \infty) \times M)$ converges to 0 in norm for $R \rightarrow \infty$, where P_R is the orthogonal projection onto $L^2([R, \infty) \times M)$.*

Proof. By assumption, $D + E$ is invertible, i.e., there is $\epsilon > 0$ such that $(-\epsilon, \epsilon) \cap \text{Spec}(D + E) = \emptyset$. We can write $P_R \circ V(u)(t) = f_t^R(D + E)u$ with

$$f_t^R(\lambda) = \begin{cases} 0; & t > R, \\ \sqrt{2|\lambda|} \exp(-t|\lambda|); & t \leq R. \end{cases}$$

For $|\lambda| \geq \epsilon$, the L^2 -norm of $t \mapsto f_t^R(\lambda)$ uniformly tends to 0 as $R \rightarrow \infty$. By Lemma 5.15, $\|P_R V\| \rightarrow 0$ as $R \rightarrow \infty$. \square

We now treat $(1 - P_R)V$. Define

$$V_D : L^2(M) \rightarrow L^2([0, \infty) \times M), \quad V_D u(t) = \sqrt{2|D|} e^{-t|D|} u,$$

i.e., exactly as V , but with $D + E$ replaced by D . We want to prove the properties for V by a comparison with V_D , where corresponding properties have been established in [PS 2014]. For the comparison we use a resolvent trick, which requires some preparation.

Lemma 5.17. *The function $\sqrt{|\lambda|}e^{-|\lambda|}$ can on \mathbb{R} be approximated in supremum norm by a function $f(\lambda)$ which is a polynomial in $1/(\lambda^2 + 1)$ (without constant term, i.e., vanishing at infinity) in such a way that $e^{-|\lambda|}$ is approximated in $L^2(\mathbb{R})$ by $f(\lambda)/\sqrt{|\lambda|}$.*

Proof. Consider the subalgebra of $C_0(\mathbb{R})$ generated by $w(t) := 1/(1 + t^2)$ and $v(t) := t/(1 + t^2)$, which separates points of \mathbb{R} . Notice the relation

$$v^2 = w - w^2. \quad (5.18)$$

Observe that the image of this subalgebra under the operator of differentiation contains for $n \geq 1$ the function $w^n v$; indeed, $w' = -2wv$, so that $(w^n)' = -2nw^n v$ for all $n \geq 1$.

Consider now the subalgebra of $C_0(\mathbb{R})$ generated by w and wv . This subalgebra also separates points; thus, by Stone–Weierstrass, every function in $C_0(\mathbb{R})$ can be approximated by a polynomial $P(w, wv)$ with $P(0, 0) = 0$ (elements in $C_0(\mathbb{R})$ vanish at infinity). Because of (5.18) we see that any continuous even function in $C_0(\mathbb{R})$ can be approximated by a polynomial in w without constant term, whereas any continuous odd function in $C_0(\mathbb{R})$ can be approximated by a polynomial of the form $P(w)v$, with $P(0) = 0$. Notice, finally, that because of the remark following (5.18) the antiderivative of a function of the form $P(w)v$, $P(0) = 0$ belongs to the algebra generated by w and v . Given $\epsilon > 0$, we apply all this to $h'(t)$, with

$$h(t) := u(t)e^{-|t|}(1 + t^2) \quad \text{and} \quad u(t) := \begin{cases} \sqrt{|t|}; & |t| \geq \epsilon, \\ \frac{7}{4}\epsilon^{-3/2}t^2 - \frac{3}{4}\epsilon^{-7/2}t^4; & |t| \leq \epsilon. \end{cases}$$

Note that $|u(t)e^{-|t|} - \sqrt{|t|}e^{-|t|}| \leq 4\sqrt{\epsilon}$.

As h is even and continuously differentiable, h' is odd and continuous. Since $e^{-|t|}$ decays exponentially, $h'(t)$ vanishes at ∞ . Therefore, we can find an odd function $Q_\epsilon(t) = R(w(t))v(t)$, with R a polynomial without constant term, such that $|h'(t) - Q_\epsilon(t)| < \epsilon$ for all $t \in \mathbb{R}$.

Then the even function $P_\epsilon(t) := \int_0^t Q_\epsilon(\tau) d\tau$ is a polynomial in $w(t) = 1/(1 + t^2)$, without constant term, such that $|h(t) - P_\epsilon(t)| \leq \epsilon|t|$ for all $t \in \mathbb{R}$, and therefore we have

$$\left| u(t)e^{-|t|} - \frac{P_\epsilon(t)}{(1 + t^2)} \right| \leq \epsilon \frac{|t|}{1 + t^2} \quad \implies \quad \left| u(t)e^{-|t|} - \frac{P_\epsilon(t)}{1 + t^2} \right| \leq \epsilon \cdot C.$$

This shows that $(1 + t^2)^{-1}P_\epsilon(t)$ approximates $\sqrt{|t|}e^{-|t|}$ in supremum norm.

Finally,

$$\begin{aligned}
\left| \frac{u(t)}{\sqrt{|t|}} e^{-|t|} - \frac{P_\epsilon(t)}{\sqrt{|t|(1+t^2)}} \right|_{L^2(\mathbb{R})} &\leq \epsilon \left(\int_{-\infty}^{\infty} \frac{|t|}{(1+t^2)^2} dt \right)^{1/2} \leq \epsilon C, \\
\left| \frac{|u(t)e^{-|t|}|}{\sqrt{|t|}} - e^{-|t|} \right|_{L^2(\mathbb{R})} &\leq \left| \frac{7}{4} \epsilon^{-3/2} t^2 - \frac{3}{4} \epsilon^{-7/4} t^4 - 1 \right|_{L^2([-\epsilon, \epsilon])} \\
&\leq \left(\int_{-\epsilon}^{\epsilon} \left| \frac{7}{4} \frac{1}{\epsilon^{3/2}} t^2 \right|^2 dt \right)^{1/2} \\
&\quad + \left(\int_{-\epsilon}^{\epsilon} \left| \frac{3}{4} \frac{1}{\epsilon^{7/2}} t^4 \right|^2 dt \right)^{1/2} + \left(\int_{-\epsilon}^{\epsilon} 1 dt \right)^{1/2} \\
&\leq C\epsilon,
\end{aligned}$$

with a constant C independent of ϵ . \square

Definition 5.19. For $f : \mathbb{R} \rightarrow \mathbb{R}$ continuous and sufficiently decaying at ∞ , define

$$V_f : L^2(M) \rightarrow L^2([0, \infty) \times M) \quad \text{by} \quad V_f u(t) := \frac{1}{\sqrt{|t|}} f(t|D + E)u,$$

and correspondingly define $V_{f,D}$ with $D + E$ replaced by D .

Lemma 5.20. *In the situation above,*

$$\max\{\|V_D - V_{f,D}\|^2, \|V - V_f\|^2\} \leq \int_0^\infty |f(t)/\sqrt{t} - e^{-t}|^2 dt.$$

Proof. By definition, $(V - V_f)u(t) = g_t(D + E)u$ with

$$g_t(\lambda) = \frac{\sqrt{2|t\lambda|} e^{-t|\lambda|} - f(t|\lambda)}{\sqrt{t}}.$$

Then

$$\begin{aligned}
\int_0^\infty |g_t(\lambda)|^2 dt &= \int_0^\infty \left| \frac{1}{\sqrt{t}} (\sqrt{2|t\lambda|} e^{-t|\lambda|} - f(t|\lambda)) \right|^2 dt \\
&\stackrel{s=t|\lambda|}{=} \int_0^\infty \frac{1}{s} |\sqrt{s} e^{-s} - f(s)|^2 ds.
\end{aligned}$$

By Lemma 5.15,

$$\max\{\|V_D - V_{f,D}\|^2, \|V - V_f\|^2\} \leq \int_0^\infty |g_t(\lambda)|^2 dt = \int_0^\infty |e^{-t} - f(t)/\sqrt{t}|^2 dt,$$

as claimed. \square

In light of this and Lemma 5.15 we can find a polynomial $f(t)$ in $(1+t^2)^{-1}$, vanishing at ∞ , such that $V - V_f$ and $V_D - V_{f,D}$ are arbitrarily small in norm. The same applies then of course also to $(1 - P_R)(V - V_f)$ and $(1 - P_R)(V_D - V_{f,D})$.

Lemma 5.21. *For a fixed polynomial $f(t)$ in $(1+t^2)^{-1}$, $V_f - V_{f,D}$ is a linear combination of operators of the form*

$$\begin{aligned} u \mapsto \left(t \mapsto \frac{1}{\sqrt{|t|}} \frac{t(D+E)}{(1+t^2(D+E)^2)^l} tE \frac{1}{(1+t^2D^2)^k} u \right), \\ u \mapsto \left(t \mapsto \frac{1}{\sqrt{|t|}} \frac{1}{(1+t^2(D+E)^2)^l} tE \frac{tD}{(1+t^2D^2)^k} u \right), \end{aligned} \quad (5.22)$$

with $k, l \geq 1$.

Proof. First, expand $a^n - b^n = \sum_{k=1}^n a^{k-1}(a-b)b^{n-k}$. Apply this to

$$a = (1+t^2(D+E)^2)^{-1} \quad \text{and} \quad b = (1+t^2D^2)^{-1}.$$

Then

$$a - b = \frac{1}{1+t^2(D+E)^2} (1+t^2D^2 - 1 - t^2(D+E)^2) \frac{1}{1+t^2D^2},$$

and $1+t^2D^2 - 1 - t^2(D+E)^2 = -t(D+E)tE - tEtD$.

Recall that for $g(t) = 1/(1+t^2)^n$ we have

$$((V_g - V_{g,D})u)(t) = \frac{1}{\sqrt{|t|}} \left(\left(\frac{1}{1+t^2(D+E)^2} \right)^n - \left(\frac{1}{1+D^2} \right)^n \right) u.$$

Application of the formulas just derived immediately gives the result. \square

Lemma 5.23. *If $f(t)$ is a polynomial in $(1+t^2)^{-1}$, then*

$$\lim_{r \rightarrow 0} \|(1 - P_r)(V_f - V_{f,D})\| = 0.$$

Proof. Because of Lemma 5.21 it suffices to consider the operators given in (5.22). Because of the factor t in front of tE which is always present, they are of the form

$$u \mapsto (t \mapsto \sqrt{t}A(t)u),$$

where $A(t)$ is a uniformly norm-bounded family of operators: it is a composition of E and of functions of $(D+E)$ and of D that are bounded in operator norm uniformly in t . The statement follows immediately from the definition of the norm on $L^2([0, \infty), L^2(M))$. \square

Lemma 5.24. *If $f(t)$ is a polynomial in $(1+t^2)^{-1}$ and $0 < r < R < \infty$, then $(P_r - P_R)(V_f - V_{f,D})$ is a norm limit of operators of finite propagation.*

Proof. By Lemma 5.21, it suffices to compose the operators in (5.22) with $(P_r - P_R)$ and prove the statement for these compositions. Now observe that these operators have the form $u \mapsto (t \mapsto A(t)u)$, where

$$A(t) = \begin{cases} 0; & t > R \text{ or } t < r, \\ \phi_t(D+E)E\psi_t(D); & r \leq t \leq R. \end{cases}$$

Then $A(t)$ is a norm-continuous function with values in operators which are norm limits of finite-propagation operators: indeed, $\phi_t(\lambda)$, $\psi_t(\lambda)$ tend to 0 for $\lambda \rightarrow \pm\infty$ and depend continuously in supremum norm on t , so that $\phi_t(D + E)$, $\psi_t(D)$ are really limits of finite-propagation operators and depend norm-continuously on t . Therefore we can — up to an arbitrarily small error in norm — replace the function $A(t)$ by a (say, piecewise constant) function $B(t)$ of operators with fixed finite propagation S . It follows that $u \mapsto (t \mapsto B(t)u)$ has finite propagation of at most $\max\{R, S\}$. \square

Lemma 5.25. *If $f(t)$ is a polynomial in $(1 + t^2)^{-1}$ and $0 < r < R < \infty$ then $(P_r - P_R)(V_f - V_{f,D}) = B(i + rD)^{-1}$ for a bounded operator B .*

Proof. Because of Lemma 5.21, we have to show that

$$u \mapsto \left(t \mapsto \begin{cases} 0; & t > R \text{ or } t < r, \\ \frac{(t(D + E))^\eta}{(1 + t^2(D + E)^2)^k} \sqrt{|t|} E \frac{(tD)^{1-\eta}(i + rD)}{(1 + t^2D^2)^l} u; & r \leq t \leq R, \end{cases} \right)$$

with $k, l \geq 1$ and $\eta \in \{0, 1\}$ is bounded. Using that $\eta = 0$ or $\eta = 1$ and $l \geq 1$, this follows from the fact that

$$\lambda \mapsto \frac{r\lambda}{1 + t^2\lambda^2} \quad \text{and} \quad \lambda \mapsto \frac{t\lambda r\lambda}{1 + t^2\lambda^2}$$

are uniformly (in $t \geq r$) bounded functions of λ (note that we substitute D into these functions to obtain one factor making up the operator we have to consider, the remaining factors being controlled by the previous considerations). \square

Theorem 5.26. $V : L^2(M) \rightarrow L^2([0, \infty) \times M)$ covers the inclusion $\{0\} \times M \rightarrow [0, \infty) \times M$ in the D^* -sense.

Proof. Choose a polynomial $f(t)$ in $(1 + t^2)^{-1}$ as above such that $(1 - P_R)(V - V_f)$ and $(1 - P_R)(V_{f,D} - V_D)$ have small norm for all $R > 0$. We write

$$\begin{aligned} V &= P_R V + (1 - P_R)(V - V_f) + (1 - P_r)(V_f - V_{f,D}) \\ &\quad + (P_r - P_R)(V_f - V_{f,D}) + (1 - P_R)(V_{f,D} - V_D) + (1 - P_R)V_D \end{aligned}$$

for a suitable choice of R such that, using Lemma 5.16, $P_R V$ has small norm, and of r such that, using Lemma 5.23, $(1 - P_r)(V_f - V_{f,D})$ has small norm.

By Lemma 5.24, $(P_r - P_R)(V_f - V_{f,D})$ is a norm limit of finite-propagation operators. Let ϕ be a compactly supported continuous function on M and ψ a compactly supported continuous function on $[0, \infty) \times M$ with $\text{supp}(\psi) \cap (\{0\} \times \text{supp}(\phi)) = \emptyset$. Then $(i + rD)^{-1}\phi : L^2(M) \rightarrow L^2(M)$ is compact and therefore, by Lemma 5.25, also $\psi(P_r - P_R)(V_f - V_{f,D})\phi$ is compact (pseudolocal condition). Finally, in [PS 2014] we have shown, using unit propagation speed of the wave operator of D on M , that $(1 - P_R)V_D$ is a norm limit of finite-propagation operators that satisfy, in

addition, the pseudolocal condition. Note that these derivations did not use invertibility of D and therefore are valid in the present context. Summarizing, we have shown that V is a norm limit of finite-propagation operators F_ϵ with the additional property that $\psi F_\epsilon \phi$ is compact for any $\phi \in C_0(M)$ and $\psi \in C_0([0, \infty) \times M)$ with $\text{supp}(\psi) \cap (\{0\} \times \text{supp}(\phi)) = \emptyset$. This proves that V covers the inclusion in the D^* -sense. \square

The proof of Proposition 5.7 is now complete. \square

5B2. *Proof of Proposition 5.9: the operator $\frac{|D+C|+\partial_t}{D+C-\partial_t}$ belongs to $D^*(\tilde{M} \times \mathbb{R})^\Gamma$.*

Notation 5.27. In all this subsection we shall denote by M the total space of a Riemannian Γ -Galois covering with compact base M/Γ . We also consider a Riemannian manifold N . (In the application we have in mind $N = \mathbb{R}$.)

We consider a Γ -equivariant Dirac-type operator D on M acting on the sections of a Clifford module bundle; as before, we will in the notation ignore this bundle. Similarly, we consider a Dirac-type operator ∂ on N (and in the applications we have in mind we shall in fact take $\partial = i\partial_t$ on $N = \mathbb{R}$). We recall that D and ∂ are essentially self-adjoint; we shall not distinguish notationally between D , ∂ and their unique self-adjoint extensions.

We wish to prove that the operator $(|D + C| + \partial_t)/(D + C - \partial_t)$ belongs to $D^*(M \times \mathbb{R})^\Gamma$. In the course of the argument, it turns out that it is useful to work not only with L^2 , but also with the Sobolev spaces H^1, H^2 on our complete manifold M , or $M \times N$. We have to understand mapping properties for perturbed Dirac operators and functions of those, acting on these Sobolev spaces. Because we don't want to assume that our perturbation is a pseudodifferential operator, we can't use standard mapping properties here; instead we will rely on abstract functional analysis of unbounded operators on Hilbert spaces.

Definition 5.28. For a manifold like M , $H^1(M)$ is defined as the domain of the (unique self-adjoint extension of the) Dirac-type operator D endowed with the graph norm $(|Ds|^2 + |s|^2)^{1/2}$. Similarly, $H^2(M)$ is the domain of D^2 , endowed with the corresponding graph norm.

Proposition 5.29. *Let E be a self-adjoint bounded equivariant operator. Then $H^1(M)$ coincides with the domain of $D + E$ and its norm is equivalent to the graph norm $(|(D + E)s|^2 + |s|^2)^{1/2}$. If $D + E$ is invertible, this norm is also equivalent to $|(D + E)s|$. In this case, $(D + E)^{-1} : L^2 \rightarrow H^1$ is bounded, even an isometry.*

Similarly, $H^2(M)$ is equal to the domain of $D^2 + 1$ endowed with the graph norm; in particular, $(D^2 + 1) : L^2 \rightarrow H^2$ is an isometry for the appropriate choice of norm on H^2 .

Any suitably equivariant differential operator of order 1 is a bounded operator $H^1 \rightarrow L^2$. In particular, on $M \times N$, both D and ∂ are bounded operators $H^1(M \times N) \rightarrow L^2(M \times N)$.

Finally, $|D|$ and $|D + E|$ are bounded maps $H^1 \rightarrow L^2$.

Proof. Only the statements about $|D|$, $|D + E|$ are not easy or standard. For those, we can write

$$|D + E| = \frac{|D + E|}{D + E + i}(D + E + i),$$

where the bounded function $(|D + E|)/(D + E + i)$ of $D + E$ is a bounded operator on L^2 and $D + E + i$ is bounded from $H^1 \rightarrow L^2$. \square

Proposition 5.30. *Given the Riemannian product $M \times N$, a Dirac-type operator D on M with bounded equivariant self-adjoint perturbation $E : L^2(M) \rightarrow L^2(M)$, a compactly supported function ϕ on N (acting by pointwise multiplication) and a compact operator $K : L^2(M) \rightarrow L^2(M)$, the composition*

$$H^1(M \times N) \xrightarrow{D+E} L^2(M \times N) = L^2(M) \otimes L^2(N) \xrightarrow{K \otimes \phi} L^2(M \times N)$$

is compact.

Proof. Using the usual reduction techniques (write ϕ as a finite sum of functions with support in a coordinate neighborhood, use charts to plant these coordinate neighborhoods into T^l) one reduces to the case where $N = T^l$. Then, the operator with general ϕ is the composition of the bounded operator on $L^2(M \times T^l)$ given by multiplication with ϕ with the special operator where $\phi = 1$. It therefore suffices to show that the latter one is compact, and we set $\phi = 1$.

We now apply the strategy of the proof of the Rellich lemma. We have to understand a bit better the domain $H^1(M \times T^l)$, which, as the domain of the (perturbed) Dirac operator on $M \times T^l$, is a subspace of $L^2(M \times T^l)$ (with its own norm). We write $L^2(M \times T^l) = L^2(M) \otimes L^2(T^l)$. Using the Fourier transform, we unitarily identify $L^2(T^l)$ with $l^2(\mathbb{Z}^l)$. Using the Browder–Gårding spectral decomposition for the self-adjoint unbounded operator $D + E$, we write as a direct integral $L^2(M) = \int d\mu(\lambda) H_\lambda$. By the definition of compact operators as norm limits of finite-rank operators, we can replace $K : L^2(M) \rightarrow L^2(M)$ up to an error of arbitrarily small norm by a finite-rank operator K_Λ such that K_Λ maps $\int_{-\Lambda}^\Lambda H_\lambda$ to itself and is zero on the complement. By definition of the spectral decomposition, $D + E$ acts on the direct integral by multiplication with the spectral parameter λ . In particular, $\int_{-\Lambda}^\Lambda H_\lambda$ is entirely contained in the domain of $D + E$ (i.e., in $H^1(M)$) and restricted to this subspace the norm of $D + E$ is certainly bounded by the norm of $D + E$ as a bounded operator from H^1 to L^2 . We denote this norm by C .

Thus, the Hilbert space $H^1(M \times T^l)$ has the direct summand $\int_{-\Lambda}^\Lambda H_\lambda \otimes H^1(T^l)$. Here, after Fourier transform $L^2(T^l) \cong l^2(\mathbb{Z}^l)$, we identify $H^1(T^l)$ with the domain

of the operator

$$l^2(\mathbb{Z}^l) \rightarrow l^2(\mathbb{Z}^l); \quad (\lambda_n)_{n \in \mathbb{Z}^k} \mapsto (|n|\lambda_n)_{n \in \mathbb{Z}}, \quad \text{with } |n_1, \dots, n_l| = |n_1| + \dots + |n_l|,$$

endowed with the graph norm. For $R > 0$, split

$$H^1(T^l) := V_R \oplus V_R^\perp,$$

where

$$V_R = \{(\lambda_n)_{n \in \mathbb{Z}^l} \mid \lambda_n = 0 \text{ if } |n| > R\}$$

is finite-dimensional. Note that the inclusion $V_R^\perp \hookrightarrow l^2(\mathbb{Z}^l)$ has norm $< R^{-1}$. We now conclude the following:

- (1) The operator $(K \otimes \text{id}_{L^2(T^l)}) \circ (D + E) : H^1(M \times T^k) \rightarrow L^2(M) \otimes L^2(T^k)$ is norm-close to $(K_\Lambda \otimes \text{id}_{L^2(T^l)}) \circ (D + E)$.
- (2) Restricted to the direct summand $\int_{-\Lambda}^\Lambda H_\lambda \otimes V_R$ of $H^1(M \times T^l)$, $(K_\Lambda \otimes \text{id}) \circ (D + E)$ has finite rank, with image $\text{im}(K_\Lambda) \otimes V_R$.
- (3) Restricted to the direct summand $\int_{-\Lambda}^\Lambda H_\lambda \otimes V_R^\perp$, the operator

$$D + E : H^1(M \times T^k) \rightarrow L^2(M \times T^k)$$

has norm $\leq C \cdot R^{-1}$, where R^{-1} comes from the ratio of the H^1 -norm and the L^2 -norm on V_R^\perp .

Finally, on the orthogonal complement of $\int_{-\Lambda}^\Lambda H_\lambda \otimes H^1(T^k)$ in $H^1(M \times T^k)$, by the choice of K_Λ , the operator $(K_\Lambda \otimes \text{id}) \circ (D + E)$ vanishes.

- (4) It follows that $(K_\Lambda \otimes \text{id}) \circ (D + E)$ is, up to an error of norm C/R , a finite-rank operator.
- (5) All together, $(K \otimes \text{id}) \circ (D + E) : H^1(M \times T^l) \rightarrow L^2(M \times T^l)$ is a norm limit of finite-rank operators, i.e., is compact. \square

Proposition 5.31. *If ϕ stands for the multiplication operator with the compactly supported C^1 -function ϕ and P for any first-order equivariant differential operator, the commutator $[\phi, P]$ is compact as an operator from $H^1(M)$ to $L^2(M)$.*

The same applies if P is replaced by E or $P + E$ for any equivariant self-adjoint bounded operator E which is a norm limit of operators with finite propagation.

Proof. The commutator $[\phi, P]$ is a multiplication operator with a derivative of ϕ , a compactly supported function. The latter ones are compact as maps from H^1 to L^2 .

For the perturbation, we have to consider also the commutator $\phi E - E\phi$. By finite propagation, up to a norm-small error we can write $\phi E = \phi E\psi$ with a compactly supported ψ . Then we only need to use that

$$H^1 \rightarrow L^2 \xrightarrow{\phi} L^2 \quad \text{and} \quad H^1 \rightarrow L^2 \xrightarrow{\psi} L^2$$

are compact by the Rellich lemma and that $E : L^2 \rightarrow L^2$ is bounded. \square

Proposition 5.32. *Let D be an equivariant Dirac-type operator on M and $E : L^2(M) \rightarrow L^2(M)$ an L^2 -bounded self-adjoint perturbation which is a norm limit of equivariant finite-propagation operators such that $D + E$ is invertible. Assume that ϕ is a compactly supported C^1 -function on $M \times N$. Then $|D + E|$, as a map from $H^1(M \times N)$ to $L^2(M \times N)$, is a limit of equivariant bounded operators $F_\epsilon : H^1(M \times N) \rightarrow L^2(M \times N)$ that are of finite propagation and such that $[F_\epsilon, \phi] : H^1(M \times N) \rightarrow L^2(M \times N)$ is compact.*

Proof. We can reduce to $\phi = \alpha(x)\beta(y)$ with $\alpha : M \rightarrow \mathbb{C}$, $\beta : N \rightarrow \mathbb{C}$ compactly supported C^1 -functions. We write $|D + E| : H^1(M \times N) \rightarrow L^2(M \times N)$ as

$$H^1(M \times N) \xrightarrow{D+E} L^2(M \times N) = L^2(M) \otimes L^2(N) \xrightarrow{(|D+E|/(D+E)) \otimes \text{id}} L^2(M) \otimes L^2(N).$$

Using Lemma 2.25, write $|D + E|/(D + E)$ on M as the limit of equivariant finite-propagation pseudolocal operators κ_ϵ and consider F_ϵ given by the composition

$$F_\epsilon : H^1(M \times N) \xrightarrow{D+E} L^2(M \times N) = L^2(M) \otimes L^2(N) \xrightarrow{\kappa_\epsilon \otimes \text{id}} L^2(M) \otimes L^2(N).$$

Observe that F_ϵ is of finite propagation as a bounded operator $H^1(M \times N) \rightarrow L^2(M \times N)$. We now show that it is also pseudolocal as a bounded operator $H^1(M \times N) \rightarrow L^2(M \times N)$. We have

$$[\phi, F_\epsilon] = ([\alpha, \kappa_\epsilon](D + E) \otimes \beta + \kappa_\epsilon([\alpha, (D + E)]) \otimes \beta).$$

Use here that β commutes with all the other operators, which allows us to split off the tensor factor β in $\phi = \beta\alpha$ throughout. The second summand is compact by Proposition 5.31. The first summand is compact by Proposition 5.30. \square

Proposition 5.33. *Given a Dirac-type operator D as above, the operator*

$$(1 + D^2)^{-1} : L^2 \rightarrow H^2$$

is a norm limit of equivariant finite-propagation operators $G_\epsilon : L^2 \rightarrow H^2$ with the property that $[\phi, G_\epsilon] : L^2 \rightarrow H^2$ is compact for any compactly supported smooth function on M .

Proof. We follow the idea of the proof of Proposition 4.19 in [PS 2014]. Consider $f(x) := (1 + x^2)^{-1}$, and arrange that $f(x) = g_\epsilon(x) + h_\epsilon(x)$, where $g_\epsilon(x)$ has compactly supported Fourier transform and the Fourier transforms of $h_\epsilon(x)$ together with its second derivative have small L^1 -norm; use here that $\hat{f}(\xi)$, even if not smooth at 0, is smooth away from 0 and all its derivatives are rapidly decreasing. Set $G_\epsilon := g_\epsilon(D)$. Then, by unit propagation of the wave operator, G_ϵ has finite

propagation, and

$$\begin{aligned}
 \|(1 + D^2)^{-1} - G_\epsilon\|_{L^2 \rightarrow H^2} &= \|h_\epsilon(D)\|_{L^2 \rightarrow H^2} \\
 &= \|(1 + D^2)h_\epsilon(D)\|_{L^2 \rightarrow L^2} \\
 &\leq |x| \mapsto (1 + x^2)h_\epsilon(x)_\infty \\
 &\leq |\hat{h}|_1 + |\hat{h}''|_1 \ll 1.
 \end{aligned}$$

The fact that G_ϵ is pseudolocal as a map $L^2 \rightarrow H^2$ is proved by employing a doubling trick and (standard) microlocal techniques, as in [PS 2014]. \square

A purely functional-analytic argument, which applies to Lipschitz manifolds without a pseudodifferential calculus, is given in [Zenobi 2015].

Proposition 5.34. *Let D be a Dirac-type operator on M with equivariant self-adjoint bounded perturbation $E : L^2(M) \rightarrow L^2(M)$ as above such that $D + E$ is invertible and such that E is the norm limit of equivariant finite-propagation operators. Then $(D + E)^{-1} : L^2 \rightarrow H^1$ is a norm limit of equivariant finite-propagation operators $A_\epsilon : L^2 \rightarrow H^1$ with the property that for any compactly supported smooth function ϕ the operator $[\phi, A_\epsilon] : L^2 \rightarrow H^1$ is compact.*

Proof. Write

$$(D + E)^{-1} = (D + i)(1 + D^2)^{-1}((D + E - i)(D + E)^{-1} - E(D + E)^{-1}).$$

Here, $(D + E - i)(D + E)^{-1} : L^2 \rightarrow L^2$ is, by Lemma 2.25, an element in D^* , thus a norm limit of equivariant finite-propagation pseudolocal operators h_ϵ . Similarly $(D + E)^{-1} : L^2 \rightarrow L^2$ and therefore $E(D + E)^{-1} : L^2 \rightarrow L^2$ are elements in C^* , and so the latter is a norm limit of equivariant finite-propagation locally compact operators λ_ϵ . Next, $D + i : H^2 \rightarrow H^1$ is bounded with propagation 0 and has the property that $[\phi, D + i] : H^2 \rightarrow H^1$ is compact by the Rellich lemma. Finally, by Proposition 5.33 above, $(1 + D^2)^{-1} : L^2 \rightarrow H^2$ is the norm limit of equivariant finite-propagation pseudolocal operators $G_\epsilon : L^2 \rightarrow H^2$. Using the derivation property of $[\phi, \cdot]$ and approximating $(D + E)^{-1}$ by $A_\epsilon := (D + i)G_\epsilon(h_\epsilon - \lambda_\epsilon)$, we easily conclude the proof. \square

Corollary 5.35. *If $E : L^2(M) \rightarrow L^2(M)$ is a bounded self-adjoint equivariant operator and a norm limit of equivariant finite-propagation operators and D is a Dirac-type operator on M as above such that $D + E$ is invertible, then the operator*

$$\frac{|D + E| + \partial_t}{D + E - \partial_t} : L^2 \rightarrow L^2$$

belongs to $D^(M \times \mathbb{R})^\Gamma$.*

Proof. We must prove that $(|D + E| + \partial_t)/(D + E - \partial_t)$ is the norm limit of a sequence of equivariant bounded operators $H_\epsilon : L^2 \rightarrow L^2$ that are of finite propagation and pseudolocal. To check pseudolocality, by a density argument it suffices to consider the commutator with compactly supported smooth functions. We choose $N = \mathbb{R}$ in the previous propositions. We observe that $(D + E - \partial_t)$ is a summand of the perturbed Dirac-type operator

$$\begin{pmatrix} 0 & D + E - \partial_t \\ D + E + \partial_t & 0 \end{pmatrix}$$

on $M \times \mathbb{R}$; thus $(D + E - \partial_t)^{-1} : L^2 \rightarrow H^1$ is a norm limit of equivariant operators $A_\epsilon : L^2 \rightarrow H^1$ that are of finite propagation and commute up to compact operators with multiplication by compactly supported smooth functions. Here we have used Proposition 5.34 with $M \times \mathbb{R}$ instead of M . By Proposition 5.32, $|D + E|$, as a map from $H^1(M \times \mathbb{R})$ to $L^2(M \times \mathbb{R})$, is a limit of bounded equivariant operators $F_\epsilon : H^1(M \times \mathbb{R}) \rightarrow L^2(M \times \mathbb{R})$ that are of finite propagation and such that $[F_\epsilon, \phi] : H^1(M \times \mathbb{R}) \rightarrow L^2(M \times \mathbb{R})$ is compact. Finally, $\partial_t : H^1 \rightarrow L^2$ has propagation zero and $[\phi, \partial_t]$ is clearly compact as a map $H^1 \rightarrow L^2$ (Rellich lemma). Summarizing, by writing

$$\frac{|D + E| + \partial_t}{D + E - \partial_t} = \frac{|D + E|}{D + E - \partial_t} + \frac{\partial_t}{D + E - \partial_t}$$

we see that $(|D + E| + \partial_t)/(D + E - \partial_t)$ is a sum of two elements in $D^*(M \times \mathbb{R})^\Gamma$ and it is therefore in $D^*(M \times \mathbb{R})^\Gamma$, as required. \square

Acknowledgments

We are glad to thank Georges Skandalis, Stephane Vassout, Charlotte Wahl, Rudolf Zeidler and Vito Felice Zenobi for interesting discussions. We thank the anonymous referee for a very careful reading of the original manuscript and for valuable suggestions. P.P. thanks the *Projet Algèbres d'Opérateurs* of *Institut de Mathématiques de Jussieu* for hospitality and financial support while part of this research was carried out; he also thanks Mathematisches Institut and the Courant Research Center “Higher order structures in mathematics” for their hospitality for several visits to Göttingen; the financial support of *Ministero dell'Università e della Ricerca Scientifica* (through the project “Spazi di Moduli e Teoria di Lie”) is also gratefully acknowledged. T. S. acknowledges the support of the Courant Research Center “Higher order structures in mathematics”.

References

- [Deeley and Goffeng 2014] R. Deeley and M. Goffeng, “Realizing the analytic surgery group of Higson and Roe geometrically, Part I: The geometric model”, preprint, 2014. To appear in *J. Homotopy Relat. Struct.* arXiv 1308.5990
- [Higson and Roe 2005a] N. Higson and J. Roe, “Mapping surgery to analysis, I: Analytic signatures”, *K-Theory* **33**:4 (2005), 277–299. MR 2007c:19004 Zbl 1083.19002
- [Higson and Roe 2005b] N. Higson and J. Roe, “Mapping surgery to analysis, II: Geometric signatures”, *K-Theory* **33**:4 (2005), 301–324. MR 2007c:19005 Zbl 1083.19003
- [Higson and Roe 2005c] N. Higson and J. Roe, “Mapping surgery to analysis, III: Exact sequences”, *K-Theory* **33**:4 (2005), 325–346. MR 2007c:19006 Zbl 1085.19002
- [Hilsum and Skandalis 1992] M. Hilsum and G. Skandalis, “Invariance par homotopie de la signature à coefficients dans un fibré presque plat”, *J. Reine Angew. Math.* **423** (1992), 73–99. MR 93b:46137 Zbl 0731.55013
- [Kasparov 1995] G. G. Kasparov, “ K -theory, group C^* -algebras, and higher signatures (conspicuous)”, pp. 101–146 in *Novikov conjectures, index theorems and rigidity, Vol. 1* (Oberwolfach, 1993), edited by S. C. Ferry et al., London Math. Soc. Lecture Note Ser. **226**, Cambridge Univ. Press, 1995. MR 97j:58153 Zbl 0957.58020
- [Leichtnam and Piazza 1998] E. Leichtnam and P. Piazza, “Spectral sections and higher Atiyah–Patodi–Singer index theory on Galois coverings”, *Geom. Funct. Anal.* **8**:1 (1998), 17–58. MR 2000b:58043 Zbl 0913.58055
- [Leichtnam and Piazza 2000] E. Leichtnam and P. Piazza, “A higher Atiyah–Patodi–Singer index theorem for the signature operator on Galois coverings”, *Ann. Global Anal. Geom.* **18**:2 (2000), 171–189. MR 2001i:58045 Zbl 0957.58017
- [Leichtnam and Piazza 2003] E. Leichtnam and P. Piazza, “Dirac index classes and the noncommutative spectral flow”, *J. Funct. Anal.* **200**:2 (2003), 348–400. MR 2004k:58043 Zbl 1030.58018
- [Leichtnam et al. 2000] E. Leichtnam, J. Lott, and P. Piazza, “On the homotopy invariance of higher signatures for manifolds with boundary”, *J. Differential Geom.* **54**:3 (2000), 561–633. MR 2002e:58040 Zbl 1032.58012
- [Lott 1992] J. Lott, “Superconnections and higher index theory”, *Geom. Funct. Anal.* **2**:4 (1992), 421–454. MR 93j:58128 Zbl 0777.58038
- [Lück 2002] W. Lück, “A basic introduction to surgery theory”, pp. 1–224 in *Topology of high-dimensional manifolds, Vol. 2* (Trieste, 2001), edited by F. T. Farrell et al., ICTP Lect. Notes **9**, Abdus Salam Int. Cent. Theoret. Phys., Trieste, 2002. MR 2004a:57041 Zbl 1045.57020
- [Melrose and Piazza 1997a] R. B. Melrose and P. Piazza, “Families of Dirac operators, boundaries and the b -calculus”, *J. Differential Geom.* **46**:1 (1997), 99–180. MR 99a:58144 Zbl 0955.58020
- [Melrose and Piazza 1997b] R. B. Melrose and P. Piazza, “An index theorem for families of Dirac operators on odd-dimensional manifolds with boundary”, *J. Differential Geom.* **46**:2 (1997), 287–334. MR 99a:58145 Zbl 0920.58053
- [PS 2007] P. Piazza and T. Schick, “Bordism, rho-invariants and the Baum–Connes conjecture”, *J. Noncommut. Geom.* **1**:1 (2007), 27–111. MR 2008a:58025 Zbl 1158.58012
- [PS 2014] P. Piazza and T. Schick, “Rho-classes, index theory and Stolz’ positive scalar curvature sequence”, *J. Topol.* **7**:4 (2014), 965–1004. MR 3286895 Zbl 06382027
- [Ranicki 2002] A. Ranicki, *Algebraic and geometric surgery*, Oxford University Press, Oxford, 2002. MR 2005e:57075 Zbl 1003.57001

- [Roe 2002] J. Roe, “Comparing analytic assembly maps”, *Q. J. Math.* **53**:2 (2002), 241–248. MR 2003c:46083 Zbl 1014.46045
- [Rosenberg and Weinberger 2006] J. Rosenberg and S. Weinberger, “The signature operator at 2”, *Topology* **45**:1 (2006), 47–63. MR 2006i:19007 Zbl 1103.58012
- [Schick 2005] T. Schick, “ L^2 -index theorems, KK -theory, and connections”, *New York J. Math.* **11** (2005), 387–443. MR 2006h:19007 Zbl 1093.58009
- [Siegel 2012a] P. Siegel, *Homological calculations with the analytic structure group*, Ph.D. thesis, The Pennsylvania State University, 2012, <http://search.proquest.com/docview/1356843373>. MR 3153339
- [Siegel 2012b] P. Siegel, “The Mayer–Vietoris sequence for the analytic structure group”, preprint, 2012. arXiv 1212.0241
- [Wahl 2010] C. Wahl, “Product formula for Atiyah–Patodi–Singer index classes and higher signatures”, *J. K-Theory* **6**:2 (2010), 285–337. MR 2011j:58030 Zbl 1220.19005
- [Wahl 2013a] C. Wahl, “The Atiyah–Patodi–Singer index theorem for Dirac operators over C^* -algebras”, *Asian J. Math.* **17**:2 (2013), 265–319. MR 3078932 Zbl 1283.58017
- [Wahl 2013b] C. Wahl, “Higher ρ -invariants and the surgery structure set”, *J. Topol.* **6**:1 (2013), 154–192. MR 3029424 Zbl 1314.19006
- [Wall 1999] C. T. C. Wall, *Surgery on compact manifolds*, 2nd ed., Mathematical Surveys and Monographs **69**, American Mathematical Society, Providence, RI, 1999. MR 2000a:57089 Zbl 0935.57003
- [Wegge-Olsen 1993] N. E. Wegge-Olsen, *K-theory and C^* -algebras, a friendly approach*, Oxford University Press, New York, 1993. MR 95c:46116 Zbl 0780.46038
- [Xie and Yu 2014] Z. Xie and G. Yu, “Positive scalar curvature, higher rho invariants and localization algebras”, *Adv. Math.* **262** (2014), 823–866. MR 3228443 Zbl 1306.58008
- [Yu 1997] G. Yu, “Localization algebras and the coarse Baum–Connes conjecture”, *K-Theory* **11**:4 (1997), 307–318. MR 98i:19004 Zbl 0888.46047
- [Zeidler 2015] R. Zeidler, “Positive scalar curvature and product formulas for secondary index invariants”, preprint, 2015. arXiv 1412.0685
- [Zenobi 2015] V. F. Zenobi, “Mapping the surgery exact sequence for topological manifolds to analysis”, preprint, 2015. To appear in *J. Homotopy Relat. Struct.* arXiv 1405.6323

Received 9 Jan 2015. Accepted 7 Apr 2015.

PAOLO PIAZZA: piazza@mat.uniroma1.it

Dipartimento di Matematica, Università degli Studi di Roma "La Sapienza", P.le Aldo Moro 5, I-00185 Roma, Italy

THOMAS SCHICK: thomas.schick@math.uni-goettingen.de

Mathematisches Institut, Georg-August-Universität Göttingen, Bunsenstr. 3, D-37073 Göttingen, Germany