

ANNALS OF K-THEORY

vol. 4 no. 3 2019

Vanishing theorems for the negative K -theory of stacks

Marc Hoyois and Amalendu Krishna



A JOURNAL OF THE K-THEORY FOUNDATION

Vanishing theorems for the negative K -theory of stacks

Marc Hoyois and Amalendu Krishna

We prove that the homotopy algebraic K -theory of tame quasi-DM stacks satisfies cdh-descent. We apply this descent result to prove that if \mathcal{X} is a Noetherian tame quasi-DM stack and $i < -\dim(\mathcal{X})$, then $K_i(\mathcal{X})[1/n] = 0$ if n is nilpotent on \mathcal{X} and $K_i(\mathcal{X}, \mathbb{Z}/n) = 0$ if n is invertible on \mathcal{X} . Our descent and vanishing results apply more generally to certain Artin stacks whose stabilizers are extensions of finite group schemes by group schemes of multiplicative type.

1. Introduction	439
2. Preliminaries on algebraic stacks	441
3. Perfect complexes on algebraic stacks	445
4. K -theory of perfect stacks	451
5. G -theory and the case of regular stacks	457
6. Cdh-descent for homotopy K -theory	461
7. The vanishing theorems	463
Acknowledgments	470
References	470

1. Introduction

The negative K -theory of rings was defined by Bass [1968] and later generalized to all schemes by Thomason and Trobaugh [1990], who established its fundamental properties such as localization, excision, Mayer–Vietoris, and the projective bundle formula.

As explained in [Thomason and Trobaugh 1990], these properties of K -theory give rise to the Bass–Thomason–Trobaugh nonconnective K -theory, or K^B -theory, which is usually nontrivial in negative degrees for singular schemes. A famous conjecture of Weibel asserts that for a Noetherian scheme X of Krull dimension d , the group $K_i(X)$ vanishes for $i < -d$. This conjecture was settled by Weibel [2001] for excellent surfaces, by Cortiñas, Haesemeyer, Schlichting and Weibel [Cortiñas et al. 2008] for schemes essentially of finite type over a field of characteristic zero, and recently by Kerz, Strunk and Tamme [Kerz et al. 2018] for all Noetherian schemes.

MSC2010: primary 19D35; secondary 14D23.

Keywords: algebraic K -theory, negative K -theory, algebraic stacks.

Before a complete proof of Weibel’s conjecture for schemes appeared in [Kerz et al. 2018], Kelly [2014] used the alteration methods of de Jong and Gabber to show that the vanishing conjecture for negative K -theory holds in characteristic $p > 0$ if one is allowed to invert p . Later, Kerz and Strunk [2017] gave a different proof of Kelly’s theorem by proving Weibel’s conjecture for negative homotopy K -theory, or KH -theory, a variant of K -theory introduced by Weibel [1989]. In their proof, Kerz and Strunk used the method of flatification by blow-up instead of alterations.

It is natural to ask for an extension of Weibel’s conjecture to algebraic stacks. The algebraic K -theory of quotient stacks was introduced by Thomason [1987a] in order to study algebraic K -theory of a scheme which can be equipped with an action of a group scheme. The localization, excision, and Mayer–Vietoris properties for the algebraic K -theory of tame Deligne–Mumford stacks were proven by the second author and Østvær [Krishna and Østvær 2012], and together with the projective bundle formula they were established for more general quotient stacks by the second author and Ravi [Krishna and Ravi 2018]. The K^B -theory of Bass–Thomason–Trobaugh and the KH -theory of Weibel were also generalized to such quotient stacks in [Krishna and Ravi 2018].

The purpose of this paper is to show that the approach of Kerz and Strunk can be generalized to a large class of algebraic stacks, including all tame Artin stacks in the sense of [Abramovich et al. 2008]. As a consequence, we obtain a generalization of Kelly’s vanishing theorem for the negative K -theory of such stacks.

1A. Vanishing of negative K -theory of stacks. Our main results apply to certain algebraic stacks with finite or multiplicative type stabilizers. More precisely, let \mathbf{Stk}' be the category consisting of the following algebraic stacks:

- stacks with separated diagonal and linearly reductive finite stabilizers;
- stacks with affine diagonal whose stabilizers are extensions of linearly reductive finite groups by groups of multiplicative type.

Note that \mathbf{Stk}' contains tame Artin stacks with separated diagonal in the sense of [Abramovich et al. 2008]. The *blow-up dimension* of a Noetherian stack \mathcal{X} is a modification of the Krull dimension which is invariant under blow-ups (see Definition 7.7); it coincides with the usual dimension when \mathcal{X} is a quasi-DM stack.

Theorem 1.1 (see Theorems 7.10, 7.14, and 7.16). *Let \mathcal{X} be a stack in \mathbf{Stk}' satisfying the resolution property or having finite inertia. Assume that \mathcal{X} is Noetherian of blow-up dimension d . Then the following hold.*

- (1) $KH_i(\mathcal{X}) = 0$ for $i < -d$.
- (2) If n is nilpotent on \mathcal{X} , $K_i(\mathcal{X})[1/n] = 0$ for $i < -d$.
- (3) If n is invertible on \mathcal{X} , $K_i(\mathcal{X}, \mathbb{Z}/n) = 0$ for $i < -d$.

1B. *Cdh-descent for the homotopy K -theory of stacks.* Cdh-descent plays a key role in all the existing vanishing theorems for negative K -theory. In the recent proof of Weibel’s conjecture in [Kerz et al. 2018], the central result is pro-cdh-descent for nonconnective algebraic K -theory. Earlier results towards Weibel’s conjecture used instead cdh-descent for homotopy K -theory KH . For schemes over a field of characteristic zero, this descent result was proven by Haesemeyer [2004], and in arbitrary characteristic, it was shown by Cisinski [2013]. For the equivariant KH -theory of quasiprojective schemes acted on by a diagonalizable or finite linearly reductive group over an arbitrary base, cdh-descent was proven by the first author [Hoyois 2016]. A key step in the proof of Theorem 1.1 is a generalization of the latter to more general algebraic stacks:

Theorem 1.2 (see Theorem 6.2). *The presheaf of homotopy K -theory spectra KH satisfies cdh-descent on the category \mathbf{Stk}' .*

Cdh-descent is the combination of two descent properties: descent for the Nisnevich topology and descent for abstract blow-ups. Descent for the Nisnevich topology holds much more generally (see Corollary 4.10) and in fact it holds for nonconnective K -theory as well (see Corollary 4.6). Descent for abstract blow-ups is more difficult and uses several nontrivial properties of the category \mathbf{Stk}' . The proof ultimately relies on the proper base change theorem in stable equivariant motivic homotopy theory, proved in [Hoyois 2017].

2. Preliminaries on algebraic stacks

A *stack* in this text means a quasicompact and quasiseparated algebraic stack. Note that all morphisms between such stacks are quasicompact and quasiseparated. Similarly, algebraic spaces and schemes are always assumed to be quasicompact and quasiseparated. We say that a morphism of stacks is *representable* if it is representable by algebraic spaces, and *schematic* if it is representable by schemes. Recall that the diagonal of a stack is representable by definition; see [Stacks 2005–, Tag 026N]. If \mathcal{X} is a stack, k is a field, and $x : \mathrm{Spec}(k) \rightarrow \mathcal{X}$ is a k -point, then the stabilizer $G_x \rightarrow \mathrm{Spec}(k)$ is a flat separated group scheme of finite type [Stacks 2005–, Tag 0B8D].

All group schemes are assumed flat and finitely presented. With this convention, if G is a group scheme over a scheme S , then $\mathcal{B}G = [S/G]$ is a stack. Recall that G is called *linearly reductive* if the pushforward functor $\mathrm{QCoh}(\mathcal{B}G) \rightarrow \mathrm{QCoh}(S)$ on quasicohherent sheaves is exact. One knows from [Abramovich et al. 2008, Theorem 2.16] that a finite étale group scheme G over S is linearly reductive if and only if its degree at each point of S is prime to the residual characteristic. Diagonalizable group schemes are also linearly reductive by [SGA 3₁, 1970, Exposé I, Théorème 5.3.3]. As linear reductivity is an fpqc-local property on S [Abramovich

et al. 2008, Proposition 2.4], every group scheme of multiplicative type is linearly reductive.

We say that a group scheme G is *almost multiplicative* if it is an extension of a finite étale group scheme by a group scheme of multiplicative type. Since the class of linearly reductive group schemes is closed under quotients and extensions [Alper 2013, Proposition 12.17], an almost multiplicative group scheme G over S is linearly reductive if and only if, for every $s \in S$, the number of geometric components of G_s is invertible in $\kappa(s)$.

2A. Quasiprojective morphisms. Recall from [Laumon and Moret-Bailly 2000, §14.3] that if \mathcal{X} is a stack and \mathcal{A}^\bullet is a quasicoherent sheaf of graded $\mathcal{O}_{\mathcal{X}}$ -algebras, then $\text{Proj}(\mathcal{A}^\bullet)$ is a local construction on the fppf site of \mathcal{X} just like for schemes and hence defines a schematic morphism of stacks $q : \text{Proj}(\mathcal{A}^\bullet) \rightarrow \mathcal{X}$. A morphism $f : \mathcal{Y} \rightarrow \mathcal{X}$ is called *quasiprojective* [Laumon and Moret-Bailly 2000, §14.3.4; Rydh 2016, Theorem 8.6] if there is a finitely generated quasicoherent sheaf \mathcal{E} on \mathcal{X} and a factorization

$$\mathcal{Y} \xrightarrow{\iota} \mathbb{P}(\mathcal{E}) \xrightarrow{q} \mathcal{X}$$

of f , where $\mathbb{P}(\mathcal{E}) = \text{Proj}(\text{Sym}^\bullet(\mathcal{E}))$ and ι is a quasicompact immersion. We say that f is *projective* if it is quasiprojective and proper. It is clear that a quasiprojective morphism of stacks is schematic and hence representable.

Lemma 2.1. *If $f : \mathcal{X} \rightarrow \mathcal{Y}$ and $g : \mathcal{Y} \rightarrow \mathcal{Z}$ are quasiprojective (resp. projective) morphisms of stacks, then $g \circ f$ is quasiprojective (resp. projective).*

Proof. The proof is the same as [Hoyois 2017, Lemma 2.13], the key point being that every quasicoherent sheaf on a quasicompact quasiseparated stack is the colimit of its finitely generated quasicoherent subsheaves [Rydh 2016]. □

If $\mathcal{I} \subset \mathcal{O}_{\mathcal{X}}$ is a finitely generated quasicoherent sheaf of ideals, defining a finitely presented closed substack $\mathcal{Z} \subset \mathcal{X}$, then $\text{Proj}(\bigoplus_{i \geq 0} \mathcal{I}^i) = \text{Bl}_{\mathcal{Z}}(\mathcal{X})$ is called the blow-up of \mathcal{X} with center \mathcal{Z} . Note that $\text{Bl}_{\mathcal{Z}}(\mathcal{X})$ is a closed substack of $\mathbb{P}(\mathcal{I})$. Since \mathcal{I} is finitely generated, it follows that the structure map $\text{Bl}_{\mathcal{Z}}(\mathcal{X}) \rightarrow \mathcal{X}$ is projective. If $\mathcal{U} \subset \mathcal{X}$ is an open substack, we say that a blow-up of \mathcal{X} is \mathcal{U} -admissible if its center is disjoint from \mathcal{U} .

2B. Flatification by blow-ups.

Theorem 2.2 (Rydh). *Let \mathcal{S} be a quasicompact and quasiseparated algebraic stack and let $f : \mathcal{X} \rightarrow \mathcal{S}$ be a morphism of finite type. Let \mathcal{F} be a finitely generated quasicoherent $\mathcal{O}_{\mathcal{X}}$ -module. Let $\mathcal{U} \subseteq \mathcal{S}$ be an open substack such that $f|_{\mathcal{U}}$ is of finite presentation and $\mathcal{F}|_{f^{-1}(\mathcal{U})}$ is of finite presentation and flat over \mathcal{U} . Then there exists a sequence of \mathcal{U} -admissible blow-ups $\tilde{\mathcal{S}} \rightarrow \mathcal{S}$ such that the strict transform of \mathcal{F} is of finite presentation and flat over $\tilde{\mathcal{S}}$.*

Proof. This is proved in [Rydh \geq 2019, Theorem 4.2]. \square

Lemma 2.3. *Let $f : \mathcal{Y} \rightarrow \mathcal{X}$ be a flat, proper, finitely presented, representable, and birational morphism of stacks. Then f is an isomorphism.*

Proof. We can assume that \mathcal{X} and hence \mathcal{Y} are algebraic spaces. Since f is flat, proper, and finitely presented, its fibers have locally constant dimension [Stacks 2005–, Tag 0D4R]. Since f is birational, its fibers must have dimension 0, so f is quasifinite [Stacks 2005–, Tag 04NV]. By Zariski’s main theorem [Stacks 2005–, Tag 082K], we deduce that f is in fact finite. Being finite, flat, and finitely presented, f is locally free, and it must be of rank 0. \square

Corollary 2.4 (Rydh). *Let $f : \mathcal{Y} \rightarrow \mathcal{X}$ be a proper representable morphism of stacks that is an isomorphism over some quasicompact open substack $\mathcal{U} \subset \mathcal{X}$. Then there exists a projective morphism $g : \tilde{\mathcal{Y}} \rightarrow \mathcal{Y}$ that is an isomorphism over \mathcal{U} such that $f \circ g$ is also projective.*

Proof. By first blowing up a finitely presented complement of \mathcal{U} in \mathcal{X} (which exists by [Rydh 2016, Proposition 8.2]) and replacing \mathcal{Y} by its strict transform, we may assume that \mathcal{U} is dense in \mathcal{X} . By Theorem 2.2, we can find a sequence of \mathcal{U} -admissible blow-ups $\tilde{\mathcal{X}} \rightarrow \mathcal{X}$ such that the strict transform $\tilde{f} : \tilde{\mathcal{Y}} \rightarrow \tilde{\mathcal{X}}$ is flat and of finite presentation. Let $g : \tilde{\mathcal{Y}} \rightarrow \mathcal{Y}$ be the induced map:

$$\begin{array}{ccccc} \mathcal{U} & \hookrightarrow & \tilde{\mathcal{Y}} & \xrightarrow{g} & \mathcal{Y} \\ \parallel & & \downarrow \tilde{f} & & \downarrow f \\ \mathcal{U} & \hookrightarrow & \tilde{\mathcal{X}} & \longrightarrow & \mathcal{X} \end{array}$$

Then g is a sequence of \mathcal{U} -admissible blow-ups and hence it is projective by Lemma 2.1. Moreover, \tilde{f} is flat, proper, finitely presented, representable, and birational, whence an isomorphism (Lemma 2.3). Thus, $f \circ g$ is the composition of an isomorphism and the sequence of blow-ups $\tilde{\mathcal{X}} \rightarrow \mathcal{X}$, so it is projective by Lemma 2.1. \square

2C. Nisnevich coverings of stacks. The following definition appears in [Hall and Rydh 2018, Definition 3.1] and, for Deligne–Mumford stacks, in [Krishna and Østvær 2012, Definition 6.3].

Definition 2.5. Let \mathcal{X} be a stack. A family of étale morphisms $\{\mathcal{U}_i \rightarrow \mathcal{X}\}_{i \in I}$ is called a *Nisnevich covering* if, for every $x \in \mathcal{X}$, there exists $i \in I$ and $u \in \mathcal{U}_i$ above x such that the induced morphism of residual gerbes $\eta_u \rightarrow \eta_x$ is an isomorphism.

Let $f : \mathcal{Y} \rightarrow \mathcal{X}$ be a morphism of stacks. A *monomorphic splitting sequence* for f is a sequence of quasicompact open substacks

$$\emptyset = \mathcal{U}_0 \subset \mathcal{U}_1 \subset \cdots \subset \mathcal{U}_n = \mathcal{X}$$

such that f admits a monomorphic section over the reduced substack $\mathcal{U}_i \setminus \mathcal{U}_{i-1}$ for all i . Note that if f is étale, such a section is an open immersion

$$\mathcal{U}_i \setminus \mathcal{U}_{i-1} \hookrightarrow \mathcal{Y} \times_{\mathcal{X}} (\mathcal{U}_i \setminus \mathcal{U}_{i-1}).$$

Proposition 2.6. *Let \mathcal{X} be a stack. A family of étale morphisms $\{\mathcal{U}_i \rightarrow \mathcal{X}\}_i$ is a Nisnevich covering if and only if the morphism $\coprod_i \mathcal{U}_i \rightarrow \mathcal{X}$ admits a monomorphic splitting sequence.*

Proof. See [Hall and Rydh 2018, Proposition 3.3]. □

Corollary 2.7. *Let \mathcal{X} be a stack and let $\{\mathcal{U}_i \rightarrow \mathcal{X}\}_{i \in I}$ be a Nisnevich covering. Then there exists a finite subset $J \subset I$ such that $\{\mathcal{U}_i \rightarrow \mathcal{X}\}_{i \in J}$ is a Nisnevich covering.*

Proof. This follows at once from Proposition 2.6. □

A Nisnevich square in the category of stacks is a Cartesian square of the form

$$\begin{array}{ccc} \mathcal{W} & \hookrightarrow & \mathcal{V} \\ \downarrow & & \downarrow f \\ \mathcal{U} & \xrightarrow{e} & \mathcal{X} \end{array} \tag{2.8}$$

where f is an étale morphism (not necessarily representable) and e is an open immersion with reduced complement \mathcal{Z} such that the induced map $\mathcal{Z} \times_{\mathcal{X}} \mathcal{V} \rightarrow \mathcal{Z}$ is an isomorphism. Nisnevich squares form a cd-structure on the category of stacks, in the sense of [Voevodsky 2010].

Proposition 2.9. *Let $f : \mathcal{Y} \rightarrow \mathcal{X}$ be a Nisnevich covering. Then there exist sequences of quasicompact open substacks*

$$\mathcal{Y}_1 \subset \cdots \subset \mathcal{Y}_n \subset \mathcal{Y}, \quad \emptyset = \mathcal{X}_0 \subset \mathcal{X}_1 \subset \cdots \subset \mathcal{X}_n = \mathcal{X},$$

such that $f(\mathcal{Y}_i) \subset \mathcal{X}_i$ and such that each square

$$\begin{array}{ccc} \mathcal{X}_{i-1} \times_{\mathcal{X}} \mathcal{Y}_i & \hookrightarrow & \mathcal{Y}_i \\ \downarrow & & \downarrow f \\ \mathcal{X}_{i-1} & \hookrightarrow & \mathcal{X}_i \end{array}$$

is a Nisnevich square.

Proof. The proof is exactly the same as [Morel and Voevodsky 1999, Proposition 1.4]. Let $\mathcal{X}_0 \subset \cdots \subset \mathcal{X}_n$ be a monomorphic splitting sequence for f (see Proposition 2.6), and $s_i : \mathcal{X}_i \setminus \mathcal{X}_{i-1} \rightarrow \mathcal{Y} \times_{\mathcal{X}} (\mathcal{X}_i \setminus \mathcal{X}_{i-1})$ a monomorphic section of the projection. Then s_i is an open immersion, so the complement of the image of s_i is a closed substack $\mathcal{Z}_i \subset \mathcal{Y} \times_{\mathcal{X}} \mathcal{X}_i$. We can then take $\mathcal{Y}_i = (\mathcal{Y} \times_{\mathcal{X}} \mathcal{X}_i) \setminus \mathcal{Z}_i$. □

Proposition 2.9 implies that the Grothendieck topology associated with the Nisnevich cd-structure is exactly the topology generated by Nisnevich coverings. The Nisnevich cd-structure on the category of stacks clearly satisfies the assumptions of Voevodsky’s descent criterion [Asok et al. 2017, Theorem 3.2.5]. It follows that a presheaf of spaces or spectra \mathcal{F} satisfies descent for Nisnevich coverings if and only if, for every Nisnevich square (2.8), the induced square

$$\begin{array}{ccc} \mathcal{F}(\mathcal{X}) & \xrightarrow{e^*} & \mathcal{F}(\mathcal{U}) \\ f^* \downarrow & & \downarrow \\ \mathcal{F}(\mathcal{V}) & \longrightarrow & \mathcal{F}(\mathcal{W}) \end{array}$$

is homotopy Cartesian.

The following recent result of Alper, Hall, and Rydh [Alper et al. ≥ 2019] on the Nisnevich-local structure of some stacks plays an important role in the proof of our cdh-descent theorem.

Theorem 2.10 (Alper–Hall–Rydh). *Let \mathcal{X} be a stack, let $x \in \mathcal{X}$ be a point, and let η_x be its residual gerbe. Suppose that the stabilizer of \mathcal{X} at a representative of x is a linearly reductive almost multiplicative group scheme. Then there exists*

- a morphism of affine schemes $U \rightarrow S$,
- a linearly reductive almost multiplicative group scheme G over S acting on U ,
- a commutative diagram of stacks

$$\begin{array}{ccc} \eta_x & \hookrightarrow & [U/G] \\ \parallel & & \downarrow f \\ \eta_x & \hookrightarrow & \mathcal{X} \end{array}$$

where f is étale.

If \mathcal{X} has affine diagonal, we can moreover choose f affine. If \mathcal{X} has finite inertia and coarse moduli space $\pi : \mathcal{X} \rightarrow X$, we can take S to be an étale neighborhood of $\pi(x)$ in X .

Remark 2.11. Linearly reductive almost multiplicative group schemes are called *nice* in [Hall and Rydh 2015] and [Alper et al. ≥ 2019], but this terminology is used differently in [Krishna and Ravi 2018], so we avoid using it. □

3. Perfect complexes on algebraic stacks

3A. Sheaves on stacks. Let \mathcal{X} be a stack. Let $\text{Lis-Ét}(\mathcal{X})$ denote the *lisse-étale site* of \mathcal{X} . Its objects are smooth morphisms $X \rightarrow \mathcal{X}$, where X is a quasicompact quasiseparated scheme. The coverings are generated by the étale covers of schemes. Let $\text{Mod}(\mathcal{X})$ denote the abelian category of sheaves of $\mathcal{O}_{\mathcal{X}}$ -modules, and $\text{QCoh}(\mathcal{X})$

that of quasicoherent sheaves, on $\text{Lis-}\acute{\text{E}}\text{t}(\mathcal{X})$. It is well known that $\text{QCoh}(\mathcal{X})$ and $\text{Mod}(\mathcal{X})$ are Grothendieck abelian categories and hence have enough injectives and all limits.

Let $\text{Ch}(\mathcal{X})$ denote the category of all (possibly unbounded) chain complexes over $\text{Mod}(\mathcal{X})$, and $\text{Ch}_{\text{qc}}(\mathcal{X})$ the full subcategory of $\text{Ch}(\mathcal{X})$ consisting of those chain complexes whose cohomology lies in $\text{QCoh}(\mathcal{X})$. Let $D(\mathcal{X})$ and $D_{\text{qc}}(\mathcal{X})$ denote their corresponding derived categories, obtained by inverting quasi-isomorphisms. If $\mathcal{Z} \hookrightarrow \mathcal{X}$ is a closed substack with open complement $j : \mathcal{U} \hookrightarrow \mathcal{X}$, we let

$$\text{Ch}_{\text{qc}, \mathcal{Z}}(\mathcal{X}) = \{\mathcal{F} \in \text{Ch}_{\text{qc}}(\mathcal{X}) \mid j^*(\mathcal{F}) \text{ is quasi-isomorphic to } 0\}.$$

The derived category of $\text{Ch}_{\text{qc}, \mathcal{Z}}(\mathcal{X})$ is denoted by $D_{\text{qc}, \mathcal{Z}}(\mathcal{X})$.

Let $j : \mathcal{X} \rightarrow \mathcal{Y}$ be a smooth morphism of algebraic stacks. We then have the pullback functor $j^* : \text{Mod}(\mathcal{Y}) \rightarrow \text{Mod}(\mathcal{X})$, which preserves quasicoherent sheaves. Since j is smooth, the functor j^* is simply the restriction functor under the inclusion $\text{Lis-}\acute{\text{E}}\text{t}(\mathcal{X}) \subset \text{Lis-}\acute{\text{E}}\text{t}(\mathcal{Y})$.

Recall from [SGA 6 1971, Definition I.4.2] that a complex of \mathcal{O}_X -modules on a scheme X is perfect if it is locally quasi-isomorphic to a bounded complex of locally free sheaves.

Definition 3.1. Let \mathcal{X} be a stack. A chain complex $P \in \text{Ch}_{\text{qc}}(\mathcal{X})$ is called *perfect* if for any affine scheme $U = \text{Spec}(A)$ with a smooth morphism $s : U \rightarrow \mathcal{X}$, the complex of A -modules $s^*(P) \in \text{Ch}(\text{Mod}(A))$ is quasi-isomorphic to a bounded complex of finitely generated projective A -modules. Equivalently, $s^*(P)$ is a perfect complex in $\text{Ch}(\text{Mod}(A))$ in the sense of [Thomason and Trobaugh 1990].

It follows from [Krishna and Ravi 2018, Lemma 2.5] that the above definition coincides with that of [Thomason and Trobaugh 1990] if \mathcal{X} is a scheme. We denote the derived category of perfect complexes on \mathcal{X} by $D_{\text{perf}}(\mathcal{X})$. The derived category of perfect complexes on \mathcal{X} whose cohomology is supported on a closed substack \mathcal{Z} is denoted by $D_{\text{perf}, \mathcal{Z}}(\mathcal{X})$.

We also need to use the canonical dg-enhancements of the triangulated categories $D_{\text{qc}}(\mathcal{X})$ and $D_{\text{perf}}(\mathcal{X})$, denoted by $\text{D}_{\text{qc}}(\mathcal{X})$ and $\text{D}_{\text{perf}}(\mathcal{X})$, respectively, whose construction we now recall. If \mathcal{X} is an affine scheme, $\text{D}_{\text{qc}}(\mathcal{X})$ is the usual symmetric monoidal derived dg-category of $\mathcal{O}(\mathcal{X})$. The 2-category of stacks embeds fully faithfully in the 2-category of presheaves of groupoids on affine schemes, which further embeds in the ∞ -category $\text{sPre}(\mathbf{Aff})$ of simplicial presheaves on affine schemes. Then one defines D_{qc} as a presheaf of symmetric monoidal dg-categories on $\text{sPre}(\mathbf{Aff})$ to be the homotopy right Kan extension of $\text{D}_{\text{qc}}|_{\mathbf{Aff}}$; see [Lurie 2018, §6.2]. In other words, it is the unique extension of $\text{D}_{\text{qc}}|_{\mathbf{Aff}}$ that transforms homotopy colimits into homotopy limits. One can show that D_{qc} satisfies descent for the fpqc topology on $\text{sPre}(\mathbf{Aff})$ [Lurie 2018, Proposition 6.2.3.1]. For \mathcal{X} a stack, the

homotopy category of $D_{\text{qc}}(\mathcal{X})$ is then equivalent to $D_{\text{qc}}(\mathcal{X})$. If \mathcal{X} is an algebraic space (or more generally a Deligne–Mumford stack), this is proved in [Lurie 2018, Proposition 6.2.4.1]. In general, this follows from the description of $D_{\text{qc}}(\mathcal{X})$ in terms of a smooth representable cover of \mathcal{X} by an algebraic space; see for instance [Hall and Rydh 2017, §1.1]. Finally, $D_{\text{perf}} \subset D_{\text{qc}}$ is the full symmetric monoidal dg-subcategory spanned by the dualizable objects. Since the process of passing to dualizable objects preserves homotopy limits of dg-categories [Lurie 2017, Proposition 4.6.1.11], D_{perf} is similarly the unique extension of $D_{\text{perf}}|_{\mathbf{Aff}}$ to $\mathbf{sPre}(\mathbf{Aff})$ that transforms homotopy colimits into homotopy limits, and it satisfies fpqc descent.

Proposition 3.2. *Let $f : \mathcal{X}' \rightarrow \mathcal{X}$ be an étale morphism of stacks and let $\mathcal{Z} \subset \mathcal{X}$ be a closed substack with quasicompact open complement such that the projection $\mathcal{Z} \times_{\mathcal{X}} \mathcal{X}' \rightarrow \mathcal{Z}$ is an isomorphism of associated reduced stacks. Then the functor*

$$f^* : D_{\text{perf}, \mathcal{Z}}(\mathcal{X}) \rightarrow D_{\text{perf}, \mathcal{Z} \times_{\mathcal{X}} \mathcal{X}'}(\mathcal{X}')$$

is an equivalence of triangulated categories.

Proof. The presheaf of dg-categories $\mathcal{X} \mapsto D_{\text{perf}}(\mathcal{X})$ satisfies descent for the fpqc topology on stacks. In particular, it satisfies Nisnevich descent, so that the square of dg-categories

$$\begin{array}{ccc} D_{\text{perf}}(\mathcal{X}) & \longrightarrow & D_{\text{perf}}(\mathcal{X} \setminus \mathcal{Z}) \\ f^* \downarrow & & \downarrow \\ D_{\text{perf}}(\mathcal{X}') & \longrightarrow & D_{\text{perf}}(\mathcal{X}' \setminus (\mathcal{Z} \times_{\mathcal{X}} \mathcal{X}')) \end{array}$$

is homotopy Cartesian. It follows that f^* induces an equivalence between the kernels of the horizontal functors. \square

3B. Perfect stacks.

Definition 3.3. Let \mathcal{X} be a stack. We say that \mathcal{X} is *perfect* if the triangulated category $D_{\text{qc}}(\mathcal{X})$ is compactly generated and $\mathcal{O}_{\mathcal{X}}$ is compact in $D_{\text{qc}}(\mathcal{X})$.

If $\mathcal{Z} \subset \mathcal{X}$ is a closed substack with quasicompact open complement, we say that the pair $(\mathcal{X}, \mathcal{Z})$ is *perfect* if \mathcal{X} is perfect and there exists a perfect complex on \mathcal{X} with support $|\mathcal{Z}|$.

We will see in Proposition 3.5 below that our notion of perfect stack agrees with the one introduced in [Ben-Zvi et al. 2010], except that we do not require perfect stacks to have affine diagonal.

Let $f : \mathcal{X}' \rightarrow \mathcal{X}$ be a morphism of stacks. We say that f is *concentrated* if for every morphism $g : \mathcal{Z} \rightarrow \mathcal{X}$, the morphism $f' : \mathcal{X}' \times_{\mathcal{X}} \mathcal{Z} \rightarrow \mathcal{Z}$ has finite cohomological dimension for quasicohherent sheaves.

Lemma 3.4. *Let $f : \mathcal{X}' \rightarrow \mathcal{X}$ be a representable morphism of stacks. Then f is concentrated. In particular, if $\mathcal{O}_{\mathcal{X}}$ is compact, then $\mathcal{O}_{\mathcal{X}'}$ is compact.*

Proof. Since f is representable, and since concentrated morphisms have faithfully flat descent by [Hall and Rydh 2017, Lemma 2.5(2)], we can assume that f is a morphism of algebraic spaces. Now the result follows because any quasi-compact and quasiseparated morphism of algebraic spaces is concentrated [Stacks 2005–, Tag 073G]. For the second statement, it suffices to show using [Neeman 1996, Theorem 5.1] that the right adjoint $f_* : D_{\text{qc}}(\mathcal{X}') \rightarrow D_{\text{qc}}(\mathcal{X})$ of f^* preserves small coproducts. This follows from the first statement and [Hall and Rydh 2017, Theorem 2.6(3)]. \square

Proposition 3.5. *Let $(\mathcal{X}, \mathcal{Z})$ be a perfect pair. Then the triangulated category $D_{\text{qc}, \mathcal{Z}}(\mathcal{X})$ is compactly generated. Moreover, an object of $D_{\text{qc}, \mathcal{Z}}(\mathcal{X})$ is compact if and only if it is perfect.*

Proof. Since $\mathcal{O}_{\mathcal{X}} \in D_{\text{qc}}(\mathcal{X})$ is compact and since a perfect complex on \mathcal{X} is dualizable, it follows that every perfect complex on \mathcal{X} is compact. On the other hand, it follows from the proofs of [Krishna and Ravi 2018, Proposition 2.7, Lemma 2.8] that compact objects of $D_{\text{qc}}(\mathcal{X})$ and $D_{\text{qc}, \mathcal{Z}}(\mathcal{X})$ are perfect. The only remark we need to make here is that the proofs in [loc. cit.] assume that \mathcal{X} is a quotient stack. However, this assumption is used only to ensure that if we choose an atlas $u : U \rightarrow \mathcal{X}$, then u has finite cohomological dimension for quasicoherent sheaves. But this follows from Lemma 3.4 because \mathcal{X} has representable diagonal and hence u is representable. Finally, the existence of a perfect complex with support $|\mathcal{Z}|$ implies, by [Hall and Rydh 2017, Lemma 4.10], that $D_{\text{qc}, \mathcal{Z}}(\mathcal{X})$ is compactly generated. \square

Lemma 3.6. *Let $f : \mathcal{Y} \rightarrow \mathcal{X}$ be a schematic morphism of stacks with a relatively ample family of line bundles. If $D_{\text{qc}}(\mathcal{X})$ is compactly generated, so is $D_{\text{qc}}(\mathcal{Y})$.*

Proof. Let $\{\mathcal{L}_i\}_{i \in I}$ be an f -ample family of line bundles on \mathcal{Y} . By Lemma 3.4, f is a concentrated morphism. It follows from [Hall and Rydh 2017, Theorem 2.6(3)] that $f_* : D_{\text{qc}}(\mathcal{Y}) \rightarrow D_{\text{qc}}(\mathcal{X})$ preserves small coproducts, and hence that its left adjoint f^* preserves compact objects. It therefore suffices to show that $D_{\text{qc}}(\mathcal{Y})$ is generated by the objects $f^*(\mathcal{F}) \otimes \mathcal{L}_i^{\otimes -n}$, for $\mathcal{F} \in D_{\text{qc}}(\mathcal{X})$ compact, $i \in I$, and $n \geq 1$. So let $\mathcal{G} \in D_{\text{qc}}(\mathcal{Y})$ be such that $\text{Hom}(f^*(\mathcal{F}) \otimes \mathcal{L}_i^{\otimes -n}, \mathcal{G}) = 0$ for every \mathcal{F} compact, $i \in I$, and $n \geq 1$. By adjunction, we have $\text{Hom}(\mathcal{F}, f_*(\mathcal{G} \otimes \mathcal{L}_i^{\otimes n})) = 0$. Since $D_{\text{qc}}(\mathcal{X})$ is compactly generated, it follows that

$$f_*(\mathcal{G} \otimes \mathcal{L}_i^{\otimes n}) = 0 \tag{3.7}$$

for every $i \in I$ and $n \geq 1$.

To show that $\mathcal{G} = 0$ in $D_{\text{qc}}(\mathcal{Y})$, we let $u : U \rightarrow \mathcal{X}$ be a smooth surjective morphism such that U is affine. This gives rise to a Cartesian square

$$\begin{array}{ccc} V & \xrightarrow{v} & \mathcal{Y} \\ g \downarrow & & \downarrow f \\ U & \xrightarrow{u} & \mathcal{X} \end{array}$$

where V is a scheme. As v is faithfully flat, it suffices to show $v^*(\mathcal{G}) = 0$. It follows from [Hall and Rydh 2017, Corollary 4.13] and (3.7) that $g_*(v^*\mathcal{G} \otimes v^*(\mathcal{L}_i)^{\otimes n}) = 0$ for all $i \in I$. Replacing \mathcal{Y} by V and \mathcal{L}_i by $v^*(\mathcal{L}_i)$, we can assume that \mathcal{X} is an affine scheme, so that \mathcal{Y} is a scheme and $\{\mathcal{L}_i\}_{i \in I}$ is an ample family of line bundles on \mathcal{Y} . In this case, (3.7) says that $\mathrm{Hom}(\mathcal{L}_i^{\otimes -n}[m], \mathcal{G}) = 0$ for all $i \in I$, $n \geq 1$, and $m \in \mathbb{Z}$. But this implies that \mathcal{G} is acyclic because $D_{\mathrm{qc}}(\mathcal{Y})$ is generated by $\{\mathcal{L}_i^{\otimes -n}\}_{i \in I, n \geq 1}$. Indeed, $D_{\mathrm{qc}}(\mathcal{Y})$ is compactly generated by bounded complexes of vector bundles [Thomason and Trobaugh 1990, Theorem 2.3.1(d)], and every vector bundle admits an epimorphism from a sum of line bundles of the form $\mathcal{L}_i^{\otimes -n}$. \square

Proposition 3.8. *Let $(\mathcal{X}, \mathcal{Z})$ be a perfect pair.*

- (1) *For every algebraic space Y and closed subspace $W \subset Y$ with quasicompact open complement, $(\mathcal{X} \times Y, \mathcal{Z} \times W)$ is perfect.*
- (2) *For every schematic morphism $f : \mathcal{Y} \rightarrow \mathcal{X}$ with a relatively ample family of line bundles, $(\mathcal{Y}, \mathcal{Y} \times_{\mathcal{X}} \mathcal{Z})$ is perfect.*

Proof. Let \mathcal{P} be a perfect complex on \mathcal{X} with support $|\mathcal{Z}|$.

(1) By [Hall and Rydh 2017, Theorem A], there exists a perfect complex \mathcal{Q} on Y with support $|W|$. Then $\pi_1^*(\mathcal{P}) \otimes \pi_2^*(\mathcal{Q})$ is a perfect complex on $\mathcal{X} \times Y$ with support $|\mathcal{Z} \times W|$. Since the projection $\pi_1 : \mathcal{X} \times Y \rightarrow \mathcal{X}$ is representable, $\mathcal{O}_{\mathcal{X} \times Y}$ is compact by Lemma 3.4. It remains to show that $D_{\mathrm{qc}}(\mathcal{X} \times Y)$ is compactly generated. We claim that there is an equivalence of presentable dg-categories

$$D_{\mathrm{qc}}(\mathcal{X} \times Y) \simeq D_{\mathrm{qc}}(\mathcal{X}) \otimes D_{\mathrm{qc}}(Y). \quad (3.9)$$

Since the tensor product of compactly generated dg-categories is compactly generated, this will complete the proof. Since Y is a quasicompact and quasiseparated algebraic space, the dg-category $D_{\mathrm{qc}}(Y)$ is dualizable [Lurie 2018, §9.4], and hence tensoring with $D_{\mathrm{qc}}(Y)$ preserves homotopy limits. Since $D_{\mathrm{qc}}(-)$ is the homotopy right Kan extension of its restriction to affine schemes, we are reduced to proving (3.9) when \mathcal{X} is an affine scheme, in which case it is a special case of [Lurie 2018, Corollary 9.4.2.4].

(2) The perfect complex $f^*(\mathcal{P})$ has support $|\mathcal{Y} \times_{\mathcal{X}} \mathcal{Z}|$. By Lemma 3.4, $\mathcal{O}_{\mathcal{Y}}$ is compact. It remains to show that $D_{\mathrm{qc}}(\mathcal{Y})$ is compactly generated, but this follows from Lemma 3.6. \square

Proposition 3.10. *Let $(\mathcal{X}, \mathcal{Z})$ be a perfect pair and let $j : \mathcal{U} \hookrightarrow \mathcal{X}$ be the open immersion complement to \mathcal{Z} . Then*

$$j^* : \frac{D_{\mathrm{perf}}(\mathcal{X})}{D_{\mathrm{perf}, \mathcal{Z}}(\mathcal{X})} \rightarrow D_{\mathrm{perf}}(\mathcal{U})$$

is an equivalence of triangulated categories, up to direct factors.

Proof. For any pair $(\mathcal{X}, \mathcal{Z})$, we have an equivalence of triangulated categories

$$j^* : \frac{D_{\text{qc}}(\mathcal{X})}{D_{\text{qc}, \mathcal{Z}}(\mathcal{X})} \rightarrow D_{\text{qc}}(\mathcal{U}).$$

Indeed, the functor $j_* : D_{\text{qc}}(\mathcal{U}) \rightarrow D_{\text{qc}}(\mathcal{X})$ is fully faithful by flat base change, so j_*j^* is a localization endofunctor of $D_{\text{qc}}(\mathcal{X})$ whose kernel is $D_{\text{qc}, \mathcal{Z}}(\mathcal{X})$ by definition. The claim now follows from [Krause 2010, Proposition 4.9.1]. If $(\mathcal{X}, \mathcal{Z})$ is perfect, then \mathcal{U} is also perfect by Proposition 3.8(2). By Proposition 3.5, all three categories are compactly generated and their subcategories of compact and perfect objects coincide. We conclude using [Krause 2010, Theorem 5.6.1]. \square

Proposition 3.11. *Suppose that \mathcal{X} is the limit of a filtered diagram (\mathcal{X}_α) of perfect stacks with affine transition morphisms. Then \mathcal{X} is perfect and the canonical map*

$$\text{hocolim}_\alpha D_{\text{perf}}(\mathcal{X}_\alpha) \rightarrow D_{\text{perf}}(\mathcal{X}) \tag{3.12}$$

is a weak equivalence of dg-categories.

Proof. It follows from Proposition 3.8(2) that \mathcal{X} is perfect. By Proposition 3.5, $D_{\text{qc}}(\mathcal{X})$ is compactly generated and $D_{\text{qc}}(\mathcal{X})^c = D_{\text{perf}}(\mathcal{X})$, and similarly for each \mathcal{X}_α . Since the pullback functors $D_{\text{qc}}(\mathcal{X}_\alpha) \rightarrow D_{\text{qc}}(\mathcal{X}_\beta)$ preserve compact objects, it follows from [Lurie 2009, Propositions 5.5.7.6 and 5.5.7.8] and [Lurie 2017, Lemma 7.3.5.10] that (3.12) is a weak equivalence if and only if the canonical map

$$D_{\text{qc}}(\mathcal{X}) \rightarrow \text{holim}_\alpha D_{\text{qc}}(\mathcal{X}_\alpha) \tag{3.13}$$

is a weak equivalence. Choosing a smooth hypercover of some \mathcal{X}_α by schemes and using flat base change, we see that the map (3.13) is the homotopy limit of a cosimplicial diagram of similar maps with \mathcal{X}_α replaced by a scheme. Hence, it suffices to prove that (3.12) is a weak equivalence when \mathcal{X}_α is a scheme, but this follows from [Thomason and Trobaugh 1990, Proposition 3.20]. \square

We now state the following two results of Hall and Rydh, which provide many examples of perfect stacks.

Theorem 3.14 (Hall–Rydh). *Let \mathcal{X} be a stack satisfying one of the following properties.*

- (1) \mathcal{X} has characteristic zero.
- (2) \mathcal{X} has linearly reductive almost multiplicative stabilizers.
- (3) \mathcal{X} has finitely presented inertia and linearly reductive almost multiplicative stabilizers at points of positive characteristic.

Then $\mathcal{O}_\mathcal{X}$ is compact in $D_{\text{qc}}(\mathcal{X})$.

Proof. See [Hall and Rydh 2015, Theorem 2.1]. \square

Theorem 3.15 (Hall–Rydh). *Let \mathcal{X} be a stack satisfying the following properties.*

- (1) $\mathcal{O}_{\mathcal{X}}$ is compact in $D_{\text{qc}}(\mathcal{X})$.
- (2) *There exists a faithfully flat, representable, separated, and quasifinite morphism $f : \mathcal{X}' \rightarrow \mathcal{X}$ of finite presentation such that \mathcal{X}' has affine stabilizers and satisfies the resolution property.*

Then, for every closed substack $\mathcal{Z} \subset \mathcal{X}$, the pair $(\mathcal{X}, \mathcal{Z})$ is perfect.

Proof. By Lemma 3.4, $\mathcal{O}_{\mathcal{X}'} = f^*(\mathcal{O}_{\mathcal{X}})$ is compact in $D_{\text{qc}}(\mathcal{X}')$. Since \mathcal{X}' has affine stabilizers and satisfies the resolution property, it has affine diagonal by [Gross 2017, Theorem 1.1]. Since moreover $\mathcal{O}_{\mathcal{X}'}$ is compact, it follows from [Hall and Rydh 2017, Proposition 8.4] that \mathcal{X}' is *crisp*. We now apply [Hall and Rydh 2017, Theorem C] to conclude that \mathcal{X} is also *crisp*. By definition of crispness, this implies that $(\mathcal{X}, \mathcal{Z})$ is perfect. \square

Corollary 3.16. *Let \mathcal{X} be a quasi-DM stack with separated diagonal and linearly reductive stabilizers. Then, for every closed substack $\mathcal{Z} \subset \mathcal{X}$, the pair $(\mathcal{X}, \mathcal{Z})$ is perfect.*

Proof. Recall that a quasi-DM stack is a stack whose diagonal is quasifinite. It follows from [Stacks 2005–, Tag 06MC] that a stack \mathcal{X} is quasi-DM if and only if there exists an affine scheme X and a faithfully flat map $f : X \rightarrow \mathcal{X}$ of finite presentation which is quasifinite. Since the diagonal of \mathcal{X} is representable and separated, it follows that f is representable and separated. Since X is affine and hence has the resolution property, the corollary follows from Theorems 3.14 and 3.15. \square

Corollary 3.17. *Let \mathcal{X} be a stack with affine diagonal and linearly reductive almost multiplicative stabilizers. Then, for every closed substack $\mathcal{Z} \subset \mathcal{X}$, $(\mathcal{X}, \mathcal{Z})$ is perfect.*

Proof. By Theorem 2.10, there exists a Nisnevich covering $\{f_i : [U_i/G_i] \rightarrow \mathcal{X}\}_{i \in I}$, where f_i is affine, U_i is affine over an affine scheme S_i , and G_i is a linearly reductive almost multiplicative group scheme over S_i . By taking a further affine Nisnevich covering of S_i , we can ensure that G_i is almost isotrivial and hence that $[U_i/G_i]$ has the resolution property; see [Hoyois 2017, Example 2.8 and Remark 2.9]. By Corollary 2.7, we can also assume that I is finite. Let $\mathcal{X}' = \coprod_i [U_i/G_i]$. Then the induced map $\mathcal{X}' \rightarrow \mathcal{X}$ is faithfully flat, quasifinite, and affine. Since \mathcal{X}' has the resolution property, we conclude that $(\mathcal{X}, \mathcal{Z})$ is perfect by Theorems 3.14 and 3.15. \square

4. K -theory of perfect stacks

In this section, we establish some descent properties of the K -theory, negative K -theory, and homotopy K -theory of stacks. Special cases of these results were earlier proven in [Krishna and Østvær 2012; Krishna and Ravi 2018; Hoyois 2016].

4A. Localization, excision, and continuity. Let \mathcal{X} be an algebraic stack. The algebraic K -theory spectrum of \mathcal{X} is defined to be the K -theory spectrum of the complicial bi-Waldhausen category of perfect complexes in $\mathrm{Ch}_{\mathrm{qc}}(\mathcal{X})$ in the sense of [Thomason and Trobaugh 1990, §1.5.2]. Here, the complicial bi-Waldhausen category structure is given with respect to the degreewise split monomorphisms as cofibrations and quasi-isomorphisms as weak equivalences. This K -theory spectrum is denoted by $K(\mathcal{X})$. Equivalently, one may define $K(\mathcal{X})$ as the K -theory spectrum of the dg-category $\mathrm{D}_{\mathrm{perf}}(\mathcal{X})$; see [Blumberg et al. 2013, Corollary 7.12]. Note that the negative homotopy groups of $K(\mathcal{X})$ are zero [Thomason and Trobaugh 1990, §1.5.3]. We shall extend this definition to negative integers in the next section. For a closed substack \mathcal{Z} of \mathcal{X} , $K(\mathcal{X} \text{ on } \mathcal{Z})$ is the K -theory spectrum of the complicial bi-Waldhausen category of those perfect complexes on \mathcal{X} which are acyclic on $\mathcal{X} \setminus \mathcal{Z}$.

Theorem 4.1 (localization). *Let $(\mathcal{X}, \mathcal{Z})$ be a perfect pair and let $j : \mathcal{U} \hookrightarrow \mathcal{X}$ be the open immersion complement to \mathcal{Z} . Then the morphisms of spectra*

$$K(\mathcal{X} \text{ on } \mathcal{Z}) \rightarrow K(\mathcal{X}) \xrightarrow{j^*} K(\mathcal{U})$$

induce a long exact sequence

$$\begin{aligned} \cdots \rightarrow K_i(\mathcal{X} \text{ on } \mathcal{Z}) \rightarrow K_i(\mathcal{X}) \rightarrow K_i(\mathcal{U}) \rightarrow K_{i-1}(\mathcal{X} \text{ on } \mathcal{Z}) \rightarrow \cdots \\ \rightarrow K_0(\mathcal{X} \text{ on } \mathcal{Z}) \rightarrow K_0(\mathcal{X}) \rightarrow K_0(\mathcal{U}). \end{aligned}$$

Proof. This follows from Proposition 3.10 as in [Krishna and Ravi 2018, Theorem 3.4]. \square

Theorem 4.2 (excision). *Let $f : \mathcal{X}' \rightarrow \mathcal{X}$ be an étale morphism of stacks and let $\mathcal{Z} \subset \mathcal{X}$ be a closed substack with quasicompact open complement such that the projection $\mathcal{Z} \times_{\mathcal{X}} \mathcal{X}' \rightarrow \mathcal{Z}$ is an isomorphism of associated reduced stacks. Then the map $f^* : K(\mathcal{X} \text{ on } \mathcal{Z}) \rightarrow K(\mathcal{X}' \text{ on } \mathcal{Z} \times_{\mathcal{X}} \mathcal{X}')$ is a homotopy equivalence of spectra.*

Proof. This follows from Proposition 3.2 using [Thomason and Trobaugh 1990, Theorem 1.9.8]. \square

Theorem 4.3 (continuity). *Let \mathcal{X} be the limit of a filtered diagram (\mathcal{X}_α) of perfect stacks with affine transition morphisms. Then the canonical map*

$$\mathrm{hocolim}_{\alpha} K(\mathcal{X}_\alpha) \rightarrow K(\mathcal{X})$$

is a homotopy equivalence.

Proof. This follows from Proposition 3.11 and the fact that K preserves filtered homotopy colimits of dg-categories. \square

4B. The Bass construction and negative K -theory. The nonconnective K -theory spectrum of any stack may be defined from the complicial bi-Waldhausen category of perfect complexes, following [Schlichting 2006], or from the dg-category $D_{\text{perf}}(\mathcal{X})$, following [Cisinski and Tabuada 2011]. This allows one to define the negative K -theory of stacks.

In this subsection, we will see that for perfect stacks a nonconnective K -theory spectrum K^B can be defined much more explicitly using the construction of Bass–Thomason–Trobaugh. One may prove that this construction agrees with those of Schlichting and Cisinski–Tabuada exactly as in [Krishna and Ravi 2018, Theorem 3.21].

The K^B -theory spectrum $K^B(\mathcal{X})$ was constructed in [Krishna and Ravi 2018, §3E] based on the following two assumptions.

- (1) \mathcal{X} is a quotient stack of the form $[X/G]$ over a field, where G is a linearly reductive group scheme.
- (2) \mathcal{X} satisfies the resolution property.

Since perfect stacks need not satisfy these conditions, we cannot directly quote the results of [Krishna and Ravi 2018] for the construction of the K^B -theory of stacks. But the proofs are identical to those in [Krishna and Ravi 2018] using Theorems 4.1 and 4.2, so we only give a brief sketch of the construction. The existence of the K^B -theory is based on the following version of the fundamental theorem of Bass.

Theorem 4.4 (Bass fundamental theorem). *Let \mathcal{X} be a perfect stack and let $\mathcal{X}[T]$ denote the stack $\mathcal{X} \times \text{Spec}(\mathbb{Z}[T])$. Then the following hold.*

- (1) For $n \geq 1$, there is an exact sequence

$$0 \rightarrow K_n(\mathcal{X}) \xrightarrow{(p_1^*, -p_2^*)} K_n(\mathcal{X}[T]) \oplus K_n(\mathcal{X}[T^{-1}]) \xrightarrow{(j_1^*, j_2^*)} K_n(\mathcal{X}[T, T^{-1}]) \xrightarrow{\partial_T} K_{n-1}(\mathcal{X}) \rightarrow 0.$$

Here p_1^*, p_2^* are induced by the projections $\mathcal{X}[T] \rightarrow \mathcal{X}$, etc. and j_1^*, j_2^* are induced by the open immersions $\mathcal{X}[T^{\pm 1}] = \mathcal{X}[T, T^{-1}] \rightarrow \mathcal{X}[T]$, etc. The sum of these exact sequences for $n = 1, 2, \dots$ is an exact sequence of graded $K_*(\mathcal{X})$ -modules.

- (2) For $n \geq 0$, $\partial_T : K_{n+1}(\mathcal{X}[T^{\pm 1}]) \rightarrow K_n(\mathcal{X})$ is naturally split by a map h_T of $K_*(\mathcal{X})$ -modules. Indeed, the cup product with $T \in K_1(\mathbb{Z}[T^{\pm 1}])$ splits ∂_T up to a natural automorphism of $K_n(\mathcal{X})$.

- (3) There is an exact sequence

$$0 \rightarrow K_0(\mathcal{X}) \xrightarrow{(p_1^*, -p_2^*)} K_0(\mathcal{X}[T]) \oplus K_0(\mathcal{X}[T^{-1}]) \xrightarrow{(j_1^*, j_2^*)} K_0(\mathcal{X}[T^{\pm 1}]).$$

Proof. The proof of this theorem is word for word identical to the proof of its scheme version given in [Thomason and Trobaugh 1990, Theorem 6.1], once we know that the algebraic K -theory spectrum satisfies the following properties:

- (1) The projective bundle formula for the projective line $\mathbb{P}^1_{\mathcal{X}}$.
- (2) Localization for the pairs $(\mathbb{P}^1_{\mathcal{X}}, \mathcal{X}[T^{-1}])$ and $(\mathcal{X}[T], \mathcal{X}[T^{\pm 1}])$.
- (3) Excision.

Property (1) follows from [Krishna and Ravi 2018, Theorem 3.8], which holds for any algebraic stack. Property (2) follows from Proposition 3.8(1) and Theorem 4.1, and property (3) is Theorem 4.2. \square

As an immediate consequence of Theorem 4.4, one obtains the following.

Theorem 4.5. *Let \mathcal{X} be a perfect stack. Then there exists a spectrum $K^B(\mathcal{X})$, together with a natural map $K(\mathcal{X}) \rightarrow K^B(\mathcal{X})$ of spectra inducing isomorphisms $\pi_i K(\mathcal{X}) \cong \pi_i K^B(\mathcal{X})$ for $i \geq 0$, which satisfies the following properties.*

- (1) *Let $\mathcal{Z} \subset \mathcal{X}$ be a closed substack with quasicompact open complement $j: \mathcal{U} \hookrightarrow \mathcal{X}$ such that $(\mathcal{X}, \mathcal{Z})$ is perfect. Then there is a homotopy fiber sequence of spectra*

$$K^B(\mathcal{X} \text{ on } \mathcal{Z}) \rightarrow K^B(\mathcal{X}) \xrightarrow{j^*} K^B(\mathcal{U}).$$

- (2) *Let $f: \mathcal{Y} \rightarrow \mathcal{X}$ be an étale map between perfect stacks such that the projection $\mathcal{Z} \times_{\mathcal{X}} \mathcal{Y} \rightarrow \mathcal{Z}$ is an isomorphism on the associated reduced stacks. Then the map $f^*: K^B(\mathcal{X} \text{ on } \mathcal{Z}) \rightarrow K^B(\mathcal{Y} \text{ on } \mathcal{Z} \times_{\mathcal{X}} \mathcal{Y})$ is a homotopy equivalence.*
- (3) *Let $\pi: \mathbb{P}(\mathcal{E}) \rightarrow \mathcal{X}$ be the projective bundle associated to a vector bundle \mathcal{E} on \mathcal{X} of rank r . Then the map*

$$\prod_0^{r-1} K^B(\mathcal{X}) \rightarrow K^B(\mathbb{P}(\mathcal{E}))$$

that sends (a_0, \dots, a_{r-1}) to $\sum_i \mathcal{O}(-i) \otimes \pi^(a_i)$ is a homotopy equivalence.*

- (4) *Let $i: \mathcal{Y} \hookrightarrow \mathcal{X}$ be a regular closed immersion and let $p: \mathcal{X}' \rightarrow \mathcal{X}$ be the blow-up of \mathcal{X} with center \mathcal{Y} . Then the square of spectra*

$$\begin{array}{ccc} K^B(\mathcal{X}) & \xrightarrow{i^*} & K^B(\mathcal{Y}) \\ p^* \downarrow & & \downarrow \\ K^B(\mathcal{X}') & \longrightarrow & K^B(\mathcal{X}' \times_{\mathcal{X}} \mathcal{Y}) \end{array}$$

is homotopy Cartesian.

- (5) *Suppose \mathcal{X} is the limit of a filtered diagram (\mathcal{X}_α) of perfect stacks with affine transition morphisms. Then the canonical map $\text{hocolim}_\alpha K^B(\mathcal{X}_\alpha) \rightarrow K^B(\mathcal{X})$ is a homotopy equivalence.*

Proof. The spectrum $K^B(\mathcal{X})$ is constructed word for word using [Theorem 4.4](#) and the formalism given in (6.2)–(6.4) of [\[Thomason and Trobaugh 1990\]](#) for the case of schemes. The proof of the asserted properties is a standard deduction from the analogous properties of $K(\mathcal{X})$. The sketch of this deduction for (1)–(4) can be found in [\[Krishna and Ravi 2018, Theorem 3.20\]](#). Note that quasicompact open substacks of \mathcal{X} , projective bundles over \mathcal{X} , and blow-ups of \mathcal{X} are perfect stacks by [Proposition 3.8\(2\)](#). For (5), it suffices to check $\text{colim}_\alpha \pi_n K^B(\mathcal{X}_\alpha) \cong \pi_n K^B(\mathcal{X})$ for all $n \in \mathbb{Z}$. This follows from [Theorem 4.3](#), since $\pi_{-n} K^B(\mathcal{X})$, for $n > 0$, is a natural retract of $K_0(\mathbb{G}_m^n \times \mathcal{X})$. □

Corollary 4.6. *Let*

$$\begin{array}{ccc} \mathcal{W} & \hookrightarrow & \mathcal{V} \\ \downarrow & & \downarrow f \\ \mathcal{U} & \xrightarrow{e} & \mathcal{X} \end{array}$$

be a Nisnevich square of stacks, and suppose that the pairs $(\mathcal{X}, \mathcal{X} \setminus \mathcal{U})$ and $(\mathcal{V}, \mathcal{V} \setminus \mathcal{W})$ are perfect. Then the induced square of spectra

$$\begin{array}{ccc} K^B(\mathcal{X}) & \xrightarrow{f^*} & K^B(\mathcal{V}) \\ e^* \downarrow & & \downarrow \\ K^B(\mathcal{U}) & \longrightarrow & K^B(\mathcal{W}) \end{array}$$

is homotopy Cartesian.

Proof. This follows immediately from [Theorem 4.5\(1\)](#) and (2). □

Remark 4.7. We remark that if $(\mathcal{X}, \mathcal{X} \setminus \mathcal{U})$ is a perfect pair and if the map $f : \mathcal{V} \rightarrow \mathcal{X}$ in [Corollary 4.6](#) is representable and separated, then $(\mathcal{V}, \mathcal{V} \setminus \mathcal{W})$ is automatically a perfect pair. The reason is that in this case, f is quasi-affine by Zariski’s main theorem for stacks [\[Laumon and Moret-Bailly 2000, Theorem 16.5\]](#) and one can apply [Proposition 3.8\(2\)](#).

Since the homotopy groups of the two spectra $K(\mathcal{X})$ and $K^B(\mathcal{X})$ agree in non-negative degrees by [Theorem 4.5](#), we make the following definition.

Definition 4.8. Let \mathcal{X} be a perfect stack and $i \in \mathbb{Z}$. We let $K_i(\mathcal{X})$ denote the i -th homotopy group of the spectrum $K^B(\mathcal{X})$.

4C. The homotopy K -theory of perfect stacks. For $n \in \mathbb{N}$, let

$$\Delta^n = \text{Spec} \left(\frac{\mathbb{Z}[t_0, \dots, t_n]}{(\sum_i t_i - 1)} \right).$$

Recall that Δ^\bullet is a cosimplicial scheme. For a perfect stack \mathcal{X} , the homotopy

K -theory of \mathcal{X} is defined as

$$KH(\mathcal{X}) = \operatorname{hocolim}_{n \in \Delta^{\text{op}}} K^B(\mathcal{X} \times \Delta^n).$$

There is a natural map $K^B(\mathcal{X}) \rightarrow KH(\mathcal{X})$ induced by $0 \in \Delta^{\text{op}}$.

Theorem 4.9. *Let \mathcal{X} be a perfect stack.*

- (1) *Let $\mathcal{Z} \subset \mathcal{X}$ be a closed substack with quasicompact open complement $j: \mathcal{U} \hookrightarrow \mathcal{X}$ such that $(\mathcal{X}, \mathcal{Z})$ is perfect. Then there is a homotopy fiber sequence of spectra*

$$KH(\mathcal{X} \text{ on } \mathcal{Z}) \rightarrow KH(\mathcal{X}) \xrightarrow{j^*} KH(\mathcal{U}).$$

- (2) *Let $f: \mathcal{Y} \rightarrow \mathcal{X}$ be an étale map between perfect stacks such that the projection $\mathcal{Z} \times_{\mathcal{X}} \mathcal{Y} \rightarrow \mathcal{Z}$ is an isomorphism on the associated reduced stacks. Then the map $f^*: KH(\mathcal{X} \text{ on } \mathcal{Z}) \rightarrow KH(\mathcal{Y} \text{ on } \mathcal{Z} \times_{\mathcal{X}} \mathcal{Y})$ is a homotopy equivalence.*
- (3) *Let $\pi: \mathbb{P}(\mathcal{E}) \rightarrow \mathcal{X}$ be the projective bundle associated to a vector bundle \mathcal{E} on \mathcal{X} of rank r . Then the map*

$$\prod_0^{r-1} KH(\mathcal{X}) \rightarrow KH(\mathbb{P}(\mathcal{E}))$$

that sends (a_0, \dots, a_{r-1}) to $\sum_i \mathcal{O}(-i) \otimes \pi^(a_i)$ is a homotopy equivalence.*

- (4) *Let $i: \mathcal{Y} \hookrightarrow \mathcal{X}$ be a regular closed immersion and let $p: \mathcal{X}' \rightarrow \mathcal{X}$ be the blow-up of \mathcal{X} with center \mathcal{Y} . Then the square of spectra*

$$\begin{CD} KH(\mathcal{X}) @>i^*>> KH(\mathcal{Y}) \\ @Vp^*VV @VVV \\ KH(\mathcal{X}') @>>> KH(\mathcal{X}' \times_{\mathcal{X}} \mathcal{Y}) \end{CD}$$

is homotopy Cartesian.

- (5) *Suppose that \mathcal{X} is the limit of a filtered diagram (\mathcal{X}_α) of perfect stacks with affine transition morphisms. Then the canonical map*

$$\operatorname{hocolim}_{\alpha} KH(\mathcal{X}_\alpha) \rightarrow KH(\mathcal{X})$$

is a homotopy equivalence.

- (6) *Suppose that $u: \mathcal{E} \rightarrow \mathcal{X}$ is a vector bundle over \mathcal{X} . Then the induced map $u^*: KH(\mathcal{X}) \rightarrow KH(\mathcal{E})$ is a homotopy equivalence.*

Proof. Properties (1)–(5) follow immediately from the definition of $KH(\mathcal{X})$ and [Theorem 4.5](#). The proof of (6) for quotient stacks is given in [[Krishna and Ravi 2018](#), Theorem 5.2] and the same proof is valid for perfect stacks. □

Corollary 4.10. *Let*

$$\begin{array}{ccc} \mathcal{W} & \hookrightarrow & \mathcal{V} \\ \downarrow & & \downarrow f \\ \mathcal{U} & \xrightarrow{e} & \mathcal{X} \end{array}$$

be a Nisnevich square of stacks, and suppose that the pairs $(\mathcal{X}, \mathcal{X} \setminus \mathcal{U})$ and $(\mathcal{V}, \mathcal{V} \setminus \mathcal{W})$ are perfect. Then the induced square of spectra

$$\begin{array}{ccc} KH(\mathcal{X}) & \xrightarrow{f^*} & KH(\mathcal{V}) \\ e^* \downarrow & & \downarrow \\ KH(\mathcal{U}) & \longrightarrow & KH(\mathcal{W}) \end{array}$$

is homotopy Cartesian.

Proof. This follows immediately from [Theorem 4.9](#)(1) and (2). □

Remark 4.11. In [\[Hoyois 2016\]](#), a potentially different definition of KH is given for certain quotient stacks, which forces KH to be invariant with respect to vector bundle *torsors* and not just vector bundles. The two definitions agree for quotients of schemes by finite or multiplicative type groups, as we will show in [Lemma 6.1](#), but they may differ in general. We do not know if the above definition of KH has good properties for general perfect stacks. □

5. G -theory and the case of regular stacks

Our goal in this section is to show that perfect stacks that are Noetherian and regular have no negative K -groups. We do this by comparing the K -theory and G -theory of such stacks.

Let \mathcal{X} be a stack. Recall that $\text{Mod}(\mathcal{X})$ is the abelian category of $\mathcal{O}_{\mathcal{X}}$ -modules on the lisse-étale site of \mathcal{X} and $\text{QCoh}(\mathcal{X}) \subset \text{Mod}(\mathcal{X})$ is the abelian subcategory of quasicoherent sheaves.

Lemma 5.1. *Assume that \mathcal{X} is a Noetherian stack. Then the inclusion*

$$\iota_{\mathcal{X}} : \text{QCoh}(\mathcal{X}) \hookrightarrow \text{Mod}(\mathcal{X})$$

induces an equivalence of the derived categories $D^+(\text{QCoh}(\mathcal{X})) \xrightarrow{\sim} D_{\text{qc}}^+(\mathcal{X})$.

Proof. To show that $D^+(\text{QCoh}(\mathcal{X})) \rightarrow D_{\text{qc}}^+(\mathcal{X})$ is full and faithful, it suffices, using standard reduction, to show that the natural map

$$\text{Ext}_{\text{QCoh}(\mathcal{X})}^i(N, M) \rightarrow \text{Ext}_{\text{Mod}(\mathcal{X})}^i(N, M)$$

is an isomorphism for all $i \in \mathbb{Z}$ for $N, M \in \text{QCoh}(\mathcal{X})$. Since this is clearly true for $i \leq 0$, and since $\iota_{\mathcal{X}} : \text{QCoh}(\mathcal{X}) \rightarrow \text{Mod}(\mathcal{X})$ is exact, it suffices to show that this functor preserves injective objects.

Let \mathcal{F} be an injective quasicoherent sheaf on \mathcal{X} . Since a direct summand of an injective object in $\text{Mod}(\mathcal{X})$ is injective and since a quasicoherent sheaf which is injective as a sheaf of $\mathcal{O}_{\mathcal{X}}$ -modules is also an injective quasicoherent sheaf, it suffices to show that there is an inclusion $\mathcal{F} \hookrightarrow \mathcal{G}$ in $\text{QCoh}(\mathcal{X})$ such that $\iota_{\mathcal{X}}(\mathcal{G})$ is injective in $\text{Mod}(\mathcal{X})$.

Since \mathcal{X} is Noetherian, we can find a smooth atlas $u : U \rightarrow \mathcal{X}$, where U is a Noetherian scheme. We can now find an inclusion $u^*(\mathcal{F}) \hookrightarrow \mathcal{H}$ in $\text{QCoh}(U)$ such that \mathcal{H} is injective as a sheaf of \mathcal{O}_U -modules, by [Thomason and Trobaugh 1990, B.4]. We now consider the maps

$$\mathcal{F} \rightarrow u_*u^*(\mathcal{F}) \rightarrow u_*(\mathcal{H}). \tag{5.2}$$

As U is Noetherian, it is clear that $u_*(\mathcal{H})$ is a quasicoherent sheaf on \mathcal{X} . Furthermore, u_* has a left adjoint $u^* : \text{Mod}(\mathcal{X}) \rightarrow \text{Mod}(U)$ which preserves quasicoherent sheaves. Since $(u : U \rightarrow \mathcal{X})$ is an object of $\text{Lis-Ét}(\mathcal{X})$, it follows that $u^* : \text{Mod}(\mathcal{X}) \rightarrow \text{Mod}(U)$ is exact. In particular, $u_* : \text{Mod}(U) \rightarrow \text{Mod}(\mathcal{X})$ has an exact left adjoint. This implies that it must preserve injective sheaves. It follows that $u_*(\mathcal{H})$ is a quasicoherent sheaf on \mathcal{X} which is injective as a sheaf of $\mathcal{O}_{\mathcal{X}}$ -modules.

Letting $\mathcal{G} = u_*(\mathcal{H})$, we are now left with showing that the two maps in (5.2) are injective. The first map is injective because u is faithfully flat and \mathcal{F} is quasicoherent. The second map is injective because $u_* : \text{QCoh}(U) \rightarrow \text{QCoh}(\mathcal{X})$ is left exact and hence preserves injections.

To show that the functor $D^+(\text{QCoh}(\mathcal{X})) \rightarrow D_{\text{qc}}^+(\mathcal{X})$ is essentially surjective, we can use its full and faithfulness shown above and an induction on the length to first see that $D^b(\text{QCoh}(\mathcal{X})) \xrightarrow{\simeq} D_{\text{qc}}^b(\mathcal{X})$. Since every object of $D_{\text{qc}}^+(\mathcal{X})$ is a colimit of objects in $D_{\text{qc}}^b(\mathcal{X})$ (using good truncations), a limit argument concludes the proof. \square

Lemma 5.3. *Let \mathcal{X} be as in Lemma 5.1 and let $P \in D(\text{QCoh}(\mathcal{X}))$ be a compact object. Then the following hold.*

- (1) *There exists an integer $r \geq 0$ such that $\text{Hom}_{D_{\text{qc}}(\mathcal{X})}(P, N[i]) = 0$ for all $N \in \text{QCoh}(\mathcal{X})$ and $i > r$.*
- (2) *There exists an integer $r \geq 0$ such that the natural map*

$$\tau^{\geq j} \text{RHom}_{D_{\text{qc}}(\mathcal{X})}(P, M) \rightarrow \tau^{\geq j} \text{RHom}_{D_{\text{qc}}(\mathcal{X})}(P, \tau^{\geq j-r} M)$$

is a quasi-isomorphism for all $M \in D_{\text{qc}}(\mathcal{X})$ and integers j .

- (3) *There exists an integer $r \geq 0$ such that the natural map*

$$\tau^{\geq j} \text{RHom}_{D(\text{QCoh}(\mathcal{X}))}(P, M) \rightarrow \tau^{\geq j} \text{RHom}_{D(\text{QCoh}(\mathcal{X}))}(P, \tau^{\geq j-r} M)$$

is a quasi-isomorphism for all $M \in D(\text{QCoh}(\mathcal{X}))$ and integers j .

Proof. It follows from [Lemma 5.1](#) that $\iota_{\mathcal{X}}$ induces an equivalence between the derived categories of perfect complexes of quasicoherent sheaves and perfect complexes of $\mathcal{O}_{\mathcal{X}}$ -modules. Since the compact objects of $D_{\text{qc}}(\mathcal{X})$ are perfect [[Krishna and Ravi 2018](#), Proposition 2.7], it follows that $D(\text{QCoh}(\mathcal{X}))$ and $D_{\text{qc}}(\mathcal{X})$ have equivalent full subcategories of compact objects. The parts (1) and (2) now follow from [[Hall and Rydh 2017](#), Lemma 4.5] and the proof of [[Hall et al. 2014](#), Lemma 2.4] shows that (1) implies (3) for any stack. \square

Lemma 5.4. *Let \mathcal{X} be a Noetherian stack such that $D_{\text{qc}}(\mathcal{X})$ is compactly generated. Then $\iota_{\mathcal{X}} : \text{QCoh}(\mathcal{X}) \rightarrow \text{Mod}(\mathcal{X})$ induces an equivalence of the unbounded derived categories $D(\text{QCoh}(\mathcal{X})) \xrightarrow{\simeq} D_{\text{qc}}(\mathcal{X})$.*

Proof. Let $\Psi : D(\text{QCoh}(\mathcal{X})) \rightarrow D_{\text{qc}}(\mathcal{X})$ denote the derived functor induced by $\iota_{\mathcal{X}}$. We have shown in the proof of [Lemma 5.3](#) that Ψ restricts to an equivalence between the full subcategories of compact objects. Using [[Benson et al. 2011](#), Lemma 4.5], it thus suffices to show that $D(\text{QCoh}(\mathcal{X}))$ is compactly generated.

So let $M \in D(\text{QCoh}(\mathcal{X}))$ be such that $\text{Hom}_{D(\text{QCoh}(\mathcal{X}))}(P, M) = 0$ for every compact object P . We need to show that $M = 0$. Since any compact object of $D(\text{QCoh}(\mathcal{X}))$ is perfect, and Ψ is conservative and induces equivalence of compact objects, it suffices to show that $\text{RHom}(P, M) \xrightarrow{\simeq} \text{RHom}(\Psi(P), \Psi(M))$ for every perfect complex P . Equivalently, we need to show that for every integer j , the map $\tau^{\geq j} \text{RHom}(P, M) \rightarrow \tau^{\geq j} \text{RHom}(\Psi(P), \Psi(M))$ is a quasi-isomorphism. [Lemma 5.3](#) now allows us to assume that $M \in D^+(\text{QCoh}(\mathcal{X}))$. But in this case, the result follows from [Lemma 5.1](#). \square

For a Noetherian stack \mathcal{X} , let $G^{\text{naive}}(\mathcal{X})$ denote the K -theory spectrum of the exact category of coherent $\mathcal{O}_{\mathcal{X}}$ -modules in the sense of Quillen, and let $G(\mathcal{X})$ be the K -theory spectrum of the complicial bi-Waldhausen category of cohomologically bounded pseudocoherent complexes in $\text{Ch}_{\text{qc}}(\mathcal{X})$, in the sense of [[Thomason and Trobaugh 1990](#), §1.5.2]. We have a natural map of spectra $G^{\text{naive}}(\mathcal{X}) \rightarrow G(\mathcal{X})$.

Lemma 5.5. *Let \mathcal{X} be a Noetherian stack such that $D_{\text{qc}}(\mathcal{X})$ is compactly generated. Then the map $G^{\text{naive}}(\mathcal{X}) \rightarrow G(\mathcal{X})$ is a homotopy equivalence.*

Proof. It follows from [Lemma 5.4](#) that $G(\mathcal{X})$ is homotopy equivalent to the K -theory of the Waldhausen category $\text{Ch}_{\text{pc}}(\text{QCoh}(\mathcal{X}))$ of cohomologically bounded pseudocoherent chain complexes of quasicoherent sheaves on \mathcal{X} . Let $\text{Ch}^b(\text{Coh}(\mathcal{X}))$ denote the Waldhausen category of bounded complexes of coherent $\mathcal{O}_{\mathcal{X}}$ -modules.

Using the fact that every quasicoherent sheaf on \mathcal{X} is a filtered colimit of coherent subsheaves [[Laumon and Moret-Bailly 2000](#), Proposition 15.4], we can mimic the proof of [[Thomason and Trobaugh 1990](#), Lemma 3.12] to conclude that the inclusion $\text{Ch}^b(\text{Coh}(\mathcal{X})) \hookrightarrow \text{Ch}_{\text{pc}}(\text{QCoh}(\mathcal{X}))$ induces a homotopy equivalence between the associated K -theory spectra. By induction on the length of complexes in

$\text{Ch}^b(\text{Coh}(\mathcal{X}))$, the map $G^{\text{naive}}(\mathcal{X}) \rightarrow K(\text{Ch}^b(\text{Coh}(\mathcal{X})))$ is also a homotopy equivalence, so we conclude the proof. \square

Lemma 5.6. *Let \mathcal{X} be a Noetherian regular stack. Then the canonical map of spectra $K(\mathcal{X}) \rightarrow G(\mathcal{X})$ is a homotopy equivalence.*

Proof. As for schemes [Thomason and Trobaugh 1990, Theorem 3.21], it suffices to show that every cohomologically bounded pseudocoherent complex E^\bullet on \mathcal{X} is perfect. Let $u : U \rightarrow \mathcal{X}$ be a smooth atlas such that U is affine. Since $(u : U \rightarrow \mathcal{X})$ is an object of $\text{Lis-Ét}(\mathcal{X})$, the functor $u^* : \text{Mod}(\mathcal{X}) \rightarrow \text{Mod}(U)$ is exact and preserves coherent sheaves. It follows that $u^*(E^\bullet)$ is a cohomologically bounded pseudocoherent complex on U . Since U is a regular scheme, we conclude from the proof of [Thomason and Trobaugh 1990, Theorem 3.21] that $u^*(E^\bullet)$ is perfect. But this implies that E^\bullet is perfect on \mathcal{X} . \square

Theorem 5.7. *Let \mathcal{X} be a Noetherian regular stack such that $D_{\text{qc}}(\mathcal{X})$ is compactly generated. Then the following hold.*

- (1) *The canonical maps $K(\mathcal{X}) \rightarrow G(\mathcal{X}) \leftarrow G^{\text{naive}}(\mathcal{X})$ are homotopy equivalences.*
- (2) *For any vector bundle \mathcal{E} on \mathcal{X} and any \mathcal{E} -torsor $\pi : \mathcal{Y} \rightarrow \mathcal{X}$, the pullback map $K(\mathcal{X}) \rightarrow K(\mathcal{Y})$ is a homotopy equivalence.*
- (3) *The canonical morphisms of spectra*

$$K(\mathcal{X}) \rightarrow K^B(\mathcal{X}) \rightarrow KH(\mathcal{X})$$

are homotopy equivalences. In particular, $K_i(\mathcal{X}) = KH_i(\mathcal{X}) = 0$ for $i < 0$.

Proof. Part (1) of the theorem follows directly from Lemmas 5.5 and 5.6. As shown in [Merkurjev 2005, Theorem 2.11], there exists a short exact sequence of vector bundles

$$0 \rightarrow \mathcal{E} \rightarrow \mathcal{W} \xrightarrow{\phi} \mathbb{A}_{\mathcal{X}}^1 \rightarrow 0$$

such that $\mathcal{Y} = \phi^{-1}(1)$. In particular, \mathcal{Y} is the complement of the projective bundle $\mathbb{P}(\mathcal{E})$ in $\mathbb{P}(\mathcal{W})$. It follows from our hypothesis and Lemma 3.6 that $\mathbb{P}(\mathcal{W})$ is a Noetherian regular stack such that $D_{\text{qc}}(\mathbb{P}(\mathcal{W}))$ is compactly generated. The same holds for $\mathbb{P}(\mathcal{E})$ as well. The Quillen localization sequence

$$G^{\text{naive}}(\mathbb{P}(\mathcal{E})) \rightarrow G^{\text{naive}}(\mathbb{P}(\mathcal{W})) \rightarrow G^{\text{naive}}(\mathcal{Y}) \tag{5.8}$$

and the projective bundle formula [Krishna and Ravi 2018, Theorem 3.8] now prove (2).

By (1) and the Quillen localization sequence for $G^{\text{naive}}(-)$ associated to the inclusions $\mathcal{X}[T] \hookrightarrow \mathbb{P}_{\mathcal{X}}^1$ and $\mathcal{X}[T^{\pm 1}] \hookrightarrow \mathcal{X}[T]$, we see that the Bass fundamental theorem holds for \mathcal{X} and moreover that the sequence (3) of Theorem 4.4 is a short exact sequence. This implies that one can define $K^B(\mathcal{X})$ as in Section 4B, and moreover that $K(\mathcal{X}) \xrightarrow{\cong} K^B(\mathcal{X})$. On the other hand, it follows from (2) that

$K^B(\mathcal{X}) \xrightarrow{\cong} K^B(\mathcal{X} \times \Delta^n)$ for every $n \geq 0$, which implies that $K^B(\mathcal{X}) \xrightarrow{\cong} KH(\mathcal{X})$. The last assertion of (3) holds because $K(\mathcal{X})$ has no negative homotopy groups. \square

6. Cdh-descent for homotopy K -theory

We denote by \mathbf{Stk}' the category of stacks \mathcal{X} satisfying one of the following conditions:

- \mathcal{X} has separated diagonal and linearly reductive finite stabilizers.
- \mathcal{X} has affine diagonal and linearly reductive almost multiplicative stabilizers.

By Corollaries 3.16 and 3.17, for every $\mathcal{X} \in \mathbf{Stk}'$ and every closed substack $\mathcal{Z} \subset \mathcal{X}$ with quasicompact open complement, the pair $(\mathcal{X}, \mathcal{Z})$ is perfect. Furthermore, by Theorem 2.10, \mathcal{X} admits a Nisnevich covering by quotient stacks $[U/G]$ where U is affine over an affine scheme S and G is a linearly reductive almost multiplicative group scheme over S .

Note that if $\mathcal{X} \in \mathbf{Stk}'$ and $\mathcal{Y} \rightarrow \mathcal{X}$ is a representable morphism with affine diagonal, then also $\mathcal{Y} \in \mathbf{Stk}'$, since the stabilizers of \mathcal{Y} are subgroups of the stabilizers of \mathcal{X} .

Lemma 6.1. *Let \mathcal{X} be a stack in \mathbf{Stk}' and let $f : \mathcal{Y} \rightarrow \mathcal{X}$ be a torsor under a vector bundle. Then*

$$f^* : KH(\mathcal{X}) \rightarrow KH(\mathcal{Y})$$

is a homotopy equivalence.

Proof. By Theorem 2.10, there exists a Nisnevich covering $[U/G] \rightarrow \mathcal{X}$, where U is affine over an affine scheme S and G is a linearly reductive S -group scheme. By Proposition 2.9 and Corollary 4.10, we are reduced to showing that

$$KH([U/G]) \rightarrow KH([U/G] \times_{\mathcal{X}} \mathcal{Y})$$

is a homotopy equivalence. But since U and S are affine and G is linearly reductive, the vector bundle torsor $[U/G] \times_{\mathcal{X}} \mathcal{Y} \rightarrow [U/G]$ has a section and hence is a vector bundle. The result now follows from Theorem 4.9(6). \square

The following theorem is our cdh-descent result for the homotopy K -theory of stacks.

Theorem 6.2. *Let \mathcal{X} be a stack in \mathbf{Stk}' and let*

$$\begin{array}{ccc} \mathcal{E} & \hookrightarrow & \mathcal{Y} \\ \downarrow & & \downarrow p \\ \mathcal{Z} & \xhookrightarrow{e} & \mathcal{X} \end{array}$$

be a Cartesian square where p is a proper representable morphism, e is a closed immersion, and p induces an isomorphism $\mathcal{Y} \setminus \mathcal{E} \cong \mathcal{X} \setminus \mathcal{Z}$. Then the induced square

of spectra

$$\begin{array}{ccc}
 KH(\mathcal{X}) & \xrightarrow{p^*} & KH(\mathcal{Y}) \\
 e^* \downarrow & & \downarrow \\
 KH(\mathcal{Z}) & \longrightarrow & KH(\mathcal{E})
 \end{array} \tag{6.3}$$

is homotopy Cartesian.

Proof. We proceed in several steps.

Step 1. We prove the result under the assumptions that p is projective, that p and e are of finite presentation, and that $\mathcal{X} = [U/G]$ is quasi-affine over $[S/G]$ for some affine scheme S and some linearly reductive isotrivial almost multiplicative group scheme G over S (note that such a stack belongs to \mathbf{Stk}' and has the resolution property). Since G is finitely presented, we can write U as an inverse limit of quasi-affine G -schemes of finite presentation over S . By [Theorem 4.9\(5\)](#), KH transforms such limits into homotopy colimits. Since homotopy colimits of spectra commute with homotopy pullbacks, we can assume that U is finitely presented over S . We are now in the situation of [\[Hoyois 2016, Theorem 1.3\]](#), and we deduce that (6.3) is a homotopy Cartesian square for the KH -theory defined in [\[Hoyois 2016\]](#) (more precisely, for the presheaf of spectra $KH_{[S/G]}$ defined in [\[Hoyois 2016, §4\]](#)). But the latter agrees with the KH -theory defined in this paper, by [Lemma 6.1](#).

Step 2. We prove the result under the assumption that p is projective and that \mathcal{X} is as in Step 1. Since every quasicoherent sheaf on \mathcal{X} is the union of its finitely generated quasicoherent subsheaves [\[Rydh 2016\]](#), we can write \mathcal{Z} as a filtered intersection of finitely presented closed substacks of \mathcal{X} . By continuity of KH ([Theorem 4.9\(5\)](#)), we can therefore assume that e is finitely presented. In particular, $\mathcal{U} = \mathcal{X} \setminus \mathcal{Z}$ is quasicompact. Since \mathcal{Y} is projective over \mathcal{X} , it is a closed substack of $\mathbb{P}(\mathcal{F})$ for some finitely generated quasicoherent sheaf \mathcal{F} on \mathcal{X} . Since \mathcal{X} has the resolution property and affine stabilizers, we can write $\mathcal{X} = [V/GL_n]$ for some quasi-affine scheme V [\[Gross 2017, Theorem 1.1\]](#). On such stacks, it is known that every quasicoherent sheaf is a filtered colimit of finitely presented quasicoherent sheaves [\[Rydh 2015, Theorem A and Proposition 2.10\(iii\)\]](#). In particular, \mathcal{F} is a quotient of a finitely presented sheaf, so we can assume without loss of generality that \mathcal{F} is finitely presented. We can again write \mathcal{Y} as a filtered intersection of finitely presented closed substacks $\mathcal{Y}_i \subset \mathbb{P}(\mathcal{F})$. By [\[Rydh 2015, Theorem C\(ii\)\]](#), the projection $\mathcal{Y}_i \times_{\mathcal{X}} \mathcal{U} \rightarrow \mathcal{U}$ is a closed immersion for sufficiently large i . But since it has a section, it must be an isomorphism. By continuity of KH , we can therefore assume that p is finitely presented, and we are thus reduced to Step 1.

Step 3. We prove the result assuming only that p is projective. By [Theorem 2.10](#) and the fact that groups of multiplicative type are isotrivial locally in the Nisnevich topology [\[Hoyois 2017, Remark 2.9\]](#), there exists a Nisnevich covering $\{\mathcal{U}_i \rightarrow \mathcal{X}\}$

where each U_i is as in Step 1. By [Proposition 2.9](#), there is a sequence of quasi-compact open substacks $\emptyset = \mathcal{X}_0 \subset \cdots \subset \mathcal{X}_n = \mathcal{X}$ together with Nisnevich squares

$$\begin{array}{ccc} \mathcal{W}_j & \hookrightarrow & \mathcal{V}_j \\ \downarrow & & \downarrow \\ \mathcal{X}_{j-1} & \hookrightarrow & \mathcal{X}_j \end{array}$$

where each \mathcal{V}_j is a quasicompact open substack of $\coprod_i U_i$. In particular, each \mathcal{V}_j and each \mathcal{W}_j is as in Step 1. Since KH satisfies Nisnevich descent ([Corollary 4.10](#)), we deduce from Step 2 by a straightforward induction on j that [\(6.3\)](#) is a homotopy Cartesian square.

Step 4. We prove the result in general. As in Step 2, we can assume that the complement of e is quasicompact. By [Corollary 2.4](#), there exists a projective morphism $\mathcal{Y}' \rightarrow \mathcal{Y}$ which is an isomorphism over the complement of e and such that the composite $\mathcal{Y}' \rightarrow \mathcal{Y} \rightarrow \mathcal{X}$ is projective. Consider the squares

$$\begin{array}{ccccc} KH(\mathcal{X}) & \xrightarrow{p^*} & KH(\mathcal{Y}) & \longrightarrow & KH(\mathcal{Y}') \\ e^* \downarrow & & \downarrow & & \downarrow \\ KH(\mathcal{Z}) & \longrightarrow & KH(\mathcal{E}) & \longrightarrow & KH(\mathcal{E}') \end{array}$$

where $\mathcal{E}' = \mathcal{E} \times_{\mathcal{Y}} \mathcal{Y}'$. The right-hand square and the total square are both homotopy Cartesian by Step 3. Hence, the left-hand square is also homotopy Cartesian, as desired. □

7. The vanishing theorems

Our goal now is to use the cdh-descent for homotopy K -theory to prove the vanishing theorems for negative K -theory. In order to apply cdh-descent, Kerz and Strunk [\[2017\]](#) used the idea of killing classes in the negative K -theory of schemes using Gruson–Raynaud flatification [\[Raynaud and Gruson 1971\]](#). In [Section 7A](#), we prove an analog of this result for stacks. This is done essentially like in the case of schemes, where we replace Gruson–Raynaud flatification with Rydh’s flatification theorem for algebraic stacks ([Theorem 2.2](#)). The vanishing results are proven in [Sections 7B](#) and [7C](#).

7A. Killing by flatification. We need the following two preparatory results about quasicoherent sheaves on stacks.

Lemma 7.1. *Let $f : \mathcal{Y} \rightarrow \mathcal{X}$ be a quasi-affine morphism of stacks. If \mathcal{X} satisfies the resolution property, so does \mathcal{Y} .*

Proof. This is [\[Hall and Rydh 2017, Lemma 7.1\]](#). □

Lemma 7.2. *Let $f : \mathcal{Y} \rightarrow \mathcal{X}$ be a smooth morphism of Noetherian stacks and let \mathcal{F} be a coherent sheaf on \mathcal{Y} which is flat over \mathcal{X} . Then \mathcal{F} has finite tor-dimension over \mathcal{Y} .*

Proof. Since the question is smooth-local on \mathcal{X} and \mathcal{Y} , we can assume that \mathcal{X} and \mathcal{Y} are Noetherian schemes. In this case, the result is [Kerz and Strunk 2017, Lemma 6]. □

Proposition 7.3 (the killing lemma). *Let \mathcal{X} be a reduced Noetherian stack and let $f : \mathcal{Y} \rightarrow \mathcal{X}$ be a smooth morphism of finite type such that \mathcal{Y} is perfect and satisfies the resolution property. Let $n > 0$ be an integer and let $\xi \in K_{-n}(\mathcal{Y})$. Then there exists a sequence of blow-ups $u : \mathcal{X}' \rightarrow \mathcal{X}$ with nowhere dense centers such that for the induced map $u_{\mathcal{Y}} : \mathcal{Y}' := \mathcal{X}' \times_{\mathcal{X}} \mathcal{Y} \rightarrow \mathcal{Y}$, one has $u_{\mathcal{Y}}^*(\xi) = 0$ in $K_{-n}(\mathcal{Y}')$.*

Proof. We repeat the proof of [Kerz and Strunk 2017, Proposition 5] with minor modifications. By the construction of negative K -theory of perfect stacks (see Definition 4.8), there exists a surjection

$$\text{Coker}(K_0(\mathcal{Y} \times \mathbb{A}^n) \rightarrow K_0(\mathcal{Y} \times \mathbb{G}_m^n)) \twoheadrightarrow K_{-n}(\mathcal{Y}), \tag{7.4}$$

natural in \mathcal{Y} . It therefore suffices to prove that, for any $\xi \in K_0(\mathcal{Y} \times \mathbb{G}_m^n)$, there exists a sequence of blow-ups $u : \mathcal{X}' \rightarrow \mathcal{X}$ with nowhere dense centers such that $u_{\mathcal{Y} \times \mathbb{G}_m^n}^*(\xi)$ lies in the image of the restriction map

$$j^* : K_0(\mathcal{Y}' \times \mathbb{A}^n) \rightarrow K_0(\mathcal{Y}' \times \mathbb{G}_m^n). \tag{7.5}$$

Since \mathcal{Y} satisfies the resolution property, it follows from Lemma 7.1 that $\mathcal{Y} \times \mathbb{G}_m^n$ also satisfies the resolution property. In particular, $K_0(\mathcal{Y} \times \mathbb{G}_m^n)$ is generated by classes of vector bundles on $\mathcal{Y} \times \mathbb{G}_m^n$. Since any finite collection of sequences of blow-ups of \mathcal{X} can be refined by a single such sequence, we can assume that ξ is represented by a vector bundle \mathcal{E} on $\mathcal{Y} \times \mathbb{G}_m^n$. We can now extend \mathcal{E} to a coherent sheaf \mathcal{F} on $\mathcal{Y} \times \mathbb{A}^n$ by [Gross 2017, Theorem 1.1; Thomason 1987b, Lemma 1.4].

Choose a commutative square

$$\begin{array}{ccc} Y & \xrightarrow{g} & X \\ p \downarrow & & \downarrow q \\ \mathcal{Y} & \xrightarrow{f} & \mathcal{X} \end{array}$$

where X and Y are algebraic spaces, p and q are smooth surjective maps, and g is smooth of finite type. By generic flatness [Stacks 2005–, Tag 06QR], we can find a dense open subspace $U \subset X$ such that $(q \times \text{id}_{\mathbb{A}^n})^*(\mathcal{F})$ is flat over U under the composite map $Y \times \mathbb{A}^n \rightarrow Y \rightarrow X$. Then U induces a dense open substack $\mathcal{U} \subset \mathcal{X}$ such that \mathcal{F} is flat over \mathcal{U} . We now apply Theorem 2.2 to find a sequence of blow-ups $u : \mathcal{X}' \rightarrow \mathcal{X}$ whose centers are disjoint from \mathcal{U} such that the strict transform $\tilde{\mathcal{F}}$ of \mathcal{F} on $\mathcal{Y}' \times \mathbb{A}^n$ is flat over \mathcal{X}' .

We consider the commutative diagram of Cartesian squares

$$\begin{array}{ccccccc}
 \mathcal{Y}' \times \mathbb{G}_m^n & \xrightarrow{j} & \mathcal{Y}' \times \mathbb{A}^n & \xrightarrow{e'} & \mathcal{Y}' & \longrightarrow & \mathcal{X}' \\
 q \downarrow & & \downarrow w & & \downarrow v & & \downarrow u \\
 \mathcal{Y} \times \mathbb{G}_m^n & \longrightarrow & \mathcal{Y} \times \mathbb{A}^n & \xrightarrow{e} & \mathcal{Y} & \longrightarrow & \mathcal{X}
 \end{array} \tag{7.6}$$

in which the vertical arrows are blow-ups and the horizontal arrows are smooth. We next recall that the strict transform $\tilde{\mathcal{F}}$ is defined by the cokernel of the map $\mathcal{H}_{E \times \mathbb{A}^n}^0(w^*(\mathcal{F})) \hookrightarrow w^*(\mathcal{F})$, where $E \hookrightarrow \mathcal{Y}'$ is the exceptional locus of the blow-up and $\mathcal{H}_{(-)}^0$ is the sheaf of sections with support. Since \mathcal{F} restricts to the vector bundle \mathcal{E} over $\mathcal{Y} \times \mathbb{G}_m^n$, which in turn is smooth over \mathcal{X} , it follows that $j^*(\tilde{\mathcal{F}}) = q^*(\mathcal{E})$ by [Stacks 2005–, Tag 080F].

Lemma 7.2 says that $\tilde{\mathcal{F}}$ has finite tor-dimension over $\mathcal{Y}' \times \mathbb{A}^n$. In particular, it defines a class $[\tilde{\mathcal{F}}] \in K_0(\mathcal{Y}' \times \mathbb{A}^n)$. Moreover, we have $[q^*(\mathcal{E})] = [j^*(\tilde{\mathcal{F}})] = j^*([\tilde{\mathcal{F}}])$. This finishes the proof. \square

7B. Vanishing of negative homotopy K -theory. We now use the techniques of cdh-descent and killing by flatification to prove our main results on the vanishing of negative K -theory of stacks.

Definition 7.7. Let \mathcal{X} be a Noetherian stack.

- (1) The *Krull dimension* $\text{Kr dim}(\mathcal{X}) \in \mathbb{N} \cup \{\pm\infty\}$ is the Krull dimension of the underlying topological space $|\mathcal{X}|$; see [Laumon and Moret-Bailly 2000, Chapter 5] for the definition of $|\mathcal{X}|$.
- (2) The *blow-up dimension* $\text{bl dim}(\mathcal{X}) \in \mathbb{N} \cup \{\pm\infty\}$ is the supremum of the integers $n \geq 0$ such that there exists a sequence $\mathcal{X}_n \rightarrow \mathcal{X}_{n-1} \rightarrow \cdots \rightarrow \mathcal{X}_0 = \mathcal{X}$ of nonempty stacks where each \mathcal{X}_i is a nowhere dense closed substack of an iterated blow-up of \mathcal{X}_{i-1} .
- (3) The *covering dimension* $\text{cov dim}(\mathcal{X}) \in \mathbb{N} \cup \{\pm\infty\}$ is the least dimension of a scheme X admitting a faithfully flat finitely presented morphism $X \rightarrow \mathcal{X}$.

Lemma 7.8. *Let \mathcal{X} be a Noetherian stack. Then*

$$\text{Kr dim}(\mathcal{X}) \leq \text{bl dim}(\mathcal{X}) \leq \text{cov dim}(\mathcal{X}).$$

If \mathcal{X} is a quasi-DM stack, all three are equal to $\text{dim}(\mathcal{X})$.

Proof. The inequality $\text{Kr dim}(\mathcal{X}) \leq \text{bl dim}(\mathcal{X})$ follows directly from the definitions, since $\text{Kr dim}(\mathcal{X})$ is the supremum of a subset of the set of integers described in Definition 7.7(2). For the inequality $\text{bl dim}(\mathcal{X}) \leq \text{cov dim}(\mathcal{X})$, it suffices to prove the following:

- (i) If $\mathcal{Y} \rightarrow \mathcal{X}$ is a blow-up, then $\text{cov dim}(\mathcal{Y}) \leq \text{cov dim}(\mathcal{X})$.
- (ii) If $\mathcal{Z} \subset \mathcal{X}$ is a nowhere dense closed substack, then $\text{cov dim}(\mathcal{Z}) \leq \text{cov dim}(\mathcal{X}) - 1$.

Let $f : X \rightarrow \mathcal{X}$ be an fppf cover, where X is a scheme. Then X is Noetherian and $X \times_{\mathcal{X}} \mathcal{Y} \rightarrow X$ is a blow-up of X . It follows that $\dim(X \times_{\mathcal{X}} \mathcal{Y}) \leq \dim(X)$, whence (i). By [Stacks 2005–, Tag 04XL], the induced map of topological spaces $|f| : |X| \rightarrow |\mathcal{X}|$ is continuous and open. Using [Stacks 2005–, Tag 03HR], we deduce that $X \times_{\mathcal{X}} \mathcal{Z}$ is a nowhere dense closed subscheme of X . It follows that $\dim(X \times_{\mathcal{X}} \mathcal{Z}) \leq \dim(X) - 1$, whence (ii).

For the last statement, we prove more generally that the following hold for every faithfully flat representable quasifinite morphism of Noetherian stacks $f : \mathcal{Y} \rightarrow \mathcal{X}$:

- (i) $\dim(\mathcal{Y}) = \dim(\mathcal{X})$.
- (ii) $\text{Kr dim}(\mathcal{Y}) \leq \text{Kr dim}(\mathcal{X})$.

If \mathcal{X} is quasi-DM, we can take \mathcal{Y} to be a scheme and we deduce $\text{cov dim}(\mathcal{X}) \leq \dim(\mathcal{X})$ and $\dim(\mathcal{X}) \leq \text{Kr dim}(\mathcal{X})$, as desired. To prove (i), by definition of the dimension of a stack [Stacks 2005–, Tag 0AFL], we are immediately reduced to the case where \mathcal{X} is an algebraic space. In this case, the claim follows from [Stacks 2005–, Tags 04NV and 0AFH]. If $Z_0 \subset \dots \subset Z_n$ is a strictly increasing sequence of irreducible closed subsets of $|\mathcal{Y}|$, then $\overline{f(Z_0)} \subset \dots \subset \overline{f(Z_n)}$ is a sequence of irreducible closed subsets of $|\mathcal{X}|$. To check that it is strictly increasing, we may again assume that \mathcal{X} is an algebraic space. If the sequence were not strictly increasing, we would have a nontrivial specialization in a fiber of $|f|$, which is a discrete space [Stacks 2005–, Tag 06RW]. This proves (ii). □

Example 7.9. Let k be a field, let $n \geq 1$, and let \mathcal{X} be the stack quotient of \mathbb{A}_k^n by the standard action of the general linear group GL_n . Then $\text{Kr dim}(\mathcal{X}) = \text{bl dim}(\mathcal{X}) = 1$, $\text{cov dim}(\mathcal{X}) = n$, and $\dim(\mathcal{X}) = n - n^2$. We do not know an example where $\text{Kr dim}(\mathcal{X}) \neq \text{bl dim}(\mathcal{X})$. □

See Section 6 for the definition of the category \mathbf{Stk}' appearing in the next theorem.

Theorem 7.10. *Let \mathcal{X} be a stack in \mathbf{Stk}' satisfying the resolution property. If \mathcal{X} is Noetherian of blow-up dimension d , then $KH_i(\mathcal{X}) = 0$ for $i < -d$.*

Proof. We prove the theorem by induction on d . Since KH is nil-invariant (take $\mathcal{Y} = \emptyset$ in Theorem 6.2), we can assume that \mathcal{X} is reduced. We can write $KH(\mathcal{X}) = \text{hocolim}_n F_n(\mathcal{X})$, where

$$F_n(\mathcal{X}) = \text{hocolim}_{\Delta_{\leq n}^{\text{op}}} K^B(\mathcal{X} \times \Delta^\bullet).$$

It suffices to show inductively on n that the canonical map $\pi_i F_n(\mathcal{X}) \rightarrow KH_i(\mathcal{X})$ is zero for all $i < -d$. This is trivial if $n < 0$, so assume $n \geq 0$.

Let $C_{i,n}(\mathcal{X})$ denote the cokernel of $\pi_i F_{n-1}(\mathcal{X}) \rightarrow \pi_i F_n(\mathcal{X})$. Since the cofiber of the map $F_{n-1}(\mathcal{X}) \rightarrow F_n(\mathcal{X})$ is canonically a direct summand of $\Sigma^n K^B(\mathcal{X} \times \Delta^n)$ (see

for instance [Lurie 2017, Remark 1.2.4.7]), we may identify $C_{i,n}(\mathcal{X})$ with a subgroup of $K_{i-n}(\mathcal{X} \times \Delta^n)$. By the induction hypothesis, the map $\pi_i F_n(\mathcal{X}) \rightarrow KH_i(\mathcal{X})$ factors through $C_{i,n}(\mathcal{X})$:

$$\begin{array}{ccccccc}
 \pi_i F_{n-1}(\mathcal{X}) & \longrightarrow & \pi_i F_n(\mathcal{X}) & \twoheadrightarrow & C_{i,n}(\mathcal{X}) & \hookrightarrow & K_{i-n}(\mathcal{X} \times \Delta^n) \\
 & & \downarrow & & \swarrow & & \\
 & & KH_i(\mathcal{X}) & & & & \\
 & \searrow 0 & & \swarrow \phi_{i,n} & & & \\
 & & & & & &
 \end{array} \tag{7.11}$$

Hence, it suffices to show that $\phi_{i,n} : C_{i,n}(\mathcal{X}) \rightarrow KH_i(\mathcal{X})$ is zero. Let ξ be in $C_{i,n}(\mathcal{X}) \subset K_{i-n}(\mathcal{X} \times \Delta^n)$. By Lemma 7.1, $\mathcal{X} \times \Delta^n$ satisfies the resolution property. By Proposition 7.3, there exists a sequence of blow-ups $u : \mathcal{X}' \rightarrow \mathcal{X}$ with nowhere dense centers such that $u^*(\xi) = 0$ in $C_{i,n}(\mathcal{X}') \subset K_{i-n}(\mathcal{X}' \times \Delta^n)$ (note that $i - n < 0$). Let $\mathcal{Z} \subset \mathcal{X}$ be a nowhere dense closed substack of \mathcal{X} such that u is an isomorphism over the complement of \mathcal{Z} . By Theorem 6.2, we have a long exact sequence

$$\cdots \rightarrow KH_{i+1}(u^{-1}(\mathcal{Z})) \rightarrow KH_i(\mathcal{X}) \rightarrow KH_i(\mathcal{X}') \oplus KH_i(\mathcal{Z}) \rightarrow \cdots .$$

Note that both \mathcal{Z} and $u^{-1}(\mathcal{Z})$ have blow-up dimension strictly less than d . By the induction hypothesis, $KH_{i+1}(u^{-1}(\mathcal{Z}))$ and $KH_i(\mathcal{Z})$ are both zero, so the map $u^* : KH_i(\mathcal{X}) \rightarrow KH_i(\mathcal{X}')$ is injective. Since $\phi_{i,n}$ is natural in \mathcal{X} , we have $u^* \phi_{i,n}(\xi) = \phi_{i,n} u^*(\xi) = 0$, and we conclude that $\phi_{i,n}(\xi) = 0$. This finishes the proof. \square

Our next goal is to remove the resolution property assumption from Theorem 7.10. We are able to do so under the additional assumption that \mathcal{X} has finite inertia. If X is a Noetherian algebraic space, we denote by $\mathbf{\acute{E}t}_X$ the category of algebraic spaces over X that are étale, separated, and of finite type. The following lemma is a Nisnevich variant of [Kerz and Strunk 2017, Proposition 3].

Lemma 7.12. *Let X be a Noetherian algebraic space, let \mathcal{F} be a Nisnevich sheaf of abelian groups on $\mathbf{\acute{E}t}_X$, and let r be an integer. Suppose that $\mathcal{F}(\mathcal{O}_{Y,y}^h) = 0$ for every point $y \in Y \in \mathbf{\acute{E}t}_X$ with $\dim \overline{\{y\}} > r$. Then $H_{\text{Nis}}^i(X, \mathcal{F}) = 0$ for all $i > r$.*

Proof. Let $s \in \mathcal{F}(X)$ be a section, and let $i : Z \hookrightarrow X$ be a closed immersion such that the support of s is $|Z|$, i.e., $|Z|$ is the closed subset of points $x \in X$ such that s is nonzero in every open neighborhood of x . We claim that

$$\dim(Z) \leq r.$$

Otherwise, let $y \in Z$ be a generic point such that $\dim \overline{\{y\}} > r$. Then $i^*(\mathcal{F})(\mathcal{O}_{Z,y}) \cong \mathcal{F}(\mathcal{O}_{X,y}^h) = 0$, so the section $i^*(s)$ of $i^*(\mathcal{F})$ vanishes on an open neighborhood Y of y in Z . This means that s itself vanishes on an étale neighborhood of Y . Since it also vanishes on $X \setminus Z$ and \mathcal{F} is a Nisnevich sheaf, it follows that s vanishes on the open $(X \setminus Z) \cup Y$, which is a contradiction.

Let S be a finite set of local sections of \mathcal{F} , and let $\mathcal{F}_S \subset \mathcal{F}$ be the subsheaf generated by S . Let $i_S : X_S \hookrightarrow X$ be a closed immersion such that $|X_S|$ is the union of the closures of the images of the supports of the sections in S , and let $j_S : X \setminus X_S \hookrightarrow X$ be the complementary open immersion. Then $j_S^*(\mathcal{F}_S) = 0$ since every $s \in S$ is zero over $X \setminus X_S$. Using the gluing short exact sequence

$$0 \rightarrow (j_S)_! j_S^*(\mathcal{F}_S) \rightarrow \mathcal{F}_S \rightarrow (i_S)_* i_S^*(\mathcal{F}_S) \rightarrow 0,$$

we deduce $\mathcal{F}_S \cong (i_S)_* i_S^*(\mathcal{F}_S)$. If we now write \mathcal{F} as a filtered colimit $\mathcal{F} \cong \text{colim}_S b\mathcal{F}_S$, we obtain

$$H_{\text{Nis}}^i(X, \mathcal{F}) \cong \text{colim}_S H_{\text{Nis}}^i(X, \mathcal{F}_S) \cong \text{colim}_S H_{\text{Nis}}^i(X_S, i_S^*(\mathcal{F}_S)).$$

The last isomorphism holds because $(i_S)_*$ is an exact functor on Nisnevich sheaves of abelian groups. By our preliminary result, $\dim(X_S) \leq r$. Since X_S is a Noetherian algebraic space, its Nisnevich cohomological dimension is bounded by its Krull dimension [Lurie 2018, Theorem 3.7.7.1]. We therefore have $H_{\text{Nis}}^i(X_S, i_S^*(\mathcal{F}_S)) = 0$ for $i > r$, whence $H_{\text{Nis}}^i(X, \mathcal{F}) = 0$ for $i > r$. □

Lemma 7.13. *Let X be a Noetherian algebraic space of finite Krull dimension, let \mathcal{F} be a presheaf of spectra on $\mathbf{\acute{E}t}_X$ satisfying Nisnevich descent, and let n be an integer. Suppose that, for every point $y \in Y \in \mathbf{\acute{E}t}_X$, $\mathcal{F}(\mathcal{O}_{Y,y}^h)$ is $(n + \dim \overline{\{y\}})$ -connective. Then the spectrum $\mathcal{F}(X)$ is n -connective.*

Proof. We can assume without loss of generality that $n = 0$. Let $\pi_* \mathcal{F}$ denote the Nisnevich sheaves of homotopy groups of \mathcal{F} . Since X is a Noetherian algebraic space of finite Krull dimension, its Nisnevich topos has finite homotopy dimension [Lurie 2018, Theorem 3.7.7.1], so that the descent spectral sequence

$$H_{\text{Nis}}^p(X, \pi_q \mathcal{F}) \Rightarrow \pi_{q-p} \mathcal{F}(X)$$

is strongly convergent. Applying Lemma 7.12 to $\pi_q \mathcal{F}$, we deduce that

$$H_{\text{Nis}}^p(X, \pi_q \mathcal{F}) = 0$$

for all $p > q$, and we conclude using the above spectral sequence. □

Theorem 7.14. *Let \mathcal{X} be a stack in \mathbf{Stk}' with finite inertia, e.g., a separated quasi-DM stack with linearly reductive stabilizers. Assume that \mathcal{X} is Noetherian of dimension d . Then $KH_i(\mathcal{X}) = 0$ for $i < -d$.*

Proof. Let X be the coarse moduli space of \mathcal{X} . Note that X is a Noetherian algebraic space of dimension d . Let \mathcal{F} be the presheaf of spectra on $\mathbf{\acute{E}t}_X$ defined by

$$\mathcal{F}(Y) = KH(\mathcal{X} \times_X Y).$$

By [Corollary 4.10](#), \mathcal{F} satisfies Nisnevich descent on $\hat{\mathbf{E}}\mathbf{t}_X$. For $y \in Y \in \hat{\mathbf{E}}\mathbf{t}_X$, let $\mathcal{X}_y^h = \mathcal{X} \times_X \mathrm{Spec}(\mathcal{O}_{Y,y}^h)$. By continuity of KH ([Theorem 4.9\(5\)](#)), we have

$$\mathcal{F}(\mathcal{O}_{Y,y}^h) \simeq KH(\mathcal{X}_y^h).$$

By [Theorem 2.10](#), the stack \mathcal{X}_y^h has the form $[U/G]$, where U is affine and G is a finite group scheme over $\mathrm{Spec}(\mathcal{O}_{Y,y}^h)$. In particular, \mathcal{X}_y^h belongs to \mathbf{Stk}' and satisfies the resolution property. Moreover, the dimension of \mathcal{X}_y^h is at most $d - \dim \overline{\{y\}}$, and it equals its blow-up dimension by [Lemma 7.8](#). It follows from [Theorem 7.10](#) that $\mathcal{F}(\mathcal{O}_{Y,y}^h)$ is $(-d + \dim \overline{\{y\})}$ -connective. By [Lemma 7.13](#), we deduce that $\mathcal{F}(X)$ is $(-d)$ -connective, i.e., that $KH_i(\mathcal{X}) = 0$ for $i < -d$. \square

7C. Vanishing of negative K -theory with coefficients. Let \mathcal{X} be a perfect stack and let $n \in \mathbb{Z}$. Recall from [[Krishna and Ravi 2018](#), §5C] that the algebraic K -theory of \mathcal{X} with coefficients is defined by

$$\begin{aligned} K^B(\mathcal{X})[1/n] &:= \mathrm{hocolim}(K^B(\mathcal{X}) \xrightarrow{n} K^B(\mathcal{X}) \xrightarrow{n} \cdots), \\ K^B(\mathcal{X}, \mathbb{Z}/n) &:= K^B(\mathcal{X}) \wedge \mathbb{S}/n, \end{aligned}$$

where \mathbb{S}/n is the mod- n Moore spectrum, and similarly for KH .

Proposition 7.15. *Let \mathcal{X} be a perfect stack.*

- (1) *If n is nilpotent on \mathcal{X} , then the canonical map $K^B(\mathcal{X})[1/n] \rightarrow KH(\mathcal{X})[1/n]$ is a homotopy equivalence.*
- (2) *If n is invertible on \mathcal{X} , then the canonical map $K^B(\mathcal{X}, \mathbb{Z}/n) \rightarrow KH(\mathcal{X}, \mathbb{Z}/n)$ is a homotopy equivalence.*

Proof. We have shown in the proof of [Proposition 3.8\(1\)](#) that there is a weak equivalence of dg-categories

$$\mathrm{D}_{\mathrm{perf}}(\mathcal{X} \times \mathbb{A}^1) \simeq \mathrm{D}_{\mathrm{perf}}(\mathcal{X}) \otimes \mathrm{D}_{\mathrm{perf}}(\mathbb{A}^1).$$

Given this, the proposition follows immediately from [[Tabuada 2017](#), Theorem 1.2]. \square

Theorem 7.16. *Let \mathcal{X} be a stack in \mathbf{Stk}' satisfying the resolution property or having finite inertia. Assume that \mathcal{X} is Noetherian of blow-up dimension d . Then the following hold.*

- (1) *If n is nilpotent on \mathcal{X} , then $K_i(\mathcal{X})[1/n] = 0$ for any $i < -d$.*
- (2) *If n is invertible on \mathcal{X} , then $K_i(\mathcal{X}, \mathbb{Z}/n) = 0$ for any $i < -d$.*

Proof. This follows from [Theorems 7.10](#) and [7.14](#) and [Proposition 7.15](#). \square

Acknowledgments

We are very grateful to David Rydh for several fruitful discussions about his recent work, which allowed us to significantly enhance the scope of this paper.

The bulk of this work was completed during the authors' stay at the Mittag-Leffler Institute as part of the research program "Algebraic-geometric and homotopical methods", and we would like to thank the institute and the organizers, Eric Friedlander, Lars Hesselholt, and Paul Arne Østvær, for this opportunity.

References

- [Abramovich et al. 2008] D. Abramovich, M. Olsson, and A. Vistoli, "Tame stacks in positive characteristic", *Ann. Inst. Fourier (Grenoble)* **58**:4 (2008), 1057–1091. [MR](#) [Zbl](#)
- [Alper 2013] J. Alper, "Good moduli spaces for Artin stacks", *Ann. Inst. Fourier (Grenoble)* **63**:6 (2013), 2349–2402. [MR](#) [Zbl](#)
- [Alper et al. \geq 2019] J. Alper, J. Hall, and D. Rydh, "The étale local structure of algebraic stacks", in preparation.
- [Asok et al. 2017] A. Asok, M. Hoyois, and M. Wendt, "Affine representability results in \mathbb{A}^1 -homotopy theory, I: Vector bundles", *Duke Math. J.* **166**:10 (2017), 1923–1953. [MR](#) [Zbl](#)
- [Bass 1968] H. Bass, *Algebraic K-theory*, W. A. Benjamin, New York, 1968. [MR](#) [Zbl](#)
- [Ben-Zvi et al. 2010] D. Ben-Zvi, J. Francis, and D. Nadler, "Integral transforms and Drinfeld centers in derived algebraic geometry", *J. Amer. Math. Soc.* **23**:4 (2010), 909–966. [MR](#) [Zbl](#)
- [Benson et al. 2011] D. J. Benson, S. B. Iyengar, and H. Krause, "Stratifying modular representations of finite groups", *Ann. of Math. (2)* **174**:3 (2011), 1643–1684. [MR](#) [Zbl](#)
- [Blumberg et al. 2013] A. J. Blumberg, D. Gepner, and G. Tabuada, "A universal characterization of higher algebraic K-theory", *Geom. Topol.* **17**:2 (2013), 733–838. [MR](#) [Zbl](#)
- [Cisinski 2013] D.-C. Cisinski, "Descente par éclatements en K-théorie invariante par homotopie", *Ann. of Math. (2)* **177**:2 (2013), 425–448. [MR](#) [Zbl](#)
- [Cisinski and Tabuada 2011] D.-C. Cisinski and G. Tabuada, "Non-connective K-theory via universal invariants", *Compos. Math.* **147**:4 (2011), 1281–1320. [MR](#) [Zbl](#)
- [Cortiñas et al. 2008] G. Cortiñas, C. Haesemeyer, M. Schlichting, and C. Weibel, "Cyclic homology, cdh-cohomology and negative K-theory", *Ann. of Math. (2)* **167**:2 (2008), 549–573. [MR](#) [Zbl](#)
- [Gross 2017] P. Gross, "Tensor generators on schemes and stacks", *Algebr. Geom.* **4**:4 (2017), 501–522. [MR](#) [Zbl](#)
- [Haesemeyer 2004] C. Haesemeyer, "Descent properties of homotopy K-theory", *Duke Math. J.* **125**:3 (2004), 589–620. [MR](#) [Zbl](#)
- [Hall and Rydh 2015] J. Hall and D. Rydh, "Algebraic groups and compact generation of their derived categories of representations", *Indiana Univ. Math. J.* **64**:6 (2015), 1903–1923. [MR](#) [Zbl](#)
- [Hall and Rydh 2017] J. Hall and D. Rydh, "Perfect complexes on algebraic stacks", *Compos. Math.* **153**:11 (2017), 2318–2367. [MR](#) [Zbl](#)
- [Hall and Rydh 2018] J. Hall and D. Rydh, "Addendum to "Étale dévissage, descent and pushouts of stacks" [*J. Algebra* 331 (1) (2011) 194–223]", *J. Algebra* **498** (2018), 398–412. [MR](#) [Zbl](#)
- [Hall et al. 2014] J. Hall, A. Neeman, and D. Rydh, "One positive and two negative results for derived categories of algebraic stacks", preprint, 2014. [arXiv](#)

- [Hoyois 2016] M. Hoyois, “Cdh descent in equivariant homotopy K -theory”, preprint, 2016. [arXiv](#)
- [Hoyois 2017] M. Hoyois, “The six operations in equivariant motivic homotopy theory”, *Adv. Math.* **305** (2017), 197–279. [MR](#) [Zbl](#)
- [Kelly 2014] S. Kelly, “Vanishing of negative K -theory in positive characteristic”, *Compos. Math.* **150**:8 (2014), 1425–1434. [MR](#) [Zbl](#)
- [Kerz and Strunk 2017] M. Kerz and F. Strunk, “On the vanishing of negative homotopy K -theory”, *J. Pure Appl. Algebra* **221**:7 (2017), 1641–1644. [MR](#) [Zbl](#)
- [Kerz et al. 2018] M. Kerz, F. Strunk, and G. Tamme, “Algebraic K -theory and descent for blow-ups”, *Invent. Math.* **211**:2 (2018), 523–577. [MR](#) [Zbl](#)
- [Krause 2010] H. Krause, “Localization theory for triangulated categories”, pp. 161–235 in *Triangulated categories*, edited by T. Holm et al., London Math. Soc. Lecture Note Ser. **375**, Cambridge Univ. Press, 2010. [MR](#) [Zbl](#)
- [Krishna and Østvær 2012] A. Krishna and P. A. Østvær, “Nisnevich descent for K -theory of Deligne–Mumford stacks”, *J. K-Theory* **9**:2 (2012), 291–331. [MR](#) [Zbl](#)
- [Krishna and Ravi 2018] A. Krishna and C. Ravi, “Algebraic K -theory of quotient stacks”, *Ann. K-Theory* **3**:2 (2018), 207–233. [MR](#) [Zbl](#)
- [Laumon and Moret-Bailly 2000] G. Laumon and L. Moret-Bailly, *Champs algébriques*, *Ergebnisse der Math.* (3) **39**, Springer, 2000. [MR](#) [Zbl](#)
- [Lurie 2009] J. Lurie, *Higher topos theory*, *Annals of Mathematics Studies* **170**, Princeton Univ. Press, 2009. [MR](#) [Zbl](#)
- [Lurie 2017] J. Lurie, “Higher algebra”, preprint, 2017, available at <http://www.math.harvard.edu/~lurie/papers/HA.pdf>.
- [Lurie 2018] J. Lurie, “Spectral algebraic geometry”, book in progress, 2018, available at <http://www.math.harvard.edu/~lurie/papers/SAG-rootfile.pdf>.
- [Merkurjev 2005] A. S. Merkurjev, “Equivariant K -theory”, pp. 925–954 in *Handbook of K -theory*, vol. 2, edited by E. M. Friedlander and D. R. Grayson, Springer, 2005. [MR](#) [Zbl](#)
- [Morel and Voevodsky 1999] F. Morel and V. Voevodsky, “ \mathbb{A}^1 -homotopy theory of schemes”, *Inst. Hautes Études Sci. Publ. Math.* **90** (1999), 45–143. [MR](#) [Zbl](#)
- [Neeman 1996] A. Neeman, “The Grothendieck duality theorem via Bousfield’s techniques and Brown representability”, *J. Amer. Math. Soc.* **9**:1 (1996), 205–236. [MR](#) [Zbl](#)
- [Raynaud and Gruson 1971] M. Raynaud and L. Gruson, “Critères de platitude et de projectivité: techniques de “platification” d’un module”, *Invent. Math.* **13** (1971), 1–89. [MR](#) [Zbl](#)
- [Rydh 2015] D. Rydh, “Noetherian approximation of algebraic spaces and stacks”, *J. Algebra* **422** (2015), 105–147. [MR](#) [Zbl](#)
- [Rydh 2016] D. Rydh, “Approximation of sheaves on algebraic stacks”, *Int. Math. Res. Not.* **2016**:3 (2016), 717–737. [MR](#) [Zbl](#)
- [Rydh \geq 2019] D. Rydh, “Equivariant flatification, étalification and compactification”, in preparation.
- [Schlichting 2006] M. Schlichting, “Negative K -theory of derived categories”, *Math. Z.* **253**:1 (2006), 97–134. [MR](#) [Zbl](#)
- [SGA 3_I 1970] M. Demazure and A. Grothendieck, *Schémas en groupes, I: Propriétés générales des schémas en groupes* (Séminaire de Géométrie Algébrique du Bois Marie 1962–64), *Lecture Notes in Mathematics* **151**, Springer, Berlin, 1970. [MR](#) [Zbl](#)

- [SGA 6 1971] A. Grothendieck, P. Berthelot, and L. Illusie, *Théorie des intersections et théorème de Riemann–Roch* (Séminaire de Géométrie Algébrique du Bois Marie 1966–1967), Lecture Notes in Mathematics **225**, Springer, Berlin, 1971. [MR](#) [Zbl](#)
- [Stacks 2005–] P. Belmans, A. J. de Jong, et al., “The Stacks project”, electronic reference, 2005–, available at <http://stacks.math.columbia.edu>.
- [Tabuada 2017] G. Tabuada, “ \mathbb{A}^1 -homotopy invariance of algebraic K -theory with coefficients and du Val singularities”, *Ann. K-Theory* **2**:1 (2017), 1–25. [MR](#) [Zbl](#)
- [Thomason 1987a] R. W. Thomason, “Algebraic K -theory of group scheme actions”, pp. 539–563 in *Algebraic topology and algebraic K-theory* (Princeton, NJ, 1983), edited by W. Browder, Ann. of Math. Stud. **113**, Princeton Univ. Press, 1987. [MR](#) [Zbl](#)
- [Thomason 1987b] R. W. Thomason, “Equivariant resolution, linearization, and Hilbert’s fourteenth problem over arbitrary base schemes”, *Adv. in Math.* **65**:1 (1987), 16–34. [MR](#) [Zbl](#)
- [Thomason and Trobaugh 1990] R. W. Thomason and T. Trobaugh, “Higher algebraic K -theory of schemes and of derived categories”, pp. 247–435 in *The Grothendieck Festschrift*, vol. III, edited by P. Cartier et al., Progr. Math. **88**, Birkhäuser, Boston, 1990. [MR](#) [Zbl](#)
- [Voevodsky 2010] V. Voevodsky, “Homotopy theory of simplicial sheaves in completely decomposable topologies”, *J. Pure Appl. Algebra* **214**:8 (2010), 1384–1398. [MR](#) [Zbl](#)
- [Weibel 1989] C. A. Weibel, “Homotopy algebraic K -theory”, pp. 461–488 in *Algebraic K-theory and algebraic number theory* (Honolulu, HI, 1987), edited by M. R. Stein and R. K. Dennis, Contemp. Math. **83**, Amer. Math. Soc., Providence, RI, 1989. [MR](#) [Zbl](#)
- [Weibel 2001] C. Weibel, “The negative K -theory of normal surfaces”, *Duke Math. J.* **108**:1 (2001), 1–35. [MR](#) [Zbl](#)

Received 3 May 2018. Accepted 29 Jan 2019.

MARC HOYOIS: hoyois@usc.edu

Department of Mathematics, University of Southern California, Los Angeles, CA, United States

AMALENDU KRISHNA: amal@math.tifr.res.in

School of Mathematics, Tata Institute of Fundamental Research, Mumbai, India

ANNALS OF K-THEORY

msp.org/akt

EDITORIAL BOARD

Joseph Ayoub	ETH Zürich, Switzerland joseph.ayoub@math.uzh.ch
Paul Balmer	University of California, Los Angeles, USA balmer@math.ucla.edu
Guillermo Cortiñas	Universidad de Buenos Aires and CONICET, Argentina gcorti@dm.uba.ar
Hélène Esnault	Freie Universität Berlin, Germany liveesnault@math.fu-berlin.de
Eric Friedlander	University of Southern California, USA ericmf@usc.edu
Max Karoubi	Institut de Mathématiques de Jussieu – Paris Rive Gauche, France max.karoubi@imj-prg.fr
Moritz Kerz	Universität Regensburg, Germany moritz.kerz@mathematik.uni-regensburg.de
Huaxin Lin	University of Oregon, USA livehlin@uoregon.edu
Alexander Merkurjev	University of California, Los Angeles, USA merkurev@math.ucla.edu
Birgit Richter	Universität Hamburg, Germany birgit.richter@uni-hamburg.de
Jonathan Rosenberg	(Managing Editor) University of Maryland, USA jmr@math.umd.edu
Marco Schlichting	University of Warwick, UK schlichting@warwick.ac.uk
Charles Weibel	(Managing Editor) Rutgers University, USA weibel@math.rutgers.edu
Guoliang Yu	Texas A&M University, USA guoliangyu@math.tamu.edu

PRODUCTION

Silvio Levy (Scientific Editor)
production@msp.org

Annals of K-Theory is a journal of the [K-Theory Foundation](http://ktheoryfoundation.org) (ktheoryfoundation.org). The K-Theory Foundation acknowledges the precious support of [Foundation Compositio Mathematica](http://foundationcompositio.com), whose help has been instrumental in the launch of the Annals of K-Theory.

See inside back cover or msp.org/akt for submission instructions.

The subscription price for 2019 is US \$490/year for the electronic version, and \$550/year (+\$25, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Annals of K-Theory (ISSN 2379-1681 electronic, 2379-1683 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

AKT peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing
<http://msp.org/>

© 2019 Mathematical Sciences Publishers

ANNALS OF K-THEORY

2019

vol. 4

no. 3

Motivic analogues of MO and MSO DONDI ELLIS	345
The IA-congruence kernel of high rank free metabelian groups DAVID EL-CHAI BEN-EZRA	383
Vanishing theorems for the negative K -theory of stacks MARC HOYOIS and AMALENDU KRISHNA	439
Higher genera for proper actions of Lie groups PAOLO PIAZZA and HESSEL B. POSTHUMA	473
Periodic cyclic homology and derived de Rham cohomology BENJAMIN ANTIEAU	505
Linkage of Pfister forms over $\mathbb{C}(x_1, \dots, x_n)$ ADAM CHAPMAN and JEAN-PIERRE TIGNOL	521