

ANNALS OF K-THEORY

vol. 5 no. 1 2020

**Twisted iterated algebraic K -theory
and topological T-duality for sphere bundles**

John A. Lind, Hisham Sati and Craig Westerland



A JOURNAL OF THE K-THEORY FOUNDATION

Twisted iterated algebraic K -theory and topological T-duality for sphere bundles

John A. Lind, Hisham Sati and Craig Westerland

We introduce a periodic form of the iterated algebraic K -theory of ku , the (connective) complex K -theory spectrum, as well as a natural twisting of this cohomology theory by higher gerbes. Furthermore, we prove a form of topological T-duality for sphere bundles oriented with respect to this theory.

1. Iterated algebraic K -theory and its twistings by n -gerbes	5
2. Chern characters, $\mathfrak{A}$ -orientations, and Umkehr maps	12
3. T-duality for sphere bundles in theories twisted by n -gerbes	17
4. Classifying spaces for T-dual pairs	23
5. Higher categories, n -vector spaces, iterated algebraic K -theory	32
Acknowledgements	38
References	39

Let ku be the connective complex K -theory spectrum. The underlying infinite loop space $\Omega^\infty ku = \mathbb{Z} \times BU$ of ku classifies virtual complex vector bundles. The cohomology theory associated to the algebraic K -theory spectrum $K(ku)$, the subject of much recent research in homotopy theory [Ausoni and Rognes 2002], has a geometric interpretation as a Grothendieck group of 2-vector bundles [Baas et al. 2004; 2011; Lind 2016]. A 2-vector bundle is a bundle whose fiber is a 2-vector space, which is a categorified form of a vector space introduced by Kapranov and Voevodsky [1994]. Forming equivalence classes of 2-vector bundles over X leads to a bimonoidal category $2\text{Vect}(X)$. By [Baas et al. 2011], the Grothendieck group completion of $2\text{Vect}(X)$ is represented by the infinite loop space $\Omega^\infty K(ku)$ underlying the algebraic K -theory of ku .

Applying the functor $K(\cdot)$ again, one is naturally led to imagine that the iterated algebraic K -theory spectrum

$$\alpha_n := K^{(n-1)}(ku) = \underbrace{K(K(\cdots K(ku)\cdots))}_{n-1}$$

MSC2010: primary 19D99, 55N20, 55R15, 55R65; secondary 18D05, 19L50, 55P43.

Keywords: T-duality, iterated algebraic K -theory, twisted K -theory, n -gerbes.

has an interpretation in terms of categorified bundles native to n -category theory. It is expected that algebraic K -theory in many cases increases chromatic complexity by one, i.e., that it produces a constant “red-shift” by one chromatic layer in stable homotopy theory [Ausoni and Rognes 2002]. In this paper, we study a Bott-periodic form $\mathfrak{A}_n := K^{(n-1)}(ku)[\beta_n^{-1}]$ of iterated algebraic K -theory. While our results do not provide direct evidence either for or against the Ausoni–Rognes red-shift conjectures [2008], our interest in the relationship between the geometric content of iterated algebraic K -theory and chromatic homotopy theory is a primary motivation for the study of T-duality in $K^{(n-1)}(ku)[\beta_n^{-1}]$ -theory.

Much as line bundles are the fundamental building blocks of vector bundles, and hence play an essential role in the K -theory of vector bundles, $(n-1)$ -gerbes are the simplest forms of n -vector bundles. For a general definition of n -gerbes as n -truncated and n -connected objects, see [Lurie 2009a, §7.2.2]. For example, when $n = 2$, a 1-gerbe (also known as a gerbe with band $U(1)$) gives rise to a rank-one 2-vector bundle. This fact is witnessed at the level of classifying spaces by a map

$$K(\mathbb{Z}, 3) \rightarrow BGL_1(ku) \rightarrow \Omega^\infty K(ku).$$

This is the 2-categorical analog of the map $\mathbb{C}P^\infty \rightarrow \mathbb{Z} \times BU$ representing the inclusion of line bundles into the Grothendieck group of vector bundles. The adjoint map $\Sigma^\infty \mathbb{C}P_+^\infty \rightarrow ku$ is a map of E_∞ ring spectra. We study here a family of analogous E_∞ ring maps

$$\Sigma^\infty K(\mathbb{Z}, n+1)_+ \rightarrow \mathfrak{a}_n := K^{(n-1)}(ku) \quad (0.1)$$

that we think of as representing the inclusion of $(n-1)$ -gerbes into the Grothendieck group of n -vector bundles.

These maps are adjoint (under the adjunction described in [Ando et al. 2014b; May 2009]) to maps of spectra $\Sigma^{n+1}H\mathbb{Z} \rightarrow gl_1(\mathfrak{a}_n)$ or, equivalently, maps of E_∞ spaces $K(\mathbb{Z}, n+1) \rightarrow GL_1(\mathfrak{a}_n)$. Delooping once, we obtain E_∞ -twistings of the cohomology theory $\mathfrak{a}_n$ by $(n+2)$ -dimensional cohomology¹ classes: each class $H \in H^{n+2}(X; \mathbb{Z})$ in the cohomology of a topological space X gives rise to twisted cohomology groups $\mathfrak{a}_n^*(X; H)$. When $n = 1$, this returns the usual notion of connective complex K -theory twisted by a gerbe (or rather, a representative of its Dixmier–Douady class in H^3). The new twists that we study have the same degree as the twists of Morava K -theory and E -theory studied previously [Sati and Westerland 2015]; we hope that further understanding of the red-shift conjecture will relate the results of that paper and this one.

¹A caveat is in order: the twisted cohomology group $\mathfrak{a}_n^*(X; H)$ depends upon the representative map $H : X \rightarrow K(\mathbb{Z}, n+2)$. Just as is the case for twisted K -theory, homotopic maps (i.e., cohomologous classes) yield isomorphic twisted cohomology groups; however, the isomorphism is not canonical.

Notice that the element $\beta \in \pi_2 ku$, given as the composite $S^2 \rightarrow \mathbb{C}P^\infty \rightarrow \mathbb{Z} \times BU$ of the fundamental class of $\mathbb{C}P^\infty = K(\mathbb{Z}, 2)$ and its inclusion as $BU(1) \times \{1\}$, is the Bott class. Localizing ku at this class yields periodic complex K -theory: $KU := ku[\beta^{-1}]$. Our first goal is an analog of this construction for $\mathfrak{a}_n$:

Theorem 0.2. *The composite of the fundamental class of $K(\mathbb{Z}, n+1)$ with the map in (0.1) yields an essential class $\beta_n \in \pi_{n+1} \mathfrak{a}_n$. Further, when n is odd, β_n is nonnilpotent, and none of its powers are torsion.*

This is proven as [Corollary 1.7](#) and [Theorem 1.10](#), below. For odd n , we will notate the localized spectrum $\mathfrak{a}_n[\beta_n^{-1}]$, i.e., the Bott-inverted iterated algebraic K -theory, by $\mathfrak{A}_n$. [Theorem 0.2](#) suggests that this is a nontrivial object of study. However, it is a consequence of a theorem of Arthan [\[1983\]](#) that $\mathfrak{A}_n$ is a *rational* spectrum whenever $n > 1$. We may regard it as a higher categorical analog of periodic topological K -theory, or perhaps an étale form of iterated K -theory.

The twisting of $\mathfrak{a}_n$ by cohomology classes in degree $n+2$ extends in a natural way to a twisting of $\mathfrak{A}_n$ via the composite

$$\Sigma^{n+1} H\mathbb{Z} \rightarrow \mathrm{gl}_1 \mathfrak{a}_n \rightarrow \mathrm{gl}_1 \mathfrak{A}_n,$$

where the second map is induced by the E_∞ ring map given by localization $\mathfrak{a}_n \rightarrow \mathfrak{A}_n$. Our second main purpose in this paper is to prove an analog of the topological T-duality isomorphism of [\[Bouwknegt et al. 2004; Bunke and Schick 2005\]](#) in this context.

Given a base space X , Bouwknegt, Evslin, and Mathai give a criterion for two sets of data (E, H) and $(\widehat{E}, \widehat{H})$ to be *T-dual* [\[Bouwknegt et al. 2004\]](#). Here E and $\widehat{E}$ are principal S^1 -bundles over X , and H and $\widehat{H}$ are cocycles representing cohomology classes in $H^3(E)$ and $H^3(\widehat{E})$, respectively. The criterion in [\[Bouwknegt et al. 2004\]](#) is given in terms of relations among various characteristic classes. Bunke and Schick [\[2005\]](#) reinterpret this criterion as amounting to the existence of a Thom class on a certain S^3 -bundle over X into which both E and $\widehat{E}$ embed. When (E, H) and $(\widehat{E}, \widehat{H})$ are a T-dual pair, there is an isomorphism of twisted K -theory groups $K^*(E; H) \cong K^{*-1}(\widehat{E}, \widehat{H})$ called the T-duality isomorphism.

Let E and $\widehat{E}$ be fiber bundles over X with fiber the q -sphere S^q , and let H and $\widehat{H}$ be classes in $H^{2q+1}(E)$ and $H^{2q+1}(\widehat{E})$, respectively. In [Definition 3.2](#), we will give a Thom class criterion for (E, H) and $(\widehat{E}, \widehat{H})$ to be T-dual in the higher (dimensional and categorical) context, similar to Bunke and Schick's. We then prove the following extension of the T-duality isomorphism in this setting.

Theorem 0.3. *Let $n = 2q - 1$, and assume that (E, H) and $(\widehat{E}, \widehat{H})$ are a T-dual pair. Let $\tau_E : E \rightarrow B \mathrm{GL}_1 \mathfrak{A}_n$ denote the orientation twisting determined by the*

vertical tangent bundle of $E \rightarrow X$. Then there is an isomorphism of twisted cohomology groups

$$T := \hat{p}_! \circ \Lambda \circ p^* : \mathfrak{A}_n^*(E; \tau_E \otimes H) \rightarrow \mathfrak{A}_n^*(\widehat{E}; \widehat{H})$$

given in terms of a Fourier–Mukai push-pull construction on the correspondence space $E \times_X \widehat{E}$. Given an $\mathfrak{A}$ -orientation of the fiber bundle $E \rightarrow X$, the T-duality isomorphism takes the form

$$T : \mathfrak{A}_n^{*+q}(E; H) \rightarrow \mathfrak{A}_n^*(\widehat{E}; \widehat{H}).$$

This is proven as [Theorem 3.3](#). In fact, we prove the result for a larger class of cohomology theories R than just $\mathfrak{A}_n$, namely those which may be twisted by n -gerbes in such a fashion that the analog of the class β_n is invertible in $\pi_* R$. One consequence of this invertibility assumption (again via [\[Arthan 1983\]](#)) is that R is a rational spectrum when $n > 1$. We will show in [Theorem 3.9](#) that this is no accident: every cohomology theory R for which the T-duality map T of the previous theorem is an isomorphism must be rational.

Additionally, we study criteria to ensure the orientability requirement of the previous result, and analyze the homotopy type of the classifying spaces for T-dual pairs, much as in [\[Bunke and Schick 2005\]](#) in the case $q = n = 1$.

For $q = 3$, T-duality for S^3 -bundles in rational cohomology, and in twisted K -theory under some conditions on the cohomology and dimensions of the underlying manifolds, was considered recently in [\[Bouwknegt et al. 2015a\]](#). The authors discuss twistings of topological K -theory by a 7-dimensional class. It is known that $BGL_1(ku)$ splits as $B\mathbb{Z}/2 \times K(\mathbb{Z}, 3) \times BB\mathrm{SU}_\otimes$. Furthermore, $BB\mathrm{SU}_\otimes$ has a 7-dimensional homotopy group isomorphic to $\mathbb{Z}$; this is visible via the k -invariant $BB\mathrm{SU}_\otimes(6) \rightarrow K(\mathbb{Z}, 7)$ in a Postnikov tower for $BB\mathrm{SU}_\otimes$. An essential map in the other direction — which is necessary for there to be a twist of K -theory by H^7 — does not exist [\[Antieau et al. 2014\]](#); hence, that degree cannot be “isolated” in general, as highlighted in [\[Sati 2009\]](#) and clarified further in [\[Sati and Westerland 2015\]](#). The conditions imposed by [\[Bouwknegt et al. 2015a\]](#) evade the obstructions for dimension reasons and allow one to isolate such a twist in special cases. If one rationalizes, as done for the most part in [\[Bouwknegt et al. 2015a\]](#), one gets periodic cohomology, which does have a degree-seven twist. We recover this result and analogs for all sphere bundles as a consequence of the variant of [Theorem 0.3](#) for periodic rational cohomology. Indeed, it is a consequence of [Theorem 3.9](#) that such a T-duality result *must* be rational outside of the case $q = 1$.

Our general result also provides a recipe for determining whether there exists a T-dual for (E, H) , where E is an S^q -bundle with structure group $G \rightarrow \mathrm{Homeo}(S^q)$ equipped with an n -gerbe H , and whether the T-dual is unique. This is determined by analyzing the connectivity of the Euler class map $BG \rightarrow K(\mathbb{Z}, q + 1)$. For the

case $q = 1$, this recovers some of the results of Baraglia [2014] and Mathai and Rosenberg [2014] for existence and uniqueness of T-dual bundles for nonprincipal circle bundles.

The paper is organized as follows. In Section 1A, we describe the multiplicative behavior of the algebraic K -theory functor and consider generalities about twisted cohomology theories. In Section 1B, we construct and analyze the twisting of iterated algebraic K -theory $K^{(n-1)}(ku)$ by higher gerbes. We prolong some computations of the homotopy of $K(ku)$ due to [Ausoni et al. 2008] to the higher setting in Section 1C. Then in Section 1D we consider a higher analog of Snaith's construction of KU , which in turn admits the universal periodic twist.

In order to obtain concrete expressions and with an eye for applications to T-duality, we study Chern characters and orientations in Section 2. We introduce two Chern characters associated to the new twisted theories, one in Section 2A and one in Section 2B, and then we describe orientations with respect the new theories in Section 2C.

The general setup and the proof of the main T-duality theorem and its converse are given in Section 3. Section 4 is concerned with the question of the existence and the uniqueness of T-dual pairs, which we analyze homotopy-theoretically in terms of classifying spaces. In particular, in Section 4A we introduce a space $P_n(G)$ which classifies the possible T-dual pairs. Then in Section 4B we classify, via a space $R_n(G)$, bundles and Thom classes that can arise in our context of T-duality. The two constructions are related in Section 4C by constructing a forgetful map from $P_n(G)$ to $R_n(G)$, where we also show how our results reproduce earlier results on T-duality.

The final section, Section 5, is a speculative account of the relationship between the iterated algebraic K -theory considered earlier in the paper and a proposed model for the algebraic K -theory of n -vector spaces defined in terms of enriched higher category theory.

1. Iterated algebraic K -theory and its twistings by n -gerbes

We recall here the multiplicative behavior of the algebraic K -theory functor and describe the twisting of iterated algebraic K -theory by higher gerbes.

1A. Generalities on twistings of algebraic K -theory. Given an E_∞ ring spectrum A , the algebraic K -theory spectrum $K(A)$ is once again an E_∞ ring spectrum. There is a natural map $BGL_1 A \rightarrow \Omega^\infty K(A)$ coming from the inclusion of A -lines into all cell A -modules. It is not the case that this is an infinite loop map: the multiplication on $BGL_1 A$ is by tensor products of A -lines, whereas that in $\Omega^\infty K(A)$ is from the sum of modules. However, this map has image in $GL_1 K(A)$

(since A -lines are invertible A -modules), and in fact the induced map $B \mathrm{GL}_1 A \rightarrow \mathrm{GL}_1 K(A)$ is an infinite loop map.

We explain the above in detail. Let A be a connective commutative ring spectrum and let $K(A)$ denote the connective algebraic K -theory spectrum of A . The underlying infinite loop space of the spectrum $K(A)$ receives a map

$$B \mathrm{GL}_1 A \rightarrow \coprod_{n \geq 0} B \mathrm{GL}_n A \rightarrow \Omega^\infty K(A) \quad (1.1)$$

from the classifying space of the space of units $\mathrm{GL}_1 A$ via the classifying space of finite-rank free A -module spectra. This is the analog in algebraic K -theory of the map $\mathbb{C}P^\infty \rightarrow \mathbb{Z} \times B \mathrm{U}$ into topological K -theory classifying the homomorphism from the Picard group into the Grothendieck group of vector bundles. As in the classical case, the source and target both inherit E_∞ space structures from the multiplication on A .

Proposition 1.2. *The map (1.1) has image in $\mathrm{SL}_1 K(A)$; the result is a map of E_∞ spaces and so lifts to a map of spectra*

$$\mu : \Sigma \mathrm{gl}_1 A \rightarrow \mathrm{gl}_1 K(A).$$

Proof. We employ the ∞ -categorical model for algebraic K -theory developed by Gepner, Groth, and Nikolaus [Gepner et al. 2015] using the language of quasi-categories. Let Perf_A denote the stable ∞ -category of compact A -module spectra. Let Line_A denote the ∞ -category of rank-one A -module spectra. Both of these ∞ -categories admit symmetric monoidal structures under the smash product $\wedge_A$ of A -modules. Writing ι_0 for the groupoid core functor which takes a quasicategory to its maximal sub-Kan complex, the inclusion of Line_A into Perf_A induces a symmetric monoidal functor $i : \iota_0 \mathrm{Line}_A \rightarrow \iota_0 \mathrm{Perf}_A$ of symmetric monoidal ∞ -groupoids. The ∞ -category Perf_A admits an additional symmetric monoidal structure under the coproduct of A -modules and the two monoidal structures combine to give the Kan complex $\iota_0 \mathrm{Perf}_A$ the structure of an E_∞ ring space [Gepner et al. 2015, Corollary 8.11]. The algebraic K -theory $K(A)$ is the connective E_∞ ring spectrum for which $\Omega^\infty K(A)$ is the group completion of $\iota_0 \mathrm{Perf}_A$ as an E_∞ ring space [Gepner et al. 2015, Definition 8.3 and Theorem 8.6]. The resulting composite

$$\iota_0 \mathrm{Line}_A \rightarrow \iota_0 \mathrm{Perf}_A \rightarrow \Omega^\infty K(A)$$

is a map of E_∞ spaces, where we use the multiplicative structure on $\iota_0 \mathrm{Perf}_A$ and $\Omega^\infty K(A)$. This map is a model for the map (1.1), as can be verified along the lines of [Ando et al. 2014a, Proposition 2.9], so the map in question is E_∞ .

The source of the map is a connected Kan complex and its image lies in the component of the unit object $A \in \iota_0 \mathrm{Perf}_A$, hence in the component $\mathrm{SL}_1 K(A)$ of the

unit in the E_∞ ring space $\Omega^\infty K(A)$. In particular, the map factors as an E_∞ map through the space $\mathrm{GL}_1 K(A)$ of units, which deloops to the map of spectra μ . $\square$

We now recall how to construct twisted forms of the cohomology theory represented by the ring spectrum A , following [Ando et al. 2010; 2018; 2014b]. Given a space X and a map $\omega : X \rightarrow B \mathrm{GL}_1 A$, let $E \rightarrow X$ be the $\mathrm{GL}_1(A)$ -fibration pulled back from $B \mathrm{GL}_1 A$ via ω . Define an A -module spectrum

$$\mathbb{A}(X; \omega) := A \wedge_{\Sigma^\infty_{\mathrm{GL}_1 A_+}} \Sigma^\infty E_+.$$

This is the Thom spectrum associated to the parametrized spectrum of A -lines over X classified by ω .

Definition 1.3. The ω -twisted A -homology and A -cohomology groups of X are defined by

$$A_q(X; \omega) := \pi_q \mathbb{A}(X; \omega) \quad \text{and} \quad A^q(X; \omega) := \pi_{-q} F_A(\mathbb{A}(X; -\omega), A).$$

Notice that in the definition of twisted cohomology we use the inverse twist $-\omega$ under the monoidal structure induced by the smash product of invertible A -modules. Sometimes the opposite convention is used, but we prefer this choice because it ensures that twisted cohomology agrees with homotopy classes of sections of the parametrized spectrum associated to ω [Lind 2016], and conforms with grading conventions when ω carries a topological dimension, for example when it arises from a classifying map for a virtual vector bundle via the J-homomorphism.

Suppose that h is a spectrum, and that $\tau : h \rightarrow \Sigma \mathrm{gl}_1 A$ is a map of spectra. We regard the map τ as an E_∞ twisting of A , because the infinite loop map $\Omega^\infty \tau : \Omega^\infty h \rightarrow B \mathrm{GL}_1 A$ allows us to twist the A -cohomology of a space X by elements $[H] \in h^0(X)$. To do so, we represent the class $[H]$ as a map $H : X \rightarrow \Omega^\infty h$ and define the H -twisted cohomology of X to be the twisted A -homology and A -cohomology groups associated to the twist $\Omega^\infty \tau \circ H : X \rightarrow B \mathrm{GL}_1 A$, i.e., we make the abbreviation $A^q(X; H) = A^q(X; \Omega^\infty \tau \circ H)$. Different representatives for the same cohomology class give isomorphic twisted cohomology groups, so the twisted cohomology theory associated to H only depends on the underlying cohomology class $[H]$, but only up to noncanonical isomorphism.

We can use the map $\mu : \Sigma \mathrm{gl}_1 A \rightarrow \mathrm{gl}_1 K(A)$ considered in Proposition 1.2 to construct E_∞ twists of the algebraic K -theory spectrum $K(A)$ from (shifts of) E_∞ twists of A , and we would like to know whether or not the twists constructed in this way are essential, i.e., homotopically nontrivial.

Proposition 1.4. For every E_∞ twisting $\tau : h \rightarrow \Sigma \mathrm{gl}_1 A$, the composite

$$\Sigma(\mu \circ \tau) : \Sigma h \rightarrow \Sigma \mathrm{gl}_1 K(A)$$

is an E_∞ twisting of $K(A)$. Further, if the rationalization of τ is essential, so too is the rationalization of $\Sigma(\mu \circ \tau)$.

Proof. Only the latter statement needs to be proved. One can see this via the rational determinant $\det_{\mathbb{Q}}$, defined by Ausoni and Rognes [2012, Proposition 5.4]. This is a map of spaces

$$\det_{\mathbb{Q}} : \Omega^\infty K(A) \rightarrow (B \mathrm{GL}_1 A)_{\mathbb{Q}}$$

whose composite with $\Omega^\infty \mu$ is the rationalization map of $B \mathrm{GL}_1 A$. Therefore, $\Omega^\infty \Sigma(\mu \circ \tau)$ factorizes the rationalization of $\Omega^\infty \tau$, which yields the result. $\square$

Remark 1.5. There is not an integral determinant map lifting $\det_{\mathbb{Q}}$. In fact, the obstruction to the existence of a continuous map $\Omega^\infty K(ku) \rightarrow B \mathrm{GL}_1(ku)$ with determinant-like properties leads to the notion of oriented 2-vector bundles, and gives rise to an oriented version of K -theory of 2-vector bundles with a lift of the natural map from $K(\mathbb{Z}, 3)$, called the determinant gerbe map [Kragh 2013].

1B. Twisting the iterated algebraic K -theory of ku . Our main example arises via connective complex topological K -theory, ku . There is a well known E_∞ twisting of ku by 3-dimensional cohomology classes; the map

$$\tau : \Sigma^3 H\mathbb{Z} \rightarrow \Sigma \mathrm{gl}_1 ku$$

is the delooping of the map of E_∞ -spaces $\mathbb{C}P^\infty \rightarrow \mathrm{GL}_1 ku$ that regards a complex line as an invertible $\mathbb{C}$ -module.

Definition 1.6. Write $\mathfrak{a}_n$ for the iterated algebraic K -theory spectrum $K^{(n-1)}(ku)$. Let

$$\tau_n : \Sigma^{n+2} H\mathbb{Z} \rightarrow \Sigma \mathrm{gl}_1 \mathfrak{a}_n$$

be the E_∞ twisting of $\mathfrak{a}_n$ obtained by an $(n-1)$ -fold iteration of the procedure in Proposition 1.4 applied to $\tau = \tau_1$.

Since the original map τ was rationally essential in homotopy, so too are the maps τ_n :

Corollary 1.7. *The composite of τ_n and the fundamental class ι of $\Sigma^{n+2} H\mathbb{Z}$ defines a nontorsion element $\tau_n \circ \iota \in \pi_{n+2} \Sigma \mathrm{gl}_1 \mathfrak{a}_n$.*

Definition 1.8. We write $\Omega^{\infty+1} \tau_n \circ \iota : S^{n+1} \rightarrow \mathrm{GL}_1 \mathfrak{a}_n$ for the map of spaces representing the image of the class $\tau_n \circ \iota$ under the canonical isomorphism $\pi_{n+2} \Sigma \mathrm{gl}_1 \mathfrak{a}_n \cong \pi_{n+1} \mathrm{gl}_1 \mathfrak{a}_n$. The map $\Omega^{\infty+1} \tau_n$ carries S^{n+1} into the component of $\mathrm{GL}_1 \mathfrak{a}_n$ corresponding to $1 \in \pi_0 \mathfrak{a}_n$. Subtracting 1 gives a based map into $\Omega^\infty \mathfrak{a}_n$, and thus a class $\beta_n \in \pi_{n+1} \mathfrak{a}_n$. In other words, we define $\beta_n := [\Omega \tau_n \circ \iota] - 1$, where $[\Omega \tau_n \circ \iota]$ denotes the image of $\Omega \tau_n \circ \iota$ under the isomorphism $\pi_{n+1} \mathrm{gl}_1 \mathfrak{a}_n \cong \pi_{n+1} \mathfrak{a}_n$ induced by the

inclusion of components $\mathrm{GL}_1 \mathfrak{a}_n \subset \Omega^\infty \mathfrak{a}_n$. Equivalently, β_n is represented by the composite map of spectra

$$\beta_n : S^{n+1} \xrightarrow{\iota} \Sigma^\infty K(\mathbb{Z}, n+1)_+ \xrightarrow{t_n} \mathfrak{a}_n, \quad (1.9)$$

where ι is the fundamental class and t_n is adjoint to $\Omega\tau_n$ in the adjunction

$$[\Sigma^\infty K(\mathbb{Z}, n+1)_+, \mathfrak{a}_n]_{E_\infty \text{ring}} \cong [\Sigma^{n+1} H\mathbb{Z}, \mathrm{gl}_1 \mathfrak{a}_n]_{\mathrm{Sp}}.$$

Note that $\beta_1 \in \pi_2 \mathfrak{a}_1 = \pi_2 ku$ is the usual Bott class which we invert to obtain KU .

Theorem 1.10. *If n is even, $2\beta_n^2 = 0$. However, if n is odd, then β_n is not nilpotent and all powers β_n^m are nontorsion, for any positive integer m .*

Proof. The first claim is simply that β_n is an odd-dimensional element of a graded-commutative ring. For the second, consider the composite map (1.9) and note that the Hurewicz image of $\iota : S^{n+1} \rightarrow \Sigma^\infty K(\mathbb{Z}, n+1)_+$ is essential. In fact, whenever n is odd, the homology ring $H_*(K(\mathbb{Z}, n+1); \mathbb{Q})$ is the rational polynomial ring on the Hurewicz image of ι . In particular, all powers of ι are rationally essential.

We recall from [Segal 1973] that the natural map of ring spectra $\Sigma^\infty \mathbb{C}P_+^\infty \rightarrow ku$ adjoint to τ is a rational equivalence. Therefore, the iterated Dennis trace map, which is a transformation from algebraic K -theory to topological Hochschild homology THH [Bökstedt et al. 1993; Dundas et al. 2013], followed by rationalization, may be written as

$$\begin{aligned} \mathfrak{a}_n &= K^{(n-1)}(ku) \xrightarrow{\mathrm{tr}^{n-1}} THH^{(n-1)}(ku) \\ &\rightarrow THH^{(n-1)}(ku_{\mathbb{Q}}) \xleftarrow{\sim} THH^{(n-1)}((\Sigma^\infty \mathbb{C}P_+^\infty)_{\mathbb{Q}}). \end{aligned}$$

However, for a loop space G , one has $THH(\Sigma^\infty G_+) \simeq \Sigma^\infty LBG_+$. When G is an m -fold loop space, BG (and hence LBG) is an $(m-1)$ -fold loop space. We iteratively observe that

$$THH^{(m)}(\Sigma^\infty G_+) \simeq \Sigma^\infty (LB)^m(G)_+ \simeq \Sigma^\infty \mathrm{Map}(T^m, B^m G)_+$$

if G is connected. In the second equivalence, we employ the fact that for connected G , $LBG \simeq BLG$. The space of constant functions $T^m \rightarrow B^m G$ is homeomorphic to $B^m G$, and so $THH^{(m)}(\Sigma^\infty G_+)$ contains a copy of $\Sigma^\infty B^m G_+$ as a wedge summand.

In the case $G = \mathbb{C}P^\infty = K(\mathbb{Z}, 2)$, it follows that $\Sigma^\infty K(\mathbb{Z}, n+1)_+$ is a summand of $THH^{(n-1)}(\Sigma^\infty \mathbb{C}P_+^\infty)$. Since rationalization is a smashing localization,

$$THH(A_{\mathbb{Q}}) \simeq (THH(A))_{\mathbb{Q}}$$

for ring spectra A . Thus, the target of the iterated trace map above splits off a wedge factor of $(\Sigma^\infty K(\mathbb{Z}, n+1)_+)_{\mathbb{Q}}$.

We claim that the composite of t_n with the iterated trace map and the projection to this wedge summand is the map from $\Sigma^\infty K(\mathbb{Z}, n+1)_+$ to its rationalization; then we may conclude that t_n is an injection on rational homotopy. To see the claim, note that if G is an infinite loop space, there is a homotopy commutative diagram

$$\begin{array}{ccccc}
 \Sigma^\infty BG_+ & \xrightarrow{\quad} & \Sigma^\infty BGL_1(\Sigma^\infty G_+)_+ & \xrightarrow{(1.1)} & K(\Sigma^\infty G_+) \xrightarrow{\text{tr}} THH(\Sigma^\infty G_+) \\
 & & & & \simeq \Sigma^\infty LBG_+ \\
 & \searrow \text{id} & & & \downarrow \text{ev} \\
 & & & & \Sigma^\infty BG_+
 \end{array}$$

yielding Waldhausen's splitting [1979] of $Q(BG_+)$ off of $A(BG) = \Omega^\infty K(\Sigma^\infty G_+)$. Then the claim follows by iteration (taking $G = K(\mathbb{Z}, m)$ for $m = 1, \dots, n$) and rationalization (to accommodate the rational Segal equivalence). $\square$

Definition 1.11. When n is odd, define $\mathfrak{A}_n := \mathfrak{a}_n[\beta_n^{-1}]$.

One could of course make this construction for n even, but the nilpotence of β_n away from the prime 2 in that case will force the resulting spectrum to have 2-torsion homotopy. In fact, these spectra are contractible when n is even; this follows from Theorem 1.14, below.

The E_∞ twisting τ_n of $\mathfrak{a}_n$ induces an E_∞ twisting $K(\mathbb{Z}, n+2) \rightarrow BGL_1 \mathfrak{A}_n$ by composition with the localization map. By the phrase “ n -gerbe” we mean a generic term for any geometric structure classified by the Eilenberg–MacLane space $K(\mathbb{Z}, n+2)$ (for a more general discussion, see [Lurie 2009a, Example 7.2.2.2]). Thus, a complex line bundle is a 0-gerbe, and a gerbe with band $U(1)$ is a 1-gerbe. We say that the E_∞ twisting of $\mathfrak{a}_n$ and $\mathfrak{A}_n$ constructed in this section are *twistings by n -gerbes*, since an n -gerbe over X gives rise to a map $H : X \rightarrow K(\mathbb{Z}, n+2)$ by which we can twist these theories.

1C. Some relations in the homotopy of $\mathfrak{a}_n$. Although the following observations will not be used in our work with the periodic spectrum $\mathfrak{A}_n$, they provide the first known results about the homotopy type of iterated algebraic K -theory of ku . In [Ausoni et al. 2008], it is shown that for $n = 2$ there is a class $\zeta \in \pi_3 K(ku)$ with the property that

$$\beta_2 = 2\zeta - \nu,$$

where ν is the image of the quaternionic Hopf fibration $\nu \in \pi_3(S^0)$ under the unit map to $K(ku) = \mathfrak{a}_2$. One may iteratively prolong this equation to one in $\pi_{n+1} \mathfrak{a}_n$ in a natural fashion, as we now explain. Following the recipe given in Proposition 1.2,

we have a homomorphism²

$$\pi_{n+1} \mathfrak{a}_n \cong \pi_{n+1} \mathfrak{gl}_1 \mathfrak{a}_n \rightarrow \pi_{n+1} \Sigma^{-1} \mathfrak{gl}_1 K(\mathfrak{a}_n) \cong \pi_{n+2} \mathfrak{a}_{n+1}$$

for each $n > 0$. By construction, this map carries β_n to β_{n+1} ; it also allows us to define classes Z_n and N_n in $\pi_{n+1} \mathfrak{a}_n$ as the iterated images of ζ and ν . Since this is a group homomorphism, the relation $\beta_2 = 2\zeta - \nu$ persists:

Proposition 1.12. *The class β_n is given in terms of the iteratively defined classes Z_n and N_n by*

$$\beta_n = 2Z_n - N_n.$$

Furthermore, since $24\nu = 0$, we also have $24N_n = 0$. Additionally, ν is nilpotent. Since the map above is not a ring homomorphism (in fact, it behaves in some sense more like a derivation), we cannot conclude the same for N_n . However, if N_n is actually nilpotent, then a sufficiently large power of β_n is 2-divisible (at least after inverting 3).

1D. Higher Snaith spectra. When $n = 1$, [Definition 1.11](#) tautologically yields periodic complex K -theory: $\mathfrak{A}_1 = KU$. In this case, there is an alternative construction of KU due to Snaith [\[1981\]](#). The map $\Sigma^\infty K(\mathbb{Z}, 2)_+[\iota^{-1}] \rightarrow KU$ induced by the localization of the map $t_1 : \Sigma^\infty K(\mathbb{Z}, 2)_+ \rightarrow ku$ studied in the proof of [Theorem 1.10](#) is an equivalence of spectra. In this section, we consider generalizations of this construction using the domain of the map $t_n : \Sigma^\infty K(\mathbb{Z}, n+1)_+ \rightarrow \mathfrak{a}_n$ and discuss their relationship with the twistings from the previous section.

Guided by Snaith's theorem, we make the following definition.

Definition 1.13. For n odd, define the *Arthan spectrum* $\mathfrak{S}_n$ to be the localization

$$\mathfrak{S}_n := \Sigma^\infty K(\mathbb{Z}, n+1)_+[\iota^{-1}]$$

of the suspension spectrum of $K(\mathbb{Z}, n+1)$ at its fundamental class.

The E_∞ twisting map $t_n : \Sigma^\infty K(\mathbb{Z}, n+1)_+ \rightarrow \mathfrak{a}_n$ carries ι to β_n , and so descends to a map between the periodic spectra $T_n : \mathfrak{S}_n \rightarrow \mathfrak{A}_n$. Snaith's theorem indicates that this is an equivalence when $n = 1$. When $n > 1$, these spectra were studied by Arthan [\[1983\]](#), where they were found to display remarkably different behavior:

Theorem 1.14 (Arthan). *If n is even, $\mathfrak{S}_n$ is contractible. If n is odd and greater than 1, $\mathfrak{S}_n$ is $(n+1)$ -periodic rational cohomology:*

$$\mathfrak{S}_n \cong H\mathbb{Q}[\iota^{\pm 1}].$$

²Here we identify $\pi_k R = \pi_k \Omega_0^\infty R$ with $\pi_k \mathfrak{gl}_1 R = \pi_k \mathrm{GL}_1 R$ for $k > 0$ via the shift in components by 1.

Let R be an A_∞ ring spectrum which receives an A_∞ ring map $\phi : \mathfrak{S}_n \rightarrow R$. The case $R = \mathfrak{A}_n$ is our main example. From ϕ , we obtain a natural composite of maps of spaces

$$K(\mathbb{Z}, n+2) \rightarrow BGL_1(\Sigma^\infty K(\mathbb{Z}, n+1)_+) \rightarrow BGL_1(\mathfrak{S}_n) \rightarrow BGL_1(R),$$

and so, for a topological space X , we may form the twisted cohomology $R^*(X; H)$ associated to a class $[H] \in H^{n+2}(X)$. Furthermore, since ι is invertible in $\mathfrak{S}_n$, the element $\phi(\iota)$ must be invertible in R .

Definition 1.15. We will call the data of the A_∞ map $\phi : \mathfrak{S}_n \rightarrow R$ a *periodic twisting of R by n -gerbes*. An equivalent (if less colorful) term is: R is an $\mathfrak{S}_n$ -*algebra spectrum*.

The spectrum $R = \mathfrak{S}_n$ obviously admits the universal periodic twisting by n -gerbes. [Theorem 1.10](#) ensures that $\mathfrak{A}_n$ is another nontrivial example. An immediate consequence of being an algebra spectrum over a rational algebra is the following:

Proposition 1.16. *If R admits a periodic twisting by n -gerbes for $n > 1$, then it is rational.*

2. Chern characters, $\mathfrak{A}$ -orientations, and Umkehr maps

Many approaches to T-duality involve cohomological expressions. In order to deduce T-duality isomorphisms in ordinary cohomology, we investigate Chern characters and orientations associated with the spectra $\mathfrak{a}_n$ and $\mathfrak{A}_n$ introduced in [Section 1B](#). These are, of course, also interesting in their own right.

2A. The Chern–Dold character. The Chern character in complex topological K -theory of a manifold X is a ring homomorphism from $KU^0(X)$ to even rational cohomology $H^{\text{even}}(X; \mathbb{Q})$ and from $KU^1(X)$ to odd rational cohomology $H^{\text{odd}}(X; \mathbb{Q})$. This periodicity can be encoded by the homomorphism of $\mathbb{Z}$ -graded rings

$$KU^*(X) \rightarrow H^*(X; \mathbb{Q}[u^{\pm 1}]),$$

where the latter is rational cohomology with coefficients in $KU^*(\text{pt}) \otimes \mathbb{Q} \cong \mathbb{Q}[u^{\pm 1}]$, and u^{-1} is the Bott generator. The Chern character may be regarded as a map of spectra

$$\text{ch} : KU \rightarrow H\mathbb{Q}[u^{\pm 1}],$$

where $H\mathbb{Q}[u^{\pm 1}]$ is 2-periodic rational cohomology. The target is a priori formal power series over $\mathbb{Q}$, but becomes polynomials when evaluating on finite-dimensional manifolds.

The above Chern character has a generalization to any generalized cohomology theory. The standard Chern–Dold character (see, e.g., [\[Adams 1972, Chapter 14\]](#))

is based upon the Eilenberg–MacLane spectrum for the rationalization of the coefficients of our theory. Let $\mathfrak{a}_n^*(\text{pt}) = \pi_*(\mathfrak{a}_n) = R^* = \sum_j R^j$. Unfortunately, we have little understanding of an explicit formula for these coefficients. Further, the results of [Ausoni 2010] suggest that R^* is most likely a very complicated ring even for $n = 2$. Nonetheless, we may make the following construction.

Definition 2.1. The *Chern–Dold character for the theory $\mathfrak{a}_n$* is the map of cohomology theories

$$\text{ch}_{\mathfrak{a}_n} : \mathfrak{a}_n^* \rightarrow H^*(\cdot; R^* \otimes \mathbb{Q})$$

induced by the rationalization map

$$\text{ch}_{\mathfrak{a}_n} : \mathfrak{a}_n \rightarrow (\mathfrak{a}_n)_{\mathbb{Q}} \simeq H(R_* \otimes \mathbb{Q}),$$

where we identify the rationalization of $\mathfrak{a}_n$ with the generalized Eilenberg–MacLane spectrum associated to the graded ring $R_* \otimes \mathbb{Q}$.

The following properties of the above character are evident from the definition and follow similarly to those of the standard Chern–Dold character [Buhštaber 1970; Adams 1972].

Lemma 2.2 (properties of the Chern–Dold character). (1) *Over a point, $\text{ch}_{\mathfrak{a}_n}$ is the canonical homomorphism $R^* \rightarrow R^* \otimes \mathbb{Q}$.*

(2) *For a finite complex X the homomorphism*

$$\text{ch}_{\mathfrak{a}_n} \otimes \mathbb{Q} : \mathfrak{a}_n^*(X) \otimes \mathbb{Q} \rightarrow H^*(X; R^* \otimes \mathbb{Q})$$

is an isomorphism.

(3) *$\text{ch}_{\mathfrak{a}_n}$ is a ring homomorphism.*

2B. A (twisted) higher Chern character via THH. Although we do not have a good understanding of the ring R^* or its rationalization, there is a stand-in for the Chern–Dold character with target a recognizable generalized Eilenberg–MacLane spectrum.

Let n be odd, and consider the composite $\text{ch}_n^{\geq 0}$:

$$\begin{array}{ccc} \mathfrak{a}_n = K^{(n-1)}(ku) & \xrightarrow{\text{tr}^{n-1}} & THH^{(n-1)}(ku) \longrightarrow THH^{(n-1)}(ku_{\mathbb{Q}}) \\ & \searrow \text{ch}_n^{\geq 0} & \uparrow \simeq \\ & & THH^{(n-1)}((\Sigma^{\infty} \mathbb{C}P_+^{\infty})_{\mathbb{Q}}) \\ & & \downarrow \text{ev} \\ & & (\Sigma^{\infty} K(\mathbb{Z}, n+1)_+)_{\mathbb{Q}} \end{array}$$

described in the proof of [Theorem 1.10](#) with ev the basepoint evaluation projection from $THH^{(n-1)}(\Sigma^\infty \mathbb{C}P_+^\infty) = \Sigma^\infty \text{Map}(T^{n-1}, K(\mathbb{Z}, n+1))_+$ to $\Sigma^\infty K(\mathbb{Z}, n+1)_+$. The target of this map is the generalized rational Eilenberg–MacLane ring spectrum $(\Sigma^\infty K(\mathbb{Z}, n+1)_+)_\mathbb{Q} \simeq H\mathbb{Q}[\beta_n]$ whose homotopy is the ring $\mathbb{Q}[\beta_n]$.

Definition 2.3. For n odd, define the *higher Chern character* $\text{ch}_n : \mathfrak{A}_n \rightarrow H\mathbb{Q}[\beta_n^{\pm 1}]$ as the localization (at β_n) of the map $\text{ch}_n^{\geq 0} : \mathfrak{a}_n \rightarrow H\mathbb{Q}[\beta_n]$.

We note that when $n = 1$, this is the usual Chern character, which may be alternatively described as the map $KU \rightarrow KU_\mathbb{Q} \simeq (\Sigma^\infty \mathbb{C}P_+^\infty[\beta^{\pm 1}])_\mathbb{Q}$ induced by rationalization and Snaith’s (or Segal’s) theorem. More generally, ch_n factors the Chern–Dold character and evidently forgets some information regarding the rational homotopy of $\mathfrak{A}_n = \mathfrak{a}_n[\beta_n^{-1}]$:

$$\begin{array}{ccc} \mathfrak{a}_n & \xrightarrow{\text{ch}_{\mathfrak{a}_n}} & H(R_* \otimes \mathbb{Q}) \\ \downarrow & & \downarrow \\ \mathfrak{A}_n & \xrightarrow{\text{ch}_n} & H\mathbb{Q}[\beta_n^{\pm 1}] \end{array}$$

(the right vertical map is induced by $\text{ch}_n^{\geq 0}$ in rational homotopy). Nonetheless, it is useful on account of the fact that the coefficients of the codomain of ch_n are computed. Furthermore, this map is highly structured:

Proposition 2.4. *The higher Chern character $\text{ch}_n : \mathfrak{A}_n \rightarrow H\mathbb{Q}[\beta_n^{\pm 1}]$ is a map of E_∞ ring spectra. Further, the composite $\mathfrak{S}_n \rightarrow \mathfrak{A}_n \rightarrow H\mathbb{Q}[\beta_n^{\pm 1}]$ is an equivalence, and splits $H\mathbb{Q}[\beta_n^{\pm 1}]$ off of $\mathfrak{A}_n$.*

Proof. The work of [\[Blumberg et al. 2014\]](#) ensures that the Dennis trace is an E_∞ map. The original map³ $\text{ch}_n^{\geq 0} : \mathfrak{a}_n \rightarrow (\Sigma^\infty K(\mathbb{Z}, n+1)_+)_\mathbb{Q}$ in the definition of ch_n may then be seen to be E_∞ : it is a composite of iterates of tr , rationalizations, an equivalence induced by a map of ring spectra, and the map ev , which is easily seen to be E_∞ . Finally ch_n is also E_∞ , being obtained from this by localization at β_n .

To obtain the second statement, it suffices to note that $\iota \in \pi_{n+1}\mathfrak{A}_n$ is carried to β_n , which was shown in the proof of [Theorem 1.10](#). $\square$

Consequently ch_n induces a map of spectra $\text{ch}_n : \text{gl}_1(\mathfrak{A}_n) \rightarrow \text{gl}_1(H\mathbb{Q}[\beta_n^{\pm 1}])$. Further, since the composite

$$\Sigma^\infty K(\mathbb{Z}, n+1)_+ \xrightarrow{t_n} \mathfrak{a}_n \xrightarrow{\text{ch}_n^{\geq 0}} (\Sigma^\infty K(\mathbb{Z}, n+1)_+)_\mathbb{Q}$$

³This need not agree with the connective cover of the map ch_n !

is the rationalization map, it follows that ch_n carries the twisting t_n of $\mathfrak{A}_n$ by H^{n+2} to the standard⁴ twisting of periodic cohomology by H^{n+2} .

Corollary 2.5. *For each $H \in H^{n+2}(X)$, the higher Chern character ch_n is a natural transformation of twisted cohomology theories,*

$$\text{ch}_n : \mathfrak{A}_n^*(X; H) \rightarrow H\mathbb{Q}^*(X; H)[\beta_n^{\pm 1}],$$

which splits the target off of the source.

2C. $\mathfrak{A}$ -orientations and Umkehr maps. In order to construct the T-duality isomorphism in twisted $\mathfrak{A}_n$ -theory via a Fourier–Mukai push-pull formula, we will need to have an Umkehr map in $\mathfrak{A}_n$ -theory. Here we set out a framework for constructing and analyzing the Umkehr map associated to a sphere bundle. We will work with an arbitrary ring spectrum R equipped with a periodic twisting of n -gerbes $\phi : \mathfrak{S}_n \rightarrow R$, but we have $R = \mathfrak{A}_n$ in mind throughout.

Let $\pi : E \rightarrow X$ be a smooth fiber bundle with fiber the q -sphere S^q and $V \rightarrow E$ denote the vertical tangent bundle of the fiber bundle π . Consider the unit map $\eta : S \rightarrow R$ from the sphere spectrum, and the J -homomorphism $J : \mathcal{O} \rightarrow \text{BGL}_1(S)$ to the space of units of S , with BJ its delooping. Choose a map $V : E \rightarrow \text{BO}$ that classifies the vertical tangent bundle and let

$$\tau_E : E \xrightarrow{V} \text{BO} \xrightarrow{BJ} \text{BGL}_1(S) \xrightarrow{\text{BGL}_1(\eta)} \text{BGL}_1(R)$$

be the twist of R -theory that it determines. The structure of an R -orientation of V , i.e., a Thom class in R -theory for the Thom spectrum E^V , is equivalent to the datum of the homotopy class of a nullhomotopy of the map τ_E .

Definition 2.6. We call τ_E the *orientation twist of E* , since it is the obstruction to the R -orientability of the vertical tangent bundle V .

Associated to the sphere bundle $\pi : E \rightarrow X$ is a Pontrjagin–Thom collapse map $\pi^! : \Sigma^\infty X_+ \rightarrow E^{-V}$. Applying R -cohomology, we get a twisted Umkehr map $\pi_! : R^*(E^{-V}) \rightarrow R^*(X)$. Since the Thom spectrum associated to the twist $-\tau_E$ is the R -module spectrum $R \wedge E^{-V}$, we may rewrite the twisted Umkehr map as

$$\pi_! : R^*(E; \tau_E) \rightarrow R^*(X).$$

More generally, if $\omega : X \rightarrow \text{BGL}_1(R)$ is a twist of R over X , then there is a product twist $\tau_E \otimes \pi^* \omega$ induced by the action of $\text{BGL}_1 S$ on $\text{BGL}_1 R$ coming from the

⁴Traditionally when considering twisted periodic cohomology, the period of the theory is 2, but of course higher degrees also arise [Sati 2010]. In this setting, the period is somewhat longer, since β_n has dimension $n + 1$. Since n is odd, the period is still even, so the two are naturally comparable.

S -algebra structure of R . The Pontrjagin–Thom collapse map induces a twisted Umkehr map of the form

$$\pi_! : R^*(E; \tau_E \otimes \pi^* \omega) \rightarrow R^*(X; \omega).$$

In the presence of an orientation, we can recover the usual Umkehr map along π .

Proposition 2.7. *A trivialization of τ_E , or equivalently an R -orientation of V , determines an (untwisted) Umkehr map*

$$\pi_! : R^*(E) \cong R^{*-q}(E; \tau_E) \rightarrow R^{*-q}(X),$$

defined as the composite of the resulting Thom isomorphism in R -theory for E^{-V} with the twisted Umkehr map.

Remark 2.8. If R admits a periodic twisting by n -gerbes for $n = 1$, then V is R -orientable if and only if it admits a $\text{spin}_{\mathbb{C}}$ structure. In the case of odd $n > 1$, the spectrum R is an $H\mathbb{Q}$ -algebra (Proposition 1.16) and the virtual tangent bundle V is R -orientable if and only if it is an orientable vector bundle in the usual sense.

The twisted Umkehr map $\pi_!$ is natural in the variable X , meaning that if

$$\begin{array}{ccc} E' & \xrightarrow{\tilde{f}} & E \\ \pi' \downarrow & & \downarrow \pi \\ X' & \xrightarrow{f} & X \end{array}$$

is a pullback diagram of bundles, then there is a commutative diagram

$$\begin{array}{ccc} R^*(E'; \tau_{E'} \otimes \pi'^* f^* \omega) & \xrightarrow{\pi'_!} & R^*(X'; f^* \omega) \\ \tilde{f}^* \downarrow & & \downarrow f^* \\ R^*(E; \tau_E \otimes \pi^* \omega) & \xrightarrow{\pi_!} & R^*(X; \omega) \end{array} \quad (2.9)$$

relating the Umkehr maps along π and π' with the contravariant functoriality of the twisted cohomology theory $R^*(\cdot; \omega)$. Notice that we use the canonical equivalence of twists

$$\tilde{f}^*(\tau_E \otimes \pi^* \omega) \simeq \tilde{f}^* \tau_E \otimes \tilde{f}^* \pi^* \omega \simeq \tau_{E'} \otimes \pi'^* f^* \omega$$

to identify the target of $\tilde{f}^*$ with the source of $\pi_!$.

We will also need the following lemma on Mayer–Vietoris sequences.

Lemma 2.10. *The Umkehr map $\pi_!$ is natural for the boundary operator δ in Mayer–Vietoris sequences for twisted R -theory; i.e., if $X = U \cup V$ is a decomposition of X*

into open subsets, and we write $E_{U \cap V}$ for the restriction of the bundle E to the open subset $U \cap V$, then the following diagram commutes:

$$\begin{array}{ccc}
 R^*(E_{U \cap V}; \tau_{E_{U \cap V}} \otimes \pi^* \omega) & \xrightarrow{\pi|_{U \cap V}!} & R^*(U \cap V; \omega) \\
 \delta \downarrow & & \downarrow \delta \\
 R^{*+1}(E; \tau_E \otimes \pi^* \omega) & \xrightarrow{\pi_!} & R^{*+1}(X; \omega)
 \end{array}$$

Proof. We first construct the Mayer–Vietoris boundary operator δ in R -theory twisted by ω . We write $X^{-\omega}$ and $U^{-\omega}$, for the R -module Thom spectra associated to the inverse of the twist $\omega: X \rightarrow B \mathrm{GL}_1(R)$ and to the restriction of ω to U , where the latter is a slight abuse of notation. The functoriality of the Thom spectrum functor gives the commutative square of spectra on the left, which we extend to a morphism of cofiber sequences

$$\begin{array}{ccccccc}
 (U \cap V)^{-\omega} & \longrightarrow & U^{-\omega} & \xrightarrow{c_1} & C_1 & \xrightarrow{\partial} & \Sigma(U \cap V)^{-\omega} \\
 \downarrow & & \downarrow & & \downarrow \phi & & \downarrow \\
 V^{-\omega} & \longrightarrow & X^{-\omega} & \xrightarrow{c_2} & C_2 & \xrightarrow{\partial} & \Sigma V^{-\omega}
 \end{array}$$

The induced morphism ϕ on cofibers is an equivalence by the excision property of the parametrized cohomology theory associated to the parametrized R -module spectrum over X determined by ω [May and Sigurdsson 2006, §20]. Another point of view is that the Thom spectrum functor $(\cdot)^{-\omega}$ preserves homotopy pushouts [Ando et al. 2014a, Corollary 3.13]⁵, such as $X = U \cup_{U \cap V} V$, and thus the induced map on cofibers is an equivalence. The Mayer–Vietoris sequence boundary operator δ is defined to be the composite $\delta = c_2^* \circ (\phi^{-1})^* \circ \partial^*$ of the induced maps in R -cohomology, where ϕ^{-1} is a homotopy inverse to the equivalence ϕ .

There is a similar diagram defining the Mayer–Vietoris boundary operator associated to the decomposition $E = E_U \cup_{E_{U \cap V}} E_V$ and the naturality of the Pontrjagin–Thom construction constructs a commutative square of cofiber sequences relating the displayed diagram to the one for E . This gives the desired relation between δ and the twisted Umkehr map $\pi_!$. $\square$

3. T-duality for sphere bundles in theories twisted by n -gerbes

In this section, we define T-duality for sphere bundles and prove the T-duality isomorphism for cohomology theories admitting periodic twisting by n -gerbes. The

⁵The cited work uses an ∞ -categorical model for the Thom spectrum functor and so the relevant result is expressed using the language of colimits in ∞ -categories. It is a consequence of their work and the identification of homotopy colimits with ∞ -colimits that any model for the Thom spectrum functor preserves homotopy colimits.

material closely follows that of [Bunke and Schick 2005] in the setting of T-duality for circle bundles. We will also prove that when $n > 1$, any cohomology theory for which such an isomorphism holds must be rational.

3A. Formulation of T-duality for sphere bundles. Let X be a topological space, and $\pi : E \rightarrow X$ and $\hat{\pi} : \hat{E} \rightarrow X$ be S^q -bundles over X . For simplicity, we work with smooth fiber bundles, but the arguments can be adapted to the case of topological fiber bundles. Note that we do not require the sphere bundles to be orientable. In the analysis of classifying spaces in Section 4, we will work with a chosen structure group for our bundles, but for now that choice is not relevant.

Define $E *_X \hat{E}$ to be the *fiberwise join* of E and $\hat{E}$ over X . Note that this is a bundle with fiber the join $S^q * S^q = S^{2q+1}$, and that there are natural fiberwise embeddings

$$i : E \hookrightarrow E *_X \hat{E} \quad \text{and} \quad \hat{i} : \hat{E} \hookrightarrow E *_X \hat{E}$$

given by inclusion of each factor in the join.

Definition 3.1. A *bundle Thom class*⁶ for an S^m -bundle $p : Y \rightarrow X$ is an element $\text{Th} \in H^m(Y; \mathbb{Z})$ with the property that its restriction to each fiber is a generator of $H^m(S^m; \mathbb{Z})$. Equivalently, $p_!(\text{Th}) = \pm 1 \in H^0(X; \mathbb{Z})$.

We let $n = 2q - 1$, which is evidently odd, and consider representatives

$$H : E \rightarrow K(\mathbb{Z}, n+2) \quad \text{and} \quad \hat{H} : \hat{E} \rightarrow K(\mathbb{Z}, n+2)$$

for cohomology classes

$$[H] \in H^{n+2}(E; \mathbb{Z}) \quad \text{and} \quad [\hat{H}] \in H^{n+2}(\hat{E}; \mathbb{Z}).$$

Definition 3.2. We say that the pairs (E, H) and $(\hat{E}, \hat{H})$ are *T-dual* if there exists a bundle Thom class $\text{Th} \in H^{n+2}(E *_X \hat{E})$ with $i^* \text{Th} = [H]$ and $\hat{i}^* \text{Th} = [\hat{H}]$.

Consider the *correspondence space* $E \times_X \hat{E}$, which is an $S^q \times S^q$ -bundle over X :

$$\begin{array}{ccc} & E \times_X \hat{E} & \\ p \swarrow & & \searrow \hat{p} \\ E & & \hat{E} \\ \pi \searrow & & \swarrow \hat{\pi} \\ & X & \end{array}$$

⁶This is simply called a *Thom class* in [Bunke and Schick 2005]; this usage conflicts with the usual notion of a Thom class as a generator θ for the cohomology of a Thom space. The two notions are closely related, since θ is carried to Th under the collapse map from a sectioned sphere bundle to its Thom space.

There is a tautological homotopy $h : I \times (E \times_X \widehat{E}) \rightarrow E *_X \widehat{E}$ from $i \circ p$ to $\hat{i} \circ \hat{p}$ that is given by the formula $h_t(e, \hat{e}) = (t, e, \hat{e})$ and recognizes the fiberwise join $E *_X \widehat{E}$ as a quotient of $I \times (E \times_X \widehat{E})$.

Pulling the twisting classes back over p and $\hat{p}$ gives cohomology classes $p^*[H]$ and $\hat{p}^*[\widehat{H}]$, respectively, on the correspondence space. But since $i^*\text{Th} = [H]$ and $\hat{i}^*\text{Th} = [\widehat{H}]$, we have a homotopy between the maps representing $p^*[H]$ and $\hat{p}^*[\widehat{H}]$:

$$\Lambda = \text{Th} \circ h : H \circ p = \text{Th} \circ i \circ p \rightarrow \text{Th} \circ \hat{i} \circ \hat{p} = \widehat{H} \circ \hat{p}.$$

The T-duality transformation T is the composite

$$\begin{array}{ccc} R^*(E; \tau_E \otimes H) & \xrightarrow{p^*} & R^*(E \times_X \widehat{E}; p^*\tau_E \otimes p^*H) \\ & & \cong \downarrow \Lambda \\ & & R^*(E \times_X \widehat{E}; \tau_{E \times_X \widehat{E}} \otimes \hat{p}^*\widehat{H}) \xrightarrow{\hat{p}!} R^*(\widehat{E}; \widehat{H}) \end{array}$$

where the middle isomorphism is induced by the homotopy Λ and the canonical isomorphism of the pullback along p of the vertical tangent bundle of $\pi : E \rightarrow X$ with the vertical tangent bundle of $\hat{p} : E \times_X \widehat{E} \rightarrow \widehat{E}$. We may now state our main theorem.

Theorem 3.3. *Let R be an A_∞ ring spectrum, and let $\phi : \mathfrak{S}_n \rightarrow R$ be a periodic twisting of R by n -gerbes. Suppose that X is homotopy equivalent to a finite CW complex, that (E, H) and $(\widehat{E}, \widehat{H})$ are T-dual, and write $\tau_E : X \rightarrow B\text{GL}_1(R)$ for the orientation twist of the vertical tangent bundle of E . Then the T-duality transformation*

$$T := \hat{p}! \circ \Lambda \circ p^* : R^*(E; \tau_E \otimes H) \rightarrow R^*(\widehat{E}; \widehat{H})$$

is an isomorphism.

Recall that our main examples of such an R are $\mathfrak{S}_n$ itself (Definition 1.13), and $\mathfrak{A}_n$ (Definition 1.11).

3B. Proof of Theorem 3.3. We will establish Theorem 3.3 through a series of lemmata.

Lemma 3.4. *Theorem 3.3 holds when X is a point.*

Proof. In this case, the Atiyah–Hirzebruch spectral sequence for the R -cohomology of $E \times \widehat{E} = S^q \times S^q$ contains $R^*(E)$ and $R^*(\widehat{E})$ as direct summands on every page, and thus cannot have any nonzero differentials. This gives an isomorphism of rings

$$R^*(E \times \widehat{E}) \cong R^*[x, \hat{x}]/(x^2, \hat{x}^2),$$

where x and $\hat{x}$ are represented by singular cohomology classes of dimension q . Write $\beta = \phi_*(\iota) \in \pi_{n+1}(R)$ for the periodicity class; when $R = \mathfrak{A}_n$, $\beta = \beta_n$. We claim that Λ is given by multiplication by the degree-0 class $1 \pm \beta x \hat{x}$, where we are regarding β as a $-2q = -(n+1)$ -dimensional cohomology class via the identification $R^*(\text{pt}) = \pi_{-*}R$.

Granting that, the result follows: $\hat{p}_! \circ \Lambda \circ p^*(1) = \hat{p}_!(1 \pm \beta x \hat{x}) = \pm \beta \hat{x}$, and

$$\hat{p}_! \circ \Lambda \circ p^*(x) = \hat{p}_![(1 \pm \beta x \hat{x})x] = \hat{p}_!(x) = 1$$

since the Umkehr map $\hat{p}_!$ is given by division by the class x if possible, and is 0 otherwise.

To see the claim, we first recall that for a trivial cohomology class $\alpha = 0 \in H^m(Y; \mathbb{Z})$, represented by a map $\alpha : Y \rightarrow K(\mathbb{Z}, m)$, the set of homotopy classes of nullhomotopies of α is a torsor for

$$[Y, \Omega K(\mathbb{Z}, m)] = H^{m-1}(Y; \mathbb{Z}) = H^m(\Sigma Y, \mathbb{Z}),$$

by concatenation of homotopies. In particular, it is in bijection with this group.

We note that since $H^{n+2}(E) = 0 = H^{n+2}(\hat{E})$, we have $H = \hat{H} = 0$. So p^*H and $\hat{p}^*\hat{H}$ are necessarily null. Further, since $H^{n+1}(E) = 0 = H^{n+1}(\hat{E})$, there is a unique homotopy class of trivialization (nullhomotopy) of H and $\hat{H}$, yielding preferred nullhomotopies of p^*H and $\hat{p}^*\hat{H}$. Note that the nullhomotopies of the pulled-back classes are not unique, since the group $H^{n+1}(E \times \hat{E}) = \mathbb{Z}$ affords many possibilities.

Indeed, the homotopy Λ of [Theorem 3.3](#) is a nontrivial homotopy between the trivialized twists p^*H and $\hat{p}^*\hat{H}$. We may therefore regard it as being represented through the action of an element of the group $H^{n+1}(E \times \hat{E}) \cong H^{n+2}\Sigma(E \times \hat{E})$ on the trivial twist. But by construction, Λ is given as a composite

$$\Sigma(E \times \hat{E}) \rightarrow E * \hat{E} = S^{n+2} \xrightarrow{\text{Th}} K(\mathbb{Z}, n+2).$$

Here the first map arises from the tautological homotopy $I \times E \times \hat{E} \rightarrow E * \hat{E}$; the factorization through $\Sigma(E \times \hat{E})$ uses the fact that the inclusions of E and $\hat{E}$ into $E * \hat{E}$ are nullhomotopic. In particular, this map induces an isomorphism in H^{n+2} . By assumption, this composite is a generator of $H^{n+2}\Sigma(E \times \hat{E})$. The map that Λ induces in R^* is therefore given by multiplication by the invertible class

$$\begin{array}{ccccccc} E \times \hat{E} & \xrightarrow{\quad} & \Omega S^{n+2} & \xrightarrow{\quad \Omega \text{Th} \quad} & K(\mathbb{Z}, n+1) & \xrightarrow{\quad \text{GL}_1 \phi \quad} & \text{GL}_1 R \\ & \searrow & \nearrow & \nearrow & \nearrow & \nearrow & \\ & & S^{n+1} & & & & \end{array}$$

$\pm \iota$ $1 \pm \beta$

The result follows, since the top-dimensional cohomology class of S^{n+1} pulls back to $x\hat{x}$ on $E \times \widehat{E}$. $\square$

We now establish the naturality properties of the T-duality transformation that we need to prove [Theorem 3.3](#) by cellular induction.

Lemma 3.5. *Let $f : Y \rightarrow X$ be a continuous map, and define $F : f^*E \rightarrow E$ and $\widehat{F} : f^*\widehat{E} \rightarrow \widehat{E}$ to be induced maps from the pullbacks of E and $\widehat{E}$ along f . Then the T-duality map for the bundles E and $\widehat{E}$ on X pulls back over f ; that is, the following diagram commutes:*

$$\begin{array}{ccc} R^*(E, \tau_E \otimes H) & \xrightarrow{T} & R^*(\widehat{E}, \widehat{H}) \\ F^* \downarrow & & \downarrow \widehat{F}^* \\ R^*(f^*E, F^*(\tau_E \otimes H)) & \xrightarrow{T} & R^*(f^*\widehat{E}, \widehat{F}^*\widehat{H}) \end{array}$$

Proof. The maps p^* and Λ commute with F^* and $\widehat{F}^*$ by the contravariant functoriality of twisted cohomology theories. The twisted Umkehr map $\hat{p}_!$ also commutes with F^* and $\widehat{F}^*$ by diagram (2.9). $\square$

Lemma 3.6. *If X is decomposed as the union $X = U \cup V$ of open subsets, we may restrict E and $\widehat{E}$ to these subsets, obtaining $E = E_U \cup E_V$ and $\widehat{E} = \widehat{E}_U \cup \widehat{E}_V$. Further, let δ and $\hat{\delta}$ be the boundary operators in the Mayer–Vietoris sequences for these decompositions of E and $\widehat{E}$, respectively. Then the T-duality transformation preserves these Mayer–Vietoris sequences; that is,*

$$T \circ \delta = \hat{\delta} \circ T : R^*(E_{U \cap V}, (\tau_E \otimes H)|_{E_{U \cap V}}) \rightarrow R^{*+1}(\widehat{E}, \widehat{H}).$$

Proof. We argue that each component of $T = \hat{p}_! \circ \Lambda \circ p^*$ commutes with the relevant boundary operator. The pullback map p^* is induced by a map of spectra over E and hence extends to a natural transformation of parametrized cohomology theories on spaces over E [[May and Sigurdsson 2006](#), §20]. The data of the boundary map ∂ in the cofiber sequence associated to a pair of spaces over E , such as in a Mayer–Vietoris sequence, is preserved by such a natural transformation. Similarly, the map Λ is induced by a map of spectra over $E \times_X \widehat{E}$, so commutes with the boundary operators. The twisted Umkehr map $\hat{p}_!$ commutes with the boundary operators by [Lemma 2.10](#). $\square$

The proof of [Theorem 3.3](#) immediately follows by cellular induction, using the previous three results, [Lemmas 3.4–3.6](#).

Remark 3.7. Note that if R' is an A_∞ ring spectrum and $\psi : R \rightarrow R'$ an A_∞ map, the composite $\psi \circ \phi$ is a periodic twisting of R' by n -gerbes, and so the results of this theorem also hold for R' . Further, it is evident from the proof that ψ throws the T-duality isomorphism for R onto that for R' .

Summarizing this result in the case that $R = \mathfrak{A}_n$, $R' = H\mathbb{Q}[\beta_n^{\pm 1}]$, and $\psi = \text{ch}_n$, we have:

Corollary 3.8. *The higher Chern character ch_n throws the T-duality isomorphism for $\mathfrak{A}_n$ onto that of periodic rational cohomology, in the sense that the following diagram commutes:*

$$\begin{array}{ccc} \mathfrak{A}_n^*(E; \tau_E \otimes H) & \xrightarrow[\cong]{T_{\mathfrak{A}_n}} & \mathfrak{A}_n^*(\widehat{E}; \widehat{H}) \\ \text{ch}_n \downarrow & & \downarrow \text{ch}_n \\ H\mathbb{Q}^*(E; \tau_E \otimes H)[\beta_n^{\pm 1}] & \xrightarrow[\cong]{T_{H\mathbb{Q}[\beta_n^{\pm 1}]}} & H\mathbb{Q}^*(\widehat{E}; \widehat{H})[\beta_n^{\pm 1}] \end{array}$$

3C. A T-duality isomorphism implies rationality. Let R be an A_∞ ring spectrum which admits a twisting by $n + 2$ -dimensional cohomology classes. That is, we take as given a map

$$\tau : K(\mathbb{Z}, n + 2) \rightarrow B\text{GL}_1(R),$$

whence we may define twisted cohomology groups $R^*(X, H)$ for $H \in H^{n+2}(X, \mathbb{Z})$. Note we do not assume that this twisting is periodic in the sense of [Definition 1.15](#). In other words, the map

$$t : \Sigma_+^\infty K(\mathbb{Z}, n + 1) \rightarrow R$$

of ring spectra adjoint to τ is not assumed to factor through the Arthan spectrum $\mathfrak{S}_n = \Sigma_+^\infty K(\mathbb{Z}, n + 1)[t^{-1}]$. In particular, we may not immediately conclude from [Proposition 1.16](#) that R is a rational spectrum.

One may ask whether a T-duality isomorphism holds in this setting. That is, if (E, H) and $(\widehat{E}, \widehat{H})$ are T-dual S^q -bundles⁷ over X in the sense of [Definition 3.2](#), we may, as before, form the T-duality transformation T as the composite

$$\begin{array}{ccc} R^*(E; \tau_E \otimes H) & \xrightarrow{p^*} & R^*(E \times_X \widehat{E}; p^* \tau_E \otimes p^* H) \\ & \cong \downarrow \Lambda & \\ R^*(E \times_X \widehat{E}; \tau_{E \times_X \widehat{E}} \otimes \widehat{p}^* \widehat{H}) & \xrightarrow{\widehat{p}_!} & R^*(\widehat{E}; \widehat{H}) \end{array}$$

and ask whether T is an isomorphism.

Theorem 3.9. *The following are equivalent:*

- (1) *The T-duality transformation T is an isomorphism for all base spaces X of the homotopy type of a finite CW complex.*
- (2) *The T-duality transformation T is an isomorphism for X a point.*

⁷We still require $n = 2q + 1$.

(3) *The twisting τ is periodic.*

In particular, any cohomology theory which admits a T-duality isomorphism for S^q -bundles with $q > 1$ must be a rational cohomology theory.

Proof. Clearly the first statement implies the second. Conversely, the proof of [Theorem 3.3](#) showed that the general case reduced to the case X a point using naturality ([Lemma 3.5](#)) and the Mayer–Vietoris sequence ([Lemma 3.6](#)) — note that these results do not require the twisting to be periodic. We have shown that the T-duality isomorphism holds when X is a point under the assumption of a periodic twisting ([Lemma 3.4](#)), so the third statement implies the second. So we need only show that the second statement implies the third.

Let us define $\beta : S^{n+1} \rightarrow R$ as we have done in [Section 1B](#), via the composite

$$S^{n+1} \xrightarrow{\iota} \Sigma^\infty K(\mathbb{Z}, n+1)_+ \xrightarrow{t} R$$

where t is adjoint to τ . We will show that β is an invertible element of $\pi_* R$ under the assumption that T-duality holds over a point. This then implies that the map of ring spectra t factors through the Arthan spectrum $\mathfrak{S}_n$, and in particular makes R an algebra spectrum over $\mathfrak{S}_n$.

As in the proof of [Lemma 3.4](#), we may write $R^*(E \times \widehat{E}) = R^*[x, \hat{x}]/(x^2, \hat{x}^2)$ and compute the T-duality map to be the R_* -linear operation given by

$$1 \mapsto \pm\beta\hat{x} \quad \text{and} \quad x \mapsto 1.$$

This can only be an isomorphism if β is invertible. □

4. Classifying spaces for T-dual pairs

We continue to insist that $n = 2q - 1$ be odd. In this section, we introduce spaces $R_n(G)$ and $P_n(G)$ that classify the objects considered in the previous section, at least in the case where the bundles E and $\widehat{E}$ are oriented. More precisely, let G be a topological group equipped with a homomorphism to $\text{Homeo}^+(S^q)$, the group of orientation-preserving homeomorphisms of the q -sphere S^q .

We will construct natural isomorphisms that give the following identifications:

- $[X, R_n(G)]$ is the set of equivalence classes of pairs (E, H) , where $E \rightarrow X$ is an S^q -bundle with structure group G , and H is a map representing a twisting class $[H] \in H^{n+2}(E)$.
- $[X, P_n(G)]$ is the set of equivalence classes of triples $(E, \widehat{E}, \text{Th})$, where E and $\widehat{E}$ are S^q -bundles with structure group G , and $\text{Th} \in H^{n+2}(E *_X \widehat{E})$ is a bundle Thom class.

There is a map $P_n(G) \rightarrow R_n(G)$ that induces the natural transformation which carries $(E, \widehat{E}, \text{Th})$ to $(E, i^* \text{Th})$ (where we make a slight abuse of notation and

write $i^* \text{Th}$ for a choice of representing map). We will analyze the homotopy type of the spaces involved and show that this map is an equivalence precisely when $q = 1$.

Remark 4.1. We may regard $[X, P_n(G)]$ as the set of T-dual pairs over X . The failure of this map to be an equivalence for $q > 1$ shows that it is not the case that every pair (E, H) has a T-dual $(\widehat{E}, \widehat{H})$, and that when a T-dual exists, there is no guarantee that it is unique. However, we will encounter special cases towards the end of [Section 4C](#) where T-duals do exist and can be unique.

4A. Euler classes and the classifying space $P_n(G)$. Let $p : E \rightarrow X$ be a fiber bundle with fiber S^m . If E is the subsphere bundle $E = S(V)$ of an oriented real vector bundle V over X of rank $m + 1$, then there is an *Euler class*

$$e(E) = e(V) \in H^{m+1}(X)$$

which is the pullback of the (usual notion of the) Thom class of X^V under the zero-section.

More generally, if E is an oriented S^m -bundle, then the fiberwise unreduced suspension

$$\Sigma_X E := I \times E / \sim, \quad \text{where } (t, e) \sim (t, e') \text{ for } t = 0, 1 \text{ when } p(e) = p(e'),$$

is a topological S^{m+1} -bundle over X which inherits an orientation from E . There are two canonical sections $s_t : X \rightarrow \Sigma_X E$, where $t = 0, 1$, given by $s_t(x) = [t, e]$ for any e in the fiber over x . The associated Thom space

$$X^{E \oplus 1} := \Sigma_X E / s_1(X)$$

is obtained by collapsing the image of s_1 to a point. Since $\Sigma_X E$ is oriented, there is a Thom class $\theta_E \in H^{m+1}(X^{E \oplus 1})$ in integral cohomology with the property that the map

$$H^*(X) \rightarrow \widetilde{H}^{*+m+1}(X^{E \oplus 1}) \quad \text{given by} \quad \alpha \mapsto (\Sigma_X p)^*(\alpha) \cup \theta_E$$

is an isomorphism. We define the Euler class of E by the formula $e(E) = s_0^*(\theta_E)$. The following is a straightforward application of the Leray–Hirsch theorem and the Gysin sequence; in the case $m = 1$, see [\[Bunke and Schick 2005, §2.2; Bouwknegt et al. 2004, §3.1\]](#).

Proposition 4.2. *The following are equivalent:*

- (1) E admits a bundle Thom class $\text{Th} \in H^m(E)$.
- (2) $H^*(E)$ is a free $H^*(X)$ -module on $1, \text{Th}$.
- (3) $e(E) = 0$.

The set of homotopy classes of nullhomotopies of a representative cocycle for $e(E) \in H^{m+1}(X)$, if trivial, is a torsor under $H^m(X)$. The same is true of bundle Thom classes $\text{Th} \in H^m(E)$: for any $\alpha \in H^m(X)$, $\text{Th} + p^*(\alpha)$ is another such bundle Thom class. This structure is indeed a torsor (as it arises from a free and transitive action of $H^m(X)$) by virtue of point (2) in the above proposition. Thus we may in fact conclude:

the set of homotopy classes of trivializations of $e(E)$ is naturally in bijection with the set of bundle Thom classes $\text{Th} \in H^m(E)$.

The next result implies that our definition of T-duality may be described in terms of characteristic classes, as in the cases $q = 1, 3$ [Bouwknegt et al. 2004; Bunke and Schick 2005; Bouwknegt et al. 2015a].

Proposition 4.3. *Suppose that E and $\widehat{E}$ are orientable S^q -bundles over X , and that (E, H) and $(\widehat{E}, \widehat{H})$ are T-dual. Then the equations*

$$e(E) = \pm \pi_![\widehat{H}] \quad \text{and} \quad e(\widehat{E}) = \pm \pi_![H]$$

hold in $H^{q+1}(X)$.

Proof. Let $\Sigma_X E$ and $\Sigma_X \widehat{E}$ be the oriented S^{q+1} -bundles over X formed by taking the fiberwise unreduced suspension. Form the fiberwise smash product $\Sigma_X E \wedge_X \Sigma_X \widehat{E}$ with respect to the canonical sections s_1 on either side. This is an S^{2q+2} -bundle and that inherits an orientation from E and $\widehat{E}$. In other words, we equip the Thom space

$$X^{E \oplus \widehat{E} \oplus 2} := (\Sigma_X E \wedge_X \Sigma_X \widehat{E}) / s_1(X)$$

with the Thom class $\theta_E \cdot \theta_{\widehat{E}} \in H^{2q+2}(X^{E \oplus \widehat{E} \oplus 2})$. Consequently, the associated Euler class satisfies

$$e(E *_X \widehat{E}) = e(E) \cdot e(\widehat{E}).$$

The S^{2q+1} -bundle $E *_X \widehat{E}$ also inherits an orientation from E and $\widehat{E}$; the Thom class on the associated Thom space agrees with $\theta_E \cdot \theta_{\widehat{E}}$ under the homeomorphism

$$X^{(E *_X \widehat{E}) \oplus 1} \cong X^{E \oplus \widehat{E} \oplus 2}.$$

Using these orientations to define Thom isomorphisms and their associated Umkehr maps, the inclusion $i : E \rightarrow E *_X \widehat{E}$ induces a commutative diagram relating the Gysin sequences for $\rho : E *_X \widehat{E} \rightarrow X$ and $\pi : E \rightarrow X$:

$$\begin{array}{ccccccc} \dots & \longrightarrow & H^{2q+1}(X) & \xrightarrow{\rho^*} & H^{2q+1}(E *_X \widehat{E}) & \xrightarrow{\rho_!} & H^0(X) \xrightarrow{e(E *_X \widehat{E})} H^{2q+2}(X) \longrightarrow \dots \\ & & \parallel & & \downarrow i^* & & \downarrow e(\widehat{E}) \\ \dots & \longrightarrow & H^{2q+1}(X) & \xrightarrow{\pi^*} & H^{2q+1}(E) & \xrightarrow{\pi_!} & H^{q+1}(X) \xrightarrow{e(E)} H^{2q+2}(X) \longrightarrow \dots \end{array}$$

Now suppose that (E, H) and $(\widehat{E}, \widehat{H})$ are T-dual. In particular, we are given a bundle Thom class $\text{Th} \in H^{2q+1}(E *_X \widehat{E})$ satisfying $i^* \text{Th} = [H]$ and $\hat{i}^* \text{Th} = [\widehat{H}]$. We do not know if (the suspension of) Th is the pullback of our preferred Thom class $\theta_E \cdot \theta_{\widehat{E}}$, but we do know that $\rho_! \text{Th} = \pm 1$, and consequently that $\pi_! [H] = \pm e(\widehat{E})$. The relation $\hat{\pi}_! [\widehat{H}] = \pm e(E)$ follows from the analogous diagram involving the Gysin sequence for $\hat{\pi}$. $\square$

We now construct the space $P_n(G)$ which parametrizes triples $(E, \widehat{E}, \text{Th})$, where E and $\widehat{E}$ are S^q -bundles with structure group G , and Th is a bundle Thom class on $E *_X \widehat{E}$.

Definition 4.4. Let $P_n(G)$ be the homotopy fiber of the composite

$$BG \times BG \xrightarrow{B\text{join}} B\text{Homeo}^+(S^q * S^q) \xrightarrow{e} K(\mathbb{Z}, 2q+2) = K(\mathbb{Z}, n+3).$$

Here $\text{join} : G \times G \rightarrow \text{Homeo}^+(S^q * S^q)$ is the join of a pair of orientation-preserving homeomorphisms and e classifies the Euler class.

By construction, homotopy classes of maps $X \rightarrow P_n(G)$ are in bijection with the set of equivalence classes of the following data:

- a pair $(E, \widehat{E})$ of S^q -bundles over X with structure group G , and
- the homotopy class of a trivialization of the Euler class of the fiberwise join $E *_X \widehat{E}$.

Applying [Proposition 4.2](#) (or rather, its subsequent refinement) in this setting yields:

Proposition 4.5. *There is a bijection between $[X, P_n(G)]$ and the set of equivalence classes of triples $(E, \widehat{E}, \text{Th})$, where $\text{Th} \in H^{n+2}(E \times_X \widehat{E})$ is a bundle Thom class.*

Remark 4.6. Up to equivalence, the data of a triple $(E, \widehat{E}, \text{Th})$ may be identified with the data of two T-dual pairs (E, H) and $(\widehat{E}, \widehat{H})$ via $(E, \widehat{E}, \text{Th}) \mapsto (E, i^* \text{Th}), (E, \hat{i}^* \text{Th})$. Thus, $P_n(G)$ is the classifying space for T-dual pairs.

4B. The classifying space $R_n(G)$. We now work with slightly more generality. Let F be a topological space and suppose that G is a topological group equipped with a continuous homomorphism $G \rightarrow \text{Homeo}(F)$. Consider pairs (E, H) consisting of a fiber bundle $\pi : E \rightarrow X$ with fiber F and structure group G , along with a map $H : E \rightarrow K(\mathbb{Z}, n+2)$ representing a cohomology class $[H] \in H^{n+2}(E; \mathbb{Z})$ on the total space that is fiberwise trivial, i.e., for every $x \in X$, the restriction of $[H]$ to $H^{n+2}(E_x; \mathbb{Z})$ is zero. We will construct a classifying space $R_n(G)$ for such pairs (E, H) . Our construction is a generalization of Bunke and Schick's classifying space for principal circle bundles equipped with a $U(1)$ -gerbe [\[2005\]](#).

Let $\text{Map}_0(F, K(\mathbb{Z}, n+2))$ be the connected component of $\text{Map}(F, K(\mathbb{Z}, n+2))$ containing the basepoint. In other words, $\text{Map}_0(F, K(\mathbb{Z}, n+2))$ is the subspace of

nullhomotopic maps. The group G acts on $\text{Map}(F, K(\mathbb{Z}, n+2))$ by $(g \cdot f)(x) = f(g^{-1}x)$.

Proposition 4.7. *Equivalence classes of pairs (E, H) are classified by the space*

$$R_n(G) := EG \times_G \text{Map}_0(F, K(\mathbb{Z}, n+2)).$$

Proof. Starting with a group G , one can extend the G -Borel construction $EG \times_G (\cdot)$ to the two-sided bar construction $B(\cdot, G, \cdot)$, where the first and last entries are right and left G -spaces, respectively. Applying this to the space of functions from F to $K = K(\mathbb{Z}, n+2)$ gives $R = B(*, G, \text{Map}_0(F, K))$. The total space of the universal pair over R is

$$E_{\text{univ}} = B(F, G, \text{Map}_0(F, K)) \rightarrow B(*, G, \text{Map}_0(F, K)) = R$$

with projection to R induced by $F \rightarrow *$. Note the evaluation map $F \times \text{Map}_0(F, K) \rightarrow K$ is G -invariant and hence descends to the homotopy quotient:

$$H_{\text{univ}} : B(F, G, \text{Map}_0(F, K)) \rightarrow K.$$

This map represents the universal cohomology class $[H_{\text{univ}}] \in H^{n+2}(E_{\text{univ}}; \mathbb{Z})$. The pair $(E_{\text{univ}}, H_{\text{univ}})$ is the universal pair.

Given a pair (E, H) , we now construct a classifying map $X \rightarrow R$. Let $P \rightarrow X$ be the principal G -bundle associated to E . The usual contracting simplicial homotopy shows that the projection $B(G, G, P) \rightarrow P$ induced by the action of G on P is a homotopy equivalence. Now form the following diagram of principal G -bundles:

$$\begin{array}{ccccc} P & \xleftarrow{\cong} & B(G, G, P) & \longrightarrow & B(G, G, \text{Map}_0(F, K)) \\ \downarrow & & \downarrow & & \downarrow \\ X & \xleftarrow{\cong} & B(*, G, P) & \longrightarrow & B(*, G, \text{Map}_0(F, K)) = R \end{array}$$

The map on the lower left is an equivalence by the five lemma, so we may choose a wrong-way homotopy inverse as indicated by the dashed arrow. The maps pointed to the right are induced by the G -map $P \rightarrow \text{Map}_0(F, K)$ adjoint to the map $H : P \times_G F = E \rightarrow K$. The composite along the bottom gives the classifying map for the pair (E, H) .

To see that the fiber bundle $E \rightarrow X$ is the pullback of the fiber bundle $E_{\text{univ}} \rightarrow R$ along this classifying map, apply $F \times_G (\cdot)$ to the total spaces of the principal G -bundles in the diagram. $\square$

Remark 4.8. More generally, if we replace $K(\mathbb{Z}, n+2)$ by any space K , then R classifies equivalence classes of pairs (E, H) consisting of a fiber bundle $E \rightarrow X$ with fiber F and structure group G and a map $H : E \rightarrow K$ that is fiberwise nullhomotopic.

4C. Comparing $R_n(G)$ and $P_n(G)$. We now return to the usual setting of oriented S^q -bundles with structure group G and set $n = 2q - 1$. We will analyze the map $f : P_n(G) \rightarrow R_n(G)$ that represents the forgetful functor taking a T-dual pair (E, H) and $(\widehat{E}, \widehat{H})$ to the first item (E, H) . Abbreviate $P = P_n(G)$ and $R = R_n(G)$ and note that by construction R fits into a fiber sequence

$$\mathrm{Map}(S^q, K(\mathbb{Z}, n+2)) \rightarrow R \rightarrow BG.$$

Also note that, by the connectivity of the Eilenberg–MacLane space, $\mathrm{Map}(S^q, K(\mathbb{Z}, n+2))$ is connected, so we are automatically restricting to nullhomotopic maps as in the construction of R .

The classifying space P also participates in a fiber sequence, which we extend to the left by one entry:

$$\cdots \rightarrow K(\mathbb{Z}, n+2) \rightarrow P \rightarrow BG \times BG \xrightarrow{e(\cdot, \cdot)} K(\mathbb{Z}, n+3).$$

The composite of the map from P with the projection $\mathrm{pr}_i : BG \times BG \rightarrow BG$ to the first or second factor classifies a principal G -bundle with fiber S^q over P , which we call E and $\widehat{E}$, respectively. The fiberwise join $E * \widehat{E}$ carries a canonical bundle Thom class by [Proposition 4.5](#). Pulling it back along the inclusion $i_1 : E \rightarrow E * \widehat{E}$ gives a class $[H] \in H^{n+2}(E)$. Choose a representing map $H : E \rightarrow K(\mathbb{Z}, n+2)$. By [Proposition 4.7](#), the pair (E, H) over P is classified by a map $f : P \rightarrow R$ which fits into a map of fiber sequences

$$\begin{array}{ccccc} K(\mathbb{Z}, n+2) & \longrightarrow & P & \longrightarrow & BG \times BG \\ \downarrow & & \downarrow f & & \downarrow \mathrm{pr}_1 \\ \mathrm{Map}(S^q, K(\mathbb{Z}, n+2)) & \longrightarrow & R & \longrightarrow & BG \end{array} \quad (4.9)$$

Consider the new fibration $P \rightarrow BG$ given by passing from the upper middle to lower right of this diagram; this now has fiber $BG \times K(\mathbb{Z}, n+2)$. There is a morphism of fiber sequences over BG :

$$\begin{array}{ccccc} BG \times K(\mathbb{Z}, n+2) & \longrightarrow & P & \longrightarrow & BG \\ \downarrow & & \downarrow f & & \parallel \\ \mathrm{Map}(S^q, K(\mathbb{Z}, n+2)) & \longrightarrow & R & \longrightarrow & BG \end{array} \quad (4.10)$$

To compare the homotopy types of P and R , we may then compare the homotopy type of their fibers. Define a function $\tilde{e} : BG \rightarrow \mathrm{Map}(S^q, K(\mathbb{Z}, n+2))$ as the composite

$$BG \xrightarrow{e} K(\mathbb{Z}, q+1) \simeq \Omega^q K(\mathbb{Z}, n+2) \rightarrow \mathrm{Map}(S^q, K(\mathbb{Z}, n+2)).$$

Here, e is the Euler class of an S^q bundle with structure group G , and the second map is the inclusion of the space of based maps into all maps. Note that the fiber $\text{Map}(S^q, K(\mathbb{Z}, n+2))$ of the bottom fibration in (4.10) is an H-space, via multiplication of maps in the target.

Lemma 4.11. *In the comparison between the fibers over P and R :*

- (1) *The induced map of fibers $K(\mathbb{Z}, n+2) \rightarrow \text{Map}(S^q, K(\mathbb{Z}, n+2))$ in (4.9) is homotopic to the inclusion $x \mapsto \text{const}_x$ of constant maps.*
- (2) *The fiber map $BG \times K(\mathbb{Z}, n+2) \rightarrow \text{Map}(S^q, K(\mathbb{Z}, n+2))$ in (4.10) is homotopic to the product of the maps $\tilde{e}$ and const .*

Proof. The fundamental class $\iota : S^{n+2} \rightarrow K(\mathbb{Z}, n+2)$ represents the bundle Thom class of the trivial sphere bundle over a point. Thus, the induced map of fibers in (4.9) is adjoint to the composite

$$S^q \times K(\mathbb{Z}, n+2) \xrightarrow{i_1 \times 1} S^{2q+1} \times K(\mathbb{Z}, n+2) \xrightarrow{\iota \times 1} K(\mathbb{Z}, n+2) \times K(\mathbb{Z}, n+2) \xrightarrow{+} K(\mathbb{Z}, n+2).$$

This composite is homotopic to the projection to the second factor, which proves the first claim.

The second claim is a parametrized form of the first. The identity map of the space P classifies a triple $(E, \widehat{E}, \text{Th})$ over P of S^q -bundles E and $\widehat{E}$, and a bundle Thom class $\text{Th} \in H^{n+2}(E *_P \widehat{E})$. In these terms, f carries this to the map $f : P \rightarrow R$ representing the pair $(E, i_1^*(\text{Th}))$, where $i_1 : E \rightarrow E *_P \widehat{E}$ is the natural inclusion.

Write $X_b \cong BG \times K(\mathbb{Z}, n+2)$ for the fiber over b , and J_b for the fiber inclusion $J_b : X_b \rightarrow P$ in the top row of (4.10). Then the composition $f \circ J_b$ represents the pair $(J_b^*E, i_1^*J_b^*(\text{Th}))$ over X_b . However, J_b^*E is trivial, since its classifying map factors through $\{b\}$. Therefore, the fiberwise join $J_b^*(E *_P \widehat{E}) = \Sigma_{X_b}^{q+1} \widehat{E}$ is the $(q+1)$ -th fiberwise suspension of $\widehat{E}$. Then the pullback of the bundle Thom class $i_1^*J_b^*(\text{Th})$ is the composite

$$S^q \times BG \times K(\mathbb{Z}, n+2) = S^q \times X_b \cong J_b^*E \xrightarrow{i_1} \Sigma_{X_b}^{q+1} \widehat{E} \xrightarrow{J_b} E *_P \widehat{E} \xrightarrow{\text{Th}} K(\mathbb{Z}, n+2).$$

Now, the Euler class of the S^{n+2} -bundle $\Sigma_{X_b}^{q+1} \widehat{E}$ is precisely the restriction of the bundle Thom class $J_b^*\text{Th}$ along the zero section. Restricting along i_1 gives an S^q -parametrized form of this fact: $i_1 \circ J_b \circ \text{Th} : S^q \times X_b \rightarrow K(\mathbb{Z}, n+2)$ is the Euler class of $\Sigma_{X_b}^{q+1} \widehat{E}$ when restricted to any point in S^q . Via adjunction, this is precisely $\tilde{e}$ in the BG variable. $\square$

Now the following result provides special cases when T-duals exist and are unique.

Corollary 4.12. *If the Euler class map $e : BG \rightarrow K(\mathbb{Z}, q + 1)$ is m -connected, so too is the comparison map $f : P \rightarrow R$. Thus, over complexes of dimension less than m , there exists a unique T-dual $(\widehat{E}, \widehat{H})$ for any (E, H) . In dimension m , such duals exist, but are not necessarily unique.*

Proof. The result then follows from [Lemma 4.11](#) and the fact that for an H-space such as $K(\mathbb{Z}, n + 2)$, the product of the constant maps and based maps gives an equivalence $K(\mathbb{Z}, n + 2) \times \Omega^q K(\mathbb{Z}, n + 2) \simeq \text{Map}(S^q, K(\mathbb{Z}, n + 2))$. $\square$

Example 4.13 (principal circle bundles). When $n = q = 1$ and $G = U(1)$, the Euler class $e : BU(1) \rightarrow K(\mathbb{Z}, 2)$ is a weak homotopy equivalence; therefore, the map f is as well. This recovers the result of Bunke and Schick [\[2005\]](#) that in the case of circle bundles and $U(1)$ -gerbes, T-dual pairs exist and are unique up to equivalence.

Example 4.14 (principal $SU(2)$ -bundles). Suppose that $n = 5$, i.e., $q = 3$, and $G = SU(2)$, and consider pairs (E, H) consisting of a principal $SU(2)$ -bundle $\pi : E \rightarrow X$ and a 5-gerbe $H : E \rightarrow K(\mathbb{Z}, 7)$ on E . The Euler class $e : BSU(2) \rightarrow K(\mathbb{Z}, 4)$ is 5-connected, and so the same holds for f . This recovers the result of Bouwknegt et al. [\[2015a\]](#) that when the base X has $\dim \leq 4$, the T-dual of (E, H) exists and is unique.

We can, furthermore, lift this statement to one of our newly constructed theories. Still when $n = 5$, the spectrum $\mathfrak{a}_5 = K^{(4)}(ku)$ is the fourth iterated algebraic K -theory of ku and $\mathfrak{A}_5 = K^{(4)}(ku)[\beta_5^{-1}]$ is the localization at the higher Bott element β_5 of degree 6. The map

$$\tau_5 : K(\mathbb{Z}, 7) \rightarrow BGL_1 K^{(4)}(ku)[\beta_5^{-1}] = BGL_1 \mathfrak{A}_5$$

defines a twisting of $\mathfrak{A}_5$ by 5-gerbes and we now consider the T-duality isomorphism. Since the total space of the universal principal $SU(2)$ -bundle is contractible, the vertical tangent bundle of E is trivializable. Hence, the associated twist τ_E of $\mathfrak{A}_5$ is trivial and the T-duality isomorphism takes the form

$$T = \hat{p}_! \circ \Lambda \circ p^* : \mathfrak{A}_5^*(E, H) \rightarrow \mathfrak{A}_5^{*-3}(\widehat{E}, \widehat{H}).$$

By [Corollary 3.8](#), the Chern character throws this T-duality isomorphism onto a T-duality isomorphism

$$T : H\mathbb{Q}^*(E; H)[\beta_5^{-1}] \xrightarrow{\cong} H\mathbb{Q}^{*-3}(\widehat{E}; \widehat{H})[\beta_5^{-1}]$$

in ordinary cohomology. This recovers the T-duality isomorphism for principal $SU(2)$ -bundles studied by Bouwknegt et al. [\[2015a, Theorem 5.2\]](#).

Example 4.15 (nonprincipal $SU(2)$ -bundles). We again suppose that $n = 5$, i.e., $q = 3$, and that $G = SO(4)$ acts on S^3 via the inclusion

$$SO(4) = \text{Isom}^+(SU(2)) \rightarrow \text{Homeo}^+(SU(2)) \cong \text{Homeo}^+(S^3).$$

In this case, the map on fibers in the map of fiber sequences comparing P and R (4.10) is

$$\tilde{e} \cdot \text{const} : B SO(4) \times K(\mathbb{Z}, 7) \rightarrow \text{Map}(S^3, K(\mathbb{Z}, 7)) \simeq K(\mathbb{Z}, 4) \times K(\mathbb{Z}, 7).$$

The effect on π_4 is the projection $+: \mathbb{Z} \oplus \mathbb{Z} \rightarrow \mathbb{Z}$. This means that over S^4 , each pair (E, H) of an S^3 bundle with structure group $SO(4)$ and a 5-gerbe H has an infinite rank-one lattice of possible T-duals $(\widehat{E}, \widehat{H})$. In particular, spherical T-duals are not unique in this case. This analysis recovers the construction of spherical T-duals given by Bouwknegt et al. [2015b] in the case of $M = S^4$. They construct an infinite family of T-duals over any compact, oriented, simply connected, 4-dimensional manifold M , but in this case the classification of all possible T-dual pairs is a more intricate problem. Given a T-dual pair (E, H) and $(\widehat{E}, \widehat{H})$, we have a T-duality isomorphism in $\mathfrak{A}_5$ -cohomology and Bott-inverted rational cohomology as in the previous example.

Example 4.16 (nonorientable bundles). Baraglia [2014] extended topological T-duality, as an isomorphism in twisted K -theory, to the case of general circle bundles. Via a geometric approach, he proves existence and uniqueness of T-duals. This requires considering a twist using real line bundles, i.e., arising from the factor $K(\mathbb{Z}/2, 1) \subset BGL_1(K)$, in addition to the usual determinantal factor $K(\mathbb{Z}, 3)$. Specifically: the map $O(2) \rightarrow \text{Homeo}(S^1)$ is a homotopy equivalence (this is a theorem of Kneser [1926]), so a general S^1 -bundle fails to be equivalent to a principal bundle only on the basis of orientability.

This was put in the homotopy-theoretic framework of Bunke and Schick by Mathai and Rosenberg [2014, Theorem 2.2], who constructed a classifying space for T-dual pairs in the not necessarily oriented setting. When we defined the classifying space $P_n(G)$, we needed our sphere bundles to be oriented to define the Euler class. We may extend Mathai and Rosenberg's construction of the classifying space for T-dual pairs to the case of higher dimensional sphere bundles by considering not necessarily oriented S^q -bundles E and $\widehat{E}$ with structure group G over a space X . Let $\tilde{X}$ be the double cover of X which orients $E *_X \widehat{E}$, and Q the oriented S^{n+1} -bundle over $\tilde{X}$ to which $E *_X \widehat{E}$ pulls back. One may show that triples $(E, \widehat{E}, \text{Th})$, where $\text{Th} \in H^{n+2}(Q)$, are classified by a space $\tilde{P}_n(G)$, which is defined as the homotopy fiber of the composite

$$BG \times BG \xrightarrow{B \text{ join}} B \text{Homeo}(S^q * S^q) \xrightarrow{(w_1, e)} E\mathbb{Z}/2 \times_{\mathbb{Z}/2} K(\mathbb{Z}, n+3).$$

Here $\mathbb{Z}/2$ acts on $K(\mathbb{Z}, n+3)$ by negating the fundamental class. The Borel construction fibers over $K(\mathbb{Z}/2, 1)$, with fiber $K(\mathbb{Z}, n+3)$; (w_1, e) is the data of the first Stiefel–Whitney class of the $S^q * S^q$ bundle $E *_X \widehat{E}$ (i.e., w_1 maps into $K(\mathbb{Z}/2, 1)$), and e is the Euler class of the oriented cover Q .

The construction of the classifying space $R_n(G)$ for pairs (E, H) is applicable in the case where E is a not necessarily oriented sphere bundle with structure group G . As in the oriented case, there is a comparison map $f : \widetilde{P}_n(G) \rightarrow R_n(G)$, and we may study its connectivity to determine the existence and uniqueness of T-dual pairs in this setting. In the case where $n = q = 1$ and $G = O(2)$, the map f is an equivalence, so we recover Baraglia’s theorem on the existence and uniqueness of T-duality for nonprincipal circle bundles.

Our proof of the T-duality isomorphism applies to nonorientable bundles: looking back to the definition of what it means for (E, H) and $(\widehat{E}, \widehat{H})$ to be T-dual, we allow E and $\widehat{E}$ to be arbitrary smooth bundles with fiber S^q . This is enough to prove the T-duality isomorphism; indeed, the orientation twist τ_E precisely carries the correction term for nonorientable sphere bundles.

Lastly, we recall Smale’s conjecture that the inclusion $O(q+1) \rightarrow \text{Diff}(S^q)$ is a homotopy equivalence; this has been verified by Smale [1959] for $q = 2$ and Hatcher [1983] for $q = 3$. These results allow us to extend to dimensions two and three the results described above for S^1 -bundles, yielding a similar construction and analysis of the classifying space $\widetilde{P}_n(G)$ of pairs (E, H) , where E is a not necessarily orientable S^q -bundle with structure group G , and $H \in H^{n+2}(E)$.

5. Higher categories, n -vector spaces, and iterated algebraic K -theory

In this section we return to the question raised in the introduction on the relationship between the iterated algebraic K -theory $\mathfrak{a}_n = K^{(n-1)}(ku)$, the Bott-inverted iterated algebraic K -theory $\mathfrak{A}_n = K^{(n-1)}(ku)[\beta_n^{-1}]$, and n -vector spaces. The discussion is entirely speculative and is independent of the results proved in the paper. Our aim is to sketch a conceptual framework that might be useful for analyzing the geometric content of T-duality for iterated algebraic K -theory. Making this material rigorous would, at a minimum, require a good theory of weighted colimits for enriched ∞ -categories. Also, for the sake of clarity, we ignore the size issues that arise.

We start with the concept that one should be able to associate an object like a vector bundle to a principal $K(\mathbb{Z}, n)$ -bundle. We call such an object a (rank-one) n -vector bundle. Often the structure group $K(\mathbb{Z}, n)$ is written $B^{n-1}U(1)$ to highlight the categorical level: the notion of an n -vector bundle is an $(n-1)$ -fold categorification of the notion of a vector bundle. The association of an n -vector bundle to a principal $K(\mathbb{Z}, n)$ -bundle works similarly to the classical theory, via an action of the group $K(\mathbb{Z}, n)$ on some structure. We take the point of view that the

canonical object to act on is the linear $(n-1)$ -category $(n-1)\mathcal{Vect}_{\mathbb{C}}$ of $(n-1)$ -vector spaces. The iterative definition of n -vector space given below is arranged so that the $(n-1)$ -category $(n-1)\mathcal{Vect}_{\mathbb{C}}$ is the basic example of a rank-one n -vector space.

For small values of n this is completely rigorous. When $n = 1$ one has the canonical action of $K(\mathbb{Z}, 1) = U(1)$ on the complex vector space $\mathbb{C}$. When $n = 2$ the group $K(\mathbb{Z}, 2) = BU(1)$ is equivalent to the Picard groupoid of the category of vector spaces, i.e., to the symmetric monoidal groupoid of 1-dimensional complex vector spaces under the tensor product. As such, $BU(1)$ acts on the category $\mathcal{Vect}_{\mathbb{C}}$ of finite-rank complex vector spaces via the tensor product. So to a $BU(1)$ -principal bundle is canonically associated a bundle with fiber $\mathcal{Vect}_{\mathbb{C}}$. The 2-category $2\mathcal{Vect}_{\mathbb{C}}$ of 2-vector spaces is defined to be the 2-category of finite-rank $\mathcal{Vect}_{\mathbb{C}}$ -modules in $\mathbb{C}$ -linear categories, as in [Kapranov and Voevodsky 1994]. Since $\mathcal{Vect}_{\mathbb{C}}$ is a rank-one object in $2\mathcal{Vect}_{\mathbb{C}}$, the bundle associated to a principal $K(\mathbb{Z}, 2)$ -bundle is a line 2-vector bundle.

The first few values of n have explicit applications [Kapranov and Voevodsky 1994; Sati et al. 2009; Freed et al. 2010]. Intuitively, the pattern continues to higher n , but the situation becomes successively more complicated and less clear as n grows.

We will sketch an approach to n -vector spaces using enriched higher categories. If $\mathcal{V}$ is a monoidal ∞ -category, let us write $\mathcal{Cat}_{\infty}^{\mathcal{V}}$ for the $(\infty, 1)$ -category of $\mathcal{V}$ -enriched ∞ -categories and $\mathcal{V}$ -functors. Although our arguments are merely heuristics, much of what we say can be implemented in the model developed by Gepner and Haugseng [2015]. For example, $\mathcal{Cat}_{\infty}^{\mathcal{Sp}}$ denotes the ∞ -category of ∞ -categories enriched in the monoidal ∞ -category $\mathcal{Sp}$ of spectra. Since $\mathcal{Sp}$ carries the additional structure of a symmetric monoidal ∞ -category, the ∞ -category $\mathcal{Cat}_{\infty}^{\mathcal{Sp}}$ is also symmetric monoidal. We can iterate this procedure further. Let

$$\mathcal{Cat}_{(\infty, n)}^{\mathcal{Sp}} = \mathcal{Cat}_{\infty}^{\mathcal{Cat}_{\infty}^{\mathcal{Sp}} \cdots \mathcal{Cat}_{\infty}^{\mathcal{Sp}}}$$

be the ∞ -category of ∞ -categories enriched in $\mathcal{Cat}_{(\infty, n-1)}^{\mathcal{Sp}}$, where $\mathcal{Cat}_{(\infty, 1)}^{\mathcal{Sp}} = \mathcal{Cat}_{\infty}^{\mathcal{Sp}}$ [Gepner and Haugseng 2015, Remark 5.7.13]. We think of objects of $\mathcal{Cat}_{(\infty, n)}^{\mathcal{Sp}}$ as (∞, n) -categories where the collection of n -morphisms between a pair of $(n-1)$ -morphisms forms a spectrum.

There is not currently a well developed theory of weighted limits and weighted colimits in enriched ∞ -categories, but we proceed as if there were, using the corresponding notions from ordinary enriched category theory as a guide for our intuition. We write $\mathcal{St}_{\infty}^{(n)}$ for the full subcategory of $\mathcal{Cat}_{(\infty, n)}^{\mathcal{Sp}}$ consisting of those objects admitting all $\mathcal{Cat}_{(\infty, n-1)}^{\mathcal{Sp}}$ -weighted colimits. Assuming that this is a sensible notion, there is a localization adjunction

$$\mathcal{Cat}_{(\infty, n)}^{\mathcal{Sp}} \xrightleftharpoons{L^{(n)}} \mathcal{St}_{\infty}^{(n)} \quad (5.1)$$

where the left adjoint freely adds all such colimits and the right adjoint is the inclusion. The case $n = 1$ was studied by Blumberg, Gepner, and Tabuada [Blumberg et al. 2013] and takes the form

$$\mathcal{Cat}_\infty^{\text{Sp}} \xrightleftharpoons{L^{(1)}} \mathcal{St}_\infty,$$

where $\mathcal{St}_\infty = \mathcal{St}_\infty^{(1)}$ is the full subcategory of $\mathcal{Cat}_\infty^{\text{Sp}}$ consisting of those spectrally enriched ∞ -categories admitting small Sp -weighted colimits, i.e., the cocomplete stable ∞ -categories. This explains the notation $\mathcal{St}_\infty^{(n)}$.

Suppose that R is an E_∞ ring spectrum. There is an object $\mathfrak{b}R$ of $\mathcal{Cat}_\infty^{\text{Sp}}$ with a single object $\bullet_0$ whose endomorphism spectrum is $\text{Hom}(\bullet_0, \bullet_0) = R$. The image of the object $\mathfrak{b}R$ under the localization functor $L^{(1)}$ is the presentable stable ∞ -category

$$L^{(1)}\mathfrak{b}R = \text{Mod}_R$$

of R -modules. The functor $L^{(1)}$ admits a symmetric monoidal structure and we write $(\text{Mod}_R, \otimes)$ for the induced symmetric monoidal structure on the image of $\mathfrak{b}R$. For the purposes of defining K -theory, we restrict to the full subcategory $(\text{Mod}_R, \otimes)^\circ$ of dualizable R -modules under the symmetric monoidal structure. It is here that we use the commutativity assumption. We could equivalently restrict to the subcategory of perfect R -modules, meaning the objects which span the thick subcategory generated by R after passage to the homotopy category. It is the dualizability notion that we will generalize below, so we concentrate on that here.

The $(\infty, 1)$ -category $(\text{Mod}_R, \otimes)^\circ$ is pointed and admits finite colimits, so we may use the ∞ -categorical version of Waldhausen’s K -theory construction [Blumberg et al. 2013; Barwick 2016] to define its algebraic K -theory:

$$K(R) = K(\text{Mod}_R) := K((L^{(1)}\mathfrak{b}R, \otimes)^\circ).$$

Notice that “ $K(R)$ ” and “ $K(\text{Mod}_R)$ ” are synonyms. Soon we will consider the K -theory of other module categories, but in that case we will never write the K -theory in terms of the underlying “ring”, only in terms of the category used to construct the K -theory.

When $n = 2$, adjunction (5.1) takes the form

$$\mathcal{Cat}_{(\infty, 2)}^{\text{Sp}} = \mathcal{Cat}_\infty^{\mathcal{Cat}_\infty^{\text{Sp}}} \xrightleftharpoons{L^{(2)}} \mathcal{St}_\infty^{(2)}$$

where the left adjoint $L^{(2)}$ freely adjoins all $\mathcal{Cat}_\infty^{\text{Sp}}$ -weighted colimits. There is an object $\mathfrak{b}^2 R$ of $\mathcal{Cat}_\infty^{\mathcal{Cat}_\infty^{\text{Sp}}}$ which has a single object $\bullet_0$ and a single 1-morphism $\bullet_1$ whose endomorphism spectrum is $\text{Hom}(\bullet_1, \bullet_1) = R$. The image of $\mathfrak{b}^2 R$ under the

localization functor $L^{(2)}$ is the $(\infty, 2)$ -category

$$L^{(2)}\mathfrak{b}^2 R = \mathrm{Mod}_{\mathrm{Mod}_R}$$

of Mod_R -module categories. An object of $\mathrm{Mod}_{\mathrm{Mod}_R}$ may be described as a presentable stable ∞ -category $\mathcal{C}$ equipped with a suitable action $\mathrm{Mod}_R \otimes \mathcal{C} \rightarrow \mathcal{C}$ of Mod_R , defined with respect to the symmetric monoidal structure $\otimes$ on stable presentable ∞ -categories. When considering dualizable Mod_R -modules, as we will do below, it is natural to restrict to the action of the subcategory Mod_R° of perfect R -modules. If $\mathcal{A}$ and $\mathcal{B}$ are Mod_R -module categories, the $(\infty, 1)$ -category of morphisms from $\mathcal{A}$ to $\mathcal{B}$ is the category of exact R -linear functors

$$\mathrm{Mod}_{\mathrm{Mod}_R}(\mathcal{A}, \mathcal{B}) \simeq \mathrm{Fun}_R^{\mathrm{ex}}(\mathcal{A}, \mathcal{B}).$$

So far, the constructions we have made would still make sense if R were an E_2 ring spectrum, but as in the $n = 1$ case, we now restrict to the full subcategory spanned by the dualizable objects. The spectral $(\infty, 2)$ -category $\mathrm{Mod}_{\mathrm{Mod}_R}$ inherits a symmetric monoidal structure by virtue of the symmetric monoidal structure on the functor $L^{(2)}$. We write $(L^{(2)}\mathfrak{b}^2 R, \otimes)^\circ \rightarrow L^{(2)}\mathfrak{b}^2 R$ for the inclusion of the subcategory spanned by the fully dualizable objects, as defined by Lurie [2009b, Claim 2.3.19]. This has the effect of discarding 1-morphisms that do not admit adjoints in the underlying homotopy bicategory and then discarding objects that are not dualizable in the underlying symmetric monoidal homotopy category. We define the K -theory of Mod_R -module categories to be the Waldhausen K -theory of the $(\infty, 1)$ -category truncation of the $(\infty, 2)$ -category of the fully dualizable objects of $\mathrm{Mod}_{\mathrm{Mod}_R}$

$$K(\mathrm{Mod}_{\mathrm{Mod}_R}) := K(\iota_1(L^{(2)}\mathfrak{b}^2 R, \otimes)^\circ).$$

In the case of symmetric monoidal bicategories with no higher morphisms, the analogous K -theory functor was constructed explicitly by Osorno [2012].

When $R = Hk$ is the Eilenberg–MacLane spectrum associated to a field k , the $(\infty, 2)$ -category $\mathrm{Mod}_{\mathrm{Mod}_{Hk}}$ is equivalent to the $(\infty, 2)$ -category of k -linear dg-categories. The fully dualizable objects of $\mathrm{Mod}_{\mathrm{Mod}_{Hk}}$ are precisely the smooth and proper dg-categories [Toën and Vaquié 2007; Antieau and Gepner 2014], and $\pi_0 K(\mathrm{Mod}_{\mathrm{Mod}_{Hk}})$ is isomorphic to the secondary K -theory $K_0^{[2]}(k)$ of the field k defined by Toën [2011; Tabuada 2016]. We also expect that $K(\mathrm{Mod}_{\mathrm{Mod}_{Hk}})$ is related to the K -theory of varieties studied by Campbell, Wolfson, and Zakharevich [Campbell 2019; Campbell et al. 2019].

Remark 5.2. A slight variation allows us to consider 2-vector spaces instead of dg-categories. We replace Mod_R with the category $\mathrm{Vect}_{\mathbb{C}}$ of finite-dimensional complex vector spaces, considered as an $(\infty, 1)$ -category enriched in spectra by de-looping the usual enrichment in topological abelian groups. In this case, $\mathrm{Mod}_{\mathrm{Vect}_{\mathbb{C}}}$

is the $(\infty, 2)$ -category of $\mathcal{V}\text{ect}_{\mathbb{C}}$ -module categories. In particular, the 2-category $2\mathcal{V}\text{ect}_{\mathbb{C}}$ of 2-vector spaces, as defined by Kapranov and Voevodsky [1994], embeds in the subcategory $(\text{Mod}_{\mathcal{V}\text{ect}_{\mathbb{C}}}, \otimes)^{\circ}$ of dualizable objects, and $K(\text{Mod}_{\mathcal{V}\text{ect}_{\mathbb{C}}})$ is equivalent to the bicategorical K -theory of 2-vector spaces $K(2\mathcal{V}\text{ect}_{\mathbb{C}})$ studied by Baas, Dundas, and Rognes [Baas et al. 2004; Osorno 2012].

Instead of applying $L^{(2)}$ to $\mathfrak{b}^2 R$, we could add colimits in a two-step process. Write $\text{Cat}_{\infty}^{\text{St}\infty}$ for the ∞ -category of ∞ -categories enriched in stable ∞ -categories. There is a localization adjunction

$$\text{Cat}_{\infty}^{\text{Cat}_{\infty}^{\text{Sp}}} \xrightleftharpoons{\mathfrak{b}L^{(1)}} \text{Cat}_{\infty}^{\text{St}\infty}$$

where the left adjoint applies $L^{(1)}$ to the hom objects. In other words, we've only added colimits at the top level. In the case of $\mathfrak{b}^2 R$, we get the category

$$(\mathfrak{b}L^{(1)})(\mathfrak{b}^2 R) = \mathfrak{b}\text{Mod}_R$$

with a single 0-cell whose endomorphism object is the stable ∞ -category of R -modules. Notice that composition in this category uses the symmetric monoidal structure $\otimes$ on Mod_R . Next, we apply $K(\cdot)$, taking care to first restrict to the dualizable objects. Formally, we are using the fact that $\mathfrak{b}R$ is a commutative monoid in $\text{Cat}_{\infty}^{\text{Sp}}$ and that the functors $L^{(1)}$ and K induce functors on categories enriched in commutative monoid objects. We have now associated to $\mathfrak{b}^2 R$ the spectral category

$$\mathfrak{b}K(R) = (\mathfrak{b}K)(\iota_1((\mathfrak{b}L^{(1)})\mathfrak{b}^2 R, \otimes)^{\circ})$$

with one object and $K(R)$ as its endomorphism spectrum. Here ι_1 denotes the truncation functor $\mathfrak{b}\iota_0$ taking an $(\infty, 2)$ -category to the underlying $(\infty, 1)$ -category that removes the noninvertible 2-morphisms. After applying the same procedure one categorical level down by adding all spectral colimits, restricting to dualizable objects and taking K -theory, we get the twice-iterated algebraic K -theory spectrum

$$K(K(R)) = K((L^{(1)}\mathfrak{b}K(R), \otimes)^{\circ}).$$

Both the K -theory of Mod_R -modules $K(\text{Mod}_{\text{Mod}_R})$ and the iterated K -theory $K(K(R))$ map to the spectral enhancement $K^{\{2\}}(R)$ of Toën's secondary K -theory defined by Hoyois, Scherotzke, and Sibilla [Hoyois et al. 2017], as we now explain. They consider a variant of the Blumberg–Gepner–Tabuada category of noncommutative motives [Blumberg et al. 2013] that is the recipient of the universal additive invariant of stable Mod_R° -module categories. More precisely, there is a functor

$$\mathcal{U} : \langle \text{Mod}_R^{\circ}\text{-modules in } \text{Cat}_{\infty}^{\text{perf}} \rangle \rightarrow \text{Mot}_R$$

from the ∞ -category of small, stable, idempotent complete Mod_R° -module categories to the ∞ -category of R -linear motives that preserves filtered colimits, preserves zero objects, sends split exact sequences to cofiber sequences, and is universal among functors with these three properties. The $(\infty, 1)$ -category Mot_R and the functor $\mathcal{U}$ admit compatible symmetric monoidal structures, and we define the K -theory of Mot_R to be the Waldhausen K -theory of the full subcategory of Mot_R generated under finite colimits and retracts by the images of dualizable objects under $\mathcal{U}$:

$$K(\mathrm{Mot}_R) := K((\mathrm{Mot}_R, \otimes)^\circ).$$

K -theory becomes corepresentable after passing to the ∞ -category Mot_R , meaning that if $\mathcal{A}$ and $\mathcal{B}$ are Mod_R° -module categories, and $\mathcal{A}$ is compact, then there is a natural equivalence of spectra

$$\mathrm{Mot}_R(\mathcal{U}(\mathcal{A}), \mathcal{U}(\mathcal{B})) \simeq K(\mathrm{Fun}_R^{\mathrm{ex}}(\mathcal{A}, \mathcal{B})).$$

When $\mathcal{A}$ and $\mathcal{B}$ are both fully dualizable as Mod_R -module categories, which in particular implies that they are compact [Hoyois et al. 2017, Proposition 4.19], then their associated motives are dualizable in Mot_R . Applying the natural transformation $\iota_0 \rightarrow K$ from the groupoid core functor to Waldhausen K -theory to the spectrally enriched $(\infty, 1)$ -categories $\mathrm{Fun}_R^{\mathrm{ex}}(\mathcal{A}, \mathcal{B})$ gives a map

$$\iota_1 \mathrm{Mod}_{\mathrm{Mod}_R}(\mathcal{A}, \mathcal{B}) \simeq \iota_0 \mathrm{Fun}_R^{\mathrm{ex}}(\mathcal{A}, \mathcal{B}) \rightarrow K(\mathrm{Fun}_R^{\mathrm{ex}}(\mathcal{A}, \mathcal{B})) \simeq \mathrm{Mot}_R(\mathcal{U}(\mathcal{A}), \mathcal{U}(\mathcal{B}))$$

which assembles into a symmetric monoidal functor of $(\infty, 1)$ -categories

$$\iota_1(\mathrm{Mod}_{\mathrm{Mod}_R})^\circ \rightarrow \mathrm{Mot}_R^\circ.$$

Applying Waldhausen K -theory gives a map of E_∞ ring spectra

$$K(\mathrm{Mod}_{\mathrm{Mod}_R}) \rightarrow K(\mathrm{Mot}_R). \quad (5.3)$$

On the other hand, the motive associated to the category Mod_R° of perfect R -modules is the unit of the symmetric monoidal category Mot_R of R -linear motives, and so the thick subcategory that it generates is equivalent to the category of modules over the endomorphism ring

$$\mathrm{Mot}_R(\mathcal{U}(\mathrm{Mod}_R^\circ), \mathcal{U}(\mathrm{Mod}_R^\circ)) \simeq K(\mathrm{Mod}_R^\circ) = K(R).$$

The inclusion of this thick subcategory induces a map of E_∞ ring spectra

$$K(K(R)) \rightarrow K(\mathrm{Mot}_R), \quad (5.4)$$

as observed by Hoyois et al. [2017, Remark 6.23].

Question 5.5. To what extent do the images of the maps (5.3) and (5.4) coincide in $K(\mathrm{Mot}_R)$?

When $R = Hk$ is the Eilenberg–MacLane spectrum associated to a field k , the work of Baas, Dundas, Richter, and Rognes [Baas et al. 2011] implies that $K(K(k))$ is equivalent to the K -theory of the full subcategory of $\text{Mod}_{\text{Mod}_{Hk}}$ spanned by the free 2-vector spaces of finite rank. Equivalently, these are the dg-categories with a single object and the free algebras k^n as the ring of morphisms. Similarly, if we replace Mod_R with $\mathcal{V}\text{ect}_{\mathbb{C}}$, as in Remark 5.2, then $K(ku)$ is equivalent to the K -theory $K(2\mathcal{V}\text{ect}_{\mathbb{C}}) \simeq K(\text{Mod}_{\mathcal{V}\text{ect}_{\mathbb{C}}})$ of 2-vector spaces.

We would like to ask a similar question about the iterated algebraic K -theory spectrum $K^{(n)}(R)$ for higher values of n . Let $\mathfrak{b}^n R \in \mathcal{C}\text{at}_{(\infty, n)}^{\text{Sp}}$ be the (∞, n) -category enriched in spectra with a single k -morphism for $0 \leq k \leq n-1$ and for which the endomorphism spectrum of the $(n-1)$ -morphism is the ring R . Let

$$\text{Mod}_R^{(n)} = L^{(n)} \mathfrak{b}^n R = \text{Mod}_{\text{Mod} \dots \text{Mod}_R}$$

denote the image of $\mathfrak{b}^n R$ under the localization functor $L^{(n)}$. The fully dualizable part

$$\text{Mod}_{\text{Mod} \dots \text{Mod}_R}^{\circ} = (\text{Mod}_R^{(n)}, \otimes)^{\circ}$$

of the symmetric monoidal (∞, n) -category $\text{Mod}_R^{(n)}$ is our higher categorical analog of the category of perfect R -modules, and in the case of $R = Hk$ we consider this to be a reasonable definition of the (∞, n) -category of n -chain complexes over k . Similarly, we define the (∞, n) -category of complex n -vector spaces to be the symmetric monoidal (∞, n) -category

$$n\mathcal{V}\text{ect}_{\mathbb{C}} = (L^{(n)} \mathfrak{b}^{n-1} \mathcal{V}\text{ect}_{\mathbb{C}}, \otimes)^{\circ} = \text{Mod}_{\text{Mod} \dots \mathcal{V}\text{ect}_{\mathbb{C}}}^{\circ}.$$

Remark 5.6. We do not know how to construct the appropriate analog of the category of noncommutative R -local motives Mot_R that would accept maps from the spectra $K^{(n)}(R)$ and $K((\text{Mod}_R^{(n)}, \otimes)^{\circ})$, and we do not know how to ask the analog of Question 5.5 when $n > 2$ (compare with [Hoyois et al. 2017, Remark 6.22]). Our results on T-duality for $\mathfrak{A}_n$ -algebras suggest that a good understanding of the relationship between $K^{(n)}(R)$ and $K((\text{Mod}_R^{(n)}, \otimes)^{\circ})$ would connect T -duality for S^q -bundles and the theory of n -vector spaces for $n = 2q + 1$.

Acknowledgements

We thank Ben Antieau, Justin Noel, and Thomas Schick for helpful comments on a previous draft and Drew Heard for pointing us to the paper of Arthan [1983]. We also thank Rune Haugseng for sharing his perspectives on higher category theory and iterated K -theory with us. Lind was supported in part by the DFG through SFB1085. Sati thanks the Erwin Schrödinger Institute for Mathematical Physics, Vienna, for hospitality and the organizers of the program “Higher Structures in

String Theory” for the opportunity to present the results of this project. Westerland was supported in part by the NSF through DMS-1406162.

References

- [Adams 1972] J. F. Adams, *Algebraic topology: a student’s guide*, London Math. Soc. Lecture Note Ser. **4**, Cambridge Univ., 1972. [MR](#) [Zbl](#)
- [Ando et al. 2010] M. Ando, A. J. Blumberg, and D. Gepner, “Twists of K -theory and TMF ”, pp. 27–63 in *Superstrings, geometry, topology, and C^* -algebras*, edited by R. S. Doran et al., Proc. Sympos. Pure Math. **81**, Amer. Math. Soc., Providence, RI, 2010. [MR](#) [Zbl](#)
- [Ando et al. 2014a] M. Ando, A. J. Blumberg, D. Gepner, M. J. Hopkins, and C. Rezk, “An ∞ -categorical approach to R -line bundles, R -module Thom spectra, and twisted R -homology”, *J. Topol.* **7**:3 (2014), 869–893. [MR](#) [Zbl](#)
- [Ando et al. 2014b] M. Ando, A. J. Blumberg, D. Gepner, M. J. Hopkins, and C. Rezk, “Units of ring spectra, orientations and Thom spectra via rigid infinite loop space theory”, *J. Topol.* **7**:4 (2014), 1077–1117. [MR](#) [Zbl](#)
- [Ando et al. 2018] M. Ando, A. J. Blumberg, and D. Gepner, “Parametrized spectra, multiplicative Thom spectra and the twisted Umkehr map”, *Geom. Topol.* **22**:7 (2018), 3761–3825. [MR](#) [Zbl](#)
- [Antieau and Gepner 2014] B. Antieau and D. Gepner, “Brauer groups and étale cohomology in derived algebraic geometry”, *Geom. Topol.* **18**:2 (2014), 1149–1244. [MR](#) [Zbl](#)
- [Antieau et al. 2014] B. Antieau, D. Gepner, and J. M. Gómez, “Actions of $K(\pi, n)$ spaces on K -theory and uniqueness of twisted K -theory”, *Trans. Amer. Math. Soc.* **366**:7 (2014), 3631–3648. [MR](#) [Zbl](#)
- [Arthan 1983] R. D. Arthan, “Localization of stable homotopy rings”, *Math. Proc. Cambridge Philos. Soc.* **93**:2 (1983), 295–302. [MR](#) [Zbl](#)
- [Ausoni 2010] C. Ausoni, “On the algebraic K -theory of the complex K -theory spectrum”, *Invent. Math.* **180**:3 (2010), 611–668. [MR](#) [Zbl](#)
- [Ausoni and Rognes 2002] C. Ausoni and J. Rognes, “Algebraic K -theory of topological K -theory”, *Acta Math.* **188**:1 (2002), 1–39. [MR](#) [Zbl](#)
- [Ausoni and Rognes 2008] C. Ausoni and J. Rognes, “The chromatic red-shift in algebraic K -theory”, *Enseign. Math.* **54**:1–2 (2008), 13–15.
- [Ausoni and Rognes 2012] C. Ausoni and J. Rognes, “Rational algebraic K -theory of topological K -theory”, *Geom. Topol.* **16**:4 (2012), 2037–2065. [MR](#) [Zbl](#)
- [Ausoni et al. 2008] C. Ausoni, B. r. I. Dundas, and J. Rognes, “Divisibility of the Dirac magnetic monopole as a two-vector bundle over the three-sphere”, *Doc. Math.* **13** (2008), 795–801. [MR](#) [Zbl](#)
- [Baas et al. 2004] N. A. Baas, B. r. I. Dundas, and J. Rognes, “Two-vector bundles and forms of elliptic cohomology”, pp. 18–45 in *Topology, geometry and quantum field theory*, edited by U. Tillmann, London Math. Soc. Lecture Note Ser. **308**, Cambridge Univ., 2004. [MR](#) [Zbl](#)
- [Baas et al. 2011] N. A. Baas, B. r. I. Dundas, B. Richter, and J. Rognes, “Stable bundles over rigid categories”, *J. Topol.* **4**:3 (2011), 623–640. [MR](#) [Zbl](#)
- [Baraglia 2014] D. Baraglia, “Topological T-duality for general circle bundles”, *Pure Appl. Math. Q.* **10**:3 (2014), 367–438. [MR](#) [Zbl](#)
- [Barwick 2016] C. Barwick, “On the algebraic K -theory of higher categories”, *J. Topol.* **9**:1 (2016), 245–347. [MR](#) [Zbl](#)

- [Blumberg et al. 2013] A. J. Blumberg, D. Gepner, and G. Tabuada, “A universal characterization of higher algebraic K -theory”, *Geom. Topol.* **17**:2 (2013), 733–838. [MR](#) [Zbl](#)
- [Blumberg et al. 2014] A. J. Blumberg, D. Gepner, and G. Tabuada, “Uniqueness of the multiplicative cyclotomic trace”, *Adv. Math.* **260** (2014), 191–232. [MR](#) [Zbl](#)
- [Bökstedt et al. 1993] M. Bökstedt, W. C. Hsiang, and I. Madsen, “The cyclotomic trace and algebraic K -theory of spaces”, *Invent. Math.* **111**:3 (1993), 465–539. [MR](#) [Zbl](#)
- [Bouwknegt et al. 2004] P. Bouwknegt, J. Evslin, and V. Mathai, “ T -duality: topology change from H -flux”, *Comm. Math. Phys.* **249**:2 (2004), 383–415. [MR](#) [Zbl](#)
- [Bouwknegt et al. 2015a] P. Bouwknegt, J. Evslin, and V. Mathai, “Spherical T -duality”, *Comm. Math. Phys.* **337**:2 (2015), 909–954. [MR](#) [Zbl](#)
- [Bouwknegt et al. 2015b] P. Bouwknegt, J. Evslin, and V. Mathai, “Spherical T -duality, II: An infinity of spherical T -duals for non-principal $SU(2)$ -bundles”, *J. Geom. Phys.* **92** (2015), 46–54. [MR](#) [Zbl](#)
- [Buhštaber 1970] V. M. Buhštaber, “The Chern–Dold character in cobordisms, I”, *Math. USSR Sb.* **12**:4 (1970), 573–594. [MR](#) [Zbl](#)
- [Bunke and Schick 2005] U. Bunke and T. Schick, “On the topology of T -duality”, *Rev. Math. Phys.* **17**:1 (2005), 77–112. [MR](#) [Zbl](#)
- [Campbell 2019] J. A. Campbell, “The K -theory spectrum of varieties”, *Trans. Amer. Math. Soc.* **371**:11 (2019), 7845–7884. [MR](#) [Zbl](#)
- [Campbell et al. 2019] J. Campbell, J. Wolfson, and I. Zakharevich, “Derived ℓ -adic zeta functions”, *Adv. Math.* **354** (2019), art. id. 106760. [MR](#) [Zbl](#)
- [Dundas et al. 2013] B. I. Dundas, T. G. Goodwillie, and R. McCarthy, *The local structure of algebraic K -theory*, Algebra and Applications **18**, Springer, 2013. [MR](#) [Zbl](#)
- [Freed et al. 2010] D. S. Freed, M. J. Hopkins, J. Lurie, and C. Teleman, “Topological quantum field theories from compact Lie groups”, pp. 367–403 in *A celebration of the mathematical legacy of Raoul Bott*, edited by P. R. Kotiuga, CRM Proc. Lecture Notes **50**, Amer. Math. Soc., Providence, RI, 2010. [MR](#) [Zbl](#)
- [Gepner and Haugseng 2015] D. Gepner and R. Haugseng, “Enriched ∞ -categories via non-symmetric ∞ -operads”, *Adv. Math.* **279** (2015), 575–716. [MR](#) [Zbl](#)
- [Gepner et al. 2015] D. Gepner, M. Groth, and T. Nikolaus, “Universality of multiplicative infinite loop space machines”, *Algebr. Geom. Topol.* **15**:6 (2015), 3107–3153. [MR](#) [Zbl](#)
- [Hatcher 1983] A. E. Hatcher, “A proof of the Smale conjecture, $\text{Diff}(S^3) \simeq O(4)$ ”, *Ann. of Math.* (2) **117**:3 (1983), 553–607. [MR](#) [Zbl](#)
- [Hoyois et al. 2017] M. Hoyois, S. Scherotzke, and N. Sibilla, “Higher traces, noncommutative motives, and the categorified Chern character”, *Adv. Math.* **309** (2017), 97–154. [MR](#) [Zbl](#)
- [Kapranov and Voevodsky 1994] M. M. Kapranov and V. A. Voevodsky, “2-categories and Zamolodchikov tetrahedra equations”, pp. 177–259 in *Algebraic groups and their generalizations: quantum and infinite-dimensional methods* (University Park, PA, 1991), edited by W. J. Haboush and B. J. Parshall, Proc. Sympos. Pure Math. **56**, Amer. Math. Soc., Providence, RI, 1994. [MR](#) [Zbl](#)
- [Kneser 1926] H. Kneser, “Die Deformationssätze der einfach zusammenhängenden Flächen”, *Math. Z.* **25**:1 (1926), 362–372. [MR](#) [Zbl](#)
- [Kragh 2013] T. Kragh, “Orientations on 2-vector bundles and determinant gerbes”, *Math. Scand.* **113**:1 (2013), 63–82. [MR](#) [Zbl](#)
- [Lind 2016] J. A. Lind, “Bundles of spectra and algebraic K -theory”, *Pacific J. Math.* **285**:2 (2016), 427–452. [MR](#) [Zbl](#)

- [Lurie 2009a] J. Lurie, *Higher topos theory*, Annals of Math. Studies **170**, Princeton Univ., 2009. [MR](#) [Zbl](#)
- [Lurie 2009b] J. Lurie, “On the classification of topological field theories”, pp. 129–280 in *Current developments in mathematics, 2008*, edited by D. Jerison et al., International, Somerville, MA, 2009. [MR](#) [Zbl](#)
- [Mathai and Rosenberg 2014] V. Mathai and J. Rosenberg, “T-duality for circle bundles via noncommutative geometry”, *Adv. Theor. Math. Phys.* **18**:6 (2014), 1437–1462. [MR](#) [Zbl](#)
- [May 2009] J. P. May, “What are E_∞ ring spaces good for?”, pp. 331–365 in *New topological contexts for Galois theory and algebraic geometry (BIRS 2008)*, edited by A. Baker and B. Richter, Geom. Topol. Monogr. **16**, Geom. Topol. Publ., Coventry, 2009. [MR](#) [Zbl](#)
- [May and Sigurdsson 2006] J. P. May and J. Sigurdsson, *Parametrized homotopy theory*, Math. Surveys and Monographs **132**, Amer. Math. Soc., Providence, RI, 2006. [MR](#) [Zbl](#)
- [Osorno 2012] A. M. Osorno, “Spectra associated to symmetric monoidal bicategories”, *Algebr. Geom. Topol.* **12**:1 (2012), 307–342. [MR](#) [Zbl](#)
- [Sati 2009] H. Sati, “A higher twist in string theory”, *J. Geom. Phys.* **59**:3 (2009), 369–373. [MR](#) [Zbl](#)
- [Sati 2010] H. Sati, “Geometric and topological structures related to M-branes”, pp. 181–236 in *Superstrings, geometry, topology, and C^* -algebras*, edited by R. S. Doran et al., Proc. Sympos. Pure Math. **81**, Amer. Math. Soc., Providence, RI, 2010. [MR](#) [Zbl](#)
- [Sati and Westerland 2015] H. Sati and C. Westerland, “Twisted Morava K -theory and E-theory”, *J. Topol.* **8**:4 (2015), 887–916. [MR](#) [Zbl](#)
- [Sati et al. 2009] H. Sati, U. Schreiber, and J. Stasheff, “ L_∞ -algebra connections and applications to String- and Chern–Simons n -transport”, pp. 303–424 in *Quantum field theory*, edited by B. Fauser et al., Birkhäuser, Basel, 2009. [MR](#) [Zbl](#)
- [Segal 1973] G. Segal, “The stable homotopy of complex projective space”, *Quart. J. Math. Oxford Ser. (2)* **24** (1973), 1–5. [MR](#) [Zbl](#)
- [Smale 1959] S. Smale, “Diffeomorphisms of the 2-sphere”, *Proc. Amer. Math. Soc.* **10** (1959), 621–626. [MR](#) [Zbl](#)
- [Snaith 1981] V. Snaith, “Localized stable homotopy of some classifying spaces”, *Math. Proc. Cambridge Philos. Soc.* **89**:2 (1981), 325–330. [MR](#) [Zbl](#)
- [Tabuada 2016] G. Tabuada, “A note on secondary K -theory”, *Algebra Number Theory* **10**:4 (2016), 887–906. [MR](#) [Zbl](#)
- [Toën 2011] B. Toën, “Lectures on dg-categories”, pp. 243–302 in *Topics in algebraic and topological K -theory*, edited by G. Cortiñas, Lecture Notes in Math. **2008**, Springer, 2011. [MR](#) [Zbl](#)
- [Toën and Vaquié 2007] B. Toën and M. Vaquié, “Moduli of objects in dg-categories”, *Ann. Sci. École Norm. Sup. (4)* **40**:3 (2007), 387–444. [MR](#) [Zbl](#)
- [Waldhausen 1979] F. Waldhausen, “Algebraic K -theory of topological spaces, II”, pp. 356–394 in *Algebraic topology* (Aarhus, Denmark, 1978), edited by A. Dold and B. Eckmann, Lecture Notes in Math. **763**, Springer, 1979. [MR](#) [Zbl](#)

Received 8 Dec 2017. Revised 12 Aug 2019. Accepted 28 Aug 2019.

JOHN A. LIND: john.alexander.lind@gmail.com

Department of Mathematics and Statistics, California State University, Chico, Chico, CA, United States

HISHAM SATI: hsati@nyu.edu

*Department of Mathematics, New York University Abu Dhabi, Saadiyat Island, Abu Dhabi,
United Arab Emirates*

CRAIG WESTERLAND: cwesterl@umn.edu

School of Mathematics, University of Minnesota, Minneapolis, MN, United States

ANNALS OF K-THEORY

msp.org/akt

EDITORIAL BOARD

Joseph Ayoub	ETH Zürich, Switzerland joseph.ayoub@math.uzh.ch
Paul Balmer	University of California, Los Angeles, USA balmer@math.ucla.edu
Guillermo Cortiñas	Universidad de Buenos Aires and CONICET, Argentina gcorti@dm.uba.ar
Hélène Esnault	Freie Universität Berlin, Germany liveesnault@math.fu-berlin.de
Eric Friedlander	University of Southern California, USA ericmf@usc.edu
Max Karoubi	Institut de Mathématiques de Jussieu – Paris Rive Gauche, France max.karoubi@imj-prg.fr
Moritz Kerz	Universität Regensburg, Germany moritz.kerz@mathematik.uni-regensburg.de
Huaxin Lin	University of Oregon, USA livehlin@uoregon.edu
Alexander Merkurjev	University of California, Los Angeles, USA merkurev@math.ucla.edu
Birgit Richter	Universität Hamburg, Germany birgit.richter@uni-hamburg.de
Jonathan Rosenberg	(Managing Editor) University of Maryland, USA jmr@math.umd.edu
Marco Schlichting	University of Warwick, UK schlichting@warwick.ac.uk
Charles Weibel	(Managing Editor) Rutgers University, USA weibel@math.rutgers.edu
Guoliang Yu	Texas A&M University, USA guoliangyu@math.tamu.edu

PRODUCTION

Silvio Levy (Scientific Editor)
production@msp.org

Annals of K-Theory is a journal of the [K-Theory Foundation](http://ktheoryfoundation.org) (ktheoryfoundation.org). The K-Theory Foundation acknowledges the precious support of [Foundation Compositio Mathematica](http://foundationcompositio.com), whose help has been instrumental in the launch of the Annals of K-Theory.

See inside back cover or msp.org/akt for submission instructions.

The subscription price for 2020 is US \$510/year for the electronic version, and \$575/year (+\$25, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Annals of K-Theory (ISSN 2379-1681 electronic, 2379-1683 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

AKT peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing
<http://msp.org/>

© 2020 Mathematical Sciences Publishers

ANNALS OF K-THEORY

2020

vol. 5

no. 1

Twisted iterated algebraic K -theory and topological T-duality for sphere bundles	1
JOHN A. LIND, HISHAM SATI and CRAIG WESTERLAND	
Hausdorffified algebraic K_1 -groups and invariants for C^* -algebras with the ideal property	43
GUIHUA GONG, CHUNLAN JIANG and LIANGQING LI	
On refined metric and hermitian structures in arithmetic, I: Galois–Gauss sums and weak ramification	79
WERNER BLEY, DAVID BURNS and CARL HAHN	
Rigidity in equivariant algebraic K -theory	141
NIKO NAUMANN and CHARANYA RAVI	
Functorality of the gamma filtration and computations for some twisted flag varieties	159
EOIN MACKALL	