

Algebra & Number Theory

Volume 5

2011

No. 5

**Kazhdan–Lusztig polynomials and drift
configurations**

Li Li and Alexander Yong



mathematical sciences publishers

Kazhdan–Lusztig polynomials and drift configurations

Li Li and Alexander Yong

The coefficients of the Kazhdan–Lusztig polynomials $P_{v,w}(q)$ are nonnegative integers that are upper semicontinuous relative to Bruhat order. Conjecturally, the same properties hold for h -polynomials $H_{v,w}(q)$ of local rings of Schubert varieties. This suggests a parallel between the two families of polynomials. We prove our conjectures for Grassmannians, and more generally, covexillary Schubert varieties in complete flag varieties, by deriving a combinatorial formula for $H_{v,w}(q)$. We introduce *drift configurations* to formulate a new and compatible combinatorial rule for $P_{v,w}(q)$. From our rules we deduce, for these cases, the coefficient-wise inequality $P_{v,w}(q) \leq H_{v,w}(q)$.

1. Introduction

Overview. This paper studies two families of polynomials $\{P_{v,w}(q)\}$ and $\{H_{v,w}(q)\}$ defined for pairs of permutations v, w in the symmetric group S_n (more generally, any Weyl group W). The former family consists of the celebrated *Kazhdan–Lusztig polynomials*, introduced in [Kazhdan and Lusztig 1979] to study representations of Hecke algebras. There it was conjectured that $P_{v,w}(q) \in \mathbb{Z}_{\geq 0}[q]$. This was later established by the same authors [1980] by interpreting $P_{v,w}(q)$ as the Poincaré polynomial for Goresky–MacPherson’s local intersection cohomology for the torus fixed point e_v of the Schubert variety X_w in the complete flag variety $\text{Flags}(\mathbb{C}^n)$.

A key contribution to the theory is R. Irving’s theorem [1988] that the $P_{v,w}(q)$ are *upper semicontinuous*: if $v' \leq v \leq w$ in Bruhat order, then $P_{v,w}(q) \leq P_{v',w}(q)$, where “ \leq ” means that, for each i , the coefficient of q^i in $P_{v,w}(q)$ is weakly smaller than the coefficient of q^i in $P_{v',w}(q)$. Thus, the Kazhdan–Lusztig polynomials are measures of the singularities of Schubert varieties whose coefficient growth tracks the worsening pathology of singularities as one moves along torus invariant \mathbb{P}^1 ’s towards the “most singular” point $e_{id} \in X_w$. In particular, $P_{v,w}(q) = 1$ if and only if $e_v \in X_w$ is a (rationally) smooth point.

MSC2000: primary 14M15; secondary 05E15, 20F55.

Keywords: Kazhdan–Lusztig polynomials, Hilbert series, Schubert varieties.

Conversely, the desire for insight into the combinatorics of Kazhdan–Lusztig polynomials naturally leads to the basic problem of understanding where and how the singularities of Schubert varieties worsen. In view of this converse problem, the growth of any semicontinuous singularity measure of Schubert varieties is of interest. One seeks concrete comparisons of different measures; see, e.g., [Woo and Yong 2008] and the references therein.

Specifically, a well-studied semicontinuous measure is given by the *Hilbert–Samuel multiplicity* $\text{mult}_{e_v}(X_w)$. However, while this contains useful local data about X_w , even more is carried by the \mathbb{Z} -graded Hilbert series of $\text{gr}_{\mathfrak{m}_{e_v}} \mathbb{O}_{e_v, X_w}$, the associated graded ring of the local ring \mathbb{O}_{e_v, X_w} ,

$$\text{Hilb}(\text{gr}_{\mathfrak{m}_{e_v}} \mathbb{O}_{e_v, X_w}, q) = \frac{H_{v,w}(q)}{(1 - q)^{\ell(w)}},$$

where $\ell(w) = \dim(X_w)$ is the *Coxeter length* of w . In particular, $\text{mult}_{e_v}(X_w) = H_{v,w}(1)$.

Conjecturally, each *h-polynomial* $H_{v,w}(q)$ is also in $\mathbb{Z}_{\geq 0}[q]$, and moreover is upper semicontinuous, just as is the case for Kazhdan–Lusztig polynomials. These conjectures suggest that the growth of the coefficients of the two families of polynomials is somehow correlated. In this paper, we offer an examination in the Grassmannian case, and more generally in the case of covexillary Schubert varieties inside $\text{Flags}(\mathbb{C}^n)$. There the nonnegativity and semicontinuity conjectures are proved by deriving a new combinatorial rule for $H_{v,w}(q)$. In addition, by introducing *drift configurations* as a model for the Kazhdan–Lusztig polynomials in these settings (after [Lascoux and Schützenberger 1981] and [Lascoux 1995]), we prove the inequality $P_{v,w}(q) \leq H_{v,w}(q)$. This combinatorial discovery further indicates the link between the two families; no alternative explanation via algebraic or geometric methods seems available at present.

Summarizing, the main thesis of this paper is that there exists a parallel between $\{P_{v,w}(q)\}$ and $\{H_{v,w}(q)\}$. Our basis for this perspective comes from proofs of compatible and positive combinatorial rules for the two families of polynomials.

Statements of the main conjecture and theorems. Recapitulating, this paper formulates, and constructs supporting combinatorics for, the following conjecture:

Conjecture 1.1. *The h-polynomials $H_{v,w}(q)$ have nonnegative integral coefficients. In addition, they are upper semicontinuous; i.e., if $v' \leq v$ in Bruhat order then $H_{v,w}(q) \leq H_{v',w}(q)$.*

The nonnegativity claim would actually be immediate if $\text{gr}_{\mathfrak{m}_{e_v}} \mathbb{O}_{e_v, X_w}$ is Cohen–Macaulay (see page 604). However, this latter assertion seems to be a folklore conjecture. Although \mathbb{O}_{e_v, X_w} is itself Cohen–Macaulay [Ramanathan 1985], this property might be lost when degenerating to $\text{gr}_{\mathfrak{m}_{e_v}} \mathbb{O}_{e_v, X_w}$. On the other hand, the

results detailed in this paper and in [Li and Yong 2011] also support the Cohen–Macaulayness conjecture. In particular, the latter would follow from that paper’s Conjecture 8.5, a stronger claim asserting that Stanley–Reisner simplicial complexes of certain Gröbner degenerations of Kazhdan–Lusztig varieties are vertex decomposable.

The semicontinuity claim is itself a strengthening of the nonnegativity claim since the smoothness of X_w at e_w implies $H_{w,w}(q) = 1$. Furthermore, although the betti numbers of $\mathrm{gr}_{\mathfrak{m}_{e_v}} \mathbb{C}_{e_v, X_w}$ are semicontinuous, the coefficients of $H_{v,w}(q)$ are an involved, signed expression in terms of those numbers. Therefore, this semicontinuity phenomenon seems substantive.

The natural projection map

$$\pi : \mathrm{Flags}(\mathbb{C}^n) \rightarrow \mathrm{Gr}_k(\mathbb{C}^n) : (\langle 0 \rangle \subset F_1 \subset \cdots \subset F_k \subset \cdots \subset F_{n-1} \subset \mathbb{C}^n) \mapsto F_k,$$

where $\mathrm{Gr}_k(\mathbb{C}^n)$ is the Grassmannian of k -dimensional planes in \mathbb{C}^n , is a fibration: local properties of torus fixed points $e_\mu \in X_\lambda \subseteq \mathrm{Gr}_k(\mathbb{C}^n)$ for Young diagrams $\lambda, \mu \subseteq k \times (n-k)$, are equivalent to local properties of $e_v \in X_w \subseteq \mathrm{Flags}(\mathbb{C}^n)$ where $v, w \in S_n$ are maximal Coxeter length representatives of λ, μ where the latter are thought of as cosets of $S_n / (S_k \times S_{n-k})$; see, e.g., [Brion 2004, Example 1.2.3]. These v and w are *cograssmannian*, i.e., they have a unique ascent, at position k : $v(k) < v(k+1)$ and $w(k) < w(k+1)$.

Lifting Grassmannian problems to $\mathrm{Flags}(\mathbb{C}^n)$ has the advantage of allowing one to embed them within the wider class of *covexillary Schubert varieties* X_w , i.e., where w is 3412-avoiding: there are no indices $i_1 < i_2 < i_3 < i_4$ such that $w(i_1), w(i_2), w(i_3), w(i_4)$ are in the same relative order as 3412. This class appears more tractable than general flag Schubert varieties since it shares many of the same features as Grassmannian Schubert varieties. However, there is a salient difference: Grassmannian Schubert varieties are locally defined by equations that are homogeneous with respect to the standard grading that assigns each variable degree one. In general, this is not true in the covexillary case. This homogeneity means that taking associated graded of the local ring essentially does nothing, and so $\mathrm{gr}_{\mathfrak{m}_{e_v}} \mathbb{C}_{e_v, X_w}$ is automatically Cohen–Macaulay; see, e.g., [Li and Yong 2011, Section 1] and page 604.

The covexillary condition has already attracted significant attention; see, e.g., [Lakshmibai and Sandhya 1990; Lascoux 1995; Manivel 2001; Knutson and Miller 2005; Knutson et al. 2008; Knutson et al. 2009; Li and Yong 2011]. In particular, Section 2.4 of [Knutson and Miller 2005] connects the condition to *ladder determinantal ideals* studied in commutative algebra. Our three main theorems below concern the covexillary setting, providing our main cases of support towards both our main thesis and Conjecture 1.1.

One of our results is to prove the following link between $H_{v,w}(q)$ and $P_{v,w}(q)$:

Theorem 1.2. *For w covexillary,*

$$P_{v,w}(q) \preceq H_{v,w}(q) \quad \text{and} \quad \deg P_{v,w}(q) = \deg H_{v,w}(q).$$

While the Grassmannian case *per se* is new and supports our thesis, the covexillary generality also further highlights the amenability of covexillary Schubert varieties. Our proof of [Theorem 1.2](#) is based on a new formula for covexillary Kazhdan–Lusztig polynomials. An earlier rule was given by A. Lascoux [[1995](#)], generalizing his earlier Grassmannian rule with M.-P. Schützenberger [[Lascoux and Schützenberger 1981](#)]. (For more recent treatments of the Grassmannian case see [[Shigechi and Zinn-Justin 2010](#); [Jones and Woo 2010](#)], for example.) Our formulation of a covexillary rule is in terms of *drift configurations*. It is entirely graphical and is perhaps more handy to compute.

To state our rule we use standard combinatorics of the symmetric group (see, e.g., [[Manivel 2001](#), Chapter 2]) as well as some terminology introduced in [[Li and Yong 2011](#)]. (The reader may wish to compare [Examples 1.5](#) and [1.6](#) below with what follows.) Let $w \in S_n$ be covexillary. Superimpose the graph $G(v)$ of v drawn with dots \circ in positions $(n - v(j) + 1, j)$ on top of the *diagram*

$$D(w) = \{(i, j) : i < n - w(j) + 1 \text{ and } j < w^{-1}(n - i + 1)\} \subset [n] \times [n].$$

Throughout, we use the convention that rows are indexed from bottom to top, and columns are indexed from left to right. Move each box ϵ of the *essential set*

$$\mathcal{E}(w) = \{(i, j) \in D(w) : (i + 1, j), (i, j + 1) \notin D(w)\}$$

diagonally southwest by the number of dots of $G(v)$ weakly southwest of ϵ . Call the resulting boxes $\{\epsilon'\}$, and define $B(v, w)$ to be the smallest Young diagram that contains $\{\epsilon'\}$ and $(1, 1)$ (we use French convention for our Young diagrams). The *shape* $\lambda(w)$ of w is obtained by sorting the vector counting the number of boxes in nonempty rows of $D(w)$ into decreasing order. Now, draw $\lambda(w)$ in the southwest corner of $B(v, w)$.

Declare that any corner of $\lambda(w)$ is *0-special*. Let $\text{arm}(b)$ (respectively, $\text{leg}(b)$) refer to the boxes in $\lambda(w)$ strictly to the right (above) of b and in the same row (column). Inductively, a box $b \in \lambda(w)$ is *z -special*, for $z \in \mathbb{N}$ if it is maximally northeast subject to

- $|\text{leg}(b)| = |\text{arm}(b)|$; and
- none of the boxes of $\{b\} \cup \text{arm}(b) \cup \text{leg}(b)$ are y -special for any $y < z$.

A box is *special* if it is z -special for some z . The *continent* of a special box b is the set of $x \in \lambda(w)$ such that b is the maximally northeast special box that is weakly southwest of x . The union of continents is

$$\text{Pangaea}(v, w) \subseteq \lambda(w)$$

(the set difference being an immovable reference continent).

Definition 1.3. A drift configuration \mathcal{D} is a nonoverlapping configuration of continents inside $B(v, w)$, such that

- each special box is diagonally weakly northeast of its position in $\text{Pangaea}(v, w)$, and
- relative southwest-northeast positions of special cells are maintained.

Let $\text{drift}(v, w)$ be the set of all such \mathcal{D} and let $\text{wt}(\mathcal{D})$ be the total distance traveled by the continents from $\text{Pangaea}(v, w)$. Consider the generating series

$$Q_{v,w}(q) = \sum_{\mathcal{D} \in \text{drift}(v,w)} q^{\text{wt}(\mathcal{D})}.$$

Theorem 1.4. Suppose that $v, w \in S_n$ and w is covexillary. Then:

- (I) $P_{v,w}(q) = Q_{v,w}(q)$.
- (II) If we instead take every box of $\lambda(w)$ to be a separate “country”, each of which “drifts” according to the rules of [Definition 1.3](#), the total number of drift configurations is $\text{mult}_{e_v}(X_w)$; hence

$$P_{v,w}(1) \leq \text{mult}_{e_v}(X_w),$$

as is manifest from (I).

- (III) There is a vertex decomposable (thus shellable) simplicial complex $\text{KL}_{v,w}$ that is homeomorphic to a ball or a sphere, and whose facets are labeled by $\mathcal{D} \in \text{drift}(v, w)$.


Our proof of (I) is a bijection with A. Lascoux’s rule (which descends to a bijection with the rule of [[Lascoux and Schützenberger 1981](#)] for Grassmannians). The multiplicity rule from (II) just restates the theorem from [[Li and Yong 2011](#)] (compare the Grassmannian rule of [[Ikeda and Naruse 2009](#)]). Although the inequality of (II) is a consequence of [Theorem 1.2](#), we are emphasizing that our rule from (I) is compatible with our multiplicity rule and makes the inequality transparent. Actually, whether such an inequality might exist was first asked to us (independently) by S. Billey and A. Woo. Afterwards, H. Naruse informed us that he has a proof for all cominuscule G/P . These questions and results provided us initial motivation for our work towards [Theorem 1.4](#). Note that as with the more general inequality of [Theorem 1.2](#), this inequality is not true in general. For example, $P_{13425,34512}(1) = 3$ while $\text{mult}_{e_{13425}}(X_{34512}) = 2$.

Statement (III) is derived from [[Knutson et al. 2008](#)]. It points out a further resemblance to the combinatorics of $\text{mult}_{e_v}(X_w)$ in [[Li and Yong 2011](#)], where a similar complex also appears.

Example 1.5. The left diagram depicts $\text{Pangaea}(v, w)$, where $v = \text{id}$ and


$$w = \underline{20} \ \underline{19} \ \underline{18} \ \underline{11} \ \underline{10} \ 9 \ 8 \ \underline{12} \ \underline{17} \ \underline{16} \ 7 \ 6 \ \underline{15} \ \underline{14} \ \underline{13} \ 5 \ 4 \ 3 \ 2 \ 1.$$

It has six continents, shown in different colors. The right diagram shows a particular drift $\mathcal{D} \in \text{drift}(v, w)$; its weight is 14.




Example 1.6. Let $w = \underline{10} \ 9 \ 5 \ 4 \ 3 \ 8 \ 2 \ 7 \ 6 \ 1$, $v = 2 \ 3 \ 4 \ 6 \ 5 \ 1 \ 7 \ 8 \ 9 \ \underline{10}$. Here $\lambda(w) = (4, 4, 3)$.

The left figure shows $D(w)$, with $G(w)$ overlaid as black dots and $G(v)$ as open circles.



Starting from $D(w)$ and the overlaid \circ 's of $G(v)$, we derive $B(v, w)$, shown on the right. The special boxes are marked by '+'s. We have $\mathcal{E}(w) = \{e_1, e_2\}$ (being the maximally northeast boxes of each connected component of $D(w)$) move to $\{e'_1, e'_2\}$, as determined by the \circ 's of $G(v)$. These are the five drift configurations:



We can write $Q_{v,w}(q) = 1 + 2q + q^2 + q^3$. □

Our proof of [Theorem 1.2](#) also depends on a new (and the first manifestly positive) combinatorial rule for covexillary $H_{v,w}(q)$. It additionally implies special

cases of the nonnegativity and upper semicontinuity conjectures. Identify a partition $\lambda = (\lambda_1 \geq \dots \geq \lambda_\ell > 0)$ with its Young diagram (in French notation). Recall that a Young tableau T of shape λ is *semistandard* if it is weakly increasing along rows and strictly increasing up columns. Given a vector $\mathbf{b} = (b_1, \dots, b_\ell)$, we say T is *flagged* by \mathbf{b} if each entry in row i is at most b_i . Let $\text{SSYT}(\lambda, \mathbf{b})$ denote the set of semistandard Young tableaux flagged by \mathbf{b} . A (nonempty) set-valued filling is semistandard if each tableau obtained by choosing a singleton from each set is semistandard [Buch 2002]. Similarly, we define *flagged set-valued semistandard tableaux*, and the set $\text{SetSSYT}(\lambda, \mathbf{b})$ [Knutson et al. 2008].

Define $U \in \text{SetSSYT}(\lambda, \mathbf{b})$ to be *lower saturated* if no smaller number can be added to any box $U(i, j)$ while maintaining semistandardness. In symbols, each $U(i, j)$ is of the form

$$[\alpha, \beta] := \{\alpha, \alpha + 1, \dots, \beta - 1, \beta\},$$

for some α, β (depending on i, j), where

$$\alpha = \max\{\max U(i, j - 1), 1 + \max U(i - 1, j)\}.$$

Our convention for lower saturated tableaux is that $U(i, 0) = 1$ for all $i > 0$ and $U(0, j) = 0$ for all $j > 0$. Let

$$\text{Lower}(\lambda, \mathbf{b}) \subseteq \text{SetSSYT}(\lambda, \mathbf{b})$$

denote this subset of lower saturated tableaux.

Define the *saturation* $\text{sat}(T) \in \text{Lower}(\lambda, \mathbf{b})$ of $T \in \text{SSYT}(\lambda, \mathbf{b})$ to be

$$\text{sat}(T)(i, j) = [\max\{T(i, j - 1), 1 + T(i - 1, j)\}, T(i, j)].$$

For $U \in \text{SetSSYT}(\lambda, \mathbf{b})$, let

$$\text{ex}(U) = |U| - |\lambda|,$$

where $|U|$ refers to the number of entries of U and $|\lambda| = \lambda_1 + \lambda_2 + \dots$.

Finally, if $T \in \text{SSYT}(\lambda, \mathbf{b})$ set

$$\text{depth}(T) := \text{ex}(\text{sat}(T)) = |\text{sat}(T)| - |T|. \tag{1-1}$$

If $\lambda(w) = (\lambda(w)_1 \geq \dots \geq \lambda(w)_\ell > 0)$, define

$$\mathbf{b} = \mathbf{b}(\Theta_{v,w}) = (b_1, \dots, b_\ell) \tag{1-2a}$$

by

$$b_i = \max\{m : B(v, w)_m \geq \lambda(w)_i + m - i\}. \tag{1-2b}$$

This is the maximum distance that the rightmost box in row i can drift diagonally northeast within $B(v, w)$ (ignoring the presence of other boxes).

Theorem 1.7. *Let $w \in S_n$ be covexillary. Then*

$$H_{v,w}(q) = \sum_{T \in \text{SSYT}(\lambda(w), \mathbf{b}(\Theta_{v,w}))} q^{\text{depth}(T)} = \sum_{U \in \text{Lower}(\lambda(w), \mathbf{b}(\Theta_{v,w}))} q^{\text{ex}(U)}.$$

Moreover, [Conjecture 1.1](#) is true under the hypothesis.

Example 1.8. For $n = 5$, $w = 52341$, $v = 12345$. There are five semistandard tableaux of shape $(2, 1)$ and flagged by $(2, 3)$:

2	3	2	3	3
1	1	1	2	2
1	1	2	2	2

Their saturations are


2	2,3	2	2, 3	3
1	1	1	1	1,2
1	1	1,2	1,2	2

The corresponding ex values are

$$0, 1, 1, 2, 1.$$

Thus by [Theorem 1.7](#), $H_{v,w}(q) = 1 + 3q + q^2$. □

Example 1.9. Continuing [Example 1.8](#), there are four drift configurations of the two continents:



The Kazhdan–Lusztig polynomial is $P_{v,w}(q) = 1 + 2q + q^2$. We see that $P_{v,w}(q) \leq H_{v,w}(q)$, in agreement with [Theorem 1.2](#). □

Organization and contents. In [Section 2](#), we state some preliminaries and further discuss [Conjecture 1.1](#). We then prove [Theorem 1.7](#). In [Section 3](#), we briefly recall, for comparison, basics about Kazhdan–Lusztig theory. We then prove [Theorem 1.2](#) while temporarily assuming [Theorem 1.4\(I\)](#). [Section 4](#) is devoted to the construction of the simplicial complex of [Theorem 1.4\(II\)](#) and proof of its asserted properties. We furthermore define polynomials generalizing $Q_{v,w}(q)$ that naturally arise from this complex. In [Section 5](#) we prove [Theorem 1.4\(I\)](#). We end that section with two comments ([Remarks 5.5](#) and [5.6](#)) about further properties of $P_{v,w}(q)$ that can be deduced from the rule. In [Section 6](#), we give a formula for a different “ q -analogue” of $\text{mult}_{e_v}(X_w)$ than $H_{v,w}(q)$. In [Section 7](#), we offer some final remarks.

2. Hilbert series of the local ring \mathbb{O}_{e_v, X_w}

2.1. Preliminaries. We use the usual identification $\text{Flags}(\mathbb{C}^n) = GL_n/B$ where B is the Borel subgroup consisting of invertible upper triangular matrices. Thus GL_n acts on $\text{Flags}(\mathbb{C}^n)$ by left multiplication, as does B , and the torus T of invertible diagonal matrices. For each $v \in S_n$, let e_v denote the associated T -fixed point. The *Schubert cell* is $X_w^\circ := Be_w$, while its Zariski closure is the *Schubert variety* $X_w = \overline{X_w^\circ}$, an irreducible variety of dimension $\ell(w)$. We have that $e_v \in X_w$ if and only if $v \leq w$ in *Bruhat order*. A neighborhood of each point $p \in X_w$ is isomorphic to a neighborhood of some e_v , by the action of B . Hence, it suffices to restrict attention to T -fixed points. Let B_- be the opposite Borel subgroup of invertible lower triangular matrices. If we set $\Omega_v^\circ = B_-vB/B$ to be the *opposite Schubert cell*, then up to crossing by affine space, a local neighborhood of $e_v \in X_w$ is given by the *Kazhdan–Lusztig variety* $\mathcal{N}_{v,w} = X_w \cap \Omega_v^\circ$ [Kazhdan and Lusztig 1979, Lemma A.4].

Suppose p is a point on a scheme Y . Let $\text{gr}_{\mathfrak{m}_p} \mathbb{O}_{p,Y}$ denote the *associated graded ring* of the local ring $\mathbb{O}_{p,Y}$ with respect to its maximal ideal \mathfrak{m}_p , i.e.,

$$\text{gr}_{\mathfrak{m}_p} \mathbb{O}_{p,Y} = \bigoplus_{i \geq 0} \mathfrak{m}_p^i / \mathfrak{m}_p^{i+1}.$$

Since $\text{gr}_{\mathfrak{m}_p} \mathbb{O}_{p,Y}$ picks up a \mathbb{Z} -grading, it now makes sense to discuss its Hilbert series. One can always express this series in the form

$$\text{Hilb}(\text{gr}_{\mathfrak{m}_p} \mathbb{O}_{p,Y}, q) = \frac{H_{p,Y}(q)}{(1-q)^{\dim Y}}$$

where $H_{p,Y}(q) \in \mathbb{Z}[q]$ is the *h-polynomial* associated to $p \in Y$. It follows from standard facts that $H_{p,Y}(1) = \text{mult}_p(Y)$; see, e.g., [Kreuzer and Robbiano 2005, Theorem 5.4.15]. Hence $H_{p,Y}(q) = 1$ if and only if Y is smooth at p . In addition, note $H_{p,Y}(0) = 1$, since this is the dimension of the zero graded piece of $\text{gr}_{\mathfrak{m}_p} \mathbb{O}_{p,Y}$, i.e., the dimension of the field $\mathbb{O}_{p,Y}/\mathfrak{m}_p$.

Now, for any $v, w \in S_n$, we define $H_{v,w}(q) \in \mathbb{Z}[q]$ to be the *h-polynomial* associated to $e_v \in X_w$. At present, there is no purely combinatorial formula (even nonpositive or recursive) for computing $H_{v,w}(q)$. However, instead one can utilize the explicit coordinates and equations for the ideal $I_{v,w}$ to define $\mathcal{N}_{v,w} = \text{Spec}(\mathbb{C}[z^{(v)}]/I_{v,w})$, as done in [Woo and Yong 2008, Section 3.2]. Then one can Gröbner degenerate $\mathcal{N}_{v,w}$ to a scheme theoretic union of coordinate subspaces $\mathcal{N}'_{v,w}$, using any of the term orders $\prec_{v,w,\pi}$ from [Li and Yong 2011, Section 3]. As explained in Theorem 3.1 (and its proof) of that reference, the stated Gröbner degenerations degenerate not only $\mathcal{N}_{v,w}$ but also its projectivized tangent cone $\text{Proj}(\text{gr}_{\mathfrak{m}_{e_v}} \mathbb{O}_{e_v, X_w})$. Therefore the *h-polynomial* of $\mathcal{N}'_{v,w}$ equals $H_{v,w}(q)$.

2.2. Conjectures. Let us now return to the discussion of [Conjecture 1.1](#). Using the method for computing $H_{v,w}(q)$ summarized above, we obtained exhaustive checks for $n \leq 7$ of the following claim, restated from the introduction:

Nonnegativity conjecture. $H_{v,w}(q) \in \mathbb{Z}_{\geq 0}[q]$.

In [[Li and Yong 2011](#), Conjecture 8.5] we conjectured that within the family of term orders $<_{v,w,\pi}$, at least one gives a Gröbner limit scheme $\mathcal{N}'_{v,w}$ that is reduced, equidimensional and whose Stanley–Reisner simplicial complex $\Delta_{v,w}$ is a vertex-decomposable ball or sphere. This implies in particular that $\Delta_{v,w}$ is shellable and thus Cohen–Macaulay. If this conjecture were true, it would follow that $\text{gr}_{\mathfrak{m}_v} \mathbb{C}_{e_v, X_w}$ is Cohen–Macaulay. Thus the nonnegativity conjecture would hold by, e.g., [[Bruns and Herzog 1993](#), Corollary 4.1.10].

In the case that $I_{v,w}$ is a homogeneous ideal, with respect to the standard grading that assigns each variable degree 1, since \mathbb{C}_{e_v, X_w} is Cohen–Macaulay [[Ramanathan 1985](#)], it follows that the associated graded ring is Cohen–Macaulay; see [[Bruns and Herzog 1993](#), Exercise 2.1.27(c)], for example. Hence nonnegativity follows in this case. A. Knutson [[2009](#), p. 25] has shown that this homogeneity occurs whenever w is 321-avoiding. Moreover, in [[Woo and Yong 2009](#), Section 5] it was explained how “parabolic moving” reduces a large percentage of cases (for $n \leq 10$) to the homogeneous case. However, not every case can be so reduced, including those in the covexillary class. Thus, these cases provide further support for the nonnegativity conjecture, separate from [Theorem 1.7](#).

Upper semicontinuity conjecture. *If $v' \leq v \leq w$ in Bruhat order, then*

$$H_{v,w}(q) \leq H_{v',w}(q).$$

Unfortunately, even if we knew $\text{gr}_{\mathfrak{m}_v} \mathbb{C}_{e_v, X_w}$ to be Cohen–Macaulay, we do not know any way to express these coefficients in homological terms that would make the upper semicontinuity conjecture transparent. It should be noted that the proof of this property for Kazhdan–Lusztig polynomials in [[Irving 1988](#)] was not achieved using the geometry of Schubert varieties. However, see the geometric argument for the more general result [[Braden and MacPherson 2001](#), Theorem 3.6].

Although any proof of the above conjectures is desired, ideally one would also like combinatorial explanations of the properties.

Let us pause to collect some further facts for small n in the following computational result. For (D) below we refer the reader to [[Woo and Yong 2008](#), Section 2.1] for the definition of *interval pattern avoidance* of $[x, y] \in S_\infty \times S_\infty$. There we explain that the existence of an *interval pattern embedding* guarantees $\mathcal{N}_{x,y} \cong \mathcal{N}_{\tilde{w},w}$, where $[x, y] \cong [\tilde{w}, w]$ is an isomorphism of posets of Bruhat intervals in S_∞ . Thus, if the inequality $P_{x,y}(q) \leq H_{x,y}(q)$ fails, so must $P_{\tilde{w},w}(q) \leq H_{\tilde{w},w}(q)$.

Proposition 2.1. (A) $\deg H_{v,w}(q) \leq \deg P_{v,w}(q)$ for $v \leq w \in S_n$ and $n \leq 6$.

- (B) $\deg H_{v,w}(q) \leq \frac{1}{2}(\ell(w) - \ell(v) - 1)$ for $v < w \in S_n$ and $n \leq 7$.
- (C) The coefficients of $H_{v,w}(q)$ form a unimodal sequence for $v, w \in S_n$ and $n \leq 7$.
- (D) $P_{v,w}(q) \leq H_{v,w}(q)$ holds for all $v \leq w \in S_n$ and $n \leq 6$, if and only if w interval pattern avoids

$$[14235, 45123], \quad [31524, 53412], \quad [14325, 45312], \\ [13425, 34512], \quad [24153, 45231], \quad [154326, 564312].$$

(Note that the first and fourth intervals, and the second and fifth intervals are related by taking inverses. For all $n \geq 1$, the inequality fails whenever w contains one of these intervals.)

Proof and discussion. Each of the assertions were verified using Macaulay 2. For (A) and (B) note that $\deg P_{v,w}(q) \leq \frac{1}{2}(\ell(w) - \ell(v) - 1)$ is a standard fact about Kazhdan–Lusztig polynomials; see item (iii) on page 608.

For (D), computation shows that $P_{v,w}(q) = H_{v,w}(q)$ for $n \leq 4$, so the inequality holds in that situation. We checked that each of the intervals $[x, y]$ listed corresponds to a failure of the inequality for $n \leq 5$. For $n = 6$ we computationally verified the claim (there are 36 cases $w \in S_6$ where the inequality fails for some $v \leq w$, and of those only one cannot be blamed on the $n = 5$ cases). The $n > 6$ case follows from general properties of interval pattern embeddings recalled above. \square

One might conjecture that both (A) and its weak form (B) hold for all n . However with (A), experience has shown that data for $n \leq 6$ is soft evidence for any conjecture that involves Kazhdan–Lusztig polynomials. Note that if (A) is true, one cannot have $P_{v,w}(q) \leq H_{v,w}(q)$ unless $\deg H_{v,w}(q) = \deg P_{v,w}(q)$, which is indeed what we show when w is covexillary.

In view of (C), it is also natural to guess that unimodality is true in general. One warning however is that the stronger assertion that the coefficients of $H_{v,w}(q)$ are *log-concave* is false, as the example below shows:

Example 2.2. Let $w = 5671234$ and $v = 1352476$. Computation using Macaulay 2 shows there is a choice of $\prec_{v,w,\pi}$ such that $\mathcal{N}'_{v,w}$ is Cohen–Macaulay (but not Gorenstein), and that $H_{1352476,5671234}(q) = 1 + 2q + q^2 + q^3$, which is not log-concave. \square

By contrast, see the related work [Rubey 2005], which shows log-concavity holds in a special ladder determinantal case (note that w is not covexillary in our counterexample).

Even knowing Cohen–Macaulayness of $\text{gr}_{\mathfrak{m}_v} \mathbb{C}_{e_v, X_w}$ does not, in and of itself, prove unimodality. In fact, R. Stanley [1989, Conjecture 4(a)] had conjectured unimodality for a general graded Cohen–Macaulay domain R over a field which is generated by R_1 . Actually, he even conjectured the stronger claim of log-concavity,

although counterexamples to the stronger claim were later found by G. Niesi and L. Robbiano; see [Brenti 1994, Section 5]. (Example 2.2 gives a different counterexample to Stanley’s log-concavity conjecture.)

It should also be mentioned that in contrast, the Kazhdan–Lusztig polynomials are not in general unimodal and in fact P. Polo [1999] proved that every nonnegative integral polynomial with constant coefficient 1 is some $P_{v,w}(q)$.

While Theorem 1.7 allows us to prove the nonnegativity, upper semicontinuity and degree properties for covexillary X_w , a solution to the following problem has eluded us:

Problem 2.3. Give a combinatorial proof (e.g., using Theorem 1.7) for the unimodality conjecture, when w is covexillary (or even cograssmannian) by establishing a sequence of explicit injections and surjections of the relevant Young tableaux.

Concerning (D), we do not expect the characterization to be valid for all n . Instead, one aims to expand this list into a (human-readable) classification, via a finite list of families of patterns to avoid, as is the case for many other properties studied in [Woo and Yong 2008].

Using the analogy with Kazhdan–Lusztig theory, numerous further problems, which had been previously considered for $P_{v,w}(q)$ but not $H_{v,w}(q)$, make sense. To name a few: Is $H_{v,w}(q)$ determined by the poset isomorphism class of the interval $[v, w]$ in Bruhat order? (This is an analogue of a conjecture of G. Lusztig.) Can one give a combinatorial algorithm for computing $H_{v,w}(q)$? Better yet, can one find a positive combinatorial rule for $H_{v,w}(q)$, thus establishing the nonnegativity conjecture?

2.3. Proof of Theorem 1.7. Continuing the definitions before the statement of Theorem 1.7, set

$$\text{sup} : \text{SetSSYT}(\lambda, \mathbf{b}) \rightarrow \text{SSYT}(\lambda, \mathbf{b})$$

by sending U to T where $T(i, j) = \max U(i, j)$. The following is clear:

Lemma 2.4. *The maps*

$$\text{sat} : \text{SSYT}(\lambda, \mathbf{b}) \rightarrow \text{Lower}(\lambda, \mathbf{b}) \quad \text{and} \quad \text{sup} |_{\text{Lower}(\lambda, \mathbf{b})} : \text{Lower}(\lambda, \mathbf{b}) \rightarrow \text{SSYT}(\lambda, \mathbf{b})$$

are mutually inverse bijections.

Let us recall some definitions and terminology utilized in [Li and Yong 2011]. Define $r_b^w = r_{(i,j)}^w$ to be the number of \bullet of $G(w)$ weakly southwest of the box $b = (i, j)$. Given $v \leq w$ and w covexillary, $\Theta_{v,w} \in S_n$ is defined there to be the unique permutation such that $\lambda(w) = \lambda(\Theta_{v,w})$ and

$$\mathcal{E}(\Theta_{v,w}) = \left\{ \mathbf{e}' : \mathbf{e}' \text{ is obtained by moving each } \mathbf{e} \text{ in } \left. \begin{array}{l} \mathcal{E}(w) \\ \text{diagonally southwest by } r_e^v \text{ units.} \end{array} \right\} \right.$$

The permutation $\Theta_{v,w}$ was proved to be itself covexillary.

Define $B(w)$ to be the smallest Young diagram with southwest corner in position $(1, 1)$ that contains all of $\mathcal{C}(w)$. Set

$$B(v, w) = B(\Theta_{v,w}).$$

If $\lambda(w) = (\lambda(w)_1 \geq \dots \geq \lambda(w)_\ell > 0)$, define $\mathbf{b} = \mathbf{b}(w) = (b_1, \dots, b_\ell)$ by

$$b_i = \max\{m : B(w)_m \geq \lambda(w)_i + m - i\}.$$

This agrees with, and slightly reformulates, the definitions of $B(v, w)$ and \mathbf{b} from the introduction.

In [Li and Yong 2011, Theorem 6.6] we proved that

$$\text{Hilb}(\text{gr}_{\mathfrak{m}_{e_v}} \mathbb{O}_{e_v, X_w}, q) = G_{\lambda(w)}(q) / (1 - q)^{\binom{n}{2}},$$

where

$$G_{\lambda(w)}(q) = \sum_{k \geq |\lambda(w)|} (-1)^{k - |\lambda(w)|} (1 - q)^k \times \#\text{SetSSYT}(\lambda(w), \mathbf{b}, k)$$

and $\#\text{SetSSYT}(\lambda(w), \mathbf{b}, k)$ is the number of flagged set-valued semistandard Young tableaux of shape $\lambda(w)$ with flag $\mathbf{b} = \mathbf{b}(\Theta_{v,w})$ which use exactly k entries.

Since the local ring \mathbb{O}_{e_v, X_w} is of dimension $\ell(w) = \binom{n}{2} - |\lambda(w)|$, we rewrite

$$\text{Hilb}(\text{gr}_{\mathfrak{m}_{e_v}} \mathbb{O}_{e_v, X_w}, q) = \frac{H_{v,w}(q)}{(1 - q)^{\ell(w)}},$$

where

$$H_{v,w}(q) = \sum_{U \in \text{SetSSYT}(\lambda(w), \mathbf{b})} (q - 1)^{\text{ex}(U)}.$$

We need to show that

$$\sum_{U \in \text{SetSSYT}(\lambda(w), \mathbf{b})} (q - 1)^{\text{ex}(U)} = \sum_{T \in \text{SSYT}(\lambda(w), \mathbf{b})} q^{\text{depth}(T)} \quad (2-1)$$

by proving that, for every $T \in \text{SSYT}(\lambda(w), \mathbf{b})$,

$$\sum_{U \in \text{sup}^{-1}(T)} (q - 1)^{\text{ex}(U)} = q^{\text{depth}(T)}.$$

There are $\text{depth}(T)$ elements in $\text{sat}(T)$ but not in T . We can delete any subset of those elements from $\text{sat}(T)$ and obtain $T' \in \text{sup}^{-1}(T)$ (so $\#\text{sup}^{-1}(T) = 2^{\text{depth}(T)}$). Hence the left-hand side is equal to

$$(1 + (q - 1))^{\text{depth}(T)} = q^{\text{depth}(T)},$$

and therefore the equality (2-1) follows. Thus, the first equality of the theorem holds and the second is clear from Lemma 2.4.

The nonnegativity claim is manifest from the combinatorial rule; however, let us also give a geometric proof. In [Li and Yong 2011] we proved that for covexillary w , $\mathcal{N}_{v,w}$ degenerates, under a choice of $\prec_{v,w,\pi}$ to a Cohen–Macaulay limit scheme $\mathcal{N}'_{v,w}$. Hence, nonnegativity of $H_{v,w}(q)$ follows from [Bruns and Herzog 1993, Corollary 4.1.10] and the discussion on page 603.

For the upper semicontinuity claim, fix $w \in S_n$ and suppose $v' \leq v \leq w$. Consider an essential box $\epsilon \in \mathcal{E}(w)$. In the construction of $\mathcal{E}(\Theta_{v,w})$, the essential box ϵ is moved diagonally southwest by r_ϵ^v units. Since $v' \leq v$, a standard characterization of Bruhat order shows $r_\epsilon^{v'} \leq r_\epsilon^v$. Thus, each essential box ϵ moves further southwest in to its position in $\mathcal{E}(\Theta_{v,w})$ than it does for $\mathcal{E}(\Theta_{v',w})$. Therefore,

$$B(v, w) \subseteq B(v', w),$$

and hence,

$$\mathbf{b}(\Theta_{v,w}) = (b_1, \dots, b_\ell) \leq \mathbf{b}(\Theta_{v',w}) = (b'_1, \dots, b'_\ell),$$

in the sense that $b_i \leq b'_i$ for every i . Consequently, $\text{SSYT}(\lambda, \mathbf{b}) \subseteq \text{SSYT}(\lambda, \mathbf{b}')$, which clearly implies $H_{v,w}(q) \leq H_{v',w}(q)$, as desired. \square

3. Kazhdan–Lusztig theory

The Hecke algebra. Let $R = \mathbb{Z}[q^{\frac{1}{2}}, q^{-\frac{1}{2}}]$ be the ring of Laurent polynomials over \mathbb{Z} in the indeterminate $q^{\frac{1}{2}}$. The Hecke algebra \mathcal{H}_{n-1} of S_n is the algebra over R with basis $\{T_w : w \in S_n\}$ and relations

$$\begin{aligned} T_{s_i} T_w &= T_{s_i w} \quad \text{if } \ell(s_i w) > \ell(w), \\ T_{s_i}^2 &= (q - 1)T_{s_i} + qT_{\text{id}}. \end{aligned}$$

There is an involution $\iota : \mathcal{H}_{n-1} \rightarrow \mathcal{H}_{n-1}$ defined by $\iota(q^{\frac{1}{2}}) = q^{-\frac{1}{2}}$ and $\iota(T_w) = T_{w^{-1}}$.

It was proved in [Kazhdan and Lusztig 1979] that there exists a basis $\{\mathcal{C}'_w\}$ of \mathcal{H}_{n-1} that is uniquely determined by the conditions

$$\iota(\mathcal{C}'_w) = \mathcal{C}'_w \quad \text{and} \quad \mathcal{C}'_w = (q^{-\frac{1}{2}})^{\ell(w)} \sum_{v \leq w} P_{v,w}(q) T_v,$$

where

- (i) $P_{w,w}(q) = 1$,
- (ii) $P_{v,w}(q) = 0$ if $v \not\leq w$, and
- (iii) $P_{v,w}(q) \in \mathbb{Z}[q]$ is of degree at most $\frac{1}{2}(\ell(w) - \ell(v) - 1)$ if $v < w$.

The existence of this basis was established by an explicit recursion for the Kazhdan–Lusztig polynomials $P_{v,w}(q)$, which we omit. Our source for these facts is [Billiey and Lakshmibai 2000, Chapter 6], to which we refer the reader to for further details.

Conditions (i) and (ii) also hold for the $H_{v,w}(q)$, while (iii) conjecturally holds (compare [Proposition 2.1](#) and the discussion thereafter). It is mildly tempting to think about another basis of the Hecke algebra defined by replacing $P_{v,w}(q)$ by $H_{v,w}(q)$ in the above definition of C'_w . While this other basis has a unimodular transition matrix with the Kazhdan–Lusztig basis, it doesn't possess any of the other nice properties, such as positive structure constants or invariance under the involution ι .

Proof of Theorem 1.2. Recall that in what follows, we are assuming the formula for $P_{v,w}(q)$ from [Theorem 1.4](#) that we prove in [Section 5](#).

Given any box $(i, j) \in \lambda(w)$, let (\hat{i}, j) be the topmost box in column j .

Let $\mathbf{b} = \mathbf{b}(\Theta_{v,w})$ be defined by Equations (1-2) (or see the proof of [Theorem 1.7](#), page 606). Define

$$\Psi : \text{drift}(v, w) \rightarrow \text{SSYT}(\lambda(w), \mathbf{b})$$

by sending a drift configuration \mathcal{D} to the semistandard tableau T , as follows. For each special box $(i, j) \in \lambda(w)$ we fill (\hat{i}, j) with the entry $(\hat{i} + d)$, where d is the distance moved in \mathcal{D} by the continent associated to (i, j) , from $\text{Pangaea}(v, w)$. Note that the value of this entry is the height of the box (\hat{i}, j) after drifting in the drift configuration \mathcal{D} . Now fill in the remaining empty boxes of $\lambda(w)$ by working down columns, from right to left, according to the prescription

$$T(i, j) = \min\{T(i+1, j) - 1, T(i-1, j+1) + 1\}. \tag{3-1}$$

By convention, set

$$T(i, j) = \begin{cases} \infty & \text{if } i > 0 \text{ and } (i, j) \notin \lambda(w), \text{ or if } j > m, \\ 0 & \text{if } i = 0 \text{ and } j \leq m, \end{cases} \tag{3-2}$$

where m is the number of columns in $\lambda(w)$.

Example 3.1. For the five drift configurations \mathcal{D} in [Example 1.6](#), the corresponding $\Psi(\mathcal{D})$ are as follows, where the boxes (\hat{i}, j) corresponding to special boxes are underlined.

3	<u>3</u>	<u>3</u>			3	<u>3</u>	<u>4</u>			3	<u>3</u>	<u>3</u>			3	<u>3</u>	<u>4</u>			3	<u>4</u>	<u>4</u>		
2	2	2	<u>2</u>		2	2	2	<u>2</u>		2	2	2	<u>3</u>		2	2	3	<u>3</u>		2	2	3	<u>3</u>	
1	1	1	1		1	1	1	1		1	1	1	2		1	1	1	2		1	1	1	2	

We will also need the $\text{sat}(\Psi(\mathcal{D}))$, which here are

3	3	3			3	3	3,4			3	3	3			3	3	4			3	3,4	4		
2	2	2	2		2	2	2	2		2	2	2	3		2	2	2,3	3		2	2	2,3	3	
1	1	1	1		1	1	1	1		1	1	1	1,2		1	1	1	1,2		1	1	1	1,2	

Lemma 3.2. *Suppose $\mathfrak{D} \in \text{drift}(v, w)$ and $T = \Psi(\mathfrak{D})$. Then:*

- (i) *T is a semistandard Young tableau (i.e., Ψ is well-defined).*
- (ii) *Ψ is an injection.*
- (iii) *If the j -th column of $\lambda(w)$ has no special box, then $T(i, j) = i$ for all $1 \leq i \leq \widehat{i}$.*
- (iv) *$\text{wt}(\mathfrak{D}) = \text{ex}(\text{sat}(T)) = \text{depth}(T)$.*

Proof. (i) Since each corner of $\lambda(w)$ is special, it is assigned a finite number. Hence (3-1) assigns each box of $\lambda(w)$ a finite number. The column semistandardness conditions are immediate from (3-1). We now establish the row semistandardness condition $T(i, j) \leq T(i, j+1)$, considering the two cases that can occur.

Case 1: (i, j) is atop a special box. That is, there is a special box (i_0, j) with $i = \widehat{i}_0$. Then if $(i, j+1)$ is in $\lambda(w)$, it is atop another special box: Suppose not. Then let the arm and leg length of (i, j) be \mathcal{L} . Note that since $\lambda(w)$ is a Young diagram, $(i - \mathcal{L} + 1, j + \mathcal{L} + 1) \notin \lambda(w)$. Thus there is a smallest integer k such that $1 \leq k \leq \mathcal{L}$ and $(i - k + 1, j + k + 1) \notin \lambda(w)$. For this k note that $(i - k + 1, j + 1)$ has equal arm and leg length equal, no other special boxes are above it (by assumption) and no boxes to strictly to its right can be special (their leg lengths are strictly longer than their arm lengths). Hence $(i - k + 1, j + 1)$ is special, but this is a contradiction.

Now that we know that both (i, j) and $(i, j+1)$ are atop special boxes, hence $T(i, j)$ and $T(i, j+1)$ are the heights of the boxes (i, j) and $(i, j+1)$ in the drift configuration \mathfrak{D} . From this interpretation, it is clear that $T(i, j) \leq T(i, j+1)$.

Case 2: (i, j) is not atop a special box. In this situation, by (3-1),

$$T(i, j) \leq T(i-1, j+1) + 1 \leq T(i, j+1).$$

(ii) This is immediate since different drift configurations will lead to different initial fillings, of the boxes (\widehat{i}, j) where (i, j) is a special box.

(iii) First note that $(\widehat{i}, j+1), (\widehat{i}-1, j+2), (\widehat{i}-2, j+3), \dots, (1, j+\widehat{i})$ must lie in $\lambda(w)$. Otherwise suppose $k \in \mathbb{Z}_{\geq 0}$ is the smallest integer that $(\widehat{i}-k, j+k+1)$ is not in $\lambda(w)$. Since the j -th column does not contain a special box, (\widehat{i}, j) is not a corner, so $(\widehat{i}, j+1)$ must lie in $\lambda(w)$, and we have $k \geq 1$. Since k is the smallest integer where the failure occurs, $(\widehat{i}-k+1, j+k)$ must lie in $\lambda(w)$, and therefore $(\widehat{i}-k, j+k)$ lies in $\lambda(w)$. The conclusion that $(\widehat{i}-k, j)$ is deduced is a similar manner as in Case 1 of (i).

Now applying (3-1) repeatedly, we have

$$T(\widehat{i}, j) \leq T(\widehat{i}-1, j+1) + 1 \leq T(\widehat{i}-2, j+2) + 2 \leq \dots \leq T(1, j+\widehat{i}-1) + \widehat{i}-1,$$

and each of the boxes being considered actually lie in $\lambda(w)$, because of what we just argued. Since $T(1, j+\widehat{i}-1) = 1$ (which holds because $(1, j+\widehat{i}) \in \lambda(w)$ so (3-1) is assigned using the boundary value $T(0, j+\widehat{i}) = 0$), we have $T(\widehat{i}, j) \leq \widehat{i}$, which forces by the fact T is semistandard that $T(i, j) = i$ for $1 \leq i \leq \widehat{i}$.

(iv) The second equality here is just the definition; see (1-1). We establish the first equality. Consider the j -th column of $\lambda(w)$.

Case 1: this column contains a special box (i, j) . The column contains \widehat{i} boxes and so each of the numbers $1, 2, \dots, \widehat{i} + d$ appears exactly once in this column of $\text{sat}(T)$, by the definition of sat and Ψ . Hence the number of extra entries of $\text{sat}(T)$ in column j is equal to $(\widehat{i} + d) - \widehat{i} = d$, which is the same as the distance moved by the continent of (i, j) .

Case 2: the column contains no special box. By (iii), there are no extra entries in this column.

Summing up the number of extra entries in each column j of $\text{sat}(T)$, we conclude that $\text{ex}(\text{sat}(T))$ is equal to $\text{wt}(\mathcal{D})$, as desired. □

Therefore,

$$P_{v,w}(q) = \sum_{\mathcal{D} \in \text{drift}(v,w)} q^{\text{wt}(\mathcal{D})} = \sum_{\mathcal{D} \in \text{drift}(v,w)} q^{\text{depth}(\Psi(\mathcal{D}))} \leq \sum_{T \in \text{SSYT}(\lambda(w), \mathbf{b})} q^{\text{depth}(T)} = H_{v,w}(q).$$

Here the first equality holds by [Theorem 1.4\(I\)](#), the second equality is by (iv), the “ \leq ” is by (ii), and the final equality is by [Theorem 1.7](#).

It remains to prove that

$$\deg H_{v,w}(q) = \deg P_{v,w}(q).$$

Since we have already proved that $P_{v,w}(q) \leq H_{v,w}(q)$ which implies $\deg P_{v,w}(q) \leq \deg H_{v,w}(q)$, we need only to prove that $\deg H_{v,w}(q) \leq \deg P_{v,w}(q)$. To do so, we will need the following lemma.

Lemma 3.3. *An element $T \in \text{SSYT}(\lambda(w), \mathbf{b})$ is in the image of $\Psi : \text{drift}(v, w) \rightarrow \text{SSYT}(\lambda(w), \mathbf{b})$ if and only if both of the following conditions are true:*

- (a) *For any box (i, j) that is not equal to (\widehat{i}', j) for a special box (i', j) , (3-1) holds under the convention (3-2).*
- (b) *If (i, j) and (i', j') are any two special boxes with (i, j) weakly southwest of (i', j') , then*

$$T(\widehat{i}, j) - \widehat{i} \leq T(\widehat{i}', j') - \widehat{i}'.$$

Proof. Let $\mathcal{D} \in \text{drift}(v, w)$. We show that $\Psi(\mathcal{D})$ satisfies (a) and (b). The condition (a) holds by the definition of Ψ . The condition (b) follows since $T(\widehat{i}, j) - \widehat{i}$ equals the distance drifted by the continent containing (i, j) , $T(\widehat{i}', j') - \widehat{i}'$ equals the distance drifted by the continent containing (i', j') , and the continent associated to (i, j) cannot move further northeast than the continent associated to (i', j') .

Conversely, we now show that every $T \in \text{SSYT}(\lambda(w), \mathbf{b})$ satisfying (a) and (b) is in the image of Ψ . Consider the (putative) drift configuration \mathcal{D} defined as follows. To each continent of \mathcal{D} associated to a special box (i, j) , shift it northeast

by $T(\widehat{i}, j) - \widehat{i}$ units. We first prove that each continent fits inside $B(v, w)$: Consider the continent with special box (i, j) . If part of the continent is shifted out of the boundary $B(v, w)$, then by (b) there is some northeast corner of $\lambda(w)$ (i.e., a 1×1 continent) that has been pushed out of $B(v, w)$ by that part of the continent. Hence the corresponding T is not in $\text{SSYT}(\lambda(w), \mathbf{b})$, a contradiction.

Now, condition (b) guarantees that \mathcal{D} can in fact be obtained without continents overlapping. Hence $\mathcal{D} \in \text{drift}(v, w)$. Finally, by (a), we have $\Psi(\mathcal{D}) = T$. □

Given $T \in \text{SSYT}(\lambda(w), \mathbf{b})$, consider this condition:

$$\begin{aligned} &\text{Some box } (i, j) \text{ in } \lambda(w) \text{ is not a northeast} \\ &\text{corner and is such that (3-1) does not hold.} \end{aligned} \tag{3-3}$$

Suppose (3-3) holds for $T = T_0$. Suppose also that (i, j) is chosen such that j is smallest, with ties broken by taking i smallest.

A brief outline of the remainder of the proof follows. Starting from T_0 , we construct a sequence $T_1, T_2, \dots \in \text{SSYT}(\lambda(w), \mathbf{b})$ with increasing depth until we arrive at a T_k that fails (3-3). This T_k is proved to be in the image of Ψ . Then we show that $\mathcal{D} := \Psi^{-1}(T_k) \in \text{drift}(v, w)$ satisfies $\text{wt}(\mathcal{D}) \geq \text{depth}(T_0)$. From this the result follows; see (3-8).

So let $T_1 \in \text{SSYT}(\lambda(w), \mathbf{b})$ be the augmentation of T_0 obtained by setting

$$T_1(i, j) = \min\{T_0(i+1, j) - 1, T_0(i-1, j+1) + 1\} \tag{3-4}$$

and letting all other entries in T_1 be the same as in T_0 .

Now we show that $T_1 \in \text{SSYT}(\lambda(w), \mathbf{b})$. To do this, we need to check the semistandardness conditions

$$T_1(i, j-1) \leq T_1(i, j) \leq T_1(i, j+1), \tag{3-5}$$

$$T_1(i-1, j) < T_1(i, j) < T_1(i+1, j). \tag{3-6}$$

We first check (3-5). The second inequality is trivial from (3-4). For the first inequality, we have

$$\begin{aligned} T_0(i, j-1) &\leq T_0(i+1, j-1) - 1 \leq T_0(i+1, j) - 1, \\ T_0(i, j-1) &\leq T_0(i-1, j) + 1 \leq T_0(i-1, j+1) + 1. \end{aligned}$$

(The second of those lines uses the minimality of our choice of (i, j) .) Hence

$$T_1(i, j-1) = T_0(i, j-1) \leq \min\{T_0(i+1, j) - 1, T_0(i-1, j+1) + 1\} = T_1(i, j).$$

Similarly for (3-6): the second inequality is trivial from (3-4), whereas for the first inequality, we have

$$\begin{aligned} T_0(i-1, j) &< T_0(i, j) \leq T_0(i+1, j) - 1, \\ T_0(i-1, j) &\leq T_0(i-1, j+1) < T_0(i-1, j+1) + 1, \end{aligned}$$

and hence

$$T_1(i-1, j) = T_0(i-1, j) < \min\{T_0(i+1, j) - 1, T_0(i-1, j+1) + 1\} = T_1(i, j).$$

Next, we claim that

$$\text{depth}(T_1) \geq \text{depth}(T_0).$$

The difference in depth between T_1 and T_0 can only be blamed on the boxes in positions (i, j) , $(i, j+1)$ and $(i+1, j)$. Without loss of generality, let us assume that each of the latter two boxes actually lie in $\lambda(w)$ (at least one of $(i, j+1)$ or $(i+1, j)$ is in $\lambda(w)$ since (i, j) is assumed to not be a northeast corner; analyzing the resulting cases is similar and easier). Taking this into account leads to

$$\begin{aligned} \text{depth}(T_1) - \text{depth}(T_0) = & T_1(i, j) - T_0(i, j) + \min\{T_1(i, j+1) - T_1(i, j), T_1(i, j+1) - T_1(i-1, j+1) - 1\} \\ & - \min\{T_0(i, j+1) - T_0(i, j), T_0(i, j+1) - T_0(i-1, j+1) - 1\} \\ & + \min\{T_1(i+1, j) - T_1(i+1, j-1), T_1(i+1, j) - T_1(i, j) - 1\} \\ & - \min\{T_0(i+1, j) - T_0(i+1, j-1), T_0(i+1, j) - T_0(i, j) - 1\}. \end{aligned}$$

Recall that T_0 and T_1 coincide outside of (i, j) . For simplicity, set

$$y := T_r(i+1, j), \quad z := T_r(i, j+1), \quad u := T_r(i+1, j-1), \quad v := T_r(i-1, j+1),$$

for $r = 0, 1$. Also let

$$x := T_0(i, j), \quad x' := T_1(i, j) = \min(y-1, v+1).$$

Using $\min(a, b) = \frac{1}{2}(a+b-|a-b|)$, this gives

$$\begin{aligned} \text{depth}(T_1) - \text{depth}(T_0) &= x' - x + \min(z - x', z - v - 1) - \min(z - x, z - v - 1) \\ &\quad + \min(y - x' - 1, y - u) - \min(y - x - 1, y - u) \\ &= x' - x + \frac{1}{2}(2z - x' - v - 1 - |x' - v - 1|) - \frac{1}{2}(2z - x - v - 1 - |x - v - 1|) \\ &\quad + \frac{1}{2}(2y - x' - u - 1 - |x' - u + 1|) - \frac{1}{2}(2y - x - u - 1 - |x - u + 1|) \\ &= \frac{1}{2}((|x - u + 1| + |x - v - 1|) - (|x' - u + 1| + |x' - v - 1|)) \\ &= \frac{1}{2}(f(x) - f(x')), \end{aligned}$$

where

$$f(a) := |a - u + 1| + |a - v - 1|.$$

It is elementary that $f(a)$ takes the minimal value throughout the (real) interval

$$[\min(v+1, u-1), \max(v+1, u-1)].$$

Notice that x' is in this interval: $x' \geq \min(v+1, u-1)$ since $y \geq u$. On the other hand, $x' \leq v+1 \leq \max(v+1, u-1)$. Since f attains its minimum at x' , we have $f(x) - f(x') \geq 0$, so $\text{depth}(T_1) \geq \text{depth}(T_0)$ as required.

Repeating this procedure so long as the undesirable property (3-3) still holds, we obtain successively $T_0, T_1, T_2, T_3, \dots$. We claim that after a finite number of iterations (3-3) finally fails for some $T_k, k \geq 0$. To see this, let the vector

$$\mathbf{u}(T) = (u_1, u_2, \dots, u_{|\lambda(w)|})$$

measure how far $T \in \text{SSYT}(\lambda(w), \mathbf{b})$ is from failing (3-3): Order the boxes in $\lambda(w)$ from left to right, and in each column from bottom up. For example, in Example 1.6, the order is

3	6	9	
2	5	8	11
1	4	7	10

For each $1 \leq i \leq |\lambda(w)|$, define u_i to be 0 if the i -th box is a northeast corner or if (3-1) holds; otherwise let $u_i = 1$. Then $\mathbf{u}(T) = (0, 0, \dots, 0)$ means that we are in the good case that (3-3) fails. We define a pure reverse lex order on $\{0, 1\}^{|\lambda(w)|}$: given $\mathbf{u}, \mathbf{u}' \in \{0, 1\}^{|\lambda(w)|}$, we say that $\mathbf{u} > \mathbf{u}'$ if

$$u_{|\lambda(w)|} = u'_{|\lambda(w)|}, \quad u_{|\lambda(w)|-1} = u'_{|\lambda(w)|-1}, \quad \dots, \quad u_{i+1} = u'_{i+1}, \quad u_i > u'_i,$$

for some i . It is straightforward to check that $\mathbf{u}(T_t) > \mathbf{u}(T_{t+1})$ at each step t , so the procedure must eventually terminate, say at step k , with $\mathbf{u}(T_k) = (0, 0, \dots, 0)$, as desired.

Let $T = T_k$ be the output of the procedure above. We want to apply Lemma 3.3 to conclude that $T_k(i, j)$ is in the image of Ψ . We must verify conditions (a) and (b) of the lemma.

Since (3-3) fails, every box that is not a northeast corner has (3-1) holding. In particular, this includes every box described by (a), and so (a) holds.

To check (b), let $\mathcal{L} := \widehat{i} - i$ be the leg length of (i, j) . Since (i, j) is special, $\mathcal{L} = |\text{arm}(i, j)|$; moreover, we can apply the argument in the proof of Lemma 3.2(iii) to the subset of the Young diagram $\lambda(w)$ consisting of those boxes strictly above row i and weakly to the right of column j , and conclude that the following boxes lie in $\lambda(w)$:

$$(\widehat{i}, j+1), \quad (\widehat{i}-1, j+2), \quad \dots, \quad (\widehat{i}-\mathcal{L}+1, j+\mathcal{L}).$$

In particular, the boxes

$$(\widehat{i}, j), \quad (\widehat{i}-1, j+1), \quad (\widehat{i}-2, j+2), \quad \dots, \quad (\widehat{i}-\mathcal{L}, j+\mathcal{L})$$

are not the northeast corners of $\lambda(w)$; hence (3-1) holds for them by the construction of $T = T_k$. By (3-1), we have

$$T(\widehat{i}-m, j+m) \geq T(\widehat{i}, j) - m, \quad \text{for } m = 0, 1, \dots, \mathcal{L}. \quad (3-7)$$

Since (\widehat{i}', j') is to the right of $(\widehat{i}, j + (\widehat{i} - \widehat{i}'))$, we have

$$T(\widehat{i}', j') \geq T(\widehat{i}', j + (\widehat{i} - \widehat{i}')) = T(\widehat{i} - (\widehat{i} - \widehat{i}'), j + (\widehat{i} - \widehat{i}')) \geq T(\widehat{i}, j) - (\widehat{i} - \widehat{i}'),$$

where the last inequality holds because of (3-7) for $m = \widehat{i} - \widehat{i}'$, and since the hypothesis that (i, j) is weakly southwest of (i', j') implies $\widehat{i} - \widehat{i}' \leq \mathcal{L} - 1$. Thus,

$$T(\widehat{i}, j) - \widehat{i} \leq T(\widehat{i}', j') - \widehat{i}'.$$

Therefore condition (b) holds.

Concluding, there exists $\mathcal{D} \in \text{drift}(v, w)$ such that $\Psi(\mathcal{D}) = T_k$ and $\text{wt}(\mathcal{D}) = \text{depth}(T_k)$. Then

$$\text{wt}(\mathcal{D}) = \text{depth}(T_k) \geq \text{depth}(T_{k-1}) \geq \dots \geq \text{depth}(T_0) \quad (3-8)$$

and so $\deg P_{v,w}(q) \geq \deg H_{v,w}(q)$, as was to be shown. \square

4. A ball of drift configurations


Construction of $\text{KL}_{v,w}$. In order to emphasize the combinatorial relations of drift configurations to Young tableaux, consider an equivalent formulation of drift configurations: A *semistandard (ordinary) drift tableau* T bijectively associated to \mathcal{D} is a filling of each continent C of $\text{Pangaea}(v, w)$ by the distance C has moved from $\text{Pangaea}(v, w)$.

Similarly, a *set-valued drift tableau* is a filling of each continent by some non-empty set of nonnegative integers; it is *semistandard* if any ordinary drift tableau it contains (in the obvious sense) is semistandard. It is *limit semistandard* if it contains at least one semistandard (ordinary) drift tableau. The *empty-face drift tableau* $\mathcal{E}_{v,w}$ is the set-valued drift tableau that is the union of all semistandard ordinary ones.

Define $\text{KL}_{v,w}$ to be the simplicial complex whose faces are indexed by limit semistandard drift tableau and where face containment is by reverse containment of drift tableau. In particular, the vertices are labeled by limit semistandard tableaux $(b \not\mapsto y)$ obtained by removing precisely one entry y from a set $\mathcal{E}_{v,w}(b)$ of the box $b \in \lambda(w)$, provided $|\mathcal{E}_{v,w}(b)| > 1$. (It will be convenient to also consider *phantom vertices* which are those $(b \not\mapsto y)$ where $|\mathcal{E}_{v,w}(b)| = 1$; these become honest vertices after coning over $\text{KL}_{v,w}$.)

This gives an example of a tableau complex in the sense of [Knutson et al. 2008]. We illustrate the case discussed in Example 1.6, showing the interior faces of the

2-dimensional complex $\text{KL}_{23465178910,10954382761}$:



The claims in [Theorem 1.4](#) about the structure of $\text{KL}_{v,w}$ then follow immediately from [[Knutson et al. 2008](#), Theorem 2.8]. We conclude that the interior faces of $\text{KL}_{v,w}$ are labeled by semistandard set-valued drift tableaux while the exterior faces are labeled by nonsemistandard but limit semistandard tableaux. Also the codimension of a face \mathcal{D} is $|\mathcal{D}| - \#\text{continents}$, the number of “extra” entries of \mathcal{D} .

K-polynomials of $\text{KL}_{v,w}$. Let us take this opportunity to formalize a connection between the K -polynomials of $\text{KL}_{v,w}$ and $P_{v,w}(q)$. We will utilize facts collected about general tableau complexes from [[Knutson et al. 2008](#), Section 4]. Let V be the set of vertices of a simplicial complex Δ and set $R = \mathbb{k}[\Delta]$ to be the polynomial ring in variables $x_{\mathfrak{v}}$ for $\mathfrak{v} \in V$. This is the ambient ring for the *Stanley–Reisner ideal* $I_{\Delta} = \langle \prod_{\mathfrak{v} \in F} x_{\mathfrak{v}} : F \text{ is not a face of } \Delta \rangle$ of Δ , and R/I_{Δ} is the *Stanley–Reisner ring*. We use the alphabet $\mathbf{t}_{\mathfrak{v}} = \{t_{\mathfrak{v}} : \mathfrak{v} \in V\}$ for the finely graded Hilbert series $\text{Hilb}(R/I_{\Delta}; \mathbf{t})$ and K -polynomials $\mathcal{K}(R/I_{\Delta}, \mathbf{t})$.

Let us define a family of polynomials for $v \leq w$, where w is covexillary. We will see this is a hybrid of the K -polynomial of $\text{KL}_{v,w}$ and the Kazhdan–Lusztig polynomial $P_{v,w}(q)$:

$$\mathfrak{P}_{v,w}(\beta; \mathbf{t}) = \sum_{\mathcal{D} \in \text{SVDT}(v,w)} \beta^{|\mathcal{D}| - \#\text{continents}(v,w)} \prod_{b \in \lambda(w)} \prod_{y \in \mathcal{D}(b)} (1 - t_{(b \not\rightarrow y)}), \quad (4-1)$$

where $\text{SVDT}(v,w)$ is the set of set-valued drift tableaux associated to drift configurations in $\text{drift}(v,w)$, $|\mathcal{D}|$ is the number of entries in \mathcal{D} , and $\#\text{continents}(v,w)$ is the number of continents in $\text{Pangaea}(v,w)$. There are a number of interesting specializations of this polynomial. Here we do not assume $|\mathcal{E}_{v,w}(b)| > 1$, i.e., $(b \not\rightarrow y)$ might be a phantom vertex.

By the ballness/sphereness claim of $\text{KL}_{v,w}$ from [Theorem 1.4](#), together with [[Knutson et al. 2008](#), Theorem 4.3], it follows that

$$\mathfrak{P}_{v,w}(-1; \mathbf{t}) = \mathfrak{H}(R/I_{\text{KL}_{v,w}}; \mathbf{t}). \tag{4-2}$$

One can consider a vertex decomposition of any complex Δ at a vertex v . This is given by $\Delta = \text{del}_v(\Delta) \cup \text{star}_v(\Delta)$, where $\text{del}_v(\Delta) = \{F \in \Delta : v \notin F\}$ is the *deletion* of v and $\text{star}_v(\Delta) = \{F \in \Delta : F \cup \{v\} \in \Delta\}$ is the *star* of v . Automatically one has, for $v = (b \not\rightarrow y)$,

$$\begin{aligned} &\mathfrak{H}(R/I_{\text{KL}_{v,w}}; \mathbf{t}) \\ &= t_{(b \not\rightarrow y)} \mathfrak{H}(R/I_{\text{del}_{(b \not\rightarrow y)}(\text{KL}_{v,w})}; \mathbf{t}) + (1 - t_{(b \not\rightarrow y)}) \mathfrak{H}(R/I_{\text{star}_{(b \not\rightarrow y)}(\text{KL}_{v,w})}; \mathbf{t}). \end{aligned} \tag{4-3}$$

By tracing the specializations below, one should eventually interpret recursions from [Lascoux and Schützenberger 1981] for $P_{v,w}(q)$ using (4-3) and thus vertex decompositions of $\text{KL}_{v,w}$. We do not pursue this here.

Consider

$$\mathfrak{P}_{v,w}(-1; t_{(b \not\rightarrow y)} \mapsto 1 - x_y) = \sum_{\mathfrak{D} \in \text{SVDT}(v,w)} (-1)^{|\mathfrak{D}| - \#\text{continents}(v,w)} \mathbf{x}^{\mathfrak{D}}, \tag{4-4}$$

where

$$\mathbf{x}^{\mathfrak{D}} = \prod_{i \geq 0} x_i^{\#\text{'s appearing in } \mathfrak{D}}.$$

Another specialization is given by

$$\mathfrak{P}_{v,w}(0; t_{(b \not\rightarrow y)} \mapsto 1 - x_y) = \sum_{\mathfrak{D} \in \text{SSDT}(v,w)} \mathbf{x}^{\mathfrak{D}}, \tag{4-5}$$

where $\text{SSDT}(v, w)$ is the set of ordinary, semistandard drift tableau associated to v, w . (In setting $\beta = 0$ we take the convention that $0^0 = 1$ in (4-1).)

Finally, by considering the principal specialization of (4-5) we have

$$\mathfrak{P}_{v,w}(0; t_{(b \not\rightarrow y)} \mapsto 1 - q^y) = P_{v,w}(q).$$

5. The proof of Theorem 1.4(I)

Proof that $Q_{v,w}(q) = P_{v,w}(q)$. We give a weight-preserving bijection between $\text{drift}(v, w)$ and the trees weight-enumerated by Lascoux’s rule [1995] for $P_{v,w}(q)$. We mostly follow the presentation of his rule found in [Billiey and Lakshmibai 2000, 6.3.29].

Given $\mathfrak{D} \in \text{drift}(v, w)$, construct a rooted, edge-labeled tree \mathcal{T} as follows. Associate to each continent C a non-root vertex of \mathcal{T} . Moreover if the special box b of C is southwest of the special box b' of an adjacent continent C' , then we draw an edge between the corresponding vertices. If there is no special box strictly southwest of b , then the corresponding vertex is joined to the root of \mathcal{T} .


Thus, each 1×1 continent $C = \{(h, \lambda(w)_h)\}$ (equivalently, those that come from northeast corners of $\lambda(w)$) corresponds to a leaf p of \mathcal{T} . Now we bound the edge

incident to p by $b_h - h$, where

$$b_h = \max\{m : B(v, w)_m \geq \lambda(w)_h + m - h\}.$$

Let $DL(\mathcal{T})$ be the set of all edge labelings of \mathcal{T} by nonnegative integers such that the labels weakly increase from root to leaf. For any edge labeled tree \mathcal{G} let $|\mathcal{G}|$ be the sum of the edge labels of \mathcal{G} .

As an example, here are the edge-labeled trees for the drift configurations in [Example 1.6](#). (The framed number below each leaf is the bound for that leaf.)



Lemma 5.1. *There is a bijection $\Phi : \text{drift}(v, w) \rightarrow DL(\mathcal{T})$ such that*

$$\text{wt}(\mathcal{D}) = |\Phi(\mathcal{D})|.$$

Proof. Define $\Phi(\mathcal{D})$ to be the edge labeling of \mathcal{T} such that the edge associated to a continent C (i.e., the edge whose child end is the vertex associated to C) is labeled by the distance that C has drifted in \mathcal{D} . That the labels are weakly increasing in $\Phi(\mathcal{D})$ is implied by the condition that the continents do not overlap in \mathcal{D} . Note that if C is a 1×1 continent then $b_h - h$ is the largest distance that C can drift inside $B(v, w)$; this accounts for the leaf bound. (For an example, see diagram immediately above.) It is then easy to check that Φ is the desired bijection. \square

Lascoux’s rule constructs a tree \mathcal{T}' as follows: For the partition $\lambda(w)$, the *parenthesis-word* is a word using “(” and “)” and obtained by walking with east and south steps along the northeast border of $\lambda(w)$. We record a “(” for each east step and a “)” for each south step. Now pair left and right parentheses starting from the closest pairs “()”. Each pair corresponds to a vertex of the tree; the closest pairs are associated to leaves and a pair encloses its children. Unpaired parentheses do not contribute to the tree. This process results in a directed forest. Finally, we introduce an additional root and attach an edge to the root of each tree in the forest.

Lemma 5.2. *There is a graph isomorphism $\delta : \mathcal{T} \rightarrow \mathcal{T}'$. Under this isomorphism, if v corresponds to a 1×1 continent associated to a corner c of $\lambda(w)$, then $\delta(v)$ corresponds to a closest parenthesis pair associated to the same corner c .*


Proof. Each leaf of \mathcal{T} corresponds to a corner c of $\lambda(w)$. On the other hand, this corner gives rise to a closest pair “()” in Lascoux’s construction, which corresponds to a leaf of \mathcal{T}' . Thus we can construct a bijection between the leaves of the

two trees, which we now argue extends to the bijection δ between the two trees themselves.

A continent C is a z -continent if it is defined by a z -special box b . Fix a vertex $v \in \mathcal{T}$ associated to such a continent. By construction, each child of v is a vertex $\{v'\}$ associated to a y -continent C' adjacent and northeast of C in $\text{Pangaea}(v, w)$, where $y < z$. Since $b \in C$ is a special box, by using the fact that $|\text{arm}(b)| = |\text{leg}(b)|$ we have that the column b is in corresponds to a “(” and the row b is in corresponds to a “)”, where these two parentheses are paired with one another in the parenthesis word. Clearly, this pair gives a vertex $v' \in \mathcal{T}'$, and all vertices of \mathcal{T}' arise this way. That is, there is a bijection at the level of vertices $\delta : \mathcal{T} \rightarrow \mathcal{T}'$. Moreover, that the children of $\delta(v)$ are exactly $\{\delta(v')\}$ (for children v' of v) is also immediate from the constructions of \mathcal{T} and \mathcal{T}' \square

Lascoux’s rule similarly defines increasing edge labelings $EL(\mathcal{T})$ on \mathcal{T} as we did for $DL(\mathcal{T})$. It remains to check that these labelings are the same as the ones in $DL(\mathcal{T})$. For this, we only need to show that the bound attached to the leaves are the same. In [Billey and Lakshmibai 2000, 6.3.29, Step 2], for each given leaf, a bigrassmannian permutation is determined in three substeps, from which Lascoux’s leaf bounds are determined. We now explain these steps. (For readers comparing what follows with that reference, note that Billey and Lakshmibai’s x is our $\tilde{w} = w^{-1}w_0$, while their w is our $\tilde{v} = v^{-1}w_0$.)

The reader may find the following diagram useful for the description of Lascoux’s labeling process:



Substep (1): leaves p of \mathcal{T} correspond to distinct numbers in the code of \tilde{w} . The code (c_1, \dots, c_n) of \tilde{w} is given by

$$c_i = \#\{j > i \mid \tilde{w}_j < \tilde{w}_i\} = \#\{\text{boxes of } D(w) \text{ in row } i\}.$$

Recall $\lambda(w)$ is the result of sorting this code into decreasing order. A leaf p of \mathcal{T} corresponds to a corner $e'' = (h, \lambda(w)_h)$ of $\lambda(w)$. Associate $\lambda(w)_h$ to p . This $\lambda(w)_h$ is equal to c_i for some i . Clearly a different c_i is assigned to each p .

Substep (2): $\lambda(w)_h$ gives a crossing of \tilde{w} . By definition, a *crossing* of \tilde{w} is a 4-tuple $(i, j, j + 1, k)$ satisfying

$$\tilde{w}_{j+1} \leq \tilde{w}_k < \tilde{w}_i \leq \tilde{w}_j, \quad \tilde{w}_i = \tilde{w}_k + 1 \quad \text{for } i \leq j < k; \tag{5-1}$$

see [Lascoux and Schützenberger 1996]. Now given the ϵ'' associated to p , there is a unique essential box ϵ in $D(w)$ that is diagonally northeast of ϵ'' . We define j and k by declaring that the coordinates of ϵ are (j, \tilde{w}_k) . Let i be such that $\tilde{w}_i = \tilde{w}_k + 1$.

We claim that $(i, j, j + 1, k)$ forms a crossing. Let us first check the weak inequalities of $\tilde{w}_{j+1} \leq \tilde{w}_k < \tilde{w}_i \leq \tilde{w}_j$ (the strict inequality being true by definition). For the rightmost inequality, we have $\tilde{w}_j = w^{-1}w_0(j) = w_{n-j+1}^{-1}$, which in words is the column position of the \bullet of $G(w)$ that necessarily must be to the right of ϵ , which itself is in column \tilde{w}_k . In other words $\tilde{w}_k \leq \tilde{w}_j$. Now, for the leftmost inequality, note $\tilde{w}_{j+1} = w^{-1}w_0(j + 1) = w^{-1}(n - j)$ which is the column position of the \bullet of $G(w)$ in row $j + 1$. Since ϵ is an essential box, that \bullet must be weakly to the left, i.e., $\tilde{w}_{j+1} \leq \tilde{w}_k$, as desired. It remains to check $i \leq j$ and $j < k$. For the former inequality, we compute $w\tilde{w}_i = n - i + 1$ which is the row position of the \bullet of $G(w)$ in column \tilde{w}_i . Since ϵ is an essential box, the \bullet is weakly below the ϵ , i.e., $i \leq j$. Similarly, for the latter inequality, we consider $w\tilde{w}_k = n - k + 1$, which is the position of the \bullet of $G(w)$ in column \tilde{w}_k . This must be strictly above the ϵ , i.e., $j < k$.

Now associate the crossing $(i, j, j + 1, k)$ to p (and hence $\lambda(w)_h$). Actually, the description in [Billey and Lakshmibai 2000] gives a different way to assign a crossing to p . However, it is straightforward to check that their crossing is same as the one described above.

Substep (3): each crossing gives a maximal bigrassmannian $[a, b, c, d]$ below \tilde{w} . Here $[a, b, c, d]$ denotes

$$(1, \dots, a, a + c + 1, \dots, a + c + b, a + 1, \dots, a + c, a + c + b + 1, \dots, a + b + c + d) \in S_n.$$

Lascoux’s rule associates to $(i, j, j + 1, k)$ a maximal bigrassmannian

$$[z, j - z, \tilde{w}_k - z, n - \tilde{w}_k - j + z],$$

where

$$z = \#\{p < j : \tilde{w}_p < \tilde{w}_k\}.$$

Notice that z is the number of \bullet ’s in $G(w)$ weakly southwest of $\epsilon = (j, \tilde{w}_k)$, i.e.,

$$z = r_\epsilon^w. \tag{5-2}$$

This concludes substep (3) of step 2 of [Billey and Lakshmibai 2000].

Lascoux’s rule then assigns to p the leaf bound

$$\text{distance}([z, j - z, \tilde{w}_k - z, n - \tilde{w}_k - j + z], \tilde{v}),$$

where

$$\text{distance}([a, b, c, d], \tilde{v}) = \max\{r \geq 0 \mid [a - r, b + r, c + r, d - r] \leq \tilde{v}\},$$

and where “ \leq ” refers to Bruhat order on S_n . This completes the description of Lascoux’s algorithm.

Recall that $r_{(a+b, a+c)}^v$ equals the number of dots of $G(v)$ weakly southwest of $(a + b, a + c)$. The proof of the following fact is straightforward to argue (and also follows from the deeper developments in [Lascoux and Schützenberger 1996]):

Lemma 5.3. *For any bigrassmannian permutation $[a, b, c, d]$ and permutation \tilde{v} in S_n , the inequality $[a, b, c, d] \leq \tilde{v}$ is equivalent to $r_{(a+b, a+c)}^v \leq a$, where $v = w_0 \tilde{v}^{-1}$. \square*

Proposition 5.4. *The leaf bounds on $DL(\mathcal{F})$ and $EL(\mathcal{F})$ are the same.*

Proof. By Lemma 5.3,

$$\begin{aligned} [z - r, j - z + r, \tilde{w}_k - z + r, n - \tilde{w}_k - j + z - r] \leq \tilde{v} &\iff \\ r_{(z-r)+(j-z+r), (z-r)+(\tilde{w}_k-z+r)}^v \leq z - r &\iff \\ r_{(j, \tilde{w}_k)}^v \leq z - r &\iff \\ r_{\epsilon}^v \leq z - r. & \end{aligned} \tag{5-3}$$

Hence, the maximal r such that any of the inequalities (5-3) hold is

$$r = z - r_{\epsilon}^v = r_{\epsilon}^w - r_{\epsilon}^v,$$

where we have used (5-2).

In terms of drift configurations, r is the largest distance that a corner $\epsilon'' = (h, \lambda(w)_h)$ can be moved diagonally northeast and remain in $B(v, w)$ (see [Li and Yong 2011, Lemma 5.7]). By the definition of $B(v, w)$, $b_h = j - r_{\epsilon}^v$. It is also easy to check that $j = h + r_{\epsilon}^w$ (again by the same lemma). Then

$$b_h - h = j - r_{\epsilon}^v - h = (j - h) - r_{\epsilon}^v = r_{\epsilon}^w - r_{\epsilon}^v = r.$$

This completes the proof of the proposition. \square

By Lascoux’s rule,

$$P_{w_0 \tilde{v}, w_0 \tilde{w}}(q) (= P_{w_0 v^{-1} w_0, w_0 w^{-1} w_0}(q) = P_{v, w}(q)) = \sum q^{|T|},$$

where the sum is over $EL(T)$ and $|T|$ is the total sum of the edge labels. Since we have established the desired weight-preserving bijection, the claim $Q_{v, q}(q) = P_{v, w}(q)$ then follows.

Remark 5.5. There are two basic symmetries of Kazhdan–Lusztig polynomials: $P_{v,w}(q) = P_{w_0v^{-1}w_0, w_0w^{-1}w_0}(q)$ and $P_{v,w}(q) = P_{v^{-1}, w^{-1}}(q)$. The first symmetry is manifest in our rule and $\text{drift}(w_0v^{-1}w_0, w_0w^{-1}w_0)$ is obtained by transposing the drift configurations of $\text{drift}(v, w)$. For the second, it is an exercise to prove that $\lambda(w) = \lambda(w^{-1})$ and $B(v, w) = B(v^{-1}, w^{-1})$, so $\text{drift}(v^{-1}, w^{-1}) = \text{drift}(v, w)$.

Remark 5.6. From [Theorem 1.4\(I\)](#) it is not hard to show the following. For $w, v \in S_n$ where w is covexillary and $v \leq w$, let k be the number of special boxes of $\lambda(w)$ and let $m = \lfloor \frac{n-k+1}{2} \rfloor$. If $[m]_q = 1 + q + \dots + q^{m-1}$, then $[q^i]P_{v,w}(q) \leq [q^i]([m]_q)^k$ for all i . In particular, $P_{v,w}(1) \leq m^k$.

6. Another q -analogue of multiplicity

We can think of $H_{v,w}(q)$ as a q -analogue of Hilbert–Samuel multiplicity, in the sense that $H_{v,w}(1) = \text{mult}_{e_v}(X_w)$. Let us point out that in the covexillary setting, there is another q -analogue available. As in [Theorem 1.4\(II\)](#), regard each box of $\lambda(w)$ as a separate country; the “drift configurations” are precisely the *pipe dreams* $P \in \text{Pipes}(v, w)$ in [[Li and Yong 2011](#)]. Now let

$$\widetilde{\text{wt}}(P) = q^d,$$

where d is the total of the distance drifted by the countries, and set

$$\widetilde{H}_{v,w}(q) = \sum_{P \in \text{Pipes}(v,w)} \widetilde{\text{wt}}(P).$$

In the following theorem we use the standard q -notation:

$$[a]_q = 1 + q + \dots + q^{a-1} \quad \text{and} \quad \binom{a}{b}_q = \frac{[a]_q [a-1]_q \dots [a-b+1]_q}{[b]_q \dots [1]_q}.$$

Theorem 6.1. $\widetilde{H}_{v,w}(q) = q^{-\sum_{i \geq 1} (i-1)\lambda_i} \det \left(\binom{b_i + \lambda_i - i + j - 1}{\lambda_i - i + j} \right)_{1 \leq i, j \leq \ell(\lambda)}$,
 where $\ell(\lambda)$ is the number of nonzero parts of λ and $\mathbf{b} = \mathbf{b}(\Theta_{v,w})$.

Proof. For brevity, we refer the reader to the setup of [[Li and Yong 2011](#), Sections 5.2 and 6.2]. Notice that

$$s_{\lambda, \mathbf{b}}(1, q, q^2, q^3, \dots) = \det \left(\binom{b_i + \lambda_i - i + j - 1}{\lambda_i - i + j} \right)_{1 \leq i, j \leq \ell(\lambda)}$$

where the left-hand side of the equality is the principal specialization of the (single) flagged Schur polynomial for shape $\lambda(w)$ with flag $\mathbf{b} = \mathbf{b}(\Theta_{v,w})$.

Given a pipe dream $P \in \text{Pipes}(v, w)$ that corresponds to a flagged semistandard Young tableau T , write

$$\text{wt}_x(P) := \text{wt}_x(T)$$

to mean the usual multivariate weight assigned to T (that is, the one such that $s_{\lambda, \mathbf{b}}(x_1, x_2, x_3, \dots) = \sum_T \text{wt}_x(T)$). Let $\text{wt}'_q(P)$ be the principal specialization of $\text{wt}_x(P)$ given by $x_i \mapsto q^{i-1}$ and finally set

$$\text{wt}_q(P) = q^{-\sum_{i \geq 1} (i-1)\lambda_i} \times \text{wt}'_q(P).$$

It remains to show that $\text{wt}_q(P) = \tilde{\text{wt}}(P)$ for each P . To do this, let us induct on $\tilde{\text{wt}}(P) \geq 0$. The base case that $\tilde{\text{wt}}(P) = 0$, i.e., where P is the starting configuration holds since $\text{wt}'_q(P) = q^{\sum_{i \geq 1} (i-1)\lambda_i}$.

Now suppose $\tilde{\text{wt}}(P) > 0$. Then there is a P' such that a move of the form

$$\begin{array}{ccc} \cdot & \cdot & \\ + & \cdot & \mapsto & \cdot & + \\ & & & \cdot & \cdot \end{array}$$

in some 2×2 subsquare of $[n] \times [n]$ brought us to P (and no other $+$ in P' has changed). Thus, we can compare $\text{wt}_x(P')$ and $\text{wt}_x(P)$: the latter only differs from the former in that some factor of x_i changed to x_{i+1} (where i and $i+1$ are the rows changed by the move above). Hence applying induction we have

$$\text{wt}_q(P) = \text{wt}_q(P') \times q = \tilde{\text{wt}}(P') \times q = \tilde{\text{wt}}(P),$$

as desired. □

It is clear from [Theorem 1.4](#) that

$$P_{v,w}(q) \leq \tilde{H}_{v,w}(q).$$

With the same proof that we used for $H_{v,w}(q)$, one shows that $\tilde{H}_{v,w}(q)$ is upper semicontinuous. However, in general $\tilde{H}_{v,w}(q) \neq H_{v,w}(q)$. Moreover, we do not know any algebraic/geometric measure for general Schubert varieties that specializes to $\tilde{H}_{v,w}(q)$.

7. Concluding remarks

We are presently unaware of any geometric proof of the inequality of [Theorem 1.2](#). For general Y , let us assume, for simplicity of our discussion, that all odd local intersection cohomology groups vanish, and set

$$P_{p,Y}(q) = \sum_{i \geq 0} \dim(\mathcal{H}_p^{2i}(Y))q^i.$$

Question 7.1. Under what assumptions is either the inequality $P_{p,Y}(q) \leq H_{p,Y}(q)$ and/or the weaker inequality $P_{p,Y}(1) \leq H_{p,Y}(1) (= \text{mult}_p(Y))$ true?

Our results on $H_{v,w}(q)$ are based on the degeneration, flat over $\text{Spec}(\mathbb{Z})$, given in [Li and Yong 2011]. Hence Theorem 1.7 is valid over a field \mathbb{k} of arbitrary characteristic and Conjecture 1.1 seems similarly valid. However, the arguments of [Li and Yong 2011] also prove that the projectivized tangent cones of the Kazhdan–Lusztig varieties $\mathcal{N}_{v,w}$ are isomorphic to those for $\mathcal{N}_{\text{id},\Theta_{v,w}}$. It is then not hard to construct some cograssmannian v', w' with the same property. We do not know if $\mathcal{N}_{v,w}$ and any such $\mathcal{N}_{v',w'}$ are actually isomorphic, although a number of useful implications would be a consequence of this fact.

A number of formulae have been obtained for $P_{v,w}(q)$. For example, general, nonpositive formulae have been obtained in [Billera and Brenti 2011] and [Brenti 1998]. Beyond the covexillary case, few positive formulae are known; see, e.g., [Billey and Warrington 2001] (which treats the 321-hexagon avoiding case) and the references therein. It would be interesting to try to extend our main theorems to these other contexts as well.

Finally, we believe many of the ideas of this paper can be extended to other Lie groups. In particular, we expect Theorems 1.2, 1.4 and 1.7 to have analogues for (co)minuscule G/P ; cf. [Boe 1988]. However, this requires sufficient technicalities that it is better left to a separate treatment.

Acknowledgements

We thank Sara Billey, Xuhua He, Hiroshi Naruse and Alexander Woo for useful suggestions and questions that inspired this work. We also thank Jonah Blasiak, Allen Knutson, Venkatramani Lakshmibai, Ezra Miller, Greg Warrington and the anonymous referee for helpful comments. AY is partially supported by NSF grants DMS-0601010 and DMS-0901331.

References

- [Billera and Brenti 2011] L. J. Billera and F. Brenti, “Quasisymmetric functions and Kazhdan–Lusztig polynomials”, *Israel J. Math.* **184** (2011), 317–348. MR 2823980
- [Billey and Lakshmibai 2000] S. Billey and V. Lakshmibai, *Singular loci of Schubert varieties*, Progress in Mathematics **182**, Birkhäuser, Boston, MA, 2000. MR 2001j:14065 Zbl 0959.14032
- [Billey and Warrington 2001] S. C. Billey and G. S. Warrington, “Kazhdan–Lusztig polynomials for 321-hexagon-avoiding permutations”, *J. Algebraic Combin.* **13**:2 (2001), 111–136. MR 2002f:05161 Zbl 0979.05109
- [Boe 1988] B. D. Boe, “Kazhdan–Lusztig polynomials for Hermitian symmetric spaces”, *Trans. Amer. Math. Soc.* **309**:1 (1988), 279–294. MR 89i:22024 Zbl 0669.17009
- [Braden and MacPherson 2001] T. Braden and R. MacPherson, “From moment graphs to intersection cohomology”, *Math. Ann.* **321**:3 (2001), 533–551. MR 2003g:14030 Zbl 1077.14522
- [Brenti 1994] F. Brenti, “Log-concave and unimodal sequences in algebra, combinatorics, and geometry: an update”, pp. 71–89 in *Jerusalem combinatorics '93*, edited by H. Barcelo and G. Kalai, Contemp. Math. **178**, Amer. Math. Soc., Providence, RI, 1994. MR 95j:05026 Zbl 0813.05007

- [Brenti 1998] F. Brenti, “Lattice paths and Kazhdan–Lusztig polynomials”, *J. Amer. Math. Soc.* **11**:2 (1998), 229–259. [MR 98h:20072](#) [Zbl 0904.20033](#)
- [Brion 2004] M. Brion, “Lectures on the geometry of flag varieties”, lecture notes, Summer school on Schubert Varieties, (Warsaw, 2003), 2004. [arXiv math/0410240](#)
- [Bruns and Herzog 1993] W. Bruns and J. Herzog, *Cohen–Macaulay rings*, Cambridge Studies in Advanced Mathematics **39**, Cambridge University Press, Cambridge, 1993. [MR 95h:13020](#) [Zbl 0788.13005](#)
- [Buch 2002] A. S. Buch, “A Littlewood–Richardson rule for the K -theory of Grassmannians”, *Acta Math.* **189**:1 (2002), 37–78. [MR 2003j:14062](#) [Zbl 1090.14015](#)
- [Ikeda and Naruse 2009] T. Ikeda and H. Naruse, “Excited Young diagrams and equivariant Schubert calculus”, *Trans. Amer. Math. Soc.* **361**:10 (2009), 5193–5221. [MR 2010i:05351](#) [Zbl 05613028](#)
- [Irving 1988] R. S. Irving, “The socle filtration of a Verma module”, *Ann. Sci. École Norm. Sup.* (4) **21**:1 (1988), 47–65. [MR 89h:17015](#) [Zbl 0706.17006](#)
- [Jones and Woo 2010] B. Jones and A. Woo, “Kazhdan–Lusztig polynomials for cograssmannian permutations”, preprint, 2010.
- [Kazhdan and Lusztig 1979] D. Kazhdan and G. Lusztig, “Representations of Coxeter groups and Hecke algebras”, *Invent. Math.* **53**:2 (1979), 165–184. [MR 81j:20066](#) [Zbl 0499.20035](#)
- [Kazhdan and Lusztig 1980] D. Kazhdan and G. Lusztig, “Schubert varieties and Poincaré duality”, pp. 185–203 in *Geometry of the Laplace operator* (Honolulu, 1979), Proc. Sympos. Pure Math. **36**, Amer. Math. Soc., Providence, R.I., 1980. [MR 84g:14054](#) [Zbl 0461.14015](#)
- [Knutson 2009] A. Knutson, “Frobenius splitting, point counting, and degeneration”, preprint, 2009. [arXiv 0911.4941](#)
- [Knutson and Miller 2005] A. Knutson and E. Miller, “Gröbner geometry of Schubert polynomials”, *Ann. of Math.* (2) **161**:3 (2005), 1245–1318. [MR 2006i:05177](#)
- [Knutson et al. 2008] A. Knutson, E. Miller, and A. Yong, “Tableau complexes”, *Israel J. Math.* **163** (2008), 317–343. [MR 2009b:05264](#) [Zbl 1145.05055](#)
- [Knutson et al. 2009] A. Knutson, E. Miller, and A. Yong, “Gröbner geometry of vertex decompositions and of flagged tableaux”, *J. Reine Angew. Math.* **630** (2009), 1–31. [MR 2010m:13017](#)
- [Kreuzer and Robbiano 2005] M. Kreuzer and L. Robbiano, *Computational commutative algebra*, vol. 2, Springer, Berlin, 2005. [MR 2006h:13036](#) [Zbl 1090.13021](#)
- [Lakshmibai and Sandhya 1990] V. Lakshmibai and B. Sandhya, “Criterion for smoothness of Schubert varieties in $Sl(n)/B$ ”, *Proc. Indian Acad. Sci. Math. Sci.* **100**:1 (1990), 45–52. [MR 91c:14061](#) [Zbl 0714.14033](#)
- [Lascoux 1995] A. Lascoux, “Polynômes de Kazhdan–Lusztig pour les variétés de Schubert vexillaires”, *C. R. Acad. Sci. Paris Sér. I Math.* **321**:6 (1995), 667–670. [MR 96g:05144](#) [Zbl 0886.05117](#)
- [Lascoux and Schützenberger 1981] A. Lascoux and M.-P. Schützenberger, “Polynômes de Kazhdan & Lusztig pour les grassmanniennes”, pp. 249–266 in *Young tableaux and Schur functors in algebra and geometry* (Toruń, 1980), Astérisque **87**, Soc. Math. France, Paris, 1981. [MR 83i:14045](#)
- [Lascoux and Schützenberger 1996] A. Lascoux and M.-P. Schützenberger, “Treillis et bases des groupes de Coxeter”, *Electron. J. Combin.* **3**:2 (1996), Res. paper 27. [MR 98c:05168](#)
- [Li and Yong 2011] L. Li and A. Yong, “Some degenerations of Kazhdan–Lusztig polynomials and multiplicities of Schubert varieties”, *Adv. Math.* **229**:1 (2011), 633–667.
- [Manivel 2001] L. Manivel, *Symmetric functions, Schubert polynomials and degeneracy loci*, SMF/AMS Texts and Monographs **6**, Amer. Math. Soc., Providence, RI, 2001. [MR 2002h:05161](#) [Zbl 0998.14023](#)

- [Polo 1999] P. Polo, “Construction of arbitrary Kazhdan–Lusztig polynomials in symmetric groups”, *Represent. Theory* **3** (1999), 90–104. [MR 2000j:14079](#) [Zbl 0968.14029](#)
- [Ramanathan 1985] A. Ramanathan, “Schubert varieties are arithmetically Cohen–Macaulay”, *Invent. Math.* **80**:2 (1985), 283–294. [MR 87d:14044](#) [Zbl 0541.14039](#)
- [Rubey 2005] M. Rubey, “The h -vector of a ladder determinantal ring cogenerated by 2×2 minors is log-concave”, *J. Algebra* **292**:2 (2005), 303–323. [MR 2007a:05147](#) [Zbl 1127.13009](#)
- [Shigechi and Zinn-Justin 2010] K. Shigechi and P. Zinn-Justin, “Path representation of maximal parabolic Kazhdan–Lusztig polynomials”, preprint, 2010. [arXiv 1001.1080](#)
- [Stanley 1989] R. P. Stanley, “Log-concave and unimodal sequences in algebra, combinatorics, and geometry”, pp. 500–535 in *Graph theory and its applications: East and West* (Jinan, 1986), edited by M. F. Capobianco et al., Ann. New York Acad. Sci. **576**, New York Acad. Sci., New York, 1989. [MR 92e:05124](#) [Zbl 0792.05008](#)
- [Woo and Yong 2008] A. Woo and A. Yong, “Governing singularities of Schubert varieties”, *J. Algebra* **320**:2 (2008), 495–520. [MR 2009j:14063](#) [Zbl 1152.14046](#)
- [Woo and Yong 2009] A. Woo and A. Yong, “A Gröbner basis for Kazhdan–Lusztig ideals”, preprint, 2009. To appear in *Amer. J. Math.* [arXiv 0909.0564](#)

Communicated by Victor Reiner

Received 2010-06-19

Revised 2010-08-22

Accepted 2010-10-01

li2345@oakland.edu

*Department of Mathematics and Statistics,
Oakland University, Rochester, MI 48309, United States*

ayong@math.uiuc.edu

*Department of Mathematics, University of Illinois at
Urbana–Champaign, Urbana, IL 61801, United States*

Algebra & Number Theory

www.jant.org

EDITORS

MANAGING EDITOR

Bjorn Poonen
Massachusetts Institute of Technology
Cambridge, USA

EDITORIAL BOARD CHAIR

David Eisenbud
University of California
Berkeley, USA

BOARD OF EDITORS

Georgia Benkart	University of Wisconsin, Madison, USA	Shigefumi Mori	RIMS, Kyoto University, Japan
Dave Benson	University of Aberdeen, Scotland	Raman Parimala	Emory University, USA
Richard E. Borcherds	University of California, Berkeley, USA	Jonathan Pila	University of Oxford, UK
John H. Coates	University of Cambridge, UK	Victor Reiner	University of Minnesota, USA
J-L. Colliot-Thélène	CNRS, Université Paris-Sud, France	Karl Rubin	University of California, Irvine, USA
Brian D. Conrad	University of Michigan, USA	Peter Sarnak	Princeton University, USA
Hélène Esnault	Universität Duisburg-Essen, Germany	Joseph H. Silverman	Brown University, USA
Hubert Flenner	Ruhr-Universität, Germany	Michael Singer	North Carolina State University, USA
Edward Frenkel	University of California, Berkeley, USA	Ronald Solomon	Ohio State University, USA
Andrew Granville	Université de Montréal, Canada	Vasudevan Srinivas	Tata Inst. of Fund. Research, India
Joseph Gubeladze	San Francisco State University, USA	J. Toby Stafford	University of Michigan, USA
Ehud Hrushovski	Hebrew University, Israel	Bernd Sturmfels	University of California, Berkeley, USA
Craig Huneke	University of Kansas, USA	Richard Taylor	Harvard University, USA
Mikhail Kapranov	Yale University, USA	Ravi Vakil	Stanford University, USA
Yujiro Kawamata	University of Tokyo, Japan	Michel van den Bergh	Hasselt University, Belgium
János Kollár	Princeton University, USA	Marie-France Vignéras	Université Paris VII, France
Yuri Manin	Northwestern University, USA	Kei-Ichi Watanabe	Nihon University, Japan
Barry Mazur	Harvard University, USA	Andrei Zelevinsky	Northeastern University, USA
Philippe Michel	École Polytechnique Fédérale de Lausanne	Efim Zelmanov	University of California, San Diego, USA
Susan Montgomery	University of Southern California, USA		

PRODUCTION

contact@msp.org

Silvio Levy, Scientific Editor

See inside back cover or www.jant.org for submission instructions.

The subscription price for 2011 is US \$150/year for the electronic version, and \$210/year (+\$35 shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Mathematical Sciences Publishers, Department of Mathematics, University of California, Berkeley, CA 94720-3840, USA.

Algebra & Number Theory (ISSN 1937-0652) at Mathematical Sciences Publishers, Department of Mathematics, University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

ANT peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
<http://msp.org/>

A NON-PROFIT CORPORATION

Typeset in L^AT_EX

Copyright ©2011 by Mathematical Sciences Publishers

Algebra & Number Theory

Volume 5 No. 5 2011

Mutation classes of \tilde{A}_n -quivers and derived equivalence classification of cluster tilted algebras of type \tilde{A}_n JANINE BASTIAN	567
Kazhdan–Lusztig polynomials and drift configurations LI LI and ALEXANDER YONG	595
Renormalization and quantum field theory RICHARD E. BORCHERDS	627
Density of rational points on isotrivial rational elliptic surfaces ANTHONY VÁRILLY-ALVARADO	659