

Algebra & Number Theory

Volume 8

2014

No. 3

**Lefschetz theorem for
abelian fundamental group with modulus**

Moritz Kerz and Shuji Saito



Lefschetz theorem for abelian fundamental group with modulus

Moritz Kerz and Shuji Saito

We prove a Lefschetz hypersurface theorem for abelian fundamental groups allowing wild ramification along some divisor. In fact, we show that isomorphism holds if the degree of the hypersurface is large relative to the ramification along the divisor.

1. Statement of main results

Lefschetz hyperplane theorems represent an important technique in the study of Grothendieck's fundamental group $\pi_1(X)$ of an algebraic variety X (we omit base points for simplicity). Roughly speaking, one gets an isomorphism of the form

$$\iota_{Y/X} : \pi_1(Y) \xrightarrow{\sim} \pi_1(X)$$

for a suitable hypersurface section $Y \rightarrow X$ if $\dim(X) \geq 3$. Purely algebraic Lefschetz theorems for projective varieties satisfying certain regularity assumptions were developed in [SGA 2 1968]. The case of nonproper varieties X and Y is more intricate because one needs a precise control of the ramification at the infinite locus. We show in the present note that for the abelian quotient of the fundamental group a Lefschetz hyperplane theorem does in fact hold. Our basic technical ingredient is the higher-dimensional ramification theory of Brylinski, Kato and Matsuda, which is recalled in Section 2. We expect that there is a noncommutative analog of our Lefschetz theorem, which should have applications to ℓ -adic representations of fundamental groups, especially over finite fields as studied in [Esnault and Kerz 2012].

To formulate our main result, let X be a normal variety over a perfect field k , and let $U \subset X$ be an open subset such that $X \setminus U$ is the support of an effective Cartier divisor on X . Let D be an effective Cartier divisor on X with support in $X \setminus U$. We introduce the abelian fundamental group $\pi_1^{\text{ab}}(X, D)$ as a quotient of $\pi_1^{\text{ab}}(U)$ classifying abelian étale coverings of U with ramification bounded by D . More precisely, for an integral curve $Z \subset U$, let Z^N be the normalization of the closure of Z in X with $\phi_Z : Z^N \rightarrow X$, the natural map. Let $Z_\infty \subset Z^N$ be the finite set

MSC2010: primary 14H30; secondary 14E22.

Keywords: fundamental group, ramification, Lefschetz theorem .

of points x such that $\phi_Z(x) \notin U$. Then $\pi_1^{\text{ab}}(X, D)$ is defined as the Pontryagin dual of the group $\text{fil}_D H^1(U)$ of continuous characters $\chi : \pi_1^{\text{ab}}(U) \rightarrow \mathbb{Q}/\mathbb{Z}$ such that, for any integral curve $Z \subset U$, its restriction $\chi|_Z : \pi_1^{\text{ab}}(Z) \rightarrow \mathbb{Q}/\mathbb{Z}$ satisfies the following inequality of Cartier divisors on Z^N :

$$\sum_{y \in Z_\infty} \text{art}_y(\chi|_Z)[y] \leq \phi_Z^* D,$$

where $\text{art}_y(\chi|_Z) \in \mathbb{Z}_{\geq 0}$ is the Artin conductor of $\chi|_Z$ at $y \in Z_\infty$ and $\phi_Z^* D$ is the pullback of D by the natural map $\phi_Z : Z^N \rightarrow X$.

Such a global measure of ramification in terms of curves has been first considered by Deligne and Laumon; see [Laumon 1981].

Now assume that X is smooth projective over k (we fix a projective embedding) and that $C = X \setminus U$ is a simple normal crossing divisor. Let Y be a smooth hypersurface section such that $Y \times_X C$ is a reduced simple normal crossing divisor on Y , and write $\text{deg}(Y)$ for the degree of Y with respect to the fixed projective embedding of X . Set $E = Y \times_X D$. Then one sees from the definition that the map $Y \cap U \rightarrow U$ induces a natural map

$$\iota_{Y/X} : \pi_1^{\text{ab}}(Y, E) \rightarrow \pi_1^{\text{ab}}(X, D).$$

Our main theorem says:

Theorem 1.1. *Assume that Y is sufficiently ample for (X, D) (see Definition 3.1). If $d := \dim(X) \geq 3$, $\iota_{Y/X}$ is an isomorphism. If $d = 2$, $\iota_{Y/X}$ is surjective.*

The prime-to- p part of the theorem is due to [Schmidt and Spieß 2000], where $p = \text{ch}(k)$. Below we see that Y is sufficiently ample if $\text{deg}(Y) \gg 0$.

Corollary 1.2. *Let X be a normal proper variety over a finite field k . Then $\pi_1^{\text{ab}}(X, D)^0$ is finite, where*

$$\pi_1^{\text{ab}}(X, D)^0 = \text{Ker}(\pi_1^{\text{ab}}(X, D) \rightarrow \pi_1^{\text{ab}}(\text{Spec}(k))).$$

Proof. In case X and $X \setminus U$ satisfy the assumption of Theorem 1.1, the corollary follows from the corresponding statement for curves. The finiteness in the curves case is a consequence of class field theory. For the general case, one can take by [de Jong 1996] an alteration $f : X' \rightarrow X$ such that X' and $X' \setminus U'$ with $U' = f^{-1}(U)$ satisfy the assumption of Theorem 1.1. Then the assertion follows from the fact that the map $f_* : \pi_1^{\text{ab}}(U') \rightarrow \pi_1^{\text{ab}}(U)$ has a finite cokernel. \square

Corollary 1.2 can also be deduced from [Raskind 1995, Theorem 6.2]. It has recently been generalized to the noncommutative setting by Deligne; see [Esnault and Kerz 2012].

Theorem 1.1 is a central ingredient in our paper [Kerz and Saito 2013]. There we use it to construct a reciprocity isomorphism between a Chow group of zero

cycles with modulus and the abelian fundamental group with bounded ramification. In fact, [Theorem 1.1](#) allows us to restrict to surfaces in this construction.

2. Review of ramification theory

First we review local ramification theory. Let K denote a henselian discrete valuation field of $\text{ch}(K) = p > 0$ with the ring \mathbb{O}_K of integers and residue field κ . Let π be a prime element of \mathbb{O}_K and $\mathfrak{m}_K = (\pi) \subset \mathbb{O}_K$ the maximal ideal. By the Artin–Schreier–Witt theory, we have a natural isomorphism for $s \in \mathbb{Z}_{\geq 1}$,

$$\delta_s : W_s(K)/(1 - F)W_s(K) \xrightarrow{\sim} H^1(K, \mathbb{Z}/p^s\mathbb{Z}), \quad (2-1)$$

where $W_s(K)$ is the ring of Witt vectors of length s and F is the Frobenius. We have the Brylinski–Kato filtration indexed by integers $m \geq 0$

$$\text{fil}_m^{\text{log}} W_s(K) = \{(a_{s-1}, \dots, a_1, a_0) \in W_s(K) \mid p^i v_K(a_i) \geq -m\},$$

where v_K is the normalized valuation of K . In this paper, we use its nonlog version introduced by Matsuda [\[1997\]](#):

$$\text{fil}_m W_s(K) = \text{fil}_{m-1}^{\text{log}} W_s(K) + V^{s-s'} \text{fil}_m^{\text{log}} W_{s'}(K),$$

where $s' = \min\{s, \text{ord}_p(m)\}$. We define ramification filtrations on $H^1(K) := H^1(K, \mathbb{Q}/\mathbb{Z})$ as

$$\begin{aligned} \text{fil}_m^{\text{log}} H^1(K) &= H^1(K)\{p'\} \oplus \bigcup_{s \geq 1} \delta_s(\text{fil}_m^{\text{log}} W_s(K)) \quad (m \geq 0), \\ \text{fil}_m H^1(K) &= H^1(K)\{p'\} \oplus \bigcup_{s \geq 1} \delta_s(\text{fil}_m W_s(K)) \quad (m \geq 1), \end{aligned}$$

where $H^1(K)\{p'\}$ is the prime-to- p part of $H^1(K)$. We note that this filtration is shifted by one from the filtration of Matsuda [\[1997, Definition 3.1.1\]](#). We also let $\text{fil}_0 H^1(K)$ be the subgroup of all unramified characters.

Definition 2.1. For $\chi \in H^1(K)$, we denote the minimal m with $\chi \in \text{fil}_m H^1(K)$ by $\text{art}_K(\chi)$ and call it the Artin conductor of χ .

We have the following facts (cf. [\[Kato 1989; Matsuda 1997\]](#)):

Lemma 2.2. (1) $\text{fil}_1 H^1(K)$ is the subgroup of tamely ramified characters.

$$(2) \text{fil}_m H^1(K) \subset \text{fil}_m^{\text{log}} H^1(K) \subset \text{fil}_{m+1} H^1(K).$$

$$(3) \text{fil}_m H^1(K) = \text{fil}_{m-1}^{\text{log}} H^1(K) \text{ if } (m, p) = 1.$$

The structure of graded quotients

$$\text{gr}_m H^1(K) = \text{fil}_m H^1(K) / \text{fil}_{m-1} H^1(K) \quad (m > 1)$$

is described as follows. Let Ω_K^1 be the absolute Kähler differential module, and put

$$\text{fil}_m \Omega_K^1 = \mathfrak{m}_K^{-m} \otimes_{\mathbb{O}_K} \Omega_{\mathbb{O}_K}^1.$$

We have an isomorphism

$$\text{gr}_m \Omega_K^1 = \text{fil}_m \Omega_K^1 / \text{fil}_{m-1} \Omega_K^1 \simeq \mathfrak{m}_K^{-m} \Omega_{\mathbb{O}_K}^1 \otimes_{\mathbb{O}_K} \kappa. \tag{2-2}$$

We have the maps

$$F^s d : W_s(K) \rightarrow \Omega_K^1, \quad (a_{s-1}, \dots, a_1, a_0) \mapsto \sum_{i=0}^{s-1} a_i^{p^i-1} da_i,$$

and one can check $F^s d(\text{fil}_n W_s(K)) \subset \text{fil}_n \Omega_K^1$.

Theorem 2.3 [Matsuda 1997]. *The maps $F^s d$ factor through δ_s and induce a natural map*

$$\text{fil}_n H^1(K) \rightarrow \text{fil}_n \Omega_K^1,$$

which induces for $m > 1$ an injective map (called the refined Artin conductor for K)

$$\text{art}_K : \text{gr}_n H^1(K) \hookrightarrow \text{gr}_n \Omega_K^1. \tag{2-3}$$

Next we review global ramification theory. Let X and C be as in the introduction, and fix a Cartier divisor D with $|D| \subset C$. We recall the definition of $\pi_1^{\text{ab}}(X, D)$. We write $H^1(U)$ for the étale cohomology group $H^1(U, \mathbb{Q}/\mathbb{Z})$, which is identified with the group of continuous characters $\pi_1^{\text{ab}}(U) \rightarrow \mathbb{Q}/\mathbb{Z}$.

Definition 2.4. We define $\text{fil}_D H^1(U)$ to be the subgroup of $\chi \in H^1(U)$ satisfying this condition. For all integral curves $Z \subset X$ not contained in C , its restriction $\chi|_Z : \pi_1^{\text{ab}}(Z) \rightarrow \mathbb{Q}/\mathbb{Z}$ satisfies the following inequality of Cartier divisors on Z^N :

$$\sum_{y \in Z_\infty} \text{art}_y(\chi|_Z)[y] \leq \phi_Z^* D,$$

where $\text{art}_y(\chi|_Z) \in \mathbb{Z}_{\geq 0}$ is the Artin conductor of $\chi|_Z$ at $y \in Z_\infty$ and $\phi_Z^* D$ is the pullback of D by the natural map $\phi_Z : Z^N \rightarrow X$. Define

$$\pi_1^{\text{ab}}(X, D) = \text{Hom}(\text{fil}_D H^1(U), \mathbb{Q}/\mathbb{Z}), \tag{2-4}$$

endowed with the usual profinite topology of the dual.

For the rest of this section, we assume that X is smooth and C is a simple normal crossing. Let I be the set of generic points of C , and let $C_\lambda = \overline{\{\lambda\}}$ for $\lambda \in I$. Write

$$D = \sum_{\lambda \in I} m_\lambda C_\lambda. \tag{2-5}$$

For $\lambda \in I$, let K_λ be the henselization of $K = k(X)$ at λ . Note that K_λ is a henselian discrete valuation field with residue field $k(C_\lambda)$.

Proposition 2.5. *We have*

$$\mathrm{fil}_D H^1(U) = \mathrm{Ker} \left(H^1(U) \rightarrow \bigoplus_{\lambda \in I} H^1(K_\lambda) / \mathrm{fil}_{m_\lambda} H^1(K_\lambda) \right).$$

Proof. This is a consequence of ramification theory developed in [Kato 1989; Matsuda 1997]. See [Kerz and Saito 2013, Corollary 2.7] for a proof. \square

Proposition 2.6. *Fix $\lambda \in I$ such that $m_\lambda > 1$ in (2-5). The refined Artin conductor art_{K_λ} (cf. Theorem 2.3) induces a natural injective map*

$$\mathrm{art}_{C_\lambda} : \mathrm{fil}_D H^1(U) / \mathrm{fil}_{D-C_\lambda} H^1(U) \hookrightarrow H^0(C_\lambda, \Omega_X^1(D) \otimes_{\mathbb{O}_X} \mathbb{O}_{C_\lambda}),$$

which is compatible with pullback along maps $f : X' \rightarrow X$ of smooth varieties with the property that $f^{-1}(C)$ is a reduced simple normal crossing divisor.

Proof. This follows from the integrality result [Matsuda 1997, Corollary 4.2.2] of the refined Artin conductor. \square

Proposition 2.6 motivates us to introduce the following log-variant of $\mathrm{fil}_D H^1(U)$:

Definition 2.7. We define $\mathrm{fil}_D^{\log} H^1(U)$ as

$$\mathrm{fil}_D^{\log} H^1(U) = \mathrm{Ker} \left(H^1(U) \rightarrow \bigoplus_{\lambda \in I} H^1(K_\lambda) / \mathrm{fil}_{m_\lambda}^{\log} H^1(K_\lambda) \right).$$

Lemma 2.8. (1) $\mathrm{fil}_C H^1(U)$ is the subgroup of tamely ramified characters.

$$(2) \quad \mathrm{fil}_D H^1(U) \subset \mathrm{fil}_D^{\log} H^1(U) \subset \mathrm{fil}_{D+C} H^1(U).$$

$$(3) \quad \mathrm{fil}_D H^1(U) = \mathrm{fil}_{D-C}^{\log} H^1(U) \text{ if } (m_\lambda, p) = 1 \text{ for all } \lambda \in I.$$

Proof. This is a direct consequence of Lemma 2.2. \square

3. Proof of the main theorem

Let X be a smooth projective variety over a perfect field of characteristic $p > 0$ and $C \subset X$ a reduced simple normal crossing divisor on X . Let $j : U = X \setminus C \subset X$ be the open immersion. We use the same notation as in the last part of the previous section. Take an effective Cartier divisor

$$D = \sum_{\lambda \in I} m_\lambda C_\lambda \quad \text{with } m_\lambda \geq 0.$$

Let $I' = \{\lambda \in I \mid p \mid m_\lambda\}$, and put

$$D' = \sum_{\lambda \in I'} (m_\lambda + 1) C_\lambda + \sum_{\lambda \in I \setminus I'} m_\lambda C_\lambda.$$

Let Y be a smooth hypersurface section such that $Y \times_X C$ is a reduced simple normal crossing divisor on Y .

Definition 3.1. (1) Assuming $\dim(X) \geq 3$, we say that Y is sufficiently ample for (X, D) if the following conditions hold:

- (A1) $H^i(X, \Omega_X^d(-\Xi + Y)) = 0$ for any effective Cartier divisor $\Xi \leq D$ and for $i = d, d - 1, d - 2$.
- (A2) For any $\lambda \in I'$, we have

$$\begin{aligned} H^0(C_\lambda, \Omega_X^1(D' - Y) \otimes \mathcal{O}_{C_\lambda}) &= H^0(C_\lambda, \mathcal{O}_{C_\lambda}(D' - Y)) \\ &= H^1(C_\lambda, \mathcal{O}_{C_\lambda}(D' - 2Y)) = 0. \end{aligned}$$

(2) Assuming $\dim(X) = 2$, we say that Y is sufficiently ample for (X, D) if the following condition holds:

- (B) $H^i(X, \Omega_X^d(-\Xi + Y)) = 0$ for any effective Cartier divisor $\Xi \leq D$ and for $i = 1, 2$.

We remark that there is an integer N such that any smooth Y of degree $\geq N$ is sufficiently ample for (X, D) .

[Theorem 1.1](#) is a direct consequence of the following:

Theorem 3.2. *Let Y be sufficiently ample for (X, D) . Write $E = Y \times_X D$.*

(1) Assuming $d := \dim(X) \geq 3$, we have isomorphisms

$$\text{fil}_D H^1(U) \xrightarrow{\sim} \text{fil}_E H^1(U \cap Y) \quad \text{and} \quad \text{fil}_D^{\log} H^1(U) \xrightarrow{\sim} \text{fil}_E^{\log} H^1(U \cap Y).$$

(2) Assuming $d = 2$, we have injections

$$\text{fil}_D H^1(U) \hookrightarrow \text{fil}_E H^1(U \cap Y) \quad \text{and} \quad \text{fil}_D^{\log} H^1(U) \hookrightarrow \text{fil}_E^{\log} H^1(U \cap Y).$$

For an abelian group M , we let $M\{p'\}$ denote the prime-to- p torsion part of M .

Lemma 3.3. (1) Assuming $d := \dim(X) \geq 3$, we have an isomorphism

$$\text{fil}_D H^1(U)\{p'\} \xrightarrow{\sim} \text{fil}_E H^1(U \cap Y)\{p'\}$$

and the same isomorphism for fil_D^{\log} .

(2) Assuming $d = 2$, we have an injection

$$\text{fil}_D H^1(U)\{p'\} \hookrightarrow \text{fil}_E H^1(U \cap Y)\{p'\}$$

and the same injection for fil_D^{\log} .

Proof. Noting

$$\text{fil}_D H^1(U)\{p'\} = \text{fil}_C H^1(U)\{p'\} = \text{fil}_C^{\log} H^1(U)\{p'\} = \text{fil}_D^{\log} H^1(U)\{p'\},$$

this follows from the tame case of [Theorem 1.1](#) due to [\[Schmidt and Spieß 2000\]](#). \square

By the above lemma, [Theorem 3.2](#) is reduced to the following:

Theorem 3.4. *Let the assumption be as in [Theorem 3.2](#). Take an integer $n > 0$.*

(1) *Assuming $d := \dim(X) \geq 3$, we have isomorphisms*

$$\mathrm{fil}_D H^1(U)[p^n] \xrightarrow{\sim} \mathrm{fil}_E H^1(U \cap Y)[p^n]$$

and the same isomorphism for $\mathrm{fil}_D^{\mathrm{log}}$.

(2) *Assuming $d = 2$, we have an injection*

$$\mathrm{fil}_D H^1(U)[p^n] \hookrightarrow \mathrm{fil}_E H^1(U \cap Y)[p^n]$$

and the same injection for $\mathrm{fil}_D^{\mathrm{log}}$.

In what follows, we consider an effective Cartier divisor with $\mathbb{Z}[1/p]$ -coefficient:

$$D = \sum_{\lambda \in I} m_\lambda C_\lambda, \quad m_\lambda \in \mathbb{Z}[1/p]_{\geq 0}.$$

We put

$$[D] = \sum_{\lambda \in I} [m_\lambda] C_\lambda \quad \text{with } [m_\lambda] = \max\{i \in \mathbb{Z} \mid i \leq m_\lambda\}$$

and $\mathcal{F}(\pm D) = \mathcal{F} \otimes_{\mathbb{C}_X} \mathbb{C}_X(\pm[D])$ for an \mathbb{C}_X -module. For D as above, let $\mathrm{fil}_D^{\mathrm{log}} W_n \mathbb{C}_X$ be the subsheaf of $j_* W_n \mathbb{C}_U$ of local sections

$$\underline{a} \in W_n \mathbb{C}_U \quad \text{such that } \underline{a} \in \mathrm{fil}_{m_\lambda}^{\mathrm{log}} W_n(K_\lambda) \text{ for any } \lambda \in I,$$

where $\mathrm{fil}_{m_\lambda}^{\mathrm{log}} W_n(K_\lambda) := \mathrm{fil}_{[m_\lambda]}^{\mathrm{log}} W_n(K_\lambda)$ is defined in [Section 2](#) for the henselization K_λ of $K = k(X)$ at λ . We note

$$\mathbb{C}_X(D) = \mathrm{fil}_D^{\mathrm{log}} W_n \mathbb{C}_X \quad \text{for } n = 1.$$

The following facts are easily checked:

- The Frobenius F induces $F : \mathrm{fil}_{D/p}^{\mathrm{log}} W_n \mathbb{C}_X \rightarrow \mathrm{fil}_D^{\mathrm{log}} W_n \mathbb{C}_X$.
- The Verschiebung V induces $V : \mathrm{fil}_D^{\mathrm{log}} W_{n-1} \mathbb{C}_X \rightarrow \mathrm{fil}_{D/p}^{\mathrm{log}} W_n \mathbb{C}_X$.
- The restriction R induces $R : \mathrm{fil}_D^{\mathrm{log}} W_n \mathbb{C}_X \rightarrow \mathrm{fil}_{D/p}^{\mathrm{log}} W_{n-1} \mathbb{C}_X$.
- The following sequence is exact:

$$0 \rightarrow \mathbb{C}_X(D) \xrightarrow{V^{n-1}} \mathrm{fil}_D^{\mathrm{log}} W_n \mathbb{C}_X \xrightarrow{R} \mathrm{fil}_{D/p}^{\mathrm{log}} W_{n-1} \mathbb{C}_X \rightarrow 0. \quad (3-1)$$

We define an object $(\mathbb{Z}/p^n \mathbb{Z})_{X|D}$ of the derived category $D^b(X)$ of bounded complexes of étale sheaves on X :

$$(\mathbb{Z}/p^n \mathbb{Z})_{X|D} = \mathrm{Cone}(\mathrm{fil}_{D/p}^{\mathrm{log}} W_n \mathbb{C}_X \xrightarrow{1-F} \mathrm{fil}_D^{\mathrm{log}} W_n \mathbb{C}_X)[-1].$$

We have a distinguished triangle in $D^b(X)$:

$$(\mathbb{Z}/p^n\mathbb{Z})_{X|D} \rightarrow \mathrm{fil}_{D/p}^{\log} W_n \mathbb{O}_X \xrightarrow{1-F} \mathrm{fil}_D^{\log} W_n \mathbb{O}_X \xrightarrow{+}. \tag{3-2}$$

Lemma 3.5. *There is a distinguished triangle*

$$(\mathbb{Z}/p\mathbb{Z})_{X|D} \rightarrow (\mathbb{Z}/p^n\mathbb{Z})_{X|D} \rightarrow (\mathbb{Z}/p^{n-1}\mathbb{Z})_{X|D/p} \xrightarrow{+}.$$

Proof. The lemma follows from the commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathbb{O}_X(D/p) & \xrightarrow{V^{n-1}} & \mathrm{fil}_{D/p}^{\log} W_n \mathbb{O}_X & \xrightarrow{R} & \mathrm{fil}_{D/p^2}^{\log} W_{n-1} \mathbb{O}_X & \longrightarrow & 0 \\ & & \downarrow 1-F & & \downarrow 1-F & & \downarrow 1-F & & \\ 0 & \longrightarrow & \mathbb{O}_X(D) & \xrightarrow{V^{n-1}} & \mathrm{fil}_D^{\log} W_n \mathbb{O}_X & \xrightarrow{R} & \mathrm{fil}_{D/p}^{\log} W_{n-1} \mathbb{O}_X & \longrightarrow & 0 \quad \square \end{array}$$

Lemma 3.6. *There is a canonical isomorphism*

$$\mathrm{fil}_D^{\log} H^1(U)[p^n] \simeq H^1(X, (\mathbb{Z}/p^n\mathbb{Z})_{X|D}).$$

Proof. Noting that the restriction of $(\mathbb{Z}/p^n\mathbb{Z})_{X|D}$ to U is $\mathbb{Z}/p^n\mathbb{Z}$ on U , we have the localization exact sequence

$$H^1(X, (\mathbb{Z}/p^n\mathbb{Z})_{X|D}) \rightarrow H^1(U, \mathbb{Z}/p^n\mathbb{Z}) \rightarrow H_C^2(X, (\mathbb{Z}/p^n\mathbb{Z})_{X|D}). \tag{3-3}$$

For the generic point λ of C_λ , (3-2) gives us an exact sequence

$$\begin{aligned} H_\lambda^1(X, \mathrm{fil}_{D/p}^{\log} W_n \mathbb{O}_X) &\xrightarrow{1-F} H_\lambda^1(X, \mathrm{fil}_D^{\log} W_n \mathbb{O}_X) \\ &\rightarrow H_\lambda^2(X, (\mathbb{Z}/p^n\mathbb{Z})_{X|D}) \rightarrow H_\lambda^2(X, \mathrm{fil}_{D/p}^{\log} W_n \mathbb{O}_X). \end{aligned}$$

By [Grothendieck 1967, Corollary 3.10] and (3-1), we have

$$H_\lambda^i(X, \mathrm{fil}_{D/p}^{\log} W_n \mathbb{O}_X) = H_\lambda^i(X, \mathrm{fil}_D^{\log} W_n \mathbb{O}_X) = 0 \quad \text{for } i \geq 2$$

and

$$\begin{aligned} H_\lambda^1(X, \mathrm{fil}_{D/p}^{\log} W_n \mathbb{O}_X) &\simeq W_n(K_\lambda) / \mathrm{fil}_{m_\lambda/p}^{\log} W_n(K_\lambda), \\ H_\lambda^1(X, \mathrm{fil}_D^{\log} W_n \mathbb{O}_X) &\simeq W_n(K_\lambda) / \mathrm{fil}_{m_\lambda}^{\log} W_n(K_\lambda). \end{aligned}$$

Thus, we get

$$H_\lambda^2(X, (\mathbb{Z}/p^n\mathbb{Z})_{X|D}) \simeq H^1(K_\lambda)[p^n] / \mathrm{fil}_{m_\lambda}^{\log} H^1(K_\lambda)[p^n].$$

Hence, Lemma 3.6 follows from (3-3) and the injectivity of

$$H_C^2(X, (\mathbb{Z}/p^n\mathbb{Z})_{X|D}) \rightarrow \bigoplus_{\lambda \in I} H_\lambda^2(X, (\mathbb{Z}/p^n\mathbb{Z})_{X|D}).$$

This injectivity is a consequence of:

Claim 3.7. For $x \in C$ with $\dim(\mathbb{O}_{X,x}) \geq 2$, we have

$$H_x^2(X, (\mathbb{Z}/p^n\mathbb{Z})_{X|D}) = 0.$$

By Lemma 3.5, it suffices to show Claim 3.7 in case $n = 1$. Triangle (3-2) gives us an exact sequence

$$H_x^1(X, \mathbb{O}_X(D)) \rightarrow H_x^2(X, (\mathbb{Z}/p\mathbb{Z})_{X|D}) \rightarrow H_x^2(X, \mathbb{O}_X(D/p)) \xrightarrow{1-F} H_x^2(X, \mathbb{O}_X(D)).$$

If $\dim(\mathbb{O}_{X,x}) > 2$, $H_x^1(X, \mathbb{O}_X(D)) = 0$ and $H_x^2(X, \mathbb{O}_X(D/p)) = 0$ by [Grothendieck 1967, Corollary 3.10], which implies $H_x^2(X, (\mathbb{Z}/p\mathbb{Z})_{X|D}) = 0$ as desired.

We now assume $\dim(\mathbb{O}_{X,x}) = 2$. Let $(\mathbb{Z}/p\mathbb{Z})_X$ denote the constant sheaf $\mathbb{Z}/p\mathbb{Z}$ on X , and put

$$\mathcal{F}_{X|D} = \text{Coker}(\mathbb{O}_X(D/p) \xrightarrow{1-F} \mathbb{O}_X(D)).$$

Note that $\mathcal{F}_{X|D} = 0$ for $D = 0$. By definition, we have a distinguished triangle

$$(\mathbb{Z}/p\mathbb{Z})_X \rightarrow (\mathbb{Z}/p\mathbb{Z})_{X|D} \rightarrow \mathcal{F}_{X|D} \xrightarrow{+}.$$

By [SGA 1 1971, Exposé X, Théorème 3.1], we have $H_x^2(X, (\mathbb{Z}/p\mathbb{Z})_X) = 0$. Hence, we are reduced to showing

$$H_x^2(X, \mathcal{F}_{X|D}) = 0. \tag{3-4}$$

Without loss of generality, we can assume that D has integral coefficients. We prove (3-4) by induction on multiplicities of D reducing to the case $D = 0$. Fix an irreducible component C_λ of C with the multiplicity $m_\lambda \geq 1$ in D , and put $D' = D - C_\lambda$. We have a commutative diagram with exact rows and columns

$$\begin{array}{ccccccc} & & (\mathbb{Z}/p\mathbb{Z})_X & & (\mathbb{Z}/p\mathbb{Z})_X & & \\ & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & \mathbb{O}_X(D'/p) & \longrightarrow & \mathbb{O}_X(D/p) & \longrightarrow & \mathcal{L} \longrightarrow 0 \\ & & \downarrow 1-F & & \downarrow 1-F & & \downarrow F \\ 0 & \longrightarrow & \mathbb{O}_X(D') & \longrightarrow & \mathbb{O}_X(D) & \longrightarrow & \mathbb{O}_{C_\lambda}(D) \longrightarrow 0 \end{array}$$

Here $\mathbb{O}_{C_\lambda}(D) = \mathbb{O}_X(D) \otimes \mathbb{O}_{C_\lambda}$, and $\mathcal{L} = \mathbb{O}_{C_\lambda}(D/p)$ if $p \mid m_\lambda$, and $\mathcal{L} = 0$ otherwise. Thus, we get short exact sequences

$$\begin{aligned} 0 \rightarrow \mathcal{F}_{X|D'} \rightarrow \mathcal{F}_{X|D} \rightarrow \mathbb{O}_{C_\lambda}(D) \rightarrow 0 & \quad \text{if } p \nmid m_\lambda, \\ 0 \rightarrow \mathcal{F}_{X|D'} \rightarrow \mathcal{F}_{X|D} \rightarrow \mathbb{O}_{C_\lambda}(D)/\mathbb{O}_{C_\lambda}(D/p)^p \rightarrow 0 & \quad \text{if } p \mid m_\lambda. \end{aligned}$$

We may assume $H_x^2(X, \mathcal{F}_{X|D'}) = 0$ by the induction hypothesis. Hence, (3-4) follows from

$$H_x^2(C_\lambda, \mathbb{O}_{C_\lambda}(D)) = 0, \tag{3-5}$$

$$H_x^2(C_\lambda, \mathbb{O}_{C_\lambda}(D)/\mathbb{O}_{C_\lambda}(E)^p) = 0, \tag{3-6}$$

where we put $E = [D/p]$. We may assume $x \in C_\lambda$ so that $\dim(\mathbb{O}_{C_\lambda, x}) = 1$ by the assumption $\dim(\mathbb{O}_{X, x}) = 2$. Equation (3-5) is a consequence of [Grothendieck 1967, Corollary 3.10]. In view of an exact sequence

$$0 \rightarrow \mathbb{O}_{C_\lambda}(pE)/\mathbb{O}_{C_\lambda}(E)^p \rightarrow \mathbb{O}_{C_\lambda}(D)/\mathbb{O}_{C_\lambda}(E)^p \rightarrow \mathbb{O}_{C_\lambda}(D)/\mathbb{O}_{C_\lambda}(pE) \rightarrow 0,$$

(3-6) follows from

$$H_x^2(C_\lambda, \mathbb{O}_{C_\lambda}(pE)/\mathbb{O}_{C_\lambda}(E)^p) = 0 \quad \text{and} \quad H_x^2(C_\lambda, \mathbb{O}_{C_\lambda}(D)/\mathbb{O}_{C_\lambda}(pE)) = 0.$$

The first assertion follows from [Grothendieck 1967, Corollary 3.10] noting that $\mathbb{O}_{C_\lambda}(pE)/\mathbb{O}_{C_\lambda}(E)^p$ is a locally free $\mathbb{O}_{C_\lambda}^p$ -module. The second assertion holds since $\mathbb{O}_{C_\lambda}(D)/\mathbb{O}_{C_\lambda}(pE)$ is supported in a proper closed subscheme T of C_λ and x is a generic point of T if $x \in T$. This completes the proof of Lemma 3.6. \square

Proof of Theorem 3.4. In view of the above results, the assertions for fil_D^{\log} of Theorem 3.4(1) and (2) follow from the following:

Theorem 3.8. *Let the assumption be as in Theorem 3.2. The natural map*

$$H^1(X, (\mathbb{Z}/p^n\mathbb{Z})_{X|D}) \rightarrow H^1(Y, (\mathbb{Z}/p^n\mathbb{Z})_{Y|E})$$

is an isomorphism for $d := \dim(X) \geq 3$, and it is injective for $d = 2$.

Proof. By Lemma 3.5, we have a commutative diagram:

$$\begin{array}{ccc}
 0 & & 0 \\
 \downarrow & & \downarrow \\
 H^1(X, (\mathbb{Z}/p\mathbb{Z})_{X|D}) & \longrightarrow & H^1(Y, (\mathbb{Z}/p\mathbb{Z})_{Y|E}) \\
 \downarrow & & \downarrow \\
 H^1(X, (\mathbb{Z}/p^n\mathbb{Z})_{X|D}) & \longrightarrow & H^1(Y, (\mathbb{Z}/p^n\mathbb{Z})_{Y|D}) \\
 \downarrow & & \downarrow \\
 H^1(X, (\mathbb{Z}/p^{n-1}\mathbb{Z})_{X|D/p}) & \longrightarrow & H^1(Y, (\mathbb{Z}/p^{n-1}\mathbb{Z})_{Y|E/p}) \\
 \downarrow & & \downarrow \\
 H^2(X, (\mathbb{Z}/p\mathbb{Z})_{X|D}) & \longrightarrow & H^2(Y, (\mathbb{Z}/p\mathbb{Z})_{Y|E})
 \end{array}$$

The theorem follows by the induction on n from the following lemma. \square

Lemma 3.9. *Let the assumption be as in [Theorem 3.2](#).*

(1) *Assuming $d \geq 3$, the natural map*

$$H^i(X, (\mathbb{Z}/p\mathbb{Z})_{X|D}) \rightarrow H^i(Y, (\mathbb{Z}/p\mathbb{Z})_{Y|E})$$

is an isomorphism for $i = 1$ and injective for $i = 2$.

(2) *Assuming $d = 2$, the natural map*

$$H^1(X, (\mathbb{Z}/p\mathbb{Z})_{X|D}) \rightarrow H^1(Y, (\mathbb{Z}/p\mathbb{Z})_{Y|E})$$

is injective.

Proof. We define an object \mathcal{K} of $D^b(X)$:

$$\mathcal{K} = \text{Cone}(\mathbb{O}_X(D/p - Y) \xrightarrow{1-F} \mathbb{O}_X(D - Y))[-1].$$

By the commutative diagram with exact horizontal sequences

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \mathbb{O}_X(D/p - Y) & \longrightarrow & \mathbb{O}_X(D/p) & \longrightarrow & \mathbb{O}_Y(E/p) & \longrightarrow & 0 \\ & & \downarrow 1-F & & \downarrow 1-F & & \downarrow 1-F & & \\ 0 & \longrightarrow & \mathbb{O}_X(D - Y) & \longrightarrow & \mathbb{O}_X(D) & \longrightarrow & \mathbb{O}_Y(E) & \longrightarrow & 0 \end{array}$$

we have a distinguished triangle in $D^b(X)$:

$$\mathcal{K} \rightarrow (\mathbb{Z}/p\mathbb{Z})_{X|D} \rightarrow (\mathbb{Z}/p\mathbb{Z})_{Y|E} \xrightarrow{\pm}.$$

Hence, it suffices to show $H^i(X, \mathcal{K}) = 0$ for $i = 1, 2$ in case $d \geq 3$ and $H^1(X, \mathcal{K}) = 0$ in case $d = 2$. We have an exact sequence

$$\begin{aligned} H^0(\mathbb{O}_X(D - Y)) &\rightarrow H^1(X, \mathcal{K}) \rightarrow H^1(\mathbb{O}_X(D/p - Y)) \\ &\rightarrow H^1(\mathbb{O}_X(D - Y)) \rightarrow H^2(X, \mathcal{K}) \rightarrow H^2(\mathbb{O}_X(D/p - Y)). \end{aligned}$$

By Serre duality, for a divisor Ξ on X , we have

$$H^i(X, \mathbb{O}_X(\Xi - Y)) = H^{d-i}(X, \Omega_X^d(-\Xi + Y))^\vee.$$

Thus, the desired assertion follows from [Definition 3.1\(A1\)](#) and [\(B\)](#). \square

It remains to deduce the assertions for fil_D of [Theorem 3.4\(1\)](#) and [\(2\)](#) from those for fil_D^{\log} . Let D' be as in the beginning of this section and $E' = D' \times_X Y$. Noting that the multiplicities of D' are prime to p , we have by [Lemma 2.8\(3\)](#)

$$\text{fil}_{D'} H^1(U) = \text{fil}_{D'-C}^{\log} H^1(U) \quad \text{and} \quad \text{fil}_{E'} H^1(U \cap Y) = \text{fil}_{E'-C \cap Y}^{\log} H^1(U \cap Y).$$

Thus, the assertions for $\text{fil}_{D'-C}^{\log}$ of [Theorem 3.4](#) imply that for $\text{fil}_{D'}$. Since $\text{fil}_D \subset \text{fil}_{D'}$, it immediately implies the injectivity of

$$\text{fil}_D H^1(U) \rightarrow \text{fil}_E H^1(U \cap Y).$$

It remains to deduce its surjectivity from that of

$$\text{fil}_{D'} H^1(U) \rightarrow \text{fil}_{E'} H^1(U \cap Y)$$

assuming $d \geq 3$. For this it, suffices to show the injectivity of

$$\text{fil}_{D'} H^1(U) / \text{fil}_D H^1(U) \rightarrow \text{fil}_{E'} H^1(U \cap Y) / \text{fil}_E H^1(U \cap Y).$$

By [Proposition 2.6](#), we have a commutative diagram

$$\begin{array}{ccc} \text{fil}_{D'} H^1(U) / \text{fil}_D H^1(U) & \hookrightarrow & \bigoplus_{\lambda \in I'} H^0(C_\lambda, \Omega_X^1(D')) \otimes_{\mathbb{O}_X} \mathbb{O}_{C_\lambda} \\ \downarrow & & \downarrow \\ \text{fil}_{E'} H^1(U \cap Y) / \text{fil}_E H^1(U \cap Y) & \hookrightarrow & \bigoplus_{\lambda \in I'} H^0(C_\lambda \cap Y, \Omega_Y^1(D')) \otimes_{\mathbb{O}_Y} \mathbb{O}_{C_\lambda \cap Y} \end{array}$$

Thus, we are reduced to showing the injectivity of the right vertical map. Putting $\mathcal{L} = \text{Ker}(\Omega_X^1 \rightarrow i_* \Omega_Y^1)$ where $i : Y \subset X$, the assertion follows from

$$H^0(C_\lambda, \mathcal{L}(D')) \otimes_{\mathbb{O}_X} \mathbb{O}_{C_\lambda} = 0.$$

Note that we used the fact that Y and C_λ intersect transversally. We have an exact sequence

$$0 \rightarrow \Omega_X^1(-Y) \rightarrow \mathcal{L} \rightarrow \mathbb{O}_X(-Y) \otimes \mathbb{O}_Y \rightarrow 0.$$

From this, we get an exact sequence

$$0 \rightarrow \Omega_X^1(D' - Y) \otimes_{\mathbb{O}_X} \mathbb{O}_{C_\lambda} \rightarrow \mathcal{L}(D') \otimes_{\mathbb{O}_X} \mathbb{O}_{C_\lambda} \rightarrow \mathbb{O}_{C_\lambda}(D' - Y) \otimes \mathbb{O}_{C_\lambda \cap Y} \rightarrow 0.$$

We also have an exact sequence

$$0 \rightarrow \mathbb{O}_{C_\lambda}(D' - 2Y) \rightarrow \mathbb{O}_{C_\lambda}(D' - Y) \rightarrow \mathbb{O}_{C_\lambda}(D' - Y) \otimes \mathbb{O}_{C_\lambda \cap Y} \rightarrow 0.$$

Therefore, the desired assertion follows from [Definition 3.1\(A2\)](#). This completes the proof of [Theorem 3.4](#). □

References

[Esnault and Kerz 2012] H. Esnault and M. Kerz, “A finiteness theorem for Galois representations of function fields over finite fields (after Deligne)”, *Acta Math. Vietnam.* **37**:4 (2012), 531–562. [MR 3058662](#) [Zbl 06133191](#)

[Grothendieck 1967] A. Grothendieck, *Local cohomology*, Lecture Notes in Mathematics **41**, Springer, Berlin, 1967. [MR 37 #219](#) [Zbl 0185.49202](#)

- [de Jong 1996] A. J. de Jong, “Smoothness, semi-stability and alterations”, *Inst. Hautes Études Sci. Publ. Math.* **83** (1996), 51–93. [MR 98e:14011](#) [Zbl 0916.14005](#)
- [Kato 1989] K. Kato, “Swan conductors for characters of degree one in the imperfect residue field case”, pp. 101–131 in *Algebraic K-theory and algebraic number theory* (Honolulu, 1987), edited by M. R. Stein and R. K. Dennis, *Contemp. Math.* **83**, Amer. Math. Soc., Providence, RI, 1989. [MR 90g:11164](#) [Zbl 0716.12006](#)
- [Kerz and Saito 2013] M. Kerz and S. Saito, “Chow group of 0-cycles with modulus and higher dimensional class field theory”, preprint, 2013. [arXiv 1304.4400](#)
- [Laumon 1981] G. Laumon, “Semi-continuité du conducteur de Swan (d’après P. Deligne)”, pp. 173–219 in *The Euler–Poincaré characteristic*, edited by J.-L. Verdier, *Astérisque* **83**, Soc. Math. France, Paris, 1981. [MR 83g:14007](#) [Zbl 0504.14013](#)
- [Matsuda 1997] S. Matsuda, “On the Swan conductor in positive characteristic”, *Amer. J. Math.* **119**:4 (1997), 705–739. [MR 99h:14022](#) [Zbl 0928.14017](#)
- [Raskind 1995] W. Raskind, “Abelian class field theory of arithmetic schemes”, pp. 85–187 in *K-theory and algebraic geometry: connections with quadratic forms and division algebras* (Santa Barbara, 1992), part 1, edited by B. Jacob and A. Rosenberg, *Proc. Sympos. Pure Math.* **58**, Amer. Math. Soc., Providence, RI, 1995. [MR 96b:11089](#) [Zbl 0832.19004](#)
- [Schmidt and Spieß 2000] A. Schmidt and M. Spieß, “Singular homology and class field theory of varieties over finite fields”, *J. Reine Angew. Math.* **527** (2000), 13–36. [MR 2001m:11108](#) [Zbl 0961.14013](#)
- [SGA 1 1971] A. Grothendieck, *Séminaire de Géométrie Algébrique du Bois Marie 1960–1961: revêtements étales et groupe fondamental*, *Lecture Notes in Mathematics* **224**, Springer, Berlin, 1971. [MR 50 #7129](#) [Zbl 0234.14002](#)
- [SGA 2 1968] A. Grothendieck, *Séminaire de Géométrie Algébrique du Bois-Marie 1962: cohomologie locale des faisceaux cohérents et théorèmes de Lefschetz locaux et globaux*, *Advanced Studies in Pure Mathematics* **2**, North-Holland, Amsterdam, 1968. [MR 57 #16294](#) [Zbl 0197.47202](#)

Communicated by Jean-Louis Colliot-Thélène

Received 2013-04-27

Revised 2013-11-12

Accepted 2013-12-10

moritz.kerz@mathematik.uni-regensburg.de

*Naturwissenschaftliche Fakultät I – Mathematik,
Universität Regensburg, 93040 Regensburg, Germany*

sshuji@msb.biglobe.ne.jp

*Interactive Research Center of Science, Graduate School
of Science and Engineering, Tokyo Institute of Technology,
Ookayama, Meguro, Tokyo 152-8551, Japan*

Algebra & Number Theory

msp.org/ant

EDITORS

MANAGING EDITOR

Bjorn Poonen
Massachusetts Institute of Technology
Cambridge, USA

EDITORIAL BOARD CHAIR

David Eisenbud
University of California
Berkeley, USA

BOARD OF EDITORS

Georgia Benkart	University of Wisconsin, Madison, USA	Susan Montgomery	University of Southern California, USA
Dave Benson	University of Aberdeen, Scotland	Shigefumi Mori	RIMS, Kyoto University, Japan
Richard E. Borcherds	University of California, Berkeley, USA	Raman Parimala	Emory University, USA
John H. Coates	University of Cambridge, UK	Jonathan Pila	University of Oxford, UK
J-L. Colliot-Thélène	CNRS, Université Paris-Sud, France	Victor Reiner	University of Minnesota, USA
Brian D. Conrad	University of Michigan, USA	Karl Rubin	University of California, Irvine, USA
Hélène Esnault	Freie Universität Berlin, Germany	Peter Sarnak	Princeton University, USA
Hubert Flenner	Ruhr-Universität, Germany	Joseph H. Silverman	Brown University, USA
Edward Frenkel	University of California, Berkeley, USA	Michael Singer	North Carolina State University, USA
Andrew Granville	Université de Montréal, Canada	Vasudevan Srinivas	Tata Inst. of Fund. Research, India
Joseph Gubeladze	San Francisco State University, USA	J. Toby Stafford	University of Michigan, USA
Roger Heath-Brown	Oxford University, UK	Bernd Sturmfels	University of California, Berkeley, USA
Ehud Hrushovski	Hebrew University, Israel	Richard Taylor	Harvard University, USA
Craig Huneke	University of Virginia, USA	Ravi Vakil	Stanford University, USA
Mikhail Kapranov	Yale University, USA	Michel van den Bergh	Hasselt University, Belgium
Yujiro Kawamata	University of Tokyo, Japan	Marie-France Vignéras	Université Paris VII, France
János Kollár	Princeton University, USA	Kei-Ichi Watanabe	Nihon University, Japan
Yuri Manin	Northwestern University, USA	Efim Zelmanov	University of California, San Diego, USA
Barry Mazur	Harvard University, USA	Shou-Wu Zhang	Princeton University, USA
Philippe Michel	École Polytechnique Fédérale de Lausanne		

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

See inside back cover or msp.org/ant for submission instructions.

The subscription price for 2014 is US \$225/year for the electronic version, and \$400/year (+\$55, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscribers address should be sent to MSP.

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

ANT peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2014 Mathematical Sciences Publishers

Algebra & Number Theory

Volume 8 No. 3 2014

Derived invariants of irregular varieties and Hochschild homology	513
LUIGI LOMBARDI	
Sato–Tate distributions of twists of $y^2 = x^5 - x$ and $y^2 = x^6 + 1$	543
FRANCESC FITÉ and ANDREW V. SUTHERLAND	
The algebraic dynamics of generic endomorphisms of \mathbb{P}^n	587
NAJMUDDIN FAKHRUDDIN	
The tame-wild principle for discriminant relations for number fields	609
JOHN W. JONES and DAVID P. ROBERTS	
Linear forms in logarithms and integral points on higher-dimensional varieties	647
AARON LEVIN	
Lefschetz theorem for abelian fundamental group with modulus	689
MORITZ KERZ and SHUJI SAITO	
Localization of spherical varieties	703
FRIEDRICH KNOP	
Lefschetz operator and local Langlands modulo ℓ : the limit case	729
JEAN-FRANÇOIS DAT	
Splitting tower and degree of tt-rings	767
PAUL BALMER	