

# *Algebra & Number Theory*

Volume 8

2014

No. 7

**Monodromy and local-global compatibility for  $l = p$**

Ana Caraiani





# Monodromy and local-global compatibility for $l = p$

Ana Caraiani

We strengthen the compatibility between local and global Langlands correspondences for  $\mathrm{GL}_n$  when  $n$  is even and  $l = p$ . Let  $L$  be a CM field and  $\Pi$  a cuspidal automorphic representation of  $\mathrm{GL}_n(\mathbb{A}_L)$  which is conjugate self-dual and regular algebraic. In this case, there is an  $l$ -adic Galois representation associated to  $\Pi$ , which is known to be compatible with local Langlands in almost all cases when  $l = p$  by recent work of Barnet-Lamb, Gee, Geraghty and Taylor. The compatibility was proved only up to semisimplification unless  $\Pi$  has Shin-regular weight. We extend the compatibility to Frobenius semisimplification in all cases by identifying the monodromy operator on the global side. To achieve this, we derive a generalization of Mokrane’s weight spectral sequence for log crystalline cohomology.

1. Introduction	1597
2. Shimura varieties	1600
3. Log crystalline cohomology	1609
4. Generalizing the Mokrane spectral sequence	1635
5. Proof of the main theorem	1643
Acknowledgements	1645
References	1645

## 1. Introduction

This paper is a continuation of [Caraiani 2012]. Here we extend our local-global compatibility result to the case  $l = p$ .

**Theorem 1.1.** *Let  $n \in \mathbb{Z}_{\geq 2}$  be an integer and  $L$  be a CM field with complex conjugation  $c$ . Let  $l$  be a prime of  $\mathbb{Q}$  and  $\iota_l : \overline{\mathbb{Q}}_l \rightarrow \mathbb{C}$  be an isomorphism. Let  $\Pi$  be a cuspidal automorphic representation of  $\mathrm{GL}_n(\mathbb{A}_L)$  satisfying*

- $\Pi^\vee \simeq \Pi \circ c$ ,

Partially supported by NSF Postdoctoral Fellowship DMS-1204465.

MSC2010: primary 11F80; secondary 11G18, 11R39.

Keywords: Galois representations, automorphic forms, local-global compatibility, monodromy operator, crystalline cohomology.

- $\Pi$  is cohomological for some irreducible algebraic representation  $\Xi$  of  $\mathrm{GL}_n(L \otimes_{\mathbb{Q}} \mathbb{C})$ .

Let

$$R_l(\Pi) : \mathrm{Gal}(\bar{L}/L) \rightarrow \mathrm{GL}_n(\bar{\mathbb{Q}}_l)$$

be the Galois representation associated to  $\Pi$  by [Shin 2011; Chenevier and Harris 2013]. Let  $y$  be a place of  $L$  above  $l$ . Then we have the isomorphism of Weil–Deligne representations

$$WD(R_l(\Pi)|_{\mathrm{Gal}(\bar{L}_y/L_y)})^{\mathrm{F-ss}} \simeq \iota_l^{-1} \mathcal{L}_{n,L_y}(\Pi_y).$$

Here  $\mathcal{L}_{n,L_y}(\Pi_y)$  is the image of  $\Pi_y$  under the local Langlands correspondence, using the geometric normalization; i.e.,  $\mathcal{L}_{n,L_y}(\Pi_y) := \mathrm{rec}(\Pi_y^\vee \otimes |\det|^{\frac{1-n}{2}})$ , where  $\mathrm{rec}$  is the local Langlands correspondence compatible with  $L$ - and  $\epsilon$ -factors (see the introduction to [Harris and Taylor 2001] for more details).  $WD(r)$  is the Weil–Deligne representation attached to a de Rham  $l$ -adic representation  $r$  of the absolute Galois group of an  $l$ -adic field; F-ss denotes Frobenius semisimplification. Note that we are assuming throughout that  $n \geq 2$ . The local-global compatibility of Langlands correspondences for  $\mathrm{GL}_1$  follows from the compatibility between local and global class field theory.

This theorem is proved in [Barnet-Lamb et al. 2012; Barnet-Lamb et al. 2011] in the case when  $\Pi$  has Shin-regular weight (either  $n$  is odd or if  $n$  is even then  $\Pi$  satisfies an additional regularity condition) and in general up to semisimplification. The strategy for obtaining the local-global compatibility of monodromy operators in these cases is to make use of the fact that the  $l$ -adic Galois representation associated to  $\Pi$  occurs in the cohomology of certain very special unitary Shimura varieties. These are associated to unitary similitude groups with signature  $(1, n-1)$  (respectively,  $(1, n)$  if  $n$  is even) at exactly one infinite place and signature  $(0, n)$  (respectively,  $(0, n+1)$ ) at all the other infinite places. The problem can be reduced to the case when  $\Pi_y$  has an Iwahori-fixed vector, in which case one has to compute the crystalline cohomology of a compact Shimura variety which is strictly semistable. This computation makes use of the weight spectral sequence for crystalline cohomology due to Mokrane [1993], which is shown to degenerate at the first page. We remark that the  $l$ -adic Galois representation associated to  $\Pi$  is only known to occur in the cohomology of a proper, smooth variety in the case when  $\Pi$  has Shin-regular weight.

Our goal in this paper is to match up the monodromy operators in the case when  $n$  is even and  $\Pi$  does not necessarily have Shin-regular weight. Following the conventions of [Taylor and Yoshida 2007], we call a Weil–Deligne representation *pure of weight  $k$*  if it admits a weight filtration, with all the weights in  $k + \mathbb{Z}$ , such that the (iterated) monodromy operator induces an isomorphism of the  $(k+i)$ -th

and  $(k - i)$ -th graded pieces for all positive integers  $i$ . By Lemma 1.4 (4) of [Taylor and Yoshida 2007], given a semisimple representation of the Weil group of some  $l$ -adic field, there is at most one way to choose the monodromy operator such that the resulting Weil–Deligne representation is pure of some weight. By Theorem 1.2 of [Caraiani 2012],  $\Pi_y$  is tempered, so we know that  $\iota_l^{-1} \mathcal{L}_{n, L_y}(\Pi_y)$  is pure of some weight.

By Theorem A of [Barnet-Lamb et al. 2011], we also know that we have an isomorphism up to semisimplification:

$$WD(R_l(\Pi)|_{\text{Gal}(\bar{L}_y/L_y)})^{\text{ss}} \simeq \iota_l^{-1} \mathcal{L}_{n, L_y}(\Pi_y)^{\text{ss}}.$$

We note that Theorem A of [Barnet-Lamb et al. 2011] is stated for an imaginary CM field  $F$ . For our CM field  $L$  we proceed as on pages 230–231 of [Harris and Taylor 2001] to find a quadratic extension  $F/L$  which is an imaginary CM field, in which  $y = y'y''$  splits, such that

$$[R_l(\Pi)|_{\text{Gal}(\bar{L}/F)}] = [R_l(BC_{F/L}(\Pi))].$$

This together with Theorem A of [Barnet-Lamb et al. 2011] gives the compatibility up to semisimplification for the place  $y$  of  $L$ . Therefore, in order to complete the proof of Theorem 1.1, it suffices to show that  $W := WD(R_l(\Pi)|_{\text{Gal}(\bar{L}_y/L_y)})^{\text{F-ss}}$  is pure of some weight when  $n$  is even. From now on we will let  $n \in \mathbb{Z}_{\geq 2}$  be an even integer.

Our argument will follow the same general lines as that of [Taylor and Yoshida 2007], which is also the strategy followed by [Barnet-Lamb et al. 2012; Barnet-Lamb et al. 2011]. We reduce the problem to the case when  $\Pi_y$  has an Iwahori-fixed vector. In this case, we find not  $W$  itself, but rather the tensor square of  $W$  in the log crystalline cohomology of a compact Shimura variety with Iwahori-level structure, and finally compute a part of this cohomology explicitly. For the last step, however, we can not make use of the Mokrane spectral sequence, since our Iwahori-level Shimura variety is no longer semistable, but rather Zariski-locally étale over a product of strictly semistable schemes. Therefore, we need to derive a formula for the log crystalline cohomology of the special fiber of this Shimura variety in terms of the crystalline cohomology of closed Newton polygon strata in the special fiber. Deriving this formula constitutes the heart of this paper; we obtain it in the form of a generalization of the Mokrane spectral sequence or as a crystalline analogue of Corollary 4.28 of [Caraiani 2012].

We briefly outline the structure of our paper. In Section 2 we reduce to the case where  $\Pi$  has an Iwahori-fixed vector, we define an inverse system of compact Shimura varieties associated to a unitary group and we show that the crystalline cohomology of the Iwahori-level Shimura variety realizes the tensor square of  $W$ . The Shimura varieties we work with are the same as those studied in [Caraiani 2012],

so in Section 2 we also recall the main results from [Caraiani 2012] concerning them. In Section 3 we recall and adapt to our situation some standard results from the theory of log crystalline cohomology and the de Rham–Witt complex; we define and study some slight generalizations of the logarithmic de Rham–Witt complex. In Section 4 we generalize the Mokrane spectral sequence to our geometric setting. The main technical result is Theorem 4.6. In Section 5 we prove Theorem 1.1.

## 2. Shimura varieties

Let  $L, \Pi, R_l(\Pi)$  and  $y$  be as described in the introduction. Below, we show that we can understand the Weil–Deligne representation  $W = WD(R_l(\Pi)_{\text{Gal}(\bar{L}_y/L_y)})^{\text{F-ss}}$  by computing a part of the crystalline cohomology of an inverse system of Shimura varieties. In the first part we closely follow Sections 2 and 7 of [Caraiani 2012] and afterwards we use some results from Section 5 of the same work.

We claim first that we can reduce the problem to the case when  $\Pi$  has Iwahori-fixed vectors at  $y$ , and we can also put ourselves in a situation where the base change from unitary groups to  $\text{GL}_n$  is well understood. This means that we can reduce the problem to understanding the cohomology of certain Iwahori-level unitary Shimura varieties. More precisely, we can find a CM field extension  $F'$  of  $L$  such that:

- $F' = EF_1$ , where  $E$  is an imaginary quadratic field in which  $l$  splits and  $F_1 = (F')^{c=1}$  has  $[F_1 : \mathbb{Q}] \geq 2$ ;
- $F'$  is soluble and Galois over  $L$ ;
- $\Pi_{F'}^0 := BC_{F'/L}(\Pi)$  is a cuspidal automorphic representation of  $\text{GL}_n(\mathbb{A}_{F'})$ ; and
- there is a place  $\mathfrak{p}$  above the place  $y$  of  $L$  such that  $\Pi_{F',\mathfrak{p}}^0$  has a nonzero Iwahori-fixed vector;

and a CM field  $F$  which is a quadratic extension of  $F'$ , such that:

- $\mathfrak{p} = \mathfrak{p}_1\mathfrak{p}_2$  splits in  $F$ ;
- $\text{Ram}_{F/\mathbb{Q}} \cup \text{Ram}_{\mathbb{Q}}(\Pi) \subset \text{Spl}_{F/F_2,\mathbb{Q}}$ , where  $F_2 := (F)^{c=1}$ ; and
- $\Pi_F^0 = BC_{F/F'}(\Pi_{F'}^0)$  is a cuspidal automorphic representation of  $\text{GL}_n(\mathbb{A}_F)$ .

We can find  $F$  and  $F'$  as in the proof of Corollary 5.9 of [Caraiani 2012]. Since purity is preserved under finite extensions by Lemma 1.4 of [Taylor and Yoshida 2007], to show that  $W$  is pure it suffices to show that

$$W_{F'} := WD(R_l(\Pi_{F'}^0)|_{\text{Gal}(\bar{F}'_{\mathfrak{p}}/F'_{\mathfrak{p}})})^{\text{F-ss}}$$

is pure. Note that in this new situation  $\Pi_{F',\mathfrak{p}}^0$  has a nonzero Iwahori-fixed vector.

We can define an algebraic group  $G$  over  $\mathbb{Q}$  and an inverse system of Shimura varieties over  $F'$  corresponding to a PEL Shimura datum  $(F, *, V, \langle \cdot, \cdot \rangle, h)$ . Here

$F$  is the CM field defined above and  $* = c$  is the involution corresponding to complex conjugation. We take  $V$  to be the  $F$ -vector space  $F^n$ . The pairing

$$\langle \cdot, \cdot \rangle : V \times V \rightarrow \mathbb{Q}$$

is a nondegenerate Hermitian pairing such that  $\langle f v_1, v_2 \rangle = \langle v_1, f^* v_2 \rangle$  for all  $f \in F$  and  $v_1, v_2 \in V$ . The last element we need is an  $\mathbb{R}$ -algebra homomorphism  $h : \mathbb{C} \rightarrow \text{End}_F(V) \otimes_{\mathbb{Q}} \mathbb{R}$  such that the bilinear pairing

$$(v_1, v_2) \rightarrow \langle v_1, h(i)v_2 \rangle$$

is symmetric and positive definite. We define the algebraic group  $G$  over  $\mathbb{Q}$  by

$$G(R) = \{ (g, \lambda) \in \text{End}_{F \otimes_{\mathbb{Q}} \mathbb{R}}(V \otimes_{\mathbb{Q}} R)^\times \times R^\times \mid \langle g v_1, g v_2 \rangle = \lambda \langle v_1, v_2 \rangle \}$$

for any  $\mathbb{Q}$ -algebra  $R$ .

We choose embeddings  $\tau_i : F \hookrightarrow \mathbb{C}$  such that  $\tau_2 = \tau_1 \circ \sigma$ , where  $\sigma$  is an element of  $\text{Gal}(F/F')$  which takes  $\mathfrak{p}_1$  to  $\mathfrak{p}_2$ . For  $\sigma \in \text{Hom}_{E, \tau_E}(F, \mathbb{C})$  we let  $(p_\sigma, q_\sigma)$  be the signature at  $\sigma$  of the pairing  $\langle \cdot, \cdot \rangle$  on  $V \otimes_{\mathbb{Q}} \mathbb{R}$ . In particular,  $\tau_E := \tau_1|_E = \tau_2|_E$  is well-defined. We claim that it is possible to choose a PEL datum as above such that  $(p_\tau, q_\tau) = (1, n - 1)$  for  $\tau = \tau_1$  or  $\tau_2$  and  $(p_\tau, q_\tau) = (0, n)$  otherwise and such that  $G_{\mathbb{Q}_v}$  is quasisplit at every finite place  $v$  of  $\mathbb{Q}$ . This follows from Lemma 2.1 of [Caraiani 2012] and the discussion following it, and it depends crucially on the fact that  $n$  is even. We choose such a PEL datum and we let  $G$  be the corresponding algebraic group over  $\mathbb{Q}$  with the prescribed signature at infinity and quasisplit at all the finite places.

Let  $\Xi_F^0 := BC_{F/L}(\Xi)$  and  $F_2 = F^{c=1}$ . Lemma 7.2 of [Shin 2011] says that we can find a character  $\psi : \mathbb{A}_E^\times / E^\times \rightarrow \mathbb{C}^\times$  and an algebraic representation  $\xi_{\mathbb{C}}$  of  $G$  over  $\mathbb{C}$  satisfying the following conditions:

- $\psi_{\Pi_F^0} = \psi^c / \psi$ .
- $\Xi_F^0$  is isomorphic to the restriction of  $\Xi'$  to  $\text{Res}_{F/\mathbb{Q}}(\text{GL}_n) \times_{\mathbb{Q}} \mathbb{C}$ , where  $\Xi'$  is obtained from  $\xi_{\mathbb{C}}$  by base change from  $G$  to  $\mathbb{G}_n := \text{Res}_{E/\mathbb{Q}}(G \times_{\mathbb{Q}} E)$ .
- $\xi_{\mathbb{C}}|_{E_\infty^\times}^{-1} = \psi_\infty^c$ .
- $\text{Ram}_{\mathbb{Q}}(\psi) \subset \text{Spl}_{F/F_2, \mathbb{Q}}$ .
- $\psi|_{\mathcal{O}_{E_u}^\times} = 1$ , where  $u$  is the place above  $l$  induced by  $\iota_l^{-1} \tau_E$ .

Define  $\xi := \iota_l \xi_{\mathbb{C}}$ , and define  $\Pi^1 := \psi \otimes \Pi_F^0$ , which is a cuspidal automorphic representation of  $\text{GL}_1(\mathbb{A}_E) \times \text{GL}_n(\mathbb{A}_F)$ .

Corresponding to the PEL datum  $(F, *, V, \langle \cdot, \cdot \rangle, h)$ , we have a PEL-type moduli problem of abelian varieties. This moduli problem is defined in Section 2.1 of [Caraiani 2012], and here we recall some facts about it. Since the reflex field of the PEL datum is  $F'$ , the moduli problem for an open compact subgroup  $U \subset G(\mathbb{A}^\infty)$  is

representable by a Shimura variety  $X_U/F'$ , which is a smooth and quasiprojective scheme of dimension  $2n - 2$ . The inverse system of Shimura varieties  $X_U$  as  $U$  varies has an action of  $G(\mathbb{A}^\infty)$ . As in Section III.2 of [Harris and Taylor 2001], starting with  $\xi$ , which is an irreducible algebraic representation of  $G$  over  $\overline{\mathbb{Q}}_l$ , we can define a lisse  $\overline{\mathbb{Q}}_l$ -sheaf  $\mathcal{L}_\xi$  over each  $X_U$ , and the action of  $G(\mathbb{A}^\infty)$  extends to the inverse system of sheaves. The direct limit

$$H^i(X, \mathcal{L}_\xi) := \varinjlim H^i(X_U \times_{F'} \overline{F'}, \mathcal{L}_\xi)$$

is a semisimple admissible representation of  $G(\mathbb{A}^\infty)$  with a continuous action of  $\text{Gal}(\overline{F'}/F')$ . It can be decomposed as

$$H^i(X, \mathcal{L}_\xi) = \bigoplus_{\pi} \pi \otimes R_{\xi,l}^i(\pi),$$

where the sum runs over irreducible admissible representations  $\pi$  of  $G(\mathbb{A}^\infty)$  over  $\overline{\mathbb{Q}}_l$ . The  $R_{\xi,l}^i(\pi)$  are finite-dimensional continuous representations of  $\text{Gal}(\overline{F'}/F')$  over  $\overline{\mathbb{Q}}_l$ . Let  $\mathcal{A}_U$  be the universal abelian variety over  $X_U$ , to the inverse system of which the action of  $G(\mathbb{A}^\infty)$  extends. To the irreducible representation  $\xi$  of  $G$  we can associate as in Section III.2 of [Harris and Taylor 2001] nonnegative integers  $m_\xi$  and  $t_\xi$  as well as an idempotent  $a_\xi$  of  $H^*(\mathcal{A}_U^{m_\xi} \times_{F'} \overline{F'}, \overline{\mathbb{Q}}_l(t_\xi))$ . (Here  $\mathcal{A}_U^{m_\xi}$  denotes the  $m_\xi$ -fold product of  $\mathcal{A}_U$  with itself over  $X_U$  and  $\overline{\mathbb{Q}}_l(t_\xi)$  is a Tate twist.) We have an isomorphism

$$H^i(X_U \times_{F'} \overline{F'}, \mathcal{L}_\xi) \simeq a_\xi H^{i+m_\xi}(\mathcal{A}_U^{m_\xi} \times_{F'} \overline{F'}, \overline{\mathbb{Q}}_l(t_\xi)),$$

which commutes with the  $G(\mathbb{A}^\infty)$ -action.

For every finite place  $v$  of  $\mathbb{Q}$  we can define a base-change morphism taking certain admissible  $G(\mathbb{Q}_v)$ -representations to admissible  $\mathbb{G}(\mathbb{Q}_v)$ -representations, as in Section 4.2 of [Shin 2011]. Recall that  $\text{Ram}_{F/\mathbb{Q}} \cup \text{Ram}_{\mathbb{Q}} \Pi^1 \subset \text{Spl}_{F/F_2, \mathbb{Q}}$ . If  $v \notin \text{Spl}_{F/F_2, \mathbb{Q}}$  then we can define the morphism

$$BC : \text{Irr}_{(l)}^{\text{ur}}(G(\mathbb{Q}_v)) \rightarrow \text{Irr}_{(l)}^{\text{ur}, \theta\text{-st}}(\mathbb{G}(\mathbb{Q}_v)),$$

taking unramified representations of  $G(\mathbb{Q}_v)$  to unramified,  $\theta$ -stable representations of  $\mathbb{G}(\mathbb{Q}_v)$ . If  $v \in \text{Spl}_{F/F_2, \mathbb{Q}}$  then the morphism

$$BC : \text{Irr}_{(l)}(G(\mathbb{Q}_v)) \rightarrow \text{Irr}_{(l)}^{\theta\text{-st}}(\mathbb{G}(\mathbb{Q}_v))$$

can be defined explicitly since  $G(\mathbb{Q}_v)$  is split. Putting these maps together, we get, for any finite set of primes  $\mathfrak{S}_{\text{fin}}$  such that

$$\text{Ram}_{F/\mathbb{Q}} \cup \text{Ram}_{\mathbb{Q}}(\Pi) \subset \mathfrak{S}_{\text{fin}} \subset \text{Spl}_{F/F_2, \mathbb{Q}},$$

a base-change morphism

$$BC : \text{Irr}_{(l)}^{\text{ur}}(G(\mathbb{A}^{\mathfrak{S}_{\text{fin}} \cup \{\infty\}})) \otimes \text{Irr}_{(l)}(G(\mathbb{A}_{\mathfrak{S}_{\text{fin}}})) \\ \rightarrow \text{Irr}_{(l)}^{\text{ur}, \theta\text{-st}}(\mathbb{G}(\mathbb{A}^{\mathfrak{S}_{\text{fin}} \cup \{\infty\}})) \otimes \text{Irr}_{(l)}^{\theta\text{-st}}(\mathbb{G}(\mathbb{A}_{\mathfrak{S}_{\text{fin}}})) .$$

Let  $p$  be a prime of  $\mathbb{Q}$  which splits in  $E$  and such that there is a place of  $F'$  above  $p$  which splits in  $F$ . Let  $\mathfrak{S}_{\text{fin}}$  be a finite set of primes such that

$$\text{Ram}_{F/\mathbb{Q}} \cup \text{Ram}_{\mathbb{Q}}(\Pi) \cup \{p\} \subset \mathfrak{S}_{\text{fin}} \subset \text{Spl}_{F/F_2, \mathbb{Q}},$$

and set  $\mathfrak{S} := \mathfrak{S}_{\text{fin}} \cup \{\infty\}$ . For any  $R \in \text{Groth}(G(\mathbb{A}^{\mathfrak{S}}) \times G(\mathbb{A}_{\mathfrak{S}_{\text{fin}}}) \times \text{Gal}(\overline{F'}/F'))$  (over  $\overline{\mathbb{Q}}_l$ ) and  $\pi^{\mathfrak{S}} \in \text{Irr}_{(l)}^{\text{ur}}(G(\mathbb{A}^{\mathfrak{S}}))$  define the  $\pi^{\mathfrak{S}}$ -isotypic part of  $R$  to be

$$R\{\pi^{\mathfrak{S}}\} := \sum_{\rho} n(\pi^{\mathfrak{S}} \otimes \rho)[\pi^{\mathfrak{S}}][\rho],$$

where  $\rho$  runs over  $\text{Irr}_l(G(\mathbb{A}_{\mathfrak{S}}) \times \text{Gal}(\overline{F'}/F'))$ . Also define

$$R[\Pi^{1, \mathfrak{S}}] := \sum_{\pi^{\mathfrak{S}}} R[\pi^{\mathfrak{S}}],$$

where each sum runs over  $\pi^{\mathfrak{S}} \in \text{Irr}_{(l)}^{\text{ur}}(G(\mathbb{A}^{\mathfrak{S}}))$  such that  $BC(l_l \pi^{\mathfrak{S}}) \simeq \Pi^{1, \mathfrak{S}}$ .

**Proposition 2.1.** *Let  $\mathfrak{S} = \mathfrak{S}_{\text{fin}} \cup \{\infty\}$  be as above. We have the equality*

$$BC(H^{2n-2}(X, \mathcal{L}_{\xi})[\Pi^{1, \mathfrak{S}}]) \simeq C_G[l_l^{-1} \Pi^{1, \infty}][R_l(\Pi_{F'}^0)^{\otimes 2} \otimes \text{rec}_{l, l_l}(\psi)]$$

of elements of  $\text{Groth}(G(\mathbb{A}^{\infty}) \times \text{Gal}(\overline{F'}/F))$ . Here  $C_G$  is a positive integer and  $\text{rec}_{l, l_l}(\psi)$  is the continuous  $l$ -adic character  $\text{Gal}(\overline{E}/E) \rightarrow \overline{\mathbb{Q}}_l^{\times}$  associated to  $\psi$  by global class field theory, normalized so that it matches uniformizers with geometric Frobenius elements.

**Remark.** Unlike in the classical situation of modular forms or in the case of Harris–Taylor-type Shimura varieties [Harris and Taylor 2001; Shin 2011], the cohomology of our inverse system of Shimura varieties realizes a twist of the tensor square of the  $l$ -adic Galois representation associated to  $\Pi$ , because we have chosen our unitary similitude groups to have signature  $(1, n - 1)$  two infinite places. One could use Matsushima’s formula and  $(\mathfrak{g}, K)$ -cohomology to check that the dimension of the Galois representation seen by this cohomology is  $n^2$ , as predicted by the statement.

*Proof.* Let  $p \in \mathfrak{S}_{\text{fin}}$  be a prime which splits in  $E$  such that there is a place  $w$  of  $F'$  above the place induced by  $\tau_E$  over  $p$  which splits in  $F$ ,  $w = w_1 w_2$ . We start by recalling some constructions and results from Sections 2 and 5 of [Caraiani 2012]. It is possible to define an integral model of each  $X_U$  over the ring of integers  $\mathbb{O}_K$  in  $K := F_{w_1} \simeq F_{w_2}$ , which itself represents a moduli problem of abelian varieties and to which the sheaf  $\mathcal{L}_{\xi}$  extends. The special fiber  $Y_U$  of this integral model has a stratification by open Newton polygon strata  $Y_{U, S, T}^{\circ}$ , according to the formal

(or étale) height of the  $p$ -divisible group of the abelian variety at  $w_1$  and  $w_2$ . Each open Newton polygon stratum is covered by a tower of Igusa varieties  $\text{Ig}_{U^p, \vec{m}}^{(h_1, h_2)}$ , where  $0 \leq h_1, h_2 \leq n - 1$  represent the étale heights of the  $p$ -divisible groups at  $w_1$  and  $w_2$ , and  $\vec{m}$  is a tuple of positive integers describing the level structure at  $p$ .

Define

$$J^{(h_1, h_2)}(\mathbb{Q}_p) := \mathbb{Q}_p^\times \times D_{K, n-h_1}^\times \times \text{GL}_{h_1}(K) \times D_{K, n-h_2}^\times \times \text{GL}_{h_2}(K) \times \prod_w \text{GL}_n(F_w),$$

where  $D_{K, n-h}$  is the division algebra over  $K$  of invariant  $1/(n-h)$  and  $w$  runs over places of  $F$  above  $\tau_E$  other than  $w_1$  and  $w_2$ . The group  $J^{(h_1, h_2)}(\mathbb{Q}_p)$  acts on the directed system of  $H_c^j(\text{Ig}_{U^p, \vec{m}}^{(h_1, h_2)}, \mathcal{L}_\xi)$ , as  $U^p$  and  $\vec{m}$  vary. Let

$$H_c(\text{Ig}^{(h_1, h_2)}, \mathcal{L}_\xi) \in \text{Groth}(G(\mathbb{A}^{\infty, p}) \times J^{(h_1, h_2)})$$

be the alternating sum of the direct limit of  $H_c^j(\text{Ig}_{U^p, \vec{m}}^{(h_1, h_2)}, \mathcal{L}_\xi)$  as in Section 5.1 of [Caraiani 2012]. Let  $\pi_p \in \text{Irr}_l(G(\mathbb{Q}_p))$  be a representation such that  $BC(\pi_p) \simeq \iota_l^{-1} \Pi_p^1$  (such a  $\pi_p$  is unique up to isomorphism since  $p$  splits in  $E$ ). Theorem 5.6 of [Caraiani 2012] gives a formula for computing the cohomology of Igusa varieties, as elements of  $\text{Groth}(\mathbb{G}(\mathbb{A}^\mathfrak{S}) \times \mathbb{G}(\mathbb{A}_{\mathfrak{S}_{\text{fin}} \setminus \{p\}}) \times J^{(h_1, h_2)}(\mathbb{Q}_p))$ :

$$BC^p(H_c(\text{Ig}^{(h_1, h_2)}, \mathcal{L}_\xi)[\Pi^1, \mathfrak{S}]) = e_0(-1)^{h_1+h_2} C_G[\iota_l^{-1} \Pi^1, \mathfrak{S}][\iota_l^{-1} \Pi_{\mathfrak{S}_{\text{fin}} \setminus \{p\}}^1][\text{Red}_n^{(h_1, h_2)}(\pi_p)] \quad (2.1)$$

Here  $e_0 = \pm 1$  independently of  $h_1, h_2$  and  $\text{Red}_n^{(h_1, h_2)}$  is a group morphism from  $\text{Groth}(G(\mathbb{Q}_p))$  to  $\text{Groth}(J^{(h_1, h_2)}(\mathbb{Q}_p))$ , defined explicitly above Theorem 5.6 of [Caraiani 2012].

We can combine the above formula with Mantovan’s formula for the cohomology of Shimura varieties. This is the equality

$$H(X, \mathcal{L}_\xi) = \sum_{0 \leq h_1, h_2 \leq n-1} (-1)^{h_1+h_2} \text{Mant}_{(h_1, h_2)}(H_c(\text{Ig}^{(h_1, h_2)}, \mathcal{L}_\xi)) \quad (2.2)$$

of elements of  $\text{Groth}(G(\mathbb{A}^\infty) \times W_K)$ . Here  $H(X, \mathcal{L}_\xi)$  is the alternating sum of the direct limit of the cohomology of the Shimura fibers (generic fibers) and

$$\text{Mant}_{(h_1, h_2)} : \text{Groth}(J^{(h_1, h_2)}(\mathbb{Q}_p)) \rightarrow \text{Groth}(G(\mathbb{Q}_p) \times W_K)$$

is the functor defined in [Mantovan 2005]. The formula (2.2) is what Theorem 22 of [Mantovan 2005] amounts to in our situation, where  $h_1$  and  $h_2$  are the parameters for the Newton stratification. The extra term  $(-1)^{h_1+h_2}$  occurs on the right-hand side because we use the same convention for the alternating sum of cohomology as in [Caraiani 2012], which differs by a sign from the conventions used in [Mantovan 2005] and [Shin 2011].

By combining formulas (2.1) and (2.2) we get

$$BC^p(H(X, \mathcal{L}_\xi)[\Pi^{1, \mathbb{S}}]) = e_0 C_G[l_l^{-1} \Pi^{1, \infty, p}] \left( \sum_{0 \leq h_1, h_2 \leq n-1} [\text{Mant}_{(h_1, h_2)}(\text{Red}_n^{(h_1, h_2)}(\pi_p))] \right)$$

in  $\text{Groth}(\mathbb{G}(\mathbb{A}^{\infty, p}) \times G(\mathbb{Q}_p) \times W_K)$ . We claim that

$$\sum_{0 \leq h_1, h_2 \leq n-1} [\text{Mant}_{(h_1, h_2)}(\text{Red}_n^{(h_1, h_2)}(\pi_p))] = [\pi_p][(\pi_{p,0} \circ \text{Art}_{\mathbb{Q}_p}^{-1})|_{W_K} \otimes l_l^{-1} \mathcal{L}_{K,n}(\Pi_{F',w}^0)]. \quad (2.3)$$

By its definition above Theorem 5.6 of [Caraiani 2012], the morphism  $\text{Red}_n^{(h_1, h_2)}(\pi_p)$  breaks down as a product

$$(-1)^{h_1+h_2} \pi_{p,0} \otimes \text{Red}^{n-h_1, h_1}(\pi_{w_1}) \otimes \text{Red}^{n-h_2, h_2}(\pi_{w_2}) \otimes \bigotimes_{w \neq w_1, w_2} \pi_w,$$

where  $w$  runs over places above the place of  $p$  induced by  $\tau_E$  other than  $w_1$  and  $w_2$ . The morphism

$$\text{Red}^{n-h, h} : \text{Groth}(\text{GL}_n(K)) \rightarrow \text{Groth}(D_{K,n-h}^\times \times \text{GL}_h(K))$$

is also defined above Theorem 5.6 of [Caraiani 2012]. On the other hand, the functor  $\text{Mant}_{(h_1, h_2)}$  also decomposes as a product (see [Shin 2011, Formula 5.6]), into

$$\text{Mant}_{(h_1, h_2)}(\rho) = \text{Mant}_{1,0}(\rho_0) \otimes \text{Mant}_{n-h_1, h_1}(\rho_{w_1}) \otimes \text{Mant}_{n-h_2, h_2}(\rho_{w_2}) \otimes \bigotimes_{w \neq w_1, w_2} \text{Mant}_{0,m}(\rho_w),$$

where  $w$  again runs over places above the place of  $p$  induced by  $\tau_E$  other than  $w_1$  and  $w_2$ . So

$$\begin{aligned} & \sum_{0 \leq h_1, h_2 \leq n-1} [\text{Mant}_{(h_1, h_2)}(\text{Red}_n^{(h_1, h_2)}(\pi_p))] \\ &= [\text{Mant}_{1,0}(\pi_{p,0})] \otimes \sum_{h_1=0}^{n-1} (-1)^{h_1} [\text{Mant}_{n-h_1, h_1}(\text{Red}^{n-h_1, h_1}(\pi_{w_1}))] \\ & \quad \otimes \sum_{h_2=0}^{n-1} (-1)^{h_2} [\text{Mant}_{n-h_2, h_2}(\text{Red}^{n-h_2, h_2}(\pi_{w_2}))] \otimes \bigotimes_{w \neq w_1, w_2} [\pi_w]. \end{aligned}$$

Now by applying Propositions 2.2(i) and 2.3 of [Shin 2011] we get the desired result (note that the normalization used in their statements is slightly different than

ours, but the relation between the two different normalizations is explained above the statement of Proposition 2.3).

Applying Equation (2.3), we first see that

$$BC(H(X, \mathcal{L}_\xi)[\Pi^1, \mathfrak{S}]) = e_0 C_G[l_l^{-1} \Pi^1, \infty][(\pi_{p,0} \circ \text{Art}_{\mathbb{Q}_p}^{-1})|_{W_K} \otimes l_l^{-1} \mathcal{L}_{K,n}(\Pi_{F',w}^0)] \quad (2.4)$$

in  $\text{Groth}(\mathbb{G}(\mathbb{A}^\infty) \times W_K)$ , which means that

$$BC(H(X, \mathcal{L}_\xi)[\Pi^1, \mathfrak{S}]) = e_0 [l_l^{-1} \Pi^1, \infty][R'(\Pi^1)],$$

for some  $[R'(\Pi^1)] \in \text{Groth}(\text{Gal}(\bar{F}'/F))$ . We show now that

$$[R'(\Pi^1)] = C_G[R(\Pi_{F'}^0)^{\otimes 2} \otimes \text{rec}_{l,l}(\psi)]$$

in  $\text{Groth}(\text{Gal}(\bar{F}'/F'))$ , using the Chebotarev density theorem. Note first that  $R'(\Pi^1)$  is simply the sum of (the alternating sum of)  $R_{\xi,l}^k(\pi^\infty)$ , where  $\pi^\infty$  runs over  $\text{Irr}_l(G(\mathbb{A}^\infty))$  such that

- $BC(l_l \pi^\mathfrak{S}) \simeq \Pi^1, \mathfrak{S}$ ,
- $BC(l_l \pi_{\mathfrak{S}_{\text{fin}}}) \simeq \Pi_{\mathfrak{S}_{\text{fin}}}$ ,
- $R_{\xi,l}^k(\pi^\infty) \neq 0$  for some  $k$ .

The set of such  $\pi$  doesn't depend on  $\mathfrak{S}$  if  $\mathfrak{S}$  is chosen as described above this proposition, so the Galois representation  $R'(\Pi^1)$  is also independent of  $\mathfrak{S}$ . Therefore, for any prime  $w_1$  of  $F$  where  $\Pi^1$  is unramified and which is above a prime  $w$  of  $F'$  which splits in  $F$  and above a prime  $p \neq l$  of  $\mathbb{Q}$  which splits in  $E$ , we can choose a finite set of places  $\mathfrak{S}$  containing  $p$  such that we get from Equation (2.4)

$$[R'(\Pi^1)|_{W_{Fw_1}}] = C_G[(R(\Pi_{F'}^0)^{\otimes 2} \otimes \text{rec}_{l,l}(\psi))_{W_{Fw_1}}].$$

By the Chebotarev density theorem (which tells us the Frobenius elements of primes  $w_1$  are dense in  $\text{Gal}(\bar{F}'/F)$ ) we conclude that

$$[R'(\Pi^1)] = C_G[R(\Pi_{F'}^0)^{\otimes 2} \otimes \text{rec}_{l,l}(\psi)]$$

in  $\text{Groth}(\text{Gal}(\bar{F}'/F'))$ .

It remains to see that  $e_0 = 1$  and that  $H^k(X, \mathcal{L}_\xi)[\Pi^1, \mathfrak{S}] = 0$  unless  $k = 2n - 2$ . In fact, it suffices to show the latter, since then  $H(X, \mathcal{L}_\xi)[\Pi^1, \mathfrak{S}]$  will have to be an actual representation, so that would force  $e_0 = 1$ . The fact that  $H^k(X, \mathcal{L}_\xi)[\Pi^1, \mathfrak{S}] = 0$  for  $k \neq 2n - 2$  can be seen, as in the proof of Corollary 7.3 of [Caraiani 2012], by choosing a prime  $p \neq l$  to work with and applying the spectral sequences in Proposition 7.2 of [ibid.], and noting that the terms of those spectral sequence are 0 outside the diagonal corresponding to  $k = 2n - 2$ . □

**Corollary 2.2.** *By Lemmas 1.4 and 1.7 of [Taylor and Yoshida 2007] and by the same argument as in the proof of Theorem 7.4 of [Caraiani 2012], in order to show that*

$$WD(R_l(\Pi_{F'}^0)|_{\text{Gal}(\overline{F_p}/F_p)})^{\text{F-ss}}$$

*is pure, it suffices to show that*

$$WD(BC(H^{2n-2}(X, \mathcal{L}_\xi)[\Pi^\mathfrak{S}]|_{\text{Gal}(\overline{F_p}/F_p)})^{\text{F-ss}}$$

*is pure, where  $\mathfrak{S}$  is chosen such that it contains  $l$ .*

At this point, we've reduced the question of proving the local-global compatibility of monodromy operators when  $l = p$  to proving that the  $\Pi^\mathfrak{S}$ -part of the cohomology of a system of proper, smooth Shimura varieties over  $F'$  gives rise to a pure Weil–Deligne representation. In the rest of this section, we shall describe integral models of these Shimura varieties which are no longer smooth but are log smooth and of Cartier type. We shall relate their log crystalline cohomology to the Weil–Deligne representation we are interested in. The upshot is that we reduce the question of local-global compatibility to proving the purity of (the  $\Pi^\mathfrak{S}$ -part of) certain log crystalline cohomology groups. This statement is made precise in Corollary 2.3 below.

Recall that  $\mathfrak{p}$  is a place of  $F'$  above  $l$  such that  $\mathfrak{p} = \mathfrak{p}_1\mathfrak{p}_2$ . From now on, set  $K := F_{\mathfrak{p}_1} \simeq F_{\mathfrak{p}_2}$ , where the isomorphism is via  $\sigma$ . Let  $\mathbb{O}_K$  be the ring of integers in  $K$  with uniformizer  $\varpi$  and residue field  $k$ . For  $i = 1, 2$  let  $\text{Iw}_{n, \mathfrak{p}_i}$  be the subgroup of matrices in  $\text{GL}_n(\mathbb{O}_K)$  which reduce modulo  $\mathfrak{p}_i$  to the Borel subgroup  $B_n(k)$ . Now we set

$$U_{\text{Iw}} = U^l \times U_l^{\mathfrak{p}_1, \mathfrak{p}_2}(m) \times \text{Iw}_{n, \mathfrak{p}_1} \times \text{Iw}_{n, \mathfrak{p}_2} \subset G(\mathbb{A}^\infty),$$

for some  $U^l \subset G(\mathbb{A}^\infty)$  compact open and  $U_l^{\mathfrak{p}_1, \mathfrak{p}_2}$  a congruence subgroup at  $l$  away from  $\mathfrak{p}_1$  and  $\mathfrak{p}_2$ . In Section 2.2 of [Caraiani 2012], an integral model for  $X_{U_{\text{Iw}}}/\mathbb{O}_K$  is defined. This is a proper scheme of dimension  $2n - 1$  with smooth generic fiber. The special fiber  $Y_{U_{\text{Iw}}}$  has a stratification by closed Newton polygon strata  $Y_{U_{\text{Iw}}, S, T}$  with  $S, T \subseteq \{1, \dots, n\}$  nonempty subsets. These strata are proper, smooth schemes over  $k$  of dimension  $2n - \#S - \#T$ . In fact,

$$Y_{U_{\text{Iw}}, S, T} = \left( \bigcap_{i \in S} Y_{1, i} \right) \cap \left( \bigcap_{j \in T} Y_{2, j} \right),$$

where each  $Y_{i, j}$  for  $i = 1, 2$  and  $j = 1, \dots, n$  is cut out by one local equation. We can also define

$$Y_{U_{\text{Iw}}}^{(l_1, l_2)} = \bigsqcup_{\substack{S, T \subseteq \{1, \dots, n\} \\ \#S = l_1 \\ \#T = l_2}} Y_{U_{\text{Iw}}, S, T}.$$

By Proposition 2.8 of [Caraiani 2012], the completed local rings of  $X_{U_{\text{Iw}}}$  at closed geometric points  $s$  of  $X_{U_{\text{Iw}}}$  are isomorphic to

$$\mathbb{C}_{X_{U_{\text{Iw}}},s}^{\wedge} \simeq W_{(K)} \llbracket X_1, \dots, X_n, Y_1, \dots, Y_n \rrbracket / (X_{i_1} \cdots X_{i_r} - \varpi, Y_{j_1} \cdots Y_{j_s} - \varpi),$$

where  $\{i_1, \dots, i_r\} \subseteq \{1, \dots, n\}$ ,  $\{j_1, \dots, j_s\} \subseteq \{1, \dots, n\}$  and  $W_{(K)}$  is the ring of integers in the completion of the maximal unramified extension of  $K$ . The closed subscheme  $Y_{1,i_l}$  is cut out in  $\mathbb{C}_{X_{U_{\text{Iw}}},s}^{\wedge}$  by  $X_{i_l} = 0$  and  $Y_{2,j_l}$  is cut out by  $Y_{j_l} = 0$ .

The action of  $G(\mathbb{A}^{\infty,p})$  extends to the inverse system  $X_{U_{\text{Iw}}}/\mathbb{C}_K$ . There is a universal abelian variety  $\mathcal{A}_{U_{\text{Iw}}}/\mathbb{C}_K$  and the actions of  $G(\mathbb{A}^{\infty})$  and  $a_{\xi}$  extend to it. We can define a stratification of the special fiber of  $\mathcal{A}_{U_{\text{Iw}}}$  by

$$\mathcal{A}_{U_{\text{Iw}},S,T} = \mathcal{A}_{U_{\text{Iw}}} \times_{X_{U_{\text{Iw}}}} X_{U_{\text{Iw}},S,T}.$$

Moreover,  $\mathcal{A}_{U_{\text{Iw}}}$  satisfies the same geometric properties as  $X_{U_{\text{Iw}}}$  with respect to the above stratification of its special fiber and the analogous statement holds for  $\mathcal{A}_{U_{\text{Iw}}}^{m_{\xi}}$ . In particular, we shall see in the next section (or it follows from Section 3 of [Caraiani 2012]) that  $\mathcal{A}_{U_{\text{Iw}}}^{m_{\xi}}$  can be endowed with a vertical logarithmic structure  $M$  such that

$$(\mathcal{A}_{U_{\text{Iw}}}^{m_{\xi}}, M) \rightarrow (\text{Spec } \mathbb{C}_K, \mathbb{N})$$

is log smooth, where  $(\text{Spec } \mathbb{C}_K, \mathbb{N})$  is the canonical log structure associated to the closed point. Also, we will see that its special fiber is of Cartier type. This means that we can define the log crystalline cohomology of  $(\mathcal{A}_{U_{\text{Iw}}}^{m_{\xi}}, M)$ . Indeed, if  $W = W(k)$  is the ring of Witt vectors of  $k$ , then we let

$$H_{\text{cris}}^*(\mathcal{A}_{U_{\text{Iw}}}^{m_{\xi}}/W)$$

be the log crystalline cohomology of  $(\mathcal{A}_{U_{\text{Iw}}}^{m_{\xi}} \times_{\mathbb{C}_K} k, M)$  (here we suppress  $M$  from the notation). This also has an action of  $a_{\xi}$  as an idempotent and of  $G(\mathbb{A}^{\infty})$ .

From the isomorphism

$$H^{2n-2}(X, \mathcal{L}_{\xi}) \simeq a_{\xi} H^{2n-2+m_{\xi}}(\mathcal{A}^{m_{\xi}}, \bar{\mathbb{Q}}_l(t_{\xi}))$$

and Corollary 2.2, we see that it is enough to show that

$$a_{\xi} \text{WD}(H^{2n-2+m_{\xi}}(\mathcal{A}^{m_{\xi}}, \bar{\mathbb{Q}}_l(t_{\xi})|_{\text{Gal}(\bar{K}/K)})[\Pi^1, \mathbb{S}])$$

is pure. Let  $\tau_0 : W \hookrightarrow \bar{\mathbb{Q}}_l$  be an embedding over  $\mathbb{Z}_l$ . By the semistable comparison theorem of [Nizioł 2008], we have

$$\begin{aligned} \lim_{\substack{\rightarrow \\ U_{\text{Iw}}}} a_{\xi} (H_{\text{cris}}^{2n-2+m_{\xi}}(\mathcal{A}_{U_{\text{Iw}}}^{m_{\xi}} \times_{\mathbb{C}_K} k/W) \otimes_{W, \tau_0} \bar{\mathbb{Q}}_l(t_{\xi}))[\Pi^1, \mathbb{S}] \\ \simeq \lim_{\substack{\rightarrow \\ U_{\text{Iw}}}} a_{\xi} \text{WD}(H^{2n-2+m_{\xi}}(\mathcal{A}^{m_{\xi}} \times_{\mathbb{C}_K} \bar{K}, \bar{\mathbb{Q}}_l(t_{\xi})|_{\text{Gal}(\bar{K}/K)})[\Pi^1, \mathbb{S}]), \end{aligned}$$

where the crystalline cohomology on the left-hand side as constructed in [Hyodo and Kato 1994] has *a priori* the structure of a  $(\varphi, N)$ -module over  $W$ , but which gives rise to a Weil–Deligne representation  $(r, N)$  of  $W_K$  by setting  $r(\sigma) := \varphi^{n[k:\mathbb{F}_p]}$  whenever  $\sigma \in W_K$  is a lift of  $\text{Frob}_k^n$ . Therefore, it suffices to understand the (direct limit of the) log crystalline cohomology of the special fiber of  $\mathcal{A}_{U_{\text{Iw}}}^{m_\xi}$ . Note that the semistable comparison theorem was first proved by Kato [1994a] and Tsuji [1999] for proper, vertical log schemes with semistable reduction; the reason for citing Niziol’s work is that her main theorem applies to a general fine and saturated, log-smooth, proper, vertical  $(\text{Spec } \mathbb{O}_K, \mathbb{N})$ -scheme with special fiber of Cartier type. The fact that  $(\mathcal{A}_{U_{\text{Iw}}}^{m_\xi}, M)$  satisfies all these properties follows immediately from the explicit description of the log structure  $M$  in Section 3.

We summarize the above discussion in the following corollary:

**Corollary 2.3.** *The Weil–Deligne representation*

$$WD(R_l(\Pi_{F'}^0)|_{\text{Gal}(\overline{F_p}/F_p)})^{\text{F-ss}}$$

is pure if

$$\lim_{\rightarrow U_{\text{Iw}}} a_\xi(H_{\text{cris}}^{2n-2+m_\xi}(\mathcal{A}_{U_{\text{Iw}}}^{m_\xi} \times_{\mathbb{O}_K} k/W) \otimes_{W, \tau_0} \overline{\mathbb{Q}}_l(t_\xi))[\Pi^{1, \mathfrak{S}}]$$

is pure, where  $\mathfrak{S}$  is chosen such that it contains  $l$ .

### 3. Log crystalline cohomology

**3A. Log structures.** Let  $\mathbb{O}_K$  be the ring of integers in a finite extension  $K$  of  $\mathbb{Q}_p$  with uniformizer  $\varpi$  and residue field  $k$ . (Here,  $p$  is some prime number, which will be taken to equal  $l$  for our applications to local-global compatibility.) Let  $W = W(k)$  be the ring of Witt vectors of  $k$ , with  $W_n = W_n(k)$  referring to the Witt vectors of length  $n$  over  $k$ . Let  $W_{(K)}$  be the ring of integers in the completion of the maximal unramified extension of  $K$ .

Let  $X/\mathbb{O}_K$  be a scheme locally of finite type such that the completions of the strict henselizations  $\mathbb{O}_{X, s}^\wedge$  at closed geometric points  $s$  of  $X$  are isomorphic to

$$W_{(K)}\llbracket X_1, \dots, X_n, Y_1, \dots, Y_n, Z_1, \dots, Z_m \rrbracket / (X_{i_1} \cdots X_{i_r} - \varpi, Y_{j_1} \cdots Y_{j_s} - \varpi)$$

for some indices  $i_1, \dots, i_r, j_1, \dots, j_s \in \{1, \dots, n\}$  and some  $1 \leq r, s \leq n$ . Also assume that the special fiber  $Y$  is a union of closed subschemes  $Y_{1, j}$  with  $j \in \{1, \dots, n\}$  which are cut out by one local equation in  $\mathbb{O}_X$ , such that if  $s$  is a closed geometric point of  $Y_{1, j}$ , then  $j \in \{i_1, \dots, i_r\}$  and  $Y_{1, j}$  is cut out in  $\mathbb{O}_{X, s}^\wedge$  by the equation  $X_j = 0$ . Similarly, assume that  $Y$  is a union of closed subschemes  $Y_{2, j}$  with  $j \in \{1, \dots, n\}$ , which are cut out by one local equation in  $\mathbb{O}_X$  such that if  $s$  is a closed geometric point of  $Y_{2, j}$  then  $j \in \{j_1, \dots, j_s\}$  and  $Y_{2, j}$  is cut out in  $\mathbb{O}_{X, s}^\wedge$  by

the equation  $Y_j = 0$ . Then, by Lemma 2.9 of [Caraiani 2012],  $X$  is Zariski-locally étale over

$$X_{r,s,m} = \text{Spec } \mathbb{O}_K[X_1, \dots, X_n, Y_1, \dots, Y_n, Z_1, \dots, Z_m]/(X_1 \cdots \varpi, Y_1 \cdots Y_s - \varpi).$$

The closed subschemes  $Y_{i,j}$  for  $i = 1, 2$  and  $j = 1, \dots, n$  are Cartier divisors, which in the local model  $X_{r,s,m}$  correspond to the divisors  $X_j = 0$  or  $Y_j = 0$ .

Let  $Y/k$  be the special fiber of  $X$ . For  $1 \leq i, j \leq n$  we define  $Y^{(i,j)}$  to be the disjoint union of the closed subschemes of  $Y$

$$(Y_{1,l_1} \cap \cdots \cap Y_{1,l_i}) \cap (Y_{2,m_1} \cap \cdots \cap Y_{2,m_j}),$$

as  $\{l_1, \dots, l_i\}$  (resp.  $\{m_1, \dots, m_j\}$ ) range over subsets of  $\{1, \dots, n\}$  of cardinality  $i$  (resp.  $j$ ). Each  $Y^{(i,j)}$  is a proper smooth scheme over  $k$  of dimension  $2n - i - j$ .

**Remark 3.1.** Even though this section is general, we will only apply the results of this section in the case when  $X$  is  $\mathcal{A}_{U_{\text{Iw}}}$  for some compact open subgroup  $U_{\text{Iw}} \subset G(\mathbb{A}^\infty)$  with Iwahori-level structure at  $\mathfrak{p}_1$  and  $\mathfrak{p}_2$ .  $X_{U_{\text{Iw}}}$  (and therefore  $\mathcal{A}_{U_{\text{Iw}}}$  as well) satisfies the above conditions by Proposition 2.8 of [Caraiani 2012]. The prime  $p$  is meant to be identified with  $l$ .

Let  $(\text{Spec } \mathbb{O}_K, \mathbb{N})$  be the log scheme corresponding to  $\text{Spec } \mathbb{O}_K$  endowed with the canonical log structure associated to the special fiber. This is given by the map  $1 \in \mathbb{N} \mapsto \varpi \in \mathbb{O}_K$ . We endow  $X$  with the log structure  $M$  associated to the special fiber  $Y$ . Let  $j : X_K \rightarrow X$  be the open immersion and  $i : Y \rightarrow X$  be the closed immersion. This log structure is defined by

$$M = j_*(\mathbb{O}_{X_K}^\times) \cap \mathbb{O}_X \rightarrow \mathbb{O}_X.$$

We have a map of log schemes  $(X, M) \rightarrow (\text{Spec } \mathbb{O}_K, \mathbb{N})$ , given by sending  $1 \in \mathbb{N}$  to  $\varpi \in M$ . Locally, we have a chart for this map, given by

$$\mathbb{N} \rightarrow \mathbb{N}^r \oplus \mathbb{N}^s / (1, \dots, 1, 0, \dots, 0) = (0, \dots, 0, 1, \dots, 1),$$

$$1 \mapsto (1, \dots, 1, 0, \dots, 0) = (0, \dots, 0, 1, \dots, 1).$$

It is easy to see from this that  $(X, M)/(\text{Spec } \mathbb{O}_K, \mathbb{N})$  is log smooth and that the log structure  $M$  on  $X$  is fine, saturated and vertical. We can pull back  $M$  to a log structure on  $Y$ , which we still denote  $M$  and then we get a log-smooth map of log schemes

$$(Y, M) \rightarrow (\text{Spec } k, \mathbb{N}).$$

(Here we have the canonical log structure on  $k$  associated to  $1 \in \mathbb{N} \mapsto 0 \in k$ , which is the same as the pullback of the canonical log structure on  $\text{Spec } \mathbb{O}_K$ .) Note that, since  $(X, M)$  is saturated over  $(\text{Spec } \mathbb{O}_K, \mathbb{N})$ , its special fiber is of Cartier type (see [Tsuji 1997]).

We can also endow  $X$  with log structures  $\tilde{M}_1$ ,  $\tilde{M}_2$  and  $\tilde{M}$ . Let  $U_{i,j}$  be the complement of  $Y_{i,,j}$  in  $X$  for  $i = 1, 2$  and  $j = 1, \dots, n$ . Let

$$j_{i,j} : U_{i,j} \rightarrow X$$

denote the open immersion. We define  $\tilde{M}_1$ ,  $\tilde{M}_2$  and  $\tilde{M}$  as follows

$$\begin{aligned} \tilde{M}_1 &= \left( \bigoplus_{j=1}^n (j_{1,j*}(\mathbb{O}_{U_{1,j}}^\times) \cap \mathbb{O}_X) \right) / \sim, \\ \tilde{M}_2 &= \left( \bigoplus_{j=1}^n (j_{1,j*}(\mathbb{O}_{U_{1,j}}^\times) \cap \mathbb{O}_X) \right) / \sim, \\ \tilde{M} &= \left( \bigoplus_{j=1}^n (j_{1,j*}(\mathbb{O}_{U_{1,j}}^\times) \cap \mathbb{O}_X) \oplus \bigoplus_{j=1}^n (j_{2,j*}(\mathbb{O}_{U_{2,j}}^\times) \cap \mathbb{O}_X) \right) / \sim, \end{aligned}$$

where  $\sim$  signifies that we have identified the image of  $\mathbb{O}_X^\times$  in all the terms of the direct sums (in other words, we are taking an amalgamated sum of the log structures associated to each of the  $Y_{i,j}$ ). We have a map  $\tilde{M} \rightarrow M$  given by inclusion on each  $\mathbb{O}_{U_{i,j}}^\times$ .

**Lemma 3.2.** *Locally on  $X$ , we have a chart for  $\tilde{M}$  given by*

$$\begin{aligned} X &\rightarrow \text{Spec } \mathbb{O}_K[X_1, \dots, X_n, Y_1, \dots, Y_n, Z_1, \dots, Z_m] / (X_1 \cdots X_r - \varpi, Y_1 \cdots Y_r - \varpi) \\ &\rightarrow \text{Spec } \mathbb{Z}[\mathbb{N}^r \oplus \mathbb{N}^s], \end{aligned}$$

where  $(0, \dots, 0, 1, 0, \dots, 0) \mapsto X_i$  if the 1 is in the  $i$ -th position and  $1 \leq i \leq r$  and  $(0, \dots, 0, 1, 0, \dots, 0) \mapsto Y_{i-r}$  if the 1 is in the  $i$ -th position and  $r + 1 \leq i \leq r + s$ .

*Proof.* We shall make use of Kato and Niziol's results on log smoothness and log regularity, namely:

- If  $f : T \rightarrow S$  is a log smooth morphism of fs log schemes with  $S$  log regular then  $T$  is log regular (see 8.2 of [Kato 1994b]).
- If  $T$  is log regular, then  $M_T = j_* \mathbb{O}_U^\times \cap \mathbb{O}_X$ , where  $j : U \hookrightarrow T$  is the inclusion of the open subset of triviality of  $T$  (see 8.6 of [Niziol 2006]).

Let us define the following log schemes over  $(\text{Spec } \mathbb{O}_K, \text{triv})$ :

$$\begin{aligned} \tilde{U} &:= \text{Spec } \mathbb{O}_K[X_1, \dots, X_n, \sigma] / (X_1 \cdots X_r - \sigma), \\ \tilde{V} &:= \text{Spec } \mathbb{O}_K[Y_1, \dots, Y_n, \tau] / (Y_1 \cdots Y_s - \tau), \\ W &:= \text{Spec } \mathbb{O}_K[Z_1, \dots, Z_m], \\ Z &:= \tilde{U} \times_{(\text{Spec } \mathbb{O}_K, \text{triv})} \tilde{V} \times_{(\text{Spec } \mathbb{O}_K, \text{triv})} W. \end{aligned}$$

Then  $Z$ , equipped with the product log structure  $L$ , is smooth over  $\mathbb{O}_K$  and log smooth over  $(\text{Spec } \mathbb{O}_K[\sigma, \tau], \text{triv})$ . Therefore,  $Z$  is regular. The log structure  $L$  is given by the simple normal crossings divisor

$$D := \left( \bigcup_{j=1}^r (X_j = 0) \right) \cup \left( \bigcup_{j=1}^s (Y_j = 0) \right).$$

Since  $Z$  is regular, the log structure  $L$  is the same as the amalgamation of the log structures defined by the smooth divisors  $(X_j = 0)$ ,  $(Y_j = 0)$ . Locally on  $X$ , we have a commutative diagram of schemes with a cartesian square

$$\begin{array}{ccccc} X & \longrightarrow & X_{r,s,m} & \longrightarrow & Z \\ & & \downarrow & & \downarrow \\ & & \text{Spec } \mathbb{O}_K & \longrightarrow & \text{Spec } \mathbb{O}_K[\tau, \sigma] \end{array} \tag{3A.1}$$

where the inverse image of  $(X_j = 0)$  in  $X$  is  $Y_j^1$  and the inverse image of  $(Y_j = 0)$  in  $X$  is  $Y_j^2$ . Therefore, the log structure on  $X$  induced by that of  $Z$  coincides with the log structure  $\tilde{M}$  defined as the amalgamated sum of the log structures induced by the  $Y_j^1$  and  $Y_j^2$ .  $\square$

If we endow  $\text{Spec } \mathbb{O}_K$  with the log structure  $\mathbb{N}^2$  associated to  $(a, b) \in \mathbb{N}^2 \mapsto \pi^{a+b} \in \mathbb{O}_K$ , then we claim that we have a log-smooth map of log schemes

$$(X, \tilde{M}) \rightarrow (\text{Spec } \mathbb{O}_K, \mathbb{N}^2) \tag{3A.2}$$

whose chart is given locally by

$$(a, b) \in \mathbb{N}^2 \mapsto (a, \dots, a, b, \dots, b) \in \mathbb{N}^r \oplus \mathbb{N}^s.$$

By definition,  $\tilde{M}$  is the amalgamated sum of  $\tilde{M}_1$  and  $\tilde{M}_2$  as log structures on  $X$  (or, in other words,  $\tilde{M}$  is the log structure associated to the prelog structure  $\tilde{M}_1 \oplus \tilde{M}_2 \rightarrow \mathbb{O}_X$ ). Therefore, it suffices to prove the following lemma:

**Lemma 3.3.** *We can define a global map of log schemes  $(X, \tilde{M}_1) \rightarrow (\text{Spec } \mathbb{O}_K, \mathbb{N})$  which locally admits the chart given by the diagonal embedding  $\mathbb{N} \rightarrow \mathbb{N}^r$ .*

*Proof.* It suffices to show that  $\varpi$  is a global section of  $\tilde{M}_1$ , since then we can simply map  $1 \in \mathbb{N}$  to  $\varpi \in \tilde{M}_1$ . For this, note that we have a natural map of log structures on  $X$

$$\tilde{M}_1 \rightarrow M,$$

since the open subset of triviality of  $\tilde{M}_1$  is the generic fiber of  $X$  and  $M$  is the log structure defined by the inclusion of the generic fiber. Moreover, we can check locally that this map is injective, since it can be described by the chart  $\mathbb{N}^r \rightarrow \mathbb{N}^r \oplus \mathbb{N}^s \rightarrow (\mathbb{N}^r \oplus \mathbb{N}^s)/\mathbb{N}$  for  $r, s \geq 1$ , where the first map is the identity on

the first factor. Now, locally on  $X$  we have the equation  $X_1 \cdots X_r = \varpi$ , where  $X_i$  are local equations defining the closed subschemes  $Y_i^1$  of  $X$ . By definition, the  $X_i$  are local sections of  $\tilde{M}_1$ , so  $\varpi$  is a local section of  $\tilde{M}_1$ . But  $\varpi$  is also a global section of  $M$  and  $\tilde{M}_1 \hookrightarrow M$ , so  $\varpi$  is a global section of  $\tilde{M}_1$ .  $\square$

**Lemma 3.4.** *We have a cartesian diagram of maps of log schemes*

$$\begin{array}{ccc} (X, M) & \longrightarrow & (X, \tilde{M}) \\ \downarrow & & \downarrow \\ (\mathrm{Spec} \mathbb{O}_K, \mathbb{N}) & \longrightarrow & (\mathrm{Spec} \mathbb{O}_K, \mathbb{N}^2) \end{array}$$

where the bottom horizontal arrow is the identity on the underlying schemes and maps  $(a, b) \in \mathbb{N}^2$  to  $a + b \in \mathbb{N}$ .

*Proof.* We go back to the notation used in the proof of Lemma 3.2. Locally on  $X$ , we have the following commutative diagram of log schemes

$$\begin{array}{ccccc} (X, M) & \longrightarrow & \tilde{U} \times_{\mathrm{Spec} \mathbb{O}_K[u]} \tilde{V} \times W & \longrightarrow & Z \\ \downarrow & & \downarrow & & \downarrow \\ (\mathrm{Spec} \mathbb{O}_K, \mathbb{N}) & \longrightarrow & (\mathrm{Spec} \mathbb{O}_K[u], \mathbb{N}) & \longrightarrow & (\mathrm{Spec} \mathbb{O}_K[\tau, \sigma], \mathbb{N}^2) \end{array}$$

where in the bottom row both  $\tau$  and  $\sigma$  are mapped to  $u$ , which is in turn mapped to 0. The second square is cartesian and the horizontal maps in it are closed, but not exact, immersions. The first bottom map is an exact closed immersion, while the first top map is the composition of an étale morphism with an exact closed immersion. The lemma follows from the commutative diagram (3A.1) and the above diagram.  $\square$

**3B. Variations on the logarithmic de Rham–Witt complex.** Define the prelog structure  $\mathbb{N}^2 \rightarrow W_n[\tau, \sigma]$  given by  $(a, b) \mapsto \tau^a \sigma^b$ . By abuse of notation, we write  $(\mathrm{Spec} W_n[\tau, \sigma], \mathbb{N}^2)$  for the log scheme endowed with the associated log structure. We have the composite map of log schemes

$$(Y, \tilde{M}) \rightarrow (\mathrm{Spec} k, \mathbb{N}^2) \rightarrow (\mathrm{Spec} W_n[\tau, \sigma], \mathbb{N}^2),$$

where  $\mathbb{N}^2 \rightarrow \mathbb{N}^2$  is the obvious isomorphism. We shall call  $(Z, \tilde{N})$  a *lifting* for this morphism if  $(Z, \tilde{N})$  is a fine log scheme such that the composite map  $(Y, \tilde{M}) \rightarrow (\mathrm{Spec} W_n[\tau, \sigma], \mathbb{N}^2)$  factors through  $f : (Y, \tilde{M}) \rightarrow (Z, \tilde{N})$  which is a closed immersion, and a map  $(Z, \tilde{N}) \rightarrow (\mathrm{Spec} W_n[\tau, \sigma], \mathbb{N}^2)$  which is log smooth. Such liftings always exists locally on  $Y$  and give rise to embedding systems as defined in Paragraph 2.18 of [Hyodo and Kato 1994]. If  $(U, \tilde{M}_U) \rightarrow (Y, \tilde{M})$  is a covering and  $(Z, \tilde{N})$  is a lifting for  $(U, \tilde{M}_U) \rightarrow (\mathrm{Spec} W_n[\tau, \sigma], \mathbb{N}^2)$ , then we may define

an embedding system  $((U^i, \tilde{M}_U^i), (Z^i, \tilde{N}^i))$  for  $(Y, \tilde{M}) \rightarrow (\text{Spec } W_n[\tau, \sigma], \mathbb{N}^2)$  by taking the fiber product of  $i + 1$  copies of  $U$  over  $Y$  and of  $i + 1$  copies of  $(Z, \tilde{N})$  over  $(\text{Spec } W_n[\tau, \sigma], \mathbb{N}^2)$ . Since  $(Y, \tilde{M})$  is an fs log scheme, we may assume the same for the local lifting  $(Z, \tilde{N})$ .

Let  $C_{(Y, \tilde{M})/(W_n, \text{triv})}^\bullet$  be the crystalline complex associated to the embedding system obtained from local liftings  $(Z^\bullet, \tilde{N}^\bullet)$ , and define

$$\tilde{C}_Y^\bullet := C_{(Y, \tilde{M})^\bullet/(W_n, \text{triv})} \otimes_{W_n(\tau, \sigma)} W_n.$$

Let  $\text{Spec } W_n[u]$  be endowed with the log structure associated to  $1 \in \mathbb{N} \mapsto u \in W_n[u]$ . Consider the map of log schemes  $G : (\text{Spec } W_n[u], \mathbb{N}) \rightarrow (\text{Spec } W_n[\tau, \sigma], \mathbb{N}^2)$  given by  $\tau, \sigma \mapsto u$  and  $(a, b) \in \mathbb{N}^2 \mapsto a + b \in \mathbb{N}$ . The pullback of  $(Y, \tilde{M})$  along  $G$  is just  $(Y, M)$ . Let  $(Z', N')$  be the pullback of  $(Z, \tilde{N})$  along  $G$ , equipped with a map  $f' : (Y', M') \rightarrow (Z', N')$  which is the pullback of  $f$ . Then  $(Z', N')$  is a (local) lifting for  $(Y, M) \rightarrow (\text{Spec } W_n[u], \mathbb{N})$ , and gives rise to an embedding system for this morphism. Indeed, what we need to check is that  $(Z', N') \rightarrow (\text{Spec } W_n[u], \mathbb{N})$  is log smooth and that  $f'$  is a closed immersion of log schemes. For the first we note that log-smoothness is preserved under base change in the category of log schemes, and that

$$(Z', N') = (((Z, \tilde{N}) \times_G (\text{Spec } W_n[u], \mathbb{N}))^{\text{int}})^{\text{sat}} \rightarrow (Z, \tilde{N}) \times_G (\text{Spec } W_n[u], \mathbb{N})$$

is log smooth. We also note that  $g : Y \rightarrow (Z \times_{\text{Spec } W_n[\tau, \sigma]} \text{Spec } W_n[u])$  is a closed immersion, since  $Y \rightarrow Z$  is a closed immersion. The morphism of schemes  $Z' \rightarrow (Z \times_{\text{Spec } W_n[\tau, \sigma]} \text{Spec } W_n[u])$  is a composition of a finite morphism with a closed immersion, so  $Y \rightarrow Z'$  is a closed immersion as well. Also,  $g^*(\tilde{N} \oplus_{\mathbb{N}^2} \mathbb{N}) \rightarrow M$  is surjective and factors through  $(f')^*(N') \rightarrow M$ , so  $(f')^*(N') \rightarrow M$  is surjective as well.

We now follow the constructions in Section 3.6 of [Hyodo and Kato 1994] using the embedding system obtained from the liftings  $(Z', N')$ . Let  $C_{(Y, M)/(W_n, \text{triv})}^\bullet$  be the crystalline complex associated to the composite  $(Z', N') \rightarrow (W_n, \text{triv})$ . Define

$$\tilde{C}_Y^\bullet := C_{(Y, M)/(W_n, \text{triv})}^\bullet \otimes_{W_n(u)} W_n.$$

On the other hand, let  $Z'' = Z' \times_{\text{Spec } W_n[u]} \text{Spec } W_n\langle u \rangle$  be endowed with  $N''$  the inverse image of the log structure  $N'$ . Let  $\mathcal{L}$  be the log structure on  $\text{Spec } W_n\langle u \rangle$  obtained by taking the inverse image of (the log structure associated to)  $\mathbb{N}$  on  $\text{Spec } W_n[u]$ . Then  $(Z'', N'')$  gives rise to an embedding system for

$$(Y, M) \rightarrow (\text{Spec } W_n\langle u \rangle, \mathcal{L}),$$

with crystalline complex  $C_{(Y, M)/(\text{Spec } W_n\langle u \rangle, \mathcal{L})}^\bullet$ . Define

$$C_Y^\bullet := C_{(Y, M)/(\text{Spec } W_n\langle u \rangle, \mathcal{L})}^\bullet \otimes_{W_n(u)} W_n.$$

Note that  $C_Y^\bullet$  is the crystalline complex  $C_{(Y,M)/(W_n, \mathbb{N})}^\bullet$  with respect to the embedding system obtained from  $(Z' \times_{\text{Spec } W_n[u]} \text{Spec } W_n, N''')$ . As in Section 3.6 of [Hyodo and Kato 1994], we have an exact sequence of complexes

$$0 \rightarrow C_Y^\bullet[-1] \rightarrow \tilde{C}_Y^\bullet \rightarrow C_Y^\bullet \rightarrow 0, \tag{3B.1}$$

where the second arrow is  $\wedge (du/u)$  and the third arrow is the canonical projection. The monodromy operator on the crystalline cohomology of  $(Y, M)$  is induced by the connecting homomorphism of this exact sequence.

**Lemma 3.5.** *Let  $C_Z^\bullet$  be one of the complexes  $\tilde{\tilde{C}}_Y^\bullet, \tilde{C}_Y^\bullet$  or  $C_Y^\bullet$  obtained with respect to a lifting  $(Z, \tilde{N})$  of some cover  $U \rightarrow Y$ . In the derived category,  $C_Z^\bullet$  is independent of the choice of lifting  $(Z, \tilde{N})$ .*

*Proof.* We may work étale locally on  $Y$ , in which case we have to show that for any two liftings  $(Z_1, \tilde{N}_1)$  and  $(Z_2, \tilde{N}_2)$  we have a canonical quasi-isomorphism between the corresponding complexes and moreover, that these quasi-isomorphisms satisfy the obvious cocycle condition for three different liftings.

First, we show that the complexes corresponding to  $(Z_1, \tilde{N}_1)$  and  $(Z_2, \tilde{N}_2)$  are quasi-isomorphic. We may assume that  $i_i : (Y, \tilde{M}) \rightarrow (Z_i, \tilde{N}_i)$  is an exact closed immersion for  $i = 1, 2$ . Let  $i_{12} : (Y, \tilde{M}) \rightarrow (Z_1 \times_{W_n} Z_2, \tilde{N}_{1 \times 2})$  be the diagonal immersion of  $(Y, \tilde{M})$  into the fiber product of  $(Z_1, \tilde{N}_1)$  and  $(Z_2, \tilde{N}_2)$  as fs log schemes over  $(W_n, \text{triv})$ . Let  $(Z_{12}, \tilde{N}_{12})$  be a log scheme such that étale locally on  $Y$  we have a factorization of  $i_{12}$

$$(Y, \tilde{M}) \xrightarrow{f} (Z_{12}, \tilde{N}_{12}) \xrightarrow{g} (Z_1 \times Z_2, \tilde{N}_{1 \times 2}),$$

with  $g$  log étale and  $f$  an exact closed immersion. This factorization is possible by Lemma 4.10 of [Kato 1989]. Let  $D_i$  be the PD-envelope of  $Y$  in  $Z_i$  (again, for  $i = 1, 2$  or  $12$ ). (Since we have exact closed immersions, the logarithmic PD-envelope coincides with the usual PD-envelope in these cases.) It suffices to show that the canonical map

$$\omega_{(Z_1, \tilde{N}_1)/W_n, \text{triv}}^\bullet \otimes_{\mathbb{C}_{Z_1}} \mathbb{C}_{D_1} \rightarrow \omega_{(Z_{12}, \tilde{N}_{12})/W_n, \text{triv}}^\bullet \otimes_{\mathbb{C}_{Z_{12}}} \mathbb{C}_{D_{12}} \tag{3B.2}$$

is a quasi-isomorphism. This follows from Paragraph 2.21 of [Hyodo and Kato 1994]. For completeness, we sketch the proof here. Let  $p_1 : (Z_{12}, N_{12}) \rightarrow (Z_1, N_1)$  be the log-smooth map induced by projection onto the first factor. For any geometric point  $\bar{y}$  of  $Y$ , the stalks at  $\bar{y}$  of  $N_{12}$  and  $p_1^* N_1$  coincide, so by replacing  $(Z_{12}, N_{12})$  with an étale neighborhood of  $\bar{y} \rightarrow Z_{12}$ , we may assume that  $N_{12} = p_1^* N_1$ . Then the map  $p_1 : Z_{12} \rightarrow Z_1$  is smooth in the usual sense. Since

the problem is étale local on  $Y$ , we may assume that  $Z_{12} \simeq Z_1 \otimes_{W_n} W_n[t_1, \dots, t_r]$  for some positive integer  $r$  such that  $Y$  is contained in the closed subscheme of  $Z_{12}$  defined by  $t_1 = \dots = t_r = 0$ . As in Proposition 6.5 of [Kato 1989], we also have  $\mathbb{O}_{D_{12}} \simeq \mathbb{O}_{D_1}\langle t_1, \dots, t_r \rangle$ , the PD-polynomial ring over  $\mathbb{O}_{D_1}$  in  $r$  variables. The quasi-isomorphism (3B.2) is reduced then to the standard quasi-isomorphism

$$W_n \rightarrow \Omega_{W_n[t_1, \dots, t_r]} \otimes_{W_n[t_1, \dots, t_r]} W_n\langle t_1, \dots, t_r \rangle.$$

The quasi-isomorphism (3B.2) commutes with  $\otimes_{W_n\langle \tau, \sigma \rangle} W_n$ , so it induces a quasi-isomorphism

$$\tilde{C}_{Z_1}^\bullet \xrightarrow{\sim} \tilde{C}_{Z_{12}}^\bullet.$$


Now consider the morphism  $Z'_{12} \rightarrow Z'_1$  obtained by pulling back  $Z_{12} \rightarrow Z_1$  along  $G$ . We claim that the canonical morphisms  $\tilde{C}_{Z'_{12}}^\bullet \rightarrow \tilde{C}_{Z'_1}^\bullet$  and  $C_{Z'_{12}}^\bullet \rightarrow C_{Z'_1}^\bullet$  are quasi-isomorphisms as well. This is proved in the same way as in the case of  $\tilde{C}^\bullet$  (for  $C_{Z'_{12}}^\bullet \rightarrow C_{Z'_1}^\bullet$  this amounts to proving that the logarithmic de Rham–Witt complex is independent of the choice of embedding system). The quasi-isomorphisms are also compatible with the canonical maps  $\tilde{C}_Z^\bullet \rightarrow \tilde{C}_{Z'}^\bullet \rightarrow C_{Z'}^\bullet$ .

Note that the above result also implies that in the derived category,  $C^\bullet$  commutes with étale base change. Indeed, if  $Y_2/Y_1$  is étale and  $(Z_1, \tilde{N}_1)$  is a lifting for  $(Y_1, \tilde{M}) \rightarrow (\text{Spec } W_n[\tau, \sigma], \mathbb{N}^2)$ , then by [Grothendieck 1967, 18.1.1] we can find, Zariski locally on  $Y_2$ , an étale morphism  $Z_2 \rightarrow Z_1$  such that the following diagram is cartesian

$$\begin{array}{ccc} Y_2 & \longrightarrow & Z_2 \\ \downarrow & & \downarrow \\ Y_1 & \longrightarrow & Z_1 \end{array}$$

We take  $\tilde{N}_2$  on  $Z_2$  to be the inverse image of  $\tilde{N}_1$ . Then  $(Z_2, \tilde{N}_2)$  is a lifting for  $(Y_2, \tilde{M}) \rightarrow (\text{Spec } W_n[\tau, \sigma], \mathbb{N}^2)$  and, since log differentials commute with étale base change [Kato 1989, Proposition 3.12],  $C_{(Z_2)}^\bullet$  on  $Y_2$  is just the pullback of  $C_{(Z_1)}^\bullet$  on  $Y_1$ .

We are left with verifying the cocycle condition. The canonical quasi-isomorphism  $\gamma_{12} : C_{Z_1}^\bullet \xrightarrow{\sim} C_{\bullet, Z_2}^\bullet$  factors through  $C_{Z_1 \times Z_2}^\bullet$ , since by construction  $Z_{12}$  is log étale over  $Z_1 \times Z_2$  and so we have a quasi-isomorphism  $C_{Z_1 \times Z_2}^\bullet \xrightarrow{\sim} C_{Z_{12}}^\bullet$ . Let  $(Z_3, \tilde{N}_3)$  be another lifting. Then we have the following commutative diagram of complexes:



where all the maps are quasi-isomorphisms. This proves the cocycle condition.  $\square$

**Corollary 3.6.** *The following sheaves on  $Y$  are well-defined and commute with étale base change:*

$$W_n \tilde{\omega}_Y^q := \mathcal{H}^q(\tilde{\mathcal{C}}_Y^\bullet), \quad W_n \tilde{\omega}_Y^q := \mathcal{H}^q(\tilde{\mathcal{C}}_Y^\bullet) \quad \text{and} \quad W_n \omega_Y^q := \mathcal{H}^q(C_Y^\bullet).$$

The sheaves  $W_n \omega_Y^q$  make up the  $q$ -th terms of the log de Rham–Witt complex associated to  $(Y, M)$ . We have canonical morphisms of sheaves on  $Y$ :

$$W_n \tilde{\omega}_Y^q \rightarrow W_n \tilde{\omega}_Y^q \rightarrow W_n \omega_Y^q.$$

In order to understand the monodromy  $N$ , we will study the short exact sequence of complexes

$$0 \rightarrow W_n \omega_Y^\bullet[-1] \rightarrow W_n \tilde{\omega}_Y^\bullet \rightarrow W_n \omega_Y^\bullet \rightarrow 0,$$

which we obtain below from the short exact sequence (3B.1). In Section 4 we will construct a resolution of this short exact sequence in terms of some subquotients of  $W_n \tilde{\omega}_Y^\bullet$ . For now, since these complexes are independent of the choice of lifting, we will fix some specific kinds of liftings of  $(Y, \tilde{M})$  over  $(W[\tau, \sigma], \mathbb{N}^2)$ , which we call *admissible liftings*, following the terminology used in [Hyodo 1991] and [Mokrane 1993]. Since  $Y$  is locally étale over

$$Y_{r,s,m} = \text{Spec } k[X_1, \dots, X_n, Y_1, \dots, Y_n, Z_1, \dots, Z_m]/(X_1 \cdots X_r, Y_1 \cdots Y_s),$$

we consider the lifting

$$\begin{aligned} Z_{r,s,m} & \\ &= \text{Spec } W[X_1, \dots, X_n, Y_1, \dots, Y_n, Z_1, \dots, Z_m, \tau, \sigma]/(X_1 \cdots X_r - \tau, Y_1 \cdots Y_s - \sigma) \end{aligned}$$

of  $(Y_{r,s,m}, \mathbb{N}^r \oplus \mathbb{N}^s)/(W[\tau, \sigma], \mathbb{N}^2)$ . The log structure on  $Z_{r,s,m}$  is also induced from  $\mathbb{N}^r \oplus \mathbb{N}^s$  (with the obvious structure map sending  $\mathbb{N}^r$  to products of the  $X_i$

and  $\mathbb{N}^s$  to products of the  $Y_j$ ). We let  $Z/Z_{r,s,m}$  be étale and such that the diagram

$$\begin{array}{ccc} (Y, \tilde{M}) & \longrightarrow & (Z, \tilde{N}) \\ \downarrow & & \downarrow \\ (Y_{r,s,m}, \mathbb{N}^r \oplus \mathbb{N}^s) & \longrightarrow & (Z_{r,s,m}, \mathbb{N}^r \oplus \mathbb{N}^s) \end{array}$$

is cartesian, with the log structures on top obtained by pullback from the ones on the bottom. Then, locally on  $Y$ , the complexes  $W_n \tilde{\omega}_Y^\bullet$ ,  $W_n \tilde{\omega}_Y^\bullet$  and  $W_n \omega_Y^\bullet$  are just pullbacks of the corresponding complexes on  $Y_{r,s,m}$  with respect to the lifting  $(Z_{r,s,m}, \mathbb{N}^r \oplus \mathbb{N}^s)$ . Note that admissible liftings exist locally on  $Y$ .

Now we will explain the relationships between  $\tilde{C}_Y^\bullet$ ,  $\tilde{C}_Y^\bullet$  and  $C_Y^\bullet$ . First, note that we have the functoriality map  $G^* \omega_{(Z, \tilde{N})/(W_n, \text{triv})} \rightarrow \omega_{(Z', N')/(W_n, \text{triv})}$ , which induces a canonical map

$$C_{(Y, \tilde{M})/(W_n, \text{triv})}^\bullet \otimes_{W_n \langle \tau, \sigma \rangle} W_n \langle u \rangle \rightarrow C_{(Y, M)/(W_n, \text{triv})}^\bullet$$

which in turn induces a canonical map  $\tilde{C}_Y^\bullet \rightarrow \tilde{C}_Y^\bullet$ . By composition, we also get a map  $\tilde{C}_Y^\bullet \rightarrow C_Y^\bullet$ . We claim that we can identify  $\tilde{C}_Y^\bullet$  with  $\tilde{C}_Y^\bullet / (d\tau/\tau - d\sigma/\sigma) \wedge \tilde{C}_Y^\bullet$  and  $C_Y^\bullet$  with  $\tilde{C}_Y^\bullet / ((d\tau/\tau) \wedge \tilde{C}_Y^\bullet + (d\sigma/\sigma) \wedge \tilde{C}_Y^\bullet)$ . We explain this in the case of  $\tilde{C}_Y^\bullet$ :

**Lemma 3.7.** *We have an isomorphism*

$$\tilde{C}_Y^\bullet / \left( \frac{d\tau}{\tau} - \frac{d\sigma}{\sigma} \right) \wedge \tilde{C}_Y^{\bullet-1} \xrightarrow{\sim} \tilde{C}_Y^\bullet.$$

*Proof.* Let  $(Z, \tilde{N})$  be an admissible lifting of  $(Y, \tilde{M})$  over  $(\text{Spec } W_n[\tau, \sigma], \mathbb{N}^2)$ . Let  $(D, \tilde{M}_D)$  be the divided power envelope of  $(Y, \tilde{M})$  in  $(Z, \tilde{N})$ . Note that the kernel of the map  $\mathcal{O}_D \rightarrow \mathcal{O}_Y$  is generated by  $\tau^{[n]}$  and  $\sigma^{[n]}$ . The divided power envelope  $(D', M'_D)$  of  $(Y, M)$  in  $(Z', N')$  satisfies the property

$$\mathcal{O}_{D'} \simeq \mathcal{O}_D \otimes_{W_n \langle \tau, \sigma \rangle} W_n \langle u \rangle,$$

where the map  $W_n \langle \tau, \sigma \rangle \rightarrow W_n \langle u \rangle$  is given by  $\tau^{[n]}, \sigma^{[n]} \mapsto u^{[n]}$ . The complexes  $\tilde{C}_Y^\bullet$  and  $\tilde{C}_Y^\bullet$  are defined by

$$\begin{aligned} \tilde{C}_Y^\bullet &:= (\omega_{(Z, \tilde{N})/(W_n, \text{triv})}^\bullet \otimes_{\mathcal{O}_Z} \mathcal{O}_D) \otimes_{W_n \langle \tau, \sigma \rangle} W_n \langle u \rangle \otimes_{W_n \langle u \rangle} W_n \\ &= (\omega_{(Z, \tilde{N})/(W_n, \text{triv})}^\bullet \otimes_{W_n[\tau, \sigma]} W_n[u]) \otimes_{\mathcal{O}_{Z'}} \mathcal{O}_{D'} \otimes_{W_n \langle u \rangle} W_n \end{aligned}$$

and

$$\tilde{C}_Y^\bullet = (\omega_{Z', N'/W_n, \text{triv}}^\bullet) \otimes_{\mathcal{O}_{Z'}} \mathcal{O}_{D'} \otimes_{W_n \langle u \rangle} W_n.$$

Note that since we have chosen an admissible lifting,  $(Z', N')$  has  $Z \times_{W_n[\tau, \sigma]} W_n[u]$  as its underlying scheme because  $\tilde{N} \oplus_{\mathbb{N}^2} \mathbb{N}$  is already fine and saturated. It is enough

to show that the sequence

$$\begin{aligned} \omega_{(Z, \tilde{N})/(W_n, \text{triv})}^{\bullet-1} \otimes_{W_n[\tau, \sigma]} W_n[u] &\rightarrow \omega_{(Z, \tilde{N})/(W_n, \text{triv})}^{\bullet} \otimes_{W_n[\tau, \sigma]} W_n[u] \\ &\rightarrow \omega_{(Z', N')/(W_n, \text{triv})}^{\bullet} \rightarrow 0 \end{aligned} \quad (3B.3)$$

is exact, where the first map is  $\wedge(d\tau/\tau - d\sigma/\sigma)$  and the second map is induced by functoriality. We denote by  $G^*$  the pullback along  $\text{Spec } W_n[u] \rightarrow \text{Spec } W_n[\tau, \sigma]$  or along  $Z' \rightarrow Z$ . By Proposition 3.12 of [Kato 1989], we have the following diagram of (vertical) exact sequences of sheaves on  $Z'$ :

$$\begin{array}{ccc} & & 0 \\ & & \downarrow \\ G^* \omega_{(\text{Spec } W_n[\tau, \sigma], \mathbb{N}^2)/(W_n, \text{triv})}^1 \otimes_{W_n[u]} \mathbb{C}_{Z'} & \longrightarrow & \omega_{(\text{Spec } W_n[u], \mathbb{N})/(W_n, \text{triv})}^1 \otimes_{W_n[u]} \mathbb{C}_{Z'} \\ \downarrow & & \downarrow \\ G^* \omega_{(Z, \tilde{N})/(W_n, \text{triv})}^1 & \longrightarrow & \omega_{(Z', N')/(W_n, \text{triv})}^1 \\ \downarrow & & \downarrow \\ G^* \omega_{(Z, \tilde{N})/(\text{Spec } W_n[\tau, \sigma], \mathbb{N}^2)}^1 & \longrightarrow & \omega_{(Z', N')/(\text{Spec } W_n[u], \mathbb{N})}^1 \\ \downarrow & & \downarrow \\ 0 & & 0 \end{array}$$

The bottom horizontal arrow is an isomorphism, since  $(Z', N')$  was obtained by pullback from  $(Z, \tilde{N})$ . In order to show that the middle horizontal arrow is a surjection, it is enough to check that  $du/u$  is in its image, but both  $d\sigma/\sigma$  and  $d\tau/\tau$  map to  $du/u$ . We also see similarly that the kernel of the middle horizontal arrow is generated by  $d\tau/\tau - d\sigma/\sigma$ . The exactness of (3B.3) follows.  $\square$

**Corollary 3.8.** *We have an isomorphism*

$$\tilde{C}_Y^{\bullet} / \left( \frac{d\tau}{\tau} \wedge \tilde{C}_Y^{\bullet-1} + \frac{d\sigma}{\sigma} \wedge \tilde{C}_Y^{\bullet-1} \right) \simeq C_Y^{\bullet}.$$

*Proof.* This follows from the exact sequence (3B.1) and Lemma 3.7.  $\square$

**Lemma 3.9.** *The sections  $d\tau/\tau, d\sigma/\sigma \in W_n \tilde{\omega}_Y^1$  are global sections, independent of the choice of admissible lifting. The same holds for  $du/u \in W_n \tilde{\omega}_Y^1$ .*

*Proof.* We will explain the proof only for  $d\tau/\tau$  since the same proof also works for  $d\sigma/\sigma$  and  $du/u$ . We use basically the same argument as for part 3 of Lemma 3.4 of [Mokrane 1993]. We consider two admissible liftings of  $(Y, \tilde{M})$ ,  $(Z_1, \tilde{N}_1)$  and

$(Z_2, \tilde{N}_2)$ , and we let  $(Z_{12}, \tilde{N}_{12})$  be defined as in Lemma 3.5. It is enough to show that locally on  $Y$

$$\frac{d\tau}{\tau} \in \omega^1_{(Z_1, \tilde{N}_1)/(W_n, \text{triv})} \otimes_{\mathbb{O}_{Z_1}} \mathbb{O}_{D_1}$$

and

$$\frac{d\tau'}{\tau'} \in \omega^1_{(Z_2, \tilde{N}_2)/(W_n, \text{triv})} \otimes_{\mathbb{O}_{Z_2}} \mathbb{O}_{D_2}$$

have the same image in  $\mathcal{H}^1(\omega^{\bullet}_{(Z_{12}, \tilde{N}_{12})/(W_n, \text{triv})} \otimes_{\mathbb{O}_{Z_{12}}} \mathbb{O}_{D_{12}})$ .

Note that  $d\tau/\tau \in \tilde{N}_1$  and  $d\tau'/d\tau' \in \tilde{N}_2$  have the same image in  $\tilde{M}$ . This is because locally on  $Y$  we have commutative diagrams

$$\begin{CD} (Y, \tilde{M}) @>>> (Z_i, \tilde{N}_i) \\ @VVV @VVV \\ (k, \mathbb{N}^2) @>>> (W_n[\tau, \sigma], \mathbb{N}^2) \end{CD}$$

for  $i = 1, 2$ , so both  $d\tau/\tau$  and  $d\tau'/\tau'$  map to the image of  $(1, 0) \in \mathbb{N}^2$  in  $\tilde{M}$ . By the construction of  $(Z_{12}, \tilde{N}_{12})$  (see the proof of Proposition 4.10 of [Kato 1989]), we know that  $d\tau/\tau - d\tau'/\tau' = m \in \tilde{N}_{12}$ . Moreover, if  $\alpha_{12} : N_{12} \rightarrow \mathbb{O}_{Z_{12}}$  is the map defining the log structure of  $Z_{12}$  then  $m$  maps to  $0 \in \tilde{M}$ , so  $v = \alpha_{12}(m) \in \mathbb{O}_{Z_{12}}^{\times}$  maps to  $1 \in \mathbb{O}_Y$ . Therefore,

$$\frac{d\tau}{\tau} - \frac{d\tau'}{\tau'} = \frac{dv}{v}$$

for some  $v \in \mathbb{O}_{D_{12}}$  for which  $W_n\langle v - 1 \rangle \subseteq \mathbb{O}_{D_{12}}$ . But then we see that  $dv/v \in d(W_n\langle v - 1 \rangle)$ , using the fact that the power series expansion of  $\log(v)$  around 1 belongs to  $W_n\langle v - 1 \rangle$ . Therefore,  $d\tau/\tau - d\tau'/\tau'$  is exact and the lemma follows.  $\square$

As in the classical case [Illusie and Raynaud 1983; Hyodo and Kato 1994], we can define operators  $F : W_{n+1}\tilde{\omega}^q \rightarrow W_n\tilde{\omega}^q$ ,  $V : W_n\tilde{\omega}^q \rightarrow W_{n+1}\tilde{\omega}^q$  and the differential  $d : W_n\tilde{\omega}^q \rightarrow W_n\tilde{\omega}^{q+1}$ , which satisfy

$$d^2 = 0, \quad FV = VF = p, \quad dF = pFd, \quad Vd = pdV \quad \text{and} \quad FdV = d.$$

Indeed, fix local liftings  $(Z_n, \tilde{N}_n)$  of  $(Y, \tilde{M}) \rightarrow (\text{Spec } W_n[\tau, \sigma], \mathbb{N}^2)$  and denote the crystalline complex  $\tilde{C}_{Z_n}^{\bullet}$  by  $\tilde{C}_n^{\bullet}$ . We can see that  $\tilde{C}_n^{\bullet}$  is flat over  $W_n$  in the same way as in Lemma 2.22 of [Hyodo and Kato 1994] (using an admissible lifting), and we have

$$\tilde{C}_n^{\bullet} \otimes_{\mathbb{Z}/p^n\mathbb{Z}} \mathbb{Z}/p^m\mathbb{Z} \xrightarrow{\sim} \tilde{C}_m^{\bullet}$$

for  $m \leq n$ . We let  $F : W_{n+1}\tilde{\omega}^{\bullet} \rightarrow W_n\tilde{\omega}^{\bullet}$  be the map induced by  $\tilde{C}_{n+1}^{\bullet} \rightarrow \tilde{C}_n^{\bullet}$  and  $V : W_n\tilde{\omega}^{\bullet} \rightarrow W_{n+1}\tilde{\omega}^{\bullet}$  be the map induced by  $p : \tilde{C}_n^{\bullet} \rightarrow \tilde{C}_{n+1}^{\bullet}$ . We define  $d$  to

be the connecting homomorphism in the exact sequence of cohomology sheaves associated to the exact sequence of crystalline complexes

$$0 \rightarrow \tilde{C}_n^\bullet \xrightarrow{p^n} \tilde{C}_{2n}^\bullet \rightarrow \tilde{C}_n^\bullet \rightarrow 0.$$

The same operators can be defined for  $W_\bullet \tilde{\omega}_Y^\bullet$  and  $W_\bullet \omega_Y^\bullet$ .

**Lemma 3.10.** *Let  $n = 1$ . Locally, fix an admissible lifting  $(Z, \tilde{N})$  as above. Let  $\text{Fr}$  be the relative Frobenius of  $Y/k$ . We have Cartier isomorphisms*

$$\begin{aligned} C^{-1} &: \omega_Y^q \xrightarrow{\sim} \mathcal{H}^q(\text{Fr}_* \omega_Y^\bullet), \\ \tilde{C}^{-1} &: \omega_{(Z', N')/(k, \text{triv})}^q \otimes_{k[u]} k \xrightarrow{\sim} \mathcal{H}^q(\text{Fr}_*(\omega_{(Z', N')/k, \text{triv}}^\bullet \otimes_{k[u]} k)), \\ \tilde{\tilde{C}}^{-1} &: \omega_{(Z, \tilde{N})/(k, \text{triv})}^q \otimes_{k[t, s]} k \xrightarrow{\sim} \mathcal{H}^q(\text{Fr}_*(\omega_{(Z, \tilde{N})/k, \text{triv}}^\bullet \otimes_{k[t, s]} k)). \end{aligned}$$

*Proof.* Note that  $(Y, M)/(\text{Spec } k, \mathbb{N})$  is log smooth of Cartier type. The Cartier isomorphism for  $W_1 \omega_Y^q$  is then defined in Section 2.12 of [Hyodo and Kato 1994]. Similarly,  $(Z', N')/(\text{Spec } k, \text{triv})$  and  $(Z, \tilde{N})/(\text{Spec } k, \text{triv})$  are log smooth and of Cartier type. Thus, the morphisms  $\tilde{C}^{-1}$  and  $\tilde{\tilde{C}}^{-1}$  for  $\tilde{C}_Y^q$  and  $\tilde{\tilde{C}}_Y^q$  are induced from the Cartier isomorphisms for these schemes.

Since we are working locally on  $Y$ , we may assume that  $Y = Y_1 \times_k Y_2$  and that the lifting  $Z = Z_1 \times Z_2$ , where  $Z_1, Z_2$  are smooth over  $k$  and  $Y_i$  is a reduced normal crossings divisor in  $Z_i$ . Let  $\mathcal{I}_i$  be the ideal defining  $Y_i \times_k Z_{3-i}$  in  $Z$  for  $i = 1, 2$ . Define  $\omega_{1,2}^\bullet := \omega_{(Z, \tilde{N})/k}^\bullet \otimes \mathcal{I}_1 \oplus \omega_{(Z, \tilde{N})/k}^\bullet \otimes \mathcal{I}_2$ . To check that  $\tilde{C}^{-1}$  is an isomorphism, we use the following commutative diagram of exact sequences:

$$\begin{array}{ccccccc} \omega_{(Z, \tilde{N})/k}^q \otimes \mathcal{I}_1 \mathcal{I}_2 & \longrightarrow & \omega_{1,2}^q & \longrightarrow & \omega_{(Z, \tilde{N})/k}^q & \longrightarrow & \tilde{C}_Y^q \longrightarrow 0 \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ \mathcal{H}^q(\text{Fr}_* \omega_{(Z, \tilde{N})/k}^\bullet \otimes \mathcal{I}_1 \mathcal{I}_2) & \longrightarrow & \omega_{1,2}^q & \longrightarrow & \mathcal{H}^q(\text{Fr}_* \omega_{(Z, \tilde{N})/k}^\bullet) & \longrightarrow & \mathcal{H}^q(F_* \tilde{\tilde{C}}_Y^\bullet) \longrightarrow 0 \end{array}$$

The complex  $\omega_{(Z, \tilde{M})/k, \text{triv}}^\bullet$  is the same as  $\Omega_{Z_1/k}^\bullet(\log Y_1) \otimes_k \Omega_{Z_2/k}^\bullet(\log Y_2)$ , so it does satisfy a Cartier isomorphism, by [Deligne and Illusie 1987, 4.2.1.1]. Similarly, the complexes on its left are (sums of) products of complexes of the form  $\Omega_{Z_i/k}^\bullet(\pm \log Y_i)$  for  $i = 1, 2$ , which also satisfy a Cartier isomorphism, by [Deligne and Illusie 1987, 4.2.1.3]. Therefore, the first three vertical arrows are isomorphisms. Once we know the exactness of the top and bottom sequence we can also deduce that the rightmost vertical arrow is an isomorphism. The exactness of the top row follows from the definition of  $\tilde{\tilde{C}}_Y^q$ .

The exactness of the bottom row follows from the cohomology long exact sequence associated to the short exact sequences obtained from the top row combined

with the Cartier isomorphisms for the first three vertical arrows, which tell us that the coboundary morphisms of these long exact sequences are all 0. Indeed, if we let  $\bar{\omega}_{(Z, \tilde{N})}^\bullet$  be the complex obtained by completing the inclusion of complexes

$$\omega_{(Z, \tilde{N})/k}^\bullet \otimes \mathcal{F}_1 \mathcal{F}_2 \rightarrow \omega_{(Z, \tilde{N})/k}^\bullet \otimes \mathcal{F}_1 \oplus \omega_{(Z, \tilde{N})/k}^\bullet \otimes \mathcal{F}_2$$

to a distinguished triangle, then we get a long exact sequence

$$\begin{aligned} \cdots \rightarrow \mathcal{H}^q(\omega_{(Z, \tilde{N})/k}^\bullet \otimes \mathcal{F}_1 \mathcal{F}_2) &\rightarrow \mathcal{H}^q(\omega_{(Z, \tilde{N})/k}^\bullet \otimes \mathcal{F}_1) \oplus \mathcal{H}^q(\omega_{(Z, \tilde{N})/k}^\bullet \otimes \mathcal{F}_2) \\ &\rightarrow \mathcal{H}^q(\bar{\omega}_{(Z, \tilde{N})}^\bullet) \rightarrow \cdots \end{aligned}$$

From the Cartier isomorphisms for  $\omega_{(Z, \tilde{N})/k}^\bullet \otimes \mathcal{F}_1 \mathcal{F}_2$  and  $\omega_{1,2}^\bullet$ , we deduce that

$$\mathcal{H}^q(\omega_{(Z, \tilde{N})/k}^\bullet \otimes \mathcal{F}_1 \mathcal{F}_2) \hookrightarrow \mathcal{H}^q(\omega_{(Z, \tilde{N})/k}^\bullet \otimes \mathcal{F}_1) \oplus \mathcal{H}^q(\omega_{(Z, \tilde{N})/k}^\bullet \otimes \mathcal{F}_2),$$

so the coboundaries of the long exact sequence are all 0. By continuing this argument, we deduce the exactness of the entire bottom row, and this proves that  $\tilde{C}^{-1}$  is an isomorphism.

Now we prove that  $\tilde{C}^{-1}$  is an isomorphism. We will show that  $\tilde{C}^{-1}$  is an isomorphism in degree  $q$  as well. From the short exact sequence (3B.1), we get the following commutative diagram with exact rows:

$$\begin{array}{ccccccc} 0 & \longrightarrow & C_Y^{q-1} & \longrightarrow & \tilde{C}_Y^q & \longrightarrow & C_Y^q \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \mathcal{H}^{q-1}(\text{Fr}_* C_Y^\bullet) & \longrightarrow & \mathcal{H}^q(\text{Fr}_* \tilde{C}_Y^\bullet) & \longrightarrow & \mathcal{H}^q(\text{Fr}_* C_Y^\bullet) \longrightarrow 0 \end{array}$$

To see that the bottom row is exact, we have to check that in the long exact cohomology sequence associated to the top row the coboundaries are all 0, which is equivalent to showing surjectivity of  $\mathcal{H}^q(\text{Fr}_* \tilde{C}_Y^\bullet) \rightarrow \mathcal{H}^q(\text{Fr}_* C_Y^\bullet)$ . However, by the top row and the Cartier isomorphism  $C^{-1}$ , the composite

$$\tilde{C}_Y^q \rightarrow C_Y^q \rightarrow \mathcal{H}^q(\text{Fr}_* C_Y^\bullet)$$

is surjective, so the desired map is surjective as well. Now we have a map of short exact sequences, where the left and right vertical maps are isomorphisms, so the middle one must be as well.  $\square$

We can define canonical projections  $\pi : W_{n+1} \tilde{\omega}_Y^\bullet \rightarrow W_n \tilde{\omega}_Y^\bullet$  using the Cartier isomorphisms. The construction works in the same way for  $W_n \tilde{\omega}_Y^\bullet$ . The definition of  $\pi$  for  $W_n \omega_Y^\bullet$  can be found in Section 1 of [Hyodo 1991] in the semistable case and in Section 4 of [Hyodo and Kato 1994] in general. The constructions in [Hyodo 1991] and in [Hyodo and Kato 1994] are the same, although they are formulated slightly differently. Our construction follows that in Section 1 of [Hyodo 1991],

by first defining a map  $p : W_n \tilde{\omega}_Y^\bullet \rightarrow W_{n+1} \tilde{\omega}_Y^\bullet$  and then showing that  $p$  is injective and its image coincides with the image of multiplication by  $p$  on  $W_{n+1} \tilde{\omega}_Y^\bullet$ . The projection  $\pi$  will then be the unique map which makes the following diagram commute:

$$\begin{array}{ccc}
 W_n \tilde{\omega}_Y^\bullet & \xleftarrow{\pi} & W_{n+1} \tilde{\omega}_Y^\bullet \\
 \downarrow p & & \searrow p \\
 & & W_{n+1} \tilde{\omega}_Y^\bullet
 \end{array}$$

The map  $p : W_n \tilde{\omega}_Y^i \rightarrow W_{n+1} \tilde{\omega}_Y^i$  is induced from  $p^{-i+1} \text{Fr}^* : \tilde{C}_Y^i \rightarrow \tilde{C}_Y^i$ , where  $\text{Fr} : (Z, \tilde{N}) \rightarrow (Z, \tilde{N})$  is a lifting of the Frobenius endomorphism of  $(Z, \tilde{N}) \times_W k$  such that  $\text{Fr}^*(W[\tau, \sigma]) \subset W[\tau, \sigma]$ . The injectivity of  $p$  and the fact that its image coincides with that of multiplication by  $p$  are deduced as in Section 2 of [Hyodo 1991] (or as in Lemma 6.8 of [Nakkajima 2005]) from the Cartier isomorphism and from the fact that  $\tilde{C}_Y^\bullet$  is  $W$ -torsion-free (when we take  $\tilde{C}_Y^\bullet$  to be the crystalline complex associated to an embedding system for  $(Y, \tilde{M})$  over  $W$ ).

Now we will consider a different interpretation of the monodromy operator  $N$ . Taking the cohomology sheaves of the short exact sequence

$$0 \rightarrow C_Y^\bullet[-1] \rightarrow \tilde{C}_Y^\bullet \rightarrow C_Y^\bullet \rightarrow 0,$$

we get a long exact sequence of sheaves on  $Y$

$$\dots \rightarrow W_n \omega_Y^{q-1} \rightarrow W_n \tilde{\omega}_Y^q \rightarrow W_n \omega_Y^q \rightarrow \dots$$

whose coboundaries are actually all 0. This can be checked as in Lemma 1.4.3 of [Hyodo 1991], since it suffices to see that the induced map on cocycles  $Z^q(\tilde{C}_Y) \rightarrow Z^q(C_Y)$  modulo  $p^n$  is surjective, and we can use the Cartier isomorphisms in Lemma 3.10 to give an explicit formula for cocycles modulo  $p^n$ . So we have a short exact sequence of sheaves on  $Y$

$$0 \rightarrow W_n \omega_Y^{q-1} \rightarrow W_n \tilde{\omega}_Y^q \rightarrow W_n \omega_Y^q \rightarrow 0, \tag{3B.4}$$

which is compatible with operators  $\pi, F, V$  and  $d$ . We have a morphism of distinguished triangles in the derived category  $D(Y_{\text{ét}}, W)$  of sheaves of  $W$ -modules on  $Y$ :

$$\begin{array}{ccccccc}
 C_Y^\bullet[-1] & \longrightarrow & \tilde{C}_Y^\bullet & \longrightarrow & C_Y^\bullet & \longrightarrow & C_Y^\bullet \\
 \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 W_n \omega_Y^\bullet[-1] & \longrightarrow & W_n \tilde{\omega}_Y^\bullet & \longrightarrow & W_n \omega_Y^\bullet & \longrightarrow & W_n \omega_Y^\bullet
 \end{array}$$

The left and right vertical maps are defined in the proof of Theorem 4.19 of [Hyodo and Kato 1994], and the middle one can be defined in exactly the same way. Note that the definition of the maps in Theorem 4.19 has a gap which is corrected in Lemma 7.18 of [Nakkajima 2005], namely, checking that they commute with the transition morphisms  $\pi : W_{n+1}\omega_Y^\bullet \rightarrow W_n\omega_Y^\bullet$ . The fact that the middle map commutes with the transition morphisms  $\pi : W_{n+1}\tilde{\omega}_Y^\bullet \rightarrow W_n\tilde{\omega}_Y^\bullet$  can be checked in the same way as in Lemma 7.18 of [Nakkajima 2005], using the corresponding Cartier isomorphism to check that the complexes  $W_n\tilde{\omega}_Y^\bullet$  give rise to formal de Rham–Witt complexes as in Definition 6.1 of [ibid.] and thus applying Corollary 6.28(8). We also need to check that  $\varprojlim W_n\tilde{\omega}_Y^1$  is torsion-free, but we can use the fact that this is known for  $\varprojlim W_n\omega_Y^1$  and the exact sequence (3B.4). The first and third vertical maps are quasi-isomorphisms by Theorem 4.19 of [Hyodo and Kato 1994], so we get an isomorphism of distinguished triangles. Thus, the exact sequence (3B.4) induces the monodromy operator  $N$  on cohomology.

Assume that  $Y$  has an admissible lifting  $\underline{Z}$  over  $(W[t, s], \mathbb{N}^2)$ , and set  $Z = \underline{Z} \otimes_W k$ . We consider a few more variations on the de Rham–Witt complex, which we will only define locally on  $Z$ . Let  $W_n\Omega_Z^\bullet$  be the de Rham–Witt complex of  $Z$ . Let

$$Y^1 = \text{Spec } k[X_1, \dots, X_n, Y_1, \dots, Y_n, Z_1, \dots, Z_m]/X_1 \cdots X_r$$

and

$$Y^2 = \text{Spec } k[X_1, \dots, X_n, Y_1, \dots, Y_n, Z_1, \dots, Z_m]/Y_1 \cdots Y_s.$$

Each  $Y^i$  is a normal crossings divisor in  $Z_{r,s,m} \times_W k$ . Let  $\mathcal{D}_n^i$  be the structure sheaf of the divided power envelope of  $Y^i$  in  $Z_{r,s,m}$  and  $\mathcal{F}\mathcal{D}_n^i = \ker(\mathcal{D}_n^i \rightarrow \mathcal{O}_{Y^i})$ . For  $i = 1, 2$ , let  $W_n\Omega_Z^\bullet(-\log Y^i)$  be the (pullback to  $Z$ ) of the “compact support” version of the de Rham–Witt complex of  $Z_{r,s,m}$  with respect to  $Y^i$ . This complex was introduced by Hyodo [1991, Section 1] and is defined by

$$W_n\Omega_{Z_{r,s,m}}^q(-\log Y^i) = \mathcal{H}^q(\Omega_{Z/W_n}^\bullet(\log Y^i) \otimes_{\mathcal{O}_{Z_{r,s,m}}} \mathcal{F}\mathcal{D}_n^i).$$

Let  $W_n\Omega_Z^\bullet(-\log Y^1 - \log Y^2)$  be the pullback from  $Z_{r,s}$  to  $Z$  of the complex defined by

$$W_n\Omega_{Z_{r,s,m}}^q(-\log Y^1 - \log Y^2) := \mathcal{H}^q(\omega_{Z_{r,s,m}, \mathbb{N}^r \oplus \mathbb{N}^s/W_n}^\bullet \otimes_{\mathcal{O}_Z} \mathcal{F}\mathcal{D}_1\mathcal{F}\mathcal{D}_2).$$

This third complex is meant to approximate a product of complexes of the form  $W_n\Omega_Z^\bullet(-\log Y)$ . When  $n = 1$ , consider  $Z^1 = \text{Spec } k[X_1, \dots, X_n, t]/(X_1 \cdots X_r - t)$ ,  $Z^2 = \text{Spec } k[Y_1, \dots, Y_n, u]/(Y_1 \cdots Y_s - u)$  and  $Z^3 = \text{Spec } k[Z_1, \dots, Z_m]$ . Then

$$\begin{aligned} W_1\Omega_{Z_{r,s,m}}^\bullet(-\log Y^1 - \log Y^2) \\ \simeq \Omega_{Z^1/k}^\bullet(-\log Y^1) \otimes_k \Omega_{Z^2/k}^\bullet(-\log Y^2) \otimes_k \Omega_{Z^3/k}^\bullet. \end{aligned} \tag{3B.5}$$

All these also are endowed with operators  $F, V$ , differential  $d$  and projection  $\pi$ , and they also satisfy a Cartier isomorphism.

Let  $\mathbb{R}$  be Raynaud’s ring, introduced in [Illusie and Raynaud 1983], i.e., the graded  $W$ -algebra generated by  $F, V$  in degree 0 and  $d$  in degree 1, subject to the usual relations

$$d^2 = 0, \quad FV = VF = p, \quad dF = pFd, \quad Vd = pdV \quad \text{and} \quad FdV = d.$$

Let  $\mathbb{R}_n$  be the right  $\mathbb{R}$ -module  $\mathbb{R}/(V^n\mathbb{R} + dV^n\mathbb{R})$ .

**Lemma 3.11.** *Let  $W_n\Omega^\bullet$  be one of the complexes  $W_n\Omega_Z^\bullet, W_n\Omega_Z^\bullet(-\log Y^i)$  for  $i = 1, 2$  or  $W_n\Omega_Z^\bullet(-\log Y^1 - \log Y^2)$ . Let*

$$W\Omega^\bullet = \varprojlim W_n\Omega^\bullet.$$

Then  $W\Omega^\bullet \otimes_{\mathbb{R}}^L \mathbb{R}_n = W_n\Omega^\bullet$ .

*Proof.* For  $n = 1$ , and  $W_n\Omega_Z^\bullet$  and  $W_n\Omega_Z^\bullet(-\log Y^i)$ , we have Cartier isomorphisms

$$W_1\Omega^i \xrightarrow{\sim} \mathcal{H}^i(F_*W_1\Omega^\bullet),$$

by [Deligne and Illusie 1987, Result 4.2.1.3]. For  $W_n\Omega_Z^\bullet(-\log Y^1 - \log Y^2)$  the Cartier isomorphism follows from the product formula (3B.5) and from the Cartier isomorphisms above. Let  $\mathcal{X}_n = \underline{Z} \times_W W_n$ . By abuse of notation, we write  $\Omega_{\mathcal{X}_n}^\bullet$  for the complex of sheaves of  $W_n$ -modules such that

$$W_n\Omega^i = \mathcal{H}^i(\Omega_{\mathcal{X}_n}^\bullet).$$

In fact, we have complexes  $\Omega_Z^\bullet, \Omega_Z^\bullet(-\log Y^i)$  or  $\Omega_Z^\bullet(-\log Y^1 - \log Y^2)$  which give the corresponding complexes  $\Omega_{\mathcal{X}_n}^\bullet, \Omega_{\mathcal{X}_n}^\bullet(-\log Y^i)$  or  $\Omega_{\mathcal{X}_n}^\bullet(-\log Y^1 - \log Y^2)$  when reduced modulo  $p^n$ . We also denote any of the initial complexes over  $W$  as  $\Omega_Z^\bullet$ . Then there is an explicit description of cocycles modulo  $p^n$  given by

$$d^{-1}(p^n\Omega_Z^{i+1}) = \sum_{k=0}^n p^k f^{n-k} \Omega_Z^i + \sum_{k=0}^{n-1} f^k d\Omega_Z^{i-1},$$

where  $f : \Omega_Z^i \rightarrow \Omega_Z^i$  is defined by  $f = \text{Fr}/p^i$ . This is the same as Formula A from Editorial Comment 11 in [Hyodo 1991] and is proven in the same way as in that paper and in the same way as in the classical crystalline cohomology case (see [Illusie 1979, 0.2.3.13]).

As in the case of  $W_n\omega_Y, W_\bullet\Omega^\bullet$  (and  $W\Omega^\bullet$ ) is endowed with a differential  $d$ , operators  $F, V$  satisfying the usual relations and a canonical projection  $\pi_n : W_{n+1}\Omega^\bullet \rightarrow W_n\Omega^\bullet$  such that  $p \circ \pi_n$  coincides with multiplication by  $p$  on  $W_{n+1}\Omega^\bullet$ .

We claim that the lemma follows from the Cartier isomorphism, from the description of cocycles modulo  $p^n$  in  $\Omega_Z^\bullet$  and from the formal properties of  $W_n\Omega^\bullet$ . The

proof is the same as for Lemma 1.3.3 of [Mokrane 1993]. We outline the argument in order to show that it applies to our case as well. To prove the desired result, we use the flat resolution of  $\mathbb{R}_n$  as an  $\mathbb{R}$ -module given by

$$0 \rightarrow \mathbb{R} \xrightarrow{(F^n, -F^n d)} \mathbb{R} \oplus \mathbb{R} \xrightarrow{dV^n + V^n} \mathbb{R} \rightarrow \mathbb{R}_n \rightarrow 0,$$

and it suffices by Corollary 1.3.3 of [Illusie and Raynaud 1983] to prove that the sequence

$$0 \rightarrow W\Omega^{i-1} \xrightarrow{(F^n, -F^n d)} W\Omega^{i-1} \oplus W\Omega^i \xrightarrow{dV^n + V^n} W\Omega^i \rightarrow W_n\Omega^i \rightarrow 0$$

is exact. The last map is the canonical projection  $\pi : W\Omega^i \rightarrow W_n\Omega^i$ .

Exactness at the first term follows from the fact that multiplication by  $p$  (and hence also  $F$ ) is injective on  $W\Omega^\bullet$ . Indeed, multiplication by  $p$  on  $W_n\Omega^\bullet$  factors as  $p \circ \pi_n$  and  $p$  is injective by definition, so if  $p(x_n) = 0$  for all  $n$  then  $\pi_n(x_n) = x_{n-1} = 0$  for all  $n$ , so  $x = (x_n) = 0$ .

Exactness at the last term is the statement that  $\pi$  is surjective, which follows by construction, since  $p = p \circ \pi$ ,  $p$  is injective and the image of  $p : W_n\Omega^\bullet \rightarrow W_{n+1}\Omega^\bullet$  coincides with the image of multiplication by  $p$ .

Now we check that  $\ker \pi = dV^n W\Omega^\bullet + V^n W\Omega^\bullet$ . Recall that  $\pi_n : W_{n+1} \rightarrow W_n$  is the canonical projection. It is enough to show that  $\ker \pi_n = dV^n W_1\Omega^\bullet + V^n W_1\Omega^\bullet$ . First, if  $x = V^n a + dV^n b \in W_{n+1}\Omega$ , it suffices to check that  $px = 0$  and indeed  $px = FV^{n+1}a + dFV^{n+1}b = 0$ . Now, let  $[x]_{n+1} \in \ker \pi_n$ , where  $x$  is an element of  $\Omega_{\mathbb{Z}}^\bullet$  modulo  $p^{n+1}$ . Then  $[px]_{n+1} = p[x]_{n+1} = 0$ , so it must be the case that  $px = p^{n+1}a + db$ . We get  $db = 0 \pmod p$ , so by the description of cocycles mod  $p$  we have  $b = pb' + Fb'' + db''$ , so that  $db = pdb' + pFdb''$ . Thus,

$$\begin{aligned} [x]_{n+1} &= [p^n a]_{n+1} + [db']_{n+1} + [Fdb'']_{n+1} \\ &= V^n [a]_{n+1} + d[p^n Fb'']_{n+1} = V^n [a] + dV^n [Fb'']. \end{aligned}$$

Now we check exactness at the second term. First, note that the sequence

$$W_{2n}\Omega^{q-1} \xrightarrow{F^n} W_n\Omega^{q-1} \xrightarrow{d} W_n\Omega^q$$

is exact, which is proved in the same way as Lemma 1.3.4 of [Mokrane 1993], by taking the long exact sequence of cohomology sheaves of the short exact sequence

$$0 \rightarrow \Omega_{\mathbb{Z}}^\bullet / p^n \Omega_{\mathbb{Z}}^\bullet \xrightarrow{p^n} \Omega_{\mathbb{Z}}^\bullet / p^{2n} \Omega_{\mathbb{Z}}^\bullet \rightarrow \Omega_{\mathbb{Z}}^\bullet / p^n \Omega_{\mathbb{Z}}^\bullet \rightarrow 0.$$

We note that the proof of the analogous statement in the classical case in [Illusie 1979, I(3.21)] is wrong and corrected in [Illusie and Raynaud 1983, II(1.3)]. Nakkajima [2005, 6.28(6)] proves this statement for formal de Rham–Witt complexes, using the same argument as Lemma 1.3.4 of [Mokrane 1993].

We now claim that the projection

$$W\Omega^\bullet/p^n W\Omega^\bullet \rightarrow W_n\Omega^\bullet$$

is a quasi-isomorphism. This implies that

$$d^{-1}(p^n W\Omega^q) = F^n W\Omega^{q-1},$$

so if  $dV^n x + V^n y = 0$ , then  $dx + p^n y = 0$ , which in turn implies  $x = F^n z$  and  $y = -F^n dz$  for some  $z \in W\Omega^{q-1}$ . This checks exactness at the second term. Moreover, the fact that

$$W\Omega^\bullet/p^n W\Omega^\bullet \rightarrow W_n\Omega^\bullet$$

is a quasi-isomorphism follows in the same way as Corollary 3.17 of [Illusie 1979], boiling down to the Cartier isomorphism and to the description of  $\ker \pi$  as  $dV^n + V^n$ .  $\square$

**Remark 3.12.** One can use the Cartier isomorphisms to check Properties 6.0.1–6.0.5 of [Nakkajima 2005] for  $\Omega_{\mathbb{Z}}^\bullet, \Omega_{\mathbb{Z}}^\bullet(-\log Y^i)$  and  $\Omega_{\mathbb{Z}}^\bullet(-\log Y^1 - \log Y^2)$ , thus proving the analogue of Proposition 6.27 there for all three complexes. Then Theorem 6.24 of [Nakkajima 2005] also implies Lemma 3.11.

**3C. The weight filtration.** The goal of this section is to define a double filtration  $P_{k,i}$  on  $W\tilde{\omega}_Y^\bullet$ , which will be an analogue of the weight filtration defined by Mokrane on  $W_n\tilde{\omega}_Y^\bullet$  in the semistable case (see [1993, Section 3]).

Let  $(Z, \tilde{N})$  be an admissible lifting of  $(Y, \tilde{M})$  over  $(W[\tau, \sigma], \mathbb{N}^2)$ . We know that such liftings exist étale locally. Let  $\mathcal{X}_n = Z \times_W W_n$ . Let  $\tilde{N}_1$  be the log structure on  $Z$  (or  $\mathcal{X}_n$ ) obtained by pulling back the log structure on  $Z_{r,s,m}$  associated to

$$\mathbb{N}^r \rightarrow W[X_1, \dots, X_n, Y_1, \dots, Y_n, Z_1, \dots, Z_m],$$

$$(0, \dots, 0, 1, 0, \dots, 0) \mapsto X_i,$$

where 1 is in the  $i$ -th position. Define  $\tilde{N}_2$  analogously. The pullback of  $\tilde{N}_i$  to  $Y$  is the same as  $\tilde{M}_i$ . For  $i = 1, 2$ , we have maps of sheaves of monoids  $\tilde{N}_i \rightarrow \tilde{N}$ .

We define the following filtration on  $\omega_{(\mathcal{X}_n, \tilde{N})/(W_n, \text{triv})}^\bullet$ :

$$P_{i,j}\omega_{(\mathcal{X}_n, \tilde{N})/(W_n, \text{triv})}^q := \text{Im}\left(\omega_{(\mathcal{X}_n, \tilde{N}_1)/(W_n, \text{triv})}^i \otimes \omega_{(\mathcal{X}_n, \tilde{N}_2)/(W_n, \text{triv})}^j \otimes \Omega_{\mathcal{X}_n/k}^{q-i-j} \rightarrow \omega_{(\mathcal{X}_n, \tilde{N})/(W_n, \text{triv})}^q\right)$$

for  $i, j \geq 0$  and  $i + j \leq q$ . This filtration respects the differential and induces a filtration  $P_{i,j}\tilde{\mathcal{C}}_Y^\bullet$  on  $\tilde{\mathcal{C}}_Y^\bullet$  (which can be thought of as a quotient of  $\omega_{(\mathcal{X}_n, \tilde{M})/(W_n, \text{triv})}^\bullet$ , as in the proof of Lemma 3.10). Note that if we let

$$P_k\omega_{(\mathcal{X}_n, \tilde{N})/(W_n, \text{triv})}^q = \text{Im}\left(\omega_{(\mathcal{X}_n, \tilde{N})/(W_n, \text{triv})}^k \otimes \Omega_{\mathcal{X}_n/k}^{q-k} \rightarrow \omega_{(\mathcal{X}_n, \tilde{N})/(W_n, \text{triv})}^q\right),$$

then  $P_k$  is the weight filtration defined in [Mokrane 1993, 1.1.1], and

$$P_{i,j} \omega^{\bullet}_{(\mathcal{E}_n, \tilde{N})/(W_n, \text{triv})} \subset P_{i+j} \omega^{\bullet}_{(\mathcal{E}_n, \tilde{N})/(W_n, \text{triv})}.$$

For  $i = 1, \dots, r$ , let  $D_{1,i}$  be the pullback to  $Z$  of the divisor of  $Z_{r,s,m}$  obtained by setting  $X_i = 0$ . Similarly, for  $i = 1, \dots, s$ , let  $D_{2,i}$  be the pullback to  $Z$  of the divisor of  $Z_{r,s,m}$  obtained by setting  $Y_i = 0$ . For  $i, j \geq 0$  let  $D^{(i,j)}$  be the disjoint union of

$$D_{1,k_1} \times_Z \dots \times_Z D_{1,k_i} \times_Z D_{2,l_1} \times_Z \dots \times_Z D_{2,l_j},$$

over all  $k_1, \dots, k_i \in \{1, \dots, r\}$  and  $l_1, \dots, l_j \in \{1, \dots, s\}$ . And let  $\tau_{i,j} : D^{(i,j)} \rightarrow Z$  be the obvious morphism, with  $\mathcal{D}_n^{(i,j)}$ ,  $\tau_{i,j}$  the pullbacks to  $\mathcal{E}_n$ . Let

$$\begin{aligned} & \text{Gr}_{i,j} \omega^q_{(\mathcal{E}_n, \tilde{N})/(W_n, \text{triv})} \\ & := P_{i,j} \omega^q_{(\mathcal{E}_n, \tilde{N})/(W_n, \text{triv})} / (P_{i-1,j} \omega^q_{(\mathcal{E}_n, \tilde{N})/(W_n, \text{triv})} + P_{i,j-1} \omega^q_{(\mathcal{E}_n, \tilde{N})/(W_n, \text{triv})}). \end{aligned}$$

For  $i, j \geq 1$ , we will define a morphism of sheaves

$$\text{Res} : \text{Gr}_{i,j} \omega^q_{(\mathcal{E}_n, \tilde{N})/(W_n, \text{triv})} \rightarrow (\tau_{i,j})_* \Omega_{\mathcal{D}_n^{(i,j)}/W_n}^{q-i-j},$$

which extends to a morphism of complexes. If

$$\omega = \alpha \wedge \frac{dX_{k_1}}{X_{k_1}} \wedge \dots \wedge \frac{dX_{k_i}}{X_{k_i}} \wedge \frac{dY_{l_1}}{Y_{l_1}} \wedge \dots \wedge \frac{dY_{l_j}}{Y_{l_j}}$$

is a local section of  $P_{i,j} \omega^q_{(\mathcal{E}_n, \tilde{N})/(W_n, \text{triv})}$  with  $k_1 < \dots < k_i$  and  $l_1 < \dots < l_j$ , then

$$\text{Res}(\omega) := \alpha|_{D_{1,k_1} \times_Z \dots \times_Z D_{1,k_i} \times_Z D_{2,l_1} \times_Z \dots \times_Z D_{2,l_j}}.$$

This factors through  $P_{i-1,j} + P_{i,j-1}$  and extends to a global map of sheaves.

Alternatively, we can follow the construction in Section 3 of Chapter II of [Deligne 1970]. Let  $\mathcal{D}_n^k$  be the disjoint union of intersections of  $k$  divisors  $D_{j,k_i}$  with  $j = 1, 2$  and  $k_i \in \{1, \dots, n\}$ . These intersections are in one-to-one correspondence with images of injections

$$f : \{1, \dots, k\} \rightarrow \{1, \dots, n\} \cup \{1, \dots, n\},$$

and so we denote one of these  $k$  intersections by  $\mathcal{D}_n^f$  (even though it only really depends on  $\text{Im } f$ ). We have

$$\mathcal{D}_n^k = \bigsqcup_{\substack{i+j=k \\ i,j \geq 0}} \mathcal{D}_n^{i,j} = \bigsqcup_{\text{Im } f} \mathcal{D}_n^f.$$

Let  $\tau_f : \mathcal{D}_n^f \rightarrow \mathcal{X}_n$  be the closed immersion. In [Deligne 1970, 3.5.2], a morphism

$$\rho_1 : (\tau_f)_* \Omega_{\mathcal{D}_n^f}^{q-k} \rightarrow P_k \omega_{(\mathcal{X}_n, \tilde{N})/(W_n, \text{triv})}^q / P_{k-1}$$

(and then a morphism  $\rho_2$ , which depends on an ordering of  $\{1, \dots, n\} \cup \{1, \dots, n\}$ ) is associated to each such injection, and the sum of  $\rho_2$  over all injections  $f$  determines an isomorphism

$$\rho : (\tau_k)_* \Omega_{\mathcal{D}_n^k/W_n}^\bullet[-k] \xrightarrow{\simeq} P_k \omega_{(\mathcal{X}_n, \tilde{N})/(W_n, \text{triv})}^q / P_{k-1}$$

by Proposition 3.6 of Chapter II of [Deligne 1970].

We are only interested in injections  $q_{i,j} : \{1, \dots, i+j\} \rightarrow \{1, \dots, n\} \cup \{1, \dots, n\}$  with image of cardinality  $i$  in the first  $\{1, \dots, n\}$  term and cardinality  $j$  in the second  $\{1, \dots, n\}$  term. We let  $\text{Res}^{-1}$  be the sum of the morphisms  $\rho_2$  over all injections  $q_{i,j}$ . When we have an injection of type  $q_{i,j}$ , the image of the morphism  $\rho_2$  defined by Deligne falls in

$$P_{i,j} \omega_{(\mathcal{X}_n, \tilde{N})/(W_n, \text{triv})}^q / (P_{i-1,j} + P_{i,j-1}) \subset P_{i+j} \omega_{(\mathcal{X}_n, \tilde{N})/(W_n, \text{triv})}^q / P_{i+j-1}.$$

For  $k \geq 1$ , we have the direct sum decompositions

$$\begin{aligned} P_k \omega_{(\mathcal{X}_n, \tilde{N})/(W_n, \text{triv})}^\bullet / P_{k-1} &= \bigoplus_{\substack{i+j=k \\ i,j \geq 0}} \text{Gr}_{i,j} \omega_{(\mathcal{X}_n, \tilde{N})/(W_n, \text{triv})}^\bullet, \\ (\tau_k)_* \Omega_{\mathcal{D}_n^{(k)}/W_n}^{q-k} &= \bigoplus_{\substack{i+j=k \\ i,j \geq 0}} (\tau_{i,j})_* \Omega_{\mathcal{D}_n^{(i,j)}/W_n}^{q-i-j}. \end{aligned}$$

It is easy to check that the isomorphism  $\rho$  matches up the  $(i, j)$  terms in each decomposition. Putting this discussion together, we get the following:

**Lemma 3.13.** *For  $i, j \geq 1$ , the map*

$$\text{Res}^{-1} : (\tau_{i,j})_* \Omega_{\mathcal{D}_n^{(i,j)}/W_n}^{q-i-j} \rightarrow \text{Gr}_{i,j} \omega_{(\mathcal{X}_n, \tilde{N})/(W_n, \text{triv})}^q$$

*is an isomorphism.*

We also have the following analogue of Lemma 1.2 of [Mokrane 1993].

**Lemma 3.14.** *We have an exact sequence of complexes*

$$\begin{aligned} 0 \rightarrow P_{i-1,j-1} \omega_{(\mathcal{X}_n, \tilde{N})/(W_n, \text{triv})}^\bullet \\ \rightarrow P_{i-1,j} \omega_{(\mathcal{X}_n, \tilde{N})/(W_n, \text{triv})}^\bullet \oplus P_{i,j-1} \omega_{(\mathcal{X}_n, \tilde{N})/(W_n, \text{triv})}^\bullet \\ \rightarrow P_{i,j} \omega_{(\mathcal{X}_n, \tilde{N})/(W_n, \text{triv})}^\bullet \rightarrow \text{Gr}_{i,j} \omega_{(\mathcal{X}_n, \tilde{N})/(W_n, \text{triv})}^\bullet \rightarrow 0. \end{aligned}$$

The long exact cohomology sequence(s) associated to this have all coboundaries 0, so we get the exact sequence

$$\begin{aligned} 0 \rightarrow \mathcal{H}^q(P_{i-1, j-1} \omega_{(\mathcal{X}_n, \tilde{N})/(W_n, \text{triv})}^\bullet) \\ \rightarrow \mathcal{H}^q(P_{i-1, j} \omega_{(\mathcal{X}_n, \tilde{N})/(W_n, \text{triv})}^\bullet) \oplus \mathcal{H}^q(P_{i, j-1} \omega_{(\mathcal{X}_n, \tilde{N})/(W_n, \text{triv})}^\bullet) \\ \rightarrow \mathcal{H}^q(P_{i, j} \omega_{(\mathcal{X}_n, \tilde{N})/(W_n, \text{triv})}^\bullet) \rightarrow \mathcal{H}^q(\Omega_{\mathbb{P}_n^{(i, j)}/W_n}^\bullet[-i-j]) \rightarrow 0. \end{aligned}$$

*Proof.* The first assertion is clear. In order to show that the second sequence is exact, it suffices to show the following two statements about cocycles:

- (1)  $Z P_{i, j} \omega_{(\mathcal{X}_n, \tilde{N})/(W_n, \text{triv})}^q \twoheadrightarrow Z \Omega_{\mathbb{P}_n^{(i, j)}/W_n}^{q-i-j}$ .
- (2)  $Z P_{i-1, j} \omega_{(\mathcal{X}_n, \tilde{N})/(W_n, \text{triv})}^q \oplus Z P_{j, i-1} \omega_{(\mathcal{X}_n, \tilde{N})/(W_n, \text{triv})}^q$   
 $\twoheadrightarrow Z(P_{i-1, j} \omega_{(\mathcal{X}_n, \tilde{N})/(W_n, \text{triv})}^q + P_{i, j-1} \omega_{(\mathcal{X}_n, \tilde{N})/(W_n, \text{triv})}^q)$ .

The first statement is proved in the same way as the main step in Lemma 1.1.2 of [Mokrane 1993]. If  $\alpha$  is a local section of  $Z \Omega_{\mathbb{P}_n^{(i, j)}/W_n}^{q-i-j}$ , assume that  $\alpha$  is supported on some

$$D_{1, k_1} \times_Z \cdots \times_Z D_{1, k_i} \times_Z D_{2, l_1} \times_Z \cdots \times_Z D_{2, l_j},$$

for some  $k_1, \dots, k_i, l_1, \dots, l_j \in \{1, \dots, n\}$ . Let

$$\rho: \mathcal{X}_n \rightarrow D_{1, k_1} \times_Z \cdots \times_Z D_{1, k_i} \times_Z D_{2, l_1} \times_Z \cdots \times_Z D_{2, l_j}$$

be the retraction associated to the immersion

$$D_{1, k_1} \times_Z \cdots \times_Z D_{1, k_i} \times_Z D_{2, l_1} \times_Z \cdots \times_Z D_{2, l_j} \rightarrow \mathcal{X}_n.$$

Then  $\rho^* \alpha$  lifts  $\alpha$  to a section of  $Z \Omega_{\mathcal{X}_n/W_n}^{q-i-j}$  and the section

$$\omega_\alpha = \rho^* \alpha \wedge \frac{dX_{k_1}}{X_{k_1}} \wedge \cdots \wedge \frac{dX_{k_i}}{X_{k_i}} \wedge \frac{dY_{l_1}}{Y_{l_1}} \wedge \cdots \wedge \frac{dY_{l_j}}{Y_{l_j}} \in P_{i, j} \omega_{(\mathcal{X}_n, \tilde{N})/(W_n, \text{triv})}^q$$

satisfies  $d\omega = 0$  and  $\text{Res}(\omega) = \alpha$ . From this, we know that the coboundaries of the long exact sequence associated to

$$\begin{aligned} 0 \rightarrow P_{i-1, j} \omega_{(\mathcal{X}_n, \tilde{N})/(W_n, \text{triv})}^\bullet + P_{i, j-1} \omega_{(\mathcal{X}_n, \tilde{N})/(W_n, \text{triv})}^\bullet \\ \rightarrow P_{i, j} \omega_{(\mathcal{X}_n, \tilde{N})/(W_n, \text{triv})}^\bullet \rightarrow \text{Gr}_{i, j} \omega_{(\mathcal{X}_n, \tilde{N})/(W_n, \text{triv})}^\bullet \rightarrow 0 \end{aligned}$$

are 0, so we also know that

$$\mathcal{H}^q(P_{i-1, j} \omega_{(\mathcal{X}_n, \tilde{N})/(W_n, \text{triv})}^\bullet + P_{i, j-1} \omega_{(\mathcal{X}_n, \tilde{N})/(W_n, \text{triv})}^\bullet) \hookrightarrow \mathcal{H}^q(P_{i, j} \omega_{(\mathcal{X}_n, \tilde{N})/(W_n, \text{triv})}^\bullet)$$

for every  $i, j \geq 1$ .

For the second statement, we have to prove that if  $\alpha \in P_{i-1,j}\omega_{(\mathcal{X}_n,\tilde{N})/(W_n,\text{triv})}^q$  and  $\beta \in P_{i,j-1}\omega_{(\mathcal{X}_n,\tilde{N})/(W_n,\text{triv})}^q$  satisfy  $d(\alpha + \beta) = 0$ , then we can find  $\alpha' \in ZP_{i-1,j}\omega_{(\mathcal{X}_n,\tilde{N})/(W_n,\text{triv})}^q$  and  $\beta' \in ZP_{i,j-1}\omega_{(\mathcal{X}_n,\tilde{N})/(W_n,\text{triv})}^q$  such that  $\alpha' + \beta' = \alpha + \beta$ . If  $\alpha \in P_{i-1,j-1}\omega_{(\mathcal{X}_n,\tilde{N})/(W_n,\text{triv})}^q$ , then we are done, since we can just take  $\alpha' = 0, \beta' = \alpha + \beta$ . The same holds for  $\beta$ . Otherwise, we have  $d\alpha \in P_{i-1,j-1}$ , so by the injectivity proved in statement (1) for  $(i-1, j)$ , we know that  $d\alpha = d\alpha_1 + d\alpha_2$  for some  $\alpha_1 \in P_{i-1,j-1}$  and  $\alpha_2 \in P_{i-2,j}$ . Thus, we've reduced our problem from  $(i-1, j)$  to  $(i-2, j)$ . Proceeding by induction, we may assume that  $i = 0$ . In that case  $d\alpha_{2i} \in P_{0,j-1}$ . By (the same argument as in the proof of) Lemma 1.1.2 of [Mokrane 1993], we have an injection

$$\mathcal{H}^q(P_{0,j-1}\omega_{(\mathcal{X}_n,\tilde{N})/(W_n,\text{triv})}^\bullet) \hookrightarrow \mathcal{H}^q(P_{0,j}\omega_{(\mathcal{X}_n,\tilde{N})/(W_n,\text{triv})}^\bullet),$$

which implies  $d\alpha_{2i} = d\alpha_{2i+1}$  for some  $\alpha_{2i+1} \in P_{0,j-1}$ . Then

$$\alpha' := \alpha - \sum_{i'=0}^i \alpha_{2i'+1} \in ZP_{i-1,j}, \quad \beta' := \beta + \sum_{i'=0}^i \alpha_{2i'+1} \in ZP_{i,j-1}$$

satisfy the desired relations. □

The double filtration  $P_{i,j}$  on  $\omega_{(\mathcal{X}_n,\tilde{N})/(W_n,\text{triv})}^\bullet$  induces a double filtration  $P_{i,j}$  on  $\tilde{\mathcal{C}}_{\mathcal{X}_n}$ , and, for  $i, j \geq 1$ , the residue morphism  $\text{Res} : P_{i,j}\omega_{(\mathcal{X}_n,\tilde{N})/(W_n,\text{triv})}^q \rightarrow \Omega_{\mathbb{Q}_n^{(i,j)}/W_n}^{q-i-j}$  factors through  $P_{i,j}\tilde{\mathcal{C}}_{\mathcal{X}_n}$ .

**Lemma 3.15.** *For any two admissible liftings  $(Z_1, \tilde{N})$  and  $(Z_2, \tilde{N})$  of  $(Y, \tilde{M})$ , we have a canonical isomorphism*

$$\alpha_{Z_1 Z_2} : \mathcal{H}^q(P_{i,j}\tilde{\mathcal{C}}_{\mathcal{X}_{1,n}}^\bullet) \rightarrow \mathcal{H}^q(P_{i,j}\tilde{\mathcal{C}}_{\mathcal{X}_{2,n}}^\bullet)$$

satisfying the cocycle condition for any three admissible liftings.

Moreover, the residue morphism

$$\text{Res}_Z : \mathcal{H}^q(P_{i,j}\tilde{\mathcal{C}}_{\mathcal{X}_n}^\bullet) \rightarrow \mathcal{H}^{q-i-j}(\Omega_{\mathbb{Q}_n^{(i,j)}/W_n}^\bullet) \simeq W_n \Omega_Y^q_{(i,j)}$$

induced on cohomology satisfies the compatibility

$$\text{Res}_{Z_1} = \text{Res}_{Z_2} \circ \alpha_{Z_1 Z_2}.$$

*Proof.* The proof of the first part is basically the same as the proof of Lemma 3.5. We take admissible lifts  $(Z_1, \tilde{N})$  and  $(Z_2, \tilde{N})$  (we denote the log structures on both simply by  $\tilde{N}$ , as it will be understood from the context which is the underlying

scheme). As in the proof of Lemma 3.5, we form  $(Z_{12}, \tilde{N})$ , which is smooth over  $(Z_i, \tilde{N})$ , even though it is not quite an admissible lift. However,  $Z_{12}$  is étale over  $\text{Spec } W[X_1, \dots, X_n, Y_1, \dots, Y_n, X'_1, \dots, X'_n, Y'_1, \dots, Y'_n, v_1^{\pm 1}, \dots, v_r^{\pm 1}, u_1^{\pm 1}, \dots, u_s^{\pm 1}] / (X_i v_i - X'_i Y_j v_j - Y'_j)$ .

So we can endow  $\tilde{C}_{\mathcal{E}_{12,n}}^\bullet$  with a filtration  $P_{i,j} \tilde{C}_{\mathcal{E}_{12,n}}^\bullet$  defined as above, in terms of log structures  $\tilde{N}_1$  and  $\tilde{N}_2$  (which come from formally “inverting” the  $X_i$  and  $X'_i$  or the  $Y_i$  and  $Y'_i$ ). Then the same argument used in the proof of Lemma 3.5 gives us quasi-isomorphisms

$$P_{i,j} \tilde{C}_{\mathcal{E}_{i,n}} \rightarrow P_{i,j} \tilde{C}_{\mathcal{E}_{12,n}}$$

for  $i = 1, 2$ , which satisfy the right compatibility condition for three admissible lifts.

For the second part, we follow the argument in Lemma 3.4(2) of [Mokrane 1993]. We let

$$\omega = \alpha \wedge \frac{dX_{k_1}}{X_{k_1}} \wedge \dots \wedge \frac{dX_{k_i}}{X_{k_i}} \wedge \frac{dY_{l_1}}{Y_{l_1}} \wedge \dots \wedge \frac{dY_{l_i}}{Y_{l_i}}$$

be a section of  $P_{i,j} \omega_{(\mathcal{E}_{1,n}, \tilde{N}) / (W_n, \text{triv})}^q$  and

$$\omega' = \alpha' \wedge \frac{dX'_{k_1}}{X'_{k_1}} \wedge \dots \wedge \frac{dX'_{k_i}}{X'_{k_i}} \wedge \frac{dY'_{l_1}}{Y'_{l_1}} \wedge \dots \wedge \frac{dY'_{l_i}}{Y'_{l_i}}$$

be a section of  $P_{i,j} \omega_{(\mathcal{E}_{2,n}, \tilde{N}) / (W_n, \text{triv})}^q$  such that  $\omega = \omega'$  in  $P_{i,j} \omega_{(\mathcal{E}_{12,n}, \tilde{N}) / (W_n, \text{triv})}^q$ .

We have to check that  $\alpha|_{\mathcal{D}_{12,n}^{(i,j)}} = \alpha'|_{\mathcal{D}_{12,n}^{(i,j)}}$ . But

$$\omega - \omega' = (\alpha - \alpha') \wedge \frac{dX_{k_1}}{X_{k_1}} \wedge \dots \wedge \frac{dX_{k_i}}{X_{k_i}} \wedge \frac{dY_{l_1}}{Y_{l_1}} \wedge \dots \wedge \frac{dY_{l_i}}{Y_{l_i}} + \Psi,$$

where  $\Psi \in P_{i,j-1} \omega_{(\mathcal{E}_{2,n}, \tilde{N}) / (W_n, \text{triv})}^q + P_{i-1,j} \omega_{(\mathcal{E}_{2,n}, \tilde{N}) / (W_n, \text{triv})}^q$ . This means that

$$(\alpha - \alpha') \wedge \frac{dX_{k_1}}{X_{k_1}} \wedge \dots \wedge \frac{dX_{k_i}}{X_{k_i}} \wedge \frac{dY_{l_1}}{Y_{l_1}} \wedge \dots \wedge \frac{dY_{l_i}}{Y_{l_i}}$$

is also a section of  $P_{i,j-1} \omega_{(\mathcal{E}_{2,n}, \tilde{N}) / (W_n, \text{triv})}^q + P_{i-1,j} \omega_{(\mathcal{E}_{2,n}, \tilde{N}) / (W_n, \text{triv})}^q$ , and so  $(\alpha - \alpha')|_{\mathcal{D}_{12,n}^{(i,j)}} = 0$ . □

**Corollary 3.16.** *We can define the sheaves*

$$P_{i,j} W_n \tilde{\omega}_Y^q := \mathcal{H}^q(P_{i,j} \tilde{C}_Y^\bullet).$$

The complexes  $P_{i,j}W_n\tilde{\omega}_Y^\bullet$  form an increasing double filtration of  $W_n\tilde{\omega}_Y^\bullet$  such that the graded pieces for  $i, j \geq 1$

$$\mathrm{Gr}_{i,j}W_n\tilde{\omega}_Y^\bullet := P_{i,j}W_n\tilde{\omega}_Y^\bullet / P_{i,j-1} + P_{i-1,j}$$

are canonically isomorphic to the de Rham–Witt complexes of the smooth subschemes  $Y^{(i,j)}$ :

$$\mathrm{Res} : \mathrm{Gr}_{i,j}W_n\tilde{\omega}_Y^\bullet \xrightarrow{\sim} W_n\Omega_{Y^{(i,j)}}^\bullet[-i-j](-i-j).$$

**Lemma 3.17.** *The constructions in Section 3C are compatible with the transition morphisms  $\pi$ , in the following way:*

(1) *The following diagrams are commutative:*

$$\begin{array}{ccc} W_{n+1}\tilde{\omega}_Y^q & \xrightarrow{\pi} & W_n\tilde{\omega}_Y^q \\ \wedge \frac{d\tau}{\tau} \downarrow & & \downarrow \wedge \frac{d\tau}{\tau} \\ W_{n+1}\tilde{\omega}_Y^q & \xrightarrow{\pi} & W_n\tilde{\omega}_Y^q \end{array} \quad \text{and} \quad \begin{array}{ccc} W_{n+1}\tilde{\omega}_Y^q & \xrightarrow{\pi} & W_n\tilde{\omega}_Y^q \\ \wedge \frac{d\sigma}{\sigma} \downarrow & & \downarrow \wedge \frac{d\sigma}{\sigma} \\ W_{n+1}\tilde{\omega}_Y^q & \xrightarrow{\pi} & W_n\tilde{\omega}_Y^q \end{array}$$

(2) *The projection  $\pi : W_{n+1}\tilde{\omega}_Y^q \rightarrow W_n\tilde{\omega}_Y^q$  preserves the weight filtration  $P_{i,j}$  on  $W_m\tilde{\omega}_Y^q$  for  $m = n, n + 1$ .*

(3) *The morphism  $\pi : P_{i,j}W_{n+1}\tilde{\omega}_Y^q \rightarrow P_{i,j}W_n\tilde{\omega}_Y^q$  is surjective.*

*Proof.* The first part follows in the same way as Proposition 8.1 of [Nakkajima 2005], by using a local admissible lifting  $(Z, \tilde{N})$  of  $(Y, \tilde{M})$  together with a lift of the Frobenius  $\Phi$ . Then  $\Phi^*(\tau) = \tau^p(1 + pu)$  for some

$$u \in \mathbb{O}_Z \otimes_{W[\tau,\sigma]} W_n\langle \tau, \sigma \rangle$$

and so  $\Phi^*(d \log \tau)$  is equivalent to  $pd \log \tau$  modulo an exact form. The same holds for  $\sigma$ .

The second part follows in the same way as Proposition 8.4 of [Nakkajima 2005]. The question is local, so we may assume that the admissible lift  $(Z, \tilde{N})$  is étale over  $\mathrm{Spec} W[X_1, \dots, X_n, Y_1, \dots, Y_n], \mathbb{N}^r \oplus \mathbb{N}^s$ . First we see that, for a lift  $\Phi$  of Frobenius we have that  $\Phi^*(d \log X_i)$  is equivalent modulo an exact form to  $pd \log X_i$  for  $1 \leq i \leq r$  and that  $\Phi^*(d \log Y_j)$  is equivalent modulo an exact form to  $pd \log Y_j$  for  $1 \leq j \leq s$ . This implies that the map  $\mathfrak{p} : W_n\tilde{\omega}_Y^q \rightarrow W_{n+1}\tilde{\omega}_Y^q$  preserves the weight filtration  $P_{i,j}$ .

In order to see that  $\pi : W_{n+1}\tilde{\omega}_Y^q \rightarrow W_n\tilde{\omega}_Y^q$  also preserves  $P_{i,j}$  we use a descending induction on  $(i, j)$  in lexicographic order. Note that  $P_{r,s}W_n\tilde{\omega}_Y^q = W_n\tilde{\omega}_Y^q$ , so there is nothing to prove in this case. We can prove the result for  $(r, s - 1)$  in the same way as Proposition 8.4(2) of [Nakkajima 2005], using the commutative diagrams

$$\begin{array}{ccc}
 P_{i,j} W_{n+1} \tilde{\omega}_Y^q & \xrightarrow{\text{Res}} & W_{n+1} \Omega_{Y^{(i,j)}}^{q-i-j} \\
 \pi \downarrow & & \downarrow \pi \\
 P_{i,j} W_n \tilde{\omega}_Y^q & \xrightarrow{\text{Res}} & W_n \Omega_{Y^{(i,j)}}^{q-i-j}
 \end{array}$$

for  $(i, j)$  successively equal to  $(r, s), (r - 1, s), \dots, (1, s)$ . At the last step we get a commutative diagram of (vertical) exact sequences

$$\begin{array}{ccc}
 0 & & 0 \\
 \downarrow & & \downarrow \\
 P_{r,s-1} W_{n+1} \tilde{\omega}_Y^q + P_{0,s} W_{n+1} \tilde{\omega}_Y^q & & P_{r,s-1} W_n \tilde{\omega}_Y^q + P_{0,s} W_n \tilde{\omega}_Y^q \\
 \downarrow & & \downarrow \\
 P_{r,s-1} W_{n+1} \tilde{\omega}_Y^q + P_{1,s} W_{n+1} \tilde{\omega}_Y^q & \xrightarrow{\pi} & P_{r,s-1} W_n \tilde{\omega}_Y^q + P_{1,s} W_n \tilde{\omega}_Y^q \\
 \downarrow & & \downarrow \\
 W_{n+1} \Omega_{Y^{(1,s)}}^{q-s-1} & \xrightarrow{\pi} & W_n \Omega_{Y^{(1,s)}}^{q-s-1} \\
 \downarrow & & \downarrow \\
 0 & & 0
 \end{array}$$

which means there is an induced morphism  $\pi : P_{r,s-1} W_{n+1} \tilde{\omega}_Y^q + P_{0,s} W_{n+1} \tilde{\omega}_Y^q \rightarrow P_{r,s-1} W_n \tilde{\omega}_Y^q + P_{0,s} W_n \tilde{\omega}_Y^q$ .

At this stage, we note that we can define

$$Y^{(0,s)} = \bigsqcup_{\substack{T \subseteq \{1, \dots, n\} \\ \#T=s}} \left( \bigcap_{i \in T} Y_i^2 \right).$$

This will be a simple reduced normal crossings divisor over  $k$ , and we can endow it with the pullback of the log structure  $\tilde{M}_1$  so that  $(Y, \tilde{M})$  is a  $(k, \mathbb{N})$ -semistable log scheme, in the terminology of Section 2.4 of [Mokrane 1993]. There is a surjective residue morphism obtained via the restriction

$$P_{i,j} W_n \tilde{\omega}_Y^q \xrightarrow{\text{Res}} P_i W_n \tilde{\omega}_{Y^{(0,j)}}^{q-j},$$

which respects the weight filtrations. Just as the commutative diagram 8.4.3 of [Nakkajima 2005] is obtained, we can use the injectivity of  $\mathfrak{p} : W_n \tilde{\omega}_{Y^{(0,s)}}^q \rightarrow W_{n+1} \tilde{\omega}_{Y^{(0,s)}}^q$  for  $Y^{(0,s)}/k$  [Nakkajima 2005, Corollary 6.28(2)] to see that there is

a commutative diagram

$$\begin{array}{ccc}
 P_{0,s}W_{n+1}\tilde{\omega}_Y^q & \xrightarrow{\text{Res}} & P_0W_{n+1}\tilde{\omega}_{Y^{(0,s)}}^{q-s} \\
 \pi \downarrow & & \downarrow \pi \\
 P_{0,s}W_n\tilde{\omega}_Y^q & \xrightarrow{\text{Res}} & P_0W_n\tilde{\omega}_{Y^{(0,s)}}^{q-s}
 \end{array}$$

We therefore get a commutative diagram of (vertical) exact sequences:

$$\begin{array}{ccc}
 0 & & 0 \\
 \downarrow & & \downarrow \\
 P_{r,s-1}W_{n+1}\tilde{\omega}_Y^q & & P_{r,s-1}W_n\tilde{\omega}_Y^q \\
 \downarrow & & \downarrow \\
 P_{r,s-1}W_{n+1}\tilde{\omega}_Y^q + P_{0,s}W_{n+1}\tilde{\omega}_Y^q & \xrightarrow{\pi} & P_{r,s-1}W_n\tilde{\omega}_Y^q + P_{0,s}W_n\tilde{\omega}_Y^q \\
 \downarrow & & \downarrow \\
 P_0W_{n+1}\tilde{\omega}_{Y^{(0,s)}}^{q-s} & \xrightarrow{\pi} & P_0W_n\tilde{\omega}_{Y^{(0,s)}}^{q-s} \\
 \downarrow & & \downarrow \\
 0 & & 0
 \end{array}$$

so there is an induced morphism  $\pi : P_{r,s-1}W_{n+1}\tilde{\omega}_Y^q \rightarrow P_{r,s-1}W_n\tilde{\omega}_Y^q$ .

Finally, the third part follows in the same way as Corollary 8.6.4 of [Nakkajima 2005]. For an admissible lift  $(Z, \tilde{N})$ , let  $Z_1 := Z \times_W k$ . We have surjective morphisms  $W_n\Omega_{Z_1}^q \rightarrow P_{0,0}W_n\tilde{\omega}_Y^q$ , which commute with the transition morphisms  $\pi$ . So  $\pi$  is surjective for  $P_{0,0}$ . Using the exact sequences of the form

$$0 \rightarrow P_{0,j-1}W_n\tilde{\omega}_Y^q \rightarrow P_{0,j}W_n\tilde{\omega}_Y^q \rightarrow P_0W_n\tilde{\omega}_{Y^{(0,j)}}^{q-j} \rightarrow 0$$

and the surjectivity of  $\pi$  on the third term, we prove by induction on  $j$  that  $\pi$  is surjective for  $P_{0,j}$ . The same statement holds for  $P_{i,0}$ . Then, we prove that  $\pi$  is surjective for a general  $P_{i,j}$  by induction on  $i + j$ , using the exact sequences of the form

$$0 \rightarrow P_{i-1,j}W_n\tilde{\omega}_Y^q + P_{i,j-1}W_n\tilde{\omega}_Y^q \rightarrow P_{i,j}W_n\tilde{\omega}_Y^q \rightarrow W_n\Omega_{Y^{(i,j)}}^{q-i-j} \rightarrow 0. \quad \square$$

#### 4. Generalizing the Mokrane spectral sequence

In this section, we derive a generalization of the Mokrane spectral sequence which will allow us to compute the log crystalline cohomology of the Shimura varieties we are interested in terms of the crystalline cohomology of certain proper smooth

Newton polygon strata in the special fiber. Mokrane’s spectral sequence applies to the case of semistable reduction. Here we treat the case of a scheme whose singularities are locally those of a product of semistable schemes which is no longer semistable.

We define a double complex  $W_n A^{\bullet\bullet}$  as follows. Its terms are

$$W_n A^{ij} := \bigoplus_{k=0}^j W_n \tilde{\omega}_Y^{i+j+2} / (P_{k,i+j+2} + P_{i+j+2,j-k}) \quad \text{for } i, j \geq 0,$$

and  $W_n A^{ij} := 0$  otherwise. The operators  $d, \pi, F, V$  of  $W_\bullet \tilde{\omega}^\bullet$  induce operators  $d', \pi, F, V$  of the procomplexes  $W_\bullet A^{\bullet j}$ . For  $x$  in the direct summand  $W_n \tilde{\omega}_Y^{i+j+2} / (P_{k,i+j+2} + P_{i+j+2,j-k})$  of  $W_n A^{ij}$ ,  $d'x$  is the class of  $(-1)^j d\tilde{x}$ , where  $\tilde{x}$  is a lift of  $x$  in  $W_n \tilde{\omega}_Y^{i+j+2}$ . We also have a differential  $d'' : W_n A^{ij} \rightarrow W_n A^{i+1,j}$  given by

$$d''x = (-1)^i \left( \frac{d\tau}{\tau} \wedge x + \frac{d\sigma}{\sigma} \wedge x \right),$$

where  $d\tau/\tau$  and  $d\sigma/\sigma$  are the global sections of  $W_n \tilde{\omega}_Y^1$  defined in Lemma 3.9. We have  $d'd'' = d''d'$ , so we indeed get a double procomplex  $(W_\bullet A^{\bullet\bullet}, d', d'')$ . As in Lemma 3.9 of [Mokrane 1993], we can use dévissage by weights to see that the components of this procomplex are  $p$ -torsion-free. Let  $W_\bullet A^\bullet$  be the simple procomplex associated to the double procomplex  $W_\bullet A^{\bullet\bullet}$ .

We define now an endomorphism  $\nu$  of bidegree  $(-1, 1)$  of  $W_n A^{\bullet\bullet}$  which will induce the monodromy operator on cohomology. For each  $k \in \{0, \dots, j\}$  we have natural maps

$$\begin{aligned} W_n \tilde{\omega}_Y^{i+j+2} / (P_{k,i+j+2} + P_{i+j+2,j-k}) \\ \rightarrow W_n \tilde{\omega}_Y^{i+j+2} / (P_{k,i+j+2} + P_{i+j+2,j+1-k}) \\ \oplus W_n \tilde{\omega}_Y^{i+j+2} / (P_{k+1,i+j+2} + P_{i+j+2,j-k}), \end{aligned}$$

which are sums of  $(-1)^{i+j+1}$   $\text{proj}$  on each factor. Summing over  $k$  we get maps  $\nu : W_n A^{ij} \rightarrow W_n A^{i-1,j+1}$ , which induce an endomorphism  $\nu$  of bidegree  $(-1, 1)$ .

The morphism of complexes  $W_n \tilde{\omega}_Y^\bullet \rightarrow W_n A^{\bullet 0}$  given by

$$x \mapsto \frac{d\tau}{\tau} \wedge \frac{d\sigma}{\sigma} \wedge x$$

factors through  $W_n \omega_Y^\bullet$ . We get a morphism of complexes

$$\Theta : W_n \omega_Y^\bullet \rightarrow W_n A^\bullet.$$

The following lemma is analogous to Theorem 9.9 of [Nakkajima 2005]. It ensures that the resulting spectral sequence will be compatible with the Frobenius endomorphism (defined as an endomorphism of  $W_n$ -modules). We let  $\Phi_n : W_n\omega_Y \rightarrow W_n\omega_Y$  be the Frobenius endomorphism induced by the absolute Frobenius endomorphism of  $(Y, M)$ .

**Lemma 4.1.** *Let  $n$  be a positive integer. Then the following hold:*

- (1) *There exists a unique endomorphism  $\tilde{\Phi}_n^{\bullet, \bullet}$  of  $W_n A^{\bullet, \bullet}$  of double complexes, making the following diagram commutative:*

$$\begin{array}{ccc} W_{n+1}A^{qm} & \xrightarrow{\pi} & W_nA^{qm} \\ p^q F \downarrow & & \downarrow \tilde{\Phi}_n^{qm} \\ W_nA^{qm} & \xrightarrow{\text{id}} & W_nA^{qm} \end{array}$$

- (2) *The endomorphism  $\tilde{\Phi}_n^{\bullet, \bullet}$  induces an endomorphism  $\tilde{\Phi}_n$  of the complex  $W_n A^\bullet$ , fitting in a commutative diagram*

$$\begin{array}{ccc} W_n\omega_Y^\bullet & \xrightarrow{\Phi_n} & W_n\omega_Y^\bullet \\ \Theta \downarrow & & \downarrow \Theta \\ W_nA^\bullet & \xrightarrow{\tilde{\Phi}_n} & W_nA^\bullet \end{array}$$

- (3) *Finally, the Poincaré residue isomorphism Res fits in the following commutative diagrams for  $i, j \geq 1$ :*

$$\begin{array}{ccc} Gr_{i,j}W_n\tilde{\omega}_Y^q & \xrightarrow{\text{Res}} & W_n\Omega_{Y(i,j)}^{q-i-j} \\ \Psi_n \downarrow & & \downarrow p^{i+j}\Phi_n \\ Gr_{i,j}W_n\tilde{\omega}_Y^q & \xrightarrow{\text{Res}} & W_n\Omega_{Y(i,j)}^{q-i-j} \end{array}$$

where  $\Psi_n$  is an endomorphism of  $W_n\tilde{\omega}_Y^\bullet$  which respects the weight filtration  $P_{i,j}$  and which induces  $\tilde{\Phi}_n^{\bullet, \bullet}$  on  $W_n A^{\bullet, \bullet}$ .

*Proof.* The proof is essentially the same as that of Theorem 9.9 of [Nakkajima 2005]. We emphasize only the key points. We can define a morphism

$$\Psi_n^{j,q} : W_n\tilde{\omega}_Y^q \rightarrow W_n\tilde{\omega}_Y^q$$

via the composition

$$W_n \tilde{\omega}_Y^q \xrightarrow{p} W_{n+1} \tilde{\omega}_Y^q \xrightarrow{p^{j-1}} W_{n+1} \tilde{\omega}_Y^q \xrightarrow{F} W_n \tilde{\omega}_Y^q.$$

The fact that these morphisms commute with the maps  $(d\tau/\tau)\wedge$  and  $(d\sigma/\sigma)\wedge$  follows from the proof of the first part of Lemma 3.17. This implies that the second diagram is commutative. The fact that the  $\Psi_n^{\bullet\bullet}$  respect the weight filtration follows from the analogous statement for  $p$ , which is proved in Lemma 3.17 as well. This means that we can use  $\Psi_n^{jj+q+2}$  to define endomorphisms  $\tilde{\Phi}_n^{jq}$  of  $W_n A^{jq}$ , at least for  $j \geq 1$ . For  $j = 0$  we use the Frobenius endomorphism  $\Phi_n$  of  $W_n(\mathbb{C}_{Y^{(k+1, j-k+1)}})$  together with the residue isomorphisms to define  $\tilde{\Phi}_n^{0q}$ . The commutativity of the first diagram now follows from the definitions, from the commutative diagram

$$\begin{array}{ccc} W_{n+1} \tilde{\omega}_Y^{qm} & \xrightarrow{\pi} & W_n \tilde{\omega}_Y^{qm} \\ p^q F \downarrow & & \downarrow \Psi_n^{qm} \\ W_n A^{qm} & \xrightarrow{\text{id}} & W_n A^{qm} \end{array}$$

(which is deduced from  $pd = dp$  and  $dF = pFd$ ) and from Diagram 9.2.2 of [Nakkajima 2005] in the case of a smooth morphism. The fact that the first diagram is commutative ensures the uniqueness of  $\Phi_n^{q,m}$ . Finally, the third commutative diagram follows from the surjectivity of  $\pi$  proved in Lemma 3.17, from Diagram 9.2.2 of [Nakkajima 2005] in the case of a smooth morphism and from the commutative diagrams

$$\begin{array}{ccc} P_{i,j} W_{n+1} \tilde{\omega}_Y^q & \xrightarrow{\text{Res}} & W_{n+1} \Omega_{Y^{(i,j)}}^{q-i-j} \\ \pi \downarrow & & \downarrow \pi \\ P_{i,j} W_n \tilde{\omega}_Y^q & \xrightarrow{\text{Res}} & W_n \Omega_{Y^{(i,j)}}^{q-i-j} \end{array}$$

for  $i, j \geq 1$ . □

**Proposition 4.2.** *The sequence*

$$0 \rightarrow W_n \omega_Y^\bullet \xrightarrow{\Theta} W_n A^{\bullet 0} \xrightarrow{d''} W_n A^{\bullet 1} \xrightarrow{d''} \dots$$

is exact.

*Proof.* We follow the proof of Proposition 3.15 of [Mokrane 1993]. Let  $\theta : W_n \tilde{\omega}_Y^{i-1} \oplus W_n \tilde{\omega}_Y^{i-1} \rightarrow W_n \tilde{\omega}_Y^i$  be defined by

$$(x, y) \mapsto \frac{d\tau}{\tau} \wedge x + \frac{d\sigma}{\sigma} \wedge y.$$

It suffices to check that the sequence

$$\begin{aligned}
 W_n \tilde{\omega}_Y^{i-2} &\xrightarrow{(\frac{d\sigma}{\sigma} \wedge, \frac{d\tau}{\tau} \wedge)} W_n \tilde{\omega}_Y^{i-1} \oplus W_n \tilde{\omega}_Y^{i-1} \xrightarrow{\theta} W_n \tilde{\omega}_Y^i \\
 &\xrightarrow{\frac{d\tau}{\tau} \wedge \frac{d\sigma}{\sigma} \wedge} W_n \tilde{\omega}_Y^{i+2} / (P_{0,i+2} + P_{i+2,0}) \\
 &\xrightarrow{d''} W_n \tilde{\omega}_Y^{i+3} / (P_{1,i+3} + P_{i+3,0}) \oplus W_n \tilde{\omega}_Y^{i+3} / (P_{0,i+3} + P_{i+3,1}) \\
 &\xrightarrow{d''} \dots
 \end{aligned} \tag{4.1}$$

is exact. We do this by using first a dévissage by weights, reducing to the case  $n = 1$  and then using the fact that the scheme  $Y$  is Zariski-locally étale over a product of (the special fibers of) strictly semistable schemes.

We let

$$\begin{aligned}
 K_{-4} &= W_n \tilde{\omega}_Y^{i-2}, \\
 K_{-3} &= W_n \tilde{\omega}_Y^{i-1} \oplus W_n \tilde{\omega}_Y^{i-1}, \\
 K_{-2} &= W_n \tilde{\omega}_Y^i, \\
 K_j &= \bigoplus_{k=0}^j W_n \tilde{\omega}_Y^{i+j+2} / (P_{k,i+j+2} + P_{i+j+2,j-k}), \quad j \geq 0.
 \end{aligned}$$

For  $j \geq -4, j \neq -1$  we define a double filtration of  $K_j$  as follows:

$$\begin{aligned}
 P_{l,m} K_{-4} &= P_{l-2,m-2} W_n \tilde{\omega}_Y^{i-2}, \\
 P_{l,m} K_{-3} &= P_{l-2,m-1} W_n \tilde{\omega}_Y^{i-1} \oplus P_{l-1,m-2} W_n \tilde{\omega}_Y^{i-1}, \\
 P_{l,m} K_{-2} &:= P_{l-1,m-1} W_n \tilde{\omega}_Y^i, \\
 P_{l,m} K_j &:= \bigoplus_{k=0}^j P_{l+k,m+j-k} W_n \tilde{\omega}_Y^{i+j+2} / (P_{k,i+j+2} + P_{i+j+2,j-k}), \quad j \geq 0.
 \end{aligned}$$

Here we set the convention  $P_{l,m} W_n \tilde{\omega}_Y^i = 0$  if either  $l < 0$  or  $m < 0$ . The sequence (4.1) is a filtered sequence and to prove exactness it suffices to prove exactness for each graded piece

$$\text{Gr}_{l,m} K_j := P_{l,m} K_j / (P_{l,m-1} K_j + P_{l-1,m} K_j).$$

For  $l, m \geq 0$  we can rewrite the sequences of graded pieces as:

$$\begin{aligned}
 \text{Gr}_{l-2,m-2} W_n \tilde{\omega}_Y^{i-2} &\rightarrow \text{Gr}_{l-2,m-1} W_n \tilde{\omega}_Y^{i-1} \oplus \text{Gr}_{l-1,m-2} W_n \tilde{\omega}_Y^{i-1} \\
 &\rightarrow \text{Gr}_{l-1,m-1} W_n \tilde{\omega}_Y^i \rightarrow \text{Gr}_{l,m} W_n \tilde{\omega}_Y^{i+2} \\
 &\rightarrow \text{Gr}_{l+1,m} W_n \tilde{\omega}_Y^{i+3} \oplus \text{Gr}_{l,m+1} W_n \tilde{\omega}_Y^{i+3} \rightarrow \dots
 \end{aligned}$$

For  $l < 0$  or  $m < 0$  the sequence is trivial.

It suffices to show that the sequence of complexes

$$\begin{aligned} \mathrm{Gr}_{l-2,m-2} W_n \tilde{\omega}_Y^\bullet[-2] &\rightarrow \mathrm{Gr}_{l-2,m-1} W_n \tilde{\omega}_Y^\bullet[-1] \oplus \mathrm{Gr}_{l-1,m-2} W_n \tilde{\omega}_Y^\bullet[-1] \\ &\rightarrow \mathrm{Gr}_{l-1,m-1} W_n \tilde{\omega}_Y^\bullet \xrightarrow{\iota} \mathrm{Gr}_{l,m} W_n \tilde{\omega}_Y^\bullet[2] \\ &\rightarrow \mathrm{Gr}_{l+1,m} W_n \tilde{\omega}_Y^\bullet[3] \oplus \mathrm{Gr}_{l,m+1} W_n \tilde{\omega}_Y^\bullet[3] \rightarrow \dots \end{aligned} \quad (4.2)$$

is exact. Note that we can check this locally. When  $l, m \geq 1$ , we know by Corollary 3.16 that

$$\mathrm{Gr}_{l,m} W_n \tilde{\omega}_Y^\bullet \simeq W_n \Omega_{Y^{(l,m)}}^\bullet[-l-m](-l-m).$$

For  $l = 0$  and  $m \geq 1$  let  $Y_{D^{0,m}}$  be the normal crossing divisor of  $D^{0,m}$  corresponding to  $s = 0$ . In this case we have

$$\mathrm{Gr}_{l,m} W_n \tilde{\omega}_Y^\bullet \simeq [W_n \Omega_{D^{0,m}}(-\log Y_{D^{0,m}}) \rightarrow W_n \Omega_{D^{0,m}}],$$

and for  $l = 0, m = 0$  we have the quasi-isomorphism

$$\begin{aligned} \mathrm{Gr}_{l,m} W_n \tilde{\omega}_Y^\bullet &\simeq [W_n \Omega_Z^\bullet(-\log Y^1 - \log Y^2) \\ &\rightarrow W_n \Omega_Z^\bullet(-\log Y^1) \oplus W_n \Omega_Z^\bullet(-\log Y^2) \rightarrow W_n \Omega_Z^\bullet], \end{aligned}$$

where  $Z = \underline{Z} \otimes_W k$ . In any case,  $\mathrm{Gr}_{l,m} W_n \tilde{\omega}_Y^\bullet$  satisfies the property

$$(\lim \mathrm{Gr}_{l,m} W_n \tilde{\omega}_Y^\bullet) \otimes_{\mathbb{R}}^L \mathbb{R}_n \simeq \mathrm{Gr}_{l,m} W_n \tilde{\omega}_Y^\bullet$$

by Lemma 1.3.3 of [Mokrane 1993] and Lemma 3.11. By Proposition 2.3.7 of [Illusie 1983], it suffices to check exactness of the sequence (4.2) for  $n = 1$ .

For  $n = 1$  and working locally with our admissible lifts, we know that the exact sequence (4.2) is the pullback to  $Y$  of the corresponding exact sequence on  $Y_1 \times_k Y_2$ . We can assume that  $Y = Y_1 \times_k Y_2$  and  $Z = Z_1 \times_k Z_2$ . Each  $Y_i$  for  $i = 1, 2$  is a reduced normal crossings divisor in  $Z_i$ , for which we know that

$$\mathrm{Gr}_{i-2} W_1 \tilde{\omega}_{Y_i}^\bullet[-1] \rightarrow \mathrm{Gr}_{i-1} W_1 \tilde{\omega}_{Y_i}^\bullet \rightarrow \mathrm{Gr}_i W_1 \tilde{\omega}_{Y_i}^\bullet[1] \rightarrow \mathrm{Gr}_{i+1} W_1 \tilde{\omega}_{Y_i}^\bullet[2] \rightarrow \dots$$

is exact, by the proof of Proposition 3.15 of [Mokrane 1993]. In other words, for  $i = 1, 2$  we have quasi-isomorphisms between the top row and the bottom row. Multiplying the quasi-isomorphisms for  $i = 1, 2$  gives us exactly the quasi-isomorphism  $\iota$  needed to prove the exactness of (4.2) in the case  $n = 1$ . Here, we use the Cartier isomorphisms for  $W_1 \tilde{\omega}_{Y_i}$  and for  $W_1 \tilde{\omega}_Y$  and the fact that

$$(\omega_{(Z_1, \tilde{N}_1)/k}^\bullet \otimes_{\mathbb{O}_{Z_1}} \mathbb{O}_{Y_1}) \otimes_k (\omega_{(Z_2, \tilde{N}_2)/k}^\bullet \otimes_{\mathbb{O}_{Z_2}} \mathbb{O}_{Y_2}) \simeq \omega_{(Z, \tilde{N})}^\bullet \otimes_{\mathbb{O}_Z} \mathbb{O}_Y,$$

where the two complexes on the left determine  $W_1 \tilde{\omega}_{Y_i}^\bullet$  for  $i = 1, 2$  and the one on the right determines  $W_1 \tilde{\omega}_Y^\bullet$ . □

**Corollary 4.3.** *The morphism of complexes  $\Theta : W_n \omega_Y^\bullet \rightarrow W_n A^\bullet$  is a quasi-isomorphism. It induces a quasi-isomorphism  $\Theta : W \omega_Y^\bullet \xrightarrow{\sim} W A^\bullet$ .*

**Proposition 4.4.** *The endomorphism  $\nu$  of  $W_\bullet A^{\bullet\bullet}$  induces the monodromy operator  $N$  over  $H_{\text{cris}}^*((Y, M)/(W, \mathbb{N}))$ .*

*Proof.* We define the double complex  $B_n^{\bullet\bullet}$  as follows:

$$\begin{aligned} B_n^{\bullet\bullet} &= W_n A^{i-1j} \oplus W_n A^{ij}, \quad i, j \geq 0, \\ d'(x_1, x_2) &= (d'x_1, d'x_2), \\ d''(x_1, x_2) &= (d''x_1 + \nu(x_2), d''x_2). \end{aligned}$$

We have a morphism of complexes  $\Psi : W_n \tilde{\omega}_Y^\bullet \rightarrow B_n^{\bullet\bullet}$  defined as follows: for  $x \in W_n \tilde{\omega}_Y^i$ ,

$$\begin{aligned} &\Psi(x) \\ &= \left( \left( \frac{d\sigma}{\sigma} - \frac{d\tau}{\tau} \right) \wedge x \pmod{P_{0,i+1} + P_{i+1,0}}, \frac{d\tau}{\tau} \wedge \frac{d\sigma}{\sigma} \wedge x \pmod{P_{0,i+2} + P_{i+2,0}} \right). \end{aligned}$$

Thus we have a commutative diagram of exact sequences of complexes:

$$\begin{array}{ccccccc} 0 & \longrightarrow & W_n \omega_Y^\bullet[-1] & \longrightarrow & W_n \tilde{\omega}_Y^\bullet & \longrightarrow & W_n \omega_Y^\bullet \longrightarrow 0 \\ & & \downarrow \Theta[-1] & & \downarrow \Psi & & \downarrow \Theta \\ 0 & \longrightarrow & W_n A^\bullet[-1] & \longrightarrow & B_n^{\bullet\bullet} & \longrightarrow & W_n A^\bullet \longrightarrow 0 \end{array}$$

where the left and right downward arrows are quasi-isomorphisms. Thus,  $\Psi$  is also a quasi-isomorphism and the commutative diagram defines an isomorphism of distinguished triangles. Thus the monodromy operator  $N$  on cohomology is induced by the coboundary operator of the bottom exact sequence, which by construction is  $\nu$ . □

We can compute the monodromy filtration of the nilpotent operator  $N$  on cohomology from the monodromy filtration of  $\nu$  on  $W_n A^\bullet$ . We will exhibit a filtration  $P_k(W_n A^\bullet) = \bigoplus_{i,j \geq 0} P_k(W_n A^{ij})$  which satisfies the following:

- (1)  $\nu(P_k(W_\bullet A^\bullet)) \subset P_{k-2}(W_\bullet A^\bullet)(-1)$ .
- (2) For  $k \geq 0$ , the induced map  $\nu^k : \text{Gr}_k(W_\bullet A^\bullet) \rightarrow \text{Gr}_{-k}(W_\bullet A^\bullet)(-k)$  is an isomorphism.

A filtration satisfying these two properties must be the monodromy filtration of  $\nu$ .

**Note 4.5.** From now on, we will not work in the category  $\mathcal{C}$  of complexes of sheaves of  $W$ -modules but rather in  $\mathbb{Q} \otimes \mathcal{C}$ , which is the category with the same set of objects as  $\mathcal{C}$ , but with morphisms  $\mathbb{Q} \otimes \text{Hom}_{\mathcal{C}}(A, B)$ . We will in fact identify the monodromy filtration of  $\nu$  on  $\mathbb{Q} \otimes W_n A^\bullet$ , but for simplicity of notation we still denote an object  $A$  of  $\mathcal{C}$  as  $A$  when we regard it as an object of  $\mathbb{Q} \otimes \mathcal{C}$ .

Define  $P_l(W_n A^{\bullet\bullet}) := \bigoplus_{i,j \geq 0} P_l(W_n A^{ij})$  for  $l \geq 0$ , where  $P_l(W_n A^{ij})$  is 0 if  $l < 2n - 2 - j$  and

$$\bigoplus_{k=0}^j \left( \sum_{m=0}^{l-2n+2+j} P_{k+m+1, 2j-k+l-2n-m+3} W_n \tilde{\omega}_Y^{i+j+2} / (P_{k,i+j+2} + P_{j-k,i+j+2}) \right)$$

if  $l \geq 2n - 2 - j$ . It is easy to check that  $v(P_l(W_n A^{ij})) \subset P_{l-2} W_n A^{i+1, j-1}$ . Moreover, we can also compute the graded pieces  $\text{Gr}_l(W_n A^{\bullet\bullet}) = \bigoplus_{i,j \geq 0} \text{Gr}_l(W_n A^{ij})$ , where

$$\begin{aligned} & \text{Gr}_l(W_n A^{ij}) \\ &= \begin{cases} 0 & \text{if } l < 2n - 2 - j, \\ \bigoplus_{k=0}^j \bigoplus_{m=0}^{l-2n+2+j} \text{Gr}_{k+m+1, 2j-k+l-2n-m+3} W_n \tilde{\omega}_Y^{i+j+2} & \text{if } l \geq 2n - 2 - j. \end{cases} \end{aligned}$$

For  $l = 2n - 2 + h$ , with  $h > 0$ , we claim that  $v$  induces an injection  $\text{Gr}_l(W_n A^{ij}) \hookrightarrow \text{Gr}_{l-1}(W_n A^{ij})$ . This can be verified through a standard combinatorial argument. We have

$$\text{Gr}_l(W_n A^{ij}) = \bigoplus_{k=0}^j \bigoplus_{m=0}^{h+j} \text{Gr}_{(k+m)+1, 2j+h+1-(k+m)} W_n \tilde{\omega}_Y^{i+j+2}$$

and

$$\text{Gr}_{l-1}(W_n A^{ij}) = \bigoplus_{k=0}^{j+1} \bigoplus_{m=0}^{h+j-1} \text{Gr}_{(k+m)+1, 2j+h+1-(k+m)} W_n \tilde{\omega}_Y^{i+j+2}.$$

The map  $v$  sends the term corresponding to a pair  $(k, m)$  to the direct sum of terms corresponding to  $(k, m)$  and to  $(k + 1, m - 1)$ . Therefore, it is easy to see that  $v$  restricted to the direct sum of terms for which  $k + m$  is constant is injective, so  $v$  is injective. Moreover, we see that  $v^h$  induces an isomorphism  $\text{Gr}_{2n-2+h}(W_n A^{ij}) \simeq \text{Gr}_{2n-2-h}(W_n A^{i-h, j+h})$ , since the terms on the right-hand side are of the form

$$\bigoplus_{k=0}^j \bigoplus_{m=0}^{h+j} \text{Gr}_{(k+m)+1, 2j+h+1-(k+m)} W_n \tilde{\omega}_Y^{i+j+2}$$

and the terms on the left-hand side are of the form

$$\bigoplus_{m=0}^j \bigoplus_{k=0}^{h+j} \text{Gr}_{(k+m)+1, 2j+h+1-(k+m)} W_n \tilde{\omega}_Y^{i+j+2},$$

so on either side we have the same number of terms corresponding to  $k + m$ . Since the filtration  $P_l(W_n A^{\bullet\bullet})$  satisfies the two properties above, it must be the monodromy filtration of  $v$ .

Note that the differentials  $d''$  on  $\text{Gr}_l(W_\bullet A^{\bullet\bullet})$  are always 0. Using the isomorphisms in Corollary 3.16 we can rewrite

$$\text{Gr}_{2n-2+h}(W_\bullet A^\bullet) \simeq \bigoplus_{j \geq 0, j \geq -h} \bigoplus_{k=0}^j \bigoplus_{m=0}^{j+h-k} (W_\bullet \Omega_{Y^{(k+m+1, 2j+h+1-(k+m))}}^\bullet)[-2j-h](-j-h).$$

This leads to the main geometric result of the paper. Recall from Section 3.1 that  $Y/k$  is the special fiber of  $X/\mathbb{O}_K$ , which is Zariski-locally étale over a product of strictly semistable schemes. Recall also that  $Y$  is globally the union of certain proper, smooth  $(2n-2)$ -dimensional subschemes  $Y_{i,j}$  with  $i = 1, 2, j = 1, \dots, n$ . Taking disjoint unions of intersections of these subschemes gives rise to schemes  $Y^{(l_1, l_2)}/k$  for  $1 \leq l_1, l_2 \leq n$ , which cover closed strata in  $Y$ . Each  $Y^{(l_1, l_2)}$  is proper, smooth and has dimension  $2n - l_1 - l_2$ .

**Theorem 4.6.** *There is a spectral sequence*

$$E_1^{-h, i+h} = \bigoplus_{j \geq 0, j \geq -h} \bigoplus_{k=0}^j \bigoplus_{m=0}^{j+h-k} H_{\text{cris}}^{i-2j-h}(Y^{(k+m+1, 2j+h+1-(k+m))}/W)(-j-h) \Rightarrow H_{\text{cris}}^i(Y/W).$$

**Remark 4.7.** Note that the schemes  $Y^{(l_1, l_2)}$  are proper and smooth so the  $E^{-h, i+h}$  terms of the spectral sequence are strictly pure of weight  $i+h$ . If the above spectral sequence degenerates at the first page, then  $H_{\text{cris}}^i(Y/W)$  is pure of weight  $i$ .

### 5. Proof of the main theorem

In this section we prove the main theorem. By Corollary 2.3, its proof reduces to the following proposition.

**Proposition 5.1.** *Let  $\mathcal{A}_{U_{\text{Iw}}}^{m_\xi}$  be the universal abelian variety over  $X_{U_{\text{Iw}}}^{m_\xi}$ . The direct limit of log crystalline cohomologies*

$$\lim_{\substack{\rightarrow \\ U_{\text{Iw}}}} a_\xi(H_{\text{cris}}^{2n-2+m_\xi}(\mathcal{A}_{U_{\text{Iw}}}^{m_\xi} \times_{\mathbb{O}_K} k/W) \otimes_W \overline{\mathbb{Q}}_l(t_\xi))[\Pi^1, \mathfrak{S}]$$

*is pure of a certain weight.*

*Proof.* Recall that we have chosen

$$U_{\text{Iw}} = U^l \times U_l^{p_1, p_2}(m) \times \text{Iw}_{n, p_1} \times \text{Iw}_{n, p_2} \subset G(\mathbb{A}^\infty).$$

Pick  $m$  large enough such that  $(\pi_l) U_l^{p_1, p_2}(m) \times \text{Iw}_{n, p_1} \times \text{Iw}_{n, p_2}$  is nonzero, where  $\pi_l \in \text{Irr}_l(G(\mathbb{Q}_l))$  is such that  $BC(\pi_l) = \iota_l^{-1} \Pi_l$ . The results of Sections 3 and 4 apply to  $\mathcal{A}_{U_{\text{Iw}}}^{m_\xi}$ . We have a stratification of its special fiber by closed Newton polygon

strata  $\mathcal{A}_{U_{\text{Iw}}, S, T}^{m_\xi}$  with  $S, T \subseteq \{1, \dots, n\}$  nonempty. For brevity, let  $K_1 := k + m + 1$  and  $K_2 := 2j + h + 1 - (k + m)$ . By Theorem 4.6, we have a spectral sequence

$$E_1^{-h, i+h} = \bigoplus_{\substack{j \geq 0 \\ j \geq -h}} \bigoplus_{k=0}^j \bigoplus_{m=0}^{j+h-k} \bigoplus_{\substack{\#S=K_1 \\ \#T=K_2}} H_{\text{cris}}^{i-2j-h}((\mathcal{A}_{U_{\text{Iw}}, S, T}^{m_\xi}/W)(-j-h)) \Rightarrow H_{\text{cris}}^i(\mathcal{A}_{U_{\text{Iw}}}^{m_\xi} \times_{\mathbb{O}_K} k/W).$$

We replace the cohomology degree  $i$  by  $i + m_\xi$ , tensor with  $\overline{\mathbb{Q}}_l(t_\xi)$ , apply  $a_\xi$  (which is obtained from a linear combination of étale morphisms); passing to a direct limit over  $U^l$  and taking the  $\Pi^{1, \mathfrak{S}}$ -isotypic components we get a spectral sequence

$$E_1^{-h, i+h} = \bigoplus_{\substack{j \geq 0 \\ j \geq -h}} \bigoplus_{k=0}^j \bigoplus_{m=0}^{j+h-k} \bigoplus_{\substack{\#S=K_1 \\ \#T=K_1}} \lim_{\substack{\rightarrow \\ U^l}} (a_\xi H_{\text{cris}}^{i+m_\xi-2j-h}((\mathcal{A}_{U_{\text{Iw}}, S, T}^{m_\xi}/W)(-j-h) \otimes_{W, \tau_0} \overline{\mathbb{Q}}_l(t_\xi))[\Pi^{1, \mathfrak{S}}]) \Rightarrow \lim_{\substack{\rightarrow \\ U^l}} (a_\xi H_{\text{cris}}^{i+m_\xi}(\mathcal{A}_{U_{\text{Iw}}}^{m_\xi} \times_{\mathbb{O}_K} k/W) \otimes_{W, \tau_0} \overline{\mathbb{Q}}_l(t_\xi))[\Pi^{1, \mathfrak{S}}].$$

For any compact open subgroup  $U^l \subset G(\mathbb{A}^{\infty, l})$  and any prime  $p \neq l$  with isomorphism  $\iota_p : \overline{\mathbb{Q}}_p \xrightarrow{\sim} \mathbb{C}$ , set  $\xi' := (\iota_p)^{-1} \iota_l \xi$  and  $\Pi' := (\iota_p)^{-1} \Pi^1$ .

We have

$$\begin{aligned} \dim_{\overline{\mathbb{Q}}_l} (\lim_{\substack{\rightarrow \\ U^l}} a_\xi H_{\text{cris}}^{i+m_\xi-2j-h}((\mathcal{A}_{U_{\text{Iw}}, S, T}^{m_\xi}/W)(-j-h) \otimes_{W, \tau_0} \overline{\mathbb{Q}}_l))[\Pi^{1, \mathfrak{S}}]^{U^l} \\ = \dim_{\overline{\mathbb{Q}}_p} (\lim_{\substack{\rightarrow \\ U^l}} a_{\xi'} H_{\text{cris}}^{i+m_{\xi'}-2j-h}(\mathcal{A}_{U_{\text{Iw}}, S, T}^{m_\xi}, \overline{\mathbb{Q}}_p))[(\Pi')^{\mathfrak{S}}]^{U^l} \\ = \dim_{\overline{\mathbb{Q}}_p} (\lim_{\substack{\rightarrow \\ U^l}} H^{i-2j-h}(X_{U_{\text{Iw}}, S, T}, \mathcal{L}_{\xi'}))[(\Pi')^{\mathfrak{S}}]^{U^l}. \end{aligned}$$

The first equality is a consequence of the main theorem of [Gillet and Messing 1987] and of Theorem 2(2) of [Katz and Messing 1974]. The former proves that crystalline cohomology is a Weil cohomology theory in the strong sense. The latter is the statement that the characteristic polynomial on  $H^i(X)$  of an integrally algebraic cycle on  $X \times X$  of codimension  $n$ , for a projective smooth variety  $X/k$  of dimension  $n$ , is independent of the Weil cohomology theory  $H$ .

The dimension in the third row is equal to 0 unless  $i = 2n - 2$  by Proposition 5.10 of [Caraiani 2012]. Therefore,  $E_1^{-h, i+h} = 0$  unless  $i = 2n - 2$ , so the  $E_1$  page of the spectral sequence is concentrated on a diagonal. The spectral sequence degenerates at the  $E_1$  page and the term corresponding to  $E_1^{h, 2n-2+h}$  is strictly pure of weight  $h + 2n - 2 + m_\xi - 2t_\xi$ , which shows that the abutment is pure.  $\square$

### Acknowledgements

I am very grateful to my advisor, Richard Taylor, for suggesting this problem and for his constant encouragement and advice. I would like to thank Kazuya Kato, Mark Kisin, Jay Pottharst and Claus Sorensen for many useful conversations and comments. I would also like to thank Luc Illusie for his detailed comments on an earlier draft of this paper and in particular for pointing me to [Nakkajima 2005].

### References

- [Barnet-Lamb et al. 2011] T. Barnet-Lamb, T. Gee, D. Geraghty, and R. Taylor, “Local-global compatibility for  $l = p$ , II”, preprint, 2011, Available at <https://www2.bc.edu/david-geraghty/files/lgcII.pdf>. To appear in *Ann. scient. de l’E.N.S.*
- [Barnet-Lamb et al. 2012] T. Barnet-Lamb, T. Gee, D. Geraghty, and R. Taylor, “Local-global compatibility for  $l = p$ , I”, *Ann. Fac. Sci. Toulouse Math.* (6) **21**:1 (2012), 57–92. MR 2954105 Zbl 1259.11057
- [Caraiani 2012] A. Caraiani, “Local-global compatibility and the action of monodromy on nearby cycles”, *Duke Math. J.* **161**:12 (2012), 2311–2413. MR 2972460 Zbl 06095601
- [Chenevier and Harris 2013] G. Chenevier and M. Harris, “Construction of automorphic Galois representations, II”, *Cambridge Math. J.* **1**:1 (2013), 53–73. Zbl 06266535
- [Deligne 1970] P. Deligne, *Équations différentielles à points singuliers réguliers*, Lecture Notes in Mathematics **163**, Springer, Berlin, 1970. MR 54 #5232 Zbl 0244.14004
- [Deligne and Illusie 1987] P. Deligne and L. Illusie, “Relèvements modulo  $p^2$  et décomposition du complexe de de Rham”, *Invent. Math.* **89**:2 (1987), 247–270. MR 88j:14029 Zbl 0632.14017
- [Gillet and Messing 1987] H. Gillet and W. Messing, “Cycle classes and Riemann–Roch for crystalline cohomology”, *Duke Math. J.* **55**:3 (1987), 501–538. MR 89c:14025 Zbl 0651.14014
- [Grothendieck 1967] A. Grothendieck, “Éléments de géométrie algébrique, IV: Étude locale des schémas et des morphismes de schémas, IV”, *Inst. Hautes Études Sci. Publ. Math.* **32** (1967), 361. MR 39 #220 Zbl 0153.22301
- [Harris and Taylor 2001] M. Harris and R. Taylor, *The geometry and cohomology of some simple Shimura varieties*, Annals of Mathematics Studies **151**, Princeton University Press, 2001. MR 2002m:11050 Zbl 1036.11027
- [Hyodo 1991] O. Hyodo, “On the de Rham–Witt complex attached to a semi-stable family”, *Compositio Math.* **78**:3 (1991), 241–260. MR 93c:14022 Zbl 0742.14015
- [Hyodo and Kato 1994] O. Hyodo and K. Kato, “Semi-stable reduction and crystalline cohomology with logarithmic poles”, pp. 221–268 in *Périodes  $p$ -adiques* (Bures-sur-Yvette, 1988), Astérisque **223**, Société Mathématique de France, Paris, 1994. MR 95k:14034 Zbl 0852.14004
- [Illusie 1979] L. Illusie, “Complexe de de Rham–Witt et cohomologie cristalline”, *Ann. Sci. École Norm. Sup.* (4) **12**:4 (1979), 501–661. MR 82d:14013 Zbl 0436.14007
- [Illusie 1983] L. Illusie, “Finiteness, duality, and Künneth theorems in the cohomology of the de Rham–Witt complex”, pp. 20–72 in *Algebraic geometry* (Tokyo/Kyoto, 1982), edited by M. Raynaud and T. Shioda, Lecture Notes in Math. **1016**, Springer, Berlin, 1983. MR 85m:14033 Zbl 0538.14013
- [Illusie and Raynaud 1983] L. Illusie and M. Raynaud, “Les suites spectrales associées au complexe de de Rham–Witt”, *Inst. Hautes Études Sci. Publ. Math.* **57** (1983), 73–212. MR 85e:14026 Zbl 0538.14012

- [Kato 1989] K. Kato, “Logarithmic structures of Fontaine–Illusie”, pp. 191–224 in *Algebraic analysis, geometry, and number theory* (Baltimore, MD, 1988), edited by J.-I. Igusa, Johns Hopkins Univ. Press, Baltimore, MD, 1989. MR 99b:14020 Zbl 0776.14004
- [Kato 1994a] K. Kato, “Semi-stable reduction and  $p$ -adic étale cohomology”, pp. 269–293 in *Périodes  $p$ -adiques* (Bures-sur-Yvette, 1988), Astérisque **223**, Société Mathématique de France, Paris, 1994. MR 95i:14020 Zbl 0847.14009
- [Kato 1994b] K. Kato, “Toric singularities”, *Amer. J. Math.* **116**:5 (1994), 1073–1099. MR 95g:14056 Zbl 0832.14002
- [Katz and Messing 1974] N. M. Katz and W. Messing, “Some consequences of the Riemann hypothesis for varieties over finite fields”, *Invent. Math.* **23** (1974), 73–77. MR 48 #11117 Zbl 0275.14011
- [Mantovan 2005] E. Mantovan, “On the cohomology of certain PEL-type Shimura varieties”, *Duke Math. J.* **129**:3 (2005), 573–610. MR 2006g:11122 Zbl 1112.11033
- [Mokrane 1993] A. Mokrane, “La suite spectrale des poids en cohomologie de Hyodo–Kato”, *Duke Math. J.* **72**:2 (1993), 301–337. MR 95a:14022 Zbl 0834.14010
- [Nakkajima 2005] Y. Nakkajima, “ $p$ -adic weight spectral sequences of log varieties”, *J. Math. Sci. Univ. Tokyo* **12**:4 (2005), 513–661. MR 2007e:14030 Zbl 1108.14015
- [Nizioł 2006] W. Nizioł, “Toric singularities: Log-blow-ups and global resolutions”, *J. Algebraic Geom.* **15**:1 (2006), 1–29. MR 2006i:14015 Zbl 1100.14011
- [Nizioł 2008] W. Nizioł, “Semistable conjecture via  $K$ -theory”, *Duke Math. J.* **141**:1 (2008), 151–178. MR 2008m:14035 Zbl 1157.14009
- [Shin 2011] S. W. Shin, “Galois representations arising from some compact Shimura varieties”, *Ann. of Math. (2)* **173**:3 (2011), 1645–1741. MR 2800722 Zbl 1269.11053
- [Taylor and Yoshida 2007] R. Taylor and T. Yoshida, “Compatibility of local and global Langlands correspondences”, *J. Amer. Math. Soc.* **20**:2 (2007), 467–493. MR 2007k:11193 Zbl 1210.11118
- [Tsuji 1997] T. Tsuji, “Saturated morphisms of logarithmic schemes”, preprint, 1997.
- [Tsuji 1999] T. Tsuji, “ $p$ -adic étale cohomology and crystalline cohomology in the semi-stable reduction case”, *Invent. Math.* **137**:2 (1999), 233–411. MR 2000m:14024 Zbl 0945.14008

Communicated by Brian Conrad

Received 2013-04-27

Revised 2014-03-31

Accepted 2014-05-18

caraiani@princeton.edu

*Department of Mathematics, Princeton University, Fine Hall,  
Washington Road, Princeton, NJ 08544, United States*

# Algebra & Number Theory

msp.org/ant

## EDITORS

### MANAGING EDITOR

Bjorn Poonen  
Massachusetts Institute of Technology  
Cambridge, USA

### EDITORIAL BOARD CHAIR

David Eisenbud  
University of California  
Berkeley, USA

## BOARD OF EDITORS

Georgia Benkart	University of Wisconsin, Madison, USA	Shigefumi Mori	RIMS, Kyoto University, Japan
Dave Benson	University of Aberdeen, Scotland	Raman Parimala	Emory University, USA
Richard E. Borcherds	University of California, Berkeley, USA	Jonathan Pila	University of Oxford, UK
John H. Coates	University of Cambridge, UK	Anand Pillay	University of Notre Dame, USA
J-L. Colliot-Thélène	CNRS, Université Paris-Sud, France	Victor Reiner	University of Minnesota, USA
Brian D. Conrad	University of Michigan, USA	Peter Sarnak	Princeton University, USA
Hélène Esnault	Freie Universität Berlin, Germany	Joseph H. Silverman	Brown University, USA
Hubert Flenner	Ruhr-Universität, Germany	Michael Singer	North Carolina State University, USA
Edward Frenkel	University of California, Berkeley, USA	Vasudevan Srinivas	Tata Inst. of Fund. Research, India
Andrew Granville	Université de Montréal, Canada	J. Toby Stafford	University of Michigan, USA
Joseph Gubeladze	San Francisco State University, USA	Bernd Sturmfels	University of California, Berkeley, USA
Roger Heath-Brown	Oxford University, UK	Richard Taylor	Harvard University, USA
Craig Huneke	University of Virginia, USA	Ravi Vakil	Stanford University, USA
Yujiro Kawamata	University of Tokyo, Japan	Michel van den Bergh	Hasselt University, Belgium
János Kollár	Princeton University, USA	Marie-France Vignéras	Université Paris VII, France
Yuri Manin	Northwestern University, USA	Kei-Ichi Watanabe	Nihon University, Japan
Barry Mazur	Harvard University, USA	Efim Zelmanov	University of California, San Diego, USA
Philippe Michel	École Polytechnique Fédérale de Lausanne	Shou-Wu Zhang	Princeton University, USA
Susan Montgomery	University of Southern California, USA		

## PRODUCTION

production@msp.org  
Silvio Levy, Scientific Editor

---

See inside back cover or [msp.org/ant](http://msp.org/ant) for submission instructions.

The subscription price for 2014 is US \$225/year for the electronic version, and \$400/year (+\$55, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscribers address should be sent to MSP.

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

---

ANT peer review and production are managed by EditFLOW<sup>®</sup> from MSP.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2014 Mathematical Sciences Publishers

# Algebra & Number Theory

Volume 8 No. 7 2014

---

Double Dirichlet series and quantum unique ergodicity of weight one-half Eisenstein series	1539
YIANNIS N. PETRIDIS, NICOLE RAULF and MORTEN S. RISAGER	
Monodromy and local-global compatibility for $l = p$	1597
ANA CARAIANI	
Finite generation of the cohomology of some skew group algebras	1647
VAN C. NGUYEN and SARAH WITHERSPOON	
On the supersingular locus of the $\mathrm{GU}(2,2)$ Shimura variety	1659
BENJAMIN HOWARD and GEORGIOS PAPPAS	
Poincaré–Birkhoff–Witt deformations of smash product algebras from Hopf actions on Koszul algebras	1701
CHELSEA WALTON and SARAH WITHERSPOON	
Highly biased prime number races	1733
DANIEL FIORILLI	
Bounded gaps between primes with a given primitive root	1769
PAUL POLLACK	



1937-0652(2014)8:7;1-4