

Algebra & Number Theory

Volume 9

2015

No. 6

**Schubert decompositions for quiver Grassmannians
of tree modules**

Oliver Lorscheid
Appendix by Thorsten Weist



Schubert decompositions for quiver Grassmannians of tree modules

Oliver Lorscheid
Appendix by Thorsten Weist

Let Q be a quiver, M a representation of Q with an ordered basis \mathcal{B} and \underline{e} a dimension vector for Q . In this note we extend the methods of Lorscheid (2014) to establish Schubert decompositions of quiver Grassmannians $\text{Gr}_{\underline{e}}(M)$ into affine spaces to the ramified case, i.e., the canonical morphism $F : T \rightarrow Q$ from the coefficient quiver T of M w.r.t. \mathcal{B} is not necessarily unramified.

In particular, we determine the Euler characteristic of $\text{Gr}_{\underline{e}}(M)$ as the number of *extremal successor closed subsets* of T_0 , which extends the results of Cerulli Irelli (2011) and Haupt (2012) (under certain additional assumptions on \mathcal{B}).

Introduction

The recent interest in quiver Grassmannians stems from a formula of Caldero and Chapoton [2006] that relates cluster variables of a quiver Q with the Euler characteristics of the quiver Grassmannians of exceptional modules of Q . Formulas for the Euler characteristics for a given quiver yield a description of the associated cluster algebra in terms of generators and relations. This opened a way to understand cluster algebras, which are defined by an infinite recursive procedure, in terms of closed formulas—provided one knows the Euler characteristics of the associated quiver Grassmannians.

Torus actions and cluster algebras associated with string algebras. While the classification of all cluster algebras seems to be as much out of reach as a classification of wild algebras, there is some hope to understand and classify cluster algebras that are associated with tame algebras. A first step in this direction has been realized by Cerulli Irelli [2011] and Haupt [2012] who established a formula for the Euler characteristics of quiver Grassmannians in the so-called unramified case. These results sufficed to understand all cluster algebras associated with string algebras.

MSC2010: primary 14M15; secondary 16G20, 05C05.

Keywords: quiver Grassmannian, Schubert decompositions, Euler characteristics, tree modules.

We review the method of Cerulli Irelli and Haupt in brevity: following Ringel [1998], we note that every exceptional representation M of a quiver Q has a tree basis \mathcal{B} , meaning that the coefficient quiver $T = \Gamma(M, \mathcal{B})$ is a tree. A subset β of $T_0 = \mathcal{B}$ is *successor closed* if for all $i \in \beta$ and all arrows $\alpha : i \rightarrow j$ in T , also $j \in \beta$. A subset β of $T_0 = \mathcal{B}$ is *of type $\underline{e} = (e_p)_{p \in Q_0}$* if $\#\beta \cap M_p = e_p$ for all $p \in Q_0$.

If the canonical morphism $F : T \rightarrow Q$ is *unramified*, i.e., the morphism of the underlying CW-complexes is locally injective, then one can define a (piecewise continuous) action of the torus \mathbb{G}_m on $\text{Gr}_{\underline{e}}(M)$ that has only finitely many fixed points. This yields the formula

$$\chi(\text{Gr}_{\underline{e}}(M)) = \#\{\text{fixed points}\} = \#\{\text{successor closed } \beta \subset T_0 \text{ of type } \underline{e}\}.$$

For other types of cluster algebras, the exceptional modules are in general not unramified tree modules. This is, for instance, the case for cluster algebras associated with exceptional Dynkin quivers of types \tilde{D} and \tilde{E} , or, more general, for cluster algebras associated with clannish algebras or exceptional tame algebras. Therefore other methods are required to treat ramified tree modules.

Cluster algebras from marked surfaces. Fomin, Shapiro and Thurston explore in [Fomin et al. 2008] the connection between cluster algebras and marked surfaces. Namely, to each surface with boundary and finitely many marked points such that each boundary component contains at least one marked point, one can associate a cluster algebra. It is shown in that paper that all cluster algebras associated with quivers of extended Dynkin types \tilde{A} and \tilde{D} come from marked surfaces.

This connection with marked surfaces yields a description of the cluster variables in terms of triangulations of the surface, which leads to a combinatorial description of the algebra. For unpunctured surfaces, i.e., all marked points are contained in the boundary, Musiker, Schiffler and Williams [Musiker et al. 2013] construct a basis for the associated cluster algebra.

Cluster algebras of punctured surfaces, which includes type D algebras, are more difficult to treat since not all mutations of clusters come from flips of triangulations. For recent results in this direction, see [Qiu and Zhou 2013]. However, these methods do not suffice yet for a complete understanding of the associated cluster algebras.

Schubert decompositions and ramified tree modules. Caldero and Reineke [2008] show that $\text{Gr}_{\underline{e}}(M)$ is smooth projective if M is exceptional. If M is an equi-oriented string module, i.e., the coefficient quiver T is an equi-oriented Dynkin quiver of type A_n , then $\text{Gr}_{\underline{e}}(M)$ has a continuous torus action with finitely many fixed points, see [Cerulli Irelli 2011]. Thus if M is an exceptional equi-oriented string module, then the Białyński–Birula decomposition [1973, Theorem 4.3] yields a decomposition of $\text{Gr}_{\underline{e}}(M)$ into affine spaces.

While a torus action with finitely many fixed points determines the Euler characteristic, a decomposition of $\text{Gr}_{\underline{e}}(M)$ into affine spaces determines the (additive structure of the) cohomology of $\text{Gr}_{\underline{e}}(M)$, which is a much stronger result. In particular, we re-obtain the Euler characteristic as the number of affine spaces occurring in the decomposition. However, the class of exceptional equi-oriented string modules is very limited. In particular, most exceptional modules of affine type D are not of this kind.

In [Lorscheid 2014], we extend decompositions of $\text{Gr}_{\underline{e}}(M)$ into affine spaces to a larger class of quiver Grassmannians by a different method. Namely, the choice of an ordered basis \mathcal{B} of M defines a decomposition of $\text{Gr}_{\underline{e}}(M)$ into Schubert cells, which are, in general, merely closed subsets of affine spaces. In certain cases, however, these Schubert cells are affine spaces themselves. The method of proof is to exhibit explicit presentations of Schubert cells in terms of generators and relations.

One requirement of [Lorscheid 2014] is that the morphism $F : T \rightarrow Q$ is unramified. It is the purpose of this note to extend the methods of that paper to ramified $F : T \rightarrow Q$. In particular, this extends, under the given additional assumptions, the formula of Cerulli Irelli and Haupt to the ramified case.

As will be shown in joint work with Thorsten Weist, the results of this text are indeed applicable to all exceptional modules of affine type \tilde{D}_n , which yields combinatorial formulas for the Euler characteristics of $\text{Gr}_{\underline{e}}(M)$.

The main result of this text. An arrow α of T is *extremal* if for every other arrow α' with $F(\alpha') = F(\alpha)$ either $s(\alpha) < s(\alpha')$ or $t(\alpha') < t(\alpha)$. A subset β of T_0 is *extremal successor closed* if for every $i \in \beta$ and every extremal arrow $\alpha : i \rightarrow j$ in T , we also have $j \in \beta$.

Under certain additional assumptions on \mathcal{B} , the quiver Grassmannian $\text{Gr}_{\underline{e}}(M)$ decomposes into affine spaces (Theorem 4.1), and the parametrization of the nonempty Schubert cells yields the formula

$$\chi(\text{Gr}_{\underline{e}}(M)) = \#\{\text{extremal successor closed } \beta \subset T_0 \text{ of type } \underline{e}\},$$

see Corollary 4.4.

Content overview. To keep the technical complexity as low as possible, we restrict ourselves in this text to tree modules over the complex numbers, though the methods work in the more general context of modules of tree extensions over arbitrary rings as considered in [Lorscheid 2014]. The technique of proof in the ramified case is essentially the same as the one used in [Lorscheid 2014]. But since the presentation of our results is different and simplified, we include all details.

This text is organized as follows. In Section 1, we review basic facts about quiver Grassmannians, their Schubert decompositions and tree modules. In Section 2,

we describe generators and relations for a Schubert cell, which are labeled by *relevant pairs* and *relevant triples*, respectively. In [Section 3](#), we introduce *extremal successor closed subsets*, *polarizations* and *maximal relevant pairs*, and we establish preliminary facts. In [Section 4](#), we state the main results and conclude with several remarks and examples.

The [Appendix](#), by Thorsten Weist, shows how to establish polarizations for exceptional modules along Schofield induction.

1. Setup

To start with, let us explain the notation and terminology that we use. By a variety we understand the space of complex points of an underlying scheme, and we broadly ignore the schematic structure of quiver Grassmannians. For more details on the notions in this section, see Sections 1 and 2 of [\[Lorscheid 2014\]](#).

1A. Quiver Grassmannians. Let $Q = (Q_0, Q_1, s, t)$ be a quiver with (complex) representation $M = (\{M_i\}_{i \in Q_0}, \{M_\alpha\}_{\alpha \in Q_1})$, with dimension vector $\underline{d} = \underline{\dim} M$, and let $\underline{e} \leq \underline{d}$ be another dimension vector for Q . The *quiver Grassmannian* $\text{Gr}_{\underline{e}}(M)$ is the set of subrepresentations N of M with $\underline{\dim} N = \underline{e}$. A *basis* \mathcal{B} for M is the union $\bigcup_{p \in Q_0} \mathcal{B}_p$ of bases \mathcal{B}_p for the vector spaces M_p . An *ordered basis* of M is a basis \mathcal{B} of M whose elements b_1, \dots, b_n are linearly ordered. The choice of an ordered basis yields an inclusion

$$\text{Gr}_{\underline{e}}(M) \rightarrow \prod_{p \in Q_0} \text{Gr}(e_p, d_p)$$

that sends N to $(N_p)_{p \in Q_0}$, which endows $\text{Gr}_{\underline{e}}(M)$ with the structure of a projective variety.

1B. Schubert decompositions. A point of the Grassmannian $\text{Gr}(e, d)$ is an e -dimensional subspace V of \mathbb{C}^d . Let V be spanned by vectors $w_1, \dots, w_e \in \mathbb{C}^d$. We write $w = (w_{i,j})_{i=1 \dots d, j=1 \dots e}$ for the matrix of all coordinates of w_1, \dots, w_e . For a subset I of $\{1, \dots, d\}$ of cardinality e , the Plücker coordinates

$$\Delta_I(V) = \det(w_{i,j})_{i \in I, j=1 \dots e}$$

define a point $(\Delta_I(V))_I$ in $\mathbb{P}(\wedge^e \mathbb{C}^d)$. For two ordered subsets $I = \{i_1, \dots, i_e\}$ and $J = \{j_1, \dots, j_e\}$ of $\{1, \dots, d\}$, we define $I \leq J$ if $i_l \leq j_l$ for all $l = 1 \dots, e$. The Schubert cell $C_I(d)$ of $\text{Gr}(e, d)$ is defined as the locally closed subvariety of all subspaces V such that $\Delta_I(V) \neq 0$ and $\Delta_J(V) = 0$ for all $J > I$.

Given a quiver Q , a representation M with ordered basis \mathcal{B} and a dimension vector \underline{e} , we say that a subset β of \mathcal{B} is *of type* \underline{e} if $\beta_p = \beta \cap \mathcal{B}_p$ is of cardinality e_p for every $p \in Q_0$. For $\underline{d} = \underline{\dim} M$, the Schubert cell $C_{\beta}(\underline{d})$ is defined as the

locally closed subset $\prod_{p \in Q_0} C_{\beta_p}(d_p)$ of $\prod_{p \in Q_0} \text{Gr}(e_p, d_p)$. The *Schubert cell* C_{β}^M is defined as the intersection of $C_{\beta}(\underline{d})$ with $\text{Gr}_{\underline{e}}(M)$ inside $\prod_{p \in Q_0} C_{\beta_p}(d_p)$. The *Schubert decomposition* of $\text{Gr}_{\underline{e}}(M)$ (w.r.t. the ordered basis \mathcal{B}) is the decomposition

$$\text{Gr}_{\underline{e}}(M) = \coprod_{\substack{\beta \subset \mathcal{B} \\ \text{of type } \underline{e}}} C_{\beta}^M$$

into locally closed subvarieties. Note that the Schubert cells C_{β}^M are affine varieties, but that they are, in general, not affine spaces. In particular, a Schubert cell C_{β}^M might be empty. We say that $\text{Gr}_{\underline{e}}(M) = \coprod C_{\beta}^M$ is a *decomposition into affine spaces* if every Schubert cell C_{β}^M is either an affine space or empty.

1C. Tree modules. Let M be a representation of a quiver Q with basis \mathcal{B} . Let $\alpha : s \rightarrow t$ be an arrow of Q and $b \in \mathcal{B}_s$. Then we have the equations

$$M_{\alpha}(b) = \sum_{b' \in \mathcal{B}_t} \lambda_{\alpha, b, c} c$$

with uniquely determined coefficients $\lambda_{\alpha, b, c} \in \mathbb{C}$. The *coefficient quiver* of M w.r.t. \mathcal{B} is the quiver $T = \Gamma(M, \mathcal{B})$ with vertex set $T_0 = \mathcal{B}$ and with arrow set

$$T_1 = \{(\alpha, b, c) \in Q_1 \times \mathcal{B} \times \mathcal{B} \mid b \in \mathcal{B}_s(\alpha), c \in \mathcal{B}_t(\alpha) \text{ and } \lambda_{\alpha, b, c} \neq 0\}.$$

It comes equipped with a morphism $F : T \rightarrow Q$ that sends $b \in \mathcal{B}_p$ to p and (α, b, c) to α , and with a thin sincere representation $N = N(M, \mathcal{B})$ of T with basis \mathcal{B} and 1×1 -matrices $N_{(\alpha, b, c)} = (\lambda_{\alpha, b, c})$. Note that M is canonically isomorphic to the pushforward F_*N (see [Lorscheid 2014, Section 4]).

The representation M is called a *tree module* if there exists a basis \mathcal{B} of M such that the coefficient quiver $T = \Gamma(M, \mathcal{B})$ is a tree. We call such a basis a *tree basis* for M .

If T is a tree, we can replace the basis elements b by certain nonzero multiples b' such that all $\lambda_{\alpha, b, c}$ equal 1. We refer to this assumption by the expression $M = F_*T$ where we identify T , by abuse of notation, with its thin sincere representation with basis $T_0 = \mathcal{B}$ and matrices (1). In this case, M and \mathcal{B} are determined as the pushforward of this thin sincere representation of T along $F : T \rightarrow Q$. In general, T is not determined by M : there are examples of tree modules M and bases \mathcal{B} and \mathcal{B}' such that $\Gamma(M, \mathcal{B})$ and $\Gamma(M, \mathcal{B}')$ are nonisomorphic trees.

2. Presentations of Schubert cells

Let Q be a quiver and M a representation with ordered basis \mathcal{B} and dimension vector \underline{d} . Let \underline{e} be another dimension vector for Q and $\beta \subset \mathcal{B}$ of type \underline{e} . In this section, we will describe coordinates and relations for the Schubert cell C_{β}^M of $\text{Gr}_{\underline{e}}(M)$.

2A. Normal form for matrix representations. Let N be a point of C_β^M . Then N_p is an e_p -dimensional subspace of M_p for every $p \in Q_0$ and has a basis $(w_j)_{j \in \beta_p}$ where $w_j = (w_{i,j})_{i \in \mathcal{B}_p}$ are column vectors in M_p . If we define $w_{i,j} = 0$ for $i, j \in \mathcal{B}$ whenever $j \in \beta$, or $i \in \mathcal{B}_p$ and $j \in \mathcal{B}_q$ with $p \neq q$, then we obtain a matrix $w = (w_{i,j})_{i,j \in \mathcal{B}}$. We call w a *matrix representation* of N . Note that N is determined by the matrix representation w , but there are in general many different matrix representations of N .

We say that a matrix $w = (w_{i,j})_{i,j \in \mathcal{B}}$ in $\text{Mat}_{\mathcal{B} \times \mathcal{B}}$ is in β -normal form if

- (i) $w_{i,i} = 1$ for all $i \in \beta$,
- (ii) $w_{i,j} = 0$ for all $i, j \in \beta$ with $j \neq i$,
- (iii) $w_{i,j} = 0$ for all $i \in \mathcal{B}$ and $j \in \beta$ with $j < i$,
- (iv) $w_{i,j} = 0$ for all $i \in \mathcal{B}$ and $j \in \mathcal{B} - \beta$, and
- (v) $w_{i,j} = 0$ for all $i \in \mathcal{B}_p$ and $j \in \beta_q$ with $p \neq q$.

Lemma 2.1. Every $N \in C_\beta^M$ has a unique matrix representation $w = (w_{i,j})_{i,j \in \mathcal{B}}$ in β -normal form.

Proof. The uniqueness follows from the fact that a matrix w in β -normal form is in reduced column echelon form by (i)–(iv). The vanishing of the Plücker coordinates $\Delta_J(N_p)$ for $J > \beta_p$ and the nonvanishing of $\Delta_{\beta_p}(N_p)$ implies that we find pivot elements in the rows $i \in \beta_p$ for each $p \in Q_0$ for a matrix presentation w of N in reduced echelon form. This shows that there is a matrix presentation w of N that satisfies (i)–(iv). Since $\mathcal{B}_p \subset N_p$, the matrix w is a block matrix and satisfies (v). \square

2B. Defining equations. Lemma 2.1 identifies C_β^M with a subset of the affine matrix space $\text{Mat}_{\mathcal{B} \times \mathcal{B}}$. The following lemma determines defining equations (along with equations (i)–(v) from Section 2A) for C_β^M , which shows that C_β^M is a closed subvariety of $\text{Mat}_{\mathcal{B} \times \mathcal{B}}$.

Lemma 2.2. Let $T = \Gamma(M, \mathcal{B})$ be the coefficient quiver of M w.r.t. \mathcal{B} and $F : T \rightarrow Q$ the canonical morphism, and recall that $T_0 = \mathcal{B}$. A matrix $w = (w_{i,j})_{i,j \in \mathcal{B}}$ in β -normal form is the matrix representation of a point N of C_β^M if and only if w satisfies the equation

$$E(\bar{\alpha}, t, s) : \sum_{\substack{\alpha \in F^{-1}(\bar{\alpha}) \\ \text{with } t(\alpha)=t}} w_{s(\alpha),s} = \sum_{\alpha \in F^{-1}(\bar{\alpha})} w_{t,t(\alpha)} w_{s(\alpha),s}.$$

for all arrows $\bar{\alpha} \in Q_1$ and all vertices $s \in F^{-1}(s(\bar{\alpha}))$ and $t \in F^{-1}(t(\bar{\alpha}))$.

If $t \in \beta$ or $s \notin \beta$, then $E(\bar{\alpha}, t, s)$ is satisfied for any w in β -normal form.

Proof. Given a matrix $w = (w_{i,j})_{i,j \in \mathcal{B}}$ in β -normal form, we write w_i for the column vector $(w_{i,j})_{j \in \mathcal{B}_p}$ where $p \in Q_0$ and $i \in \mathcal{B}_p$. The matrix w represents a

point N of $\text{Gr}_{\underline{e}}(M)$ if and only if for all $\bar{\alpha} \in Q_1$ and all $s \in \beta_s(\bar{\alpha})$, there are $\lambda_k \in \mathbb{C}$ for $k \in \beta_t(\bar{\alpha})$ such that

$$M_{\bar{\alpha}} w_s = \sum_{k \in \beta_t(\bar{\alpha})} \lambda_k w_k.$$

This means that for all $t \in F^{-1}(t(\bar{\alpha}))$,

$$\sum_{\substack{\alpha \in F^{-1}(\bar{\alpha}) \\ \text{with } t(\alpha)=t}} w_{s(\alpha),s} = [M_{\bar{\alpha}} w_s]_t \text{ equals } \sum_{k \in \beta_t(\bar{\alpha})} \lambda_k w_{t,k}.$$

For $t \in \beta_t(\bar{\alpha})$, we obtain

$$\sum_{\substack{\alpha \in F^{-1}(\bar{\alpha}) \\ \text{with } t(\alpha)=t}} w_{s(\alpha),s} = \sum_{k \in \beta_t(\bar{\alpha})} \lambda_k w_{t,k} = \sum_{k \in \beta_t(\bar{\alpha})} \lambda_k \delta_{t,k} = \lambda_t$$

by (i) and (ii) for w in β -normal form. Therefore, for arbitrary $t \in F^{-1}(t(\bar{\alpha}))$ we obtain that

$$\sum_{\substack{\alpha \in F^{-1}(\bar{\alpha}) \\ \text{with } t(\alpha)=t}} w_{s(\alpha),s} = \sum_{k \in \beta_t(\bar{\alpha})} \left(\sum_{\substack{\alpha \in F^{-1}(\bar{\alpha}) \\ \text{with } t(\alpha)=k}} w_{s(\alpha),s} \right) w_{t,k} = \sum_{\alpha \in F^{-1}(\bar{\alpha})} w_{t,t(\alpha)} w_{s(\alpha),s},$$

as claimed. If $t \in \beta$, then this equation is satisfied for all w in β -normal form by the definition of the λ_k and since $w_{t,k} = \delta_{t,k}$ for $t \in \beta$. If $s \notin \beta$, then all coefficients $w_{s(\alpha),s}$ are 0, i.e., we obtain the tautological equation $0 = 0$. This proves the latter claim of the lemma. □

2C. Relevant pairs and relevant triples. A *relevant pair* is an element of the set

$$\text{Rel}^2 = \{(i, j) \in T_0 \times T_0 \mid F(i) = F(j) \text{ and } i \leq j\}$$

and a *relevant triple* is an element of the set

$$\text{Rel}^3 = \left\{ (\bar{\alpha}, t, s) \in Q_1 \times T_0 \times T_0 \mid \begin{array}{l} \text{there is an } \alpha' : s' \rightarrow t' \text{ in } T \text{ with } F(\alpha') = \bar{\alpha}, \\ F(s') = F(s), F(t') = F(t), s' \leq s \text{ and } t \leq t' \end{array} \right\}.$$

Given a matrix $w = (w_{i,j})$ in β -normal form, we say that $w_{i,j}$ is a *constant coefficient* (w.r.t. β) if it appears in one of the equations (i)–(v) from Section 2A, and otherwise we say that $w_{i,j}$ is a *free coefficient* (w.r.t. β), which is the case if and only if there is a $p \in Q_0$ such that $i \in \mathcal{B}_p - \beta_p$, $j \in \beta_p$ and $i < j$. The significance of Rel^2 is that if $w_{i,j}$ is not constant equal to 0 w.r.t. β (for any β), then (i, j) is a relevant pair.

If we substitute for a given β all constant coefficients $w_{i,j}$ with $i \neq j$ by 0, then we obtain the β -reduced form of $E(\bar{\alpha}, t, s)$:

$$\sum_{\substack{\alpha \in F^{-1}(\bar{\alpha}) \text{ with} \\ t(\alpha)=t, s(\alpha) \leq s, \\ s(\alpha) \notin \beta \text{ or } s(\alpha)=s}} w_{s(\alpha),s} = \sum_{\substack{\alpha \in F^{-1}(\bar{\alpha}) \text{ with} \\ t < t(\alpha), s(\alpha) < s, \\ t(\alpha) \in \beta, s(\alpha) \notin \beta}} w_{t,t(\alpha)} w_{s(\alpha),s} + \sum_{\substack{\alpha \in F^{-1}(\bar{\alpha}) \text{ with} \\ s(\alpha)=s, t \leq t(\alpha), \\ t(\alpha) \in \beta \text{ or } t(\alpha)=t}} w_{t,t(\alpha)}. \quad (1)$$

The significance of Rel^3 is that if $E(\bar{\alpha}, t, s)$ is a nontrivial equation in the coefficients of a matrix w in β -normal form (for any β), then $(\bar{\alpha}, t, s)$ is a relevant triple.

In the following, we will associate certain values with relevant pairs and relevant triples. Since $T_0 = \mathcal{B}$ is linearly ordered, we can identify it order-preservingly with $\{1, \dots, n\}$. We define the root of a connected component of T as its smallest vertex, and we denote by $r(i)$ the root of the component that contains the vertex i . In particular, if T is connected, then 1 is the only root and $r(i) = 1$ for all $i \in T_0$. Let $d(i, j)$ denote the graph distance of two vertices $i, j \in T_0$. We define the *root distance of a relevant pair* (i, j) as

$$\delta(i, j) = \max\{d(i, r(i)), d(j, r(j))\}.$$

We define the *fiber length of a relevant pair* (i, j) as

$$\epsilon(i, j) = \#\{k \in T_0 \mid F(k) = F(i) \text{ and } i \leq k < j\}.$$

We consider $\mathbb{N} \times \mathbb{N} \times T_0$ with its lexicographical order, i.e., $(i, j, k) < (i', j', k')$ if $i < i'$, or $i = i'$ and $j < j'$, or $i = i', j = j'$ and $k < k'$. The inclusion

$$\begin{aligned} \Psi : \text{Rel}^2 &\rightarrow \mathbb{N} \times \mathbb{N} \times T_0 \\ (i, j) &\mapsto (\epsilon(i, j), \delta(i, j), j) \end{aligned}$$


induces a linear order on Rel^2 , i.e., $(i, j) < (i', j')$ if $\Psi(i, j) < \Psi(i', j')$.

Let $(\bar{\alpha}, t, s)$ be a relevant triple. We define $\Psi(\bar{\alpha}, t, s)$ as the maximum of $\Psi(s_{\min}, s)$ and $\Psi(t, t_{\max})$, where s_{\min} is the smallest vertex that is the source of an arrow $\alpha \in F^{-1}(\bar{\alpha})$ with $t \leq t(\alpha)$ and t_{\max} is the largest vertex that is the target of an arrow $\alpha \in F^{-1}(\bar{\alpha})$ with $s(\alpha) \leq s$.

For a relevant triple $(\bar{\alpha}, t, s)$ with $t \notin \beta$ and $s \in \beta$, we define $\Psi_\beta(\bar{\alpha}, t, s)$ as $\Psi(i, j)$ where (i, j) is the largest relevant pair that appears as an index in the β -reduced form (1) of $E(\bar{\alpha}, t, s)$. Note that $E(\bar{\alpha}, t, s)$ contains at least one nontrivial term by the definition of a relevant triple. Note further that if there is an arrow $\alpha : s \rightarrow t$ in $F^{-1}(\bar{\alpha})$ and every other arrow $\alpha' \in F^{-1}(\bar{\alpha})$ satisfies either $s < s(\alpha')$ or $t(\alpha') < t$, then the only nontrivial terms in (1) are the constant coefficients $w_{s,s}$ and $w_{t,t}$. Thus in this case $\Psi_\beta(\bar{\alpha}, t, s) = \max\{\Psi(s, s), \Psi(t, t)\}$.

Since $w_{i,j} = 0$ if $j < i$ for w in β -normal form, we have $\Psi_\beta(\bar{\alpha}, t, s) \leq \Psi(\bar{\alpha}, t, s)$. In Section 3D, we consider cases in which $\Psi_\beta(\bar{\alpha}, t, s)$ and $\Psi(\bar{\alpha}, t, s)$ are equal.

Example 2.3. A good example to illustrate the roles of relevant pairs, relevant triples and the function Ψ is the following. Let M be the preinjective representation of the Kronecker quiver $Q = K(2)$ with dimension vector $(3, 4)$. Denote the two arrows of Q by $\bar{\alpha}$ and $\bar{\gamma}$. Then there exists an ordered basis \mathcal{B} of M such that the coefficient quiver $T = \Gamma(M, \mathcal{B})$ looks like



where we label the arrows by their image under F . We investigate the Schubert cell C_β^M for $\beta = \{3, 6, 7\}$. A matrix $w = (w_{i,j})_{i,j \in \mathcal{B}}$ in β -normal form has the six free coefficients $w_{1,3}, w_{2,3}, w_{4,6}, w_{5,6}, w_{4,7}, w_{5,7}$, and $w_{3,3} = w_{6,6} = w_{7,7} = 1$. All other coefficients vanish. The nontrivial equations on the free coefficients are labeled by the relevant triples $(\bar{\alpha}, 5, 3), (\bar{\alpha}, 4, 3), (\bar{\gamma}, 5, 3)$ and $(\bar{\gamma}, 4, 3)$, and their respective β -reduced forms are

$$w_{2,3} = w_{5,6}, \quad w_{1,3} = w_{4,6}, \quad w_{1,3} = w_{2,3}w_{5,6} + w_{5,7} \quad \text{and} \quad 0 = w_{2,3}w_{4,6} + w_{4,7}.$$

It is easy to see that these equations can be solved successively in linear terms. We show how these equations are organized by the ordering of Rel^2 defined by Ψ . The relevant pairs that appear as indices of free coefficients are ordered as follows:

$$(5, 6) < (2, 3) < (4, 6) < (1, 3) < (5, 7) < (4, 7).$$

Ordered by size, we have

$$\begin{aligned} \Psi_\beta(\bar{\alpha}, 5, 3) &= (2, 3), & \Psi_\beta(\bar{\alpha}, 4, 3) &= (1, 3), \\ \Psi_\beta(\bar{\gamma}, 5, 3) &= (5, 7), & \Psi_\beta(\bar{\gamma}, 4, 3) &= (4, 7), \end{aligned}$$

which correspond to the indices of linear terms in each of the corresponding equations. Therefore, we find a unique solution in $w_{2,3}, w_{1,3}, w_{5,7}$ and $w_{4,7}$ for every $w_{5,6}$ and $w_{4,6}$, which shows that C_β^M is isomorphic to \mathbb{A}^2 .

This demonstrates how the ordering of relevant pairs organizes the defining equations for C_β^M in such a way that they are successively solvable in variables that appear in linear terms. In the next section, we will develop criteria under which this example generalizes to other representations M and ordered bases \mathcal{B} .

Remark 2.4. The definition of Ψ is based on heuristics with random examples of tree modules with ordered $F : T \rightarrow Q$. It is possible that different orders of Rel^2

lead to analogues of [Theorem 4.1](#) that include quiver Grassmannians not covered in this text. Interesting variants might include the graph distance $d(i, j)$ of i and j as an ordering criterion; e.g., consider the ordering of Rel^2 given by the map $\tilde{\Psi} : \text{Rel}^2 \rightarrow \mathbb{N} \times \mathbb{N} \rightarrow T_0$ with $\tilde{\Psi}(i, j) = (d(i, j), \epsilon(i, j), j)$. This might be of particular interest for exceptional modules that do not have an ordered tree basis such that $F : T \rightarrow Q$ is ordered. See, however, [Section 4B](#) for some limiting examples.

3. Preliminaries for the main theorem

In this section, we develop the terminology and establish preliminary facts to formulate and prove the main theorem in [Section 4](#). As before, we let Q be a quiver and M a representation with ordered basis \mathcal{B} and dimension vector \underline{d} . Let \underline{e} be another dimension vector for Q and $\beta \subset \mathcal{B}$ of type \underline{e} . Let $T = \Gamma(M, \mathcal{B})$ be the coefficient quiver of M w.r.t. \mathcal{B} and $F : T \rightarrow Q$ the canonical morphism. We identify the linearly ordered set $T_0 = \mathcal{B}$ with $\{1, \dots, n\}$.

3A. Extremal successor closed subsets. An arrow $\alpha : s \rightarrow t$ in T is called *extremal (with respect to F)* if all other arrows $\alpha' : s' \rightarrow t'$ with $F(\alpha') = F(\alpha)$ satisfy either $s < s'$ or $t' < t$. Note that if F is ordered and unramified, then every arrow of T is extremal.

Recall that $T_0 = \mathcal{B}$, which allows us to consider β as a subset of T_0 . We say that β is *extremal successor closed* if for all extremal arrows $\alpha : s \rightarrow t$ of T , either $s \notin \beta$ or $t \in \beta$. Note that if F is ordered and unramified, then β is extremal successor closed if and only if β is successor closed in the sense of [\[Cerulli Irelli 2011\]](#) and [\[Haupt 2012\]](#).

Lemma 3.1. *If β is not extremal successor closed, then C_β^M is empty.*

Proof. We assume that C_β^M is nonempty and prove the lemma by contraposition. Let $\alpha : s \rightarrow t$ be an extremal arrow in T and $\bar{\alpha} = F(\alpha)$. Let $N \in C_\beta^M$ have the matrix representation w in β -normal form. The β -reduced form of $E(\bar{\alpha}, t, s)$ is

$$w_{s,s} = w_{t,t} w_{s,s}$$

since $\alpha : s \rightarrow t$ is extremal and thus for every other $\alpha' : s' \rightarrow t'$ in $F^{-1}(\bar{\alpha})$ either $s < s'$, and thus $w_{s',s} = 0$, or $t' < t$, and thus $w_{t',t} = 0$. Since $w_{s,s} = 1$ if $s \in \beta$ (by (i)) and $w_{t,t} = 0$ if $t \notin \beta$ (by (iv)), equation $E(\bar{\alpha}, t, s)$ would be $1 = 0$ if $s \in \beta$ and $t \notin \beta$. This is not possible since we assumed that C_β is nonempty. Therefore $s \notin \beta$ or $t \in \beta$, which shows that β is extremal successor closed. □

3B. Ordered and ramified morphisms. The morphism $F : T \rightarrow Q$ is *ordered* if for all arrows $\alpha : s \rightarrow t$ and $\alpha' : s' \rightarrow t'$ of T with $F(\alpha) = F(\alpha')$, we have $s \leq s'$ if and only if $t \leq t'$.

Consider an arrow $\bar{\alpha} \in Q_1$ and a vertex $i \in T_0$. The *ramification index* $r_{\bar{\alpha}}(i)$ at i in direction $\bar{\alpha}$ is the number of arrows $\alpha \in F^{-1}(\bar{\alpha})$ with source or target i . If $r_{\bar{\alpha}}(i) > 1$, we say that F branches at i in direction $\bar{\alpha}$ and that F ramifies above $F(i)$. The morphism $F : T \rightarrow Q$ is *unramified* or a *winding* if for all $\bar{\alpha} \in Q_1$ and all $i \in T_0$, we have $r_{\bar{\alpha}}(i) \leq 1$. In other words, $F : T \rightarrow Q$ is unramified if and only if the associated map of CW-complexes is unramified.

Note that F is strictly ordered (in the sense of [Lorscheid 2014, Section 4.2]) if and only if F is ordered and unramified. From this viewpoint, we can say that we extend Theorem 4.2 of [Lorscheid 2014] from unramified morphisms $F : T \rightarrow Q$ to ramified F in this text.

3C. Polarizations. Let $I = \{i_1, \dots, i_r\}$ be a finite ordered set with $i_1 < \dots < i_r$. A *sorting* of I is a decomposition $I = I^< \sqcup I^>$ such that $I^< = \{i_1, \dots, i_s\}$ and $I^> = \{i_{s+1}, \dots, i_r\}$ for some $s \in \{1, \dots, r-1\}$. A *polarization* for a linear map $M_{\bar{\alpha}} : M_p \rightarrow M_q$ (between finite dimensional complex vector spaces) are ordered bases \mathcal{B}_p and \mathcal{B}_q for M_p and M_q , respectively, that admit sortings $\mathcal{B}_p = \mathcal{B}_{p,\bar{\alpha}}^< \sqcup \mathcal{B}_{p,\bar{\alpha}}^>$ and $\mathcal{B}_q = \mathcal{B}_{q,\bar{\alpha}}^< \sqcup \mathcal{B}_{q,\bar{\alpha}}^>$ such that $M_{\bar{\alpha}}$ restricts to a surjection $\mathcal{B}_{p,\bar{\alpha}}^< \cup \{0\} \rightarrow \mathcal{B}_{q,\bar{\alpha}}^< \cup \{0\}$ and its adjoint map $M_{\bar{\alpha}}^{\text{ad}}$ restricts to a surjection $\mathcal{B}_{q,\bar{\alpha}}^> \cup \{0\} \rightarrow \mathcal{B}_{p,\bar{\alpha}}^> \cup \{0\}$. We call these decompositions of \mathcal{B}_p and \mathcal{B}_q a *sorting* for $M_{\bar{\alpha}}$.

Let M be a representation of Q . A *polarization* for M is an ordered basis \mathcal{B} of M such that \mathcal{B}_p and \mathcal{B}_q are a polarization for every arrow $\bar{\alpha} : p \rightarrow q$ in Q . In this case, we also say that M is *polarized* by \mathcal{B} . An *ordered polarization* of M is a polarization \mathcal{B} such that the canonical morphism $F : T \rightarrow Q$ from the coefficient quiver is ordered.

In other words, M is polarized by \mathcal{B} if and only if there are, for all arrows $\bar{\alpha} : p \rightarrow q$ in Q , sortings $\mathcal{B}_p = \mathcal{B}_{p,\bar{\alpha}}^< \sqcup \mathcal{B}_{p,\bar{\alpha}}^>$ and $\mathcal{B}_q = \mathcal{B}_{q,\bar{\alpha}}^< \sqcup \mathcal{B}_{q,\bar{\alpha}}^>$ such that $r_{\bar{\alpha}}(i) \leq 1$ for all $i \in \mathcal{B}_{p,\bar{\alpha}}^< \sqcup \mathcal{B}_{q,\bar{\alpha}}^>$ and $r_{\bar{\alpha}}(i) \geq 1$ for all $i \in \mathcal{B}_{q,\bar{\alpha}}^< \sqcup \mathcal{B}_{p,\bar{\alpha}}^>$. This means that the nonzero matrix coefficients of $M_{\bar{\alpha}}$ w.r.t. \mathcal{B}_p and \mathcal{B}_q can be covered by an upper left submatrix $M_{\bar{\alpha}}^<$ and a lower right submatrix $M_{\bar{\alpha}}^>$. Here $M_{\bar{\alpha}}^<$ has at most one nonzero entry in each column and at least one nonzero entry in each row; $M_{\bar{\alpha}}^>$ has at least one nonzero entry in each column and at most one nonzero entry in each row.

Figure 1 illustrates the typical shape of a fiber of an arrow $\bar{\alpha} : p \rightarrow q$ of Q in the coefficient quiver $T = \Gamma(M, \mathcal{B})$ where \mathcal{B} is an ordered polarization for M . We use the convention that we order the vertices from left to right in growing order. The property that \mathcal{B} is a polarization is visible by the number of arrows connecting




Figure 1

to a vertex in the upper left / lower left / upper right / lower right of the picture, and the property that $F : T \rightarrow Q$ is ordered is visible from the fact that the arrows do not cross each other.

Lemma 3.2. *Let \mathcal{B} be an ordered polarization for M . Let $\bar{\alpha} : p \rightarrow q$ be an arrow in Q and $\mathcal{B}_p = \mathcal{B}_{p,\bar{\alpha}}^< \sqcup \mathcal{B}_{p,\bar{\alpha}}^>$ and $\mathcal{B}_q = \mathcal{B}_{q,\bar{\alpha}}^< \sqcup \mathcal{B}_{q,\bar{\alpha}}^>$ a sorting for $M_{\bar{\alpha}}$. Then every i in $\mathcal{B}_{q,\bar{\alpha}}^< \sqcup \mathcal{B}_{p,\bar{\alpha}}^>$ connects to a unique extremal arrow.*

Proof. It is clear that every vertex i connects to at most one extremal arrow in $F^{-1}(\bar{\alpha})$. Since M is polarized by \mathcal{B} , we have that if $r_{\alpha}(i) \geq 1$, then $r_{\alpha}(j) = 1$ for all j such that there is an arrow $\alpha : i \rightarrow j$ or $\alpha : j \rightarrow i$ in $F^{-1}(\bar{\alpha})$. In case $i = s(\alpha)$, this means that $\alpha : i \rightarrow j_0$ is extremal where j_0 is minimal among the targets of arrows in $F^{-1}(\bar{\alpha})$ with source i . In case $i = t(\alpha)$, this means that $\alpha : j_0 \rightarrow i$ is extremal where j_0 is maximal among the sources of arrows in $F^{-1}(\bar{\alpha})$ with target i . This establishes the lemma. □

Remark 3.3. Ringel [2013] develops the notion of a radiation basis in order to exhibit distinguished tree bases for exceptional modules. By Proposition 3 of that paper, a radiation basis \mathcal{B} is a polarization of M (w.r.t. any ordering of \mathcal{B}). Examples of representations with radiation basis are indecomposable representations of Dynkin quivers (with an exception for E_8) and the pullback of preinjective or preprojective modules of the Kronecker quiver $K(n)$ with n arrows to its universal covering graph. Since the coefficient quiver of a pullback is the same as the coefficient quiver of the original representation, it follows that every preinjective or preprojective representation of the Kronecker quiver $K(n)$ is polarized by some ordered basis.

The Appendix gives a general strategy to establish polarizations of exceptional modules along Schofield induction. In [Lorscheid and Weist \geq 2015], we will show that every exceptional representation M of a quiver of affine Dynkin type \tilde{D}_n has a polarization which yields a Schubert decomposition of $\text{Gr}_{\underline{e}}(M)$ into affine spaces.

3D. Maximal relevant pairs. Let $\bar{\alpha} \in Q_1$. A relevant pair (i, j) is maximal for $\bar{\alpha}$ if there exists a relevant triple $(\bar{\alpha}, t, s)$ such that $\Psi(i, j) = \Psi(\bar{\alpha}, t, s)$.

Lemma 3.4. *Assume that M is polarized by \mathcal{B} and that $\beta \subset \mathcal{B}$ is extremal successor closed. Let $(\bar{\alpha}, t, s)$ be a relevant triple with $s \in \beta$ and $t \notin \beta$. Then one of the following holds.*

- (i) *There is an extremal arrow $\alpha' : s' \rightarrow t$ in $F^{-1}(\bar{\alpha})$ such that $s' \notin \beta$ and*

$$\Psi_{\beta}(\bar{\alpha}, t, s) = \Psi(s', s) = \Psi(\bar{\alpha}, t, s).$$

In this case, the β -reduced form of $E(\bar{\alpha}, t, s)$ is

$$w_{s',s} = - \sum_{\substack{\alpha \in F^{-1}(\bar{\alpha}) \text{ with} \\ t(\alpha)=t, s(\alpha) \notin \beta}} w_{s(\alpha),s} + \sum_{\substack{\alpha \in F^{-1}(\bar{\alpha}) \text{ with} \\ s' < s(\alpha) < s, \\ s(\alpha) \notin \beta, t(\alpha) \in \beta}} w_{t,t(\alpha)} w_{s(\alpha),s} + \sum_{\substack{\alpha \in F^{-1}(\bar{\alpha}) \\ \text{with } s(\alpha)=s, \text{ and} \\ t(\alpha) \in \beta \text{ or } t(\alpha)=t}} w_{t,t(\alpha)}.$$

(ii) *There is an extremal arrow $\alpha' : s \rightarrow t'$ in $F^{-1}(\bar{\alpha})$ such that $t' \in \beta$ and*

$$\Psi_\beta(\bar{\alpha}, t, s) = \Psi(t, t') = \Psi(\bar{\alpha}, t, s).$$

In this case, the β -reduced form of $E(\bar{\alpha}, t, s)$ is

$$w_{t,t'} = \sum_{\substack{\alpha \in F^{-1}(\bar{\alpha}) \\ \text{with } t(\alpha)=t, \text{ and} \\ s(\alpha) \notin \beta \text{ or } s(\alpha)=s}} w_{s(\alpha),s} - \sum_{\substack{\alpha \in F^{-1}(\bar{\alpha}) \text{ with} \\ t < t(\alpha) < t', \\ s(\alpha) \notin \beta, t(\alpha) \in \beta}} w_{t,t(\alpha)} w_{s(\alpha),s} - \sum_{\substack{\alpha \in F^{-1}(\bar{\alpha}) \text{ with} \\ s(\alpha)=s, t(\alpha) \in \beta}} w_{t,t(\alpha)}.$$

Proof. Once we know that there is an extremal arrow $\alpha' : s' \rightarrow t$ (or $\alpha' : s \rightarrow t'$), it is clear that $s' \notin \beta$ (or $t \in \beta$), that $w_{s',s}$ (or $w_{t,t'}$) is a free coefficient and that the β -reduced form of $E(\bar{\alpha}, t, s)$ looks as described in (i) (or (ii)).

If there are extremal arrows $\alpha' : s' \rightarrow t$ and $\alpha'' : s \rightarrow t''$, then s' is minimal among the sources of arrows in $F^{-1}(\bar{\alpha})$ with target t , and t'' is maximal among the targets of arrows in $F^{-1}(\bar{\alpha})$ with source s . Clearly, we have $\Psi(\bar{\alpha}, t, s) = \max\{\Psi(s', s), \Psi(t, t'')\}$. By the definition of a relevant triple, we have $s' \leq s$ and $t \leq t''$. Since β is extremal successor closed, $s' \notin \beta$ and $t'' \in \beta$. In particular, this means that $s' \neq s$ and $t \neq t''$, and thus $w_{s',s}$ and $w_{t,t''}$ are free coefficients. By the minimality of s' and the maximality of t'' , every other free coefficient $w_{i,j}$ in the β -reduced form of $E(\bar{\alpha}, t, s)$ must satisfy $\epsilon(i, j) < \max\{\epsilon(s', s), \epsilon(t, t'')\}$. Therefore also $\Psi_\beta(\bar{\alpha}, t, s) = \max\{\Psi(s', s), \Psi(t, t'')\}$, which establishes the proposition in the case that both s and t connect to extremal arrows in the fiber of $\bar{\alpha}$.

Let $p = s(\bar{\alpha})$ and $q = t(\bar{\alpha})$. Let $\mathcal{B}_p = \mathcal{B}_{p,\bar{\alpha}}^< \amalg \mathcal{B}_{p,\bar{\alpha}}^>$ and $\mathcal{B}_q = \mathcal{B}_{q,\bar{\alpha}}^< \amalg \mathcal{B}_{q,\bar{\alpha}}^>$ be sortings for $M_{\bar{\alpha}}$. If s is not the source of any extremal arrow in the fiber of $\bar{\alpha}$, then Lemma 3.2 implies that $s \in \mathcal{B}_{p,\bar{\alpha}}^<$. By the definition of a relevant triple, there is an arrow $\alpha \in F^{-1}(\bar{\alpha})$ with $s(\alpha) \leq s$ and $t(\alpha) \leq t$. This implies that $t \in \mathcal{B}_{q,\bar{\alpha}}^<$ and, by Lemma 3.2, that there is an extremal arrow $\alpha' : s' \rightarrow t$. Since β is extremely successor closed, $s' \notin \beta$ and $w_{s',s}$ is a free coefficient.

We claim that in this situation $\Psi_\beta(\bar{\alpha}, t, s) = \Psi(s', s) = \Psi(\bar{\alpha}, t, s)$. Since α' is extremal, all $s'' \in F^{-1}(p)$ appearing in an index of the β -reduced form of $E(\bar{\alpha}, t, s)$ must lie between s' and s . This means that $\epsilon(s', s)$ is larger than $\epsilon(s', s'')$ and $\epsilon(s'', s)$ if s'' is different from both s and s' . Similarly, the largest relevant pair (t'', t') with $F(t'') = F(t') = q$ satisfies $t'' = t$ and that t' is maximal among the targets of arrows in $F^{-1}(\bar{\alpha})$ whose source is less or equal to s . Since $t, t' \in \mathcal{B}_{q,\bar{\alpha}}^<$, we have $\epsilon(t, t') \leq \epsilon(s', s)$. Equality can only hold if every s'' between s' and s is the source of precisely one arrow in $F^{-1}(\bar{\alpha})$. But then there would be such a unique arrow with source s , which is necessarily extremal. Since this contradicts the assumption that there is no extremal arrow with source s in $F^{-1}(\bar{\alpha})$, we see that $\Psi(s', s) > \Psi(t, t')$. This shows that $\Psi_\beta(\bar{\alpha}, t, s) = \Psi(s', s) = \Psi(\bar{\alpha}, t, s) = \Psi(\bar{\alpha}, t, s)$, which means that (i) is satisfied.

If t is not the target of any extremal arrow in the fiber of $\bar{\alpha}$, then we conclude analogously to the previous case that there is an extremal arrow $\alpha' : s \rightarrow t'$ with $t' \in \beta$ such that $\Psi_\beta(\bar{\alpha}, t, s) = \Psi(t, t')$. Thus in this case, (ii) is satisfied. \square

Lemma 3.5. *Let $\bar{\alpha} \in Q_1$ and $(i, j) \in \text{Rel}^2$. Assume that M is polarized by \mathcal{B} .*

- (i) *If $F(i) = s(\bar{\alpha})$, then there is at most one $\alpha : i \rightarrow t$ in $F^{-1}(\bar{\alpha})$ such that $\Psi(i, j) = \Psi(\bar{\alpha}, t, j)$.*
- (ii) *If $F(j) = t(\bar{\alpha})$, then there is at most one $\alpha : s \rightarrow j$ in $F^{-1}(\bar{\alpha})$ such that $\Psi(i, j) = \Psi(\bar{\alpha}, i, s)$.*

Proof. We prove (i). If there is only one arrow α in $F^{-1}(\bar{\alpha})$ with source i , then (i) is clear. Assume that there are two different arrows $\alpha : i \rightarrow t$ and $\alpha' : i \rightarrow t'$ in $F^{-1}(\bar{\alpha})$ with $t' < t$. Since M is polarized, we have $r_{\bar{\alpha}}(k) \leq 1$ for all $k \geq i$ and $r_{\bar{\alpha}}(l) \geq 1$ for all $l \geq t'$. This means that there is an arrow $\alpha'' : j \rightarrow t''$ and that $\epsilon(t', t'') > \epsilon(t, t'') \geq \epsilon(i, j)$. An equality $\epsilon(t, t'') = \epsilon(i, j)$ is only possible if t is maximal among the targets of arrows in $F^{-1}(\bar{\alpha})$ with source i .

This shows (i). The proof of (ii) is analogous. \square

Corollary 3.6. *Assume that M is polarized by \mathcal{B} and that $\beta \subset \mathcal{B}$ is extremal successor closed. Let $\bar{\alpha} \in Q_1$. If (i, j) is maximal for $\bar{\alpha}$, then there is a unique $(\bar{\alpha}, t, s) \in \text{Rel}^3$ such that $\Psi_\beta(\bar{\alpha}, t, s) = \Psi(i, j) = \Psi(\bar{\alpha}, t, s)$. If (i, j) is not maximal for $\bar{\alpha}$, then there is no relevant triple $(\bar{\alpha}, t, s)$ with $\Psi(i, j) = \Psi(\bar{\alpha}, t, s)$.*

Proof. This is an immediate consequence of Lemmas 3.4 and 3.5. \square

4. Schubert decompositions for tree modules

Theorem 4.1. *Let M be a representation of Q and \mathcal{B} an ordered polarization for M . Let \underline{e} be a dimension vector for Q . Assume that every $(i, j) \in \text{Rel}^2$ is maximal for at most one $\bar{\alpha} \in Q_1$. Then*

$$\text{Gr}_{\underline{e}}(M) = \coprod_{\substack{\beta \subset \mathcal{B} \\ \text{of type } \underline{e}}} C_\beta^M$$

is a decomposition into affine spaces. Moreover, C_β^M is not empty if and only if β is extremal successor closed.

Proof. By Lemma 3.1, C_β^M is empty if β is not extremal successor closed. Let β be extremal successor closed. The theorem is proven once we have shown that C_β^M is an affine space. As before, we identify T_0 order-preservingly with $\{1, \dots, n\}$. For $\psi \in \mathbb{N} \times \mathbb{N} \times T_0$, we denote by $C_\beta^M(\psi)$ the solution space of all coefficients $w_{i,j}$ with $\Psi(i, j) \leq \psi$ in all equations $E(\bar{\alpha}, t, s)$ where $(\bar{\alpha}, t, s)$ is a relevant triple with $\Psi_\beta(\bar{\alpha}, t, s) < \psi$. We show by induction over $\psi \in \Psi(\text{Rel}^2)$ that $C_\beta^M(\psi)$ is an affine space. Since $\Psi(\text{Rel}^2)$ is finite, this implies that C_β^M is an affine space as required.

As base case, consider $\psi = \Psi(n, n)$. By Lemma 2.2, only those relevant triples $(\bar{\alpha}, t, s)$ with $t \notin \beta$ and $s \in \beta$ lead to nontrivial equations $E(\bar{\alpha}, t, s)$. For such a relevant triple, $\Psi_\beta(\bar{\alpha}, t, s) \leq \psi$ if and only if $E(\bar{\alpha}, t, s)$ does not contain any free coefficient and thus is of the form $w_{s,s} = w_{t,t}w_{s,s}$. This is the case if and only if there is an extremal arrow $\alpha : s \rightarrow t$ in $F^{-1}(\bar{\alpha})$. Since β is extremal successor closed, $w_{s,s} = w_{t,t}w_{s,s}$ is satisfied. This means that $C_\beta^M(\psi) = \mathbb{A}^0$ is a point.

Consider $\psi > \Psi(n, n)$ and let ψ' be its predecessor in $\Psi(\text{Rel}^2)$. We assume that $C_\beta^M(\psi')$ is an affine space. By the assumption of the theorem, (i, j) is maximal for at most one $\bar{\alpha} \in Q_1$. If there is no such $\bar{\alpha}$, then there is no relevant triple $(\bar{\alpha}, t, s)$ with $\Psi_\beta(\bar{\alpha}, t, s) = \Psi(i, j)$, which means that $w_{i,j}$ does not appear as a maximal coefficient of an equation $E(\bar{\alpha}, t, s)$. If $i \in \beta$ or $j \notin \beta$, then $w_{i,j} = 0$ and $C_\beta^M(\psi) = C_\beta^M(\psi')$. Otherwise $w_{i,j}$ is free and $C_\beta^M(\psi) = C_\beta^M(\psi') \times \mathbb{A}^1$.

If there is an arrow $\bar{\alpha} \in Q_1$ such that (i, j) is maximal for $\bar{\alpha}$, then there exists a unique relevant triple $(\bar{\alpha}, t, s)$ such that $\Psi_\beta(\bar{\alpha}, t, s) = \Psi(i, j)$ by Corollary 3.6. If $w_{i,j}$ is not free, then $i \in \beta$ or $j \notin \beta$. By Lemma 3.4, either $t = i$ and there is an extremal arrow $\alpha : s \rightarrow j$ in $F^{-1}(\bar{\alpha})$ or $s = j$ and there is an extremal arrow $\alpha : i \rightarrow t$ in $F^{-1}(\bar{\alpha})$. In either case, if $i \in \beta$ or $j \notin \beta$, then $t \in \beta$ or $s \notin \beta$ since β is extremal successor closed. This means that $E(\bar{\alpha}, t, s)$ is trivial and thus $C_\beta^M(\psi) = C_\beta^M(\psi')$. If $w_{i,j}$ is free, but $E(\bar{\alpha}, t, s)$ is trivial, then $C_\beta^M(\psi) = C_\beta^M(\psi') \times \mathbb{A}^1$. If finally $w_{i,j}$ is free and $E(\bar{\alpha}, t, s)$ is nontrivial, then $w_{i,j}$ is determined by all coefficients $w_{i',j'}$ with $\Psi(i', j') < \Psi(i, j)$ by one of the formulas in Lemma 3.4. This means that $C_\beta^M(\psi) = C_\beta^M(\psi')$.

Thus we have shown that in all possible cases, $C_\beta^M(\psi)$ equals either $C_\beta^M(\psi')$ or $C_\beta^M(\psi') \times \mathbb{A}^1$, which are both affine spaces by the inductive hypothesis. This finishes the proof of the theorem. □

Remark 4.2. Though the assumptions of Theorem 4.1 come in a different shape than Hypothesis (H) in Section 4.5 of [Lorscheid 2014], they are indeed equivalent to Hypothesis (H) if $F : Q \rightarrow T$ is unramified.

Remark 4.3. Though we do not explicitly require that \mathcal{B} is a tree basis, it follows from the other assumptions of the theorem that M is a tree module. Indeed, if the coefficient quiver T had a loop and i was the largest vertex of this loop at maximal distance to 1, then the relevant pair (i, i) would be maximal for the two connecting arrows of the loop. Note that if M is not indecomposable, then $T = \Gamma(M, \mathcal{B})$ is not necessarily connected (cf. Example 4.7).

By [Ringel 1998], every exceptional module is a tree module. But it is clear that not every exceptional module admits an ordered tree basis such that the canonical morphism $F : T \rightarrow Q$ from the coefficient quiver is ordered. For instance, there are exceptional representations of the Kronecker quiver $K(3)$ with three arrows that attest to this fact: see the example $P(x, 3)$ in [Ringel 2013, page 15].

However, if M has a radiation basis \mathcal{B} , then we can order \mathcal{B} inductively along the construction of M by smaller radiation modules in such a way that \mathcal{B} satisfies the assumptions of the theorem. In particular, this includes all exceptional representations of Dynkin type, with an exception for E_8 . We see that the class of modules that admit an ordered basis to that we can apply the theorem lies somewhere between radiation modules and tree modules.

Corollary 4.4. *Under the assumptions of Theorem 4.1, the Euler characteristic of $\text{Gr}_e(M)$ equals the number of extremal successor closed subsets $\beta \subset \mathcal{B}$ of type e .*

Proof. Since the Euler characteristic is additive under decompositions into locally closed subsets,

$$\chi(\text{Gr}_e(M)) = \sum_{\substack{\beta \subset \mathcal{B} \\ \text{of type } e}} \chi(C_\beta^M).$$


The Euler characteristic of an affine space is 1 and the Euler characteristic of the empty set is 0. Therefore the corollary follows immediately from Theorem 4.1. \square

Corollary 4.5. *If $\text{Gr}_e(M)$ is smooth and the assumptions of Theorem 4.1 are satisfied, then the closures of the nonempty Schubert cells C_β^M of $\text{Gr}_e(M)$ represent an additive basis for the cohomology ring $H^*(\text{Gr}_e(M))$. If $n = \dim \text{Gr}_e(M)$ and $d = \dim C_\beta^M$, then the class of the closure of C_β^M is in $H^{n-2d}(\text{Gr}_e(M))$.*


Proof. This follows immediately from [Lorscheid 2014, Cor. 6.2]. \square

4A. Two examples for type D_4 .

Example 4.6 (quiver Grassmannian of a ramified tree module). We give an instance of a ramified tree module to which the methods of this text apply. Let Q be the quiver




of type D_4 and let M be the exceptional module



of Q . We can order the obvious basis \mathcal{B} in such a way that the coefficient quiver T


looks like




where we label the arrows by their images under F . For the dimension vector \underline{e} with $e_x = e_z = 0$ and $e_y = e_t = 1$, we obtain precisely one subrepresentation N of M with $\underline{\dim} N = \underline{e}$. This means that $\text{Gr}_{\underline{e}}(M)$ is a point. Therefore, the Euler characteristic of $\text{Gr}_{\underline{e}}(M)$ equals 1.

There is precisely one extremal successor closed subset of type \underline{e} , namely $\beta = \{2, 3\}$, which accounts for the Euler characteristic. It is indeed easily verified that the assumptions of [Theorem 4.1](#) are satisfied. Note that β is not successor closed, which shows that the number of successor closed subsets does not coincide with the Euler characteristic in this example.

Example 4.7 (a del Pezzo surface of degree 6). The previous representation appears as a subrepresentation of the following unramified representation. This example arose from discussions with Markus Reineke. Let Q be the same quiver as in the previous example and M the representation



of Q . We can order the obvious basis \mathcal{B} in such a way that the coefficient quiver T is



where we label the arrows by their images under F . It is clear from this picture that \mathcal{B} is an ordered polarization, and it is easily verified that every relevant pair is maximal for at most one arrow. Thus [Theorem 4.1](#) implies that the nonempty Schubert cells are affine spaces and that they are indexed by the extremal successor closed subsets β of T_0 . For type $\underline{e} = (2, 1, 1, 1)$, we obtain the nonempty Schubert cells

$$\begin{aligned}
 C_{\{1,2,4,6,8\}}^M &\simeq \mathbb{A}^0, & C_{\{1,2,5,6,8\}}^M &\simeq \mathbb{A}^1, & C_{\{1,3,4,6,9\}}^M &\simeq \mathbb{A}^1, \\
 C_{\{1,3,4,7,9\}}^M &\simeq \mathbb{A}^1, & C_{\{2,3,5,7,8\}}^M &\simeq \mathbb{A}^1, & C_{\{2,3,5,7,9\}}^M &\simeq \mathbb{A}^2.
 \end{aligned}$$

Therefore the Euler characteristic of $X = \text{Gr}_{\underline{e}}(M)$ is 6 and since X is smooth (as we will see in a moment), [Corollary 4.5](#) tells us that $H^0(X) = \mathbb{Z}$, $H^1(X) = \mathbb{Z}^4$ and $H^2(X) = \mathbb{Z}$ are additively generated by the closures of the Schubert cells.

To show that X is smooth, we consider X as a closed subvariety of $\text{Gr}(2, 3) \times \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$. Note that for a subrepresentation N of M with dimension vector \underline{e} , the 1-dimensional subspaces N_x, N_y and N_z of M_x, M_y and M_z , respectively, determine the 2-dimensional subspace N_t of M_t uniquely. The images of $N_x = \langle \binom{x_0}{x_1} \rangle$, $N_y = \langle \binom{y_0}{y_1} \rangle$ and $N_z = \langle \binom{z_0}{z_1} \rangle$ in M_t lie in a plane if and only if

$$\det \begin{bmatrix} x_0 & y_0 & 0 \\ x_1 & 0 & z_0 \\ 0 & y_1 & z_1 \end{bmatrix} = -x_0 y_1 z_0 - x_1 y_0 z_1 = 0.$$

Therefore the projection $\text{Gr}(2, 3) \times \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1 \rightarrow \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ yields an isomorphism

$$\text{Gr}_{\underline{e}}(M) \xrightarrow{\sim} \{[x_0 : x_1 \mid y_0 : y_1 \mid z_0 : z_1] \in \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1 \mid x_0 y_1 z_0 + x_1 y_0 z_1 = 0\}.$$

Since there is no point in $\text{Gr}_{\underline{e}}(M)$ that vanishes for all derivatives of the defining equation, $\text{Gr}_{\underline{e}}(M)$ is smooth.

The projection $\pi_{1,3} : \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1 \rightarrow \mathbb{P}^1 \times \mathbb{P}^1$ to the first and third coordinate restricts to a surjective morphism $\pi_{1,3} : \text{Gr}_{\underline{e}}(M) \rightarrow \mathbb{P}^1 \times \mathbb{P}^1$. It is bijective outside the fibers of $[1 : 0 \mid 0 : 1]$ and $[0 : 1 \mid 1 : 0]$, and these two fibers are

$$\pi_{1,3}^{-1}([1 : 0 \mid 0 : 1]) = \{[1 : 0 \mid y_0 : y_1 \mid 0 : 1]\} \simeq \mathbb{P}^1$$

and

$$\pi_{1,3}^{-1}([0 : 1 \mid 1 : 0]) = \{[0 : 1 \mid y_0 : y_1 \mid 1 : 0]\} \simeq \mathbb{P}^1.$$

This shows that $\text{Gr}_{\underline{e}}(M)$ is the blow-up of $\mathbb{P}^1 \times \mathbb{P}^1$ in two points, which is a del Pezzo surface of degree 6. Note that the closure of the Schubert cells $C_{\{1,2,5,6,8\}}^M$, $C_{\{1,3,4,6,9\}}^M$, $C_{\{1,3,4,7,9\}}^M$ and $C_{\{2,3,5,7,8\}}^M$ are four of the six curves on $\text{Gr}_{\underline{e}}(M)$ with self-intersection -1 . In particular, the closures of the latter two cells are the two connected components of the exceptional divisor w.r.t. the blow-up $\pi_{1,3} : \text{Gr}_{\underline{e}}(M) \rightarrow \mathbb{P}^1 \times \mathbb{P}^1$.

To return to the opening remark of this example, we see that every point of $\text{Gr}_{\underline{e}}(M)$, apart from the intersection points of pairs of (-1) -curves, is a subrepresentation of M that is isomorphic to the representation of [Example 4.6](#). There are six intersection points of pairs of (-1) -curves on $\text{Gr}_{\underline{e}}(M)$, whose coordinates in $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ are


$$\begin{aligned} & [1 : 0 \mid 1 : 0 \mid 1 : 0], \quad [0 : 1 \mid 1 : 0 \mid 1 : 0], \quad [0 : 1 \mid 0 : 1 \mid 1 : 0], \\ & [0 : 1 \mid 0 : 1 \mid 0 : 1], \quad [1 : 0 \mid 0 : 1 \mid 0 : 1], \quad [1 : 0 \mid 1 : 0 \mid 0 : 1]. \end{aligned}$$

Note that each Schubert cell contains precisely one of these points, and that these points coincide with the subrepresentations N of M that are spanned by the successor closed subsets β of \mathcal{B} .

This exemplifies the idea that the Euler characteristic of a projective variety should equal the number of \mathbb{F}_1 -points. The naive definition of the \mathbb{F}_1 -points as the points with coordinates in $\mathbb{F}_1 = \{0, 1\}$ yields the right outcome in this case. The more elaborate definition of the \mathbb{F}_1 -points as the Weyl extension $\mathscr{W}(X_{\mathbb{F}_1})$ of the blue scheme $X_{\mathbb{F}_1}$ associated with $X = \text{Gr}_{\underline{e}}(M)$ and \mathcal{B} yields a intrinsic bijection between the elements of $\mathscr{W}(X_{\mathbb{F}_1})$ and the above points. This definition of \mathbb{F}_1 -points generalizes the connection between Euler characteristics and \mathbb{F}_1 -points to a larger class of quiver Grassmannians than the naive definition. See [Lorscheid 2013, Section 4] for more details.

4B. Limiting examples. As already mentioned in Remark 2.4, there are different possible choices to order Rel^2 , which might lead to different generalities of analogues of Theorem 4.1. The following examples show, however, that we cannot simply drop an assumption in Theorem 4.1.

Example 4.8 (nonordered F). Consider the representation $M = \left[\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} : \mathbb{C}^2 \rightarrow \mathbb{C}^2 \right]$ of the quiver $Q = [\bullet \rightarrow \bullet]$. With the obvious choice of the ordered basis $\mathcal{B} = \{1, 2, 3, 4\}$ of M , the coefficient quiver $T = \Gamma(M, \mathcal{B})$ looks as follows:



The Schubert cells in the decomposition

$$\text{Gr}_{(1,1)}(M) = C_{\{1,3\}}^M \amalg C_{\{1,4\}}^M \amalg C_{\{2,3\}}^M \amalg C_{\{2,4\}}^M$$


are easily determined to be

$$C_{\{1,3\}}^M = \emptyset, \quad C_{\{1,4\}}^M \simeq \mathbb{A}^0, \quad C_{\{2,3\}}^M \simeq \mathbb{A}^0 \quad \text{and} \quad C_{\{2,4\}}^M \simeq \mathbb{G}_m.$$

In this example, we come across a Schubert cell that is isomorphic to $\mathbb{G}_m = \mathbb{A}^1 - \mathbb{A}^0$. Theorem 4.1 does indeed not apply since $F : T \rightarrow Q$ is not ordered. However, the other conditions of Theorem 4.1 are satisfied: \mathcal{B} is a polarization and every relevant pair is maximal for at most one arrow (since Q has only one arrow).

Note that the indices of the nonempty Schubert cells are precisely the extremely successor closed subsets $\beta \subset \mathcal{B}$ of type \underline{e} . However, only $\{1, 4\}$ and $\{2, 3\}$ contribute to the Euler characteristic of $\text{Gr}_{\underline{e}}(M) \simeq \mathbb{P}^1$, which is 2. These two subsets are precisely the successor closed subsets of \mathcal{B} , in coherence with the methods of [Cerulli Irelli 2011] and [Haupt 2012], which apply to this example.

Example 4.9 (nonpolarized basis). Consider the representation $M = \left[\begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} : \mathbb{C}^2 \rightarrow \mathbb{C}^2 \right]$ of the quiver $Q = [\bullet \rightarrow \bullet]$. With the obvious choice of ordered basis $\mathcal{B} = \{1, 2, 3, 4\}$ of M , the coefficient quiver $T = \Gamma(M, \mathcal{B})$ looks as follows:



The Schubert cells in the decomposition

$$\text{Gr}_{(1,1)}(M) = C_{\{1,3\}}^M \amalg C_{\{1,4\}}^M \amalg C_{\{2,3\}}^M \amalg C_{\{2,4\}}^M$$

are easily determined to be

$$C_{\{1,3\}}^M = \emptyset, \quad C_{\{1,4\}}^M \simeq \mathbb{A}^0, \quad C_{\{2,3\}}^M \simeq \mathbb{A}^0 \quad \text{and} \quad C_{\{2,4\}}^M \simeq \mathbb{G}_m.$$

The Schubert cell $C_{\{2,4\}}^M \simeq \mathbb{G}_m$ does not contradict [Theorem 4.1](#) since \mathcal{B} is not a polarization, though the canonical morphism $F : T \rightarrow Q$ is ordered and every relevant pair is maximal for at most one arrow (as Q has only one arrow).

Acknowledgements. I would like to thank Dave Anderson, Giovanni Cerulli Irelli, Markus Reineke, Cecília Salgado and Jan Schroer for helpful discussions. I would like to thank Thorsten Weist for including his ideas on polarizations as an appendix to this text.

Appendix: Tree modules with polarizations by Thorsten Weist

Let Q be a quiver without loops and oriented cycles. The aim of this appendix is to investigate under which conditions we can construct indecomposable tree modules X such that the basis \mathcal{B} of the respective coefficient quiver $T_X := \Gamma(X, \mathcal{B})$ is a polarization for X . In many cases, the question whether there exists a polarization for X is closely related to the question whether there exists a coefficient quiver without a subdiagram of the form

$$s_1 \xrightarrow{a} t_1 \xleftarrow{a} s_2 \xrightarrow{a} t_2.$$

We call a coefficient quiver without such a subdiagram a weak polarization for X . Clearly, a polarization does not have such a subdiagram. But we will see that in many cases these two conditions are already equivalent, for instance for exceptional representations. In the following, we will not always distinguish between an arrow a of the coefficient quiver and its color $F(a)$. Moreover, we will often label the arrows of the coefficient quiver by its color.

One of the main tools which can be used to construct tree modules is Schofield induction, see [Schofield 1991] and [Weist 2012] for an application to tree modules. A direct consequence is that, fixing an exceptional sequence (Y, X) with $\text{Hom}(X, Y) = 0$ and a basis (e_1, \dots, e_m) of $\text{Ext}(X, Y)$, representations appearing as the middle terms of exact sequences

$$0 \rightarrow Y^e \rightarrow Z \rightarrow X^d \rightarrow 0$$

give rise to a full subcategory $\mathcal{F}(X, Y)$ of $\text{Rep}(Q)$, the category of representations of Q . Moreover, we obtain that $\mathcal{F}(X, Y)$ is equivalent to the category of representations of the generalized Kronecker quiver $K(m)$ with $K(m)_0 = \{q_0, q_1\}$ and $K(m)_1 = \{\rho_i : q_0 \rightarrow q_1 \mid i \in \{1, \dots, m\}\}$ where $m = \dim \text{Ext}(X, Y)$. Fixing a real root α of Q , we denote by X_α the indecomposable representation of dimension α , which is unique up to isomorphism. By Schofield induction, we also know that, if α is an exceptional root of Q , there already exist exceptional roots β and γ such that $X_\beta \in X_\gamma^\perp$, $\text{Hom}(X_\beta, X_\gamma) = 0$ and $\alpha = \beta^d + \gamma^e$, where (d, e) is a real root of the generalized Kronecker quiver $K(\dim \text{Ext}(X_\beta, X_\gamma))$.

Let X and Y be two representations of a quiver Q . Then we can consider the linear map

$$\gamma_{X,Y} : \bigoplus_{q \in Q_0} \text{Hom}_k(X_q, Y_q) \rightarrow \bigoplus_{a:s \rightarrow t \in Q_1} \text{Hom}_k(X_s, Y_t)$$

defined by $\gamma_{X,Y}((f_q)_{q \in Q_0}) = (Y_a f_s - f_t X_a)_{a:s \rightarrow t \in Q_1}$.

It is well-known that we have $\ker(\gamma_{X,Y}) = \text{Hom}(X, Y)$ and $\text{coker}(\gamma_{X,Y}) = \text{Ext}(X, Y)$. The first statement is straightforward. The second statement follows because every morphism $f \in \bigoplus_{a:s \rightarrow t \in Q_1} \text{Hom}_k(X_s, Y_t)$ defines an exact sequence

$$0 \rightarrow Y \rightarrow \left((Y_q \oplus X_q)_{q \in Q_0}, \left(\left(\begin{matrix} Y_a & f_a \\ 0 & X_a \end{matrix} \right) \right)_{a \in Q_1} \right) \rightarrow X \rightarrow 0$$

with the canonical inclusion on the left hand side and the canonical projection on the right hand side.

Assume that the representations X and Y are tree modules and let $T_X = \Gamma(X, \mathcal{B}_X)$ and $T_Y = \Gamma(Y, \mathcal{B}_Y)$ be the corresponding coefficient quivers. Let $x = \underline{\dim} X$, $y = \underline{\dim} Y$. Fixing a vertex q , from now on we will denote the corresponding vertices of the coefficient quivers by $(\mathcal{B}_X)_q = \{b_1^q, \dots, b_{x_q}^q\}$ and $(\mathcal{B}_Y)_q = \{c_1^q, \dots, c_{y_q}^q\}$. Let $e_{k,l}^a$, where $a : s \rightarrow t \in Q_1$, $k = 1, \dots, x_s$ and $l = 1, \dots, y_t$, be the canonical basis of $\bigoplus_{a:s \rightarrow t} \text{Hom}_k(X_s, Y_t)$ with respect to \mathcal{B}_X and \mathcal{B}_Y , i.e., $e_{k,l}^a(b_i^s) = \delta_{i,k} \delta_{j,l} c_j^t$.

This means that the coefficient quiver $\Gamma(Z, \mathcal{B}_X \cup \mathcal{B}_Y)$ of the middle term of the exact sequence

$$E(e_{k,l}^a) : 0 \rightarrow Y \rightarrow Z \rightarrow X \rightarrow 0$$

is obtained by adding an extra arrow with color a from b_k^s to c_l^t to $T_X \cup T_Y$.

Following [Weist 2012] we call a basis of $\mathcal{E}(X, Y)$ of $\text{Ext}(X, Y)$, which solely consists of elements of the form $\overline{e_{i_k, j_k}^{a_k}}$ with $k = 1, \dots, \dim \text{Ext}(X, Y)$, $a_k \in \mathcal{Q}_1$, $1 \leq i_k \leq x_s$ and $1 \leq j_k \leq y_t$, tree-shaped. In abuse of notation, we will not always distinguish between $\overline{e_{i_k, j_k}^{a_k}}$ and $e_{i_k, j_k}^{a_k}$.

Let X be a tree module. For a vertex b_i^s and an arrow $a : s \rightarrow t \in \mathcal{Q}_1$ we define

$$N(a, b_i^s) := \{b_j^t \in (T_X)_0 \mid b_i^s \xrightarrow{a} b_j^t \in (T_X)_1\}.$$

Analogously, we define $N(a, b_i^t)$. If T_X is a weak polarization for X , we say that it is strict if we have, for all arrows $s \xrightarrow{a} t \in \mathcal{Q}_1$, that $|N(a, b_i^s)| \leq 1$ for all $1 \leq i \leq x_s$ or $|N(a, b_i^t)| \leq 1$ for all $1 \leq i \leq x_t$. Clearly, a weak polarization which is strict is a polarization as defined in Section 3C. Note that we can always assume that \mathcal{B} is ordered.

For a vertex q of T_X let $S(q) = \{F(a) \in \mathcal{Q}_1 \mid a \in (T_X)_1, s(a) = q\}$ and $T(q) = \{F(a) \in \mathcal{Q}_1 \mid a \in (T_X)_1, t(a) = q\}$.

Lemma A.1. *Let X be a tree module with coefficient quiver T_X such that for every $a \in \mathcal{Q}_1$, the map X_a is of maximal rank. Then T_X is a polarization if and only if T_X is a weak polarization.*

Proof. Since X_a is of maximal rank, X_a is either surjective or injective. Thus if, in addition, T_X is a weak polarization, this means that $|N(a, b_i^s)| \leq 1$ for all $1 \leq i \leq x_s$ or $|N(a, b_i^t)| \leq 1$ for all $1 \leq i \leq x_t$. It follows that T_X is a polarization. \square

Remark A.2. For general representations of a fixed dimension, and thus in particular for exceptional representations, it is true that all linear maps appearing are of maximal rank.

Using the notation from above we introduce the following definition:

Definition A.3. (i) Let X and Y be two tree modules with coefficient quivers T_X and T_Y . Moreover, let $\mathcal{E}(X, Y) = (e_{i_k, j_k}^{a_k})_k$ with $s_k \xrightarrow{a_k} t_k \in \mathcal{Q}_1$ be a tree-shaped basis of $\text{Ext}(X, Y)$, i.e., $e_{i_k, j_k}^{a_k} (b_{i_k}^{s_k}) = c_{j_k}^{t_k}$. Then we call $\mathcal{E}(X, Y)$ a polarization if

- (a) we have that $a_k \notin S(b_{i_k}^{s_k})$ or $a_k \notin T(c_{j_k}^{t_k})$ for all k ,
- (b) if $a_k = a_l$ and $b_{i_k}^{s_k} = b_{i_l}^{s_l}$ (resp. $c_{j_k}^{t_k} = c_{j_l}^{t_l}$) for $k \neq l$, we have $a_k \notin T(c_{j_k}^{t_k})$ (resp. $a_k \notin S(b_{i_k}^{s_k})$),
- (c) for all $b_{i_k}^{s_k} \xrightarrow{a_k} b_j^t \in (T_X)_1$ we have $|N(a_k, b_j^t)| = 1$ and for all $c_i^s \xrightarrow{a_k} c_{j_k}^{t_k} \in (T_Y)_1$ we have $|N(a_k, c_i^s)| = 1$.

(ii) If we have $a_k \notin S(b_{i_k}^{s_k})$ and $a_k \notin T(c_{j_k}^{t_k})$ for all k in the first condition and if we also have $a_k \neq a_l$ if $k \neq l$, we say that the basis is a strong polarization.

Remark A.4. Roughly speaking condition (c) ensures that $b_{i_k}^{s_k}$ is the only neighbor which is connected to b_j^t by an arrow with color a_k .

Condition (a) means that either $b_{i_k}^{s_k}$ is not the source of an arrow with color a_k (when only the coefficient quiver T_X is considered) or $c_{j_k}^{t_k}$ is not the target of an arrow with color a_k (when only the coefficient quiver T_Y is considered). In particular, if we have $a_k \notin S(b_{i_k}^{s_k})$ and $a_k \notin T(c_{j_k}^{t_k})$ for all k in the first condition, the second and third conditions are clearly satisfied.

Now we are in a position to state under which conditions an exceptional sequence together with a tree-shaped basis of the Ext-group gives rise to indecomposable representations such that, in addition, there exists a coefficient quiver which is a (weak) polarization:

Theorem A.5. *Let (Y, X) be an exceptional sequence (of tree modules) such that the coefficient quivers T_X and T_Y are weak polarizations. Moreover, let $\mathcal{E}(X, Y) = (e_{i_1, j_1}^{a_1}, \dots, e_{i_m, j_m}^{a_m})$ be a basis of $\text{Ext}(X, Y)$ which is a polarization and let M be an indecomposable tree module of $K(m)$.*

- (i) *If T_M is unramified, then the induced coefficient quiver T_Z of the middle term Z of the corresponding exact sequence*

$$e_M : 0 \rightarrow Y^e \rightarrow Z \rightarrow X^d \rightarrow 0$$

is a weak polarization for Z . Moreover, Z is indecomposable.

- (ii) *If the polarization of the basis is strong and T_M is a weak polarization, then the induced coefficient quiver T_Z of the middle term Z of the corresponding exact sequence*

$$e_M : 0 \rightarrow Y^e \rightarrow Z \rightarrow X^d \rightarrow 0$$

is a weak polarization for Z . Moreover, Z is indecomposable.

- (iii) *If X_a is injective (resp. surjective) if and only if Y_a is injective (resp. surjective) for all arrows $a \in Q_1$, then T_Z is a weak polarization if and only if T_Z is a polarization.*
- (iv) *If M , and thus also Z , is exceptional, the polarization is strict and thus T_Z is a polarization for Z .*

Proof. By simply counting arrows and vertices of the induced coefficient quiver T_Z it follows that Z is a tree module, see also [Weist 2012, Proposition 3.9]. Moreover, since M is indecomposable, by Schofield induction we know that Z is indecomposable.

Thus we only need to check that T_Z is a weak polarization for Z . We first consider the case when T_M is unramified and $\mathcal{E}(X, Y)$ not necessarily a strong polarization. Clearly, in this case T_M is a weak polarization for M . Moreover, note that, since $\mathcal{E}(X, Y)$ is a basis, if $a_k = a_l$ for $k \neq l$, we either have $j_k \neq j_l$ or $i_k \neq i_l$.

The coefficient quiver could contradict the polarization property if $b_{i_k}^{s_k} = b_{i_l}^{s_l}$ (resp. $c_{j_k}^{t_k} = c_{j_l}^{t_l}$), $a_k = a_l$ for $l \neq k$ and $a_k \in T(c_{j_k}^{t_k}) \cap (T_Y)_1$ (resp. $a_k \in S(b_{i_k}^{s_k}) \cap (T_X)_1$). But this is not possible because $\mathcal{E}(X, Y)$ is a polarization. Indeed, this would contradict condition (b).

Another possibility for T_Z not being a weak polarization is if T_M had a subdiagram

$$b_j^{q_0} \xrightarrow{a_i} b_k^{q_1} \xleftarrow{a_i} b_l^{q_0} \text{ or } b_j^{q_1} \xleftarrow{a_i} b_k^{q_0} \xrightarrow{a_i} b_l^{q_1}$$

for some $i \in \{1, \dots, m\}$. But since T_M is unramified, this is not possible.

The last possibility for T_Z not being a weak polarization is if the basis contradicts condition (c) of [Definition A.3](#).

Next we consider the case that the polarization is strong, the representation M is a weak polarization and the representation is not forced to be unramified. But in this case it is straightforward to check that the induced coefficient quiver is a weak polarization. Indeed, for two basis elements $a_k : b_{i_k}^{s_k} \rightarrow c_{j_k}^{t_k}$ and $a_l : b_{i_l}^{s_l} \rightarrow c_{j_l}^{t_l}$ with $k \neq l$, we have $a_k \neq a_l$ and, moreover, considering the original coefficient quiver T_X and T_Y , we have $|N(a_r, q)| = 0$ for $q \in \{b_{i_k}^{s_k}, c_{j_k}^{t_k}, b_{i_l}^{s_l}, c_{j_l}^{t_l}\}$ and $r \in \{k, l\}$. Thus all subdiagrams which could prevent T_Z from being a weak polarization are forced to be induced from T_M . But since T_M is a weak polarization, this cannot happen.

The third claim is straightforward because, in general, for an exact sequence $e \in \text{Ext}(X, Y)$ with middle term Z , the matrix Z_a is a block matrix with diagonal blocks X_a and Y_a for every arrow.

The last claim follows from [Lemma A.1](#), see also [Remark A.2](#). □


Remark A.6. (i) If we are only interested in (weak) polarizations, we can drop the condition that X and Y are exceptional. But in this case it is far more complicated or even impossible to say anything concerning the indecomposability of Z .

(ii) If Q is of extended Dynkin type and, moreover, (Y, X) is an exceptional sequence, we have $\dim \text{Ext}(X, Y) \leq 2$ because otherwise there would be a root d of Q having an n -parameter family of indecomposables for $n \geq 2$. Then things become easier because every indecomposable tree module of $K(2)$ is unramified.

Let $S(n)$ be the n -subspace quiver with vertices $S(n)_0 = \{q_0, q_1, \dots, q_n\}$ and arrows $S(n)_1 = \{q_i \xrightarrow{a_i} q_0 \mid i = 1, \dots, n\}$. Let us consider two examples:


Example A.7. First let $n = 4$ and consider the exceptional sequence induced by the roots $\alpha = (2, 1, 1, 1, 0)$ and $\beta = (0, 0, 0, 0, 1)$. Then coefficient quivers of X_α ,

X_β and a basis of $\text{Ext}(X_\beta, X_\alpha)$ are for instance given by



Here the dotted arrows correspond to the tree-shaped basis of $\text{Ext}(X_\beta, X_\alpha)$ under consideration, whence the remaining vertices and arrows correspond to the two coefficient quivers.

Since the basis of $\text{Ext}(X_\beta, X_\alpha)$ is a polarization, which is not strong, and since we have $\dim \text{Ext}(X_\beta, X_\alpha) \leq 2$, the first part of [Theorem A.5](#) applies. For instance, considering the exceptional representation of dimension $(1, 2)$ of $K(2)$, we obtain



on the $S(4)$ -side. This is obviously a (strict) polarization.

Example A.8. An example for a basis which is a strong polarization can be obtained when considering $S(n)$ with $n \geq 3$ and the exceptional sequence induced by the roots $\alpha = (1, 1, 0, \dots, 0)$ and $\beta = (1, 0, 1, \dots, 1)$. In this case such a basis of $\text{Ext}(X_\beta, X_\alpha)$ is given by choosing $n - 2$ out of the $n - 1$ maps mapping the one-dimensional subspace $(X_\beta)_{q_i}$ to $(X_\alpha)_{q_0}$ for $i = 2, \dots, n$.

References

- [Białynicki-Birula 1973] A. Białynicki-Birula, “Some theorems on actions of algebraic groups”, *Ann. of Math.* (2) **98** (1973), 480–497. [MR 51 #3186](#) [Zbl 0275.14007](#)
- [Caldero and Chapoton 2006] P. Caldero and F. Chapoton, “Cluster algebras as Hall algebras of quiver representations”, *Comment. Math. Helv.* **81**:3 (2006), 595–616. [MR 2008b:16015](#) [Zbl 1119.16013](#)
- [Caldero and Reineke 2008] P. Caldero and M. Reineke, “On the quiver Grassmannian in the acyclic case”, *J. Pure Appl. Algebra* **212**:11 (2008), 2369–2380. [MR 2009f:14102](#) [Zbl 1153.14032](#)
- [Cerulli Irelli 2011] G. Cerulli Irelli, “Quiver Grassmannians associated with string modules”, *J. Algebraic Combin.* **33**:2 (2011), 259–276. [MR 2012c:16043](#) [Zbl 1243.16013](#)
- [Fomin et al. 2008] S. Fomin, M. Shapiro, and D. Thurston, “Cluster algebras and triangulated surfaces. I. Cluster complexes”, *Acta Math.* **201**:1 (2008), 83–146. [MR 2010b:57032](#)

- [Haupt 2012] N. Haupt, “Euler characteristics of quiver Grassmannians and Ringel–Hall algebras of string algebras”, *Algebr. Represent. Theory* **15**:4 (2012), 755–793. [MR 2944441](#) [Zbl 1275.16016](#)
- [Lorscheid 2013] O. Lorscheid, “A blueprinted view on \mathbb{F}_1 -geometry”, 2013. To appear in *Absolute arithmetic and \mathbb{F}_1 -geometry*, edited by K. Thas, Eur. Math. Soc., Zürich. [arXiv 1301.0083](#)
- [Lorscheid 2014] O. Lorscheid, “On Schubert decompositions of quiver Grassmannians”, *J. Geom. Phys.* **76** (2014), 169–191. [MR 3144366](#) [Zbl 1297.13027](#)
- [Lorscheid and Weist \geq 2015] O. Lorscheid and T. Weist, “Quiver Grassmannians of type \tilde{D}_n ”, Split into [arXiv 1507.00392](#) and [arXiv 1507.00395](#).
- [Musiker et al. 2013] G. Musiker, R. Schiffler, and L. Williams, “Bases for cluster algebras from surfaces”, *Compos. Math.* **149**:2 (2013), 217–263. [MR 3020308](#) [Zbl 1263.13024](#)
- [Qiu and Zhou 2013] Y. Qiu and Y. Zhou, “Cluster categories for marked surfaces: punctured case”, 2013. [arXiv 1311.0010](#)
- [Ringel 1998] C. M. Ringel, “Exceptional modules are tree modules”, pp. 471–493 in *Proceedings of the Sixth Conference of the International Linear Algebra Society* (Technical University of Chemnitz, 1996), vol. 275/276, edited by S. Kirkland et al., 1998. [MR 2000c:16020](#) [Zbl 0964.16014](#)
- [Ringel 2013] C. M. Ringel, “Distinguished bases of exceptional modules”, pp. 253–274 in *Algebras, quivers and representations*, edited by A. B. Buan et al., Abel Symp. **8**, Springer, Heidelberg, 2013. [MR 3183888](#)
- [Schofield 1991] A. Schofield, “Semi-invariants of quivers”, *J. London Math. Soc.* (2) **43**:3 (1991), 385–395. [MR 92g:16019](#) [Zbl 0779.16005](#)
- [Weist 2012] T. Weist, “Tree modules”, *Bull. Lond. Math. Soc.* **44**:5 (2012), 882–898. [MR 2975149](#) [Zbl 1279.16015](#)

Communicated by Mikhail Kapranov

Received 2013-08-22

Revised 2015-02-23

Accepted 2015-06-17

oliver@impa.br

Instituto Nacional de Matemática Pura e Aplicada, Estrada Dona Castorina 110, 22460-320 Rio de Janeiro-RJ, Brazil

weist@uni-wuppertal.de

Bergische Universität Wuppertal, Gausstr. 20, 42097 Wuppertal, Germany

Algebra & Number Theory

msp.org/ant

EDITORS

MANAGING EDITOR

Bjorn Poonen
Massachusetts Institute of Technology
Cambridge, USA

EDITORIAL BOARD CHAIR

David Eisenbud
University of California
Berkeley, USA

BOARD OF EDITORS

Georgia Benkart	University of Wisconsin, Madison, USA	Susan Montgomery	University of Southern California, USA
Dave Benson	University of Aberdeen, Scotland	Shigefumi Mori	RIMS, Kyoto University, Japan
Richard E. Borcherds	University of California, Berkeley, USA	Raman Parimala	Emory University, USA
John H. Coates	University of Cambridge, UK	Jonathan Pila	University of Oxford, UK
J-L. Colliot-Thélène	CNRS, Université Paris-Sud, France	Anand Pillay	University of Notre Dame, USA
Brian D. Conrad	Stanford University, USA	Victor Reiner	University of Minnesota, USA
Hélène Esnault	Freie Universität Berlin, Germany	Peter Sarnak	Princeton University, USA
Hubert Flenner	Ruhr-Universität, Germany	Joseph H. Silverman	Brown University, USA
Sergey Fomin	University of Michigan, USA	Michael Singer	North Carolina State University, USA
Edward Frenkel	University of California, Berkeley, USA	Vasudevan Srinivas	Tata Inst. of Fund. Research, India
Andrew Granville	Université de Montréal, Canada	J. Toby Stafford	University of Michigan, USA
Joseph Gubeladze	San Francisco State University, USA	Ravi Vakil	Stanford University, USA
Roger Heath-Brown	Oxford University, UK	Michel van den Bergh	Hasselt University, Belgium
Craig Huneke	University of Virginia, USA	Marie-France Vignéras	Université Paris VII, France
Kiran S. Kedlaya	Univ. of California, San Diego, USA	Kei-Ichi Watanabe	Nihon University, Japan
János Kollár	Princeton University, USA	Efim Zelmanov	University of California, San Diego, USA
Yuri Manin	Northwestern University, USA	Shou-Wu Zhang	Princeton University, USA
Philippe Michel	École Polytechnique Fédérale de Lausanne		

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

See inside back cover or msp.org/ant for submission instructions.

The subscription price for 2015 is US \$255/year for the electronic version, and \$440/year (+\$55, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscribers address should be sent to MSP.

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

ANT peer review and production are managed by EditFLOW[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2015 Mathematical Sciences Publishers

Algebra & Number Theory

Volume 9 No. 6 2015

Bivariant algebraic cobordism	1293
JOSÉ LUIS GONZÁLEZ and KALLE KARU	
Schubert decompositions for quiver Grassmannians of tree modules	1337
OLIVER LORSCHIED	
Noncommutative geometry and Painlevé equations	1363
ANDREI OKOUNKOV and ERIC RAINS	
Electrical networks and Lie theory	1401
THOMAS LAM and PAVLO PYLYAVSKYY	
The Kac–Wakimoto character formula for the general linear Lie superalgebra	1419
MICHAEL CHMUTOV, CRYSTAL HOYT and SHIFRA REIF	
Effective Matsusaka’s theorem for surfaces in characteristic p	1453
GABRIELE DI CERBO and ANDREA FANELLI	
Adams operations and Galois structure	1477
GEORGIOS PAPPAS	