

Algebra & Number Theory

Volume 9

2015

No. 6

**Effective Matsusaka's theorem
for surfaces in characteristic p**

Gabriele Di Cerbo and Andrea Fanelli



Effective Matsusaka's theorem for surfaces in characteristic p

Gabriele Di Cerbo and Andrea Fanelli

We obtain an effective version of Matsusaka's theorem for arbitrary smooth algebraic surfaces in positive characteristic, which provides an effective bound on the multiple that makes an ample line bundle D very ample. The proof for pathological surfaces is based on a Reider-type theorem. As a consequence, a Kawamata–Viehweg-type vanishing theorem is proved for arbitrary smooth algebraic surfaces in positive characteristic.

1. Introduction	1453
2. Preliminary results	1455
3. An effective Matsusaka's theorem	1460
4. Effective very-ampleness in positive characteristic	1462
5. A Kawamata–Viehweg-type vanishing theorem in positive characteristic	1468
Acknowledgements	1473
References	1474

1. Introduction

A celebrated theorem of Matsusaka [1972] states that for a smooth n -dimensional complex projective variety X and an ample divisor D on it, there exists a positive integer M , depending only on the Hilbert polynomial $\chi(X, \mathcal{O}_X(kD))$, such that mD is very ample for all $m \geq M$. Kollár and Matsusaka [1983] improved the result, showing that the integer M only depends on the intersection numbers (D^n) and $(K_X \cdot D^{n-1})$.

The first effective versions of this result are due to Siu [2002a; 2002b] and Demailly [1996a; 1996b]; their methods are cohomological and rely on vanishing theorems. See also [Lazarsfeld 2004b] for a full account of this approach.

Although the minimal model program for surfaces in positive characteristic has recently been established, thanks to the work of Tanaka [2014; 2012], some

MSC2010: 14J25.

Keywords: effective Matsusaka, surfaces in positive characteristic, Fujita's conjectures, Bogomolov's stability, Reider's theorem, bend-and-break, effective Kawamata–Viehweg vanishing.

interesting effectivity questions remain open in this setting, after the influential papers [Ekedahl 1988] and [Shepherd-Barron 1991a].

The purpose of this paper is to present a complete solution for the following problem:

Question 1.1. Let X be a smooth surface over an algebraically closed field of positive characteristic, and let D and B be an ample and a nef divisor on X respectively. Then there exists an integer M depending only on (D^2) , $(K_X \cdot D)$ and $(D \cdot B)$ such that

$$mD - B$$

is very ample for all $m \geq M$.

The analogous question in characteristic zero with $B = 0$ was totally solved in [Fernández del Busto 1996], and a modified technique allows one to partially extend the result in positive characteristic [Ballico 1996].

The main result of this paper is the following:

Theorem 1.2. *Let D and B be respectively an ample divisor and a nef divisor on a smooth surface X over an algebraically closed field k , with $\text{char } k = p > 0$. Then $mD - B$ is very ample for any*

$$m > \frac{2D \cdot (H + B)}{D^2} ((K_X + 2D) \cdot D + 1),$$

where:

- $H := K_X + 4D$ if X is neither quasielliptic with $\kappa(X) = 1$ nor of general type.
- $H := K_X + 8D$ if X is quasielliptic with $\kappa(X) = 1$ and $p = 3$.
- $H := K_X + 19D$ if X is quasielliptic with $\kappa(X) = 1$ and $p = 2$.
- $H := 2K_X + 4D$ if X is of general type and $p \geq 3$.
- $H := 2K_X + 19D$ if X is of general type and $p = 2$.

The effective bound obtained with $H = K_X + 4D$ is expected to hold for all surfaces. Note that this bound is not far from being sharp even in characteristic zero [Fernández del Busto 1996].

The proof of Theorem 1.2 does not rely directly on vanishing theorems, but rather on Fujita's conjecture on basepoint-freeness and very-ampleness of adjoint divisors, which is known to hold for smooth surfaces in characteristic zero [Reider 1988] and for smooth surfaces in positive characteristic which are neither quasielliptic with $\kappa(X) = 1$ nor of general type [Shepherd-Barron 1991a; Terakawa 1999].

Conjecture 1.3 (Fujita). *Let X be a smooth n -dimensional projective variety and let D be an ample divisor on it. Then $K_X + kD$ is basepoint free for $k \geq n + 1$ and very ample for $k \geq n + 2$.*

If Fujita’s conjecture on very-ampleness holds then the bound of Theorem 1.2 with $H = K_X + 4D$ would work for arbitrary smooth surfaces in positive characteristic.

For surfaces which are quasielliptic with $\kappa(X) = 1$ or of general type we can prove the following effective result in the spirit of Fujita’s conjecture (see Section 4):

Theorem 1.4. *Let X a smooth surface over an algebraically closed field of characteristic $p > 0$, let D an ample Cartier divisor on X and let $L(a, b) := aK_X + bD$ for positive integers a and b . Then $L(a, b)$ is very ample for the following values of a and b :*

- (1) *If X is quasielliptic with $\kappa(X) = 1$ and $p = 3$, $a = 1$ and $b \geq 8$.*
- (2) *If X is quasielliptic with $\kappa(X) = 1$ and $p = 2$, $a = 1$ and $b \geq 19$.*
- (3) *If X is of general type with $p \geq 3$, $a = 2$ and $b \geq 4$.*
- (4) *If X is of general type with $p = 2$, $a = 2$ and $b \geq 19$.*

The key ingredient of Theorem 1.4 is a combination of a Reider-type result due to Shepherd-Barron and bend-and-break techniques.

For other results on the geography of pathological surfaces of Kodaira dimension smaller than two, see [Langer 2014].

In Section 5, a Kawamata–Viehweg-type vanishing theorem is proved for surfaces that are quasielliptic with $\kappa(X) = 1$ or of general type (see Theorem 5.7 and Corollary 5.9); this generalizes the vanishing result in [Terakawa 1999].

The core of our approach is a beautiful construction first introduced by Tango [1972] for the case of curves and Ekedahl [1988] and Shepherd-Barron [1991a] for surfaces. The same strategy was generalized by Kollár [1996] in order to investigate the geography of varieties where Kodaira-type vanishing theorems fail, via bend-and-break techniques.

2. Preliminary results

In this section we recall some techniques we will need later in this paper.

2A. Volume of divisors. Let D be a Cartier divisor on a normal variety X , not necessarily a surface. The volume of D measures the asymptotic growth of the space of global sections of multiples of D . We will recall here few properties of the volume, and we refer to [Lazarsfeld 2004b] for more details.

Definition 2.1. Let D be a Cartier divisor on X , with $\dim(X) = n$. The volume of D is defined by

$$\text{vol}(D) := \limsup_{m \rightarrow \infty} \frac{h^0(X, \mathbb{O}_X(mD))}{m^n/n!}.$$

The volume of X is defined as $\text{vol}(X) := \text{vol}(K_X)$.

It is easy to show that if D is big and nef then $\text{vol}(D) = D^n$. In general it is a hard invariant to compute, but thanks to Fujita's approximation theorem some of its properties can be deduced from the case where D is ample. For a proof of the theorem in characteristic zero we refer to [Lazarsfeld 2004b]. More recently, Takagi [2007] gave a proof of the same theorem in positive characteristic. In particular, we can deduce the log-concavity of the volume function even in positive characteristic. The proof is exactly the same as Theorem 11.4.9 in [Lazarsfeld 2004b].

Theorem 2.2. *Let D and D' be big Cartier divisors on a normal variety X defined over an algebraically closed field. Then*

$$\text{vol}(D + D')^{1/n} \geq \text{vol}(D)^{1/n} + \text{vol}(D')^{1/n}.$$

2B. Bogomolov's inequality and Sakai's theorems. We start with the notion of semistability for rank-two vector bundles on surfaces. Let X be a smooth surface defined over an algebraically closed field.

Definition 2.3. A rank-two vector bundle \mathcal{E} on X is *unstable* if it fits in a short exact sequence

$$0 \longrightarrow \mathcal{O}_X(D_1) \longrightarrow \mathcal{E} \longrightarrow \mathcal{I}_Z \cdot \mathcal{O}_X(D_2) \longrightarrow 0,$$

where D_1 and D_2 are Cartier divisors such that $D' := D_1 - D_2$ is big with $(D'^2) > 0$ and Z is an effective 0-cycle on X .

The vector bundle \mathcal{E} is *semistable* if it is not unstable.

In characteristic zero, the following celebrated result holds:

Theorem 2.4 [Bogomolov 1978]. *Let X be defined over a field of characteristic zero. Then every rank-two vector bundle \mathcal{E} for which $c_1^2(\mathcal{E}) > 4c_2(\mathcal{E})$ is unstable.*

As a consequence, one can deduce the following theorem, due to Sakai [1990, Proposition 1], which turns out to be equivalent to Theorem 2.4. This equivalence was shown in [Di Cerbo 2013].

Theorem 2.5. *Let D be a nonzero big divisor with $D^2 > 0$ on a smooth projective surface X over a field of characteristic zero. If $H^1(X, \mathcal{O}_X(K_X + D)) \neq 0$, then there exists a nonzero effective divisor E such that:*

- $D - 2E$ is big.
- $(D - E) \cdot E \leq 0$.

The previous result easily implies a weaker version of Reider's theorem:

Theorem 2.6. *Let D be a nef divisor with $D^2 > 4$ on a smooth projective surface X over a field of characteristic zero. Then $K_X + D$ has no basepoint unless there exists a nonzero effective divisor E such that $D \cdot E = 0$ and $(E^2) = -1$ or $D \cdot E = 1$ and $(E^2) = 0$.*

The following result, conjectured by Fujita, can be deduced for smooth surfaces in characteristic zero:

Corollary 2.7 (Fujita conjectures for surfaces in char 0). *Let D_1, \dots, D_k be ample divisors on a smooth surface X over a field of characteristic zero. Then $K_X + D_1 + \dots + D_k$ is basepoint free if $k \geq 3$ and very ample if $k \geq 4$.*

We remark that Theorem 2.5 is not known in general for smooth surfaces in positive characteristic, although Fujita’s conjectures are expected to hold.

2C. Ekedahl’s construction and Shepherd-Barron’s theorem. In this section we recall some classical results on the geography of smooth surfaces in positive characteristic (see [Ekedahl 1988; Shepherd-Barron 1991a; 1991b]).

For a good overview on the geography of surfaces in positive characteristic, see [Liedtke 2013].

We discuss here a construction which is due to Tango [1972] for the case of curves and Ekedahl [1988] for surfaces. There are many variations on the same theme, but we will focus on the one which is more related to the stability of vector bundles. We need this fundamental result:

Theorem 2.8 (Bogomolov). *Let \mathcal{E} be a rank-two vector bundle on a smooth projective surface X over a field of positive characteristic such that Bogomolov’s inequality does not hold (i.e., such that $c_1^2(\mathcal{E}) > 4c_2(\mathcal{E})$). Then there exists a reduced and irreducible surface Y contained in the ruled threefold $\mathbb{P}(\mathcal{E})$ such that:*

- *The restriction $\rho : Y \rightarrow X$ is p^e -purely inseparable for some $e > 0$.*
- *$(F^*)^e(\mathcal{E})$ is unstable.*

Proof. See [Shepherd-Barron 1991a, Theorem 1] □

The previous result also provides an explicit construction of the purely inseparable cover (see [Shepherd-Barron 1991a]).

Construction 2.9. Take a rank-two vector bundle \mathcal{E} such that Bogomolov’s inequality does not hold, and let e be an integer such that $F^{e*}\mathcal{E}$ is unstable. We have the commutative diagram

$$\begin{array}{ccc}
 \mathbb{P}(F^{e*}\mathcal{E}) & \xrightarrow{G} & \mathbb{P}(\mathcal{E}) \\
 p' \downarrow & & \downarrow p \\
 X & \xrightarrow{F^e} & X
 \end{array}$$

The fact that $F^{e*}\mathcal{E}$ is unstable gives an exact sequence

$$0 \longrightarrow \mathcal{O}_X(D_1) \longrightarrow F^{e*}\mathcal{E} \longrightarrow \mathcal{I}_Z \cdot \mathcal{O}_X(D_2) \longrightarrow 0$$

and a quasisection X_0 of $\mathbb{P}(F^{e*}\mathcal{E})$ (i.e., $p'_{X_0} : X_0 \rightarrow X$ is birational). Let Y be the image of X_0 via G . One can show that the induced morphism

$$\rho : Y \rightarrow X$$

is p^e -purely inseparable. Let us define $D' := D_1 - D_2$. One can show (see [Shepherd-Barron 1991a, Corollary 5]) that

$$K_Y \equiv \rho^* \left(K_X - \frac{p^e - 1}{p^e} D' \right).$$

Remark 2.10. We will be particularly interested in the case when the rank-two vector bundle \mathcal{E} comes as a nontrivial extension

$$0 \longrightarrow \mathcal{O}_X \longrightarrow \mathcal{E} \longrightarrow \mathcal{O}_X(D) \longrightarrow 0$$

associated to a nonzero element $\gamma \in H^1(X, \mathcal{O}_X(-D))$, where D is a big Cartier divisor such that $(D^2) > 0$. Indeed, the instability of $F^{e*}\mathcal{E}$ guarantees the existence of a diagram (keeping the notation as in Definition 2.3)

$$\begin{array}{ccccccc}
 & & & 0 & & & \\
 & & & \downarrow & & & \\
 & & & \mathcal{O}_X(D' + D_2) & & & \\
 & & & \downarrow f_1 & & & \\
 0 & \longrightarrow & \mathcal{O}_X & \xrightarrow{g_1} & F^{e*}\mathcal{E} & \xrightarrow{g_2} & \mathcal{O}_X(p^e D) \longrightarrow 0 \\
 & & \searrow \sigma & & \downarrow f_2 & & \\
 & & & & \mathcal{I}_Z \cdot \mathcal{O}_X(D_2) & & \\
 & & & & \downarrow & & \\
 & & & & 0 & &
 \end{array}$$

First, we claim that the composition map σ is nonzero. Assume for a contradiction that $\sigma \equiv 0$. This gives a nonzero section $\sigma' : \mathcal{O}_X \rightarrow \mathcal{O}_X(D' + D_2)$. This forces the composition $\tau := g_2 \circ f_1$ to be zero. But this implies that $D' + D_2 \leq 0$. This is a contradiction (see the proof of [Sakai 1990, Proposition 1] and [Shepherd-Barron 1991a, Lemma 16]).

This implies that $D_2 \simeq E \geq 0$; one can then rewrite the vertical exact sequence as

$$0 \longrightarrow \mathcal{O}_X(p^e D - E) \longrightarrow F^{e*}\mathcal{E} \longrightarrow \mathcal{I}_Z \cdot \mathcal{O}_X(E) \longrightarrow 0.$$

Since [Shepherd-Barron 1991a, Corollary 8] guarantees that Corollary 2.7 holds true for smooth surfaces in positive characteristic which are neither quasielliptic

with $\kappa(X) = 1$ nor of general type, we need to deduce effective basepoint-freeness and very-ampleness results only for these two classes of surfaces.

We recall here the following key result from [Shepherd-Barron 1991a]:

Theorem 2.11. *Let \mathcal{E} be a rank-two vector bundle on a smooth projective surface X over an algebraically closed field of positive characteristic such that Bogomolov’s inequality does not hold and \mathcal{E} is semistable.*

- *If X is not of general type, then X is quasielliptic with $\kappa(X) = 1$.*
- *If X is of general type and*

$$c_1^2(\mathcal{E}) - 4c_2(\mathcal{E}) > \frac{\text{vol}(X)}{(p-1)^2},$$

then X is purely inseparably uniruled. More precisely, in the notation of Theorem 2.8, Y is uniruled.

Proof. This is [Shepherd-Barron 1991a, Theorem 7], since the volume of a surface X with minimal model X' equals $(K_{X'}^2)$. □

Corollary 2.12 [Shepherd-Barron 1991a, Corollary 8]. *Corollary 2.7 holds in positive characteristic if X is neither of general type nor quasielliptic.*

2D. Bend-and-break lemmas. We recall here a well-known result in birational geometry, based on a celebrated method due to Mori (see [Kollár 1996] for an insight into these techniques).

First we need to recall some notation. Mori theory deals with effective 1-cycles in a variety X ; more specifically, we will consider nonconstant morphisms $h : C \rightarrow X$, where C is a smooth curve. In particular, these techniques allow us to deform curves for which

$$(K_X \cdot C) := \deg_C h^* K_X < 0.$$

In what follows, we will denote by $\overset{e}{\approx}$ the *effective algebraic equivalence* defined on the space of effective 1-cycles $Z_1(X)$ (see [Kollár 1996, Definition II.4.1]).

Theorem 2.13 (bend-and-break). *Let X be a variety over an algebraically closed field, and let C be a smooth, projective and irreducible curve with a morphism $h : C \rightarrow X$ such that X has local complete intersection singularities along $h(C)$ and $h(C)$ intersects the smooth locus of X . Assume the numerical condition*

$$(K_X \cdot C) < 0$$

holds. Then, for every point $x \in C$, there exists a rational curve C_x in X passing through x such that

$$h_*[C] \overset{e}{\approx} k_0[C_x] + \sum_{i \neq 0} k_i[C_i] \tag{1}$$

(as algebraic cycles), with $k_i \geq 0$ for all i and

$$-(K_X \cdot C_x) \leq \dim X + 1.$$

Proof. See [Kollár 1996, Theorem II.5.14 and Remark II.5.15]. The relation (1) can be deduced by looking directly at the proofs of the bend-and-break lemmas [Kollár 1996, Corollary II.5.6 and Theorem II.5.7]; our notation is slightly different, since in (1) we have isolated a rational curve with the required intersection properties. \square

In this paper we will need the following consequence of the previous theorem:

Corollary 2.14. *Let X be a surface which fibers over a curve C via $f : X \rightarrow C$ and let F be the general fiber of f . Assume that X has only local complete intersection singularities along F and that F is a (possibly singular) rational curve such that*

$$(K_X \cdot F) < 0.$$

Then

$$-(K_X \cdot F) \leq 3.$$

Proof. The hypotheses of Theorem 2.13 hold here, so we can take a point x in the smooth locus of X and deduce the existence of a rational curve C' passing through x such that

$$-(K_X \cdot C') \leq 3 \quad \text{and} \quad [F] \stackrel{e}{\approx} k_0[C'] + \sum_{i \neq 0} k_i[C_i].$$

By Exercise II.4.1.10 in [Kollár 1996], the curves appearing on the right hand side of the previous equation must be contained in the fibers of f . Since F is the general fiber, the second relation implies that $k_0 = 1$ and $k_i = 0$ for all $i \neq 0$, and so $C' = F$. \square

3. An effective Matsusaka's theorem

In this section we prove Theorem 1.2 assuming the results on effective very-ampleness that we will prove in the next section. If not specified, X will denote a smooth surface over an algebraically closed field of arbitrary characteristic.

First we recall the following numerical criterion for bigness, whose characteristic-free proof is based on Riemann–Roch [Lazarsfeld 2004a, Theorem 2.2.15].

Theorem 3.1. *Let D and E be nef \mathbb{Q} -divisors on X and assume that*

$$D^2 > 2(D \cdot E).$$

Then $D - E$ is big.

Before proving Theorem 1.2, we need some lemmas.

Lemma 3.2. *Let D be an ample divisor on X . Then $K_X + 2D + C$ is nef for any irreducible curve $C \subset X$.*

Proof. If $X = \mathbb{P}^2$ then the lemma is trivial. By the cone theorem and the classification of surfaces with extremal rays of maximal length, we have that $K_X + 2D$ is always a nef divisor. This implies that $K_X + 2D + C$ may have negative intersection number only when intersected with C . On the other hand, by adjunction, $(K_X + C) \cdot C = 2g - 2 \geq -2$, where g is the arithmetic genus of C . Since D is ample, the result follows. \square

We can now prove one of the main results of this section (see [Lazarsfeld 2004b, Theorem 10.2.4]).

Theorem 3.3. *Let D be an ample divisor and let B be a nef divisor on X . Then $mD - B$ is nef for any*

$$m \geq \frac{2D \cdot B}{D^2} ((K_X + 2D) \cdot D + 1) + 1.$$

Proof. To simplify the notation in the proof let us define the following numbers:

$$\begin{aligned} \eta &= \eta(D, B) := \inf\{t \in \mathbb{R}_{>0} \mid tD - B \text{ is nef}\}, \\ \gamma &= \gamma(D, B) := \inf\{t \in \mathbb{R}_{>0} \mid tD - B \text{ is pseudoeffective}\}. \end{aligned}$$

The theorem will follow if we find an upper bound on η . Note that $\gamma \leq \eta$ since a nef divisor is also pseudoeffective.

By definition $\eta D - B$ is in the boundary of the nef cone and by Nakai’s theorem we have two possible cases: either

- $(\eta D - B)^2 = 0$, or
- there exists an irreducible curve C such that $(\eta D - B) \cdot C = 0$.

If $(\eta D - B)^2 = 0$, then it is easy to see that

$$\eta \leq 2 \frac{D \cdot B}{D^2}.$$

So we can assume that there exists an irreducible curve C such that $\eta D \cdot C = B \cdot C$. Let us define $G := \gamma D - B$. Then

$$G \cdot C = (\gamma - \eta)D \cdot C \leq (\gamma - \eta).$$

Let us define $A := K_X + 2D$. By Lemma 3.2 and the definition of G , we have that $(A + C) \cdot G \geq 0$. Combining with the previous inequality we get

$$(\eta - \gamma) \leq -G \cdot C \leq A \cdot G = \gamma A \cdot D - A \cdot B.$$

In particular,

$$\eta \leq \gamma(A \cdot D + 1) - A \cdot B \leq \gamma(A \cdot D + 1).$$

The statement of our result follows from Theorem 3.1, which guarantees that $\gamma < (2D \cdot B)/(D^2)$. \square

Remark 3.4. The previous proof is characteristic-free, although the new result is for surfaces in positive characteristic.

We can now prove our main theorem, assuming the results in the next section.

Proof of Theorem 1.2. By Corollary 2.12, if X is neither of general type nor quasielliptic and $H = K_X + 4D$ then $H + N$ is very ample for any nef divisor N . By Theorem 3.3, $mD - (H + B)$ is nef for any m as in the statement. Then $K_X + 4D + (mD - K_X - 4D - B)$ is very ample. For surfaces in the other classes use Theorem 4.10 and Theorem 4.12 to obtain the desired very ample divisor H . \square

4. Effective very-ampleness in positive characteristic

The aim of this section is to complete the proof of Theorem 1.2 for quasielliptic surfaces of Kodaira dimension one and for surfaces of general type. Our ultimate goal is to prove Theorem 1.4 via a case-by-case analysis.

First we need some notation (cf. Theorem 2.5).

Definition 4.1. A big divisor D on a smooth surface X with $(D^2) > 0$ is m -unstable for a positive integer m if either:

- $H^1(X, \mathbb{C}_X(-D)) = 0$.
- $H^1(X, \mathbb{C}_X(-D)) \neq 0$ and there exists a nonzero effective divisor E such that:
 - $mD - 2E$ is big.
 - $(mD - E) \cdot E \leq 0$.

Remark 4.2. Theorem 2.5 tells us that in characteristic zero every big divisor D on a smooth surface X with $(D^2) > 0$ is 1-unstable. The same holds in positive characteristic, if we assume that the surface is neither of general type nor quasielliptic of maximal Kodaira dimension; this is a consequence of Corollary 2.12. Our goal here is to clarify the picture in the remaining cases.

We start our analysis with quasielliptic surfaces of maximal Kodaira dimension.


Proposition 4.3. *Let X be a quasielliptic surface with $\kappa(X) = 1$ and let D be a big divisor on X with $(D^2) > 0$.*

- (1) *if $p = 3$, then D is 3-unstable.*
- (2) *if $p = 2$, then D is 4-unstable.*

Proof. Assume that $p = 3$ and $H^1(X, \mathbb{C}_X(-D)) \neq 0$. This nonzero element gives a nonsplit extension

$$0 \rightarrow \mathbb{C}_X \rightarrow \mathcal{E} \rightarrow \mathbb{C}_X(D) \rightarrow 0.$$

Theorem 2.8 implies that $(F^*)^{e\mathcal{C}}$ is unstable for e sufficiently large. To prove the proposition in this case we need to show that $e = 1$. Assume $e \geq 2$ and let F be the general element of the pencil which gives the fibration in cuspidal curves $f : X \rightarrow B$. Let $\rho : Y \rightarrow X$ be the p^e -purely inseparable morphism of Construction 2.9. Then $\{C_i := \rho^* F\}$ is a family of movable rational curves in Y . Let us define $g := f \circ \rho$ and consider its Stein factorization:



Since the curves in the family $\{C_i\}$ are precisely the fibers of h , we can use Corollary 2.14 on $h : Y \rightarrow B'$ (since Y is defined via a quasisection in a \mathbb{P}^1 -bundle over X , it has hypersurface singularities along the general element of $\{C_i\}$) and deduce that

$$0 < -(K_Y \cdot C_i) \leq 3.$$

This gives a contradiction, since

$$\begin{aligned}
 3 &\geq -(K_Y \cdot C_i) = \left(\rho^* \left(\frac{p^e - 1}{p^e} (p^e D - 2E) - K_X \right) \cdot C_i \right) \\
 &= p^e \left(\left(\frac{p^e - 1}{p^e} (p^e D - 2E) - K_X \right) \cdot F \right) \\
 &= ((p^e - 1)(p^e D - 2E) \cdot F) \\
 &\geq p^e - 1 \geq 8,
 \end{aligned}$$

where E is the divisor appearing in Remark 2.10.

The same proof works for $p = 2$, although in this case we can only prove that $e \leq 2$. □

We can now focus on the general type case. We need the following theorem of Shepherd-Barron [1991a, Theorem 12].

Theorem 4.4. *Let D be a big Cartier divisor on a smooth surface X of general type which satisfies one of the following hypotheses:*

- $p \geq 3$ and $(D^2) > \text{vol}(X)$.
- $p = 2$ and $(D^2) > \max\{\text{vol}(X), \text{vol}(X) - 3\chi(\mathcal{O}_X) + 2\}$.

Then D is 1-unstable.

Since the bound of the previous theorem depends on $\chi(\mathbb{O}_X)$ if $p = 2$, we need an additional result for this case. First we recall a result by Shepherd-Barron [1991b, Theorem 8].

Theorem 4.5. *Let X be a surface in characteristic $p = 2$ of general type with $\chi(\mathbb{O}_X) < 0$. Then there is a fibration $f : X \rightarrow C$ over a smooth curve C whose generic fiber is a singular rational curve with arithmetic genus $2 \leq g \leq 4$.*

We can prove now our result.

Proposition 4.6. *Let D be a big Cartier divisor on a surface in characteristic $p = 2$ of general type with $\chi(\mathbb{O}_X) < 0$ such that $(D^2) > \text{vol}(X)$. Then D is 4-unstable.*

Proof. Assume that $H^1(X, \mathbb{O}_X(-D)) \neq 0$. As in the proof of Proposition 4.3 we have a nonsplit extension

$$0 \rightarrow \mathbb{O}_X \rightarrow \mathcal{E} \rightarrow \mathbb{O}_X(D) \rightarrow 0.$$

Using Theorem 2.8 we deduce the instability of $(F^*)^e \mathcal{E}$ for e sufficiently large. Let F be the general element of the pencil which gives the fibration in singular rational curves given by Theorem 4.5. Let $\rho : Y \rightarrow X$ be the p^e -purely inseparable morphism of Construction 2.9. As in the proof of Proposition 4.3, we use Corollary 2.14 on Y and deduce that $0 < -(K_Y \cdot C_i) \leq 3$. This gives

$$\begin{aligned} 3 &\geq -(K_Y \cdot C_i) = \left(\rho^* \left(\frac{2^e - 1}{2^e} (2^e D - 2E) - K_X \right) \cdot C_i \right) \\ &= 2^e \left(\left(\frac{2^e - 1}{2^e} (2^e D - 2E) - K_X \right) \cdot F \right) \\ &= ((2^e - 1)(2^e D - 2E) - 2^e K_X) \cdot F \geq 1. \end{aligned}$$

This implies that

$$((2^e - 1)(2^{e-1} D - E) - 2^{e-1} K_X) \cdot F = 1.$$

As a consequence, we apply Theorem 4.5 to bound the intersection $(K_X \cdot F)$:

$$(2^e - 1)((2^{e-1} D - E) \cdot F) = 2^e (g - 1) + 1,$$

where g is the arithmetic genus of F . By some basic arithmetic the only possibilities for the pair (g, e) are $(2, 1)$, $(3, 1)$, $(3, 2)$ and $(4, 1)$. □

We will use Theorem 4.4 to prove a variant of Reider’s theorem in positive characteristic. We state a technical proposition that we will need later (see [Sakai 1990, Proposition 2]).

Proposition 4.7. *Let $\pi : Y \rightarrow X$ be a birational morphism between two normal surfaces. Let \tilde{D} be a Cartier divisor on Y such that $\tilde{D}^2 > 0$. Assume there is a nonzero effective divisor \tilde{E} such that*

- $\tilde{D} - 2\tilde{E}$ is big,
- $(\tilde{D} - \tilde{E}) \cdot \tilde{E} \leq 0$.

Set $D := \pi_*\tilde{D}$, $E := \pi_*\tilde{E}$ and $\alpha = D^2 - \tilde{D}^2$. If D is nef and E is a nonzero effective divisor, then

- $0 \leq D \cdot E < \alpha/2$,
- $D \cdot E - \alpha/4 \leq E^2 \leq (D \cdot E)^2/D^2$.

The corollary we need is the following.

Corollary 4.8. *Let $\pi : Y \rightarrow X$ be a birational morphism between two smooth surfaces and let \tilde{D} be a big Cartier divisor on Y such that $(\tilde{D}^2) > 0$. Assume that*

- $H^1(X, \mathcal{O}_X(-\tilde{D})) \neq 0$,
- \tilde{D} is m -unstable for some $m > 0$.

Set $D := \pi_*\tilde{D}$ and $\alpha = D^2 - \tilde{D}^2$. Then if D is nef, there exists a nonzero effective divisor E on X such that

- $0 \leq D \cdot E < m\alpha/2$,
- $mD \cdot E - m^2\alpha/4 \leq E^2 \leq (D \cdot E)^2/D^2$.

We can now derive our effective basepoint-freeness results. We will start with quasielliptic surfaces, applying Proposition 4.3 and the previous corollary.

Proposition 4.9. *Let X be a quasielliptic surface with maximal Kodaira dimension. Let D be a big and nef divisor on X . Then the following hold.*

- For $p = 3$:
 - If $D^2 > 4$ and $|K_X + D|$ has a basepoint at $x \in X$, there exists a curve C such that $D \cdot C \leq 5$.
 - If $D^2 > 9$ and $|K_X + D|$ does not separate any two points $x, y \in X$, there exists a curve C such that $D \cdot C \leq 13$.
- For $p = 2$:
 - If $D^2 > 4$ and $|K_X + D|$ has a basepoint at $x \in X$, there exists a curve C such that $D \cdot C \leq 7$.
 - If $D^2 > 9$ and $|K_X + D|$ does not separate any two points $x, y \in X$, there exists a curve C such that $D \cdot C \leq 17$.

Proof. We start with the case $p = 3$. Assume that $|K_X + D|$ has a basepoint at $x \in X$. Let $\pi : Y \rightarrow X$ be the blowup at x . Since x is a basepoint we have that $H^1(Y, \mathcal{O}_Y(K_Y + \pi^*D - 2F)) \neq 0$, where F is the exceptional divisor of π . Let $\tilde{D} := \pi^*D - 2F$. By assumption we have that $(\tilde{D}^2) > 0$. By Proposition 4.3 we can find an effective divisor \tilde{E} such that $p\tilde{D} - 2\tilde{E}$ is big and $(p\tilde{D} - \tilde{E}) \cdot \tilde{E} \leq 0$. The previous inequality easily implies that \tilde{E} is not a positive multiple of the

exceptional divisor and in particular $E := \pi_* \tilde{E}$ is a nonzero effective divisor. Moreover, $D = \pi_* \tilde{D}$ is nef by assumption, thus we can apply Corollary 4.8. Since $\alpha = (D^2 - \tilde{D}^2) = 4$, the first inequality of the corollary implies that $D \cdot E \leq 5$.

The statement on separation of points follows in exactly the same way. Note that we allow the case $x = y$.

The bounds for the case $p = 2$ can be obtained the same way, remarking that \tilde{D} is p^2 -unstable in this case. □

The previous results can be used to derive effective very-ampleness statements for quasielliptic surfaces when D is an ample divisor.

Theorem 4.10. *Let D be an ample Cartier divisor on a smooth quasielliptic surface X with $\kappa(X) = 1$.*

- *If $p = 3$, the divisor $K_X + kD$ is basepoint-free for any $k \geq 4$ and it is very ample for any $k \geq 8$.*
- *If $p = 2$, the divisor $K_X + kD$ is basepoint-free for any $k \geq 5$ and it is very ample for any $k \geq 19$.*

In particular, if N is any nef divisor, $K_X + kD + N$ is always very ample for any $k \geq 8$ (resp. $k \geq 19$) in characteristic 3 (resp. 2).

Proof. The proof consists of explicitly computing the minimal multiple of D which contradicts the second inequality of Corollary 4.8.

Let us start with basepoint-freeness for $p = 3$. Assume that $k \geq 5$, $K_X + kD$ has a basepoint and define $D' := kD$. Then, by Proposition 4.9, we know that there exists an effective divisor E such that $(D' \cdot E) \leq 5$. This implies

$$(D \cdot E) \leq 1.$$

Now use the second inequality of Corollary 4.8 on D' to deduce

$$15 - 9 \leq 3(D' \cdot E) - 9 \leq \frac{(D' \cdot E)^2}{(D'^2)} \leq 1.$$

This is a contradiction.

Similar computations give the other bounds. □

We now deal with surfaces of general type. The analogue of Proposition 4.9 is the following.

Proposition 4.11. *Let X be a surface of general type and let D be a big and nef divisor on X . Then the following hold.*

- *For $p \geq 3$:*
 - *If $D^2 > \text{vol}(X) + 4$ and $|K_X + D|$ has a basepoint at $x \in X$, there exists a curve C such that $D \cdot C \leq 1$.*

- If $D^2 > \text{vol}(X) + 9$ and $|K_X + D|$ does not separate any two points $x, y \in X$, there exists a curve C such that $D \cdot C \leq 2$.
- For $p = 2$:
 - If $D^2 > \text{vol}(X) + 6$ and $|K_X + D|$ has a basepoint at $x \in X$, there exists a curve C such that $D \cdot C \leq 7$.
 - If $D^2 > \text{vol}(X) + 11$ and $|K_X + D|$ does not separate any two points $x, y \in X$, there exists a curve C such that $D \cdot C \leq 17$.

Proof. The proof is basically the same as Proposition 4.9. Let $p \geq 3$ and assume that $|K_X + D|$ has a basepoint at $x \in X$. Using the same notation as Proposition 4.9, we can blow up x and deduce the existence of an effective divisor \tilde{E} such that $\tilde{D} - 2\tilde{E}$ is big and $(\tilde{D} - \tilde{E}) \cdot \tilde{E} \leq 0$ (in order to deduce 1-instability we use Theorem 4.4). Also here, the first inequality of Corollary 4.8 implies that $(D \cdot E) \leq 1$.

The statement on separation of points follows in the same way.

For the bounds in the case $p = 2$ we use the same strategy, using a combination of Theorem 4.5 and Proposition 4.6. □

The following effective very-ampleness statement can be deduced. Applying Proposition 4.11 directly would provide bounds that depend on the volume. It is possible to get a uniform bound if we work with linear systems of the type $|2K_X + mD|$. Note that we get sharp statements for those linear systems.

Theorem 4.12. *Let D be an ample Cartier divisor on a smooth surface X of general type.*

- If $p \geq 3$, the divisor $2K_X + kD$ is basepoint free for any $k \geq 3$ and it is very ample for any $k \geq 4$.
- If $p = 2$ the divisor $2K_X + kD$ is basepoint free for any $k \geq 5$ and it is very ample for any $k \geq 19$.

In particular, if N is any nef divisor, $2K_X + kD + N$ is always very ample for any $k \geq 4$ (resp. $k \geq 19$) in characteristic $p \geq 3$ (resp. $p = 2$).

Proof. Since negative extremal rays of general type surfaces have length 1, if $m \geq 3$, we know that $L := K_X + mD$ is an ample divisor and $L \cdot C \geq 2$ for any irreducible curve $C \subset X$. Moreover, by log-concavity of the volume function (see Theorem 2.2) we have that

$$L^2 = \text{vol}(L) \geq \text{vol}(K_X) + 9D^2 > \text{vol}(X) + 4.$$

Proposition 4.11 implies that $K_X + L = 2K_X + kD$ is basepoint free for any $k \geq 4$. A similar computation allows us to derive very-ampleness.

The same strategy gives the result for $p = 2$. □

Proof of Theorem 1.4. This is simply given by Theorem 4.10 and Theorem 4.12. □

Remark 4.13. In [Terakawa 1999], similar results can be found. Nonetheless our approach allows us to deduce effective basepoint-freeness and very-ampleness also on quasielliptic surfaces and arbitrary surfaces of general type.

5. A Kawamata–Viehweg-type vanishing theorem in positive characteristic

In this section we give an extension of the results in [Terakawa 1999]. There, Terakawa used the results in [Shepherd-Barron 1991a] to deduce a Kawamata–Viehweg-type theorem for nonpathological surfaces. Using our methods we are able to discuss pathological surfaces and obtain an effective Kawamata–Viehweg-type theorem in positive characteristic.

Let us first recall the classical Kawamata–Viehweg vanishing theorem in its general version (see [Kollár and Mori 1998] for the general notation).

Theorem 5.1. *Let (X, B) be a klt pair over an algebraically closed field of characteristic zero and let D be a Cartier divisor on X such that $D - (K_X + B)$ is big and nef. Then*

$$H^i(X, \mathcal{O}_X(D)) = 0$$

for any $i > 0$.

In positive characteristic, even for nonpathological smooth surfaces, there are counterexamples to Theorem 5.1: Xie [2010] provided examples of relatively minimal irregular ruled surfaces in every characteristic where Theorem 5.1 fails.

Nonetheless, assuming $B = 0$, we have the following result (see [Mukai 2013]).

Theorem 5.2. *Let X be a smooth surface in positive characteristic. Assume that there exists a big and nef Cartier divisor D on X such that*

$$H^1(X, \mathcal{O}_X(K_X + D)) \neq 0.$$

Then:

- X is either quasielliptic of Kodaira dimension one or of general type.
- Up to a sequence of blowups, X has the structure of a fibered surface over a smooth curve such that every fiber is connected and singular.

Furthermore, Terakawa [1999] deduced the following vanishing result using the techniques in [Shepherd-Barron 1991a].

Theorem 5.3. *Let X be a smooth projective surface over an algebraically closed field of characteristic $p > 0$ and let D be a big and nef Cartier divisor on X . Assume that either:*

- (1) $\kappa(X) \neq 2$ and X is not quasielliptic with $\kappa(X) = 1$.
- (2) X is of general type with

- $p \geq 3$ and $(D^2) > \text{vol}(X)$; or
- $p = 2$ and $(D^2) > \max\{\text{vol}(X), \text{vol}(X) - 3\chi(\mathbb{O}_X) + 2\}$.

Then

$$H^i(X, \mathbb{O}_X(K_X + D)) = 0$$

for all $i > 0$.

Our aim is to improve this theorem for arbitrary surfaces, via bend-and-break techniques.

More generally, we want to deduce some results on the injectivity of cohomological maps

$$H^1(X, \mathbb{O}_X(-D)) \xrightarrow{F^*} H^1(X, \mathbb{O}_X(-pD))$$

where D is a big divisor on X .

The following result by Kollár is an application of bend-and-break lemmas (cf. Theorem 2.13), specialized to our two-dimensional setting.

Theorem 5.4. *Let X be a smooth projective variety over a field of positive characteristic and let D be a Cartier divisor on X such that:*

- (1) $H^1(X, \mathbb{O}_X(-mD)) \xrightarrow{F^*} H^1(X, \mathbb{O}_X(-pmD))$ is not injective for some integer $m > 0$.
- (2) *There exists a curve C on X such that*

$$(p - 1)(D \cdot C) - (K_X \cdot C) > 0.$$

Then through every point x of C there is a rational curve C_x such that

$$[C] \stackrel{e}{\approx} k_0[C_x] + \sum_{i \neq 0} k_i[C_i] \tag{2}$$

(as algebraic cycles), with $k_i \geq 0$ for all i and

$$(p - 1)(D \cdot C_x) - (K_X \cdot C_x) \leq \dim(X) + 1.$$

Proof. This is essentially a slight modification of [Kollár 1996, Theorem II.6.2]. For the reader’s convenience, we sketch it ab initio. Assumption (1) allows us to construct a finite morphism

$$\pi : Y \rightarrow X,$$

where Y is defined as a Cartier divisor in the projectivization of a nonsplit rank-two bundle over X (see [Kollár 1996, Construction II.6.1.6], which is a slight modification of Construction 2.9). Furthermore, the following property holds:

$$K_Y = \pi^*(K_X + (k(1 - p)D)),$$

where k is the largest integer for which $H^1(X, -kD) \neq 0$.

Now take the curve given in (2) and consider $C' := \text{red } \pi^{-1}(C)$. The hypothesis on the intersection numbers and the formula for the canonical divisor of Y guarantee that $(K_Y \cdot C') < 0$. Let $y \in C'$ be a preimage of x in Y . So we can apply Theorem 2.13 and deduce the existence of a rational curve C'_y passing through y . Using the projection formula, we obtain a curve C_x on X for which

$$(p-1)(D \cdot C_x) - (K_X \cdot C_x) \leq \dim(X) + 1. \quad \square$$

If we assume the dimension to be two and the divisor D to be big and nef, the asymptotic condition

$$H^1(X, \mathcal{O}_X(-mD)) = 0$$

for m sufficiently large is guaranteed by [Szpiro 1979].

This remark gives us the following corollary.

Corollary 5.5. *Let X be a smooth projective surface over a field of positive characteristic and let D be a big and nef Cartier divisor on X such that $H^1(X, \mathcal{O}_X(-D)) \neq 0$. Assume there exists a curve C on X such that*

$$(p-1)(D \cdot C) - (K_X \cdot C) > 0.$$

Then through every point x of C there is a rational curve C_x such that

$$(p-1)(D \cdot C_x) - (K_X \cdot C_x) \leq 3.$$

We will show later how Corollary 5.5 can be used to deduce an effective version of Kawamata–Viehweg-type vanishing for arbitrary smooth surfaces.

In what follows, we will also need the following lemma on fibered surfaces, which explicitly gives a bound on the genus of the fiber with respect to the volume of the surface.

Lemma 5.6. *Let $f : X \rightarrow C$ be a fibered surface of general type and let g be the arithmetic genus of the general fiber F . Then*

$$\text{vol}(X) \geq g - 4.$$

Proof. We divide our analysis into cases according to the genus b of the base, after having assumed the fibration is relatively minimal (i.e., that $K_{X/C}$ is nef).

$b \geq 2$: In this case we can deduce a better estimate. Indeed,

$$\text{vol}(X) \geq (K_X^2) = (K_{X/C}^2) + 8(g-1)(b-1) \geq 8g - 8.$$

$b = 1$: In this case we need a more careful analysis, since in positive characteristic we cannot assume the semipositivity of $f_* K_{X/C}$. Nonetheless the following general

formula holds:

$$\deg(f_*K_{X/C}) = \chi(\mathbb{O}_X) - (g - 1)(b - 1), \tag{3}$$

which specializes to

$$\deg(f_*K_{X/C}) = \chi(\mathbb{O}_X) \geq 0.$$

Formula (3) can be obtained via Riemann–Roch, since we know that $R^1 f_*K_{X/C} = \mathbb{O}_C$ and that $R^1 f_*nK_{X/C} = 0$ for $n \geq 2$ by relative minimality. The last inequality can be assumed by [Shepherd-Barron 1991b, Theorem 8]. Furthermore, one can apply the following formula

$$\deg(f_*(nK_{X/C})) = \deg(f_*K_{X/C}) + \frac{n(n-1)}{2}(K_{X/C}^2).$$

Since $K_{X/C}$ is big, we deduce that

$$\deg(f_*(2K_{X/C})) \geq 1.$$

As a consequence, we can apply the results of [Atiyah 1957] and deduce a decomposition of $f_*(2K_{X/C})$ into indecomposable vector bundles

$$f_*(2K_{X/C}) = \bigoplus_i E_i,$$

where we can assume that $\deg(E_1) \geq 1$. This implies that all quotient bundles of E_1 have positive degree. We want to show now that there exists a degree-one divisor L_1 on C such that $h^0(C, f_*(2K_{X/C}) \otimes \mathbb{O}_C(-L_1)) \neq 0$.

But this is clear, since, for any degree-one divisor L on C , one has that all quotient bundles of $f_*(2K_{X/C}) \otimes \mathbb{O}_C(-L)$ have degree zero and, up to a twisting by a degree-zero divisor on C , one can assume there exists a quotient

$$f_*(2K_{X/C} \otimes \mathbb{O}_C(-L_1)) \rightarrow \mathbb{O}_C \rightarrow 0.$$

This implies that $h^0(X, \mathbb{O}_X(2K_{X/C} - F)) (= h^0(C, f_*(2K_{X/C}) \otimes \mathbb{O}_C(-L_1))) \neq 0$, where F is the general fiber of f and, since $K_X = K_{X/C}$ is nef, that

$$(K_X \cdot (2K_X - F)) \geq 0.$$

This gives the bound

$$\text{vol}(X) \geq (K_X^2) \geq g - 1.$$

$b = 0$: Also in this case we can assume that $\chi(\mathbb{O}_X) \geq 0$ and, as a consequence, that

$$\deg(f_*K_{X/\mathbb{P}^1}) = \chi(\mathbb{O}_X) + g - 1 \geq g - 1.$$

If $g \geq 6$,

$$\deg(f_*K_{X/\mathbb{P}^1}) \geq 5.$$

This implies that $\text{deg}(f_*K_X \otimes \mathcal{O}_{\mathbb{P}^1}(3)) \geq 0$ and, as a consequence of Grothendieck’s theorem on vector bundles on \mathbb{P}^1 ,

$$h^0(X, \mathcal{O}_X(K_X - f^*\mathcal{O}_C(-3))) \neq 0.$$

As before, we have assumed that $K_{X/\mathbb{P}^1} = K_X + f^*\mathcal{O}_C(2)$ is nef, so

$$((K_X + f^*\mathcal{O}_C(2)) \cdot (K_X - f^*\mathcal{O}_C(-3))) \geq 0.$$

So in this case

$$\text{vol}(X) \geq (K_X^2) \geq 2g - 2.$$

If $g \leq 5$, we simply use the trivial inequality $\text{vol}(X) \geq 1$ to deduce

$$\text{vol}(X) \geq g - 4. \quad \square$$

Our result in this setting is an effective bound, depending only on the birational geometry of X , that guarantees the injectivity of the induced Frobenius map on the H^1 s.

Theorem 5.7. *Let X be a smooth surface in characteristic $p > 0$ and let D be a big Cartier divisor on X . Then, for all integers*

$$m > m_0 = \frac{2 \text{vol}(X) + 9}{p - 1},$$

the induced Frobenius map

$$H^1(X, \mathcal{O}_X(-mD)) \xrightarrow{F^*} H^1(X, \mathcal{O}_X(-pmD))$$

is injective. (If $\kappa(X) \neq 2$, the volume $\text{vol}(X) = 0$).

Remark 5.8. The previous result is trivial if $H^1(X, \mathcal{O}_X(-D)) = 0$. Furthermore, combined with Corollary 5.5 it gives an effective version of the Kawamata–Viehweg theorem (cf. Corollary 5.9) in the case of big and nef divisors. Our hope is to generalize this strategy in order to deduce effective vanishing theorems also in higher dimension.

Proof. Assume, for the sake of contradiction, that

$$H^1(X, \mathcal{O}_X(-\lceil m_0 \rceil D)) \xrightarrow{F^*} H^1(X, \mathcal{O}_X(-p\lceil m_0 \rceil D))$$

has a nontrivial kernel. Then, after a sequence of blowups $f : X' \rightarrow X$, we can assume the existence of a (relatively minimal) fibration (possibly with singular general fiber) of arithmetic genus g

$$\pi : X' \rightarrow C.$$

We remark that we can reduce to proving our result on X' , since $D' := f^*D$ is a big divisor and we have the following commutative diagram

$$\begin{CD}
 H^1(X', \mathcal{O}_{X'}(-\lceil m_0 \rceil D')) @>F^*>> H^1(X', \mathcal{O}_{X'}(-p\lceil m_0 \rceil D')) \\
 @V \simeq VV @VV \simeq V \\
 H^1(X, \mathcal{O}_X(-\lceil m_0 \rceil D)) @>F^*>> H^1(X, \mathcal{O}_X(-p\lceil m_0 \rceil D))
 \end{CD}$$

where the vertical isomorphisms hold because of $R^1 f_* \mathcal{O}_{X'} = 0$. We can now apply Theorem 5.4 to $\lceil m_0 \rceil D'$: we can choose C to be a general fiber F of π , which certainly intersects D' positively, and we can use Lemma 5.6 to obtain

$$(p - 1)\lceil m_0 \rceil(D' \cdot F) - (K_{X'} \cdot F) \geq (p - 1)\lceil m_0 \rceil - (2g - 2) > 3. \tag{4}$$

So we can apply Theorem 5.4: fix a point $x \in F$ and find a rational curve C_x such that

$$(p - 1)m_0(D \cdot C_x) - (K_X \cdot C_x) \leq 3.$$

Notice that, by construction, $F = C_x$, because of (2) in Theorem 5.4. But this is a contradiction, because of (4). □

We finally obtain our effective vanishing theorem.

Corollary 5.9. *Let X be a smooth surface in characteristic $p > 0$ and let D be a big and nef Cartier divisor D on X . Then*

$$H^1(X, \mathcal{O}_X(K_X + mD)) = 0$$

for all integers $m > m_0$, where:

- $m_0 = 3/(p - 1)$ if X is quasielliptic with $\kappa(X) = 1$.
- $m_0 = (2 \operatorname{vol}(X) + 9)/(p - 1)$ if X is of general type.

Proof. For surfaces of general type, one simply applies the previous result. For quasielliptic surfaces, a better bound can be obtained, since in this case $(K_X \cdot F) = 0$, where F is the general fiber. □

Acknowledgements

We would like to thank Paolo Cascini, Yoshinori Gongyo and Hiromu Tanaka for useful comments and the referee for helping us to improve our exposition.

Fanelli is funded by the *Roth Scholarship* grant of the Department of Mathematics at Imperial College London.

References

- [Atiyah 1957] M. F. Atiyah, “Vector bundles over an elliptic curve”, *Proc. London Math. Soc.* (3) **7** (1957), 414–452. MR 24 #A1274 Zbl 0084.17305
- [Ballico 1996] E. Ballico, “A positive characteristic extension of a result of del Busto on line bundles on an algebraic surface”, *Rend. Circ. Mat. Palermo* (2) **45**:3 (1996), 473–478. MR 99f:14007 Zbl 0910.14014
- [Bogomolov 1978] F. A. Bogomolov, “Holomorphic tensors and vector bundles on projective manifolds”, *Izv. Akad. Nauk SSSR Ser. Mat.* **42**:6 (1978), 1227–1287. In Russian; translated in *Math. USSR Izv.* **13**:3 (1979) 499–555. MR 80j:14014
- [Fernández del Busto 1996] G. Fernández del Busto, “A Matsusaka-type theorem on surfaces”, *J. Algebraic Geom.* **5**:3 (1996), 513–520. MR 98d:14007 Zbl 0871.14006
- [Demailly 1996a] J.-P. Demailly, “Effective bounds for very ample line bundles”, *Invent. Math.* **124**:1-3 (1996), 243–261. MR 97a:32035 Zbl 0862.14004
- [Demailly 1996b] J.-P. Demailly, “ L^2 vanishing theorems for positive line bundles and adjunction theory”, pp. 1–97 in *Transcendental methods in algebraic geometry* (Cetraro, 1994), edited by F. Catanese and C. Ciliberto, Lecture Notes in Math. **1646**, Springer, Berlin, 1996. MR 99k:32051 Zbl 0883.14005
- [Di Cerbo 2013] G. Di Cerbo, “A cohomological interpretation of Bogomolov’s instability”, *Proc. Amer. Math. Soc.* **141**:9 (2013), 3049–3053. MR 3068958 Zbl 1282.14038
- [Ekedahl 1988] T. Ekedahl, “Canonical models of surfaces of general type in positive characteristic”, *Inst. Hautes Études Sci. Publ. Math.* **67** (1988), 97–144. MR 89k:14069 Zbl 0674.14028
- [Kollár 1996] J. Kollár, *Rational curves on algebraic varieties*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) **32**, Springer, Berlin, 1996. MR 98c:14001 Zbl 0877.14012
- [Kollár and Matsusaka 1983] J. Kollár and T. Matsusaka, “Riemann–Roch type inequalities”, *Amer. J. Math.* **105**:1 (1983), 229–252. MR 85c:14007 Zbl 0538.14006
- [Kollár and Mori 1998] J. Kollár and S. Mori, *Birational geometry of algebraic varieties*, Cambridge Tracts in Mathematics **134**, Cambridge University Press, 1998. MR 2000b:14018 Zbl 0926.14003
- [Langer 2014] A. Langer, “The Bogomolov–Miyaoaka–Yau inequality for logarithmic surfaces in positive characteristic”, preprint, 2014.
- [Lazarsfeld 2004a] R. Lazarsfeld, *Positivity in algebraic geometry, I: Classical setting: line bundles and linear series*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) **48**, Springer, Berlin, 2004. MR 2005k:14001a Zbl 1093.14501
- [Lazarsfeld 2004b] R. Lazarsfeld, *Positivity in algebraic geometry, II: Positivity for vector bundles, and multiplier ideals*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) **49**, Springer, Berlin, 2004. MR 2005k:14001b Zbl 1066.14021
- [Liedtke 2013] C. Liedtke, “Algebraic surfaces in positive characteristic”, pp. 229–292 in *Birational geometry, rational curves, and arithmetic*, edited by F. Bogomolov et al., Springer, New York, 2013. MR 3114931 Zbl 06211443
- [Matsusaka 1972] T. Matsusaka, “Polarized varieties with a given Hilbert polynomial”, *Amer. J. Math.* **94** (1972), 1027–1077. MR 49 #2729 Zbl 0256.14004
- [Mukai 2013] S. Mukai, “Counterexamples to Kodaira’s vanishing and Yau’s inequality in positive characteristics”, *Kyoto J. Math.* **53**:2 (2013), 515–532. MR 3079312 Zbl 1291.14038
- [Reider 1988] I. Reider, “Vector bundles of rank 2 and linear systems on algebraic surfaces”, *Ann. of Math.* (2) **127**:2 (1988), 309–316. MR 89e:14038 Zbl 0663.14010

- [Sakai 1990] F. Sakai, "Reider–Serrano's method on normal surfaces", pp. 301–319 in *Algebraic geometry* (L'Aquila, 1988), edited by A. J. Sommese et al., Lecture Notes in Math. **1417**, Springer, Berlin, 1990. MR 91d:14001 Zbl 0732.14004
- [Shepherd-Barron 1991a] N. I. Shepherd-Barron, "Unstable vector bundles and linear systems on surfaces in characteristic p ", *Invent. Math.* **106**:2 (1991), 243–262. MR 92h:14027 Zbl 0769.14006
- [Shepherd-Barron 1991b] N. I. Shepherd-Barron, "Geography for surfaces of general type in positive characteristic", *Invent. Math.* **106**:2 (1991), 263–274. MR 92k:14033 Zbl 0813.14025
- [Siu 2002a] Y.-T. Siu, "Extension of twisted pluricanonical sections with plurisubharmonic weight and invariance of semipositively twisted plurigenera for manifolds not necessarily of general type", pp. 223–277 in *Complex geometry* (Göttingen, 2000), edited by I. Bauer et al., Springer, Berlin, 2002. MR 2003j:32027a Zbl 1007.32010
- [Siu 2002b] Y.-T. Siu, "A new bound for the effective Matsusaka big theorem", *Houston J. Math.* **28**:2 (2002), 389–409. MR 2003i:32038 Zbl 1021.32007
- [Szpiro 1979] L. Szpiro, "Sur le théorème de rigidité de Parsin et Arakelov", pp. 169–202 in *Journées de Géométrie Algébrique de Rennes, II* (Rennes, 1978), Astérisque **64**, Soc. Math. France, Paris, 1979. MR 81f:14004 Zbl 0425.14005
- [Takagi 2007] S. Takagi, "Fujita's approximation theorem in positive characteristics", *J. Math. Kyoto Univ.* **47**:1 (2007), 179–202. MR 2008i:14014 Zbl 1136.14004
- [Tanaka 2012] H. Tanaka, "The X-method for klt surfaces in positive characteristic", preprint, 2012. arXiv 1202.2497v2
- [Tanaka 2014] H. Tanaka, "Minimal models and abundance for positive characteristic log surfaces", *Nagoya Math. J.* **216** (2014), 1–70. MR 3319838 Zbl 06427229
- [Tango 1972] H. Tango, "On the behavior of extensions of vector bundles under the Frobenius map", *Nagoya Math. J.* **48** (1972), 73–89. MR 47 #3401 Zbl 0239.14007
- [Terakawa 1999] H. Terakawa, "The d -very ampleness on a projective surface in positive characteristic", *Pacific J. Math.* **187**:1 (1999), 187–199. MR 99m:14014 Zbl 0967.14008
- [Xie 2010] Q. Xie, "Counterexamples to the Kawamata–Viehweg vanishing on ruled surfaces in positive characteristic", *J. Algebra* **324**:12 (2010), 3494–3506. MR 2012b:14047 Zbl 1211.14025

Communicated by Hélène Esnault

Received 2015-02-24

Revised 2015-04-16

Accepted 2015-05-17

dicerbo@math.columbia.edu

*Department of Mathematics, Columbia University,
New York, NY 10027, United States*

a.fanelli11@imperial.ac.uk

*Department of Mathematics, Imperial College London,
180 Queen's Gate, London SW7 2AZ, United Kingdom*

Algebra & Number Theory

msp.org/ant

EDITORS

MANAGING EDITOR

Bjorn Poonen
Massachusetts Institute of Technology
Cambridge, USA

EDITORIAL BOARD CHAIR

David Eisenbud
University of California
Berkeley, USA

BOARD OF EDITORS

Georgia Benkart	University of Wisconsin, Madison, USA	Susan Montgomery	University of Southern California, USA
Dave Benson	University of Aberdeen, Scotland	Shigefumi Mori	RIMS, Kyoto University, Japan
Richard E. Borcherds	University of California, Berkeley, USA	Raman Parimala	Emory University, USA
John H. Coates	University of Cambridge, UK	Jonathan Pila	University of Oxford, UK
J-L. Colliot-Thélène	CNRS, Université Paris-Sud, France	Anand Pillay	University of Notre Dame, USA
Brian D. Conrad	Stanford University, USA	Victor Reiner	University of Minnesota, USA
Hélène Esnault	Freie Universität Berlin, Germany	Peter Sarnak	Princeton University, USA
Hubert Flenner	Ruhr-Universität, Germany	Joseph H. Silverman	Brown University, USA
Sergey Fomin	University of Michigan, USA	Michael Singer	North Carolina State University, USA
Edward Frenkel	University of California, Berkeley, USA	Vasudevan Srinivas	Tata Inst. of Fund. Research, India
Andrew Granville	Université de Montréal, Canada	J. Toby Stafford	University of Michigan, USA
Joseph Gubeladze	San Francisco State University, USA	Ravi Vakil	Stanford University, USA
Roger Heath-Brown	Oxford University, UK	Michel van den Bergh	Hasselt University, Belgium
Craig Huneke	University of Virginia, USA	Marie-France Vignéras	Université Paris VII, France
Kiran S. Kedlaya	Univ. of California, San Diego, USA	Kei-Ichi Watanabe	Nihon University, Japan
János Kollár	Princeton University, USA	Efim Zelmanov	University of California, San Diego, USA
Yuri Manin	Northwestern University, USA	Shou-Wu Zhang	Princeton University, USA
Philippe Michel	École Polytechnique Fédérale de Lausanne		

PRODUCTION

production@msp.org
Silvio Levy, Scientific Editor

See inside back cover or msp.org/ant for submission instructions.

The subscription price for 2015 is US \$255/year for the electronic version, and \$440/year (+\$55, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscribers address should be sent to MSP.

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

ANT peer review and production are managed by EditFLOW[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2015 Mathematical Sciences Publishers

Algebra & Number Theory

Volume 9 No. 6 2015

Bivariant algebraic cobordism	1293
JOSÉ LUIS GONZÁLEZ and KALLE KARU	
Schubert decompositions for quiver Grassmannians of tree modules	1337
OLIVER LORSCHIED	
Noncommutative geometry and Painlevé equations	1363
ANDREI OKOUNKOV and ERIC RAINS	
Electrical networks and Lie theory	1401
THOMAS LAM and PAVLO PYLYAVSKYY	
The Kac–Wakimoto character formula for the general linear Lie superalgebra	1419
MICHAEL CHMUTOV, CRYSTAL HOYT and SHIFRA REIF	
Effective Matsusaka’s theorem for surfaces in characteristic p	1453
GABRIELE DI CERBO and ANDREA FANELLI	
Adams operations and Galois structure	1477
GEORGIOS PAPPAS	



1937-0652(2015)9:6;1-3